

**COMPOSITIONS AND METHODS FOR PRODUCING AND OPTIMIZING VIRAL
VECTOR PRODUCER CELLS FOR CELL AND GENE THERAPY**

CROSS-REFERENCE TO RELATED APPLICATIONS

[001] This application claims priority from U.S. Provisional Patent Application No.
5 63/158,844, filed March 9, 2021, which is herein incorporated by reference in its entirety.

SEQUENCE LISTING

[002] The instant application contains a Sequence Listing which has been submitted
electronically in ASCII format and is hereby incorporated by reference in its entirety. Said
ASCII copy, created on March 1, 2022, is named P35061WO00_SL.txt and is 611 bytes in
10 size.

FIELD

[003] The present disclosure relates to the field of production and optimization of viral
vectors for cell and gene therapy.

BACKGROUND

15 [004] The growing number of gene therapy candidates combined with rapid progression
through the clinical development has created a world-wide shortage of gene therapy vectors.
More than 500 gene therapy and 100 cell therapy candidates are in different stages of
development. Greater than 2200 clinical studies are ongoing across the globe. The strong and
proven safety profile of viral vectors (*e.g.*, lentiviral vectors) has underpinned a robust clinical
20 development pipeline. However, the clinical manufacture and use of viral vectors, especially
lentiviral vectors also comes with several limitations. For example, conventional
manufacturing methods/technologies are outdated and not scalable, provide low downstream
process yields (~20%), require significant upfront capital and on-going operational costs to
setup. Furthermore, traditionally, viral vector manufacturing is seen as unpredictable and
25 highly risky, resulting in demand greatly exceeding supply which in turn drives up prices.
There is a need to identify new methods and improvement for manufacturing viral vectors by
generating stable producer lines with high titer at high volumes.

SUMMARY

[005] In an aspect, the present disclosure provides a construct comprising a promoter sequence operably linked to a first set of one or more copies of coding sequences each encoding a viral accessory protein or a recombinant viral vector genome, wherein the construct further
5 comprises two recombinase recognition sites in opposite orientations.

[006] In another aspect, the present disclosure provides a construct comprising a promoter sequence flanked by two sets of oppositely oriented coding sequences each encoding a viral accessory protein or a recombinant viral vector genome, wherein the promoter is capable of being inverted upon a recombination event inducible by a recombinase, wherein the promoter
10 is operably linked to a first set of the two sets of oppositely oriented coding sequences prior to recombination and becomes operably linked to a second set of the two sets of oppositely oriented coding sequences post recombination. In another aspect, the construct of the present disclosure may further comprise one or more endonuclease recognition motifs.

[007] In another aspect, the present disclosure provides a method of producing a viral
15 vector producer cell with an optimized stoichiometric ratio of viral vector genome and viral accessory proteins, wherein the method comprises:

- a. introducing a construct described here into a first clonal population of cells;
- b. transiently providing a recombinase to the first clonal population; and
- c. producing a second clonal population by inverting the promoter

20 [008] In another aspect, the present disclosure provides a method comprising:

- a. obtaining a first stable viral vector producer cell line;
- b. determining a viral titer from the first stable viral vector producer cell line;
and
- c. engineering one or more elements in the stable viral vector producer cell line
25 to produce a second stable viral vector producer cell line with a stoichiometric ratio of viral vector genome and one or more viral accessory proteins different from the corresponding stoichiometric ratio in the first stable viral vector producer cell line.

BRIEF DESCRIPTION OF THE DRAWINGS

30 [009] **Figure 1** provides an illustration of the genome organization of the HIV-1 Virus. The HIV-1 genome contains 9,749 bp. In addition to the *gag*, *pol*, and *env* genes common to

all retroviruses, HIV-1 contains a regulatory gene - *rev* - that is indispensable for virus replication, and five accessory genes - *tat*, *vif*, *vpr*, *vpu* and *nef* - that, while dispensable for *in vitro* virus growth, are key for *in vivo* replication and pathogenesis. Further information about the biological functions of each of the HIV-encoded proteins is provided in Table 1.

5 [010] **Figure 2** provides an exemplary illustration of how a stable vector producer cell line can be further optimized via recombination-based genome reorganization.

[011] **Figure 3** provides further optimization of an initial vector producer cell clone to provide higher viral titers.

10 [012] **Figure 4** illustrates another exemplary approach to optimize accessory gene component ratio via subtractive subcloning. For example, both a regular copy and a codon-optimized ("co") copy of the *rev* gene are used in an initial cell clone. Subsequently, one of the two *rev* gene copies are edited via, *e.g.*, CRISPR-mediated gene editing, such that only one copy remains functional and therefore the *rev* gene expression level is reduced.

15 [013] **Figure 5** provides an illustration of an exemplary construct with exemplary recombinase sites, *loxP*, in opposing orientations, flanking a promoter. The construct may comprise one or two copies, *e.g.*, of a viral helper, or viral accessory, gene.

[014] **Figure 6** provides an illustration of an exemplary construct with exemplary recombinase sites, *loxP*, in opposing orientations, flanking an exemplary expression cassette. Depending on the orientation of the coding sequences within the expression cassette, upon
20 addition of a recombinase, the cassette may comprise one or two copies, *e.g.*, of a viral helper, or viral accessory, gene.

[015] **Figure 7** provides an illustration of an exemplary construct with exemplary recombinase sites, *loxP*, in opposing orientations, flanking an exemplary expression cassette. Depending on the orientation of the coding sequences within the expression cassette, upon
25 addition of a recombinase, the cassette may comprise one or two copies, *e.g.*, of a viral helper, or viral accessory, gene.

[016] **Figure 8** provides an illustration of an exemplary construct with exemplary recombinase sites, *loxP*, in opposing orientations, flanking a promoter and an exemplary expression cassette. Depending on the orientation of the coding sequences within the
30 expression cassette, upon addition of a recombinase, the cassette may comprise two or three copies, *e.g.*, of a viral helper, or viral accessory, gene.

[017] Figure 9 provides an exemplary illustration of an alternative method for altering copy number expression from a given construct using a combination of recombination-based and gene-editing based approaches.

DETAILED DESCRIPTION

5 [018] Unless defined otherwise, technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art. One skilled in the art will recognize many methods can be used in the practice of the present disclosure. Indeed, the present disclosure is in no way limited to the methods and materials described. Where a term is provided in the singular, the inventors also contemplate aspects of the disclosure described
10 by the plural of that term, and vice versa. Where there are discrepancies in terms and definitions used in references that are incorporated by reference, the terms used in this application shall have the definitions given herein. Other technical terms used have their ordinary meaning in the art in which they are used, as exemplified by various art-specific dictionaries, for example, "The American Heritage® Science Dictionary" (Editors of the American Heritage
15 Dictionaries, 2011, Houghton Mifflin Harcourt, Boston and New York), the "McGraw-Hill Dictionary of Scientific and Technical Terms" (6th edition, 2002, McGraw-Hill, New York), or the "Oxford Dictionary of Biology" (6th edition, 2008, Oxford University Press, Oxford and New York).

[019] Any references cited herein, including, *e.g.*, all patents and publications are
20 incorporated by reference in their entirety.

[020] When a grouping of alternatives is presented, any and all combinations of the members that make up that grouping of alternatives is specifically envisioned. For example, if an item is selected from a group consisting of A, B, C, and D, the inventors specifically envision each alternative individually (*e.g.*, A alone, B alone, etc.), as well as combinations such as A,
25 B, and D; A and C; B and C; etc. The term "and/or" when used in a list of two or more items means any one of the listed items by itself or in combination with any one or more of the other listed items. For example, the expression "A and/or B" is intended to mean either or both of A and B – *i.e.*, A alone, B alone, or A and B in combination. The expression "A, B and/or C" is intended to mean A alone, B alone, C alone, A and B in combination, A and C in combination,
30 B and C in combination, or A, B, and C in combination.

[021] When a range of numbers is provided herein, the range is understood to inclusive of the edges of the range as well as any number between the defined edges of the range. For example, "between 1 and 10" includes any number between 1 and 10, as well as the number 1 and the number 10.

5 [022] As used herein, the singular form "a," "an," and "the" include plural references unless the context clearly dictates otherwise. For example, the term "a compound" or "at least one compound" may include a plurality of compounds, including mixtures thereof.

[023] The term "about" is used herein to mean approximately, roughly, around, or in the region of. When the term "about" is used in conjunction with a numerical range, it modifies
10 that range by extending the boundaries above and below the numerical values set forth and is understood to mean plus or minus 10%. For example, "about 100" would include from 90 to 110.

[024] As used herein, the term "substantially", when used to modify a quality, generally allows certain degree of variation without that quality being lost. For example, in certain
15 aspects such degree of variation can be less than 0.1%, about 0.1%, about 0.2%, about 0.3%, about 0.4%, about 0.5%, about 0.6%, about 0.7%, about 0.8%, about 0.9%, about 1%, between 1-2%, between 2-3%, between 3-4%, between 4-5%, or greater than 5%.

[025] To avoid any doubt, used herein, terms or phrases such as "about", "at least", "at least about", "at most", "less than", "greater than", "within" or alike, when followed by a series
20 of list of numbers of percentages, such terms or phrases are deemed to modify each and every number of percentage in the series or list, regardless whether the adverb, preposition, or other modifier phrase is reproduced prior to each and every member.

[026] As used herein, a "viral vector producer cell" refers to a cell which contains all the elements necessary for production of recombinant viral vector particles (including *e.g.*,
25 retroviral delivery systems). Typically, such viral vector producer cell contains one or more expression cassettes which are capable of expressing viral structural proteins (such as Gag, Pol and Env). A "stable viral vector producer cell" refers to a viral vector producer cell that contains in its nuclear genome, maintains episomally, or combination thereof, all the elements necessary for production of recombinant viral vector particles. A "stable viral vector producer
30 cell line" refers a permanently established cell culture of stable viral vector producer cells that will proliferate indefinitely given appropriate fresh medium and space.

[027] As used herein, a “recombinant viral vector” is an enveloped virion particle that contains an expressible polynucleotide sequence, and which is capable of penetrating a target host cell, thereby carrying the expressible sequence into the cell. In an aspect, an expressible polynucleotide sequence comprises or encodes a gene of interest (GOI). The enveloped particle is preferably pseudotyped with an engineered or native viral envelope or capsid protein from another viral species, including lentiviruses or non-lentiviruses, which alters the host range and infectivity of a native virus.

[028] As used herein, a “viral vector genome construct” is a construct which contains polynucleotide sequences which are packaged into a transducing recombinant viral vector. In an aspect, a viral vector genome construct, when comprising 5' LTR and 3' LTR and packaged with a functional integrase enzyme, can be used for the production of recombinant viral vectors that are capable of integrating into the host genome. In another aspect, a viral vector genome construct produces a recombinant viral vector comprising 5' LTR and 3' LTR and not capable of integrating into a host genome due to the lack of a functional integrase enzyme, which is also known as integrase-defective lentiviral vectors (IDLV).

[029] As used herein, a “viral accessory construct” refers to a construct, plasmid or isolated nucleic acid molecule containing or encoding one or more elements that are useful for producing a functional recombinant viral vector in a compatible host cell, and packaging into it an expressible heterologous sequence.

[030] As used herein, a “viral accessory protein” refers to a protein that is useful or necessary for producing a functional recombinant viral vector in a compatible host cell, and packaging into the viral vector an expressible heterologous sequence.

[031] As used herein, a “viral vector construct” refers to either a viral vector genome construct or a viral accessory construct.

[032] As used herein, the term “operably linked” describes a spatial and mechanistic relationship of two or more pieces of DNA such that one piece is capable of effecting an intended genetic outcome of another piece. For example, “operably linked” can denote a relationship between a regulatory region (typically a promoter element but may include an enhancer element) and the coding region of a gene, whereby the transcription of the coding region is under the control of the regulatory region.

[033] As used herein, the term “flanking” refers to an arrangement of nucleic acids that are adjacent to, or surround (but are not immediately adjacent to), a given region.

[034] As used herein, a “concatemer” is defined as a continuous DNA molecule that contains multiple copies of the same or substantially same DNA sequence linked in series. In an aspect, a concatemer may also contain one or more selection genes.

5 [035] As used herein, the term “trans” refers to mechanisms acting from different molecules.

[036] As used herein, the term “promoter” includes nucleic acid regions ranging in complexity and size from minimal promoters to promoters including upstream elements and enhancers.

10 [037] As used herein, the term “transduction” refers to the delivery of a nucleic acid segment using a viral vector by means of viral vector.

[038] As used herein, the term “transfection” refers to the introduction of foreign DNA into eukaryotic cells.

15 [039] Without being bound to any theory, quality and quantity of infectious vector particles derived from a viral vector producer cell line is directly affected by the stoichiometric ratio of the lentiviral vector genomic RNA to the trans expressed accessory proteins. For any given lentiviral vector genome, the optimal ratio is not known a priori, and must be determined empirically through trial and error. As this biological fact is not often appreciated, the construction of stable cell lines has historically been accomplished by the addition of accessory genes one at a time in a serial fashion. This has assured progeny clones that had and expressed
20 the accessory protein but limited the ability of the ultimate cell line to produce vector for lentiviral vector genomes with suboptimal ratios. The solution offered to this problem is to add all of the accessory elements at once in such a manner as to encourage multiple introductions of each of the elements. This not only speeds the development time of any given producer clone by collapsing the accessory gene introductions from multiple rounds of subcloning to a single
25 round, it also allows for the generation of a diverse set of clones, each with different ratios, such that when the clones are screened the likelihood that we can find a clone that produces vector of the desired quality and quantity is increased without having to know a priori what that ratio would be.

30 [040] In an aspect, this disclosure provides a method to produce a viral vector producer cell with an optimized stoichiometric ratio of viral vector genome and viral accessory proteins. In an aspect, this application provides a stable lentiviral vector producer cell line.

[041] In an aspect, a vector producer cell line is produced from a parental cell line derived from an immortalized human cell line. In another aspect, a vector producer cell line grows in defined media either with or without human/animal derived serum. In another aspect, a vector producer cell line grows in an adherent or suspension adapted manner.

5 **Recombinant viral vectors**

[042] This disclosure relates to the manufacturing and/or production of recombinant viral vectors (also known as recombinant viral particles). The present disclosure relates to recombinant viral vectors, and constructs for their manufacture, which can be utilized to introduce expressible polynucleotide sequences of interest into host cells.

10 [043] In an aspect, a viral vector producer cell disclosed herein comprises a retroviral production system, wherein the viral vector is derived from a retrovirus. Retroviruses comprise a family of enveloped viruses with a 7-12kb single-stranded positive sense RNA genome. The retrovirus family includes five groups of oncogenic retroviruses, lentiviruses and spumaviruses.

15 [044] Retroviral vector production systems typically involve separation of viral genome from viral packaging functions. Viral accessory proteins or viral accessory protein domains may be introduced via separate expression cassettes, or in trans. In an aspect, a viral accessory construct encodes or provides one or more viral accessory proteins involved in viral packaging.

[045] In an aspect, the present disclosure relates to lentiviral vectors, and constructs for
20 their manufacture, which can be utilized to introduce expressible polynucleotide sequences of interest into host cells. In an aspect, a lentiviral vector is an enveloped virion particle that contains an expressible polynucleotide sequence, and which is capable of penetrating a target host cell, thereby carrying the expressible sequence into the cell. The enveloped particle is preferably pseudotyped with an engineered or native viral envelope protein from another viral
25 species, including non-lentiviruses, which alters the host range and infectivity of the native lentivirus.

[046] Viral vectors described here can be utilized in a wide range of applications, including, *e.g.*, for protein production (including vaccine production), for gene therapy (including gene replacement, gene editing, and synthetic biology), to deliver therapeutic
30 polypeptides, to deliver siRNA, ribozymes, anti-sense, and other functional polynucleotides, etc. Such transduction vectors have the ability to carry single or dual genes, and to include inhibitory sequences (*e.g.*, RNAi or antisense). In certain aspects, the transduction vector also

carries a nucleic acid which comprises a modified 3' LTR having reduced, but not absent, transcriptional activity.

[047] Lentivirus is a group of retroviruses characterized for a long incubation period. They are classified into five serogroups according to the vertebrate hosts they infect: bovine, equine, feline, ovine/caprine and primate. Some examples of lentiviruses are human (HIV), simian (SIV) and feline (FIV) immunodeficiency viruses.

[048] Lentiviruses can deliver large amounts of genetic information into the DNA of host cells and can integrate in both dividing and non-dividing cells. The viral genome is passed onto daughter cells during division, making it one of the most efficient gene delivery vectors.

10 [049] The structure of HIV is different from that of other retroviruses. HIV is roughly spherical with a diameter of ~120 nm. HIV is composed of two copies of positive ssRNA that code for nine genes enclosed by a conical capsid containing 2,000 copies of the p24 protein. The ssRNA is tightly bound to nucleocapsid proteins, p7, and enzymes needed for the development of the virion: Reverse transcriptase (RT), Protease (PR), Ribonuclease and
15 Integrase (IN). A matrix composed of p17 surrounds the capsid ensuring the integrity of the virion. This, in turn, is surrounded by an envelope composed of two layers of phospholipids taken from the membrane of a human cell when a newly formed virus particle buds from the cell. Embedded in the viral envelope are proteins from the host cell and about 70 copies of a complex HIV protein, known as Env, that protrudes through the surface of the virus particle.
20 Env consists of a cap made of three gp120 molecules, and a stem consisting of three gp41 molecules that anchor the structure into the viral envelope. The glycoprotein complex enables the virus to attach to and fuse with target cells to initiate the infectious cycle. Further information about the biological functions of each of the HIV-encoded proteins is provided in Table 1.

Table 1: Summary of the biological functions of HIV-encoded proteins.

	Gene		Precursor proteins → products
Essential Genes for Vectorized Lentivirus	<i>gag</i>	Group-specific antigen	<i>gag</i> → MA, CA, SP1, NC, SP2, P6
	<i>pol</i>	Polymerase	<i>pol</i> → RT, RNase H, IN, PR
	<i>env</i>	Envelope	gp160 → gp120, gp41
	<i>rev</i>	Regulator of expression of virion proteins	Important for major viral protein synthesis and essential for viral replication
Additional Genes found in Wild-Type HIV	<i>tat</i>	HIV transactivator	Positive transcription regulator
	<i>vif</i>	Viral infectivity	Required for infectivity in some cell types
	<i>vpr</i>	Virus protein R	Nuclear import of pre-integration complex and host cell cycle arrest
	<i>vpu</i>	Virus protein U	Proteasomal degradation of CD44 and virion release from infected cells
	<i>nef</i>	Negative factor	Roles in apoptosis and virus infectivity

[050] In an aspect, a viral vector producer cell disclosed herein comprises a lentiviral vector production system, wherein the viral vector is derived from a lentivirus. A lentivirus is a group of retroviruses that causes slow, gradual disease. A lentiviral vector particle produced by the lentiviral vector production system disclosed herein will be capable of transducing slowly-dividing cells, whereas standard retroviruses (gamma retroviruses) can infect only mitotically active cells. “Slowly dividing” cell types may divide approximately once every three to four days.

[051] In the production of lentiviral vectors, multiple plasmids are used, one encoding envelope proteins (*env* plasmid), one or more plasmids encoding viral accessory proteins, and one plasmid comprising a gene of interest expression cassette between a lentiviral 3'-LTR and a lentiviral 5'-LTR to facilitate integration of the encoded gene(s) of interest into the host genome.

[052] In an aspect, a viral vector may be a hybrid viral vector. The term “hybrid” as used herein refers to a vector, or nucleic acid component of a vector, that contains both lentiviral sequences and non-lentiviral sequences.

[053] In an aspect, a viral vector producer cell disclosed herein comprises a herpesvirus vector production system, wherein the viral vector is derived from a herpesvirus.

[054] In an aspect, a viral vector producer cell disclosed herein comprises an adenoviral vector production system, wherein the viral vector is derived from an adenovirus. Adenovirus is a nonenveloped virus with a 36-kilobase double-stranded DNA genome. Adenovirus is an attractive gene delivery vehicle candidate for its ability to grow as a high-titer recombinant

virus, large transgene capacity, and efficient transduction of dividing and non-dividing cells. More than 50 human and nonhuman serotypes of adenovirus have been found to mediate gene delivery to a wide range of tissues.

5 [055] In an aspect, a viral vector producer cell disclosed herein comprises an adeno-associated viral vector production system, wherein the viral vector is derived from an adeno-associated virus. Adeno-associated virus (AAV) is a nonenveloped virus with a 4.7kb single-stranded DNA genome. More than 100 serotypes of AAV have been isolated from human and nonhuman tissues.

10 [056] In a further aspect, a recombinant viral vector disclosed herein is derived from a virus comprising a mosaic genome structure. In a further aspect, recombinant viral vectors disclosed herein are target-specific. In a further aspect, target-specific viral vectors are receptor-targeted. In a further aspect, target-specific viral vectors comprise recombinant antibody molecules. Methods to produce target-specific viral vectors are known in the art. In a further aspect, a recombinant vector is derived from a partially or fully synthetic nucleic acid
15 sequence.

[057] Recombinant viral vectors disclosed herein may have one or more selectable, traceable or otherwise detectable marker elements. In an aspect, a selectable element is a reporter gene. In a further aspect, a selectable element is an epitope tag. In a further aspect, a
20 viral vector may contain both a reporter gene and an epitope tag. In an aspect, an epitope tag may be selected or detected by methods known in the art, including but not limited to chromatography, enzyme assays, fluorescence assays, and immunodetection assays. In an aspect, immunodetection assays may include, but are not limited to immunoblotting, immunofluorescence, immunocytochemistry, and enzyme-linked immunosorbent assay (ELISA).

25 [058] In a further aspect, a reporter gene may be detected by methods to detect absorbance. Methods to detect absorbance are known in the art. In an aspect, a reporter gene may be detected by methods to detect fluorescence. Methods to detect fluorescence are known in the art. In a further aspect, a reporter gene may be detected by methods to detect luminescence. Methods to detect luminescence are known in the art. In an aspect, a selectable
30 marker gene is an antibiotic resistance gene. In a further aspect, an antibiotic gene encodes neomycin resistance. In a further aspect, an antibiotic gene encodes puromycin resistance.

[059] In a further aspect, traceable marker genes may include genes encoding fluorescent proteins. Methods to select fluorescent proteins with different chromophores are known in the art. In a further aspect, fluorescent proteins may be Green fluorescent protein (GFP) or variants thereof, including, but not limited to Ultramarine, Blue, and Cyan fluorescent proteins. In a further aspect, a variant of a fluorescent protein may be an optimized variant. Methods to optimize traits of fluorescent proteins are known in the art and include but are not limited to methods to improve chromophore maturation, folding kinetics, and thermostability, among other traits.

[060] In an aspect, a recombinant viral vector may be self-inactivating. The terms “self-inactivating” refer to a vector which is modified, such that the modification reduces the ability of the vector to mobilize once it has integrated into the genome of a target or host cell. For example, the modification may include deletions in the 3' long terminal repeat (LTR) region. SIN vectors possess safety advantages over non-SIN vectors for gene delivery applications.

[061] In another aspect, a recombinant viral vector produced here is a self-inactivating lentiviral vector (SIN vectors). In this kind of SIN vectors, the deletion of lentiviral enhancer and promoter sequences from the 3' LTR results in the generation of vectors which, on infection of target cells, are incapable of transcribing vector-length RNA. Because of this modification, integrated SIN vectors are incapable of further replication thus reducing the likelihood of generating replication-competent viruses as well as the danger of inadvertently influencing transcription activity of nearby endogenous promoters.

[062] In another aspect, a recombinant viral vector produced here is a conditional SIN vector. For example, in an exemplary conditional SIN vector, the 3' LTR U3 transcription regulatory elements can be replaced with an inducible promoter (*e.g.*, Tet- responsive element).

Viral vector genome construct

[063] In an aspect, disclosed herein is a viral vector genome construct encoding a recombinant viral vector genome. As used herein, a “recombinant viral vector genome” refers to a viral genome sequence engineered to become replication incompetent while harboring one or more additional sequence of interest that is typically not present in a natural form of the corresponding virus. In an aspect, a viral vector genome construct encodes a gene of interest. In an aspect, a gene of interest is operably linked to a promoter.

[064] In an aspect, a gene of interest may be a candidate gene which is of known or potential significance in the pathophysiology of a disease. In a further aspect, a gene of interest

may have a known or potential therapeutic or diagnostic application. In an aspect, a gene of interest may comprise a coding region. In a further aspect, a gene of interest may comprise a partial coding region. A gene of interest can be obtained for insertion into the viral vectors disclosed herein through a variety of techniques known in the art.

5 [065] In a further aspect, a viral vector genome construct disclosed herein comprises one or more selectable or detectable reporter element(s). In an aspect, a selectable or detectable element is a reporter. In a further aspect, a selectable or detectable reporter element is an epitope tag. In an aspect, a selectable or detectable reporter element may be selected or detected by methods known in the art including, but not limited to luminescence, absorbance,
10 fluorescence, antibiotics, antigen-antibody interactions, or a combination thereof.

[066] In an aspect, a viral vector genome construct disclosed herein comprises one or more elements selected from the group consisting of a promoter, 5' and 3' long terminal repeats, a packaging signal, a central polypurine tract, and a polyadenylation (p(A)) sequence. In another aspect, a viral vector genome construct disclosed herein comprises all the elements
15 in the preceding sentence. In a further aspect, a long terminal repeat is a self-inactivating long terminal repeat.

[067] In an aspect, a viral vector genome construct disclosed herein can be used to produce a viral like particle. In a further aspect, a viral vector genome construct disclosed herein does not comprise a promoter, a 5' long terminal repeat, a 3' long terminal repeat, a
20 packaging signal, a central polypurine tract, or a polyadenylation sequence.

[068] A viral vector genome construct of the disclosure disclosed herein may be in the form of a concatemer. In an aspect, a concatemer may contain one or more transcription factors. In a further aspect, a transcription factor may be a ligand-responsive transcription factor. In an aspect, a concatemer is made and used as described in Throm *et al.* Blood,
25 2009;113(21):5104-10. For example, a stable viral producer cell line can contain fully SIN lentiviral genome and viral accessory constructs stably integrated into the genome by concatemeric array transfection. Such array can be obtained through the ligation of DNA fragments encoding the SIN lentiviral vector genome, with drug resistance and/or other selection/reporter cassettes included into the array.

30 **Viral accessory genes/proteins/constructs**

[069] In an aspect, a viral accessory construct encodes one or more accessory proteins including for example, structural proteins (*e.g.*, the Gag precursor), processing proteins (*e.g.*,

the Pol precursor), and other proteins such as proteases, envelope protein. In another aspect, a viral accessory vector comprises sequences that provide the expression and regulatory signals needed to manufacture one or more accessory proteins in host cells and assemble functional viral particles. In one aspect, coding sequences for an Env, a Rev, and a Gag-Pol precursor are
5 on the same plasmid or viral accessory construct. In another aspect, coding sequences for an Env, a Rev, and a Gag-Pol precursor are placed on separate plasmids or viral accessory constructs. In a further aspect, separate plasmids or viral accessory constructs are used for each coding sequence of the Gag, Pol, Rev, and Envelope proteins. In an aspect, a viral accessory construct may encode one or more structural and/or regulatory viral proteins, or functional
10 fragments or domains thereof, selected from the group consisting of Group-specific antigen (Gag), RNA-dependent DNA polymerase (Pol), Regulator of expression of viral protein (Rev), Envelope (Env), Transactivator (Tat), Negative regulatory factor (Nef), Viral protein R (Vpr), Virus infectivity factor (Vif), Viral protein U (Vpu), and Viral protein X (Vpx). In another aspect, a functional fragment or domain can comprise one or more proteins selected from the
15 group consisting of MA (Matrix [p17]), CA (Capsid [p24]), NC (Nucleocapsid [p9]), p6, Protease (p10), RT (p50), RNase H (p15), and Integrase (p31). In an aspect, coding sequences of one or more viral accessory proteins are operably linked. In an aspect, coding sequences of one or more viral accessory proteins are present on separate viral accessory constructs.

[070] In an aspect, a viral accessory construct used here is for producing a recombinant
20 lentiviral vector. In an aspect, a viral accessory construct used in the present disclosure can comprise one or more of the following elements, separately or collectively, in any suitable order or position, *e.g.*, a) a heterologous promoter operably linked to a polynucleotide sequence coding for lentivirus Gag and Pol (*e.g.*, a lentivirus Gag-Pol precursor); and b) a heterologous promoter operably linked to an *env* coding sequence.

25 [071] Any suitable lentiviral 5' LTR can be utilized in accordance with the present disclosure, including an LTR obtained from any lentivirus species, sub-species, strain or clade. This includes primate and non-primate lentiviruses. Specific examples of species, etc., include, but are not limited to, *e.g.*, human immunodeficiency virus (HIV)-I (including subspecies, clades, or strains, such as A, B, C, D, E, F, and G, R5 and R5X4 viruses, etc.), HIV-2 (including
30 subspecies, clades, or strains, such as, R5 and R5X4 viruses, etc.), simian immunodeficiency virus (SIV), simian-human immunodeficiency virus (SHIV), feline immunodeficiency virus (FIV), bovine immunodeficiency virus (BIV), caprine-arthritis-encephalitis virus, Jembrana disease virus, ovine lentivirus, visna/maedi virus, and equine infectious anemia virus.

[072] Genomic reference sequences for such viruses are widely available, *e.g.*, HIV-I (NC_001802), HIV-2 (NC_001722), SIV (NC_001549), SIV-2 (NC_004455), caprine arthritis-encephalitis virus (NC_001463), feline immunodeficiency virus (NC_001482), Jembrana disease virus (NC_001654), ovine lentivirus (NC_001511), visna/maedi virus (NC_001452), equine infectious anemia virus (NC_001450), and bovine immunodeficiency virus (NC_001413).

[073] In an aspect, a lentiviral 5' LTR used here comprises signals utilized in gene expression, including enhancer, promoter, transcription initiation (capping), transcription terminator, and polyadenylation. They are typically described as having U3, R, and U5 regions. The U3 region of the LTR contains enhancer, promoter and transcriptional regulatory signals, including RBEIII, NF-kB, SpI, AP-I and/or GABP motifs. The TATA box is located about 25 base pairs from the beginning of the R sequence, depending on the species and strain from which the 5' LTR was obtained. A completely intact 5' LTR can be utilized, or a modified copy can be utilized. Modifications preferably involve the R region, where a TAR sequence is substituted (see below), and/or deletion of all or part of a U5 region. The modified 5' LTR preferably comprises promoter and enhancer activity, *e.g.*, preferably native U3, modified R with a substituted TAR, and native U5.

[074] In an aspect, a heterologous or non-viral promoter can be operably linked to a polynucleotide sequence coding for lentivirus Gag and Pol. By the term "operably linked," it is meant that a promoter is positioned in such a way that it can drive transcription of the recited coding sequences. In an aspect, *gag* and *pol* coding sequences are organized as the *gag-pol* precursor in native lentivirus. The *gag* sequence codes for a 55-kD Gag precursor protein, also called p55. The p55 is cleaved by the virally encoded Protease 4 (a product of the *pol* gene) during the process of maturation into four smaller proteins designated MA (matrix [p17]), CA (capsid [p24]), NC (nucleocapsid [p9]), and p6. The Pol precursor protein is cleaved away from Gag by a virally encoded protease, and further digested to separate the Protease (p10), RT (p50), RNase H (p15), and Integrase (p31) activities.

[075] In an aspect, one or more splice donor (SD) sites can be present in a viral vector genome construct or a viral accessory construct. A splice donor site is typically present between the 3' end of the 5'LTR and the packaging sequence. A downstream splice acceptor (SA) can also be present, *e.g.*, at the 3' end of the *pol* sequences. The SD site can be present in multiple copies at any effective locations in the vector. The SD can have a native or mutated copy of a lentiviral sequence.

[076] Native *gag-pol* sequences can be utilized in a viral accessory construct, or modifications can be made. These modifications can include, chimeric *gag-pol*, where the *gag* and *pol* sequences are obtained from different viruses (*e.g.*, different species, subspecies, strains, clades, etc.), and/or where the sequences have been modified to improve transcription and/or translation, and/or reduce recombination. In other aspects of the present disclosure, the sequences coding for the Gag and Pol precursors (or parts thereof, *e.g.*, one or more of MA (matrix [p17]), CA (capsid [p24]), NC (nucleocapsid [p9]), p6, protease (p10), RT (p50), RNase H (p15), and integrase (p31)) can be separated and placed on different vector constructs, where each sequence has its own expression signals.

10 [077] The RNA genome of HIV-I contains an approximately 120 nucleotide *psi*-packaging signal that is recognized by the nucleocapsid (NC) domain of the Gag polyprotein during virus assembly. The critical portions of the packaging signal are between the major splice donor (SD) site and the *gag* initiation codon of the HIV provirus, about distal to the U5 region of the 5' LTR. In an aspect, a packaging signal is functionally absent from the accessory
15 construct to avoid packaging of functionally active Gag-Pol precursor into the viral transduction vector. See, *e.g.*, U.S. Pat. No. 5,981,276 (Sodroski *et al.*), which describes vectors containing *gag*, but which lack the packaging signal.

[078] Additional promoter and enhancer sequences can be placed upstream of the 5' LTR in order to increase, improve, enhance, etc., transcription of the *gag-pol* precursor. Examples
20 of useful promoters include, mammalian promoters (*e.g.*, constitutive, inducible, tissue-specific), CMV, RSV, LTR from other lentiviral species, and other promoters as mentioned above and below. In addition, the construct can further comprise transcription termination signals, such as a polyA signal that is effective to terminate transcription driven by the promoter sequence. Any suitable polyA sequence can be utilized, *e.g.*, sequences from beta globin
25 (mammalian, human, rabbit, etc.), thymidine kinase, growth hormone, SV40, and many others.

[079] In an aspect, *gag-pol* sequences are placed in opposite transcriptional orientations from the envelope sequences in a single viral accessory construct. By the latter, it is meant that the direction of transcription is opposite or reversed. This can be achieved by placing the corresponding promoters in opposite directions (*i.e.*, facing each other) or using bi-directional
30 promoters (*e.g.*, Trinklein *et al.*, Genome Re-search 14:62-66, 2004). This arrangement can be utilized for safety purposes, *e.g.*, to reduce the risk of recombination and/or the production of functional recombinant HIV genomes. Safety is increased with such vectors as there is no possibility that transcriptional read-through would result in an RNA that contains both

functional *gag-pol* and *env* sequences. Transcriptional interference can be prevented by utilizing strong polyadenylation sequences that terminate transcription. Examples of strong transcription termination sequences are known in the art, including, *e.g.*, rabbit beta-globin polyadenylation signal (Lanoix and Acheson, EMBO J. 1988 Aug;7(8):2515-22), See, also

5 Plant *et al.*, Molecular and Cellular Biology, April 2005, p. 3276-3285, Vol. 25, No. 8. In addition, other elements can be inserted between the *gag-pol* and *env* coding sequences to facilitate transcriptional termination, such as a cis-acting ribozyme, or an RNAi sequence which are targeted to any putative read-through sequence. Similarly, instability sequences, termination sequences, and pause sites can be placed between the coding sequences.

10 [080] In an aspect, a viral accessory construct may encode structural viral proteins. In an aspect, a viral accessory construct may encode regulatory viral proteins. In an aspect, a viral accessory construct may encode both structural and regulatory viral proteins.

[081] In an aspect, a viral accessory construct may encode structural and/or regulatory viral proteins that include, but are not limited to Group-specific antigen (Gag), RNA-dependent

15 DNA polymerase (Pol), Regulator of expression of viral protein (Rev), Envelope (Env), Transactivator (Tat), Negative regulatory factor (Nef), Viral protein R (Vpr), Virus infectivity factor (Vif), Viral protein U (Vpu), and Viral protein X (Vpx).

[082] *Gag* encodes structural proteins such as Matrix protein (MA), Capsid protein (CA), and Nucleocapsid protein (NC). *Pol* encodes proteins such as Protease (PR), Reverse

20 transcriptase (RT), and Integrase (IN). *Env* encodes surface and transmembrane units of envelope protein.

[083] In an aspect, encoded viral accessory proteins are fusion proteins. In an aspect, encoded viral accessory proteins are partial viral accessory proteins, such as protein domains. In an aspect, viral accessory protein domains may include, but are not limited to Capsid protein

25 (CA), Matrix protein (MA), Nucleocapsid protein (NC), p6, Transcription factor specificity protein 1 (SP1), Reverse transcriptase (RT), Integrase (IN), Protease (PR), and Deoxyuridine triphosphatase (dUTPase or DU). In a further aspect, encoded viral accessory proteins include at least one full length protein or at least one protein domain.

[084] In an aspect, a viral construct can further comprise an RRE element, including an

30 RRE element which is obtained from a different lentiviral species than the 5' LTR or *gag* and *pol* sequences. The RRE element is the binding site for the Rev polypeptide which is a 13-kD sequence-specific RNA binding protein. Constructs which contain the RRE sequence depend

on the Rev polypeptide for efficient expression. Rev binds to a 240-base region of complex RNA secondary structure of the Rev response element ("RRE") that is located within the second intron of HIV, distal to the *pol* and *gag* coding sequences. The binding of Rev to RRE facilitates the export of unspliced and incompletely spliced viral RNAs from the nucleus to the cytoplasm, thereby regulating the expression of HIV proteins. The RRE element can be in any suitable position on the construct, preferably following the Gag-Pol precursor in its approximate native position. Similarly, for the Tat polypeptide, any suitable Rev polypeptide can be utilized as long as it retains the ability to bind to RRE.

Viral capsids/envelopes

[085] Virus particles contain a viral genome packaged in a protein coat called the capsid. For some viruses, the capsid is surrounded by lipid bilayer that contains viral proteins, usually including the proteins that enable the virus to bind to the host cells. This lipid and protein structure is called the virus envelope and is derived from the host cell membranes. The capsid and envelope play many roles in viral infection, including virus attachment to cells, entry into cells, release of the capsid contents into the cells, and packaging of newly formed viral particles. The capsid and envelope are also responsible for transfer of the viral genetic material from one cell to another. These structures also determine the stability characteristics of the virus particle, such as resistance to chemical or physical inactivation.

[086] In an aspect, a stable viral vector producer cell line produces an envelope protein. In an aspect, envelope protein(s) employed in this cell line system use either the native HIV *env* gene (wild-type or codon optimized ("co")) or generate a pseudotyped particle using a biocompatible substitute including, but not limited to, amphotropic envelope protein, vesicular stomatitis vector (Indiana or other strain), measles or bioengineered chimeric measles envelope proteins, gibbon ape leukemia virus, or feline leukemia virus or bioengineered FLV chimeras.

[087] In an aspect, viral vectors disclosed herein contain one or more capsid proteins. In an aspect, capsid proteins may be heterologous. Capsid proteins may be modified in order to alter vector biodistribution. In an aspect, capsid proteins may be genetically modified. In a further aspect, capsid proteins may be chemically modified. Strategies to genetically and chemically modify capsid proteins are known in the art.

[088] In an aspect, viral vectors disclosed herein may have sequences encoding for one or more envelope ("Env") proteins. Viral vector tropism is determined by the ability of the viral envelope protein to interact with molecules (proteins, lipids, or sugars) on the host cell.

[089] In an aspect, a viral accessory construct can comprise an envelope module or expression cassette comprising a heterologous promoter operably linked to an envelope coding sequence. The Env polypeptide is displayed on the viral surface and is involved in the recognition and infection of host cells by a virus particle. The host range and specificity can be changed by modifying or substituting the envelope polypeptide, *e.g.*, with an envelope expressed by a different (heterologous) viral species or which has otherwise been modified. This is called pseudotyping. See, *e.g.*, Yee *et al.*, Proc. Natl. Acad. Sci. USA 91: 9564-9568, 1994. Vesicular stomatitis virus (VSV) protein G (VSV G) has been used extensively because of its broad species and tissue tropism and its ability to confer physical stability and high infectivity to vector particles. See, *e.g.*, Yee *et al.*, Methods Cell Biol., (1994) 43:99-112.

[090] An envelope polypeptide can be utilized without limitation, including, *e.g.*, HIV gpl20 (including native and modified forms), Moloney murine leukemia virus (MoMuLV or MMLV), Harvey murine sarcoma virus (HaMuSV or HSV), murine mammary tumor virus (MuMTV or MMTV), gibbon ape leukemia virus (GALV), Rous sarcoma virus (RSV), hepatitis viruses, influenza viruses (VSV-G), Mokola virus, rabies, filovirus (*e.g.*, Ebola and Marburg, such as GP1/GP2 envelope, including NP_066246 and Q05320), amphotropic, alphavirus, etc. Other examples include, *e.g.*, envelope proteins from Togaviridae, Rhabdoviridae, Retroviridae, Poxviridae, Paramyxoviridae, and other enveloped virus families. Other example envelopes are from viruses listed in the following database located on the worldwide web at ncbi.nlm.nih.gov/genome/viruses.

[091] Furthermore, a viral envelope protein can be modified or engineered to contain polypeptide sequences that allow the transduction vector to target and infect host cells outside its normal range or more specifically limit transduction to a cell or tissue type. For example, the envelope protein can be joined in-frame with targeting sequences, such as receptor ligands, antibodies (using an antigen-binding portion of an antibody or a recombinant antibody-type molecule, such as a single chain antibody), and polypeptide moieties or modifications thereof (*e.g.*, where a glycosylation site is present in the targeting sequence) that, when displayed on the transduction vector coat, facilitate directed delivery of the virion particle to a target cell of interest. Furthermore, envelope proteins can further comprise sequences that modulate cell function. Modulating cell function with a transducing vector may increase or decrease transduction efficiency for certain cell types in a mixed population of cells. For example, stem cells could be transduced more specifically with envelope sequences containing ligands or binding partners that bind specifically to stem cells, rather than other cell types that are found

in the blood or bone marrow. Such ligands are known in the art. Non-limiting examples are stem cell factor (SCF) and Flt-3 ligand. Other examples, include, *e.g.*, antibodies (*e.g.*, single-chain antibodies that are specific for a cell-type), and essentially any antigen (including receptors) that is specific for such tissues as lung, liver, pancreas, heart, endothelial, smooth muscle, breast, prostate, epithelial, etc.

[092] Any heterologous promoter can be utilized to drive expression of a viral envelope coding sequence (or another viral accessory protein) when operably linked to it. Examples include, *e.g.*, CMV, EF1 alpha, EF1 alpha-HTLV-1 hybrid promoter, ferritin promoters, inducible promoters, constitutive promoters, and other promoters mentioned herein, etc.

10 [093] In an aspect, encoded envelope proteins are endogenous. In a further aspect, encoded envelope proteins are heterologous. Heterologous envelope proteins of the viral vectors disclosed herein may be generated using any envelope protein that is biocompatible. Biocompatibility can be determined using methods known in the art.

15 [094] In an aspect, env may be derived from human immunodeficiency virus (HIV). In an aspect, a sequence encoding an HIV-derived envelope gene may be wild-type. In a further aspect, a sequence encoding an HIV-derived envelope gene may be codon-optimized.

[095] Env may also be generated as a pseudotyped particle. Pseudotyping enables the engineering of viral vector particles with different target cell specificities, to expand and/or to alter the host range of the native virus from which the envelope protein was derived.

20 [096] In an aspect, the viral vectors disclosed herein may be amphotropic pseudotyped viral vectors. In an aspect, the viral vectors disclosed herein may be ecotropic pseudotyped viral vectors. In an aspect, the viral vectors disclosed herein may be pantropic pseudotyped viral vectors. Envelope protein sequences encoded by the viral vectors disclosed herein may be derived from any species of the genera Vesiculovirus, Gammaretrovirus, or Morbillivirus.

25 [097] In an aspect, envelope proteins may be derived from a species of the Vesiculovirus genus including, but not limited to, vesicular stomatitis New Jersey virus (VSV-NJ), and vesicular stomatitis Indiana virus (VSV-IN). In a further aspect, envelope proteins may be derived from any vesicular stomatitis virus serotype. In a further aspect, envelope proteins may be truncated proteins. In a further aspect, envelope proteins may be bioengineered
30 chimeric vesiculovirus proteins.

[098] In an aspect, envelope proteins may be derived from a species of the Gammaretrovirus genus, including, but not limited to gibbon ape leukemia virus (GALV) and

feline leukemia virus (FLV). In a further aspect, envelope proteins may be bioengineered chimeric gammaretrovirus proteins, including GALV chimeras and FLV chimeras. A “chimera” as defined herein refers to a biological entity, such as a virus, that is composed of two or more genetic fragments of distinct origin or of distinct composition.

- 5 [0099] In an aspect, envelope proteins may be derived from a species of the Morbillivirus genus including, but not limited to, measles virus. In a further aspect, envelope proteins may be bioengineered chimeric morbillivirus proteins, including bioengineered chimeric measles envelope proteins. Methods of bioengineering chimeric envelope proteins are known in the art.

10 **Optional Tat**

- [0100] In an aspect, a stable viral vector producer cell line comprises or produces a Tat protein. In another aspect, a stable viral vector producer cell line does not produce a Tat protein. In the absence of a Tat protein, a lentiviral genome vector is modified such that the HIV promoter in the 5' LTR is replaced with a heterologous enhancer/promoter to ensure
15 transcription. In an aspect, such promoter could be either viral (like CMV) or cellular (like EF1- α).

- [0101] In another aspect, a viral accessory construct can further comprise a TAR element that is obtained from a different lentiviral species, group, sub-species, sub-group, strain, or clade than the 5' LTR and/or the *gag* and *pol* sequences that are present in it, i.e., it is
20 heterologous to other lentiviral elements present in the construct. The TAR is preferably present in the 5' LTR in its normal location, e.g., between the U3 and U5 elements of the LTR, e.g., where the native R is replaced by R' of a heterologous lentiviral species.

- [0102] The TAR element is a trans-activating response region or response element that is located in the 5'LTR (e.g., R) of the viral DNA and at the 5' terminus of the corresponding
25 RNA. When present in the lentiviral RNA, the transcriptional transactivator, Tat, binds to it, activating transcription from the HIV LTR many-fold. Tat is an RNA binding protein that binds to a short-stem loop structure formed by the TAR element.

- [0103] When a heterologous TAR element is utilized, the 5' LTR can be modified routinely by substituting its native TAR for a TAR sequence from another species. Examples of TAR
30 regions are widely known. See, e.g., De Areliano *et al.*, AIDS Res. Human Retro., 21 -949-954, 2005. Such a modified lentiviral 5' LTR can comprise intact U3 and U5 regions, such that the LTR is completely functional. The TAR region or the entire R can be substituted.

[0104] As indicated above, the Tat polypeptide binds to the TAR sequence. The coding sequence for Tat can be present in a viral accessory construct. Any Tat polypeptide can be utilized as long as it is capable of binding to TAR and activating transcription of the RNA. This includes native Tat sequences which are obtained from the same or different species as the cognate TAR element, as well as engineered and modified Tat sequences.

Promoters

[0105] In an aspect, a construct disclosed here contains one or more expression cassettes that express an accessory protein or RNA molecule under the control of a constitutive, inducible, switched, recombined, disrupted/edited promoter or promoter/enhancer. In an aspect, a promoter is a minimal promoter with upstream cis regulatory to determine spatio-temporal expression pattern of the promoter. Upstream regulatory elements may include cis-acting elements (or cis-acting motifs) or transcription factor binding sites. In a further aspect, the promoter comprises a combination of heterologous upstream regulatory elements.

[0106] In an aspect, a promoter is a promoter/enhancer. As used herein, the term promoter/enhancer refers to a segment of DNA that contains sequences capable of providing both promoter and enhancer functions. The promoter/enhancer may be endogenous or exogenous or heterologous. An endogenous promoter/enhancer is one which is naturally linked with a given gene in a native viral genome. An exogenous or heterologous enhancer/promoter is one which is placed in juxtaposition to a gene by means of molecular biology techniques such that the transcription of that gene is directed by the linked promoter/enhancer.

[0107] In an aspect, a promoter is an inducible promoter. In an aspect, an inducible promoter is positively inducible and regulated by positive control. In an aspect, an inducible promoter is negatively inducible, and regulated by negative control.

[0108] In a further aspect, an inducible promoter may be a chemically inducible promoter. Chemically inducible promoters are known in the art. In a further aspect, a chemically inducible promoter may be a tetracycline controllable promoter. In a further aspect, a tetracycline-controllable promoter is a natural promoter. In a further aspect, a tetracycline controllable promoter is a synthetic promoter.

[0109] In a further aspect, an inducible promoter may be a temperature inducible promoter. In a further aspect, an inducible promoter may be a light inducible promoter. In a further aspect, an inducible promoter may be a physiologically regulated promoter.

[0110] In an aspect, a promoter may be a constitutive promoter. In an aspect, a promoter may be a switched promoter. In an aspect, a promoter may be a recombined promoter. In an aspect, a promoter may be a disrupted/edited promoter.

[0111] In an aspect, a promoter element may be naturally derivable. In a further aspect, a promoter may contain sequences derived from a eukaryotic promoter including, but not limited to CMV, EF1a, SV40, PGK1, Ubc, human beta actin, CAG, TRE, CaMKIIa, Cal1, 10, H1, and U6.

[0112] In a further aspect, a promoter comprises synthetic elements. Methods to prepare synthetic promoters are known in the art. In an aspect, a synthetic promoter is a constitutive synthetic promoter. In an aspect, a synthetic promoter is an inducible synthetic promoter. In an aspect, a synthetic promoter is a tissue specific synthetic promoter.

Polyadenylation sequences

[0113] In an aspect, a viral vector genome construct or a viral accessory construct comprises one or more polyadenylation (p(A)) sequences. Expression of recombinant DNA sequences in eukaryotic cells requires expression of signals to direct termination and polyadenylation of the resulting transcript. The term “polyadenylation sequence” as used herein refers to a nucleic acid sequence that directs the termination and polyadenylation of a nascent formed RNA transcript. Transcripts lacking a poly A tail may be unstable and quickly degraded. A poly A signal utilized in a viral vector genome construct disclosed herein may be heterologous or endogenous. An endogenous poly A signal refers to a poly A sequence that is found naturally at the 3' end of the coding region of a given gene. A heterologous poly A signal refers to a poly A sequence that is isolated from one gene and placed at the 3' end of another gene.

Expression cassettes

[0114] In an aspect, a viral vector genome construct and/or a viral accessory construct described here comprise one or more expression cassettes. Expression cassettes may be a monocistronic expression cassette or a polycistronic expression cassette.

[0115] In an aspect, a polycistronic expression cassette contains one or more viral skip sequences. Viral skip sequences are “self-cleaving” 2A peptides, which are 18-22 amino acid viral oligopeptides that mediate “cleavage” of polypeptides during translation in eukaryotic cells. The “2A” designation refers to a specific region of the viral genome. The mechanism of

2A cleavage is ribosome skipping, mediated by a highly conserved C-terminal sequences essential to the creation of steric hindrance. In an aspect, viral skip sequences may include 2A peptides derived from porcine teschovirus-1 2A (P2A). In an aspect, viral skip sequences may include 2A peptides derived from Thossea asigna virus 2A (T2A). In an aspect, viral skip
5 sequences may include 2A peptides derived from equine rhinitis A virus (E2A). In an aspect, viral skip sequences may include 2A peptides derived from foot-and-mouth disease virus (F2A). In a further aspect, viral skip sequences may be derived from any virus with a 2A sequence substantially similar to the conserved "2A" C-terminal sequence GDVEXNPGP (SEQ ID NO: 1).

10 [0116] In an aspect, a polycistronic expression cassette contains one or more internal ribosome entry site elements (IRES). An IRES element is a cis-acting RNA region that promotes internal initiation of protein synthesis. An IRES sequence is recognized by a ribosome and can therefore be used to drive translation of multiple proteins off a single transcript.

15 [0117] In a further aspect, a polycistronic expression cassette contains one or more viral skip sequences and one or more internal ribosome entry site elements.

[0118] In an aspect, polycistronic expression cassettes encode for sequences providing a similar mechanism to viral skip sequences or internal ribosome entry sequences.

Codon optimization

20 [0119] Expression cassettes contain sequences that encode one or more viral accessory proteins. In an aspect, a viral accessory protein may be encoded by a wild-type sequence. In a further aspect, a viral accessory protein may be encoded by a codon optimized ("co") sequence. Codon optimization is commonly used to increase production of recombinant proteins or viral vectors. Codon optimization is a desirable molecular tool to address codon
25 usage bias. Codon usage bias is a feature of all genomes and reflects the frequency of codon distribution within a genome is referred to as codon usage bias. Codon usage is variable between species, and preferred codons are more frequently used in highly expressed genes. Transfer RNAs, or tRNAs, reflect the codon usage in a given organism, and therefore the abundance of particular tRNAs is variable between organisms. Codon optimization is a process
30 by which DNA sequences are modified by introducing silent mutations to generate synonymous codons.

[0120] In a further aspect, an expression cassette may contain sequences that are all wild-type sequences, all codon optimized sequences, or a combination of both wild type and codon optimized sequences. In an aspect, expression of Rev, Tat, Nef, Vpr, Vif, Vpu/Vpx when included, is from wild-type or codon optimized constructs which are polycistronic using viral skip sequences (such as P2A, or T2A) or internal ribosome entry sequences or other similar mechanism or as a single message per transcript. In an aspect, expression of Gag-Pol is from a wild-type or codon optimized polycistronic message, or as separate *gag* and *pol* constructs, or as further separated CA, MA SP1, NC, p6, RT, IN, PR, and/or DU constructs.

Introducing viral vectors to target or host cells

10 [0121] In an aspect, the introduction of one or more constructs into a cell is achieved using a standard chemical, biological, or physical methods including, but not limited to, lipofectamine or lipofectamine-like chemical reagents, polyethyleneimine (PEI), calcium phosphate crystals, retroviral vector, lentiviral vector, nanoparticles or nanoparticle-like reagents, or electroporation. In another aspect, incorporation of these constructs into the cell line genome is achieved using biological recombinant enzymes including, but not limited to, 15 integrase, transposase, recombinase, the CRISPR-Cas9 system, or utilizing spontaneous or targeted insertion using cellular DNA repair machinery.

[0122] In an aspect, methods of introducing viral vector constructs to a target or host cell may include transduction or transfection. Transfection and transduction may be performed using a variety of techniques known in the art and may include optimizations for enhancing 20 transfection or transduction efficiency. In an aspect, optimization may comprise freeze-thawing reagents.

[0123] In an aspect, viral vector constructs are introduced to target or host cells using chemical methods known in the art. In an aspect, viral vector constructs are introduced to target or host cells using biological methods known in the art. In an aspect, viral vector constructs 25 are introduced to target or host cells using physical methods known in the art.

[0124] In an aspect, viral vector constructs may be introduced to a target or host cell by methods comprising optical techniques. In an aspect, viral vector constructs may be introduced to a target or host cell by methods comprising magnetic techniques. In an aspect, viral vector constructs may be introduced to a target or host cell by methods comprising biolistic techniques. In an aspect, viral vector constructs may be introduced to a target or host cell by 30 methods comprising polymer-based techniques. In an aspect, viral vector constructs may be

introduced to a target or host cell by methods comprising liposome-based techniques. In an aspect, viral vector constructs may be introduced to a target or host cell by methods comprising nanoparticle-based techniques. In a further aspect, viral vector constructs may be introduced to a target or host cell by a combination of methods comprising a combination of techniques including, but not limited to optical, magnetic, biolistic, polymer-based, liposome-based, and nanoparticle-based techniques.

[0125] In a further aspect, viral vector constructs may be introduced to a target or host cell by methods comprising electroporation. In a further aspect, viral vector constructs may be introduced to a target or host cell by methods comprising sonoporation. In a further aspect, viral vector constructs may be introduced to a target or host cell by methods comprising mechanoporation. In a further aspect, viral vector constructs may be introduced to a target or host cell by methods comprising photoporation.

[0126] In a further aspect, methods of introduction may also comprise methods that involve use of a cationic polymer, calcium phosphate, cationic lipid, or a combination thereof. In an aspect, a cationic polymer is hexadimethrine bromide (commercial brand name Polybrene).

[0127] In a further aspect, methods of introduction may also comprise methods that involve use of a retrovirus, lentivirus, transposon, CRISPR/Cas9, or recombinase. In an aspect, a recombinase may be a Cre-recombinase, Flippase recombinase, or a derivative thereof.

[0128] Methods to promote the integration of nucleic acids into production cells are known in the art, and can include, but are not limited to, linearizing a nucleic acid construct.

[0129] In an aspect, one or more viral vector constructs may be stably integrated or episomally maintained within the viral vector production cell. Gene expression of sequences encoded by any of the introduced viral vectors may occur from integrated sequences or episomes.

[0130] In an aspect, a viral vector production cell stably expressing some of the components may be transfected with remaining components that are required for vector production. Transfection of the remaining components required for viral vector production may be transient.

[0131] A viral vector construct may integrate randomly or in a site-specific manner upon introduction into a host or target cell.

Viral vector production cells

[0132] The disclosure disclosed herein provides a method of making viral vector particles in vitro by introducing one or more viral vector constructs of the disclosure into a compatible target cell or host cell and growing the cell under conditions which result in cell expansion and expression of the vector components. The terms “target cell” and “host cell” as used herein are interchangeable.

[0133] A viral vector production cell is a target cell or host cell that is capable of producing a viral vector or viral vector particle upon introduction of one or more viral vector constructs.

[0134] In an aspect, a viral vector production cell is a transgenic cell. As used herein, the term “transgenic cell” refers to a cell comprising genetic material that has been transferred naturally or by any of a number of genetic engineering techniques known in the art from one cell type to another cell type. In a further aspect, a transgenic cell refers to a cell comprising experimentally constructed genetic material. In an aspect, a viral vector production cell population is polyclonal. Polyclonal cells comprise a heterogeneous population of cells with multiple clones that may have variations in the number of integration events and sites of integration across the cells. In a further aspect, a viral vector production cell population is monoclonal.

[0135] In an aspect, a viral vector production cell is from a cell line that has been expanded from a selected viral vector production cell clone.

[0136] Viral vector production cell clones may be derived from a polyclonal population by methods known in the art. Methods of selection include, but are not limited to, limiting dilution, single cell sorting, and single cell selection. Limiting dilution may be performed by methods known in the art. Single cell sorting may be performed by methods known in the art, including, but not limited to, single cell printing, fluorescence activated cell sorting (FACS), and magnetic activated cell sorting. Single cell selection may be performed by selection methods known in the art, including, but not limited to selection for an epitope, a protein, a reporter gene, or combination thereof. In a further aspect, single cell selection methods may comprise selection via one or more metabolic or antibiotic properties.

[0137] In an aspect, viral vector production cell clones or cell lines grow in an adherent manner. In an aspect, viral vector production cell clones or cell lines grow in suspension. In a further aspect, viral vector production cell clones or cell lines that are adherent may be

suspension-adapted. In a further aspect, viral vector production cell clones or cell lines that grow in an adherent form.

[0138] In an aspect, viral vector production cell clones or cell lines are cultured in serum-supplemented or serum-free media. A person of skill in the art will be able to select an appropriate media for the given viral vector production cell type, and to modify the media composition at various stages of the method disclosed herein. Media may have a selection of secreted cellular proteins, diffusible nutrients, amino acids, organic salts, inorganic salts, vitamins, trace metals, sugars, and other growth-promoting substances such as cytokines. Media may be supplemented with glutamine or an alternative thereof.

[0139] In an aspect, viral vector production cell clones or cell lines may be any eukaryotic cell that supports the lifecycle of the specific virus from which the vector is derived. In an aspect, for a retroviral vector, a production cell clones or cell lines may be any eukaryotic cell that supports a retrovirus life cycle. In an aspect, for a lentiviral vector, a production cell clones or cell lines may be any eukaryotic cell that supports a lentivirus life cycle. In an aspect, for a herpesvirus vector, a production cell clones or cell lines may be any eukaryotic cell that supports a herpesvirus life cycle. In an aspect, for an adenoviral vector, a production cell clones or cell lines may be any eukaryotic cell that supports an adenovirus life cycle. In an aspect, for an adeno-associated viral vector, a production cell clones or cell lines may be any eukaryotic cell that supports an adeno-associated virus life cycle.

[0140] In an aspect, viral vector production cell clones or cell lines are immortalized. Cell lines may be commercially available or non-commercially available laboratory-derivatives. In a further aspect, viral vector production cell clones or cell lines are of eukaryotic origin. In an aspect, viral vector production cell clones or cell lines are of mammalian origin. Mammalian cells for the production of viral vectors are known in the art. In an aspect, viral vector production cell clones or cell lines are of human origin.

[0141] In a further aspect, a viral vector producer cell line is developed in or from Human Embryonic Kidney (HEK) 293 cells, which are highly transfectable. In a further aspect, a viral vector producer cell line is a derivate of HEK293 cells, such as HEK293T or HEK293F cells. In a further aspect, cell types for viral vector production cell clones or cell lines include, but are not limited to, HeLa cells, Vero cells, Chinese Hamster Ovary (CHO) cells, A549 cells, and NIH 3T3 cells.

Characterization of produced viral vector

[0142] Viral vector particles produced by a viral vector producer cell clone or cell line may be characterized by a variety of methods known to those of skill in the art.

5 [0143] In an aspect, a viral vector particle produced by a method disclosed herein is a pseudotyped viral particle. Pseudotyped viral particles may be produced by substituting viral attachment proteins from one viral serotype with another. As used herein, a “viral attachment protein” refers to a viral capsid protein or a viral envelope protein.

10 [0144] In an aspect, a viral vector particle produced by a method disclosed herein is a mosaic viral particle. Mosaic viral particles may be produced by mixing different viral attachment proteins from different viral variants.

[0145] In an aspect, a viral vector particle produced by a method disclosed herein is a chimeric viral particle. Chimeric viral particles may be produced by methods that include swapping smaller domains of viral attachment proteins between serotypes (via rational methods or high throughput recombination techniques).

15 [0146] From a stable viral vector producing cell clone or cell line, viral vector genome and accessory proteins may be characterized quantitatively or qualitatively. In an aspect, the stoichiometric ratio of viral vector genome and one or more accessory proteins may be determined. In a further aspect, the level of viral vector genome and one or more accessory proteins may be determined.

20 [0147] An integration profile of a selected cell clone or cell line may be determined. In an aspect, an integration profile or an insertional profile may be detected by methods known in the art such as inverse PCR, linear amplification-mediated PCR or ligation-mediated PCR. Vector flanking sequences detected by such methods can then be mapped to a host cell genome and compared to a reference set. Mapping can be performed using computational tools to map
25 and analyze vector-flanking sequences, such as QuickMap.

[0148] In an aspect, recombinant viral vectors may be harvested from a cell clone or a cell line. In an aspect, the cell line is monoclonal. Harvested viral vectors may be characterized qualitatively or quantitatively. In an aspect, viral titer is expressed in transducing units per milliliter (t.u./ml).

[0149] Viral titer may be determined using physical or functional titration. In an aspect, titration methods include but are not limited to transduction of indicator cells using dose-dependent quantities of vector supernatant.

[0150] In a further aspect, viral titer is determined by assaying for viral nucleic acid or viral protein. In an aspect, viral nucleic acid can be detected using polymerase chain reaction (PCR), reverse transcription PCR (RT-PCR), dot blot hybridization, Southern blot hybridization, or northern blot hybridization. In a further aspect, viral protein can be detected via an immunoassay. Immunoassays include but are not limited to immunoblotting (Western blotting), immunofluorescence, immunocytochemistry, and enzyme-linked immunosorbent assay (ELISA).

[0151] In a further aspect, transduced indicator cells may be assessed using PCR. Quantification by PCR may be performed using relative quantification or absolute quantification. Methods for relative or absolute quantification by PCR are known in the art.

[0152] In a further aspect, methods of viral titer determination are enzyme immunoassays. Harvested viral particles may be quantified by measuring the amount of a viral capsid protein using immunoassays specific to the virus from which the viral capsid protein was derived (for example, p24 for HIV).

[0153] Viral vector particles produced by methods disclosed herein may be concentrated and/or purified using flow-through ultracentrifugation and high-speed centrifugation, and tangential flow filtration. Flow through ultracentrifugation has been used in the past for the purification of RNA tumor viruses (Toplin *et al.*, Applied Microbiology 15:582-589, 1967; Burger *et al.*, Journal of the National Cancer Institute 45: 499-503, 1970). The present disclosure provides the use of flow-through ultracentrifugation for the purification of lentiviral vectors. This method can comprise one or more of the following steps. For example, a lentiviral vector can be produced from cells using a cell factory or bioreactor system. A transient transfection system (see above) can be used or packaging or producer cell lines can also similarly be used. A pre-clarification step prior to loading the material into the ultracentrifuge could be used if desired. Flow-through ultracentrifugation can be performed using continuous flow or batch sedimentation. The materials used for sedimentation are, *e.g.*: Cesium chloride (CsCl), potassium tartrate and potassium bromide, which create high densities with low viscosity although they are all corrosive. CsCl is frequently used for process development as a high degree of purity can be achieved due to the wide density gradient that can be created (1.0

to 1.9 g/cm³). Potassium bromide can be used at high densities, but only at elevated temperatures, i.e. 25° C, which may be incompatible with stability of some proteins. Sucrose is widely used due to being inexpensive, non-toxic and can form a gradient suitable for separation of most proteins, sub-cellular fractions and whole cells. Typically, the maximum
5 density is about 1.3 g/cm³. The osmotic potential of sucrose can be toxic to cells in which case a complex gradient material can be used, *e.g.* Nycodenz. A gradient can be used with 1 or more steps in the gradient. A preferred aspect is to use a step sucrose gradient. The volume of material can be preferably from 0.5 liters to over 200 liters per run. The flow rate speed is preferably from 5 to over 25 liters per hour. The preferred operating speed is between 25,000
10 and 40,500 rpm producing a force of up to 122,000x g. The rotor can be unloaded statically in desired volume fractions. A preferred aspect is to unload the centrifuged material in 100ml fractions. The isolated fraction containing the purified and concentrated lentiviral vector can then be exchanged in a desired buffer using gel filtration or size exclusion chromatography. Anionic or cationic exchange chromatography could also be used as an alternate or additional
15 method for buffer exchange or further purification. In addition, Tangential Flow Filtration can also be used for buffer exchange and final formulation if required. Tangential Flow Filtration (TFF) can also be used as an alternative step to ultra or high-speed centrifugation, where a two-step TFF procedure would be implemented. The first step would reduce the volume of the vector supernatant, while the second step would be used for buffer exchange, final formulation
20 and some further concentration of the material. The TFF membrane should have a membrane size of between 100 and 500 kilodaltons, where the first TFF step should have a preferable membrane size of 500 kilodaltons, while the second TFF should have a preferable membrane size of between 300 to 500 kilodaltons. The final buffer should contain materials that allow the vector to be stored for long term storage.

25 [0154] The present disclosure also provides methods for the concentration and purification of lentiviral vectors using either cell factories that contains adherent cells, or a bioreactor that contains suspension cells that are either transfected or transduced with the vector and accessory constructs to produce lentiviral vector. Non limiting examples of bioreactors include the Wave bioreactor system and Xcellerex bioreactors. Both are disposable systems. However non-
30 disposable systems can also be used. The constructs can be those described herein, as well as other recombinant viral vectors. Alternatively, the cell line can be engineered to produce lentiviral vector without the need for transduction or transfection. After transfection, the lentiviral vector can be harvested and filtered to remove particulates and then is centrifuged

using continuous flow high speed or ultra-centrifugation. In an aspect, a high-speed continuous flow device like the JCF-A zonal and continuous flow rotor with a high-speed centrifuge is used. Also provided is any continuous flow centrifuge where the speed of centrifugation is greater than 5,000xg RCF and less than 26,000x g RCF. Preferably, the continuous flow centrifugal force is about 10,500x g to 23,500 x g RCF with a spin time of between 20 hours and 4 hours, with longer centrifugal times being used with slower centrifugal force. The lentiviral vector can be centrifuged on a cushion of more dense material (a non-limiting example is sucrose but other reagents can be used to form the cushion and these are well known in the art) so that the lentiviral vector does not form aggregates that are not filterable, as is the problem with straight centrifugation of the vector that results in a viral vector pellet. Continuous flow centrifugation onto a cushion allows the vector to avoid large aggregate formation yet allows the vector to be concentrated to high levels from large volumes of transfected material that produces the lentiviral vector. In addition, a second less-dense layer of sucrose can be used to band the lentiviral vector preparation. The flow rate for the continuous flow centrifuge is preferably between 1 and 100ml per minute, but higher and lower flow rates can also be used. The flow rate is adjusted to provide ample time for the vector to enter the core of the centrifuge without significant amounts of vector being lost due to the high flow rate. If a higher flow rate is desired, then the material flowing out of the continuous flow centrifuge can be re-circulated and passed through the centrifuge a second time. After the virus is concentrated using continuous flow centrifugation, the vector can be further concentrated using Tangential Flow Filtration (TFF), or the TFF system can be simply used for buffer exchange. A non-limiting example of a TFF system is the Xampler cartridge system that is produced by GF> Healthcare. Preferred cartridges are those with a MW cut-off of 500,000 MW or less. Preferably a cartridge is used with a MW cut-off of 300,000 MW. A cartridge of 100,000MW cut-off can also be used. For larger volumes, larger cartridges can be used, and those skilled in the art can find the right TFF system for this final buffer exchange and/or concentration step prior to final fill of the vector preparation. The final fill preparation may contain factors that stabilize the vector. For example, sugars are generally used and are known in the art.

Further cell line modification

[0155] In an aspect, a cell line utilized to manufacture a recombinant viral vector can be modified in any of the ways mentioned below to enhance viral vector production, *e.g.*, by the introduction of RNAi or antisense to knock-out genes that reduce the expression of genes that limit viral vector production, or by the introduction of sequences that enhance viral vector

production. Sequences that code for cellular or viral enhancers can also be engineered into cell lines (*e.g.*, using additional plasmid vectors), such as herpes virus, hepatitis B virus, which act on HIV LTRs to enhance the level of virus product, or cellular transactivator proteins. Cellular transactivation proteins include, *e.g.*, NF- κ B, UV light responsive factors, and T cell activation factors. In another aspect, a cell line utilized to manufacture a recombinant viral vector can be modified or edited by a nuclease selected from the group consisting of a meganuclease, a zinc-finger nuclease (ZFN), a transcription activator-like effector nuclease (TALEN), a CRISPR-related nuclease (*e.g.*, Cas9, Cas12a, etc.).

[0156] In a further aspect, a cell line utilized to manufacture a recombinant viral vector can be modified or optimized using a combination of recombination-based and gene editing-based approaches.

[0157] In an aspect, a cell line can be transformed routinely with construct DNA, *e.g.*, using electroporation, calcium phosphate, liposomes, etc., to introduce the DNA into cells. Cells can be co-transformed (*i.e.*, using both accessory and transfer vectors), or they can be transformed in separate steps, where each step involves the introduction of a different vector.

[0158] Cells are cultured under conditions effective to produce viral vectors. Such conditions include, *e.g.*, the particular milieu needed to achieve protein production. Such a milieu, includes, *e.g.*, appropriate buffers, oxidizing agents, reducing agents, pH, co-factors, temperature, ion concentrations, suitable age and/or stage of cell (such as, in particular part of the cell cycle, or at a particular stage where particular genes are being expressed) where cells are being used, culture conditions (including cell media, substrates, oxygen, carbon dioxide, glucose and other sugar substrates, serum, growth factors, etc.).

[0159] The present disclosure also provides for the use of cell lines that have enhanced properties for growth, reduced dependency upon expensive factors that are present in media, produce higher yields of proteins, and produce higher titers of vector particles. For example, it has recently been reported HEK 293 cells have a specific increased expression of cellular receptors and by adding the specific ligands to the medium of the cells, they demonstrated increase proliferation potential (Allison *et al.*, Bioprocess International 3:1, 38-45, 2005). A preferred aspect is a plurality of lentiviral vectors expressing an optimized combination of ligand proteins that are of relevance to HEK 293 cells after which the cells are then sorted by high throughput methods to isolate a clone of HEK 293 cells that contains multiple copies of lentiviral vectors. These cells contain a combination of HIV vectors that express different but

also multiple copies of the ligand genes that are contained in the HIV vectors. The ligand genes could be codon optimized or mutations added to further increase their expression. A preferred combination is to have multiple copies of the ligand proteins expressed in the final isolated clonal cell that could then have multiple uses. It could be used for protein or antibody (including monoclonal, humanized, single-chain) production. It could also be used for the production of a vector such as a lentiviral vector, but not limited to a lentiviral vector. Other vectors such as Adeno and Adeno-associated vectors, murine retroviral vectors, SV40 vectors and other vectors could just as easily be produced from this now optimized cell line. A list of the receptors and their ligands that show increased expression/activity in HEK 293 cells, includes, *e.g.*, AXL receptor (gas β); EGF receptor (EGF), chemokine receptor (fractalline); PDGF receptor, beta (PDGF); IL-15R-alpha; IL-2R-alpha; chemokine receptor 2 (MCP1); IL-2R, gamma; IL-1R-1; CSF-I receptor; oncostatin receptor; IL-4R; vitamin D3 receptor; neuropilin 1 (VEGF); macrophage stimulating receptor 1 (MSP); NGF-R; PDGFR-alpha receptor; IL-11-R, *e.g.*, alpha; IL-10-R, *e.g.*, beta; FGF-R-4 (aFGF); BMP receptor, *e.g.*, type II (BMP-2); TGF-R, *e.g.*, beta receptor II (TGF-beta); FGF-R-I (bFGF); chemokine receptor 4 (SFD1a); interferon gamma receptor 1 and 2. See, BioProcess International, January 2005. Table 1, "Growth factor/cytokine receptors expressed by HEK-293." Such cells will have higher protein and vector production potential and will be less dependent upon the presence of the ligand factors to be present in the medium since the cells themselves will be producing the factors and secreting them into the medium.

[0160] For other cell types, such as CHO cells, other receptor-ligand combinations may be important. For example, the insulin growth factor receptor I, insulin growth factor and insulin are thought to have anti-apoptotic activity in cells. A plurality lentiviral vectors could be constructed so that the Insulin growth factor receptor (I or II), Insulin growth factor (I or II), Insulin and the target protein for production are all contained in the vector for transduction of production cells, such as CHO cells, and an appropriate clone selected, preferably using high-throughput methods, to select the clone showing very high production of the target protein. The optimal clone may not be a cell that highly expresses all the engineered genes or inhibitors of gene expression, rather an optimal expression level of each of the genes, which for some may be a low level of expression. The value of the lentiviral vector system and using a plurality of lentiviral vectors to engineer such cell lines is that there is a random or stochastic distribution of each vector copy number in the population of cells transduced with the lentiviral vector mixture, and therefore, by varying the amount of each vector in the mixture, the number of

copies of each individual second gene or inhibitory sequence can be optimized. A preferred combination of vectors and secondary gene or gene inhibitory sequences is that each lentiviral vector expresses the protein of interest for production and optionally in addition, at least one RNAi or gene that further promotes protein yield, or vector yield, either directly, or indirectly
5 by affecting the viability or some aspect of the producing cell. However, it may also be beneficial to have at least one lentiviral vector that only expresses the secondary genes or inhibitors of gene expression in order to increase the effect of these secondary sequences.

[0161] In an aspect, the present disclosure provides for the following non-limiting embodiments:

- 10 1. A construct comprising a promoter sequence operably linked to a first set of one or more copies of coding sequences each encoding a viral accessory protein or a recombinant viral vector genome, wherein said construct further comprises two recombinase recognition sites in opposite orientations.
- 15 2. The construct of embodiment 1, wherein said promoter sequence becomes operably linked to more or fewer copies of said coding sequences relative to said first set of one or more copies of coding sequences upon a recombination event via said two recombinase recognition sites.
- 20 3. The construct of embodiment 1, wherein said recombinase recognition sites flank said promoter sequence.
4. The construct of embodiment 1, wherein said recombinase recognition sites flank at least one coding sequence encoding a viral accessory protein or a recombinant viral
25 vector genome.
5. The construct of embodiment 1, wherein said recombinase recognition sites flank said one or more copies of said viral accessory protein coding sequence.
- 30 6. The construct of embodiment 1, wherein said recombinase recognition sites flank said one or more copies of said recombinant viral vector genome coding sequence.

7. The construct of embodiment 1, wherein said recombinase recognition sites flank said promoter and said one or more copies of said viral accessory protein coding sequence.
8. The construct of embodiment 1, wherein said recombinase recognition sites flank a
5 promoter and said one or more copies of said recombinant viral vector genome coding sequence.
9. The construct of any one of embodiments 1 to 8, further comprising a second set of one or more copies of said coding sequences, wherein, upon inversion via the two
10 opposing recombinase recognition sites, said promoter becomes operably linked to said second set of coding sequences.
10. The construct of embodiment 1, further comprising one or more endonuclease recognition motifs.
15
11. The construct of embodiment 10, wherein said one or more endonuclease recognition motifs are protospacer adjacent motifs.
12. A construct comprising a promoter sequence flanked by two sets of oppositely oriented
20 coding sequences each encoding a viral accessory protein or a recombinant viral vector genome, wherein said promoter is capable of being inverted upon a recombination event inducible by a recombinase, wherein said promoter is operably linked to a first set of said two sets of oppositely oriented coding sequences prior to recombination and becomes operably linked to a second set of said two sets of oppositely oriented coding
25 sequences post recombination.
13. The construct of embodiment 12, wherein said promoter is flanked by two recombinase recognition sites in opposite orientations.
14. A construct comprising a promoter sequence operably linked to one or more coding
30 sequences each encoding a viral accessory protein, a recombinant viral vector genome or both, wherein at least one of said one or more coding sequences are flanked by two recombinase recognition sites, wherein said at least one of said one or more coding

sequences are capable of being disrupted, deleted or inverted upon a recombination event inducible by a recombinase.

- 5 15. A construct comprising (i) a promoter sequence operably linked to a first set of one or more coding sequences each encoding a viral accessory protein, a recombinant viral vector genome or both, and (ii) two recombinase recognition sites flanking said promoter sequence and said first set of one or more coding sequences, wherein, upon a recombination event inducible by a recombinase, said promoter sequence becomes operably linked to additional coding sequences besides said first set of one or more coding sequences.
- 10 16. The construct of embodiment 9 or 12, wherein said viral accessory protein or recombinant viral vector genome is from a virus selected from the group consisting of lentivirus, retrovirus, herpesvirus, adenovirus, and adeno-associated virus.
- 15 17. The construct of embodiment 9 or 12, wherein said first and second sets of one or more copies of coding sequences are capable of providing different levels of expression of said coding sequences depending on an orientation of said promoter.
- 20 18. The construct of embodiment 9 or 12, wherein said first and second sets of one or more copies of coding sequences comprise different copy numbers.
- 25 19. The construct of embodiment 1 or 12, wherein said promoter comprises an inducible, recombined, or edited promoter.
20. The construct of embodiment 1 or 13, wherein said recombinase recognition sites comprise two palindromic recognition regions flanking a spacer region.
- 30 21. The construct of embodiment 1 or 13, wherein each of said recombinase recognition sites comprises a lox site.
22. The construct of embodiment 21, wherein said lox site comprises a wildtype lox site or a mutant lox site.

23. The construct of embodiment 20, wherein said spacer region is selected from the group consisting of loxP, lox511, lox2272, lox5171, m2, m3, and m7.
24. The construct of any one of embodiments 1 to 23, wherein said recombinase is a Cre-
5 recombinase.
25. The construct of embodiment 1 or 13, wherein each of said recombinase recognition sites are a Flippase recombinase target sequence.
- 10 26. The construct of embodiment 25, wherein said recombinase is a Flippase recombinase.
27. The construct of embodiment 1 or 12, wherein said first set of said one or more copies of coding sequences comprises two or more copies of said coding sequences arranged as a polycistronic expression cassette.
- 15 28. The construct of embodiment 27, wherein said polycistronic expression cassette is in the sense orientation downstream of said promoter prior to promoter inversion.
29. The construct of embodiment 27, wherein said polycistronic expression cassette further
20 comprises one or more viral skip sequences, internal ribosome entry site elements, or both.
30. The construct of embodiment 29, wherein said viral skip sequences are selected from the group consisting of P2A, T2A, E2A, and F2A.
- 25 31. The construct of embodiment 9 or 12, wherein said second set of said one or more copies of coding sequences comprises a monocistronic expression cassette.
32. The construct of embodiment 31, wherein said monocistronic expression cassette is in
30 the antisense orientation upstream of said promoter prior to promoter inversion.
33. The construct of any of embodiments 1 to 32, wherein said viral accessory protein is a fusion protein.

34. The construct of any of embodiments 1 to 32, wherein said viral accessory protein comprises a sequence encoding a structural viral protein, a regulatory viral protein, or both.
- 5 35. The construct of embodiment 34, wherein said structural and regulatory proteins are selected from the group consisting of Gag, Pol, Rev, Env, Tat, Nef, Vpr, Vif, Vpu, and Vpx.
- 10 36. The construct of any of embodiments 1 to 32, wherein said viral accessory protein comprises a sequence encoding one or more viral accessory protein domains.
37. The construct of embodiment 36, wherein said one or more viral accessory protein domains are selected from the group consisting of CA, MA, NC, p6, SP1, RT, IN, PR, and DU.
- 15 38. The construct of any one of embodiments 1 to 37, wherein a sequence encoding said viral accessory protein or one or more viral accessory protein domains comprise a wild-type sequence, a codon optimized sequence, or both
- 20 39. A construct comprising a promoter sequence operably linked to two or more coding sequences each encoding a viral accessory protein, a recombinant viral vector genome or both, wherein at least one of said two or more coding sequences is codon optimized.
40. A cell comprising the construct of any of embodiments 1 to 39.
- 25 41. The cell of embodiment 40, wherein said cell is a eukaryotic cell.
42. The cell of embodiment 40, wherein said cell is a mammalian cell.
- 30 43. The cell of embodiment 40, wherein said cell is a human cell.
44. The cell of embodiment 40, wherein said cell is a viral vector producer cell.

45. The cell of embodiment 40, wherein said construct is integrated in the genome of said cell.
46. The cell of embodiment 44, wherein said viral vector producer cell is adherent or in suspension.
47. The cell of embodiment 44, wherein said viral vector producer cell is cultured in serum-supplemented or serum-free media.
48. The cell of embodiment 44, wherein said viral vector producer cell is immortalized.
49. The cell of embodiment 44, wherein said viral vector producer cell is a HEK293 cell, or a derivative thereof.
50. The cell of embodiment 49, wherein said HEK293 cell is a HEK293T cell.
51. A method of producing a viral vector producer cell with an optimized stoichiometric ratio of viral vector genome and viral accessory proteins, wherein said method comprises:
- introducing the construct of any of embodiments 1 to 39 into a first clonal population of cells;
 - transiently providing a recombinase and/or a CRISPR-based complex to said first clonal population; and
 - producing a second clonal population by (i) inverting an invertible sequence flanked by recombinase recognition sites; and/or (ii) editing one or more copies of coding sequences each encoding a viral accessory protein, a recombinant viral vector genome or both.
52. The method of embodiment 51, wherein said second clonal population comprises a viral vector genome construct and one or more accessory constructs encoding one or more viral accessory proteins.

53. The method of embodiment 51, wherein said method further comprises generating from said second clonal population a stable viral vector producer cell line.
54. The method of embodiment 51, wherein said invertible sequence comprises a promoter
5 sequence, a viral accessory protein sequence, a recombinant viral vector genome sequence, or a combination thereof.
55. The method of embodiment 53, wherein said stable viral vector producer cell line exhibits a higher viral titer relative to said first clonal population.
- 10 56. The method of embodiment 55, wherein said viral titer is determined by physical titration, functional titration, or both.
57. The method of embodiment 55, wherein said viral titer is determined by assaying for
15 viral nucleic acid via an assay selected from the group consisting of PCR, RT-PCR, and quantitative detection by blot hybridization, or by assaying for viral protein via an immunoassay.
58. The method of embodiment 51, wherein said method further comprises determining
20 the stoichiometric ratio of viral vector genome RNA and said viral accessory proteins in said first clonal population, second clonal population, or both.
59. The method of embodiment 51, wherein said method further comprises quantifying the
25 level of said viral vector genome and said viral accessory proteins in said first clonal population, second clonal population, or both.
60. The method of embodiment 53, wherein said method further comprises storing said stable viral vector producer cell line by cryopreservation.
- 30 61. The method of embodiment 53, wherein said method further comprises expanding cells from said cryopreserved cell line to produce viral vectors.
62. The method of embodiment 51, wherein said recombinase recognition sites comprise two palindromic recognition regions flanking a spacer region.

63. The method of embodiment 51, wherein each of said recognition sites comprises a lox site.
- 5 64. The method of embodiment 63, wherein said lox site comprises a wildtype lox site or a mutant lox site.
65. The method of embodiment 62, wherein said spacer region is selected from the group consisting of loxP, lox511, lox2272, lox5171, m2, m3, and m7.
- 10 66. The method of embodiment 51, wherein said recombinase is a Cre-recombinase.
67. The method of embodiment 51, wherein each of said recombinase recognition sites is a Flippase recombinase target sequence.
- 15 68. The method of embodiment 51, wherein said recombinase is a Flippase recombinase.
69. The method of embodiment 51, wherein said viral vector genome construct comprises one or more elements selected from the group consisting of a 5' long terminal repeat, a 20 3' long terminal repeat, a packaging signal, and a central polypurine tract.
70. The method of embodiment 51, wherein said viral vector genome construct does not comprise a 5' long terminal repeat, a 3' long terminal repeat, a packaging signal, or a 25 central polypurine tract.
71. The method of embodiment 51, wherein said viral vector genome construct comprises a self-inactivating long terminal repeat.
72. The method of embodiment 51, wherein said viral vector genome construct comprises 30 a promoter and a polyadenylation sequence.
73. The method of embodiment 72, wherein said promoter of said viral vector genome construct is constitutive or inducible.

74. The method of embodiment 51, wherein said viral vector genome construct comprises a concatemer.
75. The method of embodiment 74, wherein said concatemer comprises multiple copies of a viral vector genome construct.
76. The method of embodiment 74, wherein said concatemer comprises one or more selection genes.
77. The method of embodiment 74, wherein said concatemer comprises one or more transcription factors.
78. The method of embodiment 51, wherein said first clonal population or said second clonal population comprises viral vector producer cells.
79. The method of embodiment 51, wherein said introducing step may be chemical, biological or physical.
80. The method of embodiment 51, wherein said introducing step comprises optical, magnetic, biolistic, polymer-based, liposome-based, nanoparticle-based methods, or a combination thereof.
81. The method of embodiment 51, wherein said introducing step comprises a transduction.
82. The method of embodiment 51, wherein said introducing step comprises a transfection.
83. The method of embodiment 79, wherein said chemical introducing step comprises a cationic polymer, calcium phosphate, cationic lipid, or a combination thereof.
84. The method of embodiment 79, wherein said biological introducing step comprises introduction via retrovirus, lentivirus, transposon, CRISPR/Cas9, or recombinase.

85. The method of embodiment 79, wherein said physical introducing step is selected from the group consisting of electroporation, sonoporation, mechanoporation, and photoporation.
- 5 86. The method of embodiment 51, wherein said recombinase is an inducible recombinase.
87. The method of embodiment 51, wherein said recombinase comprises a Cre-recombinase, a Flippase recombinase, or a modified version thereof.
- 10 88. The construct of embodiment 86, wherein said inducible recombinase is chemical or light inducible.
89. The method of embodiment 51, wherein said cell is a eukaryotic cell.
- 15 90. The method of embodiment 51, wherein said cell is a mammalian cell.
91. The method of embodiment 51, wherein said cell is a human cell.
- 20 92. The method of embodiment 51, wherein said promoter comprises an inducible, recombined, or edited promoter.
93. The method of embodiment 51, wherein said viral accessory proteins comprise sequences encoding structural viral proteins, regulatory viral proteins, or both.
- 25 94. The method of embodiment 51, wherein said structural and regulatory proteins are selected from the group consisting of Gag, Pol, Rev, Env, Tat, Nef, Vpr, Vif, Vpu, and Vpx.
- 30 95. The method of embodiment 51, wherein said viral accessory proteins comprise sequences encoding a partial viral accessory protein.
96. The method of embodiment 51, wherein said partial viral accessory protein comprises one or more viral accessory protein domains.

97. The method of embodiment 96, wherein said one or more viral accessory protein domains is selected from the group consisting of CA, MA, NC, p6, SP1, RT, IN, PR, and DU.
- 5 98. The method of any one of embodiments 93 to 97, wherein a sequence encoding said one or more viral accessory proteins or domains comprises a wild-type sequence, a codon optimized sequence, or both.
99. A method comprising:
- 10 a. obtaining a first stable viral vector producer cell line;
- b. determining a viral titer from said first stable viral vector producer cell line; and
- c. engineering one or more elements in said stable viral vector producer cell line to produce a second stable viral vector producer cell line with a stoichiometric
- 15 ratio of viral vector genome and one or more viral accessory proteins different from the corresponding stoichiometric ratio in said first stable viral vector producer cell line.
100. The method of embodiment 99, wherein said engineering is via one or more
- 20 recombination events.
101. The method of embodiment 99, wherein said engineering comprises introducing a recombinase.
- 25 102. The method of embodiment 99, wherein said engineering comprises transiently expressing a recombinase.
103. The method of embodiment 99, wherein said engineering is via gene editing.
- 30 104. The method of embodiment 99, wherein said engineering comprises introducing a CRISPR-based complex.

105. The method of embodiment 99, wherein said engineering comprises introducing a CRISPR-based complex specific to some but not all copies of coding sequences for said one or more viral accessory proteins.

5 106. The method of embodiment 99, wherein said engineering comprises both a recombination event and gene editing.

[0162] Having now generally described the disclosure, the same will be more readily understood through reference to the following examples that are provided by way of illustration and are not intended to be limiting of the present disclosure, unless specified.

10

EXAMPLES

Example 1:

[0163] As an illustration of the concept described here, an HIV-based lentiviral vector producer cell line is produced (Figure 1). Typical stable cell lines include accessory gene components in a fixed ratio; however, some vector genome constructs require a different ratio of accessory gene components to achieve optimal titers. Figure 2 provides an exemplary illustration of how a stable vector producer cell line can be further optimized via recombination-based genome reorganization. For example, after initial vector producer cell clones are produced, their viral titers are determined. If no cell clones have an optimal titer, a recombinase is then introduced via for example transient expression, the ratio of accessory genes is then again randomly reassorted (Figure 3). Alternatively, Figure 4 illustrates another approach to optimize accessory gene component ratio via subtractive subcloning. For example, both a regular copy and a codon-optimized copy of the *rev* gene are used in an initial cell clone. Subsequently, one of the two *rev* gene copies are edited via, *e.g.*, CRISPR-mediated gene editing, such that only one copy remains functional and therefore the *rev* gene expression level is reduced. In a further alternative, as shown in Figure 9, viral vector producer cell lines can be further modified or optimized using a combination of recombination-based approaches (*e.g.*, Figure 2) and gene editing-based approaches (*e.g.*, Figure 4).

15

20

25

Example 2:

[0164] As an illustration of the concept described here, exemplary methods of re-assorting and altering viral accessory gene stoichiometry via use of an enzyme and without introducing new constructs are shown in Figures 5-9. Figure 5 provides an exemplary illustration of a construct with exemplary recombinase sites, *loxP*, flanking a promoter. Alternatively, Figure 6 and Figure 7 provide exemplary illustrations of constructs with exemplary recombinase sites, *loxP*, flanking an expression cassette that is operably linked to a promoter. Alternatively, Figure 8 provides an exemplary illustration of exemplary recombinase sites, *loxP*, that flank both a promoter and an expression cassette. Depending on coding sequence orientation, addition of a recombinase to invert an intervening coding sequence results in variable numbers of gene copies expressed from a given construct, as described below in Example 3.

Example 3:

[0165] As an illustration of the concept described here, addition of a recombinase to result in promoter inversion (Figure 5, *e.g.*), cassette inversion (Figures 6 and 7, *e.g.*), or promoter and cassette inversion (Figure 8, *e.g.*) results in expression of a variable number of gene copies from a given construct. In this exemplary illustration, depending on the orientation of coding sequences within the construct region flanked by recombinase sites, a promoter can drive expression of one or two copies of a gene (Figure 5 and Figure 6, *e.g.*), or two or three copies of a gene (Figure 7 and Figure 8, *e.g.*). This concept is not limited to copy numbers of consecutive integers, for example one to two, or two to three. As shown in Figure 9, *e.g.*, a method that combines recombinase-mediated promoter and/or cassette inversion with gene editing can result in expression of one to three copies, for example, of a gene from a given construct.

CLAIMS

1. A construct comprising a promoter sequence operably linked to a first set of one or more copies of coding sequences each encoding a viral accessory protein or a recombinant viral vector genome, wherein said construct further comprises two
5 recombinase recognition sites in opposite orientations.
2. The construct of claim 1, wherein said promoter sequence becomes operably linked to more or fewer copies of said coding sequences relative to said first set of one or more copies of coding sequences upon a recombination event via said two recombinase
10 recognition sites.
3. The construct of claim 1, wherein said recombinase recognition sites flank said promoter sequence.
- 15 4. The construct of claim 1, wherein said recombinase recognition sites flank at least one coding sequence encoding a viral accessory protein or a recombinant viral vector genome.
5. The construct of claim 1, wherein said recombinase recognition sites flank said one or
20 more copies of said viral accessory protein coding sequence.
6. The construct of claim 1, wherein said recombinase recognition sites flank said one or more copies of said recombinant viral vector genome coding sequence.
- 25 7. The construct of claim 1, wherein said recombinase recognition sites flank said promoter and said one or more copies of said viral accessory protein coding sequence.
8. The construct of claim 1, wherein said recombinase recognition sites flank a promoter and said one or more copies of said recombinant viral vector genome coding sequence.
30
9. The construct of any one of claims 1 to 8, further comprising a second set of one or more copies of said coding sequences, wherein, upon inversion via the two opposing

recombinase recognition sites, said promoter becomes operably linked to said second set of coding sequences.

- 5 10. The construct of claim 1, further comprising one or more endonuclease recognition motifs.
11. The construct of claim 10, wherein said one or more endonuclease recognition motifs are protospacer adjacent motifs.
- 10 12. A construct comprising a promoter sequence flanked by two sets of oppositely oriented coding sequences each encoding a viral accessory protein or a recombinant viral vector genome, wherein said promoter is capable of being inverted upon a recombination event inducible by a recombinase, wherein said promoter is operably linked to a first set of said two sets of oppositely oriented coding sequences prior to recombination and
15 becomes operably linked to a second set of said two sets of oppositely oriented coding sequences post recombination.
13. The construct of claim 12, wherein said promoter is flanked by two recombinase recognition sites in opposite orientations.
- 20 14. A construct comprising a promoter sequence operably linked to one or more coding sequences each encoding a viral accessory protein, a recombinant viral vector genome or both, wherein at least one of said one or more coding sequences are flanked by two recombinase recognition sites, wherein said at least one of said one or more coding
25 sequences are capable of being disrupted, deleted or inverted upon a recombination event inducible by a recombinase.
15. A construct comprising (i) a promoter sequence operably linked to a first set of one or more coding sequences each encoding a viral accessory protein, a recombinant viral
30 vector genome or both, and (ii) two recombinase recognition sites flanking said promoter sequence and said first set of one or more coding sequences, wherein, upon a recombination event inducible by a recombinase, said promoter sequence becomes operably linked to additional coding sequences besides said first set of one or more coding sequences.

16. The construct of claim 9 or 12, wherein said viral accessory protein or recombinant viral vector genome is from a virus selected from the group consisting of lentivirus, retrovirus, herpesvirus, adenovirus, and adeno-associated virus.
- 5 17. The construct of claim 9 or 12, wherein said first and second sets of one or more copies of coding sequences are capable of providing different levels of expression of said coding sequences depending on an orientation of said promoter.
- 10 18. The construct of claim 9 or 12, wherein said first and second sets of one or more copies of coding sequences comprise different copy numbers.
19. The construct of claim 1 or 12, wherein said promoter comprises an inducible, recombined, or edited promoter.
- 15 20. The construct of claim 1 or 13, wherein said recombinase recognition sites comprise two palindromic recognition regions flanking a spacer region.
21. The construct of claim 1 or 13, wherein each of said recombinase recognition sites comprises a lox site.
- 20 22. The construct of claim 21, wherein said lox site comprises a wildtype lox site or a mutant lox site.
- 25 23. The construct of claim 20, wherein said spacer region is selected from the group consisting of loxP, lox511, lox2272, lox5171, m2, m3, and m7.
24. The construct of any one of claims 1 to 23, wherein said recombinase is a Cre-recombinase.
- 30 25. The construct of claim 1 or 13, wherein each of said recombinase recognition sites are a Flippase recombinase target sequence.
26. The construct of claim 25, wherein said recombinase is a Flippase recombinase.

27. The construct of claim 1 or 12, wherein said first set of said one or more copies of coding sequences comprises two or more copies of said coding sequences arranged as a polycistronic expression cassette.
- 5
28. The construct of claim 27, wherein said polycistronic expression cassette is in the sense orientation downstream of said promoter prior to promoter inversion.
29. The construct of claim 27, wherein said polycistronic expression cassette further
- 10 comprises one or more viral skip sequences, internal ribosome entry site elements, or both.
30. The construct of claim 29, wherein said viral skip sequences are selected from the group consisting of P2A, T2A, E2A, and F2A.
- 15
31. The construct of claim 9 or 12, wherein said second set of said one or more copies of coding sequences comprises a monocistronic expression cassette.
32. The construct of claim 31, wherein said monocistronic expression cassette is in the
- 20 antisense orientation upstream of said promoter prior to promoter inversion.
33. The construct of any of claims 1 to 32, wherein said viral accessory protein is a fusion protein.
- 25
34. The construct of any of claims 1 to 32, wherein said viral accessory protein comprises a sequence encoding a structural viral protein, a regulatory viral protein, or both.
35. The construct of claim 34, wherein said structural and regulatory proteins are selected from the group consisting of Gag, Pol, Rev, Env, Tat, Nef, Vpr, Vif, Vpu, and Vpx.
- 30
36. The construct of any of claims 1 to 32, wherein said viral accessory protein comprises a sequence encoding one or more viral accessory protein domains.

37. The construct of claim 36, wherein said one or more viral accessory protein domains are selected from the group consisting of CA, MA, NC, p6, SP1, RT, IN, PR, and DU.
38. The construct of any one of claims 1 to 37, wherein a sequence encoding said viral
5 accessory protein or one or more viral accessory protein domains comprise a wild-type sequence, a codon optimized sequence, or both
39. A construct comprising a promoter sequence operably linked to two or more coding sequences each encoding a viral accessory protein, a recombinant viral vector genome
10 or both, wherein at least one of said two or more coding sequences is codon optimized.
40. A cell comprising the construct of any of claims 1 to 39.
41. The cell of claim 40, wherein said cell is a eukaryotic cell.
- 15 42. The cell of claim 40, wherein said cell is a mammalian cell.
43. The cell of claim 40, wherein said cell is a human cell.
- 20 44. The cell of claim 40, wherein said cell is a viral vector producer cell.
45. The cell of claim 40, wherein said construct is integrated in the genome of said cell.
46. The cell of claim 44, wherein said viral vector producer cell is adherent or in
25 suspension.
47. The cell of claim 44, wherein said viral vector producer cell is cultured in serum-supplemented or serum-free media.
- 30 48. The cell of claim 44, wherein said viral vector producer cell is immortalized.
49. The cell of claim 44, wherein said viral vector producer cell is a HEK293 cell, or a derivative thereof.

50. The cell of claim 49, wherein said HEK293 cell is a HEK293T cell.
51. A method of producing a viral vector producer cell with an optimized stoichiometric ratio of viral vector genome and viral accessory proteins, wherein said method
5 comprises:
- a. introducing the construct of any of claims 1 to 39 into a first clonal population of cells;
 - b. transiently providing a recombinase and/or a CRISPR-based complex to said first clonal population; and
 - 10 c. producing a second clonal population by (i) inverting an invertible sequence flanked by recombinase recognition sites; and/or (ii) editing one or more copies of coding sequences each encoding a viral accessory protein, a recombinant viral vector genome or both.
- 15 52. The method of claim 51, wherein said second clonal population comprises a viral vector genome construct and one or more accessory constructs encoding one or more viral accessory proteins.
53. The method of claim 51, wherein said method further comprises generating from said
20 second clonal population a stable viral vector producer cell line.
54. The method of claim 51, wherein said invertible sequence comprises a promoter sequence, a viral accessory protein sequence, a recombinant viral vector genome sequence, or a combination thereof.
- 25 55. The method of claim 53, wherein said stable viral vector producer cell line exhibits a higher viral titer relative to said first clonal population.
56. The method of claim 55, wherein said viral titer is determined by physical titration,
30 functional titration, or both.
57. The method of claim 55, wherein said viral titer is determined by assaying for viral nucleic acid via an assay selected from the group consisting of PCR, RT-PCR, and

quantitative detection by blot hybridization, or by assaying for viral protein via an immunoassay.

58. The method of claim 51, wherein said method further comprises determining the stoichiometric ratio of viral vector genome RNA and said viral accessory proteins in said first clonal population, second clonal population, or both.
59. The method of claim 51, wherein said method further comprises quantifying the level of said viral vector genome and said viral accessory proteins in said first clonal population, second clonal population, or both.
60. The method of claim 53, wherein said method further comprises storing said stable viral vector producer cell line by cryopreservation.
61. The method of claim 53, wherein said method further comprises expanding cells from said cryopreserved cell line to produce viral vectors.
62. The method of claim 51, wherein said recombinase recognition sites comprise two palindromic recognition regions flanking a spacer region.
63. The method of claim 51, wherein each of said recognition sites comprises a lox site.
64. The method of claim 63, wherein said lox site comprises a wildtype lox site or a mutant lox site.
65. The method of claim 62, wherein said spacer region is selected from the group consisting of loxP, lox511, lox2272, lox5171, m2, m3, and m7.
66. The method of claim 51, wherein said recombinase is a Cre-recombinase.
67. The method of claim 51, wherein each of said recombinase recognition sites is a Flippase recombinase target sequence.
68. The method of claim 51, wherein said recombinase is a Flippase recombinase.

69. The method of claim 51, wherein said viral vector genome construct comprises one or more elements selected from the group consisting of a 5' long terminal repeat, a 3' long terminal repeat, a packaging signal, and a central polypurine tract.
- 5
70. The method of claim 51, wherein said viral vector genome construct does not comprise a 5' long terminal repeat, a 3' long terminal repeat, a packaging signal, or a central polypurine tract.
- 10
71. The method of claim 51, wherein said viral vector genome construct comprises a self-inactivating long terminal repeat.
72. The method of claim 51, wherein said viral vector genome construct comprises a promoter and a polyadenylation sequence.
- 15
73. The method of claim 72, wherein said promoter of said viral vector genome construct is constitutive or inducible.
74. The method of claim 51, wherein said viral vector genome construct comprises a concatemer.
- 20
75. The method of claim 74, wherein said concatemer comprises multiple copies of a viral vector genome construct.
- 25
76. The method of claim 74, wherein said concatemer comprises one or more selection genes.
77. The method of claim 74, wherein said concatemer comprises one or more transcription factors.
- 30
78. The method of claim 51, wherein said first clonal population or said second clonal population comprises viral vector producer cells.

79. The method of claim 51, wherein said introducing step may be chemical, biological or physical.
80. The method of claim 51, wherein said introducing step comprises optical, magnetic,
5 biolistic, polymer-based, liposome-based, nanoparticle-based methods, or a combination thereof.
81. The method of claim 51, wherein said introducing step comprises a transduction.
- 10 82. The method of claim 51, wherein said introducing step comprises a transfection.
83. The method of claim 79, wherein said chemical introducing step comprises a cationic polymer, calcium phosphate, cationic lipid, or a combination thereof.
- 15 84. The method of claim 79, wherein said biological introducing step comprises introduction via retrovirus, lentivirus, transposon, CRISPR/Cas9, or recombinase.
85. The method of claim 79, wherein said physical introducing step is selected from the group consisting of electroporation, sonoporation, mechanoporation, and
20 photoporation.
86. The method of claim 51, wherein said recombinase is an inducible recombinase.
87. The method of claim 51, wherein said recombinase comprises a Cre-recombinase, a
25 Flippase recombinase, or a modified version thereof.
88. The construct of claim 86, wherein said inducible recombinase is chemical or light inducible.
- 30 89. The method of claim 51, wherein said cell is a eukaryotic cell.
90. The method of claim 51, wherein said cell is a mammalian cell.
91. The method of claim 51, wherein said cell is a human cell.

92. The method of claim 51, wherein said promoter comprises an inducible, recombined, or edited promoter.
- 5 93. The method of claim 51, wherein said viral accessory proteins comprise sequences encoding structural viral proteins, regulatory viral proteins, or both.
94. The method of claim 51, wherein said structural and regulatory proteins are selected from the group consisting of Gag, Pol, Rev, Env, Tat, Nef, Vpr, Vif, Vpu, and Vpx.
- 10 95. The method of claim 51, wherein said viral accessory proteins comprise sequences encoding a partial viral accessory protein.
96. The method of claim 51, wherein said partial viral accessory protein comprises one or
15 more viral accessory protein domains.
97. The method of claim 96, wherein said one or more viral accessory protein domains is selected from the group consisting of CA, MA, NC, p6, SP1, RT, IN, PR, and DU.
- 20 98. The method of any one of claims 93 to 97, wherein a sequence encoding said one or more viral accessory proteins or domains comprises a wild-type sequence, a codon optimized sequence, or both.
99. A method comprising:
- 25 a. obtaining a first stable viral vector producer cell line;
- b. determining a viral titer from said first stable viral vector producer cell line; and
- c. engineering one or more elements in said stable viral vector producer cell line to produce a second stable viral vector producer cell line with a stoichiometric
30 ratio of viral vector genome and one or more viral accessory proteins different from the corresponding stoichiometric ratio in said first stable viral vector producer cell line.

100. The method of claim 99, wherein said engineering is via one or more recombination events.

5 101. The method of claim 99, wherein said engineering comprises introducing a recombinase.

102. The method of claim 99, wherein said engineering comprises transiently expressing a recombinase.

10 103. The method of claim 99, wherein said engineering is via gene editing.

104. The method of claim 99, wherein said engineering comprises introducing a CRISPR-based complex.

15 105. The method of claim 99, wherein said engineering comprises introducing a CRISPR-based complex specific to some but not all copies of coding sequences for said one or more viral accessory proteins.

20 106. The method of claim 99, wherein said engineering comprises both a recombination event and gene editing.

25