CLAIMS

1. Use of ~~A~~an immunoconjugate comprising an anti-CEACAM5-antibody for the manufacture of a medicament for use ~~for treating cancer~~ in combination with folinic acid, 5-fluoro-uracil and oxaliplatin (FOLFOX) for treating cancer, wherein the anti-CEACAM5-antibody comprises a CDR-H1 consisting of SEQ ID NO: 1, CDR-H2 consisting of SEQ ID NO: 2, CDR-H3 consisting of SEQ ID NO: 3, CDR-L1 consisting of SEQ ID NO: 4, CDR-L2 consisting of amino acid sequence NTR, and CDR-L3 consisting of SEQ ID NO: 5.

~~2. The immunoconjugate for the use of claim 1, wherein the anti-CEACAM5-antibody comprises a CDR-H1 consisting of SEQ ID NO: 1, CDR-H2 consisting of SEQ ID NO: 2, CDR-H3 consisting of SEQ ID NO: 3, CDR-L1 consisting of SEQ ID NO: 4, CDR-L2 consisting of amino acid sequence NTR, and CDR-L3 consisting of SEQ ID NO: 5.~~

~~3.~~2. The ~~immunoconjugate for the~~ use of claim 1 ~~or 2~~, wherein the anti-CEACAM5-antibody comprises a variable domain of a heavy chain (VH) consisting of SEQ ID NO: 6 and a variable domain of a light chain (VL) consisting of SEQ ID NO: 7.

~~4.~~3. The ~~immunoconjugate for the~~ use of ~~any of~~ claim~~s~~ 1 ~~to~~ or ~~2~~3, wherein the anti-CEACAM5-antibody comprises a heavy chain (VH) consisting of SEQ ID NO: 8 and a light chain (VL) consisting of SEQ ID NO: 9.

~~5.~~4. The ~~immunoconjugate for the~~ use of any of claims 1 to ~~4~~3, wherein the immunoconjugate comprises at least one cytostatic agent.

~~6.~~5. The ~~immunoconjugate for the~~ use of claim ~~5~~4, wherein the cytostatic agent is selected from the group consisting of radioisotopes, protein toxins, small molecule toxins, and combinations thereof.

~~7.~~6. The ~~immunoconjugate for the~~ use of claim ~~6~~5, wherein the small molecule toxins are selected from antimetabolites, DNA-alkylating agents, DNA-cross-linking agents, DNA-intercalating agents, anti-microtubule agents, topoisomerase inhibitors, and combinations thereof.

8.7. The ~~immunoconjugate for the~~ use of claim ~~7~~6, wherein the anti-microtubule agent is selected from the group consisting of taxanes, vinca alkaloids, maytansinoids, colchicine, podophyllotoxin, gruseofulvin, and combinations thereof.

9.8. The ~~immunoconjugate for the~~ use of claim ~~8~~7, wherein the maytansinoids are selected from the group consisting of N2'-deacetyl-N2'-(3-mercapto-1-oxopropyl)-maytansine (DM1) or N2'-deacetyl-N-2'(4-methyl-4-mercapto-1-oxopentyl)-maytansine (DM4), and combinations thereof.

10.9. The ~~immunoconjugate for the~~ use of any of claims 1 to ~~9~~8, wherein the anti-CEACAM5-antibody is covalently attached via a cleavable or non-cleavable linker to the at least one cytotoxic agent.

11.10. The ~~immunoconjugate for the~~ use of claim ~~10~~9, wherein said linker is selected from the group consisting of N-succinimidyl pyridyldithiobutyrate (SPDB), 4-(pyridin-2-yldisulfanyl)-2-sulfo-butyric acid (sulfo-SPDB), and succinimidyl(N-maleimidomethyl) cyclohexane-1-carboxylate (SMCC).

12.11. The ~~immunoconjugate for the~~ use of any of claims 1 to ~~11~~10, comprising an CEACAM5-antibody, which comprises a heavy chain (VH) consisting of SEQ ID NO: 8 and a light chain (VL) consisting of SEQ ID NO: 9 (huMAb2-3), and which is covalently linked to N2'-deacetyl-N-2'(4-methyl-4-mercapto-1-oxopentyl)-maytansine (DM4) via N-succinimidyl pyridyldithiobutyrate (SPDB).

13.12. The ~~immunoconjugate for the~~ use of any of claims 1 to ~~12~~11, wherein the immunoconjugate is characterised by a drug-to-antibody ratio (DAR) ranging from 1 to 10.

14.13. The ~~immunoconjugate for the~~ use of any of claims 1 to ~~13~~12, wherein the cancer is selected from the group consisting of colorectal, stomach, pancreas, and oesophagus cancer.

~~15.~~14. The ~~immunoconjugate for the~~ use of any of claims 1 to ~~14~~13, wherein the immunoconjugate and FOLFOX are administered simultaneously to a subject in need thereof.

~~16.~~15. The ~~immunoconjugate for the~~ use of claim ~~15~~14, wherein the immunoconjugate and FOLFOX are formulated (i) in a single pharmaceutical composition comprising the immunoconjugate and FOLFOX, or (ii) in the form of at least two separate pharmaceutical compositions, wherein at least one pharmaceutical composition comprises the immunoconjugate, and one or more pharmaceutical compositions comprise folinic acid, 5-fluoro-uracil and oxaliplatin, in separate or combined formulations.

~~17.~~16. The ~~immunoconjugate for the~~ use of any of claims 1 to ~~14~~13, wherein the immunoconjugate and FOLFOX are administered separately or sequentially to a subject in need thereof.

~~18.~~17. The ~~immunoconjugate for the~~ use of claim ~~17~~16, wherein the immunoconjugate and FOLFOX are formulated in the form of at least two separate pharmaceutical compositions, wherein (i) at least one pharmaceutical composition comprises the immunoconjugate, and (ii) one or more pharmaceutical compositions comprise folinic acid, 5-fluoro-uracil and oxaliplatin, in separate or combined formulations.

~~19.~~18. The ~~immunoconjugate for the~~ use of any of claims 1 to ~~18~~17, wherein the immunoconjugate comprising an anti-CEACAM5-antibody, and folinic acid, 5-fluoro-uracil and oxaliplatin (FOLFOX) are administered in 8 to 16 cycles, wherein one cycle comprises:

administering the immunoconjugate at a dose of from 60 to 210 mg/m^2, at least once in the cycle;

administering folinic acid at a dose of from 100 to 300 mg/m^2 or L-folinic acid at a dose of 100 to 200 m/m^2, at least once in the cycle;

administering 5-fluoro-uracil at a dose of from 1000 to 2000 mg/m^2, at least once in the cycle, and

administering oxaliplatin at a dose of from 50 to 200 mg/m^2, at least once in the cycle.

~~20.~~19. A pharmaceutical composition comprising the immunoconjugate of any of claims 1 to ~~14~~13, and folinic acid, 5-fluoro-uracil and oxaliplatin.

~~21.~~20. A kit comprising (i) a pharmaceutical composition of the immunoconjugate of any of claims 1 to ~~14~~ 13and (ii) one or more pharmaceutical compositions comprising folinic acid, 5-fluoro-uracil and oxaliplatin, in separate or combined formulations.

~~22.~~21. The pharmaceutical composition according to claim ~~20~~ 19 or the kit according to claim ~~21~~ 20 for the use for treating cancer.

22. Use of an immunoconjugate comprising an anti-CEACAM5-antibody for the manufacture of a medicament for use in combination with folinic acid, 5-fluoro-uracil and oxaliplatin (FOLFOX), for treating cancer, wherein the anti-CEACAM5-antibody comprises a heavy chain (VH) consisting of SEQ ID NO: 8 and a light chain (VL) consisting of SEQ ID NO: 9
wherein the immunoconjugate comprising an anti-CEACAM5-antibody, and folinic acid, 5-fluoro-uracil and oxaliplatin (FOLFOX) are administered in 8 to 16 cycles, wherein one cycle comprises:
administering the immunoconjugate at a dose of from 60 to 210 mg/m^2, at least once in the cycle;
administering folinic acid at a dose of from 100 to 300 mg/m^2 or L-folinic acid at a dose of 100 to 200 m/m^2, at least once in the cycle;
administering 5-fluoro-uracil at a dose of from 1000 to 2000 mg/m^2, at least once in the cycle, and
administering oxaliplatin at a dose of from 50 to 200 mg/m^2, at least once in the cycle,
the cycle being selected from a 1-week cycle, a 2-week cycle or a 3-week cycle.