METHOD OF DRUG DELIVERY FOR BONE ANABOLIC PROTEIN

RELATED APPLICATION

This application claims the benefit of U.S. Provisional Application No. 60/848,960, filed on October 3, 2006. The entire teachings of the above application is incorporated herein by reference.

BACKGROUND OF THE INVENTION

Parathyroid hormone-related protein ("PTHrP") is a 139 to 173 amino acid-protein. PTHrP and certain analogs are known to be useful to improve bone mass and quality in the treatment of osteoporosis and related disorders. However, the commercial use of these proteins as pharmaceutical agents requires the development of a formulation that is acceptable in terms of storage stability and ease of preparation.

Furthermore, currently available osteoporosis drugs have limitations on suitable dosage ranges due to the unwanted side-effects, such as hypercalcemia and increased stimulation of bone resorption. These unwanted side-effects and resulting dose limitations reduce the beneficial effects which can be achieved from these drugs. Thus a need exists for compounds which can be administered at a dose which will increase the beneficial effects without an increase in the unwanted side-effects.

SUMMARY OF THE INVENTION

The present invention provides a storage-stable composition containing a parathyroid hormone-related protein (PTHrP) analogue and methods of using those analogues and compositions containing those analogues as described herein to treat osteoporosis, to increase bone mass or to increase bone quality. The composition is storage stable, in sterile form, and in general may be stored at room temperature for at least several weeks to allow convenient parenteral administration to human patients.

In one embodiment, the present invention provides a storage-stable composition suitable for administration to a subject (e.g., a human). The

composition comprises a PTHrP analogue and an effective amount of buffer to maintain the pH of the composition between 2 and 7. In a particular embodiment, the PTHrP is [Glu^{22,25}, Leu^{23,28,31}, Aib^{29}, Lys^{26,30}]hPTHrP(1-34)NH_2 (SEQ ID NO.: 2).

In another embodiment, the present invention provides a sealed container containing a storage-stable composition suitable for administration to a subject. The composition comprises PTHrP or an analog thereof and an effective amount of buffer to maintain the pH of the composition between 2 and 7. In a particular embodiment, the PTHrP analogue is [Glu^{22,25}, Leu^{23,28,31}, Aib^{29}, Lys^{26,30}]hPTHrP(1-34)NH_2 (SEQ ID NO.: 2).

In another embodiment, the present invention provides a drug delivery device comprising one or more than one single-use container which comprises a storage stable composition comprising PTHrP or an analog thereof and an effective amount of buffer to maintain the pH of the composition between 2 and 7. In a particular embodiment, the PTHrP analogue is [Glu^{22,25}, Leu^{23,28,31}, Aib^{29}, Lys^{26,30}]hPTHrP(1-34)NH_2 (SEQ ID NO.:2).

In another embodiment, the present invention provides a drug delivery device comprising one or more than one multi-use container, which comprises a storage stable composition comprising PTHrP or an analog thereof and an effective amount of buffer to maintain the pH of the composition between 2 and 7. In a particular embodiment, the PTHrP analogue is [Glu^{22,25}, Leu^{23,28,31}, Aib^{29}, Lys^{26,30}]hPTHrP(1-34)NH_2 (SEQ ID NO.: 2).

In another embodiment the present invention provides a method of treating osteoporosis in a subject in need thereof comprising administering to the subject a single daily subcutaneous dose of [Glu^{22,25}, Leu^{23,28,31}, Aib^{29}, Lys^{26,30}]hPTHrP(1-34)NH_2 (SEQ ID NO.: 2) in an amount between 40 and 160 μg for a duration of time sufficient to treat the subject, typically between about 3 months to 36 months. In some embodiments, the treatment period is between about 3 months to 18 months.

In another embodiment the present invention provides a method of increasing bone mass or increasing bone quality in a subject in need thereof comprising administering to the subject a single daily subcutaneous dose of [Glu^{22,25}, Leu^{23,28,31}, Aib^{29}, Lys^{26,30}]hPTHrP(1-34)NH_2 (SEQ ID NO.: 2) in an

amount between 40 and 160 μg for a duration of time sufficient to treat the subject, typically between 3 months and 36 months. In some embodiments, the treatment period is between about 3 months to 18 months.

The PTHrP and analogue compositions of the invention exhibit storage stability in terms of hormone composition and activity. Furthermore, these compositions can be administered, in general, in higher dosages than currently available osteoporosis drugs, with the reduction or elimination of unwanted side-effects, such as, hypercalcemia or stimulation of bone resorption. This has the advantage of an increase in beneficial physiological effects due to the increased dosages and can result in a reduction in the length of treatment time.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a graph showing the stability of SEQ ID NO. 2 over 24 months at 5 °C and 25 °C without any chemical stabilizer.

FIG. 2 is a graph showing the stability of lyophilized SEQ ID NO. 2 over 24 months at 5 °C 25 °C and 40 °C.

DETAILED DESCRIPTION OF THE INVENTION

The sequence of native hPTHrP (1-34) is as follows:

Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile Gln Asp Leu Arg Arg Arg Phe Phe Leu His His Leu Ile Ala Glu Ile His Thr Ala (SEQ ID NO:1).

In a particular embodiment, the PTHrP analogue is [$Glu^{22,25}$, $Leu^{23,28,31}$, Aib^{29}, $Lys^{26,30}$]hPTHrP(1-34)NH_2 (SEQ ID NO.: 2).

Other PTHrP analogues are described in 6,921,750, 5,955,574, 6,544,949, 5, 723,577, and 5,696,095 the entire contents of each of which are incorporated herein by reference.

A "buffer" as used herein is any acid or salt combination which is pharmaceutically acceptable and capable of maintaining the composition of the present invention within a desired pH range. Buffers in the disclosed compositions maintain the pH in a range of about 2 to about 7, about 3 to about 6, about 4 to about 6, about 4.5 to about 5.6, or about 5.1. Suitable buffers include, any pharmaceutical acceptable buffer capable of maintaining the above pH ranges, such as, for example,

acetate, tartrate phosphate or citrate buffers. In one embodiment, the buffer is an acetate or tartrate buffer. In another embodiment the buffer is an acetate buffer. In one embodiment the buffer is acetic acid and sodium acetate.

In the disclosed compositions the concentration of buffer is typically in the range of about 0.1 mM to about 1000 mM, about 0.2 mM to about 200 mM, about 0.5 mM to about 50 mM, about 1 mM to about 10 mM or about 6 mM.

As used herein, an anti-microbial agent is a pharmaceutically acceptable preservative, suitable for administration to a subject, which inhibits, prevents or delays the growth or micro organisms including, for example bacteria, viruses and fungi in the compositions of the present invention. Suitable anti-microbial agents for use in the compositions and methods of the present invention include, but are not limited to, cresols, benzyl alcohol, phenol, benzalkonium chloride, benzethonium chloride, chlorobutanol, phenylethyl alcohol, methyl paraben, propyl paraben, thiomersal and phenylmercuric nitrate and acetate. In one embodiment the anti-microbial agents is m-cresol, chlorocresol or phenol. In another embodiment the anti-microbial agents is chlorocresol or phenol. In another embodiment the anti-microbial agents is phenol.

As used herein an effective amount of an anti-microbial agent is an amount effective to inhibits, prevents or delays the growth or micro organisms including, for example bacteria, viruses and fungi in the compositions of the present invention. In the compositions of the present invention, the amount of anti-microbial agent is typically in the range from about 0.1 to about 20 mg/ml, about 0.2 to about 30 mg/ml, about 0.2 to about 10 mg/ml, about 0.25 to about 5 mg/ml, about 0.5 to about 50 mg/ml, about 1 to about 10 mg/ml, about 3 mg/ml or about 5 mg/ml.

The compositions of the present invention typically are ready to administer, aqueous solutions which are sterile, storage-stable and pharmaceutically acceptable without the need for reconstitution prior to administration. The compositions of the present invention are suitable for administration to a subject which means that they are pharmaceutically acceptable, non-toxic, do not contain any components which would adversely affect the biological or hormonal effects of the peptide. The compositions of the present invention do not, for example, comprise any cells.

As used herein a composition of the present invention is storage-stable if the amount, purity of the PTHrP remains above about 95% of the original amount under one of the following conditions: (1) storage for over 2 years at 5 °C; or (2) storage for over 30 days at 25 °C.

The compositions are typically stored in a sealed container, vial or cartridge which is typically suitable for long term storage. "Suitable for long-term storage" means that the vial, container or cartridge does not allow for the escape of components of the compositions of the present invention or the ingress of external components, such as, micro organisms when kept for at least 3 months at 25 °C.

The compositions of the present invention are preferably administered by injection, typically subcutaneous injection.

The compositions of the present invention, can be stored in single-dose or multi-dose sealed containers, vials or cartridges. The sealed container, vial or cartridge is typically suitable for use with a single or multi-dose injection pen or drug delivery device, which typically allows the patient to administer the peptide themselves. The sealed container can comprise one or more doses of the peptide of the present invention, wherein each dose comprises an effective amount of the peptide as described herein.

A single-dose injection pen, or drug delivery device is typically a disposable device which uses a sealed container which comprises a single dose of an effective amount of a PTHrP in the compositions described herein. A multi-dose injection pen or drug delivery device typically contains more than one dose of an effective amount of a PTHrP thereof in the compositions described herein. The multi-dose pen can typically be adjusted to administer the desired volume of the storage stable compositions described herein. In certain embodiment the multi-dose injection pen prevents the ingress of microbial contaminants from entering the container or cartridge which can occur through multiple uses of one needle.

Injection pens, as used herein, can also comprise two containers one of which contains a PTHrP, as described herein, in a lyophilized powder, as described below, and the second container contains a liquid for reconstitution of the lyophilized powder. The contents of the two containers can be mixed prior to administration.

As discussed above the compositions of the present invention can be administered by injection. Suitable volumes of the compositions of the present invention for injection include about 0.5 to about 1 ml, about 0.1 to about 1 ml, about 0.02- to about 0.04 ml, about 0.1- to about 5.0 μl, or about 0.1- to about 1.0 μl.

In the compositions of the present invention the concentration of the peptides is from about 20 mg/ml to about 20,000 mg/ml, from about 100 mg/ml to about 10,000 mg/ml, from about 300 mg/ml to about 300 mg/ml, from about 500 mg/ml to about 2000 mg/ml and about 2 mg/ml.

The compositions of the present invention can also be lyophilized using lyophilization techniques known in the art and stored as a powder which can be reconstituted prior to administration. The term "lyophilization" as used herein is a freeze drying or dehydration technique which involves removing a solvent, preferably a water miscible solvent, more preferably water from a composition or the present invention, typically by sublimation under high vacuum when the composition is in a frozen state. Typically, lyophilization is carried out in lyophilization equipment (a lyophilizer), which comprises a drying chamber with variable temperature controls, a condenser to collect water, and a vacuum system to reduce the pressure in the drying chamber.

The terms "lyophilized composition", as used herein mean the solid residue or powder which is produced or which remains after the lyophilization procedure as defined above. The lyophilized composition of the present invention typically further comprise a pharmaceutically acceptable excipient. The term "pharmaceutically acceptable excipient" as used herein refers to a substance which is added to a solution prior to lyophilization to enhance characteristics such as the color, texture, strength, and volume of the lyophilized cake. Pharmaceutically acceptable excipients may be, for example, buffers and pH adjusters, crystalline bulking excipients, stabilizers, and tonicity raising agents.

In certain preferred embodiments the pharmaceutically acceptable excipient is a crystalline bulking excipient. The terms "crystalline bulking excipient" or "crystalline bulking agent" as used herein means an excipient which provides bulk and structure to the lyophilization cake. These crystalline bulking agents are inert

and do not react with the peptide. In addition, the crystalline bulking agents are capable of crystallizing under lyophilization conditions.

Examples of suitable crystalline bulking agents include hydrophilic excipients, such as, water soluble polymers; sugars, such as mannitol, sorbitol, xylitol, glucitol, ducitol, inositiol, arabinitol, arabitol, galactitol, iditol, allitol, maltitol, fructose, sorbose, glucose, xylose, trehalose, allose, dextrose, altrose, lactose, glucose, fructose, gulose, idose, galactose, talose, ribose, arabinose, xylose, lyxose, sucrose, maltose, lactose, lactulose, fucose, rhamnose, melezitose, maltotriose, raffinose, altritol, their optically active forms (D- or L-forms) as well as the corresponding racemates; inorganic salts, both mineral and mineral organic, such as, calcium salts, such as the lactate, gluconate, glycerylphosphate, citrate, phosphate monobasic and dibasic, succinate, sulfate and tartrate, as well as the same salts of aluminum and magnesium; carbohydrates, such as, the conventional mono- and di-saccharides as well as the corresponding polyhydric alcohols; proteins, such as, albumin; amino acids, such as glycine; emulsifiable fats and polyvinylpyrrolidone. Preferred crystalline bulking agents are selected from the group consisting of glycine, mannitol, dextran, dextrose, lactose, sucrose, polyvinylpyrrolidone, trehalose, glucose and combinations thereof. Particularly useful bulking agents include dextran.

As used herein a stabilizer is a composition which maintains the chemical, biological or hormonal stability of the peptide. Examples of stabilizing agent include polyols which includes a saccharide, preferably a monosaccharide or disaccharide, e.g., glucose, trehalose, raffinose, or sucrose; a sugar alcohol such as, for example, mannitol, sorbitol or inositol, a polyhydric alcohol such as glycerine or propylene glycol or mixtures thereof and albumin.

The compositions described herein can be used to stimulate bone growth in a subject. Thus they are useful in the treatment of diseases or disorders associated with deficiency in bone growth such as osteoporosis and bone fractures. In one embodiment, the present invention is a method of treating osteoporosis in a subject comprising administering to the subject an effective amount of composition described herein.

As used herein, "treating" can include both prophylactic, and therapeutic treatment. For example, therapeutic treatment can include delaying inhibiting or preventing the progression of osteoporosis, the reduction or elimination of symptoms associated with osteoporosis. Prophylactic treatment can include preventing, inhibiting or delaying the onset of osteoporosis.

As used herein, an effective amount refers to an amount sufficient to elicit the desired response. In the present invention, the desired biological response is an decrease in the rate of bone loss and/or an increase in the bone mass or bone quality of a subject.

Suitable dosage for use in the compositions and methods of the present invention include from about 40 to about 160 μg, about 80 to about 120 μg about 80 to about 100 μg; or from about 40 to about 50 μg, about 50 to about 60 μg, about 60 to about 70 μg, about 70 to about 80 μg, about 80 to about 90 μg, about 90 to about 100 μg, about 100 to about 110 μg, about 110 to about 120 μg, about 120 to about 130 μg, about 130 to about 140 μg, about 140 to about 150 μg, about 150 to about 160 μg; or from 40 to about 45 μg, about 45 to about 50 μg, about 50 to about 55 μg, about 55 to about 60 μg, about 60 to about 65 μg, about 65 to about 70 μg, about 70 to about 75 μg, about 75 to about 80 μg, about 80 to about 85 μg, about 85 to about 90 μg, about 90 to about 95 μg, about 95 to about 100 μg, about 100 to about 105 μg, about 105 to about 110 μg, about 110 to about 115 μg, about 115 to about 120 μg, about 120 to about 125 μg, about 125 to about 130 μg, about 130 to about 135 μg, about 135 to about 140 μg, about 140 to about 145 μg, about 145 to about 150 μg, about 150 to about 155 μg, about 155 to about 160 μg administered once per day, once every other day, twice per week once per week, once every two weeks, once per month. The doses can be a pulsatile injection, for example, once per month which causes pulsatile release of singles doses of the composition described herein.

When the dosages described above are administered once per day, once per week etc., typically the dosages are of equal amounts.

The subject as used herein can be an animal, for example, a mammal, such as a human.

A pharmaceutically acceptable salt is a salt which is suitable for administration to a subject, such as, a human. The peptides of the present invention

can have one or more sufficiently acidic proton that can react with a suitable organic or inorganic base to form a base addition salt. Base addition salts include those derived from inorganic bases, such as ammonium or alkali or alkaline earth metal hydroxides, carbonates, bicarbonates, and the like, and organic bases such as alkoxides, alkyl amides, alkyl and aryl amines, and the like. Such bases useful in preparing the salts of this invention thus include sodium hydroxide, potassium hydroxide, ammonium hydroxide, potassium carbonate, and the like. The peptides of the present invention having a sufficiently basic group, such as an amine can react with an organic or inorganic acid to form an acid addition salt. Acids commonly employed to form acid addition salts from compounds with basic groups are inorganic acids such as hydrochloric acid, hydrobromic acid, hydroiodic acid, sulfuric acid, phosphoric acid, and the like, and organic acids such as p-toluenesulfonic acid, methanesulfonic acid, oxalic acid, p-bromophenyl-sulfonic acid, carbonic acid, succinic acid, citric acid, benzoic acid, acetic acid, and the like. Examples of such salts include the sulfate, pyrosulfate, bisulfate, sulfite, bisulfite, phosphate, monohydrogenphosphate, dihydrogenphosphate, metaphosphate, pyrophosphate, chloride, bromide, iodide, acetate, propionate, decanoate, caprylate, acrylate, formate, isobutyrate, caproate, heptanoate, propiolate, oxalate, malonate, succinate, suberate, sebacate, fumarate, maleate, butyne-1,4-dioate, hexyne-1,6-dioate, benzoate, chlorobenzoate, methylbenzoate, dinitrobenzoate, hydroxybenzoate, methoxybenzoate, phthalate, sulfonate, xylenesulfonate, phenylacetate, phenylpropionate, phenylbutyrate, citrate, lactate, gamma-hydroxybutyrate, glycolate, tartrate, methanesulfonate, propanesulfonate, naphthalene-1-sulfonate, naphthalene-2-sulfonate, mandelate, and the like.

The compositions of the present invention typically do not show any or show reduced side-effects such as hypercalcemia and typically do not increase the stimulation of bone resorption at the dosage listed above. This reduction in side effects allows for administration of higher doses than commercially available osteoporosis drugs.

The compositions of the present invention can be administered by injection as described herein.

The compositions of the present invention may be administered alone or in combination with an additional therapeutic agent, such as an antiresorptive therapy, for example, bisphonsphonates and calcitonin.

EXEMPLIFICATION

EXAMPLE 1 demonstrates $Glu^{22,25}$, $Leu^{23,28,31}$, Aib^{29}, $Lys^{26,30}$]hPTHrP(1-34)NH_2 (SEQ ID NO.: 2) stability at low acetate concentration (1mM), without stabilizer

TABLE 1

Material	Supplier	Unitary Formula (per cartridge)
(SEQ ID NO.: 2)	Ipsen Ireland	0.140 mg (free base)
Tri-hydrate sodium acetate 0.1 N	Prolabo	14.6 mg
Acetic acid 0.1 N	Prolabo	1 .9 mg qs pH 5.1
Water for Injection	Meram	qs 1.4 g
Type I clear glass Cartridge 1.5 ml, washed, siliconised and sterilised	Bünderglass via Vetter	1
Grey PTFE bromobutyl cartridge rubber stopper	Daïkyo	1
Chlorobutyl rubber-metal cartridge crimp	West Pharmaceutical	1

qs = quantity sufficient to achieve

The formulation delivered 100 mcg of (SEQ ID NO.: 2) per 0.1 ml. (SEQ ID NO.: 2) was dissolved in Water for Injection containing dilute acetate buffer to give pH 5.1 was used.

Results confirm excellent chemical stability over 24 months, at 5 °C as shown in FIG. 1. This solution contains no stabilizer or preservative and only 6 mM acetate buffer.

In summation for (SEQ ID NO.: 2), stabilizer is not needed to give good stability in solution.

EXAMPLE 2: Use of citric acid buffer in lyophilised form of (SEQ ID NO.: 2).

TABLE 2

Material	Supplier	Unitary Formula (per vial)
(SEQ ID NO.: 2)	Ipsen Ireland	0.1 mg (free base)
Dextran 70	Interchemical	50 mg
Citric acid 0.25 % (w/v)	Prolabo	qs pH 4.5*
Water for injections**	Meram	qs 1 g
Type I clear glass vial, 11-13 ml	Verretubex	1
Grey chlorobutyl PTFE stopper, 20 mm	Daïkyo	1
Flip-off metal crimp	West Pharma	1

**to get pH 5 – 5.5 after lyophilisation removed after freeze-drying step.

The solutions in TABLE 2 were reconstituted with NaCl 0.9 %, to give:

ONE vial of 2 ml (=50 µg/ml) providing 10 to 80 µg/d doses (with injections of 200 µl to 1.6 ml), or

ONE vial of 5 ml (=20 µg/ml solution) providing 5 to 40 µg/d doses (with injections of 250 µl - 2 ml).

Citric acid was used to adjust pH and Dextran was used to provide a bulking agent to aid cake formation during lyophilization.

The solutions described were lyophilized in glass vials, and stored at various temperatures for up to 24 months. The content of [$Glu^{22,25}$, $Leu^{23,28,31}$, Aib^{29}, $Lys^{26,30}$]hPTHrP(1-34)NH_2 (SEQ ID NO.: 2), purity and physical tests were conducted on samples removed from storage at different times. Results are presented in FIG. 2, for peptide concentration, as percent remaining.. The data in FIG. 2 shows excellent stability over 24 months at 2-8°C.

EXAMPLE 3: Screening of formulations for $Glu^{22,25}$, $Leu^{23,28,31}$, Aib^{29}, $Lys^{26,30}$]hPTHrP(1-34)NH_2 (SEQ ID NO.: 2) to Compare Different Preservatives

TABLE 3 below shows Methyparaben and Benzyl Alcohol are not suitable preservatives for use with $Glu^{22,25}$, $Leu^{23,28,31}$, Aib^{29}, $Lys^{26,30}$]hPTHrP(1-34)NH_2 (SEQ ID NO.: 2), as precipitation and/or inactivity in preservative activity was seen.

TABLE 3

	Example 3a	Example 3b	Example 3c	Example 3d	Example 3e
Methylparaben	1.5 mg/ mL	1.35 mg/ mL	-	-	-
Propylparaben	-	0.15 mg/ mL	-	-	-
Phenol	-	-	5 mg/ mL	-	-
Chlorocresol	-	-	-	3 mg/ mL	-
Benzyl alcohol	-	-	-	-	10 mg/ mL
Preservative effectiveness test	Failed	Pass	Pass	Pass	Pass
Observationo on Issues	Precipitation observed	-	-	-	-
Preservative effectiveness test after storage 4.5 months at 5°C	Not Tested as precipitated initially	Pass	Pass	Pass	Fail

Solutions were prepared containing [$Glu^{22,25}$, $Leu^{23,28,31}$, Aib^{29}, $Lys^{26,30}$]hPTHrP(1-34)NH_2 (SEQ ID NO.: 2) 2mg/ml, acetate buffer 6mM and water for injection, with various different preservatives added at concentrations recommended for effective antimicrobial activity. Solutions were prepared at room temperature, by dissolution of the various ingredients in water for injection, with stirring over < 30 minutes to ensure complete dissolution, Solutions were filtered through 0.2 micron filter and filled into glass vials, to which a rubber stopper was applied and crimped in place to ensure complete closure.

The solution with methylparaben was less acceptable due to precipitation and inactivity immediately after manufacture of the solution. The solutions were then stored for up to 3 months at 25°C, and up to 4.5 months at 5°C and the preservative effectiveness test repeated. as described in Example 5.

EXAMPLE 4: Evaluation of Anti-microbial Preservative Effectiveness of Various concentrations of $Glu^{22,25}$, $Leu^{23,28,31}$, Aib^{29}, $Lys^{26,30}$]hPTHrP(1-34)NH_2 (SEQ ID NO.: 2) Compositions (stability study)

TABLE 4

	P87228	P87229	P87230	P87231
(SEQ ID NO.: 2)	2 mg/ mL	2 mg/ mL	2 mg/ mL	2 mg/ mL
Anti-microbial	Phenol 5 mg/ mL	Chlorocresol 3 mg/ mL	Chlorocresol 2 mg/ mL	Benzyl alcohol 10 mg/ mL
Acetate buffer	pH 5.1	pH 5.1	pH 5.1	pH 5.1

The solutions were tested according to European Pharmacopoeia, Chapter 5.1.3 "Efficacité de la conservation anti-microbienne" (Anti-microbial effectiveness test) to prove the effectiveness of the preservative.

TABLE 5: Preservative effectiveness test after manufacturing

Organisms : Bacteria	Initial organism concentration in cfu / mL	Test interval	Nb of cfu present / preparation (plate-count method)			
			P87228	P87229	P87230	P87231
			Phenol 5mg/ml	Chlorocresol 3mg/ml	Chlorocresol 2mg/ml	Benzyl alc. 10mg/ml
Staphylococcus aureus	3.8×10^5	T 0	3.4×10^5	<5	<5	4.7×10^5
		T+6 hrs	<5	<5	<5	6.8×10^2
		T+24 hrs	<5	<5	<5	<5
		T+28 days	5 (*)	<5	<5	<5
Pseudomonas aeruginosa	1.3×10^6	T 0	5	<5	<5	1.5×10^2
		T+6 hrs	<5	<5	<5	<5
		T+24 hrs	<5	<5	<5	<5
		T+28 days	<5	<5	<5	<5
E.coli	6.7×10^5	T 0	7.2×10^3	<5	<5	1.1×10^5
		T+6 hrs	<5	<5	<5	<5
		T+24 hrs	<5	<5	<5	<5
		T+28 days	<5	<5	<5	<5
(*) Bacillus Gram +, different from St. Aureus -> result conform						
Organism : Yeast and mold	Initial organism concentration in cfu / mL	Test interval	Nb of cfu present / preparation (plate-count method)			
			P87228	P87229	P87230	P87231
			Phenol 5mg/ml	chlorocresol 3mg/ml	chlorocresol 2mg/ml	Benzyl alc. 10mg/ml
Aspergillus niger	3.4×10^5	T 0	4.0×10^5	<5	<5	4.1×10^5
		T+7 days	<5	<5	<5	<5
		T+28 days	<5	<5	<5	<5
Candida albicans	3.9×10^5	T 0	4.4×10^5	<5	<5	3.8×10^5
		T+7 days	<5	<5	<5	5
		T+28 days	<5	<5	<5	<5
Results :	Conform	-	Conform	Conform	Conform	Conform

Nb of cfu= number of colony forming units

TABLE 5 Cont'd.: Preservative effectiveness test results after 3 months storage at 25°C

Organisms : Bacteria	Initial organism concentration in cfu / mL	Test interval (days)	Nb of cfu present / preparation (plate-count method)			
			P87228	P87229	P87230	P87231
			Phenol 5mg/ml	chlorocresol 3mg/ml	chlorocresol 2mg/ml	Benzyl alc. 10mg/ml
Staphylococcus aureus	2.7×10^5 (P87228, P87229, P87231) 5.2×10^5 (P87230)	0hr	1.9×10^5	<5	<5	3.8×10^5
		6hr	30	<5	<5	5.9×10^3
		24hr	<5	<5	<5	<5
		28day	<5	<5	<5	<5
Pseudomonas aeruginosa	9.9×10^5 (P87228, P87229, P87231) 8.5×10^5 (P87230)	0hr	<5	<5	<5	<5
		6hr	<5	<5	<5	<5
		24hr	<5	<5	<5	<5
		28day	<5	<5	<5	<5
E.coli	6.8×10^5 (P87228, P87229, P87231) 9.5×10^5 (P87230)	0hr	1.7×10^5	<5	<5	8.0×10^4
		6hr	<5	<5	<5	5
		24hr	<5	<5	<5	<5
		28day	<5	<5	<5	<5
Organism : Yeast and mold	Initial organism concentration in cfu / mL	Test interval	Nb of cfu present / preparation (plate-count method)			
			P87228	P87229	P87230	P87231
			Phenol 5mg/ml	Chlorocresol 3mg/ml	Chlorocresol 2mg/ml	Benzyl alc. 10mg/ml
Aspergillus niger	3.3×10^5 (P87228, P87229, P87231) 4.1×10^5 (P87230)	0hr	3.8×10^5	55	70	4.1×10^5
		7day	<5	<5	<5	<5
		28day	<5	<5	<5	<5
Candida albicans	2.7×10^5 (P87228, P87229, P87231) 3.7×10^5 (P87230)	0hr	4.0×10^5	<5	<5	3.8×10^5
		7day	<5	<5	<5	<5
		28day	<5	<5	<5	<5
Results :	Conform	-	Conform	Conform	Conform	Not Conform

TABLE 5 Cont'd.: Preservative effectiveness test results after 4.5 months storage at 5°C

Organisms: Bacteria	Initial organism concentration in cfu / mL	Test interval (days)	Nb of cfu present / preparation (plate-count method)			
			P87228	P87229	P87230	P87231
			Phenol 5mg/ml	chlorocresol 3mg/ml	chlorocresol 2mg/ml	Benzyl alc. 10mg/ml
Staphylococcus aureus	5.4×10^5	0hr	4.1×10^5	<5	<5	5.1×10^5
		6hr	<5	<5	<5	7.2×10^3
		24hr	<5	<5	<5	<5
		28day	<5	<5	<5	<5
Pseudomonas aeruginosa	9.7×10^5	0hr	<5	<5	<5	<5
		6hr	<5	<5	<5	<5
		24hr	<5	<5	<5	<5
		28day	<5	<5	<5	<5
E.coli	6.1×10^5	0hr	7.0×10^4	5	5	4.2×10^4
		6hr	<5	<5	<5	<5
		24hr	<5	<5	<5	<5
		28day	<5	<5	<5	<5
Organism: Yeast and mold	Initial organism concentration in cfu / mL	Test interval	Nb of cfu present / preparation (plate-count method)			
			P87228	P87229	P87230	P87231
			Phenol 5mg/ml	Chlorocresol 3mg/ml	Chlorocresol 2mg/ml	Benzyl alc. 10mg/ml
Aspergillus niger	5.3×10^5	0hr	3.7×10^5	1.8×10^3	7.5×10^3	4.1×10^5
		7day	<5	<5	<5	<5
		28day	<5	<5	<5	<5
Candida albicans	4.1×10^5	0hr	4.5×10^5	<5	5	4.5×10^5
		7day	<5	<5	<5	<5
		28day	<5	<5	<5	<5
Results :	Conform	-	Conform	Conform	Conform	Not Conform

TABLE 5 shows Phenol, Chlorocresol and Benzyl Alcohol all produce compliant results immediately after manufacture for both Bacteria and Yeasts/moulds. After 3 and 4.5 months storage, the preservative efficacy is maintained for Phenol and Chlorocresol, for both Bacteria and Yeasts/moulds.

However, for Benzyl Alcohol, the efficacy against Bacteria is not compliant, as the data shows insufficient rate of kill against S Aureus (TABLE 5).

EXAMPLE 5: Chemical stability of different formulations

TABLE 6 details the chemical stability of the formulations described in Example 4.

TABLE 6: $Glu^{22,25}$, $Leu^{23,28,31}$, Aib^{29}, $Lys^{26,30}$]hPTHrP(1-34)NH_2 (SEQ ID NO.: 2) stability results

		Storage conditions : 25°C, 60% RH		
		(SEQ ID NO.: 2) content in mg/mL (% initial concentration at t=0)		
Batch	Composition	0 month	1 month	3 months
P87228	(SEQ ID NO.: 2) (2mg/ml) / Phenol (5mg/ml)	1.90 (100%)	1.88 (98.9%)	1.83 (96.3%)
P87229	(SEQ ID NO.: 2) (2mg/ml) / Chlorocresol (3mg/ml)	1.98 (100%)	1.96 (99.0%)	1.94 (98.0%)
P87231	(SEQ ID NO.: 2) (2mg/ml) / Benzyl Alcohol (10mg/ml)	1.93 (100%)	1.89 (97.9%)	1.86 (96.4%)

		Storage conditions : 5°C		
		(SEQ ID NO.: 2) content in mg/mL (% initial concentration at t=0)		
Batch	Composition	0 month	3 month	4.5 month
P87228	(SEQ ID NO.: 2) (2mg/ml) / Phenol (5mg/ml)	1.90 (100%)	1.91 (100.5%)	1.89 (99.5%)
P87229	(SEQ ID NO.: 2) (2mg/ml) / Chlorocresol (3mg/ml)	1.98 (100%)	1.96 (99.0%)	1.97 (99.5%)
P87231	(SEQ ID NO.: 2) (2mg/ml) / Benzyl Alcohol (10mg/ml)	1.93 (100%)	1.94 (100.5%)	1.92 (99.5%)

As can be seen from TABLE 6 and [$Glu^{22,25}$, $Leu^{23,28,31}$, Aib^{29}, $Lys^{26,30}$]hPTHrP(1-34)NH_2 (SEQ ID NO.: 2) solution stability is not significantly influenced by the preservative selected.. TABLE 7 details the content of each preservative for the same formulations.

TABLE 7: Preservative stability results

		Storage conditions : 25°C, 60% RH		
		Preservative content in mg/ml (% initial concentration at t=0)		
Batch	Composition	0 month	1 month	3 month
P87228	(SEQ ID NO.: 2) (2mg/ml) / Phenol (5mg/ml)	4.86 (100%)	4.82 (99.2%)	4.79 (98.6%)
P87229	(SEQ ID NO.: 2) (2mg/ml) / Chlorocresol (3mg/ml)	2.78 (100%)	2.70 (97.1%)	2.56 (92.1%)
P87231	(SEQ ID NO.: 2) (2mg/ml) / Benzyl Alcohol (10mg/ml)	9.92 (100%)	9.83 (99.1%)	9.82 (99.0%)

		Storage conditions : 5°C		
		Preservative content in mg/mL (% initial concentration at t=0)		
Batch	Composition	0 month	3 month	4.5 month
P87228	(SEQ ID NO.: 2) (2mg/ml) / Phenol (5mg/ml)	4.86 (100%)	4.83 (99.4%)	4.84 (99.6%)
P87229	(SEQ ID NO.: 2) (2mg/ml) / Chlorocresol (3mg/ml)	2.78 (100%)	2.73 (98.2%)	2.74 (98.6%)
P87231	(SEQ ID NO.: 2) (2mg/ml) / Benzyl Alcohol (10mg/ml)	9.92 (100%)	9.89 (99.7%)	9.94 (100.2%)

As can be seen from TABLE 7 chlorocresol is the preservative which has the lower stability, with greater loss in preservative content under both 5 and 25°C storage.

While this invention has been particularly shown and described with references to example embodiments thereof, it will be understood by those skilled in the art that various changes in form and details may be made therein without departing from the scope of the invention encompassed by the appended claims.

SEQUENCE LISTING

<110> Radius Health, Inc.
Societe de Conseils de Recherches et D'Applications Scientifiques (S.C.R.A.S.)
Dey, Michael J.
Mondoly, Nathalie
Rigaud, Benedicte
Henderson, Bart
Lyttle, C. Richard

<120> Method of Drug Delivery for Bone Anabolic Protein

<130> 3803.1004002

<140> PCT/US2007/021216
<141> 2007-10-03

<150> 60/848,960
<151> 2006-10-03

<160> 2

<170> FastSEQ for Windows Version 4.0

<210> 1
<211> 34
<212> PRT
<213> Homo Sapiens

<400> 1
Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile Gln
1 5 10 15
Asp Leu Arg Arg Arg Phe Phe Leu His His Leu Ile Ala Glu Ile His
20 25 30
Thr Ala

<210> 2
<211> 34
<212> PRT
<213> Artificial Sequence

<220>
<223> synthetic peptide

<220>
<221> MOD_RES
<222> 29
<223> Xaa = Aib

<220>
<221> AMIDATION
<222> 34

<400> 2
Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile Gln
1 5 10 15
Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Xaa Lys Leu His
20 25 30
Thr Ala

Claims

1. Use of a storage stable composition in the manufacture of a medicament for stimulating bone growth in a subject with a bone fracture, said storage-stable composition comprising:

   a) a PTHrP analogue whose sequence is [Glu ^{22,25}, Leu^{23,28,31}, Aib^{29},Lys^{26,30}]hPTHrP(1-34)NH_2 (SEQ ID NO.2); and
   b) an effective amount of a pH buffer to maintain the pH in a range of 4.5 to 5.6.

2. The use according to claim 1 wherein said subject is administered the medicament by single daily subcutaneous injection of an amount of said composition containing from 75 to 80 μg of [Glu ^{22,25}, Leu^{23,28,31}, Aib^{29},Lys^{26,30}]hPTHrP(1-34)NH_2 (SEQ ID NO.2).

3. The use according to claim 2 wherein said storage-stable composition further comprises phenol in a concentration from 0.25 to 5 mg/mL.

4. The use according to any one of claims 1 to 3 wherein said pH buffer is an acetate buffer.

5. The use according to claim 1 wherein said subject is administered the medicament by single daily subcutaneous injection of an amount of said composition containing from 40 to 45 μg of [Glu ^{22,25}, Leu^{23,28,31}, Aib^{29},Lys^{26,30}]hPTHrP(1-34)NH_2 (SEQ ID NO.2).

6. The use according to claim 5 wherein said storage-stable composition further comprises phenol in a concentration from 0.25 to 5 mg/mL.

7. The use according to any one of claims 5 to 6 wherein said pH buffer is an acetate buffer.

8. Use of a storage stable composition according to claim 1, wherein said pH is 5.1.

9. A storage stable composition for use as a medicament for stimulating bone growth in a subject with a bone fracture, said storage-stable composition comprising:

a) a PTHrP analogue whose sequence is [Glu ^{22,25}, Leu^{23,28,31}, Aib^{29},Lys^{26,30}]hPTHrP(1-34)NH_2 (SEQ ID NO.2); and
b) an effective amount of a pH buffer to maintain the pH in a range of 4.5 to 5.6.

10. Use of a storage stable composition according to claim 9 wherein said subject is administered the medicament by single daily subcutaneous injection of an amount of said composition containing from 75 to 80 μg of [Glu $^{22.25}$, Leu$^{23.28.31}$, Aib^{29},Lys^{26,30}]hPTHrP(1-34)NH_2 (SEQ ID NO.2).

11. Use of a storage stable composition according to claim 10 wherein said storage-stable composition further comprises phenol in a concentration from 0.25 to 5 mg/mL.

12. Use of a storage stable composition according to any one of claims 9 to 11 wherein said pH buffer is an acetate buffer.

13. Use of a storage stable composition according to claim 9 wherein said subject is administered the medicament by single daily subcutaneous injection of an amount of said composition containing from 40 to 45 μg of [Glu $^{22.25}$, Leu$^{23.28.31}$, Aib^{29},Lys$^{26.30}$]hPTHrP(1-34)NH_2 (SEQ ID NO.2).

14. Use of a storage stable composition according to claim 13 wherein said storage-stable composition further comprises phenol in a concentration from about 0.25 to about 5 mg/mL.

15. Use of a storage stable composition according to any one of claims 13 to 14 wherein said pH buffer is an acetate buffer.

16. Use of a storage stable composition according to claim 9, wherein said pH is about 5.1.