

GLUCAGON ANALOGS EXHIBITING ENHANCED SOLUBILITY IN PHYSIOLOGICAL PH BUFFERS

BACKGROUND

Hypoglycemia occurs when blood glucose levels drops too low to provide
5 enough energy for the body's activities. In adults or children older than 10 years,
hypoglycemia is uncommon except as a side effect of diabetes treatment, but it can
result from other medications or diseases, hormone or enzyme deficiencies, or tumors.
When blood glucose begins to fall, glucagon, a hormone produced by the pancreas,
signals the liver to break down glycogen and release glucose, causing blood glucose
10 levels to rise toward a normal level. However for diabetics, this glucagon response to
hypoglycemia may be impaired, making it harder for glucose levels to return to the
normal range.

Hypoglycemia is a life threatening event that requires immediate medical
attention. The administration of glucagon is an established medication for treating
15 acute hypoglycemia and it can restore normal levels of glucose within minutes of
administration. When glucagon is used in the acute medical treatment of
hypoglycemia, a crystalline form of glucagon is solubilized with a dilute acid buffer
and the solution is injected intramuscularly. While this treatment is effective, the
methodology is cumbersome and dangerous for someone that is semi-conscious.
20 Accordingly, there is a need for a glucagon analog that maintains the biological
performance of the parent molecule but is sufficiently soluble and stable, under
relevant physiological conditions, that it can be pre-formulated as a solution, ready for
injection.

Additionally, diabetics are encouraged to maintain near normal blood glucose
25 levels to delay or prevent microvascular complications. Achievement of this goal
usually requires intensive insulin therapy. In striving to achieve this goal, physicians
have encountered a substantial increase in the frequency and severity of
hypoglycemia in their diabetic patients. Accordingly, improved pharmaceuticals and
methodologies are needed for treating diabetes that are less likely to induce
30 hypoglycemia than current insulin therapies.

As described herein, high potency glucagon agonists are provided that exhibit
enhanced biophysical stability and aqueous solubility at physiological pH in

pharmaceutical compositions suitable for commercial use. Native glucagon is neither soluble, nor stable in the physiological pH range and thus must be manufactured as a dry product that requires reconstitution and immediate use. The glucagon analogs described herein have enhanced physical properties that render them superior for use in current medicinal settings where the native hormone is currently employed. These compounds can be used in accordance with one embodiment to prepare pre-formulated solutions ready for injection to treat hypoglycemia. Alternatively, the glucagon agonists can be co-administered with insulin to buffer the effects of insulin to allow for a more stable maintenance of blood glucose levels. In addition, other beneficial uses of compositions comprising the modified glucagon peptides disclosed herein are described in detail below.

SUMMARY

One embodiment of the invention provides glucagon peptides that retain glucagon receptor activity and exhibit improved solubility relative to the native glucagon peptide (SEQ ID NO: 1). Native glucagon exhibits poor solubility in aqueous solution, particularly at physiological pH, with a tendency to aggregate and precipitate over time. In contrast, the glucagon peptides of one embodiment of the invention exhibit at least 2-fold, 5-fold, or even higher solubility compared to native glucagon at a pH between 6 and 8, for example, at pH 7 after 24 hours at 25°C.

In one embodiment the glucagon peptides retain at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 75% activity, 80% activity, 85% activity, or 90% of the activity of native glucagon. In one embodiment, the glucagon peptides of the present invention have potency greater than glucagon. Any of the glucagon peptides of the invention may additionally exhibit improved stability and/or reduced degradation, for example, retaining at least 95% of the original peptide after 24 hours at 25°C.

In accordance with one embodiment a glucagon peptide is provided wherein the peptide is modified by amino acid substitutions and/or additions that introduce a charged amino acid into the C-terminal portion of the peptide, and in one embodiment at a position C-terminal to position 27 of SEQ ID NO: 1. Optionally, one, two or three charged amino acids may be introduced within the C-terminal portion, and in one embodiment C-terminal to position 27. In accordance with one embodiment the

native amino acid(s) at positions 28 and/or 29 are substituted with a charged amino acid, and/or one to three charged amino acids are added to the C-terminus of the peptide. In exemplary embodiments, one, two or all of the charged amino acids are negatively charged. Additional modifications, e.g. conservative substitutions, may be made to the glucagon peptide that still allow it to retain glucagon activity.

In accordance with one exemplary embodiment the glucagon peptide comprises an amino acid sequence of SEQ ID NO: 11, or an analog thereof that contains 1 to 3 further amino acid modifications relative to native glucagon, or a glucagon agonist analog thereof. SEQ ID NO: 11 represents a modified glucagon peptide wherein the asparagine residue at position 28 of the native protein has been substituted with an aspartic acid. In another exemplary embodiment the glucagon peptide comprises an amino acid sequence of SEQ ID NO: 38, wherein the asparagine residue at position 28 of the native protein has been substituted with glutamic acid. Other exemplary embodiments include glucagon peptides of SEQ ID NOS: 24, 25, 26, 33, 35, 36 and 37.

The solubility of any of the foregoing compounds can be further improved by attaching a hydrophilic moiety to the peptide. In one embodiment the hydrophilic moiety is a polyethylene glycol chain or other water soluble polymer that is covalently linked to the side chain of an amino acid residue at position 16, 17, 21 or 24 of said glucagon peptide. The polyethylene glycol chain in accordance with one embodiment has a molecular weight selected from the range of about 500 to about 40,000 Daltons. The present invention further encompasses pharmaceutically acceptable salts of said glucagon agonists.

In other exemplary embodiments, any of the foregoing compounds can be further modified to alter its pharmaceutical properties by the addition of a second peptide to the carboxy terminus of the glucagon peptide. In one embodiment a glucagon peptide is covalently bound through a peptide bond to a second peptide, wherein the second peptide comprises a sequence selected from the group consisting of SEQ ID NO: 20, SEQ ID NO: 21 and SEQ ID NO: 22.

In yet further exemplary embodiments, any of the foregoing compounds can be further modified to improve stability by modifying the amino acid at position 15 of SEQ ID NO: 1 to reduce degradation of the peptide over time, especially in acidic or alkaline buffers.

In accordance with one embodiment a pharmaceutical composition is provided comprising any of the novel glucagon peptides disclosed herein, preferably at a purity level of at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99%, and a pharmaceutically acceptable diluent, carrier or excipient. Such compositions may
5 contain a glucagon peptide at a concentration of at least 0.5 mg/ml, 1 mg/ml, 2 mg/ml, 3 mg/ml, 4 mg/ml, 5 mg/ml, or higher. In one embodiment the pharmaceutical compositions comprise aqueous solutions that are sterilized and optionally stored within various containers. In other embodiments the pharmaceutical compositions comprise a lyophilized powder. The pharmaceutical compositions can be further
10 packaged as part of a kit that includes a disposable device for administering the composition to a patient. The containers or kits may be labeled for storage at ambient room temperature or at refrigerated temperature.

In accordance with one embodiment a method of rapidly increasing glucose level or treating hypoglycemia using a pre-formulated aqueous composition is
15 provided. The method comprises the step of administering an effective amount of an aqueous solution comprising a novel modified glucagon peptide of the present disclosure. In another embodiment a method is provided for inducing the temporary paralysis of the intestinal tract. The method comprises the step of administering one or more of the glucagon peptides disclosed herein to a patient in need thereof.

20 In yet another embodiment a method of reducing weight gain or inducing weight loss is provided, which involves administering an effective amount of an aqueous solution comprising a glucagon peptide of the invention. In further embodiments, methods of treating diabetes involving co-administering insulin and a glucagon peptide of the invention are provided.

25 Exemplary glucagon peptides are selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11 and SEQ ID NO: 33, wherein amino acid 29 of the glucagon peptide is bound to a second peptide through a peptide bond, and said second peptide comprises the sequence of SEQ ID
30 NO: 20, SEQ ID NO: 21 or SEQ ID NO: 22. In one embodiment the glucagon peptide is pegylated. In one embodiment the method comprises the step of administering a peptide comprising the sequence of SEQ ID NO: 24, SEQ ID NO: 25,

and SEQ ID NO: 26, wherein a polyethylene chain is covalently linked to amino acid position 21 or at position 24.

BRIEF DESCRIPTION OF THE DRAWINGS

5 Fig. 1 is a bar graph representing the stability of Glucagon Cys²¹maleimidoPEG_{5K} at 37°C incubated for 24, 48, 72, 96, 144 and 166 hours, respectively.

Fig. 2 represents data generated from HPLC analysis of Glucagon Cys²¹maleimidoPEG_{5K} at pH 5 incubated at 37°C for 24, 72 or 144 hours,
10 respectively.

Fig. 3 represents data showing the solubility of glucagon analogs (D28, E29, E30) relative to native glucagon after 60 hours at 25 °C at pH of 2, 4, 5.5, 7 and 8, respectively.

Fig. 4 represents data showing the solubility of glucagon analogs (E15D28,
15 D28E29 and D28E30) relative to native glucagon after 24 hours at 25 °C and then 24 hours at 4 °C at pH of 2, 4, 5.5 and 7, respectively.

Fig. 5 represents the maximum solubility of glucagon analogs D28, D28E30 and E15,D28 after 24 hours, pH 7 at 4 °C.

Fig. 6 represents data showing a glucagon receptor mediated cAMP induction
20 by glucagon analogs (K29 ▲, K30 ▼, and K29K30 ◆) relative to native glucagon ■.

Fig. 7 represents data showing a glucagon receptor mediated cAMP induction by glucagon analogs (D28 □, E29 △, E30 ▽, K30K31 ◇ and K30, ▼) relative to native glucagon ■.

25 Fig. 8 represents data showing a glucagon receptor mediated cAMP induction by glucagon analogs (D28 □, E28 ◆ and K29, ▲) relative to native glucagon ■.

Fig. 9 represents data showing a glucagon receptor mediated cAMP induction by glucagon analogs (D28E29 +, D28E30 ×, E15D28 * and E29 △) relative to native glucagon ■.

30 Fig. 10 represents data showing the change in serum glucose concentrations in beagle dogs after intramuscular administration of glucagon and glucagon analogs. The animals were administered a 0.005 mg/kg dose of either glucagon, a glucagon analog comprising glucagon with the sequence of SEQ ID NO: 31 linked to the

carboxy terminus of glucagon (glucagon-CEX) or a glucagon analog comprising an aspartic acid substitution at amino acid 28 (glucagon-Asp28) SEQ ID NO: 11.

DETAILED DESCRIPTION

5 DEFINITIONS

In describing and claiming the invention, the following terminology will be used in accordance with the definitions set forth below.

As used herein, the term "pharmaceutically acceptable carrier" includes any of the standard pharmaceutical carriers, such as a phosphate buffered saline solution,
10 water, emulsions such as an oil/water or water/oil emulsion, and various types of wetting agents. The term also encompasses any of the agents approved by a regulatory agency of the US Federal government or listed in the US Pharmacopeia for use in animals, including humans.

As used herein the term "pharmaceutically acceptable salt" refers to salts of
15 compounds that retain the biological activity of the parent compound, and which are not biologically or otherwise undesirable. Many of the compounds disclosed herein are capable of forming acid and/or base salts by virtue of the presence of amino and/or carboxyl groups or groups similar thereto.

Pharmaceutically acceptable base addition salts can be prepared from
20 inorganic and organic bases. Salts derived from inorganic bases, include by way of example only, sodium, potassium, lithium, ammonium, calcium and magnesium salts. Salts derived from organic bases include, but are not limited to, salts of primary, secondary and tertiary amines.

Pharmaceutically acceptable acid addition salts may be prepared from
25 inorganic and organic acids. Salts derived from inorganic acids include hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like. Salts derived from organic acids include acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, malic acid, malonic acid, succinic acid, maleic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic
30 acid, ethanesulfonic acid, p-toluene-sulfonic acid, salicylic acid, and the like.

As used herein, the term "treating" includes prophylaxis of the specific disorder or condition, or alleviation of the symptoms associated with a specific disorder or condition and/or preventing or eliminating said symptoms.

As used herein an "effective" amount or a "therapeutically effective amount" of a glucagon peptide refers to a nontoxic but sufficient amount of the peptide to provide the desired effect. For example one desired effect would be the prevention or treatment of hypoglycemia, as measured, for example, by an increase in blood glucose level. The amount that is "effective" will vary from subject to subject, depending on the age and general condition of the individual, mode of administration, and the like. Thus, it is not always possible to specify an exact "effective amount." However, an appropriate "effective" amount in any individual case may be determined by one of ordinary skill in the art using routine experimentation.

10 The term, "parenteral" means not through the alimentary canal but by some other route such as subcutaneous, intramuscular, intraspinal, or intravenous.

As used herein, the term "purified" and like terms relate to the isolation of a molecule or compound in a form that is substantially free of contaminants normally associated with the molecule or compound in a native or natural environment.

15 As used herein, the term "purified" does not require absolute purity; rather, it is intended as a relative definition. The term "purified polypeptide" is used herein to describe a polypeptide which has been separated from other compounds including, but not limited to nucleic acid molecules, lipids and carbohydrates.

20 The term "isolated" requires that the referenced material be removed from its original environment (e.g., the natural environment if it is naturally occurring). For example, a naturally-occurring polynucleotide present in a living animal is not isolated, but the same polynucleotide, separated from some or all of the coexisting materials in the natural system, is isolated.

25 A "glucagon peptide" as used herein includes any peptide comprising, either the amino acid sequence of SEQ ID NO: 1, or any analog of the amino acid sequence of SEQ ID NO: 1, including amino acid substitutions, additions, or deletions, or post translational modifications (e.g. methylation, acylation, ubiquitination and the like) of the peptide, that stimulates glucagon or GLP-1 receptor activity, as measured by cAMP production using the assay described in Example 13.

30 The term "glucagon agonist" refers to a complex comprising a glucagon peptide that stimulates glucagon receptor activity, as measured by cAMP production using the assay described in Example 13.

As used herein a "glucagon agonist analog" is a glucagon peptide comprising a sequence selected from the group consisting of SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, SEQ ID NO: 12 and SEQ ID NO: 13 or a analog of such a sequence that has been modified to include one or more
5 conservative amino acid substitutions at positions 2, 5, 7, 10, 11, 12, 13, 14, 16, 17, 18, 19, 20, 21, 24, 27, 28 or 29.

As used herein an amino acid "modification" refers to a substitution, addition or deletion of an amino acid, and includes substitution with or addition of any of the 20 amino acids commonly found in human proteins, as well as atypical or non-
10 naturally occurring amino acids. Throughout the application, all references to a particular amino acid position by number (e.g. position 28) refer to the amino acid at that position in native glucagon (SEQ ID NO:1) or the corresponding amino acid position in any analogs thereof. For example, a reference herein to "position 28" would mean the corresponding position 27 for a glucagon analog in which the first
15 amino acid of SEQ ID NO: 1 has been deleted. Similarly, a reference herein to "position 28" would mean the corresponding position 29 for a glucagon analog in which one amino acid has been added before the N-terminus of SEQ ID NO: 1.

As used herein an amino acid "substitution" refers to the replacement of one amino acid residue by a different amino acid residue.

20 As used herein, the term "conservative amino acid substitution" is defined herein as exchanges within one of the following five groups:

- I. Small aliphatic, nonpolar or slightly polar residues:
Ala, Ser, Thr, Pro, Gly;
- II. Polar, negatively charged residues and their amides:
25 Asp, Asn, Glu, Gln, cysteic acid and homocysteic acid;
- III. Polar, positively charged residues:
His, Arg, Lys; Ornithine (Orn)
- IV. Large, aliphatic, nonpolar residues:
Met, Leu, Ile, Val, Cys, Norleucine (Nle), homocysteine
- 30 V. Large, aromatic residues:
Phe, Tyr, Trp, acetyl phenylalanine

As used herein the general term "polyethylene glycol" or "PEG", refers to mixtures of condensation polymers of ethylene oxide and water, in a branched or straight chain, represented by the general formula $H(OCH_2CH_2)_nOH$, wherein n is at least 9. Absent any further characterization, the term is intended to include polymers of ethylene glycol with an average total molecular weight selected from the range of 500 to 40,000 Daltons. "polyethylene glycol" or "PEG" is used in combination with a numeric suffix to indicate the approximate average molecular weight thereof. For example, PEG-5,000 refers to polyethylene glycol having a total molecular weight average of about 5,000.

As used herein the term "pegylated" and like terms refers to a compound that has been modified from its native state by linking a polyethylene glycol polymer to the compound. A "pegylated glucagon peptide" is a glucagon peptide that has a PEG chain covalently bound to the glucagon peptide.

As used herein a general reference to a peptide is intended to encompass peptides that have modified amino and carboxy termini. For example, an amino acid chain comprising an amide group in place of the terminal carboxylic acid is intended to be encompassed by an amino acid sequence designating the standard amino acids.

As used herein a "linker" is a bond, molecule or group of molecules that binds two separate entities to one another. Linkers may provide for optimal spacing of the two entities or may further supply a labile linkage that allows the two entities to be separated from each other. Labile linkages include photocleavable groups, acid-labile moieties, base-labile moieties and enzyme-cleavable groups.

As used herein a "dimer" is a complex comprising two subunits covalently bound to one another via a linker. The term dimer, when used absent any qualifying language, encompasses both homodimers and heterodimers. A homodimer comprises two identical subunits, whereas a heterodimer comprises two subunits that differ, although the two subunits are substantially similar to one another.

As used herein the term "pH stabilized glucagon peptide" refers to a glucagon agonist analog that exhibits superior stability and solubility, relative to native glucagon, in aqueous buffers in the broadest pH range used for pharmacological purposes.

As used herein the term "charged amino acid" refers to an amino acid that comprises a side chain that is negatively charged (i.e., de-protonated) or positively

charged (i.e., protonated) in aqueous solution at physiological pH. For example negatively charged amino acids include aspartic acid, glutamic acid, cysteic acid, homocysteic acid, and homoglutamic acid, whereas positively charged amino acids include arginine, lysine and histidine. Charged amino acids include the charged
5 amino acids among the 20 amino acids commonly found in human proteins, as well as atypical or non-naturally occurring amino acids.

As used herein the term "acidic amino acid" refers to an amino acid that comprises a second acidic moiety, including for example, a carboxylic acid or sulfonic acid group.

10

EMBODIMENTS

Applicants have discovered that native glucagon can be modified by introducing charge at its carboxy terminus to enhance the solubility of the peptide while retaining the agonist properties of the peptide. The enhanced solubility allows
15 for the preparation and storage of glucagon solutions at near neutral pH. Formulating glucagon solutions at relatively neutral pHs (e.g. pH of about 6.0 to about 8.0) improves the long term stability of the glucagon peptides. Accordingly, one embodiment of the present invention is directed to a glucagon agonist that has been modified relative to the wild type peptide of His-Ser-Gln-Gly-Thr-Phe- Thr-Ser-Asp-
20 Tyr-Ser-Lys-Tyr-Leu-Asp-Ser- Arg-Arg-Ala-Gln-Asp-Phe-Val-Gln-Trp-Leu- Met-Asn-Thr (SEQ ID NO: 1) to improve the peptide's solubility in aqueous solutions, particularly at a pH ranging from about 5.5 to about 8.0, while retaining the native peptide's biological activity. In one embodiment charge is added to the peptide by the substitution of native non-charged amino acids with charged amino acids selected
25 from the group consisting of lysine, arginine, histidine, aspartic acid and glutamic acid, or by the addition of charged amino acids to the amino or carboxy terminus of the peptide. Surprisingly, applicants have discovered that substituting the normally occurring amino acid at position 28 and/or 29 with charged amino acids, and/or the addition of one to two charged amino acids at the carboxy terminus of the glucagon
30 peptide, enhances the solubility and stability of the glucagon peptides in aqueous solutions at physiologically relevant pHs (i.e., a pH of about 6.5 to about 7.5) by at least 5-fold and by as much as 30-fold.

Accordingly, glucagon peptides of one embodiment of the invention retain glucagon activity and exhibit at least 2-fold, 5-fold, 10-fold, 15-fold, 25-fold, 30-fold or greater solubility relative to native glucagon at a given pH between about 5.5 and 8, e.g., pH 7, when measured after 24 hours at 25°C. Any of the glucagon peptides disclosed herein may additionally exhibit improved stability at a pH within the range of 5.5 to 8, for example, retaining at least 75%, 80%, 90%, 95%, 96%, 97%, 98% or 99% of the original peptide after 24 hours at 25°C. The glucagon peptides may include additional modifications that alter its pharmaceutical properties, e.g. increased potency, prolonged half-life in circulation, increased shelf-life, reduced precipitation or aggregation, and/or reduced degradation, e.g., reduced occurrence of cleavage or chemical modification after storage.

In one embodiment a glucagon peptide with improved solubility may be prepared, for example, by introducing one, two, three or more charged amino acid(s) to the C-terminal portion of native glucagon, and in one embodiment at a position C-terminal to position 27. Such a charged amino acid can be introduced, for example by substituting a native amino acid with a charged amino acid, e.g. at positions 28 or 29, or alternatively by adding a charged amino acid, e.g. after position 27, 28 or 29. In exemplary embodiments, one, two, three or all of the charged amino acids are negatively charged. In other embodiments, one, two, three or all of the charged amino acids are positively charged. In specific exemplary embodiments, the glucagon peptide may comprise any one or two of the following modifications: substitution of N28 with E; substitution of N28 with D; substitution of T29 with D; substitution of T29 with E; insertion of E after position 27, 28 or 29; insertion of D after position 27, 28 or 29. For example, E28E30, D28E30

Additional modifications may be made to the glucagon peptide which may further increase solubility and/or stability and/or glucagon activity. The glucagon peptide may alternatively comprise other modifications that do not substantially affect solubility or stability, and that do not substantially decrease glucagon activity. In exemplary embodiments, the glucagon peptide may comprise a total of 1, up to 2, up to 3, up to 4, up to 5, up to 6, up to 7, up to 8, up to 9, or up to 10 amino acid modifications relative to the native glucagon sequence.

Exemplary modifications include but are not limited to:

conservative substitutions, for example, conservative substitutions at one or more of positions 2, 5, 7, 10, 11, 12, 13, 14, 16, 17, 18, 19, 20, 21, 24, 27, 28 or 29;

modification of the aspartic acid at position 15, for example, by substitution with glutamic acid, homoglutamic acid, cysteic acid or homocysteic acid, which may

5 reduce degradation;

addition of a hydrophilic moiety such as the water soluble polymer polyethylene glycol, as described herein, e.g. at position 16, 17, 20, 21, 24 or 29, which may increase solubility and/or half-life;

10 modification of the amino acid at position 27, for example, by substitution with methionine, leucine or norleucine;

modifications at position 1 or 2 as described herein;

C-terminal extensions as described herein;

homodimerization or heterodimerization as described herein; and

combinations of the above.

15 In accordance with one embodiment the native glucagon peptide of SEQ ID NO: 1 is modified by the substitution of the native amino acid at position 28 and/or 29 with a negatively charged amino acid (e.g., aspartic acid or glutamic acid) and optionally the addition of a negatively charged amino acid (e.g., aspartic acid or glutamic acid) to the carboxy terminus of the peptide. In an alternative embodiment
20 the native glucagon peptide of SEQ ID NO: 1 is modified by the substitution of the native amino acid at position 29 with a positively charged amino acid (e.g., lysine, arginine or histidine) and optionally the addition of one or two positively charged amino acid (e.g., lysine, arginine or histidine) on the carboxy terminus of the peptide.

In accordance with one embodiment a glucagon analog having improved solubility
25 and stability is provided wherein the analog comprises the amino acid sequence of SEQ ID NO: 34 with the proviso that at least one amino acids at position, 28, or 29 is substituted with an acidic amino acid and/or an additional acidic amino acid is added at the carboxy terminus of SEQ ID NO: 34. In one embodiment the acidic amino acids are independently selected from the group consisting of Asp, Glu, cysteic acid
30 and homocysteic acid.

In accordance with one embodiment a glucagon agonist having improved solubility and stability is provided wherein the agonist comprises the amino acid sequence of SEQ ID NO: 33, wherein at least one of the amino acids at positions 27,

28 or 29 is substituted with a non-native amino acid residue (i.e. at least one amino acid present at position 27, 28 or 29 of the analog is an acid amino acid different from the amino acid present at the corresponding position in SEQ ID NO: 1). In accordance with one embodiment a glucagon agonist is provided comprising the
5 sequence of SEQ ID NO: 33 with the proviso that when the amino acid at position 28 is asparagine and the amino acid at position 29 is threonine, the peptide further comprises one to two amino acids, independently selected from the group consisting of Lys, Arg, His, Asp or Glu, added to the carboxy terminus of the glucagon peptide.

It has been reported that certain positions of the native glucagon peptide can
10 be modified while retaining at least some of the activity of the parent peptide. Accordingly, applicants anticipate that one or more of the amino acids located at positions at positions 2, 5, 7, 10, 11, 12, 13, 14, 16, 17, 18, 19, 20, 21, 24, 27 or 29 of the peptide of SEQ ID NO: 11 can be substituted with an amino acid different from that present in the native glucagon peptide, and still retain the enhanced potency,
15 physiological pH stability and biological activity of the parent glucagon peptide. For example, in accordance with one embodiment the methionine residue present at position 27 of the native peptide is changed to leucine or norleucine to prevent oxidative degradation of the peptide.

In one embodiment a glucagon analog of SEQ ID NO: 33 is provided wherein
20 1 to 6 amino acids, selected from positions 1, 2, 5, 7, 10, 11, 12, 13, 14, 16, 17, 18, 19, 20, 21 or 24 of the analog differ from the corresponding amino acid of SEQ ID NO: 1. In accordance with another embodiment a glucagon analog of SEQ ID NO: 33 is provided wherein 1 to 3 amino acids selected from positions 1, 2, 5, 7, 10, 11, 12, 13, 14, 16, 17, 18, 19, 20, 21 or 24 of the analog differ from the corresponding amino acid
25 of SEQ ID NO: 1. In another embodiment, a glucagon analog of SEQ ID NO: 7, SEQ ID NO: 8 or SEQ ID NO: 34 is provided wherein 1 to 2 amino acids selected from positions 1, 2, 5, 7, 10, 11, 12, 13, 14, 16, 17, 18, 19, 20, 21 or 24 of the analog differ from the corresponding amino acid of SEQ ID NO: 1, and in a further embodiment those one to two differing amino acids represent conservative amino acid substitutions
30 relative to the amino acid present in the native sequence (SEQ ID NO: 1). In one embodiment a glucagon peptide of SEQ ID NO: 11 or SEQ ID NO: 13 is provided wherein the glucagon peptide further comprises one, two or three amino acid substitutions at positions selected from positions 2, 5, 7, 10, 11, 12, 13, 14, 16, 17, 18,

19, 20, 21, 24, 27 or 29. In one embodiment the substitutions at positions 2, 5, 7, 10, 11, 12, 13, 14, 16, 17, 18, 19, 20, 27 or 29 are conservative amino acid substitutions.

In one embodiment a glucagon agonist is provided comprising an analog peptide of SEQ ID NO: 1 wherein the analog differs from SEQ ID NO: 1 by having
5 an amino acid other than serine at position 2 and by having an acidic amino acid substituted for the native amino acid at position 28 or 29 or an acidic amino acid added to the carboxy terminus of the peptide of SEQ ID NO: 1. In one embodiment the acidic amino acid is aspartic acid or glutamic acid. In one embodiment a glucagon
10 analog of SEQ ID NO: 9, SEQ ID NO: 12, SEQ ID NO: 13 or SEQ ID NO: 32 is provided wherein the analog differs from the parent molecule by a substitution at position 2. More particularly, position 2 of the analog peptide is substituted with an amino acid selected from the group consisting of d-serine, alanine, glycine, n-methyl serine and amino isobutyric acid.

In another embodiment a glucagon agonist is provided comprising an analog
15 peptide of SEQ ID NO: 1 wherein the analog differs from SEQ ID NO: 1 by having an amino acid other than histidine at position 1 and by having an acidic amino acid substituted for the native amino acid at position 28 or 29 or an acidic amino acid added to the carboxy terminus of the peptide of SEQ ID NO: 1. In one embodiment the acidic amino acid is aspartic acid or glutamic acid. In one embodiment a glucagon
20 analog of SEQ ID NO: 9, SEQ ID NO: 12, SEQ ID NO: 13 or SEQ ID NO: 32 is provided wherein the analog differs from the parent molecule by a substitution at position 1. More particularly, position 1 of the analog peptide is substituted with an amino acid selected from the group consisting of d-histidine, desaminohistidine, hydroxyl-histidine, acetyl-histidine and homo-histidine.

25 In accordance with one embodiment the modified glucagon peptide comprises a sequence selected from the group consisting of SEQ ID NO: 9, SEQ ID NO: 12, SEQ ID NO: 13 and SEQ ID NO: 32. In a further embodiment a glucagon peptide is provided comprising a sequence of SEQ ID NO: 9, SEQ ID NO: 12, SEQ ID NO: 13 or SEQ ID NO: 32 further comprising one to two amino acids, added to the c-
30 terminus of SEQ ID NO: 9, SEQ ID NO: 12, SEQ ID NO: 13 or SEQ ID NO: 32, wherein the additional amino acids are independently selected from the group consisting of Lys, Arg, His, Asp, Glu, cysteic acid or homocysteic acid. In one embodiment the additional amino acids added to the carboxy terminus are selected

from the group consisting of Lys, Arg, His, Asp or Glu or in a further embodiment the additional amino acids are Asp or Glu.

In another embodiment the glucagon peptide comprises the sequence of SEQ ID NO: 7 or a glucagon agonist analog thereof. In one embodiment the peptide comprising a sequence selected from the group consisting of SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 11, SEQ ID NO: 12 and SEQ ID NO: 13. In another embodiment the peptide comprising a sequence selected from the group consisting of SEQ ID NO: 8, SEQ ID NO: 10 and SEQ ID NO: 11. In one embodiment the glucagon peptide comprises the sequence of SEQ ID NO: 8, SEQ ID NO: 10 and SEQ ID NO: 11 further comprising an additional amino acid, selected from the group consisting of Asp and Glu, added to the c-terminus of the glucagon peptide. In one embodiment the glucagon peptide comprises the sequence of SEQ ID NO: 11 or SEQ ID NO: 13, and in a further embodiment the glucagon peptide comprises the sequence of SEQ ID NO: 11.

In accordance with one embodiment a glucagon agonist is provided comprising a modified glucagon peptide selected from the group consisting of:

NH₂-His-Ser-Gln-Gly-Thr-Phe-Thr-Ser-Asp-Tyr-Ser-Lys-Tyr-Leu-Xaa-Ser-Arg-Arg-Ala-Gln-Asp-Phe-Val-Gln-Trp-Leu-Xaa-Xaa-Xaa-R (SEQ ID NO: 34),
 NH₂-His-Ser-Gln-Gly-Thr-Phe-Thr-Ser-Asp-Tyr-Ser-Lys-Tyr-Leu-Asp-Ser-Arg-Arg-Ala-Gln-Asp-Phe-Val-Gln-Trp-Leu-Met-Asp-Thr-R (SEQ ID NO: 11) and
 NH₂-His-Ser-Gln-Gly-Thr-Phe-Thr-Ser-Asp-Tyr-Ser-Xaa-Tyr-Leu-Glu-Ser-Arg-Arg-Ala-Gln-Asp-Phe-Val-Gln-Trp-Leu-Met-Asp-Thr-R (SEQ ID NO: 13)

wherein Xaa at position 15 is Asp, Glu, cysteic acid, homoglutamic acid or homocysteic acid, the Xaa at position 28 is Asn or an acidic amino acid and the Xaa at position 29 is Thr or an acidic amino acid and R is an acidic amino acid, COOH or CONH₂, with the proviso that an acidic acid residue is present at one of positions 28, 29 or 30. In one embodiment R is COOH, and in another embodiment R is CONH₂.

The present disclosure also encompasses glucagon fusion peptides wherein a second peptide has been fused to the c-terminus of the glucagon peptide to enhance the stability and solubility of the glucagon peptide. More particularly, the fusion glucagon peptide may comprise a glucagon agonist analog comprising a glucagon peptide NH₂-His-Ser-Gln-Gly-Thr-Phe-Thr-Ser-Asp-Tyr-Ser-Lys-Tyr-Leu-Xaa-Ser-Arg-Arg-Ala-Gln-Asp-Phe-Val-Gln-Trp-Leu-Xaa-Xaa-Xaa-R (SEQ ID NO: 34),

wherein R is an acidic amino acid or a bond and an amino acid sequence of SEQ ID NO: 20 (GPSSGAPPPS), SEQ ID NO: 21 (KRNRNNIA) or SEQ ID NO: 22 (KRNR) linked to the carboxy terminal amino acid of the glucagon peptide. In one embodiment the glucagon peptide is selected from the group consisting of SEQ ID NO: 33, SEQ ID NO: 7 or SEQ ID NO: 8 further comprising an amino acid sequence of SEQ ID NO: 20 (GPSSGAPPPS), SEQ ID NO: 21 (KRNRNNIA) or SEQ ID NO: 22 (KRNR) linked to the carboxy terminal amino acid of the glucagon peptide. In one embodiment the glucagon fusion peptide comprises SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5 and SEQ ID NO: 6 or a glucagon agonist analog thereof, further comprising an amino acid sequence of SEQ ID NO: 20 (GPSSGAPPPS), SEQ ID NO: 21 (KRNRNNIA) or SEQ ID NO: 22 (KRNR) linked to amino acid 29 of the glucagon peptide. In accordance with one embodiment the fusion peptide further comprises a PEG chain linked to an amino acid at position 16, 17, 21 or 24, wherein the PEG chain is selected from the range of 500 to 40,000 Daltons. In one embodiment the amino acid sequence of SEQ ID NO: 20 (GPSSGAPPPS), SEQ ID NO: 21 (KRNRNNIA) or SEQ ID NO: 22 (KRNR) is bound to amino acid 29 of the glucagon peptide through a peptide bond. In one embodiment the glucagon peptide portion of the glucagon fusion peptide comprises a sequence selected from the group consisting of SEQ ID NO: 10, SEQ ID NO: 11 and SEQ ID NO: 13. In one embodiment the glucagon peptide portion of the glucagon fusion peptide comprises the sequence of SEQ ID NO: 11 or SEQ ID NO: 13, wherein a PEG chain is linked at position 21 or 24, respectively.

In another embodiment the glucagon peptide sequence of the fusion peptide comprises the sequence of SEQ ID NO: 11, further comprising an amino acid sequence of SEQ ID NO: 20 (GPSSGAPPPS), SEQ ID NO: 21 (KRNRNNIA) or SEQ ID NO: 22 (KRNR) linked to amino acid 29 of the glucagon peptide. In one embodiment the glucagon fusion peptide comprises a sequence selected from the group consisting of SEQ ID NO: 24, SEQ ID NO: 25 and SEQ ID NO: 26. Typically the fusion peptides of the present invention will have a c-terminal amino acid with the standard carboxylic acid group. However, analogs of those sequences wherein the C-terminal amino acid has an amide substituted for the carboxylic acid are also encompassed as embodiments. In accordance with one embodiment the fusion glucagon peptide comprises a glucagon agonist analog selected from the group

consisting of SEQ ID NO: 10, SEQ ID NO: 11 and SEQ ID NO: 13, further comprising an amino acid sequence of SEQ ID NO: 23 (GPSSGAPPPS-CONH₂) linked to amino acid 29 of the glucagon peptide.

The glucagon agonists of the present invention can be further modified to improve the peptide's solubility and stability in aqueous solutions while retaining the biological activity of the glucagon peptide. In accordance with one embodiment, introduction of hydrophilic groups at one or more positions selected from positions 16, 17, 20, 21, 24 and 29 of the peptide of SEQ ID NO: 11, or a glucagon agonist analog thereof, are anticipated to improve the solubility and stability of the pH stabilize glucagon analog. More particularly, in one embodiment the glucagon peptide of SEQ ID NO: 10, SEQ ID NO: 11, SEQ ID NO: 13, or SEQ ID NO: 32 is modified to comprise one or more hydrophilic groups covalently linked to the side chains of amino acids present at positions 21 and 24 of the glucagon peptide.

In accordance with one embodiment, the glucagon peptide of SEQ ID NO: 11 is modified to contain one or more amino acid substitution at positions 16, 17, 20, 21, 24 and/or 29, wherein the native amino acid is substituted with an amino acid having a side chain suitable for crosslinking with hydrophilic moieties, including for example, PEG. The native peptide can be substituted with a naturally occurring amino acid or a synthetic (non-naturally occurring) amino acid. Synthetic or non-naturally occurring amino acids refer to amino acids that do not naturally occur *in vivo* but which, nevertheless, can be incorporated into the peptide structures described herein.

In one embodiment, a glucagon agonist of SEQ ID NO: 10, SEQ ID NO: 11 or SEQ ID NO: 13 is provided wherein the native glucagon peptide sequence has been modified to contain a naturally occurring or synthetic amino acid in at least one of positions 16, 17, 21 and 24 of the native sequence, wherein the amino acid substitute further comprises a hydrophilic moiety. In one embodiment the substitution is at position 21 or 24, and in a further embodiment the hydrophilic moiety is a PEG chain. In one embodiment the glucagon peptide of SEQ ID NO: 11 is substituted with at least one cysteine residue, wherein the side chain of the cysteine residue is further modified with a thiol reactive reagent, including for example, maleimido, vinyl sulfone, 2-pyridylthio, haloalkyl, and haloacyl. These thiol reactive reagents may contain carboxy, keto, hydroxyl, and ether groups as well as other hydrophilic

moieties such as polyethylene glycol units. In an alternative embodiment, the native glucagon peptide is substituted with lysine, and the side chain of the substituting lysine residue is further modified using amine reactive reagents such as active esters (succinimido, anhydride, etc) of carboxylic acids or aldehydes of hydrophilic moieties such as polyethylene glycol. IN one embodiment the glucagon peptide is selected from the group consisting of SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18 and SEQ ID NO: 19.

In accordance with one embodiment the pegylated glucagon peptide comprises two or more polyethylene chains covalently bound to the glucagon peptide wherein the total molecular weight of the glucagon chains is about 1,000 to about 5,000 Daltons. In one embodiment the pegylated glucagon agonist comprises a peptide of SEQ ID NO: 6, wherein a PEG chain is covalently linked to the amino acid residue at position 21 and at position 24, and wherein the combined molecular weight of the two PEG chains is about 1,000 to about 5,000 Daltons. In another embodiment the pegylated glucagon agonist comprises a peptide of SEQ ID NO: 6, wherein a PEG chain is covalently linked to the amino acid residue at position 21 and at position 24, and wherein the combined molecular weight of the two PEG chains is about 5,000 to about 20,000 Daltons.

The polyethylene glycol chain may be in the form of a straight chain or it may be branched. In accordance with one embodiment the polyethylene glycol chain has an average molecular weight selected from the range of about 500 to about 40,000 Daltons. In one embodiment the polyethylene glycol chain has a molecular weight selected from the range of about 500 to about 5,000 Daltons. In another embodiment the polyethylene glycol chain has a molecular weight of about 20,000 to about 40,000 Daltons.

As described in detail in the Examples, the glucagon agonists of the present invention have enhanced biophysical stability and aqueous solubility in solutions of physiological pH, while retaining or demonstrating enhanced bioactivity relative to the native peptide. Accordingly, the glucagon agonists of the present invention are believed to be suitable for any use that has previously been described for the native glucagon peptide. Therefore, the modified glucagon peptides described herein can be used to treat hypoglycemia, to increase blood glucose level, to induce temporary paralysis of the gut for radiological uses, to reduce and maintain body weight, as

adjunctive therapy with insulin, or to treat other metabolic diseases that result from low blood levels of glucagon.

One aspect of the present disclosure is directed to a pre-formulated aqueous solution of the presently disclosed glucagon agonist for use in treating hypoglycemia.

5 The improved stability and/or solubility of the agonist compositions described herein allow for the preparation of pre-formulated aqueous solutions of glucagon for rapid administration and treatment of hypoglycemia. Accordingly, in one embodiment a solution comprising a glucagon agonist of the present invention is provided for administration to a patient suffering from hypoglycemia. In one embodiment a
10 solution comprising a pegylated glucagon agonist as disclosed herein is provided for administration to a patient suffering from hypoglycemia, wherein the total molecular weight of the PEG chains linked to the pegylated glucagon agonist is between about 500 to about 5,000 Daltons. In one embodiment the pegylated glucagon agonist comprises a peptide selected from the group consisting of SEQ ID NO: 14, SEQ ID
15 NO: 15, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18 and SEQ ID NO: 19, and glucagon agonist analogs of thereof, wherein the side chain of an amino acid residue at position 16, 17, 21 or 24 of said glucagon peptide is covalently bound to the polyethylene glycol chain. In one embodiment, the pegylated glucagon agonist comprises the peptide of SEQ ID NO: 16, wherein the amino acid residue at position
20 21 of the peptide is covalently linked to polyethylene glycol. In one embodiment, the pegylated glucagon agonist comprises the peptide of SEQ ID NO: 17, wherein the amino acid residue at position 24 of the peptide is covalently linked to polyethylene glycol.

The method of treating hypoglycemia in accordance with the present invention
25 comprises the steps of administering the presently disclosed glucagon agonists to a patient using any standard route of administration, including parenterally, such as intravenously, intraperitoneally, subcutaneously or intramuscularly, intrathecally, transdermally, rectally, orally, nasally or by inhalation. In one embodiment the composition is administered subcutaneously or intramuscularly. In one embodiment,
30 the composition is administered parenterally and the glucagon composition is prepackaged in a syringe. Advantageously, the aqueous stable glucagon analogs disclosed herein exhibit superior stability and solubility in aqueous buffers in the broadest pH range used for pharmacological purposes, relative to native glucagon.

The use of the stabilized glucagon analogs disclosed herein allow for the preparation and storage of glucagon agonist solutions at physiological pH for long periods of time.

Applicants have discovered that pegylated glucagon peptides can be prepared that retain the parent peptide's bioactivity and specificity. However, increasing the length of the PEG chain, or attaching multiple PEG chains to the peptide, such that the total molecular weight of the linked PEG is greater than 5,000 Daltons, begins to delay the time action of the modified glucagon. In accordance with one embodiment, a glucagon peptide of SEQ ID NO: 11 or SEQ ID NO: 13, or a glucagon agonist analog thereof, is provided wherein the peptide comprises one or more polyethylene glycol chains, wherein the total molecular weight of the linked PEG is greater than 5,000 Daltons, and in one embodiment is greater than 10,000 Daltons, but less than 40,000 Daltons. Such modified glucagon peptides have a delayed time of activity but without loss of the bioactivity. Accordingly, such compounds can be administered prophylactically to extend the effect of the administered glucagon peptide.

Glucagon peptides that have been modified to be covalently bound to a PEG chain having a molecular weight of greater than 10,000 Daltons can be administered in conjunction with insulin to buffer the actions of insulin and help to maintain stable blood glucose levels in diabetics. The modified glucagon peptides of the present disclosure can be co-administered with insulin as a single composition, simultaneously administered as separate solutions, or alternatively, the insulin and the modified glucagon peptide can be administered at different time relative to one another. In one embodiment the composition comprising insulin and the composition comprising the modified glucagon peptide are administered within 12 hours of one another. The exact ratio of the modified glucagon peptide relative to the administered insulin will be dependent in part on determining the glucagon levels of the patient, and can be determined through routine experimentation.

In accordance with one embodiment a composition is provided comprising insulin and a modified glucagon peptide selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5 and glucagon agonist analogs thereof, wherein the modified glucagon peptide further comprises a polyethylene glycol chain covalently bound to an amino acid side chain at position 16, 17, 21 or 24. In one embodiment the composition is an aqueous solution comprising

insulin and the glucagon analog. In embodiments wherein the glucagon peptide comprises the sequence of SEQ ID NO: 11 or SEQ ID NO: 13, the peptide may further comprise a polyethylene glycol chain covalently bound to an amino acid side chain at position 16, 17, 21 or 24. In one embodiment the molecular weight of the PEG chain of the modified glucagon peptide is greater than 10,000 Daltons. In one embodiment the pegylated glucagon peptide comprises a peptide selected from the group consisting of SEQ ID NO: 11 and SEQ ID NO: 13 wherein the side chain of an amino acid residue at position 21 or 24 of said glucagon peptide is covalently bound to the polyethylene glycol chain. In one embodiment the polyethylene glycol chain has a molecular weight of about 10,000 to about 40,000.

In accordance with one embodiment the modified glucagon peptides disclosed herein are used to induce temporary paralysis of the intestinal tract. This method has utility for radiological purposes and comprises the step of administering an effective amount of a pharmaceutical composition comprising a pegylated glucagon peptide, a glucagon peptide comprising a c-terminal extension or a dimer of such peptides. In one embodiment the glucagon peptide comprises a sequence selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, SEQ ID NO: 12 and SEQ ID NO: 13. In one embodiment the glucagon peptide further comprises a PEG chain, of about 1,000 to 40,000 Daltons is covalently bound to an amino acid residue at position 21 or 24. In one embodiment the glucagon peptide is selected from the group consisting of SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16 and SEQ ID NO: 17. In one embodiment the PEG chain has a molecular weight of about 500 to about 5,000 Daltons.

In a further embodiment the composition used to induce temporary paralysis of the intestinal tract comprises a first modified glucagon peptide and a second modified glucagon peptide, wherein the first modified peptide comprises a sequence selected from the group consisting of SEQ ID NO: 10, SEQ ID NO: 11 and SEQ ID NO: 13, optionally linked to a PEG chain of about 500 to about 5,000 Daltons, and the second peptide comprises a covalently linked PEG chain of about 10,000 to about 40,000 Daltons. In this embodiment the PEG chain of each peptide is covalently bound to an amino acid residue at either position 21 or 24 of the respective peptide, and independent of one another.

Oxyntomodulin, a naturally occurring digestive hormone found in the small intestine, has been reported to cause weight loss when administered to rats or humans (see Diabetes 2005;54:2390-2395). Oxyntomodulin is a 37 amino acid peptide that contains the 29 amino acid sequence of glucagon (i.e. SEQ ID NO: 1) followed by an 8 amino acid carboxy terminal extension of SEQ ID NO: 23 (KRNRNNIA).
5 Accordingly, applicants believe that the bioactivity of oxyntomodulin can be retained (i.e. appetite suppression and induced weight loss/weight maintenance), while improving the solubility and stability of the compound and improving the pharmacokinetics, by substituting the glucagon peptide portion of oxyntomodulin with the modified glucagon peptides disclosed herein. In addition applicants also believe that a truncated oxyntomodulin molecule comprising a glucagon peptide of the invention, having the terminal four amino acids of oxyntomodulin removed, will also be effective in suppressing appetite and inducing weight loss/weight maintenance.
10
15 Accordingly, the present invention also encompasses the modified glucagon peptides of the present invention that have a carboxy terminal extension of SEQ ID NO: 21 (KRNRNNIA) or SEQ ID NO: 22. In accordance with one embodiment a glucagon agonist analog of SEQ ID NO: 33, further comprising the amino acid sequence of SEQ ID NO: 21 (KRNRNNIA) or SEQ ID NO: 22 linked to amino acid 20 29 of the glucagon peptide, is administered to individuals to induce weight loss or prevent weight gain. In accordance with one embodiment a glucagon agonist analog of SEQ ID NO: 11 or SEQ ID NO: 13, further comprising the amino acid sequence of SEQ ID NO: 21 (KRNRNNIA) or SEQ ID NO: 22 linked to amino acid 25 weight gain. In another embodiment a method of reducing weight gain or inducing weight loss in an individual comprises administering an effective amount of a composition comprising a glucagon agonist comprising a glucagon peptide selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4 or SEQ ID NO: 5, wherein amino acid 29 of the glucagon peptide is bound to a second 30 peptide through a peptide bond, and said second peptide comprises the sequence of SEQ ID NO: 24 (KRNRNNIA) or SEQ ID NO: 25, and wherein a PEG chain of about 1,000 to 40,000 Daltons is covalently bound to an amino acid residue at position 21 and/or 24. In one embodiment the glucagon peptide segment of the glucagon agonist

is selected from the group consisting of SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18 and SEQ ID NO: 19, wherein a PEG chain of about 1,000 to 40,000 Daltons is covalently bound to an amino acid residue at position 16, 17, 21 or 24.

5 Exendin-4, is a peptide made up of 39 amino acids. It is a powerful stimulator of a receptor known as GLP-1. This peptide has also been reported to suppress appetite and induce weight loss. Applicants have found that the terminal sequence of Exendin-4 when added at the carboxy terminus of glucagon improves the solubility and stability of glucagon without compromising the bioactivity of glucagon. In one
10 embodiment the terminal ten amino acids of Exendin-4 (i.e. the sequence of SEQ ID NO: 20 (GPSSGAPPPS)) are linked to the carboxy terminus of a glucagon peptide of the present disclosure. These fusion proteins are anticipated to have pharmacological activity for suppressing appetite and inducing weight loss/weight maintenance. In one
15 embodiment the terminal amino acid of the SEQ ID NO: 20 extension comprises an amide group in place of the carboxy group (i.e., SEQ ID NO: 23) and this sequence is linked to the carboxy terminus of a glucagon peptide of the present disclosure.

 In one embodiment a method of reducing weight gain or inducing weight loss in an individual comprises administering an effective amount of a composition comprising a glucagon agonist comprising a glucagon peptide of SEQ ID NO: 33
20 wherein amino acid 29 of the glucagon peptide is bound to a second peptide through a peptide bond, and said second peptide comprises the sequence of SEQ ID NO: 20 (GPSSGAPPPS) or SEQ ID NO: 23. In one embodiment the glucagon peptide is selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ
25 ID NO: 10, SEQ ID NO: 11, SEQ ID NO: 12 and SEQ ID NO: 13 wherein amino acid 29 of the glucagon peptide is bound to a second peptide through a peptide bond, and said second peptide comprises the sequence of SEQ ID NO: 20 (GPSSGAPPPS) or SEQ ID NO: 23. In one embodiment the glucagon peptide of the glucagon agonist is selected from the group consisting of SEQ ID NO: 11 and SEQ ID NO: 13. In one
30 embodiment the glucagon peptide segment of the fusion peptide is selected from the group consisting of SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16 and SEQ ID NO: 17, wherein the molecular weight of the PEG chain is selected from the range of 500 to 40,000 Daltons. More particularly, in one embodiment the glucagon peptide of

the fusion peptide is selected from the group consisting of SEQ ID NO: 16 and SEQ ID NO: 17 wherein the molecular weight of the PEG chain is selected from the range of 1,000 to 5,000.

In another embodiment a composition is administered to a patient to suppress appetite, prevent weight gain and/or induce weight loss by the administration of a pharmaceutical composition comprising a first pegylated glucagon peptide and a second pegylated glucagon peptide, wherein the first and second peptide are fusion peptides comprising a c-terminal peptide extension comprising SEQ ID NO: 20 (GPSSGAPPPS) or SEQ ID NO: 23. The first pegylated glycogen peptide comprising a covalently linked PEG of about 500 to about 10,000 Daltons and the second pegylated glucagon peptide comprising a covalently linked PEG chain of about 10,000 to about 40,000 Daltons.

The present disclosure also encompasses multimers of the modified glucagon peptides disclosed herein. Two or more of the modified glucagon peptides can be linked together using standard linking agents and procedures known to those skilled in the art. For example, dimers can be formed between two modified glucagon peptides through the use of bifunctional thiol crosslinkers and bi-functional amine crosslinkers, particularly for the glucagon peptides that have been substituted with cysteine, lysine ornithine, homocysteine or acetyl phenylalanine residues (e.g. SEQ ID NO: 4 and SEQ ID NO: 5). The dimer can be a homodimer or alternatively can be a heterodimer. In one embodiment the dimer comprises a homodimer of a glucagon fusion peptide wherein the glucagon peptide portion comprises an agonist analog of SEQ ID NO: 11 and an amino acid sequence of SEQ ID NO: 20 (GPSSGAPPPS), SEQ ID NO: 21 (KRNRRNNIA) or SEQ ID NO: 22 (KRNRR) linked to amino acid 29 of the glucagon peptide. In another embodiment the dimer comprises a homodimer of a glucagon agonist analog of SEQ ID NO: 11, wherein the glucagon peptide further comprises a polyethylene glycol chain covalently bound to position 21 or 24 of the glucagon peptide.

In accordance with one embodiment a dimer is provided comprising a first glucagon peptide bound to a second glucagon peptide via a linker, wherein the first glucagon peptide comprises a peptide selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10 and SEQ ID NO: 11 and the second glucagon peptide comprises SEQ ID NO: 33. Furthermore,

with regards to the second glucagon peptide, when the amino acid at position 28 is asparagine and the amino acid at position 29 is threonine, the second glucagon peptide further comprises one to two amino acids (independently selected from the group consisting of Lys, Arg, His, Asp or Glu), added to the carboxy terminus of the second
5 glucagon glucagon peptide, and pharmaceutically acceptable salts of said glucagon polypeptides.

In accordance with another embodiment a dimer is provided comprising a first glucagon peptide bound to a second glucagon peptide via a linker, wherein said first glucagon peptide is selected from the group consisting of SEQ ID NO: 2, SEQ ID
10 NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, SEQ ID NO: 12 and SEQ ID NO: 13 and the second glucagon peptide is independently selected from the group consisting of SEQ ID NO: 11 and SEQ ID NO: 13, and pharmaceutically acceptable salts of said glucagon polypeptides. In one embodiment the first glucagon peptide is selected from the group consisting of
15 SEQ ID NO: 7 and the second glucagon peptide is independently selected from the group consisting of SEQ ID NO: 11, SEQ ID NO: 12 and SEQ ID NO: 13. In one embodiment the dimer is formed between two peptides wherein each peptide comprising the amino acid sequence of SEQ ID NO: 11.

In accordance with one embodiment a pharmaceutical composition is provided
20 wherein the composition comprises a glucagon agonist analog of the present disclosure, or pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier. In one embodiment the pharmaceutical composition comprises a 1mg/ml concentration of the glucagon agonist analog and 10-50mM Triethanolamine at pH 7.0-8.5. In one embodiment the pharmaceutical composition comprises a
25 1mg/ml concentration of the glucagon agonist analog and 20mM Triethanolamine at pH 8.5.

The modified glucagon peptides of the present invention can be provided in accordance with one embodiment as part of a kit. In one embodiment a kit for administering a glucagon agonist to a patient in need thereof is provided wherein the
30 kit comprises any of the glucagon peptides of the invention in aqueous solution. Exemplary glucagon peptides for inclusion in such kits include a glucagon peptide selected from the group consisting of 1) a glucagon peptide comprising the sequence of SEQ ID NO: 7, SEQ ID NO: 10, SEQ ID NO: 11 or SEQ ID NO: 13 or SEQ ID

NO: 33; 2) a glucagon fusion peptide comprising a glucagon agonist analog of SEQ ID NO: 11 or SEQ ID NO: 13 or SEQ ID NO: 33, and an amino acid sequence of SEQ ID NO: 20 (GPSSGAPPPS), SEQ ID NO: 21 (KRNRNNIA) or SEQ ID NO: 22 (KRNR) linked to amino acid 29 of the glucagon peptide; and 3) a pegylated glucagon peptide of SEQ ID NO: 11 or SEQ ID NO: 13 or SEQ ID NO: 33, further comprising an amino acid sequence of SEQ ID NO: 20 (GPSSGAPPPS), SEQ ID NO: 21 (KRNRNNIA) or SEQ ID NO: 22 (KRNR) linked to amino acid 29 of the glucagon peptide, wherein the PEG chain covalently bound to position 16, 17, 21 or 24 has a molecular weight of about 500 to about 40,000 Daltons. In one embodiment the kit is provided with a device for administering the glucagon composition to a patient, e.g. syringe needle, pen device, jet injector or other needle-free injector. The kit may alternatively or in addition include one or more of a variety of containers, e.g., vials, tubes, bottles, single or multi-chambered pre-filled syringes, cartridges, infusion pumps (external or implantable), jet injectors, pre-filled pen devices and the like, optionally containing the glucagon peptide. Preferably, the kits will also include instructions for use. In accordance with one embodiment the device of the kit is an aerosol dispensing device, wherein the composition is prepackaged within the aerosol device. In another embodiment the kit comprises a syringe and a needle, and in one embodiment the glucagon composition is prepackaged within the syringe.

The compounds of this invention may be prepared by standard synthetic methods, recombinant DNA techniques, or any other methods of preparing peptides and fusion proteins. Although certain non-natural amino acids cannot be expressed by standard recombinant DNA techniques, techniques for their preparation are known in the art. Compounds of this invention that encompass non-peptide portions may be synthesized by standard organic chemistry reactions, in addition to standard peptide chemistry reactions when applicable.

EXAMPLES

General Synthesis Protocol:

Glucagon analogs were synthesized using HBTU-activated "Fast Boc" single coupling starting from 0.2mmole of Boc Thr(OBzl)Pam resin on a modified Applied Biosystem 430 A peptide synthesizer. Boc amino acids and HBTU were obtained from Midwest Biotech (Fishers, IN). Side chain protecting groups used were:

Arg(Tos), Asn(Xan), Asp(OcHex), Cys(pMeBzl), His(Bom), Lys(2Cl-Z), Ser(OBzl), Thr(OBzl), Tyr(2Br-Z), and Trp(CHO). The side-chain protecting group on the N-terminal His was Boc.

Each completed peptidyl resin was treated with a solution of 20% piperidine in dimethylformamide to remove the formyl group from the tryptophan. Liquid hydrogen fluoride cleavages were performed in the presence of p-cresol and dimethyl sulfide. The cleavage was run for 1 hour in an ice bath using an HF apparatus (Penninsula Labs). After evaporation of the HF, the residue was suspended in diethyl ether and the solid materials were filtered. Each peptide was extracted into 30-70ml aqueous acetic acid and a diluted aliquot was analyzed by HPLC [Beckman System Gold, 0.46x5cm Zorbax C8, 1ml/min, 45C, 214nm, A buffer =0.1%TFA, B=0.1%TFA/90%acetonitrile, gradient of 10% to 80%B over 10min].

Purification was done on a FPLC over a 2.2 x 25 cm Kromasil C18 column while monitoring the UV at 214nm and collecting 5 minute fractions. The homogeneous fractions were combined and lyophilized to give a product purity of >95%. The correct molecular mass and purity were confirmed using MALDI-mass spectral analysis.

General Pegylation Protocol: (Cys-maleimido)

Typically, the glucagon Cys analog is dissolved in phosphate buffered saline (5-10mg/ml) and 0.01M ethylenediamine tetraacetic acid is added (10-15% of total volume). Excess (2-fold) maleimido methoxyPEG reagent (Nektar) is added and the reaction stirred at room temp while monitoring reaction progress by HPLC. After 8-24hrs, the reaction mixture, is acidified and loaded onto a preparative reverse phase column for purification using 0.1%TFA/acetonitrile gradient. The appropriate fractions were combined and lyophilized to give the desired pegylated analogs.

EXAMPLE 1

Synthesis of Glucagon Cys¹⁷(1-29) and Similar MonoCys Analogs

0.2mmole Boc Thr(OBzl) Pam resin (SynChem Inc) in a 60ml reaction vessel and the following sequence was entered and run on a modified Applied Biosystems 430A Peptide Synthesizer using FastBoc HBTU-activated single couplings.

HSQGTFTSDYSKYLDSCRAQDFVQWLMNT (SEQ ID NO: 27)

The following side chain protecting groups were used: Arg(Tos), Asp(OcHex), Asn(Xan), Cys(pMeBzl), Glu(OcHex), His(Boc), Lys(2Cl-Z), Ser(Bzl), Thr(Bzl), Trp(CHO), and Tyr(Br-Z). The completed peptidyl resin was treated with 20% piperidine/dimethylformamide to remove the Trp formyl protection then transferred to an HF reaction vessel and dried in vacuo. 1.0ml p-cresol and 0.5 ml dimehyl sulfide were added along with a magnetic stir bar. The vessel was attached to the HF apparatus (Penninsula Labs), cooled in a dry ice/methanol bath, evacuated, and aprox. 10ml liquid hydrogen fluoride was condensed in. The reaction was stirred in an ice bath for 1hr then the HF was removed in vacuo. The residue was suspended in ethyl ether; the solids were filtered, washed with ether, and the peptide extracted into 50 ml aqueous acetic acid. An analytical HPLC was run [0.46 x 5 cm Zorbax C8, 1 ml/min, 45C, 214nm, A buffer of 0.1%TFA, B buffer of 0.1%TFA/90%ACN, gradient=10%B to 80%B over 10min.] with a small sample of the cleavage extract. The remaining extract was loaded onto a 2.2 x 25cm Kromasil C18 preparative reverse phase column and an acetonitrile gradient was run using a Pharmacia FPLC system. 5min fractions were collected while monitoring the UV at 214nm (2.0A). A=0.1%TFA, B=0.1%TFA/50%acetonitrile. Gradient = 30%B to 100%B over 450min.

The fractions containing the purest product (48-52) were combined frozen, and lyophilized to give 30.1mg. An HPLC analysis of the product demonstrated a purity of >90% and MALDI mass spectral analysis demonstrated the desired mass of 3429.7. Glucagon Cys²¹, Glucagon Cys²⁴, and Glucagon Cys²⁹ were similarly prepared.

EXAMPLE 2

25 Synthesis of Glucagon-Cex and Other C-Terminal Extended Analogs.

285mg (0.2mmole) methoxybenzhydrylamine resin (Midwest Biotech) was placed in a 60ml reaction vessel and the following sequence was entered and run on a modified Applied Biosystems 430A peptide synthesizer using FastBoc HBTU-activated single couplings.

30 HSQGTFTSDYSKYLSRRAQDFVQWLMNTGPSSGAPPPS (SEQ ID NO: 28)

The following side chain protecting groups were used: Arg(Tos), Asp(OcHex), Asn(Xan), Cys(pMeBzl), Glu(OcHex), His(Boc), Lys(2Cl-Z), Ser(Bzl), Thr(Bzl),

Trp(CHO), and Tyr(Br-Z). The completed peptidyl resin was treated with 20% piperidine/dimethylformamide to remove the Trp formyl protection then transferred to HF reaction vessel and dried in vacuo. 1.0ml p-cresol and 0.5 ml dimethyl sulfide were added along with a magnetic stir bar. The vessel was attached to the HF apparatus (Pennisula Labs), cooled in a dry ice/methanol bath, evacuated, and approx. 10ml liquid hydrogen fluoride was condensed in. The reaction was stirred in an ice bath for 1hr then the HF was removed in vacuo. The residue was suspended in ethyl ether; the solids were filtered, washed with ether, and the peptide extracted into 50 ml aqueous acetic acid. An analytical HPLC was run [0.46 x 5 cm Zorbax C8, 1 ml/min, 45C, 214nm, A buffer of 0.1%TFA, B buffer of 0.1%TFA/90%ACN, gradient=10%B to 80%B over 10min.] on an aliquot of the cleavage extract. The extract was loaded onto a 2.2 x 25cm Kromasil C18 preparative reverse phase column and an acetonitrile gradient was run for elution using a Pharmacia FPLC system. 5min fractions were collected while monitoring the UV at 214nm (2.0A). A=0.1%TFA, B=0.1%TFA/50%acetonitrile. Gradient = 30%B to 100%B over 450min. Fractions 58-65 were combined, frozen and lyophilized to give 198.1mg.

HPLC analysis of the product showed a purity of greater than 95%. MALDI mass spectral analysis showed the presence of the desired theoretical mass of 4316.7 with the product as a C-terminal amide. Oxyntomodulin and oxyntomodulin-KRNR were similarly prepared as the C-terminal carboxylic acids starting with the appropriately loaded PAM-resin.

EXAMPLE 3

Glucagon Cys¹⁷ Mal-PEG-5K

15.1mg of Glucagon Cys¹⁷(1-29) and 27.3mg methoxy poly(ethyleneglycol) maleimide avg. M.W.5000 (mPEG-Mal-5000,Nektar Therapeutics) were dissolved in 3.5ml phosphate buffered saline (PBS) and 0.5ml 0.01M ethylenediamine tetraacetic acid (EDTA) was added. The reaction was stirred at room temperature and the progress of the reaction was monitored by HPLC analysis [0.46 x 5 cm Zorbax C8, 1ml/min,45C, 214nm (0.5A), A=0.1%TFA, B=0.1%TFA/90%ACN, gradient=10%B to 80%B over 10min.].

After 5 hours, the reaction mixture was loaded onto 2.2 x 25 cm Kromasil C18 preparative reverse phase column. An acetonitrile gradient was run on a Pharmacia

FPLC while monitoring the UV wavelength at 214nm and collecting 5 min fractions. A=0.1%TFA, B=0.1%TFA/50% acetonitrile, gradient= 30%B to 100%B over 450 min. The fractions corresponding to the product were combined, frozen and lyophilized to give 25.9 mg.

- 5 This product was analyzed on HPLC [0.46 x 5 cm Zorbax C8, 1 ml/min, 45C, 214nm (0.5A), A=0.1%TFA, B=0.1%TFA/90%ACN, gradient=10%B to 80%B over 10min.] which showed a purity of aprox. 90%. MALDI (matrix assisted laser desorption ionization) mass spectral analysis showed a broad mass range (typical of PEG derivatives) of 8700 to 9500. This shows an addition to the mass of the starting
10 glucagon peptide (3429) of approximately 5,000 a.m.u.

EXAMPLE 4

Glucagon Cys²¹ Mal-PEG-5K

- 21.6mg of Glucagon Cys²¹(1-29) and 24mg mPEG-MAL-5000 (Nektar
15 Therapeutics) were dissolved in 3.5ml phosphate buffered saline (PBS) and 0.5ml 0.01M ethylene diamine tetraacetic acid (EDTA) was added. The reaction was stirred at room temp. After 2hrs, another 12.7 mg of mPEG-MAL-5000 was added. After 8hrs, the reaction mixture was loaded onto a 2.2 x 25cm Vydac C18 preparative reverse phase column and an acetonitrile gradient was run on a Pharmacia FPLC at 4
20 ml/min while collecting 5min fractions. A=0.1%TFA, B=0.1%TFA/50%ACN. Gradient= 20% to 80%B over 450min.

- The fractions corresponding to the appearance of product were combined frozen and lyophilized to give 34 mg. Analysis of the product by analytical HPLC [0.46 x 5 cm Zorbax C8, 1 ml/min, 45C, 214nm (0.5A), A=0.1%TFA,
25 B=0.1%TFA/90%ACN, gradient=10%B to 80%B over 10min.] showed a homogeneous product that was different than starting glucagon peptide. MALDI (matrix assisted laser desorption ionization) mass spectral analysis showed a broad mass range (typical of PEG analogs) of 8700 to 9700. This shows an addition to the mass of the starting glucagon peptide (3470) of approximately 5,000 a.m.u.

30

EXAMPLE 5

Glucagon Cys²⁴ Mal-PEG-5K

20.1mg Glucagon C²⁴(1-29) and 39.5mg mPEG-Mal-5000 (Nektar Therapeutics) were dissolved in 3.5ml PBS with stirring and 0.5 ml 0.01M EDTA was added. The reaction was stirred at room temp for 7 hrs, then another 40 mg of mPEG-Mal-5000 was added. After approximately 15 hr, the reaction mixture was loaded
5 onto a 2.2 x 25 cm Vydac C18 preparative reverse phase column and an acetonitrile gradient was run using a Pharmacia FPLC. 5 min. fractions were collected while monitoring the UV at 214nm (2.0A). A buffer = 0.1%TFA, B buffer = 0.1%TFA/50%ACN, gradient = 30%B to 100%B over 450min. The fractions corresponding to product were combined, frozen and lyophilized to give 45.8mg.
10 MALDI mass spectral analysis showed a typical PEG broad signal with a maximum at 9175.2 which is approximately 5,000 a.m.u. more than Glucagon C²⁴ (3457.8).

EXAMPLE 6

Glucagon Cys²⁴ Mal-PEG-20K

15 25.7mg of Glucagon Cys²⁴(1-29) and 40.7mg mPEG-Mal-20K (Nektar Therapeutics) were dissolved in 3.5ml PBS with stirring at room temp. and 0.5 ml 0.01M EDTA was added. After 6hrs, the ratio of starting material to product was aprox. 60:40 as determined by HPLC. Another 25.1mg of mPEG-Mal-20K was added and the reaction allowed to stir another 16hrs. The product ratio had not
20 significantly improved, so the reaction mixture was loaded onto a 2.2 x 25 cm Kromasil C18 preparative reverse phase column and purified on a Pharmacia FPLC using a gradient of 30%B to 100%B over 450min. A buffer =0.1%TFA, B buffer = 0.1%TFA/50%ACN, flow = 4ml/min, and 5 min fractions were collected while monitoring the UV at 214nm (2.0A). The fractions containing homogeneous product
25 were combined, frozen and lyophilized to give 25.7 mg. Purity as determined by analytical HPLC was ~90%. A MALDI mass spectral analysis showed a broad peak from 23,000 to 27,000 which is approximately 20,000 a.m.u. more than starting Glucagon C²⁴ (3457.8).

30 EXAMPLE 7


Glucagon Cys²⁹ Mal-PEG-5K

20.0mg of Glucagon Cys²⁹(1-29) and 24.7 mg mPEG-Mal-5000 (Nektar Therapeutics) were dissolved in 3.5 ml PBS with stirring at room temperature and 0.5

ml 0.01M EDTA was added. After 4 hr, another 15.6 mg of mPEG-Mal-5000 was added to drive the reaction to completion. After 8 hrs, the reaction mixture was loaded onto a 2.2 x 25 cm Vydac C18 preparative reverse phase column and an acetonitrile gradient was run on a Pharmacia FPLC system. 5 min fractions were collected while monitoring the UV at 214nm (2.0A). A=0.1%TFA, B=0.1%TFA/50%ACN. Fractions 75-97 were combined frozen and lyophilized to give 40.0 mg of product that is different than recovered starting material on HPLC (fractions 58-63). Analysis of the product by analytical HPLC [0.46 x 5 cm Zorbax C8, 1 ml/min, 45C, 214nm (0.5A), A=0.1%TFA, B=0.1%TFA/90%ACN, gradient=10%B to 80%B over 10min.] showed a purity greater than 95%. MALDI mass spectral analysis showed the presence of a PEG component with a mass range of 8,000 to 10,000 (maximum at 9025.3) which is 5,540 a.m.u. greater than starting material (3484.8).

EXAMPLE 8
Glucagon Cys²⁴ (2-butyrolactone)

To 24.7mg of Glucagon Cys²⁴(1-29) was added 4ml 0.05M ammonium bicarbonate/50%acetonitrile and 5.5 ul of a solution of 2-bromo-4-hydroxybutyric acid- γ -lactone (100ul in 900ul acetonitrile). After 3hrs of stirring at room temperature, another 105 ul of lactone solution was added to the reaction mixture which was stirred another 15hrs. The reaction mixture was diluted to 10ml with 10% aqueous acetic acid and was loaded onto a 2.2 x 25 cm Kromasil C18 preparative reverse phase column. An acetonitrile gradient (20%B to 80%B over 450min) was run on a Pharmacia FPLC while collecting 5min fractions and monitoring the UV at 214nm (2.0A). Flow =4ml/min, A=0.1%TFA, B=0.1%TFA/50%ACN. Fractions 74-77 were combined frozen and lyophilized to give 7.5mg. HPLC analysis showed a purity of 95% and MALDI mass spect analysis showed a mass of 3540.7 or 84 mass units more than starting material. This result consistent with the addition of a single butyrolactone moiety.



EXAMPLE 9
Glucagon Cys²⁴(S-carboxymethyl)

18.1mg of Glucagon Cys²⁴(1-29) was dissolved in 9.4ml 0.1M sodium phosphate buffer (pH=9.2) and 0.6ml bromoacetic acid solution (1.3mg/ml in acetonitrile) was added. The reaction was stirred at room temperature and the reaction progress was followed by analytical HPLC. After 1hr another 0.1ml bromoacetic acid solution was added. The reaction was stirred another 60min. then acidified with aqueous acetic acid and was loaded onto a 2.2 x 25cm Kromasil C18 preparative reverse phase column for purification. An acetonitrile gradient was run on a Pharmacia FPLC (flow = 4ml/min) while collecting 5min fractions and monitoring the UV at 214nm (2.0A). A=0.1%TFA, B=0.1%TFA/50%ACN. Fractions 26-29 were combined frozen and lyophilized to give

several mg of product. Analytical HPLC showed a purity of 90% and MALDI mass spectral analysis confirmed a mass of 3515 for the desired product.



Molecular Weight = 3515.87
 Exact Mass = 3512
 Molecular Formula = C153H224N42O50S2


SEQ ID NO: 30

5

EXAMPLE 10

Glucagon Cys²⁴ maleimido,PEG-3.4K-dimer

- 10 16mg Glucagon Cys²⁴ and 1.02mg Mal-PEG-Mal-3400,
 poly(ethyleneglycol)-bis-maleimide avg. M.W. 3400, (Nektar Therapeutics) were
 dissolved in 3.5 phosphate buffered saline and 0.5ml 0.01M EDTA and the reaction
 was stirred at room temperature. After 16hrs, another 16mg of Glucagon Cys²⁴ was
 added and the stirring continued. After approximately 40hrs, the reaction mixture was
 15 loaded onto a Pharmacia PepRPC 16/10 column and an acetonitrile gradient was run
 on a Pharmacia FPLC while collecting 2min fractions and monitoring the UV at
 214nm (2.0A). Flow=2ml/min, A=0.1%TFA, B=0.1%TFA/50%ACN. Fractions 69-
 74 were combined frozen and lyophilized to give 10.4mg. Analytical HPLC showed
 a purity of 90% and MALDI mass spectral analysis shows a component in the 9500-
 20 11,000 range which is consistent with the desired dimer.



EXAMPLE 11

Glucagon Solubility Assays:

A solution (1mg/ml or 3mg/ml) of glucagon (or an analog) is prepared in 0.01N HCl. 100ul of stock solution is diluted to 1ml with 0.01N HCl and the UV
 5 absorbance (276nm) is determined. The pH of the remaining stock solution is adjusted to pH7 using 200-250ul 0.1M Na₂HPO₄ (pH9.2). The solution is allowed to stand overnight at 4°C then centrifuged. 100ul of supernatant is then diluted to 1ml with 0.01N HCl, and the UV absorbance is determined (in duplicate).

10 The initial absorbance reading is compensated for the increase in volume and the following calculation is used to establish percent solubility:

$$\frac{\text{Final Absorbance}}{\text{Initial Absorbance}} \times 100 = \text{percent soluble}$$

Results are shown in Table 1 wherein Glucagon-Cex represents wild type glucagon (SEQ ID NO: 1) plus a carboxy terminal addition of SEQ ID NO: 20 and Glucagon-
 15 Cex R¹² represents SEQ ID NO: 1 wherein the Lys at position 12 is substituted with Arg and a peptide of SEQ ID NO: 20 is added to the carboxy terminus.

Table 1 Solubility data for glucagon analogs

<u>Analog</u>	<u>Percent Soluble</u>
Glucagon	16
Glucagon-Cex, R12	104
Glucagon-Cex	87
Oxyntomodulin	104
Glucagon, Cys17PEG5K	94
Glucagon, Cys21PEG5K	105
Glucagon, Cys24PEG5K	133

20 The solubility the glucagon agonist analogs, D28, E29, E30, E15D28, D28E30, D28E29, was investigated using the same assay used for the compounds listed in Table 1. The data (shown in Figs. 3 & 4) demonstrates the superior solubility of the D28, E29, E30, E15D28, D28E30, D28E29 analogs relative to native glucagon
 25 at pH values of 5.5 and 7.0. The data presented in Fig. 3 represents the solubility measured after 60 hours at 25 °C, whereas the data presented in Fig. 4 represents the solubility measured after 24 hours at 25 °C and then 24 hours at 4 °C. Fig. 5

represents data regarding the maximum solubility of the glucagon analogs D28, D28E30 and E15D28.

EXAMPLE 12

5 Glucagon Receptor Binding Assay

The affinity of peptides to the glucagon receptor was measured in a competition binding assay utilizing scintillation proximity assay technology. Serial 3-fold dilutions of the peptides made in scintillation proximity assay buffer (0.05 M Tris-HCl, pH 7.5, 0.15 M NaCl, 0.1% w/v bovine serum albumin) were mixed in 96
10 well white/clear bottom plate (Corning Inc., Acton, MA) with 0.05 nM (3-[¹²⁵I]-iodotyrosyl) Tyr¹⁰ glucagon (Amersham Biosciences, Piscataway, NJ), 1-6 micrograms per well, plasma membrane fragments prepared from cells over-expressing human glucagon receptor, and 1 mg/well polyethyleneimine-treated wheat germ agglutinin type A scintillation proximity assay beads (Amersham Biosciences,
15 Piscataway, NJ). Upon 5 min shaking at 800 rpm on a rotary shaker, the plate was incubated 12h at room temperature and then read on MicroBeta1450 liquid scintillation counter (Perkin-Elmer, Wellesley, MA). Non-specifically bound (NSB) radioactivity was measured in the wells with 4 times greater concentration of “cold” native ligand than the highest concentration in test samples and total bound
20 radioactivity was detected in the wells with no competitor. Percent specific binding was calculated as following: % Specific Binding = ((Bound-NSB)/(Total bound-NSB)) X 100. IC₅₀ values were determined by using Origin software (OriginLab, Northampton, MA).

25 EXAMPLE 13

Functional Assay- cAMP Synthesis

The ability of glucagon analogs to induce cAMP was measured in a firefly luciferase-based reporter assay. HEK293 cells co-transfected with either glucagon- or GLP-1 receptor and luciferase gene linked to cAMP responsive element were serum
30 deprived by culturing 16h in DMEM (Invitrogen, Carlsbad, CA) supplemented with 0.25% Bovine Growth Serum (HyClone, Logan, UT) and then incubated with serial dilutions of either glucagon, GLP-1 or novel glucagon analogs for 5 h at 37°C, 5% CO₂ in 96 well poly-D-Lysine-coated “Biocoat” plates (BD Biosciences, San Jose,

CA). At the end of the incubation 100 microliters of LucLite luminescence substrate reagent (Perkin-Elmer, Wellesley, MA) were added to each well. The plate was shaken briefly, incubated 10 min in the dark and light output was measured on MicroBeta-1450 liquid scintillation counter (Perkin-Elmer, Wellesley, MA).

- 5 Effective 50% concentrations were calculated by using Origin software (OriginLab, Northampton, MA. Results are shown in Tables 2 and 3.

Table 2
cAMP Induction by Glucagon Analogs with C-Terminus Extension

Peptide	cAMP Induction			
	Glucagon Receptor		GLP-1 Receptor	
	EC ₅₀ , nM	N*	EC ₅₀ , nM	N
Glucagon	0.22 ± 0.09	14	3.85 ± 1.64	10
GLP-1	2214.00 ± 182.43	2	0.04 ± 0.01	14
Glucagon Cex	0.25 ± 0.15	6	2.75 ± 2.03	7
Oxyntomodulin	3.25 ± 1.65	5	2.53 ± 1.74	5
Oxyntomodulin KRNR	2.77 ± 1.74	4	3.21 ± 0.49	2
Glucagon R12	0.41 ± 0.17	6	0.48 ± 0.11	5
Glucagon R12 Cex	0.35 ± 0.23	10	1.25 ± 0.63	10
Glucagon R12 K20	0.84 ± 0.40	5	0.82 ± 0.49	5
Glucagon R12 K24	1.00 ± 0.39	4	1.25 ± 0.97	5
Glucagon R12 K29	0.81 ± 0.49	5	0.41 ± 0.24	6
Glucagon Amide	0.26 ± 0.15	3	1.90 ± 0.35	2
Oxyntomodulin C24	2.54 ± 0.63	2	5.27 ± 0.26	2
Oxyntomodulin C24 PEG 20K	0.97 ± 0.04	1	1.29 ± 0.11	1

10 * - number of experiments

Table 3
cAMP Induction by Pegylated Glucagon Analogs

Peptide	cAMP Induction			
	Glucagon Receptor		GLP-1 Receptor	
	EC ₅₀ , nM	N*	EC ₅₀ , nM	N
Glucagon	0.33 ± 0.23	18	12.71 ± 3.74	2
Glucagon C17 PEG 5K	0.82 ± 0.15	4	55.86 ± 1.13	2
Glucagon C21 PEG 5K	0.37 ± 0.16	6	11.52 ± 3.68	2
Glucagon C24 PEG 5K	0.22 ± 0.10	12	13.65 ± 2.95	4
Glucagon C29 PEG 5K	0.96 ± 0.07	2	12.71 ± 3.74	2
Glucagon C24 PEG 20K	0.08 ± 0.05	3	Not determined	
Glucagon C24 Dimer	0.10 ± 0.05	3	Not determined	
GLP-1	> 1000		0.05 ± 0.02	4

* - number of experiments

5 Data for additional glucagon analogs is presented in Figs. 6-9 and in Table 4

Table 4
Observed EC₅₀s (nM) in Cells Overexpressing the Glucagon Receptor

		<u>Test 1</u>	<u>Test 2</u>	<u>Test 3</u>	<u>Test 4</u>
	Glucagon standard	0.12	0.04	0.05	0.11
15	K29	0.35			0.22
	K30	0.22	0.06		
	K29, K30	0.89			
	K30, K31		0.12		
20	D28		0.05		0.17
	E28			0.14	
	E29		0.05	0.04	
	E30		0.04		
	D28, E29			0.03	
25	D28, E30			0.05	
	D28, E15			0.15	

30 EXAMPLE 14

Stability Assay for glucagon Cys-maleimido PEG analogs

Each glucagon analog was dissolved in water or PBS and an initial HPLC analysis was conducted. After adjusting the pH (4, 5, 6, 7), the samples were incubated over a specified time period at 37°C and re-analyzed by HPLC to determine the integrity of the peptide. The concentration of the specific peptide of interest was determined and the percent remaining intact was calculated relative to the initial analysis. Results for Glucagon Cys²¹-maleimidoPEG_{5K} are shown in Figs. 1 and 2.

EXAMPLE 15

10 Changes in Serum Glucose Concentration in Beagle Dogs after Administration of Glucagon Analogs

Canine/Beagle dogs of 8-12 kg, being of 8-16 months of age and good health were used to determine the pharmacokinetics and pharmacodynamics of glucagon action. Every animal was fasted overnight and bled at the following time points after each dose: 0 hr. (pre-dose), 5, 10, 20, 30, 45, 60, 90, 120, 240 minutes post dose. Six animals were used for each dose group and approximately 1-2ml of whole blood was withdrawn at each time point. About 1.0 ml whole blood was added to K₂ EDTA tubes containing a sufficient volume of Trasylol (aprotinin) to yield at least 500 KIU/mL of whole blood. Approximately 500 uL plasma was collected after centrifuging the samples in a refrigerated centrifuge at about 1,500 – 3,000 x g for 10-15 min. The samples were transferred to plastic vials and stored frozen at -70°C, or below. The remaining 1.0 mL of whole blood was converted into serum by placing blood sample into an empty tube, letting sit at ambient temperature for 15-20 min, then centrifuging at 1,500 – 3,000 x g for 10-15 min. in a refrigerated centrifuge. The samples were transferred to plastic vials and stored frozen at -70°C, or below. Glucagon and the analogs were dissolved in 0.01N HCl at a concentration of 0.1667 mg/ml and the animals were dosed at 0.03 ml/kg.

The animals were administered a 0.005 mg/kg dose intramuscularly of either glucagon, a glucagon analog comprising glucagon with the sequence of SEQ ID NO: 31 linked to the carboxy terminus of glucagon (glucagon-CEX) or a glucagon analog comprising an aspartic acid substitution at amino acid 28 (glucagon-Asp28) SEQ ID NO: 11. The resulting data is presented in Fig. 10.

Claims:

1. A glucagon peptide comprising the sequence of SEQ ID NO: 33 or an analog of SEQ ID NO: 33, wherein said analog differs from SEQ ID NO: 33 by one to
5 three amino acid substitutions, said amino acid substitutions located at one or more positions selected from positions 1, 2, 5, 7, 10, 11, 12, 13, 14, 16, 17, 18, 19, 20, 21 and 24 of the analog, with the proviso that said glucagon peptide comprises an amino acid at position 28 or 29 different from that found at the corresponding position in
10 SEQ ID NO: 1, or an additional acidic amino acid added to the carboxy terminus of the peptide.
2. The glucagon peptide of claim 1 wherein the glucagon peptide comprises the sequence of SEQ ID NO: 34.
- 15 3. The glucagon peptide of claim 1 wherein the peptide comprises the sequence of SEQ ID NO: 10.
4. The glucagon peptide of claim 1 wherein the glucagon peptide comprises an analog of SEQ ID NO: 34, wherein amino acid at position 2 of the
20 peptide of SEQ ID NO: 34 is other than serine
5. The glucagon peptide of claim 4 wherein position 2 of the peptide is an amino acid selected from the group consisting of d-serine, alanine, glycine, n-methyl serine and amino isobutyric acid.
25
6. The glucagon peptide of claim 1 wherein the glucagon peptide comprises an analog of SEQ ID NO: 34, wherein amino acid at position 1 of the peptide of SEQ ID NO: 34 is selected from the group consisting of d-histidine, desaminohistidine, hydroxyl-histidine, acetyl-histidine and homo-histidine.
30
7. The glucagon peptide of claim 1 further comprising a hydrophilic moiety covalently bound at position 16, 17, 21 or 24 of the glucagon peptide.

8. The glucagon peptide of claim 7 wherein said hydrophilic moiety is polyethylene glycol.

9. The glucagon peptide of claim 8 wherein the polyethylene glycol chain
5 has a molecular weight selected from the range of about 1,000 to about 5,000 Daltons.

10. The glucagon peptide of claim 8 wherein the polyethylene glycol chain has a molecular weight of at least about 20,000 Daltons.

10 11. A glucagon peptide of claim 1 wherein the glucagon peptide further comprises a sequence selected from the group consisting of SEQ ID NO: 20, SEQ ID NO: 21 and SEQ ID NO: 22 bound to the carboxy terminus of said glucagon peptide.

12. A purified glucagon peptide comprising the sequence of SEQ ID NO:
15 7 or a glucagon agonist analog thereof.

13. The glucagon peptide of claim 12 wherein the peptide comprising a sequence selected from the group consisting of SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 11 and SEQ ID NO: 38.

20

14. The glucagon peptide of claim 13 further comprising an additional amino acid, selected from the group consisting of Asp and Glu, added to the carboxy terminus of the glucagon peptide.

25 15. The glucagon peptide of claim 12 further comprising a hydrophilic moiety covalently bound to an amino acid residue at a position selected from the group consisting of 16, 17, 21 and 24, and pharmaceutically acceptable salts of said glucagon peptide.

30 16. The glucagon peptide of claim 15, wherein said hydrophilic moiety is polyethylene glycol.

17. The glucagon peptide of claim 16 wherein the polyethylene glycol chain has a molecular weight selected from the range of about 1,000 to about 5,000 Daltons.

5 18. The glucagon peptide of claim 16 wherein the polyethylene glycol chain has a molecular weight of at least about 20,000 Daltons.

19. The glucagon peptide of claim 12 wherein the glucagon peptide comprises the sequence of SEQ ID NO: 11 or SEQ ID NO: 38.

10

20. The glucagon peptide of claim 19 further comprising an extension peptide bound to amino acid 29 of said glucagon peptide, said amino acid sequence selected from the group consisting of SEQ ID NO: 20, SEQ ID NO: 21 and SEQ ID NO: 22.

15

21. A purified glucagon peptide comprising the sequence of SEQ ID NO: 33, with the proviso that when the amino acid at position 28 is asparagine and the amino acid at position 29 is threonine, the peptide further comprises one to two amino acids, independently selected from the group consisting of Lys, Arg, His, Asp or Glu, added to the c-terminus of the glucagon peptide.

20

22. The glucagon peptide of claim 21 wherein the glucagon peptide comprises a sequence selected from the group consisting of SEQ ID NO: 9, SEQ ID NO: 12, SEQ ID NO: 13 and SEQ ID NO: 32.

25

23. The glucagon peptide of claim 22, wherein the peptide further comprises one to two amino acids, independently selected from the group consisting of Lys, Arg, His, Asp or Glu, added to the c-terminus of the glucagon peptide.

30 24. The glucagon peptide of claim 21 further comprising an extension peptide bound to amino acid 29 of said glucagon peptide, said amino acid sequence selected from the group consisting of SEQ ID NO: 20, SEQ ID NO: 21 and SEQ ID NO: 22.

25. The glucagon peptide of claim 21 further comprising a polyethylene glycol chain covalently bound to an amino acid residue at a position selected from the group consisting of 16, 17, 21 and 24, and pharmaceutically acceptable salts of said glucagon peptide.

5

26. A homo dimer comprising two glucagon peptides bound to one another through a linker, wherein each of said glucagon peptides comprises the sequence of SEQ ID NO: 11.

10

27. The homo dimer of claim 26 wherein the linker is selected from the group consisting of bifunctional thiol crosslinkers and bi-functional amine crosslinkers.

15

28. A dimer comprising a first glucagon peptide bound to a second glucagon peptide via a linker, wherein said first glucagon peptide comprises a peptide selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10 and SEQ ID NO: 11 and the second glucagon peptide comprises SEQ ID NO: 33, with the proviso that with regards to the second glucagon peptide, when the amino acid at position 28 is asparagine and the amino acid at position 29 is threonine, the second glucagon peptide further comprises one to two amino acids, independently selected from the group consisting of Lys, Arg, His, Asp or Glu, added to the carboxy terminus of the second glucagon peptide, and pharmaceutically acceptable salts of said glucagon polypeptides.

20

29. A pharmaceutical composition comprising the glucagon agonist of claim 12, or pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

25

30. A pharmaceutical composition comprising a glucagon agonist that comprises the sequence of SEQ ID NO: 11 or SEQ ID NO: 38, or pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

30

31. A kit for administering a glucagon agonist to a patient in need thereof, said kit comprising
a pharmaceutical composition of claim 29; and
a device for administering said composition to a patient.

5

32. The kit of claim 31 wherein said device is an aerosol dispensing device, wherein the composition is prepackaged within the aerosol device.

33. The kit of claim 31 wherein said device comprises a syringe and a
10 needle, wherein the composition is prepackaged within the syringe.

34. A method of treating hypoglycemia using a pre-formulated aqueous composition, said method comprising the steps of administering an effective amount
of a pharmaceutical composition of claim 29.

15

35. A method of causing temporary paralysis of the intestinal tract, said method comprising administering an effective amount of a pharmaceutical composition of claim 29.

20 36. A method of reducing weight gain or inducing weight loss, said method comprising administering an effective amount of a composition comprising a glucagon agonist comprising a glucagon peptide selected from the group consisting of SEQ ID NO: 8, wherein amino acid 29 of the glucagon peptide is bound to a second peptide through a peptide bond, and said second peptide comprises sequence selected
25 from the group consisting of SEQ ID NO: 20 (GPSSGAPPPS), SEQ ID NO: 21 (KRNRNNIA) and SEQ ID NO: 22 (KRNR).

37. A method of reducing weight gain or inducing weight loss, said method comprising administering an effective amount of a composition comprising a
30 glucagon agonist comprising a glucagon peptide selected from the group consisting of SEQ ID NO: 8.