

CLAIMS

1. An anti-sclerostin antibody for use in treating osteogenesis imperfecta (OI) in a human patient, wherein a therapeutically effective amount of the anti-sclerostin antibody is to be administered to the human patient, and wherein the antisclerostin antibody comprises:
 - (a) a heavy chain variable region CDR1 comprising an amino acid sequence set forth in SEQ ID NO:4;
 - (b) a heavy chain variable region CDR2 comprising an amino acid sequence set forth in SEQ ID NO: 15;
 - (c) a heavy chain variable region CDR3 comprising an amino acid sequence set forth in SEQ ID NO:26;
 - (d) a light chain variable region CDR1 comprising an amino acid sequence set forth in SEQ ID NO:37;
 - (e) a light chain variable region CDR2 comprising an amino acid sequence set forth in SEQ ID NO:48; and
 - (f) a light chain variable region CDR3 comprising an amino acid sequence set forth in SEQ ID NO:59.

2. Use of an anti-sclerostin antibody in the manufacture of a medicament for the treatment of osteogenesis imperfecta (OI) in a human patient, wherein a therapeutically effective amount of the anti-sclerostin antibody is to be administered to the human patient, and wherein the antisclerostin antibody comprises:
 - (a) a heavy chain variable region CDR1 comprising an amino acid sequence set forth in SEQ ID NO:4;
 - (b) a heavy chain variable region CDR2 comprising an amino acid sequence set forth in SEQ ID NO: 15;
 - (c) a heavy chain variable region CDR3 comprising an amino acid sequence set forth in SEQ ID NO:26;
 - (d) a light chain variable region CDR1 comprising an amino acid sequence set forth in SEQ ID NO:37;

- (e) a light chain variable region CDR2 comprising an amino acid sequence set forth in SEQ ID NO:48; and
 - (f) a light chain variable region CDR3 comprising an amino acid sequence set forth in SEQ ID NO:59.
3. The anti-sclerostin antibody for use according to claim 1 or the use according to claim 2, wherein the OI is type I OI, type III OI or type IV OI.
 4. The anti-sclerostin antibody for use according to claim 1 or 3, or the use according to claim 2 or 3, wherein the human patient has one or more mutations in the COL1A1 and/or COL1A2 genes.
 5. The anti-sclerostin antibody for use according to any one of claims 1, 3 or 4, or the use according to any one of claims 2-4, wherein the anti-sclerostin antibody comprises:
 - a) a VH polypeptide sequence having at least 90 percent sequence identity to the amino acid sequences set forth as SEQ ID NO:70; and/ or
 - b) a VL polypeptide sequence having at least 90 percent sequence identity to the amino acid sequences set forth as SEQ ID NO:81.
 6. The anti-sclerostin antibody for use according to any one of claims 1, 3, 4 or 5, or the use according to any one of claims 2-5, wherein the anti-sclerostin antibody comprises:
 - a) a full length heavy chain amino acid sequence having at least 90 percent sequence identity to the amino acid sequence set forth as SEQ ID NO: 172; and/ or
 - b) a full length light chain amino acid sequence having at least 90 percent sequence identity to the amino acid sequence set forth as SEQ ID NO: 173.
 7. The anti-sclerostin antibody for use according to any one of claims 1, 3-5 or 6, or the use according to any one of claims 2-6, wherein the anti-sclerostin antibody comprises a VL polypeptide sequence comprising the amino acid sequence set forth as SEQ ID NO:81

and a VH polypeptide sequence comprising the amino acid sequence set forth as SEQ ID NO:70.

8. The anti-sclerostin antibody for use according to any one of claims 1, 3-6 or 7, or the use according to any one of claims 2-7, wherein the anti-sclerostin antibody comprises a full length light chain amino acid sequence comprising the amino acid sequence set forth as SEQ ID NO: 173 and a full length heavy chain amino acid sequence comprising the amino acid sequence set forth as SEQ ID NO: 172.
9. The anti-sclerostin antibody for use according to any one of claims 1, 3-7 or 8, or the use according to any one of claims 2-8, wherein the anti-sclerostin antibody is to be administered at a dose of 1-50 mg per kg body weight of the human patient.
10. The anti-sclerostin antibody for use according to any one of claims 1, 3-8 or 9, or the use according to any one of claims 2-9, wherein the anti-sclerostin antibody is to be administered:
 - (a) at a dose of 1-30 mg per kg body weight of the human patient, for example at a dose of 1-20 mg per kg of body weight of the human patient; or
 - (b) at a dose of 10-30 mg per kg body weight of the human patient; and/or
 - (c) at a dose of 10-5000 mg, for example at a dose of 100-3000 mg; and/or
 - (d) to the human patient on a daily, weekly, bi-weekly, monthly, bi-monthly or quarterly basis.
11. The anti-sclerostin antibody for use according to any one of claims 1, 3-9 or 10, or the use according to any one of claims 2-10, wherein the treatment regimen comprises a first dosing regimen optionally followed by a second dosing regimen.
12. The anti-sclerostin antibody for use according to claim 11, or the use according to claim 11, wherein:

- (a) the first dosing regimen is 1-50 mg per kg body weight of the human patient is to be administered on a monthly basis, for example the first dosing regimen is 20 mg per kg body weight of the human patient is to be administered on a monthly basis;
 - (b) the second dosing regimen is 1-50 mg per kg body weight of the human patient is to be administered on a bi-monthly or quarterly basis, for example the second dosing regimen is 20 mg per kg body weight of the human patient is to be administered on a bi-monthly or quarterly basis; and/or
 - (c) the first dosing regimen is 20 mg per kg body weight of the human patient is to be administered on a monthly basis and the second dosing regimen is 20 mg per kg body weight of the human patient is to be administered on a bi-monthly or quarterly basis, for example the first dosing regimen is 20 mg per kg body weight of the human patient is to be administered on a monthly basis for a period of 1 year, and the second dosing regimen is 20 mg per kg body weight of the human patient is to be administered on a bi-monthly or quarterly basis for a period of at least 1 year.
13. The anti-sclerostin antibody for use according to claim 11, or the use according to claim 11, wherein:
- (a) the first dosing regimen is 10-5000 mg is to be administered on a monthly basis;
 - (b) the second dosing regimen is 10-5000 mg is to be administered on a bi-monthly or quarterly basis; and/or
 - (c) the first dosing regimen is 10-5000 mg is to be administered on a monthly basis and the second dosing regimen is 10-5000 mg is to be administered on a bi-monthly or quarterly basis, for example the first dosing regimen is 10-5000 mg is to be administered on a monthly basis for a period of 1 year, and the second dosing regimen is 10-5000 mg is to be administered on a bi-monthly or quarterly basis for a period of at least 1 year.
14. The anti-sclerostin antibody for use according to any one of claims 1, 3-12 or 13, or the use according to any one of claims 2-13, wherein the anti-sclerostin antibody is to be administered intravenously.

15. The anti-sclerostin antibody for use according to any one of claims 1, 3-13 or 14, or the use according to any one of claims 2-14, wherein a further therapeutic agent, such as bisphosphonate, parathyroid hormone, calcilytics, calcimimetics (e.g., cinacalcet), statins, anabolic steroids, lanthanum and strontium salts, and/or sodium fluoride is to be administered.
16. The anti-sclerostin antibody for use according to any one of claims 1, 3-14 or 15, or the use according to any one of claims 2-15, wherein the antibody is formulated in a pharmaceutical composition.