

MICROBICIDAL AND ALGICIDAL COMPOSITION, MICROBICIDAL AND ALGICIDAL PROCESS FOR A WATER SYSTEM AND PROCESS FOR PRODUCING A MICROBICIDAL AND ALGICIDAL COMPOSITION

BACKGROUND OF THE INVENTION

1. Field of the Invention

The present invention relates to a microbicidal and algicidal composition, a microbicidal and algicidal process for a water system and a process for producing a microbicidal and algicidal composition. More particularly, the present invention relates to a microbicidal and algicidal composition, a microbicidal and algicidal process for a water system and a process for producing a microbicidal and algicidal composition, which are characterized in that decomposition of effective components by the effect of ultraviolet light and copper ion is suppressed when the composition is left standing, stored or used outdoors.

2. Description of Related Art

Various troubles caused by microorganisms and algae have been taking place in many types of water system. For example, bacteria such as zooglea-form bacteria, filamentous bacteria, iron bacteria, sulfur bacteria, nitrification bacteria and sulfate reduction bacteria, fungi such as saprolegnia and penicillium and algae such as blue algae, green algae and diatoms grow in a cooling water system of the open circulation type. Slime and sludge are formed by attachment and accumulation of dirt in a soft mud condition which is formed from a mixture containing these microorganisms as the main component, inorganic substances such as soil



159159



G00002

and sand and dusts. Slime and sludge cause not only a decrease in the heat efficiency and poor water flow but also local corrosion of instruments and piping. Occasionally, bacteria grown in the water system directly exhibit adverse effects on human health. For example, legionnaires' disease is caused by *Legionella* bacteria scattered from cooling towers. In a water system in the paper manufacturing, various types of bacteria, fungi and yeast grow and slime is formed. Slime causes not only defects in the product such as holes, spots and eyes to decrease the quality of the product but also break in the manufactured paper sheet to decrease the productivity.

To prevent the troubles caused by microorganisms and algae such as those described above, it has heretofore been conducted that a chlorine-based oxidizing agent such as a hypochlorite is added to the water system. It has been considered that, in general, the growth of microorganisms and algae can be suppressed when the concentration of residual chlorine in water is 5 mg Cl/liter or higher. However, decomposition of the chlorine-based oxidizing agent such as a hypochlorite tends to be accelerated by exposure to ultraviolet light. When a microbicidal and algicidal agent is placed into a plastic container or the like and stored or left standing outdoors, the chlorine-based oxidizing agent as the effective component is decomposed by exposure to ultraviolet light. After the microbicidal and algicidal agent is dissolved into water, it is difficult in a cooling water system of the open circulation type that the cooling water is completely shielded from light. Moreover, when copper or a copper alloy is used for the material of piping and a heat exchanger in the water system and

copper ion is eluted into the water, decomposition of the chlorine-based oxidizing agent such as a hypochlorite is further accelerated.

It has been known that the effective chlorine component can be stabilized by adding sulfamic acid or a salt thereof to a hypochlorite so that N-monochlorosulfamic acid, N,N-dichlorosulfamic acid or salts thereof are formed. However, even when the stabilization is achieved by the addition of sulfamic acid or a salt thereof, a problem remains with respect to the stability of the agent when the tank of the agent is placed outdoors and exposed to ultraviolet light. Therefore, a microbicidal and algicidal composition and a microbicidal and algicidal process for a water system in which the loss in the effective components can be suppressed even when the composition is stored or used or the process is conducted in the environment exposed to ultraviolet light or in the presence of copper ion has been desired.

SUMMARY OF THE INVENTION

The present invention has an object of providing a microbicidal and algicidal composition and a microbicidal and algicidal process for a water system in which decomposition of the effective components by the effect of ultraviolet light and copper ion can be suppressed even when the composition is left standing, stored or used outdoors or the process is conducted outdoors and a process for producing the microbicidal and algicidal composition.

As the result of extensive studies by the present inventors to achieve the above object, it was found that the decomposition of the chlorine-based oxidizing agent could be suppressed by maintaining pH of a microbicidal

and algicidal composition containing the chlorine-based oxidizing agent and sulfamic acid or a salt thereof at 13 or greater and that the decomposition of the chlorine-based oxidizing agent was more effectively suppressed when an azole compound was contained and the high concentration of residual chlorine could be maintained. The present invention has been completed based on the above knowledge.

The present invention provides:

- (1) A microbicidal and algicidal composition which comprises chlorine-based oxidizing agent and sulfamic acid or a salt thereof and has a pH of 13 or greater;
- (2) A composition described in (1), wherein the chlorine-based oxidizing agent is sodium hypochlorite;
- (3) A composition described in any one of (1) and (2), wherein a content of sulfamic acid or a salt thereof is in a range of 0.3 to 1.5 moles per 1 mole of available chlorine in the chlorine-based oxidizing agent;
- (4) A composition described in any one of (1), (2) and (3), wherein a content of the chlorine-based oxidizing agent is in a range of 3 to 12% by weight expressed as the content of available chlorine;
- (5) A microbicidal and algicidal composition which comprises a chlorine-based oxidizing agent, an azole-based compound and sulfamic acid or a salt thereof;
- (6) A composition described in (5), which has a pH of 13 or greater;
- (7) A composition described in any one of (5) and (6), wherein the chlorine-based oxidizing agent is sodium hypochlorite;
- (8) A composition described in any one of (5), (6) and (7), wherein the azole-based compound is benzotriazole or tolyltriazole;

- (9) A composition described in any one of (5), (6), (7), and (8), wherein a content of sulfamic acid or a salt thereof is in a range of 0.3 to 1.5 moles per 1 mole of available chlorine in the chlorine-based oxidizing agent;
- (10) A composition described in any one of (5), (6), (7), (8) and (9), wherein a content of the azole-based compound is in a range of 0.05 to 3% by weight;
- (11) A composition described in any one of (5), (6), (7), (8), (9) and (10), wherein a content of the chlorine-based oxidizing agent is in a range of 3 to 12% by weight expressed as the content of available chlorine;
- (12) A microbicidal and algicidal process for a water system which comprises adding a chlorine-based oxidizing agent, an azole-based compound and sulfamic acid or a salt thereof to the water system;
- (13) A process described in (12), wherein the chlorine-based oxidizing agent is sodium hypochlorite;
- (14) A process described in any one of (12) and (13), wherein the azole-based compound is benzotriazole or tolyltriazole;
- (15) A process for producing a microbicidal and algicidal composition which comprises preparing an aqueous solution comprising at least sulfamic acid or a salt thereof and a pH regulator and, thereafter, adding a chlorine-based oxidizing agent to the prepared aqueous solution;
- (16) A process described in (15), wherein the chlorine-based oxidizing agent is sodium hypochlorite;
- (17) A process described in any one of (15) and (16), wherein a content of the chlorine-based oxidizing agent is in a range of 3 to 12% by weight expressed as the content of available chlorine;
- (18) A process described in any one of (15), (16) and (17), wherein a

content of sulfamic acid or a salt thereof is in a range of 0.3 to 1.5 moles per 1 mole of available chlorine in the chlorine-based oxidizing agent;

(19) A process described in any one of (15), (16), (17) and (18), wherein the aqueous solution comprising at least sulfamic acid or a salt thereof and a pH regulator further comprises an azole-based compound; and

(20) A process described in any one of (15), (16), (17), (18) and (19), wherein a content of the azole-based compound is in a range of 0.05 to 3% by weight.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows a system diagram of a circulating water system used in Examples. The numbers in Figure 1 have the following meanings:

- 1: Water tank
- 2: A pump
- 3: A flow meter
- 4: A copper tube

DESCRIPTION OF THE PREFERRED EMBODIMENTS

The first embodiment of the present invention relates to a microbicidal and algicidal composition which comprises chlorine-based oxidizing agent and sulfamic acid or a salt thereof and has a pH of 13 or greater. The second embodiment of the present invention relates to a microbicidal and algicidal composition which comprises a chlorine-based oxidizing agent, an azole-based compound and sulfamic acid or a salt thereof. In the second embodiment of the present invention, it is preferable that the microbicidal and algicidal composition has a pH of 13 or



greater. The microbicidal and algicidal process for a water system of the present invention comprises adding a chlorine-based oxidizing agent, an azole-based compound and sulfamic acid or a salt thereof to a water system. By using the composition of the present invention or in accordance with the process of the present invention, decomposition of the effective components is suppressed even in the environment exposed to ultraviolet light. Therefore, the high concentration of residual chlorine is maintained and the microbicidal and algicidal effect is effectively exhibited. Formation of slime in a cooling water system or the like can be prevented and attached slime can be cleaved.

The chlorine-based oxidizing agent used in the present invention is not particularly limited. Examples of the chlorine-based oxidizing agent include chlorine gas, chlorine dioxide, hypochlorous acid and salts thereof, chloric acid and salts thereof, perchloric acid and salts thereof and chlorinated isocyanuric acid and salts thereof. Among these chlorine-based oxidizing agents, hypochlorous acid and salts thereof are preferable. Examples of the salt of hypochlorous acid include sodium hypochlorite, potassium hypochlorite, calcium hypochlorite and barium hypochlorite. In the composition of the present invention, the content of the chlorine-based oxidizing agent is not particularly limited. It is preferable that the content of the chlorine-based oxidizing agent is in the range of 3 to 12% by weight expressed as the content of available chlorine and more preferably in the range of 4 to 8% by weight expressed as the content of available chlorine.

The salt of sulfamic acid used in the present invention is not particularly limited. Examples of the salt of sulfamic acid include sodium

sulfamate, potassium sulfamate, calcium sulfamate, strontium sulfamate, barium sulfamate, iron sulfamate and zinc sulfamate. The sulfamic acid and the salt thereof may be used singly or in combination of two or more.

Hypochlorite ion and sulfamic acid react with each other as shown in the following equations and N-monochlorosulfamate ion or N,N-dichlorosulfamate ion is formed to stabilize the effective components of the chlorine-based oxidizing agent.



In the composition of the present invention, the content of sulfamic acid or the salt thereof is not particularly limited. It is preferable that the content of sulfamic acid or the salt thereof is in the range of 0.3 to 1.5 moles and more preferably in the range of 1 to 1.5 moles per 1 mole of available chlorine in the chlorine-based oxidizing agent.

In the first embodiment of the composition of the present invention which comprises the chlorine-based oxidizing agent and sulfamic acid or a salt thereof, the composition has a pH of 13 or greater and preferably 13.5 or greater. In the second embodiment of the composition of the present invention which comprises the chlorine-based oxidizing agent, an azole compound and sulfamic acid or a salt thereof, it is preferable that the composition has a pH of 13 or greater and more preferably 13.5 or greater. When the composition has a pH smaller than 13, the stabilizing effect of the chlorine-based oxidizing agent decreases and there is the possibility that the rate of the decrease in the concentration of residual chlorine increases.

The azole compound used in the present invention is an aromatic

compound having a five-membered ring having two or more hetero atoms. Examples of the azole compound used in the present invention include single ring azole compounds such as imidazole, pyrazole, oxazole, thiazole, triazole and tetrazole; condensed multi-ring azole compounds such as benzimidazole, benzoxazole, benzisooxazole, benzothiazole, mercapto-benzimidazole, mercaptomethylbenzimidazole, mercaptobenzothiazole, benzotriazole, totyltriazole, indazole, purine, imidazothiazole and pyrazoxazole; and salts of these azole compounds when the azole compounds form salts. The azole compound may be used singly or in combination of two or more. In the composition of the present invention, the content of the azole compound is not particularly limited. It is preferable that the content of the azole compound is in the range of 0.05 to 3% by weight and more preferably in the range of 0.1 to 2% by weight.

The water system to which the microbicidal and algicidal process of the present invention is applied is not particularly limited. Examples of the water system include cooling water systems, water systems in paper and pulp manufacturing processes, dust-collecting water systems, scrubber water systems and fountain water systems. In the microbicidal and algicidal process of the present invention, the process for adding the chlorine-based oxidizing agent, the azole compound and sulfamic acid or the salt thereof is not particularly limited. For example, the chlorine-based oxidizing agent, the azole compound and sulfamic acid or the salt thereof may be added each separately, two of these components may be added simultaneously and the remaining component may be added separately or the three components may be added simultaneously. Among these processes of addition, it is preferable that the composition

comprising the chlorine-based oxidizing agent, the azole compound and sulfamic acid or the salt thereof is prepared in advance and the prepared composition is added to the water system. By preparing the composition comprising the chlorine-based oxidizing agent, the azole compound and sulfamic acid or the salt thereof in advance, the loss in the effective components of the composition by decomposition can be prevented.

By using the composition of the present invention or in accordance with the process of the present invention, the high concentration of residual chlorine can be maintained and the microbicidal and algicidal effect can be effectively exhibited to a water system even when the water system is in the environment exposed to ultraviolet light or copper or copper alloys are used for piping and heat exchangers in the water systems.

As the pH regulator used in the present invention, conventional alkali compounds can be used and sodium hydroxide and potassium hydroxide are preferable.

In the process for producing a microbicidal and algicidal composition of the present invention, the process for preparing an aqueous solution comprising sulfamic acid or the salt thereof and the pH regulator is not particularly limited. The azole compounds and/or other agents may be present in combination. The order of addition of water and sulfamic acid or the salt thereof is not particularly limited.

In accordance with the process for producing a microbicidal and algicidal composition of the present invention, the microbicidal and algicidal composition of the present invention can be produced safely without causing a decrease in the concentration of the effective

components.

To summarize the advantages of the present invention, decomposition of the effective components is prevented and the high concentration of residual chlorine is maintained due to the presence of the combination of the chlorine-based oxidizing agent and sulfamic acid or a salt thereof and the alkaline condition maintained at a pH of 13 or greater. By further adding the azole compound, a still higher concentration of residual chlorine is maintained and the microbicidal and algicidal effect is more effectively exhibited on water systems. In accordance with the process for producing a microbicidal and algicidal composition of the present invention, the microbicidal and algicidal composition can be produced safely without causing a decrease in the concentration of residual chlorine.

EXAMPLES

The present invention will be described more specifically with reference to examples in the following. However, the present invention is not limited to the examples.

In Examples and Comparative Examples, the concentration of residual chlorine was measured in accordance with the method of titration with iodine described in Japanese Industrial Standard K 0101 28.3.

In Examples 1 to 13 and Comparative Examples 1 to 5, the microbicidal and algicidal compositions were prepared by preparing a mixed aqueous solution from water and components excluding sodium hypochlorite, followed by adding an aqueous solution of sodium hypochlorite to the prepared solution.

In Examples 1 to 4 and Comparative Examples 1 to 3, the prepared microbicidal and algicidal compositions were each placed in a white narrow-necked polyethylene bottle having a capacity of 100 ml. The bottles were left standing in a thermostatted vessel kept at 40°C or 60°C in the condition such that the bottles were shielded from light from the outside. The concentration of residual chlorine in the composition was measured after 0, 6, 14 and 20 days.

Example 1

A microbicidal and algicidal composition was prepared from 12.4 g of water, 15.6 g of a 48% aqueous solution of sodium hydroxide, 12.0 g of sulfamic acid and 60.0 g of an aqueous solution of sodium hypochlorite containing 12% by weight of available chlorine. The prepared microbicidal and algicidal composition had a pH of 13.4.

The concentration of residual chlorine in the microbicidal and algicidal composition was: 7.2% by weight immediately after the preparation, 7.0% by weight after 6 days, 6.9% by weight after 14 days and 6.8% by weight after 20 days when the composition was left standing at 40°C; and 7.2% by weight immediately after the preparation, 6.2% by weight after 6 days, 4.9% by weight after 14 days and 4.8% by weight after 20 days when the composition was left standing at 60°C.

Example 2

A microbicidal and algicidal composition was prepared in accordance with the same procedures as those conducted in Example 1 except that 8.7 g of water and 19.3 g of the 48% aqueous solution of sodium

hydroxide were used. The prepared microbicial and algicidal composition had a pH of 13.8. The concentration of residual chlorine was measured in accordance with the same method as that in Example 1.

Example 3

A microbicial and algicidal composition was prepared from 11.9 g of water, 15.6 g of a 48% aqueous solution of sodium hydroxide, 12.0 g of sulfamic acid, 60.0 g of an aqueous solution of sodium hypochlorite containing 12% by weight of available chlorine and 0.5 g of benzotriazole. The prepared microbicial and algicidal composition had a pH of 13.4. The concentration of residual chlorine was measured in accordance with the same method as that in Example 1.

Example 4

A microbicial and algicidal composition was prepared in accordance with the same procedures as those conducted in Example 3 except that 8.2 g of water and 19.3 g of the 48% aqueous solution of sodium hydroxide were used. The prepared microbicial and algicidal composition had a pH of 13.8. The concentration of residual chlorine was measured in accordance with the same method as that in Example 1.

Comparative Example 1

A microbicial and algicidal composition was prepared from 39.5 g of water, 60.0 g of an aqueous solution of sodium hypochlorite containing 12% by weight of available chlorine and 0.5 g of benzotriazole. The prepared microbicial and algicidal composition had a pH of 13.0.

The concentration of residual chlorine in the microbicidal and algicidal composition was: 7.2% by weight immediately after the preparation, 5.7% by weight after 6 days, 4.4% by weight after 14 days and 4.2% by weight after 20 days when the composition was left standing at 40°C; and 7.2% by weight immediately after the preparation, 3.7% by weight after 6 days, 2.4% by weight after 14 days and 2.4% by weight after 20 days when the composition was left standing at 60°C.

Comparative Example 2

A microbicidal and algicidal composition was prepared from 16.9 g of water, 10.6 g of a 48% by weight aqueous solution of sodium hydroxide, 12.0 g of sulfamic acid, 60.0 g of an aqueous solution of sodium hypochlorite containing 12% by weight of available chlorine and 0.5 g of benzotriazole. The prepared microbicidal and algicidal composition had a pH of 12.0. The concentration of residual chlorine was measured in accordance with the same method as that in Example 1.

Comparative Example 3

A microbicidal and algicidal composition was prepared from 17.4 g of water, 10.6 g of a 48% by weight aqueous solution of sodium hydroxide, 12.0 g of sulfamic acid and 60.0 g of an aqueous solution of sodium hypochlorite containing 12% by weight of available chlorine. The prepared microbicidal and algicidal composition had a pH of 12.0. The concentration of residual chlorine was measured in accordance with the same method as that in Example 1.

The results in Examples 1 to 4 and Comparative Examples 1 to 3 are shown in Table 1.

Table 1

Example	1	2	3	4			
Comparative Example					1	2	3
Composition of components (g)							
water	12.4	8.7	11.9	8.2	39.5	16.9	17.4
48% by wt. NaOH	15.6	19.3	15.6	19.3	-	10.6	10.6
sulfamic acid	12.0	12.0	12.0	12.0	-	12.0	12.0
12% by wt. NaClO (*)	60.0	60.0	60.0	60.0	60.0	60.0	60.0
benzotriazole	-	-	0.5	0.5	0.5	0.5	-
pH	13.4	13.8	13.4	13.8	13.0	12.0	12.0
Concentration of residual chlorine (% by weight)							
left standing at 40°C							
for 0 days	7.2	7.2	7.2	7.2	7.2	7.2	7.2
for 6 days	7.0	7.1	7.0	7.1	5.7	2.1	2.1
for 14 days	6.9	7.0	6.8	7.0	4.4	1.1	1.1
for 20 days	6.8	6.7	6.8	6.7	4.2	0	0
left standing at 60°C							
for 0 days	7.2	7.2	7.2	7.2	7.2	7.2	7.2
for 6 days	6.2	6.9	6.2	6.9	3.7	0.5	0.5
for 14 days	4.9	6.2	4.9	6.1	2.4	0	0
for 20 days	4.8	6.2	4.8	6.1	2.4	0	0

* An aqueous solution of sodium hypochlorite containing 12% by weight of available chlorine was used.

As shown in Table 1, the microbicidal and algicidal composition of Comparative Example 1 which contained benzotriazole but did not contain sulfamic acid and had a pH of 13.0 showed rapid decomposition of sodium hypochlorite and a rapid decrease in the concentration of residual chlorine.

The microbicidal and algicidal compositions of Comparative Examples 2 and 3 which contained sulfamic acid but had a pH of 12.0 showed rapid decomposition of sodium hypochlorite and rapid decreases in the concentration of residual chlorine independently of the presence or the absence of benzotriazole.

The microbicidal and algicidal compositions of Examples 1 and 3 which contained the aqueous solution of sodium hydroxide in increased amounts and had a pH of 13.4 and the microbicidal and algicidal compositions of Examples 2 and 4 which contained the aqueous solution of sodium hydroxide in increased amounts and had a pH of 13.8 showed small decreases in the concentration of residual chlorine independently of the presence or the absence of benzotriazole. It was shown that the decomposition of sodium hypochlorite was effectively suppressed by using sodium hypochlorite and sulfamic acid in combination and adjusting pH at 13 or greater. In particular, stability at a high temperature was improved by increasing the amount of sodium hydroxide.

In Examples 5 to 9 and Comparative Examples 4 and 5, the prepared microbicidal and algicidal compositions were each placed in a white narrow-necked polyethylene bottle having a capacity of 100 ml. The seven bottles arranged in a row were left standing outdoors at a place exposed to the sunlight in Atsugi City from April 10 to 30. The concentration of residual chlorine in the compositions was measured after 0, 6, 14 and 20 days.

Example 5

A microbicidal and algicidal composition was prepared from 12.4 g

of water, 15.6 g of a 48% aqueous solution of sodium hydroxide, 12.0 g of sulfamic acid and 60.0 g of an aqueous solution of sodium hypochlorite containing 12% by weight of available chlorine. The prepared microbicidal and algicidal composition had a pH of 13.4.

The concentration of residual chlorine in the microbicidal and algicidal composition was: 7.2% by weight immediately after the preparation, 6.1% by weight after 6 days, 4.0% by weight after 14 days and 3.0% by weight after 20 days.

Example 6

A microbicidal and algicidal composition was prepared from 12.4 g of water, 15.6 g of a 48% aqueous solution of sodium hydroxide, 12.0 g of sulfamic acid, 60.0 g of an aqueous solution of sodium hypochlorite containing 12% by weight of available chlorine and 0.2 g of benzotriazole. The prepared microbicidal and algicidal composition had a pH of 13.4.

The concentration of residual chlorine in the microbicidal and algicidal composition was: 7.2% by weight immediately after the preparation, 6.4% by weight after 6 days, 5.2% by weight after 14 days and 4.4% by weight after 20 days.

Example 7

A microbicidal and algicidal composition was prepared and pH and the concentration of residual chlorine were measured in accordance with the same procedures as those conducted in Example 6 except that 0.5 g of benzotriazole was used.

Example 8

A microbicidal and algicidal composition was prepared and pH and the concentration of residual chlorine were measured in accordance with the same procedures as those conducted in Example 6 except that 1.0 g of benzotriazole was used.

Example 9

A microbicidal and algicidal composition was prepared and pH and the concentration of residual chlorine were measured in accordance with the same procedures as those conducted in Example 6 except that 0.5 g of tolyltriazole was used in place of benzotriazole.

Comparative Example 4

A microbicidal and algicidal composition was prepared from 24.4 g of water, 15.6 g of a 48% aqueous solution of sodium hydroxide and 60.0 g of an aqueous solution of sodium hypochlorite containing 12% by weight of available chlorine. The prepared microbicidal and algicidal composition had a pH of 14.0.

The concentration of residual chlorine in the microbicidal and algicidal composition was: 7.2% by weight immediately after the preparation, 5.7% by weight after 6 days, 3.9% by weight after 14 days and 2.8% by weight after 20 days.

Comparative Example 5

A microbicidal and algicidal composition was prepared and pH and the concentration of residual chlorine were measured in accordance with

the same procedures as those conducted in Comparative Example 4 except that 2.0 g of benzotriazole was used.

The results of Examples 5 to 9 and Comparative Examples 4 and 5 are shown in Table 2.

Table 2

Example	5	6	7	8	9		
Comparative Example						4	5
Composition of components (g)							
water	12.4	12.4	12.4	12.4	12.4	24.4	24.4
48% by wt. NaOH	15.6	15.6	15.6	15.6	15.6	15.6	15.6
sulfamic acid	12.0	12.0	12.0	12.0	12.0	-	-
12% by wt. NaClO (*)	60.0	60.0	60.0	60.0	60.0	60.0	60.0
benzotriazole	-	0.2	0.5	1.0	-	-	2.0
tolyltriazole	-	-	-	-	0.5	-	-
pH	13.4	13.8	13.4	13.4	13.4	14.0	14.0
Concentration of residual chlorine (% by weight)							
after 0 days	7.2	7.2	7.2	7.2	7.2	7.2	7.2
after 6 days	6.1	6.4	6.9	7.0	6.8	5.7	5.3
after 14 days	4.0	5.2	6.1	6.3	6.4	3.9	2.7
after 20 days	3.0	4.4	5.7	5.9	6.2	2.8	1.1

* An aqueous solution of sodium hypochlorite containing 12% by weight of available chlorine was used.

As shown in Table 2, in Examples 5 to 8 in which sulfamic acid or a combination of sulfamic acid and benzotriazole was used, the high

concentration of residual chlorine was maintained after passage of time and decomposition of the effective components was suppressed. Since the greater the amount of benzotriazole, the higher the concentration of residual chlorine, it was shown that benzotriazole contributed to suppressing decomposition of the effective components. In Example 9 in which sulfamic acid and tolyltriazole were used, the high concentration of residual chlorine was maintained after passage of time and decomposition of the effective components was suppressed.

In contrast, in Comparative Example 4 in which none of azole compounds and sulfamic acid was used, the concentration of residual chlorine after passage of time was lower than that in Examples and the effect of suppressing decomposition of the effective components was not exhibited. In Comparative Example 5 in which benzotriazole alone was used, the concentration of residual chlorine was lower than that in Comparative Example 4 and the effect of accelerating decomposition of the effective components was exhibited when benzotriazole alone was used. Although sulfamic acid exhibited only small effect of suppressing decomposition of the effective components in the single use and the azole compound exhibited the effect of accelerating decomposition of the effective components in the single use, the specific synergistic effect could be exhibited when these components were used in combination. Thus, decomposition of the effective components could be effectively suppressed and the high concentration of residual chlorine could be maintained.

In Examples 10 to 13, a circulating water system shown in Figure 1 which was equipped with a water tank 1, a pump 2, a flow meter 3 and a copper tube 4 was used. The amount of retained water in this water

system was 15 liters. The copper tube had a length of 1,350 mm and an inner diameter of 19 mm. In this water system, city water of Atsugi city was used as the test water. A prepared microbicidal and algicidal composition was added in an amount of 3.0 g and water was circulated at a flow rate of 10 liters/minute at a temperature of 30°C. The concentration of residual chlorine in the water was measured after 24 hours and 72 hours.

Example 10

A microbicidal and algicidal composition was prepared from 12.4 g of water, 15.6 g of a 48% aqueous solution of sodium hydroxide, 12.0 g of sulfamic acid and 60.0 g of an aqueous solution of sodium hypochlorite containing 12% by weight of available chlorine. The prepared microbicidal and algicidal composition had a pH of 13.4.

The concentration of residual chlorine in the circulating water to which this microbicidal and algicidal composition was added was: 13.0 mg/liter after 24 hours and 10.5 mg/liter after 72 hours and 72.9% of the concentration of residual chlorine remained after 72 hours.

Example 11

A microbicidal and algicidal composition was prepared from 12.15 g of water, 15.6 g of a 48% aqueous solution of sodium hydroxide, 12.0 g of sulfamic acid, 60.0 g of an aqueous solution of sodium hypochlorite containing 12% by weight of available chlorine and 0.25 g of benzotriazole. The prepared microbicidal and algicidal composition had a pH of 13.4.

The concentration of residual chlorine in the circulating water to

which this microbicidal and algicidal composition was added was: 13.5 mg/liter after 24 hours and 13.0 mg/liter after 72 hours and 90.3% of the concentration of residual chlorine remained after 72 hours.

Example 12

A microbicidal and algicidal composition was prepared, pH of the prepared composition was measured, the prepared composition was added to the circulating water system and the concentration of residual chlorine in the water was measured in accordance with the same procedures as those conducted in Example 11 except that 11.9 g of water and 0.5 g of benzotriazole were used.

Example 13

A microbicidal and algicidal composition was prepared, pH of the prepared composition was measured, the prepared composition was added to the circulating water system and the concentration of residual chlorine in the water was measured in accordance with the same procedures as those conducted in Example 11 except that 11.4 g of water and 1.0 g of benzotriazole were used.

The results of Examples 10 to 13 are shown in Table 3.

Table 3

Example	10	11	12	13
Composition of components (g)				
water	12.4	12.15	11.9	11.4
48% by wt. NaOH	15.6	15.6	15.6	15.6
sulfamic acid	12.0	12.0	12.0	12.0
12% by wt. NaClO (*)	60.0	60.0	60.0	60.0
benzotriazole	-	0.25	0.5	1.0
pH	13.4	13.4	13.4	13.4
Microbicidal and algicidal composition (mg/liter)	200	200	200	200
Benzotriazole (mg/liter)	-	0.5	1.0	2.0
Concentration of residual chlorine (mg/liter)				
after 24 hours	13.0	13.5	14.4	14.4
after 72 hours	10.5	13.0	14.4	14.4
Concentration of residual chlorine remaining after 72 hours (%)	72.9	90.3	100	100

* An aqueous solution of sodium hypochlorite containing 12% by weight of available chlorine was used.

As shown in Table 3, in Examples 12 and 13 in which the microbicidal and algicidal compositions containing benzotriazole were used, the concentration of residual chlorine was higher after passage of time and decomposition of the effective components were more suppressed in comparison with the result in Example 11 in which the microbicidal and algicidal composition not containing benzotriazole was used. When the concentration of benzotriazole in the circulating water was 1.0 mg/liter or

higher, the concentration of residual chlorine did not decrease at all after 72 hours.

Example 14

Water in an amount of 6.7 g, 19.3 g of a 48% by weight aqueous solution of sodium hydroxide, 2.0 g of benzotriazole and 12.0 g of sulfamic acid were weighed and placed into a 200 ml glass beaker successively in this order. After the above components were dissolved under stirring, 60.0 g of an aqueous solution of sodium hypochlorite containing 12% by weight of available chlorine was added under stirring and a microbicidal and algicidal composition was prepared after stirring.

The preparation of the microbicidal and algicidal composition could be conducted without bumping although some bubbling was observed. The concentration of residual chlorine in the obtained microbicidal and algicidal composition was 7.2% by weight.

Comparative Example 6

Water in an amount of 6.7 g, 19.3 g of a 48% by weight aqueous solution of sodium hydroxide, 2.0 g of benzotriazole and 12.0 g of sulfamic acid were weighed and placed into a 200 ml glass beaker successively in this order. After the above components were dissolved under stirring, the prepared solution was added in small portions under stirring into a 200 ml glass beaker into which 60.0 g of an aqueous solution of sodium hypochlorite containing 12% by weight of available chlorine was weighed and placed in advance. During the addition and mixing, bubbles began to be formed and eventually bumping took place. Thus, the generation of

chlorine gas was observed.

CLAIMS

1. A microbicidal and algicidal composition which comprises chlorine-based oxidizing agent and sulfamic acid or a salt thereof and has a pH of 13 or greater.
2. A composition according to Claim 1, wherein the chlorine-based oxidizing agent is sodium hypochlorite.
3. A composition according to any one of Claims 1 and 2, wherein a content of sulfamic acid or a salt thereof is in a range of 0.3 to 1.5 moles per 1 mole of available chlorine in the chlorine-based oxidizing agent.
4. A composition according to any one of Claims 1, 2 and 3, wherein a content of the chlorine-based oxidizing agent is in a range of 3 to 12% by weight expressed as the content of available chlorine.
5. A microbicidal and algicidal composition which comprises a chlorine-based oxidizing agent, an azole-based compound and sulfamic acid or a salt thereof.
6. A composition according to Claim 5, which has a pH of 13 or greater.
7. A composition according to any one of Claims 5 and 6, wherein the chlorine-based oxidizing agent is sodium hypochlorite.

8. A composition according to any one of Claims 5, 6 and 7, wherein the azole-based compound is benzotriazole or tolyltriazole.
9. A composition according to any one of Claims 5, 6, 7, and 8, wherein a content of sulfamic acid or a salt thereof is in a range of 0.3 to 1.5 moles per 1 mole of available chlorine in the chlorine-based oxidizing agent.
10. A composition according to any one of Claims 5, 6, 7, 8 and 9, wherein a content of the azole-based compound is in a range of 0.05 to 3% by weight.
11. A composition according to any one of Claims 5, 6, 7, 8, 9 and 10, wherein a content of the chlorine-based oxidizing agent is in a range of 3 to 12% by weight expressed as the content of available chlorine.
12. A microbicidal and algicidal process for a water system which comprises adding a chlorine-based oxidizing agent, an azole-based compound and sulfamic acid or a salt thereof to the water system.
13. A process according to Claim 12, wherein the chlorine-based oxidizing agent is sodium hypochlorite.
14. A process according to any one of Claims 12 and 13, wherein the azole-based compound is benzotriazole or tolyltriazole.
15. A process for producing a microbicidal and algicidal composition

which comprises preparing an aqueous solution comprising at least sulfamic acid or a salt thereof and a pH regulator and, thereafter, adding a chlorine-based oxidizing agent to the prepared aqueous solution.

16. A process according to Claim 15, wherein the chlorine-based oxidizing agent is sodium hypochlorite.

17. A process according to any one of Claims 15 and 16, wherein a content of the chlorine-based oxidizing agent is in a range of 3 to 12% by weight expressed as the content of available chlorine.

18. A process according to any one of Claims 15, 16 and 17, wherein a content of sulfamic acid or a salt thereof is in a range of 0.3 to 1.5 moles per 1 mole of available chlorine in the chlorine-based oxidizing agent.

19. A process according to any one of Claims 15, 16, 17 and 18, wherein the aqueous solution comprising at least sulfamic acid or a salt thereof and a pH regulator further comprises an azole-based compound.

20. A process according to Claim 19, wherein a content of the azole-based compound is in a range of 0.05 to 3% by weight.