FIG. 1

Light chain variable region



FIG. 1 cont.

Heavy chain variable region



FIG. 2A


Human b1 Tryptase
% Tryptase Activity
Antibody concentration (nM)
hu31A.v11 IgG4
Human b1 Tryptase
% Tryptase Activity
Antibody concentration (nM)
hu31A.v11 IgG1
Human b1 Tryptase
% Tryptase Activity
Antibody concentration (nM)
huE104.v2 IgG4
Human b1 Tryptase
% Tryptase Activity
Antibody concentration (nM)
huE104.v2 IgG1

FIG. 2B


Human primary airway smooth muscle cell proliferation
E104.v2 IgG4
31A.v11 IgG4
H3 Thymidine (CPM)
6000
5000
4000
3000
2000
1000
0
0
3
3
30
300
Tryptase β (µg/ml)
3 µg/ml Tryptase β + anti-human tryptase (µg/ml)

FIG. 2C


Human primary airway smooth muscle cell contraction
t=0
t=3 hr
% Contraction
25.0
20.0
15.0
10.0
5.0
0.0
No stimulation
Tryptase
Tryptase + huE104.v2 IgG4
Tryptase + hu31A.v11 IgG4


Histamine (ng/ml)
120
80
40
0
PBS
Tryptase (S195A)
Tryptase
Tryptase 31A.v11

FIG. 2D


30 – 50 %
Inhibition
Histamine (ng/ml)
350
300
250
200
150
100
50
0
IgE
IgE + NP
IgE + NP
SMI
IgE + NP
31A.v11

FIG. 2E

FIG. 3A


Run 1- WT tetrameric tryptase alone (peak 1)
Run 2- WT tetrameric tryptase + Fab 31A.v11 + heparin (peaks 2, 4)
Run 3- WT tetrameric tryptase + Fab 31A.v11 w/o heparin (peaks 3, 4)
Peak 2
Monomer bound by hu31A.v11 Fab
+ heparin
Peak 3
Monomer bound by hu31A.v11 Fab
w/o heparin
Peak 4
hu31A.v11 alone
Peak 1
tetramer
mAU
300
250
200
150
100
50
11
12
13
14
15
16
17
mL

FIG.3B


Run 1- His-tagged monomeric tryptase + Fab 104.v2 (peaks 2, 6)
Run 2- WT tetrameric tryptase + Fab 104.v2 (peaks 3, 7)
Run 3- WT tetrameric tryptase + Fab 104.v2 + Heparin (peaks 1, 4, 5)
peak 2 (25.8 min)
peak 3 (26 min)
peak 7 (31.8 min)
peak 6 (31.6 min)
peak 5 (31.2 min)
peak 1 (21 min)
peak 4 (27.2 min)
350
300
250
200
150
100
50
0
8.0
10.0
12.0
14.0
16.0
18.0
ml

FIG. 4A


Lung PK
Lung tryptase
Ab concentration (µg/ml)
residual fraction
days
total Ab
free Ab
monomer
tetramer
>90% tetramer neutralization
>~80% tetramer neutralization
Destabilizing mAb (KD = 0.2 nM)
Stabilizing Tetramer-Specific mAb (KD =0.02nM)

FIG. 4B


Lung PK
Lung tryptase
Dissociating mAb
(KD = 0.2 nM)
Stabilizing
Tetramer-Specific
mAb
(KD = 0.02nM)
Ab concentration (μg/ml)
residual fraction
days
total Ab
free Ab
monomer
tetramer
>80% tetramer
neutralization
>~70% tetramer
neutralization

FIG. 4C


Base-Case Scenario
4 ng/ml serum, 10 ng/ml tissue
High Tryptase Scenario
10 ng/ml serum, 40 ng/ml tissue
Dissociating mAb
(KD = 0.2 nM)
Stabilizing
Tetramer-Specific
mAb
(KD =0.02nM)
residual tetramer (fraction)
days
300 mg
100 mg
30 mg
>40% tetramer neutralization
>80%
>90%
>10% tetramer neutralization
>40%
>80%
>20% tetramer neutralization
>50%
>80%
>10% tetramer neutralization
>30%
>70%

FIG. 5A


pH 6.0
198 -
98 -
62 -
49 -
38 -
28 -
17 -
14 -
6 -
1 2 3 4 5 6


pH 6.0
Fibrinogen alpha chain intensity
80
60
40
20
0
Fibrinogen
Fibrinogen + Tryptase
+ 31A.v11 Fab
+ B12 mIgG1
+ E104.v2 Fab

FIG. 5B


pH 7.5
198 -
98 -
62 -
49 -
38 -
28 -
17 -
14 -
6 -
*
**
***
1 2 3 4 5 6
* alpha chain
** beta chain
***gamma chain


pH 7.5
Fibrinogen alpha chain intensity
80
60
40
20
0
Fibrinogen
Fibrinogen + Tryptase
+ 31A.v11 Fab
+ B12 mIgG1
+ E104.v2 Fab

FIG. 6A


Run 1: WT tetramer + huE104.v1 Fab (ref. peak)
Run 2: tetramer Y75C variant + hu31A.v11 Fab (peaks 1, 4)
Run 3: tetramer I99C variant + hu31A.v11 Fab (peaks 2, 4)
Run 4: WT tetramer + hu31A.v11 Fab (peaks 3, 4)
peak 1
Ref. peak
peak 2
peak 3
peak 4
mAU
250
200
150
100
50
0
8.0
10.0
12.0
14.0
16.0
mL

FIG. 6B


Y75C
I99C
Fab
WT monomer
peak 3
peak 4
peak 1
peak 4
peak 2
peak 4
Run 4
Run 2
Run 3

FIG. 6C


Run 1: tetramer Y75C variant + huE104.v2 Fab (peaks 1, 4)
Run 2: tetramer I99C variant + huE104.v2 Fab (peaks 2, 5)
Run 3: WT tetramer + huE104.v2 Fab (peaks 3, 6)
peak 1 (21.6 min)
peak 2 (23.8 min)
peak 3 (26 min)
peak 4/5 (31 min)
peak 6 (31.8 min)
mAU
350
300
250
200
150
100
50
0
8.0
10.0
12.0
14.0
16.0
18.0
ml

FIG. 7

| | | |

FIG. 7 cont.

| | | |


H36
A62
A63
K60d
D60e
V60c
L61
R87
R65
P84
E109
E110
P111
Q50
V85
S86

FIG. 8

FIG. 9


CH1/CL
VH/VL
Tryptase protomer
CH1/CL
VH/VL
Tryptase protomer

FIG. 10


Val60c in unbound tryptase
Tyr173d from neighboring protomer
Val90
Val60c in tryptase bound by hu31A.v11 Fab

FIG. 11


31A.v11
His36
Pro37a
Pro152
neighboring tryptase protomer
Phe153
Pro152a
Tyr37b
tryptase protomer



FIG. 12A

FIG. 12B


V71 and F78 in the grafted HC FR3 as in huE104.v1
78
71
CDR H2
slower HDX when huE104.v2 is bound (Q81, L82, L83)
Hydrophobic interactions between neighboring protomers
Y74
Y75
tryptase protomers

FIG. 13


anti-tryptase 31A.v11 IgG4, 10 mg/kg
anti-tryptase 31A.v11 IgG4, 1 mg/kg
anti-tryptase E104.v2 IgG4, 10 mg/kg
anti-tryptase E104.v2 IgG4, 1 mg/kg
Serum Concentration (µg/mL)
1000
100
10
1
0
7
14
21
Time (day)


Serum Concentration (μg/mL)
1000
100
10
1
0
7
14
21
28
35
42
Time (day)
anti-gD IgG4
anti-Tryptase 31A.v11 IgG4
anti-Tryptase E104.v2 IgG4

FIG. 14

FIG. 15


Tryptase monomer
Tryptase tetramer
Active Tryptase Assay
+ SBTI
+ ABP: label active tryptase
Biotin Tag
Linker
+ SM inhibitor: stop labeling
Specific Tryptase reactive group
+ Ab: dissociate tetramer
Av-HRP
Biotin labeled ABP
Tryptase
Capture Ab

FIG. 15 cont.


Tryptase monomer
Tryptase tetramer
Total Tryptase Assay
+ Ab: dissociate tetramer
Av-HRP
Biotin-labeled
Detection Ab
Tryptase
Capture
Ab

FIG. 16


Study Day
-7
8hr
2
7
14
21
28
30 mg/kg IV dose
Active Tryptase in BAL
Animal 1
BAL Active Tryptase (ng/ml)
urea normalized
Study Day
Animal 2
BAL Active Tryptase (ng/ml)
urea normalized
Study Day

FIG. 17


Sensitization Phase
-7 0 7 15 29 34 50 57 71 78 85 120 140 184 198-205
ID IP/IM AI IM AI ID IP/IM AI ID AI AI
AI: inhaled ID: intra-dermal
IM: intra-muscular IP: intra-peritoneal
Experimental Phase
vehicle phase
drug phase
1 2 (30 min) 4 wks 1 2 (30 min)
vehicle AI BAL, naso anti-Tryptase AI BAL, naso

FIG. 18A


Active Tryptase in BAL
vehicle
anti-Tryptase
Active Tryptase (ng/ml)
urea normalized
100
80
60
40
20
0
1002
1005
1008
1009
1010
1011
Animal
*
*
*
*
*
*


Total Tryptase in BAL
vehicle
anti-Tryptase
Total Tryptase (ng/ml)
urea normalized
8000
6000
4000
2000
0
1002
1005
1008
1009
1010
1011
Animal

FIG. 18B

FIG. 18C


Total Tryptase in nasosorption
vehicle
anti-Tryptase
Total Tryptase (pg/ml)
5000
4000
3000
300
200
100
0
1002
1005
1008
1009
1010
1011
Animal
*

FIG. 19


NP-IgE 1.6 µg in 200 µl of saline i.v.
Anti-gD or Anti-tryptase i.p.
-24 hr
0 hr
Mouse body temperature
2 hr
200 µg of NP-BSA
in 200 µl saline i.v
Anti-gD
hu31A.v11 anti-tryptase
***
Mouse body temperature (C)
38
36
34
32
Baseline
5
10
15
20
25
30
35
40
45
50
55
60
65
75
85
95
105
115
Post injection (min)