


Antiviral Nucleoside Derivatives

The invention relates to the field of antiviral therapy and, in particular, to nucleoside derivatives for treating Hepatitis C Virus (HCV) mediated diseases. The invention provides novel chemical compounds, pharmaceutical compositions comprising these compounds, methods for treatment or prophylaxis of HCV mediated diseases employing said compounds in monotherapy or in combination therapy.

BACKGROUND

Hepatitis C virus (HCV) is responsible for a large proportion of the chronic liver disease worldwide and accounts for 70% of cases of chronic hepatitis in industrialized countries. The global proportion of hepatitis C is estimated to average 3% (ranging from 0.1% to 5.0%); there are an estimated 170 million chronic carriers throughout the world. There is a continuing need for effective therapeutic agents against HCV.


Levovirin (1-(3S,4R-dihydroxy-5S-hydroxymethyl-tetrahydro-furan-2S-yl)-1H-[1,2,4]triazole-3-carboxylic acid amide), I ($R^1=R^2=R^3=H$), is a nucleoside analog and the enantiomer of the antiviral compound, ribavirin. Unlike ribavirin, levovirin does not have detectable antiviral



activity; however, levovirin stimulates immune responses by enhancing antiviral Th1 cytokine expression. Like ribavirin, levovirin lowers serum alanine aminotransferase levels in a mouse hepatitis model. (R. Tam *et al. J. Med. Chem.* 2000 44:1276-1283; M. Assenmacher *et al. Eur. J. Immunol.* 1998 28:1534-1543). Levovirin appears to lack toxicity associated with ribavirin.

While nucleoside derivatives frequently possess high levels of biological activity, their therapeutic utility is often hampered by suboptimal physical properties and poor pharmacokinetics and bioavailability that limit the amount of the nucleoside that is absorbed. Only about 15% of the dose of levovirin is absorbed systemically after oral administration. There exists a need for therapeutic agents with improved bioavailability. Prodrugs, bioreversible chemical derivatives of poorly absorbed compounds, are one approach to optimizing physical properties to improve drug delivery. (W. N. Chapman and C. J. H. Porter, *Adv. Drug Deliv.* 1996 19:149-169; D. Fleisher *et al. Adv. Drug Deliv.* 1996 19:115-130) In one approach to prodrug design, chemical derivatives are prepared to optimize oil/water partition coefficients or other physical properties to enhance passive transport across mucosal membranes. Derivatives are chosen which are substrates for nonspecific enzymes present in the cytoplasm, blood, or serum and capable of cleaving the modifying group and reverting to the bioactive parent molecule after the compound is absorbed. An ideal oral prodrug should be stable to gastric fluid and intestinal chyme, be efficiently transported across intestinal membranes and be rapidly converted to the parent drug after absorption from the gastrointestinal tract. Thus "pronucleotides" can potentially circumvent problems such as activity, bioavailability or stability of the parent nucleotide.

An alternative approach exploits nonspecific active transport systems to move the prodrug across a membrane. The prodrug portion of the molecule is designed to confer recognition by the active transport system and is cleaved after transport is complete. The nonspecific peptide transporters PepT1 and PepT2 have been suggested to be useful for improving the bioavailability of poorly absorbed drugs. (P. Balimane *et al. Biochem. Biophys. Res. Commun.* 1998 250: 246-251; K. Sawada *et al. J. Pharmacol Exp. Ther.* 1999 291(2):705-709; I. Rubio-Aliaga and H. Daniel, *Trends Pharmacol. Sci.* 2002 23(9):434-40).



IIa: R' = H; R'' = H

IIb: R' = H; R'' = Val-H

IIc: R' = CH₂OH; R'' = H

IId: R' = CH₂OH; R'' = Val-H

Valine esters **IIb** of acyclovir (Valacyclovir) **IIa** exhibit improved absorption characteristics which have been suggested to be the result of uptake *via* peptide


transporters. (Balimane, *supra*; M. E. Ganapathy *et al.* *Biochem. Biophys. Res. Commun.* 1998 246:470-75; P. J. Sinko and P. V. Balimane, *Biopharm. Drug Dispos.* 1998 19:209-17; R. L. de Vruh *et al.* *J. Pharmacol. Exp. Ther.* 1998 286:1166-70) Mitsuru Sugara *et al.* (*J. Pharm. Sci.*, 2000 89(6):781-89) suggested the improved transport of valganciclovir IIc, the valine ester of ganciclovir IIc could be attributed to the PepT1 and PepT2 transport systems. WO 01/68034 A2 (G. Wang *et al.*) disclose bioreversible modifications of the sugar and triazole moiety of levovirin to increase drug bioavailability and to treat an infection, an infestation, a neoplasm or an autoimmune disease. WO 00/23454 (A. K. Ganguly *et al.*) disclose ribavirin derivatives for coadministration with interferon alfa to patients having chronic hepatitis C infection

While the availability of efficiently absorbed prodrugs affords a route to improve the bioavailability of levovirin, exploitation of these compounds also requires a levovirin derivative with physical properties that allow for efficient manufacture and formulation of the active ingredient. Levovirin prodrugs should possess adequate thermal stability, photostability and be non-hygroscopic. Properties relevant to the formulation chemist include particle size, polymorphic form, crystal habit, and salt form. These properties influence the aqueous solubility, dissolution profile, compatibility with other components in the formulation, route of administration and the biopharmaceutical properties. The ideal nucleoside drug candidate must then possess the physical properties which allow it to be efficiently manufactured and formulated, the pharmaceutical properties which allow it to be delivered to the absorption site and chemical properties which allow recognition and uptake by the transport system and conversion back into the desired parent compound after uptake is completed.

DETAILED DISCUSSION OF THE INVENTION

Surprisingly it has now been found that several hydrophobic amino acid ester hydrochlorides and a series of neutral mono-, di-, and triacyl derivatives of levovirin possess the requisite physical and chemical properties and exhibit improved bioavailability.

The present invention relates to nucleoside compounds according to formula I, methods



of treating a disease mediated by Hepatitis C Virus (HCV) comprising administering to a mammal a therapeutically effective amount of a compound according to formula I, and pharmaceutical compositions comprising a therapeutically effective amount of a compound of formula I and at least one pharmaceutically acceptable carrier and

5 optionally containing excipients

wherein

R^1 , R^2 and R^3 are independently selected from the group consisting of hydrogen, C_{1-10} acyl, C_{1-10} alkoxy carbonyl and COR^4 where COR^4 is an amino acid or a dipeptide; and, hydrates, solvates, clathrates of said compound and acid addition salts.

10

An embodiment of the present invention is a nucleoside compound according to formula I wherein R^1 , R^2 and R^3 are as defined hereinabove.

In another embodiment there is provided a compound according to formula I wherein

15 one of R^1 , R^2 and R^3 is COR^4 , R^4 is $CH(R^5)NH_3^+ Cl^-$ or pyrrolidin-2-yl, R^5 is the side chain of a naturally occurring hydrophobic amino acid or C_{1-6} straight or branched alkyl and the other of R^1 , R^2 and R^3 are independently selected from the group consisting of hydrogen, C_{1-10} acyl, and C_{1-10} alkoxy carbonyl.

20 In another embodiment there is provided a compound according to formula I wherein R^1 is COR^4 , and R^4 is $CH(R^5)NH_3^+ Cl^-$ or pyrrolidin-2-yl, R^5 is the side chain of a naturally occurring hydrophobic amino acid or C_{1-6} straight or branched alkyl and R^2 and R^3 are independently selected from the group consisting of hydrogen, C_{1-10} acyl, and C_{1-10} alkoxy carbonyl.

25

In another embodiment there is provided a compound according to formula I wherein R^1 is COR^4 , and R^4 is $CH(R^5)NH_3^+ Cl^-$, R^5 is selected from the group consisting of $CH(CH_3)_2$ and $CH(CH_3)CH_2CH_3$ and both R^2 and R^3 are hydrogen.

30 In an another embodiment of the present invention there is provided an acid addition salt of a compound according to formula I wherein one of R^1 , R^2 and R^3 is COR^4 , R^4 is $CH(R^5)NH_2$, R^5 is the side chain of valine and the other of R^1 , R^2 and R^3 are independently selected from the group consisting of hydrogen.

In another embodiment of the present invention there is provided an acid addition salt of a compound according to formula I wherein R^1 is COR^4 , R^4 is the side chain of valine and both R^2 and R^3 are hydrogen.

- 5 In a preferred embodiment of the present invention there is provided the hydrochloride salt of a compound according to formula I wherein R^1 is COR^4 , and R^4 is the side chain of L-valine and both R^2 and R^3 are hydrogen.

A preferred representative of the compounds of the present invention is 2S-amino-3-methyl-butiric acid 5S-(3-carbamoyl-{1,2,4}triazol-1-yl)-3R,4S-dihydroxy-tetrahydro-furan-2S-ylmethyl ester hydrochloride.

In another embodiment there is provided a compound according to formula I wherein R^1 is COR^4 , and R^4 is $CH(R^5)NH_3^+ Cl^-$, R^5 is CH_3 , and both R^2 and R^3 are hydrogen.

15

In another embodiment there is provided a compound according to formula I wherein R^1 , R^2 and R^3 are C_{1-10} acyl or C_{1-10} alkoxy-carbonyl.

A preferred representative of such compounds is propionic acid 3S,4S-*bis*-propionyloxy-5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-tetrahydro-furan-2S-ylmethyl ester.

20


In another embodiment there is provided a compound according to formula I wherein R^1 is C_{1-10} acyl or C_{1-10} alkoxy-carbonyl and both R^2 and R^3 are hydrogen.

- 25 In another embodiment there is provided a compound according to formula I wherein R^1 is hydrogen and both R^2 and R^3 are independently C_{1-10} acyl or C_{1-10} alkoxy-carbonyl.

In a preferred embodiment of the present invention the following compounds are provided:

- 30 isobutyric acid 2S-(3-carbamoyl-[1,2,4]triazol-1-yl)-5S-hydroxymethyl-4S-isobutyryloxy-tetrahydro-furan-3S-yl ester; or
2,2-dimethylpropionic acid 4S-(2,2-dimethylpropionyloxy)-5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-2S-hydroxymethyl-tetrahydro-furan-3S-yl ester.

- 35 In another embodiment of the present invention is a nucleoside compound according to formula II



wherein R^1 , R^2 and R^3 are independently selected from the group consisting of
 5 hydrogen, C_{1-10} acyl, C_{1-10} alkoxycarbonyl and COR^4 where COR^4 is an amino acid or a dipeptide; R^6 is C_{1-6} acyl; and, hydrates, solvates, clathrates of said compound and acid addition salts.

The compounds of the present invention characterized by the formula I or II can be
 10 used in therapy, especially in the therapy of diseases mediated by the Hepatitis C Virus.

The compounds of the present invention characterized by the formula I or II can be administered to a mammal in a therapeutically effective amount, as a rule at a dose of
 15 between 0.1 and 300 mg/kg of body weight of the patient per day, preferably at a dose of between 1 and 100 mg/kg of body weight of the patient per day, more preferably at a dose of between 1 and 50 mg/kg of body weight of the patient per day.

In another embodiment of the present invention the compounds of the present invention
 20 characterized by the formula I or II can be administered in combination with an immune system modulator or antiviral agent, such as an interferon, interleukin, tumor necrosis factor, colony stimulating factor, anti-inflammatory agent or with a reverse transcriptase inhibitor.

25 Said interferon can be a chemically-derivatized interferon, such as for example PEG-interferon- α -2a (PEGASYS®) or PEG-interferon- α -2b (PEG-INTRON®).

One skilled in the art of protein chemistry will recognize that the development of new methodology to link polymers and proteins is ongoing and new methods of attaching
 30 polymers to interferon would produce compounds which fall within the scope of this invention when co-administered with levovirin prodrugs to treat disease mediated by HCV.

In another embodiment of the present invention there is provided a pharmaceutical composition comprising a therapeutically effective amount of a compound of formula I and II and at least one pharmaceutically acceptable carrier and optionally containing
5 excipients.

DEFINITIONS

- 10 The phrase "a" or "an" entity as used herein refers to one or more of that entity; for example, a compound refers to one or more compounds or at least one compound. As such, the terms "a" (or "an"), "one or more", and "at least one" can be used interchangeably herein.
- 15 The phrase "as defined hereinabove" refers to the first definition provided in the Detailed Description of the Invention.

The term "alkyl" as used herein denotes an unbranched or branched chain hydrocarbon residue containing 1 to 12 carbon atoms. The term "lower alkyl" denotes an unbranched
20 or branched chain hydrocarbon residue containing 1 to 6 carbon atoms. Representative lower alkyl groups include methyl, ethyl, propyl, *i*-propyl, *n*-butyl, *i*-butyl, *t*-butyl or pentyl.

The term "acyl" means an organic radical of the formula R-C(O)-, formally derived from
25 an organic acid by the removal of the hydroxyl group; the term " C_{1-12} acyl" refers to a acyl group wherein R is alkyl or aryl of 1-12 carbon atoms; and, the term "lower acyl" as used herein refers to a acyl group wherein R is C_{1-6} straight, branched or cyclic alkyl. The term "aroyl" as used herein refers to an acyl group wherein R is an aryl group.

30 The term "alkoxy" as used herein denotes an organic radical of the formula R-O- wherein the "alkyl" portion is as defined above such as methoxy, ethoxy, *n*-propyloxy, *i*-propyloxy, *n*-butyloxy, *i*-butyloxy, *t*-butyloxy, pentyloxy, hexyloxy, heptyloxy including their isomers. "Lower alkoxy" as used herein denotes an alkoxy group with a "lower alkyl" group as previously defined.

35

The term "alkoxycarbonyl" as used herein means an organic radical of the formula R-O-C(O)- where R-O- is an alkoxy group as defined herein.

The term "naturally occurring amino acids" as used herein means the L-isomers of the naturally occurring amino acids. The naturally occurring amino acids are glycine, alanine, valine, leucine, isoleucine, serine, methionine, threonine, phenylalanine, tyrosine, tryptophan, cysteine, proline, histidine, aspartic acid, asparagine, glutamic acid, glutamine, γ -carboxyglutamic acid, arginine, ornithine and lysine. Unless specifically indicated, all amino acids referred to in this application are in the L-form. The term "hydrophobic amino acid" as used herein glycine, alanine, valine, leucine, isoleucine, methionine, phenylalanine, tryptophan, and proline.

10

Compounds of formula I which are basic can form pharmaceutically acceptable salts with inorganic acids such as hydrohalic acids (e.g. hydrochloric acid and hydrobromic acid), sulphuric acid, nitric acid and phosphoric acid, and the like, and with organic acids (e.g. with acetic acid, tartaric acid, succinic acid, fumaric acid, maleic acid, malic acid, salicylic acid, citric acid, methanesulphonic acid and *p*-toluenesulfonic acid, and the like).

15

The term "solvate" as used herein means a compound of the invention or a salt, thereof, that further includes a stoichiometric or non-stoichiometric amount of a solvent bound by non-covalent intermolecular forces. Preferred solvents are volatile, non-toxic, and/or acceptable for administration to humans in trace amounts.

20

The term "hydrate" as used herein means a compound of the invention or a salt thereof, that further includes a stoichiometric or non-stoichiometric amount of water bound by non-covalent intermolecular forces.

25

The term "clathrate" as used herein means a compound of the invention or a salt thereof in the form of a crystal lattice that contains spaces (e.g., channels) that have a guest molecule (e.g., a solvent or water) trapped within.

30 The term "immunomodulator" as used herein means a therapeutic agent that assists in or is capable of modifying or regulating immune functions. An agent that causes an immunological adjustment, regulation or potentiation

The term "interferon" as used herein means the family of proteins capable of interfering with the viral infection of cells, as well as inhibiting the proliferation of normal and transformed cells, regulating cell differentiation and modulating the immune system. The four major antigenic types of interferon (α , β , γ and ω) are defined by the cellular source

35

of their production. Type I interferons (interferon (α , β and ω) compete with each other for cellular binding to the type I interferon receptor and thus share at least some components of this multi-subunit cell surface receptor, while the receptor for type II interferon (interferon γ) is a distinct entity. Both naturally-occurring and recombinant
5 interferons may be administered in combination therapy with compounds of the invention. A consensus sequence for interferon has been described in U.S. Pat. No. 4,897,471 (Y. Stabinsky).

The term "chemically-derivatized interferon" as used herein refers to an interferon
10 molecule covalently linked to a polymer which alters the physical and/or pharmacokinetic properties of the interferon. A non-limiting list of such polymers include polyalkylene oxide homopolymers such as polyethylene glycol (PEG) or polypropylene glycol (PPG), polyoxyethylenated polyols, copolymers thereof and block copolymers thereof, provided that the water solubility of the block copolymers is
15 maintained. One skilled in the art will be aware of numerous approaches to linking the polymer and interferon (for example, see A. Kozlowski and J. M. Harris *J. Control. Release* 2001 72(1-3):217-24; C. W. Gilbert and M. Park-Cho, U.S. Pat. No. 5,951,974). A non-limiting list of chemically derivatized IFN α contemplated in the present patent include PEG-interferon- α -2a (PEGASYS®) and PEG-interferon- α -2b (PEGINTRON™).

20

ABBREVIATIONS

The following abbreviations are used throughout this application and they have the meaning listed below:

25 THF: tetrahydrofuran

DMF: N,N-dimethylformamide

CBZ: benzyloxycarbonyl

PyBOP: benzotriazol-1-yloxy *tris*-pyrrolidino phosphonium hexafluorophosphate

IPA: isopropyl alcohol

30 DMAP: 4-N,N-dimethylaminopyridine

DIPEA: N,N-diisopropylethylamine

TEA: triethylamine

DEAD: diethylazodicarboxylate

PTLC: preparative thin layer chromatography

35 TsOH: *p*-toluenesulfonic acid monohydrate

NOMENCLATURE


In general the nomenclature used in this application is based on AUTONOM™ v4.0, a Beilstein Institute computerized system for generation of IUPAC systematic

5 nomenclature.

EXAMPLES OF COMPOUNDS OF THE PRESENT INVENTION

The following examples and preparations are provided to enable those skilled in the art to more clearly understand and to practice the present invention. They should not be considered as limiting the scope of the invention, but merely as being illustrative and representative thereof. Compounds in Table 1 are examples of mono-, di- and triacyl derivatives of levovirin. Compounds in Table 2 exemplify N-acyl levovirin derivatives. Compounds in Table 3 exemplify synthetic intermediates in which the one or more hydroxyl groups are protected and acylated compounds containing a ketal or acetal with the 2', 3' hydroxy groups.

TABLE 1
Acylated Levovirin Derivatives




No	R ¹	R ²	R ³	Salt	Method ₃	m.s. ⁴	m.p. ⁶
1	MeCO	MeCO	MeCO		A	371	
2	EtCO	EtCO	EtCO		A	413	
3	H	Val	Val	HCl	C	477	202-205
4	Val	H	H	Tos	B	344	110-114.5
5	Val	H	H	HCl	B	344	154-156
6	(D)-Val	H	H	Tos	B	344	
7	Ala	H	H	Tos	B	316	108-120
8	Phe	H	H	Tos	B	392	114-136
9	Leu	H	H	Tos	B	358	112-123
10	Ile	H	H	Tos	B	358	101.8-110.8
11	<i>t</i> -BuCO	H	H	-	B	329	139-141.6
12	<i>i</i> -PrCO	H	H	-	B	315	169-171.2
13	Gly	H	H	Tos	B	302	89.3-96.4
14	MeNHCH ₂ C O	H	H	Tos	B	316	69.4-86.3
15	H	<i>t</i> -BuCO	<i>t</i> -BuCO	-	C		
16	<i>i</i> -Pr-OCO	H	H	-	B	329	46-59
17	H	<i>i</i> -PrCO	<i>i</i> -PrCO	-	C	407 ⁵	179.0-179.6

No	R ¹	R ²	R ³	Salt	Method ³	m.s. ⁴	m.p. ⁶
18	H-Val-Pro	H	H	HCl	B	441	146-149
19	H	EtCO	EtCO	-	C	357	154.2-155.6
20	<i>n</i> -PrCO	<i>n</i> -PrCO	<i>n</i> -PrCO	-	A	455	
21	Val	EtCO	EtCO	Tos	D	456	60.0-63.5
22	Val	<i>i</i> -PrCO	<i>i</i> -PrCO	Tos	D	506 ⁵	72.0-76.0
23	H-Pro-Val ¹	H	H	Tos	B	441	76-92
24	EtOCO	EtOCO	EtOCO	-	A	461	
25	PhCO	PhCO	PhCO	-	A	579	
26	<i>i</i> -PrCO	<i>i</i> -PrCO	<i>i</i> -PrCO	-	A	477 ⁵	
27	<i>t</i> -BuCO	<i>t</i> -BuCO	<i>t</i> -BuCO	-	A	519 ⁵	
28	H	PhCO	PhCO	-	C	453	
29	<i>t</i> -BuCO	H	<i>t</i> -BuCO	-	C	413	
30	H	<i>n</i> -PrCO	<i>n</i> -PrCO	-	C	407	135.3-135.9
31	<i>n</i> -C ₆ H ₁₃ CO	H	H	-	B	357	151.2-152.8
32	<i>n</i> -PrOCO	<i>n</i> -PrOCO	<i>n</i> -PrOCO	-	A	503	51.7-56.6
33	H-Pro-Val ²	H	H	Tos	B	441	120-136
34	C ₇ H ₁₅ CO	H	H	-	B	371	154.4-155.8
35	C ₈ H ₁₇ CO	H	H	-	B	385	155-157.1
36	EtCO	H	H	-	B	301	178-181.8
47	H	H	EtCO	-	C	301	
48	H	H	<i>t</i> -BuCO	-	C	329	
49	H	H	<i>i</i> -PrCO	-	C	315	
51	H	<i>n</i> -BuCO	<i>n</i> -BuCO	-	C	435 ⁵	115.0-118.1
52	H	<i>n</i> -C ₅ H ₁₁ CO	<i>n</i> -C ₅ H ₁₁ CO	-	C	463 ⁵	114.8-115.3
53	H	<i>n</i> -PrOCO	<i>n</i> -PrOCO	-	C	417	101.0-103.0
54	H	<i>c</i> -C ₆ H ₁₁ CO	<i>c</i> -C ₆ H ₁₁ CO	-	C	487	195.6-197.5
55	(<i>n</i> -Pr) ₂ CHCO	H	H	-	B	393	179.0-179.9
56	<i>c</i> -C ₆ H ₁₁ CO	H	H	-	B	355	168.5-171.9
57	<i>n</i> -C ₇ H ₁₅ OCO	H	H	-	B	387	111.1-114.5
58	H	(Et) ₂ CHC O	(Et) ₂ CHC O	-	C	463 ⁵	154.9-160.3
59	<i>n</i> -C ₈ H ₁₇ OCO	H	H	-	B	401	126.3-129.1

Val= *i*-PrCH(NH₂)COLeu
=Me₂CHCH₂CH(NH₂)COPhe= PhCH₂CH(NH₂)COIle=EtCH(Me)CHNH₂C
OAla= CH₃CH(NH₂)COGly= NH₂CH₂CO

H-Val-Pro =



H-Pro-Val =




¹ more polar isomer² less polar isomer³ The methods listed herein refer to designations in the Examples (*infra*).⁴ mass spectra (M+H)⁺⁵ (M+Na)⁺⁶ melting point (°C)

TABLE 2
Acylated N-Acyl Levovirin Derivatives



No.	R ¹	R ²	R ³	R ⁶	m.s.	Method
37	EtCO	EtCO	EtCO	EtCO	469	E
38	H	<i>n</i> -PrCO	<i>n</i> -PrCO	<i>n</i> -PrCO	477 (M+Na) ⁺	E
39	<i>n</i> -PrCO	<i>n</i> -PrCO	<i>n</i> -PrCO	<i>n</i> -PrCO	547 (M+Na) ⁺	E
40	H	H	H	<i>n</i> -PrCO	315	E
50	<i>n</i> -PrCO	H	H	<i>n</i> -PrCO	477 (M+Na) ⁺	E

TABLE 3
Protected Synthetic Intermediates



	R1	R2	R3	m.s. ¹	m.p. ⁵
41	H	C(CH ₃) ₂		285	95.1-98
42	H	CHPh		333	150-153
43		Si(<i>i</i> -Pr) ₂ O Si(<i>i</i> -Pr) ₂			
44	(<i>i</i> -Pr) ₃ Si	H	H	423 ²	174.6-175.7
45	Val	C(CH ₂) ₄			
46	<i>t</i> -BuCO	C(CH ₃) ₂			

¹ mass spectra (M+H)⁺

² (M+Na)⁺

³ melting point (°C)

5

PREPARATION OF COMPOUNDS

The compounds of formula I may be prepared by various methods known in the art of organic chemistry in general and nucleoside analogue synthesis in particular. The starting materials for the syntheses are either readily available from commercial sources or are known or may themselves be prepared by techniques known in the art. The following examples (*infra*) are given to enable those skilled in the art to more clearly understand and to practice the present invention. They should not be considered as limiting the

10

scope of the invention, but merely as being illustrative and representative thereof.

General reviews of the preparation of nucleoside analogues are included in the following publications:

- 5 A M Michelson "The Chemistry of Nucleosides and Nucleotides", Academic Press, New York 1963.
- L Goodman "Basic Principles in Nucleic Acid Chemistry" Ed P O P Ts'O, Academic Press, New York 1974, Vol. 1, chapter 2.
- "Synthetic Procedures in Nucleic acid Chemistry" Ed W W Zorbach and R S Tipson, Wiley, New York, 1973, Vol. 1 and 2.
- 10 H. Vorbrüggen and C. Ruh-Pohlenz (eds) "Handbook of Nucleoside Synthesis" Wiley, New York, 2001.

Efforts have been made to ensure accuracy with respect to numbers used (e.g., amounts, temperatures), but allowance for some experimental error and deviation, including differences in calibration, rounding of numbers, and the like, is contemplated.

15

FORMULATIONS AND ADMINISTRATION

Formulations of compounds of formula I may be prepared by processes known in the formulation art. The following examples (*infra*) are given to enable those skilled in the art to more clearly understand and to practice the present invention. They should not be considered as limiting the scope of the invention, but merely as being illustrative and representative thereof.

While nucleoside derivatives of the present invention are optimized for delivery across the gastrointestinal mucosa, these compounds can be efficacious when administered by other routes of administration including continuous (intravenous drip) topical parenteral, intramuscular, intravenous, subcutaneous, transdermal (which may include a penetration enhancement agent), buccal, nasal and suppository administration, among other routes of administration. Oral administration can be in the form of tablets, coated tablets, dragées, hard and soft gelatine capsules, solutions, emulsions, syrups, or suspensions

For the manufacture of pharmaceutical preparations, the nucleoside derivatives, as well as their pharmaceutically useable salts, can be formulated with a therapeutically inert, inorganic or organic excipient for the production of tablets, coated tablets, dragées, hard and soft gelatine capsules, solutions, emulsions or suspensions. The compounds of

formula I can be formulated in admixture with a pharmaceutically acceptable carrier. For example, the compounds of the present invention can be administered orally as pharmacologically acceptable salts. Because the compounds of the present invention are mostly water soluble, they can be administered intravenously in physiological saline solution (e.g., buffered to a pH of about 7.2 to 7.5). Conventional buffers such as phosphates, bicarbonates or citrates can be used in the present compositions. Suitable excipients for tablets, coated tablets, dragées, and hard gelatin capsules are, for example, lactose, corn starch and derivatives thereof, talc, and stearic acid or its salts. If desired, the tablets or capsules may be enteric-coated or sustained release by standard techniques. Suitable excipients for soft gelatine capsules are, for example, vegetable oils, waxes, fats, semi-solid and liquid polyols. Suitable excipients for injection solutions are, for example, water, saline, alcohols, polyols, glycerin or vegetable oils. Suitable excipients for suppositories are, for example, natural and hardened oils, waxes, fats, semi-liquid or liquid polyols. Suitable excipients for solutions and syrups for enteral use are, for example, water, polyols, saccharose, invert sugar and glucose. The pharmaceutical preparations can also contain preservatives, solubilizers, stabilizers, wetting agents, emulsifiers, sweeteners, colorants, flavorants, salts for adjustment of the osmotic pressure, buffers, masking agents or antioxidants. The pharmaceutical preparations may also contain other therapeutically active agents known in the art. Suitable pharmaceutical carriers, excipients and their formulations are described in *Remington: The Science and Practice of Pharmacy* 1995, edited by E. W. Martin, Mack Publishing Company, 19th edition, Easton, Pennsylvania. Representative pharmaceutical formulations containing a compound of the present invention are described in Examples 13-15.

One of ordinary skill in the formulations art will also take advantage of favorable physical and pharmacokinetic parameters of the prodrug forms, where in delivering the present compounds to targeted site within the host organism or patient to maximize the intended effect of the compound. A skilled formulation scientist may modify the formulations within the teachings of the specification to provide numerous formulations for a particular route of administration without rendering the compositions of the present invention unstable or compromising their therapeutic activity.

In particular, the modification of the present compounds to render them more soluble in water or other vehicle, for example, may be easily accomplished by minor modifications (salt formulation, esterification, etc.) which are well within the ordinary skill in the art. It is also well within the ordinary skill of the art to modify the route of administration and


dosage regimen of a particular compound in order to manage the pharmacokinetics of the present compounds for maximum beneficial effect in patients.

The dosage can vary within wide limits and will, of course, be adjusted to the individual requirements in each particular case. For oral administration, a daily dosage of between about 0.01 and about 100 mg/kg body weight per day should be appropriate in monotherapy and/or in combination therapy. A preferred daily dosage is between about 0.1 and about 300 mg/kg body weight, more preferred 1 and about 100 mg/kg body weight and most preferred 1.0 and about 50 mg/kg body weight per day. A typical preparation will contain from about 5% to about 95% active compound (w/w). The daily dosage can be administered as a single dosage or in divided dosages, typically between 1 and 5 dosages per day.

The pharmaceutical preparations are preferably in unit dosage forms. In such form, the preparation is subdivided into unit doses containing appropriate quantities of the active component. The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation, such as packeted tablets, capsules, and powders in vials or ampoules. Also, the unit dosage form can be a capsule, tablet, cachet, or lozenge itself, or it can be the appropriate number of any of these in packaged form.

The nucleoside derivatives or the medicaments thereof may be used in monotherapy or combination therapy, i.e. the treatment may be in conjunction with the administration of one or more additional therapeutically active substance(s), for example, an immune system modulator such as an interferon, interleukin, tumor necrosis factor or colony stimulating factor or an anti-inflammatory agent and/or an antiviral agent. When the treatment is combination therapy, such administration may be concurrent or sequential with respect to that of the nucleoside derivatives. Concurrent administration, as used herein thus includes administration of the agents at the same time or at different times.

The references herein to treatment extend to prophylaxis of Hepatitis C mediated diseases as well as to the treatment of existing conditions, and that the treatment of animals includes the treatment of humans as well as other mammals. Furthermore, treatment of an Hepatitis C Virus (HCV) infection, as used herein, also includes treatment or prophylaxis of a disease or a condition associated with or mediated by Hepatitis C Virus (HCV) infection, or the clinical symptoms thereof.

EXAMPLE 1

1-(6S-hydroxymethyl-2,2-dimethyl-tetrahydro-3aS,6aS-furo[3,4-d][1,3] dioxol-4S-yl)-
 5 1H-[1,2,4]triazole-3-carboxylic acid amide (3, R'=R''=CH₃).

Levovirin (1, 1.0 g, 4.1 mmol, Roche Carolina) was suspended in 32 mL of a 2:1 mixture of dry acetone:2,2-dimethoxypropane. The solution was stirred under N₂ in an ice bath and 7 drops of concentrated perchloric acid were added dropwise. The reaction was stirred to room temperature over 4 hours. The mixture was neutralized by addition of
 10 1M sodium hydroxide solution and evaporated to a residue. The residue was purified *via* chromatography (silica gel; 5%-10% methanol/dichloromethane) to yield 0.72 g (62%) 1-(6S-hydroxymethyl-2,2-dimethyl-tetrahydro-3aS,6aS-furo[3,4-d][1,3]dioxol-4S-yl)-1H-[1,2,4]triazole-3-carboxylic acid amide. (compound 41 (3, R' = R'' = CH₃); (M+H)⁺=285; mp=95.1-98°C).

15


EXAMPLE 2

1-(6S-hydroxymethyl-2-phenyl-tetrahydro-3aS,6aS-furo[3,4-d][1,3]dioxol-4S-yl)-1H-
 [1,2,4]triazole-3-carboxylic acid amide (3; R'=H, R''=Ph).

20 Levovirin (6.00 g, 24.5 mmol, Roche Carolina) was suspended in 60 mL of benzaldehyde. Zinc chloride (5.70 g, 41.8 mmol, Aldrich Chemical) was added to the stirred mixture. After 4 hours, the reaction mixture was added dropwise to 1 l of rapidly stirred diethyl ether. The precipitate formed was filtered, rinsed with ether and then dissolved in 350 mL of ethyl acetate and 650 mL of cold 2M sodium hydroxide solution. The layers were
 25 separated and the aqueous layer was extracted two times more with ethyl acetate. The combined ethyl acetate layers were washed with brine, dried over sodium sulfate and evaporated to a solid. The solid was triturated with ether and purified by silica gel chromatography (2%-7% methanol/dichloromethane) to yield 4.4 g (54%) 1-(6S-hydroxymethyl-2-phenyl-tetrahydro-3aS,6aS-furo[3,4-d][1,3]dioxol-4S-yl)-1H-
 30 [1,2,4]triazole-3-carboxylic acid amide (compound 42 (3, R' = Ph, R'' = H); (M+H)⁺=333; mp=150-153°C).

EXAMPLE 3

1-(3R-Hydroxy-5,5,7,7-tetraisopropyl-tetrahydro-1,4,6,8-tetraoxa-5,7-disila-3aS,9aS-cyclopentacycloocten-2S-yl)-1H-[1,2,4]triazole-3-carboxylic acid amide.




5

To a stirred slurry of levovirin (3.75 g, 15.4 mmol) in 30 mL of DMF were added 30 mL of pyridine, TEA (5.35 mL, 38.4 mmol), and 1,3-dichloro-1,1,3,3-tetraisopropyl-disiloxane (6.15 mL, 19.2 mmol) at 0°C. The reaction was allowed reaction to warm to room temperature and stirred for 24 hours. The resulting solution was partitioned
10 between 1 N HCl and ethyl acetate. Organic layers were washed with brine, dried over MgSO₄ and concentrated *in vacuo*. The residue was purified *via* chromatography (25% acetone/chloroform) to yield 3.96 g (53%) 1-(3R-Hydroxy-5,5,7,7-tetraisopropyl-tetrahydro-1,4,6,8-tetraoxa-5,7-disila-3aS,9aS-cyclopentacycloocten-2S-yl)-1H-[1,2,4]triazole-3-carboxylic acid amide (compound 43).

15

EXAMPLE 4

1-(3S,4R-dihydroxy-5S-triisopropylsilanyloxymethyl-tetrahydro-furan-2S-yl)-1H-[1,2,4]triazole-3-carboxylic acid amide




To a stirred slurry of levovirin (1, 9.22 g, 37.8 mmol) in 75 mL of DMF were added at room temperature imidazole (2.80 g, 41.1 mmol) and triisopropylsilyl chloride (8.1 mL, 38 mmol). The suspension was heated to 50°C which dissolved the solids and temperature was maintained for 6 hrs. The solution was transferred to a separatory funnel and partitioned between 400 mL of ethyl acetate and 500 mL of water. The
25 organic layer (slurry) was washed with 200 mL of water three times and precipitate was filtered off. Organic layer filtrate was dried over MgSO₄ and concentrated *in vacuo*. Filtered residue was recrystallized in 150 mL of methanol; the mother liquor was

combined with organic concentrate and recrystallized further. Four crops of crystals were collected to yield 10.47 g (69%) 1-(3*S*,4*R*-dihydroxy-5*S*-triisopropylsilanyloxymethyl-tetrahydrofuran-2*S*-yl)-1*H*-[1,2,4]triazole-3-carboxylic acid amide as a white crystalline solid (compound 44; $(M+Na)^+ = 423$; mp=174.6-175.7°C).

Method A - Preparations of trisubstituted Analogs

EXAMPLE 5

Isobutyric acid 3*S*,4*S*-bis-isobutyroxy-5*S*-(3-carbamoyl-[1,2,4]triazol-1-yl)-tetrahydro-furan-2*S*-ylmethyl ester (2; $R^1 = CH(CH_3)_2$)




To a stirred slurry of levovirin (0.46 g, 1.88 mmol) in 8 mL of THF under nitrogen were added TEA (1.31 mL, 9.40 mmol) and isobutyric anhydride (1.41 mL, 8.48 mmol). The reaction vessel was fitted with cold finger attachment and heated to 65°C for 24 hours. The reaction was partitioned between ethyl acetate and a saturated aqueous $NaHCO_3$ solution. The organic layer was washed with brine, dried over $MgSO_4$, and concentrated. The residue was purified *via* silica chromatography (3% methanol/dichloromethane) to yield 0.278 g (33%) isobutyric acid 3*S*,4*S*-bis-isobutyroxy-5*S*-(3-carbamoyl-[1,2,4]triazol-1-yl)-tetrahydro-furan-2*S*-ylmethyl ester as a gummy solid. MS: (compound 26(2, $R = CH(CH_3)_2$; $(M+Na)^+ = 477$).

Proceeding in analogous fashion with the appropriate acid anhydride there was prepared: 2,2-dimethylpropionic acid 3*S*,4*S*-bis-(2,2-dimethylpropionyloxy)-5*S*-(3-carbamoyl-[1,2,4]triazol-1-yl)-tetrahydro-furan-2*S*-ylmethyl ester (compound 27; 18%); benzoic acid 3*S*,4*S*-bis-benzoyloxy-5*S*-(3-carbamoyl-[1,2,4]triazol-1-yl)-tetrahydro-furan-2*S*-ylmethyl ester (compound 25; 66%); acetic acid 3*S*,4*S*-bis-acetoxy-5*S*-(3-carbamoyl-[1,2,4]triazol-1-yl)-tetrahydro-furan-2*S*-ylmethyl ester (compound 1; 65%; $(M+H)^+ = 371$); propionic acid 3*S*,4*S*-bis-propionyloxy-5*S*-(3-carbamoyl-[1,2,4]triazol-1-yl)-tetrahydro-furan-2*S*-ylmethyl ester as a clear oil (compound 2; 52%; $(M+H)^+ = 413$);

butyric acid 3S,4S-bis-butyryloxy-5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-tetrahydro-furan-2S-ylmethyl ester as a clear oil (compound 20; 20%; $(M+H)^+ = 455$).

EXAMPLE 6

5 Carbonic acid 2S-(3-carbamoyl-[1,2,4]triazol-1-yl)-4S-ethoxycarbonyloxy-5S-ethoxycarbonyl-oxymethyl-tetrahydro-furan-3S-yl ester ethyl ester.




24

Levovirin (1, 0.5 g, 2.04 mmol, Roche Carolina) was suspended in 3 mL of DMF and 1.5 mL of pyridine. The mixture was stirred in an ice bath and ethyl chloroformate (0.78 mL, 8.19 mmol) was added slowly in three portions over 15 minutes. The reaction was stirred at room temperature for over 2 hours. Methanol was added and the reaction was stirred for 10 minutes. After evaporation, the residue was taken up in ethyl acetate and saturated ammonium chloride solution. The layers were separated and the aqueous layer was extracted with ethyl acetate once. The combined ethyl acetate layers were washed with brine and dried over sodium sulfate and concentrated. The foamy residue was purified via chromatography (3-4% methanol/dichloromethane) and lyophilization of a methanol/water solution gave solid carbonic acid 2S-(3-carbamoyl-[1,2,4]triazol-1-yl)-4S-ethoxycarbonyloxy-5S-ethoxycarbonyloxymethyl-tetrahydro-furan-3S-yl ester ethyl ester (compound 24, 74%, $(M+H)^+ = 461$).

20

Proceeding in analogous fashion with the appropriate alkyl chloroformate there was prepared: carbonic acid 2S-(3-carbamoyl-[1,2,4]triazol-1-yl)-4S-propoxycarbonyloxy-5S-propoxycarbonyl-oxymethyl-tetrahydro-furan-3S-yl ester propyl ester (compound 32, 47, $(M+H)^+ = 503$).

Method B - Preparations of 5'-monoacyl derivatives

5

EXAMPLE 7

2S-amino-3-methyl-butyric acid 5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-3R,4S-dihydroxy-tetrahydro-furan-2S-ylmethyl ester; compound with toluene-4-sulfonic acid (4: R¹ = CH(NH₂)CH(CH₃)₂).

- 10 1-(6S-Hydroxymethyl-2-phenyl-tetrahydro-3*a*S,6*a*S-furo[3,4-*d*][1,3]dioxol-4S-yl)-1*H*-[1,2,4]triazole-3-carboxylic acid amide (0.49 g, 1.47 mmol) was dissolved in 5 mL of dry DMF. N-CBZ-L-valine (0.44 g, 1.77 mmol, Aldrich Chemical), PyBOP (0.84 g, 1.62 mmol, Nova Biochem) and DIPEA (0.51 mL, 2.94 mmol) were added sequentially. After stirring for 18 hr ethyl acetate and saturated ammonium chloride solution were added.
- 15 The layers were separated and the aqueous layer was extracted with ethyl acetate once. The combined ethyl acetate layers were washed with water, saturated sodium bicarbonate solution, brine and dried over sodium sulfate. The solvent was evaporated and the residue was purified via chromatography (silica gel; gradient 2%-5% methanol/dichloromethane) to yield 520 mg (62%) of 2S-benzyloxycarbonylamino-3-methyl-butyric acid 6S-(3-carbamoyl-[1,2,4]triazol-1-yl)-2-phenyl-tetrahydro-3*a*S,6*a*S-furo[3,4-*d*][1,3]dioxol-4S-ylmethyl ester as a foam (M+H)⁺=566.
- 20

- 2S-Benzyloxycarbonylamino-3-methyl-butyric acid 6S-(3-carbamoyl-[1,2,4]triazol-1-yl)-2-phenyl-tetrahydro-3*a*S,6*a*S-furo[3,4-*d*][1,3]dioxol-4S-ylmethyl ester (0.49 g, 0.87 mmol) was dissolved in 5 mL of methanol containing 0.36 g of 20% palladium hydroxide on carbon (50 wt % water). TsOH (0.165 g, 0.87 mmol) was added and the reaction vessel was attached to a hydrogen gas-filled balloon. The vessel was purged with H₂ gas and stirred for 4.5 hr at 35°C. The mixture was then filtered through a bed of CELITE® and rinsed through with more methanol. After evaporation of solvent the residue was
- 30 dissolved in water and lyophilized to give 0.44 g (98%) of 2S-amino-3-methyl-butyric acid 5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-3R,4S-dihydroxy-tetrahydro-furan-2S-

ylmethyl ester; compound with toluene-4-sulfonic acid. (compound 4; $(M+H)^+ = 344$; m.p. = 110-114.5 °C);

Utilizing the two step sequence described above with the appropriate carboxylic acid
 5 there was obtained: 2S-amino-propionic acid 5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-3R,4S-dihydroxy-tetrahydro-furan-2S-ylmethyl ester; compound with toluene-4-sulfonic acid (compound 7; 98%; $(M+H)^+ = 316$; m.p. = 108-120 °C); 2S-amino-3-phenyl-propionic acid 5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-3R,4S-dihydroxy-tetrahydro-furan-2S-ylmethyl ester; compound with toluene-4-sulfonic acid (compound 8; 91%; $(M+H)^+ = 392$; m.p. = 114-136 °C); 2S-amino-4-methyl-pentanoic acid 5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-3R,4S-dihydroxy-tetrahydro-furan-2S-ylmethyl ester; compound with toluene-4-sulfonic acid (compound 9; 95%; $(M+H)^+ = 358$; m.p. = 112-123 °C); 2S-amino-3S-methyl-pentanoic acid 5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-3R,4S-dihydroxy-tetrahydro-furan-2S-ylmethyl ester; compound with toluene-4-sulfonic acid (compound
 10 10; 91%; $(M+H)^+ = 358$; m.p. = 101.8-110.8 °C); 2-methyl-propionic acid 5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-3R,4S-dihydroxy-tetrahydro-furan-2S-ylmethyl ester (compound 12; 94%; $(M+H)^+ = 315$; m.p. = 169-171.2 °C); 2-amino-acetic acid 5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-3R,4S-dihydroxy-tetrahydro-furan-2S-ylmethyl ester; compound with toluene-4-sulfonic acid (compound 13; 91%; $(M+H)^+ = 302$; m.p. =
 15 89.3-96.4 °C); 2-methyl-amino-acetic acid 5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-3R,4S-dihydroxy-tetrahydro-furan-2S-ylmethyl ester; compound with toluene-4-sulfonic acid (compound 14; 83%; $(M+H)^+ = 316$; m.p. = 69.4-86.3 °C).

EXAMPLE 8

25 2S-amino-3-methyl-butyric acid 5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-3R,4S-dihydroxy-tetrahydro-furan-2S-ylmethyl ester, hydrochloride (4: $R^1 = CH(NH_2)CH(CH_3)_2$).

A suspension of levovirin (1.350 mg, 1.43 mmol) in 9.5 mL of THF was treated with L-Val-CBZ (360 mg, 1.43 mmol) and triphenylphosphine (600 mg, 2.29 mmol). The
 30 reaction was stirred at rt and DEAD (0.28 mL, 1.8 mmol) was added dropwise. The reaction was stirred overnight at rt and the resulting suspension was concentrated and chromatographed (PTLC, 7% MeOH/ CH_2Cl_2) to give 2S-benzyloxycarbonylamino-3-methyl-butyric acid 5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-3R,4S-dihydroxy-tetrahydro-furan-2S-ylmethyl ester as a white solid (16%). MS: $MH^+ = 478$ (for reference to other
 35 nucleoside Mitsunobu couplings see: Wei, Y.; Pei, D. *Bioorg. Med. Chem. Lett.* 2000, 10(10), 1073).

Methanol (10 mL) and 1M HCl (0.7 mL) were added to a mixture of 2S-benzyloxycarbonylamino-3-methyl-butyrac acid 5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-3R,4S-dihydroxy-tetrahydro-furan-2S-ylmethyl ester (170 mg, 0.35 mmol) and 50 mg of 10% Pd/carbon. The resulting suspension was stirred under a hydrogen atmosphere (about 1 atm, balloon) for 30 min and the reaction was filtered through a pad of CELITE®. The filtrate was concentrated, diluted with water, and lyophilized to give 2S-amino-3-methyl-butyrac acid 5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-3R,4S-dihydroxy-tetrahydro-furan-2S-ylmethyl ester, hydrochloride as a pale yellow hygroscopic solid (compound 5, 75%, $(M+H)^+ = 344$). Recrystallization from dilute HCl/IPA gives a white crystalline solid; mp: 154-156 °C.

Utilizing the two step sequence described above with the appropriate carboxylic acid there was obtained: 2R-amino-3-methyl-butyrac acid 5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-3R,4S-dihydroxy-tetrahydro-furan-2S-ylmethyl ester; compound with toluene-4-sulfonic acid (compound 6; 90%; $(M+H)^+ = 344$).

EXAMPLE 9

2,2-Dimethyl-propionic acid 5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-3R,4S-dihydroxy-tetrahydro-furan-2S-ylmethyl ester (4: $R^1 = C(CH_3)_3$).

To a solution of 1-(6S-hydroxymethyl-2-phenyl-tetrahydro-3aS,6aS-furo[3,4-d][1,3]dioxol-4S-yl)-1SH-[1,2,4]triazole-3-carboxylic acid amide (0.24 g, 0.72 mmol) in 3 mL of 1:1 DMF/pyridine was added 2,2-dimethylpropionic anhydride (0.36 mL, 1.8 mmol) and DMAP (0.04 g, 0.36 mmol). The resulting solution was stirred overnight at rt and partitioned between 50 mL of ethyl acetate and sat. NH_4Cl solution. The aqueous layer was extracted with a second portion of ethyl acetate and the combined organic layers were washed with brine, dried over $MgSO_4$ and concentrated. The residue was chromatographed (PTLC, 5% MeOH/ CH_2Cl_2) to give 2,2-dimethyl-propionic acid 6S-(3-carbamoyl-[1,2,4]triazol-1-yl)-2-phenyl-tetrahydro-3aS,6aS-furo[3,4-d][1,3]dioxol-4S-ylmethyl ester as a clear oil (95%).

Methanol (6 mL) was added to a mixture of 2,2-dimethyl-propionic acid 6S-(3-carbamoyl-[1,2,4]triazol-1-yl)-2-phenyl-tetrahydro-3aS,6aS-furo[3,4-d][1,3]dioxol-4S-ylmethyl ester (0.37 g, 0.88 mmol) and 50% wet 10% $Pd(OH)_2/C$ (300 mg). The resulting suspension was stirred at 40 °C under a hydrogen atmosphere (about 1 atm, balloon) for 6 h and the reaction was filtered through a pad of CELITE®. The filtrate

was concentrated and the resulting oil was dissolved in 1.5 mL of MeOH and 10 mL of CH₂Cl₂ and then 3 mL of hexane was added until the solution just became cloudy. The resulting precipitated white solid was filtered to give 2,2-dimethyl-propionic acid 5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-3R,4S-dihydroxy-tetrahydrofuran-2S-ylmethyl ester

5 (compound 11; 70%; (M+H)⁺=329; m.p: 139-141.6 °C);

Utilizing the two step sequence described above with the appropriate carboxylic anhydride there was obtained: heptanoic acid 5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-3R,4S-dihydroxy-tetrahydro-furan-2S-ylmethyl ester (compound 31; 70%); propionic
10 acid 5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-3R,4S-dihydroxy-tetrahydro-furan-2S-ylmethyl ester (compound 36; 70%).

EXAMPLE 10

15 Octanoic acid 5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-3R,4S-dihydroxy-tetrahydrofuran-2S-ylmethyl ester (4: R¹ = C₇H₁₅).

1-(6S-hydroxymethyl-2-phenyl-tetrahydro-3aS,6aS-furo[3,4-*d*][1,3]dioxol-4S-yl)-1H-[1,2,4] triazole-3-carboxylic acid amide (0.25 g, 0.75 mmol) was dissolved in 1 mL DMF
20 and 0.5 mL of pyridine. The reaction solution was stirred in an ice bath and octanoyl chloride (0.16 mL, 0.94 mmol) was added dropwise. The reaction was then stirred at room temperature for 24 hr After concentration, the residue was partitioned between ethyl acetate and saturated ammonium chloride solution. The layers were separated and the aqueous layer was extracted with ethyl acetate once. The combined ethyl acetate
25 layers were washed with brine and dried over sodium sulfate. The residue after evaporation of solvent was purified by chromatography on silica gel in 5% methanol/dichloromethane to yield 0.2g (58%) of octanoic acid, 6S-(3-carbamoyl-[1,2,4]triazol-1-yl)-2-phenyl-tetrahydro-3aS,6aS-furo[3,4-*d*][1,3]dioxol-4S-ylmethyl ester was obtained; (M+H)⁺= 459. Hydrogenolysis of the benzylidene group was
30 accomplished as described in the preparation of compound 4 (*supra*) excluding the addition of TsOH to yield 102 mg (64%) octanoic acid 5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-3R,4S-dihydroxy-tetrahydrofuran-2S-ylmethyl ester as a crystalline solid (ethyl acetate-methanol). (compound 34; (M+H)⁺=371, m.p.=154.4-155.8 °C).

35 Proceeding as described above with the appropriate acid chloride there was prepared: nonanoic acid 5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-3R,4S-dihydroxy-tetrahydrofuran-2S-ylmethyl ester (compound 35; 82%; (M+H)⁺=385; m.p.=155-157.1);

EXAMPLE 11

Carbonic acid 5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-3R,4S-dihydroxy-tetrahydrofuran-2S-ylmethyl ester isopropyl ester (4: R¹ = O-*i*-C₃H₇).

- 5 1-(6S-hydroxymethyl-2-phenyl-tetrahydro-3aS,6aS-furo[3,4-*d*][1,3]dioxol-4S-yl)-1H-[1,2,4] triazole-3-carboxylic acid amide (0.3 g, 0.90 mmol) was dissolved in 2.4 mL of a 1:1 mixture of dry DMF:pyridine. The reaction was placed in an ice/salt bath and stirred as *iso*-propylchloroformate (Aldrich 1M toluene solution) was added slowly over 20 minutes. The bath was removed and the reaction was stirred for 5 hr after which 1 mL of
- 10 methanol was added and the reaction was stirred for 5 minutes more. The reaction was evaporated and the residue taken up in ethyl acetate and saturated ammonium chloride solution. The layers were separated and the aqueous layer was extracted with ethyl acetate. The combined ethyl acetate layers were washed with brine, dried over sodium sulfate and evaporated to a residue. The residue was purified by chromatography on silica
- 15 gel in 5% methanol/dichloromethane to yield 150 mg (40%) carbonic acid, 6S-(3-carbamoyl-[1,2,4]triazol-1-yl)-2-phenyl-tetrahydro-3aS,6aS-furo [3,4-*d*][1,3]dioxol-4S-ylmethyl ester isopropyl ester (M+H)⁺=419. Deprotection of carbonic acid 6S-(3-carbamoyl-[1,2,4]triazol-1-yl)-2-phenyl-tetrahydro-3aS,6aS-furo[3,4-*d*][1,3]dioxol-4S-ylmethyl ester *iso*-propyl ester as described for compound 4, in the absence of TsOH,
- 20 gave carbonic acid 5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-3R,4S-dihydroxy-tetrahydrofuran-2S-ylmethyl ester isopropyl ester (compound 16; 92%; (M+H)⁺=329; m.p.=46-59°C).

EXAMPLE 12

- 25 1-(2S-Amino-3-methyl-butyryl)-pyrrolidine-2S-carboxylic acid 5S-(3-carbamoyl-[1,2,4]-triazol-1-yl)-3R,4S-dihydroxy-tetrahydro-furan-2S-ylmethyl ester, hydrochloride.(4: R¹ = Pro-Val-H).


- 1-(6S-hydroxymethyl-2-phenyl-tetrahydro-3aS,6aS-furo[3,4-*d*][1,3]dioxol-4S-yl)-1H-[1,2,4]triazole-3-carboxylic acid amide (0.35 g, 1.05 mmol) was dissolved in 3.5 mL of
- 30 dry DMF. CBZ-NH-Val-Pro-OH (0.45 g, 1.32 mmol, Bachem), PyBOP (0.68 g, 1.32 mmol, Nova Biochem) and DIPEA (0.27 mL, 1.58 mmol) were added in sequentially. After stirring for 18 hr at 35°C, ethyl acetate and saturated ammonium chloride solution were added. The layers were separated and the aqueous layer was extracted with ethyl
- 35 acetate once. The combined ethyl acetate layers were washed with water, saturated sodium bicarbonate solution, brine and dried over sodium sulfate. The solvent was evaporated and the residue was purified by silica gel chromatography with 2% methanol/dichloromethane. Concentration of purified fractions yielded 370 mg (53%), 1-(2S-

benzyloxycarbonylamino-3-methyl-buteryl)-pyrrolidine-2S-carboxylic acid 6S-(3-carbamoyl-[1,2,4]triazol-1-yl)-2-phenyl-tetrahydro-3aS,6aS-furo[3,4-d][1,3]dioxol-4S-ylmethyl ester was obtained as a glass $(M+H)^+=663$.

- 5 Hydrogenolysis of the benzylidene protecting group was carried out as described for compound 4 (*supra*) replacing *p*-tuenesulfonic acid with HCl/ether (Aldrich, 1M solution) to yield 1-(2S-amino-3-methyl-buteryl)-pyrrolidine-2S-carboxylic acid 5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-3R,4S-dihydroxy-tetrahydro-furan-2S-ylmethyl ester, hydrochloride (compound 18; 79%; $(M+H)^+=441$; m.p.=146-149°C).

- 10 In analogous manner were prepared two isomers of 2-(pyrrolidine-2S-carboxamidyl)-3-methyl-butyrac acid 5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-3R,4S-dihydroxy-tetrahydro-furan-2S-ylmethyl ester; compound with toluene-4-sulfonic acid (compound 23, isomer 1; 88%; $(M+H)^+=441$; m.p.=76-92°C; compound 33, isomer 2; 92%; $(M+H)^+=441$;
15 m.p.=120-136°C).

Method C – Preparations of diacyl derivatives



- 20 **EXAMPLE 13**
Butyric acid 4S-butyryloxy-5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-2S-hydroxymethyl-tetrahydro-furan-3S-yl ester (7: $R^1 = C_3H_7$).

- To a stirred slurry of 1-(3S,4R-dihydroxy-5S-triisopropylsilanyloxymethyl-tetrahydro-furan-2S-yl)-1H-[1,2,4]triazole-3-carboxylic acid amide (0.40 g, 1.00 mmol) in 3.3 mL of
 25 THF were added TEA (0.48 mL, 3.46 mmol), n-butyric anhydride (0.49 mL, 2.97 mmol). The reaction vessel was fitted with a cold finger attachment and heated to 65 °C, under nitrogen for 17 hours. The reaction was partitioned between ethyl acetate and a saturated aqueous sodium bicarbonate solution. The organic layer was washed with
 30 brine, dried over $MgSO_4$, and concentrated. The residue was purified *via* silica gel chromatography (20% acetone/chloroform) to yield 47 g (88%) butyric acid 4S-butyryloxy-5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-2S-triisopropylsilanyloxymethyl-tetrahydrofuran-3S-yl ester as a clear oil.

To a stirred solution of butyric acid 4S-butyryloxy-5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-2S-triisopropylsilanyloxymethyl-tetrahydro-furan-3S-yl ester (0.47 g, 0.88 mmol) in 5 mL of acetonitrile were added 2.5 mL of 1 N H₂SO₄ at room temperature. After 16 hours, 30 mL of a saturated aqueous NaHCO₃ solution was added and product was extracted with ethyl acetate. The organic layer was washed with brine, dried over MgSO₄ and concentrated. The resulting residue was dissolved in methanol and the product precipitated with ethyl ether yielding 18 g (53%) butyric acid 4S-butyryloxy-5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-2S-hydroxymethyl-tetrahydrofuran-3S-yl ester as a white crystalline solid. (compound 30; (M+Na)⁺=407; m.p.=135.3-135.9 °C).

Proceeding as described but using the appropriate acid anhydride there was obtained: isobutyric acid 2S-(3-carbamoyl-[1,2,4]triazol-1-yl)-5S-hydroxymethyl-4S-isobutyryloxy-tetrahydro-furan-3S-yl ester (compound 17; 38%; (M+Na)⁺=407; m.p.=179.0-179.6 °C); propionic acid 4S-propionyloxy-5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-2S-hydroxymethyl-tetrahydro-furan-3S-yl ester (compound 19; 27%; (M+H)⁺=357; m.p.=154.2-155.6 °C); 2,2-dimethylpropionic acid 4S-(2,2-dimethylpropionyloxy)-5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-2S-hydroxymethyl-tetrahydro-furan-3S-yl ester (compound 15; 57%); benzoic acid 4S-benzoyloxy-5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-2S-hydroxymethyl-tetrahydro-furan-3S-yl ester (compound 28; 67%).

EXAMPLE 14

2S-Amino-3-methyl-butyric acid 4S-(2S-amino-3-methyl-butyryloxy)-5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-2S-hydroxymethyl-tetrahydrofuran-3S-yl ester dihydrochloride (7: R¹ = CH(NH₂)-i-C₃H₇).


To a stirred slurry of 1-(3S,4R-dihydroxy-5S-triisopropylsilanyloxymethyl-tetrahydrofuran-2S-yl)-1H-[1,2,4]triazole-3-carboxylic acid amide (0.47 g, 1.16 mmol) in 6 mL of THF were added at room temperature 4S-isopropyl-2,5-dioxooxazolidine-3-carboxylic acid benzyl ester (0.77 g, 2.79 mmol) and 8 drops of TEA. The reaction was allowed to stir for 16 hr and was quenched with 100 mL of a saturated aqueous NaHCO₃ solution and extracted with three 100 mL portions of ethyl acetate. The organic layers were combined, washed with brine, dried over MgSO₄, and concentrated. The resulting film was purified *via* silica gel chromatography (15% acetone/chloroform) to yield 0.53 g (53%) 2S-benzoyloxycarbonylamino-3-methyl-butyric acid 4S-(2S-

benzyloxycarbonylamino-3-methyl-butyryloxy)-5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-2S-triisopropylsilyloxymethyl-tetrahydrofuran-3S-yl ester as a clear oil.

To an argon sparged, stirred solution of 2-benzyloxycarbonylamino-3-methyl-butyric
5 acid 4-(2-benzyloxycarbonylamino-3-methyl-butyryloxy)-5-(3-carbamoyl-[1,2,4]triazol-1-yl)-2-hydroxymethyl-tetrahydro-furan-3-yl ester (0.32 g, 0.46 mmol) in 10 mL of ethanol were added hydrochloric acid (.6 mL, 1.81 mmol) and 0.20 g of 10% Pd/C. The reaction vessel was evacuated and purged with hydrogen gas (~1 atm) three times and left to stir for 6 hr. The suspension was filtered through CELITE® and filtrate was
10 concentrated. The residue was dissolved in a methanol/dichloromethane (1:10) solution and the product precipitated with hexane to yield 0.09 g (38%) 2S-amino-3-methyl-butyric acid 4S-(2S-amino-3-methyl-butyryloxy)-5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-2S-hydroxymethyl-tetrahydrofuran-3S-yl ester dihydrochloride as a white crystalline solid. (compound 3; (M+Cl)⁻=477; m.p.=202.0-205.0 °C).

EXAMPLE 15

2,2-Dimethyl-propionic acid 2S-(3-carbamoyl-[1,2,4]triazol-1-yl)-5S-(2,2-dimethyl-propionyloxymethyl)-4S-hydroxy-tetrahydro-furan-3S-yl ester (9: $R^1 = C(CH_3)_3$).



To a stirred slurry of 1-(3R-hydroxy-5,5,7,7-tetraisopropyl-tetrahydro-1,4,6,8-tetraoxa-5,7-disila-3aS,9aS-cyclopentacycloocten-2S-yl)-1H-[1,2,4]triazole-3-carboxylic acid amide (0.92 g, 1.88 mmol) in 7 mL of 1:1 DMF:pyridine were added DMAP (0.12 g, 0.94 mmol), 2,2-dimethyl propionic anhydride (0.95 mL, 4.69 mmol) and reaction was stirred for 24 hours. The reaction mixture was partitioned between ethyl acetate and a saturated aqueous ammonium chloride solution. Organic layer was washed with brine, dried over $MgSO_4$, and concentrated yielding 2,2-dimethyl-propionic acid 2S-(3-carbamoyl-[1,2,4]triazol-1-yl)-5,5,7,7-tetraisopropyl-tetrahydro-3aS,9aS-1,4,6,8-tetraoxa-5,7-disila-cyclopentacycloocten-3S-yl ester as a clear oil (99%).

To a stirred solution of 2,2-dimethyl-propionic acid 2S-(3-carbamoyl-[1,2,4]triazol-1-yl)-5,5,7,7-tetraisopropyl-tetrahydro-3aS,9aS-1,4,6,8-tetraoxa-5,7-disila-cyclopentacycloocten-3S-yl ester (0.48 g, 0.92 mmol) in 5 mL of acetonitrile were added 2.5 mL of 1 N H_2SO_4 at room temperature. After 2 hours, 30 mL of a saturated aqueous $NaHCO_3$ solution were added and product was extracted with ethyl acetate. The organic layer was washed with brine, dried over $MgSO_4$, and concentrated to give 2,2-dimethyl-propionic acid 2S-(3-carbamoyl-[1,2,4]triazol-1-yl)-4S-(3-hydroxy-1,1,3,3-tetraisopropylidisiloxanyloxy)-5S-hydroxymethyl-tetrahydro-furan-3S-yl ester (71%).

To a stirred slurry of 2,2-dimethyl-propionic acid 2S-(3-carbamoyl-[1,2,4]triazol-1-yl)-4S-(3-hydroxy-1,1,3,3-tetraisopropylidisiloxanyloxy)-5S-hydroxymethyl-tetrahydro-furan-3S-yl ester (0.38 g, 0.65 mmol) in 2.6 mL of 1:1 DMF/pyridine were added DMAP (0.40 g, 33 mmol), 2,2-dimethyl propionic anhydride (0.33 mL, 1.63 mmol) and reaction was stirred for 24 hours. The reaction was partitioned between ethyl acetate and a saturated aqueous ammonium chloride. The organic layer was washed with brine, dried over $MgSO_4$, and concentrated and the resulting residue was purified via


chromatography (15% acetone/chloroform) yielding 2,2-dimethyl-propionic acid 2S-(3-carbamoyl-[1,2,4]triazol-1-yl)-4S-(3-hydroxy-1,1,3,3-tetraisopropyl-1,3-disiloxanyloxy)-5S-(2,2-dimethyl-propionyloxymethyl)-tetrahydro-furan-3S-yl ester (54%).

- 5 To a stirred solution of 2,2-dimethyl-propionic acid 2S-(3-carbamoyl-[1,2,4]triazol-1-yl)-4S-(3-hydroxy-1,1,3,3-tetraisopropyl-1,3-disiloxanyloxy)-5S-(2,2-dimethyl-propionyloxymethyl)-tetrahydro-furan-3S-yl ester (0.24 g, 0.35 mmol) in 5 mL of acetonitrile were added 2.5 mL of 1 N H₂SO₄ at room temperature. After 72 hours, 30 mL of a saturated aqueous NaHCO₃ solution were added and the product was extracted with ethyl acetate. The organic layer was washed with brine, dried over MgSO₄ and concentrated and the residue was purified via preparatory HPLC yielding 2,2-dimethyl-propionic acid 2S-(3-carbamoyl-[1,2,4]triazol-1-yl)-5S-(2,2-dimethyl-propionyloxymethyl)-4S-hydroxy-tetrahydro-furan-3S-yl ester (compound 29, 15%, (M+H)⁺ = 413).

15

Method D - Mixed Acyl Derivatives

EXAMPLE 16



20

2S-amino-3-methyl-butyric acid 5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-3S,4S-bis-isobutyryloxy-tetrahydro-furan-2S-ylmethyl ester; compound with toluene-4-sulfonic acid (8: R¹ = *i*-Pr, R² = CH(NH₂)CH(CH₃)CH₃)

- 25 To a stirred slurry of isobutyric acid 2S-(3-carbamoyl-[1,2,4]triazol-1-yl)-5S-hydroxymethyl-4S-isobutyryloxy-tetrahydro-furan-3S-yl ester (Example 13 *supra*, compound 17, 0.50 g, 1.29 mmol) in 6 mL of THF were added at room temperature 4S-isopropyl-2,5-dioxo-oxazolidine-3-carboxylic acid benzyl ester (0.43 g, 1.55 mmol) and

- 30 -


0.3 mL of TEA. The reaction was allowed to stir for 12 hr and was quenched with 100 mL of a saturated aqueous NaHCO_3 solution and extracted with three 100 mL portions of ethyl acetate. The combined extracts were washed with brine, dried over MgSO_4 , and concentrated. The residue was purified *via* chromatography (silica gel; 35% ethyl acetate/hexane) to yield 0.52 g (65%) 2S-benzyloxycarbonylamino-3-methyl-butylric acid 5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-3S,4S-bis-isobutyryloxy-tetrahydro-furan-2S-ylmethyl ester; $(\text{M}+\text{H})^+ = 484$.

To an argon sparged, stirred solution of 2S-benzyloxycarbonylamino-3-methyl-butylric acid 5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-3S,4S-bis-isobutyryloxy-tetrahydro-furan-2S-yl methyl ester (0.52 g, 0.84 mmol) in 10 mL of methanol were added *p*-toluenesulfonic acid (0.16 g, 0.84 mmol) and 0.15 g of 10% Pd/C. The reaction vessel was evacuated and purged three times with hydrogen gas (about 1 atm) and stirred for 3 hours. The slurry was then filtered through CELITE® and the resulting filtrate was concentrated, dissolved in a methanol/dichloromethane(1:10) solution and precipitated with hexane to yield 0.18 g (33%) 2S-amino-3-methyl-butylric acid 5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-3S,4S-bis-isobutyryloxy-tetrahydrofuran-2S-yl methyl ester compound with toluene-4-sulfonic acid as a yellow solid (compound 22; $(\text{M}+\text{Na})^+ = 506$; m.p. = 72.0-76.0 °C).

Proceeding as described above with propionic acid, 4S-propionyloxy-5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-2S-hydroxymethyl-tetrahydro-furan-3S-yl ester gave 2S-benzyloxycarbonylamino-3-methyl-butylric acid 5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-3S,4S-bispropionyloxy-tetrahydro-furan-2S-ylmethyl ester (77%; $(\text{M}+\text{H})^+ = 456$) which was deprotected to yield 2S-amino-3-methyl-butylric acid 5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-3S,4S-bis-propionyloxy-tetrahydro-furan-2S-ylmethyl ester; compound with toluene-4-sulfonic acid (compound 21; 46%).

Method E – Preparations of N-acyl analogs

Example 17



39

Butyric acid 5S-(3-butyrylcarbamoyl-[1,2,4]triazol-1-yl)-4S-butyryloxy-2S-butyryloxymethyl-tetrahydro-furan-3S-yl ester.

To a stirred slurry of levovirin (1, 0.48 g, 2.0 mmol) in 7 mL of THF were added TEA
 5 (1.65 mL, 11.8 mmol), n-butyric anhydride (1.97 mL, 10.8 mmol), and DMAP (0.24 g,
 2.0 mmol). Reaction vessel was heated at 60°C for 24 hours. The solution was
 partitioned between ethyl acetate and a saturated aqueous NaCO₃ solution and the
 organic layer was washed with brine, dried over MgSO₄, and concentrated. The residue
 was purified via chromatography (35% ethyl acetate/hexane) yielding butyric acid 5S-(3-
 10 butyrylcarbamoyl-[1,2,4]triazol-1-yl)-4S-butyryloxy-2S-butyryloxymethyl-tetrahydro-
 furan-3S-yl ester (compound 39, 71%, (M+Na)⁺ = 547).

Proceeding as described above were also prepared: propionic acid 5S-(3-
 propionylcarbamoyl-[1,2,4]triazol-1-yl)-4S-propionyloxy-2S-propionyloxymethyl-
 15 tetrahydro-furan-3S-yl ester (compound 37, 95%, (M+H)⁺ = 469).

Proceeding as described above utilizing the appropriately protected intermediates were
 prepared: butyric acid 5S-(3-butyrylcarbamoyl-[1,2,4]triazol-1-yl)-4S-butyryloxy-2S-
 hydroxymethyl-tetrahydro-furan-3S-yl ester (compound 38, 35%, (M+Na)⁺ = 477) and
 20 1-(3S,4R-Dihydroxy-5S-hydroxymethyl-tetrahydro-furan-2S-yl)-1H-[1,2,4]triazole-3-
 carboxylic acid butyryl-amide (compound 40, 58%, (M+H)⁺ = 315).

EXAMPLE 18

Caco Assay

For general discussions of the Caco Assay see: S. Yee, "In Vitro Permeability Across Caco-2
 Cells (Colonic) Can Predict In Vivo (Small Intestinal) Absorption in Man- Fact or Myth"
 Pharm. Res. 14(6):763-766 (1997) and Yamashita et. al., "Analysis of Drug Permeation
 Across Caco-2 Monolayer: Implication for Predicting In Vivo Drug Absorption" Pharm.
 30 Res. 14(4):486-491(1997). For specific technical aspects see: Grass, G.M. and Sweetana,
 S.A. "In Vitro Measurement of Gastrointestinal Tissue Permeability Using a New Diffusion
 Cell" Pharm. Res. 5(6):372-376 (1988); Rubas et. al., "Comparison of the Permeability
 Characteristics of a Human Colonic Epithelial (Caco-2) Cell Line to Colon of Rabbit,
 Monkey, and Dog Intestine and Human Drug Absorption" Pharm. Res. 10(1):113-117
 35 (1993).

Incubation medium and culture conditions:

High passage (108-120) Caco-2 cells are cultured in Dulbecco's Modified Eagle Media with high Glucose and L-Glutamine (DMEM) (Gibco/Life Technologies, Cat # 11965-084) supplemented with 10% Fetal Bovine Serum, 1X L-Glutamine (Gibco/Life Technologies, Cat # 25030-081) 1X Penicillin-streptomycin (Gibco/Life Technologies, Cat # 15140-122) 1X Non-essential Amino Acids, (Gibco/Life Technologies, Cat # 11140-019). Cells are maintained in T225 cm² Cell Culture Flask Tissue Culture Treated (Costar, Cat # 3001) at 37 °C and 5% CO₂. For transport experiments, cells are plated at 7.1×10^4 cells/well into 12-well collagen-coated PTFE membrane polystyrene plates with inserts (Costar # 3493, 12 mm diameter, 0.4 µm pore size, sterile, tissue culture treated). Cells are fed every 3 days and maintained at 37 °C and 5% CO₂ for 21 days to allow complete formation of a polarized monolayer with tight junctions.

Stock and working solutions:

Kreb's-Henseleit bicarbonate buffers, pH 6.5 and 7.4

Reagents:

Distilled Water (glass distilled or Nanopure)

Kreb's-Henseleit bicarbonate buffer mix (powder, SIGMA # K-3753)

Calcium chloride dihydrate (MW = 147.0)

Sodium bicarbonate (MW = 84.01)

Dissolve Kreb's-Henseleit bicarbonate buffer mix in about 900 mL of water. When buffer mix is dissolved, add 0.373 gm calcium chloride dihydrate. After calcium chloride dihydrate dissolves, add 2.1 gm sodium bicarbonate, after sodium bicarbonate dissolves, add water QS to 1000 mL, then sterile filter through 0.2 µm filter and store in a refrigerator

Test/standard compound solutions:

Prepare 5 mg/ml stock solution of test compound in DMSO and store at 4 °C. Dilute desired amount of 5 mg/ml stock to 10 mL with pH 6.5 Kreb's Henseleit bicarbonate buffer to give a concentration of 100 µM. Then 1 mL of 100 µM solution was further diluted to 5 mL to make the concentration of 20 µM. This 20 µM test solution was used as initial donor dosing solution (D0). Warm drug solution(s) to 37 °C before use.

Assay Procedure

1. Prewarm buffer, working solutions, and three 12-well plates containing buffer for each plate of 12 inserts. Using a millicell[®]-ERS equipped with "chopstick" electrodes (Millipore, Bedford, MA) check the TEER. This procedure should be done when the cells are at approximately 37°C, since TEER is effected by temperature. Use only those inserts that have TEER above 300 ohms.
2. Decant media and wash each insert once with warm Kreb's-Henseleit bicarbonate buffer.
3. 0.5 mL pH 6.5 Kreb's-Henseleit buffer was added to apical side of the cell monolayers and 1.25 mL pH 7.4 Kreb's-Henseleit buffer to the basolateral chamber. The cells were equilibrated in 37 °C and 5% CO₂ incubator for at least 30 minutes.
4. The apical side buffer was removed and replaced with 0.5 mL 20 µM test solutions.
5. The cells were then incubated at 37 °C and 5% CO₂.
6. At 30, 60 and 90 minutes time points, the inserts were transferred to new plates which receiver sides contained 1.25 mL warm fresh pH 7.4 Kreb's-Henseleit buffer.
7. The media from all plates were collected as receiver samples.
8. After 60 min transport studies, Lucifer Yellow (0.05mL x 1000µM) was added to the apical side of the wells. At the end of the transport studies (90 minutes), the fluorescence of the receiver side samples was measured.
- 20 Sample solutions from the donor side were collected at the end of the experiments as D90 samples.

The dC/dt of test substance was calculated from sampling data at 30 (assume 0 ng/mL) and 60 minutes. The apparent permeability coefficient (P_{app}) was calculated from the following equation,

$$P_{app} = \frac{dQ}{dt} \times \frac{1}{A \times C_o} = \frac{dC}{dt} \times \frac{V}{A \times C_o}$$

where dQ is the change in amount of compound in receiver, dC is the change in the concentration of compound in receiver, V is the volume (cm³) of the receiver solution, A is the surface area (cm²) of the insert, C_o is the 'initial' concentration of drug substance, and dC/dt is the change in drug substance concentration in the receiver solution over the 90 minute incubation time, i.e., the slope (µg/cm³/sec) of the drug substance concentration in the receiver solution vs. time.

Table 4. Caco-2 cell assay permeability of selected compounds

Compound number	Caco-2 permeability (x 10 ⁻⁶ cm/sec)
2	7.8
3	8.6
5	5.6
7	1.2
9	0.9
10	1.5
17	4.8
18	3.4
28	26.6
34	6.5

EXAMPLE 19

5

Composition for Oral Administration

Ingredient	% wt./wt.
Active ingredient	20.0%
Lactose	79.5%
Magnesium stearate	0.5%

The ingredients are mixed and dispensed into capsules containing about 100 mg each; one capsule would approximate a total daily dosage.

EXAMPLE 20

10

Composition for Oral Administration

Ingredient	% wt./wt.
Active ingredient	20.0%
Magnesium stearate	0.5%
Crosscarmellose sodium	2.0%
Lactose	76.5%
PVP (polyvinylpyrrolidone)	1.0%

The ingredients are combined and granulated using a solvent such as methanol. The formulation is then dried and formed into tablets (containing about 20 mg of active compound) with an appropriate tablet machine.

EXAMPLE 21

5

Composition for Oral Administration

Ingredient	Amount
Active compound	1.0 g
Fumaric acid	0.5 g
Sodium chloride	2.0 g
Methyl paraben	0.15 g
Propyl paraben	0.05 g
Granulated sugar	25.5 g
Sorbitol (70% solution)	12.85 g
Veegum K (Vanderbilt Co.)	1.0 g
Flavoring	0.035 mL
Colorings	0.5 mg
Distilled water	q.s. to 100 mL

The ingredients are mixed to form a suspension for oral administration.

10 The features disclosed in the foregoing description, or the following claims, or the accompanying drawings, expressed in their specific forms or in terms of a means for performing the disclosed function, or a method or process for attaining the disclosed result, as appropriate, may, separately, or in any combination of such features, be utilized for realizing the invention in diverse forms thereof.


15 The foregoing invention has been described in some detail by way of illustration and example, for purposes of clarity and understanding. It will be obvious to one of skill in the art that changes and modifications may be practiced within the scope of the appended claims. Therefore, it is to be understood that the above description is intended to be illustrative and not restrictive. The scope of the invention should, therefore, be
20 determined not with reference to the above description, but should instead be determined with reference to the following appended claims, along with the full scope of equivalents to which such claims are entitled.

All patents, patent applications and publications cited in this application are hereby
25 incorporated by reference in their entirety for all purposes to the same extent as if each

individual patent, patent application or publication were so individually denoted.


Claims

1. A compound according to formula I



- 5 wherein R^1 , R^2 and R^3 are independently selected from the group consisting of hydrogen, C_{1-10} acyl, C_{1-10} alkoxy carbonyl and COR^4 where COR^4 is an amino acid or a dipeptide; and, hydrates, solvates, clathrates of said compound and acid addition salts.
- 10 2. A compound according to claim 1 wherein one of R^1 , R^2 and R^3 is COR^4 , R^4 is $CH(R^5)NH_3^+ Cl^-$ or pyrrolidin-2-yl, R^5 is the side chain of a naturally occurring hydrophobic amino acid or C_{1-6} straight or branched alkyl and the other of R^1 , R^2 and R^3 are independently selected from the group consisting of hydrogen, C_{1-10} acyl, and C_{1-10} alkoxy carbonyl.
- 15 3. A compound according to claims 1 or 2 wherein R^1 is COR^4 , and R^4 is $CH(R^5)NH_3^+ Cl^-$ or pyrrolidin-2-yl, and R^2 and R^3 are independently selected from the group consisting of hydrogen, C_{1-10} acyl, and C_{1-10} alkoxy carbonyl.
4. A compound according to claims 1 to 3 wherein R^1 is COR^4 , and R^4 is $CH(R^5)NH_3^+ Cl^-$, R^5 is selected from the group consisting of $CH(CH_3)_2$ and $CH(CH_3)CH_2CH_3$, and both R^2 and R^3 are hydrogen.
- 20 5. A compound according to claims 1 to 4 wherein R^1 is COR^4 , and R^4 is $CH(R^5)NH_3^+ Cl^-$, R^5 is $CH(CH_3)_2$, and both R^2 and R^3 are hydrogen.
6. A compound according to claims 1 to 5, wherein the compound is 2S-amino-3-methyl-butiric acid 5S-(3-carbamoyl-{1,2,4}triazol-1-yl)-3R,4S-dihydroxy-tetrahydro-furan-2S-ylmethyl ester hydrochloride.

7. A compound according to claim 1 wherein R^1 , R^2 and R^3 are independently C_{1-10} acyl or C_{1-10} alkoxy carbonyl.
8. A compound according to claims 1 or 7 wherein the compound is: propionic acid
 5 3S,4S-bis-propionyloxy-5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-tetrahydro-furan-2S-ylmethyl ester.
9. A compound according to claims 1 or 7 wherein R^1 is C_{1-10} acyl or C_{1-10} alkoxy carbonyl and both R^2 and R^3 are hydrogen.
10. A compound according to claim 1 or 7 wherein R^1 is hydrogen and both R^2 and R^3
 10 independently are C_{1-10} acyl or C_{1-10} alkoxy carbonyl.
11. A compound according to claims 1 or 7 wherein the compound is: isobutyric acid
 2S-(3-carbamoyl-[1,2,4]triazol-1-yl)-5S-hydroxymethyl-4S-isobutyryloxy-
 tetrahydro-furan-3S-yl ester; or, 2,2-dimethylpropionic acid 4S-(2,2-
 dimethylpropionyloxy)-5S-(3-carbamoyl-[1,2,4]triazol-1-yl)-2S-hydroxymethyl-
 15 tetrahydro-furan-3S-yl ester.
12. A compound according to formula II



wherein R^1 , R^2 and R^3 are independently selected from the group consisting of
 20 hydrogen, C_{1-10} acyl, C_{1-10} alkoxy carbonyl and COR^4 where COR^4 is an amino acid
 or a dipeptide; R^6 is C_{1-6} acyl; and,

hydrates, solvates, clathrates of said compound and acid addition salts.

13. Compounds according to claims 1 to 12 for use in therapy.
14. Compounds according to claims 1 to 12 for use in the therapy of a disease
 mediated by the Hepatitis C virus.

15. Use of the compounds according to claims 1 to 12 for the manufacture of a medicament for use in therapy..
16. Use of the compounds according to claims 1 to 12 for the manufacture of a medicament for use in the therapy of a disease mediated by the Hepatitis C virus.
- 5 17. A pharmaceutical composition comprising a therapeutically effective amount of a compound according to claims 1 to 12 and at least one pharmaceutically acceptable carrier and optionally containing excipients.
