Patent Claims

1. 3-(1-{3-[5-(1-methyl-piperidin-4-ylmethoxy)-pyrimidin-2-yl]-benzyl}-6-oxo-1,6-dihydro-pyridazin-3-yl)-benzonitrile or a pharmaceutically acceptable salt and/or solvate thereof for the use for the treatment of hepatocellular carcinoma (HCC).

2. 3-(1-{3-[5-(1-methyl-piperidin-4-ylmethoxy)-pyrimidin-2-yl]-benzyl}-6-oxo-1,6-dihydro-pyridazin-3-yl)-benzonitrile hydrochloride hydrate for the use for the treatment of hepatocellular carcinoma (HCC).

3. 3-(1-{3-[5-(1-methyl-piperidin-4-ylmethoxy)-pyrimidin-2-yl]-benzyl}-6-oxo-1,6-dihydro-pyridazin-3-yl)-benzonitrile according to claim 1 or 2, wherein the compound is administered to a patient in an amount of 100 mg to 800 mg per day.

4. 3-(1-{3-[5-(1-methyl-piperidin-4-ylmethoxy)-pyrimidin-2-yl]-benzyl}-6-oxo-1,6-dihydro-pyridazin-3-yl)-benzonitrile according to claim 1, 2 or 3, wherein the compound is administered orally.

5. Use of 3-(1-{3-[5-(1-methyl-piperidin-4-ylmethoxy)-pyrimidin-2-yl]-benzyl}-6-oxo-1,6-dihydro-pyridazin-3-yl)-benzonitrile or a pharmaceutically acceptable salt and/or solvate thereof for the manufacture of a medicament for the treatment of hepatocellular carcinoma (HCC).

6. Use of 3-(1-{3-[5-(1-methyl-piperidin-4-ylmethoxy)-pyrimidin-2-yl]-benzyl}-6-oxo-1,6-dihydro-pyridazin-3-yl)-benzonitrile hydrochloride hydrate for the manufacture of a medicament for the treatment of hepatocellular carcinoma (HCC).

7. Use according to claim 5 or 6, wherein the compound is administered to a patient in an amount of 100 mg to 800 mg per day.

8. Use according to claim 5, 6 or 7, wherein the compound is administered orally.

Patent Claims

1. Use of 3-(1-{3-[5-(1-methyl-piperidin-4-ylmethoxy)-pyrimidin-2-y1]-benzy1}-6-oxo-1,6-dihydro-pyridazin-3-yl)-benzonitrile or a pharmaceutically acceptable salt and/or solvate thereof for the manufacture of a medicament for the treatment of hepatocellular carcinoma (HCC).

2. Use of 3-(1-{3-[5-(1-methyl-piperidin-4-ylmethoxy)-pyrimidin-2-yl]-benzyl}-6-oxo-1,6-dihydro-pyridazin-3-yl)-benzonitrile hydrochloride hydrate for the manufacture of a medicament for the treatment of hepatocellular carcinoma (HCC).

3. Use according to claim 1 or 2, wherein the compound is administered to a patient in an amount of 100 mg to 800 mg per day.

4. Use according to claim 1, 2 or 3, wherein the compound is administered orally.