

Treatment of intrahepatic cholestatic diseases

Technical field

[0001] This invention relates to the treatment of intrahepatic cholestatic diseases.

Background art

5 [0002] Intrahepatic cholestatic diseases

[0003] Cholestasis is a condition in which the flow of bile from the liver to the duodenum is slowed or blocked. Cholestasis may be divided conveniently into two types: intrahepatic cholestasis, inside the liver, where bile formation is disturbed by conditions such as various diseases, extended intravenous nutrition, or as a side effect of certain drugs (such as some antibiotics); and extrahepatic cholestasis, occurring outside the liver, typically where the flow of bile is obstructed by a mechanical partial or complete closure of the bile duct, such as by bile duct tumors, cysts, bile duct stones, strictures, or pressure on the bile duct; though primary sclerosing cholangitis (PSC) may be intrahepatic or extrahepatic. Common symptoms of cholestasis include fatigue, pruritus (itching), jaundice, and xanthoma (deposits of cholesterol-rich material under the skin). The effects of cholestasis are profound and widespread, leading to worsening liver disease with systemic illness, liver failure, and the need for liver transplantation.

[0004] Intrahepatic cholestatic diseases, in decreasing order of frequency, include primary biliary cirrhosis (PBC), primary sclerosing cholangitis (PSC), progressive familial intrahepatic cholestasis (PFIC), and Alagille syndrome (AS).

[0005] PBC is an autoimmune disease of the liver marked by the slow progressive destruction of the small bile ducts of the liver, with the intralobular ducts affected early in the disease. When these ducts are damaged, bile builds up in the liver (cholestasis) and over time damages the tissue, which can lead to scarring, fibrosis and cirrhosis. Recent studies have shown that it may affect up to 1 in 3,000 - 4,000 people, with a sex ratio at least 9:1 female to male. There is no cure for PBC, and liver transplantation often becomes necessary; but medication such as ursodeoxycholic acid (UDCA, ursodiol) to reduce cholestasis and improve liver function, cholestyramine to absorb bile acids, modafinil for fatigue, fat-soluble vitamins (vitamins A, D, E, and K, since reduced bile flow makes it difficult for these vitamins to be absorbed) may slow the progression to allow a normal lifespan and quality of life. UDCA is the only drug approved in the United States to treat PBC. Japanese researchers have reported that the addition of

bezafibrate, a peroxisome proliferator-activated receptor- α (PPAR α) and pregnane X receptor agonist, to UDCA is helpful in treating patients who are refractory to UDCA monotherapy, improving serum biliary enzymes, cholesterol (C), and triglycerides (TGs).

[0006] PSC is a chronic cholestatic liver disease characterized by intra- or extrahepatic biliary duct inflammation and fibrosis, eventually leading to cirrhosis. The underlying cause of the inflammation is believed to be autoimmunity; and about three-fourths of patients with PSC have inflammatory bowel disease (usually ulcerative colitis), though this is reported to vary by country, as is the prevalence (generally reported at about 1 in 10,000) and sex ratio (generally reported as predominately male). Standard treatment includes UDCA, which has been shown to lower elevated liver enzyme numbers in people with PSC, but has not improved liver- or overall survival; and also includes antipruritics, cholestyramine, fat-soluble vitamins, and antibiotics to treat infections (bacterial cholangitis). In a study reported in 2009, long-term high-dose UDCA therapy was associated with improvement in serum liver tests in PSC but did not improve survival and was associated with higher rates of serious adverse events. Liver transplantation is the only proven long-term treatment.

[0007] PFIC refers to a group of three types of autosomal recessive disorders of childhood associated with intrahepatic cholestasis: deficiency of familial intrahepatic cholestasis 1 (PFIC-1), deficiency of bile salt export pump (PFIC-2), and deficiency of multidrug resistance protein 3 (PFIC-3). They have a combined incidence of 1 in 50,000 - 100,000. The onset of the disease is usually before age 2, with PFIC-3 usually appearing earliest, but patients have been diagnosed with PFIC even into adolescence. Patients usually show cholestasis, jaundice, and failure to thrive; and intense pruritus is characteristic. Fat malabsorption and fat soluble vitamin deficiency may appear. Biochemical markers include a normal γ -glutamyl transpeptidase (GGT) in PFIC-1 and -2, but a markedly elevated GGT in PFIC-3; while serum bile acid levels are greatly elevated; though serum cholesterol levels are typically not elevated, as is seen usually in cholestasis, because the disease is due to a transporter as opposed to an anatomical problem with biliary cells. The disease is typically progressive without liver transplantation, leading to liver failure and death in childhood; and hepatocellular carcinoma may develop in PFIC-2 at a very early age. Medication with UDCA is common; supplemented by fat-soluble vitamins, cholestyramine, and pancreatic enzymes in PFIC-1.

[0008] AS, also known as Alagille-Watson syndrome, syndromic bile duct paucity, and arteriohepatic dysplasia, is an autosomal dominant disorder associated with liver, heart, eye and skeletal abnormalities, as well as characteristic facial features; with an incidence of about 1 in

100,000. The liver abnormalities are narrowed and malformed bile ducts within the liver; and these result in obstruction of bile flow, causing cirrhosis (scarring) of the liver. AS is predominately caused by changes in the *Jagged1* gene, located on chromosome 20. In 3 –5 % of cases, the entire gene is deleted (missing) from one copy of chromosome 20; in the remainder, there are changes or mutations in the *Jagged1* DNA sequence. In a very small number of cases, less than 1 percent, changes in another gene, *Notch2*, result in AS. In about one-third of the cases, the mutation is inherited; in about two-thirds, the mutation is new in that case. There is no cure for AS, though the severity of liver disease typically peaks by 3 to 5 years of age and often resolves by 7 to 8 years of age. In some people, the hepatic disease will progress to end-stage liver disease and may require liver transplantation (approximately 15 % of patients with AS require liver transplantation). A number of different medications (e.g. ursodeoxycholic acid (UDCA, ursodiol)) have been used to improve bile flow and reduce itching, and many patients are given high doses of fat-soluble vitamins.

[0009] Alkaline phosphatase (ALP) and GGT are key markers of cholestasis. While an elevation of one of them alone does not indicate cholestasis, and other parameters would be needed for confirmation, elevation in both ALP and GGT is indicative of cholestasis; and a decrease in both indicates improvement of cholestasis. Thus ALP and GGT levels serve as biochemical markers for the presence of biliary pathophysiology present in intrahepatic cholestatic diseases, and ALP level has been used as a primary outcome marker in clinical studies of intrahepatic diseases such as PBC.

[0010] Treatments for intrahepatic cholestatic diseases

[0011] As mentioned above, UDCA is a common treatment for intrahepatic cholestatic diseases, because of its action in reducing cholestasis and improving liver function. However, a Cochrane Review of UDCA in PBC in 2012 found that, although UDCA showed a reduction in biomarkers of liver pathology, jaundice, and ascites, there was no evidence in the medical literature for any benefit of UDCA on mortality or liver transplantation, while its use was associated with weight gain and costs.


[0012] Obeticholic acid (6 α -ethylchenodeoxycholic acid), a semi-synthetic bile acid analog that is a highly potent farnesoid X receptor agonist, has completed a Phase 2 study in combination with ursodeoxycholic acid for PBC, and is under study in a second Phase 2 study alone on lipoprotein metabolism in PBC and a Phase 3 placebo-controlled study in PBC.

However, the only long-term treatment for many patients with intrahepatic cholestatic diseases is liver transplantation.

[0013] It would be desirable to develop pharmacological treatments for intrahepatic cholestatic diseases.

5 [0014] MBX-8025

[0015] MBX-8025 is the compound of the formula



MBX-8025 has the chemical name (*R*)-2-(4-((2-ethoxy-3-(4-(trifluoromethyl)phenoxy)propyl)-

thio)-2-methylphenoxy)acetic acid [IUPAC name as generated by CHEMDRAW ULTRA

10 12.0]. MBX-8025 and its synthesis, formulation, and use is disclosed in, for example, US Patent No. 7301050 (compound 15 in Table 1, Example M, claim 49), US Patent No. 7635718 (compound 15 in Table 1, Example M), and US Patent No. 8106095 (compound 15 in Table 1, Example M, claim 14). Lysine (L-lysine) salts of MBX-8025 and related compounds are disclosed in US Patent No. 7709682 (MBX-8025 L-lysine salt throughout the Examples, 15 crystalline forms claimed).

[0016] MBX-8025 is an orally active, potent (2 nM) agonist of peroxisome proliferator-activated receptor- δ (PPAR δ). It is specific (>600-fold and >2500-fold compared with PPAR α and peroxisome proliferator-activated receptor- γ receptors). PPAR δ activation stimulates fatty acid oxidation and utilization, improves plasma lipid and lipoprotein metabolism, glucose 20 utilization, and mitochondrial respiration, and preserves stem cell homeostasis. According to US Patent No. 7301050, PPAR δ agonists, such as MBX-8025, are suggested to treat PPAR δ -mediated conditions, including “diabetes, cardiovascular diseases, Metabolic X syndrome, hypercholesterolemia, hypo-high density lipoprotein (HDL)-cholesterolemia, hyper-low density protein (LDL)-cholesterolemia, dyslipidemia, atherosclerosis, and obesity”, with dyslipidemia 25 said to include hypertriglyceridemia and mixed hyperlipidemia.

[0017] A Phase 2 study of MBX-8025 L-lysine dihydrate salt in mixed dyslipidemia (6 groups, 30 subjects/group: once daily placebo, atorvastatin (ATV) 20 mg, or MBX-8025 L-lysine dihydrate salt at 50 or 100 mg (calculated as the free acid) capsules alone or combined

with ATV 20 mg, for 8 weeks) has been reported by Bays et al., “MBX-8025, A Novel Peroxisome Proliferator Receptor- δ Agonist: Lipid and Other Metabolic Effects in Dyslipidemic Overweight Patients Treated with and without Atorvastatin”,

J. Clin. Endocrin. Metab., **96**(9), 2889-2897 (2011) and Choi et al., “Effects of the PPAR- δ

agonist MBX-8025 on atherogenic dyslipidemia”, *Atherosclerosis*, **220**, 470-476 (2012).

Compared to placebo, MBX-8025 alone and in combination with ATV significantly ($P < 0.05$) reduced apolipoprotein B-100 by 20-38%, LDL by 18-43%, triglycerides (TGs) by 26-30%, non-HDL-C by 18-41%, free fatty acids by 16-28%, and high-sensitivity C-reactive protein by 43-72%; it raised HDL-C by 1-12% and also reduced the number of patients with the metabolic syndrome and a preponderance of small LDL particles. MBX-8025 reduced small/very small LDL particles by 40-48% compared with a 25% decrease with ATV; and MBX-8025 increased large LDL particles by 34-44% compared with a 30% decrease with ATV. MBX-8025 significantly reduced ALP by 32-43%, compared to reductions of only 4% in the control group and 6% in the ATV group; and significantly reduced GGT by 24-28%, compared to a reduction of only 3% in the control group and an increase of 2% in the ATV group. Thus MBX-8025 corrects all three lipid abnormalities in mixed dyslipidemia – lowers TGs and LDL and raises HDL, selectively depletes small dense LDL particles (92%), reduces cardiovascular inflammation, and improves other metabolic parameters including reducing serum aminotransferases, increases insulin sensitivity (lowers homeostatic model assessment-insulin resistance, fasting plasma glucose, and insulin), lowers GGT and ALP, significantly (>2-fold) reduces the percentage of subjects meeting the criteria for metabolic syndrome, and trends towards a decrease in waist circumference and increase in lean body mass. MBX-8025 was safe and generally well-tolerated, and also reduced liver enzyme levels. As explained in US Patent Application Publication No. 2010-0152295, MBX-8025 converts LDL particle size pattern I (a predominant LDL particle size of from 25.75 nm to 26.34 nm) to pattern A (a predominant LDL particle size of greater than 26.34 nm); and from pattern B (a predominant LDL particle size of less than 25.75 nm) to pattern I or A, where the LDL particle size is measured by gradient-gel electrophoresis.

Summary of the invention

[0018] This invention is the treatment of intrahepatic cholestatic diseases, comprising therapy with MBX-8025 or an MBX-8025 salt.

[0019] Because MBX-8025 lowers alkaline phosphatase and γ -glutamyl transpeptidase, which are elevated in intrahepatic cholestatic diseases, its use will result in a reduction in cholestasis and provide an effective treatment for these diseases (other drugs, such as the fibrates, which also lower ALP and GGT in dyslipidemic patients are known to reduce cholestasis in intrahepatic cholestatic diseases).

[0020] In various aspects, this invention is:

MBX-8025 or an MBX-8025 salt for use in treating an intrahepatic cholestatic disease; the use of MBX-8025 or an MBX-8025 salt for treating an intrahepatic cholestatic disease, or in the manufacture of a medicament for treating an intrahepatic cholestatic disease; pharmaceutical compositions comprising MBX-8025 or an MBX-8025 salt for treating an intrahepatic cholestatic disease; kits comprising compositions comprising MBX-8025 or an MBX-8025 salt for treating an intrahepatic cholestatic disease; and methods of treating an intrahepatic cholestatic disease by administering MBX-8025 or an MBX-8025 salt.

[0021] Preferred embodiments of this invention are characterized by the specification and by the features of Claims 1 to 14 of this application as filed.

Detailed description of the invention

[0022] Definitions

[0023] “Intrahepatic cholestatic diseases” and their treatment are described in paragraphs [0002] through [0012].

[0024] A “therapeutically effective amount” of MBX-8025 or an MBX-8025 salt means that amount which, when administered to a human for treating an intrahepatic cholestatic disease, is sufficient to effect treatment for the intrahepatic cholestatic disease. “Treating” or “treatment” of an intrahepatic cholestatic disease in a human includes one or more of:

(1) preventing or reducing the risk of developing an intrahepatic cholestatic disease, i.e., causing the clinical symptoms of an intrahepatic cholestatic disease not to develop in a subject who may be predisposed to an intrahepatic cholestatic disease but who does not yet experience or display symptoms of the intrahepatic cholestatic disease (i.e. prophylaxis);

(2) inhibiting an intrahepatic cholestatic disease, i.e., arresting or reducing the development of the intrahepatic cholestatic disease or its clinical symptoms; and

(3) relieving an intrahepatic cholestatic disease, i.e., causing regression, reversal, or amelioration of the intrahepatic cholestatic disease or reducing the number, frequency, duration or severity of its clinical symptoms.

The therapeutically effective amount for a particular subject varies depending upon the age, health and physical condition of the subject to be treated, the intrahepatic cholestatic disease and its extent, the assessment of the medical situation, and other relevant factors. It is expected that the therapeutically effective amount will fall in a relatively broad range that can be determined through routine trial.

[0025] “MBX-8025” is described in paragraphs [0014] through [0017].

[0026] Salts (for example, pharmaceutically acceptable salts) of MBX-8025 are included in this invention and are useful in the compositions, methods, and uses described in this application. These salts are preferably formed with pharmaceutically acceptable acids. See, for example, “Handbook of Pharmaceutically Acceptable Salts”, Stahl and Wermuth, eds., Verlag Helvetica Chimica Acta, Zürich, Switzerland, for an extensive discussion of pharmaceutical salts, their selection, preparation, and use. Unless the context requires otherwise, reference to MBX-8025 is a reference both to the compound and to its salts.

[0027] Because MBX-8025 contains a carboxyl group, it may form salts when the acidic proton present reacts with inorganic or organic bases. Typically the MBX-8025 is treated with an excess of an alkaline reagent, such as hydroxide, carbonate or alkoxide, containing an appropriate cation. Cations such as Na^+ , K^+ , Ca^{2+} , Mg^{2+} , and NH_4^+ are examples of cations present in pharmaceutically acceptable salts. Suitable inorganic bases, therefore, include calcium hydroxide, potassium hydroxide, sodium carbonate and sodium hydroxide. Salts may also be prepared using organic bases, such as salts of primary, secondary and tertiary amines, substituted amines including naturally-occurring substituted amines, and cyclic amines including isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, ethanolamine, 2-dimethylaminoethanol, tromethamine, lysine, arginine, histidine, caffeine, procaine, hydrabamine, choline, betaine, ethylenediamine, glucosamine, *N*-alkylglucamines, theobromine, purines, piperazine, piperidine, *N*-ethylpiperidine, and the like. As noted in paragraph [0017], MBX-8025 is currently formulated as its *L*-lysine dihydrate salt; and MBX-8025 has also been studied in clinical trials as its calcium salt.

[0028] “Comprising” or “containing” and their grammatical variants are words of inclusion and not of limitation and mean to specify the presence of stated components, groups, steps, and

the like but not to exclude the presence or addition of other components, groups, steps, and the like. Thus “comprising” does not mean “consisting of”, “consisting substantially of”, or “consisting only of”; and, for example, a formulation “comprising” a compound must contain that compound but also may contain other active ingredients and/or excipients.

5 [0029] Formulation and administration

[0030] The MBX-8025 may be administered by any route suitable to the subject being treated and the nature of the subject’s condition. Routes of administration include administration by injection, including intravenous, intraperitoneal, intramuscular, and subcutaneous injection, by transmucosal or transdermal delivery, through topical applications, nasal spray, suppository and
10 the like or may be administered orally. Formulations may optionally be liposomal formulations, emulsions, formulations designed to administer the drug across mucosal membranes or transdermal formulations. Suitable formulations for each of these methods of administration may be found, for example, in “Remington: The Science and Practice of Pharmacy”, 20th ed., Gennaro, ed., Lippincott Williams & Wilkins, Philadelphia, Pa., U.S.A. Because MBX-8025 is
15 orally available, typical formulations will be oral, and typical dosage forms will be tablets or capsules for oral administration. As mentioned in paragraph [0017], MBX-8025 has been formulated in capsules for clinical trials.

[0031] Depending on the intended mode of administration, the pharmaceutical compositions may be in the form of solid, semi-solid or liquid dosage forms, preferably in unit dosage form
20 suitable for single administration of a precise dosage. In addition to an effective amount of the MBX-8025, the compositions may contain suitable pharmaceutically-acceptable excipients, including adjuvants which facilitate processing of the active compounds into preparations which can be used pharmaceutically. “Pharmaceutically acceptable excipient” refers to an excipient or mixture of excipients which does not interfere with the effectiveness of the
25 biological activity of the active compound(s) and which is not toxic or otherwise undesirable to the subject to which it is administered.

[0032] For solid compositions, conventional excipients include, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharin, talc, cellulose, glucose, sucrose, magnesium carbonate, and the like. Liquid pharmacologically administrable
30 compositions can, for example, be prepared by dissolving, dispersing, etc., an active compound as described herein and optional pharmaceutical adjuvants in water or an aqueous excipient, such as, for example, water, saline, aqueous dextrose, and the like, to form a solution or

suspension. If desired, the pharmaceutical composition to be administered may also contain minor amounts of nontoxic auxiliary excipients such as wetting or emulsifying agents, pH buffering agents and the like, for example, sodium acetate, sorbitan monolaurate, triethanolamine sodium acetate, triethanolamine oleate, etc.

5 [0033] For oral administration, the composition will generally take the form of a tablet or capsule; or, especially for pediatric use, it may be an aqueous or nonaqueous solution, suspension or syrup. Tablets and capsules are preferred oral administration forms. Tablets and capsules for oral use will generally include one or more commonly used excipients such as lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically
10 added. When liquid suspensions are used, the active agent may be combined with emulsifying and suspending excipients. If desired, flavoring, coloring and/or sweetening agents may be added as well. Other optional excipients for incorporation into an oral formulation include preservatives, suspending agents, thickening agents, and the like.

[0034] Typically, a pharmaceutical composition of MBX-8025, or a kit comprising
15 compositions of MBX-8025, is packaged in a container with a label, or instructions, or both, indicating use of the pharmaceutical composition or kit in the treatment of an intrahepatic cholestatic disease.

[0035] A suitable amount of MBX-8025 (calculated as the free acid) for oral dosing will be 20-200 mg/day, preferably 50-100 mg/day, for an adult subject with an intrahepatic cholestatic
20 disease, depending on the disease and stage of disease and factors such as hepatic and renal function. That is, a suitable amount of MBX-8025 for oral dosing for adults in diseases such as PSC and PBC will be similar to the amounts employed in clinical trials. Suitable reductions in dose toward the lower end of the outer range above will be made for subjects who are children in diseases such as AS and PFIC, depending on such additional factors as age and body mass.

25 [0036] A person of ordinary skill in the art of the treatment of intrahepatic cholestatic disease will be able to ascertain a therapeutically effective amount of the MBX-8025 or an MBX-8025 salt for a particular disease, stage of disease, and patient to achieve a therapeutically effective amount without undue experimentation and in reliance upon personal knowledge and the disclosure of this application.

30 [0037] Example

[0038] Adult subjects with an intrahepatic cholestatic disease such as PBC are treated with doses of 50, 100, 150, or 200 mg/day of MBX-8025. Subjects are permitted their usual other

medications, including UDCA. The subjects are assessed before the study, and at intervals during the study, such as every 4 weeks during the study and 4 weeks after the last dose of the MBX-8025 therapy, for safety and pharmacodynamic evaluations. At each visit, after a 12-hour fast, blood is drawn and urine collected; and a standard metabolic panel, complete blood count, and standard urinalysis are performed. Blood is analyzed for TC, HDL-C, TG, VLDL-C, LDL-C, and apolipoprotein B, for liver function markers such as total and bone-specific alkaline phosphatases, for γ -glutamyl transpeptidase, and also for total and conjugated bilirubin. The subjects also maintain health diaries, which are reviewed at each visit. The subjects show a dose-related improvement in their disease, as shown by, for example, a decrease in ALP and GGT.

Claims:

1. A method of treating an intrahepatic cholestatic disease by administering MBX-8025 or an MBX-8025 salt.
2. MBX-8025 or an MBX-8025 salt for use in treating an intrahepatic cholestatic disease.
- 5 3. The use of MBX-8025 or an MBX-8025 salt for treating an intrahepatic cholestatic disease.
4. The use of MBX-8025 or an MBX-8025 salt in the manufacture of a medicament for treating an intrahepatic cholestatic disease.
5. A pharmaceutical composition comprising MBX-8025 or an MBX-8025 salt for treating
10 an intrahepatic cholestatic disease.
6. A kit for treating an intrahepatic cholestatic disease comprising a composition comprising MBX-8025 or an MBX-8025 salt.
7. The method, compounds, uses, composition, or kit of any one of the previous claims where the MBX-8025 or an MBX-8025 salt is MBX-8025 L-lysine dihydrate salt.
- 15 8. The method, compounds, uses, composition, or kit of any one of the previous claims where the MBX-8025 or an MBX-8025 salt is administered orally.
9. The method, compounds, uses, composition, or kit of any one of the previous claims where the daily dose of MBX-8025 or an MBX-8025 salt (when calculated as the free acid) is 20-200 mg/day, preferably 50-100 mg/day.
- 20 10. The method, compounds, uses, composition, or kit of any one of the previous claims where the MBX-8025 or an MBX-8025 salt is administered once/day.
11. The method, compounds, uses, composition, or kit of any one of the previous claims where the intrahepatic cholestatic disease is primary biliary cirrhosis.
12. The method, compounds, uses, composition, or kit of any one of the previous claims
25 where the intrahepatic cholestatic disease is primary sclerosing cholangitis.

13. The method, compounds, uses, composition, or kit of any one of the previous claims where the intrahepatic cholestatic disease is progressive familial intrahepatic cholestasis.
14. The method, compounds, uses, composition, or kit of any one of the previous claims where the intrahepatic cholestatic disease is Alagille syndrome.