

- 1 -

33415-00

A PROCESS FOR THE PREPARATION OF PYRIDINE DICARBOXYLATE
DERIVATIVES

BACKGROUND OF THE INVENTION

Pyridine-2,3-dicarboxylic acids and esters are building blocks for numerous bio-active products, particularly imidazolinone herbicides, for example U.S. 4,758,667. Known methods to prepare pyridine-2,3-
5 dicarboxylate derivatives via the oxidation of a suitable quinoline or alkylpyridine precursor are often plagued by the use of costly oxidants such as KMnO_4 , $\text{H}_2\text{Cr}_2\text{O}_7$, SeO_2 and the like; by long reaction time cycles such as in the use of ozone, electrolysis and the like; and by undesirable
10 side reactions such as decarboxylation, N-oxide formation and the like. Therefore, new methods to construct the desired pyridine dicarboxylate product are continually being sought.

It has now been found that a rapid one vessel
15 process to prepare a pyridine-2,3-dicarboxylate derivative is readily obtained via the sequential condensation of the appropriate alkyl vinyl ether with Vilsmeier reagent, oxalacetate and an ammonia source.



G00002




166166

- 2 -

SUMMARY OF THE INVENTION

The present invention provides an efficient and effective process for the preparation of pyridine-2,3-dicarboxylate derivatives of formula I


5



(I)

wherein R₁ is H or C₁-C₄alkyl optionally substituted with C₁-C₄alkoxy or halogen; and

10 R₂ and R₃ are each independently C₁-C₆alkyl which comprises reacting an alkyl vinyl ether compound of formula II




(II)

15

wherein the formula II compound is the cis isomer, the trans isomer or a mixture thereof, R is C₁-C₄alkyl and R₁ is as defined for formula I with at least one molar equivalent of Vilsmeier reagent optionally in the presence of a first solvent to form a first intermediate;

20

2) reacting said first intermediate with at least one molar equivalent of an oxalacetate of formula III



(III)

- 3 -

wherein R_2 and R_3 are as defined for formula I in the presence of at least two molar equivalents of a base to form a second intermediate; and

- 5 3) reacting said second intermediate with an ammonia source optionally in the presence of a second solvent to form the formula I pyridine diester product.


Compounds of formula I are useful in the preparation of imidazolinone herbicidal agents.

10

DETAILED DESCRIPTION OF THE INVENTION

Among the methods known to prepare pyridine 2,3-dicarboxylate derivatives are degradative methods such as oxidation of the appropriately substituted quinoline or
15 alkylpyridine precursors. However, oxidation procedures often are costly and hazardous. Advantageously, it has now been found that pyridine-2,3-dicarboxylate derivatives of formula I may be effectively prepared in a single vessel via the sequential condensation of a
20 suitable alkyl vinyl ether of formula II with Vilsmeier reagent, an oxalacetate of formula III and an ammonia source. The reaction is shown in Flow Diagram I, wherein $X^{\ominus}$ represents $Cl^{\ominus}$ or $PO_2Cl_2^{\ominus}$.

- 4 -

Flow Diagram I


5 In accordance with the process of the invention, an alkyl vinyl ether of formula II, which may be the cis isomer or the trans isomer or a mixture thereof, may be reacted with at least one molar equivalent of Vilsmeier reagent optionally in the presence of a first solvent to

10 form a first intermediate (a vinylogous imidate salt), which may then be reacted with at least one molar equivalent of an oxalacetate of formula III in the presence of at least two molar equivalents of a base, preferably an organic amine base, to form a second

- 5 -

intermediate (an iminium salt), which may be reacted with an ammonia source optionally (and preferably) in the presence of a second solvent to form the desired formula I pyridine-2,3-dicarboxylate product.

5 The term Vilsmeier reagent, as used in the specification and claims, designates the *in situ* product of the reaction of dimethyl formamide with an activating agent such as oxalyl chloride, phosgene, phosphorous oxychloride, thionyl chloride, and the like and may be
10 illustrated as an immonium salt of formula IV or the analogues thereof wherein $X^{\ominus}$ represents $Cl^{\ominus}$ or $PO_2Cl_2^{\ominus}$.



15

The term halogen as used in the specification and claims designates Cl, Br, I or F.

Solvents suitable for use as the first solvent in the inventive process may be any inert organic solvent
20 such as a hydrocarbon, e.g. hexanes, pentanes, heptanes and the like; a halogenated hydrocarbon, e.g. methylene chloride, chloroform, dichloroethane, and the like, preferably dichloroethane or methylene chloride; an aromatic hydrocarbon, e.g. benzene, toluene, xylene and
25 the like; a halogenated aromatic hydrocarbon, e.g. chlorobenzene, o-dichlorobenzene, or mixtures thereof. Preferably the reaction is conducted with a first solvent and preferably the first solvent is a halogenated hydrocarbon such as dichloroethane or methylene chloride.

30 Bases suitable for use in the inventive process are organic amines such as triethylamine, pyridine, lutidine, N,N-dimethylpiperidine, N-methylpyrrolidine and the like, preferably triethylamine or pyridine.

- 6 -


Ammonia sources suitable for use in the process of the invention may be any of the conventional means for producing NH_3 *in situ*, including ammonia gas, ammonium salts, and the like, preferably ammonium salts such as
5 ammonium acetate, ammonium sulfamate, and the like.

Solvents suitable for use as the second solvent in the inventive process are protic solvents such as water; alcohols such as C_1 - C_4 alkanols e.g. ethanol, methanol, propanol, butanol and the like, preferably ethanol;
10 organic acids such as C_1 - C_4 carboxylic acids e.g. acetic acid, propionic acid and the like, preferably acetic acid.

The rate of formation of the reaction product is generally directly related to the reaction temperature. In general, lower temperatures will decrease the rate of
15 reaction and higher temperatures will increase the rate of reaction. However, excessively high temperatures are not desired and may lead to a decrease in product yield and purity. Preferable temperatures range from about 0°C to 120°C .

20 In order to facilitate a further understanding of the invention, the following examples are set forth primarily for the purpose of illustrating certain more specific details thereof. The invention is not to be limited thereby except as defined in the claims. The term NMR
25 designates nuclear magnetic resonance.

- 7 -

EXAMPLE 1Preparation of Diethyl Pyridine-2,3-dicarboxylate

5

A solution of dimethyl formamide (73 g, 1.00 mole) in ethylene dichloride is slowly treated with oxalyl chloride (88 mL, 1.00 mole), with cooling, stirred at ambient temperatures for 16 hours, treated with ethyl vinyl ether (72.1 g, 1.00 mole) over a 1 hour period and stirred at ambient temperatures for 16 hours. This reaction mixture is treated sequentially with diethyl oxalacetate (199.28 g, 1.06 mole) and (with cooling) triethylamine (224 g, 2.2 mole), stirred for 0.5 hours and treated with a premixed solution of concentrated HCl (200 ml) and concentrated NH_4OH (200 ml) in 100 ml of water. The reaction mixture is treated further with water (250 ml), concentrated NH_4OH (70 ml) and acetic acid (200 ml). The resultant mixture is distilled under N_2 at 88°C and atmosphere pressure to remove 1850 g of distillate. The distillation pot is then treated with absolute ethanol (1.0 L) and $\text{NH}_4\text{OCOCH}_3$ (180 g), heated at reflux temperature for 16 hours and distilled at 100°C to

- 8 -

remove 887 g of distillate. The distillation pot is cooled and the residue is partitioned between water and 2:1 ethyl acetate/hexanes. The organic phase is separated, washed sequentially with water and brine and concentrated *in vacuo* to give the title product as an oil, 179.5 g, (77% pure) 58.5% yield, identified by NMR analysis.

EXAMPLES 2-7

10

Using essentially the same procedure described in Example 1, and employing the appropriately substituted vinyl ether substrate, the results shown in Table I are obtained.

-9-

TABLE I

Example Number	$ \begin{array}{c} \text{R}_1 \\ \\ \text{CH}_2 \\ \\ \text{CH} \\ \\ \text{N}^+ \text{R}_1 \\ \\ \text{RO} \end{array} \xrightarrow[\text{(1st solvent)}]{\text{Vilsmeier reagent}} \xrightarrow[\text{base}]{\text{DOX}^1} \xrightarrow[\text{(2nd solvent)}]{\text{3) } \text{NH}_4^+ \text{Y}^-} \text{R}_1\text{-} \begin{array}{c} \text{C}_6\text{H}_3\text{N} \\ \quad \\ \text{CO}_2\text{C}_2\text{H}_5 \quad \text{CO}_2\text{C}_2\text{H}_5 \end{array} $			base (Molar-equivalents)	Y ⁻ (2 nd solvent)	% yield
	R	R ₁	acid chloride (1 st solvent)			
2	nBu	H	oxalyl chloride (ClCH ₂ CH ₂ Cl)	Triethylamine (2.2)	⁻ OCOCH ₃ (C ₂ H ₅ OH)	52.8
3	nBu	H	oxalyl chloride (ClCH ₂ CH ₂ Cl)	Pyridine (2.2)	⁻ OCOCH ₃ (C ₂ H ₅ OH)	39.6
4	C ₂ H ₅	H	POCl ₃ (CH ₂ Cl ₂)	Triethylamine (3.5)	⁻ OCOCH ₃ (C ₂ H ₅ OH)	62.3
5	C ₂ H ₅	CH ₃	oxalyl chloride (CH ₂ Cl ₂)	Triethylamine (2.5)	⁻ SO ₃ NH ₂ (CH ₃ CO ₂ H)	48
6	C ₂ H ₅	C ₂ H ₅	oxalyl chloride (CH ₂ Cl ₂)	Triethylamine (2.5)	⁻ OCOCH ₃ (CH ₃ CO ₂ H)	42
7	CH ₃	CH ₂ OCH ₃	POCl ₃ (CH ₂ Cl ₂)	Triethylamine (3.5)	⁻ OH (H ₂ O/CH ₃ CO ₂ C ₂ H ₅)	17.7

¹ Diethyl oxalacetate

ACY-33415


-10-

The derivatives of formula I obtained by the process
5 of this invention may be converted to herbicidal 2-
(imidazolin-2-yl)-3-pyridine carboxylic acids by procedures
described in US Patent No 4758667. For example a 2,3-
pyridine of formula I may be reacted in the presence of a
strong base with an 2-aminoalkane carboxamide of formula V

10




wherein R₄ and R₅ are each independently C₁-C₄ alkyl to
give after pH adjustment a 2-(imidazolin-2-yl)-3-pyridine
15 of formula:



-11-

CLAIMS:

1. A process for the preparation of a compound of formula I




(I)

wherein R_1 is H or C_1 - C_4 alkyl optionally substituted with C_1 - C_4 alkoxy or halogen; and

R_2 and R_3 are each independently C_1 - C_6 alkyl which comprises


- a) reacting an alkyl vinyl ether compound of formula II



(II)

wherein the formula II compound is the cis isomer, the trans isomer or a mixture thereof, R is C_1 - C_4 alkyl and R_1 is as defined for formula I with at least one molar equivalent of Vilsmeier reagent optionally in the presence of a first solvent to form a first intermediate;

- b) reacting said first intermediate with at least one molar equivalent of an oxalacetate of formula III



(III)


wherein R_2 and R_3 are as defined for formula I in the presence of at least two molar equivalents of an organic amine base to form a second intermediate; and

-12-

- c) reacting said second intermediate with an ammonia source optionally in the presence of a second solvent to form the formula I pyridine diester product.
2. The process according to Claim 1 wherein the first solvent is present and is a halogenated hydrocarbon.
 3. The process according to Claim 1 or Claim 2 wherein the base is triethylamine.
 4. The process according to any one of Claims 1 to 3 wherein the second solvent is present and is a C₁-C₄alkanol or a C₁-C₄carboxylic acid or a mixture thereof.
 5. The process according to any one of Claims 1 to 4 wherein the ammonia source is ammonium acetate.
 6. The process according to any one of Claims 1 to 5 wherein R₁ is H, methyl, ethyl or methoxymethyl.
 7. The process according to claim 6 wherein R₁ is H.
 8. The process according to any one of Claims 1 to 7 wherein R₃ and R₄ are each independently methyl or ethyl.
 9. A compound of formula I whenever prepared by a process as claimed in any one of claims 1 to 8.
 10. An imidazolinone herbicide whenever prepared from a compound of formula I as claimed in Claim 9.


-13-

11. A process for preparing a 2-(imidazolin-2-yl)-3-pyridine of formula (VI)



which comprises


A) the preparation of a compound of formula I



wherein R₁ is H or C₁-C₄alkyl optionally substituted with C₁-C₄alkoxy or halogen; and

R₂ and R₃ are each independently C₁-C₆alkyl which comprises

a) reacting an alkyl vinyl ether compound of formula II



wherein the formula II compound is the cis isomer, the trans isomer or a mixture thereof, R is C₁-C₄alkyl and R₁ is as defined for formula I with at least one molar equivalent of Vilsmeier reagent optionally in the presence of a first solvent to form a first intermediate;

-14-

c) reacting said second intermediate with an ammonia source optionally in the presence of a second solvent to form the formula I pyridine diester product;

B) reacting a 2,3-pyridine of formula I

in the presence of a strong base with an 2-aminoalkane carboxamide of formula V:



(V)

wherein R₄ and R₅ are each independently C₁-C₄ alkyl and adjusting the pH to give a compound of formula (VI):