

PROCESS FOR THE CARBONYLATION OF ACETYLENICALLY
UNSATURATED COMPOUNDS

The invention relates to a process for the carbonylation of acetylenically unsaturated compounds, whereby a feedstock, comprising an acetylenically unsaturated compound and a relatively minor amount of an 1,2-alkadiene compound, is contacted under carbonylation conditions with carbon monoxide and a hydroxylated co-reactant.

A problem, encountered with the processes for the carbonylation of acetylenically unsaturated compounds, consists in that the available feedstocks for these processes generally contain, in addition to the acetylenically unsaturated compound, 1,2-alkadiene compounds (so-called allenes). The presence of 1,2-alkadiene compounds, even in relatively small amounts, unfavourably affects the catalyst activity. Small quantities of 1,2-alkadiene impurities (up to 0.4%) can often be tolerated. However, the amounts commonly present in the acetylenic feedstocks require that special measures are taken, before they can be used for the carbonylation process.

In EP-A-0,441,446 a carbonylation catalyst system is disclosed that performs adequately under basic conditions. The catalyst system comprises: a) a source of a Group VIII metal (e.g., a palladium compound); b) a (mono- or bidentate) (di)phosphine having an aromatic substituent which contains an imino nitrogen atom, e.g., an optionally substituted 2-pyridyl group; c) a source of protons; and d) a tertiary amine. The illustrated basic catalyst systems (based on phosphines bearing 2-pyridyl and 6-methyl-2-pyridyl-groups) exhibit improved tolerance

- 2 -


towards allenes (up to 7%, example 18), provided a tertiary amine is present (see comparative example G versus examples 12 and 15). However, no information may be gained from this document that further improvements in respect of the carbonylation process, even carried out in the absence of the tertiary amine, could be achieved.

Surprisingly it has now been found that by employing certain specific mono- or bidentate (di)phosphine ligands, the tolerance with respect to allenes is further improved.

This selection moreover allows the carbonylation to be performed under non-basic conditions, starting from conventional feedstocks that are contaminated with 1,2-alkadiene. Specific measures for the removal of 1,2-alkadiene compounds, preceding the carbonylation reaction, are thereby made redundant.

The invention may be defined as relating to a process for the carbonylation of acetylenically unsaturated compounds, whereby a feedstock comprising an acetylenically unsaturated compound and a relatively minor amount of an 1,2-alkadiene compound is contacted under carbonylation conditions with carbon monoxide and a hydroxylated co-reactant, in the presence of a catalyst system based on:

- a) a source of cations of one or more metals of Group VIII of the Periodic Table;
- b) a phosphine of the general formula $PR^1R^2R^3$ or $R^2R^3M-R-PR^1R^2$, wherein R^1 represents a heteroaryl group of the formula



This Sheet Has Been Replaced



wherein A represents a functional group that is electron-withdrawing relative to hydrogen, and each of X, Y and Z independently represents a nitrogen atom or a C-Q group, whereby Q is a hydrogen atom or a substituent, R^2 and R^3 independently represent a substituted or non-substituted (hetero)hydrocarbyl group or have the aforesaid meaning of R^1 , M is an element of Group VIa, preferably a nitrogen or phosphorus atom, R represents a bridging (substituted) hydrocarbyl group having 1 to 4 carbon atoms in the bridge; and
c) a protonic acid.

The catalyst system used in the process of the invention is based, as regards a), on a source of cations of one or more metals of Group VIII. These metals include iron, cobalt, nickel, ruthenium, rhodium, palladium, iridium, osmium, and platinum. Preferably, the catalyst system is based on a source of palladium cations.

The source of cations of metals of Group VIII may be the metallic element or a metal compound, such as a metal salt or a complex of the metal with a phosphine, with carbon monoxide or with acetylacetonate. It is advantageously a metal compound, in particular a metal salt. Examples of suitable metal salts are salts of sulphuric acid, nitric acid, sulphonic acids, phosphonic acids, perhalic acids and carboxylic acids, such as alkane carboxylic acids with 1 to 12 carbon atoms, for example acetic acid and propionic acid, or halogenated carboxylic acids, for example trichloroacetic acid and trifluoroacetic acid. Palladium acetate has proved to be a particularly suitable source of metal cations.

In the phosphines of the catalyst system (component b), each of R^2 and R^3 preferably represents a substituted or unsubstituted pyridyl, alkyl or aryl group. Examples of suitable R^2 and R^3 groups are 2-pyridyl, phenyl, tolyl-, xylyl, and cyclohexyl groups and alkyl groups

having from 3 to 7 carbon atoms. Phosphines wherein both R^2 and R^3 represent a phenyl group are preferred.

The functional group(s) A may for instance be a trialkylammonium, a nitro, cyano, sulphonic, carbonic, carbonyl, or a haloalkyl group or a halogen atom (e.g., any of the groups listed in "Organic Chemistry", by Morrison and Boyd, 3rd ed. 1980, p. 360.). Other electron-withdrawing groups will be known to the skilled man. Preferably (for ease of manufacturing the ligand) it is selected from the group consisting of halogen atoms and haloalkyl groups with one to three carbon atoms. More preferably, A represents a halogen atom such as a bromine, fluorine or chlorine atom. Phosphines, in which (each of) the functional group(s) A is a chlorine atom, are most preferred.

Examples of phosphines, wherein X, Y and/or Z represents a nitrogen atom are phosphines substituted with a pyrimidinyl-, pyrazinyl-, s-triazinyl- and as-triazinyl group.

Preferably, the phosphine is substituted with a 2-pyridyl group, i.e. X, Y and Z each represents a group C-Q, in which Q is a hydrogen atom or a substituent.

Suitable substituents represented by Q include electron-withdrawing substituents, in particular the substituents as indicated above with respect to A. Advantageously, Q represents electron-donating substituents, such as alkyl groups, for example methyl and ethyl groups, amino and (di)alkylamino groups.

Preferably, the phosphine ligand is a monophosphine of the general formula $PR^1R^2R^3$.

The function of the protonic acid on which the catalyst system further is based (component c), is believed to provide a source of protons. The protonic acid may be generated in situ.

Preferably the protonic acid has a substantially non-coordinating anion, i.e. an anion which does not, or only to a very minor extent, coordinate with the metal of Group VIII. Preferred acids in this respect are sulphuric acid, sulphononic acids and halogenated carboxylic acids.

Examples of suitable protonic acids are optionally substituted alkylsulphonic acids, such as methanesulphonic acid, trifluoromethanesulphonic acid, tert-butylsulphonic acid, perhalic acids, such as perchloric acid, and acidic ion exchange resins, such as a sulphonated ion exchange resin.

The number of moles of phosphine and of moles of protonic acid per mole atoms of metal of Group VIII may vary considerably. Recommended phosphine amounts are in the range of 10 to 100 moles of phosphine per mole atom of metal of Group VIII and in particular in the range of 20 to 80. The amount of protonic acid is preferably selected such that per mole atom of metal of Group VIII, 2 to 500 moles of protonic acid are present.

The catalyst system of the invention may be homogeneous or heterogeneous, but is preferably homogeneous. The amount in which the catalyst is applied is suitably selected such that per mole of acetylenically unsaturated compound to be converted, from 10^{-8} to 10^{-1} mole atoms of Group VIII metal is present, preferably from 10^{-7} to 10^{-2} on the same basis.

Suitable acetylenically unsaturated compounds, to be used as starting material in the process of the invention, include optionally substituted alkynes with 2 to 20 carbon atoms per molecule. Examples are acetylene, propyne, 1-butyne, 2-butyne, 1-hexyne, phenyl acetylene and benzylethyne. Preferably, unsubstituted alkynes with 3 to 10 carbon atoms are used.

In view of the industrial outlets for the carbonylated products, propyne is a preferred starting material.

As has been stated above, a major advantage of the catalyst systems of the invention consists in their tolerance towards 1,2-alkadiene compounds in the acetylenic feedstocks. Accordingly, commercially available feedstocks may be used that containing small amounts of 1,2-alkadiene compounds, such as propadiene, in addition to the acetylenically unsaturated compounds. In general, a 1,2-alkadiene content of at most 10%, based on acetylenically unsaturated compound, can be tolerated. It is recommended to use feedstocks in which the amount of 1,2-alkadiene compounds is lower, suitably in the range of 0.002 to 0.05 moles per mole of acetylenically unsaturated compound.

The hydroxylated co-reactant may be any hydroxyl-containing compound such as a monohydric, dihydric or polyhydric alkanol, a phenol, or water.

Monohydric alkanols are preferred, in particular those having from 1 to 4 carbon atoms. Among these, methanol is most preferred.

The co-reactant is suitably used in excess, thereby avoiding the need of a separate diluent or solvent. However, a liquid diluent may be applied, if so desired. Preferably, non-alkaline diluents are used, such as ketones, e.g. methylisobutylketone, or ethers, e.g. dipropylether or 2,5,8-trioxanonane (diglyme).

Owing to the high activity of the catalysts, the process of the invention proceeds readily at moderate reaction conditions. Suitable reaction temperatures are, for instance, in the range of 20 to 100 °C, preferably in the range of 30 to 75 °C.

The reaction pressure is usually selected in the range of 1 to 100 bar. Preferably, the pressure is in the range of 5 to 70 bar.

The invention is illustrated with the following, non-limiting examples.

Examples

All experiments were carried out in a 250 ml "Hastelloy C" (trade mark) magnetically stirred autoclave. The autoclave was charged with 0.025 mmoles of palladium(II) acetate, the selected diphenyl(2-pyridyl)phosphine and protonic acid in the amounts indicated hereafter, and 50 ml of methanol.

Air was evacuated from the autoclave, whereupon 30 ml of a propadiene-containing propyne feed was added.

Subsequently, carbon monoxide was supplied up to a pressure of 60 bar. The autoclave was sealed and heated to the desired reaction temperature.

As soon as the falling pressure remained constant (marking the completion of the reaction), the contents of the autoclave were cooled and a sample was withdrawn and analysed by gas liquid chromatography.

Example I

a) An experiment was carried out in the manner as outlined above, whereby as phosphine 1 mmol of biphenyl(6-chloro-2-pyridyl)phosphine and as protonic acid 2 mmol of trifluoromethanesulphonic acid was used. The feed was propyne, containing 0.2% of propadiene. The reaction temperature was 45 °C.

The reaction time (completion) was 0.16 hour. Analysis showed that methyl methacrylate had been formed with a selectivity of 99.88% at a propyne conversion of about 100%. The average reaction rate was calculated to be 73,000 moles of product per mole atom of palladium and per hour.

b) The experiment described under a) was repeated with the difference that the propyne feed contained 2.2% of propadiene.

The reaction time was 0.5 hour. The average reaction rate was calculated to be 24,000 moles of product per mole atom of palladium and per hour.

Example II

An experiment was carried out substantially as described in Example I with the following differences:

- i) the propyne feed contained 4.5% of propadiene;
- ii) 2 mmol of biphenyl(6-chloro-2-pyridyl)phosphine were used; and
- iii) the reaction temperature was 50 °C.

The reaction time was 2.5 hours. The average reaction rate was calculated to be 5,500 moles of product per mole atom of palladium and per hour.

Example III

An experiment was carried out substantially as described in Example II, with the following differences:

- i) the propyne feed contained 4.7% of propadiene;
- ii) 3 mmol of biphenyl(6-chloro-2-pyridyl)phosphine were used; and
- iii) as protonic acid 2 mmol of tert-butylsulphonic acid was used.

The reaction time was 1 hour. The average reaction rate was calculated to be 8,900 moles of product per mole atom of palladium and per hour.

Example IV

An experiment was carried out substantially as described in Example III, with the following differences:

- i) the propyne feed contained 7.0% of propadiene;
- ii) as protonic acid 3 mmol of methanesulphonic acid was used; and
- iii) the reaction temperature was 60 °C.

The reaction time was 0.5 hour. The average reaction rate was calculated to be 20,000 moles of product per mole atom of palladium and per hour. Analysis showed that the propyne conversion was >95% and that the reaction mixture contained 5% of propadiene, calculated on residual propyne, indicating that most of the propadiene had been converted into methyl methacrylate product.

Example V

a) An experiment was carried out, substantially as described in Example I(a), with the following differences:

- i) 1 mmol of biphenyl(6-bromo-2-pyridyl)phosphine was used;
- ii) as protonic acid 2 mmol of methanesulphonic acid was used; and
- iii) the reaction temperature was 30 °C.

The reaction time was 1.5 hour. The average reaction rate was calculated to be 9,200 moles of product per mole atom of palladium and per hour.

b) The experiment described under a) was repeated with the following differences:

- i) the propyne feed contained 1.7% of propadiene;
- ii) as protonic acid 2 mmol of trifluoromethanesulphonic acid was used; and
- iii) the reaction temperature was 45 °C.

The reaction time was 2.5 hours. The average reaction rate was calculated to be 6,000 moles of product per mole atom of palladium and per hour.

Example IV

An experiment was carried out, substantially as described in Example I(a), with the following differences:

- i) 1 mmol of biphenyl(6-trifluoromethyl-2-pyridyl)phosphine and 1 mmol of biphenyl(2-pyridyl)phosphine were used;

- ii) as protonic acid 3 mmol of trifluoromethanesulphonic acid was used; and
- iii) the reaction temperature was 35 °C (with an exotherm of 60 °C).

The reaction time was 1 hour. Analysis showed that methyl methacrylate had been formed with a selectivity of 98.6%. The average reaction rate was calculated to be 100,000 moles of product per mole atom of palladium and per hour.

Example A (for comparison, not according to the invention)

a) An experiment was carried out, substantially as described in Example V(a), with the following differences:

- i) 1 mmol of biphenyl(2-pyridyl)phosphine was used; and
- ii) the reaction temperature was 30 °C.

The reaction time was 2 hours. The average reaction rate was calculated to be 2,500 moles of product per mole atom of palladium and per hour.

b) The experiment described under a) was repeated with the following differences:

- i) the propyne feed contained 2.3% of propadiene; and
- ii) the reaction temperature was 50 °C.

The reaction time was 5 hours. The average reaction rate was calculated to be 625 moles of product per mole atom of palladium and per hour.

Example B (for comparison, not according to the invention)

a) An experiment was carried out, substantially as described in Example V(a), with the following differences:

- i) 1 mmol of biphenyl(3-methyl-2-pyridyl)phosphine was used; and
- ii) the reaction temperature was 50 °C.

The reaction time was 5 hours. The average reaction rate was calculated to be 7,100 moles of product per mole atom of palladium and per hour.

b) The experiment described under a) was repeated with the difference that the propyne feed contained 2.3% of propadiene.

The average reaction rate was calculated to be 800 moles of product per mole atom of palladium and per hour.

Example C (for comparison, not according to the invention)

a) An experiment was carried out, substantially as described in Example V(a), with the following differences:

- i) 1 mmol of biphenyl(3-chloro-2-pyridyl)phosphine was used; and
- ii) the reaction temperature was 35 °C.

The reaction time was 0.25 hour. The average reaction rate was calculated to be 50,000 moles of product per mole atom of palladium and per hour.

b) The experiment described under a) was repeated with the following differences:

- i) the propyne feed contained 2.3% of propadiene; and
- ii) the reaction temperature was 50 °C.

The reaction time was 5 hours. The average reaction rate was calculated to be 2,100 moles of product per mole atom of palladium and per hour.

Example D (for comparison, not according to the invention)

a) An experiment was carried out, substantially as described in Example V(a), with the following differences:

- i) 1 mmol of biphenyl(5-chloro-2-pyridyl)phosphine was used; and
- ii) the reaction temperature was 35 °C.

The reaction time was 0.1 hour. The average reaction rate was calculated to be 175,000 moles of product per mole atom of palladium and per hour.

b) The experiment described under a) was repeated with the following differences:

- i) the propyne feed contained 2.3% of propadiene; and
- ii) the reaction temperature was 50 °C.

The reaction time was 5 hours. The average reaction rate was calculated to be 3,300 moles of product per mole atom of palladium and per hour.

Example E (for comparison, not according to the invention)

An experiment was carried out, substantially as described in Example A(b), with the difference that as phosphine biphenyl(6-methyl-2-pyridyl)phosphine was used.

The average reaction rate was calculated to be 400 moles of product per mole atom of palladium and per hour.

Conclusions

As can be seen from Example A, with a non-substituted phosphine as catalyst component, a reaction rate of only 2,500 moles per mole atom Pd per hour is obtained, even when a feedstock having a relatively low (0.2%) propadiene content is used. Contrarily, the presence of substituted phosphines in the catalyst systems, used in Examples B, C, and D, results in considerably higher reaction rates in converting feedstocks having a low propadiene content.

However, as shown in Examples B(b), C(b) and D(b), the activity of these catalysts considerably decreases by the presence of propadiene in the feedstock.

In contrast, the use of the catalysts described in Examples I to VI, for the conversion of feedstocks containing 2% or more of propadiene, still results in high reaction rates.




TS 0020 SIN

C L A I M S

1. A process for the carbonylation of acetylenically unsaturated compounds, whereby a feedstock comprising an acetylenically unsaturated compound and a relatively minor amount of a 1,2-alkadiene compound is contacted under carbonylation conditions with carbon monoxide and a hydroxylated co-reactant, in the presence of a catalyst system based on:

- a) a source of cations of one or more metals of Group VIII of the Periodic Table;
- b) a phosphine of the general formula $PR^1R^2R^3$ or $R^2R^3M-R-PR^1R^2$, wherein R^1 represents a heteroaryl group of the formula



wherein A represents a functional group that is electron-withdrawing relative to hydrogen, and each of X, Y and Z independently represents a nitrogen atom or a C-Q group, whereby Q is a hydrogen atom or a substituent, R^2 and R^3 independently represent a substituted or non-substituted (hetero)hydrocarbyl group or have the aforesaid meaning of R^1 , M is an element of Group VIa, preferably a nitrogen or phosphorus atom, R represents a bridging (substituted) hydrocarbyl group having 1 to 4 carbon atoms in the bridge; and

- c) a protonic acid.

2. A process as claimed in claim 1, wherein the catalyst system is based on a phosphine wherein R^2 and R^3





* C L M S *

THIS SHEET HAS BEEN
REPLACED

- 14 -

independently represent a substituted or non-substituted pyridyl, alkyl or aryl group, preferably a phenyl group.

3. A process as claimed in claim 1 or 2, wherein the catalyst system is based on a phosphine wherein A represents a substituent selected from the group consisting of halogen atoms and haloalkyl groups with one to three carbon atoms.

4. A process as claimed in claim 3, wherein A represents a halogen atom, preferably a chlorine atom.

5. A process as claimed in any of claims 1 to 4, wherein the catalyst system is based on a phosphine wherein R¹ represents a 6-A-2-pyridyl group.

6. A process as claimed in any of claims 1 to 5, wherein the catalyst system is based on a monophosphine, preferably on biphenyl(6-chloro-2-pyridyl)phosphine.

7. A process as claimed in any one of claims 1 to 6, carried out in the absence of a tertiary amine.

8. A process as claimed in any of claims 1 to 7, wherein the amount of 1,2-alkadiene compound in the feedstock is less than 0.1 mole per mole of acetylenically unsaturated compound.

9. A process as claimed in claim 10, wherein the molar amount of 1,2-alkadiene compound in the feedstock per mole of acetylenically unsaturated compound is in the range of 0.002 to 0.05.

10. A process as claimed in any of claims 1 to 10, wherein methyl methacrylate is prepared by reacting a feedstock comprising propyne and 1,2-propadiene, with carbon monoxide and methanol.

C21/TS0020FF


This Sheet Has Been Replaced



A B S T R A C TPROCESS FOR THE CARBONYLATION OF ACETYLENICALLY
UNSATURATED COMPOUNDS

The invention relates to a process for the carbonylation of acetylenically unsaturated compounds, whereby a feedstock comprising an acetylenically unsaturated compound and a relatively minor amount of an 1,2-alkadiene compound is contacted under carbonylation conditions with carbon monoxide and a hydroxylated co-reactant, in the presence of a catalyst system, based on:

- a) a source of cations of one or more metals of Group VIII of the Periodic Table;
- b) a phosphine of the general formula $PR^1R^2R^3$ or $R^2R^3M-R-PR^1R^2$, wherein R^1 represents a heteroaryl group of the formula



- wherein A represents a functional group that is electron-withdrawing relative to hydrogen, and each of X, Y and Z independently represents a nitrogen atom or a C-Q group, whereby Q is a hydrogen atom or a substituent, R^2 and R^3 independently represent a substituted or non-substituted (hetero)hydrocarbyl group or have the aforesaid meaning of R^1 , M is an element of Group VIa, preferably a nitrogen or phosphorus atom, R represents a bridging (substituted) hydrocarbyl group having 1 to 4 carbon atoms in the bridge; and
- c) a protonic acid.

C21/TS0020FF

Substitute sheet - 23 April 1998

- 2 -


towards allenes (up to 7%, example 18), provided a tertiary amine is present (see comparative example G versus examples 12 and 15). However, no information may be gained from this document that further improvements in respect of the carbonylation process, even carried out in the absence of the tertiary amine, could be achieved.

Surprisingly it has now been found that by employing certain specific mono- or bidentate (di)phosphine ligands, the tolerance with respect to allenes is further improved.

This selection moreover allows the carbonylation to be performed under non-basic conditions, starting from conventional feedstocks that are contaminated with 1,2-alkadiene. Specific measures for the removal of 1,2-alkadiene compounds, preceding the carbonylation reaction, are thereby made redundant.

The invention may be defined as relating to a process for the carbonylation of acetylenically unsaturated compounds, whereby a feedstock comprising an acetylenically unsaturated compound and an amount of an 1,2-alkadiene compound in a molar ratio of 0.004 to 0.1 is contacted under carbonylation conditions with carbon monoxide and hydroxylated co-reactant, in the presence of a catalyst system based on:

- a) a source of cations of one or more metals of Group VIII of the Periodic Table;
- b) a phosphine of the general formula $PR^1R^2R^3$ or $R^2R^3M-R-PR^1R^2$, wherein R^1 represents a heteroaryl group of the formula



REPLACEMENT SHEET

24 APR 1998




C L A I M S

1. A process for the carbonylation of acetylenically unsaturated compounds, whereby a feedstock comprising an acetylenically unsaturated compound and an amount of 1,2-alkadiene compound in a molar ratio of 0.004 to 0.1 is contacted under carbonylation conditions with carbon monoxide and a hydroxylated co-reactant, in the presence of a catalyst system based on:

a) a source of cations of one or more metals of Group VIII of the Periodic Table;

b) a phosphine of the general formula $PR^1R^2R^3$ or $R^2R^3M-R-PR^1R^2$, wherein R^1 represents a heteroaryl group of the formula



wherein A represents a functional group that is electron-withdrawing relative to hydrogen, and each of X, Y and Z independently represents a nitrogen atom or a C-Q group, whereby Q is a hydrogen atom or a substituent, R^2 and R^3 independently represent a substituted or non-substituted (hetero)hydrocarbyl group or have the aforesaid meaning of R^1 , M is an element of Group VIa, preferably a nitrogen or phosphorus atom, R represents a bridging (substituted) hydrocarbyl group having 1 to 4 carbon atoms in the bridge; and

c) a protonic acid.

2. A process as claimed in claim 1, wherein the catalyst system is based on a phosphine wherein R^2 and R^3

REPLACEMENT SHEET

24 APR 1998



- 14 -

independently represent a substituted or non-substituted pyridyl, alkyl or aryl group, preferably a phenyl group.

3. A process as claimed in claim 1 or 2, wherein the catalyst system is based on a phosphine wherein A represents a substituent selected from the group consisting of halogen atoms and haloalkyl groups with one to three carbon atoms.

4. A process as claimed in claim 3, wherein A represents a halogen atom, preferably a chlorine atom.

5. A process as claimed in any of claims 1 to 4, wherein the catalyst system is based on a phosphine wherein R¹ represents a 6-A-2-pyridyl group.

6. A process as claimed in any of claims 1 to 5, wherein the catalyst system is based on a monophosphine, preferably on biphenyl(6-chloro-2-pyridyl)phosphine.

7. A process as claimed in any one of claims 1 to 6, carried out in the absence of a tertiary amine.

8. A process as claimed in any of claims 1 to 7, wherein the amount of 1,2-alkadiene compound in the feedstock per mole of acetylenically unsaturated compound is in the range of 0.004 to 0.05.

9. A process as claimed in any of claims 1 to 8, wherein methyl methacrylate is prepared by reacting a feedstock comprising propyne and 1,2-propadiene, with carbon monoxide and methanol.

REPLACEMENT SHEET

24 APR 1998