

IL-34 ANTISENSE AGENTS AND METHODS OF USING SAME

CROSS REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of and priority to U.S. Provisional Patent Application No. 63/201,887, filed May 17, 2021, the disclosure of which is hereby incorporated by reference in its entirety for all purposes.

BACKGROUND

[0002] Interleukin-34 (IL-34) is a recently discovered cytokine functionally overlapping with macrophage colony stimulating factor (M-CSF, also known as MCSF1 and MCSF-1), a mediator of inflammation and osteoclastogenesis in bone-degenerative diseases such as rheumatoid arthritis. Inflammatory diseases, both acute and chronic, are an important disease category that is still not completely understood.

[0003] For example, rheumatoid arthritis (RA) is a chronic autoimmune and inflammatory disorder of the joints and bones suffered by approximately 1.3 million patients in the United States. Among those afflicted with RA, the prevalence of work disability associated with RA was around 35% in the United States, with an overall health care cost of more than \$19.3 billion. Osteoarthritis (OA) is a chronic inflammatory disease of the joints, an involves mechanical wear and destruction of the smooth cartilage joint surface that caps the bones in the joints. OA is the most common form of arthritis and affects 32.5 million adults in the United States, with an overall health care cost of more than \$136 billion. No medical cure currently exists for RA or OA.

[0004] Inflammatory bowel disease is a chronic inflammatory disorder of the gastrointestinal tract suffered by approximately 1.4 million patients in the United States. It is one of the five most prevalent gastrointestinal disease burdens in the United States, with an overall health care cost of more than \$1.7 billion. Each year in the United States, inflammatory bowel disease accounts for more than 700,000 physician visits, 100,000 hospitalizations, and disability in 119,000 patients. No medical cure currently exists, so disease management requires a lifetime of care.

[0005] The two most common forms of inflammatory bowel disease are Crohn's disease and ulcerative colitis. Although Crohn's disease can affect the entire gastrointestinal tract, it primarily affects the ileum (the distal or lower portion of the small intestine) and the large intestine. Ulcerative colitis primarily affects the colon and the rectum. The etiology of

inflammatory bowel disease is not completely understood, although both environmental and genetic factors are believed to play a role in the disease. Environmental components may include alterations in flora of the gut which are affected by exposure to ingested foods and medications.

[0006] Inflammatory bowel disease is associated with abdominal pain, vomiting, diarrhea, rectal bleeding, severe cramps, muscle spasms, weight loss, malnutrition, fever, and anemia. Patients with inflammatory bowel disease may also suffer from skin lesions, joint pain, eye inflammation, and liver disorders, and children suffering from ulcerative colitis may suffer from growth defects. Although rarely fatal, these symptoms decrease quality of life for patients.

[0007] Thus, there is a pressing need to develop reliable methods of treating inflammatory disorders such as rheumatoid arthritis, osteoarthritis, or inflammatory bowel disease. There is a further need to identify methods of treatment that provide effective and permanent relief from symptoms across a broad spectrum of patients and which are not associated with negative side effects or cycles of inflammation and remission.

SUMMARY

[0008] The present application provides compositions of polynucleotides complementary to IL-34, or a portion thereof, and methods of using the compositions.

[0009] Disclosed herein, in certain embodiments, are polynucleotides, comprising a nucleotide sequence according to: (a) 5'-CTTTGGGCCCGCACCAGCTTC-3' (SEQ ID NO: 3) or (b) 5'-TCCATGACCCGGAAGCAGTT-3' (SEQ ID NO: 5), wherein at least one cytidine of the nucleotide sequence is chemically modified, and at least one nucleoside is a 2'-O-(2-methoxyethyl) (2'-MOE) nucleoside; or a pharmaceutically acceptable salt thereof. In some embodiments, no more than 10 nucleosides of the polynucleotide are chemically modified. In some embodiments, the polynucleotide comprises six 2'-MOE nucleosides. In some embodiments, the polynucleotide comprises eight 2'-MOE nucleosides. In some embodiments, the polynucleotide comprises a nucleotide sequence is selected from the group consisting of: (a) 5'-CxTxTxTxGGGCXGCACCAGCxTxTxCx-3' (SEQ ID NO: 40), wherein Cx is 2'-O-(2-methoxyethyl)cytidine, Tx is 2'-O-(2-methoxyethyl)thymidine, and X is 5-methylcytidine; (b) 5'-CxTxTxTGGGCXGCACCAGCTxTxCx-3' (SEQ ID NO: 41), wherein Cx is 2'-O-(2-methoxyethyl)cytidine, Tx is 2'-O-(2-methoxyethyl)thymidine, and X is 5-methylcytidine; (c) 5'-TxCxCxAxTGACCXGGAAGCAxGxTxTx-3' (SEQ ID NO: 42), wherein Cx is 2'-O-(2-

methoxyethyl)cytidine, Ax is 2'-O-(2-methoxyethyl)adenosine, Tx is 2'-O-(2-methoxyethyl)thymidine, Gx is 2'-O-(2-methoxyethyl)guanosine, and X is 5-methylcytidine; and (d) 5'-TxCxCxATGACCXGGAAGCAGxTxTx-3' (SEQ ID NO: 43), wherein Cx is 2'-O-(2-methoxyethyl)cytidine, Tx is 2'-O-(2-methoxyethyl)thymidine, Gx is 2'-O-(2-methoxyethyl)guanosine, and X is 5-methylcytidine; or a pharmaceutically acceptable salt thereof. In some embodiments, the polynucleotide is complementary to IL-34. In some embodiments, the polynucleotide is an IL-34 antisense oligonucleotide. In some embodiments, the polynucleotide is an IL-34 siRNA. In some embodiments, at least one internucleoside linkage of the polynucleotide is selected from the group consisting of a phosphorothioate linkage, a phosphorodithioate linkage, a phosphotriester linkage, an alkylphosphonate linkage, an aminoalkylphosphotriester linkage, an alkylene phosphonate linkage, a phosphinate linkage, a phosphoramidate linkage, and an aminoalkylphosphoramidate linkage, a thiophosphoramidate linkage, thionoalkylphosphonate linkage, a thionoalkylphosphotriester linkage, a thiophosphate linkage, a selenophosphate linkage, and a boranophosphate linkage. In some embodiments, at least one internucleoside linkage of the polynucleotide is a phosphorothioate linkage. In some embodiments, all internucleoside linkages of the polynucleotide are phosphorothioate linkages. In some embodiments, the polynucleotide is 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 nucleotides in length. In some embodiments, the polynucleotide is 20 nucleotides in length or 20-25, 20-30, 20-35, 25-30, 25-35, or 30-35 nucleotides in length. In some embodiments, the polynucleotide is no more than 20, 25, or 30 nucleotides in length. In some embodiments, the polynucleotide is from 20 to 25 nucleotides in length. In some embodiments, the polynucleotide is 20 nucleotides in length.

[0010] Disclosed herein, in certain embodiments, are pharmaceutical compositions, comprising (a) a polynucleotide disclosed herein, or a pharmaceutically acceptable salt thereof, and (b) a pharmaceutically acceptable excipient. In some embodiments, the pharmaceutical composition is suitable for topical, parenteral, oral, pulmonary, intratracheal, intranasal, transdermal, or intraduodenal administration.

[0011] Disclosed herein, in certain embodiments, are methods of treating an inflammatory disease in a patient in need thereof, the method comprising administering to the patient in need thereof an effective amount of a polynucleotide disclosed herein, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition disclosed herein. In some embodiments,

the inflammatory disease is associated with altered IL-34 expression. In some embodiments, the method inhibits inflammatory cytokine production in cells of the patient. In some embodiments, the method reduces or inhibits an IL-34 mediated inflammatory response in cells of the patient. In some embodiments, the method reduces or inhibits IL-34-mediated macrophage colony-stimulating factor receptor (M-CSFR-1) signaling in cells of the patient. In some embodiments, the cell is an intestinal cell. In some embodiments, the cell is an intestinal stromal cell. In some embodiments, the cell forms part of an intestinal fibrostricture. In some embodiments, the polynucleotide is administered intraarticularly, rectally, topically, parenterally, orally, pulmonarily, intratracheally, intranasally, transdermally, or intraduodenally. In some embodiments, the polynucleotide is administered orally. In some embodiments, the patient is a human.

[0012] Disclosed herein, in certain embodiments, are methods of preventing or treating a cancer associated with altered IL-34 expression in a patient in need thereof, the method comprising administering to the patient in need thereof an effective amount of a polynucleotide disclosed herein, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition disclosed herein. In some embodiments, the cancer has elevated IL-34 expression. In some embodiments, the cancer is colon cancer.

[0013] Disclosed herein, in certain embodiments, are methods of reducing or eliminating a fibrotic stricture in a patient suffering from an inflammatory disease, the method comprising administering an effective amount of a polynucleotide disclosed herein, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition disclosed herein. In some embodiments, the fibrotic stricture is located in the intestine. In some embodiments, the inflammatory disease is selected from the group consisting of an inflammatory bowel disease, rheumatoid arthritis, psoriasis, osteoarthritis, pouchitis, diabetes (type I and II), tissue or organ rejection, multiple sclerosis, periodontal inflammation, periodontitis, pigmented villonodular synovitis, hepatitis, sinusitis, colon cancer, colorectal cancer, colitis-associated colon cancer, sporadic colorectal cancer, coronary artery disease, Sjogren's syndrome (SS), obesity, chronic inflammation, pulmonary sarcoidosis, skin lesions, a CNS inflammatory disease, and an autoimmune disease. In some embodiments, the inflammatory disease is an inflammatory bowel disease. In some embodiments, the inflammatory bowel disease is selected from the group consisting of Crohn's disease, gastroduodenal Crohn's disease, Crohn's (granulomatous) colitis, inflammatory Crohn's

disease, fibrostricturing Crohn's disease, ulcerative colitis, collagenous colitis, lymphocytic colitis, ischaemic colitis, diversion colitis, Behçet's disease, microscopic colitis, ulcerative proctitis, proctosigmoiditis, jejunoileitis, left-sided colitis, pancolitis, ileocolitis, ileitis, and indeterminate colitis. In some embodiments, the inflammatory bowel disease is inflammatory Crohn's disease. In some embodiments, the inflammatory bowel disease is fibrostricturing Crohn's disease. In some embodiments, the polynucleotide is administered intraarticularly, rectally, topically, parenterally, orally, pulmonarily, intratracheally, intranasally, transdermally, or intraduodenally. In some embodiments, the polynucleotide is administered orally. In some embodiments, the patient is a human.

[0014] Disclosed herein, in certain embodiments, are methods for preventing or treating fibrosis, the method comprising administering to a patient in need thereof a therapeutically effective amount of a polynucleotide disclosed herein, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition disclosed herein. In some embodiments, the fibrosis is intestinal fibrosis. In some embodiments, the fibrosis is pulmonary fibrosis. In some embodiments, the fibrosis is synovial fibrosis. In some embodiments, the fibrosis is selected from the group consisting of renal fibrosis, cardiac fibrosis, endomyocardial fibrosis, myelofibrosis, retroperitoneal fibrosis, and nephrogenic systemic fibrosis. In some embodiments, the patient is also suffering from Crohn's disease. In some embodiments, the polynucleotide is administered intraarticularly, rectally, topically, parenterally, orally, pulmonarily, intratracheally, intranasally, transdermally, or intraduodenally. In some embodiments, the polynucleotide is administered orally. In some embodiments, the patient is a human.

[0015] Disclosed herein, in certain embodiments, are methods of preventing or treating intestinal fibrosis, the method comprising administering to a patient in need thereof, a pharmaceutically effective amount of a polynucleotide disclosed herein, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition disclosed herein. In some embodiments, the patient is also suffering from Crohn's disease. In some embodiments, the polynucleotide is administered intraarticularly, rectally, topically, parenterally, orally, pulmonarily, intratracheally, intranasally, transdermally, or intraduodenally. In some embodiments, the polynucleotide is administered orally. In some embodiments, the patient is a human.

[0016] Disclosed herein, in certain embodiments, are methods of preventing or treating pulmonary fibrosis, the method comprising administering to a patient in need thereof, a

pharmaceutically effective amount of a polynucleotide disclosed herein, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition disclosed herein. In some embodiments, the patient is also suffering from Crohn's disease. In some embodiments, the polynucleotide is administered intraarticularly, rectally, topically, parenterally, orally, pulmonarily, intratracheally, intranasally, transdermally, or intraduodenally. In some embodiments, the polynucleotide is administered orally. In some embodiments, the patient is a human.

[0017] Disclosed herein, in certain embodiments, are methods of preventing or treating synovial fibrosis, the method comprising administering to a patient in need thereof, a pharmaceutically effective amount of a polynucleotide disclosed herein, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition disclosed herein. In some embodiments, the polynucleotide is administered intraarticularly, rectally, topically, parenterally, orally, pulmonarily, intratracheally, intranasally, transdermally, or intraduodenally. In some embodiments, the polynucleotide is administered orally. In some embodiments, the patient is a human.

[0018] Disclosed herein, in certain embodiments, are uses of a polynucleotide disclosed herein, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for treating an inflammatory disease. In some embodiments, the inflammatory disease is selected from the group consisting of an inflammatory bowel disease, rheumatoid arthritis, psoriasis, osteoarthritis, pouchitis, diabetes (type I and II), tissue or organ rejection, multiple sclerosis, periodontal inflammation, periodontitis, pigmented villonodular synovitis, hepatitis, sinusitis, colon cancer, colorectal cancer, colitis-associated colon cancer, sporadic colorectal cancer, coronary artery disease, or Sjogren's syndrome (SS), obesity, chronic inflammation, pulmonary sarcoidosis, skin lesions, a CNS inflammatory disease, and an autoimmune disease. In some embodiments, the inflammatory disease is rheumatoid arthritis or osteoarthritis. In some embodiments, the inflammatory disease is an inflammatory bowel disease. In some embodiments, the inflammatory bowel disease is selected from the group consisting of Crohn's disease, inflammatory Crohn's disease, fibrostricturing Crohn's disease, gastroduodenal Crohn's disease, Crohn's (granulomatous) colitis, ulcerative colitis, collagenous colitis, lymphocytic colitis, ischaemic colitis, diversion colitis, Behçet's disease, microscopic colitis, ulcerative proctitis, proctosigmoiditis, jejunoileitis, left-sided colitis, pancolitis, ileocolitis, ileitis, and indeterminate colitis. In some embodiments, the inflammatory bowel disease is inflammatory

Crohn's disease. In some embodiments, the inflammatory bowel disease is fibrostricturing Crohn's disease.

[0019] Disclosed herein, in certain embodiments, are polynucleotides disclosed herein, or a pharmaceutically-acceptable salt thereof, for use as a medicament.

[0020] Disclosed herein, in certain embodiments, are polynucleotides disclosed herein, or a pharmaceutically-acceptable salt thereof, for use in the treatment of an inflammatory disease. In some embodiments, the inflammatory disease is selected from the group consisting of an inflammatory bowel disease, rheumatoid arthritis, psoriasis, osteoarthritis, pouchitis, diabetes (type I and II), tissue or organ rejection, multiple sclerosis, periodontal inflammation, periodontitis, pigmented villonodular synovitis, hepatitis, sinusitis, colon cancer, colorectal cancer, colitis-associated colon cancer, sporadic colorectal cancer, coronary artery disease, or Sjogren's syndrome (SS), obesity, chronic inflammation, pulmonary sarcoidosis, skin lesions, a CNS inflammatory disease, and an autoimmune disease. In some embodiments, the inflammatory disease is an inflammatory bowel disease. In some embodiments, the inflammatory bowel disease is selected from the group consisting of Crohn's disease, inflammatory Crohn's disease, fibrostricturing Crohn's disease, gastroduodenal Crohn's disease, Crohn's (granulomatous) colitis, ulcerative colitis, collagenous colitis, lymphocytic colitis, ischaemic colitis, diversion colitis, Behçet's disease, microscopic colitis, ulcerative proctitis, proctosigmoiditis, jejunoileitis, left-sided colitis, pancolitis, ileocolitis, ileitis, and indeterminate colitis. In some embodiments, the inflammatory bowel disease is inflammatory Crohn's disease. In some embodiments, the inflammatory bowel disease is fibrostricturing Crohn's disease.

[0021] Disclosed herein, in certain embodiments, are polynucleotides disclosed herein, or a pharmaceutically-acceptable salt thereof, for use in the treatment of fibrosis. In some embodiments, the fibrosis is intestinal fibrosis. In some embodiments, the fibrosis is pulmonary fibrosis. In some embodiments, the fibrosis is synovial fibrosis. In some embodiments, the fibrosis is selected from the group consisting of renal fibrosis, cardiac fibrosis, endomyocardial fibrosis, myelofibrosis, retroperitoneal fibrosis, and nephrogenic systemic fibrosis.

BRIEF DESCRIPTION OF THE DRAWINGS

[0022] **FIG. 1** shows the results of the IL-34 ASOs on the modulation of IL-34 expression in cultured HT-29 cells. IL-34 RNA transcripts were evaluated by real-time PCR. Levels are

normalized to β -actin. (CTR=scrambled ASO, 1=SEQ ID NO: 17, 2=SEQ ID NO: 19, 3=SEQ ID NO: 42, 4=SEQ ID NO: 43). Values are mean $\pm$ SEM of four experiments. Differences among groups were compared using 1-way analysis of variance followed by the Dunnett's post hoc test. IL-34 ASOs vs CTR, *** $P < 0.001$.

[0023] FIG. 2 shows the results of the IL-34 ASOs on the modulation of IL-34 expression in cultured AGS cells. IL-34 RNA transcripts were evaluated by real-time PCR. Levels are normalized to β -actin. (CTR=scrambled ASO, 1=SEQ ID NO: 17, 2=SEQ ID NO: 19, 3=SEQ ID NO: 42, 4=SEQ ID NO: 43). Values are mean $\pm$ SEM of four experiments. Differences among groups were compared using 1-way analysis of variance followed by the Dunnett's post hoc test. IL-34 ASOs vs CTR, *** $P < 0.001$.

[0024] FIG. 3 shows the results of the IL-34 ASOs on the modulation of IL-34 expression in UC patient-derived LPMCs. IL-34 RNA transcripts were evaluated by real-time PCR. Levels are normalized to β -actin. (CTR=scrambled ASO, 1=SEQ ID NO: 17, 2=SEQ ID NO: 19, 3=SEQ ID NO: 42, 4=SEQ ID NO: 43). Values are mean $\pm$ SEM of three experiments. Differences among groups were compared using 1-way analysis of variance followed by the Dunnett's post hoc test. IL-34 ASOs vs CTR, ** $P < 0.01$, *** $P < 0.001$.

DETAILED DESCRIPTION

[0025] Disclosed herein, *inter alia*, are polynucleotides that are complementary to an IL-34 nucleic acid, or a portion thereof (*e.g.*, IL-34 antisense oligonucleotides) and comprising one or more chemically modified nucleoside, compositions comprising said oligonucleotides, and methods of using the compositions for treating, reducing, or inhibiting a disease or disorder, such as an autoimmune or inflammatory diseases (*e.g.*, rheumatoid arthritis (RA), osteoarthritis (OA), Crohn's disease or ulcerative colitis) or fibrosis.

Definitions

[0026] As used herein, "oligonucleotide" or "polynucleotide" are used interchangeably and refers to short DNA or RNA molecules. In some embodiments, oligonucleotides comprise 2'-deoxyribonucleotides (oligodeoxyribonucleotides), which can be modified at the phosphate backbone or at the 2' sugar position. In some embodiments, oligonucleotides of the disclosure have one or more phosphorothioate (PS) backbone modifications, where one or more of the non-bridging oxygen atoms in the phosphate backbone is replaced with a sulfur atom. In some

embodiments, a “phosphorothioate oligonucleotide” or “PS oligonucleotide” has at least one internucleoside linkage that is an *O,O*-linked phosphorothioate linkage (*i.e.*, a phosphorothioate linkage). In some embodiments, a PS oligonucleotide has all internucleoside linkages being *O,O*-linked phosphorothioate linkages (*i.e.*, a phosphorothioate linkage).

[0027] As used herein, “antisense therapeutics” are a class of nucleic acid-based compounds that can be used to inhibit gene expression. Antisense therapeutics may be single- or double-stranded deoxyribonucleic acid (DNA)-based, ribonucleic acid (RNA)-based, mixed DNA and RNA-based, or DNA/RNA chemical analogue compounds. In some embodiments, antisense therapeutics comprise antisense oligonucleotides. In general, antisense therapeutics are designed to include a nucleotide sequence that is complementary or nearly complementary to an mRNA or pre-mRNA sequence transcribed from a given gene in order to promote binding between the antisense therapeutic and the pre-mRNA or mRNA. Without being bound by theory, it is believed that in most instances antisense therapeutics act by binding to an mRNA or pre-mRNA, thereby inhibiting protein translation, altering pre-mRNA splicing into mature mRNA, and/or causing destruction of mRNA. In most instances, the antisense therapeutic nucleotide sequence is complementary to a portion of a targeted gene’s or mRNA’s sense sequence.

[0028] As used herein, “IL-34 antisense therapeutics” are oligonucleotide-based compounds that include an oligonucleotide sequence complementary to an IL-34 gene sense, IL-34 pre-mRNA sense, and/or IL-34 mRNA sense sequence, or a portion thereof. IL-34 antisense therapeutics described herein can also be nucleotide chemical analog-based compounds capable of binding to an IL-34 gene sense, IL-34 pre-mRNA sense, and/or IL-34 mRNA sense sequence, or a portion thereof. In some embodiments, IL-34 antisense therapeutics may include IL-34 antisense oligonucleotides, IL-34 shRNAs, IL-34 siRNAs, IL-34 PNAs, IL-34 LNAs, miRNAs, and IL-34 morpholino oligomers.

[0029] As used herein, “antisense oligonucleotides” (ASOs) are short synthetic oligonucleotide-based sequences comprising sequences complementary to a target RNA sequence such as a mRNA that encodes a target protein (*e.g.*, IL-34). Without being bound to a particular theory, antisense oligonucleotide sequences can hybridize to a complementary region in an mRNA molecule thereby producing a double-stranded hybrid that can lead to the activation of ubiquitous catalytic enzymes, such as RNase H, which degrade DNA/RNA hybrid strands thus preventing protein translation. Without being bound by theory, an antisense oligonucleotide

provided herein can hybridize to its target sequence as RNA or DNA. Accordingly, in some embodiments, if a DNA sequence is provided as a target, the corresponding RNA sequence (including uracil instead of thymine) is included in an ASO as described herein. In some embodiments, ASOs described herein comprise chemically modified nucleosides.

[0030] The number of nucleotides included in antisense oligonucleotides described herein can vary. ASOs are typically between 8 to 50 nucleotides in length, for example, 20 nucleotides in length. In some embodiments, the antisense oligonucleotide is from 15-20, 15-25, 15-30, 15-35, 20-25, 20-30, 20-35, 25-30, 25-35, or 30-35 nucleotides in length. In particular embodiments, the antisense oligonucleotide is from 15-25 nucleotides in length. In particular embodiments, the antisense oligonucleotide is from 20-25 nucleotides in length. In some embodiments, the antisense oligonucleotide is 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, or 35 nucleotides in length. In some embodiments an antisense oligonucleotide described herein includes a maximum number of nucleotides. In some embodiments, an antisense oligonucleotide is no more than 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, or 35 nucleotides in length.

[0031] As used herein, “chemically modified nucleoside” means any nucleoside other than adenosine, cytidine, thymidine, guanosine, or uridine. A chemically modified nucleoside may comprise a modified sugar moiety and/or a modified nucleobase. In some embodiments, a chemically modified nucleosides disclosed may be a locked nucleic acid (LNA), *e.g.*, LNA cytidine, LNA thymidine, LNA adenosine, or LNA guanosine. In some embodiments, chemically modified nucleosides comprise 2'-*O*-methyl (“2'-OMe”) ribonucleosides, for example, 2'-*O*-methylcytidine, 2'-*O*-methylguanosine, 2'-*O*-methylthymidine, 2'-*O*-methyluridine, and/or 2'-*O*-methyladenosine. In other embodiments, a chemically modified nucleoside comprises a 5-methylpyrimidine, for example, 5-methylcytosine; and/or a 5-methylpurine, for example, 5-methylguanine. In some embodiments, chemically modified nucleosides can include any of the following chemically-modified nucleosides: 5-methyl-2'-*O*-methylcytidine, 5-methyl-2'-*O*-methylthymidine, 5-methylcytidine, 5-methyluridine, and/or 5-methyl-2'-deoxycytidine. In some embodiments disclosed herein, chemically modified nucleosides can include a stabilized terminal 5'-phosphate or phosphatase-resistant analogue of 5'-phosphate, *e.g.*, 5'-methyl phosphonate, 5'-methylenephosphonate, a 5'-methylenephosphonate analog, 5'-*E*-vinyl phosphonate (5'-*E*-VP), 5'-phosphorothioate, and a 5'-*C*-methyl analog; a 2'-

O-methyl ribonucleoside, *e.g.*, 2'-*O*-methylcytidine, 2'-*O*-methylguanosine, 2'-*O*-methyluridine, 2'-*O*-methylthymidine, and 2'-*O*-methyladenosine; a 5-methylpyrimidine, *e.g.*, 5-methylcytosine; a 5-methylpurine, *e.g.*, 5-methylguanine; 5-methyl-2'-*O*-methylcytidine; 5-methyl-2'-*O*-methylthymidine; 5-methylcytidine; 5-methyluridine; or 5-methyl 2'-deoxycytidine.

[0032] Additionally, in some embodiments, chemically modified nucleosides described herein include 2'-*O*-(2-methoxyethyl) ("2'-MOE") nucleosides, 2'-deoxy-2'-fluoro nucleosides, 2'-fluoro- β -D-arabinonucleosides, bridged nucleosides, LNA nucleosides, constrained ethyl (cET) nucleosides, tricyclo-DNA (tcDNA) nucleosides, 2'-O,4'-C-ethylene linked nucleic acid (ENA) nucleosides, and/or peptide nucleic acids (PNA).

[0033] As used herein, "internucleoside linkage" means the connection between the 3' position of a nucleoside and the 5' position of an adjacent nucleoside. A "modified internucleoside linkage" refers to a connection between the 3' position of a nucleoside and the 5' position of an adjacent nucleoside that is not natural. In some embodiments, a modified internucleoside linkage includes, but is not limited to: a phosphorothioate linkage; a phosphorodithioate linkage, a phosphotriester linkage, an alkylphosphonate linkage, an aminoalkylphosphotriester linkage, an alkylene phosphonate linkage, a phosphinate linkage, a phosphoramidate linkage, an aminoalkylphosphoramidate linkage, a thiophosphoramidate linkage, a thionoalkylphosphonate linkage, a thionoalkylphosphotriester linkage, a thiophosphate linkage, a selenophosphate linkage, or a boranophosphate linkage.

[0034] Peptide nucleic acids (PNAs) are short, artificially synthesized polymers with a structure that mimics DNA or RNA. PNAs include a backbone composed of repeating N-(2-aminoethyl)-glycine units linked by peptide bonds.

[0035] Locked nucleic acids (LNAs) are oligonucleotide sequences that include one or more modified RNA nucleotides in which the ribose moiety is modified with an extra bridge connecting the 2' oxygen and 4' carbon. LNAs are believed to have higher *T_m*'s than analogous oligonucleotide sequences.

[0036] Morpholino oligomers are oligonucleotide compounds that include DNA bases attached to a backbone of methylenemorpholine rings linked through phosphorodiamidate groups. Morpholino oligomers of the present invention can be designed to bind to specific RNA sequences of interest (*e.g.*, IL-34 mRNA or IL-34 pre-mRNA sequences of interest), thereby preventing gene expression.

[0037] Small hairpin RNAs (shRNAs) are generally RNA molecules with a hairpin-like structure that can be used to silence gene expression. shRNAs are generally expressed from plasmids encoding the shRNA sequence, and can be expressed from viral vectors to allow lentiviral, adenoviral, or adeno-associated viral expression. Without being bound by theory, it is believed that shRNA inhibits gene expression by taking advantage of RNA interference (RNAi) processes. In brief, the shRNA transcript is processed by Drosha and Dicer, and then loaded onto the RNA-induced silencing complex (RISC), allowing targeting of specific mRNA, and either mRNA degradation or repression of protein translation.

[0038] Small interfering RNAs (siRNAs) are double-stranded RNA molecules of approximately 20-25 base pairs in length (but which can also be, for example, 18-30 base pairs in length) that take advantage of RNAi machinery (*e.g.*, Drosha and RISC) to bind and target mRNA for degradation. siRNAs are not dependent upon plasmids or vectors for expression, and can generally be delivered directly to a target cell, for instance, by transfection.

[0039] MicroRNAs (miRNAs) are small non-coding RNA molecule containing about 22 nucleotides that function in RNA silencing and post-transcriptional regulation of gene expression. miRNAs include sequences that are complementary to portions of an mRNA sequence. miRNAs are produced from long, single-stranded RNA molecules exhibiting highly specific stem-loop structures. Artificial miRNAs are designed by replacing the mature 21 nucleotide sequence of naturally occurring miRNA sequences with 21 nucleotide sequences from a target, for example, an IL-34 mRNA target.

[0040] As used herein, “IL-34 inhibitors” comprise, but are not limited to, nucleotide-based inhibitors of IL-34 (for example, IL-34 small hairpin RNAs (shRNAs), IL-34 microRNAs (miRNAs), IL-34 small interfering RNAs (siRNAs), and IL-34 antisense oligonucleotides, including IL-34 antisense oligonucleotides that include LNA nucleotides, peptide nucleic acids (PNAs), and morpholino oligomers), and compositions that include such compounds. In some embodiments, IL-34 levels (*e.g.*, IL-34 mRNA or protein levels) and/or activity (*e.g.*, biological activity, for example, IL-34 receptor stimulation) can be decreased using IL-34 inhibitors. In some embodiments, IL-34 inhibitors comprise IL-34 antisense oligonucleotides comprising one or more chemically modified nucleoside.

[0041] As used herein, “IL-34 antisense oligonucleotide” or “IL-34 ASO” is understood to refer to an oligonucleotide comprising a nucleic acid sequence that is complementary to a nucleic

acid sequence in an mRNA molecule transcribed from the IL-34 gene, or a portion thereof. More specifically, such an oligonucleotide can be complementary to the nucleic acid sequence in the coding region of such an mRNA, or a portion thereof. In some embodiments, an IL-34 antisense oligonucleotide possesses the inherent functional property of targeting the IL-34 gene, its RNA or protein products, or another molecular entity whose activity or expression impinges upon the activity or expression of IL-34 or its products either exclusively or with a high degree of specificity. In some embodiments, an IL-34 antisense oligonucleotide can reduce the expression of IL-34 when introduced into a cell (*e.g.*, an immune cell, such as a PBMC, dendritic cell, or B-cell). In some embodiments, an IL-34 antisense oligonucleotide can reduce expression of an mRNA transcribed from the gene. In some embodiments, an IL-34 antisense oligonucleotide can reduce expression of a protein encoded by the gene. In some embodiments, an IL-34 antisense oligonucleotide can reduce secretion of a protein encoded by the gene from the cell into which the IL-34 antisense oligonucleotide was introduced. In some embodiments, an IL-34 antisense oligonucleotide comprises the sequence of SEQ ID NOs: 40-43, or pharmaceutically acceptable salts thereof.

[0042] Throughout the description and claims of this specification the word “comprise” and other forms of the word, such as “comprising” and “comprises,” means including but not limited to, and is not intended to exclude, for example, other additives, components, integers, or steps.

[0043] “Optional” or “optionally” means that the subsequently described event or circumstance can or cannot occur, and that the description includes instances where the event or circumstance occurs and instances where it does not.

[0044] As used herein, the term “about” or “approximately,” when used in reference to a quantitative value, includes the recited quantitative value itself, unless specifically stated otherwise. As used herein, the term “about” or “approximately” refers to a $\pm 10\%$ variation from the recited quantitative value unless otherwise indicated or inferred from the context.

[0045] The terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present disclosure. As used throughout this disclosure, the singular forms “a,” “an,” and “the” include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to “a composition” includes a plurality of such compositions, as well as a single composition, and a reference to “a therapeutic agent” is a reference to one or more therapeutic and/or pharmaceutical agents and equivalents thereof

known to those skilled in the art, and so forth. All percentages and ratios used herein, unless otherwise indicated, are by weight.

[0046] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure belongs. In the case of conflict, the present specification will control. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present disclosure, suitable methods and materials are described below.

Polynucleotides complementary to IL-34

[0047] Disclosed herein, *inter alia*, are polynucleotides complementary to IL-34. In some embodiments, the polynucleotide comprises the nucleotide sequence: 5'-CTTTGGGCCGACACAGCTTC-3' (SEQ ID NO: 3), or 5'-TCCATGACCCGGAAGCAGTT-3' (SEQ ID NO: 5); 5'-CTTTGGGCXGCACACAGCTTC-3' (SEQ ID NO: 7); or 5'-TCCATGACCXGGAAGCAGTT-3' (SEQ ID NO: 8), wherein a X is 5-methylcytidine, at least one cytidine of the nucleotide sequence is chemically modified, and at least one nucleoside is a 2'-O-(2-methoxyethyl) (2'-MOE) nucleoside, or a pharmaceutically acceptable salt thereof. In some embodiments, the polynucleotide comprises a nucleotide sequence selected from the group consisting of:

- a. 5'-CxTxTxTxGGGCXGCACACAGCxTxTxCx-3' (SEQ ID NO: 40), wherein Cx is 2'-O-(2-methoxyethyl)cytidine, Tx is 2'-O-(2-methoxyethyl)thymidine, and X is 5-methylcytidine;
- b. 5'-CxTxTxTGGGCXGCACACAGCTxTxCx-3' (SEQ ID NO: 41), wherein Cx is 2'-O-(2-methoxyethyl)cytidine, Tx is 2'-O-(2-methoxyethyl)thymidine, and X is 5-methylcytidine;
- c. 5'-TxTxTxAxTGACCXGGAAGCAxGxTxTx-3' (SEQ ID NO: 42), wherein Cx is 2'-O-(2-methoxyethyl)cytidine, Ax is 2'-O-(2-methoxyethyl)adenosine, Tx is 2'-O-(2-methoxyethyl)thymidine, Gx is 2'-O-(2-methoxyethyl)guanosine, and X is 5-methylcytidine; and
- d. 5'-TxTxTxATGACCXGGAAGCAGxTxTx-3' (SEQ ID NO: 43), wherein Cx is 2'-O-(2-methoxyethyl)cytidine, Tx is 2'-O-(2-methoxyethyl)thymidine, Gx is 2'-O-(2-methoxyethyl)guanosine, and X is 5-methylcytidine;

or a pharmaceutically acceptable salt thereof.

[0048] In some embodiments, the polynucleotide is an IL-34 antisense oligonucleotide (ASO).

[0049] In some embodiments, the polynucleotide is an siRNA. In some embodiments, the siRNA is an IL-34 siRNA that comprises the nucleotide sequence of any one of SEQ ID NO: 3, 5, 7, or 8 or SEQ ID NO: 40-43. In some embodiments, the polynucleotide is a part of an siRNA complex. In some embodiments, the siRNA is an siRNA duplex.

[0050] In some embodiments, the polynucleotide is an IL-34 inhibitor. In some embodiments, the IL-34 inhibitor is an IL-34 ASO or a composition comprising such an IL-34 ASO. In some embodiments, the IL-34 ASOs are short synthetic oligonucleotide sequences complementary to an IL-34 transcript (for example, an IL-34 mRNA transcript) or a portion thereof.

[0051] In some embodiments, an IL-34 ASO comprises a nucleotide sequence complementary to a portion of an IL-34 mRNA sequence, for example, a mouse or human IL-34 mRNA sequence.

[0052] In some embodiments, an IL-34 ASO comprises a nucleotide sequence complementary to a portion of an IL-34 gene sequence, for example a mouse or human IL-34 gene sequence.

[0053] In some embodiments, an IL-34 ASO comprises a nucleotide sequence that is complementary or nearly complementary to a portion of an IL-34 mRNA transcript variant, or a portion thereof, for example, human IL-34 mRNA sequence of NCBI Reference Sequence NM_001172771.1 (SEQ ID NO: 24), NM_001172772.1 (SEQ ID NO: 25), NM_152456.2 (SEQ ID NO: 26), NM_152456.3 (SEQ ID NO: 44), NM_001172771.2 (SEQ ID NO: 45), NM_001393493.1 (SEQ ID NO: 46), NM_001393495.1 (SEQ ID NO: 47), NM_001172772.2 (SEQ ID NO: 48), NM_001393494.1 (SEQ ID NO: 49), NM_001393496.1 (SEQ ID NO: 50), NM_001393497.1 (SEQ ID NO: 51), or NM_001393498.1 (SEQ ID NO: 52); or mouse IL-34 mRNA sequence of NCBI Reference Sequence NM_001135100.2 (SEQ ID NO: 27) or NM_029646.3 (SEQ ID NO: 28).

[0054] In some embodiments, an IL-34 ASO can target IL-34 mRNAs produced from IL-34 genes of one or more species, for example, mouse and human IL-34 mRNA transcripts. For example, an IL-34 ASO can target an IL-34 mRNA of a mammalian IL-34 gene, for example, a human (*i.e.*, *Homo sapiens*) IL-34 gene or a mouse (*i.e.*, *mus musculus*) IL-34 gene. In particular embodiments, the IL-34 ASO targets a human IL-34 mRNA. In some embodiments, the IL-34

ASO includes a nucleotide sequence that is complementary to a nucleotide sequence of an IL-34 gene or an IL-34 mRNA, or a portion thereof.

[0055] In some embodiments, IL-34 ASOs include but are not limited to, an IL-34 ASO that comprises the nucleotide sequence: 5'-CTCACCAAGACCCACAG-3' (SEQ ID NO: 1); 5'-GGCTTTGGGCCGACACAGCT-3' (SEQ ID NO: 2); 5'-CTTTGGGCCGACACAGCTTC-3' (SEQ ID NO: 3); 5'-TGGGCCGACACAGCTTCAGG-3' (SEQ ID NO: 4); 5'-TCCATGACCCGGAAGCAGTT-3' (SEQ ID NO: 5); 5'-TGTTTCATGTACTGAAG-3' (SEQ ID NO: 6); 5'-CTTTGGGCXGCACACAGCTTC-3' (SEQ ID NO: 7); or 5'-TCCATGACCXGGAAGCAGTT-3' (SEQ ID NO: 8), wherein X is 5-methylcytidine.

[0056] In some embodiments, the IL-34 ASO comprises the nucleotide sequence 5'-CTTTGGGCCGACACAGCTTC-3' (SEQ ID NO: 3). In some embodiments, the IL-34 ASO comprises the nucleotide sequence 5'-TCCATGACCCGGAAGCAGTT-3' (SEQ ID NO: 5).

[0057] In some embodiments, the IL-34 ASO comprises chemically modified nucleosides. In some embodiments, a chemically modified nucleoside disclosed herein includes, but is not limited to: a locked nucleic acid (LNA) nucleoside, *e.g.*, LNA cytidine, LNA thymidine, LNA adenosine, or LNA guanosine; a nucleoside having a stabilized terminal 5'-phosphate or phosphatase-resistant analogue of 5'-phosphate, *e.g.*, 5'-methyl phosphonate, 5'-methylenephosphonate, a 5'-methylenephosphonate analog, 5'-*E*-vinyl phosphonate (5'-*E*-VP), 5'-phosphorothioate, and a 5'-*C*-methyl analog; a 2'-*O*-methyl ribonucleoside, *e.g.*, 2'-*O*-methylcytidine, 2'-*O*-methylguanosine, 2'-*O*-methyluridine, 2'-*O*-methylthymidine, and 2'-*O*-methyladenosine; a 5-methylpyrimidine, *e.g.*, 5-methylcytosine; a 5-methylpurine, *e.g.*, 5-methylguanine; 5-methyl-2'-*O*-methylcytidine; 5-methyl-2'-*O*-methylthymidine; 5-methylcytidine; 5-methyluridine; or 5-methyl 2'-deoxycytidine.

[0058] In some embodiments, the IL-34 ASO comprises a 2'-*O*-methyl ribonucleoside, *e.g.*, 2'-*O*-methylcytidine, 2'-*O*-methylguanosine, 2'-*O*-methyluridine, 2'-*O*-methylthymidine, and 2'-*O*-methyladenosine.

[0059] In some embodiments, the IL-34 ASO comprises the nucleotide sequence 5'-CTTTGGGCXGCACACAGCTTC-3' (SEQ ID NO: 7), wherein X is 5-methylcytidine. In some embodiments the IL-34 ASO comprises the nucleotide sequence 5'-TCCATGACCXGGAAGCAGTT-3' (SEQ ID NO: 8), wherein X is 5-methylcytidine.

[0060] In some embodiments, the ASO comprises a 2'-*O*-methyl ("2'-OMe") ribonucleoside.

[0061] In some embodiments, the ASO has a sequence at least 90% identical to 5'-C_xT_xT_xGGGCXGCACCAGC_xT_xT_xC_x-3' (SEQ ID NO: 40), in which C_x is 2'-O-(2-methoxyethyl)cytidine, T_x is 2'-O-(2-methoxyethyl)thymidine, and X is 5-methylcytidine. In some embodiments, the ASO has a sequence at least 95% identical to 5'-C_xT_xT_xGGGCXGCACCAGC_xT_xT_xC_x-3' (SEQ ID NO: 40), in which C_x is 2'-O-(2-methoxyethyl)cytidine, T_x is 2'-O-(2-methoxyethyl)thymidine, and X is 5-methylcytidine. In some embodiments, the ASO has a sequence according to 5'-C_xT_xT_xGGGCXGCACCAGC_xT_xT_xC_x-3' (SEQ ID NO: 40), in which C_x is 2'-O-(2-methoxyethyl)cytidine, T_x is 2'-O-(2-methoxyethyl)thymidine, and X is 5-methylcytidine.

[0062] In some embodiments, the ASO has a sequence at least 90% identical to 5'-C_xT_xT_xTGGGCXGCACCAGCT_xT_xC_x-3' (SEQ ID NO: 41) in which C_x is 2'-O-(2-methoxyethyl)cytidine, T_x is 2'-O-(2-methoxyethyl)thymidine, and X is 5-methylcytidine. In some embodiments, the ASO has a sequence at least 95% identical to 5'-C_xT_xT_xTGGGCXGCACCAGCT_xT_xC_x-3' (SEQ ID NO: 41) in which C_x is 2'-O-(2-methoxyethyl)cytidine, T_x is 2'-O-(2-methoxyethyl)thymidine, and X is 5-methylcytidine. In some embodiments, the ASO has a sequence according to 5'-C_xT_xT_xTGGGCXGCACCAGCT_xT_xC_x-3' (SEQ ID NO: 41) in which C_x is 2'-O-(2-methoxyethyl)cytidine, T_x is 2'-O-(2-methoxyethyl)thymidine, and X is 5-methylcytidine.

[0063] In some embodiments, the ASO has a sequence at least 90% identical to 5'-T_xC_xC_xA_xTGACCXGGAAGCA_xG_xT_xT_x-3' (SEQ ID NO: 42), in which C_x is 2'-O-(2-methoxyethyl)cytidine, A_x is 2'-O-(2-methoxyethyl)adenosine, T_x is 2'-O-(2-methoxyethyl)thymidine, G_x is 2'-O-(2-methoxyethyl)guanosine, and X is 5-methylcytidine. In some embodiments, the ASO has a sequence at least 95% identical to 5'-T_xC_xC_xA_xTGACCXGGAAGCA_xG_xT_xT_x-3' (SEQ ID NO: 42), in which C_x is 2'-O-(2-methoxyethyl)cytidine, A_x is 2'-O-(2-methoxyethyl)adenosine, T_x is 2'-O-(2-methoxyethyl)thymidine, G_x is 2'-O-(2-methoxyethyl)guanosine, and X is 5-methylcytidine. In some embodiments, the ASO has a sequence according to 5'-T_xC_xC_xA_xTGACCXGGAAGCA_xG_xT_xT_x-3' (SEQ ID NO: 42), in which C_x is 2'-O-(2-methoxyethyl)cytidine, A_x is 2'-O-(2-methoxyethyl)adenosine, T_x is 2'-O-(2-methoxyethyl)thymidine, G_x is 2'-O-(2-methoxyethyl)guanosine, and X is 5-methylcytidine.

[0064] In some embodiments, the ASO has a sequence at least 90% identical to 5'-TxCxCxATGACCXGGAAGCAGxTxTx-3' (SEQ ID NO: 43), in which Cx is 2'-O-(2-methoxyethyl)cytidine, Tx is 2'-O-(2-methoxyethyl)thymidine, Gx is 2'-O-(2-methoxyethyl)guanosine, and X is 5-methylcytidine. In some embodiments, the ASO has a sequence at least 95% identical to 5'-TxCxCxATGACCXGGAAGCAGxTxTx-3' (SEQ ID NO: 43), in which Cx is 2'-O-(2-methoxyethyl)cytidine, Tx is 2'-O-(2-methoxyethyl)thymidine, Gx is 2'-O-(2-methoxyethyl)guanosine, and X is 5-methylcytidine. In some embodiments, the ASO has a sequence according to 5'-TxCxCxATGACCXGGAAGCAGxTxTx-3' (SEQ ID NO: 43), in which Cx is 2'-O-(2-methoxyethyl)cytidine, Tx is 2'-O-(2-methoxyethyl)thymidine, Gx is 2'-O-(2-methoxyethyl)guanosine, and X is 5-methylcytidine. In some embodiments, contemplated compositions disclosed herein may include a pharmaceutically acceptable salt.

[0065] In some embodiments described herein, an IL-34 ASO is a gapmer compound. Gapmer compounds are oligonucleotide sequences that include 5' and 3' flanking groups of modified nucleotides (referred to as region A' and C'). These flanking groups of modified nucleotides are thought to protect the internal group of nucleotides (referred to as region B') from nuclease degradation. Together, the 5' and 3' flanking groups and the internal group form a (5' to 3') A'-B'-C' structure. The internal group of nucleotides is usually from 6 to 10 nucleotides in length. In some embodiments the internal group of oligonucleotides is 10-14 molecules in length. Each 5' or 3' group of flanking nucleotides can be 3, 4, 5, 6, or more nucleotides in length. The 5' and 3' group of flanking nucleotides in a gapmer compound can be the same number of nucleotides in length. Flanking groups of modified nucleotides include, for example, 2'-MOE, 2'-OMe, and LNA nucleotides. Gapmer compound sequences can also incorporate modifications including modified internucleoside linkages (for example, phosphorothioate linkages), 2'-MOEs, 2'-OMes, LNA nucleosides, PNAs, 5-methylcytidine, and other chemically modified nucleosides described herein.

[0066] In some embodiments, the gapmer is from 15-20, 15-25, 15-30, 15-35, 20-25, 20-30, 20-35, 25-30, 25-35, or 30-35 nucleotides in length. In particular embodiments, the antisense oligonucleotide is from 15-25 nucleotides in length. In particular embodiments, the antisense oligonucleotide is from 20-25 nucleotides in length. In some embodiments, the antisense oligonucleotide is 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, or 35 nucleotides in length. In some embodiments the gapmer is 20 nucleotides in length.

[0067] In some embodiments, region A' consists of 3 or 4 nucleotide analogues, such as 2'-MOE, region B' consists of 12 or 14 DNA bases, and region C' consists of 3 or 4 nucleotide analogues such as 2'-MOE. Such gapmer designs include (A'-B'-C') 3-12-3, 3-14-3, 4-12-4, and 4-14-4.

[0068] In some embodiments, region A' consists of 5 or 6 nucleotide analogues, such as 2'-MOE, region B' consists of 8 or 10 DNA bases, and region C' consists of 5 or 6 nucleotide analogues such as 2'-MOE. Such gapmer designs include (A'-B'-C') 5-10-5 and 6-8-6.

[0069] In some embodiments the ASO comprises a gapmer of the structure 5'-NxNxNxNNNNNNNNNNNNNNNNNxNxNx-3', wherein N comprises any nucleotide or chemically modified nucleotide and Nx comprises a 2'-O-methyl ("2'-OMe") ribonucleoside. In some embodiments the ASO comprises a gapmer of the structure 5'-NxNxNxNxNNNNNNNNNNNNNNNxNxNxNx-3', wherein N comprises any nucleotide or chemically modified nucleotide and Nx comprises a 2'-O-methyl ("2'-OMe") ribonucleoside. In some embodiments the ASO comprises a gapmer of the structure 5'-NxNxNxNxNxNNNNNNNNNNNNNxNxNxNxNx-3', wherein N comprises any nucleotide or chemically modified nucleotide and Nx comprises a 2'-O-methyl ("2'-OMe") ribonucleoside. In some embodiments the ASO comprises a gapmer of the structure 5'-NxNxNxNxNxNxNNNNNNNNNNNNNxNxNxNxNxNx-3', wherein N comprises any nucleotide or chemically modified nucleotide and Nx comprises a 2'-O-methyl ("2'-OMe") ribonucleoside.

[0070] In some embodiments the ASO comprises a gapmer of the structure 5'-NxNxNxNNNNNNXNNNNNNNNNNNxNxNx-3', wherein N comprises any nucleotide or chemically modified nucleotide, X comprises a 5-methylcytidine, and Nx comprises a 2'-O-methyl ("2'-OMe") ribonucleoside. In some embodiments the ASO comprises a gapmer of the structure 5'-NxNxNxNxNxNNNNXNNNNNNNNNNNxNxNxNx-3', wherein N comprises any nucleotide or chemically modified nucleotide, X comprises a 5-methylcytidine, and Nx comprises a 2'-O-methyl ("2'-OMe") ribonucleoside. In some embodiments the ASO comprises a gapmer of the structure 5'-NxNxNxNxNNNNNNXNNNNNNNNNNNxNxNxNx-3', wherein N comprises any nucleotide or chemically modified nucleotide, X comprises a 5-methylcytidine,

and Nx comprises a 2'-O-methyl ("2'-OMe") ribonucleoside. In some embodiments the ASO comprises a gapmer of the structure 5'-NxNxNxNxNNNNNNXNNNNNNNxNxNxNx-3', wherein N comprises any nucleotide or chemically modified nucleotide, X comprises a 5-methylcytidine, and Nx comprises a 2'-O-methyl ("2'-OMe") ribonucleoside.

[0071] In some embodiments, an IL-34 ASO gapmer includes 2'-MOE nucleosides and the sequence of the IL-34 ASO is selected from one of the following:

- 5'-CxTxTxTxGGGCXGCACCAGCxTxTxCx-3' (SEQ ID NO: 40), wherein Cx is 2'-O-(2-methoxyethyl)cytidine, Tx is 2'-O-(2-methoxyethyl)thymidine, and X is 5-methylcytidine;
- 5'-CxTxTxTGGGCXGCACCAGCTxTxTx-3' (SEQ ID NO: 41), wherein Cx is 2'-O-(2-methoxyethyl)cytidine, Tx is 2'-O-(2-methoxyethyl)thymidine, and X is 5-methylcytidine;
- 5'-TxCxCxAxTGACCXGGAAGCAxGxTxTx-3' (SEQ ID NO: 42), wherein Cx is 2'-O-(2-methoxyethyl)cytidine, Ax is 2'-O-(2-methoxyethyl)adenosine, Tx is 2'-O-(2-methoxyethyl)thymidine, Gx is 2'-O-(2-methoxyethyl)guanosine, and X is 5-methylcytidine; and
- 5'-TxCxCxATGACCXGGAAGCAGxTxTx-3' (SEQ ID NO: 43), wherein Cx is 2'-O-(2-methoxyethyl)cytidine, Tx is 2'-O-(2-methoxyethyl)thymidine, Gx is 2'-O-(2-methoxyethyl)guanosine, and X is 5-methylcytidine,

or a pharmaceutically acceptable salt thereof.

[0072] In some embodiments, the first, seventeenth, and twentieth nucleosides of SEQ ID NO: 3 (relative to the 5' end) are each independently substituted with 2'-O-(2-methoxyethyl)cytidine; the second, third, fourth, eighteenth, and nineteenth nucleosides of SEQ ID NO: 3 are each independently substituted with 2'-O-(2-methoxyethyl)thymidine; and the ninth nucleoside of SEQ ID NO: 3 is substituted with 5-methylcytidine, resulting in an IL-34 ASO comprising the nucleotide sequence of SEQ ID NO: 40.

[0073] In some embodiments, the first and twentieth nucleosides of SEQ ID NO: 3 (relative to the 5' end) are each independently substituted with 2'-O-(2-methoxyethyl)cytidine; the second, third, eighteenth, and nineteenth nucleosides of SEQ ID NO: 3 are each independently substituted with 2'-O-(2-methoxyethyl)thymidine; and the ninth nucleoside of SEQ ID NO: 3 is

substituted with 5-methylcytidine, resulting in an IL-34 ASO comprising the nucleotide sequence of SEQ ID NO: 41.

[0074] In some embodiments, an IL-34 ASO comprises the nucleotide sequence of SEQ ID NO: 5, wherein at least one nucleoside is a 2'-MOE nucleoside. For example, in some embodiments, the first, nineteenth, and twentieth nucleosides of SEQ ID NO: 5 (relative to the 5' end) are each independently substituted with 2'-*O*-(2-methoxyethyl)thymidine; the second, and third nucleosides of SEQ ID NO: 5 are each independently substituted with 2'-*O*-(2-methoxyethyl)cytidine; the fourth and seventeenth nucleosides of SEQ ID NO: 5 are each independently substituted with 2'-*O*-(2-methoxyethyl)adenosine; the eighteenth nucleoside of SEQ ID NO: 5 is substituted with 2'-*O*-(2-methoxyethyl)guanosine; and the tenth nucleoside of SEQ ID NO: 5 is substituted with 5-methylcytidine, resulting in an IL-34 ASO comprising the nucleotide sequence of SEQ ID NO: 42.

[0075] In some embodiments, an IL-34 ASO comprises the nucleotide sequence of SEQ ID NO: 5, wherein at least one nucleoside is a 2'-MOE nucleoside. For example, in some embodiments, the first, nineteenth, and twentieth nucleosides of SEQ ID NO: 5 (relative to the 5' end) are each independently substituted with 2'-*O*-(2-methoxyethyl)thymidine; the second, and third nucleosides of SEQ ID NO: 5 are each independently substituted with 2'-*O*-(2-methoxyethyl)cytidine; the eighteenth nucleoside of SEQ ID NO: 5 is substituted with 2'-*O*-(2-methoxyethyl)guanosine; and the tenth nucleoside of SEQ ID NO: 5 is substituted with 5-methylcytidine, resulting in an IL-34 ASO comprising the nucleotide sequence of SEQ ID NO: 43.

[0076] For example, in some embodiments, an IL-34 ASO comprises the nucleotide sequence of SEQ ID NO: 3, wherein at least one nucleoside is a 2'-MOE nucleoside. For example, in some embodiments, the first, seventeenth, and twentieth nucleosides of SEQ ID NO: 3 (relative to the 5' end) are each independently substituted with 2'-*O*-(2-methoxyethyl)cytidine; the second, third, fourth, eighteenth, and nineteenth nucleosides of SEQ ID NO: 3 are each independently substituted with 2'-*O*-(2-methoxyethyl)thymidine; the fifth and sixteenth nucleosides of SEQ ID NO: 3 are each independently substituted with 2'-*O*-(2-methoxyethyl)guanosine; and the ninth nucleoside of SEQ ID NO: 3 is substituted with 5-methylcytidine, resulting in an IL-34 ASO comprising the nucleotide sequence of SEQ ID NO: 16.

[0077] In some embodiments, the first, seventeenth, and twentieth nucleosides of SEQ ID NO: 3 (relative to the 5' end) are each independently substituted with 2'-*O*-(2-methoxyethyl)cytidine; the second, third, fourth, eighteenth, and nineteenth nucleosides of SEQ ID NO: 3 are each independently substituted with 2'-*O*-(2-methoxyethyl)thymidine; the fifth, sixth and sixteenth nucleosides of SEQ ID NO: 3 are each independently substituted with 2'-*O*-(2-methoxyethyl)guanosine; the fifteenth nucleoside of SEQ ID NO: 3 is substituted with 2'-*O*-(2-methoxyethyl)adenosine; and the ninth nucleoside of SEQ ID NO: 3 is substituted with 5-methylcytidine, resulting in an IL-34 ASO comprising the nucleotide sequence of SEQ ID NO: 18.

[0078] In some embodiments, the first, eighth, eleventh, thirteenth, fourteenth, seventeenth, and twentieth nucleosides of SEQ ID NO: 3 (relative to the 5' end) are each independently substituted with 2'-*O*-(2-methoxyethyl)cytidine; all thymidines of SEQ ID NO: 3 are each independently substituted with 2'-*O*-(2-methoxyethyl)thymidine; all guanosines of SEQ ID NO: 3 are each independently substituted with 2'-*O*-(2-methoxyethyl)guanosine; all adenosines of SEQ ID NO: 3 are each independently substituted with 2'-*O*-(2-methoxyethyl)adenosine; and the ninth nucleoside of SEQ ID NO: 3 is substituted with 5-methylcytidine, resulting in an IL-34 ASO comprising the nucleotide sequence of SEQ ID NO: 20.

[0079] In other embodiments, one or more nucleoside(s) in SEQ ID NO: 3 may be substituted with 2'-MOE nucleosides other than those substituted in any one of SEQ ID NO: 16, 18, 20, 40, or 41. In still other embodiments, SEQ ID NO: 3 may be substituted with 2'-MOE nucleosides at one or more positions other than those substituted in any one of SEQ ID NO: 16, 18, 20, 40, or 41.

[0080] In some embodiments, an IL-34 ASO comprises the nucleotide sequence of SEQ ID NO: 5, wherein at least one nucleoside is a 2'-MOE nucleoside. For example, in some embodiments, the first, fifth, nineteenth, and twentieth nucleosides of SEQ ID NO: 5 (relative to the 5' end) are each independently substituted with 2'-*O*-(2-methoxyethyl)thymidine; the second, third, and sixteenth nucleosides of SEQ ID NO: 5 are each independently substituted with 2'-*O*-(2-methoxyethyl)cytidine; the fourth and seventeenth nucleosides of SEQ ID NO: 5 are each independently substituted with 2'-*O*-(2-methoxyethyl)adenosine; the eighteenth nucleoside of SEQ ID NO: 5 is substituted with 2'-*O*-(2-methoxyethyl)guanosine; and the tenth nucleoside of

SEQ ID NO: 5 is substituted with 5-methylcytidine, resulting in an IL-34 ASO comprising the nucleotide sequence of SEQ ID NO: 17.

[0081] In some embodiments, the first, fifth, nineteenth, and twentieth nucleosides of SEQ ID NO: 5 (relative to the 5' end) are each independently substituted with 2'-*O*-(2-methoxyethyl)thymidine; the second, third, and sixteenth nucleosides of SEQ ID NO: 5 are each independently substituted with 2'-*O*-(2-methoxyethyl)cytidine; the fourth and seventeenth nucleosides of SEQ ID NO: 5 are each independently substituted with 2'-*O*-(2-methoxyethyl)adenosine; the sixth, fifteenth, and eighteenth nucleosides of SEQ ID NO: 5 are each independently substituted with 2'-*O*-(2-methoxyethyl)guanosine; and the tenth nucleoside of SEQ ID NO: 5 is substituted with 5-methylcytidine, resulting in an IL-34 ASO comprising the nucleotide sequence of SEQ ID NO: 19.

[0082] In some embodiments, the second, third, eighth, ninth, and sixteenth nucleosides of SEQ ID NO: 5 (relative to the 5' end) are each independently substituted with 2'-*O*-(2-methoxyethyl)cytidine; all thymidines of SEQ ID NO: 5 are each independently substituted with 2'-*O*-(2-methoxyethyl)thymidine; all guanosines of SEQ ID NO: 5 are each independently substituted with 2'-*O*-(2-methoxyethyl)guanosine; all adenosines of SEQ ID NO: 5 are each independently substituted with 2'-*O*-(2-methoxyethyl)adenosine; and the tenth nucleoside of SEQ ID NO: 5 is substituted with 5-methylcytidine, resulting in an IL-34 ASO comprising the nucleotide sequence of SEQ ID NO: 21.

[0083] In other embodiments, one or more nucleoside(s) in SEQ ID NO: 5 may be substituted with 2'-MOE nucleosides other than those substituted in any one of SEQ ID NO: 17, 19, 21, 42, or 43. In still other embodiments, SEQ ID NO: 5 may be substituted with 2'-MOE nucleosides at one or more positions other than those substituted in any one of SEQ ID NO: 17, 19, 21, 42, or 43.

[0084] In some embodiments, an ASO described herein includes at least one modified internucleoside linkage. As used herein, "internucleoside linkage" means the connection between the 3' position of a nucleoside and the 5' position of an adjacent nucleoside. A "modified internucleoside linkage" refers to a connection between the 3' position of a nucleoside and the 5' position of an adjacent nucleoside that is not natural. In some embodiments, a modified internucleoside linkage includes, but is not limited to: a phosphorothioate linkage; a phosphorodithioate linkage, a phosphotriester linkage, an alkylphosphonate linkage, an

aminoalkylphosphotriester linkage, an alkylene phosphonate linkage, a phosphinate linkage, a phosphoramidate linkage, an aminoalkylphosphoramidate linkage, a thiophosphoramidate linkage, a thionoalkylphosphonate linkage, a thionoalkylphosphotriester linkage, a thiophosphate linkage, a selenophosphate linkage, or a boranophosphate linkage.

[0085] In some embodiments, ASOs described herein includes at least one chemically modified nucleoside and at least one modified internucleoside linkage. In some embodiments, ASOs described herein includes at least one chemically modified nucleoside or at least one modified internucleoside linkage.

[0086] IL-34 ASOs described herein, can include chemically modified nucleosides, including modified ribonucleosides and modified deoxyribonucleosides. Chemically modified nucleosides include, but are not limited to, 2'-O-(2-methoxyethyl) modifications, for example, 2'-O-(2-methoxyethyl)guanosine, 2'-O-(2-methoxyethyl)adenosine, 2'-O-(2-methoxyethyl)cytosine, 2'-O-(2-methoxyethyl)uridine, and 2'-O-(2-methoxyethyl)thymidine. Chemically modified nucleosides also include, but are not limited to, locked nucleic acid (LNA), 2'-O-methyl, 2'-fluoro, and 2'-fluoro- β -D-arabinonucleotide (FANA) modifications. Chemically modified nucleosides that can be included in IL-34 ASOs described herein are described in Johannes and Lucchino, (2018) "Current Challenges in Delivery and Cytosolic Translocation of Therapeutic RNAs" *Nucleic Acid Ther.* 28(3): 178–93; Rettig and Behlke, (2012) "Progress toward in vivo use of siRNAs-II" *Mol Ther* 20:483–512; and Khvorova and Watts, (2017) "The chemical evolution of oligonucleotide therapies of clinical utility" *Nat Biotechnol.*, 35(3):238-48, the contents of each of which are incorporated by reference herein. In certain embodiments, the disclosure provides mixed modalities of polynucleotide complementary to IL-34, or a portion thereof, *e.g.*, a combination of an IL-34 peptide nucleic acid (PNA) and an IL-34 locked nucleic acid (LNA).

[0087] IL-34 ASOs described herein can include chemical modifications that promote stabilization of an oligonucleotide's terminal 5'-phosphate and phosphatase-resistant analogs of 5'-phosphate. Chemical modifications that promote oligonucleotide terminal 5'-phosphate stabilization or which are phosphatase-resistant analogs of 5'-phosphate include, but are not limited to, 5'-methyl phosphonate, 5'-methylenephosphonate, 5'-methylenephosphonate analogs, 5'-E-vinyl phosphonate (5'-E-VP), 5'-phosphorothioate, and 5'-C-methyl analogs. Chemical modifications that promote ASO terminal 5'-phosphate stabilization and phosphatase-resistant

analogues of 5'-phosphate are described in Khvorova and Watts, (2017) "The chemical evolution of oligonucleotide therapies of clinical utility" *Nat Biotechnol.*, 35(3):238-48, the contents of which are incorporated by reference herein.

[0088] In some embodiments described herein, IL-34 ASOs described herein can include chemically modified nucleosides, including 2'-*O*-methyl ribonucleosides, for example, 2'-*O*-methylcytidine, 2'-*O*-methylguanosine, 2'-*O*-methyluridine, 2'-*O*-methylthymidine, and/or 2'-*O*-methyladenosine. IL-34 ASOs described herein can include one or more chemically modified nucleosides, wherein the chemically modified nucleosides comprise bases, including a 5-methylpyrimidine, for example, 5-methylcytosine, and/or a 5-methylpurine, for example, 5-methylguanine. IL-34 ASOs described herein can include any of the following chemically modified nucleosides: 5-methyl-2'-*O*-methylcytidine, 5-methyl-2'-*O*-methylthymidine, 5-methylcytidine, 5-methyluridine, and/or 5-methyl 2'-deoxycytidine.

[0089] In some embodiments, IL-34 ASOs described herein include an IL-34 ASO that includes the nucleotide sequence: 5'-CTCACCAAGACCCACAG-3' (SEQ ID NO: 1), wherein at least one nucleoside is a chemically modified nucleoside and/or at least one linkage is a modified internucleoside linkage; 5'-GGCTTTGGGCCGCACCAGCT-3' (SEQ ID NO: 2), wherein at least one nucleoside is a chemically modified nucleoside and/or at least one linkage is a modified internucleoside linkage; 5'-CTTTGGGCCGCACCAGCTTC-3' (SEQ ID NO: 3), wherein at least one nucleoside is a chemically modified nucleoside and/or at least one linkage is a modified internucleoside linkage; 5'-TGGGCCGCACCAGCTTCAGG-3' (SEQ ID NO: 4), wherein at least one nucleoside is a chemically modified nucleoside and/or at least one linkage is a modified internucleoside linkage; 5'-TCCATGACCCGGAAGCAGTT-3' (SEQ ID NO: 5), wherein at least one nucleoside is a chemically modified nucleoside and/or at least one linkage is a modified internucleoside linkage; or 5'-TGTTTCATGTACTGAAG-3' (SEQ ID NO: 6), wherein at least one nucleoside is a chemically modified nucleoside and/or at least one linkage is a modified internucleoside linkage, or a pharmaceutically acceptable salt thereof.

[0090] In some embodiments, an IL-34 ASO comprises the nucleotide sequence of SEQ ID NO: 3, wherein at least one nucleoside is a chemically modified nucleoside. For example, in some embodiments, the third cytidine (from the 5' end) of SEQ ID NO: 3 may be substituted with a chemically modified nucleoside. In some embodiments, the third cytidine (from the 5' end) of SEQ ID NO: 3 is substituted for 5-methylcytidine, resulting in an IL-34 ASO comprising

the nucleotide sequence of SEQ ID NO: 7. In some embodiments, a nucleoside other than the third cytidine (from the 5' end) of SEQ ID NO: 3 is substituted with a chemically modified nucleoside, resulting in an IL-34 ASO having a nucleotide sequence other than SEQ ID NO: 7. In some embodiments, the third cytidine (from the 5' end) of SEQ ID NO: 3 is substituted with a chemically modified nucleoside that is not 5-methylcytidine, resulting in an IL-34 ASO having a nucleotide sequence other than SEQ ID NO: 7.

[0091] In some embodiments, an IL-34 ASO comprises the nucleotide sequence of SEQ ID NO: 5, wherein at least one nucleoside is a chemically modified nucleoside. For example, in some embodiments, the fifth cytidine (from the 5' end) of SEQ ID NO: 5 may be substituted with a chemically modified nucleoside. In some embodiments the fifth cytidine (from the 5' end) of SEQ ID NO: 5 is substituted for 5-methylcytidine, resulting in an IL-34 ASO comprising the nucleotide sequence of SEQ ID NO: 8. In some embodiments a nucleoside other than the fifth cytidine (from the 5' end) of SEQ ID NO: 3 is substituted with a chemically modified nucleoside, resulting in an IL-34 ASO having a nucleotide sequence other than SEQ ID NO: 8. In some embodiments, the fifth cytidine (from the 5' end) of SEQ ID NO: 3 is substituted with a chemically modified nucleoside that is not 5-methylcytidine, resulting in an IL-34 ASO having a nucleotide sequence other than SEQ ID NO: 8.

[0092] IL-34 ASOs described herein also include IL-34 ASOs that include one or more LNA nucleotides. For example, IL-34 ASOs described herein include an IL-34 ASO that includes the nucleotide sequence of any of the following:

- 5'-cttTGGGCXGCACCAGCttc-3' (SEQ ID NO: 9), wherein c is LNA cytidine, t is LNA thymidine, and X is 5-methylcytidine;
- 5'-ctttGGGCXGCACCAGcttc-3' (SEQ ID NO: 10), wherein c is LNA cytidine, t is LNA thymidine, and X is 5-methylcytidine;
- 5'-cttTGGGCcgCACCAGCttc-3' (SEQ ID NO: 11), wherein c is LNA cytidine, t is LNA thymidine, and g is LNA guanosine;
- 5'-cttTGGGCcGCACCAGCttc-3' (SEQ ID NO: 12), wherein c is LNA cytidine and t is LNA thymidine;
- 5'-ggcXGCACCAGCttc-3' (SEQ ID NO: 13), wherein c is LNA cytidine, t is LNA thymidine, g is LNA guanosine, and X is 5-methylcytidine;

- 5'-cttTGGGCXGCACcag-3' (SEQ ID NO: 14), wherein c is LNA cytidine, t is LNA thymidine, g is LNA guanosine, a is LNA adenosine, and X is 5-methylcytidine; and
- 5'-tgaCCXGGAAGCAgtt-3' (SEQ ID NO: 15), wherein a is LNA adenosine, t is LNA thymidine, g is LNA guanosine, and X is 5-methylcytidine,

or a pharmaceutically acceptable salt thereof.

[0093] In some embodiments, an IL-34 ASO comprises the nucleotide sequence of SEQ ID NO: 3, wherein at least one nucleoside is an LNA. For example, in some embodiments, the first and twentieth nucleosides of SEQ ID NO: 3 (relative to the 5' end) are each independently substituted with LNA cytidine; the second, third, eighteenth, and nineteenth nucleosides of SEQ ID NO: 3 are each independently substituted with LNA thymidine; and the ninth nucleoside of SEQ ID NO: 3 is substituted with 5-methylcytidine, resulting in an IL-34 ASO comprising the nucleotide sequence of SEQ ID NO: 9.

[0094] In another embodiment, the first, seventeenth, and twentieth nucleosides of SEQ ID NO: 3 (relative to the 5' end) are each independently substituted with LNA cytidine; the second, third, fourth, eighteenth, and nineteenth nucleosides of SEQ ID NO: 3 are each independently substituted with LNA thymidine; and the ninth nucleoside of SEQ ID NO: 3 is substituted with 5-methylcytidine, resulting in an IL-34 ASO comprising the nucleotide sequence of SEQ ID NO: 10.

[0095] In another embodiment, the first, ninth, and twentieth nucleosides of SEQ ID NO: 3 (relative to the 5' end) are each independently substituted with LNA cytidine; the second, third, eighteenth, and nineteenth nucleosides of SEQ ID NO: 3 are each independently substituted with LNA thymidine; and the tenth nucleoside of SEQ ID NO: 3 is substituted with LNA guanosine, resulting in an IL-34 ASO comprising the nucleotide sequence of SEQ ID NO: 11.

[0096] In yet another embodiment, the first, ninth, and twentieth nucleosides of SEQ ID NO: 3 (relative to the 5' end) are each independently substituted with LNA cytidine; and the second, third, eighteenth, and nineteenth nucleosides of SEQ ID NO: 3 are each independently substituted with LNA thymidine, resulting in an IL-34 ASO comprising the nucleotide sequence of SEQ ID NO: 12.

[0097] In other embodiments, one or more nucleoside(s) in SEQ ID NO: 3 may be substituted with LNA nucleotides other than those substituted in any one of SEQ ID NO: 9, 10,

11, or 12. In other embodiments, SEQ ID NO: 3 may be substituted with LNA nucleotides at one or more positions other than those substituted in any one of SEQ ID NO: 9, 10, 11, or 12.

[0098] In particular embodiments, an IL-34 ASO includes 2'-MOE nucleosides and the sequence of the IL-34 ASO is selected from one of the following:

- 5'-CxTxTxTxGxGGCXGCACCAGxCxTxTxTx-3' (SEQ ID NO: 16), wherein Cx is 2'-O-(2-methoxyethyl)cytidine, Tx is 2'-O-(2-methoxyethyl)thymidine, Gx is 2'-O-(2-methoxyethyl)guanosine, and X is 5-methylcytidine;
- 5'-TxCxCxAxTxGACCXGGAAGCxAxGxTxTx-3' (SEQ ID NO: 17), wherein Cx is 2'-O-(2-methoxyethyl)cytidine, Tx is 2'-O-(2-methoxyethyl)thymidine, Gx is 2'-O-(2-methoxyethyl)guanosine, Ax is 2'-O-(2-methoxyethyl)adenosine, and X is 5-methylcytidine;
- 5'-CxTxTxTxGxGxGCXGCACCAGxCxTxTxTx-3' (SEQ ID NO: 18), wherein Cx is 2'-O-(2-methoxyethyl)cytidine, Tx is 2'-O-(2-methoxyethyl)thymidine, Gx is 2'-O-(2-methoxyethyl)guanosine, Ax is 2'-O-(2-methoxyethyl)adenosine, and X is 5-methylcytidine;
- 5'-TxCxCxAxTxGxACCXGGAAGxCxAxGxTxTx-3' (SEQ ID NO: 19), wherein Cx is 2'-O-(2-methoxyethyl)cytidine, Tx is 2'-O-(2-methoxyethyl)thymidine, Gx is 2'-O-(2-methoxyethyl)guanosine, Ax is 2'-O-(2-methoxyethyl)adenosine, and X is 5-methylcytidine;
- 5'-CxTxTxTxGxGxGxCxXGxCxAxCxCxAxGxCxTxTxTx-3' (SEQ ID NO: 20), wherein Cx is 2'-O-(2-methoxyethyl)cytidine, Tx is 2'-O-(2-methoxyethyl)thymidine, Gx is 2'-O-(2-methoxyethyl)guanosine, Ax is 2'-O-(2-methoxyethyl)adenosine, and X is 5-methylcytidine; and
- 5'-TxCxCxAxTxGxGxCxXGxCxAxGxCxAxGxCxTxTxTx-3' (SEQ ID NO: 21), wherein Cx is 2'-O-(2-methoxyethyl)cytidine, Tx is 2'-O-(2-methoxyethyl)thymidine, Gx is 2'-O-(2-methoxyethyl)guanosine, Ax is 2'-O-(2-methoxyethyl)adenosine, and X is 5-methylcytidine,

or a pharmaceutically acceptable salt thereof.

[0099] Additionally, in some embodiments, an IL-34 ASO includes 2'-OMe nucleosides and the sequence of the IL-34 ASO is selected from one of the following:

- 5'-CyTyTyTyGyGGCXGCACCAGyCyTyTyCy-3' (SEQ ID NO: 22), wherein Cy is 2'-*O*-methylcytidine, Ty is 2'-*O*-methylthymidine, Gy is 2'-*O*-methylguanosine, and X is 5-methylcytidine; and
- 5'-TyCyCyAyTyGACCXGGAAGCyAyGyTyTy-3' (SEQ ID NO: 23), wherein Cy is 2'-*O*-methylcytidine, Ty is 2'-*O*-methylthymidine, Gy is 2'-*O*-methylguanosine, Ay is 2'-*O*-methyladenosine, and X is 5-methylcytidine,

or a pharmaceutically acceptable salt thereof.

[0100] For example, in some embodiments, an IL-34 ASO comprises the nucleotide sequence of SEQ ID NO: 3, wherein at least one nucleoside is a 2'-OMe nucleoside. For example, in some embodiments, the first, seventeenth and twentieth nucleosides of SEQ ID NO: 3 (relative to the 5' end) are each independently substituted with 2'-*O*-methylcytidine; the second, third, fourth, eighteenth, and nineteenth nucleosides of SEQ ID NO: 3 are each independently substituted with 2'-*O*-methylthymidine; the fifth and sixteenth nucleosides of SEQ ID NO: 3 are each independently substituted with 2'-*O*-methylguanosine; and the ninth nucleoside of SEQ ID NO: 3 is substituted with 5-methylcytidine, resulting in an IL-34 ASO comprising the nucleotide sequence of SEQ ID NO: 22.

[0101] In other embodiments, one or more nucleoside(s) in SEQ ID NO: 3 may be substituted with 2'-OMe nucleosides other than those substituted in SEQ ID NO: 22. In still other embodiments, SEQ ID NO: 3 may be substituted with 2'-OMe nucleosides at one or more positions other than those substituted in SEQ ID NO: 22.

[0102] In some embodiments, an IL-34 ASO comprises the nucleotide sequence of SEQ ID NO: 5, wherein at least one nucleoside is a 2'-OMe nucleoside. For example, in some embodiments, the first, fifth, nineteenth, and twentieth nucleosides of SEQ ID NO: 5 (relative to the 5' end) are each independently substituted with 2'-*O*-methylthymidine; the second, third, and sixteenth nucleosides of SEQ ID NO: 5 are each independently substituted with 2'-*O*-methylcytidine; the fourth and seventeenth nucleosides of SEQ ID NO: 5 are each independently substituted with 2'-*O*-methyladenosine; the eighteenth nucleoside of SEQ ID NO: 5 is substituted with 2'-*O*-methylguanosine; and the tenth nucleoside of SEQ ID NO: 5 is substituted with 5-methylcytidine, resulting in an IL-34 ASO comprising the nucleotide sequence of SEQ ID NO: 23.

[0103] In other embodiments, one or more nucleoside(s) in SEQ ID NO: 5 may be substituted with 2'-OMe nucleosides other than those substituted in SEQ ID NO: 23. In still other embodiments, SEQ ID NO: 5 may be substituted with 2'-OMe nucleosides at one or more positions other than those substituted in SEQ ID NO: 23.

[0104] IL-34 ASOs described herein can include a phosphate backbone where one or more of the oligonucleoside linkages is a modified internucleoside linkage. For example, in some embodiments, an IL-34 ASO described herein may comprise at least one phosphate linkage. IL-34 ASOs described herein may include one or more modified internucleoside linkages selected from the group consisting of a phosphorothioate linkage, a phosphorodithioate linkage, a phosphotriester linkage, an alkylphosphonate linkage, an aminoalkylphosphotriester linkage, an alkylene phosphonate linkage, a phosphinate linkage, a phosphoramidate linkage, an aminoalkylphosphoramidate linkage, a thiophosphoramidate linkage, a thionoalkylphosphonate linkage, a thionoalkylphosphotriester linkage, a thiophosphate linkage, a selenophosphate linkage, and a boranophosphate linkage. In some embodiments of IL-34 ASOs described herein, at least one internucleoside linkage of the nucleotide sequence is a phosphorothioate linkage. For example, in some embodiments of IL-34 ASOs described herein, one, two, three, or more internucleoside linkages of the nucleotide sequence is a phosphorothioate linkage. In preferred embodiments of IL-34 ASOs described herein, all internucleoside linkages of the nucleotide sequence are phosphorothioate linkages. Thus, in some embodiments, all of the nucleotide linkages of an IL-34 ASO of any of SEQ ID NOs: 1-23 or SEQ ID NOs: 40-43 are phosphorothioate linkages. For example, in some embodiments, all of the nucleotide linkages of an IL-34 ASO of SEQ ID NO: 3 are phosphorothioate linkages. In another example, in some embodiments, all of the nucleotide linkages of an IL-34 ASO of SEQ ID NO: 7 are phosphorothioate linkages. In some embodiments, one or more of the nucleotide linkages of an IL-34 ASO of any of SEQ ID NOs: 1-23 or SEQ ID NOs: 40-43 are phosphorothioate linkages. For example, in some embodiments, one or more of the nucleotide linkages of an IL-34 ASO of SEQ ID NO: 3 are phosphorothioate linkages. In another example, in some embodiments, one or more of the nucleotide linkages of an IL-34 ASO of SEQ ID NO: 7 are phosphorothioate linkages.

[0105] It is contemplated that in some embodiments, a disclosed IL-34 ASO, or a pharmaceutically acceptable salt thereof, may optionally have at least one modified nucleobase,

e.g., 5-methylcytosine, and/or at least one methylphosphonate nucleotide, which is placed, for example, either at only one of the 5' or 3' ends or at both 5' and 3' ends or along the oligonucleotide sequence.

[0106] Contemplated IL-34 ASOs may optionally include at least one modified sugar. For example, the sugar moiety of at least one nucleotide constituting the oligonucleotide is a ribose in which the 2'-OH group may be replaced by any one group selected from the group consisting of OR, R, R'OR, SH, SR, NH₂, NR₂, N₃, CN, F, Cl, Br, and I (wherein R is an alkyl or aryl and R' is an alkylene).

[0107] Antisense oligonucleotides can be designed such that the targeting portion of the incorporated nucleotide sequence of each antisense oligonucleotide is completely or almost completely complementary to the IL-34 mRNA sequence, or portion thereof; or the mRNA sequence of an IL-34 interaction partner or a portion thereof. Incorporation of such complementary or nearly complementary nucleotide sequences allows one to engineer antisense oligonucleotides with a high degree of specificity for a given target. Specificity can be assessed via measurement of parameters such as dissociation constant, or other criteria such as changes in protein or RNA expression levels or other assays that measure IL-34 activity or expression. In some embodiments, the IL-34 ASOs disclosed herein are IL-34 inhibitors.

[0108] In some embodiments, polynucleotides complementary to IL-34, or a portion thereof, disclosed herein are IL-34 inhibitors comprising IL-34 small interfering RNAs (siRNAs).

[0109] Small interfering RNAs (siRNAs) are double-stranded RNA molecules of approximately 20-25 base pairs in length (but which can also be, for example, 18-30 base pairs in length) that take advantage of RNAi machinery (*e.g.*, Drosha and RISC) to bind and target mRNA for degradation. siRNAs are not dependent upon plasmids or vectors for expression, and can generally be delivered directly to a target cell, for instance, by transfection. IL-34 siRNAs are double-stranded RNA sequences that include an RNA sequence complementary to an IL-34 mRNA sequence, and which prevent IL-34 protein translation.

[0110] In some embodiments, the IL-34 siRNA is 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 20-25, 20-30, or 25-30 nucleotides in length. In some embodiments the IL-34 siRNA is from 20-25 nucleotides in length. In some embodiments, the IL-34 siRNA is no more than 20, 25, or 30 nucleotides in length.

[0111] In some embodiments, IL-34 siRNAs described herein include siRNAs that include the nucleotide sequence of any one of SEQ ID NOs: 1-23 or SEQ ID NOs: 40-43, for example, the nucleotide sequence of any one of SEQ ID NOs: 1-8, or a pharmaceutically acceptable salt thereof.

[0112] In some embodiments, the IL-34 comprises modified nucleotides. In some embodiments, one or more cytidines are replaced with 5-methylcytidine.

[0113] In some embodiments, IL-34 siRNAs described herein can include a phosphate backbone where one or more of the oligonucleoside linkages is a modified internucleoside linkage. For example, in some embodiments, an IL-34 siRNAs described herein may comprise at least one phosphate linkage. IL-34 siRNAs described herein may include one or more modified internucleoside linkages selected from the group consisting of a phosphorothioate linkage, a phosphorodithioate linkage, a phosphotriester linkage, an alkylphosphonate linkage, an aminoalkylphosphotriester linkage, an alkylene phosphonate linkage, a phosphinate linkage, a phosphoramidate linkage, an aminoalkylphosphoramidate linkage, a thiophosphoramidate linkage, a thionoalkylphosphonate linkage, a thionoalkylphosphotriester linkage, a thiophosphate linkage, a selenophosphate linkage, and a boranophosphate linkage. In some embodiments of IL-34 siRNAs described herein, at least one internucleoside linkage of the nucleotide sequence is a phosphorothioate linkage. For example, in some embodiments of IL-34 siRNAs described herein, one, two, three, or more internucleoside linkages of the nucleotide sequence is a phosphorothioate linkage. In preferred embodiments of IL-34 siRNAs described herein, all internucleoside linkages of the nucleotide sequence are phosphorothioate linkages. Thus, in some embodiments, all of the nucleotide linkages of an IL-34 siRNAs of any of SEQ ID NOs: 1-23 or SEQ ID NOs: 40-43 are phosphorothioate linkages. For example, in some embodiments, all of the nucleotide linkages of an IL-34 siRNAs of SEQ ID NO: 3 are phosphorothioate linkages. In another example, in some embodiments, all of the nucleotide linkages of an IL-34 siRNAs of SEQ ID NO: 7 are phosphorothioate linkages. In some embodiments, one or more of the nucleotide linkages of an IL-34 siRNAs of any of SEQ ID NOs: 1-23 or SEQ ID NOs: 40-43 are phosphorothioate linkages. For example, in some embodiments, one or more of the nucleotide linkages of an IL-34 siRNAs of SEQ ID NO: 3 are phosphorothioate linkages. In another example, in some embodiments, one or more of the nucleotide linkages of an IL-34 siRNAs of SEQ ID NO: 7 are phosphorothioate linkages.

[0114] Peptide nucleic acids (PNAs) are short, artificially synthesized polymers with a structure that mimics DNA or RNA. PNAs include a backbone composed of repeating N-(2-aminoethyl)-glycine units linked by peptide bonds. IL-34 PNAs described herein can be used as antisense therapeutics that bind to IL-34 RNA sequences with high specificity and inhibit IL-34 gene expression.

[0115] Locked nucleic acids (LNAs) are oligonucleotide sequences that include one or more modified RNA nucleotides in which the ribose moiety is modified with an extra bridge connecting the 2' oxygen and 4' carbon. LNAs are believed to have higher T_m's than analogous oligonucleotide sequences. IL-34 LNAs described herein can be used as antisense therapeutics that bind to IL-34 RNA sequences with high specificity and inhibit IL-34 gene expression.

[0116] Morpholino oligomers are oligonucleotide compounds that include DNA bases attached to a backbone of methylenemorpholine rings linked through phosphorodiamidate groups. Morpholino oligomers of the present invention can be designed to bind to specific IL-34 RNA sequences of interest (for example, IL-34 mRNA or IL-34 pre-mRNA sequences of interest), thereby preventing gene expression. IL-34 morpholino oligomers described herein can be used as antisense therapeutics that bind to IL-34 mRNA sequences with high specificity and inhibit IL-34 gene expression. IL-34 morpholino oligomers described herein can also be used to bind IL-34 pre-mRNA sequences, altering IL-34 pre-mRNA splicing and IL-34 gene expression.

[0117] Small hairpin RNAs (shRNAs) are generally RNA molecules with a hairpin-like structure that can be used to silence gene expression. shRNAs are generally expressed from plasmids encoding the shRNA sequence, and can be expressed from viral vectors to allow lentiviral, adenoviral, or adeno-associated viral expression. Without being bound by theory, it is believed that shRNA inhibits gene expression by taking advantage of RNA interference (RNAi) processes. In brief, the shRNA transcript is processed by Drosha and Dicer, and then loaded onto the RNA-induced silencing complex (RISC), allowing targeting of specific mRNA, and either mRNA degradation or repression of protein translation. IL-34 shRNAs described herein can inhibit gene expression of IL-34.

[0118] MicroRNAs (miRNAs) are small non-coding RNA molecule containing about 22 nucleotides that function in RNA silencing and post-transcriptional regulation of gene expression. miRNAs include sequences that are complementary to portions of an mRNA sequence. miRNAs are produced from long, single-stranded RNA molecules exhibiting highly

specific stem-loop structures. Artificial miRNAs are designed by replacing the mature 21 nucleotide sequence of naturally occurring miRNA sequences with 21 nucleotide sequences from a target, for example, an IL-34 mRNA target.

[0119] In some embodiments, the polynucleotide complementary to IL-34, or a portion thereof, described herein can include chemical modifications that promote stabilization of an oligonucleotide's terminal 5'-phosphate and phosphatase-resistant analogs of 5'-phosphate. Chemical modifications that promote oligonucleotide terminal 5'-phosphate stabilization or which are phosphatase-resistant analogs of 5'-phosphate include, but are not limited to, 5'-methyl phosphonate, 5'-methylenephosphonate, 5'-methylenephosphonate analogs, 5'-*E*-vinyl phosphonate (5'-*E*-VP), 5'-phosphorothioate, and 5'-*C*-methyl analogs. Chemical modifications that promote ASO terminal 5'-phosphate stabilization and phosphatase-resistant analogues of 5'-phosphate are described in Khvorova and Watts, (2017) "The chemical evolution of oligonucleotide therapies of clinical utility" *Nat Biotechnol.*, 35(3):238-48, the contents of which are incorporated by reference herein.

[0120] In some embodiments described herein, polynucleotides complementary to IL-34, or a portion thereof, can include chemically modified nucleosides, for example, 2'-*O*-methyl ribonucleosides, for example, 2'-*O*-methylcytidine, 2'-*O*-methylguanosine, 2'-*O*-methyluridine, 2'-*O*-methylthymidine, and/or 2'-*O*-methyladenosine. Polynucleotides complementary to IL-34, or a portion thereof, described herein can include one or more chemically modified bases, including a 5-methylpyrimidine, for example, 5-methylcytosine, and/or a 5-methylpurine, for example, 5-methylguanine. Polynucleotides complementary to IL-34, or a portion thereof, described herein can include any of the following chemically modified nucleosides: 5-methyl-2'-*O*-methylcytidine, 5-methyl-2'-*O*-methylthymidine, 5-methylcytidine, 5-methyluridine, and/or 5-methyl 2'-deoxycytidine.

[0121] 2'-OMe nucleotides are found naturally in tRNA and other small RNAs. Incorporation of 2'-OMe nucleotides into oligonucleotide sequences prevents nuclease degradation and increases stability against hydrolysis. Incorporation of 2'-OMe modification in oligonucleotides generally increases the T_m of RNA-RNA duplexes by 1-4°C per addition.

[0122] Introduction of a 2'-MOE group generally increases the T_m of the resulting oligonucleotide by about 1.1°C and improves the resistance to degradation by nuclease.

Additionally, 2'-MOE oligos are often used in gapmer compounds to preserve the RNase H-mediated degradation of target mRNA strands.

[0123] Polynucleotides complementary to IL-34, or a portion thereof, described herein can include a phosphate backbone where one or more of the oligonucleoside linkages is a phosphate linkage. Polynucleotides complementary to IL-34, or a portion thereof, described herein may include a modified oligonucleotide backbone, where one or more of the nucleoside linkages of the nucleotide sequence is selected from the group consisting of a phosphorothioate linkage, a phosphorodithioate linkage, a phosphotriester linkage, an alkylphosphonate linkage, an aminoalkylphosphotriester linkage, an alkylene phosphonate linkage, a phosphinate linkage, a phosphoramidate linkage, an aminoalkylphosphoramidate linkage, a thiophosphoramidate linkage, a thionoalkylphosphonate linkage, a thionoalkylphosphotriester linkage, a thiophosphate linkage, a selenophosphate linkage, and a boranophosphate linkage. In some embodiments of polynucleotides complementary to IL-34, or a portion thereof, described herein, at least one internucleoside linkage of the nucleotide sequence is a phosphorothioate linkage. For example, in some embodiments of polynucleotides complementary to IL-34, or a portion thereof, described herein, one, two, three, or more internucleoside linkages of the nucleotide sequence is a phosphorothioate linkage. In some embodiments, polynucleotides complementary to IL-34, or a portion thereof, described herein, all internucleoside linkages of the nucleotide sequence are phosphorothioate linkages. Thus, in some embodiments, all of the nucleotide linkages of a polynucleotide complementary to IL-34, or a portion thereof, of any of SEQ ID NOs: 1-23 or SEQ ID NOs: 40-43 are phosphorothioate linkages. In some embodiments, one or more of the nucleotide linkages of a polynucleotide complementary to IL-34, or a portion thereof, of any of SEQ ID NOs: 1-23 or SEQ ID NOs: 40-43 are phosphorothioate linkages. For example, an IL-34 siRNA described herein, can include 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or more phosphorothioate linkages. In some embodiments, all of the nucleotide linkages of an IL-34 siRNA of any of SEQ ID NOs: 1-8 are phosphorothioate linkages.

Autoimmune and Inflammatory Disease

[0124] Provided herein, in some embodiments, are methods for treating an autoimmune or inflammatory disease, comprising administering a polynucleotide complementary to IL-34, or a portion thereof (*e.g.*, ASOs comprising SEQ ID NO: 40-43, or a pharmaceutically acceptable salt thereof). In some embodiments of the disclosure, an effective amount of a disclosed

polynucleotide complementary to IL-34, or a portion thereof, for example an IL-34 antisense oligonucleotide, may be administered to a patient in need thereof to treat an inflammatory disease, for example, to inhibit inflammatory cytokine production in cells of a patient suffering from an inflammatory disease, and/or to reduce or inhibit an IL-34 mediated inflammatory response. Disclosed herein, in certain embodiments, are methods of treating inflammatory disorders related to IL-34 activities in a patient in need thereof comprising administering to the patient an antisense compound disclosed herein.

[0125] “Inflammatory disease” as used herein, refers to a number of acute and chronic inflammatory disorders including but not limited to inflammatory bowel disease, rheumatoid arthritis, psoriasis, osteoarthritis, pouchitis, diabetes (type I and II), tissue or organ rejection, multiple sclerosis, periodontal inflammation (*e.g.*, periodontitis), pigmented villonodular synovitis, hepatitis, sinusitis, colon cancer (*e.g.*, colorectal cancer, colitis-associated colon cancer, sporadic colorectal cancer), coronary artery disease, Sjogren's syndrome (SS), obesity, chronic inflammation, pulmonary sarcoidosis, skin lesions, a CNS inflammatory disease, or an autoimmune disease.

[0126] Rheumatoid arthritis (RA) is characterized by chronic, synovial inflammation affecting multiple joints that causes inflammation, swelling, and pain in and around the joints. Left untreated, RA can lead to damage to the cartilage in a joint and bone erosion around the damaged cartilage, and leads to extra-articular lesions. RA can also affect other tissues throughout the body and cause problems in organs such as the skin, mouth, lungs, heart, and eyes. In some embodiments, IL-34 levels are elevated in patients with active RA and in experimental models of inflammatory arthritis.

[0127] Provided herein are methods of treating, reducing the risk of developing, or delaying the onset of rheumatoid arthritis in a subject in need thereof comprising administering a disclosed polynucleotide (*e.g.*, a polynucleotide complementary to IL-34, or a portion thereof) or a pharmaceutical composition thereof.

[0128] Osteoarthritis (OA) is a degenerative joint disease and is characterized by is a disease of part or the entire joint involving synovial fibrosis and inflammation. OA is associated with clinical symptoms of pain, hyperalgesia, and stiffness. Pathological changes to the synovium can accelerate the progression of OA.

[0129] Provided herein are methods of treating, reducing the risk of developing, or delaying the onset of osteoarthritis or synovial fibrosis in a subject in need thereof comprising administering a disclosed polynucleotide (*e.g.*, a polynucleotide complementary to IL-34, or a portion thereof) or a pharmaceutical composition thereof.

[0130] In some embodiments, methods of treating skin lesions associated with lupus in a patient in need thereof are provided comprising administering a disclosed polynucleotide or a pharmaceutical composition thereof. In some embodiments, treating skin lesions associated with lupus comprises at least one effect selected from reducing the number of skin lesions, reducing the rate of formation of skin lesions, and reducing the severity of skin lesions. Methods of treating lupus and/or lupus nephritis in a patient suffering therefrom are provided, that include administering a disclosed polynucleotide or a pharmaceutical composition thereof. In some embodiments, methods of slowing the progression of a kidney condition associated with lupus are provided.

[0131] Provided herein are methods of treating, reducing the risk of developing, or delaying the onset of a CNS inflammatory disease in a subject in need thereof comprising administering a disclosed polynucleotide (*e.g.*, a polynucleotide complementary to IL-34, or a portion thereof) or a pharmaceutical composition thereof. CNS inflammatory disease that can be treated in this manner include multiple sclerosis, experimental autoimmune encephalomyelitis, Alzheimer's disease, amyotrophic lateral sclerosis (ALS), and Parkinson's disease.

[0132] Also provided herein are methods of treating, reducing the risk of developing, or delaying the onset of an autoimmune disease in a subject in need thereof comprising administering a disclosed polynucleotide (*e.g.*, a polynucleotide complementary to IL-34, or a portion thereof) or a pharmaceutical composition thereof. The methods include treating a subject with or at risk of developing an autoimmune disease. Autoimmune diseases that can be treated in this manner include rheumatoid arthritis, osteoarthritis, pouchitis, type I diabetes, asthma, chronic obstructive pulmonary disease (COPD), and idiopathic pulmonary fibrosis.

Inflammatory Bowel Disease

[0133] Provided herein, in some embodiments, are methods for treating an inflammatory bowel disease in a patient in need thereof, comprising administering a disclosed polynucleotide (*e.g.*, a polynucleotide complementary to IL-34, or a portion thereof, such as SEQ ID NO: 40-43,

or a pharmaceutically acceptable salt thereof) or a pharmaceutical composition thereof. Disclosed herein, in certain embodiments, are methods comprising administering a disclosed compound. "Inflammatory bowel disease," as used herein, refers to a number of chronic inflammatory diseases including Crohn's disease, inflammatory Crohn's disease, fibrostricturing Crohn's disease, gastroduodenal Crohn's disease, Crohn's (granulomatous) colitis, ulcerative colitis, collagenous colitis, lymphocytic colitis, ischaemic colitis, diversion colitis, Behçet's disease, microscopic colitis, ulcerative proctitis, proctosigmoiditis, jejunoileitis, left-sided colitis, pancolitis, ileocolitis, ileitis, and indeterminate colitis. Crohn's disease and ulcerative colitis are the two most common forms of inflammatory bowel disease. Inflammatory bowel diseases are autoimmune diseases of the digestive system. Crohn's disease may be localized to any portion of the gastrointestinal tract, including the terminal ileum, and may impact all cell types of the gastrointestinal tract. Ulcerative colitis is localized to the colon and rectum, and affects cells of the mucosa only.

[0134] Inflammatory Crohn's disease is characterized by abnormal immune response that causes excess gastrointestinal inflammation. Inflammatory Crohn's disease most often affects the intestinal walls, including portions of the small and/or large intestine, for example, the ileum and the colon. Inflammatory Crohn's disease is also associated with development of thick and swollen tissue, as well as ulcers. Mutations in genes that play a role in immune system function (*e.g.*, NOD2, ATG16L1, IL23R, and IRGM) are associated with Crohn's disease, as are genes associated with proper autophagy.

[0135] Fibrostricturing Crohn's disease is a form of Crohn's disease that includes the occurrence of fibrostenotic lesions/fibrotic strictures of the gastrointestinal tract. Fibrotic strictures are more often associated with Crohn's disease of the ileum, as opposed to the colon. Inflammation generally precedes the development strictures. Strictures can be associated with any of the following symptoms: abdominal cramping, abdominal pain, abdominal bloating and distension, loss of appetite, fatigue, nausea, vomiting, and constipation. More than 50% of patients that suffer from Crohn's disease experience fibrostenosis.

[0136] Fibrotic strictures are associated with cell surface IL-17A receptor expression and increased collagen production, including collagen subtypes I, III, and V. Fibrotic strictures are also associated with co-occurrence of fistulas. Strictures are histologically characterized by the presence of a mixture of inflammatory and mesenchymal cells with deposition of an excess of

extracellular matrix (ECM). Activated mesenchymal cells of the intestine accumulate ECM components, including fibronectin and collagen (for example, collagen types III and IV). Disorganized smooth muscle proliferation and excess ECM deposition are believed to modify the mechanical properties of stenotic bowel tissue, resulting in increased stiffness and narrowing of the intestinal tract associated with fibrotic strictures. The presence of fibrotic strictures results in intestinal tissue abnormalities in both the mucosal, submucosal, and muscular layers of the intestinal wall.

[0137] In some embodiments the cell is an intestinal cell, for example, an intestinal stromal cell, an intestinal epithelial cell, an intestinal stem cell, a secretory cell, an enterocyte, a Goblet cell, an enteroendocrine cell, a Paneth cell, a transit amplifying cell, a microfold cell, a cup cell, or a tuft cell. In some embodiments, the cell forms part of an intestinal fibrostricture. In the aforementioned embodiments, subject can be a subject in need of treatment of an inflammatory disease or fibrosis, for example, inflammatory Crohn's disease or fibrostricturing Crohn's disease. In some embodiments, methods described herein can be used to inhibit expression of one or more collagens in a cell of a subject, for example, a type I collagen, a type II collagen, a type III collagen, a type IV collagen, a type V collagen, a type VI collagen, a type VII collagen, a type VIII collagen, a type IX collagen, a type X collagen, a type XI collagen, a type XII collagen, a type XIII collagen, a type XIV collagen, a type XV collagen, a type XVI collagen, a type XVII collagen, a type XVIII collagen, a type XIX collagen, a type XX collagen, a type XXI collagen, a type XXII collagen, a type XXIII collagen, a type XXIV collagen, a type XXV collagen, a type XXVI collagen, a type XXVII collagen, a type XXVIII collagen, or a type XXIX collagen. For example, in embodiments described herein, the collagen is COL1A1, COL1A2II, COL2A1, COL3A1, COL4A1, COL4A2, COL4A3, COL4A4, COL4A5, COL4A6, COL5A1, COL5A2, COL5A3, COL6A1, COL6A2, COL6A3, COL6A5, COL7A1, COL8A1, COL8A2, COL9A1, COL9A2, COL9A3, COL10A1, COL11A1, COL11A2, COL12A1, COL13A1, COL14A1, COL15A1, COL16A1, COL17A1, COL18A1, COL19A1, COL20A1, COL21A1, COL22A1, COL23A1, COL24A1, COL25A1, EMID2, COL27A1, COL28A1, or COL29A1. In particular embodiments, the collagen is COL1A1 (collagen 1A), COL3A1 (collagen 3A), or a mixture thereof.

[0138] Both environmental and genetic factors are believed to play a role in inflammatory bowel diseases, although the identity of such factors is not well-defined. For example, habitual

smoking doubles the risk of developing Crohn's disease. Environmental components may include alterations in flora of the gut which are affected by exposure to ingested foods and medications.

[0139] Inflammatory bowel diseases are associated with symptoms including abdominal pain, vomiting, diarrhea, rectal bleeding, severe cramps, muscle spasms, weight loss, malnutrition, fever, anemia, skin lesions, joint pain, eye inflammation, liver disorders, arthritis, pyoderma gangrenosum, primary sclerosing cholangitis, and non-thyroidal illness syndrome, and treating these symptoms using a disclosed antisense compound is also contemplated in an embodiment, for example, treating children suffering from ulcerative colitis who may also suffer from growth defects.

Intestinal Fibrosis

[0140] Provided herein, in some embodiments, are methods for treating an intestinal fibrosis in a patient in need thereof, comprising administering a disclosed polynucleotide (*e.g.*, a polynucleotide complementary to IL-34, or a portion thereof, such as SEQ ID NO: 40-43, or a pharmaceutically acceptable salt thereof) or a pharmaceutical composition thereof. Intestinal fibrosis commonly results from the reaction of intestinal tissue to inflammation, such as chronic inflammation caused by inflammatory bowel disease (IBD). In IBD, increased levels of SMAD7 block anti-inflammatory gene expression mediated via TGF- β pathway activation and Smad2/3 phosphorylation. Myofibroblasts exposed to increased TGF- β signaling produce increased amounts of collagen. In the context of Crohn's disease, persistent and chronic inflammation promotes fibrotic processes, resulting in the formation of strictures, including small bowel and colonic strictures.

[0141] Intestinal fibrosis can be identified by any of a number of imaging techniques, such as contrast-enhanced ultrasonography. See, *e.g.*, Quaia *et al.* The value of small bowel wall contrast enhancement after sulfur hexafluoride-filled microbubble injection to differentiate inflammatory from fibrotic strictures in patients with Crohn's disease. *Ultrasound Med. Biol.* 38, 1324–1332 (2012); Nylund *et al.* Quantitative contrast-enhanced ultrasound comparison between inflammatory and fibrotic lesions in patients with Crohn's disease. *Ultrasound Med. Biol.* 39, 1197–1206 (2013); Stidham *et al.* Ultrasound elasticity imaging for detecting intestinal fibrosis and inflammation in rats and humans with Crohn's disease. *Gastroenterology* 141, 819–826

(2011). MRI techniques such as magnetization transfer MRI can also be used. See, *e.g.*, Maccioni *et al.* Value of T2-weighted magnetic resonance imaging in the assessment of wall inflammation and fibrosis in Crohn's disease. *Abdom. Imaging* 37, 944–957 (2012); Adler *et al.* Magnetization transfer helps detect intestinal fibrosis in an animal model of Crohn disease. *Radiology* 259, 127–135 (2011); Pazahr *et al.* Magnetization transfer for the assessment of bowel fibrosis in patients with Crohn's disease: initial experience. *Magn. Reson. Mat. Phys. Biol. Med.* 26, 291–301 (2013).

Fibrosis

[0142] Provided herein, in some embodiments, are methods for treating a fibrosis in a patient in need thereof, comprising administering a disclosed polynucleotide (*e.g.*, a polynucleotide complementary to IL-34, or a portion thereof, such as SEQ ID NO: 40-43, or a pharmaceutically acceptable salt thereof) or a pharmaceutical composition thereof. Disclosed herein, in certain embodiments, are methods of preventing or treating fibrosis, such as hepatic fibrosis, pulmonary fibrosis, synovial fibrosis, and/or intestinal fibrosis, form part of this disclosure. For example, a method of preventing or treating hepatic fibrosis is provided comprising administering to a patient in need thereof a polynucleotide complementary to IL-34, or a portion thereof, disclosed herein. Alternatively, a method of preventing or treating intestinal fibrosis is provided comprising administering to a patient in need thereof a polynucleotide complementary to IL-34, or a portion thereof, disclosed herein. Alternatively, a method of preventing or treating pulmonary fibrosis is provided comprising administering to a patient in need thereof a polynucleotide complementary to IL-34, or a portion thereof, disclosed herein. Alternatively, a method of preventing or treating synovial fibrosis is provided comprising administering to a patient in need thereof a polynucleotide complementary to IL-34, or a portion thereof, disclosed herein.

[0143] Patients treated using an above method may or may not have detectable fibrosis. In some embodiments, the patient has at least about a 5%, 10%, 20%, 30%, 40% or even 50% or more reduction in the amount of fibrosis present in the patient after administering an IL-34 antisense oligonucleotide, after *e.g.*, 1 day, 2 days, 1 week, 1 month or 6 months or more. Administering such a polynucleotide complementary to IL-34, or a portion thereof, for example an IL-34 antisense oligonucleotide, may be on, *e.g.*, at least a daily basis. The compound may be

administered orally. The delay of clinical manifestation of fibrosis in a patient as a consequence of administering a polynucleotide complementary to IL-34, or a portion thereof, for example an IL-34 antisense oligonucleotide, disclosed here may be at least e.g., 6 months, 1 year, 18 months or even 2 years or more as compared to a patient who is not administered a polynucleotide complementary to IL-34, or a portion thereof, such as one disclosed herein.

[0144] A patient in need may have hepatic fibrosis that has developed into cirrhosis. A patient at risk of hepatic fibrosis may include those patients with hepatitis B, hepatitis C or nonalcoholic steatohepatitis (NASH). NASH is included in the spectrum of nonalcoholic fatty liver diseases, including steatosis and cirrhosis. NASH is a component of the metabolic syndrome, which is characterized by obesity, type 2 diabetes mellitus, and dyslipidemia, and can eventually lead to hepatocellular carcinoma.

[0145] Methods of treating disorders associated with hepatic fibrosis are also provided, such as the treatment of at least one of: certain storage diseases and inborn errors of metabolism, such as, alpha 1-antitrypsin deficiency, copper storage diseases (e.g., Wilson's disease), fructosemia, galactosemia, glycogen storage diseases (e.g., Types III, IV, VI, IX and X), iron-overload syndromes (e.g., hemochromatosis), lipid abnormalities (e.g. Gaucher's disease), peroxisomal disorders (e.g., Zellweger syndrome), and tyrosinemia; bacterial infections (e.g., brucellosis); parasitic infections (e.g., echinococcosis); NASH; viral infections (e.g., hepatitis B or hepatitis C, including chronic hepatitis B or C); Budd-Chiari syndrome; heart failure; hepatic veno-occlusive disease; and portal vein thrombosis. Methods of treating congenital hepatic fibrosis are also contemplated. The composition may be administered orally.

[0146] Abuse of drugs and/or alcohol has been implicated in cases of hepatic fibrosis. Contemplated herein are methods of treating hepatic fibrosis in a patient with a history of drug and/or alcohol abuse. For example, a patient with a history of abusing at least one of the following: alcohol, amiodarone, chlorpromazine, isoniazid, methotrexate, methyl dopa, oxyphenisatin and, tolbutamide.

[0147] A patient at risk of intestinal fibrosis may include those patients with ulcerative colitis, inflammatory bowel disease, or Crohn's disease. A patient at risk may also include those patients with an early age at diagnosis of Crohn's or colitis, extensive and/or severe of colonic disease, patients with the presence of primary sclerosing cholangitis, and/or patient's having a family history of cancer.

[0148] Methods of treating disorders associated with intestinal fibrosis are also provided, such as the treatment of at least one of: ulcerative colitis, an inflammatory bowel disease, or Crohn's disease.

[0149] Contemplated herein are methods of preventing or treating renal fibrosis, cardiac fibrosis, endomyocardial fibrosis, idiopathic pulmonary fibrosis, synovial fibrosis, myelofibrosis, retroperitoneal fibrosis, or nephrogenic systemic fibrosis, comprising administering to a patient in need thereof, a pharmaceutical preparation comprising a polynucleotide complementary to IL-34, or a portion thereof, for example an IL-34 antisense oligonucleotide, such as an IL-34 antisense oligonucleotide disclosed herein.

[0150] The polynucleotide complementary to IL-34, or a portion thereof, of the invention, including, for example, IL-34 ASOs, can be used alone or in combination with each other whereby at least two polynucleotides complementary to IL-34, or a portion thereof, of the invention are used together in a single composition or as part of a treatment regimen. The polynucleotides complementary to IL-34, or a portion thereof of the invention may also be used in combination with other drugs for treating drug and/or alcohol abuse, renal fibrosis, cardiac fibrosis, endomyocardial fibrosis, idiopathic pulmonary fibrosis, myelofibrosis, retroperitoneal fibrosis, or nephrogenic systemic fibrosis, drug and/or alcohol abuse.

Cancer

[0151] Provided herein, in some embodiments, are methods for treating cancer in a patient in need thereof, comprising administering a disclosed polynucleotide (*e.g.*, a polynucleotide complementary to IL-34, or a portion thereof, such as SEQ ID NO: 40-43, or a pharmaceutically acceptable salt thereof) or a pharmaceutical composition thereof. In some embodiments, the cancer may be a primary cancer or a metastatic cancer. In some embodiments, the cancer is any cancer that comprises cells expressing IL-34 at an abnormal level. In some embodiments, the cancer is any cancer that is associated with altered IL-34 expression, *e.g.*, elevated IL-34 expression.

Treatment and Evaluation

[0152] A "patient" or "subject" as described herein, refers to any animal at risk for, suffering from or diagnosed for an inflammatory and/or fibrotic disease, including, but not limited to, mammals, primates, and humans. In certain embodiments, the patient may be a non-human

mammal such as, for example, a cat, a dog, or a horse. A patient may be an individual diagnosed with a high risk of developing an inflammatory and/or fibrotic disease, someone who has been diagnosed with an inflammatory and/or fibrotic disease, someone who previously suffered from an inflammatory and/or fibrotic disease, or an individual evaluated for symptoms or indications of an inflammatory and/or fibrotic disease, for example, IL-34 expression signal.

[0153] A “patient in need,” as used herein, refers to a patient suffering from any of the symptoms or manifestations of an inflammatory and/or fibrotic disease, a patient who may suffer from any of the symptoms or manifestations of an inflammatory and/or fibrotic disease, or any patient who might benefit from a method of the disclosure for treating an inflammatory and/or fibrotic disease. A patient in need may include a patient who is diagnosed with a risk of developing an inflammatory and/or fibrotic disease, a patient who has suffered from an inflammatory and/or fibrotic disease in the past, or a patient who has previously been treated for an inflammatory and/or fibrotic disease. Of particular relevance are individuals that suffer from an inflammatory and/or fibrotic disease associated with increased levels of IL-34 expression or activity.

[0154] The terms “treat”, “treatment”, “treating” and the like are used herein to generally mean obtaining a desired pharmacological and/or physiological effect. The effect may be prophylactic in terms of completely or partially preventing a disease or symptom thereof and/or may be therapeutic in terms of partially or completely curing a disease and/or adverse effect attributed to the disease. The term “treatment” as used herein covers any treatment of a disease in a mammal, particularly a human, and includes: (a) preventing the disease from occurring in a subject which may be predisposed to the disease but has not yet been diagnosed as having it; (b) inhibiting the disease, *i.e.* preventing the disease from increasing in severity or scope; (c) relieving the disease, *i.e.* causing partial or complete amelioration of the disease; or (d) preventing relapse of the disease, *i.e.* preventing the disease from returning to an active state following previous successful treatment of symptoms of the disease or treatment of the disease.

[0155] An “effective amount” or a “pharmaceutically effective amount,” as used herein, refers to the amount of an agent that is sufficient to at least partially treat a condition when administered to a patient. The therapeutically effective amount will vary depending on the severity of the condition, the route of administration of the component, and the age, weight, etc. of the patient being treated. Accordingly, an effective amount of a disclosed IL-34 antisense

oligonucleotide is the amount of the IL-34 antisense oligonucleotide necessary to treat an inflammatory and/or fibrotic disease in a patient such that administration of the agent prevents an inflammatory and/or fibrotic disease from occurring in a subject, prevents an inflammatory and/or fibrotic disease progression (*e.g.*, prevents the onset or increased severity of symptoms of inflammatory bowel disease such as rectal bleeding, anemia, or gastrointestinal inflammation), or relieves or completely ameliorates some or all associated symptoms of an inflammatory and/or fibrotic disease, *e.g.*, causes regression of the disease.

[0156] Efficacy of treatment may be evaluated by means of evaluation of gross symptoms associated with an inflammatory and/or fibrotic disease, analysis of tissue histology, biochemical assay, imaging methods such as, for example, magnetic resonance imaging, or other known methods. For instance, efficacy of treatment may be evaluated by analyzing gross symptoms of the disease such as changes in abdominal pain, vomiting, diarrhea, rectal bleeding, cramps, muscle spasms, weight loss, malnutrition, fever, anemia or other aspects of gross pathology associated with an inflammatory disease following administration of a disclosed IL-34 antisense oligonucleotide to a patient suffering from an inflammatory disease.

[0157] Efficacy of treatment may also be evaluated at the tissue or cellular level, for example, by means of obtaining a tissue biopsy (*e.g.*, a gastrointestinal tissue biopsy) and evaluating gross tissue or cell morphology or staining properties. Biochemical assays that examine protein or RNA expression may also be used to evaluate efficacy of treatment. For instance, one may evaluate IL-34, IL-6, IL-8, TNF-alpha, or levels of another protein or gene product indicative of an inflammatory and/or fibrotic disease, inflammatory cytokine production, or an IL-34 mediated inflammatory response in dissociated cells or non-dissociated tissue via immunocytochemical, immunohistochemical, Western blotting, or Northern blotting methods, or methods useful for evaluating RNA levels such as quantitative or semi-quantitative polymerase chain reaction. One may also evaluate the presence or level of expression of useful biomarkers found in fecal matter, plasma, or serum to evaluate disease state and efficacy of treatment.

[0158] In evaluating efficacy of treatment, suitable controls may be chosen to ensure a valid assessment. For instance, one can compare symptoms evaluated in a patient with an inflammatory and/or fibrotic disease following administration of a disclosed polynucleotide complementary to IL-34, or a portion thereof, for example an IL-34 antisense oligonucleotide, to those symptoms in the same patient prior to treatment or at an earlier point in the course of

treatment or in another patient not diagnosed with the inflammatory and/or fibrotic disease. Alternatively, one may compare the results of biochemical or histological analysis of gastrointestinal tissue following administration of a disclosed polynucleotide complementary to IL-34, or a portion thereof, for example an IL-34 antisense oligonucleotide, with those of gastrointestinal tissue from the same patient or from an individual not diagnosed with the inflammatory and/or fibrotic disease or from the same patient prior to administration of the polynucleotide complementary to IL-34, or a portion thereof. Additionally, one may compare blood, serum, cell, or fecal samples following administration of the polynucleotide complementary to IL-34, or a portion thereof, for example an IL-34 antisense oligonucleotide, with comparable samples from an individual not diagnosed with the inflammatory and/or fibrotic disease or from the same patient prior to administration of the polynucleotide complementary to IL-34, or a portion thereof.

[0159] Validation of IL-34 inhibition may be determined by direct or indirect assessment of IL-34 expression levels or activity. For instance, biochemical assays that measure IL-34 protein or RNA expression may be used to evaluate overall IL-34 inhibition. For instance, one may measure IL-34 protein levels in gastrointestinal tissue by Western blot to evaluate overall IL-34 levels. One may also measure IL-34 mRNA levels by means of Northern blot or quantitative polymerase chain reaction to determine overall IL-34 inhibition. One may also evaluate IL-34 protein levels or levels of another protein indicative of IL-34 signaling activity in dissociated cells, non-dissociated tissue, blood, serum, or fecal matter via immunocytochemical or immunohistochemical methods.

[0160] IL-34 inhibition may also be evaluated indirectly by measuring parameters such as macrophage or monocyte generation or proliferation, or measuring alterations in other parameters correlated with changes in IL-34 activity, including MAP kinase phosphorylation and other indicators of signaling activation of the IL-34 receptor – the macrophage colony stimulating factor receptor (M-CSFR-1, also known as MCSFR-1, M-CSFR1, and CSF1R). For instance, one may measure levels of active MAPK1 or MAPK3 phosphorylation in cells of a patient treated with a disclosed polynucleotide complementary to IL-34, or a portion thereof, for example an IL-34 antisense oligonucleotide, as an indication of IL-34 activity in the cells. One may also evaluate the presence or level of expression of useful biomarkers found in plasma, blood, fecal matter or tissue to evaluate efficacy of IL-34 inhibition.

[0161] Methods of treatment disclosed herein include methods of inhibiting inflammatory cytokine production. “Inflammatory cytokine production” refers to the expression of cytokines that initiate and/or promote an inflammatory cytokine response. An “inflammatory cytokine response” refers to an immune response that may be characterized by granulocyte recruitment, lymphocyte recruitment, systemic inflammation (especially of the gastrointestinal tract or a portion or portions thereof), fever, tissue destruction, shock, and/or death. An inflammatory cytokine response may be characterized by binding of individual cytokines to their cognate cell surface receptor (*e.g.*, IL-34 binding to CSF1R) and subsequent cascades of intracellular signaling that alter cell functions and gene expression. Inflammatory cytokines include, but are not limited to IL-1, IL-6, IL-8, IL-34, and TNF-alpha. Expression of inflammatory cytokines may occur in, for example, macrophages, monocytes, propria lamina mononuclear cells, or other cells of the gastrointestinal tract or cells of the immune system. Methods of inhibiting inflammatory cytokine production include methods that reduce expression levels of some or all inflammatory cytokines in a patient suffering from an inflammatory disease. Methods of inhibiting inflammatory cytokine production also include methods that reduce expression levels of some or all inflammatory cytokines in cells of a patient suffering from an inflammatory disease.

[0162] Methods of the disclosure for inhibiting inflammatory cytokine production include methods of reducing or inhibiting an IL-34 mediated inflammatory response. An “IL-34 mediated inflammatory response,” as used herein, refers to an inflammatory response initiated, facilitated, or promoted by IL-34 expression or IL-34 signaling activity. An IL-34 mediated inflammatory response may result in expression of inflammatory cytokines including, but not limited to, IL-34, IL-6, IL-8, or TNF-alpha, and activation of inflammatory cytokine signaling. Additionally, an IL-34 mediated inflammatory response may be characterized by granulocyte recruitment, lymphocyte recruitment, systemic inflammation (especially of the gastrointestinal tract or a portion or portions thereof), fever, tissue destruction, shock, and/or death. An IL-34 mediated inflammatory response may also be characterized by activation of IL-34 signaling, for instance, binding of IL-34 to CSF1R and phosphorylation of downstream MAP kinases. Reducing or inhibiting an IL-34 mediated inflammatory response refers to alleviating any or all of the cellular and systemic changes associated with an IL-34 mediated inflammatory response. For example, a reduction in inflammatory cytokine production, immune cell recruitment, or

tissue inflammation would indicate reducing or inhibiting of an IL-34 mediated inflammatory response.

[0163] The disclosure also provides methods of inhibiting IL-34 in cells of a patient suffering from an inflammatory and/or fibrotic disease. IL-34 may be inhibited in any cell in which IL-34 expression or activity occurs, including cells of the gastrointestinal tract, immune system, and blood. Cells of the gastrointestinal tract (including cells of the intestinal wall, the stomach, duodenum, jejunum, large intestine, small intestine, ileum, colon, rectum and anal canal), include columnar epithelial cells, intestinal stromal cells, mucosal epithelial cells, zymogenic cells, neck mucus cells, parietal cells, gastrin cells, Goblet cells, Paneth cells, oligomucus cells, and villus absorptive cells. Cells of the immune system include leukocytes, phagocytes (*e.g.*, macrophages, neutrophils, and dendritic cells), monocytes, mast cells, eosinophils, basophils, natural killer cells, innate cells, lymphocytes, B cells, and T cells. Blood cells include red blood cells (erythrocytes) and white blood cells (leukocytes, monocytes, and platelets).

Pharmaceutical Compositions and Routes of Administration

[0164] Provided herein, in some embodiments, are pharmaceutical compositions comprising polynucleotides complementary to IL-34, or a portion thereof, disclosed herein, or a pharmaceutically acceptable salt thereof, and/or a pharmaceutically acceptable excipient.

[0165] In some embodiments, the polynucleotide is an IL-34 antisense oligonucleotide (ASO). In some embodiments, the IL-34 ASO is a gapmer ASO. In some embodiments, the IL-34 ASO gapmer includes one or more 2'-MOE nucleosides. In some embodiments, the sequence of the IL-34 ASO is selected from one of the following:

- 5'-CxTxTxTGGGCXGCACCAGCxTxTx-3' (SEQ ID NO: 40), wherein Cx is 2'-O-(2-methoxyethyl)cytidine, Tx is 2'-O-(2-methoxyethyl)thymidine, and X is 5-methylcytidine;
- 5'-CxTxTxTGGGCXGCACCAGCTxTxTx-3' (SEQ ID NO: 41), wherein Cx is 2'-O-(2-methoxyethyl)cytidine, Tx is 2'-O-(2-methoxyethyl)thymidine, and X is 5-methylcytidine;
- 5'-TxTxTxAxTGACCXGGAAGCAxGxTxTx-3' (SEQ ID NO: 42), wherein Cx is 2'-O-(2-methoxyethyl)cytidine, Ax is 2'-O-(2-methoxyethyl)adenosine, Tx is 2'-O-(2-

methoxyethyl)thymidine, Gx is 2'-O-(2-methoxyethyl)guanosine, and X is 5-methylcytidine; and

- 5'-TxCxCxATGACCXGGAAGCAGxTxTx-3' (SEQ ID NO: 43), wherein Cx is 2'-O-(2-methoxyethyl)cytidine, Tx is 2'-O-(2-methoxyethyl)thymidine, Gx is 2'-O-(2-methoxyethyl)guanosine, and X is 5-methylcytidine.

[0166] As used herein the term “pharmaceutical composition” means, for example, a mixture containing a specified amount of a therapeutic compound, *e.g.*, a therapeutically effective amount, of a therapeutic compound in a pharmaceutically acceptable excipient to be administered to a mammal, *e.g.*, a human, in order to treat an inflammatory and/or fibrotic disease.

[0167] In some embodiments, a disclosed pharmaceutical composition is useful for treating or preventing an inflammatory disease (*e.g.*, rheumatoid arthritis or osteoarthritis). In some embodiments, a disclosed pharmaceutical composition is useful for treating or preventing a fibrotic disease. In some embodiments, a disclosed pharmaceutical composition is useful for treating or preventing a cancer.

[0168] As used herein, “pharmaceutically acceptable excipient” refers to a substance that aids the administration of an active agent to and/or absorption by a subject and can be included in the compositions of the present invention without causing a significant adverse toxicological effect on the patient. Non-limiting examples of pharmaceutically acceptable excipients include water, NaCl, normal saline solutions, such as a phosphate buffered saline solution, emulsions (*e.g.*, such as an oil/water or water/oil emulsions), lactated Ringer's, normal sucrose, normal glucose, binders, fillers, disintegrants, lubricants, coatings, sweeteners, flavors, salt solutions (such as Ringer's solution), alcohols, oils, gelatins, carbohydrates such as lactose, amylose or starch, fatty acid esters, hydroxymethylcellulose, polyvinyl pyrrolidone, and colors, and the like. Such preparations can be sterilized and, if desired, mixed with auxiliary agents such as lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, coloring, and/or aromatic substances and the like that do not deleteriously react with polynucleotide complementary to IL-34, or a portion thereof, of the invention. For examples of excipients, *see* Martin, Remington's Pharmaceutical Sciences, 15th Ed., Mack Publ. Co., Easton, PA (1975).

[0169] In one embodiment, a disclosed polynucleotide or a pharmaceutical composition thereof may be administered by one or several routes, including intraarticularly, rectally, topically, parenterally, orally, pulmonarily, intratracheally, intranasally, transdermally, or intraduodenally. The term parenteral as used herein includes subcutaneous injections, intrapancreatic administration, intravenous, intramuscular, intraperitoneal, intrasternal injection or infusion techniques. In some embodiments, a disclosed polynucleotide or a pharmaceutical composition thereof is administered intraarticularly. In some embodiments, a disclosed polynucleotide or a pharmaceutical composition thereof is administered rectally. In some embodiments, a disclosed polynucleotide or a pharmaceutical composition thereof is administered subcutaneously. In some embodiments, a disclosed polynucleotide or a pharmaceutical composition thereof is administered orally. In some embodiments, a disclosed polynucleotide or a pharmaceutical composition thereof may be administered directly to the gastrointestinal system, or specific regions of the gastrointestinal system (*e.g.*, the ileum, colon, or rectum) via parenteral administration.

[0170] Pharmaceutical compositions containing a polynucleotide complementary to IL-34, or a portion thereof, disclosed herein, can be presented in a dosage unit form and can be prepared by any suitable method. A pharmaceutical composition should be formulated to be compatible with its intended route of administration. Useful formulations can be prepared by methods well known in the pharmaceutical art. For example, see *Remington's Pharmaceutical Sciences*, 18th ed. (Mack Publishing Company, 1990).

[0171] Pharmaceutical formulations, for example, are sterile. Sterilization can be accomplished, for example, by filtration through sterile filtration membranes. Where the composition is lyophilized, filter sterilization can be conducted prior to or following lyophilization and reconstitution.

Parenteral Administration

[0172] The pharmaceutical compositions of the disclosure can be formulated for parenteral administration, *e.g.*, formulated for injection via the intravenous, intramuscular, subcutaneous, intralesional, or intraperitoneal routes. The preparation of an aqueous composition, such as an aqueous pharmaceutical composition containing a disclosed polynucleotide complementary to IL-34, or a portion thereof, will be known to those of skill in the art in light of the present

disclosure. Typically, such compositions can be prepared as injectables, either as liquid solutions or suspensions; solid forms suitable for using to prepare solutions or suspensions upon the addition of a liquid prior to injection can also be prepared; and the preparations can also be emulsified.

[0173] The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions; formulations including sesame oil, peanut oil or aqueous propylene glycol; and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases the form must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms, such as bacteria and fungi.

[0174] Solutions of active compounds as free base or pharmacologically acceptable salts can be prepared in water suitably mixed with a surfactant, such as hydroxypropylcellulose. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. In addition, sterile, fixed oils may be employed as a solvent or suspending medium. For this purpose, any bland fixed oil can be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid can be used in the preparation of injectables. The sterile injectable preparation may also be a sterile injectable solution, suspension, or emulsion in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, U.S.P., and isotonic sodium chloride solution. In one embodiment, a disclosed polynucleotide complementary to IL-34, or a portion thereof, may be suspended in a carrier or excipient fluid comprising 1% (w/v) sodium carboxymethylcellulose and 0.1% (v/v) TWEEN™ 80. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

[0175] Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. Generally, dispersions are prepared by incorporating the various sterilized active ingredients into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. Sterile injectable solutions of the disclosure may be prepared by incorporating a disclosed polynucleotide complementary to IL-34, or a portion thereof, for example an IL-34 antisense oligonucleotide, in the required amount of

the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filtered sterilization. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum-drying and freeze-drying techniques which yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof. The injectable formulations can be sterilized, for example, by filtration through a bacteria-retaining filter.

[0176] The preparation of more, or highly concentrated solutions for intramuscular injection is also contemplated. In this regard, the use of DMSO as solvent is preferred as this will result in extremely rapid penetration, delivering high concentrations of the disclosed IL-34 antisense oligonucleotide to a small area.

[0177] Suitable preservatives for use in such a solution include benzalkonium chloride, benzethonium chloride, chlorobutanol, thimerosal and the like. Suitable buffers include boric acid, sodium and potassium bicarbonate, sodium and potassium borates, sodium and potassium 10 carbonate, sodium acetate, sodium biphosphate and the like, in amounts sufficient to maintain the pH at between about pH 6 and pH 8, and for example, between about pH 7 and pH 7.5. Suitable tonicity agents are dextran 40, dextran 70, dextrose, glycerin, potassium chloride, propylene glycol, sodium chloride, and the like, such that the sodium chloride equivalent of the solution is in the range 0.9 plus or minus 0.2%. Suitable antioxidants and stabilizers include sodium bisulfite, sodium metabisulfite, sodium thiosulfite, thiourea and the like. Suitable wetting and clarifying agents include polysorbate 80, polysorbate 20, poloxamer 282 and tyloxapol. Suitable viscosity-increasing agents include dextran 40, dextran 70, gelatin, glycerin, hydroxyethylcellulose, hydroxymethylpropylcellulose, lanolin, methylcellulose, petrolatum, polyethylene glycol, polyvinyl alcohol, polyvinylpyrrolidone, carboxymethylcellulose and the like.

Oral Administration

[0178] In some embodiments, contemplated herein are compositions suitable for oral delivery of a disclosed polynucleotide complementary to IL-34, or a portion thereof, *e.g.*, tablets that include an enteric coating, *e.g.*, a gastro-resistant coating, such that the compositions may deliver the polynucleotide to, *e.g.*, the gastrointestinal tract of a patient. For example, such administration may result in a topical effect, substantially topically applying the polynucleotide

complementary to IL-34, or a portion thereof, directly to an affected portion of the gastrointestinal tract of a patient. Such administration, may, in some embodiments, substantially avoid unwanted systemic absorption of the polynucleotide complementary to IL-34, or a portion thereof.

[0179] For example, a tablet for oral administration is provided that comprises granules (*e.g.*, is at least partially formed from granules) that include a disclosed polynucleotide complementary to IL-34, or a portion thereof, for example an IL-34 antisense oligonucleotide, *e.g.*, an antisense oligonucleotide that includes the nucleotide sequence of any one of SEQ ID NOs: 1-23 or SEQ ID NOs: 40-43, and pharmaceutically acceptable excipients. Such a tablet may be coated with an enteric coating. Contemplated tablets may include pharmaceutically acceptable excipients such as fillers, binders, disintegrants, and/or lubricants, as well as coloring agents, release agents, coating agents, sweetening, flavoring such as wintergreen, orange, xylitol, sorbitol, fructose, and maltodextrin, and perfuming agents, preservatives and/or antioxidants.

[0180] In some embodiments, contemplated pharmaceutical formulations include an intra-granular phase that includes a disclosed polynucleotide complementary to IL-34, or a portion thereof, for example an IL-34 antisense oligonucleotide, *e.g.*, an antisense oligonucleotide that includes the nucleotide sequence of any one of SEQ ID NOs: 1-23 or SEQ ID NOs: 40-43, and a pharmaceutically acceptable salt, and a pharmaceutically acceptable filler. For example, a disclosed polynucleotide complementary to IL-34, or a portion thereof, for example an IL-34 antisense oligonucleotide, and a filler may be blended together, optionally, with other excipients, and formed into granules. In some embodiments, the intragranular phase may be formed using wet granulation, *e.g.*, a liquid (*e.g.*, water) is added to the blended polynucleotide complementary to IL-34, or a portion thereof, and filler, and then the combination is dried, milled and/or sieved to produce granules. One of skill in the art would understand that other processes may be used to achieve an intragranular phase.

[0181] In some embodiments, contemplated formulations include an extra-granular phase, which may include one or more pharmaceutically acceptable excipients, and which may be blended with the intragranular phase to form a disclosed formulation.

[0182] A disclosed formulation may include an intragranular phase that includes a filler. Exemplary fillers include, but are not limited to, cellulose, gelatin, calcium phosphate, lactose, sucrose, glucose, mannitol, sorbitol, microcrystalline cellulose, pectin, polyacrylates, dextrose,

cellulose acetate, hydroxypropylmethyl cellulose, partially pre-gelatinized starch, calcium carbonate, and others including combinations thereof.

[0183] In some embodiments, a disclosed formulation may include an intragranular phase and/or an extragranular phase that includes a binder, which may generally function to hold the ingredients of the pharmaceutical formulation together. Exemplary binders of the disclosure may include, but are not limited to, the following: starches, sugars, cellulose or modified cellulose such as hydroxypropyl cellulose, lactose, pre-gelatinized maize starch, polyvinyl pyrrolidone, hydroxypropyl cellulose, hydroxypropylmethyl cellulose, low substituted hydroxypropyl cellulose, sodium carboxymethyl cellulose, methyl cellulose, ethyl cellulose, sugar alcohols and others including combinations thereof.

[0184] Contemplated formulations, *e.g.*, that include an intragranular phase and/or an extragranular phase, may include a disintegrant such as but are not limited to, starch, cellulose, crosslinked polyvinyl pyrrolidone, sodium starch glycolate, sodium carboxymethyl cellulose, alginates, corn starch, croscellose sodium, crosslinked carboxymethyl cellulose, low substituted hydroxypropyl cellulose, acacia, and others including combinations thereof. For example, an intragranular phase and/or an extragranular phase may include a disintegrant.

[0185] In some embodiments, a contemplated formulation includes an intra-granular phase comprising a disclosed polynucleotide complementary to IL-34, or a portion thereof, for example an IL-34 antisense oligonucleotide, and excipients chosen from: mannitol, microcrystalline cellulose, hydroxypropylmethyl cellulose, and sodium starch glycolate or combinations thereof, and an extra-granular phase comprising one or more of: microcrystalline cellulose, sodium starch glycolate, and magnesium stearate or mixtures thereof.

[0186] In some embodiments, a contemplated formulation may include a lubricant, *e.g.*, an extra-granular phase may contain a lubricant. Lubricants include but are not limited to talc, silica, fats, stearin, magnesium stearate, calcium phosphate, silicone dioxide, calcium silicate, calcium phosphate, colloidal silicon dioxide, metallic stearates, hydrogenated vegetable oil, corn starch, sodium benzoate, polyethylene glycols, sodium acetate, calcium stearate, sodium lauryl sulfate, sodium chloride, magnesium lauryl sulfate, talc, and stearic acid.

[0187] In some embodiments, the pharmaceutical formulation comprises an enteric coating. Generally, enteric coatings create a barrier for the oral medication that controls the location at which the drug is absorbed along the digestive track. Enteric coatings may include a polymer that

disintegrates at different rates according to pH. Enteric coatings may include for example, cellulose acetate phthalate, methyl acrylate-methacrylic acid copolymers, cellulose acetate succinate, hydroxylpropylmethyl cellulose phthalate, methyl methacrylate-methacrylic acid copolymers, ethylacrylate-methacrylic acid copolymers, methacrylic acid copolymer type C, polyvinyl acetate-phthalate, and cellulose acetate phthalate.

[0188] Exemplary enteric coatings include Opadry[®] AMB, Acryl-EZE[®], Eudragit[®] grades. In some embodiments, an enteric coating may comprise about 5% to about 10%, about 5% to about 20%, 8 to about 15%, about 8% to about 20%, about 10% to about 20%, or about 12 to about 20%, or about 18% of a contemplated tablet by weight. For example, enteric coatings may include an ethylacrylate-methacrylic acid copolymer.

[0189] For example, in a contemplated embodiment, a tablet is provided that comprises or consists essentially of about 0.5% to about 70%, *e.g.*, about 0.5% to about 10%, or about 1% to about 20%, by weight of a disclosed polynucleotide complementary to IL-34, or a portion thereof, for example an IL-34 antisense oligonucleotide, or a pharmaceutically acceptable salt thereof. Such a tablet may include for example, about 0.5% to about 60% by weight of mannitol, *e.g.*, about 30% to about 50% by weight mannitol, *e.g.*, about 40% by weight mannitol; and/or about 20% to about 40% by weight of microcrystalline cellulose, or about 10% to about 30% by weight of microcrystalline cellulose. For example, a disclosed tablet may comprise an intragranular phase that includes about 30% to about 60%, *e.g.* about 45% to about 65% by weight, or alternatively, about 5 to about 10% by weight of a disclosed polynucleotide complementary to IL-34, or a portion thereof, for example an IL-34 antisense oligonucleotide, about 30% to about 50%, or alternatively, about 5% to about 15% by weight mannitol, about 5% to about 15% microcrystalline cellulose, about 0% to about 4%, or about 1% to about 7% hydroxypropylmethylcellulose, and about 0% to about 4%, *e.g.* about 2% to about 4% sodium starch glycolate by weight.

[0190] In another contemplated embodiment, a pharmaceutical tablet formulation for oral administration of a disclosed polynucleotide complementary to IL-34, or a portion thereof, for example an IL-34 antisense oligonucleotide, comprises an intra-granular phase, wherein the intra-granular phase includes a disclosed polynucleotide complementary to IL-34, or a portion thereof, or a pharmaceutically acceptable salt thereof (such as a sodium salt), and a pharmaceutically acceptable filler, and which may also include an extra-granular phase, that may

include a pharmaceutically acceptable excipient such as a disintegrant. The extra-granular phase may include components chosen from microcrystalline cellulose, magnesium stearate, and mixtures thereof. The pharmaceutical composition may also include an enteric coating of about 12% to 20% by weight of the tablet. For example, a pharmaceutically acceptable tablet for oral use may comprise about .5% to 10% by weight of a disclosed polynucleotide complementary to IL-34, or a portion thereof, for example an IL-34 antisense oligonucleotide, or a pharmaceutically acceptable salt thereof, about 30% to 50% by weight mannitol, about 10% to 30% by weight microcrystalline cellulose, and an enteric coating comprising an ethylacrylate-methacrylic acid copolymer.

[0191] In another example, a pharmaceutically acceptable tablet for oral use may comprise an intra-granular phase, comprising about 5 to about 10% by weight of a disclosed polynucleotide complementary to IL-34, or a portion thereof, for example an IL-34 antisense oligonucleotide, or a pharmaceutically acceptable salt thereof, about 40% by weight mannitol, about 8% by weight microcrystalline cellulose, about 5% by weight hydroxypropylmethyl cellulose, and about 2% by weight sodium starch glycolate; an extra-granular phase comprising about 17% by weight microcrystalline cellulose, about 2% by weight sodium starch glycolate, about 0.4% by weight magnesium stearate; and an enteric coating over the tablet comprising an ethylacrylate-methacrylic acid copolymer.

[0192] In some embodiments the pharmaceutical composition may contain an enteric coating comprising about 13% or about 15%, 16%, 17% or 18% by weight, *e.g.*, AcyrLEZE® (see, *e.g.*, PCT Publication No. WO2010/054826, which is hereby incorporated by reference in its entirety).

[0193] The rate at which point the coating dissolves and the active ingredient is released is its dissolution rate. In an embodiment, a contemplated tablet may have a dissolution profile, *e.g.*, when tested in a USP/EP Type 2 apparatus (paddle) at 100 rpm and 37 °C in a phosphate buffer with a pH of 7.2, of about 50% to about 100% of the polynucleotide complementary to IL-34, or a portion thereof, releasing after about 120 minutes to about 240 minutes, for example after 180 minutes. In another embodiment, a contemplated tablet may have a dissolution profile, *e.g.*, when tested in a USP/EP Type 2 apparatus (paddle) at 100 rpm and 37°C in diluted HCl with a pH of 1.0, where substantially none of the polynucleotide complementary to IL-34, or a portion thereof, is released after 120 minutes. A contemplated tablet, in another embodiment, may have a

dissolution profile, *e.g.*, when tested in USP/EP Type 2 apparatus (paddle) at 100 rpm and 37°C in a phosphate buffer with a pH of 6.6, of about 10% to about 30%, or not more than about 50%, of the polynucleotide complementary to IL-34, or a portion thereof, releasing after 30 minutes.

[0194] Contemplated formulations, *e.g.*, tablets, in some embodiments, when orally administered to the patient may result in minimal plasma concentration of the polynucleotide complementary to IL-34, or a portion thereof, for example an IL-34 antisense oligonucleotide, in the patient. In another embodiment, disclosed formulations, when orally administered to a patient, topically deliver to the colon or rectum of a patient, *e.g.*, to an affected or diseased site of a patient.

[0195] In some embodiments, methods provided herein may further include administering at least one other agent that is directed to treatment of diseases and disorders disclosed herein. In one embodiment, contemplated other agents may be co-administered (*e.g.*, sequentially or simultaneously).

[0196] Agents contemplated include immunosuppressive agents including glucocorticoids, cytostatics, antibodies, agents acting on immunophilins, interferons, opioids, TNF binding proteins, mycophenolate, and small biological agents. For example, contemplated immunosuppressive agents include, but are not limited to: tacrolimus, cyclosporine, pimecrolimus, sirolimus, everolimus, mycophenolic acid, fingolimod, dexamethasone, fludarabine, cyclophosphamide, methotrexate, azathioprine, leflunomide, teriflunomide, anakinra, anti-thymocyte globulin, anti-lymphocyte globulin, muromonab-CD3, afutuzumab, rituximab, teplizumab, efalizumab, daclizumab, basiliximab, adalimumab, infliximab, certolizumab pegol, natalizumab, and etanercept. Other contemplated agents include antibiotics, anti-diarrheals, laxatives, pain relievers, iron supplements, and calcium or vitamin D or B-12 supplements.

Dosage and Frequency of Administration

[0197] Provided herein, in some embodiments, are exemplary dosages of pharmaceutical compositions for use in treating fibrosis, for example pulmonary fibrosis, synovial fibrosis, or intestinal fibrosis in a patient in need thereof, comprising administering a polynucleotide complementary to IL-34, or a portion thereof, for example an IL-34 antisense composition (*e.g.*, ASOs comprising SEQ ID NO: 40-43, or a pharmaceutically acceptable salt thereof). Exemplary

formulations include dosage forms that include or consist essentially of about 35 mg to about 500 mg of a disclosed polynucleotide complementary to IL-34, or a portion thereof, for example an IL-34 antisense oligonucleotide. For example, formulations that include about 35 mg, 40 mg, 50 mg, 60 mg, 70 mg, 80 mg, 90 mg, 100 mg, 110 mg, 120 mg, 130 mg, 140 mg, 150 mg, 160 mg, 170 mg, 180 mg, 190 mg, 200 mg, or 250 mg of a disclosed polynucleotide complementary to IL-34, or a portion thereof, for example an IL-34 antisense oligonucleotide, are contemplated herein. In one embodiment, a formulation may include about 40 mg, 80 mg, or 160 mg of a disclosed polynucleotide complementary to IL-34, or a portion thereof, for example an IL-34 antisense oligonucleotide. In some embodiments, a formulation may include at least 100 µg of a disclosed polynucleotide complementary to IL-34, or a portion thereof, for example an IL-34 antisense oligonucleotide. For example, formulations may include about 0.1 mg, 0.2 mg, 0.3 mg, 0.4 mg, 0.5 mg, 1 mg, 5 mg, 10 mg, 15 mg, 20 mg, or 25 mg of a disclosed polynucleotide complementary to IL-34, or a portion thereof, for example an IL-34 antisense oligonucleotide. The amount administered will depend on variables such as the type and extent of disease or indication to be treated, the overall health and size of the patient, the *in vivo* potency of the polynucleotide complementary to IL-34, or a portion thereof, the pharmaceutical formulation, and the route of administration. The initial dosage can be increased beyond the upper level in order to rapidly achieve the desired blood-level or tissue level. Alternatively, the initial dosage can be smaller than the optimum, and the dosage may be progressively increased during the course of treatment. Human dosage can be optimized, *e.g.*, in a conventional Phase I dose escalation study designed to run from 40 mg to 160 mg. Dosing frequency can vary, depending on factors such as route of administration, dosage amount and the disease being treated. Exemplary dosing frequencies are once per day, once per week and once every two weeks. In some embodiments, dosing is once per day for 7 days.

Diagnostic Methods

[0198] The disclosure also provides a method of diagnosing a patient with an inflammatory and/or fibrotic disease that relies upon detecting levels of IL-34 expression signal in one or more biological samples of a patient. As used herein, the term “IL-34 expression signal” can refer to any indication of IL-34 gene expression, or gene or gene product activity. IL-34 gene products include RNA (*e.g.*, mRNA), peptides, and proteins. Indices of IL-34 gene expression that can be assessed include, but are not limited to, IL-34 gene or chromatin state, IL-34 gene interaction

with cellular components that regulate gene expression, IL-34 gene product expression levels (e.g., IL-34 RNA expression levels, IL-34 protein expression levels), or interaction of IL-34 RNA or protein with transcriptional, translational, or post-translational processing machinery. Indices of IL-34 gene product activity include, but are not limited to, assessment of IL-34 signaling activity (e.g., assessment of CSF1R activation or MAPK1/MAPK3 phosphorylation) and assessment of IL-34 receptor binding (e.g., CSF1R binding).

[0199] Detection of IL-34 expression signal may be accomplished through *in vivo*, *in vitro*, or *ex vivo* methods. In a preferred embodiment, methods of the disclosure may be carried out *in vitro*. Methods of detecting may involve detection in blood, serum, fecal matter, tissue, or cells of a patient. Detection may be achieved by measuring IL-34 expression signal in whole tissue, tissue explants, cell cultures, dissociated cells, cell extract, or body fluids, including blood or serum. Contemplated methods of detection include assays that measure levels of IL-34 gene product expression such as Western blotting, FACS, ELISA, other quantitative binding assays, cell or tissue growth assays, Northern blots, quantitative or semi-quantitative polymerase chain reaction, medical imaging methods (e.g., MRI), or immunostaining methods (e.g., immunohistochemistry or immunocytochemistry).

Listing of Exemplary Embodiments

[0200] The invention is further described by the following non-limiting exemplary embodiments:

Embodiment 1. A polynucleotide, comprising a nucleotide sequence according to:

- a. 5'-CTTTGGGCCCGCACCAGCTTC-3' (SEQ ID NO: 3), or
- b. 5'-TCCATGACCCGGAAGCAGTT-3' (SEQ ID NO: 5),

wherein at least one cytidine of the nucleotide sequence is chemically modified, and at least one nucleoside is a 2'-O-(2-methoxyethyl) (2'-MOE) nucleoside;
or a pharmaceutically acceptable salt thereof.

Embodiment 2. The polynucleotide of embodiment 1, wherein no more than 10 nucleosides are chemically modified.

Embodiment 3. The polynucleotide of embodiment 1 or 2, comprising six 2'-MOE nucleosides.

Embodiment 4. The polynucleotide of embodiment 1 or 2, comprising eight 2'-MOE nucleosides.

Embodiment 5. The polynucleotide of any one of embodiments 1-4, comprising a nucleotide sequence selected from the group consisting of:

- a. 5'-CxTxTxTGGGCXGCACCAGCxTxTxCx-3' (SEQ ID NO: 40), wherein Cx is 2'-O-(2-methoxyethyl)cytidine, Tx is 2'-O-(2-methoxyethyl)thymidine, and X is 5-methylcytidine;
- b. 5'-CxTxTxTGGGCXGCACCAGCTxTxTx-3' (SEQ ID NO: 41), wherein Cx is 2'-O-(2-methoxyethyl)cytidine, Tx is 2'-O-(2-methoxyethyl)thymidine, and X is 5-methylcytidine;
- c. 5'-TxCxCxATGACCXGGAAGCAxGxTxTx-3' (SEQ ID NO: 42), wherein Cx is 2'-O-(2-methoxyethyl)cytidine, Ax is 2'-O-(2-methoxyethyl)adenosine, Tx is 2'-O-(2-methoxyethyl)thymidine, Gx is 2'-O-(2-methoxyethyl)guanosine, and X is 5-methylcytidine; and
- d. 5'-TxCxCxATGACCXGGAAGCAGxTxTx-3' (SEQ ID NO: 43), wherein Cx is 2'-O-(2-methoxyethyl)cytidine, Tx is 2'-O-(2-methoxyethyl)thymidine, Gx is 2'-O-(2-methoxyethyl)guanosine, and X is 5-methylcytidine;

or a pharmaceutically acceptable salt thereof.

Embodiment 6. The polynucleotide of any one of embodiments 1-5, wherein the polynucleotide is complementary to IL-34.

Embodiment 7. The polynucleotide of any one of embodiments 1-6, wherein the polynucleotide is an IL-34 antisense oligonucleotide.

Embodiment 8. The polynucleotide of any one of embodiments 1-7, wherein the polynucleotide is an IL-34 siRNA.

Embodiment 9. The polynucleotide of any one of embodiments 1-8, wherein at least one internucleoside linkage of the polynucleotide is selected from the group consisting of a phosphorothioate linkage, a phosphorodithioate linkage, a phosphotriester linkage, an alkylphosphonate linkage, an aminoalkylphosphotriester linkage, an alkylene phosphonate linkage, a phosphinate linkage, a phosphoramidate linkage, and an aminoalkylphosphoramidate linkage, a thiophosphoramidate linkage, thionoalkylphosphonate linkage, a

thionoalkylphosphotriester linkage, a thiophosphate linkage, a selenophosphate linkage, and a boranophosphate linkage.

Embodiment 10. The polynucleotide of embodiment 9, wherein at least one internucleoside linkage of the polynucleotide is a phosphorothioate linkage.

Embodiment 11. The polynucleotide of any one of embodiments 1-10, wherein all internucleoside linkages of the polynucleotide are phosphorothioate linkages.

Embodiment 12. The polynucleotide of any one of embodiments 1-11, wherein the polynucleotide is 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 nucleotides in length.

Embodiment 13. The polynucleotide of any one of embodiments 1-11, wherein the polynucleotide is 20 nucleotides in length or 20-25, 20-30, 20-35, 25-30, 25-35, or 30-35 nucleotides in length.

Embodiment 14. The polynucleotide of any one of embodiments 1-13, wherein the polynucleotide is no more than 20, 25, or 30 nucleotides in length.

Embodiment 15. The polynucleotide of any one of embodiments 1-13, wherein the polynucleotide is from 20 to 25 nucleotides in length.

Embodiment 16. The polynucleotide of any one of embodiments 1-13, wherein the polynucleotide is 20 nucleotides in length.

Embodiment 17. A pharmaceutical composition, comprising (a) a polynucleotide of any one of embodiments 1-16, or a pharmaceutically acceptable salt thereof, and (b) a pharmaceutically acceptable excipient.

Embodiment 18. The pharmaceutical composition of embodiment 17, wherein the pharmaceutical composition is suitable for topical, parenteral, oral, pulmonary, intratracheal, intranasal, transdermal, or intraduodenal administration.

Embodiment 19. A method of treating an inflammatory disease in a patient in need thereof, the method comprising administering to the patient in need thereof an effective amount of a polynucleotide of any one of embodiments 1-16, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition of any one of embodiments 17-18.

Embodiment 20. The method of embodiment 19, wherein the inflammatory disease is associated with altered IL-34 expression.

Embodiment 21. The method of embodiment 19 or 20, wherein the inflammatory disease is associated with elevated IL-34 expression.

Embodiment 22. The method any one of embodiments 19-21, wherein the inflammatory disease is rheumatoid arthritis.

Embodiment 23. The method any one of embodiments 19-21, wherein the inflammatory disease is osteoarthritis.

Embodiment 24. The method of any one of embodiments 19-23, wherein the method inhibits inflammatory cytokine production in cells of the patient.

Embodiment 25. The method of any one of embodiments 19-24, wherein the method reduces or inhibits an IL-34 mediated inflammatory response in cells of the patient.

Embodiment 26. The method of any one of embodiments 19-25, wherein the method reduces or inhibits IL-34-mediated macrophage colony-stimulating factor receptor (M-CSFR-1) signaling in cells of the patient.

Embodiment 27. The method of any one of embodiments 24-26, wherein the cell is an intestinal cell.

Embodiment 28. The method of any one of embodiments 24-26, wherein the cell is an intestinal stromal cell.

Embodiment 29. The method of any one of embodiments 24-26, wherein the cell forms part of an intestinal fibrostricture.

Embodiment 30. A method of reducing or eliminating a fibrotic stricture in a patient suffering from an inflammatory disease, the method comprising administering an effective amount of a polynucleotide of any one of embodiments 1-16, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition of any one of embodiments 17-18.

Embodiment 31. The method of embodiment 27, wherein the fibrotic stricture is located in the intestine.

Embodiment 32. The method of embodiment 30 or 31, wherein the inflammatory disease is selected from the group consisting of an inflammatory bowel disease, rheumatoid arthritis, psoriasis, osteoarthritis, pouchitis, diabetes (type I and II), tissue or organ rejection, multiple sclerosis, periodontal inflammation, periodontitis, pigmented villonodular synovitis, hepatitis, sinusitis, colon cancer, colorectal cancer, colitis-associated colon cancer, sporadic colorectal cancer, coronary artery disease, Sjogren's syndrome (SS), obesity, chronic inflammation, pulmonary sarcoidosis, skin lesions, a CNS inflammatory disease, and an autoimmune disease.

- Embodiment 33.** The method of embodiment 32, wherein the inflammatory disease is a rheumatoid arthritis.
- Embodiment 34.** The method of embodiment 32, wherein the inflammatory disease is an osteoarthritis.
- Embodiment 35.** The method of embodiment 32, wherein the inflammatory disease is an inflammatory bowel disease.
- Embodiment 36.** The method of embodiment 32, wherein the inflammatory bowel disease is selected from the group consisting of Crohn's disease, gastroduodenal Crohn's disease, Crohn's (granulomatous) colitis, inflammatory Crohn's disease, fibrostricturing Crohn's disease, ulcerative colitis, collagenous colitis, lymphocytic colitis, ischaemic colitis, diversion colitis, Behçet's disease, microscopic colitis, ulcerative proctitis, proctosigmoiditis, jejunoileitis, left-sided colitis, pancolitis, ileocolitis, ileitis, and indeterminate colitis.
- Embodiment 37.** The method of embodiment 30, wherein the inflammatory bowel disease is inflammatory Crohn's disease.
- Embodiment 38.** The method of embodiment 30, wherein the inflammatory bowel disease is fibrostricturing Crohn's disease.
- Embodiment 39.** A method for preventing or treating fibrosis, the method comprising administering to a patient in need thereof a therapeutically effective amount of a polynucleotide of any one of embodiments 1-16, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition of any one of embodiments 17-18.
- Embodiment 40.** The method of embodiment 39, wherein the fibrosis is intestinal fibrosis.
- Embodiment 41.** The method of embodiment 39, wherein the fibrosis is pulmonary fibrosis.
- Embodiment 42.** The method of embodiment 39, wherein the fibrosis is synovial fibrosis.
- Embodiment 43.** The method of embodiment 39, wherein the fibrosis is selected from the group consisting of renal fibrosis, cardiac fibrosis, endomyocardial fibrosis, myelofibrosis, retroperitoneal fibrosis, and nephrogenic systemic fibrosis.
- Embodiment 44.** The method of any one of embodiments 39-43, wherein the patient is suffering from rheumatoid arthritis.
- Embodiment 45.** The method of any one of embodiments 39-43, wherein the patient is suffering from osteoarthritis.

- Embodiment 46.** The method of any one of embodiments 39-43, wherein the patient is suffering from Crohn's disease.
- Embodiment 47.** The method of any one of embodiments 19-46, wherein the polynucleotide is administered intraarticularly, rectally, topically, parenterally, orally, pulmonarily, intratracheally, intranasally, transdermally, or intraduodenally.
- Embodiment 48.** The method of embodiment 47, wherein the polynucleotide is administered intraarticularly.
- Embodiment 49.** The method of embodiment 47, wherein the polynucleotide is administered rectally.
- Embodiment 50.** The method of embodiment 47, wherein the polynucleotide is administered orally.
- Embodiment 51.** Use of a polynucleotide of any one of embodiments 1-17, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for treating an inflammatory disease.
- Embodiment 52.** The use of embodiment 51, wherein the inflammatory disease is selected from the group consisting of an inflammatory bowel disease, rheumatoid arthritis, psoriasis, osteoarthritis, pouchitis, diabetes (type I and II), tissue or organ rejection, multiple sclerosis, periodontal inflammation, periodontitis, pigmented villonodular synovitis, hepatitis, sinusitis, colon cancer, colorectal cancer, colitis-associated colon cancer, sporadic colorectal cancer, coronary artery disease, or Sjogren's syndrome (SS), obesity, chronic inflammation, pulmonary sarcoidosis, skin lesions, a CNS inflammatory disease, and an autoimmune disease.
- Embodiment 53.** The use of any one of embodiments 51-52, wherein the patient is suffering from rheumatoid arthritis.
- Embodiment 54.** The use of any one of embodiments 51-52, wherein the patient is suffering from osteoarthritis.
- Embodiment 55.** The use of any one of embodiments 51-52, wherein the patient is suffering from Crohn's disease.
- Embodiment 56.** The use of embodiment 52, wherein the inflammatory disease is an inflammatory bowel disease.
- Embodiment 57.** The use of embodiment 56, wherein the inflammatory bowel disease is selected from the group consisting of Crohn's disease, inflammatory Crohn's disease,

fibrostricturing Crohn's disease, gastroduodenal Crohn's disease, Crohn's (granulomatous) colitis, ulcerative colitis, collagenous colitis, lymphocytic colitis, ischaemic colitis, diversion colitis, Behçet's disease, microscopic colitis, ulcerative proctitis, proctosigmoiditis, jejunoileitis, left-sided colitis, pancolitis, ileocolitis, ileitis, and indeterminate colitis.

Embodiment 58. The use of embodiment 57, wherein the inflammatory bowel disease is inflammatory Crohn's disease.

Embodiment 59. The use of embodiment 57, wherein the inflammatory bowel disease is fibrostricturing Crohn's disease.

Embodiment 60. A polynucleotide of any one of embodiments 1-17, or a pharmaceutically-acceptable salt thereof, for use as a medicament.

Embodiment 61. The polynucleotide of embodiment 60 for use in treating an inflammatory disease in a patient in need thereof.

Embodiment 62. The polynucleotide of embodiment 61, wherein the inflammatory disease is rheumatoid arthritis or osteoarthritis.

Embodiment 63. A polynucleotide of any one of embodiments 1-17, or a pharmaceutically-acceptable salt thereof, for use in the treatment of an inflammatory disease.

Embodiment 64. The polynucleotide for use as embodied in embodiment 63, wherein the inflammatory disease is selected from the group consisting of an inflammatory bowel disease, rheumatoid arthritis, psoriasis, osteoarthritis, pouchitis, diabetes (type I and II), tissue or organ rejection, multiple sclerosis, periodontal inflammation, periodontitis, pigmented villonodular synovitis, hepatitis, sinusitis, colon cancer, colorectal cancer, colitis-associated colon cancer, sporadic colorectal cancer, coronary artery disease, or Sjogren's syndrome (SS), obesity, chronic inflammation, pulmonary sarcoidosis, skin lesions, a CNS inflammatory disease, and an autoimmune disease.

Embodiment 65. The polynucleotide for use as embodied in embodiment 64, wherein the inflammatory disease is rheumatoid arthritis.

Embodiment 66. The polynucleotide for use as embodied in embodiment 64, wherein the inflammatory disease is osteoarthritis.

Embodiment 67. The polynucleotide for use as embodied in embodiment 64, wherein the inflammatory disease is an inflammatory bowel disease.

Embodiment 68. The polynucleotide for use as embodied in embodiment 67, wherein the inflammatory bowel disease is selected from the group consisting of Crohn's disease, inflammatory Crohn's disease, fibrostricturing Crohn's disease, gastroduodenal Crohn's disease, Crohn's (granulomatous) colitis, ulcerative colitis, collagenous colitis, lymphocytic colitis, ischaemic colitis, diversion colitis, Behçet's disease, microscopic colitis, ulcerative proctitis, proctosigmoiditis, jejunoileitis, left-sided colitis, pancolitis, ileocolitis, ileitis, and indeterminate colitis.

Embodiment 69. The polynucleotide for use as embodied in embodiment 68, wherein the inflammatory bowel disease is inflammatory Crohn's disease.

Embodiment 70. The polynucleotide for use as embodied in embodiment 68, wherein the inflammatory bowel disease is fibrostricturing Crohn's disease.

Embodiment 71. A polynucleotide of any one of embodiments 1-17, or a pharmaceutically-acceptable salt thereof, for use in the treatment of fibrosis.

Embodiment 72. The polynucleotide for use as embodied in embodiment 71, wherein the fibrosis is intestinal fibrosis.

Embodiment 73. The polynucleotide for use as embodied in embodiment 71, wherein the fibrosis is pulmonary fibrosis.

Embodiment 74. The polynucleotide for use as embodied in embodiment 71, wherein the fibrosis is synovial fibrosis.

Embodiment 75. The polynucleotide for use as embodied in embodiment 71, wherein the fibrosis is selected from the group consisting of renal fibrosis, cardiac fibrosis, endomyocardial fibrosis, myelofibrosis, retroperitoneal fibrosis, and nephrogenic systemic fibrosis.

Embodiment 76. The polynucleotide for use as embodied in any one of embodiments 60-75, wherein the polynucleotide is to be administered intraarticularly, rectally, topically, parenterally, orally, pulmonarily, intratracheally, intranasally, transdermally, or intraduodenally.

Embodiment 77. The polynucleotide for use as embodied in embodiments 76, wherein the polynucleotide is to be administered intraarticularly.

Embodiment 78. The polynucleotide for use as embodied in embodiments 76, wherein the polynucleotide is to be administered rectally.

Embodiment 79. The polynucleotide for use as embodied in embodiments 76, wherein the polynucleotide is to be administered orally.

Embodiment 80. The polynucleotide for use as embodied in any one of embodiments 60-79, wherein the patient is a human.

EXAMPLES

[0201] The disclosure is further illustrated by the following examples. The examples are provided for illustrative purposes only, and are not to be construed as limiting the scope or content of the disclosure in any way.

Example 1: Design and evaluation of IL-34 of 3-14-3 and 4-12-4 antisense oligonucleotides

[0202] IL-34 antisense oligonucleotides of SEQ ID NOs: 40-43 complementary to mouse and human IL-34 mRNA sequences are designed and produced by commercial manufacturers.

[0203] Knockdown of IL-34 expression are evaluated in HT-29 human and RAW 264.7 mouse cell lines. HT-29 cells (ATCC Manassas, Virginia, USA) are cultured in McCoy's 5A (Lonza, Verviers, Belgium) supplemented with 10% FBS, 1% P/S (all from Lonza) and maintained at 37 °C with 5% CO₂ in a humidified incubator. Murine RAW264.7 macrophages (ATCC Manassas, Virginia, USA) are cultured in Dulbecco's modified Eagle's medium (1 g/L of glucose) with 10% FBS, 1% P/S and 1 % non-essential amino acid (NEA) (all from Lonza) and maintained at 37 °C with 5% CO₂ in a humidified incubator. The murine colon carcinoma cell line MC-38 (ABMGood, Richmond, Canada) is cultured in DMEM medium supplemented with 10% FBS and 1% P/S and maintained at 37°C with 5% CO₂ in a humidified incubator.

[0204] Cells are either left untreated or transfected with a specific IL-34 ASO or scrambled ASO using Opti-MEM medium and Lipofectamine 3000 reagent (Life Technologies, Milan, Italy) according to the manufacturer's instructions. Total RNA and protein are extracted.

[0205] Knockdown is evaluated by real-time polymerase chain reaction (RT-PCR) in HT-29 cells. RT-PCR was performed as follows: a constant amount of RNA (0.5 µg/sample) is retro-transcribed into complementary DNA (cDNA). 1 µl of cDNA/sample is then amplified using the following protocol: denaturation for 1 minute at 95°C; annealing for 30 seconds at 60°C, followed by 30 seconds of extension at 72°C. The following primer sequences are used: IL-34 forward, 5'-ACAGGAGCCGACTTCAGTAC-3' (SEQ ID NO: 29); IL-34 reverse, 5'-ACCAAGACCCACAGATACCG-3' (SEQ ID NO: 30); β-actin forward, 5'-AAGATGACCCAGATCATGTTTGAGACC-3' (SEQ ID NO: 31); and β-actin reverse, 5'-

AGCCAGTCCAGACGCAGGAT-3' (SEQ ID NO: 32). mRNA expression is calculated relative to β -Actin mRNA expression.

[0206] HT-29 cells are transfected with human IL-34 ASOs of SEQ ID NO: 40, 41, 42, or 43, or a scrambled negative control oligonucleotide (SRC AS; 5'-AACACGTCTATACGC-3' (SEQ ID NO: 33)) for 24 hours. IL-34 transcripts are evaluated by RT-PCR and values are normalized to β -actin.

[0207] Knockdown is also evaluated by Western blotting in HT-29 cells and RAW 264.7 cells. Western blotting is performed as follows: following transfection, cells are lysed on ice in buffer containing 10 mM HEPES (pH 7.9), 10 mM KCl, 0.1 mM EDTA, 0.2 mM EGTA, and 0.5% Nonidet P40 supplemented with 1 mM dithiothreitol, 10 mg/ml aprotinin, 10 mg/ml leupeptin, 1 mM phenylmethylsulfonyl fluoride, 1 mM Na₃VO₄ and 1 mM NaF. Lysates are clarified by centrifugation at 4°C, 12,000xg for 30 minutes, and separated on 10% sodium dodecyl sulphate-polyacrylamide gel electrophoresis. IL-34 is detected using mouse anti human IL-34 antibody (1:1,000 final dilution; Abcam Cambridge, UK), in combination with horseradish peroxidase (HRP)-conjugated secondary IgG monoclonal antibody (1:20,000 final dilution; Dako, Milan, Italy). HRP reaction is detected with a sensitive enhanced chemiluminescence kit (Pierce, Rockford, IL). After the analysis, blots are stripped and probed with mouse anti-human β -actin antibody (1:5,000 final dilution; Sigma-Aldrich), followed by HRP-conjugated secondary IgG monoclonal antibody (1:20,000 final dilution; Dako).

[0208] IL-34 knockdown by IL-34 ASOs is further evaluated in fibrostricturing Crohn's disease and cancer associated fibroblasts. Intestinal fibroblasts are isolated from intestinal colon cancer or fibrostricturing CD (FS CD) explants. Fibroblasts are used freshly or between passages 3 and 8. Fibroblasts are cultured in DMEM high glucose with UltraGlutamine supplemented with 10% FBS, 1% P/S (Lonza) and maintained at 37°C with 5% CO₂ in a humidified incubator. Fibroblasts are detached using Trypsin/EDTA solution evaluated for knockdown of IL-34.

Example 2: Design and evaluation of IL-34 gapmer antisense oligonucleotides

[0209] This example describes the evaluation of the capability of antisense oligonucleotides (ASOs) complementary to the IL-34 mRNA transcripts, to knock-down IL-34 expression in the HT-29 and AGS gastrointestinal cancer cell lines as well as in Lamina Propria Mononuclear

Cells (LPMCs) isolated from the inflamed colonic tissue of patients affected by ulcerative colitis (UC).

[0210] Antisense oligonucleotides complementary to human, mouse, cynomolgus monkey (and other species) IL-34 mRNA transcripts were designed. The designed ASOs are single strand oligodeoxyribonucleotides, which target the same sequence in IL-34 mRNA and are modified to have a fully phosphorothioate backbone and 2'-O-methoxyethyl (2'-MOE) sugar modifications at the 5' and 3' ends.

[0211] Four different ASOs were evaluated (SEQ ID NOs: 17, 19, 42 and 43) as described below. The sequences of the test ASOs are shown in **TABLE 1**, with X = 5-methylcytidine; Cx = 2'-O-MOE cytidine; Tx = 2'-O-MOE thymidine; Gx = 2'-O-MOE guanosine; and Ax = 2'-O-MOE adenosine.

TABLE 1

Seq ID NO:	Sequence	Description
17	5'-TxCxCxAxTxGACCXGGAAGCxAxGxTxTx-3'	5-10-5 MOE gapmer
19	5'-TxCxCxAxTxGxACCXGGAAGxCxAxGxTxTx-3'	6-8-6 MOE gapmer
42	5'-TxCxCxAxTGACCXGGAAGCxAxGxTxTx-3'	4-12-4 MOE gapmer
43	5'-TxCxCxATGACCXGGAAGCAGxTxTx-3'	3-14-3 MOE gapmer

[0212] The colorectal cancer cell line HT-29 and the gastric cancer cell line AGS (ATCC, Manassas, USA) were cultured in McCoy's 5A and RPMI 1640 media respectively (Lonza, Verviers, Belgium), supplemented with 10% FBS, 1% P/S (Lonza), and maintained at 37°C with 5% CO₂ in a humidified incubator.

[0213] Lamina propria mononuclear cells (LPMCs) were isolated from biopsies taken from the inflamed colonic tissue of patients with ulcerative colitis (UC) as previously described in Dinallo et al., *J Crohns Colitis*. 2019 May 27;13(6):772-784. LPMCs were resuspended in RPMI 1640 supplemented with 10% FBS, 1% P/S and maintained at 37°C with 5% CO₂ in a humidified incubator. To assess the ability of IL-34 ASOs to downregulate IL-34 expression, HT-29, AGS, and UC patient-derived LPMCs were transfected with either the test IL-34 ASOs or a control of relevant scrambled antisense oligonucleotide (all used at the concentration of 200 nM) for 24 hours using Opti-MEM medium and Lipofectamine 3000 reagent (Life Technologies, Milan, Italy) according to the manufacturer's instructions. At the end, cells were collected and

total RNA extracted using a PureLink mRNA Mini Kit (Thermo Fisher Scientific, Waltham, USA), according to the manufacturer's instructions. A constant amount of RNA was reverse transcribed into complementary DNA (cDNA) and amplified by real-time PCR using the following protocol: denaturation for 1 minute at 95°C; annealing for 30 seconds at 60°C, followed by 30 seconds of extension at 72°C. Primer sequences were as follows: IL-34: forward, 5'-ACAGGAGCCGACTTCAGTAC-3' (SEQ ID NO: 29); reverse, 5'-ACCAAGACCCACAGATACCG-3' (SEQ ID NO: 30); β -actin: forward, 5'-AAGATGACCCAGATCATGTTTGAGACC-3' (SEQ ID NO: 31); reverse, 5'-AGCCAGTCCAGACGCAGGAT-3' (SEQ ID NO: 32). RNA expression was calculated relative to the housekeeping β -actin gene on the base of the $\Delta\Delta C_t$ algorithm.

[0214] FIG. 1, FIG. 2, and FIG. 3 show the analysis of IL-34 transcripts in HT-29 cells, AGS cells, and LPMCs in conditions treated with CTR=scrambled ASO, 1=SEQ ID NO: 17, 2=SEQ ID NO: 19, 3=SEQ ID NO: 42, 4=SEQ ID NO: 43. The results show that all four IL-34 ASOs significantly reduced IL-34 RNA transcripts in HT-29 cells, AGS cells, and LPMCs as compared to the control. The data indicate that the four gapmer IL-34 ASOs are capable of efficiently down-regulating IL-34 expression in human cancer and immune cells and may be useful in relevant preclinical and clinical settings.

Example 3: Knockdown of IL-34 expression by IL-34 antisense oligonucleotides

[0215] The ability of IL-34 ASOs to knockdown IL-34 expression is evaluated in HT-29 cells (human colon adenocarcinoma cell line), THP-1 cells (human leukemia monocytic cell line), RAW 264.7 cells (murine macrophage cell line), and MC-38 cells (murine colon adenocarcinoma cell line).

[0216] HT-29 cells (ATCC HTB-38, Manassas, USA) are cultured in McCoy's 5A and RPMI 1640 medium, supplemented with 10% fetal bovine serum (FBS), 1% penicillin/streptomycin (P/S), and maintained at 37 °C with 5% CO₂ in a humidified incubator.

[0217] THP-1 cells (ATCC TIB-202) are grown in RPMI 1640 medium supplemented with 10 FBS and 1% P/S. To induce macrophage differentiation, cells are treated with 100 ng/mL phorbol 12-myristate 13-acetate (PMA) for 24 hours and maintained at 37°C with 5% CO₂ in a humidified incubator.

[0218] RAW 264.7 cells (ATCC TIB-71) are cultured in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% FBS, 1% P/S and 1 % non-essential amino acids (NEAA) (Lonza) and maintained at 37°C with 5% CO₂ in a humidified incubator.

[0219] MC-38 cells (ABMGood, Richmond, Canada) are cultured in DMEM medium supplemented with 10% FBS and 1% P/S and maintained at 37°C with 5% CO₂ in a humidified incubator.

[0220] Cells are either left untreated or transfected with a test IL-34 ASO or a control of scrambled ASO using Opti-MEM medium and Lipofectamine 3000 reagent in the presence or absence of specific pro-inflammatory stimuli, such as lipopolysaccharides (LPS), peptidoglycan (PGN), and poly I:C.

[0221] Cells are then collected and total RNA extracted for analysis with RT-qPCR performed as follows: a constant amount of RNA is retro-transcribed into complementary DNA (cDNA). 1 µL of cDNA/sample is then amplified using the following protocol: denaturation for 1 minute at 95°C; annealing for 30 seconds at 60°C, followed by 30 seconds of extension at 72°C. The following primer sequences are used: IL-34 forward, 5'-ACAGGAGCCGACTTCAGTAC-3' (SEQ ID NO: 29); IL-34 reverse, 5'-ACCAAGACCCACAGATACCG-3' (SEQ ID NO: 30); β-actin forward, 5'-AAGATGACCCAGATCATGTTTGAGACC-3' (SEQ ID NO: 31); and β-actin reverse, 5'-AGCCAGTCCAGACGCAGGAT-3' (SEQ ID NO: 32). mRNA expression is calculated relative to the housekeeping gene β-Actin mRNA expression on the base of the $\Delta\Delta C_t$ algorithm.

[0222] IL-34 protein knockdown is evaluated by Western blotting performed as follows: following transfection, cells are lysed on ice in a buffer containing 10 mM HEPES (pH 7.9), 10 mM KCl, 0.1 mM EDTA, 0.2 mM EGTA, and 0.5% Nonidet P40 supplemented with 1 mM dithiothreitol, 10 mg/ml aprotinin, 10 mg/ml leupeptin, 1 mM phenylmethylsulfonyl fluoride, 1 mM Na₃VO₄ and 1 mM NaF. Lysates are clarified by centrifugation at 4°C, 12,000xg for 30 minutes, and separated on 10% sodium dodecyl sulphate-polyacrylamide gel electrophoresis. IL-34 is detected using either a mouse anti human IL-34 antibody or a rabbit anti-human IL-34 antibody, in combination with horseradish peroxidase (HRP)-conjugated secondary IgG monoclonal antibody. HRP reaction is detected with a sensitive enhanced chemiluminescence kit. After the analysis, blots are stripped and probed with mouse anti-β-actin antibody, followed

by HRP-conjugated secondary IgG monoclonal antibody. Computer-assisted scanning densitometry is used to analyze the intensity of the immunoreactive bands.

[0223] The results of the transfection of the cells with the IL-34 ASOs results in decreased IL-34 RNA and protein expression.

Example 4: Knockdown of IL-34 expression by IL-34 antisense oligonucleotides in intestinal fibroblasts

[0224] IL-34 knockdown by IL-34 ASOs is further evaluated in colorectal cancer (CRC)-associated fibroblasts and fibrostricting Crohn's disease (FS CD)-derived fibroblasts.

[0225] Human intestinal fibroblasts are isolated from explants derived from the affected tissue of patients with CRC and FS CD and phenotypically as previously described. In all the experiments, fibroblasts are used freshly or up to passage 8 and are detached using a Trypsin/EDTA solution. The human fibroblasts are cultured in DMEM high glucose with UltraGlutamine supplemented with 10% FBS, 1% P/S and maintained at 37 °C with 5% CO₂ in a humidified incubator.

[0226] Cells are either left untreated or transfected with a test IL-34 ASO or a control of scrambled antisense oligonucleotides. Total RNAs are extracted for analysis using RT-qPCR as described in **Example 3**. IL-34 ASOs result in decreased IL-34 RNA expression compared to the control.

[0227] The effect of IL-34 ASOs on collagen synthesis in the fibroblasts is also evaluated. In brief, 5x10⁴ fibroblasts are plated into each well of a 12-well plate, left to adhere for 24 hours, and then either left untreated or transfected with a test IL-34 ASO or a control of scrambled ASOs for 24-48 hours. Transfection efficiency is determined by Western blotting. After 48 hours, cell-free supernatants are collected and analyzed for collagen content by a commercially available ELISA kit. IL-34 ASOs reduces both COL1A1 and COL3A1 protein synthesis in the fibroblasts compared to the control.

[0228] The effect of IL-34 ASOs on cell death is also evaluated. To score cell death, fibroblasts are washed in PBS, stained with FITC-annexin V antibody and incubated with 5 mg/mL PI for 20 min at 4°C. Fluorescence is measured using a flow cytometer. AV-/PI-cells are considered to be viable cells, AV+/PI-cells are considered to be apoptotic, and secondary

necrotic cells are characterized by AV+/PI+ staining. Transfection of FS CD fibroblasts with the IL-34 ASOs does not affect cell viability.

[0229] IL-34 knockdown in fibroblasts by IL-34 ASOs results in decreased collagen production without having a significant effect on cell viability.

Example 5: Knockdown of IL-34 expression by IL-34 antisense oligonucleotides

[0230] This example illustrates knockdown of expression of IL-34 by IL-34 antisense oligonucleotides *in vitro* in human synovial sarcoma cell line SW982 and in human fibroblast-like synoviocytes HFSL-RA.

[0231] SW982 cells are grown in Leibovitz's L15 Medium supplemented with 100 U/mL penicillin, 100 µg/mL streptomycin, and 10% fetal bovine serum (FBS). Cell cultures are maintained in a humidified atmosphere of 95% air and 5% CO₂ at 37°. Their activation is induced by 24h of pre-treatment with 10 ng/mL TNFα.

[0232] HFSL-RA cells are cultured in Synoviocyte growth medium kit (basal medium with growth supplements) and maintained at 37°C, 5% CO₂ in a humidified incubator.

[0233] Cells are either left untreated or transfected with a test IL-34 ASO or a control of scrambled antisense oligonucleotides. Total RNAs are extracted for analysis using RT-qPCR as described in **Example 3**. IL-34 protein knockdown is evaluated by Western blotting as described in **Example 3**.

[0234] IL-34 ASOs result in decreased IL-34 RNA and protein expression compared to the control.

Example 6: Evaluation of IL-34 antisense oligonucleotides *in vivo* in colitis model

[0235] This example illustrates of IL-34 knockdown by IL-34 ASO *in vivo* in a mouse model of TNBS-induced colitis.

[0236] IL-34 ASOs are dissolved in phosphate buffered saline. For oral administration, the ASOs are diluted in bicarbonate solution pH 9.5; for i.p. and rectal administration the ASOs are diluted in PBS.

[0237] Female Balb/c mice at 7-9 weeks old are used for the TNBS-induced colitis model. To induce colitis in the mice, 2 mg of 2,4,6-trinitrobenzenesulfonic acid (TNBS) in 45% ethanol is administered to lightly anesthetized mice through a 3.5F catheter inserted into the rectum. The catheter tip is inserted 4 cm proximal to the anal verge, and 150 µL of fluid is slowly instilled

into the colon, after which the mouse is held in a vertical position for 30 seconds. After 24 h of TNBS induction, mice in the treatment group receive the test IL-34 ASO or a control.

[0238] Weight changes are recorded daily to assess the induction and progression of colitis, and tissues are collected for histologic study and protein analysis. Mice are euthanized at Day 5.

[0239] For histologic studies, tissues removed from mice after euthanasia are fixed in 10% formalin solution (Sigma Aldrich), embedded in paraffin, cut into tissue sections and stained with Haematoxylin and Eosin. Stained sections are examined for evidence of colitis using different criteria such as the presence of lymphocyte infiltration, elongation and/or distortion of crypts, frank ulceration and thickening of the bowel wall. The degree of inflammation on microscopic cross-sections of the colon is graded from 0 to 4 as follows:

0: no evidence of inflammation;

1: low level of lymphocyte infiltration with infiltration seen in a <10% high-power field (hpf), no structural changes observed;

2: moderate lymphocyte infiltration with infiltration seen in <10–25% hpf, crypt elongation, bowel wall thickening which does not extend beyond mucosal layer;

3: high level of lymphocyte infiltration with infiltration seen in <25–50% hpf, thickening of bowel wall which extends beyond mucosal layer;

4: marked degree of lymphocyte infiltration with infiltration seen in >50% hpf, high vascular density, crypt elongation with distortion, transmural bowel wall-thickening with ulceration.

[0240] Mice have reduced colitis after treatment with the IL-34 ASOs.

Example 7: Evaluation of IL-34 antisense oligonucleotides *in vivo* in arthritis model

[0241] This example illustrates knockdown of expression of IL-34 by IL-34 antisense oligonucleotides IL-34 ASO in vivo in collagen-induced arthritis (CIA) animal models.

[0242] IL-34 ASOs are dissolved in phosphate buffered saline. For oral administration the ASOs are diluted in bicarbonate solution pH 9.5; for i.p and rectal administration the ASOs are diluted in PBS.

[0243] DBA/1 mice at 8-12 weeks of age are used. The animals are challenged with bovine type-II collagen emulsified with Complete Freund's adjuvant (CFA) on Day 0 (0.1 mg/ 0.1 mL/mouse, intradermally (ID) at the base of the tail) and boosted i.d. on Day 21 with collagen emulsified with IFA instead of CFA (0.1 mg/ 0.1 mL/ mouse). Animals are stratified into groups

of 10 each, based on the volume of two hind paws on Day 29. ASOs and vehicle are administered by intra-articular injection at different dosing schedule from Day 29 to Day 42. Control article dexamethasone (0.3 mg/kg) is administered orally. Arthritis is assessed by using a qualitative severity score system and hind paw volume (mL) measured with a plethysmometer on Days 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41 and 43. Additional analyses that can be performed include hind paw harvest and weight at termination, biomarkers (protein and/or mRNA), and histopathology.

[0244] Alternatively, female Lewis rats weighing 180 ± 20 g are used. The animals are challenged with porcine type-II collagen with incomplete Freund's adjuvant on Day 1 (0.2 mg/ 0.2 mL/ rat, subcutaneously (SC) at the base of the tail) and booster on Day 7 (0.1 mg/ 0.1 mL/ rat, SC). Animals are stratified into groups of 6 each, based on the volume of two hind paws on Day 16 and then dosing begins on Day 17. Control article dexamethasone (0.3 mg/kg) is administered orally once a day for 14 consecutive days from Day 17 to Day 30. ASOs and vehicle are administered by intra-articular injection. Arthritis is assessed by using a qualitative severity score system and hind paw volume (mL) measured with a plethysmometer on Days 1, 7, 10, 14, 16, 18, 20, 22, 24, 26, 28 and 30. Additional analyses that can be performed include biomarkers (protein and/or mRNA) and histopathology.

[0245] Animals treated with IL-34 ASOs show reduced arthritis.

EQUIVALENTS

[0246] The disclosure can be embodied in other specific forms with departing from the essential characteristics thereof. The foregoing embodiments therefore are to be considered illustrative rather than limiting on the disclosure described herein. The scope of the disclosure is indicated by the appended claims rather than by the foregoing description, and all changes that come within the meaning and range of equivalency of the claims are intended to be embraced therein.

SEQUENCE LISTING

X is 5-methylcytidine; Ax is 2'-O-(2-methoxyethyl)adenosine; Cx is 2'-O-(2-methoxyethyl)cytidine; Gx is 2'-O-(2-methoxyethyl)guanosine; Tx is 2'-O-(2-

methoxyethyl)thymidine; Ay is 2'-O-methyladenosine; Cy is 2'-O-methylcytidine; Gy is 2'-O-methylguanosine; Ty is 2'-O-methylthymidine.

SEQ ID NO	Sequence
1	5'-CTCACCAAGACCCACAG-3'
2	5'-GGCTTTGGGCCGCACCAGCT-3'
3	5'-CTTTGGGCCGCACCAGCTTC-3'
4	5'-TGGGCCGCACCAGCTTCAGG-3'
5	5'-TCCATGACCCGGAAGCAGTT-3'
6	5'-TGTTTCATGTACTGAAG-3'
7	5'-CTTTGGGCXGCACCAGCTTC-3'
8	5'-TCCATGACCXGGAAGCAGTT-3'
9	5'-cttTGGGCXGCACCAGCttc-3'
10	5'-ctttGGGCXGCACCAgcttc-3'
11	5'-cttTGGGCcgCACCAGCttc-3'
12	5'-cttTGGGCcGCACCAGCttc-3'
13	5'-ggcXGCACCAGCttc-3'
14	5'-cttTGGGCXGCACcag-3'
15	5'-tgaCCXGGAAGCAgtt-3'
16	5'-CxTxTxTxGxGGCXGCACCAGxCxTxTxTx-3'
17	5'-TxCxCxTxGxGACCCXGGAAGCxAxGxTxTx-3'
18	5'-CxTxTxTxGxGxGCXGCACCAxGxCxTxTxTx-3'
19	5'-TxCxCxTxGxGxACCXGGAAGxCxAxGxTxTx-3'
20	5'-CxTxTxTxGxGxGxCxXGxCxAxGxCxTxTxTx-3'
21	5'-TxCxCxTxGxGxGxCxXGxCxAxGxCxTxTxTx-3'
22	5'-CyTyTyTyGyGGCXGCACCAGyCyTyTyCy-3'
23	5'-TyCyCyAyTyGACCCXGGAAGCyAyGyTyTy-3'
24	CATCAGACGGGAAGCCTGGACTGTGGGTTGGGGGCAGCCTCAGCCTCTCCAACCTG GCACCCACTGCCCCTGGCCCTTAGGCACCTGCTTGGGGTCCTGGAGCCCCCTTAAGG CCACCAGCAAATCCTAGGAGACCGAGTCTTGGCACGTGAACAGAGCCAGATTTC CACTGAGCAGCTGCAGTCGGAGAAATCAGAGAAAGCGTCACCCAGCCCCAGATTTC CGAGGGGCTGCCAGGGACTCTCTCCTCCTGCTCCTTGGAAGGAAGACCCCCGAAA GACCCCCAAGCCACCGGCTCAGACCTGCTTCTGGGCTGCCATGGGACTTGCGGCCA CCGCCCCCGGCTGTCTCCACGCTGCCGGGCAGATAAGGGCAGCTGTGCCCTTG GGGCACCTGCTCACTCCCGCAGCCAGCCACTCCTCCAGGGCCAGCCCTTCCCTGA CTGAGTGACCACCTCTGCTGCCCCGAGGCCATGTAGGCCGTGCTTAGGCCTCTGTG GACACACTGCTGGGGACGGCGCCTGAGCTCTCAGGGGGACGAGGAACACCACCAT GCCCCGGGGCTTACCTGGCTGCGCTATCTTGGGATCTTCCTTGGCGTGGCCTTGGG GAATGAGCCTTTGGAGATGTGGCCCTTGACGCAGAATGAGGAGTGCAGTGTACG GGTTTTCTGCGGGACAAGCTGCAGTACAGGAGCCGACTTCAGTACATGAAACACTA CTTCCCCATCAACTACAAGATCAGTGTGCCTTACGAGGGGGTGTTTCAAGATCGCCA ACGTCACCAGGCTGAGGGGCCAGGTGAGCGAGCGGGAGCTGCGGTATCTGTGGGT CTTGGTGAGCCTCAGTGCCACTGAGTCGGTGCAGGACGTGCTGCTCGAGGGCCACC CATCCTGGAAGTACCTGCAGGAGGTGGAGACGCTGCTGCTGAATGTCCAGCAGGG CCTCACGGATGTGGAGTGCAGCCCCAAGGTGGAATCCGTGTTGTCCCTCTGAATG CCCCAGGGCCAAACCTGAAGCTGGTGCGGCCCAAAGCCCTGCTGGACAACCTGCTTC

	CGGGTCATGGAGCTGCTGTACTGCTCCTGCTGTAAACAAAGCTCCGTCCTAAACTG GCAGGACTGTGAGGTGCCAAGTCTCAGTCTTGCAGCCCAGAGCCCTCATTGCAGT ATGCGGCCACCCAGCTGTACCCTCCGCCCCCGTGGTCCCCCAGCTCCCCGCCTCACT CCACGGGCTCGGTGAGGCCGGTCAGGGCACAGGGCGAGGGCCTCTTGCCCTGAGC ACCCTGGATGGTGACTGCGGATAGGGGCAGCCAGACCAGTCCCACAGGAGTTCA ACTGGGTCTGAGACTTCAAGGGGTGGTGGTGGGAGCCCCCTTGGGAGAGGACCC CTGGGAAGGGTGTTCCTTTGAGGGGGATTCTGTGCCACAGCAGGGCTCAGCTT CCTGCCTTCCATAGCTGTATGGCCTCACCTGGAGCGGAGGGGACCTGGGGACCTG AAGGTGGATGGGGACACAGCTCCTGGCTTCTCCTGGTGCTGCCCTCACTGTCCCC CGCTAAAGGGGGTACTGAGCCTCCTGTGGCCCGCAGCAGTGAGGGCACAGCTGT GGGTGTCAGGGGAGACAGCCAGCACGGCGTGGCCATTCTATGACCCCCAGCCTG GCAGACTGGGGAGCTGGGGGCAGAGGGCGGTGCCAAGTGCCACATCTTGCCATAG TGGATGCTCTTCCAGTTCTTTTTTCTATTAAACACCCCACTTCCTTTGGAAAAAAA AAAAAAAAA (NM_001172771.1 Homo sapiens interleukin 34 (IL34), transcript variant 2, mRNA)
25	CTCGAAGCTCGGCGTCTGTGATGGTCTACGGGGCTTTCGAGGTGATCAGGCAGCGT CAGTCTTCAGCCGCTAAGCCGAGAAGGAGTCAGTCAGAGAGCCTTGGGCCAGAGT TCCAGGGGCTCTGGGAGTGGCTGCCAGAGCCAGATTTCACTGAGCAGCTGCAGT CGGAGAAATCAGAGAAAGCGTCACCCAGCCCCAGATTCCGAGGGGCCTGCCAGGG ACTCTCTCCTCCTGCTCCTTGGAAAGGAAGACCCCGAAAGACCCCCAAGCCACCGG CTCAGACCTGCTTCTGGGCTGCCATGGGACTTGCGGCCACCGCCCCCGGCTGTCC TCCACGCTGCCGGGCAGATAAGGGCAGCTGCTGCCCTTGGGGCACCTGCTCACTCC CGCAGCCCAGCCACTCCTCCAGGGCCAGCCCTTCCCTGACTGAGTGACCACCTCTG CTGCCCCGAGGCCATGTAGGCCGTGCTTAGGCCTCTGTGGACACACTGCTGGGGAC GGCGCTGAGCTCTCAGGGGGACGAGGAACACCACCATGCCCGGGGCTTACCT GGCTGCGCTATCTTGGGATCTTCCTTGGCGTGGCCTTGGGGAATGAGCCTTTGGAG ATGTGGCCCTTGACGCAGAATGAGGAGTGCAGTGTACGGGTTTTCTGCGGGACAA GCTGCAGTACAGGAGCCGACTTCAGTACATGAAACACTACTTCCCCATCAACTACA AGATCAGTGTGCCTTACGAGGGGGTGTTCAGAATCGCCAACGTCACCAGGCTGCAG AGGGCCCAGGTGAGCGAGCGGGAGCTGCGGTATCTGTGGGTCTTGGTGAGCCTCA GTGCCACTGAGTCGGTGCAAGACGTGCTGCTCGAGGGCCACCCATCCTGGAAGTAC CTGCAGGAGGTGGAGACGCTGCTGCTGAATGTCCAGCAGGGCCTCACGGATGTGG AGGTCAGCCCCAAGGTGGAATCCGTGTTGTCCCTCTTGAATGCCCCAGGGCCAAAC CTGAAGCTGGTGCGGCCAAAGCCCTGCTGGACAACCTGCTTCCGGGTCACTGGAGCT GCTGTACTGCTCCTGCTGTAAACAAAGCTCCGTCCTAAACTGGCAGGACTGTGAGG TGCCAAGTCCTCAGTCTTGAGCCAGAGCCCTCATTGCAGTATGCGGCCACCCAG CTGTACCTCCGCCCCCGTGGTCCCCCAGCTCCCCGCTCACTCCACGGGCTCGGTG AGGCCGGTCAGGGCACAGGGCGAGGGCCTCTTGCCCTGAGCACCTGGATGGTGA CTGCGGATAGGGGCAGCCAGACCAGTCCCACAGGAGTTCAACTGGGTCTGAGAC TTCAAGGGGTGGTGGTGGGAGCCCCCTTGGGAGAGGACCCCTGGGAAGGGTGTT TTTCCTTTGAGGGGGATTCTGTGCCACAGCAGGGCTCAGCTTCTGCCTTCCATAGC TGTCATGGCCTCACCTGGAGCGGAGGGGACCTGGGGACCTGAAGGTGGATGGGGA CACAGCTCCTGGCTTCTCCTGGTGCTGCCCTCACTGTCCCCCGCCTAAAGGGGGTA CTGAGCCTCCTGTGGCCCGCAGCAGTGAGGGCACAGCTGTGGGTTGCAGGGGAGA CAGCCAGCACGGCGTGGCCATTCTATGACCCCCAGCCTGGCAGACTGGGGAGCTG GGGGCAGAGGGCGGTGCCAAGTGCCACATCTTGCCATAGTGGATGCTCTTCCAGTT TCTTTTTTCTATTAAACACCCCACTTCCTTTGGAAAAA (NM_001172772.1 Homo sapiens interleukin 34 (IL34), transcript variant 3, mRNA)
26	CATCAGACGGGAAGCCTGGACTGTGGGTGGGGGCAGCCTCAGCCTCTCCAACCTG GCACCCACTGCCCCTGGCCCTTAGGCACCTGCTTGGGGTCTTGAGCCCCCTTAAGG CCACCAGCAAATCCTAGGAGACCGAGTCTTGGCACGTGAACAGAGCCAGATTCA

	CACTGAGCAGCTGCAGTCGGAGAAATCAGAGAAAGCGTCACCCAGCCCCAGATTC CGAGGGGCTGCCAGGGACTCTCTCCTCCTGCTCCTTGGAAGGAAGACCCCGAAA GACCCCAAGCCACCGGCTCAGACCTGCTTCTGGGCTGCCATGGGACTTGCGGCCA CCGCCCCCGGCTGTCTCCACGCTGCCGGGCAGATAAGGGCAGCTGCTGCCCTTG GGCACCTGCTCACTCCCGCAGCCCAGCCACTCCTCCAGGGCCAGCCCTTCCCTGA CTGAGTGACCACCTCTGCTGCCCGAGGCCATGTAGGCCGTGCTTAGGCCTCTGTG GACACACTGCTGGGGACGGCGCCTGAGCTCTCAGGGGGACGAGGAACACCACCAT GCCCCGGGGCTTACCTGGCTGCGCTATCTTGGGATCTTCCTTGGCGTGGCCTTGGG GAATGAGCCTTTGGAGATGTGGCCCTTGACGCAGAATGAGGAGTGCAGTGTACG GGTTTTCTGCGGGACAAGCTGCAGTACAGGAGCCGACTTCAGTACATGAAACACTA CTTCCCCATCAACTACAAGATCAGTGTGCCTTACGAGGGGGTGTTCAGAATCGCCA ACGTCACCAGGCTGCAGAGGGGCCAGGTGAGCGAGCGGGAGCTGCGGTATCTGTG GGTCTTGGTGAGCCTCAGTGCCACTGAGTCGGTGCAGGACGTGCTGCTCGAGGGCC ACCCATCTTGGAAGTACCTGCAGGAGGTGGAGACGCTGCTGCTGAATGTCCAGCA GGCCCTCACGGATGTGGAGGTGAGCCCCAAGGTGGAATCCGTGTTGTCCCTCTTGA ATGCCCCAGGGCCAAACCTGAAGCTGGTGCGGGCCAAAGCCCTGCTGGACAACCTG CTTCCGGGTCATGGAGCTGCTGTACTGCTCCTGCTGTAAACAAAGCTCCGTCCTAA ACTGGCAGGACTGTGAGGTGCCAAGTCCTCAGTCTTGACGCCAGAGCCCTCATTG CAGTATGCGGCCACCCAGCTGTACCCTCCGCCCCCGTGGTCCCCCAGCTCCCCGCC TCACTCCACGGGCTCGGTGAGGCCGGTCAGGGCACAGGGCGAGGGCCTCTTGCCCT GAGCACCCCTGGATGGTGACTGCGGATAGGGGCAGCCAGACCAGCTCCCACAGGAG TTCAACTGGGTCTGAGACTTCAAGGGGTGGTGGTGGGAGCCCCCTTGGGAGAGG ACCCCTGGGAAGGGTGTTCCTTTGAGGGGGATTCTGTGCCACAGCAGGGCTCA GCTTCTGCCTTCCATAGCTGTATGGCCTCACCTGGAGCGGAGGGGACCTGGGGA CCTGAAGGTGGATGGGGACACAGCTCCTGGCTTCTCCTGGTGTGCTGCCCTACTGTG CCCCCGCCTAAAGGGGGTACTGAGCCTCCTGTGGCCCGCAGCAGTGAGGGCACAG CTGTGGGTGTCAGGGGAGACAGCCAGCACGGCGTGGCCATTCTATGACCCCCCAGC CTGGCAGACTGGGGAGCTGGGGGCAGAGGGCGGTGCCAAGTGCCACATCTTGCCA TAGTGATGCTCTTCCAGTTTCTTTTTTCTATTAAACACCCCACTTCCTTTGAAAAA AAAAAAAAAAAAA (NM 152456.2 Homo sapiens interleukin 34 (IL34), transcript variant 1, mRNA)
27	TCTAGAAACAAATCTGAGCTGCTGTAGCGTGTGTCCCCCGCCCTCCGAGAAGACTG TAGGCTGAACACACAACAGCCTGTCCACACAGCATTTACTGCCACAGCCCCAAG GCACCTGCTCCAGGCCTTGAGCTGCTCAAAGCTTCCGGCGACGGCGACTGTAGAAG CCCACATCCTGGCAGGTGAGCAGAGCTAGACTTCACGCTGAGCAGCTGCAGCCGA AGAAACCAGAGAAAGCGCCACCCAGCCCCAGATTCCAAGAGGCCTGCCTGTGGG GACTCCCTTCTTCTTCTCCTCCTCCTCCTCCTCCTCCTGAGACCCTCAAAGTCAC TGTCTCACACCAGTTCCTGAGCTGCAATGGGACTGGCAGCCACTGCCCCCTGGCTG TCCTCCGCCCTGCCAAGCAGATAGGGCGGTAGCTACCCCAGGAGTATCTGCCCCCT CTCAGCCTGGACCAGCCCTTCTTGACTTGGTGGCCACCTTTGCTGACCTAAGGCC ATGTAGTCCCTGGCTCAGGCCTCTGGACACACTTCTGGGGACAGTGCCTCTGCTCTC AGGGGGCACCCAGTGCACCACCATGCCCTGGGGACTCGCCTGGCTATACTGTCTTG GGATCCTACTTGACGTGGCTTTGGGAAACGAGAATTTGGAGATATGGACTCTGACC CAAGATAAGGAGTGTGACCTTACAGGCTACCTTCGGGGCAAGCTGCAGTACAAGA ACCGGCTTCAGTACATGAAACATTACTTCCCCATCAACTACAGGATTGCTGTGCCTT ATGAGGGGGTACTCAGAGTGGCCAACATCACAAGGCTGCAGAAGGCTCACGTGAG TGAGCGAGAGCTTCGGTACCTGTGGGTCTTGGTGAGTCTCAATGCCACTGAGTCTG TGATGGATGTACTTCTCGAGGGGCCACCCGTCCTGGAAGTATCTACAGGAGGTTTCA ACATTGCTGGAGAACGTACAGCGGAGCCTCATGGATGTGGAGATTGGCCCTCACGT GGAAGCTGTGTTATCTTCTGAGTACTCCAGGCCTAAGCCTGAAGCTGGTGCGGC CCAAAGCCTTGCTGGACAACCTGCTTCCGGGTCATGGAAGTGTGTACTGTTCTTGCT

	GTAAACAAAGCCCCATCTTAAAATGGCAGGACTGCGAGCTGCCAGGCTCCATCCC CACAGTCCGGGGTCTTGATGCAATGTACAGCTACAAATGTGTACCCTTTGTCTCG GCAGACCCCCACCTCCCTGCCCGGATCCCCAAGCTCAAGCCATGGCTCGTTGCCCT GAGCAATCTATGTATTGACCCTGGGTCTGGGCCACTGGATGCATTTAGAAAATGGC CTGGATCTGAGACCTTCCTGGATGGTGGGTAGGTAGCCCTTTAGGAAAGGCCTTC AGGGAAAGGCCTCCTGCCTTCTCTTCCACACCCAGGCTTCCTGCCTTCTGGTGGTAT CACGGAGCCTCAGGGGAGGGATGTGGAGGGGAGGGCTGGTTGCTATCCCTTACAG CCACCTCTGCTTGTGGCCCGGTCCGAAATCTGCAAACACTGTTGCCCCACAGGCGG GTGGCCAAGAACGACGGGGTGCCTCCTAGGATCCCAGGACTTACTGGGGCTTTTCA GTTGTGCTGACTGTGGGGTGTGGGCCGAGGACCAACCAAATGCCGTATCTTGCCA TGACGATGCTCTTCTGGTTCCTTTTTCTATTAAATGGCCATTTATTTGGTTTGGCTTG CAAAAAAA (NM_001135100.2 <i>Mus musculus</i> interleukin 34 (Il34), transcript variant 1, mRNA)
28	TCTAGAAACAAATCTGAGCTGCTGTAGCGTGTGTCCCCGCCCTCCGAGAAGACTG TAGGCTGAACACACAACAGCCTGTCCCACACAGCATTTACTGCCACAGCCCCAAG GCACCTGCTCCAGGCCTTGAGCTGCTCAAAGCTTCCGGCGACGGCGACTGTAGAAG CCCACATCCTGGCAGGTGAGCAGAGCTAGACTTCACGCTGAGCAGCTGCAGCCGA AGAAACCAGAGAAAGCGCCACCCAGCCCCAGATTCCAAGAGGCCTGCCTGCTGGG GACTCCCTTCTTCTTCTCCTCCTCCTCCTCCTCCTCCTGAGACCCTCAAAGTCAC TGTCTCACACCAGTTCCTGAGCTGCAATGGGACTGGCAGCCACTGCCCCCTGGCTG TCCTCCGCCCTGCCAAGCAGATAGGGCGGTAGCTACCCAGGAGTATCTGCCCCCT CTCAGCCTGGACCAGCCCTTTCTGGACTTGGTGGCCACCTTTGCTGACCTAAGGCC ATGTAGTCCCTGGCTCAGGCCTCTGGACACACTTCTGGGGACAGTGCCTCTGCTCTC AGGGGGCACCCAGTGCACCACCATGCCCTGGGGACTCGCTGGCTATACTGTCTTG GGATCTTACTTGACGTGGCTTTGGGAAACGAGAATTTGGAGATATGGACTCTGACC CAAGATAAGGAGTGTGACCTTACAGGCTACCTTCGGGGCAAGCTGCAGTACAAGA ACCGGCTTCAGTACATGAAACATTACTTCCCCATCAACTACAGGATTGCTGTGCCTT ATGAGGGGGTACTCAGAGTGGCCAACATCACAAGGCTGCAGAAGGCTCACGTGAG TGAGCGAGAGCTTCGGTACCTGTGGGTCTTGGTGAGTCTCAATGCCACTGAGTCTG TGATGGATGTACTTCTCGAGGGCCACCCGCTCTGGAAGTATCTACAGGAGGTTTCA ACATTGCTGGAGAACGTACAGCGGAGCCTCATGGCCGTTGGTGTACACCTGCCGGG ACACGTACTTGTGACCCTGCTCAGCCAGCTGCCTGGCCTCCCCAGCCCATGGGGCA GATCATTTGACACCAGCTGGGAGCTTCTGATGATGAAAGGATGTGGAGATTGGCCC TCACGTGGAAGCTGTGTTATCTCTTCTGAGTACTCCAGGCCTAAGCCTGAAGCTGG TGCGGCCCCAAAGCCTTGCTGGACAACCTGCTTCCGGGTCATGGAAGTGTACTGT TCTTGCTGTAAACAAAGCCCCATCTTAAAATGGCAGGACTGCGAGCTGCCAGGCT CCATCCCCACAGTCCGGGGTCTTGATGCAATGTACAGCTACAAATGTGTACCCTT TGTCTCGGCAGACCCCCACCTCCCTGCCCGGATCCCCAAGCTCAAGCCATGGCTCG TTGCCCTGAGCAATCTATGTATTGACCCTGGGTCTGGGCCACTGGATGCATTTAGAA AACTGGCCTGGATCTGAGACCTTCCTGGATGGTGGGTAGGTAGCCCTTTAGGAAA GGCCTTCAGGGAAAGGCCTCCTGCCTTCTTCCACACCCAGGCTTCCTGCCTTCTG GTGGTATCACGGAGCCTCAGGGGAGGGATGTGGAGGGGAGGGCTGGTTGCTATCC CTTACAGCCACCTCTGCTTGTGGCCCGGTCCGAAATCTGCAAACACTGTTGCCCCA CAGGCGGGTGGCCAAGAACGACGGGGTGCCTCCTAGGATCCCAGGACTTACTGGG GCTTTTCAAGTTGTGCTGACTGTGGGGTGTGGGCCGAGGACCAACCAAATGCCGTA TCTTGCCATGACGATGCTCTTCTGGTTCCTTTTTCTATTAAATGGCCATTTATTTGGT TTGGCTTGCAAAAAA (NM_029646.3 <i>Mus musculus</i> interleukin 34 (Il34), transcript variant 2, mRNA)
29	5'-ACAGGAGCCGACTTCAGTAC-3'
30	5'-ACCAAGACCCACAGATACCG-3'

31	5'-AAGATGACCCAGATCATGTTTGAGACC-3'
32	5'-AGCCAGTCCAGACGCAGGAT-3'
33	5'-AACACGTCTATACGC-3'
34	5'-CTGCTCAACTTTCTGCGAAG-3'
35	5'-CTCATCTCCACATAGGTGTC-3'
36	5'-GGACACAGAGGTTTCAGTGG-3'
37	3'-GGTGACTTTGGAGACACAGG-5'
38	5'-GGAGAATGTTGTGCAGTTTGC-3'
39	3'-CGTTTGACGTGTTGTAAGAGG-5'
40	5'-CxTxTxTxGGGCXGCACCAGCxTxTxCx-3'
41	5'-CxTxTxTGGGCXGCACCAGCTxTxCx-3'
42	5'-TxCxCxAxTGACCXGGAAGCAxGxTxTx-3'
43	5'-TxCxCxATGACCXGGAAGCAGxTxTx-3'
44	AGACGGGAAGCCTGGACTGTGGGTTGGGGGCAGCCTCAGCCTCTCCAACCTGGCA CCCACTGCCCCGTGGCCCTTAGGCACCTGCTTGGGGTCCTGGAGCCCCCTTAAGGCCA CCAGCAAATCCTAGGAGACCGAGTCTTGGCACGTGAACAGAGCCAGATTTCACACT GAGCAGCTGCAGTCGGAGAAATCAGAGAAAGCGTCACCCAGCCCCAGATTCCGAG GGCCTGCCAGGGACTCTCTCCTCCTGCTCCTTGGAAAGGAAGACCCCGAAAGACC CCCCAGCCACCGGCTCAGACCTGCTTCTGGGCTGCCATGGGACTTGCGGCCACCGC CCCCCGGTGTCTCCACGCTGCCGGGCAGATAAGGGCAGCTGCTGCCCTTGGGGC ACCTGCTCACTCCCGCAGCCCAGCCACTCCTCCAGGGCCAGCCCTTCCCTGACTGA GTGACCACCTCTGCTGCCCCGAGGCCATGTAGGCCGTGCTTAGGCCTCTGTGGACA CACTGCTGGGGACGGCGCCTGAGCTCTCAGGGGGACGAGGAACACCACCATGCCC CGGGGCTTACCTGGCTGCGCTATCTTGGGATCTTCCTTGGCGTGGCCTTGGGGAAT GAGCCTTTGGAGATGTGGCCCTTGACGCAGAATGAGGAGTGCAGTGTACCGGTTT TCTGCGGGACAAGCTGCAGTACAGGAGCCGACTTCAGTACATGAAACACTACTTCC CCATCAACTACAAGATCAGTGTGCCTTACGAGGGGGTGTTCAGAATCGCCAACGTC ACCAGGCTGCAGAGGGGCCAGGTGAGCGAGCGGGAGCTGCGGTATCTGTGGGTCT TGGTGAGCCTCAGTGCCACTGAGTCGGTGCAGGACGTGCTGCTCGAGGGCCACCCA TCCTGGAAGTACCTGCAGGAGGTGGAGACGCTGCTGCTGAATGTCCAGCAGGGCCT CACGGATGTGGAGGTCAGCCCCAAGGTGGAATCCGTGTTGTCCCTCTTGAATGCCC CAGGGCCAAACCTGAAGCTGGTGCGGCCCAAAGCCCTGCTGGACAACCTGCTTCCG GGTCATGGAGCTGCTGTACTGCTCCTGCTGTAAACAAAGCTCCGTCCTAAACTGGC AGGACTGTGAGGTGCCAAGTCCTCAGTCTTGCAGCCCAGAGCCCTCATTGCAGTAT GCGGCCACCCAGCTGTACCCTCCGCCCCCGTGGTCCCCCAGCTCCCCGCCTCACTCC ACGGGCTCGGTGAGGCCGGTCAGGGCACAGGGCGAGGGCCTCTTGCCCTGAGCAC CCTGGATGGTGACTGCGGATAGGGGCAGCCAGACCAGCTCCCACAGGAGTTCAAC TGGGTCTGAGACTTCAAGGGGTGGTGGTGGGAGCCCCCTTGGGAGAGGACCCCT GGGAAGGGTGTTTTTCTTTGAGGGGGATTCTGTGCCACAGCAGGGCTCAGCTTCC TGCTTTCCATAGCTGTCATGGCCTCACCTGGAGCGGAGGGGACCTGGGGACCTGAA GGTGGATGGGGACACAGCTCCTGGCTTCTCCTGGTGCTGCCCTCACTGTCCCCCG CCTAAAGGGGTACTGAGCCTCCTGTGGCCCGCAGCAGTGAGGGCACAGCTGTGG GTTGCAGGGGAGACAGCCAGCACGGCGTGGCCATTCTATGACCCCCAGCCTGGC AGACTGGGGAGCTGGGGGCAGAGGGCGGTGCCAAGTGCCACATCTTGCCATAGTG GATGCTCTTCCAGTTTCTTTTTTCTATTAAACACCCCACTTCCTTTGG NM 152456.3 Homo sapiens interleukin 34 (IL34), transcript variant 1, mRNA
45	AGACGGGAAGCCTGGACTGTGGGTTGGGGGCAGCCTCAGCCTCTCCAACCTGGCA CCCACTGCCCCGTGGCCCTTAGGCACCTGCTTGGGGTCCTGGAGCCCCCTTAAGGCCA CCAGCAAATCCTAGGAGACCGAGTCTTGGCACGTGAACAGAGCCAGATTTCACACT GAGCAGCTGCAGTCGGAGAAATCAGAGAAAGCGTCACCCAGCCCCAGATTCCGAG

	GGGCCTGCCAGGGACTCTCTCCTCCTGCTCCTTGGAAGGAAGACCCCGAAAGACC CCCAAGCCACCGGCTCAGACCTGCTTCTGGGCTGCCATGGGACTTGCGGCCACCGC CCCCCGGCTGTCTCCACGCTGCCGGGCAGATAAGGGCAGCTGCTGCCCTTGGGGC ACCTGCTCACTCCCGCAGCCCAGCCACTCCTCCAGGGCCAGCCCTTCCCTGACTGA GTGACCACCTCTGCTGCCCCGAGGCCATGTAGGCCGTGCTTAGGCCCTGTGGACA CACTGCTGGGGACGGCGCCTGAGCTCTCAGGGGGACGAGGAACACCACCATGCCC CGGGGCTTACCTGGCTGCGCTATCTTGGGATCTTCTTGGCGTGGCCTTGGGGAAT GAGCCTTTGGAGATGTGGCCCTTGACGCAGAATGAGGAGTGCAGTGTACGGGTTT TCTGCGGGACAAGCTGCAGTACAGGAGCCGACTTCAGTACATGAAACACTACTTCC CCATCAACTACAAGATCAGTGTGCCTTACGAGGGGGTGTTCAGAATCGCCAACGTC ACCAGGCTGAGGGCCCAGGTGAGCGAGCGGGAGCTGCGGTATCTGTGGGTCTTGG TGAGCCTCAGTGCCACTGAGTCGGTGCAGGACGTGCTGCTCGAGGGGCCACCCATCC TGAAGTACCTGCAGGAGGTGGAGACGCTGCTGCTGAATGTCCAGCAGGGCCTCA CGGATGTGGAGGTGAGCCCCAAGGTGGAATCCGTGTTGTCCCTCTTGAATGCCCA GGGGAAACCTGAAGCTGGTGCGGCCCAAAGCCCTGCTGGACAACAGCTTCCGGGT CATGGAGCTGCTGTACTGCTCCTGCTGTAAACAAAGCTCCGTCTAAACTGGCAGG ACTGTGAGGTGCCAAGTCCTCAGTCTTGACGCCAGAGCCCTCATTGCAGTATGCG GCCACCCAGCTGTACCCTCCGCCCCCGTGGTCCCCCAGCTCCCCGCCTCACTCCACG GGCTCGGTGAGGCCGGTCAGGGCACAGGGCGAGGGCCTCTTGGCCTGAGCACCCCT GGATGGTGACTGCGGATAGGGGCGAGCCAGACCAGCTCCACAGGAGTTCAACTGG GTCTGAGACTTCAAGGGGTGGTGGTGGGAGCCCCCCTTGGGAGAGGACCCCTGGG AAGGGTGTTCCTTTGAGGGGGATTCTGTGCCACAGCAGGGCTCAGTTCCTGTC CTTCCATAGCTGTCATGGCCTCACCTGGAGCGGAGGGGACCTGGGGACCTGAAGGT GGATGGGGACACAGCTCCTGGCTTCTCCTGGTGTGCCCTCACTGTCCCCCGCCTA AAGGGGTACTGAGCCTCCTGTGGCCCCGAGCAGTGAGGGCACAGCTGTGGGTG CAGGGGAGACAGCCAGCACGGCGTGGCCATTCTATGACCCCCAGCCTGGCAGAC TGGGGAGCTGGGGGCAGAGGGCGGTGCCAAGTGCCACATCTTGCCATAGTGGATG CTCTTCCAGTTTCTTTTCTATTAAACACCCCACTTCCTTTTG NM_001172771.2 Homo sapiens interleukin 34 (IL34), transcript variant 2, mRNA
46	ATGATCAGAGATGTCTGCATGAGGAACAGAGGCTGCTTTGTGGATTGGGGAGCTCC CCATTAGCGGAGGCAGAGCCAGATTTCACTGAGCAGCTGCAGTCGGAGAAATC AGAGAAAGCGTCACCCAGCCCCAGATTCCGAGGGGGCTGCCAGGGACTCTCTCCTC CTGCTCCTTGGAAGGAAGACCCCGAAAGACCCCAAGCCACCGGCTCAGACCTG CTTCTGGGCTGCCATGGGACTTGCGGCCACCGCCCCCGGCTGTCTCCACGCTGC CGGGCAGATAAGGGCAGCTGCTGCCCTTGGGGCACCTGCTCACTCCCGCAGCCCAG CCACTCCTCCAGGGCCAGCCCTTCCCTGACTGAGTGACCACCTCTGCTGCCCCGAG GCCATGTAGGCCGTGCTTAGGCCTCTGTGGACACACTGCTGGGGACGGCGCCTGAG CTCTCAGGGGGACGAGGAACACCACCATGCCCGGGGCTTACCTGGCTGCGCTAT CTTGGGATCTTCTTGGCGTGGCCTTGGGGAATGAGCCTTTGGAGATGTGGCCCTT GACGCAGAATGAGGAGTGCAGTGTACGGGTTTTCTGCGGGACAAGCTGCAGTAC AGGAGCCGACTTCAGTACATGAAACACTACTTCCCCATCAACTACAAGATCAGTGT GCCTTACGAGGGGGTGTTCAGAATCGCCAACGTCACCAGGCTGCAGAGGGGCCAG GTGAGCGAGCGGGAGCTGCGGTATCTGTGGGTCTTGGTGAGCCTCAGTGCCACTGA GTCGGTGCAGGACGTGCTGCTCGAGGGCCACCCATCCTGGAAGTACCTGCAGGAG GTGGAGACGCTGCTGCTGAATGTCCAGCAGGGCCTCACGGATGTGGAGGTGAGCC CCAAGGTGGAATCCGTGTTGTCCCTCTTGAATGCCCCAGGGCCAAACCTGAAGCTG GTGCGGCCCAAAGCCCTGCTGGACAACCTGCTTCCGGGTGATGGAGCTGCTGTACTG CTCCTGCTGTAAACAAAGCTCCGTCTAAACTGGCAGGACTGTGAGGTGCCAAGTC CTCAGTCTTGACAGCCAGAGCCCTCATTGCAGTATGCGGCCACCCAGCTGTACCCT CCGCCCCCGTGGTCCCCAGCTCCCCGCCTCACTCCACGGGCTCGGTGAGGCCGGT CAGGGCACAGGGCGAGGGCCTCTTGGCCTGAGCACCCCTGGATGGTGACTGCGGAT

	AGGGGCAGCCAGACCAGCTCCACAGGAGTTCAACTGGGTCTGAGACTTCAAGGG GTGGTGGTGGGAGCCCCCTTGGGAGAGGACCCCTGGGAAGGGTGTTCCTTTG AGGGGGATTCTGTGCCACAGCAGGGCTCAGCTTCCTGCCTTCCATAGCTGTCATGG CCTCACCTGGAGCGGAGGGGACCTGGGGACCTGAAGGTGGATGGGGACACAGCTC CTGGCTTCTCCTGGTGTGCCCTCACTGTCCCCCGCCTAAAGGGGGTACTGAGCCT CCTGTGGCCCGCAGCAGTGAGGGCACAGCTGTGGGTTCAGGGGAGACAGCCAGC ACGGCGTGGCCATTCTATGACCCCCCAGCCTGGCAGACTGGGGAGCTGGGGGCAG AGGGCGGTGCCAAGTGCCACATCTTGCCATAGTGGATGCTCTTCCAGTTTCTTTTT CTATTAAACACCCCACTTCCTTTGG NM_001393493.1 Homo sapiens interleukin 34 (IL34), transcript variant 4, mRNA
47	ATGATCAGAGATGTCTGCATGAGGAACAGAGGCTGCTTTGTGGATTGGGGAGCTCC CCATTAGCGGAGGCAGAGCCAGATTTCACTGAGCAGCTGCAGTCGGAGAAATC AGAGAAAGCGTCACCCAGCCCCAGATTCGAGGGGCCCTGCCAGGGACTCTCTCCTC CTGCTCCTTGGAAGGAAGACCCCGAAAGACCCCAAGCCACCGGCTCAGACCTG CTTCTGGGCTGCCATGGGACTTGCGGCCACCGCCCCCGGCTGTCTCCACGCTG CGGCGAGATAAGGGCAGCTGCTGCCCTTGGGGCACCTGCTCACTCCCGCAGCCAG CCTCTCTCCAGGGCCAGCCCTTCCCTGACTGAGTGACCACCTCTGCTGCCCCGAG GCCATGTAGGCCGTGCTTAGGCCTCTGTGGACACACTGCTGGGGACGGCGCTGAG CTCTCAGGGGGACGAGGAACACCACCATGCCCCGGGGCTTCACCTGGCTGCGCTAT CTTGGGATCTTCCTTGGCGTGGCCTTGGGGAATGAGCCTTTGGAGATGTGGCCCTT GACGCAGAATGAGGAGTGCACTGTACCGGTTTTCTGCGGGACAAGCTGCAGTAC AGGAGCCGACTTCAGTACATGAAACACTACTTCCCCATCAACTACAAGATCAGTGT GCCTTACGAGGGGGTGTTCAGAAATCGCCAACGTCAACAGGCTGAGGGCCCAGGTG AGCGAGCGGGAGCTGCGGTATCTGTGGGTCTTGGTGAGCCTCAGTGCCACTGAGTC GGTGACGAGCTGCTGCTCGAGGGCCACCCATCCTGGAAGTACCTGCAGGAGGTG GAGACGCTGCTGCTGAATGTCCAGCAGGGCCTACGGATGTGGAGGTGAGCCCCA AGGTGGAATCCGTGTTGTCCCTCTTGAATGCCCCAGGGCCAAACCTGAAGCTGGTG CGGCCCCAAGCCCTGCTGGACAACTGCTTCCGGGTCTATGGAGCTGCTGTACTGCTC CTGCTGTAAACAAAGCTCCGTCCTAACTGGCAGGACTGTGAGGTGCCAAGTCCTC AGTCTTGCAGCCCAGAGCCCTCATTGCAGTATGCGGCCACCCAGCTGTACCCTCCG CCCCCGTGGTCCCCCAGCTCCCCGCCTCACTCCACGGGCTCGGTGAGGCCGGTCAG GGCACAGGGCGAGGGCCTCTTGCCCTGAGCACCCCTGGATGGTGACTGCGGATAGG GGCAGCCAGACCAGCTCCACAGGAGTTCAACTGGGTCTGAGACTTCAAGGGGTG GTGGTGGGAGCCCCCTTGGGAGAGGACCCCTGGGAAGGGTGTTCCTTTGAGG GGGATTCTGTGCCACAGCAGGGCTCAGCTTCCTGCCTTCCATAGCTGTCATGGCCTC ACCTGGAGCGGAGGGGACCTGGGGACCTGAAGGTGGATGGGGACACAGCTCCTTG CTTCTCCTGGTGCTGCCCTCACTGTCCCCCGCCTAAAGGGGGTACTGAGCCTCCTG TGGCCCGCAGCAGTGAGGGCACAGCTGTGGGTTCAGGGGAGACAGCCAGCACGG CGTGGCCATTCTATGACCCCCCAGCCTGGCAGACTGGGGAGCTGGGGGCAGAGGG CGGTGCCAAGTGCCACATCTTGCCATAGTGGATGCTCTTCCAGTTTCTTTTTCTAT TAAACACCCCACTTCCTTTGG NM_001393495.1 Homo sapiens interleukin 34 (IL34), transcript variant 6, mRNA
48	CTCGAAGCTCGGCGTCTGTGATGGTCTACGGGGCTTTCGAGGTGATCAGGCAGCGT CAGTCTTCAGCCGCTAAGCCGAGAAGGAGTCAGTCAGAGAGCCTTGGGCCAGAGT TCCAGGGGCTCTGGGAGTGGCTGCCAGAGCCAGATTTCACTGAGCAGCTGCAGT CGGAGAAATCAGAGAAAGCGTCACCCAGCCCCAGATTCCGAGGGGCCCTGCCAGGG ACTCTCTCCTCCTGCTCCTTGGAAGGAAGACCCCGAAAGACCCCAAGCCACCGG CTCAGACCTGCTTCTGGGCTGCCATGGGACTTGCGGCCACCGCCCCCGGCTGTCC TCCACGCTGCCGGGCAGATAAGGGCAGCTGCTGCCCTTGGGGCACCTGCTCACTCC CGCAGCCCAGCCACTCCTCCAGGGCCAGCCCTTCCCTGACTGAGTGACCACCTCTG CTGCCCCGAGGCCATGTAGGCCGTGCTTAGGCCTCTGTGGACACACTGCTGGGGAC

	GGCGCCTGAGCTCTCAGGGGGACGAGGAACACCACCATGCCCCGGGGCTTCACCT GGCTGCGCTATCTTGGGATCTTCCTTGGCGTGGCCTTGGGGAATGAGCCTTTGGAG ATGTGGCCCTTGACGCAGAATGAGGAGTGCCTGTACGGGTTTTCTGCGGGACAA GCTGCAGTACAGGAGCCGACTTCAGTACATGAAACACTACTTCCCCATCAACTACA AGATCAGTGTGCCTTACGAGGGGGTGTTCAGAATCGCCAACGTACACAGGCTGCAG AGGGCCCAGGTGAGCGAGCGGGAGCTGCGGTATCTGTGGGTCTTGGTGAGCCTCA GTGCCACTGAGTCGGTGCAGGACGTGCTGCTCGAGGGCCACCCATCCTGGAAGTAC CTGCAGGAGGTGGAGACGCTGCTGCTGAATGTCCAGCAGGGCCTCACGGATGTGG AGGTCAGCCCCAAGGTGGAATCCGTGTTGTCCCTCTTGAATGCCCCAGGGCCAAAC CTGAAGCTGGTGCGGCCCAAAGCCCTGCTGGACAACCTGCTTCCGGGTCATGGAGCT GCTGTACTGCTCCTGCTGTAAACAAAGCTCCGTCTAAACTGGCAGGACTGTGAGG TGCCAAAGTCCTCAGTCTTGAGCCCAGAGCCCTCATTGCAGTATGCGGCCACCCAG CTGTACCCTCCGCCCCCGTGGTCCCCCAGCTCCCCGCTCACTCCACGGGCTCGGTG AGGCCGGTCAGGGCACAGGGCGAGGGCCTCTTGCCCTGAGCACCCCTGGATGGTGA CTGCGATAGGGGCAGCCAGCAGCTCCACAGGAGTTCAACTGGGTCTGAGAC TTCAAGGGGTGGTGGTGGGAGCCCCCTTGGGAGAGGACCCCTGGGAAGGGTGTT TTTTCTTTGAGGGGGATTCTGTGCCACAGCAGGGCTCAGCTTCCTGCCTTCCATAGC TGTATGGCCTCACCTGGAGCGGAGGGGACCTGGGGACCTGAAGGTGGATGGGGA CACAGCTCCTGGCTTCTCCTGGTGTGCCCTCACTGTCCCCCGCCTAAAGGGGGTA CTGAGCCTCCTGTGGCCCGCAGCAGTGAGGGCACAGCTGTGGGTGTCAGGGGAGA CAGCCAGCACGGCGTGGCCATTCTATGACCCCCCAGCCTGGCAGACTGGGGAGCTG GGGGCAGAGGGCGGTGCCAAGTGCCACATCTTGCCATAGTGGATGCTCTTCCAGTT TCTTTTTTCTATTAAACACCCCACTTCTTTGG NM_001172772.2 Homo sapiens interleukin 34 (IL34), transcript variant 3, mRNA
49	ACACTGAGCAGCTGCAGTCGGAGAAATCAGAGAAAGCGTCACCCAGCCCCAGATT CCGAGGGGCTGCCAGGGACTCTCTCCTCCTGCTCCTTGGAAGGAAGACCCCGAA AGACCCCCAAGCCACCGGCTCAGACCTGCTTCTGGGCTGCCATGGGACTTGCGGCC ACCGCCCCCGGCTGTCTCCACGCTGCCGGGCAGATAAGGGCAGCTGCTGCCCTT GGGGCACCTGCTCACTCCCGCAGCCCAGCCACTCCTCCAGGGCCAGCCCTTCCCTG ACTGAGTGACCACCTCTGCTGCCCCGAGGCCATGTAGGCCGTGCTTAGGCCTCTGT GGACACACTGCTGGGGACGGCGCTGAGCTCTCAGGGGGACGAGGAACACCACCA TGCCCCGGGGCTTACCTGGCTGCGCTATCTTGGGATCTTCCTTGGCGTGGCCTTGG GGAATGAGCCTTTGGAGATGTGGCCCTTGACGCAGAATGAGGAGTGCCTGTAC GGGTTTTCTGCGGGACAAGCTGCAGTACAGGAGCCGACTTCAGTACATGAAACACT ACTTCCCCATCAACTACAAGATCAGTGTGCCTTACGAGGGGGTGTTCAGAATCGCC AACGTCAACAGGCTGCAGAGGGGCCAGGTGAGCGAGCGGGAGCTGCGGTATCTGT GGGTCTTGGTGAGCCTCAGTGCCACTGAGTCGGTGCAGGACGTGCTGCTCGAGGGC CACCCATCCTGGAAGTACCTGCAGGAGGTGGAGACGCTGCTGCTGAATGTCCAGCA GGGCCTCACGGATGTGGAGGTGAGCCCCAAGGTGGAATCCGTGTTGTCCCTCTTGA ATGCCCCAGGGCCAAACCTGAAGCTGGTGCGGCCCAAAGCCCTGCTGGACAACCTG CTTCCGGGTCATGGAGCTGCTGTACTGCTCCTGCTGTAAACAAAGCTCCGTCTTAA ACTGGCAGGACTGTGAGGTGCCAAGTCTCAGTCTTGACGCCAGAGCCCTCATTG CAGTATGCGGCCACCCAGCTGTACCCTCCGCCCCCGTGGTCCCCCAGCTCCCCGCC TCACTCCACGGGCTCGGTGAGGCCGGTCAGGGCACAGGGCGAGGGCCTCTTGCCCT GAGCACCCCTGGATGGTGACTGCGGATAGGGGCAGCCAGACCAGCTCCCACAGGAG TTCAACTGGGTCTGAGACTTCAAGGGGTGGTGGTGGGAGCCCCCTTGGGAGAGG ACCCCTGGGAAGGGTGTTTTCTTTGAGGGGGATTCTGTGCCACAGCAGGGCTCA GCTTCTGCCTTCCATAGCTGTATGGCCTCACCTGGAGCGGAGGGGACCTGGGGA CCTGAAGGTGGATGGGGACACAGCTCCTGGCTTCTCCTGGTGTGCCCTCACTGTC CCCCCGCCTAAAGGGGGTACTGAGCCTCCTGTGGCCCGCAGCAGTGAGGGCACAG CTGTGGGTGTCAGGGGAGACAGCCAGCAGGCGTGGCCATTCTATGACCCCCCAGC

	CTGGCAGACTGGGGAGCTGGGGGCAGAGGGCGGTGCCAAGTGCCACATCTTGCCA TAGTGATGCTCTTCCAGTTTCTTTTTTCTATTAAACACCCCACTTCCTTTGG NM_001393494.1 Homo sapiens interleukin 34 (IL34), transcript variant 5, mRNA
50	ACACTGAGCAGCTGCAGTCGGAGAAATCAGAGAAAGCGTCACCCAGCCCCAGATT CCGAGGGGCTGCCAGGGACTCTCTCCTCCTGCTCCTTGGAAGGAAGACCCCGAA AGACCCCAAGCCACCGGCTCAGACCTGCTTCTGGGCTGCCATGGGACTTGCGGCC ACCGCCCCCGGCTGTCTCCACGCTGCCGGGCAGATAAGGGCAGCTGCTGCCCTT GGGGCACCTGCTCACTCCCGCAGCCCAGCCACTCCTCCAGGGCCAGCCCTTCCCTG ACTGAGTGACCACCTCTGCTGCCCCGAGGCCATGTAGGCCGTGCTTAGGCCTCTGT GGACACACTGCTGGGGACGGCGCCTGAGCTCTCAGGGGGACGAGGAACACCACCA TGCCCCGGGGCTTCACCTGGCTGCGCTATCTTGGGATCTTCCTTGGCGTGGCCTTG GGAATGAGCCTTTGGAGATGTGGCCCTTGACGCAGAATGAGGAGTGCACTGTCAC GGGTTTTCTGCGGGACAAGCTGCAGTACAGGAGCCGACTTCAGTACATGAAACACT ACTTCCCCATCAACTACAAGATCAGTGTGCCTTACGAGGGGGTGTTTCAAGATCGCC AACGTCAACAGGCTGAGGGCCCAGGTGAGCGAGCGGGAGCTGCGGTATCTGTGGG TCTTGGTGAGCCTCAGTGCCACTGAGTCGGTGCAGGACGTGCTGCTCGAGGGCCAC CCATCCTGGAAGTACCTGCAGGAGGTGGAGACGCTGCTGCTGAATGTCCAGCAGG GCCTCACGGATGTGGAGGTACAGCCCCAAGGTGGAATCCGTGTTGTCCCTCTTGAAT GCCCCAGGGCCAAACCTGAAGCTGGTGCAGGCCAAAGCCCTGCTGGACAACCTGCTT CCGGGTGCTGAGGCTGCTGTACTGCTCCTGCTGTAAACAAAGCTCCGTCCTAAACT GGCAGGACTGTGAGGTGCCAAGTCTCAGTCTTGACAGCCAGAGCCCTCATTGCAG TATGCGGCCACCCAGCTGTACCCTCCGCCCCCGTGGTCCCCCAGCTCCCCGCCTCAC TCCACGGGCTCGGTGAGGCCGGTCAGGGCACAGGGCGAGGGCCTCTTGCCCTGAG CACCTGGATGGTGACTGCGGATAGGGGCAGCCAGACCAGCTCCCACAGGAGTTC AACTGGGTCTGAGACTTCAAGGGGTGGTGGTGGGAGCCCCCTTGGGAGAGGACC CCTGGGAAGGGTGTTTTCTTTGAGGGGGATTCTGTGCCACAGCAGGGCTCAGCT TCCTGCCTTCCATAGCTGTCTATGGCCTCACCTGGAGCGGAGGGGACCTGGGGACCT GAAGGTGGATGGGGACACAGCTCCTGGCTTCTCCTGGTGCTGCCCTCACTGTCCCC CCGCCTAAAGGGGGTACTGAGCCTCCTGTGGCCCCGACAGTGAAGGGCACAGCTG TGGGTTGCAGGGGAGACAGCCAGCACGGCGTGGCCATTCTATGACCCCCCAGCCTG GCAGACTGGGGAGCTGGGGGCAGAGGGCGGTGCCAAGTGCCACATCTTGCCATAG TGGATGCTCTTCCAGTTTCTTTTTTCTATTAAACACCCCACTTCCTTTGG NM_001393496.1 Homo sapiens interleukin 34 (IL34), transcript variant 7, mRNA
51	GGAAGGCGCTACAGACCCGCTGGCATAGTTAGCTGGGAGGGACACTAAGATCAGA TAGAGCAGGAAGGAAAGTGGTGTGGGGCGTGGGTGACTGAAGAGTCATTTTATCA AGTTGAAACCAGAATCCAGGATGATTGATCTCCTGAATGCCGGGTGGAAAACCC AGCAACGAGCTTTGAAAACATATACCCGGACACCAAGGGGCAGAGGCTGTTCTGG GCGGGAGGTTGTGCCTGCCCCACGGAGCGACAGAAGCGGGGAGACCAGACGTCGA CCCTGAGGCGTGCTCCTGGGGGGCTCCAGTGGCCGGCATGGGATCTTGGGATCTT CCTTGGCGTGGCCTTGGGGAATGAGCCTTTGGAGATGTGGCCCTTGACGCAGAATG AGGAGTGCACTGTACGGGTTTTCTGCGGGACAAGCTGCAGTACAGGAGCCGACTT CAGTACATGAAACACTACTTCCCCATCAACTACAAGATCAGTGTGCCTTACGAGGG GGTGTTCAGAATCGCCAACGTCACCAGGCTGCAGAGGGGCCAGGTGAGCGAGCGG GAGCTGCGGTATCTGTGGGTCTTGGTGAGCCTCAGTGCCACTGAGTCGGTGCAGGA CGTGCTGCTCGAGGGCCACCCATCCTGGAAGTACCTGCAGGAGGTGGAGACGCTG CTGCTGAATGTCCAGCAGGGCCTCACGGATGTGGAGGTACGCCCCAAGGTGGAAT CCGTGTTGTCCCTCTTGAATGCCCCAGGGCCAAACCTGAAGCTGGTGCGGCCAAA GCCCTGCTGGACAACCTGCTTCCGGGTGCTGAGAGTGTGTACTGCTCCTGCTGTAA ACAAAGCTCCGTCTAACTGGCAGGACTGTGAGGTGCCAAGTCTCAGTCTTGCA GCCAGAGCCCTCATTGCAGTATGCGGCCACCCAGCTGTACCCTCCGCCCCCGTGG TCCCCCAGCTCCCCGCCTCACTCCACGGGCTCGGTGAGGCCGGTCAGGGCACAGGG

	CGAGGGCCTCTTGCCCTGAGCACCCCTGGATGGTGACTGCGGATAGGGGCAGCCAG ACCAGCTCCCACAGGAGTTCAACTGGGTCTGAGACTTCAAGGGGTGGTGGTGGGA GCCCCCTTGGGAGAGGACCCCTGGGAAGGGTGTTTTCTTTGAGGGGGATTCTG TGCCACAGCAGGGCTCAGCTTCCTGCCTTCCATAGCTGTCATGGCCTCACCTGGAG CGGAGGGGACCTGGGGACCTGAAGGTGGATGGGGACACAGCTCCTGGCTTCTCCT GGTGCTGCCCTCACTGTCCCCCGCCTAAAGGGGGTACTGAGCCTCCTGTGGCCCG CAGCAGTGAGGGCACAGCTGTGGGTTGCAGGGGAGACAGCCAGCACGGCGTGGCC ATTCTATGACCCCCCAGCCTGGCAGACTGGGGAGCTGGGGGCAGAGGGCGGTGCC AAGTGCCACATCTTGCCATAGTGGATGCTCTTCCAGTTTCTTTTTTCTATTAAACAC CCCACTTCCTTTGG NM_001393497.1 Homo sapiens interleukin 34 (IL34), transcript variant 8, mRNA
52	GGAAGGCGCTACAGACCCGCTGGCAGTAGTTAGCTGGGAGGGACACTAAGATCAGA TAGAGCAGGAAGGAAAGTGGTGTGGGGCGTGGGTGACTGAAGAGTCATTTTATCA AGTTGAAACCAGAAATCCAGGATGATTCGATCTCCTGAATGCCGGGCTGGAAAACCC AGCAACGAGCTTTGAAAACATATCACCCGGACACCAAGGGGCAGAGGCTGTTCTGG GCGGGAGGTTGTGCTTGGCCACGGAGCGACAGAAGCGGGGAGACCAGACGCTCGA CCCTGAGGCGTGCCTCCTGGGGGGCTCCAGTGGCCGGCATGGGATCTTGGGATCTT CCTTGGCGTGGCCTTGGGGAATGAGCCTTTGGAGATGTGGCCCTTGACGCAGAATG AGGAGTGCAGTGTACGGGTTTTCTGCGGGACAAGCTGCAGTACAGGAGCCGACTT CAGTACATGAAACACTACTTCCCCATCAACTACAAGATCAGTGTGCCTTACGAGGG GGTGTTCAGAATCGCCAACGTCACCAGGCTGAGGGCCAGGTGAGCGAGCGGGAG CTGCGGTATCTGTGGGTCTTGGTGAGCCTCAGTGCCACTGAGTCGGTGCAGGACGT GCTGCTCGAGGGCCACCCATCCTGGAAGTACCTGCAGGAGGTGGAGACGCTGCTG CTGAATGTCCAGCAGGGCCTCACGGATGTGGAGGTGAGCCCCAAGGTGGAATCCG TGTTGTCCCTCTTGAATGCCCCAGGGCCAAACCTGAAGCTGGTGCGGCCCCAAAGCC CTGCTGGACAACCTGCTTCCGGGTCATGGAGCTGCTGTACTGCTCCTGCTGTAAACA AAGCTCCGTCTTAACTGGCAGGACTGTGAGGTGCCAAGTCCTCAGTCTTGCAGCC CAGAGCCCTCATTGCAGTATGCGGCCACCCAGCTGTACCCTCCGCCCCCGTGGTCC CCCAGCTCCCCGCCTCACTCCACGGGCTCGGTGAGGCCGGTCAGGGCACAGGGCG AGGGCCTCTTGCCCTGAGCACCCCTGGATGGTGACTGCGGATAGGGGCAGCCAGAC CAGCTCCCACAGGAGTTCAACTGGGTCTGAGACTTCAAGGGGTGGTGGTGGGAGC CCCCCTTGGGAGAGGACCCCTGGGAAGGGTGTTTTCTTTGAGGGGGATTCTGTG CCACAGCAGGGCTCAGCTTCCTGCCTTCCATAGCTGTCATGGCCTCACCTGGAGCG GAGGGGACCTGGGGACCTGAAGGTGGATGGGGACACAGCTCCTGGCTTCTCCTGG TGCTGCCCTCACTGTCCCCCGCCTAAAGGGGGTACTGAGCCTCCTGTGGCCCCGA GCAGTGAGGGCACAGCTGTGGGTTGCAGGGGAGACAGCCAGCACGGCGTGGCCAT TCTATGACCCCCCAGCCTGGCAGACTGGGGAGCTGGGGGCAGAGGGCGGTGCCAA GTGCCACATCTTGCCATAGTGGATGCTCTTCCAGTTTCTTTTTTCTATTAAACACC CACTTCCTTTGG NM_001393498.1 Homo sapiens interleukin 34 (IL34), transcript variant 9, mRNA

CLAIMS

1. A polynucleotide that is complementary to IL-34, or a portion thereof, comprising a nucleotide sequence according to:
 - a. 5'-CTTTGGGCCGCACCAGCTTC-3' (SEQ ID NO: 3), or
 - b. 5'-TCCATGACCCGGAAGCAGTT-3' (SEQ ID NO: 5),
 wherein at least one cytidine of the nucleotide sequence is chemically modified, and at least one nucleoside is a 2'-*O*-(2-methoxyethyl) (2'-MOE) nucleoside;
 or a pharmaceutically acceptable salt thereof.
2. The polynucleotide of claim 1, wherein no more than 10 nucleosides are chemically modified.
3. The polynucleotide of claim 1 or 2, comprising six 2'-MOE nucleosides.
4. The polynucleotide of claim 1 or 2, comprising eight 2'-MOE nucleosides.
5. The polynucleotide of any one of claims 1-4, comprising a nucleotide sequence selected from the group consisting of:
 - a. 5'-CxTxTxTGGGCXGCACCAGCxTxTx-3' (SEQ ID NO: 40), wherein Cx is 2'-*O*-(2-methoxyethyl)cytidine, Tx is 2'-*O*-(2-methoxyethyl)thymidine, and X is 5-methylcytidine;
 - b. 5'-CxTxTxTGGGCXGCACCAGCTxTxTx-3' (SEQ ID NO: 41), wherein Cx is 2'-*O*-(2-methoxyethyl)cytidine, Tx is 2'-*O*-(2-methoxyethyl)thymidine, and X is 5-methylcytidine;
 - c. 5'-TxTxTxATGACCXGGAAGCAxGxTxTx-3' (SEQ ID NO: 42), wherein Cx is 2'-*O*-(2-methoxyethyl)cytidine, Ax is 2'-*O*-(2-methoxyethyl)adenosine, Tx is 2'-*O*-(2-methoxyethyl)thymidine, Gx is 2'-*O*-(2-methoxyethyl)guanosine, and X is 5-methylcytidine; and
 - d. 5'-TxTxTxATGACCXGGAAGCAGxTxTx-3' (SEQ ID NO: 43), wherein Cx is 2'-*O*-(2-methoxyethyl)cytidine, Tx is 2'-*O*-(2-methoxyethyl)thymidine, Gx is 2'-*O*-(2-methoxyethyl)guanosine, and X is 5-methylcytidine;
 or a pharmaceutically acceptable salt thereof.

6. The polynucleotide of any one of claims 1-5, wherein the polynucleotide is an IL-34 antisense oligonucleotide.
7. The polynucleotide of any one of claims 1-5, wherein the polynucleotide is part of an IL-34 siRNA complex.
8. The polynucleotide of any one of claims 1-7, wherein at least one internucleoside linkage of the polynucleotide is selected from the group consisting of a phosphorothioate linkage, a phosphorodithioate linkage, a phosphotriester linkage, an alkylphosphonate linkage, an aminoalkylphosphotriester linkage, an alkylene phosphonate linkage, a phosphinate linkage, a phosphoramidate linkage, and an aminoalkylphosphoramidate linkage, a thiophosphoramidate linkage, thionoalkylphosphonate linkage, a thionoalkylphosphotriester linkage, a thiophosphate linkage, a selenophosphate linkage, and a boranophosphate linkage.
9. The polynucleotide of any one of claims 1-8, wherein all internucleoside linkages of the polynucleotide are phosphorothioate linkages.
10. A pharmaceutical composition, comprising (a) a polynucleotide of any one of claims 1-9, or a pharmaceutically acceptable salt thereof, and (b) a pharmaceutically acceptable excipient.
11. A method of treating an inflammatory disease in a patient in need thereof, the method comprising administering to the patient in need thereof an effective amount of a polynucleotide of any one of claims 1-9, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition of claim 10.
12. The method of claim 11, wherein the inflammatory disease is associated with elevated IL-34 expression.
13. The method of any one of claims 11 to 13, wherein the inflammatory disease is rheumatoid arthritis.
14. The method of any one of claims 11 to 13, wherein the inflammatory disease is osteoarthritis.

15. The method of any one of claims 11 to 14, wherein the method (i) inhibits inflammatory cytokine production in cells of the patient; (ii) reduces or inhibits an IL-34 mediated inflammatory response in cells of the patient; and/or (iii) reduces or inhibits IL-34-mediated macrophage colony-stimulating factor receptor (M-CSFR-1) signaling in cells of the patient.
16. The method of any one of claims 11-15, wherein the cell is an intestinal cell (e.g., intestinal stromal cell) and/or forms part of an intestinal fibrostricture.
17. A method of reducing or eliminating a fibrotic stricture in a patient suffering from an inflammatory disease, the method comprising administering an effective amount of a polynucleotide of any one of claims 1-9, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition of claim 10.
18. The method of claim 17, wherein the fibrotic stricture is located in the intestine.
19. The method of claim 17 or 18, wherein the inflammatory disease is selected from the group consisting of an inflammatory bowel disease, rheumatoid arthritis, psoriasis, osteoarthritis, pouchitis, (type I and II), tissue or organ rejection, multiple sclerosis, periodontal inflammation, periodontitis, pigmented villonodular synovitis, hepatitis, sinusitis, colon cancer, colorectal cancer, colitis-associated colon cancer, sporadic colorectal cancer, coronary artery disease, Sjogren's syndrome (SS), obesity, chronic inflammation, pulmonary sarcoidosis, skin lesions, a CNS inflammatory disease, and an autoimmune disease.
20. The method of claim 19, wherein the inflammatory disease is rheumatoid arthritis.
21. The method of claim 19, wherein the inflammatory disease is osteoarthritis.
22. The method of claim 19, wherein the inflammatory disease is an inflammatory bowel disease (e.g., Crohn's disease, gastroduodenal Crohn's disease, Crohn's (granulomatous) colitis, inflammatory Crohn's disease, fibrostricturing Crohn's disease, ulcerative colitis, collagenous colitis, lymphocytic colitis, ischaemic colitis, diversion colitis, Behçet's disease, microscopic colitis, ulcerative proctitis, proctosigmoiditis, jejunoileitis, left-sided colitis, pancolitis, ileocolitis, ileitis, or indeterminate colitis).

23. The method of claim 22, wherein the inflammatory bowel disease is inflammatory Crohn's disease or fibrostricturing Crohn's disease.
24. A method for preventing or treating fibrosis, the method comprising administering to a patient in need thereof a therapeutically effective amount of a polynucleotide of any one of claims 1-9, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition of claim 10.
25. The method of claim 26, wherein the fibrosis is pulmonary fibrosis, synovial fibrosis, or intestinal fibrosis.
26. The method of claim 26, wherein the fibrosis is selected from the group consisting of renal fibrosis, cardiac fibrosis, endomyocardial fibrosis, myelofibrosis, retroperitoneal fibrosis, and nephrogenic systemic fibrosis.
27. The method of any one of claims 24-26, wherein the patient is suffering from rheumatoid arthritis.
28. The method of any one of claims 24-26, wherein the patient is suffering from osteoarthritis.
29. The method of any one of claims 24-26, wherein the patient is suffering from Crohn's disease.
30. The method of any one of claims 11-29, wherein the polynucleotide is administered intraarticularly, rectally, topically, parenterally, orally, pulmonarily, intratracheally, intranasally, transdermally, or intraduodenally.
31. The method of any one of claims 11-30, wherein the patient is a human.
32. Use of a polynucleotide of any one of claims 1-9, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for treating an inflammatory disease.
33. A polynucleotide of any one of claims 1-9, or a pharmaceutically-acceptable salt thereof, for use as a medicament.

34. A polynucleotide of any one of claims 1-9, or a pharmaceutically-acceptable salt thereof, for use in the treatment of an inflammatory disease.
35. A polynucleotide of any one of claims 1-9, or a pharmaceutically-acceptable salt thereof, for use in the treatment of fibrosis.