


Quinoline as allosteric enhancers of the GABA-B receptors

The present invention relates to compounds of formula I



wherein

- R¹ is hydrogen, C₁-C₇ alkyl, C₁-C₇ haloalkyl, di(C₁-C₇)alkylamino, C₃-C₈ cycloalkyl,
 5 or 5 or 6 membered heterocycloalkyl;
 R² is C₁-C₇ alkyl, aryl, C₁-C₇ alkoxy(C₁-C₇)alkyl, C₁-C₇ haloalkyl or C₃-C₈
 cycloalkyl;
 R³, R⁴ are each independently hydrogen, halo, hydroxy, C₁-C₇ alkoxy, C₁-C₇
 haloalkoxy, di(C₁-C₇)alkylamino, C₁-C₇ alkylsulfonyl, or 5 or 6 membered
 10 heterocycloalkyl;
 R⁵ is hydrogen, halo, C₁-C₇ alkyl, C₁-C₇ alkoxy, C₁-C₇ haloalkoxy, or aryloxy,
 or is -NR⁷R⁸ wherein R⁷ and R⁸ are C₁-C₇ alkyl, or R⁷ and R⁸ may, together with
 the nitrogen atom to which they are attached, form or a 4 to 8 membered
 heterocycloalkyl group which may be substituted by one or more substituent(s)
 15 selected from the group consisting of halo, C₁-C₇ alkyl, C₁-C₇ alkoxy, hydroxy,
 phenyl and di(C₁-C₇)alkylamino;
 R⁶ is hydrogen or together with R⁵ may form a 5 or 6 membered heterocycloalkyl
 group which may be substituted by one or more halogen;
 and pharmaceutically acceptable acid addition salts thereof,
 20 excluding the following compounds:
 1-(6-Chloro-2-methyl-4-phenyl-quinolin-3-yl)-ethanone;
 1-(6-Bromo-4-phenyl-2-piperidin-1-yl-quinolin-3-yl)-ethanone;
 1-[4-(4-Chloro-phenyl)-2-methyl-quinolin-3-yl]-ethanone;
 1-(6-Bromo-2-methyl-4-phenyl-quinolin-3-yl)-ethanone;

1-(2,6-Dimethyl-4-phenyl-quinolin-3-yl)-ethanone; and
1-(2-Methyl-4-phenyl-6-trifluoromethoxy-quinolin-3-yl)-ethanone.

The six compounds excluded from the scope for formula I are known from chemical libraries. Said six compounds were never disclosed in relation with GABA_B receptors.

The compounds of formula I and their salts are distinguished by valuable therapeutic properties. It has been found that the compounds are active on the GABA_B receptor.

γ -Aminobutyric acid (GABA), the most abundant inhibitory neurotransmitter, activates both ionotropic GABA_{A/C} and metabotropic GABA_B receptors (*Hill and Bowery, Nature, 290, 149-152, 1981*). GABA_B receptors that are present in most regions of the mammalian brain on presynaptic terminals and postsynaptic neurons are involved in the fine-tuning of inhibitory synaptic transmission. Presynaptic GABA_B receptors through modulation of high-voltage activated Ca²⁺ channels (P/Q- and N-type) inhibit the release of many neurotransmitters. Postsynaptic GABA_B receptors activates G-protein coupled inwardly rectifying K⁺ (GIRK) channel and regulates adenylyl cyclase (*Billinton et al., Trends Neurosci., 24, 277-282, 2001; Bowery et al., Pharmacol. Rev., 54, 247-264, 2002*). Because the GABA_B receptors are strategically located to modulate the activity of various neurotransmitter systems, GABA_B receptor ligands hence could have potential therapeutics in the treatment of anxiety, depression, epilepsy, schizophrenia and cognitive disorders (*Vacher and Bettler, Curr. Drug Target, CNS Neurol. Disord. 2, 248-259, 2003; Bettler et al., Physiol Rev. 84, 835-867, 2004*).

Native GABA_B receptors are heteromeric structures composed of two types of subunits, GABA_BR1 and GABA_BR2 subunits (*Kaupmann et al., Nature, 386, 239-246, 1997 and Nature, 396, 683-687, 1998*). The structure of GABA_BR1 and R2 show that they belong to a family of G-protein coupled receptors (GPCRs) called family 3. Other members of the family 3 GPCRs include the metabotropic glutamate (mGlu1-8), Calcium-sensing, vomeronasal, pheromone and putative taste receptors (*Pin et al., Pharmacol. Ther. 98, 325-354, 2003*). The family 3 receptors (including GABA_B receptors) are characterized by two distinctly separated topological domains: an exceptionally long extracellular amino-terminal domain (ATD, 500-600 amino acids), which contains a venus flytrap module for the agonist binding (orthosteric site) (*Galvez et al., J. Biol. Chem., 275, 41166-41174, 2000*) and the 7TM helical segments plus intracellular carboxyl-terminal domain that is involved in receptor activation and G-protein coupling. The

mechanism of receptor activation by agonist in GABA_BR1R2 heterodimer is unique among the GPCRs. In the heteromer, only GABA_BR1 subunit binds to GABA, while the GABA_BR2 is responsible for coupling and activation of G-protein (*Havlickova et al., Mol. Pharmacol.* 62, 343-350, 2002; *Kniazeff et al., J. Neurosci.*, 22, 7352-7361, 2002).

- 5 *Schuler et al., Neuron*, 31, 47-58, 2001 have demonstrated that the GABA_BR1 knock-out (KO) mice exhibit spontaneous seizures and hyperalgesia. These KO mice have lost all the biochemical and electrophysiological GABA_B responses. Interestingly, the GABA_BR1 KO mice were more anxious in two anxiety paradigm, namely the light-dark box (decreased time in light) and staircase tests (decreased rears and steps climbed). They
- 10 showed a clear impairment of passive avoidance performance model indicating impaired memory processes. The GABA_BR1 KO also displayed increased hyperlocomotion and hyperactivity in new environment. The GABA_BR1 gene is mapped to chromosome 6p21.3, which is within the HLA class I, a region with linkage for schizophrenia, epilepsy and dyslexia (*Peters et al., Neurogenetics*, 2, 47-54, 1998). *Mondabon et al., Am. J. Med. Genet* 122B/1, 134, 2003 have reported about a weak association of the Ala20Val polymorphism of GABA_BR1 gene with schizophrenia. Moreover, *Gassmann et al., J. Neurosci.* 24, 6086-6097, 2004 has shown that GABA_BR2KO mice suffer from spontaneous seizures, hyperalgesia, hyperlocomotor activity and severe memory impairment, comparable to GABA_BR1KO mice. Therefore, heteromeric GABA_B R1R2 receptors are
- 15 responsible for these phenotypes.
- 20

- Baclofen (Lioresal®; β -chlorophenyl GABA), a selective GABA_B receptor agonist with EC₅₀ = 210 nM at native receptor, is the only ligand, which has been used since 1972 in clinical study for the treatment of spasticity and skeletal muscle rigidity in patients following spinal cord injury, multiple sclerosis, amyotrophic lateral sclerosis, cerebral
- 25 palsy. Most of the preclinical and clinical studies conducted with baclofen and GABA_B receptor agonists were for the treatment of neuropathic pain and craving associated with cocaine and nicotine (*Misgeld et al., Prog. Neurobiol.* 46, 423-462, 1995; *Enna et al., Life Sci*, 62, 1525-1530, 1998; *McCarson and Enna, Neuropharmacology*, 38, 1767-1773, 1999; *Brebner et al., Neuropharmacology*, 38, 1797-1804, 1999; *Paterson et al., Psychopharmacology*, 172, 179-186, 2003). In panic disorder patients, Baclofen was shown to be significantly effective in reducing the number of panic attacks and symptoms of anxiety as assessed with the Hamilton anxiety scale, Zung anxiety scale and Katz-R nervousness subscale (*Breslow et al., Am. J. Psychiatry*, 146, 353-356, 1989). In a study with a small group of veterans with chronic, combat-related posttraumatic stress disorder
- 30 (PTSD), baclofen was found to be an effective and well-tolerated treatment. It resulted in
- 35

significant improvements in the overall symptoms of PTSD, most notably the avoidance, emotional numbing and hyperarousal symptoms and also in reduced accompanying anxiety and depression (*Drake et al., Ann. Pharmacother. 37., 1177-1181, 2003*). In preclinical study, baclofen was able to reverse the reduction in prepulse inhibition (PPI) of the acoustic startle response induced by dizocilpine, but not by apomorphine in rat PPI model of psychosis (*Bortolato et al., Psychopharmacology, 171, 322-330, 2004*). Therefore, GABA_B receptor agonist has a potential in the pharmacological therapy of psychotic disorders. Unfortunately, Baclofen has a number of side-effects including the poor blood-brain-barrier penetration, very short duration of action and narrow therapeutic window (muscle relaxation, sedation and tolerance) that limit its utility.


Urwyler et al., Mol. Pharmacol., 60, 963-971, 2001 have reported on a novel class of GABA_B receptor ligands, called positive allosteric modulators, CGP7930 [2,6-di-tert-butyl-4-(3-hydroxy-2,2-dimethyl-propyl)-phenol] and its aldehyde analogue CGP13501. These ligands have no effect on their own at GABA_B receptors, but in concert with endogenous GABA, they increase both the potency and maximal efficacy of GABA at the GABA_BR1R2 (*Pin et al., Mol. Pharmacol., 60, 881-884, 2001*). Interestingly, recent study with CGP7930 (*Binet et al., J Biol Chem., 279, 29085-29091, 2004*) has shown that this positive modulator activates directly the seven transmembrane domains (7TMD) of GABA_BR2 subunit. *Mombereau et al., Neuropsychopharmacology, 1-13, 2004* have recently reported on the anxiolytic effects of acute and chronic treatment with the GABA_B receptor positive modulator, GS39783 (*N,N*-dicyclopentyl-2-methylsulfanyl-5-nitro-pyrimidine-4,6-diamine) (*Urwyler et al., J. Pharmacol. Exp. Ther., 307, 322-330, 2003*) in the light-dark box and elevated zero maze test models of anxiety. No tolerance after chronic treatment (21 days) with GS39783 (10 mg/kg, P.O., once daily) was observed. Because the GABA_B enhancers have no effect on receptor activity in the absence of GABA, but do enhance allosterically the affinity of the GABA_B receptor for the endogenous GABA, it is expected that these ligands should have an improved side effect profile as compared to baclofen. Indeed, GS39783 at 0.1-200 mg/kg, PO had no effect on spontaneous locomotor activity, rotarod, body temperature and traction test in comparison to baclofen, which showed these side effects at 2.5-15 mg/kg, PO. GS39783 did not have any effect on cognition performance as assessed by passive avoidance behavioral test in mice and rats. Furthermore, GS39783 exhibited anxiolytic-like effects in the elevated plus maze (rat), elevated zero maze (mice and rats), and the stress-induced hyperthermia (mice) test paradigms. Therefore, GS39783 represents a novel anxiolytic without side-effects associated with baclofen or benzodiazepines (*Cryan et al., J Pharmacol Exp Ther., 310, 952-963, 2004*). The preclinical investigation with the CGP7930

and GS39783 has shown that both compounds were effective at decreasing cocaine self-administration in rats (*Smith et al., Psychopharmacology, 173, 105-111, 2004*). The positive modulator, CGP7930 has also been preclinically studied for the treatment of Gastro-Esophageal Reflux Disease (GERD) and was found to be effective (WO 03/090731, Use of GABA_B receptor positive modulators in gastro-intestinal disorders).

Positive allosteric modulators have been reported for other family 3 GPCRs including mGlu1 receptor (*Knoflach et al., Proc. Natl. Acad. Sci., USA, 98, 13402-13407, 2001; Wichmann et al., Farmaco, 57, 989-992, 2002*), Calcium-sensing receptor (NPS R-467 and NPS R-568) (*Hammerland et al., Mol. Pharmacol., 53, 1083-1088, 1998*) (US 6,313, 146), mGlu2 receptor [LY487379, N-(4-(2-methoxyphenoxy)-phenyl-N-(2,2,2-trifluoroethylsulfonyl)-pyrid-3-ylmethylamine and its analogs] (WO 01/56990, Potentiators of glutamate receptors) and mGlu5 receptor (CPPHA, N-{4-chloro-2-[(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)methyl] phenyl}-2-hydroxybenzamide) (*O'Brien et al., J. Pharmacol. Exp. Ther., 27, Jan. 27, 2004*). Interestingly, it has been demonstrated that these positive modulators bind to a novel allosteric site located within the 7TMD region, thereby enhancing the agonist affinity by stabilizing the active state of the 7TMD region (*Knoflach et al., Proc. Natl. Acad. Sci., USA 98, 13402-13407, 2001; Schaffhauser et al., Mol. Pharmacol., 64, 798-810, 2003*). Moreover, the NPS R-467, NPS R-568 (Tecalset) and related compounds represent the first positive allosteric modulators that entered the clinical trials due to their allosteric mode of action.

Objects of the invention are the compounds of formula I and pharmaceutically acceptable acid addition salts thereof, the preparation of the compounds of formula I and salts thereof, medicaments containing a compound of formula I or a pharmaceutically acceptable acid addition salts thereof.

According to a first aspect, there are provided compounds of formula I



wherein

R¹ is hydrogen, C₁-C₇ alkyl, C₁-C₇ haloalkyl, di(C₁-C₇)alkylamino, C₃-C₈ cycloalkyl, or 5 or 6 membered heterocycloalkyl;


- R^2 is C_1 - C_7 alkyl, aryl, C_1 - C_7 alkoxy(C_1 - C_7)alkyl, C_1 - C_7 haloalkyl or C_3 - C_8 cycloalkyl;
 R^3, R^4 are each independently hydrogen, halo, hydroxy, C_1 - C_7 alkoxy, C_1 - C_7 haloalkoxy, di(C_1 - C_7)alkylamino, C_1 - C_7 alkylsulfonyl, or 5 or 6 membered heterocycloalkyl;
 R^5 is hydrogen, halo, C_1 - C_7 alkyl, C_1 - C_7 alkoxy, C_1 - C_7 haloalkoxy, or aryloxy,
or is $-NR^7R^8$ wherein R^7 and R^8 are C_1 - C_7 alkyl, or R^7 and R^8 may, together with the nitrogen atom to which they are attached, form or a 4 to 7 membered heterocycloalkyl group which may be substituted by one or more substituent(s) selected from the group consisting of halo, C_1 - C_7 alkyl, C_1 - C_7 alkoxy, hydroxy, phenyl and di(C_1 - C_7)alkylamino;
 R^6 is hydrogen or together with R^5 may form a 5 or 6 membered heterocycloalkyl group which may be substituted by one or more halogen;

and pharmaceutically acceptable acid addition salts thereof,
excluding the following compounds:


- 1-(6-Chloro-2-methyl-4-phenyl-quinolin-3-yl)-ethanone;
1-(6-Bromo-4-phenyl-2-piperidin-1-yl-quinolin-3-yl)-ethanone;
1-[4-(4-Chloro-phenyl)-2-methyl-quinolin-3-yl]-ethanone;
1-(6-Bromo-2-methyl-4-phenyl-quinolin-3-yl)-ethanone;
1-(2,6-Dimethyl-4-phenyl-quinolin-3-yl)-ethanone;
1-(2-Methyl-4-phenyl-6-trifluoromethoxy-quinolin-3-yl)-ethanone
Phenyl-(4-phenyl-quinolin-3-yl)-methanone;
Phenyl-(4-phenyl-quinolin-3-yl)-(2,4,6-trimethyl-phenyl)-methanone;
(2-Methyl-4-phenyl-quinolin-3-yl)-phenyl-methanone;
1-(2-Methyl-4-phenyl-quinolin-3-yl)-ethanone;
1-(6-Chloro-2-methyl-4-phenyl-quinolin-3-yl)-ethanone;
1-(4-Phenyl-quinolin-3-yl)-ethanone;
1-(6-Chloro-4-phenyl-quinolin-3-yl)-ethanone;
1-(6-Chloro-2-dimethylamino-4-phenyl-quinolin-3-yl)-ethanone;
1-(6-Methoxy-2-methyl-4-phenyl-quinolin-3-yl)-ethanone;
2,2,2-Trifluoro-1-(2-methyl-4-phenyl-quinolin-3-yl)-ethanone;
1-(6-Bromo-2-(4-morpholinyl)-4-phenyl-quinolin-3-yl)-ethanone;
1-(2,6-Dimethyl-4-phenyl-quinolin-3-yl)-ethanone;
1-[4-(4-Chlorophenyl)-2-methyl-quinolin-3-yl]-ethanone;
1-[6-Chloro-4-phenyl-2-(1-piperidinyl)-quinolin-3-yl]-ethanone;
1-[6-Bromo-4-phenyl-2-(1-piperidinyl)-quinolin-3-yl]-ethanone;
(3-Methylphenyl) (4-phenyl-quinolin-3-yl)-methanone;

(4-Chlorophenyl) (7-chloro-4-phenyl-quinolin-3-yl)-methanone;
 (7-Chloro-4-phenyl-quinolin-3-yl) phenyl-methanone;
 (7-Chloro-4-phenyl-quinolin-3-yl) (4-methoxyphenyl)-methanone;
 (7-Chloro-4-phenyl-quinolin-3-yl) (4-ethylphenyl)-methanone;
 (4-Chlorophenyl) (4-phenyl-quinolin-3-yl)-methanone;
 (4-Ethylphenyl) (4-phenyl-quinolin-3-yl)-methanone; and
 (7-Chloro-4-phenyl-quinolin-3-yl) (4-methylphenyl)-methanone.


According to a second aspect, there is provided a process for the preparation of a compound of formula I as defined above:



comprising the step of reacting a compound of formula II



with a compound of formula III.




to give the compound of formula I;

wherein R¹ to R⁶ are as defined for formula I;


and if desired, converting the compound of formula I obtained into a pharmaceutically acceptable acid addition salt.

According to a third aspect, there is provided a process for the preparation of a compound of formula Ia


-5c-



comprising the steps of reacting a compound of formula IV



with a compound of formula V



to give the compound of formula Ia;

wherein R¹ to R⁸ are as defined in claim 1;

and if desired, converting the compound of formula Ia obtained into a pharmaceutically acceptable acid addition salt.

According to a fourth aspect, there is provided a compound of formula I, prepared according to the process of the second or third aspect.

According to a fifth aspect, there is provided a medicament containing a compound of formula I as defined above.

According to a sixth aspect, there is provided the use of a compound of formula I as defined above or of a compound selected from the group consisting of:

1-(6-Chloro-2-methyl-4-phenyl-quinolin-3-yl)-ethanone;

1-(6-Bromo-4-phenyl-2-piperidin-1-yl-quinolin-3-yl)-ethanone;

1-[4-(4-Chloro-phenyl)-2-methyl-quinolin-3-yl]-ethanone;

1-(6-Bromo-2-methyl-4-phenyl-quinolin-3-yl)-ethanone;

1-(2,6-Dimethyl-4-phenyl-quinolin-3-yl)-ethanone; and

1-(2-Methyl-4-phenyl-6-trifluoromethoxy-quinolin-3-yl)-ethanone,

or of their pharmaceutically acceptable salts, for the manufacture of a medicament.

A further object of the invention is the use of the compound of formula I or of a compound selected from the group consisting of:

1-(6-Chloro-2-methyl-4-phenyl-quinolin-3-yl)-ethanone;

1-(6-Bromo-4-phenyl-2-piperidin-1-yl-quinolin-3-yl)-ethanone;

1-[4-(4-Chloro-phenyl)-2-methyl-quinolin-3-yl]-ethanone;

1-(6-Bromo-2-methyl-4-phenyl-quinolin-3-yl)-ethanone;

1-(2,6-Dimethyl-4-phenyl-quinolin-3-yl)-ethanone; and

1-(2-Methyl-4-phenyl-6-trifluoromethoxy-quinolin-3-yl)-ethanone,

and acceptable acid addition salts thereof for the manufacture of such medicaments useful in the control or prevention of illnesses, especially of illnesses and disorders of the kind referred to earlier, such as anxiety, depression, epilepsy, schizophrenia, cognitive

disorders, spasticity and skeletal muscle rigidity, spinal cord injury, multiple sclerosis, amyotrophic lateral sclerosis, cerebral palsy, neuropathic pain and craving associated with cocaine and nicotine, psychosis, panic disorder, posttraumatic stress disorders or gastrointestinal disorders, and, respectively, for the manufacture of corresponding
5 medicaments.

The following definitions of the general terms used in the present description apply irrespective of whether the terms in question appear alone or in combination.

As used herein, the term "Aryl" means a monovalent cyclic aromatic hydrocarbon moiety. Preferred aryls include, but are not limited to, optionally substituted phenyl or
10 naphthyl, as well as those aryl groups specifically illustrated by the examples herein below. Examples of substituents for aryl groups are hydroxy, halo, C₁-C₇ alkyl, C₁-C₇ haloalkyl, C₁-C₇ alkoxy, C₁-C₇ haloalkoxy, C₁-C₇ alkoxyalkyl, C₁-C₇ alkylsulfonyl, di(C₁-C₇)alkylamino or C₃-C₈ cycloalkyl.

"aryloxy" denotes an aryl group wherein the aryl group is as defined above and the
15 aryl group is connected via an oxygen atom. Preferred aryloxy is PhO-.

"C₁-C₇ alkyl" denotes a straight- or branched-carbon chain group containing from 1 to 7 carbon atoms, for example, methyl, ethyl, propyl, isopropyl, isobutyl, sec-butyl, tert-butyl, pentyl, n-hexyl as well as those specifically illustrated by the examples herein below.

20 "C₁-C₇ haloalkyl" denotes a C₁-C₇ alkyl group as defined above which is substituted by one or more halogen. Examples of C₁-C₇ haloalkyl include but are not limited to methyl, ethyl, propyl, isopropyl, isobutyl, sec-butyl, tert-butyl, pentyl or n-hexyl substituted by one or more Cl, F, Br or I atom(s) as well as those groups specifically illustrated by the examples herein below. Preferred C₁-C₇ haloalkyl are difluoro- or
25 trifluoro-methyl or ethyl.

"C₁-C₇ alkoxy" denotes a group wherein the alkyl group is as defined above and the alkyl group is connected via an oxygen atom. Preferred alkoxy are MeO- and Et-O as well as those groups specifically illustrated by the examples herein below.

30 "C₁-C₇ haloalkoxy" denotes a C₁-C₇ alkoxy group as defined above which is substituted by one or more halogen. Examples of C₁-C₇ haloalkoxy include but are not limited to methoxy or ethoxy, substituted by one or more Cl, F, Br or I atom(s) as well as those groups specifically illustrated by the examples herein below. Preferred C₁-C₇ haloalkoxy are difluoro- or trifluoro-methoxy or ethoxy.

"Halogen" denotes chlorine, iodine, fluorine and bromine.

"C₁-C₇ alkoxyalkyl" denotes a C₁-C₇ alkyl group as defined herein above which is substituted by a C₁-C₇ alkoxy group as defined herein above.

"C₁-C₇ alkylsulfonyl" denotes a sulfonyl group which is substituted by a C₁-C₇ alkyl group as defined herein above. Examples of C₁-C₇ alkylsulfonyl include but are not limited to methylsulfonyl and ethylsulfonyl as well as those groups specifically illustrated by the examples herein below.


"di(C₁-C₇)alkylamino" denotes a -NR⁷R⁸ group, wherein R⁷ and R⁸ are C₁-C₇ alkyl groups as defined herein above. Examples of di(C₁-C₇)alkylamino groups include but are not limited to di(methyl)amino, di(ethyl)amino, methylethylamino, as well as those groups specifically illustrated by the examples herein below.

"Hydroxy" denotes a -OH group.

"C₃-C₈ cycloalkyl" denotes a saturated carbon cyclic ring having 3 to 8 carbon atoms as ring members and includes but is not limited to cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, as well as those groups specifically illustrated by the examples herein below.

"4 to 8 membered heterocycloalkyl" denote a saturated mono- or bi-cyclic ring comprising from 1 to 7 carbon atoms as ring members, the other remaining ring member atoms being selected from one or more O, N and S. Preferred 4 to 8 membered heterocycloalkyl groups are 5 or 6 membered heterocycloalkyl groups. Examples of 4 to 8 and 5 or 6 membered heterocycloalkyl groups include but are not limited to optionally substituted azetidiny, piperidiny, piperaziny, homopiperaziny, azepiny, pyrrolidiny, pyrazolidiny, imidazolinyl, imidazolidiny, pyridiny, pyridaziny, pyrimidiny, oxazolidiny, isoxazolidiny, morpholinyl, thiazolidiny, isothiazolidiny, quinuclidiny, quinolinyl, isoquinolinyl, benzimidazolyl, thiadiazolylidiny, benzothiazolidiny, benzoazolylidiny, dihydrofuryl, tetrahydrofuryl, dihydropyrany, tetrahydropyrany, thiomorpholinyl, thiomorpholinylsulfoxide, thiomorpholinylsulfone, dihydroquinolinyl, dihydrisoquinolinyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl, 1-oxo-thiomorpholin, 1,1-dioxo-thiomorpholin, 1,4-diazepane, 1,4-oxazepane and 8-oxa-3-aza-bicyclo[3.2.1]oct-3-yl, as well as those groups specifically illustrated by the examples herein below.

"R⁶ together with R⁵ form a 5 or 6 membered heterocycloalkyl group" denotes 5 or 6 membered heterocycloalkyl groups as defined above which is fused to the quinoline group via R⁵ and R⁶. An example of such group is but is not limited to the following group:



5

The term "pharmaceutically acceptable acid addition salts" embraces salts with inorganic and organic acids, which include but are not limited to hydrochloric acid, nitric acid, sulfuric acid, phosphoric acid, citric acid, formic acid, fumaric acid, maleic acid, acetic acid, succinic acid, tartaric acid, methane-sulfonic acid, p-toluenesulfonic acid.

- 10 Preferred groups for R¹ may be selected from the group consisting of methyl, ethyl, propyl, i-propyl, butyl, i-butyl, and t-butyl.

Preferred groups for R² may be selected from the group consisting of methyl, ethyl, propyl, i-propyl, butyl, i-butyl, t-butyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl, CHF₂ and CF₃.

- 15 Preferred groups for R³ may be selected from the group consisting of hydrogen, Cl and F.

Preferred groups for R⁴ may be selected from the group consisting of hydrogen, methoxy, methylsulfonyl, Cl and F.

- 20 Preferred groups for R⁵ may be selected from the group consisting of Br, methyl, ethyl, propyl, i-propyl, butyl, i-butyl, and t-butyl, CF₃O, PhO, methoxy, methylsulfonyl, Cl, F or I, and when R⁵ is -NR⁷R⁸, R⁷ and R⁸ together with the nitrogen atom to which they are attached may form a group selected from the group consisting of piperidin-1-yl, morpholin-4-yl, pyrrolidin-1-yl, piperazin-1-yl, pyrrolidin-1-yl, azetidine-1-yl, and azepan-1-yl, which may be substituted by one or more F, methyl, ethyl, propyl, i-propyl, butyl, i-butyl, t-butyl, hydroxy, methoxy, phenyl, dimethylamino and 1,4-oxazepane and
25 8-oxa-3-aza-bicyclo[3.2.1]oct-3-yl.

More preferred groups for R⁵ may be selected from the group consisting of Br, I, methyl, ethyl, propyl, i-propyl, butyl, i-butyl, and t-butyl, CF₃O, PhO, methoxy, methylsulfonyl, Cl or F, and when R⁵ is -NR⁷R⁸, R⁷ and R⁸ together with the nitrogen

atom to which they are attached may form a group selected from the group consisting of piperidin-1-yl, 3,3-difluoro-piperidin-1-yl, 4-hydroxy-4-methyl-piperidin-1-yl, 4-methoxy-piperidine-1-yl, morpholin-4-yl, pyrrolidin-1-yl, 2-methyl-pyrrolidin-1-yl, 4-methyl-piperazin-1-yl, 3-hydroxy-pyrrolidin-1-yl, 3-hydroxy-azetidine-1-yl, 4-hydroxy-4-phenyl-piperidin-1-yl, 3,3-dimethylamine-pyrrolidin-1-yl, azepan-1-yl and 1,4-oxazepane and 8-oxa-3-aza-bicyclo[3.2.1]oct-3-yl.

Preferred compounds of the invention are those compounds of formula I, wherein

- R¹ is C₁-C₇ alkyl;
- R² is C₁-C₇ alkyl, phenyl, C₁-C₇ haloalkyl or C₃-C₈ cycloalkyl;
- 10 R³, R⁴ are each independently hydrogen, halo, C₁-C₇ alkoxy, C₁-C₇ alkylsulfonyl;
- R⁵ is halo, C₁-C₇ haloalkoxy, aryloxy,
or is -NR⁷R⁸ wherein R⁷ and R⁸ are C₁-C₇ alkyl, or R⁷ and R⁸ may, together with the nitrogen atom to which they are attached, form a 4 to 8 membered heterocycloalkyl group which may be substituted by one or more substituent(s)
- 15 selected from the group consisting of halo, C₁-C₇ alkyl, hydroxy, C₁-C₇ alkoxy, phenyl and di(C₁-C₇)alkylamino;
- R⁶ is hydrogen or together with R⁵ may form a 5 or 6 membered heterocycloalkyl group which may be substituted by one or more halogen.

Also preferred compounds of the invention are those compounds of formula I
20 wherein R² is C₁-C₇ alkyl, for example the following compounds:

- 1-(6-Bromo-2-ethyl-4-phenyl-quinolin-3-yl)-propan-1-one;
- 1-(6-Bromo-2-isobutyl-4-phenyl-quinolin-3-yl)-ethanone;
- 1-(6-Bromo-2-methyl-4-phenyl-quinolin-3-yl)-3-methyl-butan-1-one;
- 1-[4-(4-Chloro-phenyl)-2-methyl-6-trifluoromethoxy-quinolin-3-yl]-ethanone;
- 25 1-(6-Bromo-2-isopropyl-4-phenyl-quinolin-3-yl)-2-methyl-propan-1-one;
- 1-[4-(3,4-Dichloro-phenyl)-2-methyl-6-trifluoromethoxy-quinolin-3-yl]-ethanone;
- 1-[4-(4-Chloro-phenyl)-2-methyl-6-phenoxy-quinolin-3-yl]-ethanone; and
- 1-[4-(3-Chloro-phenyl)-2-methyl-6-trifluoromethoxy-quinolin-3-yl]-ethanone.

Other preferred are those compounds of formula I wherein, wherein R² is C₁-C₇
30 haloalkyl, for example the following compounds:

- 1-(6-Bromo-2-methyl-4-phenyl-quinolin-3-yl)-2,2,2-trifluoro-ethanone;
- 1-(6-Bromo-2-methyl-4-phenyl-quinolin-3-yl)-2,2-difluoro-ethanone;
- 2,2,2-Trifluoro-1-(2-methyl-4-phenyl-6-trifluoromethoxy-quinolin-3-yl)-ethanone;

- 1-[4-(3-Chloro-phenyl)-2-methyl-6-trifluoromethoxy-quinolin-3-yl]-2,2,2-trifluoro-ethanone;
- 1-[4-(4-Chloro-phenyl)-2-methyl-6-trifluoromethoxy-quinolin-3-yl]-2,2,2-trifluoro-ethanone;
- 5 2,2,2-Trifluoro-1-[4-(4-methoxy-phenyl)-2-methyl-6-trifluoromethoxy-quinolin-3-yl]-ethanone;
- 2,2,2-Trifluoro-1-[4-(4-fluoro-phenyl)-2-methyl-6-trifluoromethoxy-quinolin-3-yl]-ethanone;
- 1-(6-tert-Butyl-2-methyl-4-phenyl-quinolin-3-yl)-2,2,2-trifluoro-ethanone;
- 10 1-(2,2-Difluoro-6-methyl-8-phenyl-[1,3]dioxolo[4,5-g]quinolin-7-yl)-2,2,2-trifluoro-ethanone;
- 1-[4-(3,4-Difluoro-phenyl)-2-methyl-6-trifluoromethoxy-quinolin-3-yl]-2,2,2-trifluoro-ethanone;
- 2,2,2-Trifluoro-1-[4-(4-methanesulfonyl-phenyl)-2-methyl-6-trifluoromethoxy-quinolin-3-yl]-ethanone;
- 15 2,2,2-Trifluoro-1-[4-(3-fluoro-4-methoxy-phenyl)-2-methyl-6-trifluoromethoxy-quinolin-3-yl]-ethanone;
- 2,2,2-Trifluoro-1-(2-methyl-4-phenyl-6-piperidin-1-yl-quinolin-3-yl)-ethanone;
- 2,2,2-Trifluoro-1-(2-methyl-6-morpholin-4-yl-4-phenyl-quinolin-3-yl)-ethanone;
- 20 2,2,2-Trifluoro-1-(2-methyl-4-phenyl-6-pyrrolidin-1-yl-quinolin-3-yl)-ethanone;
- 2,2,2-Trifluoro-1-[2-methyl-6-(2-methyl-pyrrolidin-1-yl)-4-phenyl-quinolin-3-yl]-ethanone;
- 2,2,2-Trifluoro-1-[2-methyl-6-(4-methyl-piperazin-1-yl)-4-phenyl-quinolin-3-yl]-ethanone;
- 25 2,2,2-Trifluoro-1-[6-(3-hydroxy-pyrrolidin-1-yl)-2-methyl-4-phenyl-quinolin-3-yl]-ethanone;
- 2,2,2-Trifluoro-1-[6-(3-hydroxy-azetidin-1-yl)-2-methyl-4-phenyl-quinolin-3-yl]-ethanone;
- 1-[6-Bromo-4-(4-methanesulfonyl-phenyl)-2-methyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone;
- 30 1-[6-Bromo-4-(4-fluoro-phenyl)-2-methyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone;
- 2,2,2-Trifluoro-1-[4-(4-methanesulfonyl-phenyl)-2-methyl-6-piperidin-1-yl-quinolin-3-yl]-ethanone;
- 2,2,2-Trifluoro-1-[4-(4-fluoro-phenyl)-2-methyl-6-piperidin-1-yl-quinolin-3-yl]-ethanone;
- 35 2,2,2-Trifluoro-1-[4-(4-fluoro-phenyl)-2-methyl-6-morpholin-4-yl-quinolin-3-yl]-ethanone;

- 2,2,2-Trifluoro-1-[4-(4-methanesulfonyl-phenyl)-2-methyl-6-pyrrolidin-1-yl-quinolin-3-yl]-ethanone;
- 2,2,2-Trifluoro-1-[4-(4-fluoro-phenyl)-2-methyl-6-pyrrolidin-1-yl-quinolin-3-yl]-ethanone;
- 5 2,2,2-Trifluoro-1-[4-(4-fluoro-phenyl)-6-(3-hydroxy-pyrrolidin-1-yl)-2-methyl-quinolin-3-yl]-ethanone;
- 2,2,2-Trifluoro-1-[6-(4-hydroxy-4-phenyl-piperidin-1-yl)-2-methyl-4-phenyl-quinolin-3-yl]-ethanone;
- 2,2,2-Trifluoro-1-[6-(4-hydroxy-4-phenyl-piperidin-1-yl)-4-(4-methanesulfonyl-phenyl)-2-methyl-quinolin-3-yl]-ethanone;
- 10 2,2,2-Trifluoro-1-[4-(4-fluoro-phenyl)-6-(4-hydroxy-4-phenyl-piperidin-1-yl)-2-methyl-quinolin-3-yl]-ethanone;
- 2,2,2-Trifluoro-1-[4-(4-fluoro-phenyl)-6-(3-hydroxy-azetidin-1-yl)-2-methyl-quinolin-3-yl]-ethanone;
- 15 1-[6-Azepan-1-yl-4-(4-fluoro-phenyl)-2-methyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone;
- 1-(6-Azepan-1-yl-2-methyl-4-phenyl-quinolin-3-yl)-2,2,2-trifluoro-ethanone;
- 1-[6-(3-Dimethylamino-pyrrolidin-1-yl)-2-methyl-4-phenyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone;
- 1-[6-(3-Dimethylamino-pyrrolidin-1-yl)-4-(4-methanesulfonyl-phenyl)-2-methyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone;
- 20 1-[6-(3-Dimethylamino-pyrrolidin-1-yl)-4-(4-fluoro-phenyl)-2-methyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone;
- 2,2,2-Trifluoro-1-[6-iodo-4-(4-methanesulfonyl-phenyl)-2-methyl-quinolin-3-yl]-ethanone;
- 25 2,2,2-Trifluoro-1-[6-(4-hydroxy-4-methyl-piperidin-1-yl)-4-(4-methanesulfonyl-phenyl)-2-methyl-quinolin-3-yl]-ethanone;
- 2,2,2-Trifluoro-1-[4-(4-methanesulfonyl-phenyl)-6-(4-methoxy-piperidin-1-yl)-2-methyl-quinolin-3-yl]-ethanone;
- 1-[6-(3,3-Difluoro-piperidin-1-yl)-4-(4-methanesulfonyl-phenyl)-2-methyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone; and
- 30 2,2,2-Trifluoro-1-[4-(4-methanesulfonyl-phenyl)-2-methyl-6-(8-oxa-3-aza-bicyclo[3.2.1]oct-3-yl)-quinolin-3-yl]-ethanone.

Still other preferred compounds of the invention are those compounds of formula I, wherein R² is C₃-C₈ cycloalkyl, for example the following compounds:

[6-Bromo-4-(4-methanesulfonyl-phenyl)-2-methyl-quinolin-3-yl]-cyclopropyl-methanone;

Cyclopropyl-[4-(4-methanesulfonyl-phenyl)-2-methyl-6-morpholin-4-yl-quinolin-3-yl]-methanone;

5 Cyclopropyl-[4-(4-methanesulfonyl-phenyl)-2-methyl-6-piperidin-1-yl-quinolin-3-yl]-methanone;

[(6-Bromo-2-methyl-4-phenyl-quinolin-3-yl)-cyclopropyl-methanone;

[6-Bromo-4-(4-fluoro-phenyl)-2-methyl-quinolin-3-yl]-cyclopropyl-methanone;

Cyclopropyl-(2-methyl-4-phenyl-6-piperidin-1-yl-quinolin-3-yl)-methanone; and


10 Cyclopropyl-(2-methyl-6-morpholin-4-yl-4-phenyl-quinolin-3-yl)-methanone.

Still other preferred compounds of the invention are those compounds of formula I, wherein R² is phenyl, for example the following compounds:


(6-Bromo-2-methyl-4-phenyl-quinolin-3-yl)-phenyl-methanone; and

[4-(3-Chloro-phenyl)-2-methyl-6-trifluoromethoxy-quinolin-3-yl]-phenyl-methanone.

15 The afore-mentioned compounds of formula I can be manufactured by the following process of the invention comprising the step of reacting a compound of formula II



with a compound of formula III



20


wherein R¹ to R⁶ are as defined in formula I.

to give the compound of formula I;


and if desired, converting the compound of formula I obtained into a pharmaceutically acceptable acid addition salt.

25


The afore-mentioned compounds of formula I can also be manufactured in accordance with the invention by the following variant process comprising the step of reacting a compound of formula IV



with a compound of formula V



to give the compound of formula Ia;



5

wherein R^1 to R^8 are as defined in formula I;

and if desired, converting the compound of formula Ia obtained into a pharmaceutically acceptable acid addition salt. It is understood that the compounds of formula Ia correspond to the compounds of formula I wherein R^5 is $-NR^7R^8$ and R^7 and R^8 are as defined for formula I.

10


The invention also encompasses a compound of formula I or Ia, whenever it is prepared according to the above-mentioned processes.

In the following the preparation of compounds of formula I is described in more detail:

15 In schemes 1 and 2 are described processes for preparation of compound of formula I or Ia.


The preparation of compounds of formula I are further described in detail in working examples 1 - 46.

Scheme 1

A) cat. (NaAuCl₄·2H₂O)Method A

- According to the procedure developed by A. Arcadi, M. Chiarini, S. Di Giuseppe, and F. Marinelli, Synlett 203-206 (2003), the 2-aminobenzophenone II is reacted with the 1,3-dione III and sodium tetrachloroaurate(III) dihydrate as catalysator. The residue can be purified by conventional methods.

Scheme 2


B) cat. Pd₂dba₃, CHCl₃, rac-BINAP, Cs₂CO₃Method B

- Following a methodology developed by J. P. Wolfe and S. L. Buchwald (J. Org. Chem. 2000, 65, 1144-1157) tris(dibenzylideneacetone)dipalladium chloroform complex is added to rac-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl, cesium carbonate, the 2-amino-4-bromo-benzoquinone IV, and the amine V. The residue can be purified by conventional methods.

- One part of the starting material used in the general procedures of schemes 1 and 2 is commercially available (e.g. some of the benzophenones of formula IV, all 1,3-

diketones of formula III, and all the amines of formula V). However the non commercially available part of said starting material can be prepared according to the general procedure of method C for compounds of formula II as outlined hereafter in scheme 3, or according to the general procedure of method A for providing suitable
 5 compounds of formula IV as outline hereinabove in scheme 1. Unless otherwise specified, the intermediate compounds described therein are novel compounds:

Scheme 3



Method C

Following a procedure developed by T. Sugasawa, T. Toyoda, M. Adachi, and K. Sasakura, J. Am. Chem. Soc. 100, 4842-4852 (1978) and improved by A. W. Douglas, N. L. Abramson, I. N. Houpis, S. Karady, A. Molina, L. C. Xavier, N. Yasuda, Tetrahedron Lett. 35, 6807-6810 (1994), either gallium (III) chloride or aluminium (III) chloride are mixed with a chlorinated solvent. Aniline VII, boron trichloride and benzonitrile VIII are then added to the cold mixture. The crude product can be purified by conventional
 15 methods.

The preparation of compounds of formula II is further described in detail in working examples A1 to A16.

As mentioned earlier, the compounds of formula I and their pharmaceutically acceptable addition salts possess valuable pharmacological properties. It has been found
 20 that the compounds of the present invention have an affinity to the GABA_B receptor.

The compounds were investigated in accordance with the tests given hereinafter.

Intracellular Ca^{2+} mobilization assay

The Chinese Hamster Ovary (CHO) cells stably expressing human $\text{GABA}_B\text{R1aR2a}$ and $\text{G}\alpha 16$ were seeded at 5×10^4 cells/well in the poly-D-lysine treated, 96-well,

- black/clear-bottomed plates (BD Biosciences, Palo Alto, CA). 24 h later, the cells were loaded for 90 min at 37°C with 4 μ M Flou-4 acetoxymethyl ester (Catalog No. F-14202, Molecular Probes, Eugene, OR) in loading buffer (1xHBSS, 20 mM HEPES, 2.5 mM Probenecid). Hanks' Balanced Salt Solution (HBSS) (10X) (catalog No. 14065-049) and
- 5 HEPES (1M) (catalog No. 15630-056) were purchased from Invitrogen, Carlsbad, CA. Probenecid (250 mM) (catalog No. P8761) was from Sigma, Buchs, Switzerland. The cells were washed five times with loading buffer to remove excess dye and intracellular calcium mobilization, $[Ca^{2+}]_i$ were measured using a Fluorometric Imaging Plate Reader (FLIPR, Molecular Devices, Menlo Park, CA) as described previously (Porter et al., Br. J.
- 10 Pharmacol., 128, 13-20, 1999). The enhancers were applied 15 min before the application of the GABA. For GABA shift assay, concentration-response curves of GABA (0.0003-30 μ M) were determined in the absence and presence of 10 μ M enhancer. The GABA-shift is defined as $\text{Log} [EC_{50}(\text{GABA} + 10 \mu\text{M enhancer}) / EC_{50}(\text{GABA alone})]$. The % maximum
- 15 enhancing effect (% E_{max}) and potency (EC_{50} value) of each enhancer was determined from concentration-response curve of the enhancer (0.001-30 μ M) in the presence of 10 nM GABA (EC_{10}). Responses were measured as peak increase in fluorescence minus basal, normalized to the maximal stimulatory effect induced by 10 μ M GABA alone (considered 100%) and 10 nM GABA alone (considered 0%). The data were fitted with
- 20 the equation $Y = 100 + (Max - 100) / (1 + (EC_{50} / [drug])^n)$ where Max is the maximum effect, EC_{50} the concentration eliciting a half-maximum effect and n the Hill slope.

Example	Intracellular Ca^{2+} mobilization Assay in CHO-GABA _B R1aR2a-G α 16 cell		GABA shift
	E_{max} (%) at 10 nM GABA alone = 0% 10 μ M GABA alone = 100%	EC_{50} (μ M) at 10nM GABA	$\text{Log} [EC_{50}(\text{GABA} + 10 \mu\text{M cp}) / EC_{50}(\text{GABA alone})]$
3	124	0.80	-1.20
11	65	0.80	-1.00
15	67	1.60	-1.15
19	63	0.90	-0.70
33	62	0.33	-0.90

56	58	2.20	-0.90
----	----	------	-------

The compounds of formula I as well as their pharmaceutically usable acid addition salts can be used as medicaments, e.g. in the form of pharmaceutical preparations. The pharmaceutical preparations can be administered orally, e.g. in the form of tablets, coated tablets, dragées, hard and soft gelatine capsules, solutions, emulsions or suspensions. The administration can, however, also be effected rectally, e.g. in the form of suppositories, or parenterally, e.g. in the form of injection solutions.

The compounds of formula I and their pharmaceutically usable acid addition salts can be processed with pharmaceutically inert, inorganic or organic excipients for the production of tablets, coated tablets, dragees and hard gelatine capsules. Lactose, corn starch or derivatives thereof, talc, stearic acid or its salts etc can be used as such excipients e.g. for tablets, dragées and hard gelatine capsules.

Suitable excipients for soft gelatine capsules are e.g. vegetable oils, waxes, fats, semi-solid and liquid polyols.

Suitable excipients for the manufacture of solutions and syrups include but are not limited to water, polyols, saccharose, invert sugar, glucose.

Suitable excipients for injection solutions include but are not limited to water, alcohols, polyols, glycerol, vegetable oils.

Suitable excipients for suppositories include but are not limited to natural or hardened oils, waxes, fats, semi-liquid or liquid polyols.

Moreover, the pharmaceutical preparations can contain preservatives, solubilizers, stabilizers, wetting agents, emulsifiers, sweeteners, colorants, flavorants, salts for varying the osmotic pressure, buffers, masking agents or antioxidants. They can also contain still other therapeutically valuable substances.

The dosage can vary within wide limits and will, of course, be fitted to the individual requirements in each particular case. In general, in the case of oral administration a daily dosage of about 10 to 1000 mg per person of a compound of general formula I should be appropriate, although the above upper limit can also be exceeded when necessary.

Tablet Formulation (Wet Granulation)

<u>Item</u>	<u>Ingredients</u>	<u>mg/tablet</u>			
		5 mg	25 mg	100 mg	500 mg
	1. Compound of formula I	5	25	100	500
5	2. Lactose Anhydrous DTG	125	105	30	150
	3. Sta-Rx 1500	6	6	6	30
	4. Microcrystalline Cellulose	30	30	30	150
	5. Magnesium Stearate	1	1	1	1
	Total	167	167	167	831

10 Manufacturing Procedure

1. Mix items 1, 2, 3 and 4 and granulate with purified water.
2. Dry the granules at 50°C.
3. Pass the granules through suitable milling equipment.
4. Add item 5 and mix for three minutes; compress on a suitable press.

15

Capsule Formulation

<u>Item</u>	<u>Ingredients</u>	<u>mg/capsule</u>			
		5 mg	25 mg	100 mg	500 mg
	1. Compound of formula I	5	25	100	500
	2. Hydrous Lactose	159	123	148	---
20	3. Corn Starch	25	35	40	70
	4. Talc	10	15	10	25
	5. Magnesium Stearate	1	2	2	5
	Total	200	200	300	600

Manufacturing Procedure

- 25 1. Mix items 1, 2 and 3 in a suitable mixer for 30 minutes.
2. Add items 4 and 5 and mix for 3 minutes.
3. Fill into a suitable capsule.

EXAMPLESSynthesis of intermediates of formula II

Example A1

(2-Amino-5-tert-butyl-phenyl)-phenyl-methanone

5 The title compound was prepared following the general procedure of method C. In a glass flask fitted with magnetic stir bar, rubber septum, thermometer, Hickmann-condenser, nitrogen-purged bubbler connected to a washing bottle containing 30 % NaOH, the content of a fresh ampoule of gallium (III) chloride (5 g, 29 mmol) was added at once and then dissolved by the addition of 1,2-dichloroethane (80 mL). This solution
10 was cooled in ice, then 4-tert-butylaniline (36 mmol) was added slowly while keeping the temperature below 5 °C. Then the solution was cooled to -10 °C and a fresh 1 M solution of boron trichloride in dichloromethane (27 mL) was added via a syringe fitted with a teflon stop-cock while keeping the temperature below -5 °C. Finally benzonitrile was added (24 mmol) and the mixture was allowed to warm to 20 °C. The Hickmann-
15 condenser was replaced by a normal reflux condenser and the reaction mixture was heated in an oil-bath (90 °C) over 1-2 h in order to distil off all the dichloromethane (a total of ca. 50 mL of distillate was collected) until the reflux temperature of 80 °C was achieved. Refluxing was continued for 14 h. The reaction mixture was cooled in ice and hydrolyzed slowly with water (40 mL) and then heated at 60-80 °C for 20-30 min. in
20 order to hydrolyze the imine. The reaction mixture was cooled again and then extracted with dichloromethane and water. The crude product was purified by chromatography on silica gel in heptane/ethylacetate (4:1) and the purified product (yield 40%) analyzed by MS: $m/z = 254$ (M+H).

Example A2

25 (2-Amino-5-bromo-phenyl)-(4-fluoro-phenyl)-methanone

The title compound was prepared by reacting 4-bromoaniline and 4-methylsulfonyl benzonitrile following to the procedure of example A1. Yield 37%; MS: $m/z = 294$ (M).

Example A3

(2-Amino-5-trifluoromethoxy-phenyl)-(4-chloro-phenyl)-methanone

30 The title compound was prepared by reacting 4-(trifluoromethoxy)aniline and 4-chloro benzonitrile following to the procedure of example A1, except that Aluminium

(III) chloride was used, that the reaction time was 4h and heptane/ethylacetate (2:1) was used for the chromatography. Yield 18%; MS: $m/z = 315$ (M).

Example A4

(2-Amino-5-trifluoromethoxy-phenyl)-(3,4-dichloro-phenyl)-methanone

5 The title compound was prepared by reacting 4-(trifluoromethoxy)aniline and 3,4-dichloro benzonitrile following to the procedure of example A3, except that the reaction time was 16h. Yield 15%; MS: $m/z = 408$ (M+OAc).

Example A5

(2-Amino-5-phenoxy-phenyl)-(4-chloro-phenyl)-methanone

10 The title compound was prepared by reacting 4-phenoxyaniline and 4-chloro benzonitrile following to the procedure of example A3, except that the reaction time was 14h and a gradient of heptane/ethylacetate was used for the chromatography. Yield 38%; MS: $m/z = 324$ (M+H).

Example A6

15 (2-Amino-5-trifluoromethoxy-phenyl)-phenyl-methanone

The title compound is known from FR 7666 and was prepared by reacting 4-(trifluoromethoxy)aniline and benzonitrile following to the procedure of example A1, except that the reaction time was 16h and heptane/ethylacetate (5:1) was used for the chromatography. Yield 40%; MS: $m/z = 282$ (M+H).

20

Example A7

(2-Amino-5-trifluoromethoxy-phenyl)-(3-chloro-phenyl)-methanone

The title compound was prepared by reacting 4-(trifluoromethoxy)aniline and 3-chloro-benzonitrile following to the procedure of example A1. Yield 19%; MS: $m/z = 315$ (M).

25

Example A8

(2-Amino-5-trifluoromethoxy-phenyl)-(4-methoxy-phenyl)-methanone

The title compound was prepared by reacting 4-phenoxyaniline and 4-methoxy benzonitrile following to the procedure of example A1, except that the reaction time was

19h and a gradient of heptane/ethylacetate was used for the chromatography. Yield 19%; MS: $m/z = 312$ (M+H).

Example A9

(2-Amino-5-trifluoromethoxy-phenyl)-(4-fluoro-phenyl)-methanone

- 5 The title compound was prepared by reacting 4-(trifluoromethoxy)aniline and 4-fluorobenzonitrile following to the procedure of example A1. Yield 26%; MS: $m/z = 299$ (M).

Example A10

(6-Amino-2,2-difluoro-benzo[1,3]dioxol-5-yl)-phenyl-methanone

- 10 The title compound was prepared by reacting 2,2-difluoro-5-aminobenzodioxole and benzonitrile following to the procedure of example A1. Yield 1.5%; MS: $m/z = 366$ (M+OAc).

Example A11

(2-Amino-5-trifluoromethoxy-phenyl)-(3-trifluoromethoxy-phenyl)-methanone

- 15 The title compound was prepared by reacting 4-(trifluoromethoxy)aniline and 3-(trifluoromethoxy)benzonitrile following to the procedure of example A1. Yield 29%; MS: $m/z = 365$ (M).

Example A12

(2-Amino-5-trifluoromethoxy-phenyl)-(3,4-difluoro-phenyl)-methanone

- 20 The title compound was prepared by reacting 4-(trifluoromethoxy)aniline and 3,4-difluorobenzonitrile following to the procedure of example A1. Yield 36%; MS: $m/z = 317$ (M).

Example A13

(2-Amino-5-trifluoromethoxy-phenyl)-(4-methanesulfonyl-phenyl)-methanone

- 25 The title compound was prepared by reacting 4-(trifluoromethoxy)aniline and 4-methylsulfonylbenzonitrile following to the procedure of example A1. Yield 71%; MS: $m/z = 359$ (M).

Example A14

(2-Amino-5-trifluoromethoxy-phenyl)-(3-fluoro-4-methoxy-phenyl)-methanone

The title compound was prepared by reacting 4-(trifluoromethoxy)aniline and 3-methoxybenzonitrile following to the procedure of example A1. Yield 19%; MS: m/z =
5 329 (M).

Example A15

(2-Amino-5-bromo-phenyl)-(4-methanesulfonyl-phenyl)-methanone

The title compound was prepared by reacting 4-bromoaniline and 4-methylsulfonylbenzonitrile following to the procedure of example A1. Yield 51%; MS:
10 m/z = 355 (M+H).

Example A16

(2-Amino-5-bromo-phenyl)-phenyl-methanone

The title compound is known from US 20040127536 A1 and was prepared according to a method developed by D. Roche, K. Prasad, O. Repic, T. J. Blacklock,
15 *Tetrahedron Lett.* 41, 2083-2085 (2000). 2-Aminobenzophenone (30g, 152mmol) was suspended in acetic acid (300mL). Potassium bromide (19.9g, 167mmol), sodium perborate tetrahydrate (28g, 183mmol) and ammonium molybdate tetrahydrate (1.5g) were added and stirring continued for 3 hours at 0°C. The dense yellow precipitate which formed was diluted with ice water (300mL) and then filtered off and washed with ice
20 water and dried. One obtained 40.3 g (96%) of a yellow solid. MS: m/z = 276 (M).

Example A17

(2-Amino-5-iodo-phenyl)-(4-methanesulfonyl-phenyl)-methanone

The title compound was prepared by reacting 4-iodoaniline and 4-methylsulfonylbenzonitrile following to the procedure of example A1. Yield 31 %; MS:
25 m/z = 402 (M + H).

Synthesis of compounds of formula I according to the invention

Example I

1-(6-Bromo-2-ethyl-4-phenyl-quinolin-3-yl)-propan-1-one

The title compound was prepared according to the general procedure of method A.
30 The (2-Amino-5-bromo-phenyl)-phenyl-methanone [example A16] (on 0.1 – 1 g scale)

and 3,5-heptanedione (1.5 equiv) and sodium tetrachloroaureate(III) dihydrate (0.025 equiv) were heated in parallel in a Radley carousel under nitrogen in ethanol (10 % w/w-solution of (2-Amino-5-bromo-phenyl)-phenyl-methanone) and reacted for 24h. The reaction mixture was evaporated to dryness and the residue purified by chromatography
5 on silica gel in heptane,/ethyl acetate (20:1). Yield: 37 %. MS: $m/z = 368$ (M).

Example 2

(6-Bromo-2-methyl-4-phenyl-quinolin-3-yl)-phenyl-methanone

The title compound was prepared from (2-Amino-5-bromo-phenyl)-phenyl-methanone [example A16] and 1-phenyl-1,3-butanedione, except that the residue was
10 purified by spontaneous crystallization from the reaction mixture. Yield: 61 %; MS: $m/z = 402$ (M+H).

Example 3

1-(6-Bromo-2-methyl-4-phenyl-quinolin-3-yl)-2,2,2-trifluoro-ethanone

The title compound was prepared from (2-Amino-5-bromo-phenyl)-phenyl-
15 methanone [example A16] and 1,1,1-trifluoro-2,4-pentanedione according to the procedure of example 1, except that heptane/ethyl acetate (10:1) was used. Yield: 50 %; MS: $m/z = 392/394$ (M).

Example 4

1-(6-Bromo-2-isobutyl-4-phenyl-quinolin-3-yl)-ethanone

The title compound was prepared from (2-Amino-5-bromo-phenyl)-phenyl-methanone [example A16] and 6-methyl-2,4-heptanedione according to the procedure of
20 example 1. Yield: 9 %. MS: $m/z = 381$ (M).

Example 5

1-(6-Bromo-2-methyl-4-phenyl-quinolin-3-yl)-3-methyl-butan-1-one

The title compound was prepared from (2-Amino-5-bromo-phenyl)-phenyl-methanone [example A16] and 6-methyl-2,4-heptanedione according to the procedure of
25 example 1. Yield: 55 %. MS: $m/z = 381$ (M).

Example 6

1-(6-Bromo-2-methyl-4-phenyl-quinolin-3-yl)-2,2-difluoro-ethanone

The title compound was prepared from (2-Amino-5-bromo-phenyl)-phenyl-methanone [example A16] and 1,1-difluoroacetylacetone according to the procedure of example 1, except that the residue was purified by chromatography on aminated silica gel with heptane/ethyl acetate (5:1). Yield: 36 %. MS: $m/z = 377$ (M).

5

Example 7

1-[4-(4-Chloro-phenyl)-2-methyl-6-trifluoromethoxy-quinolin-3-yl]-ethanone

The title compound was prepared from (2-Amino-5-trifluoromethoxy-phenyl)-(4-chloro-phenyl)-methanone [example A3] and acetylacetone according to the method of example 1, except that heptane/ethyl acetate (1:2) was used. Yield: 61 %. MS: $m/z = 379$ (M).

10

Example 8

1-(6-Bromo-2-isopropyl-4-phenyl-quinolin-3-yl)-2-methyl-propan-1-one

The title compound was prepared from (2-Amino-5-bromo-phenyl)-phenyl-methanone [example A16] and 2,6-dimethyl-3,5-heptanedione according to the method of example 1, except that the reaction time was of 96h and the residue was purified by chromatography on aminated silica gel with heptane/ethyl acetate (85:15). Yield: 46%. MS: $m/z = 395/397$ (M).

15

Example 9

1-[4-(3,4-Dichloro-phenyl)-2-methyl-6-trifluoromethoxy-quinolin-3-yl]-ethanone

20

The title compound was prepared from (2-Amino-5-trifluoromethoxy-phenyl)-(3,4-dichloro-phenyl)-methanone [example A4] and acetylacetone according to the method of example 1, except that heptane/ethyl acetate (1:2). Yield: 59%. MS: $m/z = 414$ (M).

Example 10

25

1-[4-(4-Chloro-phenyl)-2-methyl-6-phenoxy-quinolin-3-yl]-ethanone

30

The title compound was prepared from (2-Amino-5-phenoxy-phenyl)-(4-chloro-phenyl)-methanone [example A5] and acetylacetone according to the method of example 1, except that the solvent was isopropanol, the reaction time was of 16.5h and the residue was purified by spontaneous crystallization from the reaction mixture. Yield: 42 %; MS: $m/z = 387$ (M).

Example 11

2,2,2-Trifluoro-1-(2-methyl-4-phenyl-6-trifluoromethoxy-quinolin-3-yl)-ethanone

The title compound was prepared from (2-Amino-5-trifluoromethoxy-phenyl)-phenyl-methanone [example A6] and 1,1,1-trifluoro-2,4-pentanedione according to the procedure of example 1, except that the reaction time was of 44h and heptane/ethyl acetate (10:1) was used. Yield: 63 %; MS: m/z = 399 (M).

Example 12

1-[4-(3-Chloro-phenyl)-2-methyl-6-trifluoromethoxy-quinolin-3-yl]-2,2,2-trifluoro-ethanone

The title compound was prepared from (2-Amino-5-trifluoromethoxy-phenyl)-(3-chloro-phenyl)-methanone [example A7] and 1,1,1-trifluoro-2,4-pentanedione according to the procedure of example 1, except that the solvent was isopropanol, the reaction time was of 60h and heptane/ethyl acetate (1:2) was used. Yield: 58 %; MS: m/z = 433 (M).

Example 13

[4-(3-Chloro-phenyl)-2-methyl-6-trifluoromethoxy-quinolin-3-yl]-phenyl-methanone

The title compound was prepared from (2-Amino-5-trifluoromethoxy-phenyl)-(3-chloro-phenyl)-methanone [example A7] and benzoylacetone according to the procedure of example 1, except that the solvent was isopropanol, and heptane/ethyl acetate (1:2) was used. Yield: 57 %; MS: m/z = 441 (M).

Example 14

1-[4-(4-Chloro-phenyl)-2-methyl-6-trifluoromethoxy-quinolin-3-yl]-2,2,2-trifluoro-ethanone

The title compound was prepared from (2-Amino-5-trifluoromethoxy-phenyl)-(4-chloro-phenyl)-methanone [example A3] and 1,1,1-trifluoro-2,4-pentanedione according to the procedure of example 1, except that the solvent was isopropanol, the reaction time was of 16h and heptane/ethyl acetate (1:2) was used. Yield: 34 %; MS: m/z = 433 (M).

Example 15

1-[4-(3-Chloro-phenyl)-2-methyl-6-trifluoromethoxy-quinolin-3-yl]-ethanone

The title compound was prepared from (2-Amino-5-trifluoromethoxy-phenyl)-(3-chloro-phenyl)-methanone [example A7] and acetylacetone according to the procedure of example 1, except that the solvent was isopropanol, and heptane/ethyl acetate (1:2) was used. Yield: 55 %; MS: $m/z = 380$ (M+H).

5

Example 16

2,2,2-Trifluoro-1-[4-(4-methoxy-phenyl)-2-methyl-6-trifluoromethoxy-quinolin-3-yl]-ethanone

The title compound was prepared from (2-Amino-5-trifluoromethoxy-phenyl)-(4-methoxy-phenyl)-methanone [example A8] and 1,1,1-trifluoro-2,4-pentanedione according to the procedure of example 1, except that the solvent was isopropanol, the reaction time was of 17h and a gradient of heptane/ethyl acetate was used. Yield: 58 %; MS: $m/z = 429$ (M).

10

Example 17

2,2,2-Trifluoro-1-[4-(4-fluoro-phenyl)-2-methyl-6-trifluoromethoxy-quinolin-3-yl]-ethanone

15

The title compound was prepared from (2-Amino-5-trifluoromethoxy-phenyl)-(4-fluoro-phenyl)-methanone [example A9] and 1,1,1-trifluoro-2,4-pentanedione according to the procedure of example 1, except that the solvent was isopropanol, the reaction time was of 16h and heptane/ethyl acetate (1:2) was used. Yield: 96 %; MS: $m/z = 417$ (M).

20

Example 18

1-(6-tert-Butyl-2-methyl-4-phenyl-quinolin-3-yl)-2,2,2-trifluoro-ethanone

25

The title compound was prepared from (2-Amino-5-tert-butyl-phenyl)-phenyl-methanone [example A1] and 1,1,1-trifluoro-2,4-pentanedione according to the procedure of example 1, except that the solvent was isopropanol, the reaction time was of 16h and heptane/ethyl acetate (1:2) was used. Yield: 37 %; MS: $m/z = 372$ (M+H).

Example 19

1-(2,2-Difluoro-6-methyl-8-phenyl-[1,3]dioxolo[4,5-g]quinolin-7-yl)-2,2,2-trifluoro-ethanone

The title compound was prepared from (6-Amino-2,2-difluoro-benzo[1,3]dioxol-5-yl)-phenyl-methanone [example A10] and 1,1,1-trifluoro-2,4-pentanedione according

30

to the procedure of example 1, except that the solvent was isopropanol, the reaction time was of 16h and heptane/ethyl acetate (1:2) was used. Yield: 30 %; MS: $m/z = 396$ (M+H).

Example 20

1-[4-(3,4-Difluoro-phenyl)-2-methyl-6-trifluoromethoxy-quinolin-3-yl]-2,2,2-trifluoro-
ethanone

5

The title compound was prepared from (2-Amino-5-trifluoromethoxy-phenyl)-(3,4-difluoro-phenyl)-methanone [example A12] and 1,1,1-trifluoro-2,4-pentanedione according to the procedure of example 1, except that the solvent was isopropanol, the reaction time was of 16h and heptane/ethyl acetate (1:2) was used. Yield: 36 %; MS: $m/z =$
10 435 (M).

Example 21

2,2,2-Trifluoro-1-[4-(4-methanesulfonyl-phenyl)-2-methyl-6-trifluoromethoxy-quinolin-3-yl]-ethanone

The title compound was prepared from (2-Amino-5-trifluoromethoxy-phenyl)-(4-methanesulfonyl-phenyl)-methanone [example A13] and 1,1,1-trifluoro-2,4-pentanedione according to the procedure of example 1, except that the solvent was isopropanol, the reaction time was of 16h and heptane/ethyl acetate (1:2) was used. Yield: 45 %; MS: $m/z = 477$ (M).

15

Example 22

20 2,2,2-Trifluoro-1-[4-(3-fluoro-4-methoxy-phenyl)-2-methyl-6-trifluoromethoxy-quinolin-3-yl]-ethanone

The title compound was prepared from (2-Amino-5-trifluoromethoxy-phenyl)-(4-methanesulfonyl-phenyl)-methanone [example A14] and 1,1,1-trifluoro-2,4-pentanedione according to the procedure of example 1, except that the solvent was isopropanol, the reaction time was of 16h and heptane/ethyl acetate (1:2) was used. Yield: 62 %; MS: $m/z = 477$ (M).

25

Example 23

1-[6-Bromo-4-(4-methanesulfonyl-phenyl)-2-methyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone

30

The title compound was prepared from (2-Amino-5-bromo-phenyl)-(4-methanesulfonyl-phenyl)-methanone [example A15] and 1,1,1-trifluoro-2,4-

pentanedione according to the procedure of example 1, except that the solvent was isopropanol, the reaction time was of 16h and heptane/ethyl acetate (1:2) was used. Yield: 55 %; MS: $m/z = 473$ (M+H).

Example 24

5 1-[6-Bromo-4-(4-fluoro-phenyl)-2-methyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone

The title compound was prepared from (2-Amino-5-bromo-phenyl)-(4-fluoro-phenyl)-methanone [example A2] and 1,1,1-trifluoro-2,4-pentanedione according to the procedure of example 1, except that the solvent was isopropanol, the reaction time was of 16h and heptane/ethyl acetate (1:2) was used. Yield: 80 %; MS: $m/z = 411/413$ (M).

10

Example 25

2,2,2-Trifluoro-1-(2-methyl-4-phenyl-6-piperidin-1-yl-quinolin-3-yl)-ethanone

The title compound was prepared according to the general procedure of method B. Following a methodology developed by J. P. Wolfe and S. L. Buchwald (*J. Org. Chem.* 2000, 65, 1144-1157) a screw-topped pressure-resistant glass vial (50 mL) equipped with
15 a magnetic stirring bar was flushed with a stream of argon and charged with tris(dibenzylideneacetone)dipalladium chloroform complex (0.01 mmol), rac-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl (0.02 mmol), dioxane (7.5 mL) and *tert.*-butanol (7.5 mL) and then stirred for 1 min before adding cesium carbonate (1.4 mmol), 1-(6-Bromo-2-methyl-4-phenyl-quinolin-3-yl)-2,2,2-trifluoro-ethanone [example 3] (1
20 mmol), and piperidine (1.2 mmol). The glass vial was covered with a pressure-resistant seal, firmly locked by a screw-cap and heated with stirring in an oil bath at 120 °C during 2h. The glass vial was cooled in ice before opening, the reaction mixture was diluted with heptane (5 mL) and filtered through a plug of Dicalite filter-aid and rinsed with heptane. The filtrate was evaporated and the residue purified by chromatography on silica gel in
25 heptane/ethyl acetate (4:1). Yield: 74 %; MS: $m/z = 399$ (M+H).

Example 26

2,2,2-Trifluoro-1-(2-methyl-6-morpholin-4-yl-4-phenyl-quinolin-3-yl)-ethanone

The title compound was prepared from 1-(6-Bromo-2-methyl-4-phenyl-quinolin-3-yl)-2,2,2-trifluoro-ethanone [example 3] and morpholine according to the procedure
30 of example 25. Yield: 49 %; MS: $m/z = 401$ (M+H).

Example 27

2,2,2-Trifluoro-1-(2-methyl-4-phenyl-6-pyrrolidin-1-yl-quinolin-3-yl)-ethanone

The title compound was prepared from 1-(6-Bromo-2-methyl-4-phenyl-quinolin-3-yl)-2,2,2-trifluoro-ethanone [example 3] and pyrrolidine according to the procedure of example 25. Yield: 56 %; MS: m/z = 385 (M+H).

Example 28

5 2,2,2-Trifluoro-1-[2-methyl-6-(2-methyl-pyrrolidin-1-yl)-4-phenyl-quinolin-3-yl]-ethanone

The title compound was prepared from 1-(6-Bromo-2-methyl-4-phenyl-quinolin-3-yl)-2,2,2-trifluoro-ethanone [example 3] and 2-methylpyrrolidine according to the procedure of example 25, except that the reaction time was of 16h. Yield: 21 %; MS: m/z
10 = 399 (M+H).

Example 29

2,2,2-Trifluoro-1-[2-methyl-6-(4-methyl-piperazin-1-yl)-4-phenyl-quinolin-3-yl]-ethanone

The title compound was prepared from 1-(6-Bromo-2-methyl-4-phenyl-quinolin-3-yl)-2,2,2-trifluoro-ethanone [example 3] and N-methylpiperazine according to the procedure of example 25, except that the reaction time was of 16h. Yield: 73 %; MS: m/z
15 = 414 (M+H).

Example 30

2,2,2-Trifluoro-1-[6-(3-hydroxy-pyrrolidin-1-yl)-2-methyl-4-phenyl-quinolin-3-yl]-ethanone
20

The title compound was prepared from 1-(6-Bromo-2-methyl-4-phenyl-quinolin-3-yl)-2,2,2-trifluoro-ethanone [example 3] and 3-pyrrolidinol according to the procedure of example 25, except that the reaction time was of 16h. Yield: 92 %; MS: m/z = 401 (M+H).

Example 31

25 2,2,2-Trifluoro-1-[6-(3-hydroxy-azetidin-1-yl)-2-methyl-4-phenyl-quinolin-3-yl]-ethanone

The title compound was prepared from 1-(6-Bromo-2-methyl-4-phenyl-quinolin-3-yl)-2,2,2-trifluoro-ethanone [example 3] and azetidin-3-ol according to the procedure of example 25, except that the reaction time was of 16h. Yield: 22 %; MS: m/z = 387
30 (M+H).

Example 32

2,2,2-Trifluoro-1-[4-(4-methanesulfonyl-phenyl)-2-methyl-6-piperidin-1-yl-quinolin-3-yl]-ethanone

The title compound was prepared from 1-[6-Bromo-4-(4-methanesulfonyl-phenyl)-2-methyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone [example 23] and piperidine according to the procedure of example 25, except that the reaction time was of 16h. Yield: 36 %; MS: $m/z = 277$ (M+H).

Example 33

2,2,2-Trifluoro-1-[4-(4-fluoro-phenyl)-2-methyl-6-piperidin-1-yl-quinolin-3-yl]-ethanone

The title compound was prepared from 1-[6-Bromo-4-(4-fluoro-phenyl)-2-methyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone [example 24] and piperidine according to the procedure of example 25, except that the reaction time was of 16h. Yield: 84 %; MS: $m/z = 417$ (M+H).

Example 34

2,2,2-Trifluoro-1-[4-(4-fluoro-phenyl)-2-methyl-6-morpholin-4-yl-quinolin-3-yl]-ethanone

The title compound was prepared from 1-[6-Bromo-4-(4-fluoro-phenyl)-2-methyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone [example 24] and morpholine according to the procedure of example 25, except that the reaction time was of 16h. Yield: 75 %; MS: $m/z = 419$ (M+H).

Example 35

2,2,2-Trifluoro-1-[4-(4-methanesulfonyl-phenyl)-2-methyl-6-pyrrolidin-1-yl-quinolin-3-yl]-ethanone

The title compound was prepared from 1-[6-Bromo-4-(4-methanesulfonyl-phenyl)-2-methyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone [example 23] and pyrrolidine according to the procedure of example 25, except that the reaction time was of 16h. Yield: 51 %; MS: $m/z = 463$ (M+H).

Example 36

2,2,2-Trifluoro-1-[4-(4-fluoro-phenyl)-2-methyl-6-pyrrolidin-1-yl-quinolin-3-yl]-ethanone

The title compound was prepared from 1-[6-Bromo-4-(4-fluoro-phenyl)-2-methyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone [example 24] and pyrrolidine according to the procedure of example 25, except that the reaction time was of 16h. Yield: 64 %; MS: $m/z = 403$ (M+H).

5

Example 37

2,2,2-Trifluoro-1-[4-(4-fluoro-phenyl)-6-(3-hydroxy-pyrrolidin-1-yl)-2-methyl-quinolin-3-yl]-ethanone

The title compound was prepared from 1-[6-Bromo-4-(4-fluoro-phenyl)-2-methyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone [example 24] and 3-pyrrolidinol according to the procedure of example 25, except that the reaction time was of 16h. Yield: 39 %; MS: $m/z = 419$ (M+H).

10

Example 38

2,2,2-Trifluoro-1-[6-(4-hydroxy-4-phenyl-piperidin-1-yl)-2-methyl-4-phenyl-quinolin-3-yl]-ethanone

15

The title compound was prepared from 1-[6-Bromo-4-(4-phenyl)-2-methyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone [example 24] and 4-hydroxy-4-phenylpiperidine according to the procedure of example 25, except that the reaction time was of 16h. Yield: 29 %; MS: $m/z = 491$ (M+H).

Example 39

2,2,2-Trifluoro-1-[6-(4-hydroxy-4-phenyl-piperidin-1-yl)-4-(4-methanesulfonyl-phenyl)-2-methyl-quinolin-3-yl]-ethanone

20

The title compound was prepared from 1-[6-Bromo-4-(4-methanesulfonyl-phenyl)-2-methyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone [example 24] and 4-hydroxy-4-phenylpiperidine according to the procedure of example 25, except that the reaction time was of 16h. Yield: 17 %; MS: $m/z = 569$ (M+H).

25

Example 40

2,2,2-Trifluoro-1-[4-(4-fluoro-phenyl)-6-(4-hydroxy-4-phenyl-piperidin-1-yl)-2-methyl-quinolin-3-yl]-ethanone

The title compound was prepared from 1-[6-Bromo-4-(4-fluoro-phenyl)-2-methyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone [example 24] and 4-hydroxy-4-

30

phenylpiperidine according to the procedure of example 25, except that the reaction time was of 16h. Yield: 32 %; MS: $m/z = 509$ (M+H).

Example 41

2,2,2-Trifluoro-1-[4-(4-fluoro-phenyl)-6-(3-hydroxy-azetidin-1-yl)-2-methyl-quinolin-3-yl]-ethanone

The title compound was prepared from 1-[6-Bromo-4-(4-fluoro-phenyl)-2-methyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone [example 24] and azetidin-3-ol according to the procedure of example 25, except that the reaction time was of 12h. Yield: 12 %; MS: $m/z = 405$ (M+H).

Example 42

1-[6-Azepan-1-yl-4-(4-fluoro-phenyl)-2-methyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone

The title compound was prepared from 1-[6-Bromo-4-(4-fluoro-phenyl)-2-methyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone [example 24] and azepane according to the procedure of example 25, except that the reaction time was of 16h. Yield: 32 %; MS: $m/z = 431$ (M+H).

Example 43

1-(6-Azepan-1-yl-2-methyl-4-phenyl-quinolin-3-yl)-2,2,2-trifluoro-ethanone

The title compound was prepared from 1-(6-Bromo-2-methyl-4-phenyl-quinolin-3-yl)-2,2,2-trifluoro-ethanone [example 3] and azepane according to the procedure of example 25, except that the reaction time was of 16h. Yield: 14 %; MS: $m/z = 413$ (M+H).

Example 44

1-[6-(3-Dimethylamino-pyrrolidin-1-yl)-2-methyl-4-phenyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone

The title compound was prepared from 1-(6-Bromo-2-methyl-4-phenyl-quinolin-3-yl)-2,2,2-trifluoro-ethanone [example 3] and 3-(dimethylamino)pyrrolidine according to the procedure of example 25, except that the reaction time was of 16h and that ethyl acetate/methanol (9:1) was used. Yield: 37 %; MS: $m/z = 428$ (M+H).

Example 45

1-[6-(3-Dimethylamino-pyrrolidin-1-yl)-4-(4-methanesulfonyl-phenyl)-2-methyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone

The title compound was prepared from 1-[6-Bromo-4-(4-methanesulfonyl-phenyl)-2-methyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone [example 23] and 3-(dimethylamino)pyrrolidine according to the procedure of example 25, except that the reaction time was of 16h and that ethyl acetate/methanol (9:1) was used. Yield: 47 %; MS: m/z = 506 (M+H).

Example 46

1-[6-(3-Dimethylamino-pyrrolidin-1-yl)-4-(4-fluoro-phenyl)-2-methyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone

The title compound was prepared from 1-[6-Bromo-4-(4-fluoro-phenyl)-2-methyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone [example 24] and 3-(dimethylamino)pyrrolidine according to the procedure of example 25, except that the reaction time was of 16h and that ethyl acetate/methanol (9:1) was used. Yield: 44 %; MS: m/z = 446 (M+H).

Example 47

2,2,2-Trifluoro-1-[6-iodo-4-(4-methanesulfonyl-phenyl)-2-methyl-quinolin-3-yl]-ethanone

The title compound was prepared from (2-Amino-5-iodo-phenyl)-(4-methanesulfonyl-phenyl)-methanone [example A17] and 1,1,1-trifluoro-2,4-pentanedione according to the procedure of example 1, except that the solvent was isopropanol, the reaction time was of 16 h and heptane/ethyl acetate (1:2) was used. Yield: 55 %; MS: m/z = 519 (M).

Example 48

2,2,2-Trifluoro-1-[6-(4-hydroxy-4-methyl-piperidin-1-yl)-4-(4-methanesulfonyl-phenyl)-2-methyl-quinolin-3-yl]-ethanone

The title compound was prepared from 1-[6-bromo-4-(4-methanesulfonyl-phenyl)-2-methyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone [example 23] and 4-methyl-

piperidin-4-ol according to the procedure of example 25. Yield: 14 %; MS: $m/z = 507$ (M+H).

Example 49

2,2,2-Trifluoro-1-[4-(4-methanesulfonyl-phenyl)-6-(4-methoxy-piperidin-1-yl)-2-methyl-quinolin-3-yl]-ethanone

The title compound was prepared from 1-[6-bromo-4-(4-methanesulfonyl-phenyl)-2-methyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone [example 23] and 4-methoxy-piperidine according to the procedure of example 25. Yield: 14 %; MS: $m/z = 507$ (M+H).

Example 50

[6-Bromo-4-(4-methanesulfonyl-phenyl)-2-methyl-quinolin-3-yl]-cyclopropyl-methanone

The title compound was prepared from (2-Amino-5-bromo-phenyl)-(4-methanesulfonyl-phenyl)-methanone [example A15] and 1-cyclopropyl-1,3-butanedione according to the procedure of example 1, except that the solvent was isopropanol, the reaction time was of 16 h and heptane/ethyl acetate (1:2) was used. Yield: 47 %; MS: $m/z = 443$ (M+H).

Example 51

Cyclopropyl-[4-(4-methanesulfonyl-phenyl)-2-methyl-6-morpholin-4-yl-quinolin-3-yl]-methanone

A tube placed under argon was charged with tris(dibenzylideneacetone)dipalladium chloroform complex (5 mg), 2-dicyclohexylphosphino-2',4',6'-triisopropylbiphenyl (5 mg) and cesium carbonate (110 mg, 0.33 mmol). [6-Bromo-4-(4-methanesulfonyl-phenyl)-2-methyl-quinolin-3-yl]-cyclopropyl-methanone [example 50] (100 mg, 0.22 mmol) in t-BuOH (5ml) was added, followed by morpholine (24 mg, 0.27 mmol). The tube was sealed and heated at 110 °C for 6 hrs. The reaction mixture was cooled to 20 °C, diluted with heptane, and filtered through Celite and purified directly by flash chromatography on silica gel in heptane / AcOEt 80 : 20 to give a yellow solid (52 mg, 51%). MS: $m/z = 451$ (M + H).

Example 52

1-[6-(3,3-Difluoro-piperidin-1-yl)-4-(4-methanesulfonyl-phenyl)-2-methyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone

The title compound was prepared from 1-[6-Bromo-4-(4-methanesulfonyl-phenyl)-2-methyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone [example 23] and 3,3-difluoropiperidine hydrochloride according to the procedure of example 51. Yield: 50 %; MS: $m/z = 513$ (M+H).

Example 53

2,2,2-Trifluoro-1-[4-(4-methanesulfonyl-phenyl)-2-methyl-6-(8-oxa-3-aza-bicyclo[3.2.1]oct-3-yl)-quinolin-3-yl]-ethanone

The title compound was prepared from 1-[6-bromo-4-(4-methanesulfonyl-phenyl)-2-methyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone [example 23] and 8-oxa-3-aza-bicyclo[3.2.1]octane according to the procedure of example 51. Yield: 61 %; MS: $m/z = 505$ (M+H).

Example 54

Cyclopropyl-[4-(4-methanesulfonyl-phenyl)-2-methyl-6-piperidin-1-yl-quinolin-3-yl]-methanone

The title compound was prepared from [6-Bromo-4-(4-methanesulfonyl-phenyl)-2-methyl-quinolin-3-yl]-cyclopropyl-methanone [example 50] and piperidine according to the procedure of example 51. Yield: 39 %; MS: $m/z = 449$ (M+H).

Example 55

[(6-Bromo-2-methyl-4-phenyl-quinolin-3-yl)-cyclopropyl-methanone

The title compound was prepared from (2-amino-5-bromo-phenyl)-phenyl-methanone [example A16] and 1-cyclopropyl-1,3-butanedione according to the procedure of example 1, except that the solvent was isopropanol, the reaction time was of 16 h and heptane/ethyl acetate (1:2) was used. Yield: 75 %; MS: $m/z = 366$ (M).

Example 56

[6-Bromo-4-(4-fluoro-phenyl)-2-methyl-quinolin-3-yl]-cyclopropyl-methanone

The title compound was prepared from (2-amino-5-trifluoromethoxy-phenyl)-(4-fluorophenyl)-methanone [example A9] and 1-cyclopropyl-1,3-butanedione according to the procedure of example 1, except that the solvent was isopropanol, the

reaction time was of 16 h and heptane/ethyl acetate (1:2) was used. Yield: 76 %; MS: m/z = 384 (M).

Example 57

Cyclopropyl-(2-methyl-4-phenyl-6-piperidin-1-yl-quinolin-3-yl)-methanone

- 5 The title compound was prepared from [(6-Bromo-2-methyl-4-phenyl-quinolin-3-yl)-cyclopropyl-methanone [example 55] and piperidine according to the procedure of example 51. Yield: 64 %; MS: m/z = 371 (M+H).

Example 58

- 10 Cyclopropyl-(2-methyl-6-morpholin-4-yl-4-phenyl-quinolin-3-yl)-methanone

The title compound was prepared from [(6-Bromo-2-methyl-4-phenyl-quinolin-3-yl)-cyclopropyl-methanone [example 55] and morpholine according to the procedure of example 51. Yield: 67 %; MS: m/z = 373 (M+H).


International Patent Appl. No. PCT/EP2005/011403

Case 22589 WO

-37-

Claims

1. Compounds of formula I



wherein

R^1 is hydrogen, C_1 - C_7 alkyl, C_1 - C_7 haloalkyl, di(C_1 - C_7)alkylamino, C_3 - C_8 cycloalkyl, or 5 or 6 membered heterocycloalkyl;

R^2 is C_1 - C_7 alkyl, aryl, C_1 - C_7 alkoxy(C_1 - C_7)alkyl, C_1 - C_7 haloalkyl or C_3 - C_8 cycloalkyl;

R^3 , R^4 are each independently hydrogen, halo, hydroxy, C_1 - C_7 alkoxy, C_1 - C_7 haloalkoxy, di(C_1 - C_7)alkylamino, C_1 - C_7 alkylsulfonyl, or 5 or 6 membered heterocycloalkyl;

R^5 is hydrogen, halo, C_1 - C_7 alkyl, C_1 - C_7 alkoxy, C_1 - C_7 haloalkoxy, or aryloxy, or is $-NR^7R^8$ wherein R^7 and R^8 are C_1 - C_7 alkyl, or R^7 and R^8 may, together with the nitrogen atom to which they are attached, form or a 4 to 7 membered heterocycloalkyl group which may be substituted by one or more substituent(s) selected from the group consisting of halo, C_1 - C_7 alkyl, C_1 - C_7 alkoxy, hydroxy, phenyl and di(C_1 - C_7)alkylamino;

R^6 is hydrogen or together with R^5 may form a 5 or 6 membered heterocycloalkyl group which may be substituted by one or more halogen;

and pharmaceutically acceptable acid addition salts thereof,

excluding the following compounds:

1-(6-Chloro-2-methyl-4-phenyl-quinolin-3-yl)-ethanone;

1-(6-Bromo-4-phenyl-2-piperidin-1-yl-quinolin-3-yl)-ethanone;

1-[4-(4-Chloro-phenyl)-2-methyl-quinolin-3-yl]-ethanone;

1-(6-Bromo-2-methyl-4-phenyl-quinolin-3-yl)-ethanone;

1-(2,6-Dimethyl-4-phenyl-quinolin-3-yl)-ethanone;

1-(2-Methyl-4-phenyl-6-trifluoromethoxy-quinolin-3-yl)-ethanone

Phenyl-(4-phenyl-quinolin-3-yl)-methanone;

International Patent Appl. No. PCT/EP2005/011403

Case 22589 WO

-38-

Phenyl-(4-phenyl-quinolin-3-yl)-(2,4,6-trimethyl-phenyl)-methanone;
 (2-Methyl-4-phenyl-quinolin-3-yl)-phenyl-methanone;
 1-(2-Methyl-4-phenyl-quinolin-3-yl)-ethanone;
 1-(6-Chloro-2-methyl-4-phenyl-quinolin-3-yl)-ethanone;
 1-(4-Phenyl-quinolin-3-yl)-ethanone;
 1-(6-Chloro-4-phenyl-quinolin-3-yl)-ethanone;
 1-(6-Chloro-2-dimethylamino-4-phenyl-quinolin-3-yl)-ethanone;
 1-(6-Methoxy-2-methyl-4-phenyl-quinolin-3-yl)-ethanone;
 2,2,2-Trifluoro-1-(2-methyl-4-phenyl-quinolin-3-yl)-ethanone;
 1-(6-Bromo-2-(4-morpholinyl)-4-phenyl-quinolin-3-yl)-ethanone;
 1-(2,6-Dimethyl-4-phenyl-quinolin-3-yl)-ethanone;
 1-[4-(4-Chlorophenyl)-2-methyl-quinolin-3-yl]-ethanone;
 1-[6-Chloro-4-phenyl-2-(1-piperidinyl)-quinolin-3-yl]-ethanone;
 1-[6-Bromo-4-phenyl-2-(1-piperidinyl)-quinolin-3-yl]-ethanone;
 (3-Methylphenyl) (4-phenyl-quinolin-3-yl)-methanone;
 (4-Chlorophenyl) (7-chloro-4-phenyl-quinolin-3-yl)-methanone;
 (7-Chloro-4-phenyl-quinolin-3-yl) phenyl-methanone;
 (7-Chloro-4-phenyl-quinolin-3-yl) (4-methoxyphenyl)-methanone;
 (7-Chloro-4-phenyl-quinolin-3-yl) (4-ethylphenyl)-methanone;
 (4-Chlorophenyl) (4-phenyl-quinolin-3-yl)-methanone;
 (4-Ethylphenyl) (4-phenyl-quinolin-3-yl)-methanone; and
 (7-Chloro-4-phenyl-quinolin-3-yl) (4-methylphenyl)-methanone.

2. Compounds of formula I according to claim 1, wherein

- R^1 is C_1 - C_7 alkyl;
 R^2 is C_1 - C_7 alkyl, phenyl, C_1 - C_7 haloalkyl;
 R^3, R^4 are each independently hydrogen, halo, C_1 - C_7 alkoxy, C_1 - C_7 alkylsulfonyl,
 R^5 is halo, C_1 - C_7 haloalkoxy, aryloxy, or $-NR^7R^8$ wherein R^7 and R^8 are C_1 - C_7 alkyl, may, together with the nitrogen atom to which they are attached, form or a 4 to 7 membered heterocycloalkyl group which may be substituted by one or more substituent(s) selected

International Patent Appl. No. PCT/EP2005/011403

Case 22589 WO

-39-

from the group consisting of halo, C₁-C₇ alkyl, hydroxy, C₁-C₇ alkoxy, phenyl and di(C₁-C₇)alkylamino;

R⁶ is hydrogen or together with R⁵ may form a 5 or 6 membered heterocycloalkyl group which may be substituted by one or more halogen; and pharmaceutically acceptable acid addition salts thereof.

3. Compounds of formula I according to claim 1, wherein R² is C₁-C₇ alkyl.

4. Compounds of formula I according to claim 3, wherein the compounds are selected from the group consisting of:

1-(6-Bromo-2-ethyl-4-phenyl-quinolin-3-yl)-propan-1-one;
1-(6-Bromo-2-isobutyl-4-phenyl-quinolin-3-yl)-ethanone;
1-(6-Bromo-2-methyl-4-phenyl-quinolin-3-yl)-3-methyl-butan-1-one;
1-[4-(4-Chloro-phenyl)-2-methyl-6-trifluoromethoxy-quinolin-3-yl]-ethanone;
1-(6-Bromo-2-isopropyl-4-phenyl-quinolin-3-yl)-2-methyl-propan-1-one;
1-[4-(3,4-Dichloro-phenyl)-2-methyl-6-trifluoromethoxy-quinolin-3-yl]-ethanone;
1-[4-(4-Chloro-phenyl)-2-methyl-6-phenoxy-quinolin-3-yl]-ethanone; and
1-[4-(3-Chloro-phenyl)-2-methyl-6-trifluoromethoxy-quinolin-3-yl]-ethanone.

5. Compounds of formula I according to claim 1, wherein R² is C₁-C₇ haloalkyl.

6. Compounds of formula I according to claim 5, wherein the compounds are selected from the group consisting of:

1-(6-Bromo-2-methyl-4-phenyl-quinolin-3-yl)-2,2,2-trifluoro-ethanone;
1-(6-Bromo-2-methyl-4-phenyl-quinolin-3-yl)-2,2-difluoro-ethanone;
2,2,2-Trifluoro-1-(2-methyl-4-phenyl-6-trifluoromethoxy-quinolin-3-yl)-ethanone;
1-[4-(3-Chloro-phenyl)-2-methyl-6-trifluoromethoxy-quinolin-3-yl]-2,2,2-trifluoro-ethanone;
1-[4-(4-Chloro-phenyl)-2-methyl-6-trifluoromethoxy-quinolin-3-yl]-2,2,2-trifluoro-ethanone;
2,2,2-Trifluoro-1-[4-(4-methoxy-phenyl)-2-methyl-6-trifluoromethoxy-quinolin-3-yl]-ethanone;
2,2,2-Trifluoro-1-[4-(4-fluoro-phenyl)-2-methyl-6-trifluoromethoxy-quinolin-3-yl]-ethanone;
1-(6-tert-Butyl-2-methyl-4-phenyl-quinolin-3-yl)-2,2,2-trifluoro-ethanone;

International Patent Appl. No. PCT/EP2005/011403

Case 22589 WO

-40-

1-(2,2-Difluoro-6-methyl-8-phenyl-[1,3]dioxolo[4,5-g]quinolin-7-yl)-2,2,2-trifluoro-ethanone;
1-[4-(3,4-Difluoro-phenyl)-2-methyl-6-trifluoromethoxy-quinolin-3-yl]-2,2,2-trifluoro-ethanone;
2,2,2-Trifluoro-1-[4-(4-methanesulfonyl-phenyl)-2-methyl-6-trifluoromethoxy-quinolin-3-yl]-ethanone;
2,2,2-Trifluoro-1-[4-(3-fluoro-4-methoxy-phenyl)-2-methyl-6-trifluoromethoxy-quinolin-3-yl]-ethanone;
2,2,2-Trifluoro-1-(2-methyl-4-phenyl-6-piperidin-1-yl-quinolin-3-yl)-ethanone;
2,2,2-Trifluoro-1-(2-methyl-6-morpholin-4-yl-4-phenyl-quinolin-3-yl)-ethanone;
2,2,2-Trifluoro-1-(2-methyl-4-phenyl-6-pyrrolidin-1-yl-quinolin-3-yl)-ethanone;
2,2,2-Trifluoro-1-[2-methyl-6-(2-methyl-pyrrolidin-1-yl)-4-phenyl-quinolin-3-yl]-ethanone;
2,2,2-Trifluoro-1-[2-methyl-6-(4-methyl-piperazin-1-yl)-4-phenyl-quinolin-3-yl]-ethanone;
2,2,2-Trifluoro-1-[6-(3-hydroxy-pyrrolidin-1-yl)-2-methyl-4-phenyl-quinolin-3-yl]-ethanone;
2,2,2-Trifluoro-1-[6-(3-hydroxy-azetidin-1-yl)-2-methyl-4-phenyl-quinolin-3-yl]-ethanone;
1-[6-Bromo-4-(4-methanesulfonyl-phenyl)-2-methyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone;
1-[6-Bromo-4-(4-fluoro-phenyl)-2-methyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone;
2,2,2-Trifluoro-1-[4-(4-methanesulfonyl-phenyl)-2-methyl-6-piperidin-1-yl-quinolin-3-yl]-ethanone;
2,2,2-Trifluoro-1-[4-(4-fluoro-phenyl)-2-methyl-6-piperidin-1-yl-quinolin-3-yl]-ethanone;
2,2,2-Trifluoro-1-[4-(4-fluoro-phenyl)-2-methyl-6-morpholin-4-yl-quinolin-3-yl]-ethanone;
2,2,2-Trifluoro-1-[4-(4-methanesulfonyl-phenyl)-2-methyl-6-pyrrolidin-1-yl-quinolin-3-yl]-ethanone;
2,2,2-Trifluoro-1-[4-(4-fluoro-phenyl)-2-methyl-6-pyrrolidin-1-yl-quinolin-3-yl]-ethanone;
2,2,2-Trifluoro-1-[4-(4-fluoro-phenyl)-6-(3-hydroxy-pyrrolidin-1-yl)-2-methyl-quinolin-3-yl]-ethanone;
2,2,2-Trifluoro-1-[6-(4-hydroxy-4-phenyl-piperidin-1-yl)-2-methyl-4-phenyl-quinolin-3-yl]-ethanone;
2,2,2-Trifluoro-1-[6-(4-hydroxy-4-phenyl-piperidin-1-yl)-4-(4-methanesulfonyl-phenyl)-2-methyl-quinolin-3-yl]-ethanone;
2,2,2-Trifluoro-1-[4-(4-fluoro-phenyl)-6-(4-hydroxy-4-phenyl-piperidin-1-yl)-2-methyl-quinolin-3-yl]-ethanone;

International Patent Appl. No. PCT/EP2005/011403

Case 22589 WO

-41-

2,2,2-Trifluoro-1-[4-(4-fluoro-phenyl)-6-(3-hydroxy-azetidin-1-yl)-2-methyl-quinolin-3-yl]-ethanone;

1-[6-Azepan-1-yl-4-(4-fluoro-phenyl)-2-methyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone;

1-(6-Azepan-1-yl-2-methyl-4-phenyl-quinolin-3-yl)-2,2,2-trifluoro-ethanone;

1-[6-(3-Dimethylamino-pyrrolidin-1-yl)-2-methyl-4-phenyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone;

1-[6-(3-Dimethylamino-pyrrolidin-1-yl)-4-(4-methanesulfonyl-phenyl)-2-methyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone;

1-[6-(3-Dimethylamino-pyrrolidin-1-yl)-4-(4-fluoro-phenyl)-2-methyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone;

2,2,2-Trifluoro-1-[6-iodo-4-(4-methanesulfonyl-phenyl)-2-methyl-quinolin-3-yl]-ethanone;

2,2,2-Trifluoro-1-[6-(4-hydroxy-4-methyl-piperidin-1-yl)-4-(4-methanesulfonyl-phenyl)-2-methyl-quinolin-3-yl]-ethanone;

2,2,2-Trifluoro-1-[4-(4-methanesulfonyl-phenyl)-6-(4-methoxy-piperidin-1-yl)-2-methyl-quinolin-3-yl]-ethanone;

1-[6-(3,3-Difluoro-piperidin-1-yl)-4-(4-methanesulfonyl-phenyl)-2-methyl-quinolin-3-yl]-2,2,2-trifluoro-ethanone; and

2,2,2-Trifluoro-1-[4-(4-methanesulfonyl-phenyl)-2-methyl-6-(8-oxa-3-aza-bicyclo[3.2.1]oct-3-yl)-quinolin-3-yl]-ethanone.

7. Compounds of formula I according to claim 1, wherein R² is cycloalkyl.

8. Compounds of formula I according to claim 7, wherein the compounds are selected from the group consisting of:

6-Bromo-4-(4-methanesulfonyl-phenyl)-2-methyl-quinolin-3-yl]-cyclopropyl-methanone;

Cyclopropyl-[4-(4-methanesulfonyl-phenyl)-2-methyl-6-morpholin-4-yl-quinolin-3-yl]-methanone;

Cyclopropyl-[4-(4-methanesulfonyl-phenyl)-2-methyl-6-piperidin-1-yl-quinolin-3-yl]-methanone;

[(6-Bromo-2-methyl-4-phenyl-quinolin-3-yl)-cyclopropyl-methanone;

[6-Bromo-4-(4-fluoro-phenyl)-2-methyl-quinolin-3-yl]-cyclopropyl-methanone;

International Patent Appl. No. PCT/EP2005/011403

Case 22589 WO

-42-

Cyclopropyl-(2-methyl-4-phenyl-6-piperidin-1-yl-quinolin-3-yl)-methanone and
Cyclopropyl-(2-methyl-6-morpholin-4-yl-4-phenyl-quinolin-3-yl)-methanone.


9. Compounds of formula I according to claim 1, wherein R^2 is phenyl.

10. Compounds of formula I according to claim 9, wherein the compounds are selected from the group consisting of:


(6-Bromo-2-methyl-4-phenyl-quinolin-3-yl)-phenyl-methanone; and

[4-(3-Chloro-phenyl)-2-methyl-6-trifluoromethoxy-quinolin-3-yl]-phenyl-methanone.


11. A process for the preparation of a compound of formula I according to any one of claims 1 to 10:



comprising the step of reacting a compound of formula II



with a compound of formula III



to give the compound of formula I;

wherein R^1 to R^6 are as defined for formula I;


International Patent Appl. No. PCT/EP2005/011403

Case 22589 WO


-43-

and if desired, converting the compound of formula I obtained into a pharmaceutically acceptable acid addition salt.


12. A process for the preparation of a compound of formula Ia



comprising the steps of reacting a compound of formula IV



with a compound of formula V



to give the compound of formula Ia;

wherein R¹ to R⁸ are as defined in claim 1;

and if desired, converting the compound of formula Ia obtained into a pharmaceutically acceptable acid addition salt.

13. A compound of formula I, prepared according to the process of any one of claim 11 or 12.

14. A medicament containing a compound of formula I according to any one of claims 1 to 10.

15. A medicament according to claim 14, wherein the medicament is useful in the control or prevention of illnesses, especially of illnesses and disorders comprising anxiety, depression, epilepsy, schizophrenia, cognitive disorders, spasticity and skeletal muscle rigidity, spinal cord injury, multiple sclerosis, amyotrophic lateral sclerosis, cerebral palsy, neuropathic pain and craving associated with cocaine and nicotine, psychosis, panic disorder, posttraumatic stress disorders or gastro-intestinal disorders.

16. The use of a compound of formula I according to any one of claims 1 to 10 or of a compound selected from the group consisting of:

- 1-(6-Chloro-2-methyl-4-phenyl-quinolin-3-yl)-ethanone;
- 1-(6-Bromo-4-phenyl-2-piperidin-1-yl-quinolin-3-yl)-ethanone;
- 1-[4-(4-Chloro-phenyl)-2-methyl-quinolin-3-yl]-ethanone;
- 1-(6-Bromo-2-methyl-4-phenyl-quinolin-3-yl)-ethanone;
- 1-(2,6-Dimethyl-4-phenyl-quinolin-3-yl)-ethanone; and
- 1-(2-Methyl-4-phenyl-6-trifluoromethoxy-quinolin-3-yl)-ethanone,

or of their pharmaceutically acceptable salts, for the manufacture of a medicament.

17. The use according to claim 16, wherein the medicament is useful in the control or prevention of illnesses comprising anxiety, depression, epilepsy, schizophrenia, cognitive disorders, spasticity and skeletal muscle rigidity, spinal cord injury, multiple sclerosis, amyotrophic lateral sclerosis, cerebral palsy, neuropathic pain and craving associated with cocaine and nicotine, psychosis, panic disorder, posttraumatic stress disorders or gastro-intestinal disorders.