
p is an integer ranging from 57 to 1420, more preferably 85 to 850; and

q is selected from the group consisting of 1, 2, 3, 4, 5, and 6.

In a preferred embodiment p of formula (IIb) ranges from 170 to 284, even more preferably from 198 to 255 and most preferably from 215 to 238.

In an equally preferred embodiment p of formula (IIb) ranges from 340 to 568, even more preferably from 398 to 510 and most preferably from 426 to 482.

More preferably, -R^1 comprises a moiety of formula (IIc):



(IIc),

wherein

p1, p2, p3, p4 are independently an integer ranging from 57 to 1420, even more preferably from 85 to 850.

In a preferred embodiment p1, p2, p3 and p4 of formula (IIc) are an integer independently selected from 170 to 284, even more preferably from 198 to 255 and most preferably from 215 to 238.

In an equally preferred embodiment p1, p2, p3 and p4 of formula (IIc) are an integer independently selected from 340 to 568, even more preferably from 398 to 510 and most preferably from 426 to 482.

In a preferred embodiment -R^2 of formula (Ib) is selected from the group consisting of -H, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl and tert-butyl. More preferably, -R^2 of formula (Ib) is selected from the group consisting of -H, methyl, ethyl, n-