

Description

The present invention relates to a process for the production of a seasoning, more particularly to the production of seasoning by the biological hydrolysis of protein containing material.

Hydrolysed proteins have been known for use as seasonings in food systems for centuries in the Far East in the form of soya sauce which traditionally has been prepared by enzymatic hydrolysis requiring a long period of time, usually several months, for preparation. In producing soya sauce, plant protein containing materials such as cooked soya beans or defatted soya flour together with carbohydrates are inoculated with *Aspergilli* and the solid culture is fermented for 2 days to make fermented koji during which time enzymes are produced which are able to hydrolyse protein and carbohydrates in the moromi stage. The fermented koji is mixed with a solution of common salt to give moromi which is fermented for 4 to 8 months by the activity of micro-organisms such as soya lactic acid bacteria and soya yeasts from which the soya sauce is obtained by removing the solids portion from the fermented moromi.

About 100 years ago, a more rapid method of hydrolysing proteins for producing seasonings was developed using hydrochloric acid in which the time required is only a few hours. However, in recent years, the use of acid hydrolysed plant protein (HPP) in culinary applications has been under criticism due to the presence of some chloro-compounds which arises from the acid process. Therefore, attempts have been made to develop HPP replacements which can be used as body-givers in culinary applications. Soya sauce is one such suitable replacement. However, owing to the differences in the raw materials and the processing methods involved, the two products, HPP and soya sauce, have some differences in terms of chemical composition and flavour profile. Dosage of soya sauce which can be used as an HPP replacement is limited due to its "fermented" note. The different processing procedures also result in a significant variation in the degree of hydrolysis of the protein containing material to the amino acids. Soya sauce has a lower amino acid content than HPP and this leads to a significantly weaker body in soya sauce than in HPP.

We have developed a process for the production of a seasoning based on modified standard soya sauce technology in which the fermented koji is treated, before forming the moromi, by a low temperature hydrolysis whereby the resultant seasoning has a stronger body than a standard soya sauce. A similar hydrolysis during which a temperature of 45-60°C is applied is disclosed in EP-A-417 481.

Accordingly, the present invention provides a process of treating a fermented protein koji prepared from a protein containing material and a carbohydrate, for the production of a seasoning, which comprises hydrolysing the fermented protein koji at a temperature of from 2° to 25°C and a pH of from 4.5 to 10 for a period of from 6 hours to 28 days.

The fermented koji is prepared by the conventional soya sauce process which comprises, for example, inoculating a protein containing material and a carbohydrate with a culture of *Aspergillus oryzae* and/or *Aspergillus sojae* on a culture bed to form the fermented koji. The protein containing material is advantageously a plant protein material, for instance, soya beans, corn gluten or rice gluten but is preferably wheat gluten. The plant protein containing material is advantageously cooked and is preferably used in solid particulate form for enabling the mould of *Aspergillus oryzae* and/or *Aspergillus sojae* to grow on the surface of the particles and eventually penetrate into the particles. The koji is conveniently fermented in the solid state.

The hydrolysis of the fermented koji is preferably carried out in the substantial or complete absence of salt and advantageously with constant agitation.

The fermented koji is conveniently hydrolysed by mixing with from 1 to 5 parts of water per part by weight of fermented koji, most advantageously in the absence of salt. The hydrolysis is carried out preferably at a temperature from 2° to 20°C, for a period of from 12 hours to 25 days, more preferably from 3° to 15°C for a period of from 18 hours to 22 days and especially from 4° to 10°C for a period of from 24 hours to 20 days.

After hydrolysis of the fermented koji, salt and yeast are conveniently added to the hydrolysed fermented koji to form a moromi. The moromi is preferably fermented for a period of from 1 to 6 weeks, more preferably from 2 to 4 weeks. The moromi fermentation may be carried out under aerobic or anaerobic conditions.

After the fermentation, the matured moromi may be pressed to separate a liquid sauce from a solid residue. The liquid sauce is advantageously pasteurised e.g. at a temperature of from 80° to 120°C and then filtered to give a liquid seasoning. If desired, the liquid sauce may be made into a powder for instance, by concentration, then vacuum dried to a low moisture content and finally milled into a powder to give a solid seasoning.

The process according to the present invention results in a higher level or degree of release of amino acids than is generally possible with conventional soya sauce processes.

The seasoning either in liquid or powder form has a higher amino acid content than soya sauce prepared by conventional methods. Because of the higher amino acid content, the seasoning of the present invention has more body than soya sauce prepared by conventional methods. The seasoning prepared according to the present invention has excellent organoleptical stability.

The present invention will now be further illustrated by the following Examples in which parts and percentages are given by weight.

Example 1

Wheat gluten was extruded through a Clextral extruder into pieces having an average diameter of 5mm and had a porous structure.

65 kg of the extrudates were soaked in 65 kg water at 75°C for 5 minutes. The soaked extrudates were then heated to 100°C and held at the same temperature for 10 minutes and afterwards cooled to below 40°C by applying vacuum. The pasteurisation step was carried out to eliminate secondary contamination after the extrusion step. Finally the cooked extrudates were mixed with a mixture of 28 kg of roasted wheat and 20 g of TKJ (*Aspergillus oryzae* seed inoculum) to give a wheat gluten koji which was fermented for 42 hours by a procedure similar to that used in a conventional soya sauce process. The wheat gluten koji contained no added salt.

During the 42 hours of koji fermentation, the following temperature profiles were maintained for the culture bed:

0 - 25 hours 30°C

25 - 42 hours 27°C

Similar to a conventional soya sauce process, the koji was mixed at 18th and 25th hours to ensure sufficient airflow through the culture bed for good ventilation.

55 kg of the fermented wheat gluten koji was mixed with 150 kg of water at 10°C which had previously been sterilised by boiling and then cooled. The mixture was hydrolysed for 24 hours in a jacketed enclosed vessel with chilled water circulating through the jacket to maintain the desired temperature. The mixture was agitated continuously during the hydrolysis.

After the hydrolysis, 27 kg of salt and 11 kg of yeast inoculum were added to 200 kg of the hydrolysed fermented wheat gluten koji to give a moromi which was fermented aerobically by bubbling compressed air through the batch for 4 weeks.

Finally, the fermented moromi was pressed to separate a wheat gluten sauce from a solid residue. The wheat gluten sauce was treated at 90°C for 20 minutes. The liquid sauce was concentrated by

evaporation. The concentrate obtained was dried in a vacuum oven and then milled into a powder.

For organoleptic evaluation, 10g of liquid sauce or 3.5 g powder were diluted with 250 ml of boiling water. In both cases the seasoning was found to have more body and a more rounded flavour than a conventional soya sauce.

The powder was found to be shelf stable at 30°C for more than 12 months in moisture tight packaging (alu-laminated sachets) and had excellent colour stability. The seasoning was found to be microbiologically stable.

Example 2

A similar procedure to that described in Example 1 was followed except that the hydrolysis of the fermented wheat gluten koji was carried out over a period of 7 days instead of 24 hours.

The seasoning was found to have more body and a more rounded flavour profile than a conventional soya sauce. The seasoning was microbiologically stable as in Example 1 and had excellent colour stability in powder form.

Example 3

A similar procedure to that described in Example 1 was followed except that the hydrolysis of the fermented wheat gluten koji was carried out at 4°C for 14-20 days.

The seasoning was found to have more body and a more rounded flavour profile than a conventional soya sauce. The seasoning was microbiologically stable as in Example 1 and had excellent colour stability in powder form.

Claims

1. A process of treating a fermented protein koji prepared from a protein containing material and a carbohydrate, for the production of a seasoning, which comprises hydrolysing the fermented protein koji at a temperature of from 2° to 25°C and a pH of from 4.5 to 10 for a period of from 6 hours to 28 days.
2. A process according to claim 1 wherein the protein containing material is a plant protein material.
3. A process according to claim 1 wherein the protein containing material is cooked.
4. A process according to claim 1 wherein the protein containing material is used in particulate form for enabling the mould of *Aspergillus oryzae* and/or *Aspergillus sojae* to grow



The seasoning either in liquid or powder form has a higher amino acid content than soya sauce prepared by conventional methods. Because of the higher amino acid content, the seasoning of the present invention has more body than soya sauce prepared by conventional methods. The seasoning prepared according to the present invention has excellent organoleptical stability.

The present invention will now be further illustrated by the following Examples in which parts and percentages are given by weight.

Example 1

Wheat gluten was extruded through a Clextral extruder into pieces having an average diameter of 5mm and had a porous structure.

65 kg of the extrudates were soaked in 65 kg water at 75°C for 5 minutes. The soaked extrudates were then heated to 100°C and held at the same temperature for 10 minutes and afterwards cooled to below 40°C by applying vacuum. The pasteurisation step was carried out to eliminate secondary contamination after the extrusion step. Finally the cooked extrudates were mixed with a mixture of 28 kg of roasted wheat and 20 g of TKJ (*Aspergillus oryzae* seed inoculum) to give a wheat gluten koji which was fermented for 42 hours by a procedure similar to that used in a conventional soya sauce process. The wheat gluten koji contained no added salt.

During the 42 hours of koji fermentation, the following temperature profiles were maintained for the culture bed:

0 - 25 hours 30°C

25 - 42 hours 27°C

Similar to a conventional soya sauce process, the koji was mixed at 18th and 25th hours to ensure sufficient airflow through the culture bed for good ventilation.

55 kg of the fermented wheat gluten koji was mixed with 150 kg of water at 10°C which had previously been sterilised by boiling and then cooled. The mixture was hydrolysed for 24 hours in a jacketed enclosed vessel with chilled water circulating through the jacket to maintain the desired temperature. The mixture was agitated continuously during the hydrolysis.

After the hydrolysis, 27 kg of salt and 11 kg of yeast inoculum were added to 200 kg of the hydrolysed fermented wheat gluten koji to give a moromi which was fermented aerobically by bubbling compressed air through the batch for 4 weeks.

Finally, the fermented moromi was pressed to separate a wheat gluten sauce from a solid residue. The wheat gluten sauce was treated at 90°C for 20 minutes. The liquid sauce was concentrated by

evaporation. The concentrate obtained was dried in a vacuum oven and then milled into a powder.

For organoleptic evaluation, 10g of liquid sauce or 3.5 g powder were diluted with 250 ml of boiling water. In both cases the seasoning was found to have more body and a more rounded flavour than a conventional soya sauce.

The powder was found to be shelf stable at 30°C for more than 12 months in moisture tight packaging (alu-laminated sachets) and had excellent colour stability. The seasoning was found to be microbiologically stable.

Example 2

A similar procedure to that described in Example 1 was followed except that the hydrolysis of the fermented wheat gluten koji was carried out over a period of 7 days instead of 24 hours.

The seasoning was found to have more body and a more rounded flavour profile than a conventional soya sauce. The seasoning was microbiologically stable as in Example 1 and had excellent colour stability in powder form.

Example 3

A similar procedure to that described in Example 1 was followed except that the hydrolysis of the fermented wheat gluten koji was carried out at 4°C for 14-20 days.

The seasoning was found to have more body and a more rounded flavour profile than a conventional soya sauce. The seasoning was microbiologically stable as in Example 1 and had excellent colour stability in powder form.

Claims

1. A process of treating a fermented protein koji prepared from a protein containing material and a carbohydrate, for the production of a seasoning, which comprises hydrolysing the fermented protein koji at a temperature of from 2° to 25°C and a pH of from 4.5 to 10 for a period of from 6 hours to 28 days.
2. A process according to claim 1 wherein the protein containing material is a plant protein material.
3. A process according to claim 1 wherein the protein containing material is cooked.
4. A process according to claim 1 wherein the protein containing material is used in particulate form for enabling the mould of *Aspergillus oryzae* and/or *Aspergillus sojae* to grow



on the surface of the particles and eventually penetrate into the particles.

5. A process according to claim 1 wherein the hydrolysis of the koji is carried out in the substantial or complete absence of salt.
6. A process according to claim 1 wherein the hydrolysis of the koji is carried out with constant agitation.
7. A process according to claim 1 wherein the fermented koji is hydrolysed by mixing with from 1 to 5 parts of water per part by weight of fermented koji.
8. A process according to claim 1 wherein the fermented koji is hydrolysed at a temperature from 2° to 20° C for a period of from 12 hours to 25 days.
9. A process according to claim 1 wherein, after the hydrolysis of the fermented koji, salt is added to the hydrolysed fermented koji to form a moromi and the moromi is fermented to give a seasoning.
10. A seasoning preparable by a process according to any of the preceding claims having a higher amino acid content than soya sauce prepared by conventional methods.

Patentansprüche

1. Verfahren zur Behandlung eines fermentierten Koji-Proteins, das aus einem proteinhaltigen Material und einem Kohlenhydrat hergestellt wird, zur Herstellung eines Gewürzes, welches ein Hydrolysieren des fermentierten Koji-Proteins bei einer Temperatur von 2° bis 25° C und einem pH-Wert von 4,5 bis 10 während eines Zeitraums von 6 Stunden bis 28 Tagen umfaßt.
2. Verfahren nach Anspruch 1, worin das proteinhaltige Material ein pflanzliches Proteinmaterial ist.
3. Verfahren nach Anspruch 1, worin das proteinhaltige Material gekocht wird.
4. Verfahren nach Anspruch 1, worin das proteinhaltige Material in Partikelform verwendet wird, um Aspergillus oryzae- und/oder Aspergillus sojae-Schimmelpilz an der Oberfläche der Teilchen wachsen und gegebenenfalls in die Teilchen eindringen zu lassen.

5. Verfahren nach Anspruch 1, worin die Hydrolyse des Kojis bei weitgehender oder vollständiger Abwesenheit von Salz durchgeführt wird.

6. Verfahren nach Anspruch 1, worin die Hydrolyse des Kojis unter gleichmäßigem Rühren erfolgt.

7. Verfahren nach Anspruch 1, worin das fermentierte Koji durch Vermischen mit 1 bis 5 Teilen Wasser je Gewichtsteil fermentiertem Koji hydrolysiert wird.

8. Verfahren nach Anspruch 1, worin das fermentierte Koji bei einer Temperatur von 2° bis 20° C während eines Zeitraums von 12 Stunden bis 25 Tagen hydrolysiert wird.

9. Verfahren nach Anspruch 1, worin nach der Hydrolyse des fermentierten Kojis Salz zu dem hydrolysierten fermentierten Koji zugesetzt wird, um ein Moromi auszubilden, und das Moromi fermentiert wird, um ein Gewürz zu ergeben.

10. Ein durch ein Verfahren nach einem der vorstehenden Ansprüche herstellbares Gewürz, das einen höheren Aminosäuregehalt aufweist als eine nach herkömmlichen Methoden hergestellte Sojasauce.

Revendications

1. Procédé de traitement d'un koji de protéines fermenté, préparé à partir d'une substance contenant des protéines et d'un glucide, pour la production d'un assaisonnement, qui comprend l'hydrolyse du koji de protéines fermenté à une température de 2 à 25° C et un pH de 4,5 à 10 pendant un temps de 6 heures à 28 jours.
2. Procédé suivant la revendication 1, dans lequel la substance contenant des protéines est une substance végétale contenant des protéines.
3. Procédé suivant la revendication 1, dans lequel la substance contenant des protéines a subi une cuisson.
4. Procédé suivant la revendication 1, dans lequel la substance contenant des protéines est utilisée sous forme de particules pour permettre à la moisissure de l'espèce Aspergillus oryzae et/ou Aspergillus sojae de se développer sur la surface des particules et de pénétrer finalement dans les particules.



5. Procédé suivant la revendication 1, dans lequel l'hydrolyse du koji est effectuée en l'absence notable ou totale de sel.
6. Procédé suivant la revendication 1, dans lequel l'hydrolyse du koji est effectuée avec agitation constante. 5
7. Procédé suivant la revendication 1, dans lequel le koji fermenté est hydrolysé par mélange avec une quantité de 1 à 5 parties d'eau par partie en poids de koji fermenté. 10
8. Procédé suivant la revendication 1, dans lequel le koji fermenté est hydrolysé à une température de 2 à 20 °C pendant un temps de 12 à 25 jours. 15
9. Procédé suivant la revendication 1, dans lequel, après l'hydrolyse du koji fermenté, du sel est ajouté au koji fermenté hydrolysé pour former un moromi et le moromi est fermenté pour obtenir un assaisonnement. 20
10. Assaisonnement pouvant être préparé par un procédé suivant l'une quelconque des revendications précédentes, ayant une teneur en amino-acides supérieure à celle de la sauce au soja préparée par des procédés classiques. 25

30

35

40

45

50

55

5