

APPLICATION FOR PATENT

**CORN PLANT AND SEED CORRESPONDING TO TRANSGENIC EVENT
MON89034 AND METHODS FOR DETECTION AND USE THEREOF**

**HEATHER ANDERSON; JENNIFER DOUGLAS; JEANNA GROAT; SCOTT
JOHNSON; REBECCA KELLY; JOHN KORTE; JAMES RICE**

CROSS REFERENCE TO RELATED APPLICATIONS

This application claims the benefit of priority to US Provisional Application Serial No. 60/808,834 filed May 26, 2006.

FIELD OF THE INVENTION

[0001] The present invention relates to transgenic corn event MON89034 and plant parts and seed thereof. The event exhibits resistance to insect infestation from insects in the order Lepidoptera. The invention also relates to methods for using plants and seeds comprising DNA that is diagnostic for the presence of the transgenic event when probed for the presence of nucleotide sequences that are unique to the transgenic event, and to methods for detecting the presence of said corn event in a biological sample by detecting specific nucleotide sequences that are unique to the transgenic event. The invention provides nucleotide sequences that are unique to the event.

BACKGROUND OF THE INVENTION

[0002] This invention relates to the Lepidopteran resistant transgenic variety of corn (*Zea mays*) plant referred to herein as event MON89034, and to unique DNA sequences present that, when detected in any sample or variety of corn, are diagnostic for the presence of the transgenic corn plant event MON89034 in that sample or variety, and also relates to the detection of the transgene/genomic insertion region in corn MON89034, and progeny plants and seeds derived therefrom.

[0003] The corn plant event MON89034 is particularly resistant to insects in the Lepidoptera family such as Fall armyworm (*Spodoptera frugiperda*), European corn borer (*Ostrinia nubilalis*), corn earworm (*Helicoverpa zea*), southwestern corn borer (*Diatraea grandiosella*), and black cutworm (*Agrotis ipsilon*) and the like, all of which are agronomically important insect pests.

[0004] Corn is an important crop and is a primary food source in many areas of the world. Biotechnology methods have been applied to corn for the purpose of improving agronomic traits and the quality of the product. One such agronomic trait is insect resistance, for example, genetically engineered resistance to lepidopteran and coleopteran species that arises in corn plants genetically engineered to contain one or more genes encoding insecticidal agents (see for example, US Patent 6,489,542 and US Patent 6,620,988). It is advantageous

to detect the presence of a particular transgenic event in a biological sample in order to determine whether one or more progeny of a sexual cross contains the transgenic material. For example, the detection of the event in a sample is important for licensing purposes, for establishing and maintaining purity standards, important for complying with regulatory agencies, for complying with food ingredient standards, for use in legal proceedings in establishing that one or more particular individuals or entities has been using the particular event without a license from the owner or licensee of any patents directed to the transgenic event, and for insuring compliance with various government regulations and/or laws.

[0005] In addition, methods that enable the detection of a particular plant would be helpful when complying with regulations requiring the pre-market approval and labeling of foods derived from the recombinant crop plants. Individuals or entities that are resistant to the presence of a transgenic event in a sample also desire reliable methods for detecting the presence of the transgene in a sample in order for them to be able to capitalize on their business, which takes advantage of an absence of transgenes in their products.

[0006] Despite these advantages, it is possible that insects may evolve resistance to plants expressing only one *B. thuringiensis* δ -endotoxin. Such resistance, should it become widespread, would clearly limit the commercial value of germplasm containing single Bt genes.

[0007] One possible way of increasing the effectiveness of insecticidal agents provided via transgenic plants and directed at controlling target insect pests and contemporaneously reducing the likelihood of emergence of insect pests resistant to such insecticidal agents would be to ensure that transgenic crops express high levels of these insecticidal agents, such as *Bacillus thuringiensis* delta-endotoxins (McGaughey and Whalon (1992), Science 258:1451-55; Roush (1994) Biocontrol. Sci. Technol., 4:501-516). In addition, having a repository of insecticidal genes that are effective against groups of insect pests and which manifest their effects through different modes of action can safeguard against development of resistance. The onset of resistance could be substantially delayed as a result of providing a crop that expresses two or more insecticidal activities exhibiting overlapping toxicity to the same insect species. One means for achieving such dual modes of action could be to provide a plant expressing a Bt gene toxic to a particular insect species along with a dsRNA that is provided for the purpose of targeting for suppression an essential gene of the same insect species targeted by the Bt toxin, the dsRNA eliciting an RNAi response upon ingestion by the

target pest, providing a means for redundancy in the event that the insect develops resistance either to the dsRNA or to the Bt gene. Alternatively, co-expression in a plant of two or more insecticidal toxins both toxic to the same insect species but each exhibiting a different mode of effectuating its killing activity, particularly when both are expressed at high levels, provides a means for effective resistance management. Examples of such insecticides useful in such combinations include but are not limited to Bt toxins, *Xenorhabdus* sp. or *Photorhabdus* sp. insecticidal proteins, deallergized and de-glycosylated patatin proteins and/or permuteins, plant lectins, and the like.

[0008] The expression of foreign genes in plants is known to be influenced by their chromosomal position, perhaps due to chromatin structure (e.g., heterochromatin) or the proximity of transcriptional regulation elements (e.g., enhancers) close to the integration site (Weising et al.(19880 Ann. Rev. Genet 22:421-477). For this reason, it is often necessary to screen a large number of events in order to identify an event characterized by optimal expression of an introduced gene of interest. Even then, with dozens or even hundreds of different transgenic events in hand, there is no certainty of success in identifying a single transgenic event that provides the optimum levels of expression of the at least two different toxins or insecticidal agents and lacks any undesirable agronomic deficiencies or phytotoxic effects, either as a result of the insertion into some essential or partially essential region of the plant genome, or as a result of toxic effects brought about by the levels of expression of the transgenes. For example, it has been observed in plants and in other organisms that there may be wide variation in the levels of expression of an introduced gene among events. There may also be differences in spatial or temporal patterns of expression, for example, differences in the relative expression of a transgene in various plant tissues, that may not correspond to the patterns expected from transcriptional regulatory elements present in the introduced gene construct. For this reason, it is common to produce several hundreds to several thousands different events and screen the events for a single event that has the desired transgene expression levels and patterns for commercial purposes. An event that has the desired levels or patterns of transgene expression is useful for introgressing the transgene into other genetic backgrounds by sexual outcrossing using conventional breeding methods. Progeny of such crosses maintain the transgene expression characteristics of the original transformant. This strategy is used to ensure reliable gene expression in a number of varieties that are suitably adapted to specific local growing conditions.

[0009] It is possible to detect the presence of a transgene by any well known nucleic acid detection method such as the polymerase chain reaction (PCR) or DNA hybridization using nucleic acid probes. These detection methods generally focus on frequently used genetic elements, such as promoters, terminators, marker genes, or even the coding sequence encoding the protein or dsRNA of interest expressed from the transgene(s), etc. As a result, such methods may not be useful for discriminating between different events, particularly those produced using the same DNA construct, unless the sequence of chromosomal DNA adjacent to the inserted DNA ("flanking DNA") is known. Depending on the method used for introducing the transgene(s) into a plant genome, aberrant or unusual effects can be observed, which often severely complicate the identification of the plant genome sequences flanking the transgenic DNA that was intended to be introduced into the plant. Often, rearrangements of the inserted DNA, rearrangements of the flanking genome DNA, or rearrangements of both the inserted DNA and the flanking genome DNA are prevalent, and complicate the analysis of the insertional event being evaluated. Therefore, it is advantageous to have a means for selecting, for identifying, and for insuring the purity and characteristics of a particular transgenic event in a sample, and the only way to accomplish this is to identify one or more unique sequences associated only with the desired transgenic event, and the presence of such sequences in a biological sample containing DNA of the plant species into which the transgenic DNA was inserted to give rise to the event are thus diagnostic for the event in such sample.

SUMMARY OF THE INVENTION

[0010] The present invention is related to the transgenic corn plant designated MON89034 and progeny that are indistinguishable from corn event MON89034 (to the extent that they also contain at least one allele that corresponds to the inserted transgenic DNA) thereof having seed deposited on March 28, 2006 with American Type Culture Collection (ATCC) with Accession No. PTA-7455. Another aspect of the invention is the progeny plants, or seeds, or regenerable parts of the plants and seeds of the corn event MON89034 that contain a polynucleotide selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, and SEQ ID NO:5. The invention also includes plant parts of the corn event MON89034 that include, but are not limited to pollen, ovule, flowers, shoots, roots, stalks, silks, tassels, ears, and leaves, so long as these parts contain at least the

polynucleotides as set forth above. Novel genetic compositions contained in the genome of MON89034 and products such from MON89034 such as meal, flour, oil, pulp, and biomass left over in a field of corn plants corresponding to MON89034 event are an aspect of this invention.

[0011] The invention provides an insect resistant corn plant that has all of the physiological and morphological characteristics of the corn event MON89034.

[0012] According to one aspect of the invention, compositions and methods are provided for detecting the presence of the transgene/genomic insertion region from a novel corn plant designated MON89034. DNA sequences are provided that comprise at least one junction sequence of MON89034 selected from the group consisting of SEQ ID NO:1 (located at positions 2051 to 2070 on SEQ ID NO: 5) and SEQ ID NO:2 (located at positions 11367 to 11388) and complements thereof; wherein a junction sequence spans the junction between heterologous DNA inserted into the genome and the DNA from the corn cell flanking the insertion site and is diagnostic for the event (Figure 1). A corn event MON89034 and seed comprising these DNA molecules is an aspect of this invention.

[0013] DNA sequences that comprise the novel transgene/genomic insertion region, SEQ ID NO:3 and SEQ ID NO:4 (Figure 2) from corn event MON89034 are aspects of this invention. The corn plant and seed comprising these molecules are also aspects of this invention.

[0014] According to another aspect of the invention, two DNA molecules are provided for use in a DNA detection method, wherein the first DNA molecule comprises at least 11 or more contiguous polynucleotides of any portion of the transgene region of the DNA molecule of SEQ ID NO:3 and a DNA molecule of similar length of any portion of a 5' flanking corn genomic DNA region of SEQ ID NO:3, where these DNA molecules when used together are useful as DNA primers in a DNA amplification method that produces an amplicon. The amplicon produced using these DNA primers in the DNA amplification method is diagnostic for corn event MON 89304 when the amplicon contains SEQ ID NO:1. Any amplicon produced by DNA primers homologous or complementary to any portion of SEQ ID NO:3 and any amplicon that comprises SEQ ID NO:1 is an aspect of the invention.

[0015] According to another aspect of the invention, two DNA molecules are provided for use in a DNA detection method, wherein the first DNA molecule comprises at least 11 or more contiguous polynucleotides of any portion of the transgene region of the DNA molecule

of SEQ ID NO:4 and a DNA molecule of similar length of any portion of a 3' flanking corn genomic DNA of SEQ ID NO:4, where these DNA molecules are useful as DNA primers in a DNA amplification method. The amplicon produced using these DNA primers in the DNA amplification method is diagnostic for corn event MON 89304 when the amplicon contains SEQ ID NO:2. Any amplicons produced by DNA primers homologous or complementary to any portion of SEQ ID NO:4 and any amplicon that comprises SEQ ID NO:2 is an aspect of the invention.

[0016] According to another aspect of the invention, methods of detecting the presence of DNA corresponding to the corn event MON89034 in a sample are provided. Such methods comprise: (a) contacting the sample comprising DNA with a primer set that, when used in a nucleic acid amplification reaction with genomic DNA from corn event MON89034, produces an amplicon that is diagnostic for corn event MON89034; (b) performing a nucleic acid amplification reaction, thereby producing the amplicon; and (c) detecting the amplicon wherein said amplicon comprises SEQ ID NO:1 or SEQ ID NO:2.

[0017] A corn plant, or seed, or product derived from the plant or seed MON89034 wherein the genomic DNA comprising comprises a DNA molecule consisting essentially of SEQ ID NO:5 and complements thereof. A corn plant, or seed, or product derived from the plant or seed MON89034, in which the genomic DNA when isolated from the corn plant, or seed, or product comprises a DNA molecule incorporating nucleotides 2061 to 11377 of SEQ ID NO:5 and complements thereof.

[0018] A corn plant, or seed, or product derived from the plant or seed MON89034, in which the genomic DNA when isolated from the corn plant, or seed, or product produces an amplicon in a DNA amplification method, wherein DNA primer molecules SEQ ID NO:6 and SEQ ID NO:7 is used in the DNA amplification method.

[0019] According to another aspect of the invention, methods of detecting the presence of a DNA corresponding to the MON89034 event in a sample, such methods comprising: (a) contacting the sample comprising DNA with a probe that hybridizes under stringent hybridization conditions with genomic DNA from corn event MON89034 and does not hybridize under the stringent hybridization conditions with a control corn plant; (b) subjecting the sample and probe to stringent hybridization conditions; and (c) detecting hybridization of the probe to the corn event MON89034 DNA wherein said probe comprises SEQ ID NO:1 and SEQ ID NO:2.

[0020] Another aspect of the invention is a method of determining the zygosity of the progeny of corn event MON89034 comprising: (a) contacting the sample comprising corn DNA with a primer set comprising SQ2842 (SEQ ID NO:6), SQ2843(SEQ ID NO:7), SQ6523(SEQ ID NO:10), SQ6524(SEQ ID NO:11), PB880 (SEQ ID NO:14) and PB2931 (SEQ ID NO:15) that when used in a nucleic-acid amplification reaction with genomic DNA from corn event MON89034, produces a first amplicon that is diagnostic for corn event MON89034 and (b) performing a nucleic acid amplification reaction, thereby producing the first amplicon; and (c) detecting the first amplicon; and (d) contacting the sample comprising corn DNA with said primer set, that when used in a nucleic-acid amplification reaction with genomic DNA from corn plants produces a second amplicon comprising the native corn genomic DNA homologous to the corn genomic region of a transgene insertion identified as corn event MON89034; and (e) performing a nucleic acid amplification reaction, thereby producing the second amplicon and (f) detecting the second amplicon; and (g) comparing the first and second amplicons in a sample, wherein the presence of both amplicons indicates the sample is heterozygous for the transgene insertion.

[0021] One aspect of the invention is providing in the diet of a lepidopteran pest an insecticidally effective amount of corn event MON89034.

[0022] Another aspect of the present invention is providing a composition or biological sample in the form of a commodity or foodstuff that is derived from corn event MON89034, the commodity or foodstuff comprising ears of corn, shucked corn, corn silk, corn pollen, cracked corn, corn meal, crushed corn, corn flour, corn oil, corn starch, corn steep liquor, corn malt, corn sugar, corn syrup, margarine produced from corn oil, unsaturated corn oil, saturated corn oil, corn flakes, pop corn, ethanol and/or liquor produced from corn or corn products comprising DNA diagnostic for corn event MON89034, distillers dry goods solids (DDGS) produced from fermentation of such corn event, and animal feeds comprising such DDGS and/or corn, whether or not whole, cracked, or crushed, processed foodstuffs, a cosmetic, and a bulking agent in which there is found a detectable amount of a polynucleotide that is diagnostic for the presence of the transgenic corn event MON89034 in the biological sample. An alternative means for providing corn as a foodstuff is to provide corn in various forms of grain for feeding, such as whole corn, cracked corn, crushed corn, and various forms of the foregoing in a blend with milo, suet, millet, sunflower, oats, wheat, rice, beans, and the like. Detectable amounts of a nucleotide sequence in such commodity or foodstuff, such as is

set forth at SEQ ID NO:1 or SEQ ID NO:2, or the complements thereof, is diagnostic for the presence of such transgenic event MON89034 DNA in the sample, and therefore, the presence of the transgenic event cells as having originated the DNA in the sample.

[0023] The foregoing and other aspects of the invention will become more apparent from the following detailed description.

DRAWINGS

[0024] Figure 1. Organization of the transgene insert present within the genome of transgenic corn event MON89034. The central open or white bar represents the inserted DNA. Below the white bar is a diagram that represents the various elements within the inserted DNA. The ends of the inserted DNA have been arbitrarily designated as 5' (to the left side of the Figure) and 3' (to the right side of the Figure). The Right Border and Left Border sequences or segments are labeled beneath each end of the diagram illustrating the various elements within the inserted DNA. The labeled elements in the expression cassettes within the inserted DNA are, in consecutive order starting from the Right Border: e35S promoter, wheat CAB untranslated leader, rice actin intron, coding sequence for Cry1A.105, wheat HSP17 3' termination and polyadenylation sequence, FMV promoter, hsp70 intron, rubisco small subunit chloroplast targeting peptide coding sequence, Cry2Ab coding sequence, nos 3' termination and polyadenylation signal sequence, and then the Left Border. The vertically hatched bars at either end of the central open or white bar correspond to the arbitrarily labeled 5' and 3' corn genome flanking sequences. The longest black line above the hatched and open or white bar represents SEQ ID NO:5 (the full length sequence represented by the figure depicting the 5' flanking sequence, the inserted DNA sequence, and the 3' flanking sequence). The shorter black lines above and below the black line labeled as SEQ ID NO:5 represent the approximate positions within SEQ ID NO:5 in which each of the specifically labeled sequences can be found (i.e., SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, and SEQ ID NO:4). SEQ ID NO:1 and SEQ ID NO:2, and any sequence derived from corn event MON89034 containing SEQ ID NO:1 and/or SEQ ID NO:2, are diagnostic for corn event MON89034 DNA in a biological sample.

DETAILED DESCRIPTION

[0025] The following definitions and methods are provided to better define the present invention and to guide those of ordinary skill in the art in the practice of the present invention. Unless otherwise noted, terms are to be understood according to conventional usage by those of ordinary skill in the relevant art. Definitions of common terms in molecular biology may also be found in Rieger et al., *Glossary of Genetics: Classical and Molecular*, 5th edition, Springer-Verlag: New York, 1991; and Lewin, *Genes V*, Oxford University Press: New York, 1994.

[0026] As used herein, the term “corn” means *Zea mays* or maize and includes all plant varieties that can be bred with corn, including wild maize species.

[0027] As used herein, the term “comprising” means “including but not limited to”.

[0028] A transgenic “event” is produced by transformation of plant cells with heterologous DNA, i.e., a nucleic acid construct that includes a transgene of interest, regeneration of a population of plants resulting from the insertion of the transgene into the genome of the plant, and selection of a particular plant characterized by insertion into a particular genome location. The term “event” refers to the original transformant and progeny of the transformant that include the heterologous DNA. The term “event” also refers to progeny produced by a sexual outcross between the transformant and another variety that include the heterologous DNA. Even after repeated back-crossing to a recurrent parent, the inserted DNA and flanking DNA from the transformed parent is present in the progeny of the cross at the same chromosomal location. The term “event” also refers to DNA from the original transformant comprising the inserted DNA and flanking genomic sequence immediately adjacent to the inserted DNA that would be expected to be transferred to a progeny that receives inserted DNA including the transgene of interest as the result of a sexual cross of one parental line that includes the inserted DNA (e.g., the original transformant and progeny resulting from selfing) and a parental line that does not contain the inserted DNA. The present invention relates to the event MON89034 DNA, plant cells, tissues, seeds and processed products derived from MON89034.

[0029] It is also to be understood that two different transgenic plants can also be mated to produce offspring that contain two independently segregating added, exogenous genes. Selfing of appropriate progeny can produce plants that are homozygous for both added, exogenous genes. Back-crossing to a parental plant and out-crossing with a non-transgenic plant are also contemplated, as is vegetative propagation. Descriptions of other breeding

methods that are commonly used for different traits and crops can be found in one of several references, e.g., Fehr, in *Breeding Methods for Cultivar Development*, Wilcox J. ed., American Society of Agronomy, Madison WI (1987).

[0030] A "probe" is an isolated nucleic acid to which is attached a conventional detectable label or reporter molecule, e.g., a radioactive isotope, ligand, chemiluminescent agent, or enzyme. Such a probe is complementary to a strand of a target nucleic acid, in the case of the present invention, to a strand of genomic DNA from corn event MON89034 whether from a corn plant or from a sample that includes DNA from the event. Probes according to the present invention include not only deoxyribonucleic or ribonucleic acids but also polyamides and other probe materials that bind specifically to a target DNA sequence and can be used to detect the presence of that target DNA sequence.

[0031] "Primers" are isolated nucleic acids that are annealed to a complementary target DNA strand by nucleic acid hybridization to form a hybrid between the primer and the target DNA strand, then extended along the target DNA strand by a polymerase, e.g., a DNA polymerase. Primer pairs of the present invention refer to their use for amplification of a target nucleic acid sequence, e.g., by the polymerase chain reaction (PCR) or other conventional nucleic-acid amplification methods.

[0032] Probes and primers are generally 11 nucleotides or more in length, preferably 18 nucleotides or more, more preferably 24 nucleotides or more, and most preferably 30 nucleotides or more. Such probes and primers hybridize specifically to a target sequence under high stringency hybridization conditions. Preferably, probes and primers according to the present invention have complete sequence similarity with the target sequence, although probes differing from the target sequence and that retain the ability to hybridize to target sequences may be designed by conventional methods.

[0033] Methods for preparing and using probes and primers are described, for example, in *Molecular Cloning: A Laboratory Manual*, 2nd ed., vol. 1-3, ed. Sambrook et al., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989 (hereinafter, "Sambrook et al., 1989"); *Current Protocols in Molecular Biology*, ed. Ausubel et al., Greene Publishing and Wiley-Interscience, New York, 1992 (with periodic updates) (hereinafter, "Ausubel et al., 1992"); and Innis et al., *PCR Protocols: A Guide to Methods and Applications*, Academic Press: San Diego, 1990. PCR-primer pairs can be derived from a known sequence, for

example, by using computer programs intended for that purpose such as Primer (Version 0.5, © 1991, Whitehead Institute for Biomedical Research, Cambridge, MA).

[0034] Primers and probes based on the flanking DNA and insert sequences disclosed herein can be used to confirm (and, if necessary, to correct) the disclosed sequences by conventional methods, e.g., by re-cloning and sequencing such sequences.

[0035] The nucleic acid probes and primers of the present invention hybridize under stringent conditions to a target DNA sequence. Any conventional nucleic acid hybridization or amplification method can be used to identify the presence of DNA from a transgenic event in a sample. Nucleic acid molecules or fragments thereof are capable of specifically hybridizing to other nucleic acid molecules under certain circumstances. As used herein, two nucleic acid molecules are said to be capable of specifically hybridizing to one another if the two molecules are capable of forming an anti-parallel, double-stranded nucleic acid structure. A nucleic acid molecule is said to be the "complement" of another nucleic acid molecule if they exhibit complete complementarity. As used herein, molecules are said to exhibit "complete complementarity" when every nucleotide of one of the molecules is complementary to a nucleotide of the other. Two molecules are said to be "minimally complementary" if they can hybridize to one another with sufficient stability to permit them to remain annealed to one another under at least conventional "low-stringency" conditions. Similarly, the molecules are said to be "complementary" if they can hybridize to one another with sufficient stability to permit them to remain annealed to one another under conventional "high-stringency" conditions. Conventional stringency conditions are described by Sambrook et al., 1989, and by Haymes et al., In: Nucleic Acid Hybridization, A Practical Approach, IRL Press, Washington, DC (1985), Departures from complete complementarity are therefore permissible, as long as such departures do not completely preclude the capacity of the molecules to form a double-stranded structure. In order for a nucleic acid molecule to serve as a primer or probe it need only be sufficiently complementary in sequence to be able to form a stable double-stranded structure under the particular solvent and salt concentrations employed.

[0036] As used herein, a substantially homologous sequence is a nucleic acid sequence that will specifically hybridize to the complement of the nucleic acid sequence to which it is being compared under high stringency conditions. Appropriate stringency conditions which promote DNA hybridization, for example, 6.0 x sodium chloride/sodium citrate (SSC) at

about 45°C, followed by a wash of 2.0 x SSC at 50°C, are known to those skilled in the art or can be found in Current Protocols in Molecular Biology, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. For example, the salt concentration in the wash step can be selected from a low stringency of about 2.0 x SSC at 50°C to a high stringency of about 0.2 x SSC at 50°C. In addition, the temperature in the wash step can be increased from low stringency conditions at room temperature, about 22°C, to high stringency conditions at about 65°C. Both temperature and salt may be varied, or either the temperature or the salt concentration may be held constant while the other variable is changed. In a preferred embodiment, a nucleic acid of the present invention will specifically hybridize to one or more of the nucleic acid molecules set forth in SEQ ID NO: 1 and 2 or complements thereof or fragments of either under moderately stringent conditions, for example at about 2.0 x SSC and about 65°C. In a particularly preferred embodiment, a nucleic acid of the present invention will specifically hybridize to one or more of the nucleic acid molecules set forth in SEQ ID NO:1 and SEQ ID NO:2 or complements or fragments of either under high stringency conditions. In one aspect of the present invention, a preferred marker nucleic acid molecule of the present invention has the nucleic acid sequence set forth in SEQ ID NO:1 and SEQ ID NO:2 or complements thereof or fragments of either. In another aspect of the present invention, a preferred marker nucleic acid molecule of the present invention shares between 80% and 100% or 90% and 100% sequence identity with the nucleic acid sequence set forth in SEQ ID NO:1 and SEQ ID NO:2 or complement thereof or fragments of either. In a further aspect of the present invention, a preferred marker nucleic acid molecule of the present invention shares between 95% and 100% sequence identity with the sequence set forth in SEQ ID NO:1 and SEQ ID NO:2 or complement thereof or fragments of either. SEQ ID NO:1 and SEQ ID NO:2 may be used as markers in plant breeding methods to identify the progeny of genetic crosses similar to the methods described for simple sequence repeat DNA marker analysis, in "DNA markers: Protocols, applications, and overviews: (1997) 173-185, Cregan, et al., eds., Wiley-Liss NY; all of which is herein incorporated by reference in its' entirety. The hybridization of the probe to the target DNA molecule can be detected by any number of methods known to those skilled in the art, these can include, but are not limited to, fluorescent tags, radioactive tags, antibody based tags, and chemiluminescent tags.

[0037] Regarding the amplification of a target nucleic acid sequence (e.g., by PCR) using a particular amplification primer pair, "stringent conditions" are conditions that permit the

primer pair to hybridize only to the target nucleic-acid sequence to which a primer having the corresponding wild-type sequence (or its complement) would bind and preferably to produce a unique amplification product, the amplicon, in a DNA thermal amplification reaction.

[0038] The term "specific for (a target sequence)" indicates that a probe or primer hybridizes under stringent hybridization conditions only to the target sequence in a sample comprising the target sequence.

[0039] As used herein, "amplified DNA" or "amplicon" refers to the product of nucleic-acid amplification of a target nucleic acid sequence that is part of a nucleic acid template. For example, to determine whether the corn plant resulting from a sexual cross contains transgenic event genomic DNA from the corn plant of the present invention, DNA extracted from a corn plant tissue sample may be subjected to nucleic acid amplification method using a primer pair that includes a primer derived from flanking sequence in the genome of the plant adjacent to the insertion site of inserted heterologous DNA, and a second primer derived from the inserted heterologous DNA to produce an amplicon that is diagnostic for the presence of the event DNA. The amplicon is of a length and has a sequence that is also diagnostic for the event. The amplicon may range in length from the combined length of the primer pairs plus one nucleotide base pair, preferably plus about fifty nucleotide base pairs, more preferably plus about two hundred-fifty nucleotide base pairs, and even more preferably plus about four hundred-fifty nucleotide base pairs. Alternatively, a primer pair can be derived from flanking sequence on both sides of the inserted DNA so as to produce an amplicon that includes the entire insert nucleotide sequence. A member of a primer pair derived from the plant genomic sequence may be located a distance from the inserted DNA molecule, this distance can range from one nucleotide base pair up to about twenty thousand nucleotide base pairs. The use of the term "amplicon" specifically excludes primer dimers that may be formed in the DNA thermal amplification reaction.

[0040] Nucleic-acid amplification can be accomplished by any of the various nucleic-acid amplification methods known in the art, including the polymerase chain reaction (PCR). A variety of amplification methods are known in the art and are described, inter alia, in U.S. Patent Nos. 4,683,195 and 4,683,202 and in PCR Protocols: A Guide to Methods and Applications, ed. Innis et al., Academic Press, San Diego, 1990. PCR amplification methods have been developed to amplify up to 22 kb of genomic DNA and up to 42 kb of bacteriophage DNA (Cheng et al., Proc. Natl. Acad. Sci. USA 91:5695-5699, 1994). These

methods as well as other methods known in the art of DNA amplification may be used in the practice of the present invention. The sequence of the heterologous DNA insert or flanking sequence from corn event MON89034 with seed samples deposited as ATCC numbers can be verified (and corrected if necessary) by amplifying such sequences from the event using primers derived from the sequences provided herein followed by standard DNA sequencing of the PCR amplicon or of the cloned DNA.

[0041] The amplicon produced by these methods may be detected by a plurality of techniques. One such method is Genetic Bit Analysis (Nikiforov, et al. *Nucleic Acid Res.* 22:4167-4175, 1994) where an DNA oligonucleotide is designed which overlaps both the adjacent flanking genomic DNA sequence and the inserted DNA sequence. The oligonucleotide is immobilized in wells of a microwell plate. Following PCR of the region of interest (using one primer in the inserted sequence and one in the adjacent flanking genomic sequence), a single-stranded PCR product can be hybridized to the immobilized oligonucleotide and serve as a template for a single base extension reaction using a DNA polymerase and labelled ddNTPs specific for the expected next base. Readout may be fluorescent or ELISA-based. A signal indicates presence of the insert/flanking sequence due to successful amplification, hybridization, and single base extension.

[0042] Another method is the pyrosequencing technique as described by Winge (*Innov. Pharma. Tech.* 00:18-24, 2000). In this method an oligonucleotide is designed that overlaps the adjacent genomic DNA and insert DNA junction. The oligonucleotide is hybridized to single-stranded PCR product from the region of interest (one primer in the inserted sequence and one in the flanking genomic sequence) and incubated in the presence of a DNA polymerase, ATP, sulfurylase, luciferase, apyrase, adenosine 5' phosphosulfate and luciferin. dNTP's are added individually and the incorporation results in a light signal which is measured. A light signal indicates the presence of the transgene insert/flanking sequence due to successful amplification, hybridization, and single or multi-base extension.

[0043] Fluorescence polarization as described by Chen, et al., (*Genome Res.* 9:492-498, 1999) is a method that can be used to detect the amplicon of the present invention. Using this method an oligonucleotide is designed which overlaps the genomic flanking and inserted DNA junction. The oligonucleotide is hybridized to single-stranded PCR product from the region of interest (one primer in the inserted DNA and one in the flanking genomic DNA sequence) and incubated in the presence of a DNA polymerase and a fluorescent-labeled

ddNTP. Single base extension results in incorporation of the ddNTP. Incorporation can be measured as a change in polarization using a fluorimeter. A change in polarization indicates the presence of the transgene insert/flanking sequence due to successful amplification, hybridization, and single base extension.

[0044] Taqman® (PE Applied Biosystems, Foster City, CA) is described as a method of detecting and quantifying the presence of a DNA sequence and is fully understood in the instructions provided by the manufacturer. Briefly, a FRET oligonucleotide probe is designed which overlaps the genomic flanking and insert DNA junction. The FRET probe and PCR primers (one primer in the insert DNA sequence and one in the flanking genomic sequence) are cycled in the presence of a thermostable polymerase and dNTPs. Hybridization of the FRET probe results in cleavage and release of the fluorescent moiety away from the quenching moiety on the FRET probe. A fluorescent signal indicates the presence of the flanking/transgene insert sequence due to successful amplification and hybridization.

[0045] Molecular Beacons have been described for use in sequence detection as described in Tyangi, et al. (Nature Biotech.14:303-308, 1996) Briefly, a FRET oligonucleotide probe is designed that overlaps the flanking genomic and insert DNA junction. The unique structure of the FRET probe results in it containing secondary structure that keeps the fluorescent and quenching moieties in close proximity. The FRET probe and PCR primers (one primer in the insert DNA sequence and one in the flanking genomic sequence) are cycled in the presence of a thermostable polymerase and dNTPs. Following successful PCR amplification, hybridization of the FRET probe to the target sequence results in the removal of the probe secondary structure and spatial separation of the fluorescent and quenching moieties that results in the production of a fluorescent signal. The fluorescent signal indicates the presence of the flanking/transgene insert sequence due to successful amplification and hybridization.

[0046] Other described methods, such as, microfluidics (US Patent pub. 2006068398, US Patent No. 6,544,734) provide methods and devices to separate and amplify DNA samples. Optical dyes used to detect and quantitate specific DNA molecules (WO/05017181). Nanotube devices (WO/06024023) that comprise an electronic sensor for the detection of DNA molecules or nanobeads that bind specific DNA molecules and can then be detected.

[0047] DNA detection kits are provided using the compositions disclosed herein. The kits are useful for the identification of corn event MON89034 DNA in a sample and can be applied at least to methods for breeding corn plants containing the appropriate event DNA.

The kits contain DNA primers and/or probes that are homologous or complementary to segments selected from the sequences as set forth at SEQ ID NO:1-7, or DNA primers or probes homologous or complementary to DNA contained in the transgene genetic elements of DNA as set forth in the Sequence Listing. These DNA sequences can be used in DNA amplification reactions or as probes in a DNA hybridization method for detecting the presence of polynucleotides diagnostic for the presence of the target DNA in a sample. The production of a predefined amplicon in a thermal amplification reaction is diagnostic for the presence of DNA corresponding to PTO-7455 genome DNA in the sample. If hybridization is selected, detecting hybridization of the probe to the biological sample is diagnostic for the presence of the MON89034 transgenic event DNA in the sample. Typically, the sample is corn, or corn products or by-products of the use of corn.

[0048] The present invention provides a transgenic corn plant designated as corn event MON89034, progeny of the plant, and cells of the plant, as well as seed produced from the plant. Representative seed for growing the plant, for producing progeny, for obtaining cells, or for producing a crop of said seed comprising the transgenic corn event have been deposited on March 28, 2006 with the American Type Culture Collection (ATCC) and have the accession number PTA-7455.

[0049] The plant and cells and products produced from these embodiments and the like contain DNA that is diagnostic for the presence of DNA derived from any cell derived from the transgenic corn event MON89034 in any biological sample. This is because these two novel sequences are contained within the cells of the transgenic corn event MON89034. The diagnostic DNA comprises a nucleotide sequence that is selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, and SEQ ID NO:5. The relationship of these sequences is described more particularly herein and in the legend to Figure 1 and with reference to Figure 1.

[0050] Corn plants grown from seed that are homozygous for the DNA diagnostic for transgenic corn event MON89034 are also within the scope of the present invention. Corn plants grown from seed that are heterozygous for the DNA diagnostic for transgenic corn event MON89034 are also within the scope of the present invention so long as these seed also comprise the diagnostic DNA sequences. Cells, seed, and tissue produced from such plants comprising the diagnostic DNA are also within the scope of the present invention.

[0051] Corn plants and corn plant cells and the like comprising DNA diagnostic for the transgenic corn event MON89034 exhibit resistance to lepidopteran insect infestation. These cells and plants contain DNA encoding the insecticidal protein (insecticide, toxic agent) Cry2Ab and DNA having nucleotide sequences of SEQ ID NO:1 and SEQ ID NO:2 that form a part of the genome of the cells of the plant. These plants and plant cells also contain a DNA encoding insecticidal protein (insecticide, toxic agent) Cry1A.105. These proteins can be referred to as a first and a second insecticidal protein, respectively or in the inverse. Expression of these proteins is achieved from the regulatory components/genetic elements that are embedded within the expression cassettes that provide for the expression of each of the DNA sequences encoding these toxins and are fully described herein and in the legend to Figure 1 and with reference to Figure 1 and the sequence as set forth at SEQ ID NO:5. Corn plants and corn plant cells comprising these sequences are effective for protecting plants from lepidopteran pest infestation, whether heterozygous or homozygous for the alleles in which these coding sequences are present.

[0052] The present invention also provides amplicons that can be produced from the sequences described herein that are diagnostic for the presence in a biological sample of DNA derived from DNA of the transgenic corn event MON89034. An amplicon diagnostic for the presence of transgenic corn event MON89034 DNA in a biological sample contains at least one polynucleotide segment consisting of the nucleotide sequence as set forth at SEQ ID NO:1 or SEQ ID NO:2. These amplicons can be produced using primer sequences as described herein below from any biological sample that contains at least about 0.5 femtomole or about 0.5 pico-gram of DNA derived from the transgenic corn event MON89034. Such biological sample sources of DNA corresponding to the transgenic corn event MON89034 can be corn meal, corn oil, corn cake, corn seed, corn germ, corn starch, and corn flour and the like derived from that transgenic event.

[0053] The invention also provides isolated polynucleotide molecules that exhibit contiguous nucleotide sequences such as those set forth in SEQ ID NO:5. These contiguous nucleotide sequences comprise: (1) from about 11 to about 12000 nucleotides and any length in-between, and further comprise the contiguous nucleotides as set forth at nucleotide position 1-11 or 9-20 in SEQ ID NO:1 and 1-11 or 9-20 as set forth in SEQ ID NO:2; (2) any contiguous nucleotide sequence as set forth in SEQ ID NO:3 from about 11 to about 2000 nucleotides and any length in-between, and further comprise the contiguous nucleotides as set

forth at nucleotide position 1-11 and 9-20 as set forth at SEQ ID NO:1; any contiguous nucleotide sequence as set forth in SEQ ID NO:4 from about 11 to about 914 nucleotides and any length in-between, and further comprise the contiguous nucleotides as set forth at nucleotide position 1-11 and 9-20 as set forth at SEQ ID NO:2. These isolated polynucleotide molecules are useful in DNA amplification methods to produce one or more amplicons from a biological sample that contains corn DNA. The detection of such an amplicon is diagnostic for the presence of transgenic corn event MON89034 DNA in the sample. The isolated polynucleotide molecules are also useful in various nucleotide detection methods for detecting the presence of DNA derived from transgenic corn event MON89034 in a biological sample. In particular, polynucleotide probes comprising at least about 11 contiguous nucleotide as set forth in SEQ ID NO:1 or SEQ ID NO:2 are useful as probes in such methods for detecting the transgenic event MON89034 DNA in a sample. The complementary sequences of these isolated polynucleotide molecules are also useful in the same detection and/or amplification methods.

[0054] Kits for use in detecting the presence of DNA derived from the transgenic corn event MON89034 in a biological sample are also provide by the present invention. A kit uses a probe polynucleotide molecule, the probe molecule containing at least from about 11 to about 12000 contiguous nucleotides exhibiting substantial homology, or exhibiting substantial complementarity to a nucleotide segment comprising a sequence as set forth at SEQ ID NO:5, would be useful for detecting the presence of MON89034 DNA in a sample. The probe molecule should contain at least one of the sequences as set forth at SEQ ID NO:1 and SEQ ID NO:2. The sequences set forth at SEQ ID NO:1 and SEQ ID NO:2 can also be referred to as junction sequences, i.e., the sequences at either end of the transgenic DNA inserted into the corn plant to give rise to the transgenic corn event MON89034. These sequences, arbitrarily referred to as 5' and 3' ends respectively, contain part of the inserted DNA sequence and part of the flanking corn genome sequence. For example, SEQ ID NO:1 represents at its 5' half the 3' end terminus of the corn genome sequence flanking the 5' end of the inserted DNA, the 5' end of the inserted DNA being represented by the 3' end half of the sequence as set forth at SEQ ID NO:1. SEQ ID NO:2 represents at its 5' half the 3' end terminus of the inserted DNA, and at its 3' end half the 5' end of the corn genome sequence flanking the 3' end of the inserted DNA. In the naturally occurring corn genome at the position of the inserted sequence set forth in SEQ ID NO:5, the flanking sequence at the 5'

end of the inserted DNA and the flanking sequence at the 3' end of the inserted DNA are joined, and a first primer molecule that hybridizes to the sequence complementary to the sequence set forth at SEQ ID NO:3 (other than the 3' end 21 nucleotides of SEQ ID NO:3) and a second primer molecule that hybridizes to the sequence as set forth at SEQ ID NO:4 (other than the 5' end 20 nucleotides of SEQ ID NO:4) will produce an amplicon in a thermal amplification reaction with template that is DNA other than MON89034 DNA that is diagnostic for the absence of the inserted DNA in MON89034, and the same primers will produce an amplicon that is slightly larger than 12000 nucleotides (depending on the position of the primers in the flanking sequences set forth in SEQ ID NO:3 and SEQ ID NO:4) when using MON89034 DNA as a template. Other embodiments are also provided.

[0055] A kit for detecting the junction sequence SEQ ID NO:1 or SEQ ID NO:2 of corn event MON89034 in a biological sample is provided. The kit contains a polynucleotide probe which is or is fully complementary to a sequence selected from the group consisting of SEQ ID NO:1 or SEQ ID NO:2 or complements thereof, and also contains a pair of primers for use in a nucleic-acid amplification reaction. The pair of primers can be referred to as a first primer consisting of at least about 15 to about 50 contiguous nucleotides from the corn genome portion of SEQ ID NO:3 and a second primer consisting of at least about 15 to about 50 contiguous nucleotides complementary to the heterologous insert DNA portion of SEQ ID NO:5. The first primer of the pair of polynucleotide primers hybridizes specifically to the reverse complement sequence corresponding to that set forth in SEQ ID NO:3 from about nucleotide position 1 through about position 2050 and the second primer of said pair of polynucleotide primers hybridizes specifically to the sequence as set forth in SEQ ID NO:5 from about nucleotide position 2060 through about nucleotide position 12,208, and are extended toward each other to form an amplicon which comprises SEQ ID NO:1, said amplicon being diagnostic for the presence of MON89034 event DNA in the sample. A different pair of primers can be referred to as a first primer consisting of at least about 15 to about 50 contiguous nucleotides complementary to the corn genome portion of SEQ ID NO:4 and a second primer consisting of at least about 15 to about 50 contiguous nucleotides from the heterologous insert DNA portion of SEQ ID NO:5. The first primer of the pair of polynucleotide primers hybridizes specifically to the reverse complement sequence corresponding to that set forth in SEQ ID NO:5 from about nucleotide position 1 through about position 11,370 and the second primer of the pair of polynucleotide primers hybridizes

specifically to the sequence as set forth SEQ ID NO:4 from about nucleotide position 1 through about nucleotide position 914, and are extended toward each other to form an amplicon which comprises SEQ ID NO:2, said amplicon being diagnostic for the presence of MON89034 event DNA in said sample.

[0056] These primer pairs are useful in producing amplicons that comprise either SEQ ID NO:1 or SEQ ID NO:2, as the case may be, and are thus diagnostic for the presence of MON89034 DNA in a biological sample. These amplicons enable the detection of the presence of a junction sequence diagnostic corn event MON89034 in a biological sample.

[0057] A method for producing and detecting an amplicon that is diagnostic for a transgenic corn event MON89034 DNA in a biological sample comprising corn DNA is also provided. The method comprises contacting the biological sample together with two or more primers in a nucleic acid amplification reaction, performing a nucleic acid amplification reaction, then detecting the amplicon. The presence of the amplicon is diagnostic for said event DNA in the sample so long as the amplicon contains at least one of the contiguous sequences as set forth at SEQ ID NO:1 and SEQ ID NO:2, from about nucleotide position 1-11 or 9-20, or the complementary sequences corresponding to these positions.

[0058] The nucleotide sequences diagnostic for the presence of transgenic corn event MON89034 in a biological sample can also be detected using other methods. For example, contacting a biological sample suspected of containing MON89034 DNA with a probe that hybridizes under stringent hybridization conditions with one or more of the nucleotide sequences as set forth at SEQ ID NO:1 or SEQ ID NO:2, subjecting the sample and probe to stringent hybridization conditions; and then detecting hybridization of the probe to the nucleotide sequence. Detecting hybridization is diagnostic for the presence of the MON89034 DNA in the sample.

[0059] Primer polynucleotides for use in producing, in a thermal amplification reaction, an amplicon that is diagnostic for the presence of corn event MON89034 DNA in a biological sample are also provided by the present invention. Typically, the primers are provided in pairs, the members of the primer pair being referred to for convenience as a first primer and a second primer. A first primer can consist of at least about 15 contiguous nucleotides from the corn genome portion as set forth in SEQ ID NO:3 and a second primer can consist of at least about 15 contiguous nucleotides complementary to the heterologous insert DNA portion as set forth in SEQ ID NO:5. These two primers would produce an amplicon in a thermal

amplification reaction with template DNA obtained from corn event MON89034 DNA that contains a polynucleotide sequence as set forth at SEQ ID NO:1. Alternatively, a first primer can consist of at least about 15 contiguous nucleotides from the corn genome portion of SEQ ID NO:4, and a second primer can consist of at least about 15 contiguous nucleotides complementary to the heterologous insert DNA portion of SEQ ID NO:5. These two primers would produce an amplicon in a thermal amplification reaction with template DNA obtained from corn event MON89034 DNA that contains a polynucleotide sequence as set forth at SEQ ID NO:2.

[0060] An alternative method for detecting a junction sequence of corn event MON89034 in a biological sample comprising corn DNA, such as SEQ ID NO:1 or SEQ ID NO:2, consists of contacting the sample with a polynucleotide probe that hybridizes under stringent hybridization conditions with one of the junction sequences, subjecting the sample and probe to stringent hybridization conditions; and detecting hybridization of the probe to the junction sequence. Detecting the binding/hybridization of the probe to the junction sequence is indicative of the presence of the MON89034 DNA in the biological sample. A stably transformed maize plant, the DNA of which produces a DNA amplicon comprising SEQ ID NO:1 or SEQ ID NO:2 when subjected to the method set forth herein, is within the scope of the present invention. Exemplary primer sequences, in particular, a pair of primer sequences, are set forth herein in the examples and at SEQ ID NO:6 and SEQ ID NO:7.

[0061] An alternative method of detecting the presence of corn event MON89034 DNA in a biological sample can consist of the steps of contacting the sample with a probe that hybridizes under stringent hybridization conditions with MON89034 DNA and does not hybridize under stringent hybridization conditions with corn plant genomic DNA that is not MON89034 DNA, subjecting the sample and probe to stringent hybridization conditions, and then detecting hybridization of the probe to MON89034 DNA. A probe consistent with this embodiment is or is complementary to a sequence selected from the group consisting of SEQ ID NO:1 and SEQ ID NO:2. Detecting hybridization of the probe to the sample is diagnostic for the presence of the corn event MON89034 polynucleotide in the sample. The biological sample can be any sample containing MON89034 DNA including but not limited to corn oil, corn meal, corn flour, corn gluten, corn cakes, corn starch, corn steep liquor, corn tissue, corn cells, corn grain, corn pollen, corn root tissue, DDGS, and even ethanol produced as a byproduct of fermentation of such transgenic corn so long as the sample contains at least a

detectable amount of a polynucleotide diagnostic for the presence of the MON89034 event in the sample. A polynucleotide probe can be any nucleotide selected from the group consisting of a deoxyribonucleic acid, a ribonucleic acid, and a nucleotide analogue, and can be labeled with at least one fluorophores, molecule containing a radio-emitting isotope, or a hapten type molecule that can be detected specifically with an antibody or other binding type reaction.

[0062] A variety of corn comprising a DNA diagnostic for the presence of a MON89034 transgenic event DNA can be obtained by breeding a corn plant comprising transgenic corn event MON89034 DNA together with a corn plant other than event MON89034 to produce a hybrid corn plant comprising DNA diagnostic for said event. Such a hybrid corn plant comprising DNA diagnostic for the transgenic corn event MON89034 is within the scope of the present invention, as are seed produced from the hybrid (so long as the comprises DNA diagnostic for transgenic corn event MON89034), and pollen, ovule, seed, roots, or leaves of the hybrid corn plant MON89034, also to the extent these contain the diagnostic DNA sequences, and progeny produced from such embodiments.

[0063] The present invention provides a method for protecting a corn plant from lepidopteran insect infestation comprising providing in the diet of a target lepidopteran insect pest one or more transgenic corn plant cells, each corn plant cell comprising in its genome a polynucleotide corresponding to the sequence as set forth in both SEQ ID NO:1 and SEQ ID NO:2 and the contiguous nucleotide sequence as set forth in SEQ ID NO:5 between SEQ ID NO:1 and SEQ ID NO:2. The target lepidopteran insect that feeds on such transgenic corn plant cells is inhibited from further feeding on the corn plant from which the corn plant cells are derived.

[0064] Compositions are also provided by the present invention that are toxic to target lepidopteran pests of corn plants. A composition of transgenic plant cells provided in the diet of a target lepidopteran insect pest, in which each transgenic corn plant cell comprises in its genome a polynucleotide corresponding to the sequence as set forth in both SEQ ID NO:1 and SEQ ID NO:2, along with the contiguous nucleotide sequence as set forth in SEQ ID NO:5 between SEQ ID NO:1 and SEQ ID NO:2, is effective for providing protection against lepidopteran insect infestation to a corn plant or corn plant cell, so long as the corn plant or cell is expressing Cry1A.105 and/or Cry2Ab2 from the expression cassettes contained within the contiguous nucleotide sequence. Such compositions, in the form of transgenic corn seed, have been deposited with the American Type Culture Collection under accession number

PTA-9708. Such insect resistant corn plants, or parts thereof, will contain DNA in the genome of the cells of such plant that have at least one nucleotide sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, and SEQ ID NO:5. Progeny and seed of the insect resistant corn plant, in which the progeny and seed have the diagnostic sequences referred to herein, are also included within the scope of the present invention. Such insect resistant corn plants can be produced in a method comprising crossing a transgenic corn plant event MON89034 with a different corn plant, and selecting insect resistant progeny by analyzing for at least one nucleotide sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4 and SEQ ID NO:5.

[0065] The insect resistant transgenic corn event MON89034 can be combined with other transgenic varieties of corn, such as corn resistant to herbicides such as glyphosate, glufosinate, and diacamba, and the like, or corn resistant to root devouring insects as a result of insertion of sequences encoding proteins such as PS149B1 and modified Cry3Bb, or other varieties of transgenic corn resistant to lepidopteran insect infestation as a result of insertion of sequences encoding other toxin proteins such as VIP3A, Cry1Ab, and Cry1Fa and the like. Various combinations of all of these different transgenic events are bred together with the corn plants of the present invention, i.e., the MON89034 event, to provide improved varieties of hybrid transgenic corn resistant to coleopteran and lepidopteran infestation, and resistant to selective herbicides. Such varieties exhibit improved yield and drought tolerance characteristics compared to non-transgenic and individual trait transgenic varieties.

[0066] A method of producing a corn plant resistant to insect infestation is provided, wherein the corn plant comprises an insecticidally effective amount of the toxin coding sequences as set forth in SEQ ID NO:5. The method comprises extracting the toxin coding sequences from transgenic corn event MON89034 and introducing these coding sequences, alone or together, into one or more corn cells, to produce transgenic corn cells comprising these one or more toxin coding sequences. The transgenic corn cells are then grown (regenerated) into transgenic corn plants comprising the one or more coding sequences, and the transgenic plants then exhibit resistance to insect infestation.

[0067] A method for determining the zygosity of the DNA of a transgenic corn plant comprising corn event MON89034 DNA, with respect to the DNA that is diagnostic for the presence of such MON89034 DNA in a biological sample, is provided by the present

invention. The method consists of, as a first step, contacting the sample with three different primers comprising SEQ ID NO:6, SEQ ID NO:7, and SEQ ID NO:10, that when used together in a nucleic-acid amplification reaction comprising corn event MON89034 DNA, produces a first amplicon that is diagnostic for corn event MON89034, and when used in a nucleic-acid amplification reaction comprising corn genomic DNA other than MON89034 DNA, produces a second amplicon that is diagnostic for corn genomic DNA other than MON89034 DNA. The following steps consist of performing a nucleic acid amplification reaction, and comparing the amplicons produced during the thermal amplification reaction. Detecting the presence of both amplicons is diagnostic of the zygosity of the sample. Detecting only the first amplicon is indicative of the sample containing only MON89034 DNA, i.e., a homozygous sample. Detecting only the second amplicon is indicative of the sample containing no MON89034 DNA. Detecting both the first and second amplicons together in a sample is indicative of a sample containing (1) heterozygous DNA with reference to a pure sample containing only heterozygous starting material, or (2) a sample containing both homozygous and heterozygous starting sample DNA's, or (3) a sample containing some combination of homozygous, heterozygous, and/or samples other than MON89034 DNA.

[0068] The invention also provides for growing corn plants comprising DNA diagnostic for a transgenic DNA segment inserted into the genome of the cells of the corn plants. The DNA in the genome of the corn cells comprises any one or all of the sequences selected from the group consisting of:

- (a) the nucleotide sequence as set forth in SEQ ID NO:5;
- (b) both of the nucleotide sequences as set forth SEQ ID NO:1 and SEQ ID NO:2;
- (c) the nucleotide sequence as set forth at SEQ ID NO:3; and
- (d) the nucleotide sequence as set forth at SEQ ID NO:4.

[0069] The following examples are included to demonstrate examples of certain preferred embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples that follow represent approaches the inventors have found function well in the practice of the invention, and thus can be considered to constitute examples of preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific

embodiments that are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

EXAMPLES

[0070] Example 1

[0071] This Example illustrates the construction and molecular characterization of transgenic corn event MON89034.

[0072] The corn plant MON89034 was produced by an *Agrobacterium* mediated transformation process of an inbred corn line with the plasmid construct pMON38850 (the expression cassette is shown in Figure 1). The transformation method used is similar to that described in US Patent No. 6,603,061. The plasmid construct pMON38850 contains the linked plant expression cassettes with the regulatory genetic elements necessary for expression of the Cry1A.105 insecticidal protein in corn plant cells. Corn cells were regenerated into intact corn plants consisting of at least about 23,000 different transgenic events. Individual transgenic events (plants) were selected from the population of events that showed integrity of the plant expression cassettes and resistance to Lepidopteran insect larvae feeding damage. A corn plant that contains in its genome the linked plant expression cassettes of pMON38850 is an aspect of the present invention. After substantial analysis of these transgenic events, the MON89034 transgenic event was selected on the basis of its molecular characterization and the absence of any undesirable phenotypic or agronomic deficiency effects.

[0073] The sequences of the transgene genetic elements contained in MON89034 corn genome as illustrated in Figure 1 consists of the following elements each in operable linkage to each other. First at the arbitrarily defined 5' end of the sequence (i.e., near the left central portion of the segment depicted in Figure 1) is labeled a portion of the right border region (RB) from *Agrobacterium tumefaciens*. This is followed in sequence by an expression cassette consisting of an enhanced CaMV 35S promoter element (herein referred to as P-CaMV35Sen, located at positions 2350 to 2651 on SEQ ID NO:5); a wheat chlorophyll A/B binding protein untranslated leader sequence (herein referred to as L-Ta.lhcb1, located at positions 2678 to 2738 on SEQ ID NO:5); a rice actin intron sequence (herein referred to as I-Os.Act1, located at positions 2755 to 3234 on SEQ ID NO:5); a non-naturally occurring sequence encoding the chimeric gene Cry1A.105 (located at positions 3244 to 6777 on SEQ ID NO:5); and a 3' termination region from wheat (herein referred to as T-Ta.Hsp17-1:1:1, located at positions 6809 to 7018 on SEQ ID NO:5). The combination of the above referenced elements, other than the border sequence, function together when in a corn plant to

cause the expression of the Cry1A.105 insecticidal protein. These elements are then linked in sequence to another expression cassette consisting of the following elements: a Figwort mosaic promoter (located at positions 7086 to 7649 on SEQ ID NO:5), a Zea mays Hsp70 leader (herein referred to as HSP70 or I-Hsp70, located at positions 7672 to 8475 on SEQ ID NO:5), and a Zea mays chloroplast transit peptide coding sequence (herein referred to as CTP2 or TS-SSU-CTP, located at positions 8492 to 8892 on SEQ ID NO:5). These operably linked segments are then linked to a nucleotide sequence encoding the insecticidal protein Cry2Ab (located at positions 8893 to 10800 on SEQ ID NO:5), which is linked at its 3' end to a 3' non-translated region of the nopaline synthase gene of *Agrobacterium tumefaciens* (herein referred to as T-AGRtu.nos-1:1:13, located at positions 10827 to 11377 on SEQ ID NO:5). These elements flanking the Cry2Ab coding sequence function together to direct the expression of Cry2Ab when present in a corn plant. The Cry2Ab expression cassette is then followed in sequence by a nucleotide sequence consisting of a sufficient portion of the left border (LB) region from *Agrobacterium tumefaciens*.

[0074] DNA molecules useful as primers in DNA amplification methods can be derived from the sequences of the genetic elements of the transgene insert contained in the MON89034 event. These primer molecules can be used as part of a primer set that also includes a DNA primer molecule derived from the genome of event flanking the transgene insert.

[0075] The portion of the pMON38850 plasmid DNA inserted into the corn genome, giving rise to the transgenic corn plant event MON89034, consisting of the left and right border segments and the two linked plant expression cassettes (a first expression cassette encoding Cry1A.105, and a second expression cassette encoding Cry2Ab, wherein each cassette can be interchangeable as to whether one is designated as being a first or a second cassette) in between the border segments was characterized by detailed molecular analyses. These analyses were conducted to identify events that contained only a single and intact inserted segment consisting of the borders and the desired two expression cassettes in between the borders (number of integration sites within the corn genome), the copy number (the number of copies of the T-DNA within one locus), and the integrity of the inserted gene cassettes (i.e., absence of any rearrangements or sequence variation from the sequence known to be present in the plasmid pMON38850). DNA molecular probes were used that included the intact Cry1A.105 coding region and its respective regulatory elements, the promoters,

introns, and polyadenylation sequences of the plant expression cassettes, and plasmid pMON38850 backbone DNA region. The data obtained from the analyses of all events demonstrated that MON89034 contains a single T-DNA insertion with one copy of the Cry1A.105 expression cassette. No additional elements from the transformation vector pMON38850, linked or unlinked to intact gene cassettes, were detected in the genome of MON89034. Finally, PCR and DNA sequence analyses were performed to determine the 5' and 3' insert-to-plant genome junctions, confirm the organization of the elements within the insert (see for example, Figure 1), and determine the complete sequence of the DNA inserted into the corn plant genome that gave rise to the transgenic corn event MON89034. The complete inserted sequence, together with a portion of the corn genome flanking sequences at either end of the inserted DNA, is depicted in the sequence as set forth at SEQ ID NO:5.

[0076] Genomic DNA from MON89034, and non transgenic DNA from corn other than MON89034 (control DNA) was extracted from corn seed by first processing the seed (up to 200 seeds) to a fine powder in a Harbil 5G-HD paint shaker (Harbil Inc, Cincinnati, Ohio). Briefly, the powdered seed was extracted in extraction buffer (EM Science Cat. No. 3700, EM Science, Gibbstown, New Jersey, USA) and DNA precipitated from solution with isopropanol (Sigma Cat. No. I-0398, Sigma, St. Louis, MO, USA). The precipitated DNA was spooled into a microcentrifuge tube containing 70 percent ethanol. The DNA was pelleted in a microcentrifuge at maximum speed (~14,000 rpm) for ~5 minutes, vacuum-dried, and re-dissolved in TE buffer (pH 8.0). The DNA was then stored in a 4°C refrigerator. This method can be modified by one skilled in the art to extract DNA from a single corn seed.

[0077] Exemplary methods used to identify event MON89034 in a sample are described in an event specific endpoint Taqman PCR for which examples of conditions are described in Table 1 and Table 2. The DNA primers used in the assay are primers SQ2842 (SEQ ID NO:6), SQ2843 (SEQ ID NO:7), 6FAM™ labeled primer PB880 (SEQ ID NO:14) and VIC™ labeled primer PB2931 (SEQ ID NO:15), 6FAM and VIC are florescent dye products of Applied Biosystems (Foster City, CA) attached to the DNA primers. For Taqman MGB probes, the 5'-exonuclease activity of Taq DNA polymerase cleaves the probe from the 5'-end, between the fluorophore and quencher. When hybridized to the target DNA strand, quencher and fluorophore are separated sufficiently in three dimensional space to produce a fluorescent (fluorophore excitation wavelength) signal.

[0078] SQ2842 (SEQ ID NO:6), and SQ2843 (SEQ ID NO:7), when used in these reaction methods with PB880 (SEQ ID NO:14) produce a DNA amplicon that is diagnostic for event MON89034 DNA. The controls for this analysis should include a positive control from corn containing event MON89034 DNA, a negative control from non-transgenic corn or from transgenic corn other than event MON89034, and a negative control that contains no template DNA.

[0079] SQ1564 (SEQ ID NO:17) and SQ1565 (SEQ ID NO:18) when used in these reaction methods with PB351 (SEQ ID NO:21) produce an amplicon that is diagnostic of Cry1A.105 in MON89034.

[0080] These assays are optimized for use with an Applied Biosystems GeneAmp PCR System 9700 or Stratagene Robocycler, MJ Engine, Perkin-Elmer 9700, or Eppendorf Mastercycler Gradient thermocycler. Other methods and apparatus known to those skilled in the art that produce amplicons that identify the identify event MON89034 DNA is within the skill of the art.

[0081] Any probe that binds specifically to SEQ ID NO:1 or to its perfect complementary sequence in a biological sample and contains at least 11 contiguous nucleotides as set forth in SEQ ID NO:1, or as the case may be, the reverse complement of the sequence in SEQ ID NO:1, so long as the binding can be detected, is diagnostic for the presence of corn event MON89034 DNA in that sample. Any probe that binds specifically to SEQ ID NO:2 or to its perfect complementary sequence in a biological sample and contains at least 11 contiguous nucleotides as set forth in SEQ ID NO:2, or as the case may be, the reverse complement of the sequence in SEQ ID NO:2, so long as the binding can be detected, is diagnostic for the presence of corn event MON89034 DNA in that sample.

[0082] Any pair of primers that is used for or designed for use in producing an amplicon from a biological sample comprising corn DNA, and the amplicon comprises either SEQ ID NO:1 or SEQ ID NO:2, or as the case may be, comprises both sequences, is considered to be within the scope of the present invention. Any such amplicon comprising either SEQ ID NO:1 or SEQ ID NO:2 or both is considered, for the purposes of the invention disclosed herein, to be diagnostic for the presence of the corn event MON89034 DNA in such biological sample. The following example is provided as reference for one skilled in the art.

Table 1. Corn MON89034 Event Specific Endpoint Taqman PCR

Step	Reagent	Amount	Comments
1	Nuclease-free water	add to 10 μ l final volume	-
2	2X Universal Master Mix (Applied Biosystems cat. # 4304437)	5 μ l	1 X final concentration
3	Primers SQ2842 (SEQ ID NO:6), and SQ2843(SEQ ID NO:7) resuspended in nuclease-free water to a concentration of 20 μ M each)	0.5 μ l	1.0 μ M final concentration
4	Primer 6FAM™ (resuspended in nuclease-free water to a concentration of 10 μ M)	0.2 μ l	0.2 μ M final concentration
5	Internal Control Primer -SQ2842 and internal control primer SQ2843	0.2 μ l	0.2 μ M final concentration
6	Extracted DNA (template): • Samples to be analyzed (individual leaves) • Negative control • Negative control • Positive control • Positive control	3.0 μ l • 4-80 ng of genomic DNA • 4 ng of non-transgenic corn genomic DNA • no DNA template (solution in which DNA was resuspended) • 4 ng of genomic DNA from known event MON89034 heterozygous corn • 4 ng of genomic DNA from known event MON89034 homozygous corn	Diluted in water
7	Gently mix, add 1-2 drops of mineral oil on top of each reaction.		

[0083] The DNA amplification can be set up and conducted using any means for thermocycling, including manual manipulations or electronically controlled manipulations of temperature steps and cycles. Stratagene Robocycler, MJ Engine, Perkin-Elmer 9700, or Eppendorf Mastercycler Gradient thermocycler or Applied Biosystems GeneAmp PCR System 9700 or MJ Research DNA Engine PTC-225 thermal cyclers have been used successfully to conduct the following cycling parameters. When running the PCR in the

Eppendorf Mastercycler Gradient or MJ Engine, the thermocycler was cycled in the calculated mode. When using the Perkin-Elmer 9700, the cycling conditions were conducted with the ramp speed set at maximum.

Table 2. Zygoty assay thermocycler conditions

Cycle No.	Settings: Applied Biosystems GeneAmp PCR System 9700
1	50°C 2 minutes
1	95°C 10 minutes
10	95°C 15 seconds 64°C 1 minute (-1°C/ cycle)
30	95°C 15 seconds 54°C 1 minute
1	10°C soak

[0084] Example 2

[0085] This example illustrates the identification of a corn plant comprising DNA diagnostic for the transgenic corn event MON89034 in its genome and the determination of the zygoty of such corn plant.

[0086] The methods used to identify heterozygous from homozygous progeny containing event MON89034 DNA in its genome are described in a zygoty assay for which conditions are exemplified in Table 3 and Table 4. The exemplary DNA primers used in the zygoty assay are primers SQ2842 (SEQ ID NO: 6), SQ2843(SEQ ID NO:7), SQ6523(SEQ ID NO:10), SQ6524(SEQ ID NO:11), 6FAM™ labeled primer PB880(SEQ ID NO:14) and VIC™ labeled primer PB2931(SEQ ID NO:15). As indicated above, 6FAM and VIC are florescent dye products of Applied Biosystems (Foster City, CA) attached to the DNA primer.

[0087] SQ2842 (SEQ ID NO:6), SQ2843(SEQ ID NO:7), SQ6523(SEQ ID NO:10), SQ6524(SEQ ID NO:11), when used together in a thermal amplification reaction in which a biological sample containing template DNA contains DNA that is diagnostic for the presence corn event MON89034 in the sample, produces a DNA amplicon diagnostic for corn DNA other than corn event MON89034 DNA (independent of whether the corn DNA is derived from non transgenic or from some other transgenic sample). Alternatively, the reaction will

produce two different DNA amplicons from a biological sample containing DNA derived from a corn genome that is heterozygous for the allele corresponding to the inserted DNA present in transgenic corn event MON89034. These two different amplicons will correspond to a first amplicon that is derived from the wild type corn genome locus, and a second amplicon that is diagnostic for the presence of corn event MON89034 DNA. A sample of corn DNA that gives rise only to a single amplicon corresponding to the second amplicon described for the heterozygous genome is diagnostic for the presence of corn event MON89034 in the sample and is diagnostic for determining that the corn DNA used as template arise from a corn seed that is homozygous for the allele corresponding to the transgenic corn event MON89034 inserted DNA. The controls for this analysis should include a positive control from homozygous and heterozygous corn containing event MON89034 DNA, a negative control from non-transgenic corn or any other transgenic variety of corn, and a negative control that contains no template DNA. This assay is optimized for use with a Stratagene Robocycler, MJ Engine, Perkin-Elmer 9700, or Eppendorf Mastercycler Gradient thermocycler. Other methods and apparatus known to those skilled in the art that produce amplicons that identify the zygosity of the progeny of crosses made with MON89034 plants is within the skill of the art.

Table 3. Zygosity assay reaction solutions

Step	Reagent	Amount	Comments
1	Nuclease-free water	add to 5 μ l final volume	-
2	2X Universal Master Mix (Applied Biosystems cat. # 4304437)	2.5 μ l	1 X final concentration
3	Primers SEQ ID NO:6, and SEQ ID NO:7 (resuspended in nuclease-free water to a concentration of 20 μ M)	0.05 μ l	0.25 μ M final concentration
4	PB880 SEQ ID NO:14 Primer 6FAM TM (resuspended in nuclease-free water to a concentration of 10 μ M)	0.01 μ l	0.4 μ M final concentration
5	PB2931 SEQ ID NO:15 Primer VIC TM (resuspended in nuclease-free water to a concentration of 10 μ M)	0.01 μ l	0.15 μ M final concentration
6	<i>REDTaq</i> DNA polymerase (1 unit/ μ l)	1.0 μ l (recommended to switch pipets prior to next step)	1 unit/reaction
7	Extracted DNA (template): • Samples to be analyzed (individual leaves) • Negative control • Negative control • Positive control • Positive control	2.0 μ l • 4-80 ng of genomic DNA • 4 ng of non-transgenic corn genomic DNA • no DNA template (solution in which DNA was resuspended) • 4 ng of genomic DNA from known event MON89034 heterozygous corn • 4 ng of genomic DNA from known event MON89034 homozygous corn	Diluted in water
8	Gently mix, add 1-2 drops of mineral oil on top of each reaction.		

[0088] DNA amplification in a Stratagene Robocycler, MJ Engine, Perkin-Elmer 9700, or Eppendorf Mastercycler Gradient thermocycler or Applied Biosystems GeneAmp PCR System 9700 or MJ Research DNA Engine PTC-225 thermal cycler have been used

successfully to conduct the following cycling parameters. When using the Eppendorf Mastercycler Gradient or MJ Engine, the cycles were conducted in the calculated mode. When using the Perkin-Elmer 9700, the cycles were conducted with the ramp speed set at maximum.

Table 4. Zygosity assay thermocycler conditions^a

No. of Cycles in Consecutive Order	Temperature and Duration
1	50°C 2 minutes
1	95°C 10 minutes
10	95°C 15 seconds 64°C 1 minute (-1°C/ cycle)
30	95°C 15 seconds 54°C 1 minute
1	10°C soak

a: using Applied Biosystems GeneAmp PCR System 9700

[0089] Seed corresponding to the transgenic event MON89034 have been deposited on March 28, 2006 under the Budapest Treaty with the American Type Culture Collection (ATCC), 10801 University Boulevard, Manassas, Va. 20110. The ATCC accession number or patent deposit designation is PTA-7455. The deposit will be maintained in the depository for a period of 30 years, or 5 years after the last request, or for the effective life of the patent, whichever is longer, and will be replaced as necessary during that period.

[0090] Having illustrated and described the principles of the present invention, it should be apparent to persons skilled in the art that the invention can be modified in arrangement and detail without departing from such principles. We claim all modifications that are within the spirit and scope of the appended claims.

[0091] All publications and published patent documents cited in this specification are incorporated herein by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

What is claimed is:

1. A transgenic corn plant cell comprising a polynucleotide having the sequence of SEQ ID NO:1, SEQ ID NO:2, or the complement thereof.
2. The transgenic corn plant cell of claim 1, comprising a polynucleotide having the sequence of SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5 or the complement thereof.
3. The transgenic corn plant cell of claim 1 or 2, obtained from MON89034 corn event, a representative sample of seed of said corn event deposited with the American Type Culture Collection under accession number PTA-7455.
4. A transgenic corn plant, or parts thereof, comprising a polynucleotide having the sequence of SEQ ID NO:1, SEQ ID NO:2, or the complement thereof.
5. The transgenic corn plant, or parts thereof, of claim 4, comprising a polynucleotide having the sequence of SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, or the complement thereof.
6. The transgenic corn plant, or parts thereof, of claim 5, a representative sample of seed of said corn plant deposited with the American Type Culture Collection under accession number PTA-7455.
7. Seed of the transgenic corn plant of claim 4 or 5, wherein said seed comprises said polynucleotide.
8. An insect resistant corn plant, or parts thereof, wherein DNA encoding Cry2Ab and DNA having the nucleotide sequences of SEQ ID NO:1 and SEQ ID NO:2 form a part of the genome in the cells of said corn plant or parts thereof.
9. The insect resistant corn plant, or parts thereof, of claim 8, further comprising DNA encoding Cry1A.105.

10. Seed of the insect resistant corn plant of claim 8 or 9, wherein said seed comprises said DNA encoding Cry2Ab and said DNA having the nucleotide sequences of SEQ ID NO:1 and SEQ ID NO:2.
11. A composition derived from the transgenic corn plant, or parts thereof, as set forth in any one of claims 4 to 6, wherein said composition comprises a detectable amount of said DNA molecule, and wherein said composition is a commodity product selected from the group consisting of corn meal, corn oil, corn cake, corn seed, corn germ, corn starch, corn flour, corn pollen, corn silk, corn steep liquor, corn malt, corn sugar, corn syrup, margarine produced from corn oil, distillers dry goods solids (DDGS), cosmetic and bulking agent.
12. A method of producing an insect resistant corn plant comprising:
 - (a) crossing the transgenic corn plant of any one of claims 4 to 6 with a different corn plant;
 - (b) obtaining at least one progeny plant derived from the cross of (a); and
 - (c) selecting progeny that comprise the polynucleotides having the nucleotide sequences of SEQ ID NO:1 and SEQ ID NO:2, wherein said selected progeny is an insect resistant corn plant.
13. The method of claim 12, wherein said selecting step (c) includes subjecting said at least one progeny plant obtained from (b) to a nucleic acid amplification reaction, wherein progeny that produces an amplicon comprising at least one polynucleotide having the sequence of SEQ ID NO:1 or SEQ ID NO:2 is selected, or subjecting said at least one progeny plant obtained from (b) to a nucleic acid hybridization reaction, wherein progeny hybridizing to a probe that hybridizes under stringent conditions with one or more DNA sequence selected from SEQ ID NO:1 and SEQ ID NO:2 is selected.
14. A method of producing an insect resistant corn plant, comprising

- (a) transforming a corn plant cell with a polynucleotide having the sequence of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, or SEQ ID NO:5; and
 - (b) regenerating a corn plant from said transformed cell, wherein said corn plant comprises said DNA molecule and is insect resistant.
15. A method for protecting a corn plant from insect infestation, comprising providing in the diet of a Lepidopteran pest of corn an insecticidally effective amount of cell(s) or tissue(s) of the transgenic corn plant, or parts thereof, of any one of claims 4, 5, 6, 8 and 9.
16. The method of claim 15, wherein said Lepidopteran pest is selected from the group consisting of Fall armyworm (*Spodoptera frugiperda*), European corn borer (*Ostrinia nubilalis*), corn earworm (*Helicoverpa zea*), southwestern corn borer (*Diatraea grandiosella*), and black cutworm (*Agrotis ipsilon*).
17. A pair of DNA molecules comprising: a first DNA molecule and a second DNA molecule, wherein the DNA molecules are of sufficient length of contiguous nucleotides of SEQ ID NO:3 or SEQ ID NO:5 or its complement to function as DNA primers or probes diagnostic for DNA extracted from corn plant MON89034 or progeny thereof.
18. The pair of DNA molecules of claim 17, wherein said first DNA molecule comprises at least 20 contiguous nucleotides of any portion of the transgene region of SEQ ID NO:3 or SEQ ID NO:5, or the complement thereof, and said second DNA molecule comprises a similar length of a 5' flanking corn genomic DNA region of SEQ ID NO:3, or the complement thereof.
19. The pair of DNA molecules of claim 18, wherein said first DNA molecule comprises at least at least 20 contiguous nucleotides complementary to the heterologous insert DNA portion of SEQ ID NO:5, and wherein said second DNA molecule comprises at least at least 20 contiguous nucleotides from the corn genome portion of SEQ ID NO:3.

20. The pair of DNA molecule of claim 18, wherein said first DNA molecule hybridizes specifically to SEQ ID NO:5 from about nucleotide position 2060 through about nucleotide position 12,208, and wherein said second DNA molecule hybridizes specifically to the reverse complement sequence corresponding to SEQ ID NO:3 from about nucleotide position 1 through about position 2050.
21. A pair of DNA molecules comprising: a first DNA molecule and a second DNA molecule, wherein the DNA molecules are of sufficient length of contiguous nucleotides of SEQ ID NO:4 or SEQ ID NO:5 or its complement to function as DNA primers or probes diagnostic for DNA extracted from corn plant MON89034 or progeny thereof.
22. The pair of DNA molecules of claim 21, wherein said first DNA molecule comprises at least 20 contiguous nucleotides of any portion of the transgene region of SEQ ID NO:4 or SEQ ID NO:5, or the complement thereof, and said second DNA molecule comprises a similar length of a 3' flanking corn genomic DNA region of SEQ ID NO:4, or the complement thereof.
23. The pair of DNA molecules of claim 22, wherein said first DNA molecule comprises at least at least 20 contiguous nucleotides from the heterologous insert DNA portion of SEQ ID NO:5, and wherein said second DNA molecule comprises at least at least 20 contiguous nucleotides complementary to the corn genome portion of SEQ ID NO:4.
24. The pair of DNA molecules of claim 22, wherein said first DNA molecule hybridizes specifically to the reverse complement sequence corresponding to SEQ ID NO:5 from about nucleotide position 1 through about position 11305, and wherein said second DNA molecule hybridizes specifically to SEQ ID NO:4 from about nucleotide position 21 through about nucleotide position 914.
25. The pair of DNA molecules of any one of claims 17 to 24, wherein said first DNA molecule comprises SEQ ID NO:6 and said second DNA molecule comprises SEQ ID NO:7.

26. A method of detecting the presence of corn event MON89034 DNA in a biological sample, the method comprising:
- (a) contacting said biological sample with a DNA primer pair comprising DNA molecules of sufficient length of contiguous nucleotides of SEQ ID NO:3 or its complement, SEQ ID NO:4 or its complement, SEQ ID NO:5 or its complement, to function as DNA primers or probes diagnostic for DNA extracted from corn plant MON89034 or progeny thereof;
 - (b) providing a nucleic acid amplification reaction condition;
 - (c) performing said nucleic acid amplification reaction, thereby producing a DNA amplicon molecule; and
 - (d) detecting said DNA amplicon molecule, wherein detection of an amplicon comprising at least one of SEQ ID NO:1, SEQ ID NO:2 and complement thereof is indicative of the presence of said corn event MON89034 DNA in said biological sample.
27. The method of claim 26, wherein said biological sample is a DNA sample extracted from a corn plant.
28. A method of detecting the presence of corn event MON89034 DNA in a biological sample, the method comprising:
- (a) contacting said biological sample with a DNA probe that hybridizes under stringent conditions with one or more of the nucleotide sequences as set forth in SEQ ID NO:1 or SEQ ID NO:2 or the complement thereof;
 - (b) subjecting said biological sample and DNA probe to stringent hybridization conditions; and
 - (c) detecting hybridization of said DNA probe to said biological sample, wherein detection of hybridization is indicative of the presence of said corn event MON89034 DNA in said biological sample.

29. The method of claim 28, wherein said biological sample is a DNA sample extracted from a corn plant.
30. The method of claim 28 or 29, wherein said DNA probe comprises the nucleotide sequence as set forth in SEQ ID NO:1 or SEQ ID NO:2 or the complement thereof.
31. The method of claims 28 to 30, wherein said DNA probe is labeled with at least one fluorophore.
32. The method of claim 26 or 28, wherein said biological sample is selected from the group consisting of corn meal, corn oil, corn cake, corn seed, corn germ, corn starch, corn flour, corn pollen, corn silk, corn steep liquor, corn malt, corn sugar, corn syrup, margarine produced from corn oil, distillers dry goods solids (DDGS), cosmetic and bulking agent.
33. A DNA detection kit comprising: at least one DNA molecule of sufficient length of contiguous nucleotides homologous or complementary to the sequences as set forth in SEQ ID NO:3, SEQ ID NO:4, or SEQ ID NO:5 to function as a DNA primer or probe specific for corn event MON89034 and/or its progeny.
34. The DNA detection kit of claim 33, wherein said at least one DNA molecule comprises the sequence as set forth in SEQ ID NO:1, SEQ ID NO:2, or the complement thereof.
35. The DNA detection kit of claim 34, wherein said at least one DNA molecule has the nucleotide sequence as set forth in SEQ ID NO:1, SEQ ID NO:2, or the complement thereof.
36. The DNA detection kit of claim 33, wherein said kit comprises a pair of DNA molecules according to any one of claims 17 to 25.
37. A method of determining zygosity of DNA of a corn plant comprising corn event MON89034 in a biological sample comprising:

- (a) contacting said sample with a primer set comprising the nucleotide sequences as set forth in SEQ ID NO:6, SEQ ID NO:7, and SEQ ID NO:10, that (1) when used in a nucleic acid amplification reaction comprising corn event MON89034 DNA, produces a first amplicon that is diagnostic for corn event MON89034, and (2) when used in a nucleic acid amplification reaction comprising corn genomic DNA other than MON89034 DNA, produces a second amplicon that is diagnostic for corn genomic DNA other than MON89034 DNA;
 - (b) performing a nucleic acid amplification reaction; and
 - (c) detecting the amplicons so produced, wherein detection of presence of both amplicons indicates that said sample is heterozygous for corn event MON89034 DNA, wherein detection of only the first amplicon indicates that said sample is homozygous for corn event MON89034 DNA.
38. The method of claim 37, wherein said the primer set is further used together with polynucleotides having the sequence as set forth in SEQ ID NO:14 and SEQ ID NO:15.
39. An insect resistant corn plant, or a seed thereof, or a progeny thereof, the genome of said plant, seed or progeny comprises the transgene genetic elements shown in Fig. 1 and junction sequences spanning the insertion site having the sequence as set forth in SEQ ID NO:3 and SEQ ID NO:4, the transgene genetic elements shown in Fig. 1 being present in the corn event MON89034 deposited with American Type Culture Collection (ATCC) with Accession No. PTA-7455.
40. An isolated DNA polynucleotide primer molecule comprising at least 30 contiguous nucleotides of SEQ ID NO:3, or its complement that, when used in a DNA amplification method, produces an amplicon comprising SEQ ID NO:1.
41. An isolated DNA polynucleotide primer molecule comprising at least 30 contiguous nucleotides of SEQ ID NO:4, or its complement that, when used in a DNA amplification method, produces an amplicon comprising SEQ ID NO:2.

42. A DNA detection kit comprising at least one molecule of 30 or more contiguous nucleotides of SEQ ID NO:3 or SEQ ID NO:4 or a complementary sequence thereof that, when used in a DNA amplification method, produces an amplicon comprising the nucleotide sequence as set forth in SEQ ID NO:1 or SEQ ID NO:2.
43. A method of detecting the presence of DNA corresponding to corn plant MON89034 DNA in a sample, the method comprising:
- (a) contacting the sample comprising DNA with a primer pair, which when used in a nucleic acid amplification reaction with genomic DNA from corn plant MON89034, produces an amplicon comprising the nucleotide sequence as set forth in SEQ ID NO:1 or SEQ ID NO:2; and
 - (b) performing a nucleic acid amplification reaction, thereby producing the amplicon; and
 - (c) detecting the amplicon.
44. A method of detecting the presence of DNA corresponding to corn plant MON89034 DNA in a sample, the method comprising:
- (a) contacting the sample comprising DNA with a probe that hybridizes under stringent hybridization conditions with genomic DNA from corn plant MON89034 and does not hybridize under the stringent hybridization conditions with a control corn plant, wherein said probe is homologous or complementary to the nucleotide sequence as set forth in SEQ ID NO:1 or SEQ ID NO:2; and
 - (b) subjecting the sample and probe to stringent hybridization conditions; and
 - (c) detecting hybridization of the probe to the DNA.