

FIG. 1




FIG. 2





**FIG. 3**



|                                                           |        |
|-----------------------------------------------------------|--------|
| Number of unique proteins groups                          | 11,231 |
| Number of quantified proteins groups                      | 10,992 |
| Number of proteins with >1 unique peptides and quantified | 9,526  |

FIG. 4




***FIG. 5***

| <u>Cellular Viability (CTG; IC<sub>50</sub>; <math>\mu</math>M)</u> |      |
|---------------------------------------------------------------------|------|
| OCI-LY10 (MYD88 L265P):                                             | 0.19 |
| Daudi (MYD88 WT):                                                   | 0.33 |

FIG. 6





**FIG. 7**

Tumor Growth Inhibition in Implanted Ly10 (MYD88 mut) Tumor

**FIG. 8**

*In vivo* Degradation




FIG. 9




FIG. 10




FIG. 11




FIG. 12




FIG. 13






**FIG. 14**

| Dose (mg/kg) | OCI-LY10 Plasma Exposure (μM)* | Tumor (RAK4) (% Control ± SD, day 5) |
|--------------|--------------------------------|--------------------------------------|
| 25           | 0.15                           | 65.4 ± 8.2                           |
| 50           | 0.34                           | 75.3 ± 3.5                           |

**FIG. 15**

11/22



**FIG. 16****FIG. 17****MSU-Induced Gouty Arthritis in Rats**



**FIG. 18****FIG. 19**

FIG. 20

Dual-IRAK4/IMiD Degraders Show Increased IFN signaling compared to Either Drug Alone




FIG. 21




FIG. 22




FIG. 23




FIG. 24




FIG. 25




FIG. 26




**FIG. 27**




**FIG. 28**




**FIG. 29** OCI-LY10 Xenograft Studies




**FIG. 30** OCI-LY10 Xenograft Studies




FIG. 31




FIG. 32







**FIG. 33****FIG. 34**

**FIG. 35****FIG. 36**

Mouse Imiquimod (IMQ) induced skin inflammation




FIG. 37




FIG. 38

Dose-dependent IRAK4 knockdown was observed in wild-type mouse skin and spleen with treatment of I-417

