

33088-00


- 1 -

REPLACED SHEETS		SG
☒	Description : paged 14, 12-53 to paged	
☒	Claim(s) : paged 51 to paged	
☐	Drawing(s) : paged to paged	
☒	Abstract : paged 51 to paged	
DATE OF RECEIPT		

HERBICIDAL COMPOUNDS

5 This invention relates to novel oxadiazole carbonamide compounds, and to their use as herbicides. The invention also relates to a process for the preparation of such compounds, and to certain novel intermediates therefor.

10 In accordance with the present invention there is provided a compound of the general formula



wherein A represents an optionally substituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl or aralkyl group, or a group of general formula X-Y-(CH₂)_n- where X represents an optionally substituted alkyl or aryl group, n represents 1 or 2 and Y represents an oxygen atom or a group of general formula -S(O)_m- where m represents 0, 1 or 2; R¹ represents a hydrogen atom or an alkyl group; and R² represents a hydrogen atom or an optionally substituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, aralkyl or heteroaralkyl group.

Generally, when any of the above mentioned moieties comprises an alkyl group this alkyl group may be linear or branched and may suitably contain 1 to 10, preferably 1 to 6 carbon atoms. An alkenyl group may suitably contain 2 to 6 carbon atoms, preferably 2 to 4. A cycloalkyl group may have from 3 to 8 carbon atoms, most preferably 3 or, especially, 5 or 6. When any groups are designated as

This Sheet Has Been Replaced





- 2 -

being optionally substituted, the substituent groups which are optionally present may be any of those customarily employed in the development of pesticidal compounds, and/or the modification of such compounds to influence their structure/activity, persistence, penetration or other property. In relation to moieties defined above as comprising an optionally substituted alkyl, alkenyl, or cycloalkyl group, including alkyl parts of aralkyl or heteroalkyl groups, specific examples of such substituents include halogen, especially fluorine, chlorine or bromine atoms, and nitro, cyano, hydroxyl, C_{1-4} alkoxy, C_{1-4} haloalkoxy, (C_{1-4} alkoxy)carbonyl groups, amino, alkyl- and phenyl-sulphinyl, -sulphenyl and -sulphonyl groups, and C_{1-4} alkylamino groups. It is preferred, however, that such moieties are unsubstituted, or halogen-substituted. In relation to moieties defined above as comprising an optionally substituted aryl or heteroaryl group, including aryl and heteroaryl parts of aralkyl and heteroaralkyl groups, optional substituents include halogen, especially fluorine, chlorine and bromine atoms, and nitro, cyano, amino, hydroxyl, C_{1-4} alkyl, C_{1-4} haloalkyl (especially CF_3) and C_{1-4} alkoxy groups. 1 to 3 substituents may suitably be employed. A halogen atom may most suitably be a fluorine, chlorine or bromine atom.

25

Suitably, A represents an optionally substituted alkyl, cycloalkyl, alkenyl, phenyl, pyridyl or thienyl group, or a group of general formula $X-S(O)_m-(CH_2)_n-$ where X represents an optionally substituted alkyl or phenyl group, n represents 2 or, more preferably, 1, and m represents 0, 1 or, most preferably 2, or a group of general formula $X-O-(CH_2)_n-$ where X represents an optionally substituted alkyl group and n represents 1 or, preferably 2. Preferably A represents an alkyl or haloalkyl group, or an alkenyl group, an optionally

35



- 3 -

substituted phenyl group, or a pyridyl group or a thienyl group.

5 A preferred optionally substituted alkyl group A is a C_{3-6} group, preferably branched.

A preferred alkenyl group A is an allyl group.

10 A preferred optionally substituted phenyl group A is a phenyl group which is unsubstituted, or substituted by one or two moieties independently selected from halogen atoms, especially chlorine and fluorine, and alkyl, especially C_{1-4} alkyl, and haloalkyl, especially C_{1-4} haloalkyl, in particular trifluoromethyl, groups. An
15 especially preferred optionally substituted phenyl group A is unsubstituted phenyl, and phenyl mono-substituted in the 2- or 4- position, especially by chlorine, or C_{1-4} alkyl.

20 A preferred thienyl group A is 2-thienyl. Preferred pyridyl groups A are 2-pyridyl and, especially, 3-pyridyl.

Suitably, R^1 represents a hydrogen atom or a C_{1-4} alkyl group, for example ethyl or, especially, methyl.
25 Preferably R^1 represents a hydrogen atom.

Suitably, R^2 represents an optionally substituted alkyl group, an alkenyl group, or an optionally substituted aryl, aralkyl or heteroaralkyl group.

30 Preferred optionally substituted alkyl groups R^2 are C_{1-8} alkyl groups, unsubstituted or substituted by C_{3-6} cycloalkyl, especially cyclopropyl, or by alkoxy, especially methoxy, or by propargyl.

35



A preferred alkenyl group R^2 is an allyl group.

5 A preferred optionally substituted aralkyl group R^2 is phenyl-alkylene in which the phenyl group is optionally, but preferably, substituted by 1 or 2 moieties independently selected from halogen, especially fluorine, and $C_{1,4}$ alkyl, especially methyl. The alkylene group is preferably methylene or $C_{2,4}$ alkylene, straight chained or, preferably, branched, for example $-CH(CH_3)-$.


10

A preferred optionally substituted heteroaralkyl group R^2 is a furyl, pyridyl or thienyl group connected to an alkylene group, the latter suitably being as defined herein with respect to aralkyl groups R^2 . A preferred
15 heteroaryl group thereof is 2-thienyl.

A preferred optionally substituted aryl group R^2 is a phenyl which is unsubstituted, or substituted by 1 or 2 moieties selected from halogen atoms and $C_{1,4}$ alkyl groups.

20

In general preferred groups R^2 conform to the general formula



when R^3 represents a hydrogen atom or a $C_{1,2}$ alkyl group, and
25 R^4 represents a pyridyl, furyl or thienyl group, or an optionally substituted alkyl or phenyl group. Preferably, R^3 represents a hydrogen atom or a methyl group, and R^4 represents a 2-thienyl, methyl, phenyl or 3-methylphenyl group.

30

Included in the scope of the present invention are (R) and (S) isomers of compounds of general formula I and II having an optical centre, and salts, N-oxides and acid addition compounds.

35




Particularly interesting activity appears to be found in (S)-isomer compounds of general formula I and II having the R^2 group $-CH(R^3)-R^4$, wherein the C atom is optically active.


5

The invention also provides a process for the preparation of a compound of general formula I or II, which comprises reacting a respective compound of the general formula III or IV

10



(III)



(IV)

15 where X represents a halogen atom, especially chlorine, or a C_{1-4} alkoxy group, especially methoxy or ethoxy, with a compound of general formula HNR^1R^2 (V).

20 Suitably the reaction takes place at an elevated temperature, for example 50-100°C. Whilst an organic solvent may be employed, it is not necessary to add a solvent.

25 Further aspects of the invention relate to novel compounds of general formula III and IV, and to their preparation.

30 Ester compounds of general formula III may be prepared by treating a compound of general formula $A-CO-NH-CH_2-CN$ (VI) with a compound of general formula R^5-ONO (VII) where R^5 is suitably an alkyl, for example i-propyl group, and with an alkanol $X-OH$. Preferably this reaction is carried out at a lowered temperature, for example -15 to 0°C, at least at the start, although the temperature may
35 thereafter be permitted to rise, suitably to ambient



temperature. An acid chloride, for example acetyl chloride, or hydrogen chloride, is suitably present.

5 The acetonitrile compound of general formula VI may suitably be prepared by reacting an acid chloride of general formula A-COHal (VII) where Hal represents a halogen, especially chlorine, atom, with aminoacetonitrile hydrochloride, suitably at a temperature in the range 10-50°C, preferably ambient temperature, and in the presence
10 of a base. An alkali metal base, for example sodium carbonate, is suitable.

Further information applicable to each step is mentioned above for the preparation of the compounds of
15 general formula I may be found in:

- H. Brachwitz, Journal f.prakt. Chem. 314, 447 (1972) - acid chloride reaction;
- H. Brachwitz, Journal f.prakt. Chem. 314, 455 (1972) - nitrite reaction;
- 20 - Houben-Weyl 11/2, 20 ff, E5/2, 983 ff, Organikum, Deutscher Verlag der Wissenschaften, Berlin (1990), 409 f, (final step, amination).

Compounds of general formula IV may be prepared by
25 reacting an oxime amide A-C(NH₂)=NOH (IX) with a compound of general formula HalCOCOXX (X) where Hal represents a halogen, preferably chlorine atom. Suitably the reaction takes place in the presence of an organic solvent, preferably an amine, for example pyridine. Suitably an
30 elevated temperature, for example 50-100°C, is employed.

Compounds of general formula IX are suitably prepared from a nitrile compound of general formula A-CN (XI), by addition of hydroxylamine hydrochloride in the presence of



- 7 -

an alkali metal hydroxide, in the presence of water, at an elevated temperature, for example 50-100°C.

Further information applicable to each step mentioned
5 above for preparation of compounds of general formula II may be found in:

- Freund, Lenze, Ber. 24, 2154; and S. Chiou, H.G. Shine, J. Heterocycl. Chem. 26, 125 (1989) - oxime amide preparation;

10 - G. Palazzo, G. Strani, Gazz. Chim. Ital., 90, 1290-8 (1960); and S. Chiou, H.G. Shine, J. Heterocycl. Chem. 26, 125 (1989) - ester preparation;

- Houben-Weyl 11/2, 20 ff, E5/2, 983 ff, Organikum, Deutscher Verlag der Wissenschaften, Berlin 1990, 409 f.;
15 and DOS 2224338 (1972); C.A. 78, 72156p (1973); DOS 2140281 (1972); C.A. 77, 5486n (1972) - final step.

Acid chloride compounds of general formula III or IV may be prepared by converting an ester compound of general
20 formula III or IV to the acid, under standard conditions using an alkali hydroxide, and converting the acid to the acid chloride under standard conditions, suitably by means of thionyl chloride.

25 When the target compound of general formula I is an anilide, in which R¹ and R² respectively represent a hydrogen atom and an optionally substituted phenyl group, a preferred method involves reacting the appropriate ester of general formula III or IV with the appropriate
30 optionally substituted aniline in the presence of an inert solvent, for example an ether, preferably a cyclic ether, for example tetrahydrofuran, in the presence of an alkali metal base, preferably a hydride. Such a reaction is suitably effected at a temperature in the range from
35 ambient temperature to the reflux temperature.



- 8 -

Alternatively the method described above using an acid chloride of general formula III or IV may suitably be employed.

5 Further, textbook, information about the chemistry of
1,2,4-oxadiazoles can be found in Weissberger, The
Chemistry of Heterocyclic Compounds, Vol. 17, 245-262
(1962); and A.R. Katritzky, C.W. Rees, Comprehensive
Heterocyclic Chemistry, Pergamon Press, 1984 Vol. 6, 386-
10 391.

 The compounds of general formula I and II have been
found to show interesting activity as herbicides.
Accordingly, the invention further provides a herbicidal
15 composition comprising a compound of formula I or II as
defined above in association with at least one carrier,
and a method of making such a composition which comprises
bringing a compound of formula I or II into association
with at least one carrier. Preferably there are at least
20 two carriers, at least one of which is a surface-active
agent.

 The invention also provides a method of combating
undesired plant growth at a locus, comprising application
25 of such a compound or composition.

 Particularly interesting activity has been found
against grasses and broad leaf weeds, pre-and post-
emergence. Selectivity in maize and rice has also been
30 found. This activity provides a further aspect of the
present invention.

 Compounds of general formula III and IV have also
been found to have some herbicidal activity and
35 accordingly herbicidal compositions containing such



compounds, and a method of combating undesired plant growth at a locus, using such a compound or composition, is also in accordance with the present invention.

5 In a method as mentioned above, the dosage of the active ingredient, the compound of general formula I, may, for example, be from 0.01 to 10kg/ha, suitably 0.05 to 4kg/ha. The locus may be an agricultural or horticultural locus, comprising, for example, a plant or soil. In a
10 preferred method the locus contains undesired plant growth and treatment is by foliar spray application.

The invention also provides the use of a compound as defined above, as a herbicide.

15

A carrier in a composition according to the invention is any material with which the active ingredient is formulated to facilitate application to the locus to be treated, or to facilitate storage, transport or handling.
20 A carrier may be a solid or a liquid, including a material which is normally gaseous but which has been compressed to form a liquid, and any of the carriers normally used in formulating pesticidal compositions may be used. Preferably compositions according to the invention contain
25 0.5 to 95% by weight of active ingredient.

Suitable solid carriers include natural and synthetic clays and silicates, for example natural silicas such as diatomaceous earths; magnesium silicates, for example
30 talcs; magnesium aluminium silicates, for example attapulgites and vermiculites; aluminium silicates, for example kaolinities, montomorillonites and micas; calcium carbonate; calcium sulphate; ammonium sulphate; synthetic hydrated silicon oxides and synthetic calcium or aluminium
35 silicates; elements, for example carbon and sulphur;

natural and synthetic resins, for example coumarone resins, polyvinyl chloride, and styrene polymers and copolymers; solid polychlorophenols; bitumen; waxes; and solid fertilisers, for example superphosphates.

5

Suitable liquid carriers include water; alcohols, for example isopropanol and glycols; ketones, for example acetone, methyl ethyl ketone, methyl isobutyl ketone and cyclohexanone; ethers; aromatic or aliphatic hydrocarbons, for example benzene, toluene and xylene; petroleum fractions, for example kerosine and light mineral oils; chlorinated hydrocarbons, for example carbon tetrachloride, perchloroethylene and trichloroethane. Mixtures of different liquids are often suitable.

15

Agricultural compositions are often formulated and transported in a concentrated form which is subsequently diluted by the user before application. The presence of small amounts of a carrier which is a surface-active agent facilitates this process of dilution. Thus, preferably at least one carrier in a composition according to the invention is a surface-active agent. For example the composition may contain at least two carriers, at least one of which is a surface-active agent.

25

A surface-active agent may be an emulsifying agent, a dispersing agent or a wetting agent; it may be nonionic or ionic. Examples of suitable surface-active agents include the sodium or calcium salts of polyacrylic acids and lignin sulphonic acids; the condensation products of fatty acids or aliphatic amines or amides containing at least 12 carbon atoms in the molecule with ethylene oxide and/or propylene oxide; fatty acid esters of glycerol, sorbitan, sucrose or pentaerythritol; condensates of these with ethylene oxide and/or propylene oxide; condensation

35

products of fatty alcohol or alkyl phenols, for example p-octylphenol or p-octylcresol, with ethylene oxide and/or propylene oxide; sulphates or sulphonates of these condensation products; alkali or alkaline earth metal salts, preferably sodium salts, of sulphuric or sulphonic acid esters containing at least 10 carbon atoms in the molecule, for example sodium lauryl sulphate, sodium secondary alkyl sulphates, sodium salts of sulphonated castor oil, and sodium alkylaryl sulphonates such as dodecylbenzene sulphonate; and polymers of ethylene oxide and copolymers of ethylene oxide and propylene oxide.

The compositions of the invention may for example be formulated as wettable powders, dusts, granules, solutions, emulsifiable concentrates, emulsions, suspension concentrates and aerosols. Wettable powders usually contain 25, 50 or 75% w of active ingredient and usually contain in addition to solid inert carrier, 3-10% w of a dispersing agent and, where necessary, 0-10% w of stabiliser(s) and/or other additives such as penetrants or stickers. Dusts are usually formulated as a dust concentrate having a similar composition to that of a wettable powder but without a dispersant, and are diluted in the field with further solid carrier to give a composition usually containing $\frac{1}{2}$ -10% w of active ingredient. Granules are usually prepared to have a size between 10 and 100 BS mesh (1.676 - 0.152 mm), and may be manufactured by agglomeration or impregnation techniques. Generally, granules will contain $\frac{1}{2}$ -75% w active ingredient and 0-10% w of additives such as stabilisers, surfactants, slow release modifiers and binding agents. The so-called "dry flowable powders" consist of relatively small granules having a relatively higher concentration of active ingredient. Emulsifiable concentrates usually contain, in addition to a solvent and, when necessary, co-



- 12 -

solvent, 10-50% w/v active ingredient, 2-20% w/v emulsifiers and 0-20% w/v of other additives such as stabilisers, penetrants and corrosion inhibitors. Suspension concentrates are usually compounded so as to
5 obtain a stable, non-sedimenting flowable product and usually contain 10-75% w active ingredient, 0.5-15% w of dispersing agents, 0.1-10% w of suspending agents such as protective colloids and thixotropic agents, 0-10% w of
10 other additives such as defoamers, corrosion inhibitors, stabilisers, penetrants and stickers, and water or an organic liquid in which the active ingredient is substantially insoluble; certain organic solids or inorganic salts may be present dissolved in the formulation to assist in preventing sedimentation or as
15 antifreeze agents for water.

Aqueous dispersions and emulsions, for example compositions obtained by diluting a wettable powder or a concentrate according to the invention with water, also
20 lie within the scope of the invention. The said emulsions may be of the water-in-oil or of the oil-in-water type, and may have a thick "mayonnaise"-like consistency.

The invention is illustrated by the following
25 Examples. In these Examples, when an optical rotation angle is mentioned, it is the rotation in degrees caused by 1 g of the compound in 1 ml of solution, in a tube of 10 cm length, at ambient temperature, using light of the sodium D-line (589 nm).

30

EXAMPLE 1

5-(2-Fluorophenyl)-1,2,4-oxadiazole-3-carbonic acid -N-(R,S)-1-phenethyl amide

(Type I, A=2-F-Ph, R¹=H, R²=Ph-CH(CH₃)-)

35 a) 2-Fluorobenzoyl-aminoacetonitrile.



14.8 g Aminoacetonitrile hydrochloride was dissolved in 80 ml water. The solution was stirred while 25.37 g 2-fluorobenzoylchloride and a solution of 21.2 g sodium carbonate in 120 ml water were dropped simultaneously from two dropping funnels into it. The mixture was stirred overnight. The precipitate was filtered by suction and recrystallised from methanol plus water, filtered and dried.

Yield: 25.8 g beige solid (90.5%)

10 m.p. 70-72°C.

b) 5-(2-Fluorophenyl)-1,2,4-oxadiazole-3-carbonic acid methyl ester.

25.9 g Fluorobenzoyl aminoacetonitrile was dissolved in 580 ml methanol and chilled to 0-5°C. 20.5 g Isopropyl nitrite and after that 27.5 g acetyl chloride were slowly dropped into the solution while stirring in an ice bath. The mixture was stirred over a weekend at ambient temperature. The precipitate was filtered and the solution evaporated. The precipitate and residue were together heated with water for 10 minutes to 50°C, cooled and the precipitate filtered by suction and dried. The product was recrystallised from methanol.

Yield: 21.8 g white crystals (65.4%)

25 m.p. 103°C.

c) 5-(2-Fluorophenyl)-1,2,4-oxadiazole-3-carbonic acid-N-(R,S)-1-phenethyl amide.

1.6 g of 5-(2-fluorophenyl)-1,2,4-oxadiazole-3-carbonic acid methyl ester and 3.5 ml R,S-1-phenylethylamine were stirred and heated to 80°C for 3 hours. The mixture was dissolved in acetic acid ethyl ester, extracted with 2M hydrochloric acid and water, and the organic solution dried and evaporated. The residue was cleaned by



- 14 -

chromatography (silica gel, acetic acid ethyl ester/petroleum ether 4:6 v:v)

Yield: 1.7 g white solid (75.9%)

m.p. 109°C.

5 The structure was confirmed by NMR:

NMR 300 MHz, CDCl₃ ; 1.65 ppm D 3H; 5.4 ppm M 1H;

7.25-7.4 ppm M 8H; 7.65 ppm M 1H; 8.2 ppm T 1H.

EXAMPLE 2

10 5-(2-Fluorophenyl)-1,2,4-oxadiazole-3-carbonic acid -N-
(S)-1-phenethyl amide

The title compound was prepared as Example 1 above, but using (S)-1-phenylethylamine instead of the racemate.

15 Yield: 1.59 g white solid (71.1%)

m.p. 77-79°C.

Rotation: alpha (D,22): +21.87 in methanol

The structure was confirmed by NMR, 99% S-isomer.

20 EXAMPLE 3

5-t-Butyl-1,2,4-oxadiazole-3-carbonic acid-N-thiophen-
2-lylmethyl amide

(Type I, A=t-Bu, R¹=H, R²=2-thienyl-CH₂-)

25 a) Pivaloylaminoacetonitrile

The compound was prepared according to Example 1 a) from 36.17 g pivaloylchloride and 27.76 g aminoacetonitrile hydrochloride

Yield: 27.6 g white solid (65.6%)

30 m.p. 85°C.

b) 5-t-Butyl-1,2,4-oxadiazole-3-carbonic acid ethyl ester

9.5 g Pivaloylacetonitrile was dissolved in 150 ml
35 ethanol. The solution was chilled to -5°C. 7.84 g



- 15 -

Isopropyl nitrite was dropped in and hydrogen chloride gas was introduced for 5 minutes. The solution was stirred for 2 hours in an ice bath and for 2 days at ambient temperature. The solution was filtered from byproduct and

Yield: 11.5 g fluid

The structure was confirmed by NMR.

c) 5-t-Butyl-1,2,4-oxadiazole-3-carbonic acid-N-thiophen-2-ylmethyl amide.

1.5 g 5-t-Butyl-1,2,4-oxadiazole-3-carbonic acid ethyl ester was heated with 2.5 ml 2-aminomethylthiophene (excess) to 65°C for 6 hours. The mixture was dissolved in dichloromethane and washed with 2M hydrochloric acid and water. The solution was evaporated, the residue cleaned by chromatography and recrystallised with petroleum ether. Yield: 1.44 g white solid (71.4%)

m.p. 83-85°C.

The structure was confirmed by NMR:

NMR 300 MHz CDCl₃ : 1.4 ppm S 9H; 4.8 ppm D 2H; 6.95 ppm M 1H; 7.05 ppm D 1H; 7.25 ppm M 2H.

EXAMPLE 4

3-(4-Chlorophenyl)-1,2,4-oxadiazole-5-carbonic acid n-butylamide

(Type II, A=4-Cl-Ph, R¹=H, R²=n-Bu)

a) 4-Chlorobenzoyloxime amide.

10.43 g Hydroxylamine hydrochloride was dissolved in 10 ml water. 6.0 g Sodium hydroxide dissolved in 25 ml water was added. This solution was mixed with 20.6 g 4-chlorobenzonitrile in 100 ml ethanol. The charge was stirred for 70 hours at 70°C, evaporated, and the residue



- 16 -

extracted with t-butyl methyl ether. The solution was washed twice with water, dried and evaporated. The residue was cleaned by chromatography (silica gel, t-butyl methyl ether) and crystallised.

5 Yield: 18.15 g white solid (71%)
m.p. 130-132°C.

b) 3-(4-Chlorophenyl)-1,2,4-oxadiazole-5-carbonic acid ethyl ester

10 7.6 g 4-Chlorobenzoyloxime amide were dissolved in 15 ml pyridine. 6.6 g Oxalic acid monoethyl ester chloride were dropped in while chilling in an ice bath. The mixture was heated to 70°C for 20 minutes, poured onto 100 ml ice water, stirred for 15 minutes, filtered by suction and
15 dried.

Yield: 6.0 g white solid

m.p. 50-52°C

The structure was confirmed by NMR.

20 c) 3-(4-Chlorophenyl)-1,2,4-oxadiazole-5-carbonic acid-N-butyl amide.

1.5 g 3-(4-Chlorophenyl)-1,2,4-oxadiazole-5-carbonic acid ethyl ester were stirred with 1.46 g n-butyl amine at 70°C for 1 hour. The mixture was dissolved in 15 ml t-butyl
25 methyl ether and washed with 2M hydrochloric acid and water. The solution was evaporated and the residue crystallised from diisopropyl ether.

Yield: 1.1 g white solid

m.p. 87°C

30 The structure was confirmed by NMR:

NMR 300 MHz CDCl₃ : 0.9 ppm T 3H; 1.4 ppm M 2H; 1.6 ppm M 2H; 3.45 ppm Q 2H; 7.1 ppm broad 1H; 7.45 ppm D 2H; 8.0 ppm D 2H.



EXAMPLE 5

3-Isopropyl-1,2,4-oxadiazole-5-carbonic acid-N-1-
(thiophen-2-yl)-ethyl amide

5 (Type II, A=i-Pr, R¹=H, R²=2-thienyl-CH(CH₃)-)

a) Isobutyryl oxime amide.

The compound was prepared as described in Example 4 a)
from 10.35 g isobutylnitrile and hydroxylamine. The
10 reaction mixture was heated for 20 hours to 70°C and
evaporated. The residue was extracted four times with t-
butyl methyl ether, and the extract was dried and
evaporated.

Yield: 16.1 g colourless oil (98%) raw material,
15 containing 82% by weight of the desired product (assessed
by NMR).

b) 3-Isopropyl-1,2,4-oxadiazole-5-carbonic acid ethyl
ester

20 10.2 g of the raw material from Example 5 a) containing
8.36 g isobutyryloxime amide was dissolved in 30 ml
pyridine. 15 g Oxalic acid monoethyl ester chloride was
added and the mixture heated for 15 minutes to 70°C. The
mixture was poured onto 70 ml ice water, extracted with t-
25 butyl methyl ether and the solution evaporated.

Yield: 10.4 g (68.9%) colourless fluid.

c) 3-Isopropyl-1,2,4-oxadiazole-5-carbonic acid-N-1-
thiophen-2-yl)-ethyl amide

30 1.84 g 3-Isopropyl-1,2,4-oxadiazole-5-carbonic acid ethyl
ester and 2.54 g 1-thiophen-2-ylethylamine were heated for
1 hour at 80°C. The mixture was dissolved in t-butyl
methyl ether, the solution washed with 2M sulphuric acid
and water and evaporated. The product was crystallised
35 from diisopropyl ether.



- 18 -

Yield: 1.9 g white solid (70%)

m.p. 52-55°C

The structure was confirmed by NMR:

NMR 300 MHz CDCl₃ : 1.35 ppm D 6H; 1.75 ppm D 3H; 3.1 ppm

5 P 1H; 5.6 ppm P 1H; 7.0 ppm M 1H; 7.1 ppm D 1H;

7.2-7.3 ppm M 2H.

Anilides of general formula I or II ($R^1 = H$, $R^2 =$
optionally substituted phenyl) were also prepared from the
10 appropriate esters III or IV and an optionally substituted
aniline in tetrahydrofuran in the presence of sodium
hydride. An example is Example 6 hereunder.

EXAMPLE 6

15 3-Isopropenyl-1,2,4-oxadiazole-5-carbonic acid anilide
(Type II, $A = CH_2 = C(CH_3)-$, $R^1 = H$, $R^2 = Ph$)

5.46 g 3-Isopropene-1,2,4-oxadiazole-5-carbonic acid
ethyl ester and 2.79 g aniline were dissolved in 15 ml
tetrahydrofuran. A suspension of 0.79 g sodium hydride in
20 9 ml tetrahydrofuran was added dropwise with stirring.
After addition of about half of the sodium hydride the
reaction began with evolution of hydrogen. The mixture
was refluxed for two hours and was stirred overnight. 100
ml Acetic acid ethyl ester and the solution was stirred
25 with 50 ml water for a half hour. The mixture was
filtered from a byproduct, the organic solution separated,
dried and evaporated. The residue (2.8 g) was
chromatographed over a silica column using toluene as
eluant. The yield was 1.4 g of a yellow oil which was
30 crystallised from diisopropyl ether plus petrol ether to
give 0.7 g of white crystals.

m.p. 80-82°C.

The structure was confirmed by NMR:



- 19 -

¹H NMR 300 MHz: 2.2 ppm S 3H; 5.6 ppm S 1H; 6.2 ppm S 1H; 7.2 ppm T 1H; 7.4 ppm T 2H; 7.7 ppm T 2H; 8.75 ppm S broad 1H.

5 EXAMPLE 7

5-(3-Trifluoromethylphenyl)-1,2,4-oxadiazole-3-carbonic acid 3-methylanilide

(Type I, A=3-CF₃-Ph, R¹=H, R²=3-CH₃-Ph)

10 3-Trifluorobenzoyl aminoacetonitrile was prepared according to Example 1a from 14.8 g aminoacetonitrile and 33.4 g 3-trifluorobenzoylchloride. Yield: 35.4g white solid (97%), m.p. 85-87°C.

15 5-(3-Trifluoromethylphenyl)-1,2,4-oxadiazole-3-carbonic acid ethyl ester was prepared according to Example 1b from 35.4g 3-trifluoromethylbenzoyl aminonitrile and 17.58 g isopropyl nitrite in 400 ml ethanol. Yield: 32 g white solid (72.1%), m.p. 47-49°C.

20

From this ester 5-(3-trifluoromethylphenyl)-1,2,4-oxadiazole-3-carbonic acid was prepared. 32 g of the ester was dissolved in 400 ml ethanol. A solution of 5.4 g sodium hydroxide in 100 ml water was added, the mixture
25 was stirred for 1 hour at 60°C and cooled to ambient temperature. The precipitate was separated and the solution evaporated. Precipitate and residue were dissolved in water, the solution filtered and acidified with concentrated hydrogen chloride. The acid
30 precipitated and was filtered by suction and dried. Yield: 17.3 g white solid (61%), m.p. 113-114°C.

From this acid 5-(3-trifluoromethylphenyl)-1,2,4-oxadiazole-3-carbonic acid chloride was prepared. 4.6 g
35 of the acid and 12 ml thionyl chloride were refluxed for



- 20 -

1.5 hours. The mixture was evaporated. Toluene was added and the mixture was again evaporated. Yield: 4.9 g oil.

2.49 g of this raw acid chloride were dissolved in 30 ml tetrahydrofuran. 1.93 g m-toluidine and 1.82 g triethylamine were added. The mixture was refluxed for 3.5 hours, filtered and the solution was evaporated. The residue was dissolved in acetic acid ester, the solution washed with water and evaporated. The residue (2.3 g) was cleaned by flash chromatography (silica column, acetic acid ester/n-hexane 6:4). The product, the title compound, was crystallised by rubbing with petrol ether. Yield: 1.1 g (35%) white solid, m.p. 75-78°C.

The structure was confirmed by NMR:

NMR 300 MHz 1H in acetone; 2.3 ppm S 3H; 7.0 ppm D 1H; 7.3 ppm T 1H; 7.7 ppm M 2H; 8.0 ppm T 1H; 8.1 ppm D 1H; 8.5 ppm M 2H; 9.9 ppm S 1H.

Further Examples of type I and II as stated above were prepared according to the methods of Examples 1 to 7 and are mentioned in Table 1 below. Structures were checked by NMR. A symbol R or S as a superscript to a carbon atom denotes the R or S isomer as being predominant.

25

The following further compounds set out in Table 2 below are further examples of the general formula III and IV as stated above.

TABLE 1

Ex. No.	Compound Type	A	R ¹	R ²	m.p. (°C)	Rotation [α] _D ²⁵	Solvent
8	II	t-By	H	2-thienyl-CH(CH ₃)-	85-88		
9	II	Ph	H	2-thienyl-CH(CH ₃)-	129-130		
10	II	4-Cl-Ph	H	2-thienyl-CH(CH ₃)-	96-98		
11	II	4-Cl-Ph	H	2-Ph-C ⁶ H(CH ₃)-	98-99	+59.7°	CH ₃ OH
12	II	4-Cl-Ph	H	2-thienyl-CH ₂ -	116-119		
13	II	i-Pr	H	n-Bu	oil		
14	II	CH ₂ =C(CH ₃)-	H	Ph-C ⁶ H(CH ₃)-	168-170	+21.5°	CH ₃ OH
15	II	CH ₂ =C(CH ₃)-	H	n-Bu	48-51		
16	II	2,6-Cl ₂ -Ph	H	2-thienyl-CH(CH ₃)-	130-132		
17	II	3-pyridyl	H	2-thienyl-CH(CH ₃)-	124-125		
18	II	2,6-Cl ₂ -Ph	H	Ph-C ⁶ H(CH ₃)-	119-121	+21.0°	CH ₂ Cl ₂
19	II	3-pyridyl	H	Ph-C ⁶ H(CH ₃)-	75-95	+60.5°	CH ₂ Cl ₂
20	II	3-pyridyl	H	Et	138-140		
21	II	3-pyridyl	H	2-thienyl-CH ₂ -	175-177		
22	II	i-Pr	H	2-thienyl-CH(CH ₃)-	oil		
23	II	i-Bu	H	2-thienyl-CH(CH ₃)-	76-78		
24	II	Ph	H	Ph-CH(CH ₃)-	114-115		
25	II	Ph	H	cyclohexyl	109-111		



Ex. No.	Compound Type	A	R ¹	R ²	m.p. (°C)	Rotation [α] _D ²⁵	Solvent
26	II	t-Bu	H	Ph-CH(CH ₃)-	112-113		
27	II	Ph	H	Ph-C ⁸ H(CH ₃)-	glassy	+50.3°	CH ₃ OH
28	II	Ph	H	Ph-C ⁸ H(CH ₃)-	glassy	-50.5°	CH ₃ OH
29	II	Ph	H	2-thienyl-CH ₂ -	115-117		
30	I	4-Cl-Ph	H	Ph-CH ₂ -	125-130		
31	I	4-Cl-Ph	H	Ph-CH(CH ₃)-	117-120		
32	I	4-Cl-Ph	H	Ph-C ⁸ H(CH ₃)-	108-111	+33.8°	CH ₃ OH
33	I	4-i-Pr-Ph	H	Ph-CH ₂ -	126-130		
34	I	4-i-Pr-Ph	H	Ph-CH(CH ₃)-	96-100		
35	I	4-i-Pr-Ph	H	Et	84-86		
36	I	2-Cl-Ph	H	Ph-CH(CH ₃)-	77		
37	I	4-i-Pr-Ph	H	Ph-C ⁸ H(CH ₃)-	89	+28.3°	CH ₃ OH
38	I	t-Bu	H	Ph-CH(CH ₃)-	116-120		
39	I	4-i-Pr-Ph	H	(3-CH ₃ -Ph)-CH ₂ -	90.		
40	I	Ph	H	Ph-CH(CH ₃)-	110-112		
41	I	4-Cl-Ph	H	2-thienyl-CH ₂ -	150-151		
42	I	4-CH ₃ -Ph	H	Ph-CH(CH ₃)-	120		
43	I	t-Bu	H	Ph-C ⁸ H(CH ₃)-	glassy	-33.7°	CH ₃ OH
44	I	Ph	H	Ph-C ⁸ H(CH ₃)-	71-72	+32.5°	CH ₃ OH
45	I	Cl-t-Bu	H	Ph-C ⁸ H(CH ₃)-	oil	-28.3°	CH ₃ OH
46	I	i-Pr	H	Ph-C ⁸ H(CH ₃)-	oil	-35.9°	CH ₃ OH



Ex. No.	Compound Type	A	R ¹	R ²	m.p. (°C)	Rotation [α] _D ²⁰	Solvent
47	I	i-Pr	H	Ph-CH(CH ₃)-	oil		
48	I	i-Pr	H	2-thienyl-CH ₂ -	77-80		
49	I	-C(CH ₃)(C ₂ H ₅) ₂	H	Ph-CH(CH ₃)-	oil		
50	I	t-Bu	H	2-thienyl-CH(CH ₃)-	69		
51	I	2-F-Ph	H	Et	85-87		
52	I	3-CF ₃ -Ph	H	i-Pr	72-74		
53	I	3-CF ₃ -Ph	H	n-Pr	69-73		
54	I	2,4-Cl ₂ -Ph	CH ₃	CH ₃	78-80		
55	I	t-Bu	H	Ph-CH ₂	85-88		
56	I	t-Bu	H	n-Pr	oil		
57	I	t-Bu	H	i-Pr	76-79		
58	I	t-Bu	H	CH ₂ =CH-CH ₂ -	oil		
59	I	4-Cl-Ph	H	n-Pr	134-135		
60	I	4-Cl-Ph	H	i-Pr	106-108		
61	I	4-Cl-Ph	CH ₃	CH ₃	91-93		
62	I	3,4-Cl ₂ -Ph	H	n-Pr	106		
63	I	3,4-Cl ₂ -Ph	CH ₃	CH ₃	98-100		
64	I	4-Cl-Ph	CH ₃	Ph-CH ₂ -	98		
65	I	4-i-Pr-Ph	H	n-Pr	60-61		
66	I	4-i-Pr-Ph	H	CH ₂ =CH-CH ₂ -	55		
67	I	2-F-Ph	H	Ph-(CH ₂) ₃ -	77-78		



Ex. No.	Compound Type	A	R ¹	R ²	m.p. (°C)	Rotation [α] _D ²⁵	Solvent
68	I	2-Cl-Ph	H	Ph-CH ₂ -	79-85		
69	I	2-Cl-Ph	H	n-Pr	58-59		
70	I	4-1-Pr-Ph	H	Ph-C ⁶ H(CH ₃)-	88	-26.9°	CH ₃ OH
71	I	2-Cl-Ph	H	2-pyridyl-CH ₂ -	108-112		
72	I	4-CH ₃ -Ph	H	Ph-CH ₂ -	146		
73	I	t-Bu	H	Ph-CH(CH ₃)-	oil		
74	I	4-CH ₃ -Ph	H	Ph-C ⁶ H(CH ₃)-	83	+22.8°	CH ₃ OH
75	I	2-thienyl	H	Ph-C ⁶ H(CH ₃)-	110	+32.8°	CH ₃ OH
76	I	4-CH ₃ -Ph	H	2-furyl-CH ₂ -	136-138		
77	I	Ph	H	HC≡C-C(CH ₃) ₂ -	87-90		
78	I	2-Cl-Ph	H	3-pyridyl-CH ₂ -	116-120		
79	I	2-Cl-Ph	H	4-pyridyl-CH ₂ -	124-128		
80	I	t-Bu	H	CH ₃ O-(CH ₂) ₅ -	oil		
81	I	4-1-Pr-Ph	H	cyclopropyl-CH ₂ -	76		
82	I	Cl-t-Bu	H	Ph-CH(CH ₃)-	oil		
83	I	Cl-t-Bu	H	2-thienyl-CH ₂ -	oil		
84	I	1-Pr	H	Ph-CH ₂ -	53-63		
85	I	t-Bu	H	cyclopropyl-CH ₂ -	69		
86	I	t-Bu	H	(4-F-Ph)-CH ₂ -	80-83		
87	I	t-Bu	H	4-pyridyl-CH ₂ -	77-81		
88	I	t-Bu	H	CH ₃ -(CH ₂) ₅ -CH(CH ₃)-	oil		

This Sheet Has Been Replaced



Ex. No.	Compound Type	A	R ¹	R ²	m.p. (°C)	Rotation [α] _D ²⁰	Solvent
89	I	4-i-Pr-Ph	H	CH ₃ -(CH ₂) ₅ -CH(CH ₃)-	48-51		
90	I	-C(CH ₃) ₂ (C ₂ H ₅) ₂	H	Ph-C ⁵ H(CH ₃)-	oil	-28.8°	CH ₃ OH
91	I	2-Cl-Ph	H	Ph-C ⁵ H(CH ₃)-	85-87	+12.3°	CH ₃ OH
92	I	2-Cl-Ph	H	2-thienyl-C(CH ₃)-	100-102		
93	I	2-Cl-Ph	H	2-thienyl-CH ₂ -	63-65		
94	I	2-F-Ph	H	2-thienyl-CH(CH ₃)-	170-175		
95	I	i-Pr	H	2-thienyl-CH(CH ₃)-	oil		
96	I	-C(CH ₃) ₂ (C ₂ H ₅) ₂	H	2-thienyl-CH(CH ₃)-	oil		
97	II	t-Bu	H	Ph-C ⁵ H-(CH ₃)-	oil	-30.4°	CH ₃ OH
98	I	2-F-Ph	H	2-thienyl-CH ₂ -	133-135		
99	II	i-Pr	H	Ph-C ⁵ H(CH ₃)-	oil	-32.1°	CH ₂ Cl
100	II	CH ₂ =C(CH ₃)-	H	2-thienyl-CH(CH ₃)-	65-67		
101	II	CH ₂ =C(CH ₃)-	C ₂ H ₅	C ₂ H ₅	oil		
102	II	CH ₂ =C(CH ₃)-	H	2-thienyl-CH ₂ -	57-59		
103	II	2-pyridyl	H	2-thienyl-CH(CH ₃)-	117-119		
104	I	t-Bu	H	n-octyl	oil		
105	I	t-Bu	H	i-Bu	92-95		
106	I	t-Bu	H	s-Bu	44-46		
107	I	t-Bu	H	n-Bu	oil		
108	I	t-Bu	H	3-pyridyl-CH ₂ -	85-88		
109	I	cyclopentyl	H	phenyl-C ⁵ H(CH ₃)-	oil	-24.3°	CH ₃ OH

This Sheet Has Been Replaced



Ex. No.	Compound Type	A	R ¹	R ²	m.p. (°C)	Rotation [α] _D ²⁵	Solvent
110	I	cyclopentyl	H	phenyl-CH(CH ₃)-	45-48		
111	I	t-Bu	H	2-thienyl-CH(C ₂ H ₅)-	76		
112	I	cyclohexyl	H	phenyl-CH(CH ₃)-	40		
113	I	i-Bu	H	2-thienyl-CH(CH ₃)-	oil		
114	I	i-Bu	H	2-thienyl-C ⁵ H(CH ₃)-	oil	-27.6°	CH ₃ OH
115	I	cyclopentyl	H	2-thienyl-CH(CH ₃)-	oil		
116	I	n-Pr	H	2-thienyl-CH(CH ₃)-	oil		
117	I	n-Pr	H	2-thienyl-C ⁵ H(CH ₃)-	oil	-33.5°	CH ₃ OH
118	I	4-F-Ph	H	2-thienyl-CH ₂ -	106		
119	I	4-F-Ph	H	2-thienyl-C ⁵ H(CH ₃)-	75-77	+18.5°	CH ₃ OH
120	I	i-Pr	H	2-thienyl-CH(CH ₃)-	82		
121	II	Ph-SO ₂ -CH ₂	H	2-thienyl-CH(CH ₃)-	108-111		
122	II	4-CH ₃ -Ph	H	2-thienyl-CH(CH ₃)-	112-114		
123	II	6-Cl-pyrid-2-yl	H	2-thienyl-CH(CH ₃)-	147-148		
124	II	4-CH ₃ -Ph	H	phenyl-C ⁵ H(CH ₃)-	112-113	+82.2°	CH ₂ Cl ₂
125	II	6-Cl-pyrid-2-yl	H	phenyl-C ⁵ H(CH ₃)-	135-138	+73.5°	CH ₂ Cl ₂
126	I	CH ₂ Cl-C(CH ₃) ₂ -	H	2-thienyl-CH(CH ₃)-	oil		
127	I	Ph	H	2-thienyl-CH(CH ₃)-	147		
128	I	4-F-Ph	H	2-thienyl-CH(CH ₃)-	110		
129	II	Ph-CH=CH-	H	phenyl-C ⁵ H(CH ₃)-	98-115	+76.7°	CH ₂ Cl ₂
130	II	Ph-CH=CH-	H	2-thienyl-CH ₂ -	135-136		



Ex. No.	Compound Type	A	R ¹	R ²	m.p. (°C)	Rotation [α] _D ²⁵	Solvent
131	II	n-Bu	H	2-thienyl-CH(CH ₃)-	oil		
132	II	i-Bu	H	2-thienyl-CH(CH ₃)-	oil		
133	II	i-Bu	H	2-thienyl-C ⁶ H(CH ₃)-	white solid	-7.0°	CH ₂ Cl
134	II	4-Cl-Ph	H	5-chloro-thien-2-yl-CH(CH ₃)-	127-130		
135	II	4-CF ₃ -Ph	H	2-thienyl-CH(CH ₃)-	90-94		
136	II	4-CH ₃ O-Ph	H	2-thienyl-CH(CH ₃)-	88-91		
137	II	4-CH ₃ O-Ph	H	phenyl-C ⁶ H(CH ₃)-	oil	+58.0°	CH ₂ Cl ₂
138	II	4-CF ₃ -Ph	H	phenyl-C ⁶ H(CH ₃)-	oil	+38.5°	CH ₂ Cl ₂
139	II	4-CF ₃ -Ph	H	2-thienyl-CH ₂ -	102-104		
140	II	4-CF ₃ -Ph	H	2-thienyl-CH ₂ -	122-124		
141	II	4-F-Ph	H	2-thienyl-CH ₂ -	92-95		
142	I	cyclohexyl	H	2-thienyl-CH(CH ₃)-	52-55		
143	I	cyclohexyl	H	phenyl-C ⁶ H(CH ₃)-	oil	-24.9°	CH ₃ OH
144	II	t-Bu	H	2-thienyl-CH ₂ -	70-71		
145	II	i-Pr	H	2-thienyl-CH ₂ -	oil		
146	II	Ph-CH=CH-	H	2-thienyl-CH(CH ₃)-	89-91		
147	I	2-F-Ph	H	Ph-C ⁶ H(CH ₃)-	76-77	-22.1°	CH ₃ OH
148	II	CH ₃ -S-CH ₂	H	2-thienyl-CH(CH ₃)-	oil		
149	II	CH ₃ -O-(CH ₂) ₂ -	H	phenyl-C ⁶ H(CH ₃)-	oil	Rotation angle not measured	



Ex. No.	Compound Type	A	R ¹	R ²	m.p. (°C)	Rotation [α] _D ²⁵	Solvent
150	II	CH ₃ -O-(CH ₂) ₂ -	H	2-thienyl-CH(CH ₃)-	oil		

This Sheet Has Been Replaced



TABLE 2

Example No.	Compound Type	A	X	m.p. (°C)
151	IV	4-Cl-Ph	Et	50-52
152	IV	i-Pr	Et	oil
153	IV	2,6-Cl ₂ -Ph	Et	68-69
154	IV	3-pyridyl	Et	oil
155	IV	Ph	Et	53-54
156	IV	t-Bu	Et	oil
157	III	4-i-Pr-Ph	Et	oil
158	III	2-Cl-Ph	Et	oil
159	IV	CH ₂ =C(CH ₃)-	Et	oil
160	IV	2-pyridyl	Et	85-88
161	III	-C(CH ₃)(C ₂ H ₅) ₂	Et	oil
162	III	CH ₂ =CH-	Et	49-50
163	III	4-F-Ph	Et	70
164	III	2-Cl-Ph	Et	55-60
165	III	3,4-Cl ₂ -Ph	Et	94-96
166	III	4-CH ₃ -Ph	Et	69-70
167	III	2-thienyl	Et	69-70
168	IV	4-pyridyl	Et	87-88
169	IV	4-F-Ph	Et	49-51
170	IV	n-Bu	Et	oil



Example No.	Compound Type	A	X	m.p. (°C)
171	IV	benzyl	Et	oil
172	IV	4-CF ₃ -Ph	Et	58-61
173	IV	6-Cl-pyrid-2-yl	Et	96-99
174	IV	Ph-SO ₂ -CH ₃ -	Et	136-141
175	IV	4-CH ₃ O-Ph	Et	oil
176	IV	4-CH ₃ -Ph	Et	57-59
177	III	i-Pr	Et	oil

This Sheet Has Been Replaced



Ex. No.	Compound Type	A	R ¹	R ²	m.p. (°C)	Rotation [α] _D ²²	Solvent
178	II	i-Pr	H	(4-t-Bu-Ph)-CH ₂ -	94-97		
179	II	4-Cl-Ph	H	(4-4-Bu-Ph)-CH ₂ -	128-133		
180	II	2-F-Ph	H	(2-thienyl)-CH(CH ₃)-	125-128		
181	II	2-F-Ph	H	(2-thienyl)-CH ₂ -	119-121		
182	II	2-F-Ph	H	Ph-CH(CH ₃)-(S)	oil		
183	II	2-F-Ph	H	(4-t-Bu-Ph)CH ₂ -	69-77		
184	II	3-F-Ph	H	(2-thienyl)-CH(CH ₃)-	116-118		
185	II	3-F-Ph	H	(2-thienyl)-CH ₂ -	116-118		
186	II	3-F-Ph	H	Ph-CH(CH ₃)-(S)	oil		
187	II	t-Bu	H	Ph-	91-94		
188	II	t-Bu	H	Tetralin-1-yl-	128-130		
189	II	t-Bu	H	Tetralin-5-yl-	185-189		
190	II	4-F-Ph	H	Tetralin-1-yl-	142-144		
191	II	3-F-Ph	H	Tetralin-1-yl-	110-111		
192	II	2-F-Ph	H	Tetralin-1-yl-	152-155		
193	II	t-Bu	H	(1-Naphthyl)-CH(CH ₃)-(S)	glass		
194	II	4-F-Ph	H	(1-Naphthyl)-CH(CH ₃)-	119-122		
195	II	4-F-Ph	H	(1-Naphthyl)-CH(CH ₃)-(S)	135-138		
196	II	4-F-Ph	H	(3-thienyl)-CH(CH ₃)-	104-106		
197	II	2,6-F ₂ -Ph-	H	(2-thienyl)-CH ₂ -	100-102		
198	II	t-Bu	H	(3-thienyl)-CH(CH ₃)-	81-82		
199	II	2,6-F ₂ -Ph	H	(2-thienyl)-CH(CH ₃)-	97-99		

This Sheet Has Been Replaced



Ex. No.	Compound Type	A	R ¹	R ²	m.p. (°C)	Rotation [α] _D ²²	Solvent
200	II	2,6-F ₂ -Ph	H	Ph-CH(CH ₃) - (S)	87-91		
201	II	2,4-F ₂ -Ph	H	(2-thienyl) - CH(CH ₃) -	102-103		
202	II	2,4-F ₂ -Ph	H	Ph-CH(CH ₃) - (S)	44-47		
203	II	4-F-Ph	H	(2-thienyl) - CH(CH ₃) - (S)	77-80	+ 16.34°	CH ₂ Cl ₂
204	II	t-Bu	H	(2-thienyl) - CH(CH ₃) - (S)	oil	+ 35.62°	CH ₂ Cl ₂
205	II	2-F-Ph	H	2-thienyl - CH(CH ₃) - (S)	79-82	+ 2.19°	CH ₂ Cl ₂
206	II	CH ₃ -O-(CH ₂) ₂	H	(2-thienyl) - CH ₂ -	oil		
207	II	2-Cl-Pyrid-3-yl	H	(2-thienyl) - CH ₂ -	97-102		
208	I	t-Bu	H	(1-naphthyl) - CH(CH ₃) -	oil		
209	I	4-F-Ph	H	(3-thienyl) - CH(CH ₃) -	108-110		
210	I	t-Bu	H	(2-thienyl) - CH(CH ₃) - (S)	oil	-53.97°	MeOH
211	I	t-Bu	H	(3-thienyl) - CH(CH ₃) -	75		
212	I	t-Bu	H	(1-naphthyl) - CH(CH ₃) -	89		
213	I	4-F-Ph	H	(1-naphthyl) - CH(CH ₃) -	127-128		
214	II	2,4-F ₂ -Ph	H	(3-thienyl) - CH(CH ₃) -	130-131		
215	I	2-F-Ph	H	(3-thienyl) - CH(CH ₃) -		130	
216	I	4-i-Pr-Ph	H	(3-thienyl) - CH(CH ₃) -		89-91	
217	II	2-Cl-Pyrid-3-yl	H	(2-thienyl) - CH(CH ₃) -		88-9	
218	II	2-Cl-Pyrid-3-yl	H	Ph-CH(CH ₃) -		oil	
219	II	4-F-Ph	H	(2-thienyl) - CH(CH ₃) - (S)	59-60	-18.63°	CH ₂ Cl ₂
220	I	n-Heptyl	H	(2-thienyl) - CH(CH ₃) - (S)	oil		

This Sheet Has Been Replaced



Ex. No.	Compound Type	A	R ¹	R ²	m.p. (°C)	Rotation [α] _D ²²	Solvent
221	I	Cyclopentyl	H	(3-Me-Ph)-CH ₃ -	oil		
222	I	4-Cl-Ph	H	(3-thienyl)-CH(CH ₃)-	117		
223	I	Cyclopentyl	H	(3-thienyl)-CH(CH ₃)-	oil		
224	I	Ph	H	(3-thienyl)-CH(CH ₃)-	126		
225	II	i-Bu	H	(3-thienyl)-CH(CH ₃)-	68		
226	II	2,6-F ₂ -Ph	H	(3-thienyl)-CH(CH ₃)-	137-140		
227	I	4-F-Ph	H	(4-Cl-Ph)-CH(CH ₃)-	137-138		
228	II	4-F-Ph	H	(4-Cl-Ph)-CH(CH ₃)-	135		
229	I	2-Cl-Pyrid-3-yl	H	(2-thienyl)-CH ₂ -	97-98		
230	I	2-Cl-Pyrid-3-yl	H	Ph-CH(CH ₃)-(S)	110	+9.05°	MeOH
231	I	2-Cl-Pyrid-3-yl	H	(2-thienyl)-CH(CH ₃)-	71		
232	I	n-Bu	H	Ph-CH(CH ₃)-(S)	oil	-31.0	MeOH
233	I	n-Bu	H	(2-thienyl)-CH(CH ₃)-	oil		
234	I	n-Bu	H	(2-thienyl)-CH ₂ -	oil		
235	I	2-Cl-Pyrid-3-yl	H	Ph-CH(CH ₃)-	115-116		
236	I	Ph	H	(2-thienyl)-CH ₂ -	164		

This Sheet Has Been Replaced



Herbicidal Activity

To evaluate their herbicidal activity, compounds of formula I, II, III and IV according to the invention were tested using a representative range of plants: maize, Zea mays (Z); rice, Oryza sativa (R); barnyard grass, Echinochloa crusgalli (G); oat, Avena sativa (O); linseed, Linum usitatissimum (L); mustard, Sinapsis alba (M); sugar beet, Beta vulgaris (B) and soya bean, Glycine max (S).

The tests fall into two categories, pre-emergence and post-emergence. The pre-emergence tests involved spraying a liquid formulation of the compound onto the soil in which the seeds of the plant species mentioned above had recently been sown. The post-emergence tests involved two types of test, viz., soil drench and foliar spray tests. In the soil drench tests the soil in which the seedling plants of the above species were growing was drenched with a liquid formulation containing a compound of the invention, and in the foliar spray tests the seedling plants were sprayed with such a formulation.

The soil used in the tests was a prepared horticultural loam.

The formulations used in the tests were prepared from solutions of the test compounds in acetone containing 0.4% by weight of an alkylphenol/ethylene oxide condensate available under the trade mark TRITON X-155. These acetone solutions were diluted with water and the resulting formulations applied at dosage levels corresponding to 5kg and 1kg of active material per hectare in a volume equivalent to 900 litres per hectare in the soil spray and foliar spray test, and at a dosage of level equivalent to 10 kilograms of active material per hectare in a volume equivalent to approximately 3,000 litres per hectare in the soil drench tests.



In the pre-emergence tests untreated sown soil and in the post-emergence tests untreated soil bearing seedling plants were used as controls.

- 5 The herbicidal effects of the test compounds were assessed visually twelve days after spraying the foliage and the soil, and thirteen days after drenching the soil and were recorded on a 0-9 scale. A rating 0 indicates growth as untreated control, a rating 9 indicates death.
- 10 An increase of 1 unit on the linear scale approximates to a 10% increase in the level of effect.

- 15 The results of the tests are set out in Table 3 below. A blank space in Table 3 for a compound for which other results are given indicates a rating 0. Where a compound has no results and entirely blank spaces, this indicates that data is not yet available, for this compound. An asterisk denotes that a test was not undertaken, or was not completed.

20

This Sheet Has Been Replaced



TABLE 3

Ex. No.	Soil drench 10 Kg/ha								Dosage kg/ha	Foliar Spray								Pre-emergence							
	Z	R	G	O	L	M	B	S		Z	R	G	O	L	M	B	S	Z	R	G	O	L	M	B	S
1			6	5	4	6	8		5 1	3		9	6	7	9										
2			5		2	7	3	5	5 1	4		9	5	8	8	9	7			4			3	5	3
3	4	4	6	4	7	6	4	6	5 1	4	4	9	4	8	8	9	9	2		9	4	3	9	9	3
4									5 1	2		4	1	7	9	9	4						6	1	
5	7		9	8	9	9	9	8	5 1	2		9	7	9	9	9	8	4		8	7	7	8	8	5
7									5 1	2		3		1	6	4	2								
8	7	2	9	8	9	9	9	8	5 1	2	4	9	7	9	9	9	8	4	5	8	7	6	9	8	2
9					5	6	9		5 1			9	5	8	8	9	6								
10									5 1	5		8	6	9	8	9	7					4	2		
11									5 1	6		9	6	8	8	9	7					4	4	2	

This Sheet Has Been Replaced



Ex. No.	Soil drench 10 Kg/ha								Dosage kg/ha	Foliar Spray								Pre-emergence							
	Z	R	G	O	L	M	B	S		Z	R	G	O	L	M	B	S	Z	R	G	O	L	M	B	S
12									5 1	4		8	7	9	8	9	6						5	2	1
13									5 1	2		8	5	8	8	9	5						4		
14									5 1			7	1	6	7	2	4				6	2	4	2	
15									5 1	2		8	6	7	9	9	6				4				
16									5 1			6	2	2	6	5	2								
17									5 1			7		7	9	9	2				4		4	5	6
18									5 1	2		9	6	9	9	7	6				4		4	2	2
19									5 1			8		8	8	7	4						2	7	5
20									5 1	4		8	2	7	8	9	2								
21									5 1			2			2	5									
22									5 1	4		8		7	5	7	4			6			7	6	5

This Sheet Has Been Replaced



Ex. No.	Soil drench 10 Kg/ha							Dosage kg/ha	Foliar Spray							Pre-emergence									
	Z	R	G	O	L	M	B		S	Z	R	G	O	L	M	B	S	Z	R	G	O	L	M	B	S
23					5	7			5 1		8 7	4	7 6	8 7	9 8	6			6 2	2		7 2		6 5	
24					7	6	2		5 1		7 6		8 7	8 8	9 8	6 5									
25									5 1				4	7 2											
26	7	4	8	7	9	9	9	7	5 1		7 5	4 2	7 6	9 8	9 8	5 4			2			6 5	7 5		
27						6	2		5 1		8 7	6 2	5 4	7 7	9 8	2			7 2			8 5	5 2	4 2	
28									5 1		9 8	5 2	8 7	8 6	8 8	6 5			7 6			7 5	7 4		
29			4		5	6	5		5 1		8 7	4 2	5 2	4 2	8 8	4 2						6 4	2		
30									5 1		4	2	7 5	9 9	9 4	4 4									
31									5 1		9 8	4 2	8 7	9 8	9 9	7 6									
32						4	2		5 1		8 7	4 2	9 6	7 7	9 9	6 4									
33									5 1		6 2		8 4	7 7	9 9										

This Sheet Has Been Replaced



Ex. No.	Soil drench 10 Kg/ha							Dosage kg/ha	Foliar Spray							Pre-emergence								
	Z	R	G	O	L	M	S		Z	R	G	O	L	M	B	S	Z	R	G	O	L	M	B	S
34								5 1	4		9 8	4 2	9 8	8 7	9 9	4 2			5			6 2	4 5	
35								5 1			8 2		7 6	6 6	9 9						4 2	3		
36								5 1	2		7 6	4 2	9 8	8 7	9 9	4 2								
37								5 1	5 2	4 1	9 8	4 2	9 8	7 7	9 9	7 5								
38	4			6	2	5	5	5 1	3		9 7	4 2	8 5	9 8	9 9	8 7	6 5	4 8	7 6	8	9 9	9 9	6 5	
39								5 1			4		6 5	9 7	9 9	4 2								
40					2	4	3	5 1	4 2	3 1	9 8	4 2	8 6	9 8	9 9	7 6				4 1	7 4	7 2		
41								5 1		4 2	8 6	2	8 6	9 7	9 9	6 5					6 2		2	
42								5 1	3	2 2	9 8	4 2	9 8	9 8	8 8	6 5					6 5	6 2	3	
43	7	9	8	8	9	9	8	5 1	4 2	8 8	9 9	8 6	9 7	9 8	9 7	7 5	2	9 7	4 6	8 9	9 9	9 8	6 5	
44	7	6	7	8	9	9	7	5 1			9 8	4 2	9 7	7 6	9 9	8 6		6 2		2	9 7	9 7	2	

This Sheet Has Been Replaced



Ex. No.	Soil drench 10 Kg/ha								Dosage kg/ha	Foliar Spray								Pre-emergence							
	Z	R	G	O	L	M	B	S		Z	R	G	O	L	M	B	S	Z	R	G	O	L	M	B	S
45	7	8	8	8	8	9	9	8	5 1	4 2	1	9 8	7 4	8 2	9 8	9 9	7 6	7		7					
46	7	8	9	8	8	9	9	8	5 1	6 2	4	9 9	7 2	8 4	9 6	9 9	8 6	6		8 4	6		4	6	
47	7	8	9	7	9	9	8	8	5 1	7 5	4 2	9 8	8 5	8 7	8 8	9 8	7 5	5		7 5		6 2	7 5	8 4	2
48	7	5	9	6	8	9	9	8	5 1	2		9 8	4 2	7 2	9 8	9 8	7 6	5	2	8 8	6 5	7 6	9 7	7 6	2
49	6		5			8	9		5 1			9 8	6 2	7 5	8 7	9 9	4 2			2			7 2	2 1	
50	8	5	9	8	7	8	9	8	5 1	6 2		9 9	7 5	9 8	9 9	9 9	8 7	4		9 7	6 5	7 8	8 8	4 2	
51					*	3	4	1	5 1		4 1	6 4	2 2	6 2	8 6	8 6	7 5					2	5 1	3	
52									5 1	2	1	4	2		3 2		3 1								
53									5 1	2		5 2	1	2	5 3	3 1									
54									5 1		2	6 2	2	6 3	8 4	8 2	4 1								
55					4	8	6	4	5 1		2	6 2	3	6 3	9 7	9 4	5 2		6	5		7 2	7 2		



Ex. NO.	Soil drench 10 Kg/ha								Dosage kg/ha	Foliar Spray								Pre-emergence							
	Z	R	G	O	L	M	B	S		Z	R	G	O	L	M	B	S	Z	R	G	O	L	M	B	S
56					2	3	5	4	5 1			2		2	6 3	5	4 2			6			3	3	3
57						4	2	3	5 1					5	6 2	4 2	3 1								
58					4	4	4	*	5 1					4	7 3	5	5 3								
59									5 1					6 2	9 8	8 4	3								
60					4	7	2		5 1			3	2	6 4	9 7	7 5	2					6	5		
61									5 1			4	2	6 2	5 2	7 6	6	2	*	*	*	*	*	*	
62									5 1			3		4 1	5 4	8 6	4								
63									5 1			4		4 3	7 5	4 3	4 3								
64									5 1					7 6	9 8	7 7									
65									5 1					6 2	6 6	8 7						3	2		
66						5			5 1					6 2	7 5	9 7						7 2	3		

This Sheet Has Been Replaced



Ex. No.	Soil drench 10 Kg/ha.							Dosage kg/ha	Foliar Spray							Pre-emergence									
	Z	R	G	O	L	M	B		S	Z	R	G	O	L	M	B	S	Z	R	G	O	L	M	B	S
67								5 1	3				2	4	3										
68								5 1	4 2		7 5	4	7 5	8 7	9 8	3							2	3	
69								5 1			4 2	2	5 2	7 4	6 2	5 2							3		
70								5 1	2		7 2		4 2	9 5	8 4	3 2									
71								5 1			4		2	6 5	4 2	3									
72								5 1						4 2	6 5										
73	4			2	7	8	5	5 1			9 2	2	5 4	6 2	7 2	5 2			2	5 2		7 2	5 2		
74					7	7	5	5 1					8 5	8 8	9 8	4									
75	4	3	5		4	8	6	5 1			8 6		7 5	8 7	9 8	5 4									
76								5 1						5 1	1										
77					3	8	7	5 1			9 2	1	6 2	9 4	8 2	6 4						4 2			2

This Sheet Has Been Replaced



Ex. No.	Soil drench 10 Kg/ha								Dosage kg/ha	Foliar Spray								Pre-emergence							
	Z	R	G	O	L	M	B	S		Z	R	G	O	L	M	B	S	Z	R	G	O	L	M	B	S
78									5 1			4			7 5	1	2								
79									5 1	1		7 2	3	7 5	8 7	9 8	5								
80					2	1	2	4	5 1			3	1	4 2	4 2	2 3	4								
81									5 1	1		3		8 6	8 7	8 7	2								
82	6	2	7	7	6	9	8	5	5 1	4		9 4	5	7 2	8 7	9 8	8 2			2			5	8	
83	6		5	4		8	8	7	5 1			9 7	2	6 2	9 8	8 4						6	2		
84	4		7		4	7	6	5	5 1			9	2	4 2	8 4	9 2	6 2								
85	4		5	6	7	6	5	5	5 1			4	1	4 2	5 6	6 2	5 2			4 2			2	1	4
86	4		5	6	7	8	8	5	5 1			6 2	4 2	7 5	8 7	9 7	6 5	2		4 2			5	6	2
87			4	3	4	7	3		5 1		2	7	2	4 3	6 4	7 4	3			3	1		4	1	
88						2	2	4	5 1	2		7 4			2	4	2								

This Sheet Has Been Replaced



Ex. No.	Soil drench 10 Kg/ha								Dosage kg/ha	Foliar Spray								Pre-emergence							
	Z	R	G	O	L	M	B	S		Z	R	G	O	L	M	B	S	Z	R	G	O	L	M	B	S
89									5 1	1 1		3 2		1 1		4									
90	5	4	8	5	2	9	9	5	5 1	4 2		9 8	4 2	8 5	9 8	9 8	7 5	2		4 2	4 3		6 5	4 2	
91	2		5			5	6		5 1	4		9 8	7 4	8 7	8 8	8 8	7 7			3		2	3	5 4	2
92									5 1			8 8	6 5	7 6	8 7	9 8	6 5								
93									5 1	2		9 8	6 2	7 5	7 7	9 8	2								
94			2		6	8	8	4	5 1	4		9 8	4 2	9 8	8 8	9 8	6 5								
95	7	*	8	7	8	8	8	7	5 1	2		9 9	4 2	8 5	9 9	8 6		2		9 8	7 2	7 6	8 8	8 7	2
96	5		7	4	8	8	5	4	5 1	2		9 8	7 2	9 8	8 8	9 8	6 2			7 7		2	7 2	7 5	
97	4	1	6	7	7	7	8	7	5 1		3	8 7	3 2	7 6	9 9	9 9	7 3	2	2	6 2	2	3	8 4	8 6	2
98			4		7	6			5 1			9 8	6	9 8	8 8	9 8				4 2	2	4	5 4	4 4	
99	5		7		5	7	5	7	5 1	5	2	9 8	7 5	6 5	9 8	9 8	6 2	2		8 7	2	5	9 7	8 7	



Ex. No.	Soil drench 10 Kg/ha								Dosage kg/ha	Foliar Spray								Pre-emergence							
	Z	R	G	O	L	M	B	S		Z	R	G	O	L	M	B	S	Z	R	G	O	L	M	B	S
100			7		6	5	7	4	5 1			9 8	7 5	8 6	8 7	9 9	6 5			9 7	6	7 5	2		
101									5 1			7 5	2		6 2	5 2	2								
102					3	5			5 1			9 8	6	5 2	9 8	9 7	4 2		6 5		2				
103			7	4	9	8		2	5 1			8 7	6 2	8 5	4 2	9 9	2		7 2	6 2	2	9 7			
104									5 1			2			4 2	5 2	4 2								
105	2		4		2	5	4	4	5 1						5 2	4 2	4 2								
106	2		5	4	3	6	5	4	5 1				6	4	7 5	4 2	5		4		2	4 2	5 2	6 2	
107	4		5	2	7	8	6	5	5 1				5 2	4 2	8 7	6 2	5 2	2	7 5	2	6 5	2 4	2		
108						4	1	1	5 1				3 2		6 2	6 4	3				1				
109	*	*	*	*	*	*	*	*	5 1			5 2	6 2	6 4	8 8	9 9	8 6	2	7 2		6 6	8 2	2		
110	7	6	8	6	9	9	9	7	5 1					6 2	8 8	9 8	6 5								



Ex. No.	Soil drench 10 Kg/ha								Dosage kg/ha	Foliar Spray								Pre-emergence							
	Z	R	G	O	L	M	B	S		Z	R	G	O	L	M	B	S	Z	R	G	O	L	M	B	S
111	*	*	*	*	*	*	*	*	5 1	2	3	9	4	7	9	8	6			9	6	5	1	4	
112	4		5	6	7	9	7	6	5 1	3	7	9	4	8	9	7	7					2		5	
113	5		8	8	5	9	9	8	5 1	4		9	6	8	9	9	8	6		8	5	6	9	9	4
114	7		7		6	7	8	9	5 1	4		9	6	8	9	9	7	6		8		4	9	9	
115	6		8		7	9	9	8	5 1	2		9	6	9	9	9	8			7	3	2	9	8	2
116	7		8	9	9	9	9	8	5 1	5		9	7	7	8	8	4	4		9	2	5	8	8	2
117	7	7	8	7	7	9	9	8	5 1	5		7	6	5	8	9	8	5	3	8	5	7	8	7	6
118					4	6	9	4	5 1	2	5	7	5	9	9	9	6								
119			8	7	7	6	9	6	5 1	3	4	9	4	8	9	9	8			1	1		7	9	3
120					3			1	5 1	5	5	9	6	9	8	9	7								
121					6	9	5	7	5 1		2	8	2	8	9	9	6						6	4	

This Sheet Has Been Replaced



Ex. No.	Soil drench 10 Kg/ha								Dosage kg/ha	Foliar Spray								Pre-emergence							
	Z	R	G	O	L	M	B	S		Z	R	G	O	L	M	B	S	Z	R	G	O	L	M	B	S
122					5		6		5			9	7	8	9	9	8								
123			3	8	5	7	9	5	1			9	8	7	9	9	7								
124						6	4		5		6	5	7	8	9	9	7								
125	7	8	9	9	7	9	9	7	1		5	4	8	8	9	9	8					6	5		
126									5		6	3	5	8	7	9	9	3		3	4	4	9	9	8
127									1		6	1	9	6	7	9	9					4	4	3	2
128									5				8	8	9	9	7								
129							6		1				3	9	9	9	6					4	4	3	
130									5		4		8	4	8	9	7								
131	4		8	5	6	7	9	5	1				9	8	9	9	6			8	2	3	9	9	2
132	5	4	8	5	9	8	9	8	1		2		9	5	9	9	5			8		9	9	5	2

Ex. No.	Soil drench 10 Kg/ha								Dosage kg/ha	Foliar Spray								Pre-emergence							
	Z	R	G	O	L	M	B	S		Z	R	G	O	L	M	B	S	Z	R	G	O	L	M	B	S
133	7	6	8		6	4	9	8	5 1		4	9 8	3	7 5	9 8	9 9	8 5			9 8	2	6 2	9 8	8 6	2
134									5 1					6 6	9 9	8 4									
135						4	5		5 1		5 4	9 9	4 3	8 7	9 8	8 7						6 4	9 7		
136					3	6	8		5 1		6 4	8 8	6 5	9 7	9 8	7 7						7 4	6 2		
137			4	5	4	7	6		5 1		3 5	9 2	6 5	8 7	9 8	9 5						6 4	2		
138					5	4	6		5 1		4 3	9 2	6 5	7 7	9 8	7 6						8 4	8 7		
139			4		6	7	4		5 1		4 3	8 3	5 5	7 4	9 8	7 5						6 2			
140	*	*	*	*	*	*	*	*	5 1		3 3	4 3	5 5	6 5	8 7	9 5	6 2					6 5	6 2		
141					5	7			5 1			9 9	7 5	9 7	9 8	7 6						6 4	3		
142	6		7	7	9	9	9	6	5 1		3		5 2	7 6	9 8	7 6									
143			6	4	6	8	6	5	5 1			8 7	4	7 6	8 8	9 8	2			4		8 1	9 4		



Ex. No.	Soil drench 10 Kg/ha								Dosage kg/ha	Foliar Spray								Pre-emergence							
	Z	R	G	O	L	M	B	S		Z	R	G	O	L	M	B	S	Z	R	G	O	L	M	B	S
144	7	3	8	7	4	9	9	8	5 1	6		9 8	6 5	8 7	9 8	9 9	7 6			9 9	6 2	5			
145	4		8	7	6	9	9	5	5 1	2		9 8	6 2	8 6	9 9	5 4			8 6	2	1		8 2	9 2	
146									5 1	5 2		8 4	3	8 8	9 9	6 5									
147									5 1	1		4	3	5 2	8 6	4 7			2				4 3		
148				1	3	2	8		5 1		2	7 4	3	5 5	9 7	8 8	2								
149	7	6	8	5	3	9	8	8	5 1		3	9 5	4 1	6 6	9 8	8 5				7	1	2	7 2	3 3	
150	7	7	9	8	4	9	9	9	5 1			9 9	3 2	7 6	9 9	8 8				2 5	4	8 7	8 2		
151									5 1			4 2			5 2	6 2									
152	*	*	*	*	*	*	*	*	5 1			7			6 2	5 2									
153									5 1			1			4 2										
154									5 1	4 2		6			2		2								

This Sheet Has Been Replaced



Ex. No.	Soil drench 10 Kg/ha								Dosage kg/ha	Foliar Spray								Pre-emergence							
	Z	R	G	O	L	M	B	S		Z	R	G	O	L	M	B	S	Z	R	G	O	L	M	B	S
174									5 1							6 6	5 2								

This Sheet Has Been Replaced



Ex. No.	Dosage	Pre-Emergence														
		T R Z W A	H O R V W	Z E A M X	H E L A N	A L O M Y	S E T V I	G A L A P	S T E M E	V E R P E	C H E A L	M A T I N	I P O H E	A B U T H	A M B E L	A M A R E
193	0.8 0.4									1 1						
194	0.8 0.4				1 1				1	1 1						
195	0.8 0.4									2 2					2 2	
196	0.8 0.4						2		4 1	8 6	5 5	5 1				4 2
197	0.8 0.4								6	4	2 1	2			1 1	2 1
198	0.8 0.4			1	1				9 8	6 5	8 8	9 8	1	8 3	7 4	6 4
199	0.8 0.4			1					1		2 2	5	1		1	2 1
200	0.8 0.4			1	2 1				7	7 6	8 2	5		3	2 1	7 3
201	0.8 0.4			1					5 2	4	7 7	7				5 3
202	0.8 0.4						2		8 5	9 3	4 4	6 3			2	8 7
203	0.8 0.4				2				4 3	4	7 7	8 6		1	2 1	6 2
204	0.8 0.4				2 1	4 1	3 1		9 8	8 7	8 8	9 9	3	8 8	8 7	8 5

This Sheet Has Been Replaced





Ex. No.	Soil drench 10 kg/ha								Dosage kg/ha	Foliar Spray								Pre-emergence.							
	Z	R	G	O	L	M	B	S		Z	R	G	O	L	M	B	S	Z	R	G	O	L	M	B	S
206	0	0	3	2	2	9	7	4	5	0	0	7	0	3	8	8	3	0	0	0	0	0	5	4	2
207	0	0	0	0	0	7	2	0	1	1	0	4	1	0	8	9	2	0	0	0	0	0	0	0	0
217	3	0	7	7	9	9	9	1	5	3	3	9	3	8	9	9	4	1	0	0	0	0	5	8	0
218	0	0	0	0	0	6	5	1	1	3	1	5	0	3	8	8	2	0	0	0	0	0	0	0	0
219	2	0	5	2	5	8	9	2	5	4	0	7	5	8	8	9	6	4	0	0	0	0	9	8	0
229	0	0	0	0	0	8	3	0	1	0	0	7	0	5	7	7	0	0	0	0	0	0	0	0	0
230	0	0	4	5	6	7	5	0	1	4	0	5	2	6	7	8	2	0	0	5	0	0	6	8	0
231	2	1	5	7	5	9	9	4	5	5	0	6	2	8	9	9	5	0	0	0	0	0	5	7	0
232	2	6	6	2	4	8	8	7	1	1	1	9	5	4	8	8	8	0	0	2	0	2	8	4	2
233	6	2	8	8	4	9	9	6	5	5	0	9	5	8	9	9	8	2	0	6	0	5	9	7	0
234	4	0	6	5	5	8	8	4	1	4	0	9	0	6	8	9	5	0	0	0	2	0	0	0	0
235	2	0	5	4	5	8	4	1	1	5	0	6	2	4	9	6	5	0	0	0	0	0	7	5	0
236	0	0	2	1	3	5	5	0	1	0	0	0	0	0	8	6	2	0	0	0	0	0	7	2	0

This Sheet Has Been Replaced



CLAIMS

1. A compound of the general formula



10

wherein A represents an optionally substituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl or aralkyl group, or a group of general formula $\text{X}-\text{Y}-(\text{CH}_2)_n-$ where X represents an optionally substituted alkyl or aryl group, n represents 1 or 2, and Y represents an oxygen atom or a group of general formula $-\text{S}(\text{O})_m-$ where m represents 0, 1 or 2; R^1 represents a hydrogen atom or an alkyl group; and R^2 represents a hydrogen atom or an optionally substituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, aralkyl or heteroaralkyl group.

2. A compound as claimed in Claim 1, wherein A represents an optionally substituted alkyl, cycloalkyl, alkenyl, phenyl, pyridyl or thienyl group, or a group of general formula $\text{X}-\text{S}(\text{O})_m-(\text{CH}_2)_n$ where X represents an optionally substituted alkyl or phenyl group, m represents 0, 1 or 2, and n represents 1 or 2, or a group of general formula $\text{X}-\text{O}-(\text{CH}_2)_n-$ where X represents an optionally substituted alkyl group and n represents 1 or 2.

30

3. A compound as claimed in Claim 2, wherein A represents an alkyl or haloalkyl group, or an alkenyl group, an optionally substituted phenyl group, or a pyridyl group or a thienyl group.

35

4. A compound as claimed in Claim 3, wherein A represents a C_{3-6} alkyl group, an allyl group, a phenyl



group which is unsubstituted, or substituted by one or two moieties independently selected from halogen atoms, and alkyl and haloalkyl groups, or a 2-thienyl, 2-pyridyl or 3-pyridyl group.

5

5. A compound as claimed in any preceding claim, wherein R^1 represents a hydrogen atom.

10

6. A compound as claimed in any preceding claim, wherein R^2 represents an optionally substituted alkyl group, an alkenyl group, or an optionally substituted aryl, aralkyl or heteroaralkyl group.

15


7. A compound as claimed in Claim 6, wherein R^2 represents a C_{1-8} alkyl group, unsubstituted, or substituted by C_{3-6} cycloalkyl, or by C_{1-4} alkoxy or propargyl; or an allyl group; or by a phenyl group which is optionally substituted by 1 or 2 moieties independently selected from halogen atoms or C_{1-4} alkyl groups; or a group of general formula R^4-C_{1-4} alkylene where R^4 represents a pyridyl, furyl or thienyl group, or a phenyl group which is optionally substituted by one or two moieties independently selected from halogen and C_{1-4} alkyl.

20


25

8. A process for the preparation of a compound of general formula I or II, which comprises reacting a respective compound of the general formula III or IV

30



(III)



(IV)

35

where X represents a halogen atom or a C_{1-4} alkoxy group, with a compound of general formula HNR^1R^2 (V), where A, R^1 and R^2 are as defined in any preceding claim.



9. A compound of general formula I or II as defined by any of Claims 1 to 7, when prepared by a process as claimed in Claim 8.

5 10. A compound of general formula III or IV per se, as defined by Claim 8.

10 11. A process for the preparation of an ester compound of general formula III, as claimed in Claim 10, which comprises treating a compound of general formula $A-CO-NH-CH_2-CN$ (VI) with a compound of general formula R^5-ONO (VII) where R^5 is an alkyl group, and with an alkanol XOH .

15 12. A process for the preparation of an ester compound of general formula IV, as claimed in Claim 10, which comprises reacting an oxime amide $A-C(NH_2)=NOH$ (IX) with a compound of general formula $HalCOCOOR$ (X) where Hal represents a halogen atom.

20 13. An ester compound of general formula III, as defined by Claim 8, when prepared by a process as claimed in Claim 11.

25 14. An ester compound of general formula IV, as defined by Claim 8, when prepared by a process as claimed in Claim 12.

30 15. A process for the preparation of an acid halide compound of general formula III or IV, as claimed in Claim 10, by derivatising an ester compound of general formula III or IV, as claimed in Claim 10, 13 or 14.

35 16. An acid halide compound of general formula III or IV, as defined by Claim 8, when prepared by a process as claimed in Claim 15.



17. A herbicidal composition comprising a compound of general formula I or II as claimed in any of Claims 1 to 7, or Claim 9, together with a carrier.
- 5 18. A composition as claimed in Claim 17, comprising at least two carriers, at least one of which is a surface-active agent.
- 10 19. A method of combating undesired plant growth at a locus, comprising application to the locus of a compound of general formula I or II as claimed in any of Claims 1 to 7, or 9, or of a composition as claimed in Claim 17 or 18.
- 15 20. Use of a compound of general formula I or II, as claimed in any of Claims 1 to 7, or 9, or of a composition as claimed in Claim 17 or 18, as a herbicide.
- 20 21. A herbicidal composition comprising a compound of general formula III or IV as claimed in any of Claims 10, 13, 14 and 16, together with a carrier.
- 25 22. A composition as claimed in any of Claim 21 comprising at least two carriers, at least one of which is a surface-active agent.
- 30 23. A method of combating undesired plant growth at a locus, comprising application to the locus of a compound of general formula III or IV as claimed in any of Claims 10, 13, 14 and 16.
- 35 24. Use of a compound of general formula III or IV, as claimed in any of Claims 10, 13, 14 and 16, as a herbicide.
25. A compound of general formula I, II, III or IV; or a process for its preparation; or a compound so prepared; or



a herbicidal method, composition or use involving such a compound; any such aspect being substantially as hereinbefore described or defined with particular reference to the Examples.

