

This Sheet Has Been Replaced



The present invention relates to a new group of compounds with protected hydroxy groups which are precursors for organoleptic compounds (such as fragrances and masking agents) and antimicrobial compounds.

5

A principal strategy currently employed in imparting odours to consumer products is the admixing of the fragrance directly into the product. There are, however, several drawbacks to this strategy. The fragrance material
10 can be too volatile, resulting in fragrance loss during manufacturing, storage, and use. Many fragrance materials are also unstable over time. This again results in loss during storage.

15 In many consumer products it is desirable for the fragrance to be released slowly over time. Microencapsulation and inclusion complexes with cyclodextrins have been used to help decrease volatility, improve stability and provide slow-release properties. However, these methods are for a
20 number of reasons often not successful. In addition, cyclodextrins can be too expensive.

Fragrance precursors for scenting fabrics being washed in the presence of lipase-containing detergents are described
25 in WO 95/04809. The fragrance precursors contained in the detergent and/or in the softener are cleaved by the lipase and a single odoriferous compound, either an odoriferous alcohol or aldehyde or ketone is yielded. Thereby a prolonged scenting effect on the fabric is obtained.

30

REPLACED SHEETS		SG
☒	Description : paged to paged	52
☒	Claim(s) : paged to paged	53
☐	Drawing(s) : paged to paged	
☐	Abstract : paged to paged	

One object of the present invention is to provide new precursors for compounds with different activities.

It is a preferred object of the present invention to
5 provide compounds cleaved under different activating conditions.


A further object of the invention is to provide new compounds which are stable under transport and storage
10 conditions.

A further object of the present invention is to provide precursor molecules supplying different active compounds simultaneously or successively.


15

The present invention relates to compounds of formula I

I



20 wherein Y is



m is an integer of 1 or greater

25 n is 1, 2 or 3;

This Sheet Has Been Replaced




R^1, R^2, R^3, R^4, R^5 and R^6 represent independently hydrogen, substituted or unsubstituted alkyl-, alkenyl-, alkynyl-, cycloalkyl-, cycloalkenyl- or aromatic radicals which can
5 additionally contain one or more

-O- and/or $\begin{array}{c} \text{O} \\ \parallel \\ -\text{C}- \end{array}$ groups;

whereby one or two rings can be built by the combination
10 of the respective R^1 to R^6 and said ring(s) can be substituted by one or more alkyl group;

X is $-\text{OR}^7$ and R^7 is the residue of an alcohol $R^7\text{OH}$, or the residue of the enol form of an aldehyde or ketone, or X is
15 a primary, or secondary amino group forming an amide;


Z is



20 q is the same or bigger than m;

R^8 represents hydrogen, a straight or branched, unsub-
25 stituted or substituted alkyl-, alkenyl-, cycloalkyl-, cycloalkenyl- or aromatic radical which optionally comprises and/or is substituted by one or more heteroatoms,

This Sheet Has Been Replaced



and/or group(s) comprising a heteroatom preferably by
-CO-, OCOR⁷, COOR⁷, COY, Si and/or N;

5 R⁹ is the residue -OR¹² of an alcohol of formula R¹²OH or
the residue of the enol form of an aldehyde or ketone or
has the definition given for Y and R⁹ and Y can be the
same or different and optionally comprises and/or is sub-
stituted by a heteroatom, and/or group(s) comprising a
heteroatom;

10

R¹⁰ and R¹¹ represent independently hydrogen, substituted
or unsubstituted alkyl, alkenyl, cycloalkyl, cycloalkenyl
or an aromatic residue which optionally comprise and/or
are substituted by one or more heteroatoms, and/or
15 group(s) comprising a heteroatom preferred heteroatoms and
groups with heteroatoms for R⁹, R¹⁰ and R¹¹ are O, Si, N
and CO.

20 Further characteristics and advantages of the invention
are described by claims 2 to 11 and by the following
specification and examples.

The compounds of formula I are not limited to any
particular stereoisomers, all possible stereoisomers (E/Z
25 isomers, enantiomers, diastereomers) and all mixtures are
thus included within the scope of the invention.


The compounds of formula I are virtually odourless under
room temperature, atmospheric conditions and about 20 to
30 100% relative humidity. However, under activating condi-
tions, they are cleaved and one or more active compounds

This Sheet Has Been Replaced


with organoleptic and/or antimicrobial properties are generated.

The activating conditions which lead to cleavage and the
 5 desired active compounds comprise the presence of skin
 bacteria, especially axilla bacteria, of an enzyme such as
 protease or lipase, elevated temperature or acidic or
 alkaline pH-values and/or light. The compounds of formula
 I, upon cleavage, provide lactones and optionally
 10 aldehydes, ketones, and/or alcohols having organoleptic
 and/or antimicrobial activity and therefore permit the
 development of useful consumer products with enhanced
 organoleptic and/or microbiological properties. Further,
 the compounds of formula I, upon cleavage can generate
 15 fluorescent coumarins useful as optical brighteners.


The compounds I of the present invention are cleaved in
 under activating conditions in two successive steps, first
 the „protective group„ Z is removed resulting in a
 20 hydroxyester



which decomposes into one or more organoleptic lactone(s),
 25 and one or more alcohol(s), amine(s), aldehyde(s) and/or
 ketone(s).

This Sheet Has Been Replaced






5 Z may be considered a protective group which prevents the hydroxy ester Y-H from premature cyclisation to an organoleptic lactone. At the same time Z can generate one or more additional organoleptic compound(s). In the following, cleavage of a β -ketoester of formula I is shown.

10

This Sheet Has Been Replaced



This Sheet Has Been Replaced




- 1 is Z-Y, a protected hydroxy ester (β -ketoester)
- 2 is ZH, a hydroxyester
- 3 is an organoleptic lactone
- 4 is an organoleptic alcohol
- 5 5 is an β -ketoacid
- 6 is an organoleptic ketone

Thus one compound of formula I can yield under activating conditions three different organoleptic compounds.


10

If Z is



- 15 it stands preferably for the residue of an odourless or antimicrobial acid optionally substituted by groups yielding upon cleavage one or more organoleptic compounds. Examples of esters in which R^8 is an odourless acid are:
- tetradecanoic acid 1-(2-hex-3-enyloxycarbonyl-ethyl)-
- 20 heptyl ester;
- tetradecanoic acid 1-[2-(3,7-dimethyl-octa-2,6-dienyloxy-carbonyl)-ethyl]-octyl ester;
- benzoic acid 1-[2-(3,7-dimethyl-octa-2,6-dienyloxy-carbonyl)-ethyl]-octyl ester;
- 25 dodecanoic acid 1-[2-(3,7-dimethyl-oct-6-enyloxycarbonyl)-ethyl]-heptyl ester;
- succinic acid 1-[2-(1,5-dimethyl-1-vinyl-hex-4-enyloxy-carbonyl)-ethyl]-heptyl ester hex-3-enyl ester is a compound wherein R^8 is COR^7 ;


This Sheet Has Been Replaced



succinic acid bis-{1-[2-(3,7-dimethyl-octa-2,6-dienyloxy-carbonyl)-ethyl]-octyl} ester with R^8 being COY.

If Z is

5



R^9 is preferably the residue of an organoleptic compound or Y. Examples of carbonates in which R^9 is an organoleptic compound are:

10

4-phenethyloxycarbonyloxy-decanoic acid 3,7-dimethyl-oct-6-enyl-ester;

4-phenethyloxycarbonyloxy-decanoic acid hex-3-enyl ester;

15

4-hex-3-enyloxycarbonyloxy-undecanoic acid 3,7-dimethyl-octa-2,6-dienyl ester;


4-phenethyloxycarbonyloxy-undecanoic acid 3,7-dimethyl-octa-2,6-dienyl ester;

4-{1-[2-(1,1,5-trimethyl-hexyloxycarbonyl)-ethyl]octyloxy-carbonyloxy}-decanoic acid 1,1,5-trimethyl-hexyl ester is

20

a carbonate with R^9 being Y.


If Z stands for




25

This Sheet Has Been Replaced





is preferably derived from a non odorous mono- or diamine.
4-(Bis-decyl-carbamoyloxy)-undecanoic acid hex-3-enyl
5 ester is such a preferred carbamate.



10 can also be a photolabile ester.

The compounds of the present invention can act as fragrance precursors in personal care products, in laundry products, cleaning compositions, pet care products and environment scents such as air fresheners. They can also act as precursors for odour masking agents, e.g. in the same products as the fragrance precursors. They also can act as precursors for antimicrobial agents. The fragrance precursors and the precursors for odour masking agents of the invention may be used individually in an amount effective to enhance or to mask the characteristic odour of a material. More commonly, however, the compounds are mixed with other fragrance components in an amount sufficient to provide the desired odour characteristics.

25

Due to the in situ generation of the active compounds the desired effect is prolonged and the substantivity on

This Sheet Has Been Replaced




different substrates is enhanced. If two or more active compounds are provided, they can be generated, depending on the precursor and/or the activating conditions, simultaneously or successively. Further, the precursors of the
 5 invention provide slow release of the active compounds.

A broad range of known odorants or odorant mixtures can be generated from precursors of the invention.


10 Examples of aldehydes include:

- 2,6,10-trimethylundec-9-enal*
- 1,2,3,4,5,6,7,8-octahydro-8,8-dimethyl-2-naphthalenecarboxaldehyde
- 15 tridecanal
- 2-[4-(1-methylethyl)phenyl]-ethanal
- 2,4-dimethyl-cyclohex-3-ene-1-carbox-aldehyde*
- 4-carboxaldehyde-1,3,5-trimethyl-cyclohex-1-ene*
- 1-carboxaldehyde-2,4-dimethyl-cyclohex-3-ene*
- 20 1-carboxaldehyde-4-(4-hydroxy-4-methylpentyl)-cyclohex-3-ene*
- 3,5,5-trimethyl-hexanal
- heptanal*
- 2,6-dimethyl-hept-5-enal*
- 25 decanal**
- dec-9-enal
- dec-4-enal
- 2-methyldecanal*
- undec-10-enal**
- 30 undecanal*
- dodecanal**

This Sheet Has Been Replaced



- 2-methyl-undecanal**
 tridecanal
 octanal**
 nonanal*
- 5 3,5,5-trimethylhexanal
 undec-9-enal**
 2-phenyl-propanal*
 4-methyl-phenyl-acetaldehyde*
 3,7-dimethyl-octanal*
- 10 dihydrofarnesal**
 7-hydroxy-3,7-dimethyl-octanal*
 2,6-dimethyl-oct-5-enal
 2-[4-(1-methylethyl)phenyl]-ethanal*
 3-(3-isopropyl-phenyl)-butanal**
- 15 2-(3,7-dimethyloct-6-enoxy)-ethanal
 1-carboxaldehyde-4-(4-methyl-3-pentenyl)-cyclohex-3-ene*
 2,3,5,5-tetramethyl-hexanal
 longifolic aldehyde
 2-methyl-4-(2,6,6-trimethylcyclohex-2-en-1-yl)-butanal*
- 20 2-methyl-3-(4-tert-butylphenyl)-propanal**
 4-(1,1-dimethyl-ethyl)-benzene-propanal*
 2-[4-(1-methylethyl)-phenyl]-propanal
 alpha-methyl-1,3-benzodioxole-5-propanal*
 3,7-dimethyl-oct-6-enal*
- 25 2-methyl-3-(4-isopropylphenyl)-propionaldehyde*
 4-(4-hydroxy-4-methyl-pentyl)-cyclohex-3-en-1-carboxaldehyde**
 alpha-methyl-1,3-benzodioxole-5-propanal*
 1-carboxaldehyde-4-(1,1-dimethylethyl)-cyclohexane
- 30 4-(octahydro-4,7-methano-5H-inden-5-ylidene)-butanal
 [(3,7-dimethyl-6-octenyl)-oxy]-acetaldehyde**




- hex-2-enal*
- 2-nonenal*
- 2-tridecenal*
- 3,7-dimethyl-oct-2,6-dien-1-al*
- 5 2-nonadienal*
- 2,4-dimethyl-2,6-heptadienal
- trans-dec-2-en-1-al*
- 2,4-diethyl-hep-2,6-dien-1-al*
- dodec-2-en-1-al*
- 10 3,7-dimethyl-oct-2,6-dien-1-al*
- 2,4-diethyl-hepta-2,6-dienal
- 3,7-dimethyl-nona-2,6-dien-1-al*
- 3-propyl-2-hepten-1-al
- 1-carboxaldehyde-4-(prop-2-en-2-yl)-cyclohex-1-ene
- 15 undecanal

whereby * indicates the preferred aldehydes and **
indicate the more preferred aldehydes.

- 20 Examples of ketones include:


- 2-heptyl-cyclopentanone
- 2,2,6,10-tetramethyltricyclo-[5.4.0.0(6,10)]-undecan-4-one
- benzylacetone*
- 25 carvone*;
- 1,2,3,5,6,7-hexahydro-1,1,2,3,3,-pentamethyl-4H-inden-4-one*
- methyl heptenone*
- dimethyl octenone*
- 30 2,5-dimethyl-oct-2-en-6-one**
- 2-(butan-2-yl)-cyclohexanone*

This Sheet Has Been Replaced



- 2-hexyl-cyclopent-2-en-1-one*
- 2-(1-methylethyl)-5-methyl-cyclohexanone*
- 2-(2-methylethyl)-5-methyl-cyclohexanone**
- 3-methyl-cyclopentadecanone
- 5 4-(1,1-dimethylpropyl)pentyl-cyclohexanone*
- 4-tert-pentyl-cyclohexanone*
- 2-oxo-1-pentyl-cyclopentane-acetic acid methyl ester**
- 3-oxo-2-pentyl-cyclopentane-acetic acid methyl ester**
- 1-(1,2,3,4,5,6,7,8-octahydro-2,3,8,8-tetramethyl-2-
- 10 naphthalenyl)-ethanone*
- 3-methyl-5-propyl-cyclohex-2-en-1-one*
- 4-(2,6,6-trimethylcyclohex-1-en-1-yl)butan-2-one**
- 4-(2,6,6-trimethylcyclohex-2-en-1-yl)butan-2-one**
- 2-methyl-5-(1-methylethenyl)-cyclohex-2-en-1-one*
- 15 cyclopentadecanone**
- 1-(4-hydroxyphenyl)-butan-3-one**
- 4-benzo-1,3-dioxo-5-yl-but-2-one**
- 4-(1,3-benzodioxol-5-yl)-2-butanone**
- nonan-3-one*
- 20 nonan-2-one*
- octan-2-one*
- 2-heptanone*
- butan-2-one*
- 6-methyl-hept-5-en-2-one*
- 25 6,10-dimethyl-undeca-5,9-dien-2-one*
- 1-(2,4,4-trimethyl-2-cyclohexen-1-yl)-2-buten-1-one*
- 1-(2-cyclohexen)-2,4,4-trimethyl-but-2-enone*
- carvone**
- 2-hexyl-cyclo-pent-2-en-1-one**
- 30 2-pentyl-cyclopent-2-en-1-one
- 3-methyl-2-pentyl-cyclopent-2-en-1-one**

This Sheet Has Been Replaced



- 2-hexylidenecyclopentanone*
- 3,5-diethyl-5,6-dimethyl-2-cyclohexenone*
- 4,4A,5,6,7,8-hexahydro-6-isopropenyl-4,4A-dimethyl-2(3H)-naphthalenone**
- 5 3-methyl-6-propylidenecyclohexanone*
- 4-(1-methylethyl)-cyclohex-2-en-1-one
(E)-oct-3-en-2-one
- 1-(2,3,4,7,8,8A-hexahydro-3,6,8,8-tetramethyl-1H-3A,7-methanoazulen-5-yl)-ethanone*
- 10 2-hydroxy-3,5-dimethyl-cyclopent-2-ene-1-one*
- 1-(3,3-dimethyl-1-cyclohexen-1-yl)ethanone*
- 1-(2,4,6-trimethylcyclohex-3-en-1-yl)-but-1-en-3-one
acetylisolongifolene
- 2-(3-methylbut-2-en-1-yl)-3-methyl-cyclopent-2-en-1-one
- 15 2,6,6-trimethyl-1,3-cyclohexadienyl-1-carbaldehyde**
- 3-methyl-5-(2,2,3-trimethylcyclopent-3-ene-1-yl)pent-3-ene-2-one*
- 5-butylidene-2,2,4-trimethylcyclopentanone
- 1,2,3,5,6,7-hexahydro-1,1,2,3,3-pentamethyl-4H-inden-4-one**
- 20 3-methyl-5-propyl-cyclohex-2-en-1-one**
- 4,4A,5,6,7,8-hexahydro-6-isopropyl-2(3H)-naphthalenone
- 4-(2,6,6-trimethyl-1-cyclohexen-1-yl)-butan-2-one**
- 4-methoxyphenylethanone**
- 25 acetophenone*
- 1-(2-naphthalenyl)-ethanone**
- 3-methyl-4-(2,6,6-trimethyl-2-cyclohexen-1-yl)-3-buten-2-one**
- 2-acetylpyrazine*
- 30 3,5,5-trimethyl-cyclohex-2-en-1,4-dione*
- (E)-5-methyl-2-hepten-4-one

This Sheet Has Been Replaced



acetyl diisoamylene**

dec-3-en-2-one

2-ethyl-3,6,6-trimethylcyclohex-2-enyl-but-2-en-1-one

1-(5,5-dimethyl-1(6)-cyclohexen-1-yl)-4-penten-1-one**

5 1-(2,6,6-trimethyl-1-cyclohexen-1-yl)-but-2-ene-1-one**

1-(2,6,6-trimethyl-2-cyclohexen-1-yl)-but-2-ene-1-one**

1-(2,6,6-trimethyl-3-cyclohexen-1-yl)-but-2-ene-1-one**

2,4,4,5,5-pentamethyl-1-cyclopentene-1-yl-ethanone*

10 whereby * indicates the preferred ketones and ** indicate the more preferred ketones.

Examples of alcohols are primary, secondary and tertiary alcohols and phenols such as:

15

amyl alcohol

hexyl alcohol*

2-hexyl alcohol*

heptyl alcohol*

20 octyl alcohol*

nonyl alcohol*

decyl alcohol*

undecyl alcohol*

lauryl alcohol*

25 myristic alcohol

3-methyl-but-2-en-1-ol*

3-methyl-1-pentanol

cis-3-hexenol*

cis-4-hexenol*

30 3,5,5-trimethyl-hexanol


3,4,5,6,6-pentamethylheptan-2-ol*

This Sheet Has Been Replaced



- citronellol*
- geraniol*
- oct-1-en-3-ol
- 2,5,7-trimethyl-octan-3-ol
- 5 2-cis-3,7-dimethyl-2,6-octadien-1-ol
- 6-ethyl-3-methyl-5-octen-1-ol*
- 3,7-dimethyl-oct-3,6-dienol*
- 3,7-dimethyloctanol*
- 7-methoxy-3,7-dimethyl-octan-2-ol*
- 10 cis-6-nonenol*
- 5-ethyl-2-nonanol
- 6,8-dimethyl-2-nonanol*
- 2,2,8-trimethyl-7(8)-nonene-3-ol
- nona-2,6-dien-1-ol
- 15 4-methyl-3-decen-5-ol*
- dec-9-en-1-ol
- benzylalcohol
- 2-methyl-undecanol
- 10-undecen-1-ol
- 20 1-phenyl-ethanol*
- 2-phenyl-ethanol*
- 2-methyl-3-phenyl-3-propenol
- 2-phenyl-propanol*
- 3-phenyl-propanol*
- 25 4-phenyl-2-butanol
- 2-methyl-5-phenyl-pentanol*
- 2-methyl-4-phenyl-pentanol*
- 3-methyl-5-phenyl-pentanol*
- 2-(2-methylphenyl)-ethanol*
- 30 4-(1-methylethyl)-benzene-methanol
- 4-(4-hydroxyphenyl)-butan-2-one*

This Sheet Has Been Replaced



- 2-phenoxy-ethanol*
4-(1-methylethyl)-2-hydroxy-1-methyl benzene
2-methoxy-4-methyl-phenol
4-methyl-phenol
5 anisic alcohol*
p-tolyl alcohol*
cinnamic alcohol*
vanillin*
ethyl vanillin*
10 eugenol*
isoeugenol*
thymol
anethol*
decahydro-2-naphthalenol
15 borneol*
cedrenol*
farnesol*
fenchyl alcohol*
menthol*
20 3,7,11-trimethyl-2,6,10-dodecatrien-1-ol
alpha ionol*
tetrahydro ionol*
2-(1,1-dimethylethyl)cyclohexanol*
3-(1,1-dimethylethyl)cyclohexanol*
25 4-(1,1-dimethylethyl)cyclohexanol*
4-isopropyl-cyclohexanol
6,6-dimethyl-bicyclo[3.3.1]hept-2-ene-2-ethanol
6,6-dimethyl-bicyclo[3.1.1]hept-2-ene-methanol*
p-menth-8-en-3-ol*
30 3,3,5-trimethyl-cyclohexanol
2,4,6-trimethyl-3-cyclohexenyl-methanol*



- 4-(1-methylethyl)-cyclohexyl-methanol*
- 4-(1,1-dimethylethyl)-cyclohexanol
- 2-(1,1-dimethylethyl)-cyclohexanol
- 2,2,6-trimethyl- α -propyl-cyclohexane propanol*
- 5 5-(2,2,3-trimethyl-3-cyclopentenyl)-3-methylpentan-2-ol*
- 3-methyl-5-(2,2,3-trimethylcyclopentyl-3-enyl)pent-4-en-2-ol*
- 2-ethyl-4(2,2,3-trimethylcyclopentyl-3-enyl)but-2-en-1-ol*
- 4-(5,5,6-trimethylbicyclo[2.2.1]hept-2-yl)-cyclohexanol*
- 10 2-(2-methylpropyl)-4-hydroxy-4-methyl-tetrahydropyran*
- 2-cyclohexyl-propanol*
- 2-(1,1-dimethylethyl)-4-methyl-cyclohexanol*
- 1-(2-tert-butyl-cyclohexyloxy)-2-butanol*
- 1-(4-isopropyl-cyclohexyl)-ethanol*
- 15 1-(4-hydroxyphenyl)-butan-3-one
- 2,6-dimethyl-oct-7-en-2-ol**
- 2,6-dimethyl-heptan-2-ol**
- 3,7-dimethyl-octa-1,6-dien-3-ol**
- 20 * indicates preferred alcohols and ** indicate the more preferred alcohols.

Examples of lactones include:

25

6-methyl-pyran-2-one

5-heptyldihydro-2(3H)-furanone*

5-pentyldihydro-2(3H)-furanone*

5-(3-hexenyl)-dihydro-5-methyl-(Z)-2(3H)-furanone

30

5-hexyldihydro-5-methyl-2(3H)-furanone

5-hexyldihydro-2(3H)-furanone*

This Sheet Has Been Replaced



- 5-octyldihydro-2(3H)-furanone
 8-(1-methylethyl)-1-oxaspiro[4.5]-decan-2-one*
 8-methyl-1-oxaspiro[4.5]decan-2-one
 8-ethyl-1-oxaspiro[4.5]decan-2-one
 5 5-(1,5-dimethyl-4-hexenyl)-dihydro-2(3H)-furanone
 2-oxo-5-butyl-tetrahydrofuran*
 4-methyl-5-pentyl-dihydro-2(3H)-furan-2-one
 5-hexyldihydro-5-methyl-2(3H)-furanone
 dihydro-5-methyl-5-vinyl-2(3H)-furanone
 10 octahydro-2H-1-benzopyran-2-one
 tetrahydro-6-pentyl-2H-pyran-2-one
 tetrahydro-6-hexyl-2H-pyran-2-one
 tetrahydro-6-heptyl-2H-pyran-2-one
 tetrahydro-6-(3-pentenyl)-(E)-2H-pyran-2-one
 15 tetrahydro-6-(2-pentenyl)-(Z)-2H-pyran-2-one
 (E)-oxacycloheptadec-10-en-one**
 oxacyclohexadecan-2-one**
 dodeca-12-olide

20 whereby * indicates the preferred lactones.

It is a matter of course, that it is not possible to give a complete list of the organoleptic especially odoriferous and/or antimicrobial aldehydes, ketones, alcohols and
 25 lactones which are generated as a result of the desired cleavage of the esters of formula I by skin bacteria, by enzymes, by elevated temperatures, by acidic and/or alkaline pH-values or by light. The skilled person is, however, quite aware of those aldehydes, ketones, alcohols
 30 and lactones which provide the desired organoleptic, e.g. fragrance and odour masking and/or antimicrobial effects.

This Sheet Has Been Replaced



While manufacturing compositions, the precursors of the invention may be used according to methods known to the perfumer, such as e.g. from W.A. Poucher, Perfumes, Cosmetics, Soaps, 2, 7th Edition, Chapman and Hall, London
5 1974.

The compounds of formula I may preferably be used as sustained release odorants but also to mask or attenuate undesirable odours or to provide additional odours not
10 initially present in consumer products, i.e. personal care products such as cosmetic products destined for application to human skin such as underarm deodorants or antiperspirants or other deodorants contacting the body, or in hand lotions, baby powders, baby lotions, ointments, foot
15 products, facial cleansers, body wipes, facial make-up, colognes, after-shave lotions, shaving creams, etc. Additional applications include laundry detergents, fabric softeners, fabric softener sheets, (automatic) dishwasher detergents, and other enzyme-containing consumer products.
20 Further applications are air fresheners and odorants, odour masking agents and/or antimicrobial agents.

The amount required to produce the desired, overall effect varies depending upon the particular compounds of formula
25 I chosen, the product in which it will be used, and the particular effect desired.

For example, depending upon the selection and concentration of the compound chosen, when a compound of the
30 formula I is added either singly or as a mixture, e.g. to a deodorant or laundry product composition at levels

ranging from about 0.1 to about 10 % by weight, or most preferred about 0.25 to about 4 % by weight, an odorant, i.e. an odoriferous, aldehyde, ketone, alcohol or lactone in an „organoleptically effective amount,, is released when
 5 the product is used. This newly formed odorant serves to enhance the odour of the product itself or of a fragrance present in the product.

Compounds of formula I can be prepared by using a wide
 10 variety of methods known to the skilled chemist.

For example, for the synthesis of esters see Comprehensive Organic Chemistry, vol 2, D. Barton, W.D. Ollis, Pergamon Press, p. 871.

For example, for the synthesis of carbonates see
 15 Comprehensive Organic Chemistry, vol 2, D. Barton, W.D. Ollis, Pergamon Press, p. 1070.

For example, for the synthesis of carbamates see Comprehensive Organic Chemistry, vol 2, D. Barton, W.D. Ollis, Pergamon Press, p. 1083.


20 Convenient methods are outlined in the Examples without limiting the invention thereto.

Example 1

Acetic acid 3-(4-tert-butyl-phenyl)-2-methyl-propenyl
 25 ester

A solution of 200 g 2-methyl-3-(4-tert-butylphenyl)-propanal, 280 ml triethylamine and 13.4 g sodium acetate in 800 ml of acetic anhydride was stirred at 120°C for 5.5 hours. Then the solution was cooled, water was added and
 30 the water phase was extracted with hexane. The organic phase was washed with 2N NaOH and water to neutrality,

This Sheet Has Been Replaced



dried and evaporated to dryness. The residue was distilled to yield 185 g of a colourless liquid.

NMR (CDCl₃):

- 5 δ 7.35-6.97 (m, 5H), 3.43+3.21 (s, 2H, E/Z), 2.13 (s, 3H), 1.60 (s, 3H), 1.30 (s, 9H) ppm.

Example 2

Acetic acid 3-(3-isopropyl-phenyl)-but-1-enyl ester

- 10 According to the procedure of example 1, acetic acid 3-(3-isopropyl-phenyl)-but-1-enyl ester was prepared from 3-(3-isopropylphenyl)-butanal, acetic anhydride, sodium acetate and triethylamine.

15 Example 3

4-Oxo-undecanoyl chloride

- To a suspension of 50 g 4-oxo-undecanoic acid (Synthesis, 1987, 408) in 350 ml of ether, 22.15 g pyridine was dropped in at 0°C. Then a solution of 32.72 g thionyl chloride in 50 ml of ether was dropped in at 0-5°C and then the reaction mixture was stirred for 20 hours at room temperature. Then the reaction mixture was filtered and evaporated to dryness. The residue (53.83 g yellow oil) was not further purified.


25

Example 4

4-Oxo-decanoic acid 3,7-dimethyl-oct-6-enyl ester

- A solution of 20.0 g 4-oxo-1-decanoic acid (Synthesis, 1987, 408), 16.8 g citronellol extra and 0.5 g p-toluenesulfonic acid in 150 ml of cyclohexane was refluxed in a

This Sheet Has Been Replaced



flask equipped with a Dean-Stark trap for 3 hours. Then the reaction mixture was cooled, diluted with ether, washed with saturated NaHCO_3 and water. The organic phase was dried, filtered and evaporated to dryness. The
 5 resulting oil was purified by chromatography to yield 31.7 g of a yellow oil.

NMR (CDCl_3):

δ 5.04-5.19 (m, 1H), 4.11-4.17 (m, 2H), 2.68-2.79 (m, 2H),
 10 2.53-2.60 (m, 2H), 2.41-2.48 (t, 2H), 2.03-1.91 (q, 2H),
 1.75-1.44 (m, 6H), 1.43-1.08 (m, 12H), 0.92-0.84 (m, 6H)
 ppm.

Example 5

15 4-Oxo-decanoic acid hex-3-enyl ester

According to the procedure of Example 4, 4-oxo-decanoic acid hex-3-enyl ester was prepared from 4-oxo-decanoic acid, cis-3-hexenol and p-toluenesulfonic acid.

20 Example 6

4-Oxo-undecanoic acid hex-3-enyl ester

According to the procedure of Example 4, 4-oxo-undecanoic acid hex-3-enyl ester was prepared from 4-oxo-undecanoic acid, cis-3-hexenol and p-toluenesulfonic acid.


25

Example 7

4-Oxo-nonanoic acid hex-3-enyl ester

According to the procedure of Example 4, 4-oxo-nonanoic acid hex-3-enyl ester was prepared from 4-oxo-nonanoic
 30 acid (Synthesis, 1987, 408), cis-3-hexenol and p-toluene-sulfonic acid.

This Sheet Has Been Replaced



Example 84-Oxo-nonanoic acid 3,7-dimethyl-octa-2,6-dienyl ester

5 A solution of 20 g 4-oxo-nonanoic acid, 19.3 g geraniol,
 26.8 g N,N-dicyclohexyl-carbodiimide and 1.0 g 4-
 pyrrolidinopyridine in 300 ml of dichloromethane was
 stirred for 24 hours at room temperature. The precipitate
 was filtered off, the filtrate was diluted with ether,
 10 washed with aqueous hydrochloric acid, saturated NaHCO₃
 and brine. The organic phase was dried, filtered and
 evaporated to dryness. The resulting oil-crystal mixture
 was purified by chromatography to yield 21.4 g of a
 colourless oil.

15

NMR (CDCl₃):

δ 5.32 (t, 1H), 5.09 (m, 1H), 4.10 (d, 2H), 2.71 (m, 2H),
 2.52 (m, 2H), 2.43 (t, 2H), 2.07 (m, 4H), 1.77-1.50 (m,
 11H), 1.29 (m, 4H), 0.91 (t, 3H) ppm.


20

Example 94-Oxo-decanoic acid 1,5-dimethyl-1-vinyl-hex-4-enyl ester

According to the procedure of example 8, 4-oxo-decanoic
 acid 1,5-dimethyl-1-vinyl-hex-4-enyl ester was prepared
 25 from 4-oxo-decanoic acid and (±)-linalool.

Example 104-Oxo-decanoic acid 1,1,5-trimethyl-hexyl ester

According to the procedure of example 8, 4-oxo-decanoic
 acid 1,1,5-trimethyl-hexyl ester was prepared from 4-oxo-
 30 decanoic acid and 2,6-dimethyl-heptan-2-ol.



Example 114-Oxo-undecanoic acid 2-benzyloxycarbonyl-2-benzyloxy
carboxylamino-ethyl ester

- 5 According to the procedure of Example 8, 4-oxo-undecanoic
acid 2-benzyloxycarbonyl-2-benzyloxycarboxylamino-ethyl
ester was prepared from 4-oxo-undecanoic acid, 2-
benzyloxycarboxylamino-3-hydroxy propionic acid benzyl
ester, N,N'-dicyclohexyl-carbodiimide and dimethylamino-
10 pyridine.

Example 124-Oxo-undecanoic acid 3-(4-tert-butyl-phenyl)-2-methyl-
propenyl ester

- 15 To a solution of 43.79 g acetic acid 3-(4-tert-butyl-
phenyl)-2-methyl-propenyl ester in 200 ml of
tetrahydrofuran, a solution of 27.16 g potassium tert-
butoxide in 200 ml of tetrahydrofuran was dropped in at -
78°C. After stirring at this temperature for 90 minutes, a
20 solution of 53.00 g 4-oxo-undecanoyl chloride in 200 ml of
tetrahydrofuran was dropped in. After stirring at -78°C
for 2.5 hours, the solution was quenched with saturated
sodium bicarbonate solution and diluted with ether. The
organic phase was washed with saturated sodium bicarbonate
25 solution and brine, dried and evaporated to dryness. The
residue was purified by thin-film distillation and
chromatography to yield 25.73 g of a yellow oil.

NMR (CDCl₃):

This Sheet Has Been Replaced



δ 7.37-6.93 (m, 5H), 3.42+3.22 (s, 2H, E/Z), 2.82-2.61 (m, 4H), 2.52-2.38 (m, 2H), 1.70-1.48 (m, 5H), 1.47-1.15 (m, 17H), 0.98-0.80 (t, 3H) ppm.

5 Example 13

4-Oxo-undecanoic acid 3-(3-isopropyl-phenyl)-but-1-enyl ester

According to the procedure of example 12, 4-oxo-undecanoic acid 3-(3-isopropyl-phenyl)-but-1-enyl ester was prepared from acetic acid 3-(3-isopropyl-phenyl)-but-1-enyl ester, 4-oxo-undecanoyl chloride and potassium tert-butoxide.

Example 14

4-Hydroxy-decanoic acid 3,7-dimethyl-oct-6-enyl ester

15 A solution of 2.0 g sodium borohydride in 30 ml of water was cooled to 5°C. A solution of 4-oxo-decanoic acid 3,7-dimethyl-oct-6-enyl ester in 75 ml of THF was added to the reaction during 12 minutes and the resulting reaction mixture was stirred at room temperature for 5 hours. Then 20 the reaction mixture was diluted with ether, washed with saturated NaHCO₃, brine and water. The organic phase was dried, filtered and evaporated to dryness. The resulting liquid was purified by chromatography to yield 7.6 g of a liquid.

25

NMR (CDCl₃):

δ 5.09 (bt, 1H), 4.11 (t, 2H), 3.61 (m, 1H), 2.42 (t, 2H), 2.07 (m, 2H), 2.12-1.02 (m, 22H), 0.88 (m, 6H) ppm.

30 Example 15

4-Hydroxy-nonanoic acid hex-3-enyl ester

This Sheet Has Been Replaced



According to the procedure of Example 14, 4-hydroxy-nonanoic acid hex-3-enyl ester was prepared from 4-oxo-nonanoic acid hex-3-enyl ester, sodium borohydride and water.

5

Example 16

4-Hydroxy-decanoic acid hex-3-enyl ester

According to the procedure of Example 14, 4-hydroxy-decanoic acid hex-3-enyl ester was prepared from 4-oxo-decanoic acid hex-3-enyl ester, sodium borohydride and water.

10

Example 17

4-Hydroxy-decanoic acid 1,5-dimethyl-1-vinyl-hex-4-enyl ester

15

According to the procedure of Example 14, 4-hydroxy-decanoic acid 1,5-dimethyl-1-vinyl-hex-4-enyl ester was prepared from 4-oxo-decanoic acid 1,5-dimethyl-1-vinyl-hex-4-enyl ester, sodium borohydride and water.

20

Example 18

4-Hydroxy-undecanoic acid hex-3-enyl ester

According to the procedure of Example 14, 4-hydroxy-undecanoic acid hex-3-enyl ester was prepared from 4-oxo-undecanoic acid hex-3-enyl ester, sodium borohydride and water.

25

Example 19

4-Hydroxy-nonanoic acid 3,7-dimethyl-octa-2,6-dienyl ester

According to the procedure of Example 14, 4-hydroxy-nonanoic acid 3,7-dimethyl-octa-2,6-dienyl ester was

30

This Sheet Has Been Replaced



prepared from 4-oxo-nonanoic acid 3,7-dimethyl-octa-2,6-dienyl ester, sodium borohydride and water.

Example 20

5 4-Hydroxy-decanoic acid 1,1,5-trimethyl-hexyl ester

According to the procedure of Example 14, 4-hydroxy-decanoic acid 1,1,5-trimethyl-hexyl ester was prepared from 4-oxo-decanoic acid 1,1,5-trimethyl-hexyl ester, sodium borohydride and water.

10

Example 21

4-Hydroxy-undecanoic acid 3,7-dimethyl-octa-2,6-dienyl ester

15 4-Hydroxy-undecanoic acid sodium salt

To a solution of 43.6 g sodium hydroxide in 150 ml of methanol heated to reflux, 200 g gamma-undecalactone were dropped in. After stirring for 2 hours at reflux, the mixture was cooled to room temperature and evaporated to dryness. The resulting crystals were washed with hexane to yield 240 g colourless crystals.

NMR (CDCl₃):

25 δ 5.1-4.8 (br s, OH), 3.63-3.42 (m, 1H), 2.39-2.20 (t, 2H), 1.89-1.52 (m, 2H), 1.51-1.15 (m, 12H), 1.00-0.81 (t, 3H) ppm.

1-Chloro-3,7-dimethyl-octa-2,6-diene

30 To a mixture of 170 g linalool and 20 mg bismuth(III)-oxide heated to 60 °C, 130.5 g trimethylchlorosilane were dropped in. Then the mixture was cooled to room tempe-

This Sheet Has Been Replaced



rature and the organic layer was separated. The resulting oil was purified by distillation to yield 158.35 g of a colourless oil.

5 NMR (CDCl₃):

δ 5.56-5.35 (t, 1H), 5.18-4.99 (m, 1H), 4.16-4.02 (d, 2H), 2.26-1.91 (m, 4H), 1.89-1.45 (m, 9H) ppm.

4-Hydroxy-undecanoic acid 3,7-dimethyl-octa-2,6-dienyl
10 ester

A mixture of 155 g 1-chloro-3,7-dimethyl-octa-2,6-diene, 202 g 4-hydroxy-undecanoic acid sodium salt and 5 g tetrabutylammoniumbromide in 800 ml of dimethylformamide was heated to 50°C. After stirring for 24 hours, the
15 mixture was cooled to room temperature and filtered through Celite. The filtrate was diluted with ether, washed with water, 2N HCl, saturated sodium bicarbonate and brine. The organic phase was dried and evaporated to dryness. The resulting yellow oil was purified by wipe
20 film distillation to yield 96.6 g of a yellow oil.

NMR (CDCl₃):

δ 5.42-5.37 (t, 1H), 5.16-5.01 (m, 1H), 4.65-4.53 (m, 2H), 3.69-3.52 (m, 1H), 2.60-2.22 (m, 2H), 2.20-1.95 (m, 4H),
25 1.89-1.12 (m, 24H), 1.02-0.78 (t, 3H) ppm.

Example 22

4-Hydroxy-undecanoic acid dodecylamide

A solution of 19.0 g aluminium chloride, dissolved in
30 methylene chloride, was cooled to 0°C and 46.5 g dodecylamine in methylene chloride was added dropwise

This Sheet Has Been Replaced



(exothermic). At room temperature 19.3 g gamma-undecalactone was added rapidly. The temperature rose to 36°C. The reaction was stirred at room temperature for 4 hours. The mixture was quenched with water and filtered
5 over Celite. The liquid was extracted with ether and washed with water and brine. The solution was dried, filtered and evaporated to dryness. The resulting brown crystals were purified by recrystallisation from ether/ethylacetate to give pale yellow crystals.

10

NMR (CDCl₃):

δ 5.7-5.6 (s, 1H), 3.7-3.55 (m, 1H), 3.3-3.15 (m, 2H), 2.4-2.3 (m, 2H), 1.95-1.2 (m, 34H), 0.95-0.85 (m, 6H) ppm.

15 Example 23

4-Hydroxy-undecanoic acid 3-(4-tert-butyl-phenyl)-2-methyl-propenyl ester

A solution of 10.00 g 4-oxo-undecanoic acid 3-(4-tert-butyl-phenyl)-2-methyl-propenyl ester was dissolved in 60
20 ml methanol and a trace of bromocresol green was added. When 1.63 g sodium cyanoborohydride was added, the colour changed immediately from yellow to deep blue. Several drops of 2N HCl/methanol solution turned the colour of the reaction back to yellow. The reaction was stirred for
25 2 1/2 hours, with occasional addition of acid to maintain the yellow colour. The reaction mixture was evaporated to dryness and water was added to the residue. This solution was extracted with ether and washed with water. The solution was dried, filtered and evaporated to dryness to
30 yield 10.07 g of a colourless oil.

NMR (CDCl₃):

5 δ 7.35-7.25 (m, 2H), 7.15-7.03 (m, 3H), 3.72-3.57 (m, 1H),
 3.27 + 3.2 (s, 2H, E/Z), 2.62-2.5 (m, 4H), 2.5-2.4 (m, 2H),
 1.65-1.56 (m, 5H), 1.35-1.21 (m, 17H), 0.95-0.82 (m, 3H)
 ppm.

Example 24

4-Hydroxy-undecanoic acid 3-(3-isopropyl-phenyl)-but-1-enyl ester

10 According to the procedure of Example 23, 4-hydroxy-undecanoic acid 3-(3-isopropyl-phenyl)-but-1-enyl ester was prepared from 4-oxo-undecanoic acid 3-(3-isopropyl-phenyl)-but-1-enyl ester and sodium cyanoborohydride.

15 Example 25

4-Hydroxy-undecanoic acid 2-benzyloxycarbonyl-2-benzyloxycarbonylamino-ethyl ester

20 According to the procedure of Example 23, 4-hydroxy-undecanoic acid 2-benzyloxycarbonyl-2-benzyloxycarbonylamino-ethyl ester was prepared from 4-oxo-undecanoic acid 2-benzyloxycarbonyl-2-benzyloxycarbonylamino-ethyl ester and sodium cyanoborohydride.

Example 26

25 3-[2-(tert-Butyl-dimethyl-silanyloxy)-phenyl]-acryloyl chloride

3-[2-(tert-Butyl-dimethyl-silanyloxy)-phenyl]-acrylic acid tert-butyl-dimethyl-silanyl ester

30 To a solution of 10.00 g 3-(2-hydroxy-phenyl)-acrylic acid (Tiemann; Herzfeld, Chem. Ber., 10 (1877), 285) and 19.3 g



tert-butyldimethylsilyl chloride in 40 ml of dimethylformamide was added 16.59 g imidazole. After stirring at 60°C for 6 hours, the mixture was poured onto water and extracted with hexane. The organic phase was
 5 washed with saturated sodium bicarbonate solution and water, dried and evaporated to dryness. The residue was not further purified.

10 3-[2-(tert-Butyl-dimethyl-silanyloxy)-phenyl]-acryloyl chloride

To a solution of 10.01 g 3-[2-(tert-butyl-dimethyl-silanyloxy)-phenyl]-acrylic acid tert-butyl-dimethyl-silanyl ester in 25 ml of dichloromethane, three drops of dimethylformamide were added and 4.14 g oxalyl chloride
 15 was dropped in at 0°C. After stirring at 0°C for 90 minutes and at room temperature overnight, the mixture was evaporated to dryness. The resulting brown solid (7.52 g) was not further purified.

20 NMR (CDCl₃):


δ 8.09-7.96 (d, 1H), 7.33-7.25 (d, 1H), 7.15-6.98 (m, 1H), 6.80-6.56 (m, 2H), 6.42-6.30 (d, 1H), 0.80 (s, 9H), 0.01 (s, 6H) ppm.

25

Example 27

Tetradecanoic acid 1-(2-hex-3-enyloxycarbonyl-ethyl)-heptyl ester

To a mixture of 1.10 g of 4-hydroxy-decanoic acid hex-3-enyl ester, 0.64 g pyridine and 0.1 g dimethylamino-
 30 pyridine in 5 ml tetrahydrofuran was slowly added a



solution of 1.10 g myristoyl chloride in 5 ml tetrahydrofuran. The mixture was stirred for 16 hours at room temperature.

Water was added and the mixture extracted with ethyl acetate. The organic layer was washed with saturated sodium bicarbonate, 2N hydrochloric acid, water, dried over magnesium sulfate and evaporated to dryness. The residue was chromatographed on silica gel to give 1.81 g of a colourless liquid.

10

NMR (CDCl₃):

δ 5.56 (m, 1H), 5.37 (m, 1H), 4.89 (q, 1H), 4.06 (t, 2H), 2.38-2.19 (m, 6H), 2.05 (q, 2H), 1.84 (m, 2H), 1.71-1.13 (m, 4H), 1.25 (s, 28H), 0.97 (t, 3H), 0.93-0.84 (m, 6H)

15 ppm.

Example 28

Tetradecanoic acid 1-[2-(3,7-dimethyl-octa-2,6-dienyloxy carbonyl)-ethyl]-octyl ester

20 According to the procedure of Example 27, tetradecanoic acid 1-[2-(3,7-dimethyl-octa-2,6-dienyloxycarbonyl)-ethyl]-octyl ester was prepared from 4-hydroxy-undecanoic acid 3,7-dimethyl-octa-2,6-dienyl ester, pyridine, dimethylaminopyridine and myristoyl chloride.

25

Example 29

Benzoic acid 1-[2-(3,7-dimethyl-octa-2,6-dienyloxy carbonyl)-ethyl]-octyl ester

30 According to the procedure of Example 27, benzoic acid 1-[2-(3,7-dimethyl-octa-2,6-dienyloxycarbonyl)-ethyl]-octyl ester was prepared from 4-hydroxy-undecanoic acid 3,7-

This Sheet Has Been Replaced



dimethyl-octa-2,6-dienyl ester, pyridine, dimethylamino-pyridine and benzoyl chloride.

Example 30

- 5 Dodecanoic acid 1-[2-(3,7-dimethyl-oct-6-enyloxycarbonyl)-ethyl]-heptyl ester

According to the procedure of Example 27, dodecanoic acid 1-[2-(3,7-dimethyl-oct-6-enyloxycarbonyl)-ethyl]-heptyl ester was prepared from 4-hydroxy-decanoic acid 3,7-
10 dimethyl-oct-6-enyl ester, pyridine, dimethylaminopyridine and lauroyl chloride.

Example 31


- 15 Dodecanoic acid 1-{2-[3-(4-tert-butyl-phenyl)-2-methyl-propenyloxycarbonyl]-ethyl}octyl ester

According to the procedure of Example 27, dodecanoic acid 1-{2-[3-(4-tert-butyl-phenyl)-2-methyl-propenyloxy-carbonyl]-ethyl}octyl ester was prepared from 4-hydroxy-undecanoic acid 3-(4-tert-butyl-phenyl)-2-methyl-propenyl
20 ester, lauroyl chloride, pyridine and dimethylamino-pyridine.

Example 32

- 25 Dodecanoic acid 1-{2-[3-(3-isopropyl-phenyl)-but-1-enyloxycarbonyl]-ethyl}octyl ester

According to the procedure of Example 27, dodecanoic acid 1-{2-[3-(3-isopropyl-phenyl)-but-1-enyloxycarbonyl]-ethyl}octyl ester was prepared from 4-hydroxy-undecanoic acid 3-(3-isopropyl-phenyl)-but-1-enyl ester, lauroyl
30 chloride, pyridine and dimethylaminopyridine.



Example 33Dodecanoic acid 1-[2-(2-benzyloxycarbonyl-2-benzyloxy carbonylamino-ethoxycarbonyl)-ethyl]-octyl ester

According to the procedure of Example 27, dodecanoic acid
 5 1-[2-(2-benzyloxycarbonyl-2-benzyloxycarbonylamino-ethoxycarbonyl)-ethyl]-octyl ester was prepared from 4-hydroxy-undecanoic acid 2-benzyloxycarbonyl-2-benzyloxy-carbonylamino-ethyl ester, lauroyl chloride and pyridine.

10 Example 344-Phenethyloxycarbonyloxy-decanoic acid 3,7-dimethyl-oct-6-enyl-ester

To a mixture of 3 g 4-hydroxy-decanoic acid 3,7-dimethyl-oct-6-enyl ester and 1.45 g pyridine in 7 ml tetra-
 15 hydrofuran was slowly added a solution of 1.87 g chlorocarbonic acid phenethyl ester (Schiving et al., Bull. Soc. Chim. Fr. (4), 43, 1928, 858) in 7 ml tetrahydrofuran. The mixture was stirred for 16 hours.
 Water was added and the mixture extracted with ethyl
 20 acetate. The organic layer was washed with saturated sodium bicarbonate, 2N hydrochloric acid, water, dried over magnesium sulfate and evaporated to dryness. The residue was chromatographed on silica gel to give 3.91 g of a colourless liquid.

25

NMR (CDCl₃):

δ 7.35-7.18 (m, 5H), 5.08 (m, 1H), 4.70 (m, 1H), 4.34 (t, 2H), 4.10 (t, 2H), 2.98 (t, 2H), 2.34 (m, 2H), 2.04-1.74 (m, 4H), 1.72-1.08 (m, 20H), 0.96-0.84 (m, 7H) ppm.

30

Example 35

This Sheet Has Been Replaced



4-Phenethyloxycarbonyloxy-decanoic acid hex-3-enyl ester

According to the procedure of Example 34, 4-phenethyloxy-carbonyloxy-decanoic acid hex-3-enyl ester was prepared from 4-hydroxy-decanoic acid hex-3-enyl ester, pyridine
 5 and chlorocarbonic acid phenethyl ester (Schiving et al., Bull. Soc. Chim. Fr. (4), 43, 1928, 858).

Example 36

10 4-Hex-3-enyloxycarbonyloxy-undecanoic acid 3,7-dimethyl-octa-2,6-dienyl ester

According to the procedure of Example 34, 4-hex-3-enyloxycarbonyloxy-undecanoic acid 3,7-dimethyl-octa-2,6-dienyl ester was prepared from 4-hydroxy-undecanoic acid 3,7-dimethyl-octa-2,6-dienyl ester, pyridine and chloro-
 15 carbonic acid hex-3-enyl ester (K. F. Podraza, J. Heterocycl. Chem. 1984, 21(4), 1197).

Example 37


20 4-Ethoxycarbonyloxy-nonanoic acid 3,7-dimethyl-octa-2,6-dienyl ester

According to the procedure of Example 34, 4-ethoxy-carbonyloxy-nonanoic acid 3,7-dimethyl-octa-2,6-dienyl ester was prepared from 4-hydroxy-nonanoic acid 3,7-dimethyl-octa-2,6-dienyl ester, pyridine and chloro-
 25 carbonic acid ethyl ester (commercially available).

Example 38

30 4-Phenethyloxycarbonyloxy-undecanoic acid 3,7-dimethyl-octa-2,6-dienyl ester

According to the procedure of Example 34, 4-phenethyloxycarbonyloxy-undecanoic acid 3,7-dimethyl-octa-



2,6-dienyl ester was prepared from 4-hydroxy-undecanoic acid 3,7-dimethyl-octa-2,6-dienyl ester, pyridine and chlorocarbonic acid phenethyl ester (Schiving et al., Bull. Soc. Chim. Fr. (4), 43, 1928, 858).

5

Example 39

4-Hex-3Z-enyloxycarbonyloxy-undecanoic acid 3-(4-isopropyl-phenyl-but-1-enyl ester

According to the procedure of Example 34, 4-hex-3Z-
10 enyloxycarbonyloxy-undecanoic acid 3-(4-isopropyl-phenyl)but-1-enyl ester was prepared from 4-hydroxy-undecanoic acid 3-(3-isopropyl-phenyl)-but-1-enyl ester, cis-hexenol-chloroformate and pyridine.

15 Example 40

4-(3-Methyl-5-phenyl-pentyloxycarbonyloxy)-undecanoic acid 3-(3-isopropyl-phenyl)-but-1-enyl ester

According to the procedure of Example 34, 4-(3-methyl-5-phenyl-pentyloxycarbonyloxy)-undecanoic acid 3-(3-
20 isopropyl-phenyl)-but-1-enyl ester was prepared from 4-hydroxy-undecanoic acid 3-(3-isopropyl-phenyl)-but-1-enyl ester, 3-methyl-5-phenyl-pentanol-chloroformate and pyridine.

25 Example 41

3-[2-(tert-Butyl-dimethyl-silanyloxy)-phenyl]-acrylic acid 1-(2-hex-3-enyloxycarbonyl-ethyl)-hexyl ester

According to the procedure of Example 34, 3-[2-(tert-butyl-dimethyl-silanyloxy)-phenyl]-acrylic acid 1-(2-hex-
30 3-enyloxycarbonyl-ethyl)-hexyl ester was prepared from 4-hydroxy-nonanoic acid hex-3-enyl ester, 3-[2-(tert-butyl-

This Sheet Has Been Replaced



dimethyl-silanyloxy)-phenyl]-acryloyl chloride and pyridine.

Example 42

- 5 4-(4-Allyl-2-methoxy-phenoxy-carbonyloxy)-undecanoic acid
3,7-dimethyl-octa-2,6-dienyl ester

According to the procedure of Example 34, 4-(4-allyl-2-methoxy-phenoxy-carbonyloxy)-undecanoic acid 3,7-dimethyl-octa-2,6-dienyl ester was prepared from 4-hydroxy-
 10 undecanoic acid 3,7-dimethyl-octa-2,6-dienyl ester, pyridine and chlorocarbonic acid-(4-allyl-2-methoxy-phenyl) ester (Einhorn, D.R.P. 224108).

Example 43

- 15 4-(Bis-decyl-carbamoyloxy)-undecanoic acid hex-3-enyl
ester

To 6.20 ml of an ice-cooled 20% solution of phosgene in toluene was slowly added a solution of 3 g 4-hydroxy-undecanoic acid hex-3-enyl ester and 0.93 g pyridine in 3
 20 ml toluene. The reaction mixture was stirred for 16 hours at room temperature. A mixture of 3.48 g didecylamine and 0.93 g pyridine was added and the mixture was stirred for another 16 hours.

Water was added and the mixture extracted with ethyl
 25 acetate. The organic layer was washed with saturated sodium bicarbonate, 2N hydrochloric acid, water, dried over magnesium sulfate and evaporated to dryness. The residue was chromatographed on silica gel to give 2.9 g of a colourless liquid.

30

NMR (CDCl₃):

This Sheet Has Been Replaced


δ 5.0 (m, 1H), 5.32 (m, 1H), 5.28 (quint, 1H), 4.06 (t, 2H), 3.18 (m, 4H), 2.35 (m, 4H), 2.05 (quint, 2H), 2.37 (m, 2H), 1.67-1.41 (m, 10H), 1.28 (bs, 30H), 0.98 (t, 3H), 0.88 (m, 9H) ppm.

5

Example 44

Succinic acid bis-[1-[2-(3,7-dimethyl-octa-2,6-dienyloxycarbonyl)-ethyl]-octyl] ester

To an ice cooled solution of 0.91 g succinic acid dichloride in 15 ml tetrahydrofuran was slowly added a mixture of 4 g 4-hydroxy-undecanoic acid 3,7-dimethyl-octa-2,6-dienyl ester and 0.93 g pyridine in 15 ml tetrahydrofuran. The mixture was stirred for 16 hours at room temperature.

Water was added and the mixture extracted with ethyl acetate. The organic layer was washed with saturated sodium bicarbonate, 2N hydrochloric acid, water, dried over magnesium sulfate and evaporated to dryness. The residue was chromatographed on silica gel to give 2.2 g of a colourless liquid.


NMR (CDCl₃):

δ 5.33 (bt, 2H), 5.09 (m, 2H), 4.91 (quint, 2H), 4.6 (m, 4H), 2.60 (m, 5H), 2.33 (m, 4H), 2.09 (m, 8H), 1.8 (m, 4H), 1.71 (bs, 12H), 1.62 (s, 6H), 1.52 (m, 4H), 1.27 (bs, 20H), 0.91 (bt, 6H) ppm.

Example 45

Succinic acid 1-[2-(1,5-dimethyl-1-vinyl-hex-4-enyloxycarbonyl)-ethyl]-heptyl ester hex-3-enyl ester

30



Succinic acid monohex-3-enyl ester

A mixture of 100 g succinic anhydride, 100 g cis-3-hexen-1-ol, 88.5 ml pyridine and 7.3 g dimethylaminopyridine in 500 ml dichloromethane was refluxed for 4 hours. Ether was
 5 added and the mixture was acidified with HCl 2N then washed with brine. The organic layer was dried over magnesium sulfate and concentrated. The residue was distilled under vacuum to afford 185 g of product.

10 NMR (CDCl₃):

δ 11.1 (bs, 1H), 5.53 (m, 1H), 5.31 (m, 1H), 4.11 (t, 2H), 2.64 (m, 4H), 2.39 (q, 2H), 2.07 (quint, 2H), 0.97 (t, 3H) ppm.

15 2-chlorocarbonic-propionic acid hex-3-enyl ester

To an ice cooled mixture of 184 g of succinic acid monohex-3-enyl ester and 79 g pyridine in 750 ml diethylether was added a solution of 69 ml thionyl chloride in 150 ml diethylether. The mixture was stirred
 20 for 17 hours at room temperature then filtered and concentrated to afford 184 g of a brown oil.

NMR (CDCl₃):

δ 5.02 (m, 1H), 4.82 (m, 1H), 4.12 (dt, 2H), 3.21 (t, 1H),
 25 2.72 (m, 5H), 2.39 (q, 2H), 2.04 (quint, 2H), 0.98 (t, 3H) ppm.

Succinic acid 1-[2-(1,5-dimethyl-1-vinyl-hex-4-enyloxycarbonyl)-ethyl]-heptyl ester hex-3-enyl ester

30 To a mixture of 3 g 4-hydroxy-decanoic acid 1,5-dimethyl-1-vinyl-hex-4-enyl ester and 1.6 g pyridine in 15 ml

tetrahydrofuran was added a solution of 2.2 g 2-chlorocarbonic-propionic acid hex-3-enyl ester in 15 ml tetrahydrofuran. The mixture was stirred for 16 hours at room temperature.

- 5 Water was added and the mixture extracted with ethyl acetate. The organic layer was washed with saturated sodium bicarbonate, 2N hydrochloric acid, water, dried over magnesium sulfate and evaporated to dryness. The residue was chromatographed on silica gel to give 1.9 g of
10 a colourless liquid.

NMR (CDCl₃):

- δ 5.96 (dd, 1H), 5.52 (m, 1H), 5.34 (m, 1H), 5.21-5.03 (m, 2H), 4.08 (t, 2H), 2.62 (s, 4H), 2.48-2.21 (m, 4H), 2.15-
15 1.71 (m, 8H), 1.70 (s, 3H), 1.54-1.46 (m, 8H), 1.23 (bs, 8H), 0.96 (t, 3H), 0.88 (bt, 3H) ppm.

Example 46

- Succinic acid 1-[2-(3,7-dimethyl-octa-2,6-dienyloxy-carbonyl)-ethyl]-hexyl ester hex-3Z-enyl ester
20

- According to the procedure of Example 45, succinic acid 1-[2-(3,7-dimethyl-octa-2,6-dienyloxy-carbonyl)-ethyl]-hexyl ester hex-3Z-enyl ester was prepared from 4-hydroxy-nonanoic acid 3,7-dimethyl-octa-2,6-dienyl ester, 2-
25 chlorocarbonic-propionic acid hex-3Z-enyl ester and pyridine.

Example 47

- 4-{1-[2-(1,1,5-trimethyl-hexyloxy-carbonyl)-ethyl]-octyloxy-carbonyloxy}-decanoic acid 1,1,5-trimethyl-hexyl ester
30

This Sheet Has Been Replaced



To a mixture of 3 g 4-hydroxy-decanoic acid 1,1,5-trimethyl-hexyl ester and 0.75 g pyridine was slowly added 2.5 ml of a 20% solution of phosgene in toluene. The mixture was stirred for 16 hours at room temperature.

- 5 Water was added and the mixture extracted with ethyl acetate. The organic layer was washed with saturated sodium bicarbonate, 2N hydrochloric acid, water and dried and evaporated to dryness. The residue was chromatographed on silica gel to give 1.9 g of a colourless liquid.

10

NMR (CDCl₃):

δ 4.71 (quint, 2H), 2.29 (m, 4H), 1.88 (m, 4H), 1.71-1.42 (m, 12 H), 1.42 (s, 12H), 1.39-1.08 (m, 22H), 1.88 (d, 18H) ppm.

15

Example 48

4-Hex-3-enyloxycarbonyloxy-undecanoic acid 3-(4-tert-butyl-phenyl)-2-methyl-propenyl ester


- According to the procedure of Example 34, 4-hex-3-enyloxycarbonyloxy-undecanoic acid 3-(4-tert-butyl-phenyl)-2-methyl-propenyl ester was prepared from 4-hydroxy-undecanoic acid 3-(4-tert-butyl-phenyl)-2-methyl-propenyl ester, pyridine and chlorocarbonic acid hex-3-enyl ester (K. F. Podraza, J. Heterocycl. Chem. 1984, 21(4), 1197).
- 20
- 25

Example 49

Carbonic acid 1-(2-dodecylcarbamoylethyl)-octyl ester hex-3-enyl ester

- 30 According to the procedure of Example 34, carbonic acid 1-(2-dodecylcarbamoylethyl)-octyl ester hex-3-enyl ester

This Sheet Has Been Replaced



was prepared from 4-hydroxy-undecanoic acid dodecylamide, pyridine and chlorocarbonic acid hex-3-enyl ester (K. F. Podraza, J. Heterocycl. Chem. 1984, 21(4), 1197).

5 Example 50

(E)-3-(2-Hydroxy-phenyl)-acrylic acid 1-(2-hex-3Z-
enyloxy-carbonyl-ethyl)-hexyl ester

To a solution of 6.14 g 3-[2-(*tert*-butyl-dimethyl-silanyloxy)-phenyl]-acrylic acid 1-(2-hex-3-enyloxy-carbonyl-ethyl)-hexyl ester in 50 ml of tetrahydrofuran, a
10 solution of 25 ml of a 1M tetrabutylammoniumfluoride solution was dropped in at 0°C. After stirring at room temperature for 90 minutes, the solution was quenched with water and extracted with ether. The organic phase was
15 washed with water, dried and evaporated to dryness. The residue was purified by Kugelrohr-distillation and chromatography to yield 0.51 g of a colourless oil.

NMR (CDCl₃):

20 δ 8.01-7.85 (d, 1H), 7.40-7.23 (m, 3H), 6.85-6.71 (m, 2H), 6.60-6.45 (d, 1H), 5.52-5.08 (m, 2H), 5.06- 4.88 (m, 1H), 4.05-3.89 (t, 2H), 2.44-2.12 (m, 4H), 2.10-1.68 (m, 4H), 1.67-1.00 (m, 8H), 0.98-0.68 (m, 6H) ppm.

25 Example 51

Test cloth was washed with a lipase-containing detergent to which one or more of the precursors of Examples 16 to 50 had been added. Headspace analysis of the wet and dry laundry indicated the presence of the fragrances. The
30 fragrance level was higher than when the test cloth was

washed with a lipase-containing detergent to which one or more fragrances were added.

Example 52

5 Test cloth was washed with a lipase-containing detergent and then a fabric softener, containing one or more of the precursors of Examples 16 to 50 was added to the rinse cycle. The laundry was then line-dried. Headspace analysis of the wet and dry laundry indicated the presence of the
10 fragrances. The fragrance level was higher, especially in the dry phase, than when the test cloth was washed with a lipase-containing detergent and then a fabric softener, containing one or more fragrances, was added to the rinse cycle.

15

Example 53

Test cloth was washed with a detergent and then a fabric softener, containing one or more of the precursors of Examples 16 to 50 was added to the rinse cycle. The
20 laundry was then tumble-dried. Headspace analysis of the wet and dry laundry indicated the presence of the fragrances. The fragrance level was higher, especially in the dry phase, than when the test cloth was washed with a detergent and then a fabric softener, containing one or
25 more fragrances, was added to the rinse cycle.

Example 54

Axilla bacteria cultures containing 0.1 % of one or more of the precursors of Examples 16 to 50 were incubated for
30 20 hours at 30 °C. After filtration from the cells, the presence of the corresponding fragrance was in each case

This Sheet Has Been Replaced



detected by headspace-GC techniques and/or the majority of an 18 member panel.

The same tests were carried out with inactivated cultures
 5 (85°/20 min). The odour of the corresponding fragrance could not be detected after incubation, excluding therefore a hydrolysis by the medium or the culture.

The following set forth examples for the use of the
 10 compounds of the present invention in various products. The methods of forming the following compositions are well known to those skilled in the art. All formulations may contain additional ingredients known to those skilled in the art, e.g. colorants, opacifiers, buffers,
 15 antioxidants, vitamins, emulsifiers, UV absorbers, silicones and the like. All products can also be buffered to the desired pH. All values are % w/w. Delayed Release Fragrances stands in the following for compounds of Examples 16-50.

20

Example 55

a) Deo-colognes

	Delayed Release Fragrances	0.5	1.5	2.5	6.0
25	Fragrance	0.5	1.5	2.5	6.0
	Triclosan (Ciba Geigy)	1.0	-	0.75	1.0
	Alcohol to	100	100	100	100

b) Deo-Sticks


30 Antiperspirant

Ethylene Glycol Monostearate 7.0

This Sheet Has Been Replaced



	Shea butter	3.0
	Neobee 1053 (PVO International)	12.0
	Generol 122 (Henkel)	5.0
	Kesscowax B (Akzo)	17.0
5	Dimethicone Dow Corning 345	35.0
	Aluminum Sesquichlorhydrate	20.0
	Delayed Release Fragrances	0.5
	Fragrance	0.5
10	Antiperspirant	
	Steary Alcohol	17.0
	Castor Wax	3.0
	Talc	5.0
	Aluminum Zirconium Tetrachlorhydrate	20.0
15	Delayed Release Fragrances	1.0
	Fragrance	1.0
	Dimethicone Dow 245	to 100.0
	Clear Deodorant Stick	
20	Witconol APM	43.0
	Propylene Glycol	20.0
	Alcohol 39C	20.0
	Demin water	7.0
	Monamid 150ADD	5.0
25	Millithix 925	2.0
	Ottasept Extra	0.5
	Delayed Release Fragrances	0.75
	Fragrance	0.75
30	Deodorant Stick	
	Propylene Glycol	69.0



	Demin Water	21.8
	Triclosan	0.2
	Sodium Stearate	8.0
	Delayed Release Fragrances	0.5
5	Fragrance	0.5
	Alcohol free Deodorant Stick	
	PPG-3 Myristyl Ether(Witconol APM)	36.0
	Propylene Glycol	36.0
10	Demin Water	19.0
	Triclosan	0.25
	Sodium Stearate	7.75
	Delayed Release Fragrances	0.5
	Fragrance	0.5
15	Antiperspirant Aerosol	
	Absolute Ethanol	15.0
	Zirconium Aluminum tetrachlorhydrate	5.0
	Bentone 38	1.5
20	Delayed Release Fragrances	0.75
	Fragrance	0.75
	S-31 Hydrocarbon propellant	to 100.0
	Antiperspirant Pump	
25	Demin water	57.5
	Aluminum Sesquichlorhydrate	20.0
	Triton X-102 (Union Carbide)	2.0
	Dimethyl Isosorbide (ICI)	20.0
	Delayed Release Fragrances	0.25
30	Fragrance	0.25

This Sheet Has Been Replaced



	Roll-On	
	Dimethicone DC 354 (Dow Corning)	69.0
	Bentone 38	10.0
	Rezal 36 GP (Reheis Chem. Co.)	20.0
5	Delayed Release Fragrances	0.5
	Fragrance	0.5

In the above examples, the following components were used:

10	Triclosan	5-chloro-2-(2,4-dichlorophenoxy)phenol
	Neobee 1053	glycerol tricaprates/caprylates
	Generol 122	soya sterol
	Kesscowax B	cetyl alcohol and glycol polymer
	Witconol APM	polypropylene glycol-3 myristyl ether
15	Monamid 150 ADD	cocoamide diethanolamine
	Millithix 925	dibenzylidene sorbitol
	Ottasept Extra	quaternium 18 hectorite
	Bentone 38	quaternium 18 hectorite
	Triton X-102	octoxynol-13
20	Dimethicone DC 354	mixture of fully methylated linear siloxanepolymers end-blocked with trimethylsiloxy units
	Rezal 36 GP	Aluminium zirconium tetrachlorohydroxyglycine

25


Example 56

a) Fabric softener of the ester quat type (4 x concentrate):

30

INGREDIENTS	CHEMICAL NAME	%
-------------	---------------	---

This Sheet Has Been Replaced



PHASE A

DEIONISED WATER to 100.0

5 MgCl_2 (saturated sol.) Magnesium chloride 1.0

PHASE B

REWOQUAT WE 18 Di-(tallow carboxyethyl) 15.0
hydroxy ethyl methylammonium
10 methosulfate

GENAPOL O 100 Ethoxylated fatty alcohol 2.0
C16-C18 10EO

15 ANTIFOAM DB 31 0.5

PHASE C

ISOPROPYL ALCOHOL 3.0

20 PRESERVATIVE Qs

PERFUME Qs

25 PROCESS:

While stirring and heating to 65° C, mix part A, then part B preheated to 65° C. After cooling to room temperature, add part C.

30 The PH value of the finished product is 2.60.

This Sheet Has Been Replaced



Recommended level of perfume is 1.0 %. Delayed release fragrances can be any part of this 1.0 %.

b) Fabric softener of the ester quat type (1 x
5 concentrate):

	INGREDIENTS	CHEMICAL NAME	%
	<u>PHASE A</u>		
10	DEIONISED WATER		to 100.0
	<u>PHASE B</u>		
15	REWOQUAT WE 18	Di-(tallow carboxyethyl) hydroxy ethyl methyl ammonium methosulfate	6.0
	DOBANOL 25-9	Ethoxylated fatty alcohol C12-C15 9EO	0.5
20	ANTIFOAM DB 31		0.1
	<u>PHASE C</u>		
25	MYACIDE BT 30	2-bromo-2-nitropropane 1,3 diol	0.03
	PROXEL GXL	Benzisothiazolinone sodium salt	0.02
30	PERFUME		Qs

PROCESS:


This Sheet Has Been Replaced



While stirring and heating to 65° C, mix part A, then part B preheated to 65° C. After cooling to room temperature, add part C.

- 5 The pH value of the finished product is 3.50.
Recommended level of perfume: 0.3 %. Delayed release fragrances can be any part of this 0.3 %.

This Sheet Has Been Replaced




Claims


1. Compounds of Formula I

I

5



wherein Y is



10

m is an integer of 1 or greater

n is 1, 2 or 3;

15

R^1 , R^2 , R^3 , R^4 , R^5 and R^6 represent independently hydrogen, substituted or unsubstituted alkyl-, alkenyl-, alkynyl-, cycloalkyl-, cycloalkenyl- or aromatic radicals which can additionally contain one or more


20

O

-O- and/or -C- groups;




This Sheet Has Been Replaced



whereby one or two rings can be built by the combination of the respective R^1 to R^6 and said ring(s) can be substituted by one or more alkyl groups;

5 X is $-OR^7$ and R^7 is the residue of an alcohol R^7OH , or the residue of the enol form of an aldehyde or ketone, or X is a primary, or secondary amino group forming an amide;

10 Z is



q is the same or bigger than m;

15

R^8 represents hydrogen, a straight or branched, unsubstituted or substituted alkyl-, alkenyl-, cycloalkyl-, cycloalkenyl- or aromatic radical which optionally comprises and/or is substituted by one or more heteroatoms, and/or group(s) comprising a heteroatom;

20


R^9 is the residue $-OR^{12}$ of an alcohol of formula $R^{12}OH$ or the residue of the enol form of an aldehyde or ketone or has the definition given for Y and R^9 and Y can be the same or different and optionally comprises and/or is substituted by a heteroatom and/or group(s) comprising a heteroatom;

25




5 R^{10} and R^{11} represent independently hydrogen, substituted or unsubstituted alkyl, alkenyl, cycloalkyl, cycloalkenyl or an aromatic residue which optionally comprise and/or are substituted by one or more heteroatoms and/or group(s) comprising a heteroatom.

10 2. Compound according to claim 1, wherein Z is



15 and R^8 is derived from an odourless acid optionally substituted by groups yielding upon cleavage one or more organoleptic compounds.

3. Compound according to claim 1, wherein Z is




20 and R^9 is derived from an organoleptic compound or Y.

4. Compound according to claim 1, wherein Z is


This Sheet Has Been Replaced



This Sheet Has Been Replaced



and




5

is derived from a non odorous mono- or diamine.

5. Compounds according to claim 1 or 2, wherein Z is the
10 residue of a β -keto acid.

6. Compounds according to claim 1 or 2, wherein




15

is a photolabile ester.

7. Compounds of claim 6 yielding under activating
conditions, preferably light, an odoriferous or
20 fluorescent coumarin.

8. Compounds according to any one of the preceding
claims, wherein R^8 comprises and/or is substituted by
one or more





- 5 9. Compound according to any one of the preceding claims, wherein R^9 , R^{10} and/or R^{11} comprise and/or are substituted by one or more O, Si, N and/or CO.
- 10 10. Compositions comprising a compound of formula I as precursor for at least one organoleptic compound.
11. Laundry compositions according to claim 10 comprising a compound of formula I as precursor for an organoleptic compound and/or a fluorescent coumarin.



This Sheet Has Been Replaced

