CHIP-IN-A-WELL SCANNING

RELATED APPLICATION

This application claims the benefit of U.S. Provisional Application Serial Number 60/433,186, filed on December 13, 2002, which is incorporated herein by reference.

BACKGROUND OF THE INVENTION

The present invention relates to the packaging and scanning of materials, more specifically the packaging and scanning of microarrays.

SUMMARY OF THE INVENTION

One embodiment of the present invention discloses a method of packaging a microarray wherein the microarray is suspended in a well of a well plate. In one embodiment of the invention, microarrays can be selected from diced wafers and then placed in wells of a well plate. The microarrays are captured in a suspended form, with or without a buffer, in the wells of the well plate with the use of an optically clear window. In one embodiment of the invention, the microarrays are then successfully scanned with the surface of the microarray up or down.

In another embodiment of the invention, a microarray can be selected from a diced wafer and can be placed in the well of a cartridge. The microarray can be contained in the well of a well plate with or without buffer. The microarray is then successfully scanned with the surface of the microarray up or down.

BRIEF DESCRIPTION OF THE DRAWINGS

The accompanying drawings, which are incorporated in and form a part of this specification, illustrate embodiments of the invention and, together with the description, serve to explain the principles of the invention:

Figure 1 depicts a wafer, a microtiter plate well and a microarray suspended in a well of a microtiter well plate.

Figure 2 depicts a wafer, a cartridge and a microarray suspended in the well of the cartridge.

DETAILED DESCRIPTION OF THE INVENTION

I. General

The present invention has many preferred embodiments and relies on many patents, applications and other references for details known to those of the art. Therefore, when a patent, application, or other reference is cited or repeated below, it should be understood that it is incorporated by reference in its entirety for all purposes as well as for the proposition that is recited.

As used in this application, the singular form "a," "an," and "the" include plural references unless the context clearly dictates otherwise. For example, the term "an agent" includes a plurality of agents, including mixtures thereof.

An individual is not limited to a human being but may also be other organisms including but not limited to mammals, plants, bacteria, or cells derived from any of the above.

Throughout this disclosure, various aspects of this invention can be presented in a range format. It should be understood that the description in range format is merely for convenience and brevity and should not be construed as an inflexible limitation on the scope of the invention. Accordingly, the description of a range should be considered to have specifically disclosed all the possible subranges as well as individual numerical values within that range. For example, description of a range such as from 1 to 6 should be considered to have specifically disclosed subranges such as from 1 to 3, from 1 to 4, from 1 to 5, from 2 to 4, from 2 to 6, from 3 to 6 etc., as well as individual numbers within that range, for example, 1, 2, 3, 4, 5, and 6. This applies regardless of the breadth of the range.

The practice of the present invention may employ, unless otherwise indicated, conventional techniques and descriptions of organic chemistry, polymer technology, molecular biology (including recombinant techniques), cell biology, biochemistry, and immunology, which are within the skill of the art. Such conventional techniques include polymer array synthesis, hybridization, ligation, and detection of hybridization using a label. Specific illustrations of suitable techniques can be had by reference to the example herein below. However, other equivalent conventional procedures can, of course, also be used. Such conventional techniques and descriptions can be found in standard laboratory manuals such as *Genome Analysis: A Laboratory Manual Series (Vols. I-IV)*, *Using Antibodies: A Laboratory Manual*, *Cells: A Laboratory Manual*, *PCR Primer: A Laboratory Manual*, and *Molecular Cloning: A Laboratory Manual* (all from Cold Spring Harbor Laboratory Press), Stryer, L. (1995) *Biochemistry* (4th Ed.) Freeman, New York, Gait, "Oligonucleotide Synthesis: A Practical Approach" 1984, IRL Press, London, Nelson and Cox (2000), Lehninger, *Principles of Biochemistry* 3rd Ed., W.H. Freeman Pub., New York, NY and Berg et al. (2002) *Biochemistry*, 5th Ed., W.H. Freeman Pub., New York, NY, all of which are herein incorporated in their entirety by reference for all purposes.

The present invention can employ solid substrates, including arrays in some preferred embodiments. Methods and techniques applicable to polymer (including protein) array synthesis have been described in U.S. Serial No. 09/536,841, WO 00/58516, U.S. Patent Nos. 5,143,854, 5,242,974, 5,252,743, 5,324,633, 5,384,261, 5,405,783, 5,424,186, 5,451,683, 5,482,867, 5,491,074, 5,527,681, 5,550,215, 5,571,639, 5,578,832, 5,593,839, 5,599,695, 5,624,711, 5,631,734, 5,795,716, 5,831,070, 5,837,832, 5,856,101, 5,858,659, 5,936,324, 5,968,740, 5,974,164, 5,981,185, 5,981,956, 6,025,601, 6,033,860, 6,040,193, 6,090,555, 6,136,269, 6,269,846 and 6,428,752, in PCT Applications Nos. PCT/US99/00730 (International Publication Number WO 99/36760) and PCT/US01/04285 (International Publication Number WO 01/58593), which are all incorporated herein by reference in their entirety for all purposes.

Patents that describe synthesis techniques in specific embodiments include U.S. Patent Nos. 5,412,087, 6,147,205, 6,262,216, 6,310,189, 5,889,165, and

5,959,098. Nucleic acid arrays are described in many of the above patents, but the same techniques are applied to polypeptide arrays.

Nucleic acid arrays that are useful in the present invention include those that are commercially available from Affymetrix (Santa Clara, CA) under the brand name GeneChip®. Example arrays are shown on the website at affymetrix.com.

The present invention also contemplates many uses for polymers attached to solid substrates. These uses include gene expression monitoring, profiling, library screening, genotyping and diagnostics. Gene expression monitoring, and profiling methods can be shown in U.S. Patent Nos. 5,800,992, 6,013,449, 6,020,135, 6,033,860, 6,040,138, 6,177,248 and 6,309,822. Genotyping and uses therefore are shown in U.S. Serial Nos. 60/319,253, 10/013,598 (U.S. Patent Application Publication 20030036069), and U.S. Patent Nos. 5,856,092, 6,300,063, 5,858,659, 6,284,460, 6,361,947, 6,368,799 and 6,333,179. Other uses are embodied in U.S. Patent Nos. 5,871,928, 5,902,723, 6,045,996, 5,541,061, and 6,197,506.

The present invention also contemplates sample preparation methods in certain preferred embodiments. Prior to or concurrent with genotyping, the genomic sample may be amplified by a variety of mechanisms, some of which may employ PCR. See, e.g., *PCR Technology: Principles and Applications for DNA Amplification* (Ed. H.A. Erlich, Freeman Press, NY, NY, 1992); *PCR Protocols: A Guide to Methods and Applications* (Eds. Innis, et al., Academic Press, San Diego, CA, 1990); Mattila et al., *Nucleic Acids Res.* 19, 4967 (1991); Eckert et al., *PCR Methods and Applications* 1, 17 (1991); PCR (Eds. McPherson et al., IRL Press, Oxford); and U.S. Patent Nos. 4,683,202, 4,683,195, 4,800,159 4,965,188,and 5,333,675, and each of which is incorporated herein by reference in their entireties for all purposes. The sample may be amplified on the array. See, for example, U.S. Patent No. 6,300,070 and U.S. Serial No. 09/513,300, which are incorporated herein by reference.

Other suitable amplification methods include the ligase chain reaction (LCR) (e.g., Wu and Wallace, *Genomics* 4, 560 (1989), Landegren *et al.*, *Science* 241, 1077 (1988) and Barringer *et al. Gene* 89:117 (1990)), transcription amplification (Kwoh *et al., Proc. Natl. Acad. Sci.* USA 86, 1173 (1989) and WO88/10315), self sustained sequence replication (Guatelli *et al., Proc. Nat. Acad. Sci.* USA, 87, 1874 (1990) and

WO90/06995), selective amplification of target polynucleotide sequences (U.S. Patent No 6,410,276), consensus sequence primed polymerase chain reaction (CP-PCR) (U.S. Patent No. 4,437,975), arbitrarily primed polymerase chain reaction (AP-PCR) (U.S. Patent No. 5, 413,909, 5,861,245) and nucleic acid based sequence amplification (NABSA). (See, U.S. Patent Nos. 5,409,818, 5,554,517, and 6,063,603, each of which is incorporated herein by reference). Other amplification methods that may be used are described in, U.S. Patent Nos. 5,242,794, 5,494,810, 4,988,617 and in U.S. Serial No. 09/854,317, each of which is incorporated herein by reference.

Additional methods of sample preparation and techniques for reducing the complexity of a nucleic sample are described in Dong *et al.*, *Genome Research* 11, 1418 (2001), in U.S. Patent No. 6,361,947, 6,391,592 and U.S. Serial Nos. 09/916,135, 09/920,491 (U.S. Patent Application Publication 20030096235), 09/910,292 (U.S. Patent Application Publication 20030082543), and 10/013,598.

Methods for conducting polynucleotide hybridization assays have been well developed in the art. Hybridization assay procedures and conditions will vary depending on the application and are selected in accordance with the general binding methods known including those referred to in: Maniatis et al. *Molecular Cloning: A Laboratory Manual* (2nd Ed. Cold Spring Harbor, N.Y, 1989); Berger and Kimmel *Methods in Enzymology*, Vol. 152, Guide to Molecular Cloning Techniques (Academic Press, Inc., San Diego, CA, 1987); Young and Davism, *P.N.A.S*, 80: 1194 (1983). Methods and apparatus for carrying out repeated and controlled hybridization reactions have been described in U.S. Patent Nos. 5,871,928, 5,874,219, 6,045,996 and 6,386,749, 6,391,623 each of which are incorporated herein by reference

The present invention also contemplates signal detection of hybridization between ligands in certain preferred embodiments. See U.S. Patent Nos. 5,143,854, 5,578,832; 5,631,734; 5,834,758; 5,936,324; 5,981,956; 6,025,601; 6,141,096; 6,185,030; 6,201,639; 6,218,803; and 6,225,625, in U.S. Serial No. 60/364,731 and in PCT Application PCT/US99/06097 (published as WO99/47964), each of which also is hereby incorporated by reference in its entirety for all purposes.

Methods and apparatus for signal detection and processing of intensity data are disclosed in, for example, U.S. Patents Nos. 5,143,854, 5,547,839, 5,578,832, 5,631,734, 5,800,992, 5,834,758; 5,856,092, 5,902,723, 5,936,324, 5,981,956, 6,025,601, 6,090,555, 6,141,096, 6,185,030, 6,201,639; 6,218,803; and 6,225,625, in U.S. Serial No. 60/364,731 and in PCT Application PCT/US99/06097 (published as WO99/47964), each of which also is hereby incorporated by reference in its entirety for all purposes.

The practice of the present invention may also employ conventional biology methods, software and systems. Computer software products of the invention typically include computer readable medium having computer-executable instructions for performing the logic steps of the method of the invention. Suitable computer readable medium include floppy disk, CD-ROM/DVD/DVD-ROM, hard-disk drive, flash memory, ROM/RAM, magnetic tapes and etc. The computer executable instructions may be written in a suitable computer language or combination of several languages. Basic computational biology methods are described in, e.g., Setubal and Meidanis *et al.*, *Introduction to Computational Biology Methods* (PWS Publishing Company, Boston, 1997); Salzberg, Searles, Kasif, (Ed.), *Computational Methods in Molecular Biology*, (Elsevier, Amsterdam, 1998); Rashidi and Buehler, *Bioinformatics Basics: Application in Biological Science and Medicine* (CRC Press, London, 2000) and Ouelette and Bzevanis *Bioinformatics: A Practical Guide for Analysis of Gene and Proteins* (Wiley & Sons, Inc., 2nd ed., 2001). See U.S. Patent No. 6,420,108.

The present invention may also make use of various computer program products and software for a variety of purposes, such as probe design, management of data, analysis, and instrument operation. See, U.S. Patent Nos. 5,593,839, 5,795,716, 5,733,729, 5,974,164, 6,066,454, 6,090,555, 6,185,561, 6,188,783, 6,223,127, 6,229,911 and 6,308,170.

Additionally, the present invention may have preferred embodiments that include methods for providing genetic information over networks such as the Internet as shown in U.S. Serial Nos. 10/063,559 (United States Publication No. US20020183936), 60/349,546, 60/376,003, 60/394,574 and 60/403,381.

II. Glossary

The following terms are intended to have the following general meanings as used herein.

An "array" is an intentionally created collection of molecules which can be prepared either synthetically or biosynthetically. The molecules in the array can be identical or different from each other. The array can assume a variety of formats, e.g., libraries of soluble molecules; libraries of compounds tethered to resin beads, silica chips, or other solid supports.

An allele refers to one specific form of a genetic sequence (such as a gene) within a cell or within a population, the specific form differing from other forms of the same gene in the sequence of at least one, and frequently more than one, variant sites within the sequence of the gene. The sequences at these variant sites that differ between different alleles are termed "variances", "polymorphisms", or "mutations".

At each autosomal specific chromosomal location or "locus" an individual possesses two alleles, one inherited from the father and one from the mother. An individual is "heterozygous" at a locus if it has two different alleles at that locus. An individual is "homozygous" at a locus if it has two identical alleles at that locus.

Nucleic acid library or array is an intentionally created collection of nucleic acids which can be prepared either synthetically or biosynthetically and screened for biological activity in a variety of different formats (e.g., libraries of soluble molecules; and libraries of oligos tethered to resin beads, silica chips, or other solid supports). Additionally, the term "array" is meant to include those libraries of nucleic acids which can be prepared by spotting nucleic acids of essentially any length (e.g., from 1 to about 1000 nucleotide monomers in length) onto a substrate. The term "nucleic acid" as used herein refers to a polymeric form of nucleotides of any length, either ribonucleotides, deoxyribonucleotides or peptide nucleic acids (PNAs), that comprise purine and pyrimidine bases, or other natural, chemically or biochemically modified, non-natural, or derivatized nucleotide bases. The backbone of the polynucleotide can comprise sugars and phosphate groups, as may typically be found in RNA or DNA, or modified or substituted sugar or phosphate groups. A polynucleotide may comprise modified nucleotides, such as methylated nucleotides and nucleotide analogs. The sequence of nucleotides may be interrupted by non-

nucleotide components. Thus the terms nucleoside, nucleotide, deoxynucleoside and deoxynucleotide generally include analogs such as those described herein. These analogs are those molecules having some structural features in common with a naturally occurring nucleoside or nucleotide such that when incorporated into a nucleic acid or oligonucleoside sequence, they allow hybridization with a naturally occurring nucleic acid sequence in solution. Typically, these analogs are derived from naturally occurring nucleosides and nucleotides by replacing and/or modifying the base, the ribose or the phosphodiester moiety. The changes can be tailor made to stabilize or destabilize hybrid formation or enhance the specificity of hybridization with a complementary nucleic acid sequence as desired.

Biopolymer or biological polymer is intended to mean repeating units of biological or chemical moieties. Representative biopolymers include, but are not limited to, nucleic acids, oligonucleotides, amino acids, proteins, peptides, hormones, oligosaccharides, lipids, glycolipids, lipopolysaccharides, phospholipids, synthetic analogues of the foregoing, including, but not limited to, inverted nucleotides, peptide nucleic acids, Meta-DNA, and combinations of the above. "Biopolymer synthesis" is intended to encompass the synthetic production, both organic and inorganic, of a biopolymer.

Related to a bioploymer is a "biomonomer" which is intended to mean a single unit of biopolymer, or a single unit which is not part of a biopolymer. Thus, for example, a nucleotide is a biomonomer within an oligonucleotide biopolymer, and an amino acid is a biomonomer within a protein or peptide biopolymer; avidin, biotin, antibodies, antibody fragments, etc., for example, are also biomonomers. initiation Biomonomer: or "initiator biomonomer" is meant to indicate the first biomonomer which is covalently attached via reactive nucleophiles to the surface of the polymer, or the first biomonomer which is attached to a linker or spacer arm attached to the polymer, the linker or spacer arm being attached to the polymer via reactive nucleophiles.

Buffer: A substance that minimizes change in the acidity of a solution when an acid or base is added to the solution. A buffer contains either a weak acid and a soluble ionic salt of the acid or a weak base and a soluble ionic salt of the base.

Cartridge: A body forming an area or space referred to as a well wherein a microarray is contained and separated from the passage of liquids.

Complementary or substantially complementary: Refers to the hybridization or base pairing between nucleotides or nucleic acids, such as, for instance, between the two strands of a double stranded DNA molecule or between an oligonucleotide primer and a primer binding site on a single stranded nucleic acid to be sequenced or amplified. Complementary nucleotides are, generally, A and T (or A and U), or C and G. Two single stranded RNA or DNA molecules are said to be substantially complementary when the nucleotides of one strand, optimally aligned and compared and with appropriate nucleotide insertions or deletions, pair with at least about 80% of the nucleotides of the other strand, usually at least about 90% to 95%, and more preferably from about 98 to 100%. Alternatively, substantial complementary exists when an RNA or DNA strand will hybridize under selective hybridization conditions to its complement. Typically, selective hybridization will occur when there is at least about 65% complementary over a stretch of at least 14 to 25 nucleotides, preferably at least about 75%, more preferably at least about 90% complementary. See, M. Kanehisa, *Nucleic Acids Res.* 12:203 (1984), incorporated herein by reference.

Combinatorial Synthesis Strategy: A combinatorial synthesis strategy is an ordered strategy for parallel synthesis of diverse polymer sequences by sequential addition of reagents which may be represented by a reactant matrix and a switch matrix, the product of which is a product matrix. A reactant matrix is a l column by m row matrix of the building blocks to be added. The switch matrix is all or a subset of the binary numbers, preferably ordered, between l and m arranged in columns. A "binary strategy" is one in which at least two successive steps illuminate a portion, often half, of a region of interest on the substrate. In a binary synthesis strategy, all possible compounds which can be formed from an ordered set of reactants are formed. In most preferred embodiments, binary synthesis refers to a synthesis strategy which also factors a previous addition step. For example, a strategy in which a switch matrix for a masking strategy halves regions that were previously illuminated, illuminating about half of the previously illuminated region and protecting the remaining half (while also protecting about half of previously protected regions and illuminating about half of previously protected regions). It will

be recognized that binary rounds may be interspersed with non-binary rounds and that only a portion of a substrate may be subjected to a binary scheme. A combinatorial "masking" strategy is a synthesis which uses light or other spatially selective deprotecting or activating agents to remove protecting groups from materials for addition of other materials such as amino acids.

Effective amount refers to an amount sufficient to induce a desired result.

A fragment, segment, or DNA segment refers to a portion of a larger DNA polynucleotide or DNA. A polynucleotide, for example, can be broken up, or fragmented into, a plurality of segments. Various methods of fragmenting nucleic acid are well known in the art. These methods may be, for example, either chemical or physical in nature. Chemical fragmentation may include partial degradation with a DNase; partial depurination with acid; the use of restriction enzymes; intron-encoded endonucleases; DNA-based cleavage methods, such as triplex and hybrid formation methods, that rely on the specific hybridization of a nucleic acid segment to localize a cleavage agent to a specific location in the nucleic acid molecule; or other enzymes or compounds which cleave DNA at known or unknown locations. Physical fragmentation methods may involve subjecting the DNA to a high shear rate. High shear rates may be produced, for example, by moving DNA through a chamber or channel with pits or spikes, or forcing the DNA sample through a restricted size flow passage, e.g., an aperture having a cross sectional dimension in the micron or submicron scale. Other physical methods include sonication and nebulization. Combinations of physical and chemical fragmentation methods may likewise be employed such as fragmentation by heat and ion-mediated hydrolysis. See for example, Sambrook *et al.*, "*Molecular Cloning: A Laboratory Manual*," 3rd Ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (2001) ("Sambrook *et al.*) which is incorporated herein by reference for all purposes. These methods can be optimized to digest a nucleic acid into fragments of a selected size range. Useful size ranges may be from 100, 200, 400, 700 or 1000 to 500, 800, 1500, 2000, 4000 or 10,000 base pairs. However, larger size ranges such as 4000, 10,000 or 20,000 to 10,000, 20,000 or 500,000 base pairs may also be useful. See, e.g., Dong *et al.*, *Genome Research* 11, 1418 (2001), in U.S. Patent No 6,361,947, 6,391,592, incorporated herein by reference.

Genome is all the genetic material in the chromosomes of an organism. DNA derived from the genetic material in the chromosomes of a particular organism is genomic DNA. A genomic library is a collection of clones made from a set of randomly generated overlapping DNA fragments representing the entire genome of an organism.

Genotyping refers to the determination of the genetic information an individual carries at one or more positions in the genome. For example, genotyping may comprise the determination of which allele or alleles an individual carries for a single SNP or the determination of which allele or alleles an individual carries for a plurality of SNPs. A genotype may be the identity of the alleles present in an individual at one or more polymorphic sites.

Hybridization conditions will typically include salt concentrations of less than about 1M, more usually less than about 500 mM and preferably less than about 200 mM. Hybridization temperatures can be as low as 5°C., but are typically greater than 22°C., more typically greater than about 30°C., and preferably in excess of about 37°C. Longer fragments may require higher hybridization temperatures for specific hybridization. As other factors may affect the stringency of hybridization, including base composition and length of the complementary strands, presence of organic solvents and extent of base mismatching, the combination of parameters is more important than the absolute measure of any one alone.

Hybridizations, e.g., allele-specific probe hybridizations, are generally performed under stringent conditions. For example, conditions where the salt concentration is no more than about 1 Molar (M) and a temperature of at least 25 degrees-Celcius (°C), e.g., 750 mM NaCl, 50 mM NaPhosphate, 5 mM EDTA, pH 7.4 (5X SSPE)and a temperature of from about 25 to about 30°C.

Hybridizations are usually performed under stringent conditions, for example, at a salt concentration of no more than 1 M and a temperature of at least 25°C. For example, conditions of 5X SSPE (750 mM NaCl, 50 mM NaPhosphate, 5 mM EDTA, pH 7.4) and a temperature of 25-30°C are suitable for allele-specific probe hybridizations. For stringent conditions, see, for example, Sambrook, Fritsche and Maniatis. "*Molecular Cloning A laboratory Manual*" 2nd Ed. Cold Spring

Harbor Press (1989) which is hereby incorporated by reference in its entirety for all purposes above.

The term "hybridization" refers to the process in which two single-stranded polynucleotides bind non-covalently to form a stable double-stranded polynucleotide; triple-stranded hybridization is also theoretically possible. The resulting (usually) double-stranded polynucleotide is a "hybrid." The proportion of the population of polynucleotides that forms stable hybrids is referred to herein as the "degree of hybridization."

Hybridization probes are oligonucleotides capable of binding in a base-specific manner to a complementary strand of nucleic acid. Such probes include peptide nucleic acids, as described in Nielsen et al., Science 254, 1497-1500 (1991), and other nucleic acid analogs and nucleic acid mimetics. See U.S. Patent No. 6,156,501.

"Hybridizing specifically to" refers to the binding, duplexing, or hybridizing of a molecule substantially to or only to a particular nucleotide sequence or sequences under stringent conditions when that sequence is present in a complex mixture (e.g., total cellular) DNA or RNA.

Isolated nucleic acid is an object species invention that is the predominant species present (i.e., on a molar basis it is more abundant than any other individual species in the composition). Preferably, an isolated nucleic acid comprises at least about 50, 80 or 90% (on a molar basis) of all macromolecular species present. Most preferably, the object species is purified to essential homogeneity (contaminant species cannot be detected in the composition by conventional detection methods).

Ligand: A ligand is a molecule that is recognized by a particular receptor. The agent bound by or reacting with a receptor is called a "ligand," a term which is definitionally meaningful only in terms of its counterpart receptor. The term "ligand" does not imply any particular molecular size or other structural or compositional feature other than that the substance in question is capable of binding or otherwise interacting with the receptor. Also, a ligand may serve either as the natural ligand to which the receptor binds, or as a functional analogue that may act as an agonist or antagonist. Examples of ligands that can be investigated by this invention include, but are not restricted to, agonists and antagonists for cell

membrane receptors, toxins and venoms, viral epitopes, hormones (e.g., opiates, steroids, etc.), hormone receptors, peptides, enzymes, enzyme substrates, substrate analogs, transition state analogs, cofactors, drugs, proteins, and antibodies.

Linkage disequilibrium or allelic association means the preferential association of a particular allele or genetic marker with a specific allele, or genetic marker at a nearby chromosomal location more frequently than expected by chance for any particular allele frequency in the population. For example, if locus X has alleles a and b, which occur equally frequently, and linked locus Y has alleles c and d, which occur equally frequently, one would expect the combination ac to occur with a frequency of 0.25. If ac occurs more frequently, then alleles a and c are in linkage disequilibrium. Linkage disequilibrium may result from natural selection of certain combination of alleles or because an allele has been introduced into a population too recently to have reached equilibrium with linked alleles.

Microtiter plates are arrays of discrete wells that come in standard formats (96, 384 and 1536 wells) which are used for examination of the physical, chemical or biological characteristics of a quantity of samples in parallel.

"Mixed population" or "complex population" refers to any sample containing both desired and undesired nucleic acids. As a non-limiting example, a complex population of nucleic acids may be total genomic DNA, total genomic RNA or a combination thereof. Moreover, a complex population of nucleic acids may have been enriched for a given population but include other undesirable populations. For example, a complex population of nucleic acids may be a sample which has been enriched for desired messenger RNA (mRNA) sequences but still includes some undesired ribosomal RNA sequences (rRNA).

"Monomer" refers to any member of the set of molecules that can be joined together to form an oligomer or polymer. The set of monomers useful in the present invention includes, but is not restricted to, for the example of (poly)peptide synthesis, the set of L-amino acids, D-amino acids, or synthetic amino acids. As used herein, "monomer" refers to any member of a basis set for synthesis of an oligomer. For example, dimers of L-amino acids form a basis set of 400 "monomers" for synthesis of polypeptides. Different basis sets of monomers may be used at successive steps in the synthesis of a polymer. The term "monomer" also

refers to a chemical subunit that can be combined with a different chemical subunit to form a compound larger than either subunit alone.

"mRNA" or "mRNA transcripts" as used herein, include, but not limited to pre-mRNA transcript(s), transcript processing intermediates, mature mRNA(s) ready for translation and transcripts of the gene or genes, or nucleic acids derived from the mRNA transcript(s). Transcript processing may include splicing, editing and degradation. As used herein, a nucleic acid derived from an mRNA transcript refers to a nucleic acid for whose synthesis the mRNA transcript or a subsequence thereof has ultimately served as a template. Thus, a cDNA reverse transcribed from an mRNA, an RNA transcribed from that cDNA, a DNA amplified from the cDNA, an RNA transcribed from the amplified DNA, etc., are all derived from the mRNA transcript and detection of such derived products is indicative of the presence and/or abundance of the original transcript in a sample. Thus, mRNA derived samples include, but are not limited to, mRNA transcripts of the gene or genes, cDNA reverse transcribed from the mRNA, cRNA transcribed from the cDNA, DNA amplified from the genes, RNA transcribed from amplified DNA, and the like.

Nucleic acid library or array is an intentionally created collection of nucleic acids which can be prepared either synthetically or biosynthetically and screened for biological activity in a variety of different formats (e.g., libraries of soluble molecules; and libraries of oligos tethered to resin beads, silica chips, or other solid supports). Additionally, the term "array" is meant to include those libraries of nucleic acids which can be prepared by spotting nucleic acids of essentially any length (e.g., from 1 to about 1000 nucleotide monomers in length) onto a substrate. The term "nucleic acid" as used herein refers to a polymeric form of nucleotides of any length, either ribonucleotides, deoxyribonucleotides or peptide nucleic acids (PNAs), that comprise purine and pyrimidine bases, or other natural, chemically or biochemically modified, non-natural, or derivatized nucleotide bases. The backbone of the polynucleotide can comprise sugars and phosphate groups, as may typically be found in RNA or DNA, or modified or substituted sugar or phosphate groups. A polynucleotide may comprise modified nucleotides, such as methylated nucleotides and nucleotide analogs. The sequence of nucleotides may be interrupted by non-nucleotide components. Thus the terms nucleoside, nucleotide, deoxynucleoside

and deoxynucleotide generally include analogs such as those described herein. These analogs are those molecules having some structural features in common with a naturally occurring nucleoside or nucleotide such that when incorporated into a nucleic acid or oligonucleoside sequence, they allow hybridization with a naturally occurring nucleic acid sequence in solution. Typically, these analogs are derived from naturally occurring nucleosides and nucleotides by replacing and/or modifying the base, the ribose or the phosphodiester moiety. The changes can be tailor made to stabilize or destabilize hybrid formation or enhance the specificity of hybridization with a complementary nucleic acid sequence as desired.

Nucleic acids according to the present invention may include any polymer or oligomer of pyrimidine and purine bases, preferably cytosine, thymine, and uracil, and adenine and guanine, respectively. See Albert L. Lehninger, PRINCIPLES OF BIOCHEMISTRY, at 793-800 (Worth Pub. 1982). Indeed, the present invention contemplates any deoxyribonucleotide, ribonucleotide or peptide nucleic acid component, and any chemical variants thereof, such as methylated, hydroxymethylated or glucosylated forms of these bases, and the like. The polymers or oligomers may be heterogeneous or homogeneous in composition, and may be isolated from naturally-occurring sources or may be artificially or synthetically produced. In addition, the nucleic acids may be DNA or RNA, or a mixture thereof, and may exist permanently or transitionally in single-stranded or double-stranded form, including homoduplex, heteroduplex, and hybrid states.

An "oligonucleotide" or "polynucleotide" is a nucleic acid ranging from at least 2, preferable at least 8, and more preferably at least 20 nucleotides in length or a compound that specifically hybridizes to a polynucleotide. Polynucleotides of the present invention include sequences of deoxyribonucleic acid (DNA) or ribonucleic acid (RNA) which may be isolated from natural sources, recombinantly produced or artificially synthesized and mimetics thereof. A further example of a polynucleotide of the present invention may be peptide nucleic acid (PNA). The invention also encompasses situations in which there is a nontraditional base pairing such as Hoogsteen base pairing which has been identified in certain tRNA molecules and postulated to exist in a triple helix. "Polynucleotide" and "oligonucleotide" are used interchangeably in this application.

"Optically clear" refers to the property of a material for transmitting light waves with a minimum loss of intensity or attenuation of the light.

Probe: A probe is a surface-immobilized molecule that can be recognized by a particular target. Examples of probes that can be investigated by this invention include, but are not restricted to, agonists and antagonists for cell membrane receptors, toxins and venoms, viral epitopes, hormones (e.g., opioid peptides, steroids, etc.), hormone receptors, peptides, enzymes, enzyme substrates, cofactors, drugs, lectins, sugars, oligonucleotides, nucleic acids, oligosaccharides, proteins, and monoclonal antibodies.

"Primer" is a single-stranded oligonucleotide capable of acting as a point of initiation for template-directed DNA synthesis under suitable conditions e.g., buffer and temperature, in the presence of four different nucleoside triphosphates and an agent for polymerization, such as, for example, DNA or RNA polymerase or reverse transcriptase. The length of the primer, in any given case, depends on, for example, the intended use of the primer, and generally ranges from 15 to 30 nucleotides. Short primer molecules generally require cooler temperatures to form sufficiently stable hybrid complexes with the template. A primer need not reflect the exact sequence of the template but must be sufficiently complementary to hybridize with such template. The primer site is the area of the template to which a primer hybridizes. The primer pair is a set of primers including a 5' upstream primer that hybridizes with the 5' end of the sequence to be amplified and a 3' downstream primer that hybridizes with the complement of the 3' end of the sequence to be amplified.

Polymorphism refers to the occurrence of two or more genetically determined alternative sequences or alleles in a population. A polymorphic marker or site is the locus at which divergence occurs. Preferred markers have at least two alleles, each occurring at frequency of greater than 1%, and more preferably greater than 10% or 20% of a selected population. A polymorphism may comprise one or more base changes, an insertion, a repeat, or a deletion. A polymorphic locus may be as small as one base pair. Polymorphic markers include restriction fragment length polymorphisms, variable number of tandem repeats (VNTR's), hypervariable regions, minisatellites, dinucleotide repeats, trinucleotide repeats, tetranucleotide

repeats, simple sequence repeats, and insertion elements such as Alu. The first identified allelic form is arbitrarily designated as the reference form and other allelic forms are designated as alternative or variant alleles. The allelic form occurring most frequently in a selected population is sometimes referred to as the wildtype form. Diploid organisms may be homozygous or heterozygous for allelic forms. A diallelic polymorphism has two forms. A triallelic polymorphism has three forms. Single nucleotide polymorphisms (SNPs) are included in polymorphisms.

Receptor: A molecule that has an affinity for a given ligand. Receptors may be naturally-occurring or manmade molecules. Also, they can be employed in their unaltered state or as aggregates with other species. Receptors may be attached, covalently or noncovalently, to a binding member, either directly or via a specific binding substance. Examples of receptors which can be employed by this invention include, but are not restricted to, antibodies, cell membrane receptors, monoclonal antibodies and antisera reactive with specific antigenic determinants (such as on viruses, cells or other materials), drugs, polynucleotides, nucleic acids, peptides, cofactors, lectins, sugars, polysaccharides, cells, cellular membranes, and organelles. Receptors are sometimes referred to in the art as anti-ligands. As the term receptors is used herein, no difference in meaning is intended. A "Ligand Receptor Pair" is formed when two macromolecules have combined through molecular recognition to form a complex. Other examples of receptors which can be investigated by this invention include but are not restricted to those molecules shown in U.S. Patent No. 5,143,854, which is hereby incorporated by reference in its entirety.

"Solid support", "support", and "substrate" are used interchangeably and refer to a material or group of materials having a rigid or semi-rigid surface or surfaces. In many embodiments, at least one surface of the solid support will be substantially flat, although in some embodiments it may be desirable to physically separate synthesis regions for different compounds with, for example, wells, raised regions, pins, etched trenches, or the like. According to other embodiments, the solid support(s) will take the form of beads, resins, gels, microspheres, or other geometric configurations. See U.S. Patent No. 5,744,305 for exemplary substrates.

Suspended refers to the state of floating without sinking or falling.

Surface or target surface refers to the area of the microarray to be analyzed.

Target: A molecule that has an affinity for a given probe. Targets may be naturally-occurring or man-made molecules. Also, they can be employed in their unaltered state or as aggregates with other species. Targets may be attached, covalently or noncovalently, to a binding member, either directly or via a specific binding substance. Examples of targets which can be employed by this invention include, but are not restricted to, antibodies, cell membrane receptors, monoclonal antibodies and antisera reactive with specific antigenic determinants (such as on viruses, cells or other materials), drugs, oligonucleotides, nucleic acids, peptides, cofactors, lectins, sugars, polysaccharides, cells, cellular membranes, and organelles. Targets are sometimes referred to in the art as anti-probes. As the term targets is used herein, no difference in meaning is intended. A "Probe Target Pair" is formed when two macromolecules have combined through molecular recognition to form a complex.

Wafer: A substrate having surface to which a plurality of arrays are bound. In a preferred embodiment, the arrays are synthesized on the surface of the substrate to create multiple arrays that are physically separate. In one preferred embodiment of a wafer, the arrays are physically separated by a distance of at least about 0.1, 0.25, 0.5, 1 or 1.5 millimeters. The arrays that are on the wafer may be identical, each one may be different, or there may be some combination thereof. Particularly preferred wafers are about 8" x 8" and are made using the photolithographic process.

A Well Plate or Plate: A body having a plurality of arrays in which each microarray is separated by a physical barrier resistant to the passage of liquids and forming an area or space, referred to as a well.

III. Chip In a Well ™ Scanning

Figure 1 depicts an embodiment of the current application. One embodiment of the invention uses microarrays (102) that can be individual microrrrays selected from diced wafers (101). The individual microarrays (102) are then placed in a well (105) of a well plate (103) and contained in the well (105) of the well plate (103) with the use of an optically clear window (104), which can be made of, for example, fused silica, and successfully scanned. An example of a well plate used in the methods of this invention is based on a standard 96-well microtiter plates. The well plates of the current invention have a plurality of wells that can be arrayed in a

variety of ways. In one embodiment of the invention, the individual microarrays (102) do not need to be fixed in the well (105). In a further embodiment of the invention, the individual microarrays (102) can be suspended into the wells (105) with or without a buffer.

In a further embodiment of the invention depicted in Figure 2 of the current application uses microarrays (202) which can be individual microarrays selected from diced wafers (201). The individual microarrays (202) are then placed in the well (204) of a cartridge (203) and contained in the well (204) with the use of an optically clear window (205), which can be made of, for example, fused silica, and successfully scanned. In one embodiment of the invention, the individual microarrays (202) do not need to be fixed in the well (204). In a further embodiment of the invention, the individual microarrays (202) can be suspended into the wells (204) with or without a buffer.

It is to be understood that the description is intended to be illustrative and not restrictive. Many variations of the invention will be apparent to those of skill in the art upon reviewing the above description. The scope of the invention should be determined with reference to the appended claims, along with the full scope of equivalents to which such claims are entitled. All cited references, including patent and non-patent literature, are incorporated herewith by reference in their entireties for all purposes.

CLAIMS

What is claimed is:

1. A method of packaging a microarray comprising:

placing a microarray in a well of a well plate; wherein the microarray is suspended in the well.

2. The method of claim 1, wherein the well plate has one or more wells.

3. The method of claim 1, wherein the microarray is suspended in the well with buffer.

4. The method of claim 1, wherein the microarray is suspended in the well without buffer.

5. The method of claim 1, wherein the plate has an optically clear window.

6. The method of claim 5, wherein the clear window is made of fused silica.

7. A method of scanning a microarray chip comprising:

dicing one or more wafers into individual microarrays;

placing the individual microarrays into a well of a well plate; wherein the individual microarrays are contained in the well in a suspended form with an optically clear window;

scanning the microarrays through the optically clear window.

8. The method of scanning a microarray of claim 7, wherein the optically clear window is made of fused silica.

9. The method of scanning a microarray of claim 7, wherein the scanning is done with the surface of the individual microarray up.

10. The method of scanning a microarray of claim 7, wherein the scanning is done with the surface of the individual microarray down.

11. The method of scanning a microarray of claim 7, wherein the well plate has one or more wells.