SPECIFICATION

Polycyclic Carbamoylpyridone Derivative Having HIV Integrase Inhibitory Activity

[Technical Field]

[0001]

The present invention relates to novel compounds possessing an antiviral activity, in detail polycyclic carbamoylpyridone derivatives possessing an inhibitory activity against HIV integrase and a pharmaceutical composition containing the same, especially an anti-HIV agent.

[Background Art]

[0002]

Among viruses, human immunodeficiency virus (HIV), a kind of retrovirus, is known to cause acquired immunodeficiency syndrome (AIDS). The therapeutic agent for AIDS is mainly selected from a group of reverse transcriptase inhibitors (e.g., AZT, 3TC) and protease inhibitors (e.g., Indinavir), but they are proved to be accompanied by side effects such as nephropathy and the emergence of resistant viruses. Thus, the development of anti-HIV agents having the other mechanism of action has been desired.

On the other hand, a combination therapy is reported to be efficient in treatment for AIDS because of the frequent emergence of the resistant mutant. Reverse transcriptase inhibitors and protease inhibitors are clinically used as an anti-HIV agent, however agents having the same mechanism of action often exhibit cross-resistance or only an additional activity. Therefore, anti-HIV agents having the other mechanism of action are desired.

Under the circumstances above, an HIV integrase inhibitor has been focused on as an anti-HIV agent having a novel mechanism of action (Ref: Patent Documents 1 and 2) . As an anti-HIV agent having such a mechanism of action, known are carbamoyl-substituted hydroxypyrimidinone derivative (Ref: Patent Documents 3 and 4) and carbamoyl-substituted hydroxypyrrolidione derivative (Ref: Patent Document 5). Further, a patent application concerning carbamoyl-substituted hydroxypyridone derivative has been filed (Ref: Patent Document 6, Example 8) .

Other known carbamoylpyridone derivatives include 5-alkoxypyridine-3-carboxamide derivatives and γ-pyrone-3-carboxamide derivatives, which are a plant growth inhibitor or herbicide (Ref: Patent Documents 7-9).

Other HIV integrase inhibitors include N-containing condensed cyclic compounds

(Ref: Patent Document 10).

[Patent Document 1]WO03/0166275

[Patent Document 2]WO2004/024693

[Patent Document 3]WO03/035076

[Patent Document 4]WO03/035076

[Patent Document 5]WO2004/004657

[Patent Document 6]JP Patent Application 2003-32772

[Patent Document 7]JP Patent Publication 1990-108668

[Patent Document 8]JP Patent Publication 1990-108683

[Patent Document 9]JP Patent Publication 1990-96506

[Patent Document 10]WO2005/016927

[Disclosure of Invention]

[Problem to be Solved by the Invention]

[0003]

The development of a novel integrase inhibitor has been desired.


[Means to Solve the Problem]

[0004]

The present inventors have intensively studied to find that a novel polycyclic carbamoylpyridone derivative possesses a potent HIV integrase inhibitory activity.

Moreover, the present inventors have discovered that a compound of the present invention and a pharmaceutical composition containing the same are useful as an antiviral agent, an antiretroviral agent, an anti-HIV agent, an anti-HTLV-1 (Human T cell leukemia virus type 1) agent, an anti-FIV (Feline immunodeficiency virus) agent or an anti-SIV (Simian immunodeficiency virus) agent, especially an anti-HIV agent or anti-AIDS agent, to accomplish the present invention shown below.

(1)A compound of the formula:



(I)

(wherein,

Z^1 is NR^4;

R^4 is hydrogen, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted cycloalkyl lower alkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted aryl, optionally substituted aryl lower alkyl, optionally substituted aryloxy, optionally substituted heterocyclic group, optionally substituted heterocycle lower alkyl, optionally substituted heterocycleoxy, hydroxy, optionally substituted amino, optionally substituted phosphoric acid residue, aryl substituted with optionally substituted phosphoric acid residue, aralkyl substituted with optionally substituted phosphoric acid residue, hydroxy substituted with optionally substituted phosphoric acid residue, amino substituted with optionally substituted phosphoric acid residue or lower alkyl substituted with optionally substituted phosphoric acid residue (the lower alkyl may be intervened by a heteroatom group selected from CO, O, S, SO, SO_2, NR^a (R^a is hydrogen or lower alkyl), -N= and =N-)), O or CH_2;

Z^2 is optionally substituted lower alkylene or optionally substituted lower alkenylene, each may be intervened by a heteroatom group selected from O, S, SO, SO_2, NR^5 (R^5 is hydrogen, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted cycloalkyl lower alkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted aryl, optionally substituted aryl lower alkyl, optionally substituted aryloxy, optionally substituted heterocyclic group, optionally substituted heterocycle lower alkyl, optionally substituted heterocycleoxy, hydroxy or optionally substituted amino, optionally substituted phosphoric acid residue, aryl substituted with optionally substituted phosphoric acid residue, aralkyl substituted with optionally substituted phosphoric acid residue, hydroxy substituted with optionally substituted phosphoric acid residue, amino substituted with optionally substituted phosphoric acid residue or lower alkyl substituted with optionally substituted phosphoric acid residue (the lower alkyl may be intervened by a heteroatom group selected from CO, O, S, SO, SO_2, NR^5 (R^5 is selected independently from the same substituent group as R^4), -N= and =N-)), -N= or =N-

R^1 is hydrogen or lower alkyl;


X is a single bond, a heteroatom group selected from O, S, SO, SO_2 and NH, or lower alkylene or lower alkenylene each may be intervened by the heteroatom;

R^2 is optionally substituted aryl;

R^3 is hydrogen, halogen, hydroxy, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted lower alkenyl, optionally substituted

lower alkoxy, optionally substituted lower alkenyloxy, optionally substituted aryl, optionally substituted aryloxy, optionally substituted heterocyclic group, optionally substituted heterocycleoxy or optionally substituted amino;

R^4 and Z^2 part taken together forms a ring, where the compound (I) is represented by the following formula (I-1), or (I-11):



(I-1)

(wherein,

A ring is optionally substituted heterocycle;

R^{14} and R^X are independently hydrogen, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted cycloalkyl lower alkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted lower alkenyloxy, optionally substituted aryl, optionally substituted aryl lower alkyl, optionally substituted aryloxy, optionally substituted heterocyclic group, optionally substituted heterocycle lower alkyl, optionally substituted heterocycleoxy, optionally substituted phosphoric acid residue, aryl substituted with optionally substituted phosphoric acid residue, aralkyl substituted with optionally substituted phosphoric acid residue, hydroxy substituted with optionally substituted phosphoric acid residue, amino substituted with optionally substituted phosphoric acid residue or lower alkyl substituted with optionally substituted phosphoric acid residue (the lower alkyl may be intervened by a heteroatom group selected from O, S, SO, SO_2, NR^5 (R^5 is selected independently from the same substituent group as R^4), -N= and =N-), hydroxy, optionally substituted amino, optionally substituted lower alkyl carbonyl, optionally substituted cycloalkylcarbonyl, optionally substituted cycloalkyl lower alkyl carbonyl, optionally substituted lower alkoxy carbonyl, optionally substituted arylcarbonyl, optionally substituted aryl lower alkyl carbonyl, optionally substituted aryloxycarbonyl, optionally substituted heterocyclecarbonyl, optionally substituted heterocycle lower alkyl carbonyl, optionally substituted heterocycleoxy carbonyl or optionally substituted aminocarbonyl;


a broken line represents the presence or absence of a bond, provided that when the broken line represents the presence of a bond, R^X is not present;

R^1 is hydrogen or lower alkyl;

X is a single bond, a heteroatom group selected from O, S, SO, SO_2 and NH, or lower alkylene or lower alkenylene each may be intervened by the heteroatom group;

R^2 is optionally substituted aryl;

R^3 is hydrogen, halogen, hydroxy, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted lower alkenyloxy, optionally substituted aryl, optionally substituted aryloxy, optionally substituted heterocyclic group, optionally substituted heterocycleoxy or optionally substituted amino)



(I-11)

(wherein,

D ring is optionally substituted heterocycle;

R^1 is hydrogen or lower alkyl;

X is a single bond, a heteroatom group selected from O, S, SO, SO_2 and NH, or lower alkylene or lower alkenylene each may be intervened by the heteroatom group;

R^2 is optionally substituted aryl;

R^3 is hydrogen, halogen, hydroxy, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted lower alkenyloxy, optionally substituted aryl, optionally substituted aryloxy, optionally substituted heterocyclic group, optionally substituted heterocycleoxy or optionally substituted amino)), its pharmaceutically acceptable salt, or solvate thereof.

(2) A compound according to the above (1), pharmaceutically acceptable salt, or solvate thereof, wherein R^1 is hydrogen.

(3) A compound according to the above (1), pharmaceutically acceptable salt, or


solvate thereof, wherein X is lower alkylene; R^2 is phenyl or phenyl substituted with at least halogen.

(4) A compound according to the above (1), pharmaceutically acceptable salt, or solvate thereof, wherein R^3 is hydrogen, halogen, hydroxy, lower alkyl, lower alkenyl, lower alkoxy, lower alkenyloxy or optionally substituted amino.

(5) A compound according to the above (1), pharmaceutically acceptable salt, or solvate thereof, wherein R^3 is hydrogen.

(6) A compound according to the above (1), pharmaceutically acceptable salt, or solvate thereof, wherein R^1 is hydrogen or lower alkyl; X is lower alkylene; R^2 is phenyl or phenyl substituted with at least halogen; R^3 is hydrogen, halogen, hydroxy, lower alkyl, lower alkenyl, lower alkoxy, lower alkenyloxy or optionally substituted amino.

(7) A compound of the formula:



(I-1)

(wherein,

A ring is optionally substituted heterocycle;

R^{14} and R^X are independently hydrogen, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted cycloalkyl lower alkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted lower alkenyloxy, optionally substituted aryl, optionally substituted aryl lower alkyl, optionally substituted aryloxy, optionally substituted heterocyclic group, optionally substituted heterocycle lower alkyl, optionally substituted heterocycleoxy, optionally substituted phosphoric acid residue, aryl substituted with optionally substituted phosphoric acid residue, aralkyl substituted with optionally substituted phosphoric acid residue, hydroxy substituted with optionally substituted phosphoric acid residue, amino substituted with optionally substituted phosphoric acid residue or

lower alkyl substituted with optionally substituted phosphoric acid residue (the lower alkyl may be intervened by a heteroatom group selected from O, S, SO, SO_2, NR^5 (R^5 is selected independently from the same substituent group as R^4), -N= and =N-), hydroxy, optionally substituted amino, optionally substituted lower alkyl carbonyl, optionally substituted cycloalkylcarbonyl, optionally substituted cycloalkyl lower alkyl carbonyl, optionally substituted lower alkoxy carbonyl, optionally substituted arylcarbonyl, optionally substituted aryl lower alkyl carbonyl, optionally substituted aryloxycarbonyl, optionally substituted heterocyclecarbonyl, optionally substituted heterocycle lower alkyl carbonyl, optionally substituted heterocycleoxy carbonyl or optionally substituted aminocarbonyl;

a broken line represents the presence or absence of a bond, provided that when the broken line represents the presence of a bond, R^X is not present;

R^1 is hydrogen or lower alkyl;

X is a single bond, a heteroatom group selected from O, S, SO, SO_2 and NH, or lower alkylene or lower alkenylene each may be intervened by the heteroatom group;

R^2 is optionally substituted aryl;

R^3 is hydrogen, halogen, hydroxy, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted lower alkenyloxy, optionally substituted aryl, optionally substituted aryloxy, optionally substituted heterocyclic group, optionally substituted heterocycleoxy or optionally substituted amino), its pharmaceutically acceptable salt, or solvate thereof


(8) A compound according to the above (7), pharmaceutically acceptable salt, or solvate thereof, wherein R^1 is hydrogen or lower alkyl; X is lower alkylene; R^2 is phenyl or phenyl substituted with at least halogen; R^3 is hydrogen, halogen, hydroxy, lower alkyl, lower alkenyl, lower alkoxy, lower alkenyloxy or optionally substituted amino.

(9) A compound according to the above (7), pharmaceutically acceptable salt, or solvate thereof, wherein a broken line represents the absence of a bond.

(10) A compound according to the above (7), pharmaceutically acceptable salt, or solvate thereof, wherein R^X is hydrogen; R^{14} is hydrogen or optionally substituted lower alkyl.

(11) A compound according to the above (7), pharmaceutically acceptable salt, or solvate thereof, wherein A ring is an optionally substituted and optionally condensed 5- to 7- membered heterocycle containing 1 to 2 hetero atom(s).

(12) A compound of the formula:



(I-1-1)

(wherein,

A ring is an optionally substituted and optionally condensed 5- to 7- membered heterocycle containing 1 to 2 hetero atom(s);

the stereochemistry of an asymmetric carbon represented by * shows R- or S-configuration, or a mixture thereof;

R^{14} and R^X are independently hydrogen, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted cycloalkyl lower alkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted lower alkenyloxy, optionally substituted aryl, optionally substituted aryl lower alkyl, optionally substituted aryloxy, optionally substituted heterocyclic group, optionally substituted heterocycle lower alkyl, optionally substituted heterocycleoxy, optionally substituted phosphoric acid residue, aryl substituted with optionally substituted phosphoric acid residue, aralkyl substituted with optionally substituted phosphoric acid residue, hydroxy substituted with optionally substituted phosphoric acid residue, amino substituted with optionally substituted phosphoric acid residue or lower alkyl substituted with optionally substituted phosphoric acid residue (the lower alkyl may be intervened by a heteroatom group selected from O, S, SO, SO_2, NR^5 (R^5 is selected independently from the same substituent group as R^4), -N= and =N-), hydroxy, optionally substituted amino, optionally substituted lower alkyl carbonyl, optionally substituted cycloalkylcarbonyl, optionally substituted cycloalkyl lower alkyl carbonyl, optionally substituted lower alkoxy carbonyl, optionally substituted arylcarbonyl, optionally substituted aryl lower alkyl carbonyl, optionally substituted

aryloxycarbonyl, optionally substituted heterocyclecarbonyl, optionally substituted heterocycle lower alkyl carbonyl, optionally substituted heterocycleoxy carbonyl or optionally substituted aminocarbonyl;

R^8 is hydrogen, halogen, hydroxy, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted lower alkenyloxy, optionally substituted aryl, optionally substituted aryloxy, optionally substituted heterocyclic group, optionally substituted heterocycleoxy or optionally substituted amino), its pharmaceutically acceptable salt, or

R^1 is hydrogen or lower alkyl;

R is independently selected from halogen and Substituent group S1;

Substituent group S1(: optionally substituted phosphoric acid residue, aryl substituted with optionally substituted phosphoric acid residue, aralkyl substituted with optionally substituted phosphoric acid residue, hydroxy substituted with optionally substituted phosphoric acid residue, amino substituted with optionally substituted phosphoric acid residue, or lower alkyl substituted with optionally substituted phosphoric acid residue (wherein the lower alkyl may be intervened with a heteroatom group(s) selected from CO, O, O, S, SO, SO_2, NR^a (R^a is hydrogen or lower alkyl), -N= and =N-), lower alkoxy lower alkyl, amino lower alkyl optionally substituted with mono- or di- lower alkyl, halogenated lower alkyl, lower alkoxy, carbamoyl optionally substituted with mono- or di- lower alkyl, optionally substituted lower alkyl sulfonyl amino, halogenated lower alkoxy, hydroxy lower alkyl)

m is an integer of 0 to 3, its pharmaceutically acceptable salt, or solvate thereof.


(13) A compound according to the above (12), pharmaceutically acceptable salt, or solvate thereof, wherein R^x and R^{14} are independently hydrogen or optionally substituted lower alkyl.

(14) A compound according to the above (12), pharmaceutically acceptable salt, or solvate thereof, wherein R^x and R^{14} are hydrogens.

(15) A compound according to the above (12), pharmaceutically acceptable salt, or solvate thereof, wherein R^3 is hydrogen.


(16) A compound according to the above (12), pharmaceutically acceptable salt, or solvate thereof, wherein m is 0, or 1 to 3 and at least one of R is halogen.

(17) A compound according to the above (7) or (12), pharmaceutically acceptable salt, or solvate thereof, wherein A ring is any one of the following:




Z = O or NR^{26}

(A-1)



Z = O or NR^{31}

(A-2)



Z = O or NR^{40}

(A-3)

(wherein, R^{20} to R^{40} are each independently a group selected from Substituent group S2, or any two groups of R^{20} to R^{40}, which bonds to the same carbon atom, taken together with the carbon atom, may form an optionally substituted carbocyle or optionally substituted heterocycle, or each combination of (R^{20} and R^{22}), (R^{23} and R^{24}), (R^{25} and R^{26}), (R^{27} and R^{29}), (R^{30} and R^{31}), (R^{32} and R^{34}), (R^{35} and R^{36}), (R^{37} and R^{38}), and (R^{39}andR^{40}), taken together with the neighboring atom, may form an optionally substituted carbocyle or optionally substituted heterocycle.

Substituent group S2: hydrogen, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted cycloalkyl lower alkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted lower alkenyloxy, optionally substituted aryl, optionally substituted aryl lower alkyl, optionally substituted aryloxy, optionally substituted heterocycle, optionally substituted heterocycle lower alkyl, optionally substituted heterocycleoxy, hydroxy, optionally substituted amino, optionally substituted lower alkylcarbonyl, optionally substituted cycloalkylcarbonyl, optionally substituted cycloalkyl lower alkylcarbonyl, optionally substituted lower alkoxycarbonyl, optionally substituted arylcarbonyl, optionally substituted aryl lower alkylcarbonyl, optionally substituted aryl oxycarbonyl, optionally substituted heterocyclecarbonyl, optionally substituted heterocycle lower alkylcarbonyl, optionally substituted heterocycleoxycarbonyl,

optionally substituted aminocarbonyl, optionally substituted phosphoric acid residue, aryl substituted with optionally substituted phosphoric acid residue, aralkyl substituted with optionally substituted phosphoric acid residue, hydroxy substituted with optionally substituted phosphoric acid residue, amino substituted with optionally substituted phosphoric acid residue, or lower alkyl substituted with optionally substituted phosphoric acid residue (the lower alkyl may be intervened with a heteroatom group(s) selected from CO, O, S, SO, SO_2, NR^5 (R^5 is independently selected from the same Substituent group as R^4), -N= and =N-)

the stereochemistry of an asymmetric carbon represented by * shows R- or S- configuration, or a mixture thereof)

(18) A compound according to the above (17), pharmaceutically acceptable salt, or solvate thereof, wherein R^{20} to R^{40} are each independently hydrogen or substituted lower alkyl, or any two groups of R^{20} to R^{40}, which bonds to the same carbon atom, taken together with the carbon atom, may form an optionally substituted 3- to 7-membered carbocyle or optionally substituted 3- to 7- membered heterocycle, or each combination of (R^{20} and R^{22}), (R^{23} and R^{24}), (R^{25} and R^{26}), (R^{27} and R^{29}), (R^{30} and R^{31}), (R^{32} and R^{34}), (R^{35} and R^{36}), (R^{37} and R^{38}), and (R^{39}andR^{40}), taken together with the neighboring atom, may form an optionally substituted 5- to 7- membered carbocyle or optionally substituted 5- to 7- membered heterocycle.

(19) A compound according to the above (17), pharmaceutically acceptable salt, or solvate thereof, wherein A ring is a ring represented by (A-1); one of R^{20} to R^{25} is optionally substituted lower alkyl and the others are hydrogens.

(20) A compound according to the above (17), pharmaceutically acceptable salt, or solvate thereof, wherein A ring is a ring represented by (A-1); one of (R^{20} and R^{22}), (R^{23} and R^{24}), and (R^{25} and R^{26}), taken together with the neighboring atom, may form an optionally substituted 5- to 7- membered carbocyle or optionally substituted 5- to 7-membered heterocycle.

(21) A compound according to the above (17), pharmaceutically acceptable salt, or solvate thereof, wherein A ring is a ring represented by (A-1); Z=NR^{26}, and R^{25} and R^{26} taken together with the neighboring atom may form an optionally substituted 5- to 7-

membered heterocycle.

(22) A compound according to the above (17), pharmaceutically acceptable salt, or solvate thereof, wherein A ring is a ring represented by (A-2); one of R^{27} to R^{30} is optionally substituted lower alkyl and the others are hydrogens.

(23) A compound according to the above (17), pharmaceutically acceptable salt, or solvate thereof, wherein A ring is a ring represented by (A-2); one of ($R^{27}$and R^{29}) and (R^{30} and R^{31}), taken together with the neighboring atom, may form an optionally substituted 5- to 7- membered carbocyle or optionally substituted 5- to 7- membered heterocycle.

(24) A compound according to the above (17), pharmaceutically acceptable salt, or solvate thereof, wherein A ring is a ring represented by (A-2); Z=NR^{31}, and R^{30} and R^{31} taken together with the neighboring atom may form an optionally substituted 5- to 7- membered heterocycle.

(25) A compound according to the above (17), pharmaceutically acceptable salt, or solvate thereof, wherein A ring is a ring represented by (A-3); one of R^{32} to R^{39} is optionally substituted lower alkyl and the others are hydrogens.

(26) A compound according to the above (17), pharmaceutically acceptable salt, or solvate thereof, wherein A ring is a ring represented by (A-3); one of (R^{32}andR^{34}),(R^{35}andR^{36}),(R^{37}andR^{38}),and(R^{39}andR^{40}), taken together with the neighboring atom, may form an optionally substituted 5- to 7- membered carbocyle or optionally substituted 5- to 7- membered heterocycle.


(27) A compound according to the above (17), pharmaceutically acceptable salt, or solvate thereof, wherein A ring is a ring represented by (A-3); Z=NR^{40}, and R^{39} and R^{40} taken together with the neighboring atom may form an optionally substituted 5- to 7- membered heterocycle.

(28) A compound according to the above (12), pharmaceutically acceptable salt, or solvate thereof, wherein R^X is hydrogen; R^{14} is hydrogen or optionally substituted lower alkyl;R^3 is hydrogen; m is 1 to 3 and at least one of Rs is halogen; A ring is a ring

described in the above (17).

(29) A compound according to the above (12), pharmaceutically acceptable salt, or solvate thereof, wherein R^X is hydrogen; R^{14} is hydrogen; R^3 is hydrogen; m is 0, or 1 to 3 and at least one of Rs is halogen; A ring is a ring described in the above (17); R^{20} to R^{40} are each independently hydrogen or substituted lower alkyl, or any two groups of R^{20} to R^{40}, which bonds to the same carbon atom, taken together with the carbon atom, may form an optionally substituted 3- to 7- membered carbocyle or optionally substituted 3- to 7- membered heterocycle, or each combination of (R^{20} and R^{22}), (R^{23} and R^{24}), (R^{25} and R^{26}), (R^{27} and R^{29}), (R^{30} and R^{31}), (R^{32} and R^{34}), (R^{35} and R^{36}), (R^{37} and R^{38}), and (R^{39}andR^{40}), taken together with the neighboring carbon atom, may form an optionally substituted 5- to 7- membered carbocyle or optionally substituted 5- to 7- membered heterocycle.

(30) A compound of the formula:



(I-11)

(wherein,

D ring is optionally substituted heterocycle;

R^1 is hydrogen or lower alkyl;

X is a single bond, a heteroatom group selected from O, S, SO, SO_2 and NH, or lower alkylene or lower alkenylene each may be intervened by the heteroatom group;

R^2 is optionally substituted aryl;

R^3 is hydrogen, halogen, hydroxy, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted lower alkenyloxy, optionally substituted aryl, optionally substituted aryloxy, optionally substituted heterocyclic group, optionally substituted heterocycleoxy or optionally substituted amino), pharmaceutically acceptable salt, or solvate thereof

(31) A compound selected from the group consisting of:

(3*R*,11a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-methyl-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(4a*R*,13a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-10-hydroxy-9,11-dioxo-2,3,4a,5,9,11,13,13a-octahydro-1*H*-pyrido[1,2-*a*]pyrrolo[1',2':3,4]imidazo[1,2-*d*]pyrazine-8-carboxamide;

(3a*S*,13a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-8-hydroxy-7,9-dioxo-1,2,3,3a,4,5,7,9,13,13a-decahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrrolo[1,2-*c*]pyrimidine-10-carboxamide;

(4a*S*,13a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-10-hydroxy-9,11-dioxo-2,3,4a,5,9,11,13,13a-octahydro-1*H*-pyrido[1,2-*a*]pyrrolo[1',2':3,4]imidazo[1,2-*d*]pyrazine-8-carboxamide;

(4a*S*,13a*R*)-*N*-[(4-Fluorophenyl)methyl]-10-hydroxy-9,11-dioxo-2,3,4a,5,9,11,13,13a-octahydro-1*H*-pyrido[1,2-*a*]pyrrolo[1',2':3,4]imidazo[1,2-*d*]pyrazine-8-carboxamide;

(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-5,7-dioxo-3-(phenylmethyl)-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3a*S*,13a*S*)-*N*-[(4-Fluorophenyl)methyl]-8-hydroxy-7,9-dioxo-1,2,3,3a,4,5,7,9,13,13a-decahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrrolo[1,2-*c*]pyrimidine-10-carboxamide;

(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-[(1*S*)-1-methylpropyl]-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-methyl-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3*S*,11a*R*)-*N*-[(4-Fluorophenyl)methyl]-6-hydroxy-3-methyl-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-3-(1,1-dimethylethyl)-6-hydroxy-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3*S*,11a*R*)-3-(1,1-Dimethylethyl)-*N*-[(4-fluorophenyl)methyl]-6-hydroxy-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-5,7-dioxo-3-phenyl-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-(hydroxymethyl)-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(2*S*,3*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-methyl-5,7-dioxo-2-phenyl-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3*R*,11a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-5,7-dioxo-3-(phenylmethyl)-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3*R*,11a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-(2-methylpropyl)-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(5a*R*,14a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-11-hydroxy-10,12-dioxo-1,2,3,4,5a,6,10,12,14,14a-decahydropyrido[1,2-*a*]pyrido[1',2':3,4]imidazo[1,2-*d*]pyrazine-9-carboxamide;

(2*S*,3*S*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-[(methyloxy)methyl]-5,7-dioxo-2-phenyl-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3*S*,11a*R*)-3-(Cyclohexylmethyl)-*N*-[(2,4-difluorophenyl)methyl]-6-hydroxy-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-(1-methylethyl)-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(5a*R*,14a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-12-hydroxy-11,13-dioxo-5a,6a,7,11,13,14a-hexahydro-5*H*-indeno[1',2':4,5][1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-10-carboxamide;

(2*S*,3*R*,11a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-5,7-dioxo-2,3-diphenyl-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(2*S*,3*R*,11a*R*)-*N*-[(2,4-difluorophenyl)methyl]-6-hydroxy-5,7-dioxo-2,3-diphenyl-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3*R*,11a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-(1-methylethyl)-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-[2-(methylthio)ethyl]-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-[2-(methylsulfonyl)ethyl]-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-(1*H*-indol-3-ylmethyl)-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(4*R*,12a*R*)-*N*-[(4-fluorophenyl)methyl]-7-hydroxy-4-methyl-1-(2-methylpropyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(4*R*,12a*R*)-*N*-[(4-Fluorophenyl)methyl]-7-hydroxy-4-methyl-1-(1-methylethyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-4-methyl-1-(2-methylpropyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(4*S*,12a*S*)-1-(Cyclopropylmethyl)-*N*-[(2,4-difluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-1-(2-furanylmethyl)-7-hydroxy-4-methyl-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1-(1,3-thiazol-2-ylmethyl)-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(4a*R*,6a*R*,14a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-12-hydroxy-11,13-dioxo-1,3,4,4a,5,6a,7,11,13,14a-decahydro-2*H*-pyrido[1',2':4,5]pyrazino[1,2-*a*][3,1]benzoxazine-10-carboxamide;

(4a*R*,6a*R*,14a*S*)-*N*-[(4-Fluorophenyl)methyl]-12-hydroxy-11,13-dioxo-1,3,4,4a,5,6a,7,11,13,14a-decahydro-2*H*-pyrido[1',2':4,5]pyrazino[1,2-*a*][3,1]benzoxazine-10-carboxamide

(3*S*,4a*R*,6a*R*,14a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-12-hydroxy-11,13-dioxo-3-phenyl-1,3,4,4a,5,6a,7,11,13,14a-decahydro-2*H*-pyrido[1',2':4,5]pyrazino[1,2-*a*][3,1]benzoxazine-10-carboxamide;

(4aS,6aS,14aS)-N-[(2,4-Difluorophenyl)methyl]-12-hydroxy-6-(2-methylpropyl)-11,13-dioxo-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide;

(6aR,7aS,11aS)-N-[(2,4-Difluorophenyl)methyl]-1-hydroxy-2,13-dioxo-2,6a,7,7a,8,9,10,11,11a,13-decahydro-6H-pyrido[1',2':4,5]pyrazino[1,2-a]benzimidazole-3-carboxamide;

(6aS,7aS,11aS)-N-[(2,4-Difluorophenyl)methyl]-1-hydroxy-2,13-dioxo-2,6a,7,7a,8,9,10,11,11a,13-decahydro-6H-pyrido[1',2':4,5]pyrazino[1,2-a]benzimidazole-3-carboxamide;

(5aS,14aS)-N-[(2,4-Difluorophenyl)methyl]-11-hydroxy-10,12-dioxo-1,2,3,4,5a,6,10,12,14,14a-decahydropyrido[1,2-a]pyrido[1',2':3,4]imidazo[1,2-d]pyrazine-9-carboxamide;

(4aR,14aR)-N-[(2,4-Difluorophenyl)methyl]-9-hydroxy-8,10-dioxo-2,3,4,4a,5,6,8,10,14,

14a-decahydro-1H-pyrido[1,2-c]pyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-11-carboxamide;

(4*R*,12a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-4-methyl-1-(3-methylbutyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-4-methyl-1-(1-methylethyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-4-methyl-1-(3-methylbutyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1-(3-pyridinylmethyl)-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(4*S*,12a*S*)-1-Cyclopropyl-*N*-[(2,4-difluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-4-methyl-1-[2-(methyloxy)ethyl]-

6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(3aS,5aS,13aS)-N-[(2,4-Difluorophenyl)methyl]-11-hydroxy-5-(2-methylpropyl)-10,12-dioxo-2,3,3a,4,5,5a,6,10,12,13a-decahydro-1H-cyclopenta[e]pyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-9-carboxamide;

(3*R*,11a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-3-ethyl-6-hydroxy-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(4aS,6aS,14aS)-N-[(2,4-Difluorophenyl)methyl]-12-hydroxy-6-[2-(4-morpholinyl)ethyl]-11,13-dioxo-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide;

(3aR,5aR,13aS)-N-[(2,4-Difluorophenyl)methyl]-11-hydroxy-10,12-dioxo-1,2,3,3a,4,5a,6,10,12,13a-decahydrocyclopenta[d]pyrido[1',2':4,5]pyrazino[2,1-b][1,3]oxazine-9-carboxamide;

(4aS,6aS,14aS)-N-[(2,4-Difluorophenyl)methyl]-12-hydroxy-6-methyl-11,13-dioxo-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide;

(4aS,6aS,14aS)-N-[(2,4-Difluorophenyl)methyl]-12-hydroxy-6-[2-(methyloxy)ethyl]-11,

13-dioxo-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide;

(4aS,6aS,14aS)-6-[2-(Acetylamino)ethyl]-N-[(2,4-difluorophenyl)methyl]-12-hydroxy-11,13-dioxo-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide;

(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-3-ethyl-6-hydroxy-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3*S*,11a*R*)-3-Butyl-*N*-[(2,4-difluorophenyl)methyl]-6-hydroxy-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-[(4-hydroxyphenyl)methyl]-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(4S,12aS)-1-Cyclobutyl-N-[(2,4-difluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-9-carboxamide;

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1-(tetrahydro-2*H*-thiopyran-4-yl)-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-1,4-bis(2-methylpropyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(4aS,6aS,14aS)-N-[(2,4-Difluorophenyl)methyl]-12-hydroxy-6-(2-hydroxyethyl)-11,13-dioxo-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide;

(4aS,6aS,14aS)-6-Cyclopropyl-N-[(2,4-difluorophenyl)methyl]-12-hydroxy-11,13-dioxo-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide;

(4aS,6aS,14aS)-N-[(2,4-Difluorophenyl)methyl]-12-hydroxy-11,13-dioxo-6-[2-(1-pyrrolidinyl)ethyl]-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide;

(4aS,14aS)-N-[(2,4-Difluorophenyl)methyl]-9-hydroxy-8,10-dioxo-2,3,4,4a,5,6,8,10,14,14a-decahydro-1H-pyrido[1,2-c]pyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-11-carboxamide;

(4S,12aS)-N-[(4-Fluorophenyl)methyl]-7-hydroxy-4-methyl-1-[2-(methyloxy)ethyl]-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-9-carboxamide;

(4S,12aS)-1-Cyclobutyl-N-[(4-fluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-9-carboxamide;

(4S,12aS)-N-[(4-Fluorophenyl)methyl]-7-hydroxy-4-methyl-1-(2-methylpropyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-9-carboxamide;

(4S,12aS)-N-[(4-Fluorophenyl)methyl]-7-hydroxy-1,4-dimethyl-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-9-carboxamide;

(4*S*,12a*S*)-*N*-[(4-Fluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1-(tetrahydro-2*H*-thiopyran-4-yl)-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-1,4-dimethyl-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(4*S*,12a*S*)-*N*-[(4-Fluorophenyl)methyl]-7-hydroxy-4-methyl-1-(1-methylethyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(4*S*,12a*S*)-*N*-[(4-Fluorophenyl)methyl]-7-hydroxy-1,4-bis(2-methylpropyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

enantiomers thereof; diastereomers thereof; mixtures of enantiomers thereof; mixtures of diastereomers thereof; mixtures of enantiomers and diastereomers thereof; and pharmaceutically acceptable salts thereof.

(32) A compound selected from the group consisting of:

(4a*S*,13a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-10-hydroxy-9,11-dioxo-2,3,4a,5,9,11,13,13a-octahydro-1*H*-pyrido[1,2-*a*]pyrrolo[1',2':3,4]imidazo[1,2-*d*]pyrazine-8-carboxamide;

(4a*S*,13a*R*)-*N*-[(4-Fluorophenyl)methyl]-10-hydroxy-9,11-dioxo-2,3,4a,5,9,11,13,13a-octahydro-1*H*-pyrido[1,2-*a*]pyrrolo[1',2':3,4]imidazo[1,2-*d*]pyrazine-8-carboxamide;

(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-[(1*S*)-1-methylpropyl]-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-methyl-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3*S*,11a*R*)-*N*-[(4-Fluorophenyl)methyl]-6-hydroxy-3-methyl-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-4-methyl-1-(2-methylpropyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxa

mide;

(4*S*,12a*S*)-1-(Cyclopropylmethyl)-*N*-[(2,4-difluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(4a*R*,6a*R*,14a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-12-hydroxy-11,13-dioxo-1,3,4,4a,5,6a,7,11,13,14a-decahydro-2*H*-pyrido[1',2':4,5]pyrazino[1,2-*a*][3,1]benzoxazine-10-carboxamide;

(4a*R*,6a*R*,14a*S*)-*N*-[(4-Fluorophenyl)methyl]-12-hydroxy-11,13-dioxo-1,3,4,4a,5,6a,7,11,13,14a-decahydro-2*H*-pyrido[1',2':4,5]pyrazino[1,2-*a*][3,1]benzoxazine-10-carboxamide;

(4S,9aR)-5-Hydroxy-4-methyl-6,10-dioxo-3,4,6,9,9a,10-hexahydro-2H-1-oxa-4a,8a-diaza-anthracene-7-carboxylic acid 2, 4-difluorobenzylamide.

(4R,9aS)-5-Hydroxy-4-methyl-6,10-dioxo-3,4,6,9,9a,10-hexahydro-2H-1-oxa-4a,8a-diaza-anthracene-7-carboxylic acid 2, 4-difluorobenzylamide.

(2R,9aS)-5-Hydroxy-2-methyl-6,10-dioxo-3,4,6,9,9a,10-hexahydro-2H-1-oxa-4a,8a-diaza-anthracene-7-carboxylic acid 4-fluorobenzylamide.


enantiomers thereof; diastereomers thereof; mixtures of enantiomers thereof; mixtures of diastereomers thereof; mixtures of enantiomers and diastereomers thereof; and pharmaceutically acceptable salts thereof.

(33) A compound according to the above (31) or (32) wherein the pharmaceutically acceptable salt is a sodium salt.

(34) A pharmaceutical composition comprising a compound according to any one of the above (1) to (33), or a pharmaceutically acceptable salt, or solvate thereof.

(35) A pharmaceutical composition according to the above (34), which is an anti-HIV agent.


(36) A process for the preparation of a compound of formula (I-20a)



(I-20a)


wherein R^e is one or two halogen; R^z is C_{1-8}alkyl, C_{6-14}arylC_{1-8}alkyl, C_{6-14}aryl, or alkoxy; and P^1 is C_{6-14}arylC_{1-8}alkyl;

comprising condensing a compound of the formula




wherein R^e is one or two halogen; R^{50} is C_{1-8}alkyl; and P^1 is C_{6-14}arylC_{1-8}alkyl;

with a compound of the formula



wherein R^z is C_{1-8}alkyl, C_{6-14}arylC_{1-8}alkyl, C_{6-14}aryl, or alkoxy;
to form a compound of formula (I-20a).


(37) A process for the preparation of a compound of formula (I-20b)




(I-20b)

wherein R^e is one or two halogen; R^z is C_{1-8}alkyl, C_{6-14}arylC_{1-8}alkyl, C_{6-14}aryl, or alkoxy; and P^1 is C_{6-14}arylC_{1-8}alkyl;

comprising condensing a compound of the formula




wherein R^e is one or two halogen; R^{50} is C_{1-8}alkyl; and P^1 is C_{6-14}arylC_{1-8}alkyl;
with a compound of the formula



wherein R^z is C_{1-8}alkyl, C_{6-14}arylC_{1-8}alkyl, C_{6-14}aryl, or alkoxy;
to form a compound of formula (I-20b).


(38) A process for the preparation of a compound of formula (I-21a)




(I-21a)

wherein R^e is one or two halogen; and P^1 is C_{6-14}arylC$_{1-8}$alkyl;

comprising condensing a compound of the formula




wherein R^e is one or two halogen; R^{50} is C_{1-8}alkyl; and P^1 is C_{6-14}arylC$_{1-8}$alkyl; with a compound of the formula



to form a compound of formula (I-21a).


(39) A process for the preparation of a compound of formula (I-21b)




(I-21b)

wherein R^e is one or two halogen; and P^1 is C_{6-14}arylC$_{1-8}$alkyl;

comprising condensing a compound of the formula




wherein R^e is one or two halogen; R^{50} is C_{1-8}alkyl; and P^1 is C_{6-14}arylC_{1-8}alkyl; with a compound of the formula




to form a compound of formula (I-21b).

(40) A process for the preparation of a compound of formula (I-22a)




(I-22a)

wherein R^e is one or two halogen; and P^1 is C_{6-14}arylC_{1-8}alkyl;
comprising condensing a compound of the formula




wherein R^e is one or two halogen; R^{50} is C_{1-8}alkyl; and P^1 is C_{6-14}arylC_{1-8}alkyl; with a compound of the formula




to form a compound of formula (I-22a).

(41) A process for the preparation of a compound of formula (I-22b)




(I-22b)

wherein R^e is one or two halogen; and P^1 is $C_{6-14}arylC_{1-8}alkyl$;
comprising condensing a compound of the formula




wherein R^e is one or two halogen; R^{50} is $C_{1-8}alkyl$; and P^1 is $C_{6-14}arylC_{1-8}alkyl$;
with a compound of the formula



to form a compound of formula (I-22b).


(42) A process for the preparation of a compound of formula (I-23a)




(I-23a)

wherein R^e is one or two halogen; and P^1 is $C_{6-14}arylC_{1-8}alkyl$;

comprising condensing a compound of the formula




wherein R^e is one or two halogen; R^{50} is C_{1-8}alkyl; and P^1 is C_{6-14}arylC_{1-8}alkyl; with a compound of the formula




to form a compound of formula (I-23a).

(43) A process for the preparation of a compound of formula (I-23b)




(I-23b)

wherein R^e is one or two halogen; and P^1 is C_{6-14}arylC_{1-8}alkyl;
comprising condensing a compound of the formula




wherein R^e is one or two halogen; R^{50} is C_{1-8}alkyl;
with a compound of the formula



to form a compound of formula (I-23b).


(44) A process for the preparation of a compound of formula (I-24a)




(I-24a)

wherein R^e is one or two halogen; R^z is C_{1-8}alkyl; R^{z1} is hydrogen, C_{3-6}cycloalkyl, , heterocycle, or C_{1-8}alkyl optionally substituted with hydroxy, C_{3-6}cycloalkyl, alkoxy, heterocycle, heteroaryl, C_{6-14}aryl, or amino, wherein said amino may be optionally substituted with –C(O)C_{1-8}alkyl or C_{1-8}alkyl; and P^1 is C_{6-14}arylC_{1-8}alkyl;

comprising condensing a compound of the formula




wherein R^e is one or two halogen; and R^{50} is C_{1-8}alkyl; and P^1 is C_{6-14}arylC_{1-8}alkyl; with a compound of the formula



wherein R^z is C_{1-8}alkyl; R^{z1} is hydrogen, C_{3-6}cycloalkyl, , heterocycle, or C_{1-8}alkyl optionally substituted with hydroxy, C_{3-6}cycloalkyl, alkoxy, heterocycle, heteroaryl, C_{6-14}aryl, or amino, wherein said amino may be optionally substituted with –C(O)C_{1-8}alkyl or C_{1-8}alkyl;

to form a compound of the formula (I-24a).


(45) A process for the preparation of a compound of formula (I-24b)




(I-24b)

wherein R^e is one or two halogen; R^z is C_{1-8}alkyl; R^{z1} is hydrogen, C_{3-6}cycloalkyl, , heterocycle, or C_{1-8}alkyl optionally substituted with hydroxy, C_{3-6}cycloalkyl, alkoxy, heterocycle, heteroaryl, C_{6-14}aryl, or amino, wherein said amino may be optionally substituted with $-C(O)C_{1-8}$alkyl or C_{1-8}alkyl; and P^1 is C_{6-14}arylC_{1-8}alkyl;

comprising condensing a compound of the formula




wherein R^e is one or two halogen; R^{50} is C_{1-8}alkyl; and P^1 is C_{6-14}arylC_{1-8}alkyl;
with a compound of the formula



wherein R^z is C_{1-8}alkyl; and R^{z1} is hydrogen, C_{3-6}cycloalkyl, , heterocycle, or C_{1-8}alkyl optionally substituted with hydroxy, C_{3-6}cycloalkyl, alkoxy, heterocycle, heteroaryl, C_{6-14}aryl, or amino, wherein said amino may be optionally substituted with $-C(O)C_{1-8}$alkyl or C_{1-8}alkyl;

to form a compound of the formula (I-24b).


(46) A process for the preparation of a racemic compound of formula (I-25)




(I-25)

wherein R^e is one or two halogen; R^{z1} is hydrogen, C_{3-6}cycloalkyl, , heterocycle, or C_{1-8}alkyl optionally substituted with hydroxy, C_{3-6}cycloalkyl, alkoxy, heterocycle, heteroaryl, C_{6-14}aryl, or amino, wherein said amino may be optionally substituted with $-C(O)C_{1-8}$alkyl or C_{1-8}alkyl; and P^1 is is C_{6-14}arylC_{1-8}alkyl;

comprising condensing a compound of the formula




wherein R^e is one or two halogen; and R^{50} is C_{1-8}alkyl; and P^1 is C_{6-14}arylC_{1-8}alkyl; with a racemic compound of the formula



wherein R^{z1} is hydrogen, C_{3-6}cycloalkyl, , heterocycle, or C_{1-8}alkyl optionally substituted with hydroxy, C_{3-6}cycloalkyl, alkoxy, heterocycle, heteroaryl, C_{6-14}aryl, or amino, wherein said amino may be optionally substituted with $-C(O)C_{1-8}$alkyl or C_{1-8}alkyl;

to form a racemic compound of the formula (I-25).


(47) A process for the preparation of a racemic compound of formula (I-26)



(I-26)


wherein R^e is one or two halogen; R^{z1} is hydrogen, C_{3-6}cycloalkyl, , heterocycle, or C_{1-8}alkyl optionally substituted with hydroxy, C_{3-6}cycloalkyl, alkoxy, heterocycle, heteroaryl, C_{6-14}aryl, or amino, wherein said amino may be optionally substituted with $-C(O)C_{1-8}$alkyl or C_{1-8}alkyl; and P^1 is C_{6-14}arylC_{1-8}alkyl;

comprising condensing a compound of the formula



wherein R^e is one or two halogen; R^{50} is C_{1-8}alkyl; and P^1 is C_{6-14}arylC_{1-8}alkyl;


with a racemic compound of the formula



wherein R^{z1} is hydrogen, C_{3-6}cycloalkyl, , heterocycle, or C_{1-8}alkyl optionally substituted with hydroxy, C_{3-6}cycloalkyl, alkoxy, heterocycle, heteroaryl, C_{6-14}aryl, or amino, wherein said amino may be optionally substituted with $-C(O)C_{1-8}$alkyl or C_{1-8}alkyl;


to form a racemic compound of formula (I-26).

(48) A process for the preparation of a racemic compound of formula (I-27)




(I-27)

wherein R^e is halogen; and P^1 is C_{6-14}arylC_{1-8}alkyl;
comprising condensing a compound of the formula



wherein R^e is one or two halogen; R^{50} is C_{1-8}alkyl; and P^1 is C_{6-14}arylC_{1-8}alkyl;
with a racemic compound of the formula



to form a racemic compound of formula (I-27).

(49). A compound of formula (I-20a) described in above (36). formula (I-20b) described in above (37), formula (I-21a) described in above (38), formula (I-21b) described in above (39), formula (I-22a) described in above (40), formula (I-22b) described in above (41), formula (I-23a) described in above (42), formula (I-23b) described in above (43), formula (I-24a) described in above (44), formula (I-24b) described in above (45), formula (I-25) described in above (46), formula (I-26) described in above (47), or formula (I-27) described in above (48), or a pharmaceutically acceptable salt thereof.

(50) A compound of formula (I-20a) described in above (36). formula (I-20b) described in above (37), formula (I-21a) described in above (38), formula (I-21b) described in above (39), formula (I-22a) described in above (40), formula (I-22b) described in above (41), formula (I-23a) described in above (42), formula (I-23b)

described in above (43), formula (I-24a) described in above (44), formula (I-24b) described in above (45), formula (I-25) described in above (46), formula (I-26) described in above (47), or formula (I-27) described in above (48), or a pharmaceutically acceptable salt thereof, wherein each P^1 is hydrogen.

The present invention further provides a pharmaceutical composition containing any of the compounds shown above, a pharmaceutically acceptable salt or a solvate thereof, especially an anti-HIV agent.

[Effect of the Invention]

[0005]

The present invention compounds possess an integrase inhibitory activity and/or a cell-growth inhibitory activity against virus, especially HIV. Accordingly, they are useful for the prevention or treatment of various diseases mediated by integrase or virus infection diseases (e.g., AIDS). The present invention further provides a process for preparing a diastereomer, a mixture thereof, or racemate.

[Preferred Embodiment of the Invention]

[0006]

The terms used herein are explained below. Each term, alone or in combination with another tem, means as follows.

"Lower alkylene" means a straight or branched C1 to C6 alkylene such as methylene, ethylene, trimethylene, n-propylene, tetramethylene, ethylethylene, pentamethylene, or hexamethylene, preferably C1 to C4 straight alkylene such as methylene, ethylene, trimethylene, and tetramethylene, more preferably methylene or ethylene.

"Lower alkenylene" means a straight or branched C2 to C6 alkenylene, which consists of the above "Lower alkylene" having one or more double bonds, such as vinylene, propylene, or butenylene, preferably a straight C2 to C3 alkenylene such as vinylene or propylene.

"Lower alkyl" means a straight or branched C1 to C10 alkyl such as methyl, ethyl, n-propyl, i-propyl, t-butyl, isobutyl, sec-butyl, n-pentyl, and n-hexyl, and preferred is C1 to C8 alkyl, more preferred is methyl, ethyl or n-propyl, n-pentyl, isopentyl, neopentyl, tert-pentyl, n-hexyl, isohexyl, n-heptyl, n-octyl, n-nonyl, and n-decyl, preferably C1 to C6 lower alkyl, more preferably C1 to C4 lower alkyl such as methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, n-pentyl, isopentyl, neopentyl, tert-pentyl, n-hexyl, and isohexyl.

When lower alkyl is intervened with "-N=" or "=N-", the lower alkyl may have a double bond to form $-CH_2-N=CH_2$, $-CH=N-CH_3$ etc.

"Alkenyl" means a straight or branched C2 to C8 alkenyl, which consists of the above "alkyl" having one or more double bonds, such as vinyl, 1-propenyl, 2-propenyl, 1-butenyl, 2-butenyl, 3-butenyl, 1,3-butadienyl, and 3-methyl-2-butenyl, preferably C2 to C6 alkenyl, and more preferably C2 to C4 alkenyl.

"Lower alkenyloxy" means oxy attached to the above lower alkenyl, such as vinyloxy, 1-propenyloxy, 2-propenyloxy, 1-butenyloxy, 2-butenyloxy, 3-butenyloxy, 1,3-butadienyloxy, and 3-methyl-2-butenyloxy.

"Cycloalkyl" means C3 to C8 cyclic saturated hydrocarbon, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopentyl, and cyclooctyl, preferably C3 to C6 cycloalkyl.

"Cycloalkyl lower alkyl" means lower alkyl substituted with the above cycloalkyl, such as cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclopentylmethyl, cyclohexylmethyl, and cyclohexylethyl, and preferably C3 to C6 cycloalkyl lower alkyl.

"Aryl" means monocyclic aromatic hydrocarbon (e.g., phenyl) and polycyclic hydrocarbon (e.g., 1-naphthyl,2- naphthyl, 1-anthryl, 2-anthryl, 9-anthryl, 1-phenanthryl, 2-phenanthryl, 3-phenanthryl, 4-phenanthryl, 9-phenanthryl), preferably phenyl or naphthyl (e.g., 1-napthyl, 2-naphthyl).

"Aralkyl" or "aryl lower alkyl" means the above lower alkyl substituted with 1 to 3 of the above aryl, such as benzyl, diphenylmethyl, triphenylmethyl, phenethyl, 1-napthylmethyl, 2- napthylmethyl, preferably benzyl.

"Aryloxy" means oxy attached to the above aryl, such as 1-naphthyloxy, 2-naphthyloxy, 1-anthryloxy, 2-anthryloxy, 9-anthryloxy, 1-phenanthryloxy, 2-phenanthryloxy, 3-phenanthryloxy, 4-phenanthryloxy, and 9-phenanthryloxy, preferably phenyloxy or naphthyloxy (e.g., 1- napthyloxy, 2-naphthyloxy).

"Heterocyclic group" means "heteroring" or "heteroaryl".

"Heteroring" means a non-aromatic ring which has at least one of N, O and/or S in the ring and may be bonded at any substitutable position, preferably 5- to 7-membered ring, such as 1-pyrrolyl, 2-pyrrolyl, 3-pyrrolyl, 1-pyrrolidinyl, 2-pyrrolidinyl, 3-pyrrolidinyl, 1-imidazolyl, 2-imidazolyl, 4-imidazolyl, 1-imidazolidinyl, 2-imidazolidinyl, 4-imidazolidinyl, 1-pyrazolinyl, 3-pyrazolinyl, 4-pyrazolinyl, 1-pyrazolidinyl, 3-pyrazolidinyl, 4-pyrazolidinyl, piperidino, 2-piperidyl, 3-piperidyl, 4-piperidyl, 1-piperadinyl, 2-piperadinyl, 2-morpholinyl, 3-morpholinyl, morpholino, and tetrahydropyranyl. The non-aromatic ring is a

saturated or unsaturated ring.

"Heteroaryl" means monocyclic aromatic hetero-type ring or condensed aromatic hetero-type ring.

"Monocyclic aromatic hetero-type ring" means a 5- to 8- membered aromatic ring, which contains 1 to 4 of O, S, P and/ or N and may be bonded at any substitutable position.

"Condensed aromatic hetero-type ring" means a group wherein an aromatic ring containing 1 to 4 of O, S, P and/ or N is condensed with 1 to 4 of 5- to 8-membered aromatic ring(s) or the other 5- to 8-membered aromatic heteroring(s).

Examples of "heteroaryl" include furyl (e.g., 2-furyl, 3-furyl), thienyl (e.g., 2-thienyl, 3-thienyl), pyrrolyl (e.g., 1-pyrrolyl, 2-pyrrolyl, 3-pyrrolyl), imidazolyl (e.g., 1-imidazolyl, 2-imidazolyl, 4-imidazolyl), pyrazolyl (e.g., 1-pyrazolyl, 3-pyrazolyl, 4-pyrazolyl), triazolyl (e.g., 1,2,4-triazole-1-yl, 1,2,4-triazole-3-yl, 1,2,4-triazole-4-yl), tetrazolyl (e.g., 1-tetrazolyl, 2-tetrazolyl, 5-tetrazolyl), oxazolyl (e.g., 2-oxazolyl, 4-oxazolyl, 5-oxazolyl), isoxazolyl (e.g., 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl), thiazolyl (e.g., 2-thiazolyl, 4-thiazolyl, 5-thiazolyl), thiadiazolyl, isothiazolyl (e.g., 3-isothiazolyl, 4-isothiazolyl, 5-isothiazolyl), pyridil(e.g., 2-pyridil, 3-pyridil, 4-pyridil), pyridazinyl (e.g., 3-pyridazinyl, 4-pyridazinyl), pyrimidinyl (e.g., 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl), furazanyl (e.g., 3-furazanyl), pyrazinyl (e.g., 2-pyrazinyl), oxadiazolyl (e.g., 1,3,4-oxadiazole-2-yl), benzofuryl (e.g., 2-benzo[b]furyl, 3-benzo[b]furyl, 4-benzo[b]furyl, 5-benzo[b]furyl, 6-benzo[b]furyl, 7-benzo[b]furyl), benzothienyl(e.g., 2-benzo[b]thienyl, 3-benzo[b]thienyl, 4-benzo[b]thienyl, 5-benzo[b]thienyl, 6-benzo[b]thienyl, 7-benzo[b]thienyl), benzoimidazolyl (e.g., 1-benzoimidazolyl, 2-benzoimidazolyl, 4-benzoimidazolyl, 5-benzoimidazolyl), dibenzofuryl, benzooxazolyl, quinoxalinyl (e.g., 2-quinoxalinyl, 5-quinoxalinyl, 6-quinoxalinyl), cinnolinyl (e.g., 3-cinnolinyl, 4-cinnolinyl, 5-cinnolinyl, 6-cinnolinyl, 7-cinnolinyl, 8-cinnolinyl), quinazolinyl (e.g., 2-quinazolinyl, 4-quinazolinyl, 5-quinazolinyl, 6-quinazolinyl, 7-quinazolinyl, 8-quinazolinyl), quinolyl(e.g., 2-quinolyl, 3-quinolyl, 4-quinolyl, 5-quinolyl, 6-quinolyl, 7-quinolyl, 8-quinolyl), phthalazinyl (e.g., 1-phthalazinyl, 5-phthalazinyl, 6-phthalazinyl), isoquinolyl (e.g., 1-isoquinolyl, 3-isoquinolyl, 4-isoquinolyl, 5-isoquinolyl, 6-isoquinolyl, 7-isoquinolyl, 8-isoquinolyl), purinyl, pteridinyl (e.g., 2-pteridinyl, 4-pteridinyl, 6-pteridinyl, 7-pteridinyl), carbazolyl, phenanthridinyl, acridinyl (e.g., 1-acridinyl, 2-acridinyl, 3-acridinyl, 4-acridinyl, 9- acridinyl), indolyl (e.g., 1-indolyl, 2-indolyl, 3-indolyl, 4-indolyl, 5-indolyl, 6-indolyl, 7-indolyl), isoindolyl, phenazinyl (e.g., 1-phenazinyl,

2-phenazinyl) or phenothiadinyl (e.g., 1-phenothiadinyl, 2-phenothiadinyl, 3-phenothiadinyl, 4-phenothiadinyl).

"Heterocycle" means a cycle which can be lead to the above heterocyclic group.

"Heterocyclic group lower alkyl" or "Heterocycle lower alkyl" means lower alkyl substituted with the above heterocyclic group.

"Heterocyclic group oxy" or "Heterocycle oxy" means an oxy attached to the above heterocyclic group.

"Heterocyclic group carbonyl" or "Heterocyclecarbonyl" means a carbonyl attached to the above heterocyclic group

"Lower alkoxy" or "alkoxy" means an oxy attached to the above lower alkyl, such as methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, tert-butoxy.

"Lower alkylcarbonyl", "cycloalkylcarbonyl", "cycloalkyl lower alkylcarbonyl", "lower alkoxycarbonyl", "arylcarbonyl", "aryl lower alkylcarbonyl", "aryloxycarbonyl", "heterocyclecarbonyl", "heterocycle lower alkylcarbonyl", and "heterocycle oxycarbonyl", each means a carbonyl attached to the above "lower alkyl", "cycloalkyl", "cycloalkyl lower alkyl", "lower alkoxy", "aryl", "aryl lower alkyl", "aryloxy", "heterocycle", "heterocycle lower alkyl", and "heterocycleoxy", respectively.

[0007]

When a substituent(s) is/are present on "optionally substituted lower alkyl", "optionally substituted cycloalkyl", "optionally substituted cycloalkyl lower alkyl", "optionally substituted lower alkenyl", "optionally substituted lower alkoxy", "optionally substituted aryl", "optionally substituted aryl lower alkyl", "optionally substituted aryloxy", "optionally substituted aryloxy lower alkyl", "optionally substituted heterocyle, "optionally substituted heterocyclic group", "optionally substituted heterocycle lower alkyl", "optionally substituted heterocycleoxy", "optionally substituted lower alkenyloxy", "optionally substituted lower alkylcarbonyl", "optionally substituted cycloalkylcarbonyl", "optionally substituted cycloalkyl lower alkylcarbonyl", "optionally substituted lower alkoxycarbonyl", "optionally substituted arylcarbonyl", "optionally substituted aryl lower alkylcarbonyl", "optionally substituted aryloxycarbonyl", "optionally substituted heterocyclecarbonyl", "optionally substituted heterocycle lower alkylcarbonyl", "optionally substituted heterocycleoxycarbonyl", "optionally substituted lower alkylene", "optionally substituted lower alkenylene", "optionally substituted phosphoric acid residue", "optionally substituted carbocycle" or "optionally

substituted heterocycle", each may be substituted with the same or different, 1 to group(s) selected from Substituent group B at any position.


Examples of Substituent group B include hydroxy, carboxy, halogen (F,Cl,Br,I halo lower alkyl (e.g., CF_3,CH_2CF_3, CH_2CCl_3), halo lower alkoxy (e.g., OCF_3, OCH_2CF OCH_2CCl_3), lower alkyl (e.g., methyl, ehtyl, isopropyl, tert-butyl), lower alkenyl (e.g vinyl), lower alkynyl (e.g., ethynyl), cycloalkyl (e.g., cyclopropyl), cycloalkenyl (e.g cyclopropenyl), lower alkoxy (e.g., methoxy, ethoxy, propoxy, butoxy), lower alkenylox (e.g., vinyloxy, allyloxy), lower alkoxycarbonyl (e.g., methoxycarbonyl, ethoxycarbonyl tert-butoxycarbonyl), nitro, nitroso, optionally substituted amino (e.g., alkylamin (e.g., methylamino, ethylamino, dimethylamino), acylamino (e.g., acetylamino benzoylamino), aralkylamino (e.g., benzylamino, tritylamino), hydroxyamino), azido aryl (e.g., phenyl), aralkyl (e.g., benzyl), cyano, isocyano, isocyanate, thiocyanate isothiocyanate, mercapt, alkylthio (e.g., methylthio), alkylsulfonyl (e.g. methansulfonyl, ethansulfonyl), optionally substituted alkylsulfonylamino (e.g. methanesulfonylamino, ethansulfonylamino, N-methylsulfonyl-N'-methylamino), optionally substituted carbamoyl (e.g., alkylcarbamoyl (e.g., methylcarbamoyl, ethylcarbamoyl, dimethylcarbamoyl)), sulfamoyl, acyl (e.g., formyl, acetyl), formyloxy, haloformyl, oxal, thioformyl, thiocarboxy, dithiocarboxy, thiocarbamoyl, sulfino, sulfo, sulfoamino, hydrazino, azido, ureido, amizino, quanidino, phthalimide, oxo, phosphoric acid residue, lower alkyl which is substituted with a phosphoric acid residue and may be intervened with a heteroatom group(s), aryl substituted with a phosphoric acid residue, aralkyl substituted with a phosphoric acid residue, hydroxyl lower alkyl, preferably hydroxy, carboxy, halogen(F,Cl,Br,I), halo lower alkyl (e.g., CF_3,CH_2CF_3, CH_2CCl_3), halo lower alkoxy (e.g., OCF_3, OCH_2CF_3, OCH_2CCl_3), lower alkyl (e.g., methyl, ethyl, isopropyl, tert-butyl), lower alkoxy (e.g., methoxy, ethoxy, propoxy, butoxy), optionally substituted amino (e.g., alkylamino (e.g., methylamino, ethylamino, dimethylamino), oxo, or phosphoric acid residue.

Examples of a substituent of "optionally substituted amino" or "optionally substituted carbamoyl" include mono- or di- lower alkyl, lower alkylcarbonyl, lower alkylsulfonyl, optionally substituted lower alkyl (e.g., methyl, ethyl, isopropyl, benzyl, carbamoylalkyl (e.g., carbamoylmethyl), mono- or di- lower alkylcarbamoyl lower alkyl (e.g., dimethylcarbamoylethyl), hydroxyl lower alkyl, heterocycle lower alkyl (e.g., morpholinoethyl, tetrahydropyranylethyl), alkoxycarbonyl lower alkyl (e.g., ethoxycarbonylmethyl, ethoxycarbonylethyl), mono- or di- lower alkylamino lower alkyl (e.g., dimethylaminoethyl)), lower alkoxy lower alkyl (e.g., methoxyethyl,

ethoxymethyl, ethoxyethyl, isopropoxyethyl), acyl (e.g., formyl, optionally substituted lower alkylcarbonyl (e.g., acetyl, propionyl, butylyl, isobutylyl, valeryl, isovaleryl, pivaroyl, hexanoyl, octanoyl, methoxyethylcarbonyl, 2,2,2-trifluoroethylcarbonyl, ethoxycarbonylmethylcarbonyl), lower alkoxy lower alkylcarbonyl (e.g., methoxyethylcarbonyl), lower alkylcarbamoyl lower alkylcarbonyl (e.g., methylcarbamoylehtylcarbonyl), alkoxycarbonylacetyl), optionally substituted arylcarbonyl (e.g., benzoyl, toloyl), optionally substituted aralkyl (e.g., benzyl, 4-fluorobenzyl), hydroxy, optionally substituted lower alkylsulfonyl (e.g., methanesulfonyl, ethanesulfonyl, isopropylsulfonyl, 2,2,2-trifluoroethanesulfonyl, benzylsulfonyl, methoxyethylsulfonyl), lower alkyl, or arylsulfonyl optionally substituted with halogen (e.g., benzenesulfonyl, toluenesulfonyl, 4-fluorobenzenesulfonyl, fluorobenzenesulfonyl), cycloalkyl (e.g., cyclopropyl), aryl optionally substituted with lower alkyl (e.g., phenyl), lower alkylaminosulfonyl (e.g., methylaminosulfonyl, dimethylaminosulfonyl), lower alkylaminocarbonyl (e.g., dimethylaminocarbonyl), lower alkoxycarbonyl (e.g., ethoxycarbonyl), cycloalkylcarbonyl (e.g., cyclopropylcarbonyl, cyclohexylcarbonyl), optionally substituted sulfamoyl (e.g., sulfamoyl, methylsulfamoyl, dimethylsulfamoyl), lower alkylcarbonylamino (e.g., methylcarbonylamino), heterocycle (e.g., morpholino, tetrahydropyranyl), optionally substituted amino (e.g., mono- or di-alkylamino (e.g., dimethylamino), formylamino).

As to amino of "optionally substituted amino", "optionally substituted carbamoyl", or "optionally substituted carbamoylcarbonyl", two substituents on the amino together with the neighboring N atom may form an N-containing heterocycle which optionally contains S and/or O in the ring (preferably 5- to 7- membered ring or saturated ring) and is optionally substituted with oxo or hydroxy. The optional S atom in the ring may be substituted with oxo. The N-containing heterocycle is preferably a 5- or 6-membered ring such as piperadinyl, piperidino, morpholino, pyrrolidino, 2-oxopiperidino, 2-oxopyrrolidino, 4-hydroxymorpholino.

"Phosphoric acid residue" means a group shown of the formula: $-PO(OH)_2$. "Optionally substituted phosphoric acid residue" means a phosphoric acid residue wherein the OH part and/or a hydrogen of the OH is optionally substituted with a phosphoric acid residue, preferably shown by the formula:




(P-1)

(wherein, R^A and R^B each is independently OR^C or NR^DR^E (wherein R^C, R^D and R^E are each independently hydrogen, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted aryl, optionally substituted heterocyclic group, or R^D and R^E taken together with the neighboring N atom may form an optionally substituted heterocycle (preferably 5- to 6- membered ring)) or R^A and R^B taken together with the neighboring P atom may form an optionally substituted heterocycle (preferably 5- to 6- membered ring)).

Preferably, R^A and R^B are bothOR^C, or one of them is OR^C and the other is NR^DR^E.


R^C, R^D and R^E each is preferably, independently, lower alkyl (e.g., methyl, ethyl).

The optionally substituted heterocycle formed by R^A and R^B taken together with the neighboring P atom may be the following structure:




(wherein, the broken line means a part of the ring)

Hydroxy substituted with optionally substituted phosphoric acid residue is preferably hydroxy substituted with a phosphoric acid residue substituted with di lower alkyls, and more preferably a group of the formula:



Amino substituted with optionally substituted phosphoric acid residue is preferably amino substituted with a phosphoric acid residue substituted with di lower

alkyls, and more preferably a group of the formula:



[0008]

(More preferable embodiments)

R^1 is hydrogen or lower alkyl, preferably hydrogen.

X is a single bond, a heteroatom group selected from O, S, SO, SO_2 and NH (hereafter also referred to as "M"), or lower alkylene or lower alkenylene each may be intervened by the heteroatom. The term of "intervened by" means the following cases:

1) The heteroatom group is present between carbon atoms which constitutes the alkylene or alkenylene.

2) The heteroatom group is attached to the N atom of the carbamoyl group neighboring to X.

3) The heteroatom group is attached to R^2 neighboring to X.


The heteroatom group (M) may be the same or different, and one or more atoms. Examples of that lower alkylene is intervened by a heteroatom group include $-M-CH_2-$, $-CH_2-M-CH_2-$, $-CH_2-M-$, and $-CH_2-M-M-CH_2-$.

X is preferably a spacer consisting 1 to 3 joined atoms. X is more preferably lower alkylene or lower alkenylene each may be intervened by a heteroatom group, or O. X is most preferably C1 to C3 alkylene, C2 to C3 alkenylene, or O. Especially preferred is methylene or O.

R^2 is optionally substituted aryl, preferably phenyl. A substituent on the aryl is the same or different, 1 to 3, preferably 1 to 2 substituent(s), including preferably halogen, hydroxy, amino, lower alkylamino, cyano, carboxy, formyl, oxo, lower alkyl, lower alkoxy, lower alkylthio, carbamoyl, and lower alkylcarbamoyl, and Substituent group S1(: optionally substituted phosphoric acid residue, aryl substituted with optionally substituted phosphoric acid residue, aralkyl substituted with optionally substitutedphosphoric acid residue, hydroxyl substituted with optionally substitutedphosphoric acid residue, amino substituted with optionally

substitutedphosphoric acid residue, lower alkyl substituted with optionally substitutedphosphoric acid residue (said lower alkyl may be intervened with a hetero atom group(s) selected from O, S, SO, SO_2, NR^5 (R^5 is independently selected from the same substituent group for R^4), -N= and =N-), lower alkoxy lower alkyl, amino lower alkyl optionally substituted with mono- or di-lower alkyl, halogenated lower alkyl, lower alkoxy, carbamoyl optionally substituted with mono- or di-lower alkyl, optionally substituted lower alkylsulfonylamino, halogenated lower alkoxy, hydroxyl lower alkyl), more preferably halogen,hydroxy,amino,cyano,lower alkyl, lower alkoxy or Substituent group S1, and most preferred is halogen (e.g., F) and/or a group selected from Substituent group S1. A substituent on the aryl is preferably at the 4-position. R^2 is more preferably phenyl or phenyl substituted with at least halogen, and most preferably 4-halogenophenyl (e.g., 4-F-phenyl). In another embodiment, R^2 is preferably phenyl optionally substituted with 1 to 3 R(s) mentioned below.

In all compounds of the present invention, the structure of "-X-R^2" is preferably shown by the formula below:




R each is independently a group selected from halogen and Substituent group S1.

Substituent group S1: optionally substituted phosphoric acid residue, aryl substituted with optionally substituted phosphoric acid residue, aralkyl substituted with optionally substituted phosphoric acid residue, hydroxyl substituted with optionally substituted phosphoric acid residue, amino substituted with optionally substituted phosphoric acid residue, lower alkyl substituted with optionally substituted phosphoric acid residue (said lower alkyl may be intervened by a heteroatom group(s) selected from CO, O, S, SO, SO_2, NR^a (R^a is hydrogen or lower alkyl), -N= and =N-), lower alkoxy lower alkyl, optionally substituted amino lower alkyl (the substituent: mono- or di- lower alkyl, lower alkylcarbonyl, or lower alkylsulfonyl), halogenated lower alkyl, lower alkoxy, optionally substituted carbamoyl (the substituent: mono- or di- lower alkyl, lower alkylcarbonyl, or lower alkylsulfonyl), optionally substituted lower alkylsulfonylamino, halogenated lower alkoxy, and hydroxyl lower alkyl.

m is an integer of 0 to 3, preferably 0 or 1 to 2. when m is 1, R is preferably halogen. When m is 2, R is more preferably the same or different group selected from halogen, lower alkyl, lower alkoxy, lower alkoxylower alkyl, halogenated lower alkyl, halogenated lower alkoxy, lower alkylsulfonylamino, carbamoyl, and lower alkylcarbamoyl. More preferably, R is two halogens, or halogen and another group. R preferably locates at the 4-position and optional another position of the benzene ring.

R^3 can be a various substituent which does not bring a negative effect to the pharmacological activity, including hydrogen, halogen, hydroxy, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted lower alkenyloxy, optionally substituted aryl, optionally substituted aryloxy, optionally substituted heterocyclic group, optionally substituted heterocycleoxy, and optionally substituted amino. Examples of substituent of "optionally substituted" include halogen, hydroxy, amino, lower alkylamino, cyano, carboxy, formyl, oxo, lower alkyl, lower alkoxy, lower alkylthio, carbamoyl, lower alkylcarbamoyl, aryl, heterocyclic group, lower alkylcarbonyl, lower alkylcarbonyloxy, lower alkoxycarbonyl, halogenated lower alkyl, halogenated lower alkoxy, and preferably halogen, hydroxy, amino, lower alkylamino, lower alkyl, and lower alkoxy. R^3 is more preferably hydrogen, halogen, hydroxy, lower alkyl, lower alkenyl, lower alkoxy, lower alkenyloxy or optionally substituted amino, and most preferably hydrogen or lower alkyl (e.g., methyl), esp. hydrogen.


Z^2 shows C, CH, optionally substituted lower alkylene, lower alkenylene etc., and Z^2 and R^4 of Z^1 taken together form a ring, whereby compound (I) shows a tricyclic compound (I-1) or (I-11) shown below, or its derivative, tetracyclic compound.



(I-1)

A ring is optionally substituted heterocycle containing at least an N atom. The

heterocycle is a 5- to 7-membered ring which contains preferably 1 to 3, more preferably 2 to 3 atoms of O, S and/or N. The heterocycle is preferably selected from the above heterocycle. The arc optionally contains 1 to 2 heteroatom(s) at any possible position. One of preferable embodiments of A ring is an optionally substituted ring shown below.



(Z is CH_2, O, S, SO, SO_2 or NR^{19})

A ring is preferably a ring of (a), (b), or (c).


Z is preferably O or NR^{19}.

When Z is NR^{19}, examples of R^{19} include 1)hydrogen, 2)optionally substituted lower alkyl (the substituent is e.g., amino optionally substituted with mono- or di-lower alkyl; cycloalkyl; hydroxy; optionally substituted heterocyclic group (preferably 5- to 7-membered ring, e.g., furyl, thienyl, thiazolyl, pyridil, morpholino, imidazole; examples of the substituent include lower alkyl, halogen); optionally substituted heterocyclecarbonyl (the heterocycle is preferably 5- to 7-membered ring, e.g., morpholinocarbonyl); optionally substituted phenyl (the substituent is e.g., lower alkyl, amino, lower alkylamino, hydroxy, halogen, halogenated lower alkyl, lower alkoxy, halogenated lower alkoxy, lower alkylthio, lower alkylsulfonyl), acetylamino, carbamoyl, carbamoyl substituted with mono- or di- lower alkyl, lower

alkylsulfonylamino, lower alkoxy, carbonyl, halogen, thiol, lower alkylthio), 3) lower alkenyl, 4) acyl (e.g., lower alkylcarbonyl), 5) lower alkylsulfonyl. R^{19} may be selected from Substituent group S2 shown below.


The other substituent on A ring may be selected from R^{15} to R^{18} or Substituent group S2, preferably lower alkyl. Substituents on A ring may form a condensed ring or a spiro ring as mentioned below, whereby compound (I) includes a tetracyclic compound.

A ring is more preferably any of the following rings:




Z = O or NR^{26}

(A-1)



Z = O or NR^{31}

(A-2)



Z = O or NR^{40}

(A-3)

(wherein, R^{20} to R^{40} are each independently a group selected from Substituent group S2, or any two groups of R^{20} to R^{40}, which bonds to the same carbon atom, taken together with the carbon atom, may form a spiro ring, i.e., an optionally substituted carbocyle or optionally substituted heterocycle, or each combination of (R^{20} and R^{22}), (R^{23} and R^{24}), (R^{25} and R^{26}), (R^{27} and R^{29}), (R^{30} and R^{31}), (R^{32} and R^{34}), (R^{35} and R^{36}), (R^{37} and R^{38}), and (R^{39}andR^{40}), taken together with the neighboring atom, may form an optionally substituted carbocyle or optionally substituted heterocycle.

Substitution group S2: hydrogen, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted cycloalkyl lower alkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted lower alkenyloxy, optionally substituted aryl, optionally substituted aryl lower alkyl, optionally substituted aryloxy, optionally substituted heterocycle, optionally substituted heterocycle lower alkyl, optionally substituted heterocycleoxy, hydroxy, optionally substituted amino, optionally substituted lower alkylcarbonyl, optionally substituted cycloalkylcarbonyl, optionally substituted cycloalkyl lower alkylcarbonyl,

optionally substituted lower alkoxycarbonyl, optionally substituted arylcarbonyl, optionally substituted aryl lower alkylcarbonyl, optionally substituted aryl oxycarbonyl, optionally substituted heterocyclecarbonyl, optionally substituted heterocycle lower alkylcarbonyl, optionally substituted heterocycleoxycarbonyl, optionally substituted aminocarbonyl, optionally substituted phosphoric acid residue, aryl substituted with optionally substituted phosphoric acid residue, aralkyl substituted with optionally substituted phosphoric acid residue, hydroxy substituted with optionally substituted phosphoric acid residue, amino substituted with optionally substituted phosphoric acid residue, or lower alkyl substituted with optionally substituted phosphoric acid residue (the lower alkyl may be intervened with a heteroatom group(s) selected from CO, O, S, SO, SO_2, NR^5 (R^5 is independently selected from the same substitution group as R^4), -N= and =N-)

The stereochemistry of an asymmetric carbon represented by * shows the R- or S-configuration, or a mixture thereof)

In one embodiment, R^{20} to R^{40} each is preferably hydrogen, optionally substituted lower alkyl (examples of the substituent: OH, lower alkoxy, cycloalkyl, lower alkylthio, lower alkylsulfonyl, heterocyclic group, aryl, optionally substituited amino (examples of the substituent: lower alkyl, acyl)), cycloalkyl, optionally substituited aryl (examples of the substituent: OH, lower alkyl), and optionally substituited heterocyclic group.

In one embodiment, R^{20} to R^{25}, R^{27} to R^{30}, and R^{32} to R^{39}, each is preferably hydrogen, C1-C8 alkyl, C6-C14 aryl C1-C8 alkyl, C6-C14 aryl, or alkoxy.

In one embodiment, R^{26}, R^{31}, and R^{40}, each is preferably hydrogen, C3-6 cycloalkyl, heterocycle, or C1-8 alkyl optionally substituted with hydroxy, C3-6 cycloalkyl, alkoxy, heterocycle, heteroaryl, C6-14 aryl, or amino, wherein said amino may be optionally substituted with –C(O)C1-8 alkyl or C1-8 alkyl.

More Preferred embodiments are shown below for example

I) When A ring is A-1, preferred is that 1) Z is NR^{26} and R^{26} and R^{24} taken together form heterocycle, and the others are hydrogens; 2) Z is O or NR^{26}, (R^{20} and R^{22}) or

(R^{23} and R^{24}) taken together form cycloalkyl which is substituted with phenyl, the others are hydrogens or optionally substituted lower alkyl.

II) When A ring is A-2, preferred is that 1) Z is O, R^{27} or R^{28} is lower alkyl, and the others are hydrogens; 2) Z is NR^{31} and R^{30} and R^{31} taken together form heterocycle and the others are hydrogens, or R^{27} and R^{29} taken together form cycloalkyl and the others are hydrogens ; 3) Z is O, R^{27} and R^{29} taken together form cycloalkyl which may be condensed with phenyl, and the others are hydrogens

R^{14} and R^x are each independently hydrogen, optionally substitutedlower alkyl, optionally substituted cycloalkyl, optionally substituted cycloalkyl lower alkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted lower alkenyloxy, optionally substituted aryl, optionally substituted aryl lower alkyl, optionally substituted aryloxy, optionally substituted heterocyclic group, optionally substituted heterocycle lower alkyl, optionally substituted heterocycleoxy, hydroxy, optionally substituted amino,optionally substituted lower alkylcarbonyl, optionally substituted cycloalkylcarbonyl, optionally substituted cycloalkyl lower alkylcarbonyl, optionally substituted lower alkoxycarbonyl, optionally substituted arylcarbonyl, optionally substituted aryl lower alkylcarbonyl, optionally substituted aryloxycarbonyl, optionally substituted heterocyclecarbonyl, optionally substituted heterocycle lower alkylcarbonyl, optionally substituted heterocycleoxycarbonyl, optionally substituted aminocarbonyl, optionally substituted phosphoric acid residue, aryl substituted with optionally substituted phosphoric acid residue, aralkyl substituted with optionally substituted phosphoric acid residue, hydroxy optionally substituted with optionally substituted phosphoric acid residue, amino substituted with optionally substituted phosphoric acid residue, or lower alkyl substituted with optionally substituted phosphoric acid residue (the lower alkyl may be intervened with a heterotom group(s) selected from O, S, SO, SO_2, NR^a (R^a is hydrogen or lower alkyl), -N= and =N-).


R^{14} and R^x are each independently, preferably, hydrogen, hydroxyl, optionally substituted lower alkyl (the substituent is preferably, e.g., amino, lower alkyl amino, hydroxy, lower alkoxy). R^{14} and R^x are preferably hydrogens.

A broken line in the compound (I-1) represents the presence or absence of a bond,


provided that when the broken line represents the presence of a bond, R^X is not present.

[0009]

The compound (I) includes the following compounds.




(I-1)




(I-1-1)

(wherein each symbol is as defined above)



(I-8)

F ring means the same heterocycle as A ring, preferably 5- to 7-membered ring, and the substituents on F ring are the same as those for A ring. The other symbols are as defined above.



(I-3)

(wherein each symbol is as defined above; Z is O or NR^{19}; R^{15} to R^{19} are each independently hydrogen or a group selected from the above Substituent group S2, or each combination of (R^{15} and R^{16}), (R^{17} and R^{18}), (R^{16} and R^{18}), and (R^{18} and R^{19}) taken together with the neighboring atom(s), may form an optionally substituted carbocycle (preferably 5- to 6-membered ring) or an optionally substituted heterocyle (preferably 5- to 6-membered ring); or each combination of (R^{15} and R^{16}) and (R^{17} and R^{18}) taken together may fom oxo)

Compound (I-3) is preferably as follows.


(1) R^1 is hydrogen; R^3 is hydrogen; m is 1 or 2; R^{14} is hydrogen.

(2) m is 1 or 2, R is each independently halogen, halogenated lower alkyl, lower alkoxy, halogenated lower alkoxy, lower alkoxy lower alkyl, hydroxy lower alkyl, optionally substituted amino lower alkyl (the substituent is mono- or di- lower alkyl, lower alkylcarbonyl, or lower alkylsulfonyl), optionally substituted carbamoyl (the substituent is mono- or di- lower alkyl, lower alkylcarbonyl, or lower alkylsulfonyl), phosphoric acid residue, aryl substituted with optionally substituted phosphoric acid residue, aralkyl substituted with optionally substituted phosphoric acid residue or sulfonylamino optionally substituted with lower alkyl; R^1 is hydrogen; R^3 is hydrogen; R^{14} is hydrogen, hydroxyl or lower alkyl optionally substituted with mono- or di- lower alkylamino; Z is O or NR^{19} (R^{19} is hydrogen or lower alkyl, lower alkoxy lower alkyl, optionally substituted phosphoric acid residue, aryl substituted with optionally substituted phosphoric acid residue, aralkyl substituted with optionally substituted phosphoric acid residue, hydroxy substituted with optionally substituted phosphoric acid residue, amino substituted with optionally substituted phosphoric acid residue, or lower alkyl substituted with optionally substituted phosphoric acid residue).

(3)R is each independently, -F, $-CF_3$, -OMe, $-OCF_3$, $-CH_2OMe$, $-CH_2OH$, $-CH_2N(Me)_2$, -CONHMe, $-CON(Me)_2$, $-CH_2PO(OEt)_2$, $-PO(OEt)_2$, $-NHSO_2Me$, or $-NMeSO_2Me$;R^1 is hydrogen; R^3 is hydrogen; m is 1 or 2; R^{14} is hydrogen, hydroxyl or $-CH_2N(Me)_2$; Z is O or NR^{19} (R^{19} is hydrogen or $-CH(Me)_2$, $-(CH_2)_2OMe$,

-$(CH_2)_2PO(OEt)_2$).

(4)R^{15} and R^{16} are hydrogens; R^{17} and R^{18} are hydrogens or taken together with the neighboring atom form a 3- to 7-membered carbocyle; and/or Z is O or NH. This case preferably also satisfys the above (2) or (3).



(I-11)

D ring means the same heterocycle as A ring, preferably 5- to 7-membered ring, and the substituents on D ring are the same as those for A ring. The other symbols are as defined above.

The structure of compound (I) has at least the following characteristics.

(1) The main structure, condensed heterocycle, is substituted with oxo (=O), hydroxyl (OH) and oxo.

(2) A substituted carbamoyl group (-$CONR^1XR^2$) is attached to the position neighboring to the oxo group on the condensed hereocycle.

The above structure contributes to a remarkably potent integrase inhibitory activity and/or cell-growth inhibitory activity against virus including HIV. In contrast, the structures of the other parts such as Z^1, Z^2, and R^3 each may be of variety, being optionally substituted or optionally condensed, and its condensed ring is also optionally substituted.

The present invention provides a pharmaceutically acceptable salt or a solvate of compound (I). All theoretically possible tautomer, geometrical isomer, optically active compound, and racemate thereof are within the scope of the invention.

Pharmaceutically acceptable salts of a compound of the present invention include, as basic salts, for example, alkali metal salts such as sodium or potassium salts; alkaline-earth metal salts such as calcium or magnesium salts; ammonium salts; aliphatic amine salts such as trimethylamine, triethylamine, dicyclohexylamine, ethanolamine, diethanolamine, triethanolamine or procaine salts; aralkyl amine salts

such as N, N-dibenzylethylenediamine salts; heterocyclic aromatic amine salts such as pyridin salts, picoline salts, quinoline salts or isoquinoline salts; quaternary ammonium salts such as tetramethylammonium salts, tetraethylammonium salts, benzyltrimethylammonium salts, benzyltriethylammonium salts, benzyltributylammonium salts, methyltrioctylammonium salts or tetrabutylammonium salts; and basic amino acid salts such as arginine salts or lysine salts. Acid salts include, for example, mineral acid salts such as hydrochloride, sulfates salts, nitrate salts, phosphates salts, carbonates salts, hydrogencarbonates or perchlorate; organic acid salts such as acetates, propionates, lactates, maleates, fumarates, tararic acid salts, malates, citrates salts, ascorbates, formic acid; sulfonates such as methanesulfonates, isethionates, benzenesulfonates, or p-toluenesulfonates; and acidic amino acid salts such as aspartates or glutamates.


Solvates of a compound of the present invention include alcholates and hydrates.

[0012]

A general process for producing the present compound will be exemplified below.

(Method of preparing raw material)

[Chemical formula 41]



(wherein L^1 is a leaving group (e.g.; halogen); P^1 and P^2 are a hydroxy protecting group; P^3 is a carboxy protecting group (e.g.: lower alkyl); R^a and R^b are hydrogen or a substituent on an amino group)

Examples of a hydroxy protecting group (P^1, P^2) include acyl (e.g.: acetyl, pivaloyl, benzoyl), aralkyl (e.g.: benzyl), lower alkyl (e.g.: methyl), alkoxyalkyl (e.g.: methoxymethyl, methoxyethyl), lower alkylsulfonyl (e.g.: methanesulfonyl), arylsulfonyl (e.g.: benzenesulfonyl, toluenesulfonyl), alkoxycarbonyl (e.g.:

methoxycarbonyl) and the like.

As a carboxy protecting group (P^3), lower alkyl (e.g.: methyl, ethyl), and aralkyl (e.g.: benzyl) are exemplified.

[0013]

(First step)

The present step is a reaction for condensing a compound (II) and a compound (III) to synthesize a compound (IV). The reaction may be performed according to the condition for a reaction of amidating carboxylic acid which is generally performed. A compound (II) may be reacted as it is, or may be reacted after converted into corresponding acid chloride or active ester. Preferably, the reaction is performed in a suitable solvent in the presence of a condensing agent.

As a condensing agent, dicyclohexylcarbodiimide, 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride and the like may be used. If necessary, a reagent such as 1-hydroxybenzotriazole and N-hydroxysuccinimide, or a base such as triethylamine, N-methylmorpholine, and pyridine may be added.

A reaction temperature is 0 to 150°C, preferably room temperature to 70°C.

As a reaction solvent, a non-protonic solvent can be broadly used, and tetrahydrofuran (THF), 1,4-dioxane, dimethylformamide (DMF), methylene chloride, chloroform and the like are preferable.

A reaction time is a few minutes to a few tens hours, preferably 9 to 17 hours.

(Second step)

The present step is a reaction for introducing a protected hydroxy group (OP^1) into a compound (IV) to produce a compound (V). The reaction may be performed according to the condition for an alkoxylating reaction which is generally performed.

For example, a compound (V) in which P^1 is methyl can be synthesized by reacting a compound (IV) with metal alkoxide (e.g.: sodium methoxide).

A reaction temperature is 0 to 200°C, preferably 80 to 120°C.

As a reaction solvent, alcohol, dimethylformamide (DMF), and dimethyl sulfoxide (DMSO) are exemplified.

A reaction time is a few minutes to a few tens hours, preferably 5 to 10 hours.

(Third step)

The present step is a reaction for protecting a hydroxy group of a compound

(V) to produce a compound (VI). The reaction may be performed according to the condition for a reaction of protecting a hydroxy group which is generally performed. For example, by using diisopropyl azodicarboxylate or diethyl azodicarboxylate together with an alcohol and various phosphines, a compound (VI) in which P^2 is alkyl can be synthesized.

A reaction temperature is 0 to 100°C, preferably 0°C to room temperature.

As a reaction solvent, THF, toluene, dichloromethane and the like are exemplified.

A reaction time is a few minutes to a few tens hours, preferably 1 to 3 hours.

(Fourth step)

The present step is a reaction of oxidizing a nitrogen atom of a compoung (VI) to produce a compound (VII). The reaction may be performed according to the condition for an oxidation reaction using an oxidizing agent which is generally performed.

A reaction temperature is 0 to 100°C, preferably under ice-cooling to room temperature.

As a reaction solvent, chloroform, methylene chloride, acetic acid and the like are exemplified.

Examples of an oxidizing agent include metachloroperbenzoic acid, hydrogen peroxide and the like.

A reaction time is a few minutes to a few tens hours, preferably 1 to 5 hours.

(Fifth step)

The present step is a reaction for hydroxylating a methyl group of a compound (VII). Preferably, after acetoxylation by a reaction with acetic anhydride (reaction temperature: 0 to 150°C, preferably 120 to 140°C), this may be hydrolyzed (e.g.: treatment with a base (e.g.: alkali metal hydroxide)).

A reaction time is a few minutes to a few tens hours, preferably 0.5 to 2 hours for acetoxylation, and 0.5 to 1 hour for hydrolysis.

(Sixth step)

The present step is a reaction for oxidizing a hydroxy group of a compound (VIII) to synthesize a compound (IX).

A reaction temperature is 0 to 150°C, preferably room temperature to 70°C.

As a reaction solvent, chloroform and the like are exemplified.

As an oxidizing agent, dimethyl sulfoxide and the like are exemplified.

A reaction time is a few minutes to a few tens hours, preferably 0.1 to 1 hour.

(Seventh step)

The present step is a reaction for oxidizing a formyl group of a compound (IX) to synthesize a compound (X).

A reaction temperature is 0 to 150°C, preferably under ice-cooling to room temperature.

As a reaction solvent, an alcohol and the like are exemplified.

As an oxidizing agent, potassium hydroxide and iodine are exemplified.

A reaction time is a few minutes to a few tens hours, preferably 0.5 to 3 hours.

(Eighth step)

The present step is a reaction for deprotecting an OP^2 part of a compound (X) to synthesize a compound (XI). The reaction may be performed according to the condition for a reaction of deprotecting a hydroxy protecting group which is generally performed.

A reaction temperature is 0 to 150°C, preferably under ice-cooling to room temperature.

As a reaction solvent, acetonitrile, methylene chloride, THF and the like are exemplified.

A reaction time is a few minutes to a few tens hours, preferably 1 to 3 hours.

(Ninth step)

The present step is a reaction for deprotecting an OP^1 part of a compound (XI) to synthesize a compound (I-A). The reaction may be treated preferably with a Lewis acid (e.g.: aluminum chloride).

A reaction temperature is 0 to 150°C, preferably 10 to 50°C.

As a reaction solvent, methylene chloride, THF and the like are exemplified.

A reaction time is a few minutes to a few tens hours, preferably 1 to 3 hours.

(Tenth step)

The present step is a reaction for deprotecting an ester part ($COOP^3$) of a

compound (X) to synthesize carboxylic acid (XII). Preferably, hydrolysis with an alkali (e.g.: NaOH) may be performed.

A reaction temperature is 0 to 150°C, preferably 10 to 50°C.

As a reaction solvent, methanol, water and the like are exemplified.

A reaction time is a few minutes to a few tens hours, preferably a few minutes to 2 hours.

Carboxylic acid (XII) can be converted into various derivatives (e.g.; amide).

(Eleventh step)

The present step is a reaction for reacting a compound (XII) with various amines to synthesize a compound (XIII). The reaction may be performed according to the condition for a reaction of amidating carboxylic acid which is generally performed and, for example, the reaction may be performed as in the first step.

A reaction temperature is 0 to 150°C, preferably room temperature to 70°C.

As a reaction solvent, a non-protonic solvent can be broadly used, and tetrahydrofuran (THF), 1,4-dioxane, dimethylformamide (DMF), methylene chloride, chloroform and the like are preferable.

A reaction time is a few minutes to a few tens hours, preferably a few minutes to 3 hours.

An amide part of the resulting compound (XIII) may be further chemically modified (e.g.: N-alkylation).

(Twelfth step)

The present step is a reaction for deprotecting OP^1 and OP^2 parts of a compound (XIII) to synthesize a compound (I-B). The reaction may be performed according to the condition for a reaction of deprotecting a hydroxy protecting group which is generally performed.

For example, when pyridine hydrochloride is used, a reaction temperature is 0 to 200°C, preferably 150 to 180 degree.

A reaction time is a few minutes to a few tens hours, preferably 1 to 5 minutes.

(Thirteenth step)

The present step is a reaction for deprotecting an ester part ($COOP^3$) of a compound (XI) to synthesize carboxylic acid (XIV). Preferably, hydrolysis with an

alkali (e.g.: lithium hydroxide) may be performed.

A reaction temperature is 0 to 150°C, preferably 10 to 50°C.

As a reaction solvent, methanol, water and the like are exemplified.

A reaction time is a few minutes to a few tens hours, preferably a few minutes to 3 hours.

(Fourteenth step)

The present step is a reaction for deprotecting an OP^1 part of a compound (XIV) to synthesize a compound (I-C). The reaction may be treated preferably with a Lewis acid (e.g.: boron tribromide).

A reaction temperature is 0 to 150°C, preferably under ice-cooling to room temperature.

As a reaction solvent, dichloromethane and the like are exemplified.


A reaction time is a few minutes to a few tens hours, preferably a few minutes to 5 hours.

[0014]

The monocyclic carbamoylpyridone derivative obtained above is derived into a bicyclic compound by the following method.

(Process 1)

[Chemical formula 42]



(wherein R^1, X, R^2, P^1, P^3 and R^4 are as define above, and L^2 is a leaving group such as halogen etc.)

(Fifteenth step)

The present step is a reaction for reacting the compound (XI) or a compound (XI') which is a tautomer thereof with an allyl compound to synthesize a compound (XV). A compound (XI') can be synthesized, for example, according to the method of Example A-1.

The reaction is performed preferably in the presence of a base (e.g.: cesium carbonate).

A reaction temperature is 0 to 100°C, preferably 10 to 40°C.

As a reaction solvent, dimethylformamide and the like are exemplified.

A reaction time is a few minutes to a few tens hours, preferably 1 to 10 hours.

(Sixteenth step)

The present step is a reaction for oxidizing a compound (XV) to synthesize a compound (XVI). As an oxidizing agent, osmium tetraoxide and alkali metal osmium tetraoxide (e.g.: K_2OsO_4) are exemplified.

A reaction temperaturwe is 0 to 100°C, preferably 10 to 40°C.

As a reaction solvent, 1,4-dioxane, tetrahydrofuran and the like are exemplified.

A reaction time is a few minutes to a few tens hours, preferably 1 to 5 hours.

(Seventeenth step)

The present step is a reaction for reacting a compound (XVI) with amine (XVII) to perform dehydration condensation to synthesize a compound (XVIII).

A reaction temperature is 0 to 200°C, preferably 140 to 180°C.

As a reaction solvent, methylene chloride, acetonitrile and the like are exemplified.

A reaction time is a few minutes to a few tens hours, preferably 0.5 to 1.5 hours.

(Eighteenth step)

The present step is a reaction for deprotecting a compound (XVIII) preferably with an acid to synthesize a compound (XIX), and may be performed according to the condition for a conventional reaction of deprotecting a protected hydroxy group.

A reaction temperature is 0 to 200°C.

As an acid, pyridine hydrochloride, trifluoroacetic acid and the like are exemplified.

As a reaction solvent, the acid and trimethylsilyl iodide are exemplified.

A reaction time is a few minutes to a few tens hours, preferably 15 minutes to 1 hour.

(Nineteenth step)

The present step is a reaction for reducing a compound (XVIII) to synthesize a compound (XX).

As a reducing agent, H_2/Pd · C and the like are exemplified.

A reaction temperatuer is 0 to 100°C, preferably 10 to 30°C.

As a reaction sovelnt, dimethylformamide, methanol, tetrahydrofuran and the like are exemplified.


A reaction time is a few minutes to a few tens hours, preferably 5 to 20 hours.

[0015]

(Process 2)

The intermediate (XVIII) may be also synthesized by a method shown below.

[Chemical formula 43]



(Twentieth step)

The present step is a reaction for reacting a compound (XIV) with a compound (XXI) to synthesize a compound (XXII). The present reaction may be performed according to the condition for a conventional amidation reaction.

A reaction temperature is 0 to 100°C, preferably 0 to 50°C.

As a reaction solvent, dimethylformamide, methylene chloride, tetrahydrofuran and the like are exemplified.

A reaction time is a few minutes to a few tens hours, preferably 1 to 10 hours.

(Twenty-first step)

The present step is a reaction for reacting a compound (XXII) with an acid to perform deprotection and intramolecular ring closure, to synthesize a compound (XXIII). The present reaction may be performed according to the condition for a conventional reaction of deprotecting acetal.

A reaction temperature is 0 to 100°C, preferably room temperature to 80°C.

As a reaction solvent, dioxane, tetrahydrofuran and the like are exemplified.

A reaction time is a few minutes to a few tens hours, preferably 0.5 to 1 hour.

As an acid, hydrochloric acid, and paratoluenesulfonic acid are exemplified.

(Twenty-second step)

The present step is a reaction for dehydrating a compound (XXIII) to

synthesize a compound (XXIV). The present reaction may be peformed according to the condition for a conventional dehydration reaction.

A reaction temperature is 0 to 100°C, preferably room temperature to 80°C.


As a reaction solvent, acetonitrile, methylene chloride and the like are exemplified.


A reaction time is a few minutes to a few tens hours, preferably 1 to 5 hours.

[0016]

(Process 3)

[Chemical formula 44]





(Twenty-third step)

The present step is a reaction for reacting a compound (XVI) with amine (XXIV) to perform dehydration condensation to synthesize a compound (XXV) according to the seventeenth step or a method of synthesizing a compound 17-1. Preferably, as a reaction catalyst, an acid (e.g.: acetic acid) is added, and a microwave reaction apparatus is used.

A reaction temperature is 0 to 200°C, preferably 140 to 180°C.

As a reaction solvent, methylene chloride, acetonitrile and the like are exemplified.

A reaction time is a few minutes to a few tens hours, preferably 0.5 to 1.5 hours.

(Twenty-fourth step)

The present step is a reaction for deprotecting a compound (XXV) preferably with an acid to synthesize a compound (XXVI) according to the eighteenth step, and may be performed according to the condition for a conventional reaction of deprotecting a protected hydroxy group.

A reaction temprature is 0 to 200°C.

As an acid, pyridine hydrochloride, trifluoroacetic acid and the like are exemplified.


As a reaction solvent, the aforementioned acid and trimethylsilyl iodide are exemplified.

A reaction time is a few minutes to a few tens hours, preferably 15 minutes to 1 hour.

[0017]

(Process 4)

[Chemical formula 45]



(Twenty-fifth step)

The present step is a reaction for reacting a compound (XIV) with a compound (XXIV) to synthesize a compound (XXVII) according to the twentieth step. The present reaction may be performed according to the condition for a conventional amidation reaction.

A reaction temperature is 0 to 100°C, preferably 0 to 50°C.

As a reaction solvent, dimethylformamide, methylene chloride, tetrahydrofuran and the like are exemplified.

A reaction time is a few minutes to a few tens hours, preferably 1 to 10 hours.

(Twenty-sixth step)

The present step is a reaction for reacting a compound (XXVII) or a tautomer

thereof with an allyl compound to synthesize a compound (XXVIII) according to the fifteenth step.

A reaction is performed preferably in the presence of a base (e.g.: cesium carbonate).

A reaction temperature is 0 to 100°C, preferably 10 to 40°C.

As a reaction solvent, dimethylformamide and the like are exemplified.

A reaction time is a few minutes to a few tens hours, preferably 1 to 10 hours.

(Twenty-seventh step)

The present step is a reaction for oxidizing a compound (XXVIII) to synthesize a compound (XXIX) according to the sixteenth step.

As an oxidizing agent, osmium tetraoxide and alkali metal osmium tetraoxide (e.g.: K_2OsO_4) are exemplified.

A reaction temperature is 0 to 100°C, preferably 10 to 40°C.

As a reaction solvent 1,4-dioxane, tetrahydrofuran and the like are exemplified.

A reaction time is a few minutes to a few tens hours, preferablky 1 to 5 hours.

(Twenty-eighth step)

The present step is a reaction for dehydration-condensing a compound (XXIX) to synthesize a compound (XXX) according to the seventeenth step or a method of synthesizing a compound 17-1. Preferably, as a reaction catalyst, an acid (e.g.: acetic acid) is added, and a microwave reaction apparatus is used.

A reaction temperature is 0 to 200°C, preferably 140 to 180°C.

As a reaction solvent, methylene chloride, acetonitrile and the like are exemplified.

A reaction time is a few minutes to a few tens hours, preferably 0.5 to 1.5 hours.

(Twenty-ninth step)

The present step is a reaction for deprotecting a compound (XXX) preferably with an acid to synthesize a compound (XXXI) according to the eighteenth step, and may be peformed according to the condition for a conventional reaction of deprotecting a protected hydroxy group.

A reaction temperature is 0 to 200°C.

As an acid, pyridine hydrochloride, trifluoroacetic acid and the like are exemplified.

As a reaction solvent, the aforementioned acid and trimethylsilyl iodide are exemplified.


A reaction time is a few minutes to a few tens hours, preferably 15 minutes to 1 hour.

[0018]

(Process 5)

A compound (I-3) in which Z is NR^{10} can be synthesized according to the following reaction scheme, according to Process 4.


[Chemical formula 46]



[0019]

(Process 10)

[Chemical formula 51]



(wherein respective symbols are as defined above)

(Forty-ninth step)

A compound (XIV-16) is obtained by reacting a compound (XIV) with an amine reagent, according to the thirty-fifth step.

(Fiftieth step)

A compound (XIV-17) is obtained by subjecting a compound (XIV-16) to a general acetal deprotecting reaction according to the forty-fourth step.


(Fifty-first step)

A compound (XIV-18) is obtained (D ring formation) by deprotecting a P^1 part of a compound (XIV-14) according to the thirty-eighth step.

[0020]

The present invention further provides various intermediates (I-P) shown below and a process for preparing the same, as well as a process for preparing the above mentioned compound (I) comprising the deprotection of the intermediate.


(Intermediates)



(I-P)


(P^1 is a hydroxyl-protecting group; the other symbols are as defined above)

Preferred compounds are shown below. Each P^1 is a hydroxyl-protecting group, such as $C_{6-14}arylC_{1-8}alkyl$ (e.g., benzyl (=Bn)).




(I-20a)

Preferably, wherein R^e is one or two halogen; R^z is $C_{1-8}alkyl$, $C_{6-14}arylC_{1-8}alkyl$, $C_{6-14}aryl$, or alkoxy; and P^1 is $C_{6-14}arylC_{1-8}alkyl$;




(I-20b)

Preferably, wherein R^e is one or two halogen; R^z is $C_{1-8}alkyl$, $C_{6-14}arylC_{1-8}alkyl$, $C_{6-14}aryl$, or alkoxy; and P^1 is $C_{6-14}arylC_{1-8}alkyl$;




(I-21a)

Preferably, wherein R^e is one or two halogen; and P^1 is $C_{6-14}arylC_{1-8}alkyl$;




(I-21b)

Preferably, wherein R^e is one or two halogen; and P^1 is $C_{6-14}arylC_{1-8}alkyl$;




(I-22a)

Preferably, wherein R^e is one or two halogen; and P^1 is $C_{6-14}arylC_{1-8}alkyl$;




(I-22b)

Preferably, wherein R^e is one or two halogen; and P^1 is $C_{6-14}arylC_{1-8}alkyl$;




(I-23a)

Preferably, wherein R^e is one or two halogen; and P^1 is $C_{6-14}arylC_{1-8}alkyl$;




(I-23b)

Preferably, wherein R^e is one or two halogen; and P^1 is $C_{6-14}arylC_{1-8}alkyl$;




(I-24a)

Preferably, wherein R^e is one or two halogen; R^z is C_{1-8}alkyl; R^{z1} is hydrogen, C_{3-6}cycloalkyl, , heterocycle, or C_{1-8}alkyl optionally substituted with hydroxy, C_{3-6}cycloalkyl, alkoxy, heterocycle, heteroaryl, C_{6-14}aryl, or amino, wherein said amino may be optionally substituted with $-C(O)C_{1-8}$alkyl or C_{1-8}alkyl;




(I-24b)

Preferably, wherein R^e is one or two halogen; R^z is C_{1-8}alkyl; R^{z1} is hydrogen, C_{3-6}cycloalkyl, , heterocycle, or C_{1-8}alkyl optionally substituted with hydroxy, C_{3-6}cycloalkyl, alkoxy, heterocycle, heteroaryl, C_{6-14}aryl, or amino, wherein said amino may be optionally substituted with $-C(O)C_{1-8}$alkyl or C_{1-8}alkyl; and P^1 is C_{6-14}arylC_{1-8}alkyl;




(I-25)

Preferably, wherein R^e is one or two halogen; R^{z1} is hydrogen, C_{3-6}cycloalkyl, , heterocycle, or C_{1-8}alkyl optionally substituted with hydroxy, C_{3-6}cycloalkyl, alkoxy, heterocycle, heteroaryl, C_{6-14}aryl, or amino, wherein said amino may be optionally substituted with $-C(O)C_{1-8}$alkyl or C_{1-8}alkyl;and P^1 is C_{6-14}arylC_{1-8}alkyl;



(I-26)


Preferably, wherein R^e is one or two halogen; R^{z1} is hydrogen, C_{3-6}cycloalkyl, , heterocycle, or C_{1-8}alkyl optionally substituted with hydroxy, C_{3-6}cycloalkyl, alkoxy, heterocycle, heteroaryl, C_{6-14}aryl, or amino, wherein said amino may be optionally substituted with $-C(O)C_{1-8}$alkyl or C_{1-8}alkyl; and P^1 is C_{6-14}arylC_{1-8}alkyl;



(I-27)


Preferably, wherein R^e is halogen; and P^1 is C_{6-14}arylC_{1-8}alkyl;

The above intermediates, compound (I-20a), (I-20b), (I-21a), (I-21b), (I-22a), (I-22b), (I-23a), (I-23b), (I-24a), (I-24b), (I-25), (I-26), or (I-27), can be prepared by condensing a compound of the formula:




wherein R^e is one or two halogen; and R^{50} is C_{1-8}alkyl;


with each amine shown below, respectively:




wherein R^z is C_{1-8}alkyl, C_{6-14}arylC_{1-8}alkyl, C_{6-14}aryl, or alkoxy;




wherein R^z is C_{1-8}alkyl, C_{6-14}arylC_{1-8}alkyl, C_{6-14}aryl, or alkoxy;




wherein R^z is C_{1-8}alkyl; R^{z1} is hydrogen, C_{3-6}cycloalkyl, , heterocycle, or C_{1-8}alkyl optionally substituted with hydroxy, C_{3-6}cycloalkyl, alkoxy, heterocycle, heteroaryl, C_{6-14}aryl, or amino, wherein said amino may be optionally substituted with –C(O)C_{1-8}alkyl or C_{1-8}alkyl;




wherein R^z is C_{1-8}alkyl; R^{z1} is hydrogen, C_{3-6}cycloalkyl, , heterocycle, or C_{1-8}alkyl optionally substituted with hydroxy, C_{3-6}cycloalkyl, alkoxy, heterocycle, heteroaryl, C_{6-14}aryl, or amino, wherein said amino may be optionally substituted with –C(O)C_{1-8}alkyl or C_{1-8}alkyl;



wherein R^{z1} is hydrogen, C_{3-6}cycloalkyl, , heterocycle, or C_{1-8}alkyl optionally substituted with hydroxy, C_{3-6}cycloalkyl, alkoxy, heterocycle, heteroaryl, C_{6-14}aryl, or amino, wherein said amino may be optionally substituted with –C(O)C_{1-8}alkyl or C_{1-8}alkyl;



wherein R^{z1} is hydrogen, C_{3-6}cycloalkyl, , heterocycle, or C_{1-8}alkyl optionally substituted with hydroxy, C_{3-6}cycloalkyl, alkoxy, heterocycle, heteroaryl, C_{6-14}aryl, or amino, wherein said amino may be optionally substituted with $-C(O)C_{1-8}$alkyl or C_{1-8}alkyl;



The condition for the above condensation is illustrated below for example.

Examples of the solvent include halocarbons such as dichloromethane, dichloroethane, and acetic acid.

The reaction temperature is preferably, 0 to 200 ℃, more preferably, 50 to 170℃.

The reaction time is usually several minutes to several hours.

The above intermediates, compound (I-20a), (I-20b), (I-21a), (I-21b), (I-22a), (I-22b), (I-23a), (I-23b), (I-24a), (I-24b), (I-25), (I-26), or (I-27), can be deprotected to give each corresponding deprotected compound wherein P^1 is hydrogen, or its pharmaceutically acceptable salt, which are encompassed within the scope of compound (I) of the present invention.

In addition, the present compound obtained above may be further chemically modified to synthesize another compound. In addition, when there is a reactive functional group (e.g.: OH, COOH, NH_2) on a side chain part etc. in the above reaction, the group may be protected before the reaction and may be deprotected after the reaction, if desired.

The present compound is useful, for example, as a drug such as an anti-virus drug. The present compound has the remarkable inhibitory action on integrase of a virus. Therefore, the present compound can be expected to have the preventive or

therapeutic effect for various diseases derived from a virus which produces at least integrase, and is grown at infection in an animal cell, and is useful as an integrase inhibiting agent for retrovirus (e.g. HIV-1, HIV-2, HTLV-1, SIV, FIV etc.), and is useful as an anti-HIV drug etc.

In addition, the present compound may be used in joint use therapy by combining an anti-HIV drug having the different action methanism such as a reverse trascriptase inhibiter and/or a protease inhibiting agent. Particularly, currently, an integrase inhibiter is not marketed, and it is useful to use in joint use therapy by combining the present compound with a reverse transcriptase inhibiter and/or a protease inhibiter.

Further, the above use includes not only use as a medical mixture for anti-HIV, but also use as a joint use agent for increasing the anti-HIV activity of other anti-HIV drug such as cocktail therapy.

In addition, the present compound can be used in order to prevent infection with a retrovirus vector from spreading into a tissue other than an objective tissue, upon use of a retrovirus vector based on HIV or MLV in the field of gene therapy. Particularly, when a cell is infected with a vector in vitro, and the cell is returned into a body, if the present compound is administered in advance, extra infection can be prevented in a body.

The present compound can be adminitetered orally or parenterally. In the case of oral administration, the present compound can be also used as a conventional preparation, for example, as any dosage form of a solid agent such as tablets, powders, granules, capsules and the like; an aqueous agent; an oily suspension; or a liquid agent such as syrup and elixir. In the case of parenteral administration, the present compound can be used as an aqueous or oily suspension injectable, or a nasal drop. Upon preparation of it, conventional excipients, binders, lubricants, aqueous solvents, oily solvents, emulsifiers, suspending agents, preservatives, stabilizers and the like may be arbitrarily used. As an anti-HIV-drug, particularly, an oral agent is preferable. A preparation of the present invention is prepared by combining (e.g. mixing) a therapeutically effective amount of the present compound with a pharmaceutically acceptable carrier or diluent.

A dose of the present invention is different depeniding on an administration method, an age, a weight and conditionn of a patient, and a kind of a disease and, usually, in the case of oral administraton, about 0.05mg to 3000mg, preferably about 0.1mg to 1000mg may be administered per adult a day, if necessary, by dividing the

dose. In addition, in the case of parenteral administration, about 0.01mg to 1000mg, preferably about 0.05mg to 500mg is administered per adult a day.

Examples are shown below.

[0025]


Example A-1)


9-Hydroxy-2-(2-methoxy-ethyl)-1,8-dioxo-1,8-dihydro-2H-pyrid[1,2-a]pyrazine -7-carboxylic acid 4-fluoro-banzylamide

Example B-1)

9-Hydroxy-2-(2-methoxy-ethyl)-1,8-dioxo-1,3,4,8-tetrahydro-2H-pyrid[1,2-a]py razine-7-carboxylic acid 4-fluoro-benzylamide

[Chemical formula 52]





1) Maltol 1 (189g, 1.5mol) was dissolved in dimethylformamide (1890ml), and benzyl bromide (184ml, 1.5mol) was added. After the solution was stirred at 80°C for 15 minutes, potassium carbonate (228g, 1.65mol) was added, and the mixture was stirred for 1 hour. After the reaction solution was cooled to room temperature, an inorganic salt was filtered, and the filtrate was distilled off under reduced pressure. To the again precipitated inorganic salt was added tetrahydrofuran (1000ml), this was filtered, and the filtrate was distilled off under reduced pressure to obtain the crude product (329g, >100%) of 3-benzyloxy-2-methyl-pyran-4-one 2 as a brown oil.
NMR ($CDCl_3$)δ: 2.09(3H, s), 5.15(2H, s), 6.36(1H, d, J=5.6Hz), 7.29-7.41(5H, m), 7.60(1H, d, J=5.6Hz).

2) The compound 2 (162.2g, 750mmol) was dissolved in ethanol (487ml), and aqueous ammonia (28%, 974ml) and a 6N aqueous sodium hydroxide solution (150ml, 900mmol) were added. After the reaction solution was stirred at 90 °C for 1 hour, this was cooled to under ice-cooling, and ammonium chloride (58g, 1080mmol) was added. To the reaction solution was added chloroform, this was extracted, and the organic layer was washed with an aqueous saturated sodium bicarbonate solution, and dried with anhydrous sodium sulfate. The solvent was distilled off under reduced pressure, isopropyl alcohol and diethyl ether were added to the residue, and precipitated crystals were filtered to obtain 3-benzyloxy-2-methyl-1H-pyridine-4-one 3 (69.1g, 43%) as a pale yellow crystal.
NMR (DMSO-d_6)δ: 2.05(3H, s), 5.04(2H, s), 6.14(1H, d, J=7.0Hz), 7.31-7.42(5H, m), 7.46(1H, d, J=7.2Hz), 11.29(1H, brs).

3) The above compound 3 (129g, 599mmol) was suspended in acetonitrile (1300ml), and N-bromosuccinic acid imide (117g, 659mmol) was added, followed by stirring at room temperature for 90 minutes. Precipitated crystals were filtered, and washed with acetonitrile and diethyl ether to obtain 3-benzyloxy-5-bromo-2-methyl-pyridine-4-ol 4(154g, 88%) as a colorless crystal.
NMR (DMSO-d_6)δ: 2.06(3H, s), 5.04(2H, s), 7.32-7.42(5H, m), 8.03(1H, d, J=5.5Hz), 11.82(1H, brs).

4) To a solution of the compound 4 (88g, 300mmol), palladium acetate (13.4g, 60mmol) and 1,3-bis(diphenylphosphino)propane (30.8g, 516mmol) in dimethylformamide (660ml) were added methanol (264ml) and triethylamine (210ml, 1.5mol) at room temperature. The interior of a reaction vessel was replaced with carbon monoxide, and the material was stirred at room temperature for 30 minutes, and stirred at 80 degree for 18 hours. A vessel to which ethyl acetate (1500ml), an aqueous saturated ammonium chloride solution (1500ml) and water (1500ml) had been added was stirred under ice-cooling, and the reaction solution was added thereto. Precipitates were filtered, and washed with water (300ml), ethyl acetate (300ml) and diethyl ether (300ml) to obtain 5-benzyloxy-4-hydroxy-6-methyl-nicotinic acid methyl ester 5 (44.9g, 55%) as a colorless crystal.
NMR (DMSO-d_6)δ: 2.06(3H, s), 3.72(3H, s), 5.02(2H, s), 7.33-7.42(5H, m), 8.07(1H, s).

5) After a solution of the compound 5 (19.1g, 70mmol) in acetic anhydride (134ml) was stirred at 130 °C for 40 minutes, the solvent was distilled off under reduced pressure to obtain 4-acetoxy-5-benzyloxy-6-methyl-nicotinic acid methyl ester 6 (19.9g, 90%) as a flesh colored crystal.
NMR ($CDCl_3$)δ: 2.29(3H, s), 2.52(3H, s), 3.89(3H, s), 4.98(2H, s), 7.36-7.41(5H, m), 8.85(1H, s).

6) To a solution of the compound 6 (46.2g, 147mmol) in chloroform (370ml) was added metachloroperbenzoic acid (65%) (42.8g, 161mmol) in portions under ice-cooling, and this was stirred at room temperature for 90 minutes. To the reaction solution was added a 10% aqueous potassium carbonate solution, and this was stirred for 10 minutes, followed by extraction with chloroform. The organic layer was washed with successively with a 10% aqueous potassium carbonate solution, an aqueous saturated ammonium chloride solution, and an aqueous saturated sodium chloride solution, and

dried with anhydrous sodium sulfate. The solvent was distilled off under reduced pressure, and the residue was washed with diisopropyl ether to obtain 4-acetoxy-5-benzyloxy-6-methyl-1-oxy-nicotinic acid methyl ester 7 (42.6g, 87%) as a colorless crystal.
NMR ($CDCl_3$)δ: 2.30(3H, s), 2.41(3H, s), 3.90(3H, s), 5.02(2H, s), 7.37-7.39(5H, m), 8.70(1H, s).

7) To acetic anhydride (500ml) which had been heated to stir at 130 °C was added the compound 7 (42.6g, 129mmol) over 2 mintues, and this was stirred for 20 minutes. The sovent was distilled off under reduced pressure to obtain 4-acetoxy-6-acetoxymethyl-5-benzyloxy-nicotinic acid methyl ester 8 (49.6g, >100%) as a black oil.
NMR ($CDCl_3$)δ: 2.10(3H, s), 2.28(3H, s), 3.91(3H, s), 5.07(2H, s), 5.20(2H, s), 7.35-7.41(5H, m), 8.94(1H, s).

8) To a solution of the compound 8 (46.8g, 125mmol) in methanol (140ml) was added a 2N aqueous sodium hydroxide solution (376ml) under ice-cooling, and this was stirred at 50 °C for 40 minutes. To the reaction solution were added diethyl ether and 2N hydrochloric acid under ice-cooling, and precipitated crystals were filtered. Resulting crystals were washed with water and diethyl ether to obtain 5-benzyloxy-4-hydroxy-6-hydroxymethyl-nicotinic acid 9 (23.3g, 68%) as a colorless crystal.
NMR (DMSO-d_6)δ: 4.49(2H, s), 5.19(2H, s), 5.85(1H, brs), 7.14-7.20(2H, m), 7.33-7.43(7H, m), 8.30(1H, s), 10.73(1H, t, J=5.8Hz), 11.96(1H, brs).

9) To a solution of the compound 9 (131g, 475mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (219g, 1140mmol) and 1-hydroxybenzotriazole (128g, 950mmol) in dimethylformamide (1300ml) was added 4-fluorobenzylamine (109ml, 950mmol), and this was stirred at 80°C for 1.5 hours. After the reaction solution was cooled to room temperature, hydrochloric acid was added, followed by extraction with ethyl acetate. The extract was washed with a 5% aqueous potassium carbonate solution, an aqueous saturated ammonium chloride solution, and an aqueous saturated sodium chloride solution, and dried with anhydrous sodium sulfate. The solvent was distilled off under reduced pressure to obtain a mixture (175g) of 10 and 11. the resulting mixture was dissolved in acetic

acid (1050ml) and water (1050ml), and zinc (31.1g, 475mmol) was added, followed by heating to reflux for 1 hour. After the reaction solution was cooled to room tempreture, a 10% aqueous potassium carbonate solution was added, followed by extraction with ethyl acetate. The extract was washed with an aqueous saturated ammonium chloride solution, and an aqueous saturated sodium chloride solution, and dried with anhydrous sodium sulfate. After the solvent was distilled off under reduced pressure, this was washed with diethyl ether to obtain 5-benzyloxy-N-(4-fluoro-benzyl)-4-hydroxy-6-hydroxymethyl-nicotinic acid amide 10 (107g, 59%) as a colorless crystal.
NMR (DMSO-d_6)δ: 4.45(2H, d, J=4.3Hz), 4.52(2H, d, J=5.8Hz), 5.09(2H, s), 6.01(1H, brs), 7.36-7.43(5H, m), 8.31(1H, s), 12.63(1H, brs).

10) After manganese dioxide (49g) was added to a suspension of the compound 10 (9.8g, 25.6mmol) in chloroform (490ml), the mixture was stirred at room temperature for 1 hour. After the reaction solution was stirred at 60 °C for 20 minutes, Celite filtration was performed, and this was washed with chloroform heated at 50 °C. The filtrate was distilled off under reduced pressure to obtain 5-benzyloxy-N-(4-fluoro-benzyl)-6-formyl-4-hydroxy-nicotinic acid amide 12 (8.2g, 84%) as a pale yellow crystal.
NMR (DMSO-d_6)δ: 4.53(2H, d, J=5.8Hz), 5.38 (2H, s), 7.15-7.21(2H, m), 7.35-7.46(7H, m), 8.33(1H, s), 9.90(1H, s), 10.35(1H, t, J=5.8Hz), 12.49(1H, brs).

11) To an aqueous solution (105ml) of sodium chlorite (7.13g, 78.8mmol), and sulfamic acid (7.65g, 78.8mmol) was added a solution of the compound 12 (15.0g, 39.4mmol) in tetrahydrofuran (630ml) under ice-coling, and the mixture was stirred at room temperature for 1 hour. After water (2500ml) was added to the reaction solution, precipitated crystals were filtered. Washing with diethyl ether afforded 3-benzyloxy-5-(4-fluoro-benzylcarbamoyl)-4-hydroxy-pyridine-2-carboxylic acid 13 (14.0g, 90%) as a colorless crystal.
NMR (DMSO-d_6)δ: 4.52(2H, d, J=5.8Hz), 5.13 (2H, s), 7.14-7.19(2H, m), 7.31-7.40(5H, m), 7.47-7.49(2H, m), 8.31(1H, d, J=4.5Hz), 10.44(1H, t, J=5.9Hz), 12.47(1H, brs).

12) A solution of the compound 13 (198mg, 0.500mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (115mg, 0.600mmol) and 1-hydroxybenzotriazole (81mg, 0.600mmol) in dimethylformamide (3ml) was

stirred at room temperature for 1.5 hours. Then, methanol (3ml) and triethylamine (153ul, 1.10mmol) were added, and the mixture was heated to reflux for 1.5 hours. The reaction solution was diluted with ethyl acetate, washed with an aqueous saturated sodium bicarbonate solution, a 10% aqueous citric acid solution, and an aqueous saturated sodium chloride solution, and dried with anhydrous sodium sulfate. The solvent was distilled off under reduced pressure, and the residue was washed with diethyl ether to obtain
3-benzyloxy-5-(4-fluoro-benzylcarbamoyl)-4-hydroxy-pyridine-2-carboxylic acid methyl ester 14 (141mg, 69%) as a colorless crystal.
NMR (DMSO-d_6)δ: 3.85(3H, s), 4.52(2H, d, J=6.0Hz), 5.15(2H, s), 7.13-7.21(2H, m), 7.31-7.47(7H, m), 8.33(1H, s), 10.41(1H, t, J=6.0Hz), 12.59(1H, brs).

13) After 3-bromopropene (2.15ml, 24.8mmol) was added to a solution of the compound 14 (6.79g, 16.5mmol), and cesium carbonate (8.09g, 24.8mmol) in dimethylformamide (54ml), the mixture was stirred at room temperature for 4.5 hours. To the reaction solution was added an aqueous ammonium chloride solution, and this was extracted with ethyl acetate, washed with water and an aqueous saturated sodium chloride solution, and dried with anhydrous sodium sulfate. The solvent was distilled off under reduced pressure, and the residue was washed with diethyl ether to obtain
1-allyl-3-benzyloxy-5-(4-fluoro-benzylcarbamoyl)-4-oxo-1,4-dihydro-pyridine-2-carboxylic acid methyl ester 15 (6.15g, 83%) as a colorless crystal.
NMR ($CDCl_3$) δ: 3.76(3H, s), 4.54(2H, d, J=6.0Hz), 4.60(2H, d, J=6.0Hz), 5.20-5.37(2H, m), 5.25(2H, s), 5.80-5.93(1H, m), 6.98-7.04(2H, m), 7.31-7.35(7H, m), 8.45(1H, s), 10.41(1H, m).

14) To a solution of the compound 15 (7.6g, 16.9mmol) in 1,4-dioxane (228ml) was added an aqueous solution (38ml) of potassium osmate dihydrate (372mg, 1.01mmol), and sodium metaperiodate (14.5g, 67.6mmol) was further added, followed by stirring at room temperature for 2 hours. The reaction solution was added to a vessel to which ethyl acetate (300ml) and water (300ml) had been added, while stirring. The organic layer was washed with water, a 5% aqueous sodium hydrogen sulfite solution and an aqueous saturated sodium chloride solution, and dried with anhydrous sodium sulfate. The solvent was distilled off under reduced pressure, and the residue was washed with diethyl ether to obtain

3-benzyloxy-5-(4-fluoro-benzylcarbamoyl)-4-oxo-1-(2-oxo-ethyl)-1,4-dihydro-pyridine-2-carboxylic acid methyl ester 16 (5.39g, 71%) as a colorless crystal.
NMR ($CDCl_3$)δ: 3.74(3H, s), 4.60(2H, d, J=5.9Hz), 4.87(2H, s), 5.27(2H, s), 6.98-7.04(2H, m), 7.30-7.40(7H, m), 8.39(1H, s), 9.58(1H, s), 10.38(1H, s).

15) To a solution of the compound 16 (400mg, 0.884mmol) in methylene chloride (12ml) were added 2-methoxyethylamine (77ul, 0.884mmol) and acetic acid (18ul), and the mixture was stirred at room temperature for 5 minutes. Thereafter, the reaction was performed at 140 °C for 30 minutes in a microwave reaction apparatus. The solvent was distilled off under reduced pressure, the residue was subjected to silica gel column chromatography, and fractions eluting with toluene-acetone were concentrated under reduced pressure to obtain 9-benzyloxy-2-(2-methy-ethyl)-1,8-dioxo-1,8-dihydro-2H-pyrid[1,2-a]pyrazine-7-carboxylic acid 4-fluoro-benzylamide 17-1 (226mg, 54%) as a yellow solid.
NMR ($CDCl_3$)δ: 3.35(3H, s), 3.65(2H, t, J=5.1Hz), 3.97(2H, t, J=4.5Hz), 4.63(2H, d, J=5.7Hz), 5.28(2H, s), 6.56(2H, m), 7.01(2H, t, J=8.7Hz), 7.38-7.30(5H, m), 7.65(2H, d, J=6.6Hz), 10.63(1H, s).

16) To the compound 17-1 (140mg, 0.293mmol) was added trifluoroacetic acid (1.4ml) under ice-cooling, and the mixture was stirred at 0 °C for 5 minutes and, then, at room temperature for 1.5 hours. The solvent was distilled off under reduced pressure, and this was diluted with chloroform, and added to ice water. This was washed with an aqueous saturated sodium bicarbonate solution, a 10% aqueous citric acid solution and water, and dried with anhydrous sodium sulfate. The solvent was distilled off under reduced pressure, and the residue was recrystallized with methylene chloride-ethanol to obtain Example A-1 (89mg, 79%) as a yellow crystal.
melting point: 223-224 °C
NMR (DMSO-d_6)δ: 3.25(3H, s), 3.58(2H, t, J=5.4Hz), 3.92(2H, t, J=5.1Hz), 4.53(2H, d, J=5.7Hz), 6.87(1H, d, 6.3Hz), 7.14(2H, t, J=9.0Hz), 7.33-7.38(2H, m), 7.47(1H, d, J=6.0Hz), 8.77(1H, s), 10.56(1H, t, J=6.0Hz), 12.00(1H, brs).

17) The compound 17-1 (157mg, 0.329mmol) was dissolved in dimethylformamide (18ml) and methanol (1ml), 10% palladium-carbon powder (31mg) was added, and the mixture was stirred at room temperature for 20 hours under the hydrogen atmosphere. The reaction solution was filtered with Celite, and the filtrate was


concentrated under reduced pressure. The residue was dissolved in chloroform, this was filtered with Celite again, and the filtrate was concentrated under reduced pressure. The residue was recrystallized with methylene chloride-methanol to obtain Example B-1 (66mg, 52%) as a brown crystal.

melting point: 197-199 °C

NMR (DMSO-d_6)δ:3.27(3H, s), 3.55(2H, t, J=5.1Hz), 3.68(2H, t, J=5.1Hz), 3.79(2H, s), 4.36(2H, s), 4.51(2H, d, J=5.7Hz), 7.15(2H, t, J=8.7Hz), 7.32-7.37(2H, m), 8.38(1H, s), 10.46(1H, t, J=5.4Hz), 12.41(1H, s).

Example C-1

[Chemical formula 55]



1) A compound 33 was synthesized using 1-aminomethylcyclopentanol hydroxyethylamine according to the method of synthesizing a compound 17-1.

1H-NMR ($CDCl_3$)δ: 1.30-1.80(10H, m), 3.47(1H, d, J=11.4Hz), 3.61(1H, d, J=11.4Hz), 3.80-3.95(1H, m), 4.30(1H, dd, J=14.7, 3.0Hz), 4.60(2H, d, J=5.7Hz), 5.17-5.23(2H, m), 5.39(1H, d, J=9.9Hz), 6.95-7.10(2H, m), 7.20-7.40(5H, m), 7.58(2H, d, J=7.2Hz), 8.41(1H, s), 10.40(1H, s).

2) A compound 33-2 was synthesized using hydroxyethylamine according to the similar method.

Compound 33-2)

5-Benzyloxy-4,6-dioxo-2,3,4,6,9,9a-hexahydro-1-oxa-3a,8a-diaza-cyclopenta[b]naphthalene-7-carboxylic acid 4-fluorobenzylamide

1H-NMR (DMSO-d_6)δ: 3.48-3.58(1H, m), 3.73-3.86(1H, m), 3.97-4.10(2H, m), 4.20-4.30(1H, m), 4.46-4.60(2H, m), 4.85(1H, dd, J=12.3, 3.5Hz), 5.40(1H, d, J=10.2Hz),

5.18(1H, d, J=10.2Hz), 5.28(1H, dd, J=10.2, 3.2Hz), 7.10-7.20(2H, m), 7.23-7.40(5H, m), 7.50-7.73(2H, m), 8.60(1H, s), 10.22(1H, m).

3) Example C-1 was synthesized using a compound 33, according to the method of synthesizing Example A-1.

Melting point: >300°C

1H-NMR (DMSO-d_6)δ: 1.10-1.60(10H, m), 3.25(1H, d, J=11.4Hz), 3.37(1H, d, J=11.4Hz), 3.76(1H, t, J=10.5Hz), 4.30(2H, d, J=5.8Hz), 4.66(1H, dd, J=12.2, 3.8Hz), 5.22(1H, dd, J=3.8, 10.4Hz), 6.90-6.96(2H, m), 7.10-7.15(2H, m), 8.25(1H, s), 10.10(1H, brs), 11.32(1H, brs).

The following compounds were synthesized using the similar method.

Example C-2)

5-Hydroxy-4,6-dioxo-2,3,4,6,9,9a-hexahydro-1-oxa-3a,8a-diaza-cyclopenta[b]naphthalene-7-carboxylic acid 4-fluorobenzylamide

Melting point: 272-274 °C

1H-NMR (DMSO-d_6)δ: 3.59-3.67(1H, m), 3.72-3.81(1H, m), 3.98-4.10(2H, m), 4.27-4.35(1H, m), 4.52(2H, d, J=7.2Hz), 4.92(1H, dd, J=12.3, 12.3Hz), 5.27(1H, dd, J=3.6, 9.9Hz), 7.11-7.20(2H, m), 7.30-7.40(2H, m), 8.49(1H, s), 10.32(1H, t, J=5.6Hz), 11.53(1H, s).

Example C-3)

5-Hydroxy-6,10-dioxo-3,4,6,9,9a,10-hexahydro-2H-1-oxa-4a,8a-diazaanthracene-7-carboxylic acid 4-fluorobenzylamide

melting point: 259 °C

1H-NMR (DMSO-d_6)δ: 1.60-1.67(1H, m), 1.72-1.85(1H, m), 3.25(1H, td, J=12.8, 3.5Hz), 3.86-3.93(1H, m), 4.06(1H, dd, J=11.4, 4.2Hz), 4.44-4.57(5H, m), 5.28(1H, t, J=3.8Hz), 7.13-7.18(2H, m), 7.33-7.37(2H, m), 8.51(1H, s), 10.36(1H, t, J=6.0Hz), 12.47(1H, s).

Example C-4)

5-Hydroxy-1-isopropyl-4,6-dioxo-2,3,4,6,9,9a-hexahydro-1H-1,3a,8a-triaza-cyclopenta[b]naphthalene-7-carboxylic acid 4-fluoro-benzylamide

melting point: 232-234°C

NMR (DMSO-d_6)δ: 1.03(3H, d, 6.6Hz), 1.14(3H, d, 6.6Hz), 2.79-3.66(5H, m), 3.82(1H, t, 10.8Hz), 4.51(3H, m), 4.90(1H, m), 7.15(2H, t, 9.0Hz), 7.34(2H, m), 8.45(1H, s),

10.39(1H, t, 5.4Hz), 11.60(1H, s).

Example C-5)

5-Hydroxy-4,6-dioxo-2,3,4,6,9,9a-hexahydro-1H-1,3a,8a-triaza-cyclopenta[b]naphthalene-7-carboxylic acid 4-fluoro-benzylamide

melting point: 256-258 °C

NMR (DMSO-d_6)δ: 3.00-3.55(5H, m), 3.96(1H, t, 11.4Hz), 4.52(2H, d, 11.7Hz), 4.76(2H, m), 7.16(2H, t, 8.7Hz), 7.35(2H, m), 8.48(1H, s), 10.42(1H, t, 5.4Hz), 11.91(1H, s).

Example C-6)

5-Hydroxy-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthracene-7-carboxylic acid 4-fluoro-benzylamide

melting point: 255°C

NMR (DMSO-d_6)δ: 1.60(1H, s), 2.75-3.16(4H, m), 4.52(2H, d, 6.0Hz), 4.13-4.68(4H, m), 7.16(2H, 9.0Hz, t), 7.34(2H, m), 10.42(1H, s), 10.44(1H, 6.0Hz, t), 12.81(1H, s).

Example C-7)

1-(2-Diethylamino-ethyl)-5-hydroxy-4,6-dioxo-2,3,4,6,9,9a-hexahydro-1H-1,3a,8a-triaza-cyclopenta[b]naphthalene-7-carboxylic acid 4-fluoro-benzylamide

melting point: 186-187 °C

NMR (DMSO-d_6)δ: 0.97(6H, t, 7.2Hz), 2.42-2.91(10H, m), 3.44-3.87(5H, m), 4.23(1H, m), 4.51(2H, d, 5.7Hz), 5.00(1H, m), 7.16(2H, t, 9.0Hz), 7.33-7.37(2H, m), 8.43(1H, s), 10.39(1H, t, 5.7Hz), 11.81(1H, s).

Example C-8)

1-Hydroxy-2,11-dioxo-2,5,5a,7,8,9,10,11-octahydro-6-oxa-4a,10a-diaza-cyclohepta[b]naphthalene-3-carboxylic acid 4-fluoro-benzylamide

melting point: 242-244 °C

NMR (DMSO-d_6)δ: 1.40-2.00(4H, m), 3.20-3.30(1H, m), 3.66-3.77(2H, m), 4.14-4.23(1H, m), 4.38-4.41(1H, m), 4.52(2H, d, 6.3Hz), 4.58-4.63(1H, m), 5.34(1H, brs), 7.15(2H, t, 9.0Hz), 7.33-7.37(2H, m), 8.50(1H, s), 10.39(1H, brs), 12.14(1H, s).

Example C-9)

5-Hydroxy-1-(2-hydroxy-ethyl)-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthracene-7-carboxylic acid 4-fluoro-benzylamide

NMR (DMSO-d_6)δ: 1.58-1.80(1H, m), 2.70-3.60(7H, m), 4.40-4.54(6H, m), 4.77-4.82(1H, m), 7.15(2H, t, 9.0Hz), 7.33-7.38(2H, m), 8.52(1H, s), 10.43(1H, brs), 12.57(1H, s).

Example C-10)

1-Hydroxy-2,11-dioxo-2,5a,6,7,8,9,10,11-octahydro-5H-4a,6,10a-triaza-cyclohepta[b]naphthalene-3-carboxylic acid 4-fluoro-benzylamide

melting point: 256°C

NMR (DMSO-d_6)δ: 1.47-1.77(4H, m), 2.69-2.81(2H, m), 3.34-3.41(1H, m), 4.08-4.12(1H, m), 4.26-4.40(2H, m), 4.52(2H, d, J=6.0Hz), 7.15(2H, t, 8.8Hz), 7.33-7.36(2H, m), 8.43(1H, s), 10.46(1H, t, J=6.0Hz), 12.68(1H, s).

Example C-11)

5-Hydroxy-1-(2-methoxy-ethyl)-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthracene-7-carboxylic acid 4-fluoro-benzylamide

melting point: 147°C

NMR (DMSO-d_6)δ: 1.56-1.74(2H, m), 2.53-2.58(1H, m), 2.66-3.10(4H, m), 3.18(3H, s), 3.41-3.39(2H, m), 4.37-4.52(5H, m), 4.73-4.80(1H, m), 7.15(2H, t, 8.8Hz), 7.33-7.37(2H, m), 8.56(1H, s), 10.40(1H, t, J=6.0Hz), 12.62(1H, s).

Example C-12)

5-Hydroxy-1-(2-isopropoxy-ethyl)-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthracene-7-carboxylic acid 4-fluoro-benzylamide

melting point: 151 °C

NMR (DMSO-d_6)δ: 1.02(6H, dd, J=4.0, 6.0Hz), 1.56-1.67(2H, m), 2.53-2.58(1H, m), 2.74-3.04(4H, m), 3.18(3H, s), 3.41-3.52(3H, m), 4.41-4.59(5H, m), 4.79-4.83(1H, m), 7.15(2H, t, 8.8Hz), 7.34-7.36(2H, m), 8.58(1H, s), 10.40(1H, t, J=6.0Hz), 12.56(1H, s).

Example C-13)

5-Hydroxy-3,3-dimethyl-6,10-dioxo-3,4,6,9,9a,10-hexahydro-2H-1-oxa-4a,8a-diaza-anthracene-7-carboxylic acid 4-fluoro-benzylamide

melting point: 275-277 °C

NMR (DMSO-d_6)δ: 2.97(3H, s), 3.01(3H, s), 3.00-3.18(3H, m), 4.45-4.56(5H, m), 5.16(1H, s), 7.15(2H, t, J=9Hz), 7.35(2H, dd, J=5.4Hz, 8.7Hz), 8.51(1H, s), 10.36(1H, t, J=5.7Hz), 12.4(1H, s).

Example C-14)

1-Cyclohexyl-5-hydroxy-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthrac
ne-7-carboxylic acid-4-fluoro-benzylamide

melting point: 275-277 °C

NMR (DMSO-d_6)δ: 1.22-1.70(2H, m), 2.50-3.02(3H, m), 4.45(4H, m), 4.52(2H, s),
4.78(1H, d, J=13.2Hz), 7.16(2H, t, J=8.7Hz), 7.35(2H, dd, J=5.7Hz, 8.4Hz), 8.62(1H, s)
10.52(1H, s), 12.55(1H, s).

Example C-15)

5-Hydroxy-1-isopropyl-6,10-dioxo-1,2,3,4,6,9,9a, 10-octahydro-1,4a,8a-triaza-anthrace
ne-7-carboxylic acid-4-fluoro-benzylamide

melting point: 220 °C

NMR (DMSO-d_6)δ: 0.94(6H, d, J=9.6Hz), 1.53-1.67(2H, m), 2.92-3.30(3H, m),
4.32-4.40(4H, m), 4.52(2H, d, J=5.7Hz), 4.89(1H, d, J=14.1Hz), 7.16(2H, t, J=9.0Hz),
7.35(2H, dd, J=6.3Hz, 9.0Hz), 8.61(1H, s), 10.46(1H, s), 12.55(1H, s).

Example C-16)

5-Hydroxy-3,3-dimethyl-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthrac
ene-7-carboxylic acid 4-fluoro-benzylamide

melting point: 280 °C

NMR (DMSO-d_6)δ: 0.87(3H, s), 0.93(3H, s), 2.59-3.15(6H, m), 4.09-4.57(6H, m),
7.14(2H, d, J=9.0Hz), 7.34(2H, dd, J=5.4Hz, 8.4Hz), 8.42(1H, s), 10.46(1H, s),
12.77(1H, s).

Example C-17)

5-Hydroxy-1-(2-morpholin-4-yl-2-oxo-ethyl)-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4
a,8a-triaza-anthracene-7-carboxylic acid 4-fluoro-benzylamide

melting point: 140 °C

NMR (DMSO-d_6)δ: 1.60(2H, m), 2.91-3.62(13H, m), 4.41(2H, m), 4.51(2H, d, J=4.8Hz),
4.80(2H, m), 7.15(2H, t, J=8.7Hz), 7.34(2H, m), 8.44(1H, s), 10.43(1H, s), 12.54(1H, s).

Example C-18

1-(3-Acetylamino-propyl)-5-hydroxy-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-tri
aza-anthracene-7-carboxylic acid 4-fluoro-benzylamide

melting point: 177-178 °C

NMR (DMSO-d_6)δ: 1.74(3H, s), 1.49-2.98(9H, m), 3.60(1H, s), 4.25-4.65(7H, m), 7.14(2H, t, J=8.4Hz), 7.34(2H, m), 7.71(1H, s), 8.26(1H, s), 10.60(1H, s).

Example C-19)

1-Dimethycarbamoylmethyl-5-hydroxy-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthracene-7-carboxylic acid 4-fluoro-benzylamide

melting point: 190 °C

NMR (DMSO-d_6)δ: 1.60(2H, m), 2.76(3H, s), 2.83(3H, s), 2.90-3.59(5H, s), 4.40(2H, m), 4.51(2H, d, 5.7Hz), 4.80(1H, d, d=14.4Hz), 4.98(1H, s), 7.16(2H, t, J=8.4Hz), 7.34(2H, m), 8.54(1H, s), 10.42(1H, s).

Example C-20)

5-Hydroxy-1-(3-methanesulfonylamino-propyl)-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthracene-7-carboxylic acid 4-fluoro-benzylamide

melting point: 176 °C

NMR (DMSO-d_6)δ: 1.54-1.75(4H, m), 2.80(3H, s), 2.30-3.04(8H, m), 4.45(2H, m), 4.52(2H, d, J=5.6Hz), 4.75(1H, d, J=13.2Hz), 6.91(1H, t, J=5.6Hz), 7.16(2H, t, J=8.8Hz), 7.36(2H, m), 8.61(1H, s), 10.41(1H, t, J=5.6Hz), 12.58(1H, s).

Example C-21)


5-Hydroxy-2-methyl-6,10-dioxo-3,4,6,9,9a,10-hexahydro-2H-1-oxa-4a,8a-dizazaanthracene-7-carboxylic acid 4-fluorobenzylamide

NMR (CDCl3)δ: 1.27(3H, d, J=6.0Hz), 1.55-1.78(2H, m), 3.11(1H, td, J=12.9, 3.7Hz), 3.89-4.00(1H, m), 4.16(1H, dd, J=13.8, 3.9Hz), 4.34(1H, dd, J=13.8, 3.9Hz), 4.60(2H, d, J=6.0Hz), 4.71(1H, ddd, J=13.5, 4.8, 1.8Hz), 5.08(1H, t, J=3.9Hz), 6.96-7.04(2H, m), 7.26-7.35(2H, m), 8.32(1H, s), 10.41(1H, br s), 12.41(1H, br s).

Example F-1)

5-Hydroxy-1-isobutyl-4,6-dioxo-2,3,4,6,9,9a-hexahydro-1H-1,3a,8a-triazacyclopenta[b]naphthalene-7-carboxylic acid-4-fluorobenzylamide

[Chemical formula 59]



1) According to the method of synthesizing a compound 17-1, the crude purified product (503mg) of a compound 48 was obtained at a yield of 82% from a compound 16 (600mg).

2) To a solution of a compound 48 (100mg, 0.22mmol), isobutylaldehyde (39μl, 0.432mmol) and acetic acid (25μl, 0.432mmol) in dichloromethane (4ml) was added sodium triacetoxyborohydride (92mg, 0.432mmol) under ice-cooling, and the mixture was stirred at room temperature for 2 hours. Further, isobutylaldehyde (20μl) and sodium triacetoxyborohydride (46mg) were added, and the mixture was stirred for 30 minutes. To the reaction solution was added water, this was extracted with chloroform, and the organic layer was washed with an aqueous saturated sodium bicarbonate solution. After drying, the solvent was distilled off under reduced pressure, and this was purified by silica gel column chromatography. A compound 49 (87mg) was obtained as a colorless crystal at a yield of 78%.
1H-NMR ($CDCl_3$)δ: 0.96(3H, d, J=6.6Hz), 0.97(3H, d, J=6.3Hz), 1.72-1.86(1H, m), 2.25-2.41(2H, m), 2.47-2.58(1H, m), 3.39-3.46(1H, m), 3.69-3.76(2H, m), 3.85-3.93(1H, m), 4.06(1H, dd, J=9.9, 2.7Hz), 4.16-4.22(1H, m), 4.57(1H, dd, J=15.3, 5.1Hz), 4.64(1H, dd, J=14.7, 5.1Hz), 5.20(1H, d, J=9.9Hz), 5.38(1H, d, J=9.9Hz), 6.96-7.05(2H, m), 7.28-7.36(5H, m), 7.58-7.62(2H, m), 8.40(1H, s), 10.44(1H, br s).

3) According to the method of a step 17) of Example B-1, a compound F-1 (43mg) was obtained at a yield of 64% from a compound 49 (81mg).
1H-NMR (DMSO-d_6)δ: 0.90(3H, d, J=6.4Hz), 0.91(3H, d, J=6.0Hz), 1.75-1.84(1H, m), 2.24-2.39(1H, m), 2.39-2.54(2H, m), 3.36-3.43(1H, m), 3.52-3.60(1H, m), 3.67-3.73(1H, m), 3.81-3.88(1H, m), 4.19-4.23(1H, m), 4.52(2H, d, J=6.0Hz), 4.94-4.99(1H, m),

7.12-7.20(2H, m), 7.32-7.38(2H, m), 8.45(1H, s), 10.37(1H, t, J=2.0Hz), 11.74(1H, s).

According to the same manner as that of Example F-1, the following Example compounds F-2 to F-63 were synthesized.

Example F-2)

5-Hydroxy-1-isobutyl-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triazaanthracene-7-carboxylic acid 4-fluorobenzylamide

melting point: 146-148 °C

1H-NMR (DMSO-d_6)δ: 0.63(3H, d, J=6.6Hz), 0.79(3H, d, J=6.6Hz), 1.56-1.66(2H, m), 1.67-1.75(1H, m), 1.94-1.99(1H, m), 2.41-2.54(2H, m), 2.96-3.06(2H, m), 4.41-4.59(5H, m), 4.76-4.81(1H, m), 7.14-7.21(2H, m), 7.33-7.38(2H, m), 8.61(1H, s), 10.40(1H, d, J=5.8Hz), 12.56(1H, s).

Example F-3)

1-Cyclopropylmethyl-5-hydroxy-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthracene-7-carboxylic acid 4-fluoro-benzylamide

melting point: 182-184 °C

NMR (DMSO-d_6)δ: 0.06(2H, m), 0.43(2H, d, 8.4Hz), 0.80(1H, m), 1.66(2H, m), 2.28-3.30(4H, m), 4.40-4.50(4H, m), 4.52(2H, d, 6.0Hz), 4.78(2H, m), 7.15(2H, t, 8.7Hz), 7.34(2H, m), 8.55(1H, s), 10.47(1H, s), 12.55(1H, s).

Example F-4)

1-Cyclopentylmethyl-5-hydroxy-6,1-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthracene-7-carboxylic acid 4-fluoro-benzylamide

melting point: 184-185 °C°C

NMR (DMSO-d_6)δ: 0.88-2.10(1H, m), 2.60(2H, m), 2.95-3.28(2H, m), 4.38-4.53(6H, m), 4.82(1H, m), 7.15(2H, t, 9.0Hz), 7.34(2H, m), 8.57(1H, s), 10.42(1H, s), 12.45(1H, s).

Example F-5)

5-Hydroxy-1-(4-methylsulfanylbenzyl)-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triazaanthracene-7-carboxylic acid 4-fluorobenzylamide

(DMSO-d_6)δ: 1.51-1.56(1H, m), 1.69-1.74(1H, m), 2.42(3H, s), 2.55-2.62(1H, m), 2.80-2.84(1H, m), 3.00-3.08(1H, m), 3.32-3.36(1H, m), 3.93(1H, d, J=13.6Hz), 4.45-4.53(4H, m), 4.58(1H, s), 4.83(1H, d, J=15.2Hz), 7.11-7.19(6H, m), 7.33-7.40(2H, m), 8.34(1H, s), 10.38(1H, t, J=6.0Hz), 12.58(1H, s).

Example F-6)

1-(5-Chloro-1,3-dimethyl-1H-pyrazol-4-ylmethyl)-5-hydroxy-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triazaanthracene-7-carboxylic acid 4-fluorobenzylamide

(DMSO-d_6)δ: 1.56-1.59(2H, m), 1.88(3H, s), 2.37-2.45(1H, m), 2.76-2.80(1H, m), 3.00-3.06(2H, m), 3.64(3H, s), 3.87(1H, d, J=13.2Hz), 4.40-4.55(5H, m), 4.97(1H, d, J=14.4Hz), 7.13-7.19(2H, m), 7.33-7.38(2H, m), 8.56(1H, s), 10.39(1H, t, J=6.0Hz), 12.46(1H, s).

Example F-7)

5-Hydroxy-1-(3-methoxybenzyl)-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triazaanthracene-7-carboxylic acid 4-fluorobenzylamide

(DMSO-d_6)δ: 1.52-1.57(1H, m), 1.70-1.80(1H, m), 2.60-2.68(1H, m), 2.84-2.90(1H, m), 3.01-3.09(1H, m), 3.36(1H, d, J=14.0Hz), 3.61(3H, s), 3.91(1H, d, J=14.0Hz), 4.45-4.52(4H, m), 4.58(1H, s), 4.76(1H, d, J=14.8Hz), 6.68-6.73(2H, m), 6.77(1H, d, J=7.6Hz), 7.13-7.19(3H, m), 7.33-7.38(2H, m), 8.17(1H, s), 10.38(1H, t, J=6.0Hz), 12.57(1H, s).

Example F-8)

5-Hydroxy-1-(4-methanesulfonylbenzyl)-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triazaanthracene-7-carboxylic acid 4-fluorobenzylamide

(DMSO-d_6)δ: 1.54-1.58(1H, m), 1.74-1.80(1H, m), 2.67-1.74(1H, m), 2.83-2.87(1H, m), 3.05-3.12(1H, m), 3.18(3H, s), 3.52(1H, d, J=14.8Hz), 4.09(1H, d, J=14.8Hz), 4.46-4.52(4H, m), 4.67(1H, s), 4.73(1H, d, J=14.8Hz), 7.12-7.18(2H, m), 7.32-7.36(2H, m), 7.46(2H, m), 7.80(2H, d, J=8.0Hz), 8.17(1H, s), 10.37(1H, t, J=5.8Hz), 12.59(1H, s).

Example F-9)

5-Hydroxy-1-(6-methoxypyridin-3-ylmethyl)-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triazaanthracene-7-carboxylic acid 4-fluorobenzylamide

(DMSO-d_6)δ: 1.51-1.56(1H, m), 1.71-1.77(1H, m), 2.58-2.66(1H, m), 2.80-2.86(1H, m), 3.01-3.09(1H, m), 3.38(1H, d, J=13.6Hz), 3.78(3H, s), 3.87(1H, d, J=13.6Hz), 4.45-4.52(4H, m), 4.60(1H, s), 4.82(1H, d, J=13.6Hz), 6.71(1H, d, J=8.6Hz), 7.12-7.19(2H, m), 7.33-7.38(2H, m), 7.49(1H, d, J=8.6Hz), 7.98(1H, s), 8.30(1H, s), 10.37(1H, t, J=6.0Hz), 12.58(1H, s).

Example F-10)

5-Hydroxy-1-isobutyl-3,3-dimethyl-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triazaanthracene-7-carboxylic acid 4-fluorobenzylamide

(DMSO-d_6)δ: 0.64(3H, d, J=6.4Hz), 0.82(3H, d, J=6.8Hz), 0.90(3H, s), 0.91(3H, s), 1.59-1.67(1H, m), 1.92-1.97(1H, m), 2.11-2.15(1H, m), 2.51-2.57(1H, m), 2.67(1H, d, J=12.0Hz), 2.77(1H, d, J=12.8Hz), 4.13(1H, s), 4.21(1H, d, J=12.8Hz), 4.47-4.59(3H, s), 4.80(1H, dd, J=14.4, 2.8Hz), 7.14-7.19(2H, m), 7.34-7.38(2H, m), 8.66(1H, s), 10.41(1H, t, J=6.0Hz), 12.44(1H, s).

Example F-11)

5-Hydroxy-1,3,3-trimethyl-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triazaanthracene-7-carboxylic acid 4-fluorobenzylamide

(DMSO-d_6)δ: 0.89(6H, s), 2.14-2.18(1H, m), 2.24(3H, s), 2.54-2.58(1H, m), 2.74-2.78(1H, s), 3.88(1H, s), 4.21(1H, d, J=13.2Hz), 4.45-4.53(3H, m), 4.72-4.76(1H, m), 7.13-7.19(2H, m), 7.33-7.38(2H, m), 8.64(1H, s), 10.40(1H, t, J=6.0Hz), 12.46(1H, s).

Example F-12)

4-[7-(4-Fluorobenzylcarbamoyl)-5-hydroxy-6,10-dioxy-3,4,6,9,9a,10-hexahydro-2H-1,4a,8a-triazaanthracene-1-yl]butanoic acid ethyl ester

($CDCl_3$)δ: 1.23(3H, t, J=7.1Hz), 1.70-1.79(1H, m), 1.86-2.00(1H, m), 2.17-2.34(2H, m), 2.46-2.57(1H, m), 2.61-2.77(2H, m), 2.85-2.92(1H, m), 3.13-3.18(1H, m), 4.13(2H, q, J=7.1Hz), 4.27-4.34(2H, m), 4.57-4.63(3H, m), 4.66-4.73(1H, m), 6.95-7.03(2H, m), 7.29-7.36(2H, m), 8.36(1H, s), 10.48(1H, t, J=4.8Hz), 12.50(1H, s).

Example F-13)

1-(3-Dimethylcarbamoylpropyl)-5-hydroxy-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triazaanthracene-7-carboxylic acid 4-fluorobenzylamide

($CDCl_3$)δ:1.62-1.82(3H, m), 1.83-2.00(1H, m), 2.10-2.35(2H, m), 2.57-2.65(2H, m), 2.75-2.95(2H, m), 2.92(3H, s), 2.96(3H, s), 3.07-3.14(1H, m), 4.23-4.30(2H, m), 4.60(2H, d, J=6.0Hz), 4.68(1H, dd, J=13.2, 4.5Hz), 5.12(1H, d, J=12.6Hz), 6.95-7.02(2H, m), 7.28-7.35(2H, m), 8.42(1H, s), 1054(1H, t, J=5.4Hz), 12.51(1H, s).

Example F-14)

5-Hydroxy-1-(4-morpholin-4-yl-4-oxobutyl)-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triazaanthracene-7-carboxylic acid 4-fluorobenzylamide

($CDCl_3$)δ: 1.61-1.83(3H, m), 1.84-2.00(1H, m), 2.12-2.23(1H, m), 2.25-2.36(1H, m), 2.56-2.64(2H, m), 2.75-2.95(2H, m), 3.09-3.15(1H, m), 3.37(2H, t, J=4.8Hz), 3.61-3.66(6H, m), 4.26-4.32(2H, m), 4.59(2H, d, J=5.7Hz), 4.68(1H, dd, J=13.2, 4.5Hz), 4.95-5.01(1H, m), 6.95-7.03(2H, m), 7.28-7.35(2H, m), 8.40(1H, s), 10.52(1H, t, J=5.7Hz), 12.51(1H, s).

Example F-15)

5-Hydroxy-1-methyl-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triazaanthracene-7-carboxylic acid 4-fluorobenzylamide

melting point: 252-253°C

(DMSO-d_6)δ: 1.56-1.75(2H, m), 2.22(3H, s), 2.50-2.55(1H, m), 2.90-3.10(2H, m), 4.17(1H, brs), 4.39-4.42(2H, m), 4.52(2H, d, J=6.0Hz), 4.74-4.78(1H, m), 7.13-7.17(2H, m), 7.33-7.37(2H, m), 8.61(1H, s), 10.40(1H, t, J=6.0Hz), 12.54(1H, s).

Example F-16)

5-Hydroxy-6,10-dioxo-1-thiophen-3-ylmethyl-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triazaanthracene-7-carboxylic acid 4-fluorobenzylamide

melting point: 242-243°C

(DMSO-d_6)δ: 1.52-1.73(2H, m), 2.59-2.62(1H, m), 2.87-3.03(2H, m), 3.52(1H, d, J=13.6Hz), 3.90(1H, d, J=14.4Hz), 4.40-4.56(5H, m), 4.83-4.90(1H, m), 6.92(1H, d, J=5.2Hz), 7.13-7.17(2H, m), 7.28-7.37(3H, m), 7.42-7.44(1H, m), 8.46(1H, s), 10.39(1H, t, J=6.0Hz), 12.58(1H, s).

Example F-17)

5-Hydroxy-6,10-dioxo-1-thiazol-2-ylmethyl-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triazaanthracene-7-carboxylic acid 4-fluorobenzylamide

melting point 214-215°C

(DMSO-d_6)δ: 1.54-1.72(2H, m), 2.75-2.81(1H, m), 2.95-3.07(2H, m), 3.80(1H, d, J=16.0Hz), 4.37(1H, d, J=16.4Hz), 4.44-4.51(4H, m), 4.69(1H, brs), 4.89-4.93(1H, m), 7.13-7.17(2H, m), 7.32-7.35(2H, m), 7.55(1H, d, J=3.2Hz), 7.69(1H, d, J=3.2Hz), 8.37(1H, s), 10.36(1H, t, J=6.0Hz), 12.50(1H, s).

Example F-18)

5-Hydroxy-(3-methylsulfanyl-propyl)-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triazaanthracene-7-carboxylic acid 4-fluorobenzylamide

melting point: 162-164°C

(DMSO-d_6)δ: 1.50-1.82(4H, m), 2.27(3H, s), 2.32-2.44(3H, m), 2.60-2.82(2H, m), 3.00-3.14(2H, m), 4.37-4.59(5H, m), 4.75-4.79(1H, m), 7.13-7.17(2H, m), 7.33-7.35(2H, m), 8.60(1H, s), 10.40(1H, t, J=6.0Hz), 12.57(1H, s).

Example F-19)

5-Hydroxy-6,10-dioxo-1-pyridin-4-ylmethyl-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triazaanthracene-7-carboxylic acid 4-fluorobenzylamide

melting point: 180-183°C

(DMSO-d_6)δ: 1.52-1.76(2H, m), 2.62-2.80(2H, m), 3.01-3.07(1H, m), 3.42(1H, d, J=15.2Hz), 4.05(1H, d, J=15.2Hz), 4.49-4.50(4H, m), 4.64(1H, brs), 4.78-4.81(1H, m), 7.12-7.21(4H, m), 7.32-7.36(2H, m), 8.33(1H, s), 8.42(2H, d, J=4.4Hz), 10.39(1H, t, J=6.0Hz), 12.55(1H, s).

Example F-20)

1-Cyclohexylmethyl-5-hydroxy-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triazaanthracene-7-carboxylic acid 4-fluorobenzylamide

melting point: 201-202°C

(DMSO-d_6)δ: 0.56-0.59(1H, m), 0.87-0.84(1H, m), 1.02-1.13(3H, m), 1.23-1.29(1H, m), 1.49-1.70(6H, m), 1.92-1.97(1H, m), 2.52-2.55(1H, m), 2.96-3.03(2H, m), 4.40-4.43(3H, m), 4.52(2H, d, J=6.0Hz), 4.73-4.77(1H, m), 7.12-7.16(2H, m), 7.32-7.36(2H, m), 8.59(1H, s), 10.40(1H, t, J=5.2Hz), 12.58(1H, s).

Example F-21)

5-Hydroxy-6,10-dioxo-1-pyridin-2-ylmethyl-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triazaanthracene-7-carboxylic acid 4-fluorobenzylamide

melting point: 216-219°C

(DMSO-d_6)δ: 1.52-1.76(2H, m), 2.66-2.80(1H, m), 2.90-3.07(2H, m), 3.67(1H, d, J=15.2Hz), 4.01(1H, d, J=13.2Hz), 4.37-4.97(4H, m), 4.62(1H, brs), 4.85-4.88(1H, m), 7.07-7.25(4H, m), 7.33-7.36(2H, m), 7.64-7.68(1H, m), 8.26(1H, s), 8.45(1H, s), 10.36(1H, t, J=6.0Hz), 12.57(1H, s).

Example F-22)

1-(2-Ethyl-butyl)-5-hydroxy-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triazaanthracene-7-carboxylic acid 4-fluorobenzylamide

melting point: 137-140°C
(DMSO-d_6)δ: 0.62(3H, t, J=7.2Hz), 0.77(3H, t, J=7.2Hz), 0.99-1.30(5H, m), 1.57-1.71(2H, m), 1.97-2.02(1H, m), 2.44-2.58(2H, m), 3.02-3.32(2H, m), 4.34-4.57(5H, m), 4.78-4.82(1H, m), 7.13-7.17(2H, m), 7.32-7.36(2H, m), 8.60(1H, s), 10.39(1H, t, J=5.2Hz), 12.54(1H, s).

Example F-23)
5-Hydroxy-1-(2-morpholin-4-ylethyl)-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triazaanthracene-7-carboxylic acid 4-fluorobenzylamide
melting point: 254-256°C
(DMSO-d_6)δ: 1.55-1.68(2H, m), 2.28-2.39(8H, m), 2.59-2.65(1H, m), 2.82-3.09(3H, m), 3.33-3.58(5H, m), 4.34-4.50(3H, m), 4.52(2H, d, J=5.2Hz), 4.79-4.84(1H, m), 7.12-7.17(2H, m), 7.32-7.36(2H, m), 8.52(1H, s), 10.45(1H, t, J=5.2Hz), 12.55(1H, s).

Example F-24)
1-Hydroxy-6-methyl-2,11-dioxo-2,5a,6,7,8,9,10,11-octahydro-5H-4a,6,10a-triaza-cyclohepta[b]naphthalene-3-carboxylic acid 4-fluorobenzylamide
melting point: 255°C
(DMSO-d_6)δ: 1.48-1.55(1H, m), 1.67-1.80(3H, m), 2.29(3H, s), 2.75-2.80(2H, m), 3.23-3.31(1H, m), 4.07-4.09(1H, m), 4.36-4.40(1H, m), 4.45-4.59(3H, m), 4.68-4.69(1H, m), 7.13-7.17(2H, m), 7.30-7.37(2H, m), 8.50(1H, s), 10.42(1H, t, J=6.0Hz), 12.42(1H, s).

Example F-25)
1-Hydroxy-6-isobutyl-2,11-dioxo-2,5a,6,7,8,9,10,11-octahydro-5H-4a,6,10a-triaza-cyclohepta[b]naphthalene-3-carboxylic acid 4-fluorobenzylamide
melting point: 221-223°C
DMSO-d_6)δ: 0.81(3H, d, J=6.8Hz), 0.84(3H, d, J=6.4Hz), 1.45-1.78(5H, m), 2.36-2.54(2H, m), 2.27-2.93(2H, m), 3.17-3.23(1H, m), 4.03-4.06(1H, m), 4.32-4.56(4H, m), 4.82-4.85(1H, m), 7.13-7.17(2H, m), 7.30-7.37(2H, m), 8.48(1H, s), 10.42(1H, t, J=6.0Hz), 12.53(1H, s).

Example F-26)
6-Cyclopropylmethyl-1-hydroxy-2,11-dioxo-2,5a,6,7,8,9,10,11-octahydro-5H-4a,6,10a-triaza-cyclohepta[b]naphthalene-3-carboxylic acid 4-fluorobenzylamide

melting point: 213°C

DMSO-d_6)δ: 0.15-0.26(2H, m), 0.46-0.48(2H, m), 0.86-1.06(1H, m), 1.45-1.75(4H, m), 2.45-2.65(1H, m), 2.68-2.83(1H, m), 2.91-2.98(2H, m), 3.17-3.26(1H, m), 4.08-4.14(1H, m), 4.43-4.45(2H, m), 4.54(2H, d, J=5.6Hz), 4.89-4.91(1H, m), 7.15-7.19(2H, m), 7.35-7.39(2H, m), 8.50(1H, s), 10.47(1H, t, J=6.0Hz), 12.52(1H, s).

Example F-27)

1-Furan-2-ylmethyl-5-hydroxy-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthracene-7-carboxylic acid 4-fluorobenzylamide

melting point: 193-197°C

DMSO-d_6)δ: 1.67(2H, m), 2.61(1H, s), 2.93(2H, m), 3.75(1H, d, J=14.8Hz), 3.84(1H, d, J=14.8Hz), 4.34-4.47(3H, m), 4.52(2H, d, J=6.0Hz), 4.96(1H, d, J=14.8Hz), 6.36(2H, s), 7.16(2H, t, J=8.8Hz), 7.35(2H, m), 7.59(1H, s), 8.97(1H, s), 10.43(1H, s), 12.51(1H, s).

Example F-28)

1-(4-Dimethylamino-benzyl)-5-hydroxy-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthracene-7-carboxylic acid 4-fluorobenzylamide

melting point: 221-223°C

DMSO-d_6)δ: 1.55-1.99(2H, m), 2.87(6H, s), 2.87-3.06(4H, m), 3.80(1H, d, J=14.0Hz), 4.50(5H, m), 4.83(1H, d, J=14.0Hz), 6.58(2H, d, J=9.6Hz), 6.98(2H, d, J=8.8Hz), 7.15(2H, t, J=8.8Hz), 7.35(2H, m), 8.31(1H, s), 10.39(1H, s), 12.58(1H, s).

Example F-29)

5-Hydroxy-6,10-dioxo-1-(4-trifluoromethyl-benzyl)-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthracene-7-carboxylic acid 4-fluorobenzylamide

melting point: 273-277°C

DMSO-d_6)δ: 1.52-1.70(2H, m), 2.63-3.04(3H, m), 3.50(1H, d, J=14.8Hz), 4.10(1H, d, J=14.8Hz), 4.54(5H, m), 4.79(1H, d, J=14.8Hz), 7.14(2H, t, J=8.8Hz), 7.33(2H, m), 7.55(2H, d, J=6.8Hz), 7.61(2H, d, J=8.0Hz), 8.22(1H, s), 10.40(1H, s), 12.56(1H, s).

Example F-30)

5-Hydroxy-6,10-dioxo-1-pyridin-3-ylmethyl-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthracene-7-carboxylic acid 4-fluorobenzylamide

melting point: 210-212°C

DMSO-d_6)δ: 1.51-1.76(2H, m), 2.63(1H, t, J=12.8Hz), 2.80(1H, d, J=12.0Hz), 3.07(1H, t,

J=12.8Hz), 3.44(1H, d, J=13.2Hz), 4.00(1H, d, 14.0Hz), 4.47(4H, m), 4.62(1H, s), 4.84(1H, d, J=14.0Hz), 7.16(2H, t, J=8.8Hz), 7.33(2H, m), 7.58(1H, d, J=7.6Hz), 8.30(1H, s), 8.45(2H, s), 10.41(1H, s), 12.57(1H, s).

Example F-31)

1-(2-Chloro-6-fluoro-benzyl)-5-hydroxy-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthracene-7-carboxylic acid 4-fluorobenzylamide

melting point: 213-215°C

DMSO-d_6)δ:1.58(2H, 2H), 2.55-3.09(3H, m), 3.45(1H, d, J=12.4Hz), 4.16(1H, d, J=12.4Hz), 4.40-4.58(4H, m), 5.12(1H, d, J=14.4Hz), 7.15-7.38(7H, m), 8.66(1H, s), 10.41(1H, t, J=6.4Hz), 12.46(1H, s).

Example F-32)

5-Hydroxy-1-(4-methoxy-benzyl)-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthracene-7-carboxylic acid 4-fluorobenzylamide

melting point: 191-193°C

NMR (DMSO-d_6)δ:1.50-1.77(2H, m), 2.58-3.06(3H, m), 3.68(3H, s), 3.88(1H, d, J=13.6Hz), 4.41-4.55(4H, m), 4.80(2H, d, J=14.4Hz), 6.80(2H, d, J=8.8Hz), 7.09(2H, d, J=8.4Hz), 7.15(2H, t, J=8.8Hz), 7.35(2H, m), 8.28(1H, s), 10.48(1H, s), 12.58(1H, s).

Example F-33)

1-(3,5-Bis-trifluoromethyl-benzyl)-5-hydroxy-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-octahydro-1,4a,8a-triaza-anthracene-7-carboxylic acid 4-fluorobenzylamide

melting point: 275-277°C

NMR (DMSO-d_6)δ:1.58-1.88(2H, m), 2.51-3.14(3H, m), 3.33-4.10(3H, m), 4.51(2H, m), 4.73(1H, m), 7.15(2H, m), 7.34(2H, m), 7.82-7.93(4H, m), 10.31(1H, s), 12.57(1H, s).

Example F-34)

1-(4-Diethylamino-benzyl)-5-hydroxy-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthracene-7-carboxylic acid 4-fluorobenzylamide

melting point: 182°C

NMR (DMSO-d_6)δ: 1.04(6H, t, J=6.8Hz), 1.50-1.69(2H, m), 2.55-3.05(3H, m), 3.26(4H, q, J=7.2Hz), 3.80(1H, d, J=13.6Hz), 4.44-4.57(4H, m), 4.91(1H, d, J=12.4Hz), 6.52(2H, d, J=8.8Hz), 6.94(2H, d, J=8.4Hz), 7.15(2H, t, J=8.4Hz), 7.35(2H, m), 8.46(1H, s), 10.41(1H, s), 12.60(1H, s).

Example F-35)

5-Hydroxy-1-((E)-2-methyl-but-2-enyl)-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthracene-7-carboxylic acid 4-fluoro-benzylamide

melting point: 175-177°C

NMR (DMSO-d_6)δ: 1.35(3H, s), 1.51(3H, d, J=6.0Hz), 1.52-1.69(3H, m), 2.60-3.15(3H, m), 4.31-4.52(5H, m), 4.67-4.76(1H, m), 5.30-5.40(1H, m), 7.15(2H, t, J=8.4Hz), 7.28-43(2H, m), 8.46(1H, s), 10.39(1H, brs), 12.60(1H, s).

Example F-36)

1-(3-Dimethylamino-2-methyl-propyl)-5-hydroxy-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthracene-7-carboxylic acid 4-fluoro-benzylamide

NMR (DMSO-d_6)δ: 0.63-0.68(2H, m), 1.57-1.82(3H, m), 2.11-2.49(10H, m), 2.98-3.11(2H, m), 4.41-4.54(5H, m), 4.73-4.80(1H, m), 7.14-7.18(2H, m), 7.31-7.38(2H, m), 8.58(1H, s), 10.40(1H, s), 12.57(1H, s).

Example F-37)

1-(3,3-Dimethyl-butyl)-5-hydroxy-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthracene-7-carboxylic acid 4-fluoro-benzylamide

melting point: 175-177°C

NMR (DMSO-d_6)δ: 1.19-1.36(2H, m), 1.57-1.70(2H, m), 2.23-2.30(1H, m), 2.51-2.69(2H, m), 2.97-3.04(2H, m), 4.42-4.54(5H, m), 4.78(1H, d, J=14.0Hz), 7.13-7.17(2H, m), 7.33-7.36(2H, m), 8.63(1H, s), 10.39(1H, t, J=6.0Hz), 12.56(1H, s).

Example F-38)

1-Ethyl-5-hydroxy-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthracene-7-carboxylic acid 4-fluoro-benzylamide

melting point: 221°C

NMR (DMSO-d_6)δ: 0.94(3H, t, J=6.8Hz), 1.56-1.71(2H, m), 2.45-2.50(1H, m), 2.59-2.76(2H, m), 2.96-3.03(2H, m), 4.40-4.44(3H, m), 4.52(2H, d, J=6.0Hz), 4.77-4.82(1H, m), 7.14-7.18(2H, m), 7.34-7.38(2H, m), 8.62(1H, s), 10.41(1H, t, J=6.0Hz), 12.59(1H, s).

Example F-39)

5-Hydroxy-6,10-dioxo-1-(2-oxo-propyl)-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anth

racene-7-carboxylic acid 4-fluoro-benzylamide
melting point: 244-246°C
NMR (DMSO-d_6)δ: 1.54-1.61(1H, m), 1.67-1.76(1H, m), 2.22(3H, s), 2.50-2.56(1H, m), 2.91-3.02(2H, m), 4.18(1H, s), 4.38-4.45(2H, m), 4.52(2H, d, J=6.0Hz), 4.76(1H, d, J=14.4Hz), 7.13-7.18(2H, m), 7.34-7.37(2H, m), 8.61(1H, s), 10.40(1H, t, J=6.0Hz), 12.54(1H, s).

Example F-40)
5-Hydroxy-6,10-dioxo-1-(4,4,4-trifluoro-butyl)-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthracene-7-carboxylic acid 4-fluoro-benzylamide
melting point: 220°C
NMR (DMSO-d_6)δ: 1.53-1.62(2H, m), 1.67-1.75(1H, m), 2.07-2.18(2H, m), 2.40-2.47(1H, m), 2.64-2.78(2H, m), 2.96-3.04(2H, m), 4.42-4.49(2H, m), 4.53(2H, d, J=5.2Hz), 4.74(1H, d, J=12.8Hz), 7.13-7.17(2H, m), 7.33-7.37(2H, m), 8.61(1H, s), 10.40(1H, t, J=6.0Hz), 12.57(1H, s).

Example F-41)
5-Hydroxy-1-(3-methyl-butyl)-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthracene-7-carboxylic acid 4-fluoro-benzylamide
melting point: 151°C
NMR (DMSO-d_6)δ: 0.78(6H, dd, J=7.6, 16.2Hz), 1.21-1.28(2H, m), 1.41-1.48(1H, m), 1.56-1.71(2H, m), 2.22-2.31(1H, m), 2.51-2.59(1H, m), 2.66-2.73(1H, m), 2.96-3.05(2H, m), 4.41-4.55(5H, m), 4.80(1H, d, J=13.2Hz), 7.13-7.18(2H, m), 7.33-7.37(2H, m), 8.64(1H, s), 10.40(1H, t, J=6.0Hz), 12.57(1H, s).

Example F-42)
5-Hydroxy-1-isobutyl-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthracene-7-carboxylic acid 3-chloro-2-fluoro-benzylamide
melting point: 180-182°C
NMR (DMSO-d_6)δ: 0.62(3H, d, J=6.0Hz), 0.78(3H, d, J=6.4Hz), 1.55-1.69(3H, m), 1.93-1.99(1H, m), 2.97-3.08(2H, m), 4.39-4.46(3H, m), 4.59-4.64(2H, m), 4.75-4.81(1H, m), 7.16-7.23(1H, m), 7.27-7.34(1H, m), 7.47-7.53(1H, m), 8.59(1H, s), 10.44(1H, s), 12.57(1H, s).

Example F-43)

1-Cyclopropylmethyl-5-hydroxy-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthracene-7-carboxylic acid 3-chloro-2-fluoro-benzylamide
melting point: 189-192°C
NMR (DMSO-d_6)δ: 0.00-0.10(2H, m), 0.35-0.41(2H, m), 0.70-0.77(1H, m), 1.57-1.69(2H, m), 2.52-2.65(1H, m), 2.67-2.85(1H, m), 2.91-2.99(1H, m), 4.30-4.41(2H, m), 4.48-4.52(2H, m), 4.71-4.80(1H, m), 7.06-7.10(1H, m), 7.18-7.22(1H, m), 7.36-7.40(1H, m), 8.52(1H, s), 10.30(1H, s), 12.26(1H, s).

Example F-44)
1-Furan-2-ylmethyl-5-hydroxy-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthracene-7-carboxylic acid 3-chloro-2-fluoro-benzylamide
melting point: 190-192°C
NMR (DMSO-d_6)δ: 1.56-1.68(2H, m), 2.54-2.63(1H, m), 2.89-2.99(2H, m), 3.80(2H, dd, J=18.4, 33.2Hz), 4.37-4.51(3H, m), 4.62(2H, d, J=6.0Hz), 4.97(1H, d, J=15.2Hz), 6.39(2H, s), 7.18-7.22(1H, m), 7.31-7.34(1H, m), 7.48-7.51(1H, m), 7.58(1H, s), 8.64(1H, s), 10.45(1H, t, J=6.0Hz), 12.55(1H, s).

Example F-45)
5-Hydroxy-6,10-dioxo-1-thiazol-2-ylmethyl-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthracene-7-carboxylic acid 3-chloro-2-fluoro-benzylamide
melting point: 217-219°C
NMR (DMSO-d_6)δ: 1.59-1.74(2H, m), 2.76-2.83(1H, m), 2.97-3.08(2H, m), 3.90(1H, d, J=16.0Hz), 4.36(1H, d, J=16.0Hz), 4.45-4.69(5H, m), 4.89(1H, d, J=14.8Hz), 7.18-7.22(1H, m), 7.28-7.31(1H, m), 7.47-7.53(1H, m), 7.54(1H, d, J=3.2Hz), 7.68(1H, d, J=3.2Hz), 8.34(1H, s), 10.40(1H, d, J=6.0Hz), 12.52(1H, s).

Example F-46)
5-Hydroxy-6,10-dioxo-1-pyridin-2-ylmethyl-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthracene-7-carboxylic acid 3-chloro-2-fluoro-benzylamide
melting point: 190-193°C
NMR (DMSO-d_6)δ: 1.54-1.61(1H, m), 1.69-1.75(1H, m), 2.66-2.74(1H, m), 2.91-3.08(2H, m), 3.68(1H, d, J=14.4Hz), 4.02(1H, d, J=14.8Hz), 4.40-4.67(5H, m), 4.85(1H, d, J=12.4Hz), 7.16-7.35(3H, m), 7.46-7.52(1H, m), 7.61-7.69(1H, m), 8.20(1H, s), 8.43-8.47(1H, m), 10.41(1H, d, J=6.0Hz), 12.58(1H, s).

Example F-47)

5-Hydroxy-1-isobutyl-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthracene-7-carboxylic acid 2,4-difluoro-benzylamide

melting point: 194°C

NMR (DMSO-d_6)δ: 0.62(3H, d, J=6.4Hz), 0.78(3H, d, J=6.4Hz), 1.55-1.69(3H, m), 1.93-1.99(1H, m), 2.97-3.08(2H, m), 4.39-4.46(3H, m), 4.50-4.59(2H, m), 4.77(1H, d, J=14.4Hz), 7.03-7.09(1H, m), 7.20-7.28(1H, m), 7.36-7.43(1H, m), 8.59(1H, s), 10.39(1H, s), 12.56(1H, s).

Example F-48)

1-Cyclopropylmethyl-5-hydroxy-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthracene-7-carboxylic acid 2,4-difluoro-benzylamide

melting point: 169-171°C

NMR (DMSO-d_6)δ: 0.00-0.10(2H, m), 0.42-0.44(2H, m), 0.77-0.81(1H, m), 1.59-1.74(2H, m), 2.27-2.32(1H, m), 2.62-2.72(1H, m), 3.05-3.12(1H, m), 4.30-4.58(5H, m), 4.69(1H, d, J=14.8Hz), 7.03-7.11(1H, m), 7.22-7.26(1H, m), 7.37-7.40(1H, m), 8.62(1H, s), 10.40(1H, t, J=6.0Hz), 12.57(1H, s).

Example F-49)

1-Furan-2-ylmethyl-5-hydroxy-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthracene-7-carboxylic acid 2,4-difluoro-benzylamide

melting point: 186-188°C

NMR (DMSO-d_6)δ: 1.55-1.68(2H, m), 2.55-2.64(1H, m), 2.88-2.99(2H, m), 3.80(2H, dd, J=15.6, 34.8Hz), 4.36-4.56(5H, m), 4.97(1H, d, J=16.0Hz), 6.39(2H, s), 7.05-7.08(1H, m), 7.21-7.26(1H, m), 7.37-7.44(1H, m), 7.58(1H, s), 8.64(1H, s), 10.38(1H, t, J=5.6Hz), 12.53(1H, s).

Example F-50)

5-Hydroxy-6,10-dioxo-1-thiazol-2-ylmethyl-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthracene-7-carboxylic acid 2,4-difluoro-benzylamide

melting point: 168-170°C

NMR (DMSO-d_6)δ: 1.59-1.74(2H, m), 2.76-2.83(1H, m), 2.97-3.08(2H, m), 3.89(1H, d, J=16.4Hz), 4.36(1H, d, J=16.0Hz), 4.44-4.55(4H, m), 4.69(1H, s), 4.89(1H, d, J=14.8Hz), 7.03-7.09(1H, m), 7.20-7.27(1H, m), 7.34-7.41(1H, m), 7.54(1H, d, J=3.2Hz), 7.68(1H, d, J=3.2Hz), 8.34(1H, s), 10.35(1H, d, J=6.0Hz), 12.50(1H, s).

Example F-51)

5-Hydroxy-6,10-dioxo-1-pyridin-2-ylmethyl-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthracene-7-carboxylic acid 2,4-difluoro-benzylamide

melting point: 200-203°C

NMR (DMSO-d_6)δ: 1.54-1.61(1H, m), 1.69-1.78(1H, m), 2.71-2.79(1H, m), 2.91-3.09 (2H, m), 3.72(1H, d, J=14.4Hz), 4.07(1H, d, J=14.4Hz), 4.44-4.54(4H, m), 4.70(1H, s), 4.82(1H, d, J=14.4Hz), 7.04-7.10(1H, m), 7.21-7.42(4H, m), 7.74-7.80(1H, m), 8.17(1H, s), 8.47-8.49(1H, m), 10.35(1H, d, J=6.0Hz), 12.57(1H, s).

Example F-52)

1-Hydroxy-6-methyl-2,11-dioxo-2,5a,6,7,8,9,10,11-octahydro-5H-4a,6,10a-triaza-cyclohepta[b]naphthalene-3-carboxylic acid 3-chloro-2-fluoro-benzylamide

melting point: 230-231°C

NMR (DMSO-d_6)δ: 1.47-1.53(1H, m), 1.62-1.78(3H, m), 2.29(3H, s), 2.77-2.81 (2H, m), 4.05-4.10(1H, m), 4.35-4.40(1H, m), 4.54-4.64(3H, m), 4.70(1H, s), 7.18-7.22(1H, m), 7.30-7.34(1H, m), 7.47-7.52(1H, m), 8.49(1H, s), 10.47(1H, d, J=6.0Hz), 12.44(1H, s).

Example F-53)

1-Hydroxy-6-isobutyl-2,11-dioxo-2,5a,6,7,8,9,10,11-octahydro-5H-4a,6,10a-triaza-cyclohepta[b]naphthalene-3-carboxylic acid 3-chloro-2-fluoro-benzylamide

melting point: 215-216°C

NMR (DMSO-d_6)δ: 0.83(6H, dd, J=6.8, 13.6Hz), 1.45-1.80(5H, m), 2.36-2.41(1H, m), 2.77-2.93(2H, m), 3.17-3.24(1H, m), 4.02-4.09(1H, m), 4.32-4.40(2H, m), 4.61(2H, d, J=5.6Hz), 4.82-4.84(1H, m), 7.18-7.22(1H, m), 7.30-7.33(1H, m), 7.48-7.51(1H, m), 8.47(1H, s), 10.48(1H, t, J=6.0Hz), 12.55(1H, s).

Example F-54)

6-Cyclopropylmethyl-1-hydroxy-2,11-dioxo-2,5a,6,7,8,9,10,11-octahydro-5H-4a,6,10a-triaza-cyclohepta[b]naphthalene-3-carboxylic acid 3-chloro-2-fluoro-benzylamide

melting point: 212°C

NMR (DMSO-d_6)δ: 0.00-0.10(2H, m), 0.40-45(2H, m), 0.80-0.87(1H, m), 1.45-1.77(3H, m), 2.64-2.69(1H, m), 2.85-2.95(2H, m), 3.13-3.20(1H, m), 4.03-4.09(1H, m), 4.36-4.40(2H, m), 4.59(2H, d, J=5.6Hz), 4.84-4.86(1H, m), 7.16-7.20(1H, m), 7.28-7.32(1H, m), 7.46-7.50(1H, m), 8.45(1H, s), 10.46(1H, t, J=6.0Hz), 12.50(1H, s).

Example F-55)

6-Furan-2-ylmethyl-1-hydroxy-2,11-dioxo-2,5a,6,7,8,9,10,11-octahydro-5H-4a,6,10a-triaza-cyclohepta[b]naphthalene-3-carboxylic acid 3-chloro-2-fluoro-benzylamide

Melting point: 189-190°C

NMR (DMSO-d_6)δ: 1.48-1.63(3H, m), 1.70-1.77(1H, m), 2.79-2.83(2H, m), 3.90(2H, dd, J=14.8, 39.6Hz), 4.05-4.11(1H, m), 4.40-4.51(2H, m), 4.61(2H, d, J=5.6Hz), 4.89-4.91(1H, m), 6.30-6.33(1H, m), 6.38-6.40(1H, m), 7.18-7.22(1H, m), 7.30-7.34(1H, m), 7.48-7.53(1H, m), 7.57(1H, s), 8.45(1H, s), 10.45(1H, t, J=6.0Hz), 12.44(1H, s).

Example F-56)

1-Hydroxy-6-methyl-2,11-dioxo-2,5a,6,7,8,9,10,11-octahydro-5H-4a,6,10a-triaza-cycloheptа[b]naphthalene-3-carboxylic acid 2,4-difluoro-benzylamide

melting point: 241°C

NMR (DMSO-d_6)δ: 1.47-1.53(1H, m), 1.62-1.78(3H, m), 2.29(3H, s), 2.77-2.81 (2H, m), 4.05-4.10(1H, m), 4.35-4.40(1H, m), 4.53-4.61(3H, m), 4.69(1H, s), 7.03-7.08(1H, m), 7.20-7.27(1H, m), 7.37-7.43(1H, m), 8.49(1H, s), 10.42(1H, d, J=6.0Hz), 12.43(1H, s).

Example F-57)

1-Hydroxy-6-isobutyl-2,11-dioxo-2,5a,6,7,8,9,10,11-octahydro-5H-4a,6,10a-triaza-cyclohepta[b]naphthalene-3-carboxylic acid 2,4-difluoro-benzylamide

melting point: 203°C

NMR (DMSO-d_6)δ: 0.82(6H, dd, J=6.4, 13.2Hz), 1.45-1.80(5H, m), 2.36-2.42(1H, m), 2.77-2.93(2H, m), 3.15-3.23(1H, m), 4.02-4.08(1H, m), 4.32-4.41(2H, m), 4.54(2H, d, J=5.6Hz), 4.82-4.84(1H, m), 7.02-7.09(1H, m), 7.20-7.27(1H, m), 7.36-7.43(1H, m), 8.47(1H, s), 10.41(1H, t, J=6.0Hz), 12.54(1H, s).

Example F-58)

6-Cyclopropylmethyl-1-hydroxy-2,11-dioxo-2,5a,6,7,8,9,10,11-octahydro-5H-4a,6,10a-triaza-cyclohepta[b]naphthalene-3-carboxylic acid 2,4-difluoro-benzylamide

melting point: 182-183°C

NMR (DMSO-d_6)δ: 0.00-0.10(2H, m), 0.40-45(2H, m), 0.80-0.87(1H, m), 1.43-1.77(3H, m), 2.60-2.69(1H, m), 2.85-2.95(2H, m), 3.11-3.19(1H, m), 4.00-4.06(1H, m), 4.36-4.40(2H, m), 4.51(2H, d, J=5.6Hz), 4.83-4.87(1H, m), 7.00-7.07(1H, m), 7.16-7.23(1H, m), 7.34-7.38(1H, m), 8.44(1H, s), 10.39(1H, t, J=6.0Hz), 12.47(1H, s).

Example F-59)

6-Furan-2-ylmethyl-1-hydroxy-2,11-dioxo-2,5a,6,7,8,9,10,11-octahydro-5H-4a,6,10a-triaza-cyclohepta[b]naphthalene-3-carboxylic acid 2,4-difluoro-benzylamide

melting point: 171-173°C

NMR (DMSO-d_6)δ: 1.47-1.64(3H, m), 1.70-1.77(1H, m), 2.79-2.83(2H, m), 3.90(2H, dd, J=15.6, 39.6Hz), 4.05-4.11(1H, m), 4.41-4.57(4H, m), 4.90-4.92(1H, m), 6.30-6.33(1H, m), 6.38-6.40(1H, m), 7.03-7.09(1H, m), 7.20-7.27(1H, m), 7.37-7.45(1H, m), 7.57(1H, s), 8.44(1H, s), 10.41(1H, t, J=6.0Hz), 12.43(1H, s).

Example F-60)

5-Hydroxy-6,10-dioxo-3,4,6,9,9a,10-hexahydro-2H-1-oxa-4a,8a-diaza-anthracene-7-carboxylic acid 3-chloro-2-fluoro-benzylamide

melting point: 276°C

NMR (DMSO-d_6)δ: 1.60-1.68(1H, m), 1.77-1.84(1H, m), 3.85-3.93 (1H, m), 4.03-4.07(1H, m), 4.43-4.62(5H, m), 5.28(1H, s), 7.17-7.22(1H, m), 7.29-7.34(1H, m), 7.47-7.52(1H, m), 8.49(1H, s), 10.41(1H, d, J=6.0Hz), 12.48(1H, s).

Example F-61)

5-Hydroxy-6,10-dioxo-3,4,6,9,9a,10-hexahydro-2H-1-oxa-4a,8a-diaza-anthracene-7-carboxylic acid 2,4-difluoro-benzylamide

melting point: 258°C

NMR (DMSO-d_6)δ: 1.60-1.69(1H, m), 1.77-1.85(1H, m), 3.86-3.92 (1H, m), 4.04-4.08(1H, m), 4.43-4.55(5H, m), 5.28(1H, s), 7.03-7.09(1H, m), 7.21-7.27(1H, m), 7.36-7.43(1H, m), 8.50(1H, s), 10.35(1H, d, J=6.0Hz), 12.47(1H, s).

Example F-62)

5-Hydroxy-1-(2-methoxy-ethyl)-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthracene-7-carboxylic acid 3-chloro-2-fluoro-benzylamide

melting point: 193°C

NMR (DMSO-d_6)δ: 1.53-1.73(2H, m), 2.51-2.58(1H, m), 2.71-2.78(1H, m), 2.81-2.87 (1H, m), 2.95-3.08(2H, m), 3.17(3H, s), 4.40-4.52(3H, m), 4.62(1H, d, J=5.6Hz), 4.78(1H, d, J=14.4Hz), 7.18-7.22(1H, m), 7.30-7.34(1H, m), 7.47-7.52(1H, m), 8.55(1H, s), 10.45(1H, d, J=6.0Hz), 12.59(1H, s).

Example F-63)

5-Hydroxy-1-(2-methoxy-ethyl)-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthracene-7-carboxylic acid 2,4-difluoro-benzylamide

melting point: 166-168°C

NMR (DMSO-d_6)δ: 1.55-1.72(2H, m), 2.51-2.58(1H, m), 2.70-2.77(1H, m), 2.80-2.87(1H, m), 2.97-3.07(2H, m), 3.18(3H, s), 4.39-4.52(3H, m), 4.54(1H, d, J=5.2Hz), 4.78(1H, d, J=13.6Hz), 7.03-7.09(1H, m), 7.20-7.27(1H, m), 7.37-7.43(1H, m), 8.55(1H, s), 10.40(1H, d, J=6.0Hz), 12.58(1H, s).

Example F-64)


5-Hydroxy-1-(1H-imidazol-4-ylmethyl)-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triazaanthracene-7-carboxylic acid 4-fluorobenzylamide

(DMSO-d_6)δ: 1.55-1.59(1H, m), 1.64-1.70(1H, m), 2.58-2.66(1H, m), 2.87-2.95(2H, m), 3.67(1H, d, J=15.2Hz), 3.73(1H, d, J=15.2Hz), 4.34(1H, s), 4.38-4.43(1H, m), 4.47-4.54(3H, m), 5.05(1H, d, J=14.0Hz), 7.00(1H, s), 7.13-7.19(2H, m), 7.33-7.38(1H, m), 7.59(1H, s), 8.55(1H, s), 10.41(1H, t, J=5.6Hz), 11.95(1H, br s), 12.59(1H, s).

Example H-1)

1-Acetyl-5-hydroxy-4,6-dioxo-2,3,4,6,9,9a-hexahydro-1H-1,3a,8a-triaza-cyclopenta[b]naphthalene-7-carboxylic acid 4-fluoro-benzylamide

[Chemical formula 61]



1) To a solution of a compound 48 (120mg, 0.26 mmol) in methylene chloride (1.2 ml) were added triethylamine (43 μl, 0.31 mmol), acetic anhydride (29 μl, 0.31 mmol), and 4-dimethylaminopyridine (cat.) at room temperature, and the mixture was stirred for 30 minutes. Further, triethylamine (18 μl, 0.13 mmol) and acetic anhydride (12 μl, 0.13 mmol) were added, and the mixture was stirred for 4 hours. 2N hydrochloric


acid was added, this was extracted with chloroform, and the organic layer was washed with water, dried with sodium sulfate, and concentrated under reduced pressure. Diisopropyl ether was added to crystallize the material, which was filtered to obtain 53 (112 mg) as a pale orange crystal at a yield of 86 %.

2) An Example compound H-1 (71 mg) was obtained at a yield of 82 % from a compound 53 (106 mg), according to the method of Example B-1 17).
melting point 290°C
NMR (DMSO-d_6)δ: 2.08(3H, s), 3.44-4.21(5H, m), 4.51(2H, d, 5.7Hz), 4.93(1H, m), 5.46-5.62(1H, m), 7.15(2H, t, 9.0Hz), 7.34(2H, m), 8.49(1H, s), 10.40(1H, t, 5.7Hz), 11.48(1H, s).

An Example compound H-2 was synthesized according to the same manner as that of Example H-1.

Example H-2)
1-Acetyl-5-hydroxy-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthracene-7-carboxylic acid 4-fluoro-benzylamide
melting point: 290°C
NMR (DMSO-d_6)δ: 1.95(2H, m), 2.14(3H, s), 2.85(2H, m), 4.45(4H, m), 4.51(2H, d, 5.7Hz), 5.99(1H, s), 7.15(2H, t, 9.0Hz), 7.34(2H, m), 8.37(1H, s), 10.46(1H, s), 12.28(1H, s).

Example I-1)
5-Hydroxy-1-methanesulfonyl-4,6-dioxo-2,3,4,6,9,9a-hexahydro-1H-1,3a,8a-triaza-cyclopenta[b]naphthalene-7-carboxylic acid 4-fluoro-benzylamide
[Chemical formula 62]




1) To a solution of a compound 48 (140 mg, 0.30 mmol) in pyridine (1.4 ml) were added methanesulfonyl chloride (28 µl, 0.36 mmol), and 4-dimethylaminopyridine (cat.) at room temperature, and the mixture was stirred for 3 hours. After 2N hydrochloric acid was added, this was extracted with ethyl acetate, and the organic layer was washed with water, dried with sodium sulfate, and concentrated under reduced pressure. Diisopropylether was added to crystallize the material, which was filtered to obtain 54 (127 mg) as a pale orange crystal at a yield of 78 %.

2) According to the method of Example B-1 17), an Example compound I-1 (21 mg) was obtained at a yield of 21 % from a compound 54 (123 mg).
melting point: 260°C
NMR (DMSO-d_6)δ: 3.16(3H, s), 3.30-4.15(5H, m), 4.45(2H, d, 5.7Hz), 4.27(2H, m), 5.36(1H, m), 7.14(2H, t, 8.7Hz), 7.33(2H, m), 8.22(1H, s), 10.53(1H, s).

According to the same manner as that of Example I-1, an Example compound I-2 was synthesized.

Example I-2)
5-Hydroxy-1-methanesulfonyl-6,10-dioxo-1,2,3,4,6,9,9a,10-octahydro-1,4a,8a-triaza-anthracene-7-carboxylic acid 4-fluoro-benzylamide
melting point: 257-259°C
NMR (DMSO-d_6)δ: 1.80-1.96(2H, m), 3.02-3.58(2H, m), 3.16(3H, s), 4.76(2H, m), 5.56(1H, s), 7.16(2H, t, 9.0Hz), 7.35(2H, m), 8.36(1H, s), 10.39(1H, s).

Example L-1)
5,9-Dihydroxy-6,10-dioxo-3,4,6,9,9a,10-hexahydro-1H-2-oxa-4a,8a-diaza-anthracene-7-carboxylic acid 4-fluoro-benzylamide
[Chemical formula 65]




1) According to the method of synthesizing a compound 66, a compound 62 (278 mg, 57%) was obtained from a compound 13 (357 mg).

2) According to the method of synthesizing a compound 57, a compound 63 (202 mg, 79 %) was obtained from a compound 62 (278 mg).

3) To a solution of a compound 63 (200 mg, 0.403 mmol) in chloroform (2 ml) were added dimethyl sulfoxide (286 μl, 4.03 mmol), and triethylamine (337 μl, 2.42 mmol), the mixture was stirred for 10 minutes under ice-cooling, a sulfur trioxide-pyridine complex (321 mg, 2.02 mmol) was added, and the mixture was stirred at room temperature for 2 hours. To the reaction solution was added water (3 ml), and chloroform was distilled off under reduced pressure, followed by extraction with ethyl acetate. The organic layer was washed with water, dried with anhydrous sodium sulfate, and the solvent was distilled off under reduced pressure. The crystalline residue was washed with ethyl acetate to obtain a compound 64 (60 mg) at a yield of 30 %.

4) Using a compound 64, and according to the method of synthesizing Example A-1, an Example compound L-1 was synthesized.
NMR (DMSO-d_6)δ: 2.98-3.10(1H, m), 3.38-3.60(2H, m), 3.80-4.20(5H, m), 4.40-4.55(2H, m), 5.48(1H, brs), 5.85(1H, s), 7.15(2H, t, J=8.4Hz), 7.33-7.37(2H, m), 8.45(1H, s), 8.60(1H, s), 10.27-10.42(1H, m), 12.61(1H, brs).

Example M-1)
1-Hydroxy-2,10-dioxo-2,4b,5,6,7,8,9,10-octahydro-4a,9a-diaza-benzo[a]azulene-3-carboxylic acid 4-fluoro-benzylamide
[Chemical formula 66]




1) According to the method of synthesizing a compound 21, a compound 65 (207 mg) was obtained at a yield of 24 % from a compound 13 (250 mg).

2) According to the method of synthesizing a compound 64, a compound 66 (313 mg, 67 %) was obtained from a compound 65 (470 mg).

3) After trifluoroacetic acid (10 ml) was added to a compound 66 (100 mg, 0.020 mmol), the mixture was stirred at 75°C for 4 hours. The solvent was distilled off under reduced pressure, and this was diluted with chloroform, and added to ice water. This was washed with an aqueous saturated sodium bicarbonate solution, a 10 % aqueous citric acid solution, and water, and dried with anhydrous sodium sulfate, and the solvent was distilled off under reduced pressure. The residue was subjected to silica gel column chromatography, and fractions eluted with chloroform-methanol were concentrated under reduced pressure, and recrystallized with ethyl acetate-diisopropyl ether to obtain an Example compound M-1 (23 mg, 16 %).
melting point 281-283°C
NMR (DMSO-d_6)δ: 1.43-1.52(2H, m), 1.62-1.83(3H, m), 2.04-2.18(1H, m), 2.23-2.35(1H, m), 4.08-4.16(1H, m), 4.48-4.53(2H, m), 5.58-5.61(1H, m), 7.11-7.20(2H, m), 7.30-7.38(2H, m), 8.29(1H, s), 10.30-10.36(1H, m), 12.78(1H, brs).

Example X-1)
(R)-6-Hydroxy-5,7-dioxo-2,3,5,7,11,11a-hexahydro-1H-pyrido[1,2-a]pyrrolo[1,2-d]pyrazine-8-carboxylic acid 4-fluoro-benzylamide
[Chemical formula 67]



1) Selenium dioxide (666mg, 6.0mmol) was added to the solution of compound 2 (216mg, 1.0mmol) in bromobenzene (2ml). Then the mixture was heated up to 160°C, and stirred for 16h. After celite filtration the solvent was evaporate. The precipitate was purified by silicagel column chromatography, and fractions eluting with n-hexan/EtOAc were concentrated under reduced pressure to obtain compound 100 (164mg, 71%) as a yellow oil.

1H-NMR ($CDCl_3$)δ: 5.52(1H, s), 6.50(1H, d, J=6.0Hz), 7.36(5H, m), 7.74(1H, d, J=6.3Hz), 9.88(1H, s).

2) Sulfamic acid (1.50g, 15.4mmol) and $NaClO_2$ (1.05g, 11.6mmol) was added to the solution of compound 100 (2.54g, 11.0mmol) in acetone (20ml) and water (30ml). Then the mixture was stirred for 3h. The solvent was evaporated under reduced pressure to obtain compound 101 (2.18mg, 80%) as a white solid.

1H-NMR (DMSO-d_6)δ: 5.11(2H, s), 6.55(1H, d, J=5.4Hz), 7.32-7.46(5H, m), 8.21(1H, d, J=5.7Hz).

3) (R)-2-N-BOC-aminomethyl pyrrolidine (391mg, 1.95mmol) was added to the solution of compound 101 (400mg, 1.62mmol),

1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (373mg, 1.95mmol), and 1-hydroxybenzotriazole (219mg, 1.62mmol) in THF (6ml). After sttiring for 16h $NaHCO_3$ aqueous solution was added to the mixture. The mixture was extracted with EtOAc, which was washed with NH_4Cl aqueous solution and brine. The organic phase was dried over $MgSO_4$. After a filtration the solvent was removed under reduced pressure to obtain compound 102 (694mg, 100%) as a white solid.
1H-NMR ($CDCl_3$)δ: 1.46(9H, s), 1.56-2.14(4H, m), 3.29(4H, m), 4.18(1H, m), 5.24(1H, s), 5.27(1H, s), 6.46(1H, d, J=5.7Hz), 7.35(5H, m), 7.69(1H, d, J=5.7Hz).

4) The solution of compound 102 (694mg, 1.95mmol) in HCl/EtOAc (4mol/l, 8ml) was stirred for 30 min. The solvent was removed under reduced pressur, diluted with EtOH (16ml) then. A saturated $NaHCO_3$ aqueous solution was added to the solution to control pH at 9. The mixture was strred at 50 °C for 2h, then diluted with water. The mixture was extracted with $CHCl_3$, washed with brine, and dried over $MgSO_4$. The solvent was removed under reduced pressure to obtain compound 103 (413mg, 68%) as a yellow solid.
1H-NMR ($CDCl_3$)δ: 1.54-2.22(4H, m), 3.60(2H, m), 3.80(1H, t, J=12.0Hz), 4.18(1H, d, J=12.0Hz), 5.15(1H, d, J=9.9Hz), 5.35(1H, d, J=9.9Hz), 6.71(1H, d, J=5.4Hz), 7.33(3H, m), 7.50(1H, d, J=5.1Hz), 7.63(2H, d, J=7.2Hz).

5) NaOAc (118mg, 1.44mmol) and bromine (0.234ml, 2.62mmol) were added to the solution of compound 103 (408mg, 1.31mmol) in acetic acid (8ml), stirred for 30 min then. An aqueous solution of NaOH (2M) was added to the mixture, and extracted with CH_2Cl_2, washed with brine, and dried over Na_2SO_4. The solvent was removed under reduced pressure to give compound 104 (390mg, 77%) as a white solid.
1H-NMR ($CDCl_3$)δ: 1.55-2.19(4H, m), 3.55-4.02(5H, m), 5.12(1H, d, J=9.6Hz), 5.35(1H, d, J=9.9Hz), 7.29-7.38(3H, m), 7.61(1H, s), 7.67(2H, d, J=6.6Hz).

6) Tetrakis triphenylphosphine palladium (0)(77mg, 0.067mmol) and N,N-diisopropylethylamine (0.29ml, 1.67mmol) were added to the solution of compound 104 (130mg, 0,334mmol) in DMSO (2.6ml). the mixture was stirred under CO atmosphere for 2h at 80°C . The reaction mixture was diluted with a saturated NH_4Cl aqueous solution, extracted with EtOAc then. And the organic phase was washed with brine, and dried over Na_2SO_4. The precipitate was purified by silicagel column chromatography, and fractions eluting with MeOH/EtOAc were concentrated

under reduced pressure to obtain compound 105 (115mg, 75%) as a white oil.
1H-NMR ($CDCl_3$)δ: 1.56-2.33(4H, m), 3.66(2H, m), 3.90(2H, m), 4.19(1H, s), 4.66(2H, m), 5.20(1H, d, J=9.9Hz), 5.37(1H, d, J=9.9Hz), 7.00(2H, t, J=8.7Hz), 7.33(5H, m), 7.61(2H, m), 8.39(1H, m), 10.50(1H, s).

7) A mixture of compound 105 (111mg, 0.241mmol) and palladium-carbon (10%, 22 mg) in THF (8ml) and MeOH (2ml) was stirred under hydrogen atmosphere for 3h. After celite filteration the solvent was removed under reduced pressure to give the example X-1 (57mg, 64%) as a white solid.
Melting point: 274°C
1H-NMR (DMSO-d_6)δ:1.56-2.25(4H, m), 3.48-3.65(2H, m), 4.01(2H, m), 4.51(2H, d, J=5.7Hz), 4.71(1H, d, J=9.9Hz), 7.14(2H, t, J=9.0Hz), 7.33(2H, dd, J=5.7, 8.7Hz), 8.41(1H, s), 10.44(1H, t, J=6.0Hz), 12.18(1H, s).

The following compounds were synthesized using the similar method.

Example X-2)
(R)-6-Hydroxy-5,7-dioxo-2,3,5,7,11,11a-hexahydro-1H-pyrido[1,2-a]pyrrolo[1,2-d]pyrazine-8-carboxylic acid 2,4-difluoro-benzylamide
Melting point: 300°C
1H-NMR (DMSO-d_6)δ: 1.03-2.20(4H, m), 3.39-3.66(2H, m), 4.02(2H, m), 4.54(2H, d, J=6.0Hz), 4.71(1H, d, J=9.9Hz), 7.06(1H, m), 7.23(1H, m), 7.38(1H, m), 8.41(1H, s), 10.43(1H, t, J=6.0Hz), 12.19(1H, s).

Example X-3)
(R)-6-Hydroxy-5,7-dioxo-2,3,5,7,11,11a-hexahydro-1H-pyrido[1,2-a]pyrrolo[1,2-d]pyrazine-8-carboxylic acid 3-chloro-2-fluoro-benzylamide
Melting point: 304°C
1H-NMR (DMSO-d_6)δ: 3.44-3.66(2H, m), 4.01(2H, m), 4.61(2H, d, J=5.4Hz), 4.70(1H, d, J=9.0Hz), 7.20(1H, m), 7.31(1H, m), 7.49(1H, m), 8.41(1H, s), 10.49(1H, t, J=5.7Hz), 12.20(1H, s).

Example X-4)
1-Hydroxy-2,9-dioxo-2,5,6,7,8,9,10,10a-octahydro-4a,8a-diaza-anthracene-3-carboxylic acid 4-fluoro-benzylamide

Melting point: 259°C

1H-NMR (DMSO-d_6)δ:1.33-1.79(6H, m), 2.51(1H, m), 3.88(1H, m), 4.12(1H, dd, J=9.3, 14.1Hz), 4.38(1H, d, J=12.9Hz), 4.53(3H, m), 7.16(2H, t, J=9.0Hz), 7.34(2H, dd, J=5.7, 8.7Hz), 8.39(1H, s), 10.44(1H, t, J=6.3Hz), 12.84(1H, s).

According to the same manner as that of Example C-21, the following Example compounds Y-1 to Y-18 were synthesized.

Example Y-1)

(3S,9aS)-5-Hydroxy-3-methyl-6,10-dioxo-3,4,6,9,9a,10-hexahydro-2H-1-oxa-4a,8a-diaza-anthracene-7-carboxylic acid 2,4-difluoro-benzylamide

Example Y-9)

(3R,9aR)-5-Hydroxy-3-methyl-6,10-dioxo-3,4,6,9,9a,10-hexahydro-2H-1-oxa-4a,8a-diaza-anthracene-7-carboxylic acid 2,4-difluoro-benzylamide

1H-NMR ($CDCl_3$)δ: 0.90(3H, d, J=6.9Hz), 2.00-2.10(1H, m), 2.70(1H, dd, J=11.6, 13.4Hz), 3.41(1H, dd, J=11.2, 12.9Hz), 4.05-4.45(2H, m), 4.30-4.38(1H, dd, J=4.0, 14.1Hz), 4.63(2H, d, J=5.9Hz), 4.65-4.75(1H, m), 4.98(1H, t, J=3.7Hz), 6.80-6.84(2H, m), 7.32-7.40(1H, m), 8.31(1H, s), 10.38(1H, brs), 12.37(1H, s).

Example Y-2)

(4S,9aR)-5-Hydroxy-4-methyl-6,10-dioxo-3,4,6,9,9a,10-hexahydro-2H-1-oxa-4a,8a-diaza-anthracene-7-carboxylic acid 2,4-difluoro-benzylamide

Example Y-3)

(4R,9aS)-5-Hydroxy-4-methyl-6,10-dioxo-3,4,6,9,9a,10-hexahydro-2H-1-oxa-4a,8a-diaza-anthracene-7-carboxylic acid 2,4-difluoro-benzylamide

1H-NMR ($CDCl_3$)δ: 1.42(3H, d, J=7.0Hz), 1.56(1H, dd, J=2.0, 14.0Hz), 2.19-2.30(1H, m), 4.02(1H, d, J=2.2Hz), 4.05(1H, t, J=2.3Hz), 4.12(1H, dd, J=6.0, 13.6Hz), 4.27(1H, dd, J=4.2, 13.4Hz), 4.64(2H, d, J=5.9Hz), 4.95-5.05(1H, m), 5.26(2H, d, J=4.1, 5.8Hz), 6.75-6.85(2H, m), 7.30-7.40(1H, m), 8.30(1H, s), 10.38(1H, brs), 12.45(1H, s).

Example Y-4)

(2R,9aR)-5-Hydroxy-2-methoxymethyl-6,10-dioxo-3,4,6,9,9a,10-hexahydro-2H-1-oxa-4a,8a-diaza-anthracene-7-carboxylic acid 2,4-difluoro-benzylamide

Example Y-8)

(2S,9aS)-5-Hydroxy-2-methoxymethyl-6,10-dioxo-3,4,6,9,9a,10-hexahydro-2H-1-oxa-4a

,8a-diaza-anthracene-7-carboxylic acid 2,4-difluoro-benzylamide
1H-NMR ($CDCl_3$)δ: 1.60-1.80(2H, m), 3.09-3.21(1H, m), 3.37(3H, s), 3.35-3.50(2H, m), 4.00-4.11(1H, m), 4.24(1H, d, J=13.1Hz), 4.36(1H, d, J=10.1Hz), 4.64(1H, d, J=5.9Hz), 4.70-4.80(1H, m), 5.12(1H, s), 6.75-6.85(2H, m), 7.30-7.40(1H, m), 8.30(1H, s), 10.38(1H, brs), 12.33(1H, brs).

Example Y-5)
(5aR,6aS,10aR)-1-Hydroxy-2,12-dioxo-2,5,5a,7,8,9,10,10a,11,12-decahydro-6aH-6-oxa-4a,11a-diaza-naphthalene-8-carboxylic acid 2,4-difluoro-benzylamide [racemate]
1H-NMR (DMSO-d6)δ: 1.00-1.85(9H, m), 2.90(1H, t, J=4.2Hz), 4.36(1H, dd, J=4.2, 12.9Hz), 4.44-4.57(4H, m), 5.32(1H, t, J=3.9Hz), 7.03-7.09(1H, m), 7.20-7.27(1H, m), 7.35-7.43(1H, m), 8.49(1H, s), 10.34(1H, brs).

Example Y-6)
(2S,9aR)-2-Ethyl-5-hydroxy-6,10-dioxo-3,4,6,9,9a,10-hexahydro-2H-1-oxa-4a,8a-diaza-anthracene-7-carboxylic acid 2,4-difluoro-benzylamide
Example Y-7)
(2R,9aS)-2-Ethyl-5-hydroxy-6,10-dioxo-3,4,6,9,9a,10-hexahydro-2H-1-oxa-4a,8a-diaza-anthracene-7-carboxylic acid 2,4-difluoro-benzylamide
1H-NMR (DMSO-d6)δ: 0.87(3H, d, J=5.4Hz), 1.40-1.51(3H, m), 1.75(1H, d, J=10.8Hz), 3.22(1H, t, J=10.2Hz), 3.73-3.78(1H, m), 4.41-4.57(4H, m), 5.29(1H, s), 7.03-7.07(1H, m), 7.21-7.26(1H, m), 7.37-7.42(1H, m), 8.50(1H, s), 10.34(1H, brs), 12.48(1H, s).

Example Y-10)
(2S,9aS)-5-Hydroxy-6,10-dioxo-2-phenyl-3,4,6,9,9a,10-hexahydro-2H-1-oxa-4a,8a-diaza-anthracene-7-carboxylic acid 2,4-difluoro-benzylamide
1H-NMR ($CDCl_3$)δ: 1.70-1.82(1H, m), 1.98(1H, d, J=9.6Hz), 3.49(1H, t, J=9.6Hz), 4.54-4.68(5H, m), 4.98(1H, d, J=8.7Hz), 5.51(1H, s), 7.04-7.08(1H, m), 7.21-7.42(7H, m), 8.50(1H, s), 10.38(1H, s), 12.45(1H, s).

Example Y-11)
(2S,9aS)-5-Hydroxy-2-isopropyl-6,10-dioxo-3,4,6,9,9a,10-hexahydro-2H-1-oxa-4a,8a-diaza-anthracene-7-carboxylic acid 2,4-difluoro-benzylamide
Example Y-12)
(2R,9aR)-5-Hydroxy-2-isopropyl-6,10-dioxo-3,4,6,9,9a,10-hexahydro-2H-1-oxa-4a,8a-di

aza-anthracene-7-carboxylic acid 2,4-difluoro-benzylamide
1H-NMR (DMSO-d6)δ: 0.86(6H, dd, J=4.8, 13.5Hz), 1.41-1.49(1H, m), 1.57-1.69(1H, m), 1.72-1.78(1H, m), 3.20(1H, t, J=8.4Hz), 3.52-3.59(1H, m), 4.41-4.46(5H, m), 5.29(1H, s), 7.01-7.08(1H, m), 7.21-7.26(1H, m), 7.37-7.43(1H, m), 8.50(1H, s), 10.35(1H, brs), 12.48(1H, s).

Example Y-13)
(3S,9aS)-5-Hydroxy-3-methyl-6,10-dioxo-3,4,6,9,9a,10-hexahydro-2H-1-oxa-4a,8a-diaza-anthracene-7-carboxylic acid 4-fluoro-benzylamide
Example Y-14)
(3R,9aR)-5-Hydroxy-3-methyl-6,10-dioxo-3,4,6,9,9a,10-hexahydro-2H-1-oxa-4a,8a-diaza-anthracene-7-carboxylic acid 4-fluoro-benzylamide
1H-NMR (DMSO-d6)δ: 0.81(3H,d, J=6.6Hz), 1.84-1.93(1H, m), 2.86(1H, t, J=12.5Hz), 3.48(1H, t, J=11.1Hz), 3.97-4.03(1H, m), 4.41-4.60(3H, m), 4.52(2H, d, J=5.9Hz), 5.20(1H, t, J=3.8Hz), 7.12-7.20(2H, m), 7.32-7.38(2H, m), 8.52(1H, s), 10.36(1H, t, J=5.9Hz), 12.45(1H, s).

Example Y-15)
(2R,9aS)-5-Hydroxy-2-methyl-6,10-dioxo-3,4,6,9,9a,10-hexahydro-2H-1-oxa-4a,8a-diaza-anthracene-7-carboxylic acid 2,4-difluoro-benzylamide
Example Y-16)
(2S,9aR)-5-Hydroxy-2-methyl-6,10-dioxo-3,4,6,9,9a,10-hexahydro-2H-1-oxa-4a,8a-diaza-anthracene-7-carboxylic acid 2,4-difluoro-benzylamide
1H-NMR (DMSO-d6)δ: 1.14(3H, d, J=6.0Hz), 1.38(1H, m), 1.75(1H, d, J=13.8Hz), 3.18-3.29(1H, m), 3.95-4.06(1H, m), 4.42-4.58(3H, m), 4.54(2H, d, J=5.7Hz), 5.30(1H, t, J=3.9Hz), 7.03-7.10(1H, m), 7.20-7.29(1H, m), 7.35-7.44(1H, m), 8.50(1H, s), 10.35(1H, t, J=5.7Hz), 12.48(1H, s).

Example Y-17)
(2S,9aR)-5-Hydroxy-2-methyl-6,10-dioxo-3,4,6,9,9a,10-hexahydro-2H-1-oxa-4a,8a-diaza-anthracene-7-carboxylic acid 4-fluoro-benzylamide
Example Y-18)
(2R,9aS)-5-Hydroxy-2-methyl-6,10-dioxo-3,4,6,9,9a,10-hexahydro-2H-1-oxa-4a,8a-diaza-anthracene-7-carboxylic acid 4-fluoro-benzylamide
1H-NMR (DMSO-d6)δ: 1.15(3H, d, J=6.0Hz), 1.35-1.50(1H, m), 1.75(1H, d, J=12.9Hz),

3.23(1H, td, J=13.0, 2.8Hz), 3.95-4.03(1H, m),4.41-4.59(3H, m), 4.52(2H, d, J=6.0Hz), 5.30(1H, t, J=3.9Hz), 7.12-7.19(2H, m), 7.32-7.38(2H, m), 8.52(1H, s), 10.36(1H, t, J=6.0Hz), 12.48(1H, s).


Corresponding amino-alcohol derivatives used in syntheses of Y-1 to Y-18 were prepared as optically pure version using methods similar to those described in the following reports.

3-Amino-2-methyl-propan-1-ol, and 4-Amino-butan-2-ol were prepared according to the method of Russell A. Barrow (J. Am. Chem. Soc. 1995, 117, 2479-2490).
3-Amino-butan-1-ol were prepared according to the method of P. Besse (Tetrahedron Asymmetry 10(1999) 2213-2224).
1-Amino-pentan-3-ol, 1-Amino-4-methyl-pentan-3-ol, 4-Amino-1-methoxy-butan-2-ol, and 3-Amino-1-phenyl-propan-1-ol were prepared according to the method described in the following literatures, U.S. Pat. Appl. Publ., 2004133029, 08 Jul 2004, PCT Int. Appl., 2002012173, 14 Feb 2002.


All examples below consist of >95% ee and >6:1 diastereomeric purity unless indicated otherwise. The compounds shown in table ZZ consist of mixtures of diastereomers at the depicted stereocenter in ratios of 1:1 to >10:1. Stereocenters that were formed during the process' below have been assigned using NMR techniques well know in the art (1D and 2D method) and/or using vibrational circular dichroism techniques. Stereochemical assignment determinatons were performed on representative examples and closely related compounds were assigned by analogy in some cases. The schemes below are meant to be general guidance to how examples were synthesized. It will be possible that one skilled in the art may rearrange the order of steps or change substituents to apply the method described below and in the examples to construct compounds of the general formula. Additional methods known to those skilled in the art or commonly present in the literature may also be applied

to perform similar transformations and arriving at the same compounds of the general formula or amino alcohol and diamine precursors.


[Chemical formula 68]




[Chemical formula 69]




[Chemical formula 70]




[Chemical formula 71]




[Chemical formula 72]




[Chemical formula 73]




[Chemical formula 74]




[Chemical formula 74]




[Chemical formula 75]




[Chemical formula 76]




[Chemical formula 77]




[Chemical formula 78]




[Chemical formula 79]




[Chemical formula 80]




[Chemical formula 81]



[Chemical formula 82]




[Chemical formula 83]



Example Z-1:

(3*R*,11a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-methyl-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide sodium salt.



a)

(3*R*,11a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-3-methyl-5,7-dioxo-6-[(phenylmethyl)oxy]-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide. To a solution of 16a (409 mg, 0.87 mmol) in dichloroethane (20 mL) was added

(2*R*)-2-amino-1-propanol (0.14 mL, 1.74 mmol) and 10 drops of glacial acetic acid. The resultant solution was heated at reflux for 2 h. Upon cooling, Celite was added to the mixture and the solvents removed *in vacuo* and the material was purified via silica gel chromatography (2% CH_3OH/CH_2Cl_2 gradient elution) to give (3*R*,11a*S*)-*N*-[(2,4-difluorophenyl)methyl]-3-methyl-5,7-dioxo-6-[(phenylmethyl)oxy]-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (396 mg, 92%) as a glass. ^{1}H NMR ($CDCl_3$) δ 10.38 (m, 1 H), 8.42 (s, 1 H), 7.54-7.53 (m, 2 H), 7.37-7.24 (m, 4 H), 6.83-6.76 (m, 2 H), 5.40 (d, *J* = 10.0 Hz, 1 H), 5.22 (d, *J* = 10.0 Hz, 1 H), 5.16 (dd, *J* = 9.6, 6.0 Hz, 1 H), 4.62 (m, 2 H), 4.41 (m, 1 H), 4.33-4.30 (m, 2 H), 3.84 (dd, *J* = 12.0, 10.0 Hz, 1 H), 3.63 (dd, *J* = 8.4, 7.2 Hz, 1 H), 1.37 (d, *J* = 6.0 Hz, 3 H); ES$^+$ MS: 496 (M+1).


b)

(3*R*,11a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-methyl-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide sodium salt. To a solution of (3*R*,11a*S*)-*N*-[(2,4-difluorophenyl)methyl]-3-methyl-5,7-dioxo-6-[(phenylmethyl)oxy]-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (396 mg, 0.80 mmol) in methanol (30 mL) was added 10% Pd/C (25 mg). Hydrogen was bubbled through the reaction mixture via a balloon for 2 h. The resultant mixture was filtered through Celite with methanol and dichloromethane. The filtrate was concentrated *in vacuo* to give (3*R*,11a*S*)-*N*-[(2,4-difluorophenyl)methyl]-6-hydroxy-3-methyl-5,7-dioxo-2,3,5,7,11,11a-

hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide as a pink tinted white solid (278 mg, 86%). ^{1}H NMR ($CDCl_3$) δ 11.47 (m, 1 H), 10.29 (m, 1 H), 8.32 (s, 1 H), 7.36 (m, 1 H), 6.82 (m, 2 H), 5.31 (dd, *J* = 9.6, 3.6 Hz, 1 H), 4.65 (m, 2 H), 4.47-4.38 (m, 3 H), 3.93 (dd, *J* = 12.0, 10.0 Hz, 1 H), 3.75 (m, 1 H), 1.49 (d, *J* = 5.6 Hz, 3 H); ES^+ MS: 406 (M+1). The above material (278 mg, 0.66 mmol) was taken up in ethanol (10 mL) and treated with 1 *N* sodium hydroxide (aq) (0.66 mL, 0.66 mmol). The resulting suspension was stirred at room temperature for 30 min. Ether was added and the liquids were collected to provide the sodium salt of the title compound as a white powder (291 mg, 99%). ^{1}H NMR (DMSO-d_6) δ 10.68 (m, 1 H), 7.90 (s, 1 H), 7.35 (m, 1 H), 7.20 (m, 1 H), 7.01 (m, 1 H), 5.20 (m, 1 H), 4.58 (m, 1 H), 4.49 (m, 2 H), 4.22 (m, 2 H), 3.74 (dd, *J* = 11.2, 10.4 Hz, 1 H), 3.58 (m, 1 H), 1.25 (d, *J* = 4.4 Hz, 3 H).

Example Z-2:

(4a*R*,13a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-10-hydroxy-9,11-dioxo-2,3,4a,5,9,11,13,13a-octahydro-1*H*-pyrido[1,2-*a*]pyrrolo[1',2':3,4]imidazo[1,2-*d*]pyrazine-8-carboxamide.



a)

(4a*R*,13a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-9,11-dioxo-10-[(phenylmethyl)oxy]-2,3,4a,5,9,11,13,13a-octahydro-1*H*-pyrido[1,2-*a*]pyrrolo[1',2':3,4]imidazo[1,2-*d*]pyrazine-8-carboxamide. A solution of 16a (24 mg, 0.05 mmol), [(2*S*)-2-pyrrolidinylmethyl]amine (0.1 mL) and 2 drops of glacial acetic acid were heated under microwave conditions at 140

°C for 10 min. Upon cooling, Celite was added to the mixture and the solvents removed *in vacuo* and the material was purified via silica gel chromatography (2% CH_3OH/CH_2Cl_2 gradient elution) to give (4a*R*,13a*S*)-*N*-[(2,4-difluorophenyl)methyl]-9,11-dioxo-10-[(phenylmethyl)oxy]-2,3,4a,5,9,11,13,13a-octahydro-1*H*-pyrido[1,2-*a*]pyrrolo[1',2':3,4]imidazo[1,2-*d*]pyrazine-8-carboxamide (19 mg, 71%) as a white solid. ^1H NMR ($CDCl_3$) δ 10.41 (m, 1 H), 8.38 (s, 1 H), 7.56 (m, 2 H), 7.38-7.24 (m, 4 H), 6.80 (m, 2 H), 5.38 (d, *J* = 9.6 Hz, 1 H), 5.10 (d, *J* = 10.0 Hz, 1 H), 4.62 (m, 2 H), 4.40 (m, 1 H), 4.25 (dd, *J* = 12.0, 6.8 Hz, 1 H), 4.10 (d, *J* = 12.8 Hz, 1 H), 3.83 (m, 1 H), 3.71 (m, 1 H), 3.14-3.04 (m, 2 H), 2.78 (m, 1 H), 2.11-1.58 (m, 4 H); ES^+ MS: 521 (M+1).

b)


(4a*R*,13a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-10-hydroxy-9,11-dioxo-2,3,4a,5,9,11,13,13a-octahydro-1*H*-pyrido[1,2-*a*]pyrrolo[1',2':3,4]imidazo[1,2-*d*]pyrazine-8-carboxamide.

To a solution of (4a*R*,13a*S*)-*N*-[(2,4-difluorophenyl)methyl]-9,11-dioxo-10-[(phenylmethyl)oxy]-2,3,4a,5,9,11,13,13a-octahydro-1*H*-pyrido[1,2-*a*]pyrrolo[1',2':3,4]imidazo[1,2-*d*]pyrazine-8-carboxamide (19 mg, 0.04 mmol) in methanol (8 mL) was added 10% Pd/C (10 mg). Hydrogen was bubbled through the reaction mixture via a balloon for 2 h. The resultant mixture was filtered through Celite with methanol and dichloromethane. The filtrate was concentrated *in vacuo* to give the title compound (6 mg, 38%) as a white solid. ^1H NMR ($CDCl_3$) δ 11.73 (m, 1 H), 10.36 (m, 1 H), 8.31 (s, 1 H), 7.33 (m, 1 H), 6.78 (m, 2 H), 4.62 (m, 2 H), 4.50 (m, 1 H), 4.27-4.19 (m, 2 H), 3.87-3.77 (m, 2 H),

3.16-3.08 (m, 2 H), 2.83 (m, 1 H), 2.11-1.65 (m, 4 H); ES+ MS: 431 (M+1).

Example Z-3:

(3a*S*,13a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-8-hydroxy-7,9-dioxo-1,2,3,3a,4,5,7,9,13,13a-decahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrrolo[1,2-*c*]pyrimidine-10-carboxamide.



a) *N*-BOC-(2*S*)-2-(Hydroxymethyl)-1-pyrrolidine. To a solution of *N*-BOC-L-proline (4.17 g, 19.4 mmol) in THF (40 mL) at 0 °C was added BH_3-THF (21.4 mL, 1 M in THF, 21.4 mmol) dropwise. The bath was removed and the resultant solution stirred at room temperature for 2 h. Methanol was added to quench the mixture and the solvents were removed *in vacuo*. The residue was taken up in ethyl acetate and washed with sodium bicarbonate and brine. The aqueous layers were extracted twice with ethyl acetate. The combined organics were dried over Na_2SO_4, filtered and concentrated to give *N*-BOC-(2*S*)-2-(hydroxymethyl)-1-pyrrolidine (3.82 g, 98%) as a clear oil. This material was used without further purification. ^{1}H NMR ($CDCl_3$) δ 3.94 (m, 1 H), 3.62 (dd, *J* = 11.2, 3.2 Hz, 1 H), 3.56 (dd, *J* = 10.8, 7.2 Hz, 1 H), 3.44 (m, 1 H), 3.29 (m, 1 H), 2.62 (br, 1 H), 1.98 (m, 1 H), 1.85-1.72 (m, 2 H), 1.58 (m, 1 H).

b) *N*-BOC-(2*S*)-2-({[(4-Methylphenyl)sulfonyl]oxy}methyl)-1-pyrrolidine. To a cold (0 °C) solution of *N*-BOC-(2*S*)-2-(hydroxymethyl)-1-pyrrolidine (350 mg, 1.74 mmol) in dichloromethane (20 mL) was added triethylamine (0.29 mL, 2.08 mmol), and

toluenesulfonyl chloride (398 mg, 2.08 mmol). *N,N*-dimethylaminopyridine (70 mg) was added and the resultant solution was allowed to warm to rt as the bath warmed and stirred for 4 h. Water was added and the layers separated. The aqueous layer was washed with sodium bicarbonate and then with brine. The combined organics were dried over Na_2SO_4, filtered and concentrated followed by flash chromatography purification to give *N*-BOC-(2*S*)-2-({[(4-methylphenyl)sulfonyl]oxy}methyl)-1-pyrrolidine (460 mg, 75%) as a clear oil. ^1H NMR exists as rotamers ($CDCl_3$) δ 7.77 (d, 2 H), 7.33 (m, 2 H), 4.08 (m, 1 H), 3.97-3.88 (m, 1 H), 3.35-3.25 (m, 2 H), 2.43 (s, 3 H), 1.95-1.79 (m, 4 H), 1.40 and 1.35 (s, 9 H rotomeric BOC *t*-butyl).

c) *N*-BOC-(2*S*)-2-Cyano-1-pyrrolidine. A mixture of *N*-BOC-(2*S*)-2-({[(4-methylphenyl)sulfonyl]oxy}methyl)-1-pyrrolidine (460 mg, 1.29 mmol) and KCN (256 mg, 3.88 mmol) were heated at 90 °C in DMSO (10 mL) for 6.5 h. The mixture was cooled to room temperature and EtOAc and water were added. The organics were washed with water twice and then with brine. The aqueous layers were extracted with EtOAc and the combined organics dried over Na_2SO_4, filtered and concentrated followed by flash chromatography purification to give *N*-BOC-(2*S*)-2-cyano-1-pyrrolidine (179 mg, 66%) as an oil. ^1H NMR exists as rotomers ($CDCl_3$) δ 3.99 (m, 1 H), 3.43-3.37 (m, 2 H), 2.33-2.51 (m, 2 H), 2.17-1.83 (m, 4 H), 1.46 and 1.44 (s, 9 H rotomeric BOC *t*-butyl).

d) *N*-BOC-(2*S*)-2-(2-Aminoethyl)-1-pyrrolidine. A solution of *N*-BOC-

(2*S*)-2-cyano-1-pyrrolidine (179 mg, 0.85 mmol) in ethanol saturated with anhydrous ammonia was treated with Raney-Ni (1 mL of 50% aq. Suspension) and 50 psi of H_2 overnight. The mixture was filtered through Celite and the filtrate was concentrated *in vacuo*. The residue was purified by flash chromatography (10% CH_3OH/CH_2Cl_2 with 1% NH_4OH gradient elution) through a short plug of silica gel to give *N*-BOC-(2*S*)-2-(2-aminoethyl)-1-pyrrolidine (90 mg, 50%) as a clear oil. ^1H NMR exists as rotomers ($CDCl_3$) δ 3.88-3.77 (m, 1 H), 3.33-3.24 (m, 2 H), 2.66 (m, 2 H), 1.89-1.54 (m, 6 H), 1.40 (s, 9 H).

e) {2-[(2*S*)-2-Pyrrolidinyl]ethyl}amine. A solution of *N*-BOC-(2*S*)-2-(2-aminoethyl)-1-pyrrolidine (90 mg, 0.42 mmol) in THF (6 mL) was treated with 4 *N* HCl (aq) (2 mL) and stirred at room temperature for 3 h. The mixture was concentrated *in vacuo* to give the title compound as its HCl salt. A portion of this material (40 mg) was dissolved in methanol and treated with solid supported carbonate resin (MP-Carbonate, Argonaut Technologies) to freebase the amines. After 30 minutes, the solution was filtered through a fritted tube and the solvents removed carefully *in vacuo* to give {2-[(2*S*)-2-pyrrolidinyl]ethyl}amine (30 mg) as its free base. ^1H NMR ($CDCl_3$) δ 3.06 (m, 1 H), 2.94 (m, 1 H), 2.83 (m, 1 H), 2.79-2.69 (m, 2 H), 1.90-1.56 (m, 6 H).

f)

(3a*S*,13a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7,9-dioxo-8-[(phenylmethyl)oxy]-1,2,3,3a,4,5,7,9,13,13a-decahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrrolo[1,2-*d*]pyrimidine-10-carb

oxamide. A solution of 16a (30 mg, 0.06 mmol), {2-[(2*S*)-2-pyrrolidinyl]ethyl}amine (30 mg, 0.26mmol) and 2 drops of glacial acetic acid were heated under microwave conditions at 140 °C for 10 min. Upon cooling, Celite was added to the mixture and the solvents removed *in vacuo* and the material was purified via silica gel chromatography (2% CH_3OH/CH_2Cl_2 gradient elution) to give (3a*S*,13a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7,9-dioxo-8-[(phenylmethyl)oxy]-1,2,3,3a,4,5,7,9,13,13a-decahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrrolo[1,2-*c*]pyrimidine-10-carboxamide. (25 mg, 74%) as a film. ^{1}H NMR ($CDCl_3$) δ 10.44 (m, 1 H), 8.32 (s, 1 H), 7.59 (m, 2 H), 7.38-7.24 (m, 4 H), 6.80 (m, 2 H), 5.28-5.22 (m, 2 H), 4.67 (dd, *J* = 13.6, 2.8 Hz, 1 H), 4.62 (m, 2 H), 4.26 (m, 1 H), 4.11-4.03 (m, 2 H), 2.91 (m, 1 H), 2.81 (m, 1 H), 2.37 (m, 1 H), 2.24 (m, 1 H), 1.92 (m, 1 H), 1.82-1.76 (m, 3 H), 1.52-1.38 (m, 2 H); ES$^+$ MS: 535 (M+1).

g)


(3a*S*,13a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-8-hydroxy-7,9-dioxo-1,2,3,3a,4,5,7,9,13,13a-decahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrrolo[1,2-*c*]pyrimidine-10-carboxamide.

To a solution of (3a*S*,13a*S*)-*N*-[(2,4-difluorophenyl)methyl]-7,9-dioxo-8-[(phenylmethyl)oxy]-1,2,3,3a,4,5,7,9,13,13a-decahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrrolo[1,2-*c*]pyrimidine-10-carboxamide (25 mg, 0.05 mmol) in methanol (8 mL) was added 10% Pd/C (10 mg). Hydrogen was bubbled through the reaction mixture via a balloon for 18 h. The resultant mixture was filtered through Celite with methanol and dichloromethane. The filtrate was concentrated *in vacuo* to give the title compound (14 mg, 67%) as a

white solid. ^{1}H NMR ($CDCl_3$) δ 12.53 (br, 1 H), 10.44 (s, 1 H), 8.29 (s, 1 H), 7.34 (m, 1 H), 6.78 (m, 2 H), 4.71-4.58 (m, 3 H), 4.29-4.14 (m, 3 H), 2.99 (m, 1 H), 2.88 (m, 1 H), 2.44 (m, 1 H), 2.30 (m, 1 H), 1.97-1.38 (m, 6 H); ES^+ MS: 445 (M+1).

Example Z-4:

(4a*S*,13a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-10-hydroxy-9,11-dioxo-2,3,4a,5,9,11,13,13a-octahydro-1*H*-pyrido[1,2-*a*]pyrrolo[1',2':3,4]imidazo[1,2-*d*]pyrazine-8-carboxamide sodium salt.



a) [(2*R*)-2-Pyrrolidinylmethyl]amine. To a solution of *N*-BOC-(2*R*)-2-(aminomethyl)-1-pyrrolidine (1.37 g, 6.85 mmol) in THF (20 mL) was added 4 *N* HCl (aq) (8 mL). The resultant solution was stirred at room temperature overnight. The solvents were removed *in vacuo* and the residue was treated with MP-carbonate resin in methanol and dichloromethane. After 1 h, the resin was removed via filtration through a fritted tube and the volatiles were removed carefully *in vacuo* to produce the free based amine (760 mg crude >100%) as a oil. This material was used without further purification. ^{1}H NMR ($CDCl_3$) δ 3.13 (m, 1 H), 2.92 (m, 1 H), 2.82-2.62 (m, 5 H), 1.88-1.30 (m, 4 H).

b)

(4a*S*,13a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-9,11-dioxo-10-[(phenylmethyl)oxy]-2,3,4a,5

,9,11,13,13a-octahydro-1*H*-pyrido[1,2-*a*]pyrrolo[1',2':3,4]imidazo[1,2-*d*]pyrazine-8-carb oxamide. In a similar manner as described in example Z-2 from 16a (435 mg, 0.93 mmol) and [(2*R*)-2-pyrrolidinylmethyl]amine (200 mg, 2.0 mmol) in 1,2-dichloroethane (20 mL) and 15 drops of glacial acetic acid was obtained (4a*S*,13a*R*)-*N*-[(2,4-difluorophenyl)methyl]-9,11-dioxo-10-[(phenylmethyl)oxy]-2,3,4a,5, 9,11,13,13a-octahydro-1*H*-pyrido[1,2-*a*]pyrrolo[1',2':3,4]imidazo[1,2-*d*]pyrazine-8-carbo xamide (321 mg, 67%) as a white solid. ^{1}H NMR ($CDCl_3$) δ 10.41 (m, 1 H), 8.35 (s, 1 H), 7.56 (m, 2 H), 7.55-7.24 (m, 4 H), 6.80 (m, 2 H), 5.35 (d, J = 10.0 Hz, 1 H), 5.13 (d, J = 10.0 Hz, 1 H), 4.60 (m, 2 H), 4.38 (dd, J = 10.4, 3.2 Hz, 1 H), 4.21 (dd, J = 12.0, 6.8 Hz, 1 H), 4.04 (dd, J = 12.4, 2.8 Hz, 1 H), 3.77 (apparent t, J = 11.6 Hz, 1 H), 3.68 (m, 1 H), 3.11-3.00 (m, 2 H), 2.75 (m, 1 H), 2.08-1.84 (m, 3 H), 1.65 (m, 1 H); ES$^+$ MS: 521 (M+1).

c)

(4a*S*,13a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-10-hydroxy-9,11-dioxo-2,3,4a,5,9,11,13,13a -octahydro-1*H*-pyrido[1,2-*a*]pyrrolo[1',2':3,4]imidazo[1,2-*d*]pyrazine-8-carboxamide. In a similar manner as described in example Z-2 from (4a*S*,13a*R*)-*N*-[(2,4-difluorophenyl)methyl]-9,11-dioxo-10-[(phenylmethyl)oxy]-2,3,4a,5, 9,11,13,13a-octahydro-1*H*-pyrido[1,2-*a*]pyrrolo[1',2':3,4]imidazo[1,2-*d*]pyrazine-8-carbo xamide (518 mg, 0.99 mmol) and 10% Pd/C (35 mg) in methanol (40 mL) was obtained (4a*S*,13a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-10-hydroxy-9,11-dioxo-2,3,4a,5,9,11,13,13a -octahydro-1*H*-pyrido[1,2-*a*]pyrrolo[1',2':3,4]imidazo[1,2-*d*]pyrazine-8-carboxamide (430 mg, 99%) as a white solid. ^{1}H NMR ($CDCl_3$) δ 11.73 (m, 1 H), 10.36 (m, 1 H),


8.32 (s, 1 H), 7.35 (m, 1 H), 6.79 (m, 2 H), 4.64 (m, 2 H), 4.54 (dd, J = 10.8, 4.0 Hz, 1 H), 4.28-4.19 (m, 2 H), 3.90-3.79 (m, 2 H), 3.18-3.10 (m, 2 H), 2.84 (m, 1 H), 2.14-1.92 (m, 3 H), 1.72 (m, 1 H).

d)

(4a*S*,13a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-10-hydroxy-9,11-dioxo-2,3,4a,5,9,11,13,13a-octahydro-1*H*-pyrido[1,2-*a*]pyrrolo[1',2':3,4]imidazo[1,2-*d*]pyrazine-8-carboxamide sodium salt. In a similar manner as described in example Z-1 from (4a*S*,13a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-10-hydroxy-9,11-dioxo-2,3,4a,5,9,11,13,13a-octahydro-1*H*-pyrido[1,2-*a*]pyrrolo[1',2':3,4]imidazo[1,2-*d*]pyrazine-8-carboxamide (430 mg, 1.0 mmol) and sodium hydroxide (1.0 mL, 1.0 M aq, 1.0 mmol) in 20 mL of ethanol was formed the corresponding sodium salt (425 mg, 94%) as a white solid. ^{1}H NMR (D_2O) δ 7.85 (s, 1 H), 7.23 (m, 1 H), 6.82 (m, 2 H), 4.51-4.46 (m, 3 H), 4.28 (m, 1 H),3.95 (m, 1 H), 3.84 (m, 1 H), 3.62 (m, 1 H), 3.16 (m, 1 H), 2.89 (m, 1 H), 2.84 (m, 1H), 1.90 (m, 2 H), 1.73 (m, 1 H), 1.60 (m, 1 H). ES$^+$ MS: 431 (M+1).

Example Z-5:

(4a*S*,13a*R*)-*N*-[(4-Fluorophenyl)methyl]-10-hydroxy-9,11-dioxo-2,3,4a,5,9,11,13,13a-octahydro-1*H*-pyrido[1,2-*a*]pyrrolo[1',2':3,4]imidazo[1,2-*d*]pyrazine-8-carboxamide.




The title compound was made in two steps using a similar process to that described

in example Z-2. 16 (60 mg, 0.13 mmol) and [(2*R*)-2-pyrrolidinylmethyl]amine (100 mg, 1.0 mmol) were reacted in dichloromethane (2 mL) with acetic acid to give (4a*S*,13a*R*)-*N*-[(4-fluorophenyl)methyl]-9,11-dioxo-10-[(phenylmethyl)oxy]-2,3,4a,5,9,11,13,13a-octahydro-1*H*-pyrido[1,2-*a*]pyrrolo[1',2':3,4]imidazo[1,2-*d*]pyrazine-8-carboxamide (60 mg, 91%). This material was hydrogenated in a second step as described in example Z-2 to give (4a*S*,13a*R*)-*N*-[(4-fluorophenyl)methyl]-10-hydroxy-9,11-dioxo-2,3,4a,5,9,11,13,13a-octahydro-1*H*-pyrido[1,2-*a*]pyrrolo[1',2':3,4]imidazo[1,2-*d*]pyrazine-8-carboxamide (21 mg, 42%) as a white solid. ^{1}H NMR ($CDCl_3$) δ 11.72 (m, 1 H), 1.37 (m, 1 H), 8.33 (s, 1 H), 7.29 (m, 2 H), 6.97 (m, 2 H), 4.57 (m, 2 H), 4.52 (m, 1 H), 4.24-4.19 (m, 2 H), 3.87-3.76 (m, 2 H), 3.14-3.07 (m, 2 H), 2.82 (m, 1 H), 2.11-1.89 (m, 3 H), 1.68 (m, 1 H); ES$^+$ MS: 413 (M+1).

Example Z-6:

(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-5,7-dioxo-3-(phenylmethyl)-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide.




The title compound was made in two steps using a similar process to that described in example Z-2. 16a (37 mg, 0.08 mmol) and (2*S*)-2-amino-3-phenyl-1-propanol (35 mg, 0.24 mmol) were reacted in dichloromethane (2 mL) with acetic acid to give (3*S*,11a*R*)-*N*-[(2,4-difluorophenyl)methyl]-5,7-dioxo-3-(phenylmethyl)-6-[(phenylmethy

l)oxy]-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (41 mg, 91%). This material was hydrogenated in a second step as described in example Z-2 to give (3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-5,7-dioxo-3-(phenylmethyl)-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide. (25 mg, 75%) as a white solid. ^{1}H NMR ($CDCl_3$) δ 11.47 (br, 1 H), 10.28 (m, 1 H), 8.35 (m, 1 H), 7.37-7.26 (m, 4 H), 7.18 (m, 2 H), 6.79 (m, 2 H), 5.03 (m, 1 H), 4.64-4.61 (m, 3 H), 4.40 (m, 1 H), 4.23 (apparent t, J = 7.2 Hz, 1 H), 3.96 (dd, J = 8.8, 6.4 Hz, 1 H), 3.88 (apparent t, J = 11.2 Hz, 1 H), 3.37 (dd, J = 13.6, 3.2 Hz, 1 H), 2.99 (dd, J = 13.2 8.8 Hz, 1 H); ES$^+$ MS: 482 (M+1).

Example Z-7:

(3a*S*,13a*S*)-*N*-[(4-Fluorophenyl)methyl]-8-hydroxy-7,9-dioxo-1,2,3,3a,4,5,7,9,13,13a-decahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrrolo[1,2-*c*]pyrimidine-10-carboxamide.




The title compound was made in two steps using a similar process to that described in example Z-2. 16 (84 mg, 0.13 mmol) and {2-[(2*S*)-2-Pyrrolidinyl]ethyl}amine (150 mg, 1.3 mmol) were reacted in dichloromethane (2 mL) with acetic acid to give (3a*S*,13a*S*)-*N*-[(4-fluorophenyl)methyl]-7,9-dioxo-8-[(phenylmethyl)oxy]-1,2,3,3a,4,5,7,9,13,13a-decahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrrolo[1,2-*c*]pyrimidine-10-carboxamide (86 mg, 90%). This material was hydrogenated in a second step as described in

example Z-2 to give (3a*S*,13a*S*)-*N*-[(4-Fluorophenyl)methyl]-8-hydroxy-7,9-dioxo-1,2,3,3a,4,5,7,9,13,13a-decahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrrolo[1,2-*d*]pyrimidine-10-carboxamide. (63 mg, 88%) as a white solid. ^{1}H NMR ($CDCl_3/CD_3OD$) δ 10.45 (m, 1 H), 8.23 (s, 1 H), 7.35 (m, 2 H), 6.94 (t, *J* = 8.8 Hz, 2 H), 4.63 (m, 1 H), 4.58-4.48 (m, 2 H), 4.33 (dd, *J* = 13.6, 3.6 Hz, 1 H), 4.21 (m, 1 H), 4.11 (m, 1 H), 2.98 (m, 1 H), 2.85 (td, *J* = 13.2, 3.2 Hz, 1 H), 2.41 (m, 1 H), 2.29 (m, 1 H), 1.92 (m, 1 H), 1.83-1.75 (m, 3 H), 1.54-1.35 (m, 2 H); ES$^+$ MS: 427 (M+1).

Example Z-8:

(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-[(1*S*)-1-methylpropyl]-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide sodium salt.




The title compound was made in two steps using a similar process to that described in example Z-1. 16a (417 mg, 0.89 mmol) and L-isoleucinol (259 mg, 2.21 mmol) were reacted in 1,2-dichloroethane (40 mL) with acetic acid to give (3*S*,11a*R*)-*N*-[(2,4-difluorophenyl)methyl]-3-[(1*S*)-1-methylpropyl]-5,7-dioxo-6-[(phenylmethyl)oxy]-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (426 mg, 90%). This material was hydrogenated in a second step as described in example Z-1 to give

(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-[(1*S*)-1-methylpropyl]-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (376 mg, 99%) as a coarse white solid. ^{1}H NMR ($CDCl_3$) δ 11.43 (br, 1 H), 10.27 (br, 1 H), 8.32 (s, 1 H), 7.33 (m, 1 H), 6.79 (m, 2 H), 5.26 (dd, *J* = 9.6, 4.0 Hz, 1 H), 4.62 (m, 2 H), 4.42-4.35 (m, 2 H), 4.19 (dd, *J* = 8.8, 7.2 Hz, 1 H), 4.01 (dd, *J* = 8.8, 5.6 Hz, 1 H), 3.86 (dd, *J* = 12.0, 10.0 Hz, 1 H), 2.27 (m, 1 H), 1.40 (m, 1 H), 1.15 (m, 1 H), 0.97 (t, *J* = 7.2 hz, 3 H), 0.91 (d, *J* = 6.8 Hz, 3 H); ES$^+$ MS: 448 (M+1). This material (360 mg, 0.81 mmol) was treated with sodium hydroxide (0.81 mL, 1.0 M, 0.81 mmol) in ethanol (15 mL) as described in example Z-1 to provide its corresponding sodium salt (384 mg, 99%) as a white solid. ^{1}H NMR (DMSO-d_6) δ 10.82 (m, 1 H), 7.80 (m, 1 H), 7.33 (m, 1 H), 7.18 (m, 1 H), 7.00 (m, 1 H), 5.14 (m, 1 H), 4.47 (d, *J* = 5.6 Hz, 2 H), 4.31 (m, 1 H), 4.18 (m, 1 H), 3.96 (m, 1 H), 3.84 (m, 1 H), 3.71 (m, 1 H), 3.40 (m, 1 H), 1.88 (m, 1 H), 1.36 (m, 1 H), 1.04 (m, 1 H), 0.85 (t, *J* = 7.2 Hz, 3 H), 0.80 (d, *J* = 6.8 Hz, 3 H); ES$^+$ MS: 448 (M+1).

Example Z-9:

(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-methyl-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide sodium salt.




The title compound was made in two steps using a similar process to that described in example Z-1. 16a (510 mg, 1.08 mmol) and (2*S*)-2-amino-1-propanol (0.17 mL,

2.17 mmol) were reacted in 1,2-dichloroethane (20 mL) with acetic acid to give (3*S*,11a*R*)-*N*-[(2,4-difluorophenyl)methyl]-3-methyl-5,7-dioxo-6-[(phenylmethyl)oxy]-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (500 mg, 93%). This material was hydrogenated in a second step as described in example Z-1 to give (3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-methyl-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (386 mg, 94%) as a tinted white solid. ^{1}H NMR ($CDCl_3$) δ 11.46 (m, 1 H), 10.28 (m, 1 H), 8.32 (s, 1 H), 7.35 (m, 1 H), 6.80 (m, 2 H), 5.30 (dd, J = 10.0, 4.0 Hz, 1 H), 4.63 (m, 2 H), 4.48-4.37 (m, 3 H), 3.91 (dd, J = 12.0, 10.0 Hz, 1 H), 3.73 (m, 1 H), 1.48 (d, J = 6.0 Hz, 3 H); ES$^+$ MS: 406 (M+1). This material (385 mg, 0.95 mmol) was treated with sodium hydroxide (0.95 mL, 1.0 M, 0.95 mmol) in ethanol (15 mL) as described in example Z-1 to provide its corresponding sodium salt (381 mg, 94%) as a white solid. ^{1}H NMR (DMSO-*d*$_6$) δ 10.66 (m, 1 H), 7.93 (s, 1 H), 7.33 (m, 1 H), 7.20 (m, 1 H), 7.01 (m, 1 H), 5.19 (m, 1 H), 4.59 (m, 1 H), 4.48 (m, 2 H), 4.22 (m, 2 H), 3.75 (m, 1 H), 3.57 (m, 1 H), 1.24 (d, J = 5.6 Hz, 3 H).

Example Z-10:


(3*S*,11a*R*)-*N*-[(4-Fluorophenyl)methyl]-6-hydroxy-3-methyl-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide.



The title compound was made in two steps using a similar process to that described in example Z-2. 16 (100 mg, 0.22 mmol) and (2*S*)-2-amino-1-propanol (0.10 mL, 1.28 mmol) were reacted in dichloromethane (2 mL) with acetic acid to give (3*S*,11a*R*)-*N*-[(4-fluorophenyl)methyl]-3-methyl-5,7-dioxo-6-[(phenylmethyl)oxy]-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (100 mg, 95%). This material was hydrogenated in a second step as described in example Z-2 to give (3*S*,11a*R*)-*N*-[(4-Fluorophenyl)methyl]-6-hydroxy-3-methyl-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (80 mg, 99%) as a white solid. ^{1}H NMR ($CDCl_3$) δ 11.43 (br, 1 H), 10.28 (br, 1 H), 8.35 (s, 1 H), 7.28 (m, 2 H), 6.97 (m, 2 H), 5.29 (m, 1 H), 4.55-4.38 (m, 5 H), 3.89 (apparent t, *J* = 10.8 Hz, 1 H), 3.70 (m, 1 H), 1.45 (d, *J* = 5.6 Hz, 3 H); ES- MS: 386 (M-1).

Example Z-11:

(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-3-(1,1-dimethylethyl)-6-hydroxy-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide




The title compound was made in two steps using a similar process to that described in example Z-2. 16a (41 mg, 0.09 mmol) and freebased L-*tert*-leucinol (59 mg, 0.50 mmol) were reacted in dichloromethane (2 mL) with acetic acid to give (3*S*,11a*R*)-*N*-[(2,4-difluorophenyl)methyl]-3-(1,1-dimethylethyl)-5,7-dioxo-6-[(phenylm

ethyl)oxy]-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (40 mg, 86%). This material was hydrogenated in a second step as described in example Z-2 to give (3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-3-(1,1-dimethylethyl)-6-hydroxy-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (33 mg, 99%) as a tinted white solid. ^{1}H NMR ($CDCl_3$) δ 10.29 (s, 1 H), 8.37 (s, 1 H), 7.34 (m, 1 H), 6.79 (m, 2 H), 5.43 (m, 1 H), 4.62 (m, 2 H), 4.36 (m, 2 H), 4.21 (m, 1 H), 3.99 (, 1 H), 3.81 (m, 1 H), 1.03 (s, 9 H); ES$^+$ MS: 448 (M+1).

Example Z-12:

(3*S*,11a*R*)-3-(1,1-Dimethylethyl)-*N*-[(4-fluorophenyl)methyl]-6-hydroxy-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide.




The title compound was made in two steps using a similar process to that described in example Z-2. 16 (41 mg, 0.09 mmol) and freebased L-*tert*-leucinol (59 mg, 0.50 mmol) were reacted in dichloromethane (2 mL) with acetic acid to give (3*S*,11a*R*)-3-(1,1-dimethylethyl)-*N*-[(4-fluorophenyl)methyl]-5,7-dioxo-6-[(phenylmethyl)oxy]-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (40 mg, 85%). This material was hydrogenated in a second step as described in example Z-2 to give (3*S*,11a*R*)-3-(1,1-Dimethylethyl)-*N*-[(4-fluorophenyl)methyl]-6-hydroxy-5,7-dioxo-2,3,5,

7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (32 mg, 97%) as a tinted white solid. ^{1}H NMR ($CDCl_3$) δ 11.15 (br, 1 H), 10.32 (s, 1 H), 8.38 (s, 1 H), 7.29 (m, 2 H), 6.98 (m, 2 H), 5.48 (m, 1 H), 4.58 (m, 2 H), 4.36 (m, 2 H), 4.21 (m, 1H), 3.99 (m, 1H), 3.79 (m, 1 H), 1.02 (s, 9 H); ES$^+$ MS: 430 (M+1).

Example Z-13:

(8*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-5,7-dioxo-3-phenyl-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide.




The title compound was made in two steps using a similar process to that described in example Z-2. 16a (33 mg, 0.07 mmol) and L-phenylglycinol (19 mg, 0.14 mmol) were reacted in dichloromethane (2 mL) with acetic acid to give (3*S*,11a*R*)-*N*-[(4-fluorophenyl)methyl]-5,7-dioxo-3-phenyl-6-[(phenylmethyl)oxy]-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (37 mg, 95%). This material was hydrogenated in a second step as described in example Z-2 to give (3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-5,7-dioxo-3-phenyl-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (33 mg, 99%) as a tinted white solid. ^{1}H NMR ($CDCl_3$) δ 11.23 (br, 1 H), 10.27 (s, 1 H), 8.39 (s,1 H), 7.43-7.32 (m, 6 H), 6.80 (m, 2 H), 5.58 (d, J = 6.8 Hz, 1 H), 5.37 (apparent t, J = 6.8 Hz, 1 H), 4.67-4.62 (m, 3 H), 4.54 (d, J = 10.4 Hz, 1 H), 4.11 (m, 1 H), 4.01 (m, 1 H); ES$^+$

MS: 468 (M+1).

Example Z-14:


(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-(hydroxymethyl)-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide.



The title compound was made in two steps using a similar process to that described in example Z-2. 16a (50 mg, 0.10 mmol) and (2*R*)-2-amino-3-[(phenylmethyl)oxy]-1-propanol (0.1 mL) were reacted in dichloromethane (2 mL) with acetic acid to give (3*S*,11a*R*)-*N*-[(2,4-difluorophenyl)methyl]-5,7-dioxo-6-[(phenylmethyl)oxy]-3-{[(phenylmethyl)oxy]methyl}-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (61 mg, 99%). This material was hydrogenated in a second step as described in example Z-2 to give (3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-(hydroxymethyl)-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (37 mg, 87%) as a tinted white solid. ^{1}H NMR ($CDCl_3/CD_3OD$) δ 8.23 (s, 1 H), 7.32 (m, 1 H), 6.79 (m, 2 H), 5.31 (d, J = 7.6 Hz, 1 H), 4.56 (s, 2 H), 4.42-4.36 (m, 3 H), 4.17-4.11 (m, 2 H), 3.85 (m, 1 H), 3.62 (d, J = 11.2 Hz, 1 H).

Example Z-15:

(2*S*,3*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-methyl-5,7-dioxo-2-phenyl-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide.




The title compound was made in two steps using a similar process to that described in example Z-2. 16a (25 mg, 0.05 mmol) and (1*S*,2*R*)-(+)-norephedrine (0.1 mL) were reacted in dichloromethane (2 mL) with acetic acid to give (2*S*,3*R*)-*N*-[(2,4-difluorophenyl)methyl]-3-methyl-5,7-dioxo-2-phenyl-6-[(phenylmethyl)oxy]-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (30 mg, 99%). This material was hydrogenated in a second step as described in example Z-2 to give (2*S*,3*R*)-*N*-[(2,4-difluorophenyl)methyl]-6-hydroxy-3-methyl-5,7-dioxo-2-phenyl-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (25 mg, 91%) as a white solid. This material is a single diastereomer (>6:1 diastereomeric ratio but unconfirmed relative stereochemistry at the aminal center). ^{1}H NMR ($CDCl_3/CD_3OD$) δ 10.28 (m, 1 H), 8.38 (s, 1 H), 7.10-7.30 (m, 6 H), 6.78 (m, 2 H), 5.70 (d, J = 7.6 Hz, 1 H), 5.36 (d, J = 5.2 Hz, 1 H), 4.82 (m, 1 H), 4.61 (m, 2 H), 4.47 (d, J = 10.4 Hz, 1 H), 4.00 (apparent t, J = 10.4 Hz, 1 H), 0.94 (d, J = 6.4 Hz, 3 H); ES$^+$ MS: 482 (M+1).

Example Z-16:

(3*R*,11a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-5,7-dioxo-3-(phenylmethyl)-2,3,5,


7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide



The title compound was made in two steps using a similar process to that described in example Z-2. 16a (34 mg, 0.07 mmol) and (2*R*)-2-amino-3-phenyl-1-propanol (D-phenylalaninol) (50 mg, 0.33 mmol) were reacted in dichloromethane (2 mL) with acetic acid to give (3*R*,11a*S*)-*N*-[(2,4-difluorophenyl)methyl]-5,7-dioxo-3-(phenylmethyl)-6-[(phenylmethyl)oxy]-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (29 mg, 70%). This material was hydrogenated in a second step as described in example Z-2 to give (3*R*,11a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-5,7-dioxo-3-(phenylmethyl)-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (24 mg, 98%) as a white solid. ^{1}H NMR ($CDCl_3$) δ 11.46 (br, 1 H), 10.27 (m, 1 H), 8.33 (m, 1 H), 7.32-7.16 (m, 6 H), 6.78 (m, 2 H), 5.02 (m, 1 H), 4.61 (m, 3 H), 4.39 (m, 1 H), 4.22 (m, 1 H), 3.95 (m, 1 H), 3.87 (m, 1 H), 3.36 (m, 1 H), 2.97 (dd, J = 13.2 8.8 Hz, 1 H); ES$^+$ MS: 482 (M+1).

Example Z-17:


(3*R*,11a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-(2-methylpropyl)-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide.



The title compound was made in two steps using a similar process to that described in example Z-2. 16a (32 mg, 0.07 mmol) and (2*R*)-2-amino-4-methyl-1-pentanol (0.1 mL) were reacted in dichloromethane (2 mL) with acetic acid to give (3*R*,11a*S*)-*N*-[(2,4-difluorophenyl)methyl]-3-(2-methylpropyl)-5,7-dioxo-6-[(phenylmethyl)oxy]-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (43 mg, 99%). This material was hydrogenated in a second step as described in example Z-2 to give (3*R*,11a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-(2-methylpropyl)-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (32 mg, 90%) as a white solid. ^{1}H NMR ($CDCl_3$) δ 11.47 (br, 1 H), 10.29 (m, 1 H), 8.35 (s, 1 H), 7.39 (m, 1 H), 6.80 (m, 2 H), 5.31 (m, 1 H), 4.62 (m, 2 H), 4.44 (m, 2 H), 4.37 (m, 1 H), 3.88 (m, 1 H), 3.84 (dd, *J* = 8.0, 5.6 Hz, 1 H), 2.04 (m, 1 H), 1.62 (m, 1 H), 1.41 (m, 1 H), 1.00 (d, *J* = 5.6 Hz, 3 H), 0.99 (d, *J* = 6.0 Hz, 3 H); ES$^+$ MS: 448 (M+1).

Example Z-18:

(5a*R*,14a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-11-hydroxy-10,12-dioxo-1,2,3,4,5a,6,10,12,14,14a-decahydropyrido[1,2-*a*]pyrido[1',2':3,4]imidazo[1,2-*d*]pyrazine-9-carboxamide.



a) 1,1-Dimethylethyl (2*R*)-2-(aminocarbonyl)-1-piperidinecarboxylate. To a cold (0 °C) solution of (2*R*)-1-{[(1,1-dimethylethyl)oxy]carbonyl}-2-piperidinecarboxylic acid (1.0 g, 4.36 mmol) in THF (20 mL) was added triethylamine (0.60 mL, 4.36 mmol) followed by slow addition of methyl chloroformate (0.34 mL, 4.36 mmol). After a few minutes a suspension had formed. To this mixture was added concentrated NH_4OH (1.5 mL) and the solution was allowed to warm to rt as the bath warmed and stirred for a total of 4 h. The mixture was concentrated *in vacuo* and the residue was taken up in EtOAc. The organic layer was washed with citric acid, sodium bicarbonate and then brine, dried over Na_2SO_4. Filtration and concentration gave 1,1-dimethylethyl (2*R*)-2-(aminocarbonyl)-1-piperidinecarboxylate (1.0 g, 99%). ^1H NMR ($CDCl_3$) δ 6.03 (br, 1 H), 5.45 (br, 1 H), 4.77 (br, 1 H), 4.06 (br, 1 H), 2.82 (m, 1 H), 2.29 (m, 1 H), 1.67-1.43 (m, 13 H).

b) 1,1-Dimethylethyl (2*R*)-2-cyano-1-piperidinecarboxylate. To a cold (0 °C) solution of 1,1-dimethylethyl (2*R*)-2-(aminocarbonyl)-1-piperidinecarboxylate (269 mg, 1.17 mmol) in THF (10 mL) was added triethylamine (0.33 mL, 2.34 mmol) and then trifluoroacetic anhydride (0.17 mL, 1.17 mmol). The mixture was stirred at 0 °C for 1 h and concentrated *in vacuo*. The residue was taken up in EtOAc and washed successively with sodium bicarbonate, 0.5 *N* HCl and brine. The organics were dried over Na_2SO_4, filtered and concentrated to give 1,1-dimethylethyl (2*R*)-2-cyano-1-piperidinecarboxylate (255 mg, 99%) as a crystalline solid upon standing. ^1H NMR ($CDCl_3$) δ 5.23 (br, 1 H), 4.05 (br, 1 H), 2.93 (br, 1 H), 1.93-1.39 (m, 6 H), 1.46 (s, 9 H).

c) 1,1-Dimethylethyl (2*R*)-2-(aminomethyl)-1-piperidinecarboxylate. An ammonia saturated ethanol solution of 1,1-dimethylethyl (2*R*)-2-cyano-1-piperidinecarboxylate (255 mg, 1.19 mmol) was reduced with Raney-Ni in a similar manner to that described in example Z-3 to give after filtration through a short plug of silica, 1,1-dimethylethyl (2*R*)-2-(aminomethyl)-1-piperidinecarboxylate (236 mg, 91%), as an oil. ^{1}H NMR ($CDCl_3/CD_3OD$) δ 4.15 (br, 1 H), 3.97 (m, 1 h), 2.96 (m, 1 H), 2.75-2.69 (m, 2 H), 2.23-2.08 (m, 3 H), 1.59-1.55 (m, 3 H), 1.43 (s, 9 H).

d) [(2*R*)-2-Piperidinylmethyl]amine bis HCl salt. A solution of 1,1-dimethylethyl (2*R*)-2-(aminomethyl)-1-piperidinecarboxylate (236 mg, 1.08 mmol) in THF (10 mL) was treated with 4 *N* HCl (3 mL) as described in example Z-3 to give the bis HCl salt of[(2*R*)-2-Piperidinylmethyl]amine. ^{1}H NMR (DMSO-d_6) δ 9.67 (br, 1 H), 9.48 (br, 1 H), 8.48 (br, 2 H), 3.70 (br, 2 H), 3.20 (m, 1 H), 3.04 (m, 1 H), 2.86 (m, 1 H), 1.89-1.41 (m, 6 H).


e)

(5a*R*,14a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-11-hydroxy-10,12-dioxo-1,2,3,4,5a,6,10,12,14,14a-decahydropyrido[1,2-*a*]pyrido[1',2':3,4]imidazo[1,2-*d*]pyrazine-9-carboxamide. The title compound was made in two steps using a similar process to that described in example Z-2. 16a (50 mg, 0.11 mmol) and [(2*R*)-2-Piperidinylmethyl]amine (150 mg, 1.31 mmol) (free based using carbonate resin as described in example Z-3) were reacted in dichloromethane (2 mL) with acetic acid to give

(5a*R*,14a*R*)-*N*-[(2,4-difluorophenyl)methyl]-10,12-dioxo-11-[(phenylmethyl)oxy]-1,2,3,4,5a,6,10,12,14,14a-decahydropyrido[1,2-*a*]pyrido[1',2':3,4]imidazo[1,2-*d*]pyrazine-9-carboxamide (50 mg, 88%). This material was hydrogenated in a second step as described in example Z-2 to give (5a*R*,14a*R*)-*N*-[(2,4-difluorophenyl)methyl]-11-hydroxy-10,12-dioxo-1,2,3,4,5a,6,10,12,14,14a-decahydropyrido[1,2-*a*]pyrido[1',2':3,4]imidazo[1,2-*d*]pyrazine-9-carboxamide (11 mg, 44%) as a white solid. ^{1}H NMR ($CD_3OD/CDCl_3$) δ 10.46 (m, 1 H), 8.32 (s, 1 H), 7.31 (m, 1 H), 6.80 (m, 2 H), 4.64-4.52 (m, 3 H), 4.14 (dd, *J* = 10.4, 2.8 Hz, 1 H), 3.91-3.82 (m, 2 H), 3.19 (apparent t, *J* = 10.8 Hz, 1 H), 3.08 (d, *J* = 10.4 Hz, 1 H), 2.50 (m, 1 H), 2.27 (m, 1 H), 1.99-1.30 m, 6 H); ES$^+$ MS: 445 (M+1).

Example Z-19:

(2*S*,3*S*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-[(methyloxy)methyl]-5,7-dioxo-2-phenyl-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide




The title compound was made in two steps using a similar process to that described in example Z-2. 16a (36 mg, 0.07 mmol) and (2*R*)-2-amino-4-methyl-1-pentanol (0.1 mL) were reacted in dichloromethane (2 mL) with acetic acid to give (2*S*,3*S*)-*N*-[(2,4-difluorophenyl)methyl]-3-[(methyloxy)methyl]-5,7-dioxo-2-phenyl-6-[(phenylmethyl)oxy]-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-c

arboxamide. This material was hydrogenated in a second step as described in example Z-2 to give (2*S*,3*S*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-[(methyloxy)methyl]-5,7-dioxo-2-phenyl-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (25 mg, 64% for 2 steps) as a white solid. This material is a single diastereomer (>6:1 diastereomeric ratio but unconfirmed relative stereochemistry at the aminal center). ^{1}H NMR ($CDCl_3$) δ 11.48 (br, 1 H), 10.30 (m, 1 H), 8.39 (s, 1 H), 7.39-7.24 (m, 6 H), 6.78 (m, 2 H), 5.46 (dd, *J* = 10.0, 3.6 Hz, 1 H), 5.33 (d, *J* = 7.2 Hz, 1 H), 4.63 (m, 2 H), 4.54 (dd, *J* = 12.4, 4.0 Hz, 1 H), 4.19 (m, 1 H), 4.12 (dd, *J* = 10.4, 3.2 Hz, 1 H), 4.06 (m, 1 H), 3.55 (dd, *J* = 10.4, 1.6 Hz, 1 H), 3.40 (s, 3 H); ES$^+$ MS: 512 (M+1).

Example Z-20:

(3*S*,11a*R*)-3-(Cyclohexylmethyl)-*N*-[(2,4-difluorophenyl)methyl]-6-hydroxy-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide.




The title compound was made in two steps using a similar process to that described in example Z-2. 16a (36 mg, 0.08 mmol) and (2*S*)-2-amino-3-cyclohexyl-1-propanol (30 mg, 0.19 mmol) were reacted in dichloromethane (2 mL) with acetic acid to give (3*S*,11a*R*)-3-(cyclohexylmethyl)-*N*-[(2,4-difluorophenyl)methyl]-5,7-dioxo-6-[(phenylmethyl)oxy]-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (27 mg, 61%). This material was hydrogenated in a second step as described in

example Z-2 to give (3*S*,11a*R*)-3-(cyclohexylmethyl)-*N*-[(2,4-difluorophenyl)methyl]-6-hydroxy-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (25 mg, 99%) as a white solid. ^{1}H NMR ($CDCl_3$) δ 11.48 (br, 1 H), 10.28 (s, 1 H), 8.33 (s, 1 H), 7.33 (m, 1 H), 6.78 (m, 2 H), 5.29 (m, 1 H), 4.61 (m, 2 H), 4.47-4.33 (m, 3 H), 3.87-3.81 (m, 2 H), 2.05 (m, 1 H), 1.75-1.64 (m, 6 H), 1.39 (m, 1 H), 1.25-1.14 (m, 3 H), 1.02-0.97 (m, 2 H); ES$^+$ MS: 488 (M+1).

Example Z-21:

(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-(1-methylethyl)-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide.




The title compound was made in two steps using a similar process to that described in example Z-1. 16a (42 mg, 0.09 mmol) and (2*S*)-2-amino-3-methyl-1-butanol (0.1 mL) were reacted in 1,2-dichloroethane (8 mL) with acetic acid to give (3*S*,11a*R*)-*N*-[(2,4-difluorophenyl)methyl]-3-(1-methylethyl)-5,7-dioxo-6-[(phenylmethyl)oxy]-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (40 mg, 86%). This material was hydrogenated in a second step as described in example Z-1 to give (3*S*,11a*R*)-*N*-[(2,4-difluorophenyl)methyl]-6-hydroxy-3-(1-methylethyl)-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (34 mg,

99%) as a white solid. ^{1}H NMR ($CDCl_3$) δ 10.29 (br, 1 H), 8.36 (s, 1 H), 7.33 (m, 1 H), 6.79 (m, 2 H), 5.29 (d, *J* = 6.4 Hz, 1 H), 4.61 (m, 2 H), 4.44 (d, *J* = 9.6 Hz, 1 H), 4.34 (m, 1 H), 4.17 (m, 1 H), 4.02 (dd, *J* = 8.4, 5.2 Hz, 1 H), 3.86 (m, 1 H), 2.37 (m, 1 H), 0.97 (m, 6 H); ES$^+$ MS: 434 (M+1).

Example Z-22:

(5a*R*,14a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-12-hydroxy-11,13-dioxo-5a,6a,7,11,13,14a-hexahydro-5*H*-indeno[1',2':4,5][1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-10-carboxamide.




The title compound was made in two steps using a similar process to that described in example Z-1. 16a (42 mg, 0.09 mmol) and (1*S*,2*R*)-1-amino-2,3-dihydro-1*H*-inden-2-ol (100 mg, 0.67 mmol) were reacted in 1,2-dichloroethane (5 mL) with acetic acid to give (5a*R*,14a*S*)-*N*-[(2,4-difluorophenyl)methyl]-11,13-dioxo-12-[(phenylmethyl)oxy]-5a,6a,7,11,13,14a-hexahydro-5*H*-indeno[1',2':4,5][1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-10-carboxamide (55 mg, 99%). This material was hydrogenated in a second step as described in example Z-1 to give (5a*R*,14a*S*)-*N*-[(2,4-difluorophenyl)methyl]-12-hydroxy-11,13-dioxo-5a,6a,7,11,13,14a-hexahydro-5*H*-indeno[1',2':4,5][1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-10-carboxamide (45 mg, 97%) as a white solid. ^{1}H NMR ($CDCl_3$) δ 10.28 (m, 1 H), 8.33 (s, 1 H),


7.69 (d, J = 7.2 Hz, 1 H), 7.34-7.19 (m, 4 H), 6.78 (m, 2 H), 5.96 (d, J = 6.0 Hz, 1 H), 5.32 (m, 1 H), 5.22 (m, 1 H), 4.60 (m, 2 H), 4.45 (d, J = 9.2 Hz, 1 H), 3.96 (apparent t, J = 10.8 Hz, 1 H), 3.40 (dd, J = 18.0, 6.8 Hz, 1 H), 3.24 (d, J = 17.6 Hz, 1 H);); ES+ MS: 480 (M+1).

Example Z-23 & Z-24:

(2*S*,3*R*,11a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-5,7-dioxo-2,3-diphenyl-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide &

(2*S*,3*R*,11a*R*)-*N*-[(2,4-difluorophenyl)methyl]-6-hydroxy-5,7-dioxo-2,3-diphenyl-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide.






The title compounds were made in two steps using a similar process to that described in example Z-1. 16a (40 mg, 0.09 mmol) and (1*S*,2*R*)-2-amino-1,2-diphenylethanol (50 mg, 0.23 mmol) were reacted in 1,2-dichloroethane (5 mL) with acetic acid to give (2*S*,3*R*,11a*S*)-*N*-[(2,4-difluorophenyl)methyl]-5,7-dioxo-2,3-diphenyl-6-[(phenylmethyl)

oxy]-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (34 mg, 63%) and (2*S*,3*R*,11a*R*)-*N*-[(2,4-difluorophenyl)methyl]-5,7-dioxo-2,3-diphenyl-6-[(phenylmethyl) oxy]-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (13 mg, 24%). These materials were hydrogenated in a second step as described in example Z-1 to give (2*S*,3*R*,11a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-5,7-dioxo-2,3-diphenyl-2,3,5,7, 11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (example Z-23, 29 mg, 99%) as a white solid and (2*S*,3*R*,11a*R*)-*N*-[(2,4-difluorophenyl)methyl]-6-hydroxy-5,7-dioxo-2,3-diphenyl-2,3,5,7, 11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (example Z-24, 10 mg, 89%) as a white solid respectively. For example Z-23: ^{1}H NMR (DMSO-d_6) δ 10.29 (t, J = 5.6 Hz, 1 H), 8.55 (s, 1 H), 7.38 (m, 1 H), 7.22 (m, 1 H), 7.11-6.95 (m, 11 H), 6.16 (dd, J = 10.4, 3.6 Hz, 1 H), 5.71 (m, 2 H), 4.90 (m, 1 H), 4.54 (m, 2 H), 4.38 (t, J = 11.2 Hz, 1 H); ES$^+$ MS: 544 (M+1). For example Z-24: ^{1}H NMR ($CDCl_3$) δ 11.64 (br, 1 H), 10.30 (s, 1 H), 8.45 (s, 1 H), 7.34 (m, 1 H), 7.01-6.90 (m, 10 H), 6.80 (m, 2 H), 5.56 (m, 2 H), 5.42 (d, J = 6.4 Hz, 1 H), 4.73 (m, 1 H), 4.63 (m, 2 H), 4.49 (m, 1 H); ES$^+$ MS: 544 (M+1).

Example Z-25:


(3*R*,11a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-(1-methylethyl)-5,7-dioxo-2,3,5, 7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide.



The title compound was made in two steps using a similar process to that described in example Z-1. 16a (40 mg, 0.09 mmol) and (2*R*)-2-amino-3-methyl-1-butanol (0.1 mL) were reacted in 1,2-dichloroethane (8 mL) with acetic acid to give (3*R*,11a*S*)-*N*-[(2,4-difluorophenyl)methyl]-3-(1-methylethyl)-5,7-dioxo-6-[(phenylmethyl)oxy]-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (41 mg, 92%). This material was hydrogenated in a second step as described in example Z-1 to give (3*R*,11a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-(1-methylethyl)-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (32 mg, 94%) as a white solid. ^{1}H NMR ($CDCl_3$) δ 11.42 (br, 1 H), 10.27 (br, 1 H), 8.34 (s, 1 H), 7.31 (m, 1 H), 6.78 (m, 2 H), 5.28 (d, J = 6.0 Hz, 1 H), 4.60 (m, 2 H), 4.42 (m, 1 H), 4.33 (m, 1 H), 4.16 (m, 1 H), 4.01 (dd, J = 8.8, 5.2 Hz, 1 H), 3.85 (m, 1 H), 2.37 (m, 1 H), 0.97 (d, J = 6.8 Hz, 3 H), 0.95 (d, J = 6.4 Hz, 3 H); ES$^+$ MS: 434 (M+1).

Example Z-26


(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-[2-(methylthio)ethyl]-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide.



The title compound was made in two steps using a similar process to that described in example Z-1. 16a (43 mg, 0.09 mmol) and (2*S*)-2-amino-4-(methylthio)-1-butanol (0.1 mL) were reacted in 1,2-dichloroethane (5 mL) with acetic acid to give (3*S*,11a*R*)-*N*-[(2,4-difluorophenyl)methyl]-3-[2-(methylthio)ethyl]-5,7-dioxo-6-[(phenyl methyl)oxy]-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carbox amide (41 mg, 81%). This material (20 mg, 0.04 mmol) was treated with trifluoroacetic acid (1 mL) in dichloromethane (3 mL) at 0 °C to rt over 6 h. The mixture was concentrated *in vacuo* and subjected to reverse phase preparative HPLC purification to provide (3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-[2-(methylthio)ethyl]-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (12 mg, 72%) as a white solid. ^{1}H NMR ($CDCl_3$) δ 11.35 (br, 1 H), 10.25 (s, 1 H), 8.34 (s, 1 H), 7.33 (m, 1 H), 6.79 (m, 2 H), 5.32 (m, 1 H), 4.62-4.53 (m, 3 H), 4.43-4.39 (m, 2 H), 3.91-3.87 (m, 2 H), 2.63-2.53 (m, 2 H), 2.39 (m, 1 H), 2.12 (s, 3 H), 1.89 (m, 1 H); ES$^+$ MS: 466 (M+1).

Example Z-27


(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-[2-(methylsulfonyl)ethyl]-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide.



To a solution of (3*S*,11a*R*)-*N*-[(2,4-difluorophenyl)methyl]-3-[2-(methylthio)ethyl]-5,7-dioxo-6-[(phenyl methyl)oxy]-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carbox amide (20 mg, 0.04 mmol) in dichloromethane (5 mL) at 0 °C was added *m*-CPBA (20 mg, 70%, 0.082 mmol). The resultant solution was allowed to warm as the bath warmed and stirred a total of 3 h. The reaction was quenched by the addition of $Na_2S_2O_3$ (aq) and sodium bicarbonate. The layers were separated and the organic layer washed with brine. The aqueous layer was extracted with dichloromethane and the combined organics dried over Na_2SO_4. Filtration and concentration provided (3*S*,11a*R*)-*N*-[(2,4-difluorophenyl)methyl]-3-[2-(methylsulfonyl)ethyl]-5,7-dioxo-6-[(phe nylmethyl)oxy]-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-car boxamide (26 mg, 99%) as a white solid. This material was hydrogenated in a second step as described in example Z-1 to give (3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-[2-(methylsulfonyl)ethyl]-5,7-di oxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (22 mg, 99%) as a white solid. ^{1}H NMR ($CDCl_3$) δ 11.00 (br, 1 H), 10.16 (s, 1 H), 8.33 (s, 1 H), 7.36 (m, 1 H), 6.81 (m, 2 H), 5.42 (m, 1 H), 4.62 (m, 3 H), 4.41 (m,2 H), 3.93 (m, 2 H), 3.31 (m, 2 H), 2.98 (s, 3 H), 2.40 (m, 1 H), 2.28 (m, 1 H); ES$^+$ MS: 498 (M+1).

Example Z-28:

(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-(1*H*-indol-3-ylmethyl)-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide.




The title compound was made in two steps using a similar process to that described in example Z-1. 16a (43 mg, 0.09 mmol) and (2*S*)-2-amino-3-(1*H*-indol-3-yl)-1-propanol (100 mg, 0.52 mmol) were reacted in 1,2-dichloroethane (5 mL) with acetic acid to give (3*S*,11a*R*)-*N*-[(2,4-difluorophenyl)methyl]-3-(1*H*-indol-3-ylmethyl)-5,7-dioxo-6-[(phenylmethyl)oxy]-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (36 mg, 64%). This material was hydrogenated in a second step as described in example Z-1 to give (3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-(1*H*-indol-3-ylmethyl)-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (29 mg, 95%) as a white solid. ^{1}H NMR ($CDCl_3/CD_3OD$) δ 10.34 (m, 1 H), 8.98 (br, 1 H), 8.24 (s, 1 H), 7.58 (d, *J* = 8.0 Hz, 1 H), 7.32 (m, 2 H), 7.15-7.01 (m, 3 H), 6.78 (m, 2 H), 4.94 (d, *J* = 6.8 Hz, 1 H), 4.71 (d, *J* = 5.6 Hz, 1 H), 4.59 (m, 2 H), 4.35 (d, *J* = 10.4 Hz, 1 H), 4.22 (m, 1 H), 3.99 (m, 1 H), 3.81 (m, 1 H), 3.40 (dd, *J* = 13.6, 11.6 Hz, 1 H), 3.18 (dd, *J* = 14.0, 8.4 Hz, 1 H); ES$^+$ MS: 521 (M+1).

Example Z-29:

(4*R*,12a*R*)-*N*-[(4-fluorophenyl)methyl]-7-hydroxy-4-methyl-1-(2-methylpropyl)-6,8-diox

o-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamid

e.



a) (2*R*)-2-({[(1,1-Dimethylethyl)oxy]carbonyl}amino)propyl methanesulfonate. To a stirred solution of 1,1-dimethylethyl [(1*R*)-2-hydroxy-1-methylethyl]carbamate (5.00 g, 28.5 mmol) and triethylamine (5.92 mL, 42.9 mmol) in CH_2Cl_2 (30 mL) cooled to 0 °C and under a nitrogen atmosphere was added dropwise a solution of methanesulfonyl chloride (2.43 mL, 31.5 mmol) in CH_2Cl_2 (25 mL). Stirring was continued for 20 minutes at 0 °C, after which time the reaction was judged complete by TLC analysis (1:1 hexanes/EtOAc). The solution was poured into water and the layers were separated. The organic phase was washed with 0.1 N HCl and then with 5% $NaHCO_3$, dried over Na_2SO_4, filtered and concentrated to give (2*R*)-2-({[(1,1-dimethylethyl)oxy]carbonyl}amino)propyl methanesulfonate (7.08 g, 98%) as a white solid. ^1H NMR (400 MHz, $CDCl_3$) δ 1.23 (d, J = 6.8 Hz, 3H), 1.44 (s, 9H), 3.03 (s, 3H), 3.97 (m, 1H), 4.15 (dd, J = 4.2, 9.8 Hz, 1H), 4.21 (m, 1H), 4.61 (br s, 1H).

b) 1,1-Dimethylethyl [(1*R*)-2-cyano-1-methylethyl]carbamate. To a stirred solution of (2*R*)-2-({[(1,1-dimethylethyl)oxy]carbonyl}amino)propyl methanesulfonate (7.08 g, 27.9 mmol) in DMSO (50 mL) was added NaCN (3.78 g, 84.0 mmol). The solution was stirred at 70 °C for 2 hours, over which time the formation of a precipitate was

observed. After cooling at room temperature, water was added and the mixture was extracted with Et_2O. The ethereal layers were washed with a brine solution, dried over Na_2SO_4, filtered and concentrated to give 1,1-dimethylethyl [(1*R*)-2-cyano-1-methylethyl]carbamate (3.81 g, 73%) as a pale yellow solid. ^1H NMR (400 MHz, $CDCl_3$) δ 1.30 (d, J = 6.8 Hz, 3H), 1.42 (s, 9H), 2.51 (dd, J = 3.8, 16.6 Hz, 1H), 2.73 (m, 1H), 3.93 (m, 1H), 4.63 (br s, 1H).

c) 1,1-Dimethylethyl [(1*R*)-3-amino-1-methylpropyl]carbamate. A solution of 1,1-dimethylethyl [(1*R*)-2-cyano-1-methylethyl]carbamate (1.30 g, 7.1 mmol) in ethanol saturated with anhydrous ammonia was treated with Raney-Ni (1.5 mL of 50% aq. Suspension) and 55 psi of H_2 overnight. The mixture was filtered through Celite and the filtrate was concentrated *in vacuo*. The residue was purified by flash chromatography (80:19:1 CH_2Cl_2/MeOH/NH_4OH (37%) gradient elution) through a short plug of silica gel to give 1,1-dimethylethyl [(1*R*)-3-amino-1-methylpropyl]carbamate (1.37 g, 100%) as a clear oil that solidified. ^1H NMR (400 MHz, $CDCl_3$) δ 1.14 (d, J = 6.8 Hz, 3H), 1.43-1.62 (m, 13H), 2.76 (m, 2H), 3.77 (m, 1H), 4.57 (m, 1H).

d) 1,1-Dimethylethyl {(1*R*)-1-methyl-3-[(2-methylpropyl)amino]propyl}carbamate. 1,1-dimethylethyl [(1*R*)-3-amino-1-methylpropyl]carbamate (0.320 g, 1.70 mmol), isobutyraldehyde (150 µL, 1.62 mmol), and sodium triacetoxyborohydride (0.512 g, 2.42 mmol) were stirred in anhydrous dichloroethane (10 mL) at ambient temperature overnight. The reaction was quenched by the addition of saturated $NaHCO_3$ and

then extracted with dichloromethane. The combined extracts were washed with water, dried over Na_2SO_4, filtered and concentrated. The residue was purified by flash chromatography (80:19:1 CH_2Cl_2/MeOH/NH_4OH (37%) gradient elution) through a short plug of silica gel to afford 1,1-dimethylethyl {(1*R*)-1-methyl-3-[(2-methylpropyl)amino]propyl}carbamate (0.158 g, 40%) as a clear oil. ^1H NMR (400 MHz, $CDCl_3$) δ 0.90 (d, *J* = 6.4 Hz, 6H), 1.13 (d, *J* = 6.4 Hz, 3H), 1.42-1.51 (m, 11H), 1.67-1.75 (m, 2H), 2.33-2.42 (m, 2H), 2.58-2.72 (m, 2H), 3.72 (m, 1H), 5.20 (m, 1H).

e) [(3*R*)-3-Aminobutyl](2-methylpropyl)amine. An ice cold solution of 1,1-dimethylethyl {(1*R*)-1-methyl-3-[(2-methylpropyl)amino]propyl}carbamate (0.158 g, 0.65 mmol) in THF (8 mL) was treated with 4 *N* HCl (aq) (2 mL) and then stirred at room temperature for 2 h. The mixture was concentrated *in vacuo* to give [(3*R*)-3-aminobutyl](2-methylpropyl)amine dihydrochloride. The HCl salt was then dissolved in dichloromethane and a minimal amount of methanol and treated with solid supported carbonate resin (MP-Carbonate, Argonaut Technologies). After 30 minutes, the solution was filtered through a fritted tube and the solvents removed carefully *in vacuo* to give [(3*R*)-3-aminobutyl](2-methylpropyl)amine (65 mg). ^1H NMR (400 MHz, $CDCl_3$) δ 0.88 (d, *J* = 6.0 Hz, 6H), 1.06 (d, *J* = 5.6 Hz, 3H), 1.23-1.53 (m, 5H), 1.71-1.74 (m, 1H), 2.39 (m, 2H), 2.65 (m, 2H), 2.97 (m, 1H).


f)

(4*R*,12a*R*)-*N*-[(4-Fluorophenyl)methyl]-7-hydroxy-4-methyl-1-(2-methylpropyl)-6,8-dio

xo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide. The title compound was made in two steps using a similar process to that described in example Z-2. 16 (40 mg, 0.09 mmol) and [(3*R*)-3-aminobutyl](2-methylpropyl)amine (65 mg, 0.45 mmol) were reacted in dichloromethane (2 mL) with acetic acid to give (4*R*,12a*R*)-*N*-[(4-fluorophenyl)methyl]-4-methyl-1-(2-methylpropyl)-6,8-dioxo-7-[(phenylmethyl)oxy]-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide (29 mg, 60%). This material was hydrogenated in a second step as described in example Z-2 to give (4*R*,12a*R*)-*N*-[(4-fluorophenyl)methyl]-7-hydroxy-4-methyl-1-(2-methylpropyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide (18 mg, 75%) as a tan solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 0.77 (d, *J* = 6.4 Hz, 3H), 0.84 (d, *J* = 6.4 Hz, 3H), 1.32 (d, *J* = 7.2 Hz), 1.45-1.49 (m, 1H), 1.57-1.67 (m, 1H), 2.03-2.12 (m, 2H), 2.21-2.27 (m, 1H), 2.73-2.79 (m, 1H), 2.87-2.92 (m, 1H), 4.16-4.24 (m, 2H), 4.45 (s, 1H), 4.54-4.64 (m, 2H), 4.96-4.99 (m, 1H), 6.96-7.00 (m, 2H), 7.29-7.32 (m, 2H), 8.27 (s, 1H), 10.46 (s, 1H), 12.55 (s, 1H); ES$^+$ MS: 456 (M+1).

Example Z-30:

(4*R*,12a*R*)-*N*-[(4-Fluorophenyl)methyl]-7-hydroxy-4-methyl-1-(1-methylethyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide



a) [(3*R*)-3-Aminobutyl](1-methylethyl)amine. The free diamine was prepared in a similar manner as described in example Z-29. ^{1}H NMR (400 MHz, $CDCl_3$) δ 1.04 (d, J = 6.4 Hz, 6H), 1.06 (d, J = 6.4 Hz, 3H), 1.41-1.58 (m, 5H), 2.62-2.66 (m, 2H), 2.74-2.80 (m, 1H), 2.92-3.00 (m, 1H).


b)

(4*R*,12a*R*)-*N*-[(4-Fluorophenyl)methyl]-7-hydroxy-4-methyl-1-(1-methylethyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide. The title compound was made in two steps using a similar process to that described in example Z-2. 16 (40 mg, 0.088 mmol) and [(3*R*)-3-aminobutyl](1-methylethyl)amine (78 mg, 0.60 mmol) were reacted in dichloromethane (2 mL) with acetic acid to give (4*R*,12a*R*)-*N*-[(4-fluorophenyl)methyl]-4-methyl-1-(1-methylethyl)-6,8-dioxo-7-[(phenylmethyl)oxy]-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide (26 mg, 56%). This material was hydrogenated in a second step as described in example Z-2 to give (4*R*,12a*R*)-*N*-[(4-fluorophenyl)methyl]-7-hydroxy-4-methyl-1-(1-methylethyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide (21 mg, 90%) as an off-white solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 1.01 (d, J = 5.6 Hz, 3H), 1.06 (d, J = 6.0 Hz, 3H), 1.31 (d, J = 6.8 Hz, 3H), 1.57 (m, 1H), 1.98 (m, 1H),

2.70-2.82 (m, 2H), 3.15 (m, 1H), 4.15-4.19 (m, 1H), 4.30 (m, 1H), 4.48 (s, 1H), 4.54-4.59 (m, 2H), 4.97 (m, 1H), 6.98 (m, 2H), 7.29-7.32 (m, 2H), 8.27 (s, 1H), 10.49 (s, 1H), 12.52 (s, 1H).

Example Z-31:

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-4-methyl-1-(2-methylpropyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide.



a) 1,1-Dimethylethyl [(1*S*)-2-cyano-1-methylethyl]carbamate. The nitrile was prepared in two steps using a modified procedure as described in example Z-29. To a stirred solution of (2*S*)-2-({[(1,1-dimethylethyl)oxy]carbonyl}amino)propyl methanesulfonate (8.40 g, 33.2 mmol) in DMSO (50 mL) and KCN (6.51 g, 100.0 mmol) cooled to 0 °C was added 18-crown-6 (9.05 g, 34.3 mmol). The solution was allowed to warm to room temperature and then heated to 70 °C for 1 hour. After cooling at room temperature, water was added and the mixture was extracted with Et_2O. The ethereal layers were washed with a brine solution, dried over Na_2SO_4, filtered and concentrated to give 1,1-dimethylethyl [(1*S*)-2-cyano-1-methylethyl]carbamate (5.37 g, 88%) as a pale yellow solid. ^1H NMR (400 MHz, $CDCl_3$) δ 1.32 (d, J = 6.8 Hz, 3H), 1.44 (s, 9H), 2.52 (dd, J = 4.0, 16.4 Hz, 1H), 2.74 (m, 1H), 3.95 (m, 1H), 4.65 (br s, 1H).

b) [(3*S*)-3-Aminobutyl](2-methylpropyl)amine dihydrochloride was prepared in a similar manner as described in example Z-29. ^{1}H NMR (400 MHz, $CDCl_3/CD_3OD$) δ 0.99 (m, 6H), 1.34 (m, 3H), 2.13-2.27 (m, 3H), 2.76 (m, 2H), 3.07 (m, 2H), 3.47 (m, 1H), 8.22 (m, 1 H), 8.83 (m, <1 H).


c)

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-4-methyl-1-(2-methylpropyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide. The title compound was made in two steps using a similar process to that described in example Z-2. 16a (80 mg, 0.17 mmol) and free based [(3*S*)-3-aminobutyl](2-methylpropyl)amine (107 mg, 0.74 mmol) were reacted in dichloromethane (2 mL) with acetic acid to give (4*S*,12a*S*)-*N*-[(2,4-difluorophenyl)methyl]-4-methyl-1-(2-methylpropyl)-6,8-dioxo-7-[(phenylmethyl)oxy]-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide (76 mg, 76%) as a film. This material was hydrogenated in a second step as described in example Z-2 to give (4*S*,12a*S*)-*N*-[(2,4-difluorophenyl)methyl]-7-hydroxy-4-methyl-1-(2-methylpropyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide (39 mg, 80%) as an off-white solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 0.76 (d, J = 6.4 Hz, 3H), 0.84 (d, J = 6.4 Hz, 3H), 1.32 (d, J = 7.2 Hz, 3H), 1.45-1.50 (m, 1H), 1.60-1.69 (m, 1H), 2.03-2.12 (m, 2H), 2.21-2.27 (m, 1H), 2.73-2.79 (m, 1H), 2.87-2.93 (m, 1H), 4.16-4.25 (m, 2H), 4.45 (s, 1H), 4.57-4.68 (m, 2H), 4.96-5.01 (m, 1H), 6.75-6.82 (m, 2H),

7.32-7.38 (m, 1H), 8.26 (s, 1H), 10.45 (s, 1H), 12.56 (s, 1H); ES+ MS: 475 (M+1).

Example Z-32:

(4*S*,12a*S*)-1-(Cyclopropylmethyl)-*N*-[(2,4-difluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide.



a) 1,1-Dimethylethyl {(1*S*)-3-[(cyclopropylmethyl)amino]-1-methylpropyl}carbamate. The protected diamine was prepared using a modified procedure as described in example Z-29. 1,1-dimethylethyl [(1*S*)-3-amino-1-methylpropyl]carbamate (0.293 g, 1.56 mmol), cyclopropane carboxaldehyde (96 µL, 1.30 mmol), and sodium triacetoxyborohydride (0.439 g, 2.07 mmol) were stirred in a 1:1 mixture of anhydrous dichloroethane and tetrahydrofuran (10 mL) at ambient temperature overnight. The reaction was quenched by the addition of saturated $NaHCO_3$ and then extracted with EtOAc. The combined extracts were washed with saturated $NaHCO_3$, then a solution of brine, dried over Na_2SO_4, filtered and concentrated. The residue was purified by flash chromatography (80:19:1 CH_2Cl_2/MeOH/NH_4OH (37%) gradient elution) through a short plug of silica gel to afford 1,1-dimethylethyl {(1*S*)-3-[(cyclopropylmethyl)amino]-1-methylpropyl}carbamate (76 mg, 26%) as a clear oil. ^1H NMR (400 MHz, $CDCl_3$) δ 0.09-0.13 (m, 2H), 0.44-0.49 (m, 2H), 0.92-0.95 (m, 1H), 1.14 (d, J = 6.4 Hz, 3H), 1.43-1.70 (m, 12H), 2.38-2.50 (m, 2H), 2.62-2.73 (m, 2H),

3.74 (m, 1H), 4.88 (m, 1H).

b) [(3*S*)-3-Aminobutyl](cyclopropylmethyl)amine dihydrochloride was prepared in a similar manner as described in example Z-29. ^{1}H NMR (400 MHz, $CDCl_3/CD_3OD$) δ 0.40 (m, 2H), 0.64 (m, 2H), 1.15 (m, 1H), 1.34 (m, 3H), 2.12-2.25 (m, 2H), 2.82 (m, 2H), 3.08 (m, 2H), 3.47 (m, 1H), 8.25 (br, < 1H), 9.04 (br, < 1H),


c)

(4*S*,12a*S*)-1-(Cyclopropylmethyl)-*N*-[(2,4-difluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide. The title compound was made in two steps using a similar process to that described in example Z-2. 16a (50 mg, 0.106 mmol) and free based [(3*S*)-3-aminobutyl](cyclopropylmethyl)amine (44 mg, 0.31 mmol) were reacted in dichloromethane (2 mL) with acetic acid to give (4*S*,12a*S*)-1-(cyclopropylmethyl)-*N*-[(2,4-difluorophenyl)methyl]-4-methyl-6,8-dioxo-7-[(phenylmethyl)oxy]-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide (50 mg, 83%) as a film. This material was hydrogenated in a second step as described in example Z-2 to give (4*S*,12a*S*)-1-(cyclopropylmethyl)-*N*-[(2,4-difluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide (23 mg, 56%) as an off-white solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 0.11 (m, 2H), 0.56-0.59 (m, 2H), 0.77 (m, 1H), 1.34 (d, J = 7.2 Hz, 3H), 1.46-1.50 (m, 1H), 2.04-2.13 (m, 1H), 2.30-2.34 (m, 1H), 2.46-2.51 (m, 1H), 2.90-2.96 (m, 1H), 3.16-3.19 (m, 1H),

4.21-4.30 (m, 2H), 4.51 (s, 1H), 4.58-4.67 (m, 2H), 5.00-5.05 (m, 1H), 6.75-6.82 (m, 2H), 7.31-7.37 (m, 1H), 8.28 (s, 1H), 10.46 (s, 1H), 12.55 (br, 1H); ES$^+$ MS: 473 (M+1).

Example Z-33:

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-1-(2-furanylmethyl)-7-hydroxy-4-methyl-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide.



a) [(3*S*)-3-Aminobutyl](2-furanylmethyl)amine dihydrochloride was prepared in a similar manner as described in example Z-32. ^{1}H NMR (400 MHz, $CDCl_3/CD_3OD$) δ 1.27 (d, J = 6.4 Hz, 3H), 1.96-2.05 (m, 1H), 2.14-2.19 (m, 1H), 3.00-3.04 (m, 2H), 3.38-3.39 (m, 1H), 4.11-4.18 (m, 2H), 6.34 (m, 1H), 6.59 (m, 1H), 7.40 (m, 1H), 8.18 (br, <1 H), 9.41 (br, < 1 H).


b)

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-1-(2-furanylmethyl)-7-hydroxy-4-methyl-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide. The title compound was made in two steps using a similar process to that described in example Z-2. 16a (36 mg, 0.076 mmol) and free based [(3*S*)-3-aminobutyl](2-furanylmethyl)amine (70 mg, 0.42 mmol) were reacted in dichloromethane (2 mL) with acetic acid to give

(4*S*,12a*S*)-*N*-[(2,4-difluorophenyl)methyl]-1-(2-furanylmethyl)-4-methyl-6,8-dioxo-7-[(phenylmethyl)oxy]-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide (32 mg, 70%) as a film. This material was hydrogenated in a second step as described in example Z-2 to give (4*S*,12a*S*)-*N*-[(2,4-difluorophenyl)methyl]-1-(2-furanylmethyl)-7-hydroxy-4-methyl-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide (20 mg, 76%), as an off-white solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 1.24 (d, *J* = 6.8 Hz, 3H), 1.45-1.49 (m, 1H), 2.04-2.13 (m, 1H), 2.77-2.82 (m, 1H), 2.94-3.01 (m, 1H), 3.65 (d, *J* = 15.6 Hz, 1H), 3.89 (d, *J* = 16.0 Hz, 1H), 4.27-4.31 (m, 1H), 4.39-4.41 (m, 1H), 4.49-4.53 (m, 1H), 4.58-4.66 (m, 1H), 4.98-5.03 (m, 1H), 6.24 (m, 1H), 6.36 (m, 1H), 6.75-6.82 (m, 2H), 7.31-7.39 (m, 1H), 7.40 (m, 1H), 8.26 (s, 1H), 10.47 (m, 1H), 12.50 (br, 1H); ES$^+$ MS: 499 (M+1).

Example Z-34:

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1-(1,3-thiazol-2-ylmethyl)-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide.



a) [(3*S*)-3-Aminobutyl](1,3-thiazol-2-ylmethyl)amine dihydrochloride was prepared in a similar manner as described in example Z-32. ^{1}H NMR (400 MHz, $CDCl_3/CD_3OD$) δ 1.28 (d, *J* = 6.4 Hz, 3H), 2.05 (m, 1H), 2.17 (m, 1H), 3.20 (m, 2H), 3.39 (m, 1H),

4.51-4.58 (m, 2H), 7.52 (d, 1H), 7.82 (d, 1H).


b)

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1-(1,3-thiazol-2-ylmethyl)-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide. The title compound was made in two steps using a similar process to that described in example Z-2. 16a (35 mg, 0.074 mmol) and free based [(3*S*)-3-aminobutyl](1,3-thiazol-2-ylmethyl)amine were reacted in dichloromethane (2 mL) with acetic acid to give (4*S*,12a*S*)-*N*-[(2,4-difluorophenyl)methyl]-4-methyl-6,8-dioxo-7-[(phenylmethyl)oxy]-1-(1,3-thiazol-2-ylmethyl)-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide (36 mg, 80%) as a film. This material was debenzylated in a second step to in a manner similar to Z-26 to give (4*S*,12a*S*)-*N*-[(2,4-difluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1-(1,3-thiazol-2-ylmethyl)-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide (18 mg, 60%) as an off-white solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 1.30 (d, J = 7.2 Hz. 3H), 1.49-1.53 (m, 1H), 2.12-2.18 (m, 1H), 2.93-2.96 (m, 1H), 3.07-3.13 (m, 1H), 3.99-4.03 (m, 1H), 4.13-4.17 (m, 1H), 4.24-4.27 (m, 1H), 4.57-4.61 (m, 3H), 5.03-5.06 (m, 1H), 6.75-6.82 (m, 2H), 7.26 (m, 1H), 7.31-7.37 (m, 2H), 7.76 (m, 1H), 7.94 (m, 1H), 10.40 (m, 1H), 12.48 (m, 1H); ES^+ MS: 516 (M+1).

Example Z-35:

racemic-(4a*R*,6a*R*,14a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-12-hydroxy-11,13-dioxo-1,3,4,

4a,5,6a,7,11,13,14a-decahydro-2*H*-pyrido[1',2':4,5]pyrazino[1,2-*a*][3,1]benzoxazine-10-carboxamide



a)

racemic-(4a*R*,6a*R*,14a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-11,13-dioxo-12-[(phenylmethyl)oxy]-1,3,4,4a,5,6a,7,11,13,14a-decahydro-2*H*-pyrido[1',2':4,5]pyrazino[1,2-*a*][3,1]benzoxazine-10-carboxamide. *racemic-cis*-2-Hydroxymethyl-1-cyclohexylamine hydrochloride (24 mg, 0.186 mmol) was dissolved in a dichloromethane solution containing a small amount of methanol (to dissolve) and excess MP-Carbonate (Argonaut Technologies) was added, the mixture was stirred for 30 minutes, and the MP-Carbonate was removed by filtration. The free amine solution was transferred to a microwave vessel containing 16a (29 mg, 0.0617 mmol). One drop of glacial acetic acid was added and the solution was heated for 10 minutes at 140 °C. The resultant solution was absorbed on celite and the material was purified by silica gel chromatography (0-12% methanol/dichloromethane gradient elution) to yield the desired product as a white solid (18 mg, 53%). ^{1}H NMR ($CDCl_3$) δ 10.40 (m, 1 H), 8.35 (s, 1 H), 7.60 (m, 2 H), 7.34-7.26 (m, 4 H), 6.80 (m, 2 H), 5.35-5.23 (m, 2 H), 5.13 (m, 1 H), 4.77 (m, 1 H), 4.70 (m, 2 H), 4.22 (dd, J = 13.2, 3.2 Hz, 1 H), 4.07 (dd, J = 13.2, 6.4, 1 H), 3.96 (m, 1 H), 3.76 (dd, J = 11.2, 4.4, 1 H), 2.22 (m, 1 H), 1.84 (m, 1 H), 1.74-1.40 (m, 6 H), 1.17 (m, 1 H); ES$^+$ MS: 550 (M +1).


b)

racemic-(4a*R*,6a*R*,14a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-12-hydroxy-11,13-dioxo-1,3,4,4a,5,6a,7,11,13,14a-decahydro-2*H*-pyrido[1',2':4,5]pyrazino[1,2-*a*][3,1]benzoxazine-10-carboxamide.

racemic-(4a*R*,6a*R*,14a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-11,13-dioxo-12-[(phenylmethyl)oxy]-1,3,4,4a,5,6a,7,11,13,14a-decahydro-2*H*-pyrido[1',2':4,5]pyrazino[1,2-*a*][3,1]benzoxazine-10-carboxamide (13 mg, 0.0236 mmol) was dissolved in tetrahydrofuran and 10 w.t.% Pd/C (13 mg) was added. Hydrogen was passed through the solution several times and the mixture was stirred at 1 atm hydrogen for 18 hours until the reaction was determined complete by TLC (5% methanol/dichloromethane). The mixture was filtered through Celtite, eluting with methanol/chloroform and the filtrate was concentrated under reduced pressure and purified by HPLC to yield the title compound (7.3 mg, 73%) ^{1}H NMR ($CDCl_3$) δ 12.45 (m, 1 H), 10.38 (s, 1 H), 8.30 (s, 1 H), 7.32 (m, 1 H), 6.83-6.76 (m, 2 H), 5.23 (m, 1 H), 4.75 (m, 1 H), 4.63 (m, 2 H), 4.26 (m, 1 H), 4.12-4.01 (m, 2 H), 3.83 (m, 1 H), 2.30 (m, 1 H), 1.91 (m, 1 H), 1.80 (m, 1 H), 1.67-1.40 (m, 5 H), 1.20 (m, 1 H); ES$^+$ MS: 460 (M +1).

Example Z-36:

racemic-(4a*R*,6a*R*,14a*S*)-*N*-[(4-Fluorophenyl)methyl]-12-hydroxy-11,13-dioxo-1,3,4,4a,5,6a,7,11,13,14a-decahydro-2*H*-pyrido[1',2':4,5]pyrazino[1,2-*a*][3,1]benzoxazine-10-carboxamide.



a)

racemic-(4aR,6aR,14aS)-N-[(4-Fluorophenyl)methyl]-11,13-dioxo-12-[(phenyl methyl)oxy]-1,3,4,4a,5,6a,7,11,13,14a-decahydro-2H-pyrido[1',2':4,5]pyrazino[1,2-a][3, 1]benzoxazine-10-carboxamide. In a manner similar to that described in example Z-35, from *racemic-cis*-2-Hydroxymethyl-1-cyclohexylamine hydrochloride (50 mg, 0.303 mmol) and 16 (45 mg, 0.0995 mmol) was prepared *racemic*-(4aR,6aR,14aS)-N-[(4-fluorophenyl)methyl]-11,13-dioxo-12-[(phenylmethyl)ox y]-1,3,4,4a,5,6a,7,11,13,14a-decahydro-2H-pyrido[1',2':4,5]pyrazino[1,2-a][3,1]benzoxa zine-10-carboxamide (48 mg, 91%) as a white solid. ^{1}H NMR ($CDCl_3$) δ 10.42 (m, 1 H), 8.37 (s, 1 H), 7.59 (m, 2 H), 7.38-7.24 (m, 5 H), 6.98 (m, 2 H), 5.26-5.18 (m, 2 H), 5.07 (m, 1 H), 4.74 (m, 1 H), 4.62-4.51(m, 2 H), 4.20 (dd, J = 13.6, 4 Hz, 1 H), 4.04 (m, 1 H), 3.91 (m, 1 H), 3.71 (dd, J = 11.3, 4.8 Hz, 1 H), 2.18 (m, 1 H), 1.82 (m, 1 H), 1.73-1.63 (m, 2 H), 1.62-1.56 (m, 2 H), 1.48 (, 1 H), 1.38 (m, 1 H), 1.14 (m, 1 H); ES$^+$ MS: 532 (M +1).


b)

racemic-(4a*R*,6a*R*,14a*S*)-N-[(4-Fluorophenyl)methyl]-12-hydroxy-11,13-dioxo-1,3,4,4a, 5,6a,7,11,13,14a-decahydro-2H-pyrido[1',2':4,5]pyrazino[1,2-a][3,1]benzoxazine-10-car boxamide. In a manner similar to that described in example Z-37, from *racemic*-(4aR,6aR,14aS)-N-[(4-fluorophenyl)methyl]-11,13-dioxo-12-[(phenylmethyl)ox

y]-1,3,4,4a,5,6a,7,11,13,14a-decahydro-2H-pyrido[1',2':4,5]pyrazino[1,2-a][3,1]benzoxazine-10-carboxamide (37 mg, 0.0696 mmol) and 10 w.t. % Pd/C (3 mg) was prepared the title compound (18 mg, 58%) as a white solid after purification by HPLC. ^{1}H NMR ($CDCl_3$) δ 12.47 (s, 1 H), 10.39 (m, 1 H), 8.32 (s, 1 H), 7.30 (m, 2 H), 6.98 (m, 2 H), 5.22 (m, 1 H), 4.74 (m, 1 H), 4.58 (m, 2 H), 4.28 (dd, J = 13.2, 4 Hz, 1 H), 4.12-3.98 (m, 2 H), 3.81 (dd, J = 11.6, 4.8 Hz, 1 H), 2.29 (m, 1 H), 1.91-1.19 (m, 8 H); ES$^+$ MS: 442 (M +1).

Example Z-37:

racemic-(3*S*,4a*R*,6a*R*,14a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-12-hydroxy-11,13-dioxo-3-phenyl-1,3,4,4a,5,6a,7,11,13,14a-decahydro-2*H*-pyrido[1',2':4,5]pyrazino[1,2-*a*][3,1]benzoxazine-10-carboxamide.



a)

racemic-(3*S*,4a*R*,6a*R*,14a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-11,13-dioxo-3-phenyl-12-[(phenylmethyl)oxy]-1,3,4,4a,5,6a,7,11,13,14a-decahydro-2*H*-pyrido[1',2':4,5]pyrazino[1,2-*a*][3,1]benzoxazine-10-carboxamide. In a manner similar to that described in example Z-35, from *racemic*-[(1*R*,2*S*,5*S*)-2-amino-5-phenylcyclohexyl]methanol hydrochloride (32 mg, 0.160 mmol) and 16a (30 mg, 0.064 mmol) was prepared *racemic*-(3*S*,4a*R*,6a*R*,14a*S*)-*N*-[(2,4-difluorophenyl)methyl]-11,13-dioxo-3-phenyl-12-[(phenylmethyl)oxy]-1,3,4,4a,5,6a,7,11,13,14a-decahydro-2*H*-pyrido[1',2':4,5]pyrazino[1,

2-*a*][3,1]benzoxazine-10-carboxamide (35 mg, 88%) as a white solid. ^{1}H NMR ($CDCl_3$) δ 10.41 (m, 1 H), 8.38 (s, 1 H), 7.66 (m, 2 H), 7.40-7.26 (m, 6 H), 6.81 (m. 3 H), 5.32-5.25 (m, 2 H), 5.17 (m, 1 H), 4.89 (m, 1 H), 4.66-4.62 (m, 2 H), 4.26 (dd, *J* = 13.6, 4 Hz, 1 H), 4.13-4.04 (m, 2 H), 3.85 (dd, *J*=11.2, 4.4 Hz, 1 H), 2.56 (m, 1 H), 2.37 (m, 1 H), 2.03-1.64 (m, 6 H); ES^+ MS: 626 (M +1).

b)


racemic-(3*S*,4a*R*,6a*R*,14a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-12-hydroxy-11,13-dioxo-3-phenyl-1,3,4,4a,5,6a,7,11,13,14a-decahydro-2*H*-pyrido[1',2':4,5]pyrazino[1,2-*a*][3,1]benzoxazine-10-carboxamide.

racemic-(3*S*,4a*R*,6a*R*,14a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-11,13-dioxo-3-phenyl-12-[(phenylmethyl)oxy]-1,3,4,4a,5,6a,7,11,13,14a-decahydro-2*H*-pyrido[1',2':4,5]pyrazino[1,2-*a*][3,1]benzoxazine-10-carboxamide (27 mg, 0.0432 mmol) was suspended in methanol, 10 w.t. % Pd/C (3 mg) was added and hydrogen was bubbled through the system several times until the reaction was determined complete by TLC (5% methanol/dichloromethane). The suspension was filtered through Celite eluting with methanol/chloroform and the filtrate was concentrated under reduced pressure and purified by HPLC to give the title compound (13 mg, 57%) as a white solid. ^{1}H NMR ($CDCl_3$) δ 12.40 (br s, 1 H), 10.37 (m, 1 H), 8.32 (s, 1 H), 7.37-7.28 (m, 3 H), 7.24-7.15 (m, 4 H), 6.79 (m, 2 H), 5.78 (br s, 1 H), 4.85 (m, 1 H), 4.62 (m, 2 H), 4.29 (m, 1 H), 4.16-4.09 (m, 2 H), 3.92 (dd, *J* = 11.6, 4.8 Hz, 1 H), 2.58 (m, 1 H), 2.46 (m, 1 H), 2.07-1.64 (m, 7 H); ES^+ MS: 536 (M +1).

Example Z-38:

Sodium

racemic-(4aS,6aS,14aS)-10-({[(2,4-difluorophenyl)methyl]amino}carbonyl)-6-(2-methylpropyl)-11,13-dioxo-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazolin-12-olate.



a) *racemic*-1,1-Dimethylethyl [(1S,2R)-2-(hydroxymethyl)cyclohexyl]carbamate. *racemic*-[(1*R*,2*S*,5*S*)-2-Amino-5-phenylcyclohexyl]methanol hydrochloride (800 mg, 4.82 mmol) was dissolved in MeOH (40 mL) and bis(1,1-dimethylethyl) dicarbonate (1.16 g, 5.30 mmol) and triethylamine (4 mL, 28.92 mmol) were added and the mixture was stirred 18 hours at ambient temperature. The solvents were removed under reduced pressure, ethyl acetate and aqueous saturated sodium bicarbonate were added and the product was extracted with ethyl acetate. The combined organics were dried over sodium sulfate and the solvents were removed under reduced pressure. Purification by silica gel chromatography (9:1 hexanes: ethyl acetate to ethyl acetate gradient elution) gave 1,1-dimethylethyl *racemic*-[(1S,2R)-2-(hydroxymethyl)cyclohexyl]carbamate (934 mg, 85%) as a white solid. ^{1}H NMR ($CDCl_3$) δ 4.87 (m, 1H), 4.03-3.95 (m, 2 H), 3.26 (m, 1 H), 3.15 (m, 1 H), 1.73-1.48 (m, 5 H), 1.38 (s, 9 H). 1.27-1.15 (m, 3 H), 0.887 (m, 1 H).

b) *racemic*-1,1-Dimethylethyl [(1S,2R)-2-Formylcyclohexyl]carbamate. To a

solution of dimethylsulfoxide (0.2 mL, 2.88 mmol) in dichloromethane (3 mL) at -78 °C was added oxalyl chloride (0.72 mL, 1.44 mmol) dropwise. The mixture was stirred 10 minutes and *racemic*-1,1-dimethylethyl [(1S,2R)-2-(hydroxymethyl)cyclohexyl]carbamate (220 mg, 0.961 mmol) in dichlormethane was added dropwise and stirred 10 minutes. Triethylamine (0.53 mL, 3.84 mmol) was added slowly and the reaction was stirred at -78 °C for one hour and allowed to warm to ambient temperature. Water was added and product was extracted with dichloromethane. The combined organics were washed with brine and dried over sodium sulfate. Removal of solvents under reduced pressure afforded *racemic*-1,1-dimethylethyl [(1S,2R)-2-formylcyclohexyl]carbamate (223 mg, quantitative) as a yellow oil. ^{1}H NMR ($CDCl_3$) δ 9.61 (s, 1 H), 5.19 (m, 1 H), 3.88 (m, 1 H), 2.61 (m, 1 H), 1.85 (m, 1 H), 1.63-1.49 (m, 4 H), 1.37-1.16 (m, 12 H).

c) *racemic*-1,1-dimethylethyl ((1S,2S)-2-{[(2-Methylpropyl)amino]methyl} cyclohexyl)carbamate. *racemic*-1,1-Dimethylethyl [(1S,2R)-2-formylcyclohexyl]carbamate (223 mg, 0.982 mmol) was dissolved in dichloroethane and 2-methylpropyl)amine (0.15 mL, 1.47 mmol) and sodium triacetoxyborohydride (290 mg, 1.37 mmol) were added and the reaction was stirred at ambient temperature for 18 hours. Aqueous sodium bicarbonate was added and the product was extracted with dichloromethane. The combined extracts were dried over sodium sulfate and the solvents were removed under reduced pressure. Purification by silica gel chromatography (dichloromethane to 1% ammonium hydroxide 19% methanol 80% dichloromethane gradient elution) afforded

racemic-1,1-dimethylethyl ((1S,2S)-2-{[(2-methylpropyl)amino]methyl}cyclohexyl)carbamate (112 mg, 40%) as a clear colorless oil. ^{1}H NMR ($CDCl_3$) δ 6.06 (br s, 1 H), 3.76 (br s, 1 H), 2.63 (m, 1 H), 2.43-2.37 (m, 2 H), 2.25 (m, 1 H), 1.81 (m, 1 H), 1.71-1.59 (m, 3 H), 1.44-1.32 (m, 14 H), 1.27-1.19 (m, 2 H), 0.866 (m, 6 H).

d) *racemic*-(1S,2S)-2-{[(2-Methylpropyl)amino]methyl}cyclohexanamine hydrochloride.

In a manner similar to that describe in example Z-3, step e, from *racemic*-1,1-dimethylethyl ((1S,2S)-2-{[(2-methylpropyl)amino]methyl} cyclohexyl)carbamate (112 mg, 0.394 mmol) was prepared (1S,2S)-2-{[(2-methylpropyl)amino]methyl}cyclohexanamine hydrochloride (130 mg, > 100%) as a white solid. ^{1}H NMR (methanol-d_4/$CDCl_3$) δ 8.68-8.28 (m, 1 H), 3.62 (br s, 1 H), 3.26 (m, 1 H), 2.83-2.78 (m, 3 H), 2.54 (br s, 1 H), 2.12 (m, 1 H), 1.82-1.66 (m, 3 H), 1.53-1.39 (m, 5 H), 0.96 (m, 6 H). 0.766 (m, 1 H).

e)

racemic-(4aS,6aS,14aS)-N-[(2,4-Difluorophenyl)methyl]-6-(2-methylpropyl)-11,13-dioxo-12-[(phenylmethyl)oxy]-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5] pyrazino[1,2-a]quinazoline-10-carboxamide. In a manner similar to that described in Z-35, from *racemic*-(1S,2S)-2-{[(2-methylpropyl)amino]methyl}cyclohexanamine hydrochloride (130 mg, 0.508 mmol) and 16a (55 mg, 0.117 mmol) was prepared

racemic-(4aS,6aS,14aS)-N-[(2,4-difluorophenyl)methyl]-6-(2-methylpropyl)-11,13-dioxo-12-[(phenylmethyl)oxy]-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide (44 mg, 62%) with a 12: 1 d.r. ^{1}H NMR ($CDCl_3$) δ 10.46 (m, 1H), 8.33 (s, 1 H), 7.59 (m, 2 H), 7.37-7.24 (m, 4 H), 6.79 (m, 2 H), 5.30-5.23 (m, 2 H), 4.75-4.56 (m, 3 H), 4.23-4.09 (m, 3 H), 2.69-2.66 (m, 2 H), 2.21-1.98 (m, 3 H), 1.80 (m, 1 H), 1.71-1.33 (m, 6 H), 1.26-1.19 (m, 2 H), 0.810 (m, 3 H), 0.720 (m, 3 H); ES^+ MS: 605 (M +1).

f)

racemic-(4aS,6aS,14aS)-N-[(2,4-Difluorophenyl)methyl]-12-hydroxy-6-(2-methylpropyl)-11,13-dioxo-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide. In a manner similar to that described in example Z-37, from *racemic*-(4aS,6aS,14aS)-N-[(2,4-difluorophenyl)methyl]-6-(2-methylpropyl)-11,13-dioxo-12-[(phenylmethyl)oxy]-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide (39 mg, 0.064 mmol) and 10 w.t. % Pd/C (7 mg) was prepared *racemic*-(4aS,6aS,14aS)-N-[(2,4-difluorophenyl)methyl]-12-hydroxy-6-(2-methylpropyl)-11,13-dioxo-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide (36 mg, > 100%) as a tan solid. ^{1}H NMR ($CDCl_3$) δ 12.60 (br s, 1 H), 10.43 (br s, 1 H), 8.25 (s, 1 H), 7.35 (m, 1 H), 6.78 (m, 2 H), 4.77 (m, 1 H), 4.63 (m, 2 H), 4.49 (br s, 1 H), 4.30-4.13 (m, 2 H), 3.63-3.40 (m, 2 H), 2.88-2.71 (m, 2 H), 2.32-2.21 (m, 2 H), 2.05 (m, 1 H), 1.88-1.11 (m, 7 H), 0.830 (m, 3 H), 0.760 (m, 3

H); AP+ MS: 515 (M +1).


g) Sodium *racemic*-(4aS,6aS,14aS)-10-({[(2,4-Difluorophenyl)methyl]amino}carbonyl)-6-(2-methylpropyl)-11,13-dioxo-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazolin-12-olate. In a manner similar to that described in example Z-1, from *racemic*-(4aS,6aS,14aS)-N-[(2,4-difluorophenyl)methyl]-12-hydroxy-6-(2-methylpropyl)-11,13-dioxo-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide (37 mg, 0.071 mmol) and 1 N sodium hydroxide (0.07 mL) the title compound was prepared as a yellow solid (26 mg, 68 %). ^{1}H NMR (DMSO-d_6) δ 10.73 (m, 1 H), 7.94 (s, 1 H), 7.32 (m, 1 H), 7.19 (m, 1 H), 7.00 (m, 1 H), 4.59-4.41 (m, 3 H), 4.28 (m, 2 H), 4.14 (br s, 1 H), 2.63-2.60 (m, 2 H), 1.98-1.61 (m, 5 H), 1.48-1.36 (m, 4 H), 0.997 (m, 3 H), 0.760 (m, 3 H), 0.660 (m, 2 H); AP+ MS: 515 (M +1 of free acid).


Example Z-39:

(6aR,7aS,11aS)-N-[(2,4-Difluorophenyl)methyl]-1-hydroxy-2,13-dioxo-2,6a,7,7a,8,9,10,11,11a,13-decahydro-6H-pyrido[1',2':4,5]pyrazino[1,2-a]benzimidazole-3-carboxamide

& Example Z-40:

(6aS,7aS,11aS)-N-[(2,4-Difluorophenyl)methyl]-1-hydroxy-2,13-dioxo-2,6a,7,7a,8,9,10,11,11a,13-decahydro-6H-pyrido[1',2':4,5]pyrazino[1,2-a]benzimidazole-3-carboxamide.





a)

(6aR,7aS,11aS)-N-[(2,4-Difluorophenyl)methyl]-2,13-dioxo-1-[(phenylmethyl)o-xy]-2,6a,7,7a,8,9,10,11,11a,13-decahydro-6H-pyrido [1',2':4,5] pyrazino[1,2-a] benzimidazole-3-carboxamide and (6aS,7aS,11aS)-N-[(2,4-difluorophenyl)methyl]-2,13-dioxo-1-[(phenylmethyl)oxy]-2,6a,7,7a,8,9,10,11,11a,13-decahydro-6H-pyrido[1',2':4,5]pyrazino[1,2-a]benzimidazole-3-carboxamide. In a manner similar to that described in example Z-2, from [(1S,2S)-2-aminocyclohexyl]amine (122 mg, 1.07 mmol) and 16a (200 mg, 0.426 mmol) was prepared (6aR,7aS,11aS)-N-[(2,4-difluorophenyl)methyl]-2,13-dioxo-1-[(phenylmethyl)oxy]-2,6a,7,7a,8,9,10,11,11a,13-decahydro-6H-pyrido [1',2':4,5] pyrazino[1,2-a] benzimidazole-3-carboxamide (58 mg) and (6aS,7aS,11aS)-N-[(2,4-difluorophenyl)methyl]-2,13-dioxo-1-[(phenylmethyl)oxy]-2,6a,7,7a,8,9,10,11,11a,13-decahydro-6H-pyrido[1',2':4,5]pyrazino[1,2-a]benzimidazole-3-carboxamide (10.6 mg) after separation of the diastereomers using silica gel chromatography (0-12% methanol/dichloromethane). (6aR,7aS,11aS)-N-[(2,4-difluorophenyl)methyl]-2,13-dioxo-1-[(phenylmethyl)oxy]-2,6a,7,7a,8,9,10,11,11a,13-decahydro-6H-pyrido [1',2':4,5] pyrazino[1,2-a] benzimidazole-3-carboxamide (major): ^{1}H NMR ($CDCl_3$) δ 10.40 (m, 1 H), 8.33 (s, 1 H), 7.57 (m, 2 H), 7.40-7.25 (m, 4 H), 6.81 (m, 2 H), 5.32 (d, J = 10 Hz, 1 H), 5.13 (d, J

= 10 Hz, 1 H), 4.64-4.58 (m, 3 H), 4.21 (dd, J = 12.4, 3.2 Hz, 1 H), 3.79 (m, 1 H), 3.04 (m, 1 H), 2.73 (m, 1 H), 2.53 (m, 1 H), 2.01-1.79 (m, 4 H), 1.36-1.24 (m, 4 H); ES$^+$ MS: 535 (M +1).

(6aS,7aS,11aS)-N-[(2,4-difluorophenyl)methyl]-2,13-dioxo-1-[(phenylmethyl)oxy]-2,6a,7,7a,8,9,10,11,11a,13-decahydro-6H-pyrido[1',2':4,5]pyrazino[1,2-a]benzimidazole-3-carboxamide (minor diastereomer): ^{1}H NMR ($CDCl_3$) δ 10.33 (m, 1 H), 8.28 (s, 1 H), 7.61 (m, 2 H), 7.39-7.28 (m, 3 H), 6.79 (m, 2 H), 5.29 (d, J = 9.6 Hz, 1 H), 5.05 (d, J = 9.6 Hz, 1 H), 4.84 (m, 1 H), 4.60 (m, 2 H), 3.90-3.84 (m, 2 H), 3.07 (m, 1 H), 2.75 (m, 1 H), 2.49 (m, 1 H), 2.07 (m, 1 H), 1.90-1.51 (m, 4 H), 1.33-1.19 (m, 4 H); MS data matches that of its diastereomer.

b) (For example Z-39), (6aR,7aS,11aS)-N-[(2,4-Difluorophenyl)methyl]-1-hydroxy-2,13-dioxo-2,6a,7,7a,8,9,10,11,11a,13-decahydro-6H-pyrido[1',2':4,5]pyrazino[1,2-a]benzimidazole-3-carboxamide. In a manner similar to that described in example Z-37, from the minor diastereomer prepared in step a (6aS,7aS,11aS)-N-[(2,4-difluorophenyl)methyl]-2,13-dioxo-1-[(phenylmethyl)oxy]-2,6a,7,7a,8,9,10,11,11a,13-decahydro-6H-pyrido[1',2':4,5]pyrazino[1,2-a]benzimidazole-3-carboxamide (7 mg, 0.0131 mmol) and 10 w.t. % Pd/C (catalytic amount) was prepared (6aR,7aS,11aS)-N-[(2,4-difluorophenyl)methyl]-1-hydroxy-2,13-dioxo-2,6a,7,7a,8,9,10,11,11a,13-decahydro-6H-pyrido[1',2':4,5]pyrazino[1,2-a]benzimidazole-3-carboxamide (2.8 mg, 48%) after purification by HPLC. ^{1}H NMR ($CDCl_3$) δ 12.15 (br s, 1 H), 10.42 (br s, 1 H), 8.31 (s, 1 H), 7.36 (m, 1 H), 6.80 (m, 2 H), 5.01 (m, 1 H), 4.63 (m, 2


H), 4.16 (m, 1 H), 3.96 (m, 1H), 3.06-2.93 (m, 2 H), 2.61 (m, 1 H), 2.18 (m, 1 H), 1.93 (m, 1 H), 1.60-1.13 (m, 4 H), 0.893-0.840 (m, 2 H); ES^+ MS: 445 (M +1).

c) (For example Z-40).

(6aS,7aS,11aS)-N-[(2,4-Difluorophenyl)methyl]-1-hydroxy-2,13-dioxo-2,6a,7,7a,8,9,10,11,11a,13-decahydro-6H-pyrido[1',2':4,5]pyrazino[1,2-a]benzimidazole-3-carboxamide. In a manner similar to that described in example Z-37, from the major diastereomer (30 mg, 0.0561 mmol) prepared in step a and 10 w.t. % Pd/C (catalytic amount), (6aS,7aS,11aS)-N-[(2,4-Difluorophenyl)methyl]-1-hydroxy-2,13-dioxo-2,6a,7,7a,8,9,10,11,11a,13-decahydro-6H-pyrido[1',2':4,5]pyrazino[1,2-a]benzimidazole-3-carboxamide was prepared as a white solid (15 mg, 60%) after purification by HPLC. ^1H NMR (methanol-d_4/$CDCl_3$) δ 10.41 (m, 1 H), 8.25 (s, 1 H), 7.30 (m, 1 H), 6.77 (m, 2 H), 4.77 (m, 1 H), 4.57 (m, 2 H), 4.45 (m, 1 H), 3.91 (m, 1 H), 3.12 (m, 1 H), 2.67 (m, 1 H), 2.12 (m, 1 H), 1.87-1.84 (m, 2 H), 1.47-1.33 (m, 4 H); ES^+ MS: 445 (M +1).

Example Z-41:

(5aS,14aS)-N-[(2,4-Difluorophenyl)methyl]-11-hydroxy-10,12-dioxo-1,2,3,4,5a,6,10,12,14,14a-decahydropyrido[1,2-a]pyrido[1',2':3,4]imidazo[1,2-d]pyrazine-9-carboxamide.



a)

(5aS,14aS)-N-[(2,4-Difluorophenyl)methyl]-10,12-dioxo-11-[(phenylmethyl)oxy]-1,2,3,4

,5a,6,10,12,14,14a-decahydropyrido[1,2-a]pyrido[1',2':3,4]imidazo[1,2-d]pyrazine-9-carboxamide. In a manner similar to that described in example Z-18, from 16a (50 mg, 0.108 mmol) and [(2S)-2-piperidinylmethyl]amine hydrochloride (50 mg, 0.269 mmol, made in a similar manner as described in example Z-18) was prepared (5aS,14aS)-N-[(2,4-difluorophenyl)methyl]-10,12-dioxo-11-[(phenylmethyl)oxy]-1,2,3,4,5a,6,10,12,14,14a-decahydropyrido[1,2-a]pyrido[1',2':3,4]imidazo[1,2-d]pyrazine-9-carboxamide (40 mg, 78 %). ^{1}H NMR ($CDCl_3$) δ 10.43 (m, 1 H), 8.38 (s, 1 H), 7.59 (m, 2 H), 7.59-7.25 (m, 4 H), 6.81 (m, 2 H), 5.38 (d, J = 10 Hz, 1 H), 5.19 (d, J = 10 Hz, 1 H), 4.65-4.62 (m, 2 H), 4.20 (dd, J = 12, 2.8 Hz, 1 H), 4.00 (dd, J = 12.4, 2.8 Hz, 1 H), 3.85 (m, 1 H), 3.74 (m, 1 H), 3.27 (m, 1 H), 2.99 (m, 1 H), 2.43 (m, 1 H), 2.24 (m, 1 H), 1.94-1.87 (m, 2 H), 1.77-1.58 (m, 2 H), 1.39-1.24 (m, 2 H); ES$^+$ MS: 535 (M +1).


b)

(5aS,14aS)-N-[(2,4-Difluorophenyl)methyl]-11-hydroxy-10,12-dioxo-1,2,3,4,5a,6,10,12,14,14a-decahydropyrido[1,2-a]pyrido[1',2':3,4]imidazo[1,2-d]pyrazine-9-carboxamide. In a manner similar to that described in example Z-37, from (5aS,14aS)-N-[(2,4-difluorophenyl)methyl]-10,12-dioxo-11-[(phenylmethyl)oxy]-1,2,3,4,5a,6,10,12,14,14a-decahydropyrido[1,2-a]pyrido[1',2':3,4]imidazo[1,2-d]pyrazine-9-carboxamide (18 mg, 0.0337 mmol) and 10 w.t.% Pd/C (catalytic amount) was prepared the title compound as a white solid (13 mg, 87%) after purification by HPLC. ^{1}H NMR ($CDCl_3$) δ 11.71 (br s, 1 H), 10.36 (br s, 1 H), 8.31 (s, 1 H), 7.34 (m, 1 H), 6.78 (m, 2 H), 4.64-4.57 (m, 2 H), 4.28 (m, 1 H), 4.12 (m, 1 H), 3.92-3.89 (m, 2 H), 3.22 (m, 1 H), 3.04 (m, 1 H), 2.49 (m, 1 H), 2.28 (m, 1 H), 1.97-1.89 (m, 2 H), 1.78 (m, 1 H),

1.66-1.60 (m, 2 H), 1.43-1.36 (m, 2 H); ES+ MS: 445 (M +1).

Example Z-42:

(4aR,14aR)-N-[(2,4-Difluorophenyl)methyl]-9-hydroxy-8,10-dioxo-2,3,4,4a,5,6,8,10,14,14a-decahydro-1H-pyrido[1,2-c]pyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-11-carboxamide.



a) Phenylmethyl (2R)-2-(hydroxymethyl)-1-piperidinecarboxylate. In a manner similar to that described in example Z-3a, from (2R)-1-{[(phenylmethyl)oxy]carbonyl}-2-piperidinecarboxylic acid (4.93 g, 18.75 mmol) was prepared phenylmethyl (2R)-2-(hydroxymethyl)-1-piperidinecarboxylate (2.24 g, 48%) as an oil that solidified upon standing to a white solid. ^{1}H NMR ($CDCl_3$) δ 7.36-7.26 (m, 5 H), 5.18-5.10 (m, 2 H), 4.37 (m, 1 H), 4.03 (m, 1 H), 3.84 (, m, 1 H), 3.63 (m, 1 H), 2.96 (br s, 1 H), 1.71-1.42 (m, 6 H).

b) Phenylmethyl (2R)-2-(cyanomethyl)-1-piperidinecarboxylate. In a manner similar to that described in example Z-3b, from phenylmethyl (2R)-2-(hydroxymethyl)-1-piperidinecarboxylate (1.09g, 4.38 mmol) was prepared phenylmethyl (2R)-2-({[(4-methylphenyl)sulfonyl]oxy}methyl)-1-piperidinecarboxylate (1.05g, 59% impure with uncharacterized byproduct) as a clear colorless oil after purification using silica gel chromatography (10-100% ethyl acetate-hexanes). It is

necessary to use this material in the next step as soon as possible or yields deteriorate dramatically. In a manner similar to that described in example Z-3c, from phenylmethyl (2R)-2-({[(4-methylphenyl)sulfonyl]oxy}methyl)-1-piperidinecarboxylate (1.05 g, 2.61 mmol) and sodium cyanide (383 mg, 7.82 mmol) was prepared phenylmethyl (2R)-2-(cyanomethyl)-1-piperidinecarboxylate (171 mg, 25 %) as a yellow oil. ^{1}H NMR ($CDCl_3$) δ 7.35-7.29 (m, 5 H), 5.13 (s, 2 H), 4.65 (m, 1 H), 4.10 (m, 1 H), 2.96 (m, 1 H), 2.60 (m, 2 H), 1.82-1.67 (m, 4 H), 1.54-1.39 (m, 2 H).

d) Phenylmethyl (2R)-2-(2-aminoethyl)-1-piperidinecarboxylate. In a manner similar to that described in example Z-3d, from phenylmethyl (2R)-2-(cyanomethyl)-1-piperidinecarboxylate (171 mg, 0.663 mmol) was prepared phenylmethyl (2R)-2-(2-aminoethyl)-1-piperidinecarboxylate (119 mg, 68%) as a clear colorless residue. ^{1}H NMR ($CDCl_3$) δ 7.32-7.25 (m, 5 H), 5.08 (m, 2 H), 4.39 (br s, 1 H), 4.01 (br s, 1 H), 2.78 (m, 1 H), 2.60-2.56 (m, 2 H), 1.95-1.86 (m, 3 H), 1.63-1.35 (m, 6 H).

e) {2-[(2R)-2-Piperidinyl]ethyl}amine. Phenylmethyl (2R)-2-(2-aminoethyl)-1-piperidinecarboxylate (119 mg, 0.454 mmol) was dissolved in methanol and 10 w.t.% Pd/C (120 mg) was added. Hydrogen was bubbled through the solution for 15 minutes and the reaction was stirred under 1 atm hydrogen for 18 hours until determined complete by TLC (1% ammonium hydroxide 19% methanol 80% dichloromethane). The suspension was filtered through Celite eluting with

methanol and the filtrate was carefully concentrated under reduce pressure to yield a clear colorless liquid (58 mg, quantitative). ^{1}H NMR ($CDCl_3$) δ 2.99 (m, 1 H), 2.71-2.66 (m, 2 H), 2.57-2.48 (m, 2 H), 1.72 (m, 1 H), 1.61-1.52 (m, 2 H), 1.48-1.42 (m, 2 H), 1.35-1.25 (m, 2 H), 1.05 (m, 1 H).

f)

(4aR,14aR)-N-[(2,4-Difluorophenyl)methyl]-8,10-dioxo-9-[(phenylmethyl)oxy]-2,3,4,4a,5,6,8,10,14,14a-decahydro-1H-pyrido[1,2-c]pyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-11-carboxamide. In a manner similar to that described in example Z-35, from 16a (50 mg, 0.106 mmol) and {2-[(2R)-2-piperidinyl]ethyl}amine (58 mg, 0.454 mmol) was prepared (4aR,14aR)-N-[(2,4-difluorophenyl)methyl]-8,10-dioxo-9-[(phenylmethyl)oxy]-2,3,4,4a,5,6,8,10,14,14a-decahydro-1H-pyrido[1,2-c]pyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-11-carboxamide (47 mg, 81%). ^{1}H NMR ($CDCl_3$) δ 10.50 (br s, 1 H), 8.33 (s, 1 H), 7.60 (s, 2 H), 7.38-7.24 (m, 4 H), 6.80 (m, 2 H), 5.29-5.22 (m, 2 H), 4.66-4.56 (m, 3 H), 4.30 (m, 1 H), 4.19 (m, 1 H), 3.78 (br s, 1 H), 2.86-2.80 (m, 2 H), 2.18 (br s, 1 H), 1.94 (m, 1 H), 1.68-1.36 (m, 6 H), 1.23 (br s, 2 H); ES$^+$ MS: 549 (M +1).


g)

(4aR,14aR)-N-[(2,4-Difluorophenyl)methyl]-9-hydroxy-8,10-dioxo-2,3,4,4a,5,6,8,10,14,14a-decahydro-1H-pyrido[1,2-c]pyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-11-carboxamide. In a manner similar to that described in example Z-37, from (4aR,14aR)-N-[(2,4-difluorophenyl)methyl]-8,10-dioxo-9-[(phenylmethyl)oxy]-2,3,4,4a,

5,6,8,10,14,14a-decahydro-1H-pyrido[1,2-c]pyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-1 1-carboxamide (47 mg, 0.0857 mmol) and a catalytic amount of 10 w.t.% Pd/C was prepared the title compound as a white solid (19 mg, 54%) after purification by HPLC. ^{1}H NMR ($CDCl_3$) δ 10.49 (m, 1 H), 8.29 (s, 1 H), 7.34 (m, 1 H), 6.79 (m, 2 H), 4.67-4.56 (m, 3 H), 4.41 (m, 1 H), 4.20 (m, 1 H), 3.93 (s, 1 H), 2.94-2.87 (m, 2 H), 2.28 (br s, 1 H), 2.01 (m, 1 H), 1.68-1.54 (m, 4 H), 1.44 (m, 1 H), 1.29-1.23 (m, 3 H), 0.850 (m, 1 H); ES$^+$ MS: 459 (M +1).

Example Z-43:

(4*R*,12a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-4-methyl-1-(3-methylbutyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide.



a) [(3*R*)-3-Aminobutyl](3-methylbutyl)amine dihydrochloride was prepared in a similar manner as described in example Z-32. ^{1}H NMR (400 MHz, $CDCl_3/CD_3OD$) δ 0.87 (d, J = 5.2 Hz, 6H), 1.32 (m, 3H), 1.61 (m, 3H), 2.10-2.20 (m, 2H), 2.90-3.04 (m, 4H), 3.45 (m, 1H), 8.23 (br, < 1 H), 8.96 (br, < 1 H).


b)

(4*R*,12a*R*)-*N*-[(2,4-Dfluorophenyl)methyl]-7-hydroxy-4-methyl-1-(3-methylbutyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxam

ide. The title compound was made in two steps using a similar process to that described in example Z-2. 16a (40 mg, 0.085 mmol) and free [(3*R*)-3-aminobutyl](3-methylbutyl)amine (46 mg, 0.35 mmol) were reacted in dichloromethane (2 mL) with acetic acid to give (4*R*,12a*R*)-*N*-[(2,4-difluorophenyl)methyl]-4-methyl-1-(3-methylbutyl)-6,8-dioxo-7-[(phenylmethyl)oxy]-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide (44 mg, 90%) as a film. This material was hydrogenated in a second step as described in example Z-2 to give (4*R*,12a*R*)-*N*-[(2,4-difluorophenyl)methyl]-7-hydroxy-4-methyl-1-(3-methylbutyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide (11 mg, 30%) as an off-white solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 0.84 (d, *J* = 6.8 Hz, 3H), 0.86 (d, *J* = 6.8 Hz, 3H), 1.24-1.36 (m, 5H), 1.47-1.53 (m, 2H), 2.02-2.11 (m, 1H), 2.36-2.43 (m, 1H), 2.54-2.61 (m, 1H), 2.77-2.92 (m, 2H), 4.16-4.26 (m, 2H), 4.44 (m, 1H), 4.62-4.64 (m, 2H), 4.95-5.02 (m, 1H), 6.75-6.81 (m, 2H), 7.31-7.37 (m, 1H), 8.27 (s, 1H), 10.43 (m, 1H), 12.54 (s, 1H); ES$^+$ MS: 489 (M+1).

Example Z-44:

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-4-methyl-1-(1-methylethyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide.



a) [(3*S*)-3-Aminobutyl](1-methylethyl)amine dihydrochloride was prepared in a similar manner as described in example Z-29. ^{1}H NMR (400 MHz, $CDCl_3/CD_3OD$) δ 1.20-1.25 (m, 9H), 1.93-2.02 (m, 2H), 2.92 (m, 2H), 3.20-3.29 (m, 2H), 8.04 (br, < 1 H), 8.64 (br, < 1 H).


b)

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-4-methyl-1-(1-methylethyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide. The title compound was made in two steps using a similar process to that described in example Z-2. 16a (60 mg, 0.13 mmol) and free based [(3*S*)-3-aminobutyl](1-methylethyl)amine (55 mg, 0.42 mmol) were reacted in dichloromethane (2 mL) with acetic acid to give (4*S*,12a*S*)-*N*-[(2,4-difluorophenyl)methyl]-4-methyl-1-(1-methylethyl)-6,8-dioxo-7-[(phenylmethyl)oxy]-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide (40 mg, 57%) as a film. This material was hydrogenated in a second step as described in example Z-2 to give (4*S*,12a*S*)-*N*-[(2,4-difluorophenyl)methyl]-7-hydroxy-4-methyl-1-(1-methylethyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide (17 mg, 50%) as an off-white solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 1.02 (d, J = 6.4 Hz, 3H), 1.07 (d, J = 6.4 Hz, 3H), 1.33 (d, J = 7.2 Hz, 3H), 1.55-1.58 (m, 1H), 1.94-2.03 (m, 1H), 2.70-2.77 (m, 1H), 2.81-2.86 (m, 1H), 3.11-3.18 (m, 1H), 4.17 (dd, J = 3.0, 13.8 Hz, 1H), 4.32 (dd, J = 3.2, 14.0 Hz, 1H), 4.48 (m, 1H), 4.59-4.69 (m, 2H), 4.97-5.00 (m, 1H), 6.77-6.83 (m, 2H), 7.33-7.39 (m, 1H), 8.28 (s, 1H), 10.50 (m, 1H), 12.55 (s, 1H);

ES+ MS: 461 (M+1).

Example Z-45:

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-4-methyl-1-(3-methylbutyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide.



a) [(3*S*)-3-Aminobutyl](3-methylbutyl)amine dihydrochloride was prepared in a similar manner as described in example Z-32. ^{1}H NMR (400 MHz, $CDCl_3/CD_3OD$) δ 0.86 (d, J = 5.6 Hz, 6H), 1.27 (d, J = 6.0 Hz, 3H), 1.58 (m, 3H), 2.03-2.14 (m, 2H), 2.87-2.99 (m, 4H), 3.38 (m, 1H), 8.15 (br, < 1 H), 8.87 (br, < 1 H).


b)

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-4-methyl-1-(3-methylbutyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide. The title compound was made in two steps using a similar process to that described in example Z-2. 16a (0.100 g, 0.21 mmol) and free based [(3*S*)-3-aminobutyl](3-methylbutyl)amine (0.104 g, 0.66 mmol) were reacted in dichloromethane (2 mL) with acetic acid to give (4*S*,12a*S*)-*N*-[(2,4-difluorophenyl)methyl]-4-methyl-1-(3-methylbutyl)-6,8-dioxo-7-[(phenylmethyl)oxy]-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine

-9-carboxamide (88 mg, 72%) as a film. This material was hydrogenated in a second step as described in example Z-2 to give (4*S*,12a*S*)-*N*-[(2,4-difluorophenyl)methyl]-7-hydroxy-4-methyl-1-(3-methylbutyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide (55 mg, 74%). ^{1}H NMR (400 MHz, $CDCl_3$) δ 0.84 (d, *J* = 6.4 Hz, 3H), 0.85 (d, *J* = 6.4 Hz, 3H), 1.24-1.37 (m, 5H), 1.45-1.53 (m, 2H), 2.02-2.11 (m, 1H), 2.37-2.44 (m, 1H), 2.56-2.63 (m, 1H), 2.80-2.92 (m, 2H), 4.22-4.29 (m, 2H), 4.45 (s, 1H), 4.62-4.63 (m, 2H), 4.97-5.00 (m, 1H), 6.75-6.82 (m, 2H), 7.31-7.37 (m, 1H), 8.37 (s, 1H), 10.48 (m, 1H), 12.53 (br, 1H); ES$^+$ MS: 489 (M+1).

Example Z-46:

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1-(3-pyridinylmethyl)-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide.



a) 1,1-Dimethylethyl {(1*S*)-1-methyl-3-[(3-pyridinylmethyl)amino]propyl}carbamate. The protected diamine was prepared using a modified procedure as described in example Z-32. A solution of 1,1-dimethylethyl [(1*S*)-3-amino-1-methylpropyl]carbamate (0.296 g, 1.6 mmol) and 3-pyridinecarboxaldehyde (120 µL, 1.3 mmol) in a 1:1 mixture of anhydrous dichloroethane and tetrahydrofuran (10 mL) was treated with acetic acid (374 µL, 6.6

mmol) and stirred for 30 minutes. Sodium triacetoxyborohydride (0.444 g, 2.1 mmol) was added and the solution was stirred for 2 hours. The resultant was subjected to a workup and purification procedure as described in example Z-32 to give 1,1-dimethylethyl {(1*S*)-1-methyl-3-[(3-pyridinylmethyl)amino]propyl}carbamate (0.245 g, 66%) as a clear oil. ^{1}H NMR (400 MHz, $CDCl_3$) δ 1.12 (d, *J* = 6.4 Hz, 3H), 1.42 (s, 9H), 1.46-1.54 (m, 1H), 1.68 (m, 1H), 2.61-2.75 (m, 2H), 3.73- 3.80 (m, 3H), 4.86 (m, 1H), 7.22-7.24 (m, 1H), 7.68 (d, *J* = 8.0 Hz, 1H), 8.48 (m, 1H), 8.53 (m, 1H).

b) [(3*S*)-3-Aminobutyl](3-pyridinylmethyl)amine dihydrochloride was prepared in a similar manner as described in example Z-29.


c)

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1-(3-pyridinyl methyl)-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carbo xamide. The title compound was made in two steps using a similar process to that described in example Z-2. 16a (60 mg, 0.13 mmol) and free based [(3*S*)-3-aminobutyl](3-pyridinylmethyl)amine (83 mg, 0.47 mmol) were reacted in dichloromethane (2 mL) with acetic acid to give (4*S*,12a*S*)-*N*-[(2,4-difluorophenyl)methyl]-4-methyl-6,8-dioxo-7-[(phenylmethyl)oxy]-1-(3-pyridinylmethyl)-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimi dine-9-carboxamide (72 mg, 95%) as a film. This material was hydrogenated in a second step as described in example Z-2 to give (4*S*,12a*S*)-*N*-[(2,4-difluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1-(3-pyridinyl

methyl)-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide (34 mg, 56%) as an off-white solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 1.37 (d, J = 6.8 Hz, 3H), 1.43-1.47 (m, 1H), 2.12 (m, 1H), 2.60-2.92 (m, 2H), 3.53 (d, J = 14.0 Hz, 1H), 3.82 (d, J = 14.4 Hz, 1H), 4.23-4.31 (m, 2H), 4.55-4.64 (m, 3H), 5.06-5.11 (m, 1H), 6.75-6.82 (m, 2H), 7.20-7.23 (m, 1H), 7.31-7.36 (m, 1H), 7.50 (m, 1H), 7.92 (s, 1H), 8.48 (s, 1H), 10.39 (m, 1H), 12.5 (br, 1H); ES$^+$ MS: 510 (M+1).

Example Z-47:

(4*S*,12a*S*)-1-Cyclopropyl-*N*-[(2,4-difluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide.



a) 1,1-Dimethylethyl [(1*S*)-1-methyl-3-oxopropyl]carbamate. To a stirred solution of 1,1-dimethylethyl [(1*S*)-2-cyano-1-methylethyl]carbamate (0.656 g, 3.56 mmol) in anhydrous ether cooled to -40 °C was added dropwise a 1.0 M solution of diisobutylaluminum hydride in hexanes (14.2 mL, 14.2 mmol) over 20 minutes. Stirring was continued at this temperature for an additional 20 minutes. The yellow solution was quenched with Rochelle's salt and the resultant stirred at room temperature for 1 hour. The solids were filtered off through celite and rinsed with EtOAc. The organics were washed with brine, concentrated, and flash chromatographed (10-100% EtOAc/hexanes) to give 1,1-dimethylethyl [(1*S*)-1-methyl-3-oxopropyl]carbamate (0.193 g, 30 %) as a clear oil. ^{1}H NMR (400

MHz, $CDCl_3$) δ 1.22 (d, J = 6.8 Hz, 3H), 1.41 (s, 9H), 2.53-2.65 (m, 2H), 4.08-4.13 (m, 1H), 4.63 (m, 1H), 9.74-9.75 (m, 1H).

b) 1,1-Dimethylethyl [(1*S*)-3-(cyclopropylamino)-1-methylpropyl]carbamate. The protected diamine was prepared using a modified procedure as described in example Z-32. A solution of 1,1-dimethylethyl [(1*S*)-1-methyl-3-oxopropyl]carbamate (0.178 g, 0.95 mmol) and cyclopropylamine (197 µL, 2.85 mmol) in anhydrous dichloroethane (10 mL) was treated with acetic acid (272 µL, 4.8 mmol) and stirred for 30 minutes. Sodium triacetoxyborohydride (0.444 g, 2.1 mmol) was added and the solution was stirred for 20 hours. The resultant was subjected to a workup and purification procedure as described in example Z-32 to give 1,1-dimethylethyl [(1*S*)-3-(cyclopropylamino)-1-methylpropyl]carbamate (0.136 g, 63%) as a clear oil. ^{1}H NMR (400 MHz, $CDCl_3$) δ 0.32-0.42 (m, 4H), 1.12 (d, J = 6.8 Hz, 3H), 1.39-1.51 (m, 10H), 1.58-1.92 (m, 2H), 2.05-2.10 (m, 1H), 2.67-2.80 (m, 2H), 3.71 (m, 1H), 4.78 (m, 1H).

c) [(3*S*)-3-Aminobutyl]cyclopropylamine dihydrochloride was prepared in a similar manner as described in example Z-29. ^{1}H NMR (400 MHz, $CDCl_3/CD_3OD$) δ 0.70-0.75 (m, 2H), 0.90-0.94 (m, 2H), 1.18 (d, J = 6.8 Hz, 3H), 1.84-1.94 (m, 1H), 1.97-2.05 (m, 1H), 2.49-2.54 (m, 1H), 2.99-3.04 (m, 2H), 3.23-3.28 (m, 1H).


d)

(4*S*,12a*S*)-1-Cyclopropyl-*N*-[(2,4-difluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-

1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide. The title compound was made in two steps using a similar process to that described in example Z-2. 16a (80 mg, 0.17 mmol) and free based [(3*S*)-3-aminobutyl]cyclopropylamine (75 mg, 0.59 mmol) were reacted in dichloromethane (2 mL) with acetic acid to give (4*S*,12a*S*)-1-cyclopropyl-*N*-[(2,4-difluorophenyl)methyl]-4-methyl-6,8-dioxo-7-[(phenylmethyl)oxy]-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide (74 mg, 80%) as a film. This material was hydrogenated in a second step as described in example Z-2 to give (4*S*,12a*S*)-1-cyclopropyl-*N*-[(2,4-difluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide (32 mg, 52%) as an off-white solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 0.37-0.54 (m, 3H), 0.64-0.70 (m, 1H), 1.35 (d, *J* = 7.2 Hz, 3H), 1.45-1.49 (m, 1H), 1.76-1.80 (m, 1H), 2.03-2.12 (m, 1H), 2.86-2.93 (m, 1H), 2.99-3.04 (m, 1H), 4.30 (dd, *J* = 4.0, 13.6 Hz, 1H), 4.49-4.67 (m, 4H), 5.00-5.07 (m, 1H), 6.75-6.82 (m, 2H), 7.32-7.36 (m, 1H), 8.28 (s, 1H), 10.49 (m, 1H), 12.53 (s, 1H); ES$^+$ MS: 459 (M+1).

Example Z-48:

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-4-methyl-1-[2-(methyloxy)ethyl]-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide.



a) [(3*S*)-3-Aminobutyl][2-(methyloxy)ethyl]amine dihydrochloride. The protected diamine, 1,1-dimethylethyl ((1*S*)-1-methyl-3-{[2-(methyloxy)ethyl]amino}propyl)carbamate was prepared in a similar manner as described in example Z-47. Subsequently, [(3*S*)-3-aminobutyl][2-(methyloxy)ethyl]amine dihydrochloride was prepared in a similar manner as described in example Z-29. ^{1}H NMR (400 MHz, $CDCl_3/CD_3OD$) δ 1.21 (d, J = 5.6 Hz, 3H), 1.93 (m, 1H), 2.04 (m, 1H), 2.98-3.05 (m, 4H), 3.22 (m, 2H), 3.26-3.31 (m, 4H), 8.06 (br, < 1 H), 8.81 (br, < 1 H).


b)

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-4-methyl-1-[2-(methyloxy)ethyl]-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide. The title compound was made in two steps using a similar process to that described in example Z-2. 16a (60 mg, 0.13 mmol) and free based [(3*S*)-3-aminobutyl][2-(methyloxy)ethyl]amine (53 mg, 0.37 mmol) were reacted in dichloromethane (2 mL) with acetic acid to give (4*S*,12a*S*)-*N*-[(2,4-difluorophenyl)methyl]-4-methyl-1-[2-(methyloxy)ethyl]-6,8-dioxo-7-[(phenylmethyl)oxy]-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide (47 mg, 63%) as a film. This material was hydrogenated in a

second step as described in example Z-2 to give (4*S*,12a*S*)-*N*-[(2,4-difluorophenyl)methyl]-7-hydroxy-4-methyl-1-[2-(methyloxy)ethyl]-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide (38 mg, 97%) as an off-white solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 1.34 (d, *J* = 7.2 Hz, 3H), 1.49 (m, 1H), 2.03-2.12 (m, 1H), 2.67-2.70 (m, 1H), 2.81-2.92 (m, 2H), 3.06-3.15 (m, 1H), 3.30-3.37 (m, 4H), 3.58-3.63 (m, 1H), 4.20 (dd, *J* = 3.4, 14.2 Hz, 1H), 4.50-4.59 (m, 1H), 4.62-4.65 (m, 3H), 5.00-5.03 (m, 1H), 6.75-6.81 (m, 2H), 7.31-7.37 (m, 1H), 8.27 (s, 1H), 10.46 (s, 1H), 12.54 (s, 1H); ES$^+$ MS: 477 (M+1).

Example Z-49:

racemic-(3aS,5aS,13aS)-N-[(2,4-Difluorophenyl)methyl]-11-hydroxy-5-(2-methylpropyl)-10,12-dioxo-2,3,3a,4,5,5a,6,10,12,13a-decahydro-1H-cyclopenta[e]pyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-9-carboxamide.



a) *racemic*-(1S,2S)-2-{[(2-Methylpropyl)amino]methyl}cyclopentanamine hydrochloride.

In a manner similar to example Z-18a-c, from *racemic*-(1R,2S)-2-({[(1,1-dimethylethyl)oxy]carbonyl}amino)cyclopentanecarboxylic acid (255 mg, 1.11 mmol) was prepared *racemic*-1,1-dimethylethyl [(1S,2S)-2-(aminomethyl)cyclopentyl]carbamate (153 mg, 64 % over 3 steps) as a white green residue. Reductive amination with isobutyraldehyde followed by deprotection

as described in Z-38 steps c and d respectively, gave *racemic*-(1S,2S)-2-{[(2-methylpropyl)amino]methyl}cyclopentanamine hydrochloride (105 mg, 39% over 5 steps from amino acid). ^{1}H NMR (methanol-d_4/$CDCl_3$) 8.90 (br s, <1 H), 8.64 (br s, <1 H), 8.28 (m, 1 H), 3.97 (br s, 1 H), 3.37 (m, 1 H), 2.83-2.69 (m, 3 H), 2.18-1.69 (m, 7 H), 0.996 (m, 6 H).


b)

racemic-(3aS,5aS,13aS)-N-[(2,4-Difluorophenyl)methyl]-11-hydroxy-5-(2-methylpropyl)-10,12-dioxo-2,3,3a,4,5,5a,6,10,12,13a-decahydro-1H-cyclopenta[e]pyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-9-carboxamide. In a manner similar to that described in example Z-35, from *racemic*-(1S,2S)-2-{[(2-methylpropyl)amino]methyl}cyclopentanamine hydrochloride 105 mg, 0.434 mmol) and 16a (56 mg, 0.119 mmol) was prepared *racemic*-(3aS,5aS,13aS)-N-[(2,4-difluorophenyl)methyl]-5-(2-methylpropyl)-10,12-dioxo-11-[(phenylmethyl)oxy]-2,3,3a,4,5,5a,6,10,12,13a-decahydro-1H-cyclopenta[e]pyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-9-carboxamide (52 mg, 74%). This material was deprotected in a second step similar to the procedure described in example Z-37. Thus, from *racemic*-(3aS,5aS,13aS)-N-[(2,4-difluorophenyl)methyl]-5-(2-methylpropyl)-10,12-dioxo-11-[(phenylmethyl)oxy]-2,3,3a,4,5,5a,6,10,12,13a-decahydro-1H-cyclopenta[e]pyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-9-carboxamide (48 mg, 0.081 mmol) and 10% Pd/C (catalytic amount), the title compound was prepared as a white solid after purification by HPLC (30 mg, 75%). ^{1}H NMR ($CDCl_3$) 12.59 (s, 1 H), 10.42 (s, 1 H), 828 (s, 1 H),

7.34 (m, 1 H), 6.79 (m, 2 H), 4.83 (s, 1 H), 4.63-4.58 (m, 3 H), 4.29 (m, 1 H), 4.14 (m, 1 H), 2.91 (m, 1 H), 2.46-2.32 (m, 3 H), 2.15-2.09 (m, 2 H), 1.85-1.61 (m, 5 H), 1.39 (m, 1 H), 0.88 (m, 6 H); ES+ MS: 501 (M +1).

Example Z-50:


(3*R*,11a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-3-ethyl-6-hydroxy-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide.



The title compound was made in two steps using a similar process to that described in example Z-2. 16a (40 mg, 0.09 mmol) and (2*R*)-2-amino-1-butanol (0.02 mL, 0.21 mmol) were reacted in dichloromethane (2 mL) with acetic acid to give (3*R*,11a*S*)-*N*-[(2,4-difluorophenyl)methyl]-3-ethyl-5,7-dioxo-6-[(phenylmethyl)oxy]-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (40 mg, 93%). This material was hydrogenated in a second step as described in example Z-2 to give (3*R*,11a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-3-ethyl-6-hydroxy-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (30 mg, 91%) as a white solid. ^{1}H NMR ($CDCl_3$) δ 11.49 (br, 1 H), 10.28 (m, 1 H), 8.35 (s, 1 H), 7.34 (m, 1 H), 6.79 (m, 2 H), 5.30 (m, 1 H), 4.62 (m, 2 H), 4.45-4.32 (m, 3 H), 3.93-3.86 (m, 2 H), 2.11 (m, 1 H), 1.65 (m, 1 H), 0.98 (t, *J* = 7.6 Hz, 3 H); ES+ MS: 420 (M +1).

Example Z-51:

racemic-(4aS,6aS,14aS)-N-[(2,4-Difluorophenyl)methyl]-12-hydroxy-6-[2-(4-morpholinyl)ethyl]-11,13-dioxo-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide.



a) *racemic*-1,1-Dimethylethyl [(1S,2R)-2-formylcyclohexyl]carbamate. An alternative procedure from the one given in example Z-38b follows: To a solution of Dess-Martin Periodane (564 mg, 1.33 mmol) in dichloromethane was added *racemic*-1,1-dimethylethyl [(1S,2R)-2-(hydroxymethyl)cyclohexyl]carbamate (305 mg, 1.33 mmol, see example Z-38a) dropwise as a solution in dichloromethane. The reaction was stirred 1 hour at ambient temperature until judged complete by TLC (1:1 hexanes: ethyl acetate $KMnO_4$ stain). The reaction was quenched with aqueous sodium bicarbonate and sodium thiosulfate solutions, extracted with dichloromethane, and the combined organics were dried over sodium sulfate. Silica gel chromatography (0-50% ethyl acetate/ hexanes gradient elution) gave *racemic*-1,1-dimethylethyl [(1S,2R)-2-formylcyclohexyl]carbamate (280, 93%). See example Z-38b for NMR data.

b) *racemic*-{[(1S,2S)-2-Aminocyclohexyl]methyl}[2-(4-morpholinyl)ethyl]amine

hydrochloride. In a manner similar to that described in example Z-38c-d from *racemic*-1,1-dimethylethyl [(1S,2R)-2-formylcyclohexyl]carbamate (78 mg, 0.344 mmol, prepared using the procedure from example Z-38b) and [2-(4-morpholinyl)ethyl]amine (67 mg, 0.515 mmol) was prepared *racemic*-{[(1S,2S)-2-aminocyclohexyl]methyl}[2-(4-morpholinyl)ethyl]amine hydrochloride (95 mg, 78% over 2 steps) as a white solid. ^{1}H NMR (methanol-d_4/$CDCl_3$) 8.18 (br s, 1 H), 3.84-3.493 (m, 11 H), 3.19-3.119 (m, 5 H), 2.42 (m, 1 H), 2.11 (br s, 2 H), 1.87-1.17 (m, 10 H).


c)

racemic-4aS,6aS,14aS)-N-[(2,4-Difluorophenyl)methyl]-12-hydroxy-6-[2-(4-morpholinyl)ethyl]-11,13-dioxo-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide. In a manner similar to that described in example Z-35, from *racemic*-{[(1S,2S)-2-aminocyclohexyl]methyl}[2-(4-morpholinyl)ethyl]amine hydrochloride (95 mg, 0.272 mmol) and 16a (45 mg, 0.0957 mmol) was prepared *racemic*-(4aS,6aS,14aS)-N-[(2,4-difluorophenyl)methyl]-6-[2-(4-morpholinyl)ethyl]-11,13-dioxo-12-[(phenylmethyl)oxy]-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide (27 mg, 43%). This material was deprotected in a second step similar to the procedure described in example Z-37. From *racemic*-(4aS,6aS,14aS)-N-[(2,4-difluorophenyl)methyl]-6-[2-(4-morpholinyl)ethyl]-11,13-dioxo-12-[(phenylmethyl)oxy]-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':

4,5]pyrazino[1,2-a]quinazoline-10-carboxamide (27 mg, 0.0408 mmol) and 10% Pd/C (1 mg) the title compound was prepared as a white solid after purification by HPLC. ^{1}H NMR ($CDCl_3$) 12.30 (br s, <1 H), 10.41 (br s, 1 H), 8.29 (s, 1 H), 7.34 (m, 2 H), 6.78 (m, 2 H), 4.76 (m, 1 H), 4.62-4.54 (m, 3 H), 4.29 (m, 2 H), 3.65 (m, 4 H), 3.01 (m, 1 H), 2.76 (m, 2 H), 2.58-2.42 (m, 7 H), 2.21 (m, 1 H), 1.89-1.23 (m, 8 H); ES$^+$ MS: 572 (M +1).

Example Z-52:

racemic-(3aR,5aR,13aS)-N-[(2,4-Difluorophenyl)methyl]-11-hydroxy-10,12-dioxo-1,2,3,3a,4,5a,6,10,12,13a-decahydrocyclopenta[d]pyrido[1',2':4,5]pyrazino[2,1-b][1,3]oxazine-9-carboxamide.



a) *racemic*-1,1-Dimethylethyl [(1S,2R)-2-(hydroxymethyl)cyclopentyl]carbamate. *racemic*-(1R,2S)-2-({[(1,1-Dimethylethyl)oxy]carbonyl}amino)cyclopentanecarboxylic acid (22 mg, 0.096 mmol) was dissolved in tetrahydrofuran and placed in an ice-water bath. Triethylamine was added, followed by the slow addition of methyl chloroformate. The reaction was stirred ten minutes in the ice-bath and sodium borohydride was added. Methanol was then added slowly and stirring was continued for two hours while the ice-bath expired. 1 M Potassium hydrogen sulfate was added, the reaction was partially concentrated, and product was extracted with dichloromethane. The combined organics were washed with sodium bicarbonate, brine, and dried over sodium sulfate. Removal of solvents under reduced pressure

afforded *racemic*-1,1-dimethylethyl [(1S,2R)-2-(hydroxymethyl)cyclopentyl]carbamate (25 mg, >100%). ^{1}H NMR ($CDCl_3$) 4.50 (br s, 1 H), 4.06 (m, 1 H), 3.54 (m, 1 H), 3.37 (m, 1 H), 2.09 (m, 1 H), 1.96 (m, 1 H), 1.64 (m, 3 H), 1.52 (m, 1 H), 1.43 (s, 9 H), 1.11 (m, 2 H).

b) *racemic*-[(1R,2S)-2-Aminocyclopentyl]methanol hydrochloride. In a manner similar to that described in example, from *racemic*-1,1-dimethylethyl [(1S,2R)-2-(hydroxymethyl)cyclopentyl]carbamate and 4 N HCl was prepared *racemic*-[(1R,2S)-2-aminocyclopentyl]methanol hydrochloride (20 mg, quantitative). ^{1}H NMR (methanol-d4-$CDCl_3$) 7.76 (br s, <1 H), 3.73 (m, 1 H), 3.61- 3.28 (m, 3 H), 2.27 (br s, 1 H), 2.01 (m, 2.01 (m, 1 H), 1.74-1.70 (m, 2 H), 1.56-1.42 (m, 2 H), 1.16 (br s, 1 H), 1.05 (br s, 1 H).


c)

racemic-(3aR,13aS)-N-[(2,4-Difluorophenyl)methyl]-11-hydroxy-10,12-dioxo-1,2,3,3a,4,5a,6,10,12,13a-decahydrocyclopenta[d]pyrido[1',2':4,5]pyrazino[2,1-b][1,3]oxazine-9-carboxamide. In a manner similar to that described in example Z-35, from *racemic*-[(1R,2S)-2-aminocyclopentyl]methanol hydrochloride (20 mg, 0.132 mmol) and 16a (24 mg, 0.051 mmol) was prepared *racemic*-(3aR,13aS)-N-[(2,4-difluorophenyl)methyl]-10,12-dioxo-11-[(phenylmethyl)oxy]-1,2,3,3a,4,5a,6,10,12,13a-decahydrocyclopenta[d]pyrido[1',2':4,5]pyrazino[2,1-b][1,3]oxazine-9-carboxamide (7 mg, 26 %) as a white solid. This material was deprotected in a second step similar to the procedure described in example Z-37. Thus, from

racemic-(3aR,13aS)-N-[(2,4-difluorophenyl)methyl]-10,12-dioxo-11-[(phenylmethyl)oxy]-1,2,3,3a,4,5a,6,10,12,13a-decahydrocyclopenta[d]pyrido[1',2':4,5]pyrazino[2,1-b][1,3]oxazine-9-carboxamide (7 mg, 0.012 mmol) and 10% Pd/C (1 mg), was obtained *racemic*-(3aR,13aS)-N-[(2,4-difluorophenyl)methyl]-11-hydroxy-10,12-dioxo-1,2,3,3a,4,5a,6,10,12,13a-decahydrocyclopenta[d]pyrido[1',2':4,5]pyrazino[2,1-b][1,3]oxazine-9-carboxamide (4 mg, 72%) white solid. ^{1}H NMR ($CDCl_3$) 12.20 (br s, 1 H), 10.37 (br s, 1 H), 8.31 (s, 1 H), 7.35 (m, 1 H), 6.80 (m, 2 H), 5.16 (m, 1 H), 4.77 (m, 1 H), 4.64 (m, 2 H), 4.28 (m, 1 H), 4.09 (m, 1 H), 3.97 (m, 1 H), 3.45 (m, 1 H), 2.49-2.20 (m, 2 H), 1.89-1.58 (m, 4 H), 0.936-0.840 (m, 1 H); ES$^+$ MS: 446 (M +1).

Example Z-53:

racemic-(4aS,6aS,14aS)-N-[(2,4-Difluorophenyl)methyl]-12-hydroxy-6-methyl-11,13-dioxo-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide.



a) *racemic*-{[(1S,2S)-2-Aminocyclohexyl]methyl}methylamine hydrochloride. In a manner similar to that described in example Z-38c-d from *racemic*-1,1-dimethylethyl [(1S,2R)-2-formylcyclohexyl]carbamate (0.410 mmol) and methyl amine (0.5 mL of a 2 M tetrahydrofuran solution) was prepared *racemic*-{[(1S,2S)-2-aminocyclohexyl]methyl}methylamine hydrochloride in two steps as a white solid (46 mg, 53% 2 steps). ^{1}H NMR (methanol-d_4/$CDCl_3$) 9.05 (br s,<1


H), 8.72 (br s, < 1 H), 8.24 (br s, 1 H), 3.34 (m, 1 H), 3.29 (m, 1 H), 2.85 (br s, 1H), 2.66 (br s, 4 H), 2.38 (br s, 1 H), 2.07-1.83 (m, 2 H), 1.67-1.14 (m, 6 H).

b)

racemic-(4aS,6aS,14aS)-N-[(2,4-Difluorophenyl)methyl]-12-hydroxy-6-methyl-11,13-dioxo-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide. In a manner similar to that described in example Z-35, from *racemic*-{[(1S,2S)-2-aminocyclohexyl]methyl}methylamine hydrochloride (46 mg, 0.215 mmol) and 16a (35 mg, 0.0744 mmol) was prepared *racemic*-(4aS,6aS,14aS)-N-[(2,4-difluorophenyl)methyl]-6-methyl-11,13-dioxo-12-[(phenylmethyl)oxy]-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide(17 mg, 41%) as a white solid. This material was deprotected in s second step similar to the procedure described in example Z-37. Thus, from *racemic*-(4aS,6aS,14aS)-N-[(2,4-difluorophenyl)methyl]-6-methyl-11,13-dioxo-12-[(phenylmethyl)oxy]-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide (17 mg, 0.0302 mmol) and 10% Pd/C (1 mg) was prepared the title compound as a white solid (9 mg, 64%). ^{1}H NMR ($CDCl_3$) 10.44 (m, 1 H), 8.29 (s, 1 H), 7.34 (m, 1 H), 6.79 (m, 2 H), 4.78 (m, 1 H), 4.62 (br s, 2 H), 4.29 (br s, 2 H), 3.41 (s, 1 H), 2.92 (m, 1 H), 2.66 (m, 1 H), 2.35-2.25 (m, 4 H), 1.90-1.74 (m, 2 H), 1.67-1.24 (m, 6 H); ES^+ MS: 473(M +1).

Example Z-54:

racemic-(4aS,6aS,14aS)-N-[(2,4-Difluorophenyl)methyl]-12-hydroxy-6-[2-(methyloxy)ethyl]-11,13-dioxo-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide.



a) *racemic*-{[(1S,2S)-2-Aminocyclohexyl]methyl}[2-(methyloxy)ethyl]amine hydrochloride.

In a manner similar to that described in example Z-38c-d from *racemic*-1,1-dimethylethyl [(1S,2R)-2-formylcyclohexyl]carbamate (93 mg, 0.410 mmol) and [2-(methyloxy)ethyl]amine (0.05 mL, 0.615 mmol) was prepared in two steps *racemic*-{[(1S,2S)-2-aminocyclohexyl]methyl}[2-(methyloxy)ethyl]amine hydrochloride (63 mg, 60% 2 steps) as a white solid. ^{1}H NMR (methanol-d_4/$CDCl_3$) 9.02 (br s, <1 H), 8.78 (br s, <1, H), 8.29 (br s, 1 H), 3.69 (br s, 2 H), 3.46 (s, 3H), 3.36-3.18 (m, 4 H), 2.97 (br s, 1 H), 2.46 (br s, 1 H), 1.86-1.40 (m, 8 H).


b)

racemic-4aS,6aS,14aS)-N-[(2,4-Difluorophenyl)methyl]-12-hydroxy-6-[2-(methyloxy)ethyl]-11,13-dioxo-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide. In a manner similar to that described in example Z-35, from *racemic*-{[(1S,2S)-2-aminocyclohexyl]methyl}[2-(methyloxy)ethyl]amine hydrochloride (63mg, 0.244 mmol) and 16a (40 mg, 0.0851 mmol) was prepared

racemic-(4aS,6aS,14aS)-N-[(2,4-difluorophenyl)methyl]-6-[2-(methyloxy)ethyl]-11,13-dioxo-12-[(phenylmethyl)oxy]-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide (44 mg, 81%) as a white solid. This material was deprotected in a second step similar to the procedure described in example Z-37. Thus, from *racemic*-(4aS,6aS,14aS)-N-[(2,4-difluorophenyl)methyl]-6-[2-(methyloxy)ethyl]-11,13-dioxo-12-[(phenylmethyl)oxy]-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide (44 mg, 0.0726mmol) and 10% Pd/C (1 mg) the title compound was prepared as a white solid (37 mg, quantitative). ^{1}H NMR ($CDCl_3$) 12.60 (br s, 1 H), 10.47 (m, 1 H), 8.28 (s, 1 H), 7.34 (m, 1 H), 6.79 (m, 2 H), 4.81 (m, 1 H), 4.64 (m 3 H), 4.51 (m, 1 H), 4.26 (m, 1 H), 3.63 (m, 1 H), 3.31 (s, 3 H), 3.19 (m, 1 H), 2.86 (m, 1 H), 2.67 (2m, 2 H), 2.21 (m, 1 H), 1.91-1.78 (m, 2 H), 1.671.52 (m, 4 H), 1.46-1.24 (m, 3 H); ES$^+$ MS: 517 (M +1).

Example Z-55:

racemic-(4aS,6aS,14aS)-6-[2-(Acetylamino)ethyl]-N-[(2,4-difluorophenyl)methyl]-12-hydroxy-11,13-dioxo-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide.



a) *racemic*-N-[2-({[(1S,2S)-2-Aminocyclohexyl]methyl}amino)ethyl]acetamide

hydrochloride. In a manner similar to that described in example Z-38c-d from *racemic*-1,1-dimethylethyl [(1S,2R)-2-formylcyclohexyl]carbamate (93 mg, 0.41 mmol) and N-(2-aminoethyl)acetamide (63 mg, 0.615 mmol), *racemic*-N-[2-({[(1S,2S)-2-aminocyclohexyl]methyl}amino)ethyl]acetamide hydrochloride was prepared in two steps as a white solid (82 mg), 71% 2 steps). ^{1}H NMR (methanol-d_4/$CDCl_3$) 8.86 (br s, 1 H), 8.29 (br s, 1 H), 3.62-3.51 (m, 3 H), 3.40-3.28 (m, 4 H), 3.22-2.93 (m, 3 H), 2.47 (m, 1 H), 2.08-2.06 (m, 4 H), 1.83-1.75 (m, 2 H), 1.56-1.44 (m, 3 H), 1.23 (m, 1 H).


b)

racemic-4aS,6aS,14aS)-6-[2-(Acetylamino)ethyl]-N-[(2,4-difluorophenyl)methyl]-12-hydroxy-11,13-dioxo-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide. In a manner similar to that described in example Z-35, from *racemic*-N-[2-({[(1S,2S)-2-aminocyclohexyl]methyl}amino)ethyl]acetamide hydrochloride (82 mg, 0.349 mmol) and 16a (50 mg, 0.106 mmol) was prepared the title compound (24 mg, 36%). This material was deprotected in a second step similar to the procedure described in example Z-37. Thus, from *racemic*-(4aS,6aS,14aS)-6-[2-(acetylamino)ethyl]-N-[(2,4-difluorophenyl)methyl]-11,13-dioxo-12-[(phenylmethyl)oxy]-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide (24 mg, 0.0379 mmol) and 10% Pd/C (1 mg) was prepared the title compound as a white solid after purification by HPLC. ^{1}H NMR ($CDCl_3$) 12.59 (s, 1 H), 10.44 (s, 1 H), 8.35 (s, 1 H), 7.32 (m, 1 H), 6.79 (m, 2 H), 5.86

(s, 1 H), 4.78 (m, 1 H), 4.61-4.50 (m, 3 H), 4.30 (m, 1 H), 3.35 (m, 1 H), 3.18 (m, 1 H), 2.96 (m, 1 H), 2.76 (m, 2 H), 2.48 (m, 1 H), 2.19 (m, 1 H), 1.89-1.23 (m, 12 H); ES^+ MS: 544 (M +1).

Example Z-56:


(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-3-ethyl-6-hydroxy-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide.



The title compound was made in two steps using a similar process to that described in example Z-2. 16a (40 mg, 0.09 mmol) and (2*S*)-2-amino-1-butanol (0.1 mL) were reacted in dichloromethane (2 mL) with acetic acid to give (3*S*,11a*R*)-*N*-[(2,4-difluorophenyl)methyl]-3-ethyl-5,7-dioxo-6-[(phenylmethyl)oxy]-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (39 mg, 90%). This material was hydrogenated in a second step as described in example Z-2 to give (3*S*,11a*R*)-*N*-[(2,4-difluorophenyl)methyl]-3-ethyl-6-hydroxy-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (37 mg, 99%) as a tinted white solid. ^1H NMR ($CDCl_3$) δ 11.47 (br, 1 H), 10.26 (m, 1 H), 8.35 (s, 1 H), 7.32 (m, 1 H), 6.77 (m, 2), 5.29 (m, 1 H), 4.60 (m, 2 H), 4.47-4.32 (m, 3 H), 3.93-3.85 (m, 2 H), 2.08 (m, 1 H), 1.68 (m, 1 H), 0.95 (t, J = 7.6 Hz, 3 H); ES^+ MS: 420 (M +1).

Example Z-57:


(3*S*,11a*R*)-3-Butyl-*N*-[(2,4-difluorophenyl)methyl]-6-hydroxy-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide.



The title compound was made in two steps using a similar process to that described in example Z-2. 16a (40 mg, 0.09 mmol) and (2*S*)-2-amino-1-hexanol (100 mg) were reacted in dichloromethane (2 mL) with acetic acid to give (3*S*,11a*R*)-3-butyl-*N*-[(2,4-difluorophenyl)methyl]-5,7-dioxo-6-[(phenylmethyl)oxy]-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (43 mg, 94%). This material was hydrogenated in a second step as described in example Z-2 to give (3*S*,11a*R*)-3-butyl-*N*-[(2,4-difluorophenyl)methyl]-6-hydroxy-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (33 mg, 92%) as a tinted white solid. ^{1}H NMR ($CDCl_3$) δ 11.48 (br, 1 H), 10.27 (br, 1 H), 8.36 (br, 1 H), 7.31 (m, 1 H), 6.77 (m, 2), 5.28 (m, 1 H), 4.59-4.36 (m, 5 H), 3.83 (m, 2 H), 2.08 (m, 1 H), 1.58 (m, 1 H), 1.39-1.23 (m, 4 H), 0.90 (t, J = 6.8 Hz, 3 H); ES$^+$ MS: 448 (M+1).

Example Z-58:


(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-[(4-hydroxyphenyl)methyl]-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide



The title compound was made in two steps using a similar process to that described in example Z-2. 16a (40 mg, 0.09 mmol) and 4-[(2*S*)-2-amino-3-hydroxypropyl]phenol (43 mg, 0.21 mmol) were reacted in dichloromethane (2 mL) with acetic acid to give (3*S*,11a*R*)-*N*-[(2,4-difluorophenyl)methyl]-3-[(4-hydroxyphenyl)methyl]-5,7-dioxo-6-[(phenylmethyl)oxy]-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (10 mg, 20%). This material was hydrogenated in a second step as described in example Z-2 and purified via preparative HPLC to give (3*S*,11a*R*)-*N*-[(2,4-difluorophenyl)methyl]-6-hydroxy-3-[(4-hydroxyphenyl)methyl]-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide (7 mg, 63%) as a white solid. ^{1}H NMR (CD_3OD) δ 10.43 (m, 1 H), 8.34 (s, 1 H), 7.33 (m, 1 H), 7.00 (d, *J* = 8.4 Hz, 2 H), 6.82 (m, 2 H), 6.71 (d, *J* = 8.4 Hz, 2 H), 5.05 (m, 1 H), 4.67-4.57 (m, 4 H), 4.21 (dd, *J* = 8.8, 7.2 Hz, 1 H), 3.94 (dd, *J* = 8.8, 6.4 Hz, 1 H), 3.21 (dd, *J* = 13.2, 3.2 Hz, 1 H), 2.90 (dd, *J* = 13.6, 8.8 Hz, 1 H); ES$^+$ MS: 498 (M+1).

Example Z-59:

(4S,12aS)-1-Cyclobutyl-N-[(2,4-difluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-9-carboxamide.



a) [(3*S*)-3-Aminobutyl]cyclobutylamine dihydrochloride was prepared in a similar manner as described in example Z-47. ^{1}H NMR (400 MHz, $CDCl_3/CD_3OD$) δ 1.23 (d, J = 6.4 Hz, 3H), 1.69-2.26 (m, 8H), 2.83 (m, 2H), 3.31-3.33 (m, 1H), 3.55 (m, 1H), 8.08 (br, <1H), 9.07 (br, <1H).


b)

(4*S*,12a*S*)-1-Cyclobutyl-*N*-[(2,4-difluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide. The title compound was made in two steps using a similar process to that described in example Z-2. 16a (80 mg, 0.17 mmol) and free based [(3*S*)-3-aminobutyl]cyclobutylamine (96 mg, 0.68 mmol) were reacted in dichloromethane (2 mL) with acetic acid to give (4*S*,12a*S*)-1-cyclobutyl-*N*-[(2,4-difluorophenyl)methyl]-4-methyl-6,8-dioxo-7-[(phenylmethyl)oxy]-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide (68 mg, 70%) as a film. This material was hydrogenated in a second step as described in example Z-2 to give (4*S*,12a*S*)-1-cyclobutyl-*N*-[(2,4-difluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide (57 mg, 100%) as an off-white solid. ^{1}H NMR (400 MHz, CDCl3) δ 1.31 (d, J = 6.8 Hz, 3H), 1.46-1.70 (m, 4H), 1.91-2.12 (m, 4H), 2.52 (m, 1H), 2.90-2.93 (m, 1H), 3.06 (m,

1H), 4.16-4.29 (m, 3H), 4.57-4.66 (m, 2H), 4.99-5.05 (m, 1H), 6.75-6.82 (m, 2H), 7.32-7.38 (m, 1H), 8.20 (s, 1H), 10.44 (s, 1H), 12.51 (s, 1H); ES+ MS: 473 (M+1).

Example Z-60:

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1-(tetrahydro-2*H*-thiopyran-4-yl)-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide.



a) [(3*S*)-3-Aminobutyl]tetrahydro-2*H*-thiopyran-4-ylamine dihydrochloride was prepared in a similar manner as described in example Z-47. ^{1}H NMR (400 MHz, $CDCl_3/CD_3OD$) δ 1.21 (d, J = 6.4 Hz , 3H), 1.65-1.75 (m, 2H), 1.90-2.10 (m, 2H), 2.35 (m, 2H), 2.56-2.61 (m, 4H), 2.92-2.98 (m, 3H), 3.27-3.31 (m, 1H), 8.05 (br, <1H), 8.90 (br, <1H).


b)

(4*S*,12a*S*)-*N*-[(2,4-difluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1-(tetrahydro-2*H*-thiopyran-4-yl)-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide. The title compound was made in two steps using a similar process to that described in example Z-2. 16a (80 mg, 0.17 mmol) and free based [(3S)-3-aminobutyl]tetrahydro-2H-thiopyran-4-ylamine (108 mg, 0.58 mmol) were reacted in dichloromethane (2 mL) with acetic acid to give

(4*S*,12a*S*)-*N*-[(2,4-difluorophenyl)methyl]-4-methyl-6,8-dioxo-7-[(phenylmethyl)oxy]-1-(tetrahydro-2*H*-thiopyran-4-yl)-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide (56 mg, 54%) as a film. This material was debenzylated in a second step to in a manner similar to Z-26 to give (4*S*,12a*S*)-*N*-[(2,4-difluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1-(tetrahydro-2*H*-thiopyran-4-yl)-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide (56 mg, >100%) as an off-white solid. ^{1}H NMR (400 MHz, CDCl3) δ 1.30 (d, J = 6.8 Hz , 3H), 1.54-1.58 (m, 1H), 1.72-1.82 (m, 3H), 1.97-2.11 (m, 2H), 2.60-2.76 (5H), 2.86 (m, 2H), 4.17-4.30 (m, 2H), 4.62-4.66 (m, 3H), 4.92-4.96 (m, 1H), 6.75-6.82 (m, 2H), 7.32-7.38 (m, 1H), 8.31 (s, 1H), 10.46 (s, 1H), 12.48 (s, 1H); ES$^+$ MS: 519 (M+1).

Example Z-61:

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-1,4-bis(2-methylpropyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide.



a) [(3*S*)-3-Amino-5-methylhexyl](2-methylpropyl)amine dihydrochloride was prepared in a similar manner as described in example Z-32. ^{1}H NMR (400 MHz, CDCl3/CD3OD) δ 0.87 (d, J = 6.4 Hz ,6H), 0.97 (d, J = 6.8 Hz , 6H), 1.34-1.41 (m, 1H), 1.45-1.52 (m, 1H), 1.58-1.66 (m, 1H), 2.01-2.13 (m, 2H), 2.72-2.73 (m, 2H), 3.03-3.06

(m, 2H), 3.29 (m, 2H), 8.07 (br, <1H), 8.71 (br, <1H).


b)

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-1,4-bis(2-methylpropyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide. The title compound was made in two steps using a similar process to that described in example Z-2. 16a (80 mg, 0.17 mmol) and free based [(3S)-3-amino-5-methylhexyl](2-methylpropyl)amine (117 mg, 0.63 mmol) were reacted in dichloromethane (2 mL) with acetic acid to give (4*S*,12a*S*)-*N*-[(2,4-difluorophenyl)methyl]-1,4-bis(2-methylpropyl)-6,8-dioxo-7-[(phenylmethyl)oxy]-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide (68 mg, 66%) as a film. This material was hydrogenated in a second step as described in example Z-2 to give (4*S*,12a*S*)-*N*-[(2,4-difluorophenyl)methyl]-7-hydroxy-1,4-bis(2-methylpropyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide (56 mg, 97%) as an off-white solid. ^{1}H NMR (400 MHz, CDCl3) δ 0.74 (d, *J* = 6.4 Hz, 3H), 0.84 (d, *J* = 6.4 Hz, 3H), 0.97-1.00 (m, 6H), 1.37-1.83 (m, 5H), 2.03-2.12 (m, 2H), 2.21-2.28 (m, 1H), 2.77 (m, 1H), 2.90-2.93 (m, 1H), 4.19-4.40 (m, 3H), 4.59-4.70 (m, 2H), 4.96-4.97 (m, 1H), 6.77-6.83 (m, 2H), 7.33-7.39 (m, 1H), 8.28 (s, 1H), 10.47 (s, 1H), 12.59 (br, 1H); ES$^+$ MS: 517 (M+1).

Example Z-62:

racemic-(4aS,6aS,14aS)-N-[(2,4-Difluorophenyl)methyl]-12-hydroxy-6-(2-hydroxyethyl

)-11,13-dioxo-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide.



a) *racemic*-2-({[(1S,2S)-2-Aminocyclohexyl]methyl}amino)ethanol hydrochloride. In a manner similar to that described in example Z-55a, from *racemic*-1,1-dimethylethyl [(1S,2R)-2-formylcyclohexyl]carbamate (112 mg, 0.497 mmol) and 2-aminoethanol (0.04 mL, 0.746 mmol) was prepared *racemic*-2-({[(1S,2S)-2-aminocyclohexyl]methyl}amino)ethanol bis-hydrochloride in two steps (102 mg, 84% over 2 steps). ^{1}H NMR (methanol-d_4/$CDCl_3$) 8.81-8.40 (m, < 2 H), 8.16 (br s, 1 H), 4.02-3.93 (m, 2 H), 3.80 (br s, 2 H), 3.53 (m, 1 H), 3.36-2.93 (m, 6 H), 2.41 (br s, 1 H), 2.05 (m, 1 H), 1.75-1.41 (m, 4 H).


b)

racemic-(4aS,6aS,14aS)-N-[(2,4-Difluorophenyl)methyl]-12-hydroxy-6-(2-hydroxyethyl)-11,13-dioxo-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide. In a manner similar to that described in example Z-35, from 16a (45 mg, 0.0957 mmol) and *racemic*-2-({[(1S,2S)-2-aminocyclohexyl]methyl}amino)ethanol hydrochloride (102 mg, 0.418 mmol) was prepared *racemic*-(4aS,6aS,14aS)-N-[(2,4-difluorophenyl)methyl]-6-(2-hydroxyethyl)-11,13-dioxo-12-[(phenylmethyl)oxy]-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyra

zino[1,2-a]quinazoline-10-carboxamide (7 mg, 12 %) as a white solid after silica gel chromatography (1-12% methanol/dichloromethane gradient elution). This material was deprotected in a second step similar to the procedure described in example Z-37. Thus, from *racemic*-(4aS,6aS,14aS)-N-[(2,4-difluorophenyl)methyl]-6-(2-hydroxyethyl)-11,13-dioxo-12-[(phenylmethyl)oxy]-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide (7 mg, 0.0118 mmol) the title compound was prepared after purification by HPLC (3 mg, 50 %). ^{1}H NMR ($CDCl_3$) 12.57 (br s, 1 H), 10.45 (m, 1 H), 8.29 (s, 1 H), 7.34 (m, 1 H), 6.78 (m, 2 H), 4.80 (m, 1 H), 4.71 (s, 1 H), 4.62 (m, 2 H), 4.44 (m, 1 H), 4.33 (m, 1 H), 3.75 (m, 1 H), 3.62-3.20 (m, 3 H), 3.13 (m, 1 H), 2.74-2.71 (m, 2 H), 2.24 (m, 1 H), 1.90-1.37 (m, 12 H), 1.27-1.23 (m, 3 H)1.12 (m, 1 H); ES$^+$ MS: 503 (M +1).

Example Z-63:

racemic-(4aS,6aS,14aS)-6-Cyclopropyl-N-[(2,4-difluorophenyl)methyl]-12-hydroxy-11,13-dioxo-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide.



a) *racemic*-(1S,2S)-2-[(Cyclopropylamino)methyl]cyclohexanamine hydrochloride. In a manner similar to that described in example Z-55a, from *racemic*-1,1-dimethylethyl [(1S,2R)-2-formylcyclohexyl]carbamate (112 mg, 0.497

mmol) and cyclopropylamine (0.05 mL, 0.746 mmol) was prepared *racemic*-(1S,2S)-2-[(cyclopropylamino)methyl]cyclohexanamine bis hydrochloride salt in two steps (102 mg, 86% over 2 steps). This material was used without further purification. ^{1}H NMR (methanol-d_4/$CDCl_3$) 8.31 (br s, 1 H), 3.75 (br s, 1 H), 3.54 (m, 1 H), 2.96 (m, 1 H), 2.71 (m, 1 H), 2.27 (m, 1 H), 1.94 (m, 1 H), 1.76-1.15 (m, 8 H), 0.88-0.78 (m, 3 H).


b)

racemic-(4aS,6aS,14aS)-6-Cyclopropyl-N-[(2,4-difluorophenyl)methyl]-12-hydroxy-11,13-dioxo-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide. In a manner similar to that described in example Z-35, from 16a (45 mg, 0.0957 mmol) and *racemic*-(1S,2S)-2-[(cyclopropylamino)methyl]cyclohexanamine hydrochloride (102 mg, 0.425 mmol) was prepared *racemic*-(4aS,6aS,14aS)-6-cyclopropyl-N-[(2,4-difluorophenyl)methyl]-11,13-dioxo-12-[(phenylmethyl)oxy]-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide as a white solid after silica gel chromatography (1-12% methanol/dichloromethane gradient elution). This material was deprotected in a second step similar to the procedure described in example Z-37. Thus, from *racemic*-(4aS,6aS,14aS)-6-cyclopropyl-N-[(2,4-difluorophenyl)methyl]-11,13-dioxo-12-[(phenylmethyl)oxy]-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide (56 mg, 0.0949 mmol) the title compound was prepared as a white solid (41 mg, 81%). ^{1}H NMR ($CDCl_3$) 12.10 (br s, < 1 H), 10.45

(m, 1 H), 8.27 (s, 1 H), 7.33 (m, 1 H), 6.88 (m, 2 H), 4.77 (m, 1 H), 4.61-4.49 (m, 4 H), 4.33 (m, 1 H), 2.94 (m, 1 H), 2.79 (m, 1 H), 2.17 (m, 1 H), 1.86-0.86 (m, 10 H), 0.658 (m, 1 H), 0.499-0.32 (m, 2 H); ES^+ MS: 499 (M +1).

Example Z-64:

racemic-(4aS,6aS,14aS)-N-[(2,4-Difluorophenyl)methyl]-12-hydroxy-11,13-dioxo-6-[2-(1-pyrrolidinyl)ethyl]-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide formic acid salt



a) *racemic*-(1S,2S)-2-({[2-(1-Pyrrolidinyl)ethyl]amino}methyl)cyclohexanamine hydrochloride. In a manner similar to that described in example Z-55a, from *racemic*-1,1-dimethylethyl [(1S,2R)-2-formylcyclohexyl]carbamate (112 mg, 0.497 mmol) and 2-(1-pyrrolidinyl)ethanamine (0.09 mL, 0.746 mmol) was prepared *racemic*-(1S,2S)-2-({[2-(1-pyrrolidinyl)ethyl]amino}methyl)cyclohexanamine (88 mg, 60% 2 steps) as the bis hydrochloride salt in two steps as a white solid. ^1H NMR (methanol-d_4/$CDCl_3$) 9.68 (br s, < 1 H), 9.24 (br s, < 1 H), 8.25 (br s, 1 H), 3.75- 3.04 (m, 11 H), 2.37 (br s, 1 H), 2.06-1.20 (m, 12 H).

b)

racemic-(4aS,6aS,14aS)-N-[(2,4-Difluorophenyl)methyl]-12-hydroxy-11,13-dio


xo-6-[2-(1-pyrrolidinyl)ethyl]-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide formic acid salt.

In a manner similar to that described in example Z-35, from 16a (30 mg, 0.0638 mmol) and *racemic*-(1S,2S)-2-({[2-(1-pyrrolidinyl)ethyl]amino}methyl)cyclohexanamine hydrochloride (88 mg, 0.296 mmol) was prepared *racemic*-(4aS,6aS,14aS)-N-[(2,4-difluorophenyl)methyl]-11,13-dioxo-12-[(phenylmethyl)oxy]-6-[2-(1-pyrrolidinyl)ethyl]-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide as a white solid (31 mg, 76%) after silica gel chromatography (1-12% methanol/dichloromethane gradient elution). This material was deprotected in a second step similar to the procedure described in example Z-37. Thus, from *racemic*-(4aS,6aS,14aS)-N-[(2,4-difluorophenyl)methyl]-11,13-dioxo-12-[(phenylmethyl)oxy]-6-[2-(1-pyrrolidinyl)ethyl]-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide (31 mg, 0.048 mmol) the title compound was prepared as a yellow solid after purification by HPLC (18 mg, 66%). ^{1}H NMR ($CDCl_3$) 10.39 (br s, 1 H), 8.56 (br s, 1 H), 8.39 (br s, 1 H), 7.34 (m, 1 H), 6.78 (m, 2 H), 4.76-4.40 (m, 6 H), 3.26-2.89 (m, 7 H), 2.73 (m, 1 H), 2.15 (m, 1 H), 2.02-1.18 (m, 14 H); ES^+ MS: 556 (M +1).

Example Z-65:

(4aS,14aS)-N-[(2,4-Difluorophenyl)methyl]-9-hydroxy-8,10-dioxo-2,3,4,4a,5,6,8,10,14,14a-decahydro-1H-pyrido[1,2-c]pyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-11-carboxami

de.




a) {2-[(2S)-2-Piperidinyl]ethyl}amine. This compound was prepared in a similar manner as its enantiomer described in example Z-42a.

b)

(4aS,14aS)-N-[(2,4-Difluorophenyl)methyl]-9-hydroxy-8,10-dioxo-2,3,4,4a,5,6,8,10,14,14a-decahydro-1H-pyrido[1,2-c]pyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-11-carboxamide. In a manner similar to that described in example Z-35, from {2-[(2S)-2-piperidinyl]ethyl}amine (28 mg, 0.218 mmol) and 16a (30 mg, 0.0638 mmol) was prepared (4aS,14aS)-N-[(2,4-difluorophenyl)methyl]-8,10-dioxo-9-[(phenylmethyl)oxy]-2,3,4,4a,5,6,8,10,14,14a-decahydro-1H-pyrido[1,2-c]pyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-11-carboxamide (29 mg, 82%). This material was deprotected in a second step similar to that described in example Z-37 to give the title compound as a white solid (26 mg, quantitative). ^{1}H NMR ($CDCl_3$) δ 12.44 (br s, 1 H), 10.48 (s, 1 H), 8.26 (s, 1 H), 7.35 (m, 1 H), 6.80 (m, 2 H), 4.68-4.57 (m, 2 H), 4.38 (m, 1 H), 4.20 (m, 1 H), 3.93 (s, 1H), 3.63-3.39 (m, 2 H), 2.91 (m, 2 H), 2.29 (br s, 1 H), 2.02 (m, 1 H), 1.69-1.45 (m, 4 H), 1.30-1.24 (m, 2 H), 1.12 (br s, 1 H); ES$^+$ MS: 459 (M+1).

Example Z-66:

(4S,12aS)-N-[(4-Fluorophenyl)methyl]-7-hydroxy-4-methyl-1-[2-(methyloxy)ethyl]-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-9-carboxamide.



a) [(3S)-3-Aminobutyl][2-(methyloxy)ethyl]amine bis hydrochloride. In a manner similar to that described in example Z-47, from 1,1-dimethylethyl [(1S)-1-methyl-3-oxopropyl]carbamate (76 mg, 0.406 mmol) and 2-(methyloxy)ethyl]amine (0.05 mL, 0.609 mmol) was prepared [(3S)-3-aminobutyl][2-(methyloxy)ethyl]amine as the bis hydrochloride salt in two steps (19 mg, quantitative). ^{1}H NMR (methanol-d_4/$CDCl_3$) δ 9.02 (< 1 H), 8.24 (< 1 H), 3.68 (br s, 2 H), 3.49 (br s, 1 H), 3.34 (br s, 4 H), 3.15 (br s, 4 H), 2.26-2.11 (m, 2 H), 1.35 (br s, 3 H).


b)

(4S,12aS)-N-[(4-Fluorophenyl)methyl]-7-hydroxy-4-methyl-1-[2-(methyloxy)ethyl]-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-9-carboxamide. In a manner similar to that described in example Z-35, from 16 (15

mg, 0.034 mmol) and [(3S)-3-Aminobutyl][2-(methyloxy)ethyl]amine bis hydrochloride (19 mg, 0.087 mmol), (4S,12aS)-N-[(4-fluorophenyl)methyl]-4-methyl-1-[2-(methyloxy)ethyl]-6,8-dioxo-7-[(phenylmethyl)oxy]-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-9-carboxamide was prepared as a white solid after silica gel chromatography (1-12% methanol/dichloromethane). This material was deprotected in a second step similar to that described in example Z-37 to give the title compound as a yellow solid (9 mg, 60 %, 2 steps). ^{1}H NMR ($CDCl_3$) δ 12.56 (s, 1 H), 10.51 (m, 1 H), 8.29 (s, 1 H), 7.32 (m, 2 H), 6.98 (m, 2 H), 5.03 (m, 1 H), 4.65-4.59 (m, 2 H), 4.53 (m, 1 H), 4.21 (m, 1 H), 3.61-3.40 (m, 2 H), 3.34-3.13 (m, 3 H), 3.08 (m, 1 H), 2.94-2.84 (m, 2 H), 2.68 (m, 1 H), 2.07 (m, 1 H), 1.50 (m, 1 H), 1.35 (d, J = 7.2 Hz, 3 H), 1.14 (m, 1 H); ES$^+$ MS: 459 (M+1).

Example Z-67:

(4S,12aS)-1-Cyclobutyl-N-[(4-fluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-9-carboxamide.



a) [(3S)-3-Aminobutyl]cyclobutylamine bis-hydrochloride. In a manner similar to that described in example Z-47, from 1,1-dimethylethyl

[(1S)-1-methyl-3-oxopropyl]carbamate (76 mg, 0.406 mmol) and cyclobutylamine (0.05 mL, 0.609 mmol) was prepared [(3S)-3-Aminobutyl]cyclobutylamine bis-hydrochloride in two steps (23 mg, 27%). ^{1}H NMR (methanol-d_4/$CDCl_3$) δ 8.86 (s, < 1 H), 7.97 (s, < 1 H), 3.46 (m, 1 H), 3.21 (m, 1 H), 2.74 (m, 2 H), 2.14-2.08 (m, 4 H), 1.94-1.62 (m, 5 H), 1.13 (d, J = 6 Hz, 1 H).


b)

(4S,12aS)-1-Cyclobutyl-N-[(4-fluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-9-carboxamide. In a similar manner to that described in example Z-35a, from 16 (18 mg, 0.39 mmol) and [(3S)-3-Aminobutyl]cyclobutylamine bis-hydrochloride (23 mg, 0.107 mmol),

(4S,12aS)-1-cyclobutyl-N-[(4-fluorophenyl)methyl]-4-methyl-6,8-dioxo-7-[(phenylmethyl)oxy]-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-9-carboxamide was prepared as a white solid. This material was deprotected in a second step similar to that described in example Z-37 to give the title compound as a white solid after purification by HPLC (4.5 mg, 25 % 2 steps). ^{1}H NMR ($CDCl_3$) δ 12.54 (s, 1 H), 10.48 (s, 1 H), 8.20 (s, 1 H), 7.31 (m, 2 H), 6.98 (m, 2 H), 5.02 (m, 1 H), 4.61-4.57 (m, 2 H), 4.26-4.14 (m, 3 H), 3.05 (m, 1 H), 2.90 (m, 1 H), 2.49 (m, 1 H), 2.12 (m, 1 H), 2.05-1.87 (m, 3 H), 1.84-1.61 (m, 3 H), 1.46 (m, 1 H), 1.32 (m, 3 H); ES^+ MS: 455 (M+1).

Example Z-68:

(4S,12aS)-N-[(4-Fluorophenyl)methyl]-7-hydroxy-4-methyl-1-(2-methylpropyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-9-carboxamide



a) [(3S)-3-Aminobutyl](2-methylpropyl)amine bis-hydrochloride. In a manner similar to that described in example Z-47, this compound was prepared from 1,1-dimethylethyl [(1S)-1-methyl-3-oxopropyl]carbamate (76 mg, 0.406 mmol) and (2-methylpropyl)amine (0.06 mL, 0.609 mmol) in two steps as the bis-hydrochloride salt (22 mg, 25 %). ^{1}H NMR (methanol-d_4/$CDCl_3$) δ 3.25 (br s, 1 H), 2.91 (br s, 2 H), 2.64 (m, 2 H), 2.02-1.93 (m, 3 H), 1.17 (m, 3 H), 0.88 (m, 6 H).


b)

(4S,12aS)-N-[(4-Fluorophenyl)methyl]-7-hydroxy-4-methyl-1-(2-methylpropyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-9-carboxamide. In a similar manner to that described in example Z-35, from 16 (16 mg, 0.035 mmol) and [(3S)-3-Aminobutyl](2-methylpropyl)amine bis-hydrochloride (20 mg, 0.0925 mmol), (4S,12aS)-N-[(4-fluorophenyl)methyl]-4-methyl-1-(2-methylpropyl)-6,8-dioxo-7-[(phenylmethyl)oxy]-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-9-

carboxamide was prepared as a white solid. This material was deprotected in a second step similar to that described in example Z-37 to give the title compound as a tan solid (13 mg, 68% 2 steps). ^{1}H NMR ($CDCl_3$) δ 12.57 (s, 1 H), 10.46 (s, 1 H), 8.27 (s, 1 H), 7.32 (m, 2 H), 6.99 (m, 2 H), 4.98 (m, 1 H), 4.63-4.54 (m, 2 H), 4.45 (m, 1 H),4.26-4.16 (m, 2 H), 2.91 (m, 1 H),-2.77-(m, 1 H), 2.24 (m, 1 H), 2.14-2.03 (m, 2 H), 1.63 (m, 1 H), 1.48 (m, 1 H), 1.33 (m, 3 H), 1.09 (m, 1 H), 0.850 (m, 3 H), 0.789 (m, 3 H); ES$^+$ MS: 457 (M+1).

Example Z-69:

(4S,12aS)-N-[(4-Fluorophenyl)methyl]-7-hydroxy-1,4-dimethyl-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-9-carboxamide.




a) [(3S)-3-Aminobutyl]methylamine bis-hydrochloride. In a manner similar to that described in example Z-47, this compound was prepared from 1,1-dimethylethyl [(1S)-1-methyl-3-oxopropyl]carbamate (76 mg, 0.409 mmol) and excess methylamine (2 M in tetrhydrofuran) in two steps as the bis hydrochloride salt (17% 2 steps). ^{1}H NMR (methanol-d_4/$CDCl_3$) δ 3.16 (m, 1 H), 3.08 (s, 2 H), 2.83 (m, 2 H), 2.45 (s, 3 H), 1.88 (m, 1 H), 1.75 (m, 1 H), 1.09 (m, 3 H).

b)

(4S,12aS)-N-[(4-Fluorophenyl)methyl]-7-hydroxy-1,4-dimethyl-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-9-carboxamide. In a similar manner to that described in example Z-35, from 16 (18 mg, 0.0398 mmol) and [(3S)-3-aminobutyl]methylamine bis-hydrochloride (19 mg, 0.109 mmol, (4S,12aS)-N-[(4-fluorophenyl)methyl]-1,4-dimethyl-6,8-dioxo-7-[(phenylmethyl)oxy]-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-9-carboxamide was prepared as a white solid. This material was deprotected in a second step similar to that described in example Z-37 to give the title compound as a tan solid (7 mg, 44% 2 steps). ^{1}H NMR ($CDCl_3$) δ 12.53 (s, 1 H), 10.47 (s, 1 H), 8.29 (s, 1 H), 7.32 (m, 2 H), 6.99 (m, 2 H), 5.04 (1 H), 4.60 (m, 2 H), 4.23 (s, 3 H), 2.83-2.80 (m, 2 H), 2.32 (s, 3 H), 2.13 (m, 1 H), 1.48 (m, 1 H), 1.34 (m, 3 H); ES$^+$ MS: 415 (M+1).

Example Z-70:

(4*S*,12a*S*)-*N*-[(4-Fluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1-(tetrahydro-2*H*-thiopyran-4-yl)-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide.




The title compound was made in two steps using a similar process to that described in example Z-2. 16 (25 mg, 0.055 mmol) and free based [(3*S*)-3-aminobutyl]tetrahydro-2*H*-thiopyran-4-ylamine (48 mg, 0.26 mmol) were

reacted in dichloromethane (2 mL) with acetic acid to give (4*S*,12a*S*)-*N*-[(4-fluorophenyl)methyl]-4-methyl-6,8-dioxo-7-[(phenylmethyl)oxy]-1-(tetrahydro-2*H*-thiopyran-4-yl)-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide (16 mg, 49%) as a film. This material was debenzylated in a second step in a manner similar to Z-26 to give (4*S*,12a*S*)-*N*-[(4-fluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1-(tetrahydro-2*H*-thiopyran-4-yl)-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide (8 mg, 59%) as an off-white solid. ^{1}H NMR (400 MHz, CDCl3) δ 1.30 (d, *J* = 7.2 Hz, 3H), 1.53-1.58 (m, 1H), 1.72-2.10 (m, 5H), 2.56-2.76 (m, 5H), 2.84-2.87 (m, 2H), 4.18 (dd, *J* = 2.8, 14.0 Hz, 1H), 4.26 (dd, *J* = 3.4, 14.2 Hz, 1H), 4.92-4.97 (m, 1H), 6.96-7.00 (m, 2H), 7.29-7.36 (m, 2H), 8.31 (s, 1H), 10.48 (m, 1H), 12.48 (br, 1H); ES$^+$ MS: 501 (M+1).

Example Z-71:

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-1,4-dimethyl-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide.




a) [(3*S*)-3-Aminobutyl]methylamine dihydrochloride was prepared in a similar manner as described in example Z-47. ^{1}H NMR (400 MHz, CDCl3) δ 1.18 (d, *J* = 6.8 Hz, 3H), 1.82-1.91 (m, 1H), 1.94-2.03 (m, 1H), 2.53 (s, 3H), 2.89-2.93 (m, 2H), 3.22-3.30 (m, 1H), 8.02 (br, <1H), 8.81 (br, <1H).

b)

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-1,4-dimethyl-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide. The title compound was made in two steps using a similar process to that described in example Z-2. 16a (40 mg, 0.085 mmol) and free based [(3*S*)-3-aminobutyl]methylamine (24 mg, 0.23 mmol) were reacted in dichloromethane (2 mL) with acetic acid to give (4*S*,12a*S*)-*N*-[(2,4-difluorophenyl)methyl]-1,4-dimethyl-6,8-dioxo-7-[(phenylmethyl)oxy]-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide (39 mg, 89%) as a film. This material was hydrogenated in a second step as described in example Z-2 to give (4*S*,12a*S*)-*N*-[(2,4-difluorophenyl)methyl]-7-hydroxy-1,4-dimethyl-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide (32 mg, 97%) as an off-white solid. ^{1}H NMR (400 MHz, CDCl3) δ 1.33 (d, *J* = 6.4 Hz, 3H), 1.46-1.50 (m, 1H), 2.12-2.14 (m, 1H), 2.32 (s, 3H), 2.83 (m, 2H), 4.24 (m, 3H), 4.62 (m, 2H), 5.02 (m, 1H), 6.77-6.79 (m, 2H), 7.33 (m, 1H), 8.30 (s, 1H), 10.43 (s, 1H), 12.50 (br, 1H); ES$^+$ MS: 433 (M+1).

Example Z-72:


(4*S*,12a*S*)-*N*-[(4-Fluorophenyl)methyl]-7-hydroxy-4-methyl-1-(1-methylethyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide.



The title compound was made in two steps using a similar process to that described in example Z-2. 16 (27 mg, 0.060 mmol) and free based [(3S)-3-aminobutyl](1-methylethyl)amine (67 mg, 0.51 mmol) were reacted in dichloromethane (2 mL) with acetic acid to give (4*S*,12a*S*)-*N*-[(4-fluorophenyl)methyl]-4-methyl-1-(1-methylethyl)-6,8-dioxo-7-[(phenyl methyl)oxy]-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide (18 mg, 56%) as a film. This material was hydrogenated in a second step as described in example Z-2 to give (4*S*,12a*S*)-*N*-[(4-fluorophenyl)methyl]-7-hydroxy-4-methyl-1-(1-methylethyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide (15 mg, >100%) as an off-white solid. ^{1}H NMR (400 MHz, CDCl3) δ 1.02 (d, *J* = 6.4 Hz, 3H), 1.07 (d, *J* = 6.4 Hz, 3H), 1.32 (d, *J* = 6.8 Hz, 3H), 1.54-1.58 (m, 1H), 1.94-2.03 (m, 1H), 2.71-2.76 (m, 1H), 2.82-2.88 (m, 1H), 3.13-3.16 (m, 1H), 4.16-4.19 (m, 1H), 4.30-4.33 (m, 1H), 4.48 (m, 1H), 4.55-4.65 (m, 2H), 4.97-5.00 (m, 1H), 6.97-7.01 (m, 2H), 7.30-7.34 (m, 2H), 8.28 (s, 1H), 10.51 (m, 1H), 12.55 (s, 1H); ES$^+$ MS: 443 (M+1).

Example Z-73:

(4*S*,12a*S*)-*N*-[(4-Fluorophenyl)methyl]-7-hydroxy-1,4-bis(2-methylpropyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide.







The title compound was made in two steps using a similar process to that described in example Z-2. 16 (25 mg, 0.055 mmol) and free based [(3S)-3-amino-5-methylhexyl](2-methylpropyl)amine (21 mg, 0.11 mmol) were reacted in dichloromethane (2 mL) with acetic acid to give (4*S*,12a*S*)-*N*-[(4-fluorophenyl)methyl]-1,4-bis(2-methylpropyl)-6,8-dioxo-7-[(phenylmethyl)oxy]-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide (8 mg, 25%) as a film. This material was hydrogenated in a second step as described in example Z-2 to give (4*S*,12a*S*)-*N*-[(4-fluorophenyl)methyl]-7-hydroxy-1,4-bis(2-methylpropyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide (5 mg, 78%) as an off-white solid. ^{1}H NMR (400 MHz, CDCl3) δ 0.74 (d, *J* = 6.4 Hz, 3H), 0.84 (d, *J* = 6.4 Hz, 3H), 0.97-1.00 (m, 6H), 1.37-1.66 (m, 5H), 1.75-1.82 (m, 1H), 2.05-2.09 (m, 2H), 2.21-2.26 (m, 1H), 2.72-2.79 (m, 1H), 2.87-2.93 (m, 1H), 4.16-4.26 (m, 2H), 4.38 (m, 1H), 4.55-4.66 (m, 2H), 4.93-4.99 (m, 1H), 6.97-7.02 (m, 2H), 7.31-7.34 (m, 2H), 8.27 (s, 1H), 10.49 (m, 1H), 12.61 (s, 1H); ES$^+$ MS: 499 (M+1).









Example ZZ-1 to ZZ-24









Examples in table below were isolated as a mixture of diastereomers ranging from 1:1 to >10:1 ratios of stereoisomers at the center indicated as undefined.





Characterization data reported herein consists of observed mass spectral signals for molecular ions (M+1) of the compounds using electrospray ionization methods in the positive mode using LC/MS techniques well known in the field. Reported retention times refer to observed UV peaks confirmed by NMR methods for the examples below using the following gradient on a phenomenex C18 reverse phase HPLC column (150 mmX4.6 mm 5 micron). Solvent A = water w/ 0.1% formic acid, solvent B = acetonitrile w/ 0.1% formic acid. Gradient = 10%B for 1 min, gradient from 10% to 90% B from 1 to 9 min, ramping to 100% B at 9.01 min and holding at 100% B for 2 min. In several cases the diastereomers were not separable by the standard HPLC conditions reported above and thus reported as a single retention time.

[Table A]


Example No.	Structure	Observed LC/MS or HPLC data
ZZ-1		ES+ MS: 419 (M +1)
ZZ-2		ES+ MS: 406 (M +1)
ZZ-3		ES+ MS: 509 (M +1)
ZZ-4		ES+ MS: 429 (M +1)

ZZ-5		ES+ MS: 415 (M +1)
ZZ-6		ES+ MS: 491 (M +1)
ZZ-7		ES+ MS: 509 (M +1)
ZZ-8		ES+ MS: 443 (M +1)
ZZ-9		ES+ MS: 461 (M +1)
ZZ-10		ES+ MS: 501 (M +1)
ZZ-11		ES+ MS: 475 (M +1)
ZZ-12		ES+ MS: 489 (M +1)

ZZ-13		ES+ MS: 460 (M +1)
ZZ-14		ES+ MS: 442 (M +1)
ZZ-15		ES+ MS: 489 (M +1)
ZZ-16		8.174 & 8.295 min.
ZZ-17		ES+ MS: 461 (M +1)
ZZ-18		ES+ MS: 447 (M +1)
ZZ-19		ES+ MS: 446 (M +1)
ZZ-20		ES+ MS: 432 (M +1)

ZZ-21		7.368 min
ZZ-22		7.150 min
ZZ-23		ES+ MS: 447 (M +1)
ZZ-24		ES+ MS: 447 (M +1)

The present invention further includes the following compounds.



(I-7)

[Table B]

No	(R) m	R^a
1	4－F	$-CH_3$
2	4－F	$-CH(CH_3)_2$
3	4－F	$-CH_2CH_2OCH_3$
4	2，4－F	$-CH_3$
5	2，4－F	$-CH(CH_3)_2$
6	2，4－F	$-CH_2CH_2OCH_3$
7	2－F，3－Cl	$-CH_3$
8	2－F，3－Cl	$-CH(CH_3)_2$
9	2－F，3－Cl	$-CH_2CH_2OCH_3$

Experimental Example 1

The HIV integrase inhibitory activity was investigated based on the following assay method.

(1) Preparation of DNA solution

By the same method as that described in Experimental Example 1 of WO 2004/024693, a substrate DNA solution (2 pmol/μl) and a target DNA solution (5 pmol/μl) were prepared. After each target DNA solution was once boiled, a temperature was slowly lowered to anneal complementary chains, which was used. Each sequence of a substrate DNA and a target DNA is as described in the same Experimental Example.

(2) Measurement of inhibition rate (IC_{50} value)

Streptavidin (manufactured by Vector Laboratories) was dissolved in a 0.1M carbonate buffer solution (composition: 90 mM Na_2CO_3, 10 mM $NaHCO_3$) to a concentration of 40 μg/ml. Each 50 μl of this solution was added to a well of an immunoplate (manufactured by NUNC), this is allowed to stand at 4°C overnight to adsorb. Then, each well was washed with a phosphate buffer (composition: 13.7 mM NaCl, 0.27 mM KCl, 0.43 mM Na_2HPO_4, 0.14 mM KH_2PO_4) two times, and 300 μl of a phosphate buffer containing 1 % skim milk to block it for 30 minutes. Further, each well was washed with a phosphate buffer two times, 50 μl of a substrate DNA solution (2 pmol/μl) was added to adsorb at room temperature for 30 minutes while shaking, and this was washed with a phosphate buffer two times and, then, distilled water once.

Then, to each well prepared as described above were added 12 μl of a buffer (composition: 150 mM MOPS (pH7.2), 75 mM $MnCl_2$, 50 mM 2-mercaptoethanol, 25% glycerol, 500 μg/ml bovine serum albumin-fraction V), and 51 μl of a reaction solution prepared from 39 μl of distilled water. Then, 9 μl of an integrase solution (30 pmol) was added, and the mixture was mixed well. To a well as a negative control (NC) was added 9 μl of a diluting solution (composition: 20 mM MOPS (pH7.2), 400 mM potassium glutamate, 1 mM EDTA, 0.1% NP-40, 20% glycerol, 1 mM DTT, 4 M urea), and this was mixed well using a plate mixer.

After the plate was incubated at 30°C for 60 minutes, the reaction solution was discarded, followed by washing with 250 μl of a washing buffer (composition: 150 mM MOPS (pH7.2), 50 mM 2-mercaptoethanol, 25% glycerol, 500 μg/ml bovine serum albumin-fraction V) three times.

Then, to each well were added 12 μl of a buffer (composition: 150 mM MOPS

(pH7.2), 75 mM $MgCl_2$, 50 mM 2-mercaptoethanol, 25% glycerol, 500 μg/ml bovine serum albumin-fraction V), and 53 μl of a reaction solution prepared from 41 μl of distilled water. Further, 6 μl of a solution of a test compound in DMSO was added to each well, and 6 μl of DMSO was added to a well as a positive control (PC), followed by mixing well using a plate mixer. After the plate was incubated at 30°C for 30 minutes, 1 μl of a target DNA (5 pmol/μl) was added, and this was mixed well using a plate mixer.

After each plate was incubated at 30°C for 10 minutes, the reaction solution was discarded, followed by washing with a phosphate buffer two times. Then, an anti-digoxigenin antibody labeled with alkaline phosphatase (sheep Fab fragment: manufactured by Boehringer) was diluted 2000-fold with an antibody diluting solution, 100 μl of the diluent was added to bind at 30°C for 1 hour, and this was washed successively with a phosphate buffer containing 0.05 % Tween20 two times, and a phosphate buffer once. Then, 150 μl of an alkaline phosphatase coloring buffer (composition: 10 mM paranitrophenyl phosphate (manufactured by Vector Laboratories), 5 mM $MgCl_2$, 100 mM NaCl, 100 mM Tris-HCl (pH 9.5)) was added to react at 30°C for 2 hours, 50 μl of a 1N NaOH solution was added to stop the reaction, an absorbance (OD405 nm) of each well was measured, and an inhibition rate (IC_{50}) was obtained according to the following calculation equation.

Inhibition rate (%) = 100[1-{(C abs.- NC abs.) / (PC abs.- NC abs.)}]

C abs.: absorbance of well of compound

NC abs.: absorbance of NC

PC abs.: absorbance of PC

Results are shown below.

[Table 1]

Example No.	Integrase inhibitory activity (IC50, ng/ml)
C-2	3.3
F-2	3.8
H-2	3.2

The present compound showed the strong integrase inhibitory activity against HIV.

Experimental Example 2

A derivative of 293T cells expressing an attachment factor to improve adherence to plastic were used for the assay. A VSV-g pseudotyped HIV vector that expresses luciferase (herein referred to as PHIV) was produced by transfection of cells with the pGJ3-Luci vector plasmid (Jármy, G. et al., J. Medical Virology, 64:223-231, 2001) and pVSV-g (Clontech). Cells were mixed with the PHIV vector and then mixed with serially diluted compounds. After incubation at 37°C and 5% CO_2 for two days, the plates were read by using Steady Glo luciferase assay reagent (Promega) as recommended by the manufacturer. To assess non-HIV specific inhibition, a similar assay was performed, except that cell/PHIV vector mixture was replaced by cells which had been previously transduced and constitutively expressed luciferase.

[Table 2]

Example number	PHIV IC_{50} *=<10 nM, **=10-100 nM, ***>100 nM
Z-1	*
Z-2	*
Z-3	*
Z-4	*
Z-5	*
Z-6	*
Z-7	*
Z-8	**
Z-9	*
Z-10	*
Z-11	*
Z-12	*
Z-13	**
Z-14	**
Z-15	*
Z-16	*
Z-17	*
Z-18	*
Z-19	*
Z-20	**
Z-21	*
Z-22	*
Z-23	*

Z-24	*
Z-25	*
Z-26	*
Z-27	***
Z-28	*
Z-29	*
Z-30	*
Z-31	*
Z-32	*
Z-33	*
Z-34	*
Z-35	*
Z-36	*
Z-37	*
Z-38	**
Z-39	*
Z-40	*
Z-41	*
Z-42	*
Z-43	*
Z-44	*
Z-45	*
Z-46	*
Z-47	*
Z-48	*
Z-49	*
Z-50	*
Z-51	*
Z-52	*
Z-53	*
Z-54	*
Z-55	**
Z-59	*
Z-60	*

Formulation Example

A term "active ingredient" means the present compound, a tautomer thereof, a pharmaceutically acceptable salt thereof, or a solvate thereof.

(Formulation Example 1)

A hard gelatin capsule is prepared using the following ingredients:

	dose (mg/capsule)
Active ingredient	250
Starch (dried)	200
Magnesium stearate	10

Total	460mg

(Formulation Example 2)

A tablet is prepared using the following ingredients:


	dose (mg/tablet)
Active ingredient	250
Cellulose (microcrystalline)	400
Silicon dioxide (fumed)	10
Stearic acid	5
Total	665mg

Ingredients are mixed, and compressed to obtain tablets, each weighing 665 mg.

CLAIMS

[Claim 1]

A compound of the formula:



(I)

(wherein,

Z^1 is NR^4;


R^1 is hydrogen or lower alkyl;

X is a single bond, a heteroatom group selected from O, S, SO, SO_2 and NH, or lower alkylene or lower alkenylene each may be intervened by the heteroatom;

R^2 is optionally substituted aryl;

R^3 is hydrogen, halogen, hydroxy, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted lower alkenyloxy, optionally substituted aryl, optionally substituted aryloxy, optionally substituted heterocyclic group, optionally substituted heterocycleoxy or optionally substituted amino;

R^4 and Z^2 part taken together forms a ring, where the compound (I) is represented by the following formula (I-1), or (I-11):



(I-1)

(wherein,

A ring is optionally substituted 5- to 7-membered heterocycle, which contains 1 to 3 O, S, and/or N atom and the substituent is selected from Substituent group S2 and two of the substituents taken together with a neighboring atom(s) may form an optionally substituted carbocyle or an optionally substituted heterocycle

Substituent group S2: hydrogen, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted cycloalkyl lower alkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted lower alkenyloxy, optionally substituted aryl, optionally substituted aryl lower alkyl, optionally substituted aryloxy, optionally substituted heterocycle, optionally substituted heterocycle lower alkyl, optionally substituted heterocycleoxy, hydroxy, optionally substituted amino, optionally substituted lower alkylcarbonyl, optionally substituted cycloalkylcarbonyl, optionally substituted cycloalkyl lower alkylcarbonyl, optionally substituted lower alkoxycarbonyl, optionally substituted arylcarbonyl, optionally substituted aryl lower alkylcarbonyl, optionally substituted aryl oxycarbonyl, optionally substituted heterocyclecarbonyl, optionally substituted heterocycle lower alkylcarbonyl, optionally substituted heterocycleoxycarbonyl, optionally substituted aminocarbonyl, optionally substituted phosphoric acid residue, aryl substituted with optionally substituted phosphoric acid residue, aralkyl substituted with optionally substituted phosphoric acid residue, hydroxy substituted with optionally substituted phosphoric acid residue, amino substituted with optionally substituted phosphoric acid residue, or lower alkyl substituted with optionally substituted phosphoric acid residue (the lower alkyl may be intervened with a heteroatom group(s) selected from CO, O, S, SO, SO_2, NR^5 (R^5 is hydrogen, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted cycloalkyl lower alkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted aryl, optionally substituted aryl lower alkyl, optionally substituted aryloxy, optionally substituted heterocyclic group, optionally substituted heterocycle lower alkyl, optionally substituted heterocycleoxy, hydroxy, optionally substituted amino, optionally substituted phosphoric acid residue, aryl substituted with optionally substituted phosphoric acid residue, aralkyl substituted with optionally substituted phosphoric acid residue, hydroxy substituted with optionally substituted phosphoric acid residue, amino substituted with optionally substituted phosphoric acid residue or lower alkyl substituted with optionally

substituted phosphoric acid residue (the lower alkyl may be intervened by a heteroatom group selected from CO, O, S, SO, SO_2, NR^a (R^a is hydrogen or lower alkyl), -N= and =N-)), O or CH_2), -N= and =N-), and;

R^{14} and R^X are independently hydrogen, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted cycloalkyl lower alkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted lower alkenyloxy, optionally substituted aryl, optionally substituted aryl lower alkyl, optionally substituted aryloxy, optionally substituted heterocyclic group, optionally substituted heterocycle lower alkyl, optionally substituted heterocycleoxy, optionally substituted phosphoric acid residue, aryl substituted with optionally substituted phosphoric acid residue, aralkyl substituted with optionally substituted phosphoric acid residue, hydroxy substituted with optionally substituted phosphoric acid residue, amino substituted with optionally substituted phosphoric acid residue or lower alkyl substituted with optionally substituted phosphoric acid residue (the lower alkyl may be intervened by a heteroatom group selected from O, S, SO, SO_2, NR^5 (R^5 is hydrogen, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted cycloalkyl lower alkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted aryl, optionally substituted aryl lower alkyl, optionally substituted aryloxy, optionally substituted heterocyclic group, optionally substituted heterocycle lower alkyl, optionally substituted heterocycleoxy, hydroxy, optionally substituted amino, optionally substituted phosphoric acid residue, aryl substituted with optionally substituted phosphoric acid residue, aralkyl substituted with optionally substituted phosphoric acid residue, hydroxy substituted with optionally substituted phosphoric acid residue, amino substituted with optionally substituted phosphoric acid residue or lower alkyl substituted with optionally substituted phosphoric acid residue (the lower alkyl may be intervened by a heteroatom group selected from CO, O, S, SO, SO_2, NR^a (R^a is hydrogen or lower alkyl), -N= and =N-)), O or CH_2), -N= and =N-), hydroxy, optionally substituted amino, optionally substituted lower alkyl carbonyl, optionally substituted cycloalkylcarbonyl, optionally substituted cycloalkyl lower alkyl carbonyl, optionally substituted lower alkoxy carbonyl, optionally substituted arylcarbonyl, optionally substituted aryl lower alkyl carbonyl, optionally substituted aryloxycarbonyl, optionally substituted heterocyclecarbonyl, optionally substituted heterocycle lower alkyl carbonyl, optionally substituted heterocycleoxy carbonyl or optionally substituted aminocarbonyl;


a broken line represents the presence or absence of a bond, provided that when the broken line represents the presence of a bond, R^X is not present;

R^1 is hydrogen or lower alkyl;

X is a single bond, a heteroatom group selected from O, S, SO, SO_2 and NH, or lower alkylene or lower alkenylene each may be intervened by the heteroatom group;

R^2 is optionally substituted aryl;

R^3 is hydrogen, halogen, hydroxy, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted lower alkenyloxy, optionally substituted aryl, optionally substituted aryloxy, optionally substituted heterocyclic group, optionally substituted heterocycleoxy or optionally substituted amino)



(I-11)

(wherein,

D ring is optionally substituted 5- to 7-membered heterocycle; , which contains 1 to 3 O, S, and/or N atom and the substituent is selected from Substituent group S2 and two of the substituents taken together with a neighboring atom(s) may form an optionally substituted carbocyle or an optionally substituted heterocycle

Substituent group S2: hydrogen, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted cycloalkyl lower alkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted lower alkenyloxy, optionally substituted aryl, optionally substituted aryl lower alkyl, optionally substituted aryloxy, optionally substituted heterocycle, optionally substituted heterocycle lower alkyl, optionally substituted heterocycleoxy, hydroxy, optionally substituted amino, optionally substituted lower alkylcarbonyl, optionally substituted cycloalkylcarbonyl, optionally substituted cycloalkyl lower alkylcarbonyl, optionally substituted lower alkoxycarbonyl, optionally substituted arylcarbonyl, optionally substituted aryl lower alkylcarbonyl, optionally substituted aryl oxycarbonyl, optionally substituted heterocyclecarbonyl, optionally substituted

heterocycle lower alkylcarbonyl, optionally substituted heterocycleoxycarbonyl, optionally substituted aminocarbonyl, optionally substituted phosphoric acid residue, aryl substituted with optionally substituted phosphoric acid residue, aralkyl substituted with optionally substituted phosphoric acid residue, hydroxy substituted with optionally substituted phosphoric acid residue, amino substituted with optionally substituted phosphoric acid residue, or lower alkyl substituted with optionally substituted phosphoric acid residue (the lower alkyl may be intervened with a heteroatom group(s) selected from CO, O, S, SO, SO_2, NR^5 (R^5 is hydrogen, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted cycloalkyl lower alkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted aryl, optionally substituted aryl lower alkyl, optionally substituted aryloxy, optionally substituted heterocyclic group, optionally substituted heterocycle lower alkyl, optionally substituted heterocycleoxy, hydroxy, optionally substituted amino, optionally substituted phosphoric acid residue, aryl substituted with optionally substituted phosphoric acid residue, aralkyl substituted with optionally substituted phosphoric acid residue, hydroxy substituted with optionally substituted phosphoric acid residue, amino substituted with optionally substituted phosphoric acid residue or lower alkyl substituted with optionally substituted phosphoric acid residue (the lower alkyl may be intervened by a heteroatom group selected from CO, O, S, SO, SO_2, NR^a (R^a is hydrogen or lower alkyl), -N= and =N-)), O or CH_2), -N= and =N-), and;

R^1 is hydrogen or lower alkyl;

X is a single bond, a heteroatom group selected from O, S, SO, SO_2 and NH, or lower alkylene or lower alkenylene each may be intervened by the heteroatom group;

R^2 is optionally substituted aryl;

R^3 is hydrogen, halogen, hydroxy, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted lower alkenyloxy, optionally substituted aryl, optionally substituted aryloxy, optionally substituted heterocyclic group, optionally substituted heterocycleoxy or optionally substituted amino));

or a pharmaceutically acceptable salt, or solvate thereof.

[Claim 2]

A compound according to Claim 1, pharmaceutically acceptable salt, or solvate thereof, wherein R^1 is hydrogen.

[Claim 3]

A compound according to Claim 1, pharmaceutically acceptable salt, or solvate thereof, wherein X is lower alkylene; R^2 is phenyl or phenyl substituted with at least halogen.

[Claim 4]

A compound according to Claim 1, pharmaceutically acceptable salt, or solvate thereof, wherein R^3 is hydrogen, halogen, hydroxy, lower alkyl, lower alkenyl, lower alkoxy, lower alkenyloxy or optionally substituted amino.

[Claim 5]


A compound according to Claim 1, pharmaceutically acceptable salt, or solvate thereof, wherein R^3 is hydrogen.

[Claim 6]

A compound according to Claim 1, pharmaceutically acceptable salt, or solvate thereof, wherein R^1 is hydrogen or lower alkyl; X is lower alkylene; R^2 is phenyl or phenyl substituted with at least halogen; R^3 is hydrogen, halogen, hydroxy, lower alkyl, lower alkenyl, lower alkoxy, lower alkenyloxy or optionally substituted amino.

[Claim 7]

A compound of the formula:



(I-1)

(wherein,

A ring is optionally substituted 5- to 7-membered heterocycle, which contains 1 to 3 O, S, and/or N atom and the substituent is selected from Substituent group S2 and two of the substituents taken together with a neighboring atom(s) may form an optionally substituted carbocyle or an optionally substituted heterocycle

Substituent group S2: hydrogen, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted cycloalkyl lower alkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted lower alkenyloxy, optionally substituted aryl, optionally substituted aryl lower alkyl, optionally substituted aryloxy, optionally substituted heterocycle, optionally substituted heterocycle lower alkyl, optionally substituted heterocycleoxy, hydroxy, optionally substituted amino, optionally substituted lower alkylcarbonyl, optionally substituted cycloalkylcarbonyl, optionally substituted cycloalkyl lower alkylcarbonyl, optionally substituted lower alkoxycarbonyl, optionally substituted arylcarbonyl, optionally substituted aryl lower alkylcarbonyl, optionally substituted aryl oxycarbonyl, optionally substituted heterocyclecarbonyl, optionally substituted heterocycle lower alkylcarbonyl, optionally substituted heterocycleoxycarbonyl, optionally substituted aminocarbonyl, optionally substituted phosphoric acid residue, aryl substituted with optionally substituted phosphoric acid residue, aralkyl substituted with optionally substituted phosphoric acid residue, hydroxy substituted with optionally substituted phosphoric acid residue, amino substituted with optionally substituted phosphoric acid residue, or lower alkyl substituted with optionally substituted phosphoric acid residue (the lower alkyl may be intervened with a heteroatom group(s) selected from CO, O, S, SO, SO_2, NR^5 (R^5 is hydrogen, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted cycloalkyl lower alkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted aryl, optionally substituted aryl lower alkyl, optionally substituted aryloxy, optionally substituted heterocyclic group, optionally substituted heterocycle lower alkyl, optionally substituted heterocycleoxy, hydroxy, optionally substituted amino, optionally substituted phosphoric acid residue, aryl substituted with optionally substituted phosphoric acid residue, aralkyl substituted with optionally substituted phosphoric acid residue, hydroxy substituted with optionally substituted phosphoric acid residue, amino substituted with optionally substituted phosphoric acid residue or lower alkyl substituted with optionally substituted phosphoric acid residue (the lower alkyl may be intervened by a heteroatom group selected from CO, O, S, SO, SO_2, NR^a (R^a is hydrogen or lower alkyl), -N= and =N-)), O or CH_2), -N= and =N-), and;

R^{14} and R^X are independently hydrogen, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted cycloalkyl lower alkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally

substituted lower alkenyloxy, optionally substituted aryl, optionally substituted aryl lower alkyl, optionally substituted aryloxy, optionally substituted heterocyclic group, optionally substituted heterocycle lower alkyl, optionally substituted heterocycleoxy, optionally substituted phosphoric acid residue, aryl substituted with optionally substituted phosphoric acid residue, aralkyl substituted with optionally substituted phosphoric acid residue, hydroxy substituted with optionally substituted phosphoric acid residue, amino substituted with optionally substituted phosphoric acid residue or lower alkyl substituted with optionally substituted phosphoric acid residue (the lower alkyl may be intervened by a heteroatom group selected from O, S, SO, SO_2, NR^5 (R^5 is hydrogen, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted cycloalkyl lower alkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted aryl, optionally substituted aryl lower alkyl, optionally substituted aryloxy, optionally substituted heterocyclic group, optionally substituted heterocycle lower alkyl, optionally substituted heterocycleoxy, hydroxy, optionally substituted amino, optionally substituted phosphoric acid residue, aryl substituted with optionally substituted phosphoric acid residue, aralkyl substituted with optionally substituted phosphoric acid residue, hydroxy substituted with optionally substituted phosphoric acid residue, amino substituted with optionally substituted phosphoric acid residue or lower alkyl substituted with optionally substituted phosphoric acid residue (the lower alkyl may be intervened by a heteroatom group selected from CO, O, S, SO, SO_2, NR^a (R^a is hydrogen or lower alkyl), -N= and =N-)), O or CH_2), -N= and =N-), hydroxy, optionally substituted amino, optionally substituted lower alkyl carbonyl, optionally substituted cycloalkylcarbonyl, optionally substituted cycloalkyl lower alkyl carbonyl, optionally substituted lower alkoxy carbonyl, optionally substituted arylcarbonyl, optionally substituted aryl lower alkyl carbonyl, optionally substituted aryloxycarbonyl, optionally substituted heterocyclecarbonyl, optionally substituted heterocycle lower alkyl carbonyl, optionally substituted heterocycleoxy carbonyl or optionally substituted aminocarbonyl;

a broken line represents the presence or absence of a bond, provided that when the broken line represents the presence of a bond, R^X is not present;

R^1 is hydrogen or lower alkyl;

X is a single bond, a heteroatom group selected from O, S, SO, SO_2 and NH, or lower alkylene or lower alkenylene each may be intervened by the heteroatom group;

R^2 is optionally substituted aryl;

R^3 is hydrogen, halogen, hydroxy, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted lower alkenyloxy, optionally substituted aryl, optionally substituted aryloxy, optionally substituted heterocyclic group, optionally substituted heterocycleoxy or optionally substituted amino);
or a pharmaceutically acceptable salt, or solvate thereof


[Claim 8]
A compound according to Claim 7, pharmaceutically acceptable salt, or solvate thereof, wherein R^1 is hydrogen or lower alkyl; X is lower alkylene; R^2 is phenyl or phenyl substituted with at least halogen; R^3 is hydrogen, halogen, hydroxy, lower alkyl, lower alkenyl, lower alkoxy, lower alkenyloxy or optionally substituted amino.

[Claim 9]
A compound according to Claim 7, pharmaceutically acceptable salt, or solvate thereof, wherein a broken line represents the absence of a bond.

[Claim 10]
A compound according to Claim 7, pharmaceutically acceptable salt, or solvate thereof, wherein R^X is hydrogen; R^{14} is hydrogen or optionally substituted lower alkyl.

[Claim 11]
A compound according to Claim 7, pharmaceutically acceptable salt, or solvate thereof, wherein A ring is optionally substituted 5- to 7-membered heterocycle, which contains 1 to 2 O, S, and/or N atom and the substituent is selected from said Substituent group S2 and two of the substituents taken together with a neighboring atom(s) may form an optionally substituted carbocyle or an optionally substituted heterocycle.

[Claim 12]
A compound of the formula:



(I-1-1)

(wherein,

A ring is optionally substituted 5- to 7-membered heterocycle, which contains 1 to 2 O, S, and/or N atom and the substituent is selected from Substituent group S2 and two of the substituents taken together with a neighboring atom(s) may form an optionally substituted carbocyle or an optionally substituted heterocycle

Substituent group S2: hydrogen, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted cycloalkyl lower alkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted lower alkenyloxy, optionally substituted aryl, optionally substituted aryl lower alkyl, optionally substituted aryloxy, optionally substituted heterocycle, optionally substituted heterocycle lower alkyl, optionally substituted heterocycleoxy, hydroxy, optionally substituted amino, optionally substituted lower alkylcarbonyl, optionally substituted cycloalkylcarbonyl, optionally substituted cycloalkyl lower alkylcarbonyl, optionally substituted lower alkoxycarbonyl, optionally substituted arylcarbonyl, optionally substituted aryl lower alkylcarbonyl, optionally substituted aryl oxycarbonyl, optionally substituted heterocyclecarbonyl, optionally substituted heterocycle lower alkylcarbonyl, optionally substituted heterocycleoxycarbonyl, optionally substituted aminocarbonyl, optionally substituted phosphoric acid residue, aryl substituted with optionally substituted phosphoric acid residue, aralkyl substituted with optionally substituted phosphoric acid residue, hydroxy substituted with optionally substituted phosphoric acid residue, amino substituted with optionally substituted phosphoric acid residue, or lower alkyl substituted with optionally substituted phosphoric acid residue (the lower alkyl may be intervened with a heteroatom group(s) selected from CO, O, S, SO, SO_2, NR^5 (R^5 is hydrogen, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted cycloalkyl lower alkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted aryl, optionally substituted aryl lower alkyl, optionally substituted aryloxy, optionally substituted heterocyclic group, optionally

substituted heterocycle lower alkyl, optionally substituted heterocycleoxy, hydroxy, optionally substituted amino, optionally substituted phosphoric acid residue, aryl substituted with optionally substituted phosphoric acid residue, aralkyl substituted with optionally substituted phosphoric acid residue, hydroxy substituted with optionally substituted phosphoric acid residue, amino substituted with optionally substituted phosphoric acid residue or lower alkyl substituted with optionally substituted phosphoric acid residue (the lower alkyl may be intervened by a heteroatom group selected from CO, O, S, SO, SO_2, NR^a (R^a is hydrogen or lower alkyl), -N= and =N-)), O or CH_2), -N= and =N-), and;

the stereochemistry of an asymmetric carbon represented by * shows R- or S-configuration, or a mixture thereof;

R^{14} and R^X are independently hydrogen, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted cycloalkyl lower alkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted lower alkenyloxy, optionally substituted aryl, optionally substituted aryl lower alkyl, optionally substituted aryloxy, optionally substituted heterocyclic group, optionally substituted heterocycle lower alkyl, optionally substituted heterocycleoxy, optionally substituted phosphoric acid residue, aryl substituted with optionally substituted phosphoric acid residue, aralkyl substituted with optionally substituted phosphoric acid residue, hydroxy substituted with optionally substituted phosphoric acid residue, amino substituted with optionally substituted phosphoric acid residue or lower alkyl substituted with optionally substituted phosphoric acid residue (the lower alkyl may be intervened by a heteroatom group selected from O, S, SO, SO_2, NR^5 (R^5 is hydrogen, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted cycloalkyl lower alkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted aryl, optionally substituted aryl lower alkyl, optionally substituted aryloxy, optionally substituted heterocyclic group, optionally substituted heterocycle lower alkyl, optionally substituted heterocycleoxy, hydroxy, optionally substituted amino, optionally substituted phosphoric acid residue, aryl substituted with optionally substituted phosphoric acid residue, aralkyl substituted with optionally substituted phosphoric acid residue, hydroxy substituted with optionally substituted phosphoric acid residue, amino substituted with optionally substituted phosphoric acid residue or lower alkyl substituted with optionally substituted phosphoric acid residue (the lower alkyl may be intervened by a heteroatom group selected from CO, O, S, SO, SO_2, NR^a (R^a is

hydrogen or lower alkyl), -N= and =N-)), O or CH_2), -N= and =N-), hydroxy, optionally substituted amino, optionally substituted lower alkyl carbonyl, optionally substituted cycloalkylcarbonyl, optionally substituted cycloalkyl lower alkyl carbonyl, optionally substituted lower alkoxy carbonyl, optionally substituted arylcarbonyl, optionally substituted aryl lower alkyl carbonyl, optionally substituted aryloxycarbonyl, optionally substituted heterocyclecarbonyl, optionally substituted heterocycle lower alkyl carbonyl, optionally substituted heterocycleoxy carbonyl or optionally substituted aminocarbonyl;

R^3 is hydrogen, halogen, hydroxy, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted lower alkenyloxy, optionally substituted aryl, optionally substituted aryloxy, optionally substituted heterocyclic group, optionally substituted heterocycleoxy or optionally substituted amino), its pharmaceutically acceptable salt, or

R^1 is hydrogen or lower alkyl;

R is independently selected from halogen and Substituent group S1;

Substituent group S1(: optionally substituted phosphoric acid residue, aryl substituted with optionally substituted phosphoric acid residue, aralkyl substituted with optionally substituted phosphoric acid residue, hydroxy substituted with optionally substituted phosphoric acid residue, amino substituted with optionally substituted phosphoric acid residue, or lower alkyl substituted with optionally substituted phosphoric acid residue (wherein the lower alkyl may be intervened with a heteroatom group(s) selected from CO, O, O, S, SO, SO_2, NR^a (R^a is hydrogen or lower alkyl), -N= and =N-), lower alkoxy lower alkyl, amino lower alkyl optionally substituted with mono- or di- lower alkyl, halogenated lower alkyl, lower alkoxy, carbamoyl optionally substituted with mono- or di- lower alkyl, optionally substituted lower alkyl sulfonyl amino, halogenated lower alkoxy, hydroxy lower alkyl)

m is an integer of 0 to 3);

or a pharmaceutically acceptable salt, or solvate thereof.

[Claim 13]

A compound according to Claim 12, pharmaceutically acceptable salt, or solvate thereof, wherein R^X and R^{14} are independently hydrogen or optionally substituted

lower alkyl.

[Claim 14]

A compound according to Claim 12, pharmaceutically acceptable salt, or solvate thereof, wherein R^X and R^{14} are hydrogens.

[Claim 15]


A compound according to Claim 12, pharmaceutically acceptable salt, or solvate thereof, wherein R^3 is hydrogen.

[Claim 16]

A compound according to Claim 12, pharmaceutically acceptable salt, or solvate thereof, wherein m is 0, or 1 to 3 and at least one of R is halogen.


[Claim 17]

A compound according to Claim 7 or 12, pharmaceutically acceptable salt, or solvate thereof, wherein A ring is any one of the followings:




Z = O or NR^{26}

(A-1)



Z = O or NR^{31}

(A-2)



Z = O or NR^{40}

(A-3)

(wherein, R^{20} to R^{40} are each independently a group selected from Substituent group S2, or any two groups of R^{20} to R^{40}, which bonds to the same carbon atom, taken together with the carbon atom, may form an optionally substituted carbocyle or optionally substituted heterocycle, or each combination of (R^{20} and R^{22}), (R^{23} and R^{24}), (R^{25} and R^{26}), (R^{27} and R^{29}), (R^{30} and R^{31}), (R^{32} and R^{34}), (R^{35} and R^{36}), (R^{37} and R^{38}), and (R^{39}andR^{40}), taken together with the neighboring atom, may form an optionally substituted carbocyle or optionally substituted heterocycle.

Substituent group S2: hydrogen, optionally substituted lower alkyl, optionally

substituted cycloalkyl, optionally substituted cycloalkyl lower alkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted lower alkenyloxy, optionally substituted aryl, optionally substituted aryl lower alkyl, optionally substituted aryloxy, optionally substituted heterocycle, optionally substituted heterocycle lower alkyl, optionally substituted heterocycleoxy, hydroxy, optionally substituted amino, optionally substituted lower alkylcarbonyl, optionally substituted cycloalkylcarbonyl, optionally substituted cycloalkyl lower alkylcarbonyl, optionally substituted lower alkoxycarbonyl, optionally substituted arylcarbonyl, optionally substituted aryl lower alkylcarbonyl, optionally substituted aryl oxycarbonyl, optionally substituted heterocyclecarbonyl, optionally substituted heterocycle lower alkylcarbonyl, optionally substituted heterocycleoxycarbonyl, optionally substituted aminocarbonyl, optionally substituted phosphoric acid residue, aryl substituted with optionally substituted phosphoric acid residue, aralkyl substituted with optionally substituted phosphoric acid residue, hydroxy substituted with optionally substituted phosphoric acid residue, amino substituted with optionally substituted phosphoric acid residue, or lower alkyl substituted with optionally substituted phosphoric acid residue (the lower alkyl may be intervened with a heteroatom group(s) selected from CO, O, S, SO, SO_2, NR^5 (R^5 is hydrogen, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted cycloalkyl lower alkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted aryl, optionally substituted aryl lower alkyl, optionally substituted aryloxy, optionally substituted heterocyclic group, optionally substituted heterocycle lower alkyl, optionally substituted heterocycleoxy, hydroxy, optionally substituted amino, optionally substituted phosphoric acid residue, aryl substituted with optionally substituted phosphoric acid residue, aralkyl substituted with optionally substituted phosphoric acid residue, hydroxy substituted with optionally substituted phosphoric acid residue, amino substituted with optionally substituted phosphoric acid residue or lower alkyl substituted with optionally substituted phosphoric acid residue (the lower alkyl may be intervened by a heteroatom group selected from CO, O, S, SO, SO_2, NR^a (R^a is hydrogen or lower alkyl), -N= and =N-)), O or CH_2), -N= and =N-)

the stereochemistry of an asymmetric carbon represented by * shows R- or S-configuration, or a mixture thereof)

[Claim 18]

A compound according to Claim 17, pharmaceutically acceptable salt, or solvate thereof, wherein R^{20} to R^{40} are each independently hydrogen or substituted lower alkyl, or any two groups of R^{20} to R^{40}, which bonds to the same carbon atom, taken together with the carbon atom, may form an optionally substituted 3- to 7- membered carbocyle or optionally substituted 3- to 7- membered heterocycle, or each combination of (R^{20} and R^{22}), (R^{23} and R^{24}), (R^{25} and R^{26}), (R^{27} and R^{29}), (R^{30} and R^{31}), (R^{32} and R^{34}), (R^{35} and R^{36}), (R^{37} and R^{38}), and (R^{39}andR^{40}), taken together with the neighboring atom, may form an optionally substituted 5- to 7- membered carbocyle or optionally substituted 5- to 7- membered heterocycle.

[Claim 19]

A compound according to Claim 17, pharmaceutically acceptable salt, or solvate thereof, wherein A ring is a ring represented by (A-1); one of R^{20} to R^{25} is optionally substituted lower alkyl and the others are hydrogens.

[Claim 20]

A compound according to Claim 17, pharmaceutically acceptable salt, or solvate thereof, wherein A ring is a ring represented by (A-1); one of (R^{20} and R^{22}), (R^{23} and R^{24}), and (R^{25} and R^{26}), taken together with the neighboring atom, may form an optionally substituted 5- to 7- membered carbocyle or optionally substituted 5- to 7- membered heterocycle.

[Claim 21]

A compound according to Claim 17, pharmaceutically acceptable salt, or solvate thereof, wherein A ring is a ring represented by (A-1); $Z=NR^{26}$, and R^{25} and R^{26} taken together with the neighboring atom may form an optionally substituted 5- to 7- membered heterocycle.

[Claim 22]

A compound according to Claim 17, pharmaceutically acceptable salt, or solvate thereof, wherein A ring is a ring represented by (A-2); one of R^{27} to R^{30} is optionally substituted lower alkyl and the others are hydrogens.

[Claim 23]

A compound according to Claim 17, pharmaceutically acceptable salt, or solvate thereof, wherein A ring is a ring represented by (A-2); one of ($R^{27}$and R^{29}) and (R^{30} and R^{31}), taken together with the neighboring atom, may form an optionally substituted 5- to 7- membered carbocyle or optionally substituted 5- to 7- membered heterocycle.

[Claim 24]

A compound according to Claim 17, pharmaceutically acceptable salt, or solvate thereof, wherein A ring is a ring represented by (A-2); Z=NR^{31}, and R^{30} and R^{31} taken together with the neighboring atom may form an optionally substituted 5- to 7- membered heterocycle.

[Claim 25]

A compound according to Claim 17, pharmaceutically acceptable salt, or solvate thereof, wherein A ring is a ring represented by (A-3); one of R^{32} to R^{39} is optionally substituted lower alkyl and the others are hydrogens.

[Claim 26]

A compound according to Claim 17, pharmaceutically acceptable salt, or solvate thereof, wherein A ring is a ring represented by (A-3); one of (R^{32}andR^{34}),(R^{35}andR^{36}),(R^{37}andR^{38}),and(R^{39}andR^{40}), taken together with the neighboring atom, may form an optionally substituted 5- to 7- membered carbocyle or optionally substituted 5- to 7- membered heterocycle.

[Claim 27]

A compound according to Claim 17, pharmaceutically acceptable salt, or solvate thereof, wherein A ring is a ring represented by (A-3); Z=NR^{40}, and R^{39} and R^{40} taken together with the neighboring atom may form an optionally substituted 5- to 7- membered heterocycle.

[Claim 28]


A compound according to Claim 12, pharmaceutically acceptable salt, or solvate thereof, wherein R^X is hydrogen; R^{14} is hydrogen or optionally substituted lower; R^3 is hydrogen; m is 1 to 3 and at least one of Rs is halogen; A ring is a ring described in Claim 17.

[Claim 29]

A compound according to Claim 12, pharmaceutically acceptable salt, or solvate thereof, wherein R^X is hydrogen; R^{14} is hydrogen; R^3 is hydrogen; m is 0, or 1 to 3 and at least one of R is halogen; A ring is a ring described in Claim 17; R^{20} to R^{40} are each independently hydrogen or substituted lower alkyl, or any two groups of R^{20} to R^{40}, which bonds to the same carbon atom, taken together with the carbon atom, may form an optionally substituted 3- to 7- membered carbocyle or optionally substituted 3- to 7- membered heterocycle, or each combination of (R^{20} and R^{22}), (R^{23} and R^{24}), (R^{25} and R^{26}), (R^{27} and R^{29}), (R^{30} and R^{31}), (R^{32} and R^{34}), (R^{35} and R^{36}), (R^{37} and R^{38}), and (R^{39}andR^{40}), taken together with the neighboring carbon atom, may form an optionally substituted 5- to 7- membered carbocyle or optionally substituted 5- to 7- membered heterocycle.

[Claim 30]

A compound of the formula:



(I-11)

(wherein,

D ring is optionally substituted 5- to 7-membered heterocycle; , which contains 1 to 3 O, S, and/or N atom and the substituent is selected from Substituent group S2 and two of the substituents taken together with a neighboring atom(s) may form an optionally substituted carbocyle or an optionally substituted heterocycle

Substituent group S2: hydrogen, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted cycloalkyl lower alkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted lower alkenyloxy, optionally substituted aryl, optionally substituted aryl lower alkyl, optionally substituted aryloxy, optionally substituted heterocycle, optionally substituted heterocycle lower alkyl, optionally substituted heterocycleoxy, hydroxy, optionally substituted amino, optionally substituted lower alkylcarbonyl, optionally substituted cycloalkylcarbonyl, optionally substituted cycloalkyl lower alkylcarbonyl,

optionally substituted lower alkoxycarbonyl, optionally substituted arylcarbonyl, optionally substituted aryl lower alkylcarbonyl, optionally substituted aryl oxycarbonyl, optionally substituted heterocyclecarbonyl, optionally substituted heterocycle lower alkylcarbonyl, optionally substituted heterocycleoxycarbonyl, optionally substituted aminocarbonyl, optionally substituted phosphoric acid residue, aryl substituted with optionally substituted phosphoric acid residue, aralkyl substituted with optionally substituted phosphoric acid residue, hydroxy substituted with optionally substituted phosphoric acid residue, amino substituted with optionally substituted phosphoric acid residue, or lower alkyl substituted with optionally substituted phosphoric acid residue (the lower alkyl may be intervened with a heteroatom group(s) selected from CO, O, S, SO, SO_2, NR^5 (R^5 is hydrogen, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted cycloalkyl lower alkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted aryl, optionally substituted aryl lower alkyl, optionally substituted aryloxy, optionally substituted heterocyclic group, optionally substituted heterocycle lower alkyl, optionally substituted heterocycleoxy, hydroxy, optionally substituted amino, optionally substituted phosphoric acid residue, aryl substituted with optionally substituted phosphoric acid residue, aralkyl substituted with optionally substituted phosphoric acid residue, hydroxy substituted with optionally substituted phosphoric acid residue, amino substituted with optionally substituted phosphoric acid residue or lower alkyl substituted with optionally substituted phosphoric acid residue (the lower alkyl may be intervened by a heteroatom group selected from CO, O, S, SO, SO_2, NR^a (R^a is hydrogen or lower alkyl), -N= and =N-)), O or CH_2), -N= and =N-), and;

R^1 is hydrogen or lower alkyl;

X is a single bond, a heteroatom group selected from O, S, SO, SO_2 and NH, or lower alkylene or lower alkenylene each may be intervened by the heteroatom group;

R^2 is optionally substituted aryl;

R^3 is hydrogen, halogen, hydroxy, optionally substituted lower alkyl, optionally substituted cycloalkyl, optionally substituted lower alkenyl, optionally substituted lower alkoxy, optionally substituted lower alkenyloxy, optionally substituted aryl, optionally substituted aryloxy, optionally substituted heterocyclic group, optionally substituted heterocycleoxy or optionally substituted amino);

or a pharmaceutically acceptable salt, or solvate thereof

[Claim 31]

A compound selected from the group consisting of:

(3*R*,11a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-methyl-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(4a*R*,13a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-10-hydroxy-9,11-dioxo-2,3,4a,5,9,11,13,13a-octahydro-1*H*-pyrido[1,2-*a*]pyrrolo[1',2':3,4]imidazo[1,2-*d*]pyrazine-8-carboxamide;

(3a*S*,13a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-8-hydroxy-7,9-dioxo-1,2,3,3a,4,5,7,9,13,13a-decahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrrolo[1,2-*c*]pyrimidine-10-carboxamide;

(4a*S*,13a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-10-hydroxy-9,11-dioxo-2,3,4a,5,9,11,13,13a-octahydro-1*H*-pyrido[1,2-*a*]pyrrolo[1',2':3,4]imidazo[1,2-*d*]pyrazine-8-carboxamide;

(4a*S*,13a*R*)-*N*-[(4-Fluorophenyl)methyl]-10-hydroxy-9,11-dioxo-2,3,4a,5,9,11,13,13a-octahydro-1*H*-pyrido[1,2-*a*]pyrrolo[1',2':3,4]imidazo[1,2-*d*]pyrazine-8-carboxamide;

(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-5,7-dioxo-3-(phenylmethyl)-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3a*S*,13a*S*)-*N*-[(4-Fluorophenyl)methyl]-8-hydroxy-7,9-dioxo-1,2,3,3a,4,5,7,9,13,13a-decahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrrolo[1,2-*c*]pyrimidine-10-carboxamide;

(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-[(1*S*)-1-methylpropyl]-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-methyl-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3*S*,11a*R*)-*N*-[(4-Fluorophenyl)methyl]-6-hydroxy-3-methyl-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-3-(1,1-dimethylethyl)-6-hydroxy-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3*S*,11a*R*)-3-(1,1-Dimethylethyl)-*N*-[(4-fluorophenyl)methyl]-6-hydroxy-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-5,7-dioxo-3-phenyl-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-(hydroxymethyl)-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(2*S*,3*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-methyl-5,7-dioxo-2-phenyl-2,3,5,7

,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3*R*,11a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-5,7-dioxo-3-(phenylmethyl)-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3*R*,11a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-(2-methylpropyl)-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(5a*R*,14a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-11-hydroxy-10,12-dioxo-1,2,3,4,5a,6,10,12,14,14a-decahydropyrido[1,2-*a*]pyrido[1',2':3,4]imidazo[1,2-*d*]pyrazine-9-carboxamide;

(2*S*,3*S*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-[(methyloxy)methyl]-5,7-dioxo-2-phenyl-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3*S*,11a*R*)-3-(Cyclohexylmethyl)-*N*-[(2,4-difluorophenyl)methyl]-6-hydroxy-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-(1-methylethyl)-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(5a*R*,14a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-12-hydroxy-11,13-dioxo-5a,6a,7,11,13,14a-hexahydro-5*H*-indeno[1',2':4,5][1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-10-carboxamid

e;

(2*S*,3*R*,11a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-5,7-dioxo-2,3-diphenyl-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(2*S*,3*R*,11a*R*)-*N*-[(2,4-difluorophenyl)methyl]-6-hydroxy-5,7-dioxo-2,3-diphenyl-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3*R*,11a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-(1-methylethyl)-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-[2-(methylthio)ethyl]-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-[2-(methylsulfonyl)ethyl]-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-(1*H*-indol-3-ylmethyl)-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(4*R*,12a*R*)-*N*-[(4-fluorophenyl)methyl]-7-hydroxy-4-methyl-1-(2-methylpropyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(4*R*,12a*R*)-*N*-[(4-Fluorophenyl)methyl]-7-hydroxy-4-methyl-1-(1-methylethyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-4-methyl-1-(2-methylpropyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(4*S*,12a*S*)-1-(Cyclopropylmethyl)-*N*-[(2,4-difluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-1-(2-furanylmethyl)-7-hydroxy-4-methyl-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1-(1,3-thiazol-2-ylmethyl)-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(4a*R*,6a*R*,14a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-12-hydroxy-11,13-dioxo-1,3,4,4a,5,6a,7,11,13,14a-decahydro-2*H*-pyrido[1',2':4,5]pyrazino[1,2-*a*][3,1]benzoxazine-10-carboxa

mide;

(4a*R*,6a*R*,14a*S*)-*N*-[(4-Fluorophenyl)methyl]-12-hydroxy-11,13-dioxo-1,3,4,4a,5,6a,7,11,13,14a-decahydro-2*H*-pyrido[1',2':4,5]pyrazino[1,2-*a*][3,1]benzoxazine-10-carboxamide;

(3*S*,4a*R*,6a*R*,14a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-12-hydroxy-11,13-dioxo-3-phenyl-1,3,4,4a,5,6a,7,11,13,14a-decahydro-2*H*-pyrido[1',2':4,5]pyrazino[1,2-*a*][3,1]benzoxazine-10-carboxamide;

(4aS,6aS,14aS)-N-[(2,4-Difluorophenyl)methyl]-12-hydroxy-6-(2-methylpropyl)-11,13-dioxo-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide;

(6aR,7aS,11aS)-N-[(2,4-Difluorophenyl)methyl]-1-hydroxy-2,13-dioxo-2,6a,7,7a,8,9,10,11,11a,13-decahydro-6H-pyrido[1',2':4,5]pyrazino[1,2-a]benzimidazole-3-carboxamide;

(6aS,7aS,11aS)-N-[(2,4-Difluorophenyl)methyl]-1-hydroxy-2,13-dioxo-2,6a,7,7a,8,9,10,11,11a,13-decahydro-6H-pyrido[1',2':4,5]pyrazino[1,2-a]benzimidazole-3-carboxamide;

(5aS,14aS)-N-[(2,4-Difluorophenyl)methyl]-11-hydroxy-10,12-dioxo-1,2,3,4,5a,6,10,12,14,14a-decahydropyrido[1,2-a]pyrido[1',2':3,4]imidazo[1,2-d]pyrazine-9-carboxamide;

(4aR,14aR)-N-[(2,4-Difluorophenyl)methyl]-9-hydroxy-8,10-dioxo-2,3,4,4a,5,6,8,10,14,14a-decahydro-1H-pyrido[1,2-c]pyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-11-carboxamide;

(4*R*,12a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-4-methyl-1-(3-methylbutyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-4-methyl-1-(1-methylethyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-4-methyl-1-(3-methylbutyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1-(3-pyridinylmethyl)-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(4*S*,12a*S*)-1-Cyclopropyl-*N*-[(2,4-difluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-

1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-4-methyl-1-[2-(methyloxy)ethyl]-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(3aS,5aS,13aS)-N-[(2,4-Difluorophenyl)methyl]-11-hydroxy-5-(2-methylpropyl)-10,12-dioxo-2,3,3a,4,5,5a,6,10,12,13a-decahydro-1H-cyclopenta[e]pyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-9-carboxamide;

(3*R*,11a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-3-ethyl-6-hydroxy-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(4aS,6aS,14aS)-N-[(2,4-Difluorophenyl)methyl]-12-hydroxy-6-[2-(4-morpholinyl)ethyl]-11,13-dioxo-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide;

(3aR,5aR,13aS)-N-[(2,4-Difluorophenyl)methyl]-11-hydroxy-10,12-dioxo-1,2,3,3a,4,5a,6,10,12,13a-decahydrocyclopenta[d]pyrido[1',2':4,5]pyrazino[2,1-b][1,3]oxazine-9-carboxamide;

(4aS,6aS,14aS)-N-[(2,4-Difluorophenyl)methyl]-12-hydroxy-6-methyl-11,13-dioxo-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-car

boxamide;

(4aS,6aS,14aS)-N-[(2,4-Difluorophenyl)methyl]-12-hydroxy-6-[2-(methyloxy)ethyl]-11,13-dioxo-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide;

(4aS,6aS,14aS)-6-[2-(Acetylamino)ethyl]-N-[(2,4-difluorophenyl)methyl]-12-hydroxy-11,13-dioxo-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide;

(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-3-ethyl-6-hydroxy-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3*S*,11a*R*)-3-Butyl-*N*-[(2,4-difluorophenyl)methyl]-6-hydroxy-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-[(4-hydroxyphenyl)methyl]-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(4S,12aS)-1-Cyclobutyl-N-[(2,4-difluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-9-carboxamide;

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1-(tetrahydro-2*H*-thiopyran-4-yl)-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-1,4-bis(2-methylpropyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(4aS,6aS,14aS)-N-[(2,4-Difluorophenyl)methyl]-12-hydroxy-6-(2-hydroxyethyl)-11,13-dioxo-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide;

(4aS,6aS,14aS)-6-Cyclopropyl-N-[(2,4-difluorophenyl)methyl]-12-hydroxy-11,13-dioxo-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide;

(4aS,6aS,14aS)-N-[(2,4-Difluorophenyl)methyl]-12-hydroxy-11,13-dioxo-6-[2-(1-pyrrolidinyl)ethyl]-1,2,3,4,4a,5,6,6a,7,11,13,14a-dodecahydropyrido[1',2':4,5]pyrazino[1,2-a]quinazoline-10-carboxamide;

(4aS,14aS)-N-[(2,4-Difluorophenyl)methyl]-9-hydroxy-8,10-dioxo-2,3,4,4a,5,6,8,10,14,14a-decahydro-1H-pyrido[1,2-c]pyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-11-carboxamide;

(4S,12aS)-N-[(4-Fluorophenyl)methyl]-7-hydroxy-4-methyl-1-[2-(methyloxy)ethyl]-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-9-carboxamide;

(4S,12aS)-1-Cyclobutyl-N-[(4-fluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-9-carboxamide;

(4S,12aS)-N-[(4-Fluorophenyl)methyl]-7-hydroxy-4-methyl-1-(2-methylpropyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-9-carboxamide;

(4S,12aS)-N-[(4-Fluorophenyl)methyl]-7-hydroxy-1,4-dimethyl-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-a]pyrimidine-9-carboxamide;

(4*S*,12a*S*)-*N*-[(4-Fluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1-(tetrahydro-2*H*-thiopyran-4-yl)-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-1,4-dimethyl-6,8-dioxo-1,2,3,4,6,8

,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(4*S*,12a*S*)-*N*-[(4-Fluorophenyl)methyl]-7-hydroxy-4-methyl-1-(1-methylethyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(4*S*,12a*S*)-*N*-[(4-Fluorophenyl)methyl]-7-hydroxy-1,4-bis(2-methylpropyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

enantiomers thereof; diastereomers thereof; mixtures of enantiomers thereof; mixtures of diastereomers thereof; mixtures of enantiomers and diastereomers thereof; and pharmaceutically acceptable salts thereof.

[Claim 32]

A compound selected from the group consisting of:

(4a*S*,13a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-10-hydroxy-9,11-dioxo-2,3,4a,5,9,11,13,13a-octahydro-1*H*-pyrido[1,2-*a*]pyrrolo[1',2':3,4]imidazo[1,2-*d*]pyrazine-8-carboxamide;

(4a*S*,13a*R*)-*N*-[(4-Fluorophenyl)methyl]-10-hydroxy-9,11-dioxo-2,3,4a,5,9,11,13,13a-octahydro-1*H*-pyrido[1,2-*a*]pyrrolo[1',2':3,4]imidazo[1,2-*d*]pyrazine-8-carboxamide;

(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-[(1*S*)-1-methylpropyl]-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-methyl-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(3*S*,11a*R*)-*N*-[(4-Fluorophenyl)methyl]-6-hydroxy-3-methyl-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide;

(4*S*,12a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-7-hydroxy-4-methyl-1-(2-methylpropyl)-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(4*S*,12a*S*)-1-(Cyclopropylmethyl)-*N*-[(2,4-difluorophenyl)methyl]-7-hydroxy-4-methyl-6,8-dioxo-1,2,3,4,6,8,12,12a-octahydropyrido[1',2':4,5]pyrazino[1,2-*a*]pyrimidine-9-carboxamide;

(4a*R*,6a*R*,14a*S*)-*N*-[(2,4-Difluorophenyl)methyl]-12-hydroxy-11,13-dioxo-1,3,4,4a,5,6a,7,11,13,14a-decahydro-2*H*-pyrido[1',2':4,5]pyrazino[1,2-*a*][3,1]benzoxazine-10-carboxamide;

(4a*R*,6a*R*,14a*S*)-*N*-[(4-Fluorophenyl)methyl]-12-hydroxy-11,13-dioxo-1,3,4,4a,5,6a,7,11,13,14a-decahydro-2*H*-pyrido[1',2':4,5]pyrazino[1,2-*a*][3,1]benzoxazine-10-carboxamide;

4S,9aR)-5-Hydroxy-4-methyl-6,10-dioxo-3,4,6,9,9a,10-hexahydro-2H-1-oxa-4a,

8a-diaza-anthracene-7-carboxylic acid 2,4,-difluoro-benylamide;

4R,9aS)-5-Hydroxy-4-methyl-6,10-dioxo-3,4,6,9,9a,10-hexahydro-2H-1-oxa-4a,
8a-diaza-anthracene-7-carboxylic acid 2,4,-difluoro-benylamide;

2R,9aS)-5-Hydroxy-2-methyl-6,10-dioxo-3,4,6,9,9a,10-hexahydro-2H-1-oxa-4a,
8a-diaza-anthracene-7-carboxylic acid 4-fluoro-benylamide;

enantiomers thereof; diastereomers thereof; mixtures of enantiomers thereof; mixtures of diastereomers thereof; mixtures of enantiomers and diastereomers thereof; and pharmaceutically acceptable salts thereof.

[Claim 33]
A compound according to claims 31 or 32 wherein the pharmaceutically acceptable salt is a sodium salt.

[Claim 34]

(4a*S*,13a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-10-hydroxy-9,11-dioxo-2,3,4a,5,9,11,13,13a-octahydro-1*H*-pyrido[1,2-*a*]pyrrolo[1',2':3,4]imidazo[1,2-*d*]pyrazine-8-carboxamide; or the pharmaceutically acceptable salt.

[Claim 35]

(3*S*,11a*R*)-*N*-[(2,4-Difluorophenyl)methyl]-6-hydroxy-3-methyl-5,7-dioxo-2,3,5,7,11,11a-hexahydro[1,3]oxazolo[3,2-*a*]pyrido[1,2-*d*]pyrazine-8-carboxamide; or the pharmaceutically acceptable salt.

[Claim 36]

4R,9aS)-5-Hydroxy-4-methyl-6,10-dioxo-3,4,6,9,9a,10-hexahydro-2H-1-oxa-4a, 8a-diaza-anthracene-7-carboxylic acid 2,4,-difluoro-benylamide; or the pharmaceutically acceptable salt

[Claim 37]


A pharmaceutical composition comprising a compound according to any one of Claims 1 to 36, or a pharmaceutically acceptable salt, or solvate thereof.

[Claim 38]

A pharmaceutical composition according to Claim 37, which is an anti-HIV agent.

[Claim 39]


A process for the preparation of a compound of formula (I-20a)




(I-20a)

wherein R^e is one or two halogen; R^z is C_{1-8}alkyl, C_{6-14}arylC_{1-8}alkyl, C_{6-14}aryl, or alkoxy; and P^1 is C_{6-14}arylC_{1-8}alkyl;

comprising condensing a compound of the formula




wherein R^e is one or two halogen; R^{50} is C_{1-8}alkyl; and P^1 is C_{6-14}arylC_{1-8}alkyl;
with a compound of the formula



wherein R^z is C_{1-8}alkyl, C_{6-14}arylC_{1-8}alkyl, C_{6-14}aryl, or alkoxy;
to form a compound of formula (I-20a).


[Claim 40]
A process for the preparation of a compound of formula (I-20b)




(I-20b)

wherein R^e is one or two halogen; R^z is C_{1-8}alkyl, C_{6-14}arylC_{1-8}alkyl, C_{6-14}aryl, or alkoxy; and P^1 is C_{6-14}arylC_{1-8}alkyl;

comprising condensing a compound of the formula




wherein R^e is one or two halogen; R^{50} is C_{1-8}alkyl; and P^1 is C_{6-14}arylC_{1-8}alkyl;
with a compound of the formula



wherein R^z is C_{1-8}alkyl, C_{6-14}arylC_{1-8}alkyl, C_{6-14}aryl, or alkoxy;
to form a compound of formula (I-20b).


[Claim 41]
A process for the preparation of a compound of formula (I-21a)




(I-21a)

wherein R^e is one or two halogen; and P^1 is C_{6-14}arylC_{1-8}alkyl;

comprising condensing a compound of the formula




wherein R^e is one or two halogen; R^{50} is C_{1-8}alkyl; and P^1 is C_{6-14}arylC_{1-8}alkyl;
with a compound of the formula



to form a compound of formula (I-21a).

[Claim 42]


A process for the preparation of a compound of formula (I-21b)




(I-21b)

wherein R^e is one or two halogen; and P^1 is $C_{6-14}arylC_{1-8}alkyl$;

comprising condensing a compound of the formula




wherein R^e is one or two halogen; R^{50} is $C_{1-8}alkyl$; and P^1 is $C_{6-14}arylC_{1-8}alkyl$; with a compound of the formula



to form a compound of formula (I-21b).

[Claim 43]


A process for the preparation of a compound of formula (I-22a)




(I-22a)

wherein R^e is one or two halogen; and P^1 is $C_{6-14}arylC_{1-8}alkyl$;

comprising condensing a compound of the formula:




wherein R^e is one or two halogen; R^{50} is C_{1-8}alkyl; and P^1 is C_{6-14}arylC_{1-8}alkyl; with a compound of the formula



to form a compound of formula (I-22a).


[Claim 44]

A process for the preparation of a compound of formula (I-22b)




(I-22b)

wherein R^e is one or two halogen; and P^1 is C_{6-14}arylC_{1-8}alkyl; comprising condensing a compound of the formula




wherein R^e is one or two halogen; R^{50} is C_{1-8}alkyl; and P^1 is C_{6-14}arylC_{1-8}alkyl; with a compound of the formula



to form a compound of formula (I-22b).

[Claim 45]


A process for the preparation of a compound of formula (I-23a)



(I-23a)


wherein R^e is one or two halogen; and P^1 is $C_{6-14}arylC_{1-8}alkyl$;

comprising condensing a compound of the formula



wherein R^e is one or two halogen; R^{50} is $C_{1-8}alkyl$; and P^1 is $C_{6-14}arylC_{1-8}alkyl$;


with a compound of the formula



to form a compound of formula (I-23a).

[Claim 46]


A process for the preparation of a compound of formula (I-23b)




(I-23b)

wherein R^e is one or two halogen; and P^1 is $C_{6-14}arylC_{1-8}alkyl$;

comprising condensing a compound of the formula




wherein R^e is one or two halogen; R^{50} is C_{1-8}alkyl; and P^1 is C_{6-14}arylC_{1-8}alkyl; with a compound of the formula



to form a compound of formula (I-23b).

[Claim 47]

A process for the preparation of a compound of formula (I-24a)




(I-24a)


wherein R^e is one or two halogen; R^{z1} is hydrogen, C_{3-6}cycloalkyl, heterocycle, or C_{1-8}alkyl optionally substituted with hydroxy, C_{3-6}cycloalkyl, alkoxy, heterocycle, heteroaryl, C_{6-14}aryl, or amino, wherein said amino may be optionally substituted with –C(O)C_{1-8}alkyl or C_{1-8}alkyl;

and P^1 is C_{6-14}arylC_{1-8}alkyl;

comprising condensing a compound of the formula




wherein R^e is one or two halogen; R^{50} is C_{1-8}alkyl; and P^1 is C_{6-14}arylC_{1-8}alkyl;
with a compound of the formula



wherein R^z is C_{1-8}alkyl; R^{z1} is hydrogen, C_{3-6}cycloalkyl, , heterocycle, or C_{1-8}alkyl optionally substituted with hydroxy, C_{3-6}cycloalkyl, alkoxy, heterocycle, heteroaryl, C_{6-14}aryl, or amino, wherein said amino may be optionally substituted with –C(O)C_{1-8}alkyl or C_{1-8}alkyl;

to form a compound of the formula (I-24a).


[Claim 48]
A process for the preparation of a compound of formula (I-24b)




(I-24b)

wherein R^e is one or two halogen; R^{z1} is hydrogen, C_{3-6}cycloalkyl, heterocycle, or C_{1-8}alkyl optionally substituted with hydroxy, C_{3-6}cycloalkyl, alkoxy, heterocycle, heteroaryl, C_{6-14}aryl, or amino, wherein said amino may be optionally substituted with –C(O)C_{1-8}alkyl or C_{1-8}alkyl;
and P^1 is C_{6-14}arylC_{1-8}alkyl;

comprising condensing a compound of the formula




wherein R^e is one or two halogen; R^{50} is C_{1-8}alkyl; and P^1 is C_{6-14}arylC_{1-8}alkyl;
with a compound of the formula



wherein R^z is C_{1-8}alkyl; R^{z1} is hydrogen, C_{3-6}cycloalkyl, heterocycle, or C_{1-8}alkyl optionally substituted with hydroxy, C_{3-6}cycloalkyl, alkoxy, heterocycle, heteroaryl, C_{6-14}aryl, or amino, wherein said amino may be optionally substituted with –C(O)C_{1-8}alkyl or C_{1-8}alkyl;
to form a compound of the formula (I-24b).


[Claim 49]
A process for the preparation of a racemic compound of formula (I-25)




(I-25)

wherein R^e is one or two halogen; R^{z1} is hydrogen, C_{3-6}cycloalkyl, heterocycle, or C_{1-8}alkyl optionally substituted with hydroxy, C_{3-6}cycloalkyl, alkoxy, heterocycle, heteroaryl, C_{6-14}aryl, or amino, wherein said amino may be optionally substituted with –C(O)C_{1-8}alkyl or C_{1-8}alkyl; and P^1 is C_{6-14}arylC_{1-8}alkyl;

comprising condensing a compound of the formula



wherein R^e is one or two halogen; R^{50} is C_{1-8}alkyl; and P^1 is C_{6-14}arylC_{1-8}alkyl;
with a racemic compound of the formula




wherein R^{z1} is hydrogen, C_{3-6}cycloalkyl, heterocycle, or C_{1-8}alkyl optionally substituted with hydroxy, C_{3-6}cycloalkyl, alkoxy, heterocycle, heteroaryl, C_{6-14}aryl, or amino, wherein said amino may be optionally substituted with $-C(O)C_{1-8}$alkyl or C_{1-8}alkyl;

to form a racemic compound of the formula (I-25).

[Claim 50]


A process for the preparation of a racemic compound of formula (I-26)




(I-26)

wherein R^e is one or two halogen; R^{z1} is hydrogen, C_{3-6}cycloalkyl, heterocycle, or C_{1-8}alkyl optionally substituted with hydroxy, C_{3-6}cycloalkyl, alkoxy, heterocycle, heteroaryl, C_{6-14}aryl, or amino, wherein said amino may be optionally substituted with $-C(O)C_{1-8}$alkyl or C_{1-8}alkyl; and P^1 is C_{6-14}arylC_{1-8}alkyl;

comprising condensing a compound of the formula




wherein R^e is one or two halogen; R^{50} is C_{1-8}alkyl; and P^1 is C_{6-14}arylC_{1-8}alkyl;
with a racemic compound of the formula




wherein R^{z1} is hydrogen, C_{3-6}cycloalkyl, heterocycle, or C_{1-8}alkyl optionally substituted with hydroxy, C_{3-6}cycloalkyl, alkoxy, heterocycle, heteroaryl, C_{6-14}aryl, or amino, wherein said amino may be optionally substituted with $-C(O)C_{1-8}$alkyl or C_{1-8}alkyl;
to form a racemic compound of formula (I-26).

[Claim 51]
A process for the preparation of a racemic compound of formula (I-27)




(I-27)

wherein R^e is halogen; and P^1 is C_{6-14}arylC_{1-8}alkyl;
comprising condensing a compound of the formula



wherein R^e is one or two halogen; R^{50} is C_{1-8}alkyl; and P^1 is C_{6-14}arylC_{1-8}alkyl;
with a racemic compound of the formula



to form a racemic compound of formula (I-27).

[Claim 52]

A compound as claimed in any of claims 1 to 36 for use in medical therapy.

[Claim 53]

Use of a compound as claimed in any of claims 1 to 36 in the manufacture of a medicament for the treatment or prophylaxis of an HIV infection.

[Claim 54]

A compound of formula (I-20a) described in Claim 39. formula (I-20b) described in Claim 40, formula (I-21a) described in Claim 41, formula (I-21b) described in Claim 42, formula (I-22a) described in Claim 43, formula (I-22b) described in Claim 44, formula (I-23a) described in Claim 45, formula (I-23b) described in Claim 46, formula (I-24a) described in Claim 47, formula (I-24b) described in Claim 48, formula (I-25) described in Claim 49, formula (I-26) described in Claim 50, or formula (I-27) described in Claim 51, or a pharmaceutically acceptable salt thereof.

[Claim 55]

A compound of formula (I-20a) described in Claim 39. formula (I-20b) described in Claim 40, formula (I-21a) described in Claim 41, formula (I-21b) described in Claim 42, formula (I-22a) described in Claim 43, formula (I-22b) described in Claim 44, formula (I-23a) described in Claim 45, formula (I-23b) described in Claim 46, formula (I-24a) described in Claim 47, formula (I-24b) described in Claim 48, formula (I-25) described in Claim 49, formula (I-26) described in Claim 50, or formula (I-27) described in Claim 51, or a pharmaceutically acceptable salt thereof, wherein each P^1 is hydrogen.

[Claim 56]

A pharmaceutical composition according to claim 37 wherein said composition comprises at least one additional therapeutic agent selected from reverse transcriptase inhibitors and protease inhibitors.