

This Sheet Has Been Replaced



Title: SUBSTITUTED 5-(2-((2-ARYL-2-HYDROXYETHYL)AMINO) PROPYL)-1,3-BENZODIOXOLES

The present invention relates to novel 1,3-benzodioxole compounds which have antidiabetic and/or anti-hyperglycemic properties in mammals. More particularly it relates to novel substituted 5-(2-((2-aryl-2-hydroxyethyl)amino)propyl)-1,3-benzodioxoles. The present invention also relates to pharmaceutical compositions comprising these compounds, methods for the preparation of these compounds, as well as methods for the use of these compounds in treating diabetes and/or hyperglycemia and/or obesity in mammals.

THIS SHEET HAS BEEN
REPLACED




SUBSTITUTED 5-(2-((2-ARYL-2-HYDROXYETHYL)
AMINO)PROPYL)-1,3-BENZODIOXOLES

The present invention relates to novel 1,3-benzodioxole compounds which have antidiabetic and/or anti-hyperglycemic properties in mammals. More particularly it relates to novel substituted 5-(2-((2-aryl-2-hydroxyethyl)amino)propyl)-1,3-benzodioxoles. The present invention also relates to pharmaceutical compositions comprising these compounds, methods for the preparation of these compounds, as well as methods for the use of these compounds in treating diabetes and/or hyperglycemia and/or obesity in mammals.


BACKGROUND OF THE INVENTION

It is well known to employ medicinal agents in the treatment of persons suffering from diabetes, hyperglycemia and obesity.

Ainsworth et al, United States Patent No. 4,478,849, disclose secondary amine compounds having the general formula (I):




wherein R_1 is a hydrogen, fluorine or chlorine atom or a hydroxyl, hydroxymethyl, methyl, methoxyl, amido, formamido, acetamido, methylsulfonylamido, nitro, benzyloxy, methylsulfonylmethyl, ureido, trifluoromethyl, or methoxybenzylamino group; R_2 is a hydrogen, fluorine or chlorine atom or a hydroxyl group; R_3 is a hydrogen or chlorine atom or a hydroxyl group; R_4 is a carboxylic acid group or a salt, ester or amide thereof; R_5 is a hydrogen, chlorine or fluorine atom or a methyl, methoxyl, or hydroxyl group or a carboxylic acid group




wherein R_1 and R_2 are hydrogen or methyl; n is 1, 2 or 3; and Z is alkyl of 1 to 6 carbon atoms, alkoxy of 1 to 6 carbon atoms, halogen or hydrogen; useful in treating obesity and/or hyperglycemia and/or inflammation in mammals.

Ferris, United States Patent No. 4,341,793 discloses secondary amine compounds of the formula (IV):




wherein A_1 and A_2 are hydrogen or methyl; n is 1, 2 or 3; and R is hydrogen, chlorine, bromine, hydroxy, nitro, amino or trifluoromethyl; which are useful as anti-hyperglycemia agents or anti-obesity agents.

Smith et al., United States Patent No. 4,309,443, disclose cinnamic acid derivatives of the formula (V):




wherein R_1 and R_2 and R_3 are as defined in relation to formula (II); R_4 is a hydrogen, chlorine or fluorine



or a salt, ester or amide thereof; R_6 is a hydrogen atom or a methyl or propyl group; X is an oxygen atom or a bond; and Y is an alkylene group of up to 6 carbon atoms or a bond, which have anti-obesity and/or anti-hyper-glycemic activity.

Ainsworth et al, United States Patent No. 4,396,627, disclose secondary amine compounds of the formula (II):




wherein R_1 , R_2 and R_3 are as defined in relation to formula (I) or each independently represents a bromine atom; R_4 is an alkyl group of 1 to 10 carbon atoms substituted by a hydroxyl, lower alkoxyl, oxo, lower acyloxy or OCH_2CO_2H group or lower alkyl ester thereof; R_5 is a hydrogen, chlorine or fluorine atom or a methyl, methoxyl or hydroxyl group or a carboxylic acid group or a salt, ester or amide thereof; R_6 is a hydrogen atom or a methyl, ethyl or propyl group; R_7 is a hydrogen atom or a methyl, ethyl, or propyl group; X is an oxygen atom or a bond; and Y is an alkylene group of up to six carbon atoms or a bond; which are useful in reducing high blood glucose and lipid levels in humans and animals.

Ainsworth et al, United States Patent No. 4,385,066 disclose arylethanol amine derivatives of the formula (III):




atom or a methyl, methoxyl or hydroxyl group or a carboxylic acid group or a salt, ester or amide thereof; R_5 and R_6 are a hydrogen atom or methyl group; R_7 and R_8 are a hydrogen atom or methyl or ethyl group; X is an oxygen atom or a bond; and Y is an alkylene group of up to 5 carbon atoms; which have been found to possess anti-obesity and/or anti-hyperglycemia activity.

Duckworth, United States Patent No. 4,382,958, disclose secondary amine compositions of the formula (VI):



wherein each of R_1 and R_2 are hydrogen, methyl or ethyl; R_3 is hydrogen, fluorine, chlorine, bromine or trifluoromethyl; each of R_4 and R_5 is hydrogen, fluorine, chlorine, bromine, alkyl having 1 to 6 carbon atoms, or alkoxy having 1 to 6 carbon atoms; n is 1 or 2; and X is straight or branched alkylene having 1 to 12 carbon atoms; which are anti-obesity, hypoglycemia, anti-inflammatory and platelet aggregation inhibiting agents.

Ainsworth et al., United States Patent No. 4,654,371, disclose secondary amine compounds of the formula (VII):




wherein R_{15} is a hydrogen atom or a methyl group; R_{16} is a hydrogen atom or a methyl group and m is 1; useful in

treating obesity and hyperglycemia in humans or animals.

Ikezaki et al., United States Patent No. 4,490,392 disclose benzylalcohol derivatives of the formula (VIII):

5


(VIII)



wherein R is hydroxy, benzyloxy, halogen or alkoxy having 1 to 4 carbon atoms and Ring A is monomethoxyphenyl, dimethoxyphenyl, trimethoxyphenyl or 3,4-methylenedioxyphenyl; which are useful as anti-diabetic agents.

10

Ferris, United States Patent No. 4,432,993 discloses 2-(2-benzofuranyl)ethanolamine derivatives of the formula (IX):



wherein R_1 is hydrogen or methyl; R_2 is hydrogen or methyl; R_3 is hydroxy, hydroxy(C_1 - C_6)alkoxy, benzyloxy or a group X-Y-Z wherein (i) X is a bond or oxygen, Y is a C_1 - C_6 straight or branched alkylene, and Z is hydrogen or carboxy; or (ii) X is a bond or $-O-CH_2-$, Y is a C_2 - C_6 straight or branched alkenylene and Z is carboxy; R_4 is hydrogen, hydroxy, halogen, C_1 - C_6 alkyl or C_1 - C_6 alkoxy; and n is 1, 2 or 3; which have anti-obesity, hypoglycemia, anti-inflammatory, and platelet-aggregation inhibiting activity.


25

Ainsworth et al., United States Patent No.



-6-


4,338,333 disclose ethanamine derivatives of the formula (X):



wherein R_1 , R_2 and R_3 are as defined in relation to
 5 formula (I) or each independently represent a bromine
 atom; R_4 is a hydrogen atom or methyl group; R_5 is a
 hydrogen atom or methyl group; R_6 is a hydrogen,
 fluorine or chlorine atom or a methyl, methoxyl or
 hydroxy group; X is an oxygen atom or a bond; Y is an
 10 alkylene group of up to 6 carbon atoms or a bond; and Z
 is an alkylene, alkenylene or an alkynylene group of up
 to 10 carbon atoms; which possess anti-obesity and
 anti-hyperglycemia properties.

Ikezaki et al., United States Patent No.


15 4,032,575 disclose benzylalcohol amine derivatives of
 the formula (XI):



wherein Ring A is monohydroxyphenyl; which induce a
 decrease in blood sugar levels when Ring A is
 20 2-hydroxyphenyl.


Holloway et al., United States Patent No.

4,772,631, disclose phenoxyacetic acid ethers of the
 formula (XII):



wherein R_1 is hydrogen or fluorine; R_2 and R_3 are hydrogen or alkyl having 1 to 3 carbon atoms; Z is CH_2OH or a group $-\text{COR}_4$ in which R_4 is OH , NH_2 or alkoxy having 1 to 6 carbon atoms; which are useful in treating obesity and related conditions.

Philion, United States Patent No. 4,751,246 disclose benzenemethanol and ethylamine compounds of the formula (XIII):



wherein R_1 , R_2 and R_3 are independently hydrogen or lower alkyl; n is an integer from 1 to 3; Ar is phenyl, methylenedioxyphenyl or phenyl having from 1 to 3 substituents selected from the group consisting of halo, lower alkyl, hydroxy and lower alkoxy; R is lower alkyl; and Y is hydrogen, lower alkyl, lower alkoxy, lower alkyl; lower alkanoyl, aroyl, benzenesulfonyl or toluene-sulfonyl; which are useful as anti-hypertensive agents.

It has now been discovered that a group of novel substituted 5-(2-((2-aryl-2-hydroxyethyl)amino)propyl)-1,3-benzodioxoles possess vastly increased anti-hyperglycemia and anti-obesity properties with greater β_3 selectivity in comparison with the prior art compounds. The compounds are therefore useful in treating diabetes, hyperglycemia and obesity, exhibiting minimal side effects, i.e. heart rate increase and muscle tremor in humans and animals, when formulated

into pharmaceutical compositions.


BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 shows a comparison of the selectivity between a compound of the present invention and a prior art compound.


FIG. 2 shows a comparison of the selectivity between a compound of the present invention and a prior art compound.

SUMMARY OF THE INVENTION

According to the present invention there are provided new compounds of the formula (XIV):



wherein R_1 and R_4 may be one or more groups which may be the same or different and are selected from the group consisting of hydrogen, C_1 to C_4 alkyl, C_1 to C_4 alkoxy, hydroxy, halogen, trifluoromethyl, carboxy, hydroxyalkyl, alkoxy carbonyl, C_1 to C_4 thioalkyl, sulfonyl and sulfinyl; X is a divalent radical consisting of



wherein R' is selected from the group consisting of hydrogen, C_1 to C_4 alkyl and C_1 to C_4 acyl and Y is selected from the group consisting of carbonyl and thiocarbonyl; R_2 and R_3 may be the same or different and



-9-

are selected from the group consisting of hydrogen and C_1 to C_4 alkyl; R_5 and R_6 are selected from the group consisting of hydrogen, carboxy, alkoxycarbonyl, hydroxymethyl, $-CH_2OCH_2COOR_7$ and $-CH_2OCH_2CH_2OR_7$, where R_7 is hydrogen or C_1 to C_4 alkyl; with the provision that R_5 and R_6 may not both be hydrogen; and the pharmaceutically acceptable salts and esters thereof, the enantiomers thereof, the racemic mixtures thereof and the diastereomeric mixtures thereof.

The compounds of the above formula have centers of asymmetry at the carbon atoms marked with an asterisk. The compounds may, therefore, exist in at least two and often four stereoisomeric forms. The present invention encompasses all stereoisomers of the compounds whether free from other stereoisomers or admixed with other stereoisomers in any proportion and thus includes, for instance, racemic mixture of enantiomers as well as the diastereomeric mixture of isomers.

A preferred embodiment of the invention is compounds wherein R_1 is one or two groups as defined above. A more preferred embodiment is compounds wherein R_1 is two of the defined groups.

Preferably both asymmetric carbon atoms have the R absolute stereochemical configuration.

The absolute configuration of any compound may be determined by conventional X-ray crystallography.

The preferred compounds are $(\underline{R}^*, \underline{R}^*)-(+/-)-5-(2-((2-(3\text{-chlorophenyl})-2\text{-hydroxyethyl})\text{amino})\text{propyl})-1,3\text{-benzodioxole-2,2-dicarboxylic acid, dimethyl ester; } (\underline{R}^*, \underline{R}^*)-(+/-)-5-(2-((2-(3\text{-chlorophenyl})-2\text{-hydroxyethyl})\text{amino})\text{propyl})-1,3\text{-benzodioxole-2,2-dicarboxylic acid, diethyl ester; } (\underline{R}^*, \underline{R}^*)-(+/-)-5-(2-((2-(3\text{-chlorophenyl})-2\text{-hydroxyethyl})\text{amino})\text{propyl})-1,3\text{-benzodioxole-2,2-dicarboxylic acid, diisopropyl ester; } (\underline{R}^*, \underline{R}^*)-(+/-)-5-(2-((2-$



(3-chlorophenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, disodium salt; ($\underline{R}^*$, $\underline{R}^*$)-(+/-)((5-(2-(5-(3-chlorophenyl)-2-oxo-3-oxazolidinyl)propyl)-1,3-benzodioxole-2,2-diyl)-bis-(methy-
5 leneoxy)bis acetic acid, dimethyl ester; ($\underline{R}^*$, $\underline{R}^*$)-(+/-)-3-(2-(2,2-bis(2-hydroxyethoxy)methyl)-1,3-benzodioxole-5-yl)-1-methylethyl)5-(3-chlorophenyl)-2-oxazolidinone; ($\underline{R}^*$, $\underline{R}^*$)-(+/-)-((5-(2-((2-(3-chlorophenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-diyl)bis(methyleneoxy))
10 bis acetic acid, dimethyl ester; ($\underline{R}^*$, $\underline{R}^*$)-(+/-)alpha-(((2-(2,2-bis(2-hydroxyethoxyl)methyl)-1,3-benzodioxole-5-yl)-1-methylethyl)amino)methyl)-3-chlorobenzenemethanol; and the optically active derivatives thereof.

Also according to the present invention there
15 is provided a method of treating diabetes and/or hyperglycemia and/or obesity in humans or other mammals which comprises administering to a human or other mammal an antiobesity effective amount or an anti-hyperglycemia effective amount of a compound of the present invention.

Further, according to the present invention
20 there are provided pharmaceutical compositions of matter comprising an effective amount of the compounds of the present invention in combination with a pharmaceutically acceptable carrier; as well as a method for increasing
25 the content of lean meat in edible mammals, which comprises administering to edible mammals an effective amount of the compound.

Also according to the present invention there
are provided processes for producing the compounds of
30 the present invention and a process for the resolution of the optical isomers of the present invention and salts and esters thereof.

DETAILED DESCRIPTION OF THE INVENTION

The disease diabetes mellitus is characterized by metabolic defects in production and utilization of glucose which result in the failure to maintain appropriate blood sugar levels. The result of these defects is elevated blood glucose or hyperglycemia. Research on the treatment of diabetes has centered on attempts to normalize fasting and postprandial blood glucose levels. Treatments have included parenteral administration of exogenous insulin, oral administration of drugs and dietary therapies.

Two major forms of diabetes mellitus are now recognized. Type I diabetes, or insulin-dependent diabetes, is the result of an absolute deficiency of insulin, the hormone which regulates glucose utilization. Type II diabetes, or insulin-independent diabetes, often occurs in the face of normal, or even elevated levels of insulin and appears to be the result of the inability of tissues to respond appropriately to insulin. Most of the Type II diabetics are also obese.

The compounds of this invention were tested for hypoglycemic and anti-obesity activity according to the following procedure.

Obese mice (C57 B1/6J (ob/ob)), and/or diabetic mice (C57 B1/KsJ (db/db)) were obtained from Jackson Laboratories, Bar Harbor, Maine. Obese rats (fa/fa) were obtained from Charles River Laboratories, Wilmington, Massachusetts. Obese mice were 8 weeks of age and diabetic mice were 9 weeks of age at the start of the test. Obese rats were 12 - 14 weeks of age at the start.

The test compounds were dissolved in methanol, mixed with powdered Purina rodent chow on a weight of compound to weight of chow basis and thoroughly dried.



- 12 -

Groups of 6 control mice or rats received vehicle (methanol) treated chow.

Groups of 6 test mice were fed ad libitum for up to seven weeks and food consumption was measured daily by weighing the food bins before and after the addition of fresh chow. Thus, a 40 g mouse, fed the test compound at a concentration of 0.02 percent of the diet, would receive a dose of 20 mg/kg/day if it ate 4 g of chow per day.

Groups of 6 test rats were fed 25 g of chow per day for one month. Except for the first day or two, they consumed all their food each day. When these animals are fed ad libitum, they consume 26 g of chow per day and the compounds have no effect on food consumption.

The mice or rats were weighed before the first treatment and once at the end of each indicated treatment period.

Blood samples were collected before the first treatment and once at the end of each indicated treatment period by retro-orbital puncture using heparinized capillary tubes.

Plasma was separated by centrifugation in a Beckman microfuge for 5 minutes. Plasma glucose concentrations were determined with the Beckman glucose analyzer which uses a glucose oxidase method.

The results of these tests on representative compounds of this present invention and a comparative compound of the prior art appear in Table I - III.

TABLE 1
PLASMA GLUCOSE

Compound	Type of Mice	Dose (gwt)	Plasma Glucose Levels in mg/100 ml			
			0	Weeks		4
				1	5	
(R*,R*)-(+/-)-5(2-((2-(3-chlorophenyl))-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester hydrobromide	ob/ob	0(control)	233	283		275
		0.0005	270	164		179
		0.002	276	150		156
		0.005	237	143		157
	db/db		0	1	5	7
		0(control)	445	496	543	566
		0.0005	452	162	164	170
		0.002	453	151	150	161
		0.005	447	148	138	180
(R*,R*)-(+/-)-5(2-((2-(3-chlorophenyl))-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, disodium salt	ob/ob	0(control)	232	271		245
		0.0005	218	191		240
		0.002	220	162		197
		0.005	220	145		165
	db/db		0	1	5	7
		0(control)	445	496	543	566
		0.002	453	236	190	250
		0.005	452	155	145	165
(R*,S*)-(+/-)-5(2-((2-(3-chlorophenyl))-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester	ob/ob	0(control)	227	284		211
		0.005	218	162		174
		0.02	219	135		153
			0	5	10	

THIS SHEET HAS BEEN
REPLACED



TABLE II
CHANGE IN WEIGHT


Compound	Type of Mice or rat	Dose (% wt)	Change in weight in grams				Average Daily Food Consumption g/animal
			weeks				
			1	2	3	4	
(R*,R*)-(+/-)-5(2-((2-(3-(chloro-phenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester hydrobromide	ob/ob	0(control)	1.4	3.7	5.4	7.2	4.5
		0.0005	0.5	2.2	3.2	5.2	5.6
		0.002	-0.3	1.1	1.6	3.2	5.6
		0.005	-1.1	-0.7	-0.2	1.6	5.8
	fa/Ea	0(control)	18	39	57	72	25
		0.0006	14	20	27	26	25
		0.002	1	2	3	4	25
		0.006	10	4	-15	-8	25
(R*,R*)-(+/-)-5(2-((2-(3-(chloro-phenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, disodium salt	ob/ob	0(control)	3.1	5.6	7.0	9.3	4.5
		0.002	1.6	4.1	4.7	7.3	5.2
		0.005	0.2	1.6	2.4	4.7	5.9

TABLE III
PLASMA GLUCOSE COMPARATIVE

Compound	Type of Mice or rat	Dose (% wt)	Plasma Glucose Level in mg/100 ml		
			Weeks		
			0	1	2
R*,R*)-(+/-)-(4-(2-((2-(4-chlorophenyl)-2-hydroxyethyl)amino)propyl)phenoxy)-acetic acid, methyl ester**	ob/ob	0(control)	233	183	252
		0.0005	232	140	126
		0.0001	231	143	119
		0.005	231	137	138
	db/db	0(control)	444	349	604
		0.005	440	138	132
(R*,R*)-(+/-)-alpha-(((2-(1,3-benzodioxol-5-yl)-1-methylethyl)amino)methyl)-3-chlorobenzene-methanol mixture with (R*,S*)-(+/-)-alpha-(((2-(1,3-benzodioxol-5-yl)-1-methylethyl)amino)methyl)-3-chlorobenzenemethanol	ob/ob	0(control)	217	217	
		0.002	226	234	
		0.005	227	175	
		0.02	227	149	
(R*,S*)-(+/-)-alpha-(((2-(1,3-benzodioxol-5-yl)-1-methylethyl)amino)methyl)-3-chlorobenzene-methanol	ob/ob	0(control)	217	217	
		0.002	228	204	
		0.005	226	176	
		0.02	229	128	

** prior art compound





The above Table I demonstrates that compounds of the present invention and the pharmaceutically active salts thereof effectively lower blood glucose levels when administered orally to genetic strains of hyperglycemic mice which are animal models of type II diabetes. The compounds of the present invention are also seen, in Table II, to decrease weight gain when administered to genetic strains of mice and rats which are animal models of obesity. The exact mechanism by which they act is not known and the invention should not be construed as limited to any particular mechanism of action.

As effective hypoglycemic and weight loss agents, these compounds are useful for the treatment of hyperglycemia and obesity in Type II diabetes.

SELECTIVITY

β -Adrenergic receptors can be divided into β_1 , β_2 and β_3 -subtypes. Activation of β_1 -receptors invokes increases in heart rate while activation of β_2 -receptors stimulates glycogen breakdown in muscle and thereby prevents glycogen synthesis. Activation of β_3 -receptors stimulates lipolysis (the breakdown of adipose tissue triglycerides to glycerol and free fatty acids), and thereby promotes the loss of fat mass. Compounds that stimulate β_3 -receptors will have anti-obesity activity. In addition, they have hypoglycemic or anti-diabetic activity, but the mechanism of this effect is unknown. A compound that selectively stimulates β_3 -receptors, i.e., has little or no β_1 or β_2 -activity, will have the desired anti-diabetic and/or anti-obesity activity, but without the undesirable effects of increased heart rate (β_1 -effect) or muscle tremor (β_2 -effect).

Selectivity of a compound was determined using the following procedures. The effect on heart rate (β_1 -effect) was determined using isolated right atria. Guinea pig hearts were placed in a petri dish containing



-17-

aerated Krebs Henseleit bicarbonate buffer (KHB) of the following composition (mM): NaCl, 112.9; KCl, 4.7; KH_2PO_4 , 1.2; $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 1.2; NaHCO_3 , 25.0; CaCl_2 , 2.4 and glucose, 11.5. The buffer was continuously aerated with 95% O_2 - 5% CO_2 . The right atria were dissected and clamped at one end to a tissue holder and electrode assembly (MRA Corp.) and tied at the other end to a force-displacement transducer (Gould-Statham, Burco). The tissues were maintained at 32°C in a 50 ml capacity glass chamber (MRA Corp.) preloaded with 0.5g tension. Heart rate was monitored by a Grass polygraph. After equilibration for two hours, the atria were exposed to $1 \times 10^{-6}\text{M}$ isoproterenol for 5 minutes. The increase in heart rate was calculated and considered to be the maximum response for that tissue. The tissues were then washed and allowed to equilibrate for 90 minutes. Cumulative concentration-response curves were then determined for each compound or vehicle. All responses were measured 5 minutes post-exposure. The responses of each compound were expressed as a percent of the response to isoproterenol. The molar EC_{50} value is the concentration of compound that gave 50% of its own maximum increase in heart rate.

The β_2 -effect of the compounds was determined by their ability to inhibit the insulin-mediated incorporation of ^{14}C -glucose into glycogen in isolated muscle. Soleus muscle from mice were dissected, tied at each end, and placed in a clamp to maintain tension. The clamped muscle was added to a vial that contained 2 ml of KHB with 1.5% bovine serum albumin, 0.3 mU/ml insulin, 5mM (U^{-14}C) glucose, and the appropriate additions of compound or vehicle. The vial was gassed with 95% O_2 - 5% CO_2 , capped, and the tissue incubated for one hour in a rotating water bath at 37°C. The muscle was then removed, rinsed in ice-cold saline, *



blotted and weighed. It was then added to 1 ml of 30% KOH containing 5 mg/ml of oyster glycogen, boiled for 10 minutes, and 0.4 ml of 2% Na_2SO_4 and 3.2 ml of 100% ethanol added. After standing overnight at 4°C, the glycogen was pelleted by centrifugation, washed once with 66% ethanol, and repelleted. The pellet was dissolved in 1 ml of water and the amount of radioactivity present determined by liquid scintillation. The decrease in radioactivity incorporated into glycogen by each compound is expressed as a percentage of the decrease obtained in the presence of $1 \times 10^{-7} \text{M}$ isoproterenol. The molar EC_{50} value is the concentration of compound that gave 50 percent of its own maximum percentage decrease in insulin-mediated incorporation of ^{14}C -glucose into glycogen.

The β_3 -effect of the compounds was determined by their ability to stimulate adipocyte lipolysis. Rat epididymal fat pads were excised and placed in 0.9% saline. Four grams of tissue were transferred to a flask with 20 ml of aerated Krebs-Henseleit bicarbonate (KHB) buffer containing 3% fatty acid-free bovine serum albumin to which 75 mg of crude bacterial collagenase (Worthington) had been added. The tissue was incubated for about 45 minutes at 37°C with gentle shaking. The cells were then washed three times with two volumes of KHB buffer, filtered through two layers of gauze, and brought to a final volume of 80 ml with KHB buffer. One ml aliquots of the cell suspension were added to plastic test tubes containing the appropriate additions of vehicle or compound. The cells were gassed for 1 minute with 95% O_2 -5% CO_2 , capped, and incubated at 37°C with continuous shaking for a total of 30 minutes. The reaction was stopped by adding 0.1 ml of 30% perchloric acid and 0.1 ml of chloroform. After centrifugation, 0.5 ml of supernatant was transferred to another test



tube and neutralized with 0.04 ml of 3 M K_2CO_3 -0.5 M triethanolamine. The amount of glycerol generated from the hydrolysis of endogenous triglycerides was determined in a coupled-enzyme spectrophotometric assay.

5 One-tenth milliliter of the neutralized extract was added to a test tube that contained 0.91 ml of an assay mixture comprised of the following: 0.84 M glycine, 0.42 M hydrazine sulfate, 4.2 mM EDTA, 0.9 mM β -NAD, 9.9 mM $MgCl_2$, 1 mM ATP, 17 U of glycerophosphate

10 dehydrogenase, and 4.3 U of glycerokinase. The test tubes were incubated for 40 minutes at 37°C with constant shaking. The amount of NADH generated, which is proportional to the amount of glycerol, was determined from the increase in absorbance at 340 nm.

15 This value was corrected for the amount of NADH generated in the absence of glycerol by incubating another aliquot of the neutralized extract with the same assay mixture but without glycerokinase. The molar EC_{50} value is the molar concentration of compound that gave

20 50% of that compound's own maximum rate of lipolysis.

Figure 1 shows a comparison of the selectivity of a compound of the present invention CL_1 (R^*, R^*)-(+/)-5-(2-((2-(3-chlorophenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester hydrobromide with that of the corresponding prior art compound, BRL_1 (R^*, R^*)-(+/)-4-(2-((2-(4-chlorophenyl)-2-hydroxyethyl)amino)propyl)phenoxy)-acetic acid, methyl ester (Beecham).

The results indicate that prior art compound

30 BRL_1 has slightly greater β_3 potency, but that it also has more of the undesirable β_1 and β_2 activities than does CL_1 . Figure 2 shows a comparison of the selectivity of a compound of the present invention CL_2 (R^*, R^*)-(+/)-5-(2-((2-(3-chlorophenyl)-2-hydroxyethyl)amino)propyl)-1,3-

35 benzodioxole-2,2-dicarboxylic acid, disodium salt with



the corresponding prior art compound, BRL₂ (R*,R*)-(+/-)-alpha-(((2-(1,3-benzodioxol-5-yl)-1-methylethyl)amino)methyl)-3-chlorobenzenemethanol (Beecham).

5 The results indicate that CL₂ has slightly greater β_3 potency than does BRL₂. CL₂ has no measurable β_1 activity whereas BRL₂ has significant β_1 activity. CL₂ showed only minimal β_2 activity at 1×10^{-4} M whereas BRL₂ was a full agonist with measurable activity at 1×10^{-7} M. Thus, both of the prior art compounds have more
10 of the undesirable activities than do the corresponding compounds of the present invention. These results are summarized in Table IV below.

Table IV also summarizes the relative β_3 potencies of the resolved forms (RR and SS) of both the
15 salt (CL₂) and methyl ester (CL₁). It can be seen from this data that the majority of activity resides with the RR isomer in each case. In the case of the disodium salts, the RR enantiomer is 47 fold more potent than the SS enantiomer. In the case of the methyl ester, the RR
20 isomer is 36 fold more potent.



TABLE IV .

β_3 - SELECTIVITY COMPARISON

Compound	Lipolysis (β_3) (EC ₅₀ , M)	R.A. Rate (β_1) (EC ₅₀ , M)	Glycogen Synthesis (β_2) (EC ₅₀ , M)
Salts:			
CL ₂	6 x 10 ⁻⁹	>10 ⁻³	>1 x 10 ⁻⁴
CL ₃	3 x 10 ⁻⁹	-	-
CL ₄	1.4 x 10 ⁻⁷	-	-
BRL ₂ **	9 x 10 ⁻⁹	5.6 x 10 ⁻⁶	8 x 10 ⁻⁷
Esters:			
CL ₁	3 x 10 ⁻⁸	1.4 x 10 ⁻⁸	8 x 10 ⁻⁷
CL ₅	8 x 10 ⁻⁹	-	-
CL ₆	2.9 x 10 ⁻⁷	-	-
BRL ₁ **	1.3 x 10 ⁻⁸	2 x 10 ⁻⁷	1.2 x 10 ⁻⁷
Isoproterenol	1.2 x 10 ⁻⁸	1.5 x 10 ⁻⁹	8 x 10 ⁻⁹

β_3 - SELECTIVITY ((EC₅₀ - β_1 or β_2)/(EC₅₀ - β_3))

	Selectivity for Lipolysis over:		Relative Selectivity (CL/BRL)	
	Atria	Gly. Syn.	Atria	Gly. Syn.
Salts:				
CL ₂	>167,000	>16,700	>268	>198
BRL ₂ **	622	89		
Esters:				
CL ₁	47	27	3.1	3.0
BRL ₁ **	15	9		
Isoproterenol	0.1	0.7		

CL₁ = (R*,R*)-(+/-)-5-(2-((2-(3-chlorophenyl)-2-hydroxyethyl amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester hydrobromide

CL₂ = (R*,R*)-(+/-)-5-(2-((2-(3-chlorophenyl)-2-hydroxyethyl amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, disodium salt

This Sheet Has Been Replaced



- BRL₁ = (R*,R*)-(+/-)-(4-(2-((2-(4-chlorophenyl)-2-hydroxyethyl)amino)propyl)phenoxy)-acetic acid, methyl ester
- BRL₂ = (R*,R*)-(+/-)-(4-(2-((2-(3-chlorophenyl)-2-hydroxyethyl)amino)propyl)phenoxy)-acetic acid, monosodium salt
- ** = comparative example
- CL₃ = (R,R)-(+/-)-5-(2-((2-(3-chlorophenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, disodium salt.
- CL₄ = (S,S)-(+/-)-5-(2-((2-(3-chlorophenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, disodium salt.
- CL₅ = (R,R)-(+/-)-5-(2-((2-(3-chlorophenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester hydrobomide.
- CL₆ = (S,S)-(+/-)-5-(2-((2-(3-chlorophenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester hydrobomide.

The results indicate that the compounds of the present invention, CL₂ and CL₁ are 200 and 3 times more selective for lipolysis than are the prior art compounds, BRL₂ and BRL₁ respectively.


5

In addition to the abilities of the compounds of the present invention described hereinabove, some of the compounds of the present invention are useful in the preparation of other compounds of the present invention.

10


The compounds of the present invention may generally be prepared according to the following process, the process comprising:

(a) reacting a compound of the formula



15

with a ketalizing reagent of the formula




wherein R₁, R₂, R₃ and R₄ are as above-defined in relation to formula (XIV) and R₈ and R₉ are C₁ to C₄

THIS SHEET HAS BEEN
REPLACED




alkyl and Z is dihalomethylene, carbonyl or thiocarbonyl to produce a compound of the formula



5 wherein R_1 , R_2 , R_3 , R_4 and R_8 and R_9 are as above defined; and, either


(b)(i) reacting the product of step (a) with a base followed by an acid and alcohol to produce a compound of the formula



10 wherein R_1 , R_2 , R_3 and R_4 are as defined above and R_5 and R_6 are alkoxy carbonyl; and optionally

(b)(ii) reacting the product of step (b)(i) with a reagent or reagents for converting the groups represented by R_5 and R_6 to carboxy, hydroxymethyl,
 15 $-\text{CH}_2\text{OCH}_2\text{COOR}_7$ and $\text{CH}_2\text{OCH}_2-\text{CH}_2\text{OR}_7$ wherein R_7 is hydrogen or C_1 to C_4 alkyl; or


(c)(i) reacting the product of step (a) with a carbonyl reducing agent to form a compound of the formula



wherein R_1 , R_2 , R_3 and R_4 are as defined above; and

- (c)(ii) reacting the product of (c)(i) with a reagent or reagents for converting the CH_2OH groups to a $-\text{CH}_2\text{OCH}_2\text{COOR}_7$ group or a $-\text{CH}_2\text{OCH}_2\text{CH}_2\text{OR}_7$ group wherein R_7 is as defined above; or

(c)(iii) reacting the product of step (c)(ii) with a base to produce a compound of the formula




- wherein R_{10} and R_{11} are selected from a $-\text{CH}_2\text{OCH}_2\text{COOR}_7$ group, a $-\text{CH}_2\text{OCH}_2\text{CH}_2\text{OR}_7$ group and R_7 is as above defined.


Any conventional ketalizing reagents, bases, acids alcohols, reagents for converting to defined groups and carbonyl reducing agents known to those skilled in the art may be employed and are contemplated by the present invention.

In preferred embodiments of the present invention, the following reaction scheme may be employed.

This Sheet Has Been Replaced




1. carbonyldiimidazole
2. separate isomers




HEET HAS BEEN
REPLACED

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED



In accordance with the above preferred reaction scheme 2-(3-chlorophenyl)-2-hydroxyethylamine 1, and 3,4-dimethoxyphenylacetone 2 are reacted with sodium cyanoborohydride in methanol, giving 3-chloro-
 5 alpha-(((2-(3,4-dimethoxyphenyl)-1-methylethyl)amino)methyl) benzenemethanol 3 which is reacted with carbonyl diimidazole and triethylamine in tetrahydrofuran, followed by separation of isomers, giving cyclized derivative 4 which is reacted with boron tribromide in dichloromethane,
 10 giving ($\underline{R}^*$, $\underline{R}^*$)-(+/)-5-(3-chlorophenyl)-3-(2-(3,4-dihydroxy phenyl)-1-methylethyl)2-oxazolidinone 5. Compound 5 is then reacted with diethyl dibromomalonate and anhydrous potassium carbonate in acetone, giving ($\underline{R}^*$, $\underline{R}^*$)-(+/)-5-(2-(5-(3-chlorophenyl)-2-oxo-3-oxazolidinyl)propyl)-1,
 15 3-benzodioxole-2,2-dicarboxylic acid, diethyl ester 6 which is then reacted first with sodium hydroxide in ethanol and then with hydrogen chloride gas in methanol, giving the product ($\underline{R}^*$, $\underline{R}^*$)-(+/)-5-(2-((2-(3-chloro-phenyl) 2-hydroxyethyl)-amino)propyl)-1,3-benzodioxole-2,
 20 2-dicarboxylic acid, dimethyl ester 7.

Compound 6 may alternatively be reacted with lithium borohydride in anhydrous tetrahydrofuran followed by reaction with sodium hydride and methyl bromoacetate in anhydrous tetrahydrofuran, giving
 25 ($\underline{R}^*$, $\underline{R}^*$)-(+/)-((5-(2((5-(3-chlorophenyl)-2-oxo)-3-oxazolidinyl)propyl)-1,3-benzodioxole-2,2-diyl)bis(methyleneoxy)) bis acetic acid, dimethyl ester 8 which is then reacted with lithium borohydride in anhydrous tetrahydrofuran, giving ($\underline{R}^*$, $\underline{R}^*$)-(+/)-3-(2-((2,2-bis((2-hydroxyethoxy) methyl)-1,3-benzodioxole-5-yl)-1-methylethyl)-5-(3-chloro-phenyl)2-oxazolidinone 9. Compound 9 is then refluxed with sodium hydroxide in ethanol, under argon, giving
 30 ($\underline{R}^*$, $\underline{R}^*$)-(+/)-alpha-(((2-(2,2-bis((2-hydroxyethoxy)-methyl)-1,3-benzodioxole-5-yl)-1-methylethyl)amino)-methyl)-3-chlorobenzenemethanol 10.
 35




Alternatively compound 8 may be refluxed with sodium hydroxide in ethanol, then neutralized and reacted with hydrogen chloride gas in methanol to derive the product 11.


- 5 In another preferred embodiment, optical isomers of the present invention and the derivatives, salts and esters thereof may be resolved by a process comprising: (a) attaching a chiral auxiliary group at the N-9 position of a mixture of (+) and (-)
- 10 enantiomers of the compounds to form a new pair of diastereoisomers; (b) separating the new pair of diastereoisomers and recovering the new (+) or (-) enantiomer; and (c) converting the substantially pure new diastereoisomer into the corresponding desired
- 15 substantially pure (+) or (-) enantiomer of a derivative, salt or ester of the compound.

In a preferred embodiment, the following Scheme II may be employed.



SCHEME II





In accordance with the reaction sequence outlined in Scheme II, 1,3-benzodioxole-2,2-dicarboxylic acid, 5-(2-((2-(3-chlorophenyl)-2-hydroxyethyl)amino)propyl-, dimethyl ester, hydrobromide (R*, R*)-(+/-)-, 5 (12) is treated with the acid chloride derived from (S)-(-)- -methoxy- -trifluoromethylphenylacetic acid (Mosher's acid). The diastereomeric amides 13 and 14 are separated by preparative HPLC, treated with sodium hydroxide, and the disodium salts 15 and 16 (pure 10 enantiomers), respectively are isolated by reverse phase column chromatography. Treatment of each salt with hydrobromic acid in methanol yields each diester, 1,3-benzodioxole-2,2-dicarboxylic acid, 5-(2-((-3-chlorophenyl)-2-hydroxyethyl)amino)propyl-, dimethyl 15 ester, hydrobromide (R,R) (17) and 1,3-benzodioxole-2,2-dicarboxylic acid, 5-(2-((2-(3-chlorophenyl)-2-hydroxyethyl)amino)propyl-, dimethyl ester hydrobromide (S,S) (18) respectively. In a similar manner, the corresponding diethyl and diisopropyl esters can be 20 prepared.


Repeating the above sequence starting with 1,3-benzodioxole-2,2-dicarboxylic acid, 5-((2-(3-chlorophenyl)-2-hydroxyethyl)amino)propyl-, dimethyl ester, hydrobromide (R*,S*)-(+/-) yields both enantiomers 25 1,3-benzodioxole-2,2-dicarboxylic acid, 5-(2-((2-(3-chlorophenyl)-2-hydroxyethyl)amino)propyl-, dimethyl ester, hydrobromide (R,S) and 1,3-benzodioxole-2,2-carboxylic acid, 5-(2-((2-(3-chlorophenyl)-2-hydroxyethyl)amino)propyl-, dimethyl ester, hydrobromide (S,R) 30 respectively. In a similar manner, the corresponding diethyl and diisopropyl esters can be prepared.

Alternatively, the chiral synthesis of the compounds of the present invention may generally be prepared according to the following process, the process 35 comprising;




-32-


(a)(i) reacting an optically active compound of the formula



wherein R_2 , R_3 and R_4 are as above defined in relation to formula (XIV) and R'' is C_1 to C_4 alkyl with a compound of the formula



wherein R_1 is as above defined in relation to formula (XIV) to produce a compound of the formula




wherein R_1 , R_2 , R_3 , R_4 , and R'' are as defined above;

(b)(i) reacting the product of step (a)(i) with a cyclizing reagent followed by separating the diastereoisomers and recovering the new (+) or (-) enantiomer; or


15

(a)(ii) reacting an optically active compound of the formula




wherein R_2 , R_3 , R_4 and R'' are as above defined with an optically active compound of the formula

20






where R_1 is as above defined to produce a compound of the formula




wherein R_1 , R_2 , R_3 , R_4 and R'' are as defined above and
 5 (b)(ii) optionally, reacting said compound with a cyclizing reagent to produce a compound of the formula



wherein R_1 , R_2 , R_3 , R_4 , Y and R'' are as defined above,
 10 and

(c) recovering the substantially pure new enantiomers of steps (b)(i), (a)(ii) or (b)(ii) and if desired converting said substantially pure (+) or (-) enantiomer into the corresponding derivative, salt or
 15 ester of the compounds of the present invention.


Further, the present invention provides a process for producing the initial reactant of the above process, having the general formula




20 wherein R_2 and R_3 may be the same or different and are selected from the group consisting of hydrogen and $C_1 - C_4$ alkyl and R'' are independently selected from C_1 to C_4 alkyl and R^4 is selected from the group consisting of hydrogen, $C_1 - C_4$ alkyl, $C_1 - C_4$ alkoxy, hydroxy,
 25 halogen, trifluoromethyl, carboxy, hydroxyalkyl, alkoxy carbonyl, C_1 to C_4 thioalkyl, sulfonyl and

sulfinyl; the process comprising

(a) reacting a compound of the formula




with a carboxylating reagent to produce a compound of the formula




wherein R_3 and R_4 are as above defined, and

(b) reacting the product of step (a) with an alkylating agent to produce a compound of the formula




wherein R_3 , R_4 , and R' are as above defined, and

(c) reacting the product of step (b) with a selective reducing agent to produce a compound of the formula



wherein R_3 , R_4 , and R' are as above defined, and

(d) reacting the product of step (c) with a sulfonylating agent to produce a compound of the formula



wherein R_2 , R_3 , R_4 , and R' are as above defined, and

(e) reacting the product of step (d) with an



acid followed by hydrogen and palladium to produce said compound.

More specifically, the chiral synthesis may be carried out by performing the synthetic reaction sequence outlined in Scheme III. L-DOPA 19 is treated with di-t-butyl-dicarbonate in dimethylformamide, giving (S)-N-(1-(1,1-dimethylethoxy)carbonyl)-3-hydroxy-L-tyrosine 20 which is reacted with methyl iodide and anhydrous potassium carbonate in acetone, giving (S)-N-(1,1-dimethylethoxy)carbonyl)-3,4-dimethoxy-L-phenyl-alanine methyl ester 21 which is reduced with lithium borohydride giving (S)-1,1-dimethylethyl-(2-(3,4-dimethoxyphenyl)-1-(hydroxymethyl)ethyl) carbonate 22, which is reacted with methanesulfonyl chloride and triethylamine in methylene chloride, giving (S)-1,1-dimethylethyl-(2-(3,4-dimethoxyphenyl)-1-((methylsulfonyl)oxy)methyl)ethyl)carbamate 23. Compound 23 is treated with trifluoroacetic acid and then is reduced using 10% Pd/C and hydrogen giving (R)-3,4-dimethoxy--methyl-benzeneethanamine 24. The compound 24 is treated with 3-chlorostyrene oxide 25 and N-(trimethylsilyl)acetamide giving a mixture of (R,S)- and (S,S)-3-chloro--(((2-(3,4-dimethoxyphenyl)-1-methylethyl)amino)methyl)benzenemethanol 26. Reaction of 26 with carbonyl diimidazole and triethylamine in tetrahydrofuran, followed by chromatographic separation of the diastereomers gives cyclized derivative 27, which is reacted with boron tribromide in dichloromethane, giving (R,R)-5-(3-chlorophenyl)-3-(2-(3,4-dihydroxyphenyl)-1-methylethyl)-2-oxazolidinone 28. Compound 28 is then reacted with diethyl dibromomalonate and anhydrous potassium carbonate in acetone, giving (R,R)-5-(2-(5-(3-chlorophenyl)-2-oxo-3-oxazolidinyl)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, diethyl ester 29. Compound 29 is reacted with 5N sodium hydroxide in ethanol and then




isolated by reverse phase chromatography to give
(R,R)-1,3-benzodioxole-2,2-dicarboxylic acid,
5-(2-((2-(3-chlorophenyl)-2-hydroxyethyl)amino)propyl)-,
disodium salt 15.

- 5 Alternatively, reacting compound 24 with
(R)-3-chlorostyrene oxide, enantiomerically pure 15 is
prepared. Enantiomerically pure epoxides can be
prepared by asymmetric reduction of the corresponding
x-chloroketone as described by Corey, J. Am. Chem. Soc.,
10 109, 5551 (1987), followed by base catalyzed closure to
the epoxide.




SCHEME III






1) Carbonyldiimidazole


2) Separate Isomers



$\text{BBr}_3, \text{CH}_2\text{Cl}_2$




$\text{Br} \begin{array}{c} \text{CO}_2\text{C}_2\text{H}_5 \\ \text{CO}_2\text{C}_2\text{H}_5 \end{array}$



1) 5N. NaOH, EtOH

2) Reverse phase HPLC





The active compounds of the present invention may be orally administered as a pharmaceutical composition, for example, with an inert diluent, or with an assimilable edible carrier, or they may be enclosed
5 in hard or soft shell capsules, or they may be compressed into tablets, or they may be incorporated directly with the food of the diet. For oral therapeutic administration, these active compounds may be incorporated with excipients and used in the form of tablets, pills, capsules, ampules,
10 sachets, elixirs, suspensions, syrups, and the like. Such compositions and preparations should contain at least 0.1 percent of active compound. The percentage of active compound in these compositions may, of course, be varied and may conveniently be between about 2 percent to about
15 60 percent of the weight of the unit. The amount of active compound in such therapeutically useful compositions is such that an effective dosage will be obtained.

The effective dosage of active ingredient employed may vary depending on the particular compound
20 employed, the mode of administration, the condition being treated and the severity of the condition being treated.

When treating diabetes mellitus and/or hyperglycemia generally satisfactory results are
25 obtained when the compounds of the present invention are administered at a daily dosage of from about 0.1 milligram to about 1 milligram per kilogram of animal body weight, preferably given in divided doses two to six times a day, or in sustained release form. For most
30 large mammals, the total daily dosage is from about 3.5 milligrams to about 140 milligrams, preferably from about 3.5 milligrams to about 5 milligrams. In the case of a 70 kg adult human, the total daily dose will generally be from about 7 milligrams to about 70
35 milligrams. This dosage regimen may be adjusted to



provide the optimal therapeutic response.

When treating obesity, in conjunction with diabetes and/or hyperglycemia, or alone, generally satisfactory results are obtained when the compounds of the present invention are administered at a daily dosage of from 1 milligram to about 10 milligrams per kilogram of animal body weight, preferably given in divided doses two to six times a day, or in sustained release form. For most large mammals, the total daily dosage is from about 35 milligrams to about 1,400 milligrams, preferably from about 35 milligrams to about 50 milligrams. In the case of a 70 kg adult human, the total daily dose will generally be from about 70 milligrams to about 700 milligrams. This dosage regimen may be adjusted to provide the optimal therapeutic response.

The tablets, pills, capsules, and the like may also contain a binder such as gum tragacanth, acacia, corn starch or gelatin; excipients such as dicalcium phosphate; a disintegrating agent such as corn starch, potato starch, alginic acid; a lubricant such as magnesium stearate; and a sweetening agent such as sucrose, lactose or saccharin. When a dosage unit form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier such as a fatty oil.

Various other materials may be present as coatings or to modify the physical form of the dosage unit. For instance, tablets may be coated with shellac, sugar or both. A syrup or elixir may contain, in addition to the active ingredient, sucrose as a sweetening agent, methyl and propylparabens as preservatives, a dye and a flavoring such as cherry or orange flavor.

These active compounds may also be administered parenterally. Solutions or suspensions of these active



compounds can be prepared in water suitably mixed with a surfactant such as hydroxypropylcellulose. Dispersions can also be prepared in glycerol, liquid polyethylene glycols and mixtures thereof in oils. Under ordinary
5 conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or
10 dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases, the form must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of
15 manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (e.g. glycerol, propylene glycol and
20 liquid polyethylene glycol), suitable mixtures thereof, and vegetable oils.

The compounds of the present invention also possess utility for increasing lean meat deposition and/or improving lean meat to fat ratio in edible
25 animals, i.e. ungulate animals and poultry.

Animal feed compositions effective for increasing lean meat deposition and for improving lean meat to fat ratio in poultry, swine, sheep, goats, domestic pets and cattle are generally prepared by
30 mixing the compounds of the present invention with a sufficient amount of animal feed to provide from about 1 to 1000 ppm of the compound in the feed.

Animal feed supplements can be prepared by admixing about 75% to 95% by weight of a compound of the
35 present invention with about 5% to about 25% by weight



of a suitable carrier or diluent.. Carriers suitable for use to make up the feed supplement compositions include the following: alfalfa meal, soybean meal, cottonseed oil meal, linseed oil meal, sodium chloride, cornmeal, 5 cane molasses, urea, bone meal, corncob meal and the like. The carrier promotes a uniform distribution of the active ingredients in the finished feed into which the supplement is blended. It thus performs an important function by ensuring proper distribution of 10 the active ingredient throughout the feed.

If the supplement is used as a top dressing for the feed, it likewise helps to ensure uniformity of distribution of the active material across the top of the dressed feed.

15 The preferred medicated swine, cattle, sheep and goat feed generally contain from 0.01 to 400 grams of active ingredient per ton of feed, the optimum amount for these animals usually being about 50 to 300 grams per ton of feed.

20 The preferred poultry and domestic pet feeds usually contain about 0.01 to 400 grams and preferably 10 to 400 grams of active ingredient per ton of feed.

For parenteral administration the compounds of the present invention may be prepared in the form of a 25 paste or a pellet and administered as an implant, usually under the skin of the head or ear of the animal in which increase in lean meat deposition and improvement in lean mean to fat ratio is sought.

In general, parenteral administration involves 30 injection of a sufficient amount of the compounds of the present invention to provide the animal with 0.001 to 100 mg/kg/day of body weight of the active ingredient. The preferred dosage for swine, cattle, sheep and goats is in the range of from 0.001 to 50 mg/kg/day of body 35 weight of active ingredient; whereas, the preferred dose



level for poultry and domestic pets is usually in the range of from 0.001 to 35 mg/kg/day of body weight.

Paste formulations can be prepared by dispersing the active compound in a pharmaceutically acceptable oil such as peanut oil, sesame oil, corn oil or the like.

Pellets containing an effective amount of the compounds of the present invention can be prepared by admixing the compounds of the present invention with a diluent such as carbowax, carnuba wax, and the like, and a lubricant, such as magnesium or calcium stearate, can be added to improve the pelleting process.

It is, of course, recognized that more than one pellet may be administered to an animal to achieve the desired dose level which will provide the increase in lean meat deposition and improvement in lean meat to fat ratio desired. Moreover, it has been found that implants may also be made periodically during the animal treatment period in order to maintain the proper drug level in the animal's body.

The method of the present invention has several advantages; for the pet owner or veterinarian who wishes to increase leanness and trim unwanted fat from pet animals, the present invention provides the means by which this can be accomplished. For the poultry men and swine raisers, using the method of the present invention yields leaner animals which command higher prices from the meat industry.



The results of animal growth regulant evaluation of compounds of the invention appear in Table V.

Antilipogenic Evaluation of Test Compounds - Mouse Study

CFI female mice, 55 days old, are weighed in groups of 5 and allotted to cages to minimize weight variation among cages. Treatments are randomly assigned to cages.

Each of the treatments are tested in 1 to 3 replicates, i.e., in 3 cages of 5 mice each. There are 10 cages of 5 control mice each. Test compounds are mixed in the diet at the dosage level indicated. Feed and water are offered ad libitum for a 12-day test period. The mice are weighed as a group of 5 and the weight gain determined. The mice are sacrificed by cervical dislocation. The right uterine fat pad of each mouse is removed. The fat pads for each cage of 5 mice are weighed as a unit. The fat versus body weight ratio is determined using the absolute weights of the fat pads and the absolute body weight.

It can be seen that a reduction in fat pad weights of animals is generally indicative of a reduction of total body fat of the treated animals.

TABLE V

REPARTITION EFFECT OF TEST COMPOUNDS

<u>COMPOUND</u>	<u>DOSE PPM</u>	<u>% AGR* GAIN OVER CONTROL</u>	<u>% FAT REDUCTION OVER CONTROL</u>	<u>FAT VS BODY WEIGHT RATIO</u>
1,3-Benzodioxole-2,2-dicarboxylic acid, 5-[2-[2-(3-chlorophenyl)-2-hydroxyethyl]amino]propyl]-, disodium salt, (R*,R*)-(+/-).	200	-3.47	56.76	1:29
1,3-Benzodioxole-2,2-dicarboxylic acid, 5-[2-[2-(3-chlorophenyl)-2-hydroxyethyl]amino]propyl]-, dimethyl ester, hydrobromide, (R*,R*)-(+/-).	200	12.90	65.51	1:43
1,3-Benzodioxole-2,2-dicarboxylic acid, 5-[2-[2-(3-chlorophenyl)-2-hydroxyethyl]amino]propyl]-, disodium salt, [R-(R*,R*)].	200	14.30	53.65	1:33
	100	4.36	58.19	1:33
	50	-9.23	54.35	1:26
1,3-Benzodioxole-2,2-dicarboxylic acid, 5-[2-[2-(3-chlorophenyl)-2-hydroxyethyl]amino]propyl]-, bis(1-methylethyl)ester, hydrochloride, [R-(R*,R*)].	200	23.92	60.25	1:43
	100	9.74	60.90	1:40
	50	15.58	68.01	1:32
CONTROL				1:13

*AGR - animal growth regulant gain

THIS SHEET HAS BEEN
REPLACED

- 45 -



This Sheet Has Been Replaced



5 The compounds showed good fat reducing activity without interfering with normal animal growth.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

10 The following specific examples illustrate the present invention. They are not to be construed to limit the claims in any manner whatsoever.

EXAMPLE 1

15 (R*,R*)-(+/-)-5-(2-((2-(3-Chlorophenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester

20 To a mixture of 53.2 g of trimethylsilyl cyanide and 0.5 g of anhydrous aluminum chloride under argon is added dropwise 75.3 g of 3-chlorobenzaldehyde at such a rate that the reaction temperature does not exceed 80°C. The mixture is stirred for 30 minutes, then filtered and washed with ether. The combined filtrate and wash is evaporated to an oil which is distilled on a Kugelrohr giving 107.4 g of O-trimethylsilyl-3-chloromandelonitrile (70-77°C) as a pale yellow liquid.

25 To a mixture of 34.7 g of sodium borohydride and 500 ml of tetrahydrofuran is added 104.6 g of trifluoroacetic acid over 40 minutes with stirring in a 20°C water bath. A 110 g portion of O-trimethylsilyl-3-chloromandelonitrile is added over 45 minutes, followed by 150 ml of tetrahydrofuran. The mixture is stirred overnight in the 20°C water bath, then 500 ml of water is added dropwise with ice bath cooling. The mixture is then stirred at room temperature for 1 hour, filtered through diatomaceous earth, washed with tetrahydrofuran and the combined filtrate and wash are

30

35



concentrated in vacuo until most of the tetrahydrofuran is removed. A 250 ml portion of water and 150 ml of concentrated hydrochloric acid are added, the solution is heated on a steam bath for 1.5 hours, cooled and extracted twice with dichloromethane. The aqueous extracts are combined, made strongly basic with 200 ml of 10N sodium hydroxide in a cooling bath and extracted four times with dichloromethane. The dichloromethane extracts are combined, washed twice with water, dried and evaporated to a yellow oil. This oil is dissolved in ether, filtered through diatomaceous earth and evaporated to an oil. This oil is distilled on a Kugelrohr giving 58.73 g of 2-(3-chlorophenyl)-2-hydroxyethylamine as thick orange oil.

A mixture of 4 g of 2-(3-chlorophenyl)-2-hydroxyethylamine, 4.53 g of 3,4-dimethoxyphenylacetone, 2 g of sodium cyanoborohydride and 40 ml of methanol is stirred for 3 hours and then worked up giving 7.5 g of oil. This oil is purified by chromatography, eluting with hexane:ethyl acetate (1:1), then ethyl acetate, giving 5.35 g of 3-chloro- α -(((2-(3,4-dimethoxyphenyl)-1-methylethyl)amino)methyl)benzenemethanol as thick yellow syrup.

A mixture of 4.39 g of 3-chloro- α -(((2-(3,4-dimethoxyphenyl)-1-methylethyl)amino)methyl)benzenemethanol, 4.5 g of carbonyldiimidazole, 13 ml of triethylamine and 45 ml of tetrahydrofuran are stirred overnight, then poured into water and extracted twice with ethyl acetate. The extracts are combined, washed twice with 2N hydrochloric acid, once with brine, dried and evaporated to an oil. This oil is chromatographed, eluting with hexane:ethyl acetate (3:1 to 2:1). Fractions 16-19 are combined and evaporated giving 2.02 g of a colorless syrup. To an ice cold solution of 1.71 g of this syrup in 70 ml of dichloromethane is added



dropwise 1.3 ml of boron tribromide. The mixture is stirred at 0° to 5°C for 15 minutes, then at room temperature for 20 minutes, quenched with water and stirred for 20 minutes. The dichloromethane layer is separated, washed with brine, dried and evaporated, giving 1.6 g of (R*,S*)-(+/-)-5-(3-chlorophenyl)-3-(2-(3, 4-dihydroxyphenyl)-1-methylethyl)-2-oxazolidinone as a foamy solid.

A mixture of 0.8 g of the above oxazolidinone, 0.74 g of diethyl dibromomalonate, 1.2 g of anhydrous potassium carbonate and 20 ml of acetone are stirred overnight with the addition of a few drops of diethyl dibromomalonate. The mixture is filtered, washed with acetone and the combined filtrate and wash are evaporated to a yellow oil. The oil is purified by flash chromatography, eluting with 5 percent acetone in toluene. The pure fractions are combined and evaporated, giving 766 mg of (R*,S*)-(+/-)-5-(2-(5-(3-chlorophenyl)-2-oxo-3-oxazolidinyl)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, diethyl ester as a colorless oil.

A mixture of 636 mg of the above diethyl ester, 23 ml of 5N sodium hydroxide and 44 ml of ethanol are heated at reflux under argon overnight, then cooled and evaporated. The solid residue is taken up in methanol, placed in an ice bath and hydrogen chloride gas is passed through the solution for 3-4 minutes. This mixture is stirred for 1.5 hours, then poured cautiously into saturated aqueous sodium bicarbonate and extracted with ethyl acetate. This ethyl acetate extract is washed with brine, dried, filtered and evaporated to an oil. This oil is purified by flash chromatography, eluting with ethyl acetate. The pure fractions are combined and evaporated, giving 264 mg of the desired product as a colorless oil.

EXAMPLE 2

(R*,S*)-(+/-)-((5-(2-(5-(3-chlorophenyl))-2-
oxo-3-oxazolidinyl)propyl)-1,3-benzodioxole-
2,2-diyl)bis(methyleneoxy)bis acetic
acid, dimethyl ester


A mixture of 1.63 g of (R*,S*)-(+/-)-5-(2-(5-(3-chlorophenyl)-2-oxo-3-oxazolidinyl)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, diethyl ester and 2.5 g of lithium borohydride in 25 ml of anhydrous tetrahydrofuran are reacted, giving 1.26 g of the corresponding alcohol as a white foam.

A 590 mg portion of 60 percent sodium hydride is washed three times with hexane. To this is added under argon, 10 ml of dry tetrahydrofuran and, over 5 minutes, a solution of 1.24 g of (R*,R*)-(+/-)-5-(2-(5-(3-chlorophenyl)-2-oxo-3-oxazolidinyl)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid in 10 ml of dry tetrahydrofuran. The solution is stirred for 5 minutes, then 1.36 g of methyl bromoacetate is added over 10 minutes. After stirring overnight this mixture is poured into aqueous ammonium chloride and extracted twice with ethyl acetate. The extracts are combined, dried and evaporated to an oil which is purified by chromatography, giving 1.03 g of the desired product as a pale yellow oil.

EXAMPLE 3

(R*,R*)-(+/-)-3-(2-(2,2-Bis((2-hydroxyethoxy)-methyl)-1,3-benzodioxol-5-yl)-1-methylethyl)-5-(3-chlorophenyl)-2-oxazolidinone

A mixture of 420 mg of (R*,R*)-(+/-)-5-(2-(5-(3-chlorophenyl)-2-oxo-3-oxazolidinyl)propyl)-1,3-benzodioxole)-2,2-diyl)bis(methyleneoxy)bis acetic acid, dimethyl ester, 0.75 g of lithium borohydride and 10 ml of anhydrous tetrahydrofuran are allowed to react for 1.5 hours, giving 340 mg of the desired product as a milky film.

EXAMPLE 4

(R^{*},R^{*})-(+/)-((5-(2-(2-(3-chlorophenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-diyl)bis(methyleneoxy)bis acetic acid, dimethyl ester

5 A mixture of 370 mg of (R^{*},R^{*})-(+/)-((5-(2-(2-(3-chlorophenyl)-2-oxo-3-oxazolidinyl)propyl)-1,3-benzodioxole-2,2-diyl)bis(methyleneoxy)bis acetic acid, dimethyl ester, 9 ml of 5N sodium hydroxide and 19 ml of absolute ethanol are refluxed overnight, then cooled, acidified to pH 5 with concentrated hydrochloric acid and evaporated to dryness in vacuo. The residue is dissolved in 20 ml of methanol, then saturated with hydrogen chloride gas and stirred for 1.5 hours. The mixture is poured into aqueous sodium bicarbonate and extracted twice with ethyl acetate. The extracts are combined, washed with brine, dried and evaporated. The residue is purified by flash chromatography, eluting with ethyl acetate, giving 220 mg of the desired product as a pale yellow thick oil.

EXAMPLE 5

(R^{*},R^{*})-(+/)-alpha-(((2-(2,2-bis((2-hydroxyethoxy)methyl)-1,3-benzodioxole-5-yl)-1-methylethyl)-amino)methyl)-3-chlorobenzenemethanol

25 A mixture of 270 mg of (R^{*},R^{*})-(+/)-3-(2-(2,2-bis((2-hydroxyethoxy)methyl)-1,3-benzodioxole-5-yl)-1-methylethyl)-5-chlorophenyl)-2-oxazolidinone, 200 ml of ethanol and 5 ml of 5N sodium hydroxide are refluxed under argon for 8 hours, then poured into salty water and extracted twice with ethyl acetate. The extracts are combined, washed with brine, dried and evaporated. The residue is purified by flash chromatography, eluting with dichloromethane:methanol:ammonium hydroxide (250:35:5). Fractions 3 and 4 are combined and evaporated, giving 170 mg of the desired product as a thick yellow oil.

EXAMPLE 6

1,3-Benzodioxole-2,2-Dicarboxylic Acid, 5-(2-((2-(3-Chlorophenyl)-2-Hydroxyethyl)Amino)Propyl),
Disodium Salt, (R*,R*)-(+/-)-

5 A solution of (1,3-benzodioxole-2,2-di-carboxylic acid, 5-(2-(5-(3-chlorophenyl)-2-oxo-3-oxazolidinyl)propyl-, diethyl ester, (R*,R*)-(+/-)-, (550 mg 1.09 mmol), 5N NaOH (15 mL) and distilled water (15 mL) is heated at reflux under an argon atmosphere
10 for 17 hours. The solution is cooled, and then acidified to pH 9 using concentrated HCl causing a copious amount of white solid to precipitate. The mixture is filtered and the solid is washed with water. The combined filtrate and water wash (53 mL) is loaded
15 onto a column packed with C₁₈ reverse phase silica gel (75 mL) which had first been washed with MeOH (150 mL) and then a pH 8.5 solution and then the product is eluted using MeOH:water (1:1). Evaporation of the product-containing fractions yielded a yellow oil which
20 became a powdery solid upon addition of MeOH. The solid was collected and washed with ether to yield a dry white powder, 395 mg (78%).

EXAMPLE 7

25 1,3-Benzodioxole-2,2-Dicarboxylic Acid, 5-(2-((2-(3-Chlorophenyl)-2-Hydroxyethyl)Amino)Propyl)-, Dimethyl
Ester, Hydrobromide, (R*,R*)-(+/-)-

Absolute methanol (20 mL) is treated dropwise with distilled acetyl bromide (20 drops) and the solution is stirred for 5 minutes. To this solution is
30 added (R*,R*)-(+/-)-5-(2-((2-(3-chlorophenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, disodium salt (500 mg, 1.09 mmol) and the solution is stirred for 24 hours at room temperature. The solution is evaporated to give to yield a brown oil
35 which is dissolved in chloroform and filtered to remove



sodium bromide. The chloroform is evaporated and ether is added to the resulting brown oil to give a beige solid. The solid is collected and washed with ether to give a dry powder (520 mg, 90%). Recrystallization from acetonitrile-ether yielded the analytical sample, mp 145 - 147°C.

EXAMPLE 8

1,3-Benzodioxole-2,2-Dicarboxylic Acid, 5-(2-((2-(3-Chlorophenyl)-2-Hydroxyethyl)Amino)Propyl)-,

Disodium salt, (R,R) and

1,3-Benzodioxole-2,2-Dicarboxylic acid, 5-(2-((-3-Chlorophenyl)-2-Hydroxyethyl)Amino)Propyl)-,

Disodium Salt, (S,S)

A solution of (R*,R*)-(+/-)-5-(2((2-(3-chlorophenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester hydrobromide (1.00 g, 1.88 mmol), triethylamine (0.48g, 4.71 mmol) and (S)-(-)-2-trifluoromethyl-2-methoxyphenylacetyl chloride (0.54 g, 2.26 mmol) in 10 mL of methylene chloride is stirred at room temperature for 3 days. The yellow solution is diluted with 50 mL of ether and washed sequentially with 2N HCl, aqueous sodium bicarbonate, and brine. The organic solution is dried (MgSO₄), filtered and concentrated to give a yellow oil. Purification by preparative HPLC (silica, hexane: ethyl acetate 4:1) yielded: (1) the less polar isomer, 229 mg (18%) and (2) the more polar isomer, 248 mg (20%), both as dry foams.

The more polar isomer (235 mg, 0.353 mmol) is heated at reflux in a solution of ethanol (4mL) and 5N NaOH (2mL) for 24 hours under argon atmosphere. The solution is cooled and acidified to pH 9 using concentrated HCl causing some white solid to form. The entire mixture is loaded onto a flash chromatography



column packed with C₁₈ silica (40 mL) which had been prepared as described in the (R*,S*)-5-(2-((2-(3-chlorophenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, disodium salt experiment.

Isolation of the product is accomplished in the same manner as was previously described. The product (single enantiomer), a white powder (117 mg, 71%) is obtained.

The other enantiomer is obtained by a similar hydrolysis of the less polar diastereomer.

EXAMPLE 9

(R,R)-1,3-Benzodioxole-2,2-dicarboxylic acid,
5-(2-((2-(3-chlorophenyl)-2-hydroxyethyl)
amino)propyl)-, disodium salt

A mixture of 25 g of L-DOPA, 27.7 g of di-*t*-butyl dicarbonate in 300 ml of dimethylformamide is heated at 65°C for 22 hours, cooled, poured into 5 percent citric acid solution and extracted twice with ethyl acetate. The ethyl acetate extracts are washed with brine, dried, filtered and evaporated to yield 31.0 g of (S)-N-(1,1-dimethylethoxy)carbonyl-3-hydroxy-L-tyrosine, as a brown oil.

A mixture of 31.0 g of the above *t*-butyl carbonate derivative, 52 ml of iodomethane, and 145 g of anhydrous potassium carbonate in 300 ml of acetone is heated to reflux for 20 hours, cooled, filtered and the solvent evaporated to yield a residue. The residue is then partitioned between water and methylene chloride. The methylene chloride extract is washed with brine, dried, filtered and evaporated to yield 27 g of (S)-N-(1,1-dimethylethoxy)carbonyl-3,4-dimethoxy-L-phenylalanine, methyl ester as a yellow solid.

A mixture of 27.0 g of the above ester derivative, 3.7 g of lithium borohydride and 180 ml of tetrahydrofuran is allowed to react for 18 hours, giving 22.0 g of (S)-1,1-dimethylethyl-(2-(3,4-dimethoxyphenyl)-1-(hydroxymethyl)ethyl)carbamate as a white solid.



To an ice cold solution of 22.0 g of the above hydroxymethyl derivative, 15 ml of triethylamine in 140 ml of anhydrous tetrahydrofuran, is added 7.0 ml of methanesulfonylchloride, stirred at room temperature for 2 hours, diluted with water, and extracted with ethyl acetate. The ethyl acetate extract is washed with 2N hydrochloric acid, 1N sodium hydroxide, brine, and then dried, filtered and evaporated to give 25.0 g of (S)-1, 1-dimethylethyl-(2-(3,4-dimethoxyphenyl)-1-((methylsulfonyl)oxy)methyl)ethyl)carbamate as a white solid.

A mixture of 18.0 g of the above mesyl derivative and 35 ml of trifluoroacetic acid in 80 ml of methyl chloride is reacted for 18 hours and solvent is evaporated to yield a gray oil. The oil is dissolved in ethanol and 15 g of sodium acetate, and 1.8 g of 10 percent palladium on carbon is added. The resulting mixture is then hydrogenated on a Parr shaker overnight. The catalyst is filtered and the filtrate evaporated to yield an oil. The oil is purified by kugelrohr distillation to yield 5.8 g of (R)-3,4-dimethoxymethylbenzeneethanamine as a white solid.

A mixture of 420 mg of the above amine and 311 mg of N-(trimethylsilyl)acetamide in 2 ml of dimethylsulfoxide is stirred at room temperature for 1 hour. A solution of 349 mg of 3-chlorostyreneoxide in 0.3 ml dimethylsulfoxide is added to the above mixture and the resulting mixture is heated at 65°C to 70°C for 20 hours, cooled, poured into a mixture of ice and concentrated hydrochloric acid, stirred, basified with 10N NaOH and extracted with ethyl acetate. The ethyl acetate extract is washed with brine, dried, filtered and evaporated to yield an oil which is purified by flash chromatography to yield 489 mg of (R,S) and (S,S)-3-chloro- α -(((2-(3,4-dimethoxyphenyl)-1-methylethyl)-amino)methyl)benzenemethanol.



5 A mixture of 1.4 g of the above benzenemethanol derivatives, 1.39 g of carbonyldiimidazole, 3.5 ml of triethylamine and 15 ml of tetrahydrofuran is stirred overnight, then poured into water and extracted twice with ethyl acetate. The ethyl acetate extract is washed with 2N hydrochloric acid, brine, dried and evaporated to an oil. The oil is chromatographed, eluting with hexane:ethyl acetate (3:1 to 2:1). Fractions containing bottom spot are combined and evaporated giving 318 mg of colorless oil. To an ice cold solution of 300 mg of this oil is added dropwise 0.23 ml of boron tribromide. The mixture is stirred at 0°C to 5°C for 15 minutes, then at room temperature for 20 minutes, quenched with water and stirred for 20 minutes. The dichloromethane layer is separated, washed with brine, dried and evaporated giving 254 mg of (R,R)-5-(3-chlorophenyl)-3-(2-(3,4-dihydroxyphenyl)-1-methylethyl)-2-oxazolidinone as a foaming solid.

20 A mixture of 240 mg of the above oxazolidinone, 234 mg of diethyl bromomalonate, 450 mg of anhydrous potassium carbonate and 10 ml of acetone is stirred overnight, filtered, washed with acetone and evaporated to a brown oil. The oil is purified by flash chromatography, eluting with 5 percent acetone in toluene. The pure fractions are combined and evaporated giving 237 mg of (R,R)-5-(3-chlorophenyl)-2-oxo-3-oxazolidinylpropyl)-1,3-benzodioxole-2,2-dicarboxylic acid, diethyl ester as a colorless oil.

30 A mixture of 234 mg of the above diethyl ester, 8.5 ml of 5N sodium hydroxide and 17 ml of ethanol is heated at reflux under argon overnight, cooled and then acidified to pH 9 using concentrated hydrochloric acid causing a copious amount of a white solid to precipitate. The mixture is filtered and the solid is washed with water. The combined filtrate and



water wash (22 ml) is loaded onto a column packed with C₁₈ reverse phase silica gel, which was first washed with methanol (64 ml), and then a pH 8.5 solution. The product is eluted using methanol:water (1:1). Evaporation of the product containing fractions gives a yellow oil which becomes a powdery solid upon the addition of methanol. The solid is collected and washed with ether to yield 8.5 mg of the compound of this example as a white solid.

EXAMPLE 10

(R*,R*)-(+/-)-5-(2-((2-(3-chlorophenyl)-2-hydroxyethyl)-amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid,
diethyl ester, hydrobromide

The procedure to prepare the dimethyl ester of Example 7 is repeated, substituting ethanol for methanol to produce the compound of the example as a white powder, mp 174-176°C.

EXAMPLE 11

(R*,R*)-(+/-)-5-(2-((2-(3-chlorophenyl)-2-hydroxyethyl)-amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid,
diisopropyl ester, hydrobromide

The procedure to prepare the dimethyl ester of Example 7 is repeated, substituting 2-propanol for methanol to produce the compound of the example as an off-white powder, mp 170-172°C.

EXAMPLES 12 - 107

The procedure of Example 1 is repeated, except that the substituents on the starting compounds are varied. Table A sets forth the compounds produced in accordance with the present invention.



TABLE A

<u>EXAMPLE</u>	<u>COMPOUND</u>
12	5-(2-((-2,3-dichlorophenyl)-2-hydroxyethyl) amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, diethyl ester
13	5-(2-((2-(3-chlorophenyl)-2-hydroxyethyl) amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, diethyl ester
14	5-(2-((2-(3-chlorophenyl)-2-hydroxyethyl) amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dibutyl ester
15	5-(2-((2-(3-chlorophenyl)-2-hydroxyethyl) amino)butyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester
16	5-(2-((2-(3,4-dichlorophenyl)-2-hydroxyethyl) amino)butyl)-1,3-benzodioxole-2,2-dicarboxylic acid, diethyl ester
17	5-(2-((2-(3-chlorophenyl)-2-hydroxyethyl) amino)butyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dibutyl ester
18	5-(2-((2-(3-bromophenyl)-2-hydroxyethyl) amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester.
19	5-(2-((2-(3-bromophenyl)-2-hydroxyethyl) amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, diethyl ester
20	5-(2-((2-(3-bromophenyl)-2-hydroxyethyl) amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dibutyl ester
21	5-(2-((2-(3-bromophenyl)-2-hydroxyethyl) amino)butyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester
22	5-(2-((2-(3-bromophenyl)-2-hydroxyethyl) amino)butyl)-1,3-benzodioxole-2,2-dicarboxylic acid, diethyl ester



TABLE A.
CONTINUED

<u>EXAMPLE</u>	<u>COMPOUND</u>
23	5-(2-((2-(3-bromophenyl)-2-hydroxyethyl)amino)butyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dibutyl ester
24	5-(2-((2-(3-fluorophenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester
25	5-(2-((2-(3-fluorophenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, diethyl ester
26	5-(2-((2-(3-fluorophenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dibutyl ester
27	5-(2-((2-(3-fluorophenyl)-2-hydroxyethyl)amino)butyl)-1,3-benzodioxole-2,2-dicarboxylic acid, diethyl ester
28	5-(2-((2-(3-fluorophenyl)-2-hydroxyethyl)amino)butyl)-1,3-benzodioxole-2,2-dicarboxylic acid, diethyl ester
29	5-(2-((2-(3-fluorophenyl)-2-hydroxyethyl)amino)pentyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dibutyl ester
30	5-(2-((2-(3-trifluoromethylphenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester
31	5-(2-((2-(3-trifluoromethylphenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, diethyl ester
32	5-(2-((2-(3-trifluoromethylphenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dibutyl ester
33	5-(2-((2-(3-trifluoromethylphenyl)-2-hydroxyethyl)amino)butyl)-1,3-benzodioxole-2,4-dicarboxylic acid, dimethyl ester



TABLE A
CONTINUED

<u>EXAMPLE</u>	<u>COMPOUND</u>
34	5-(2-((2-(3-trifluoromethylphenyl)-2-hydroxyethyl)amino)butyl)-1,3-benzodioxole-2,2-dicarboxylic acid, diethyl ester
35	5-(2-((2-(3-trifluoromethylphenyl)-2-hydroxyethyl)amino)butyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dibutyl ester
36	5-(2-((2-(3-methoxyphenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester
37	5-(2-((2-(3-methoxy-4-chlorophenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, diethyl ester
38	5-(2-((2-(3-methoxyphenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dibutyl ester
39	5-(2-((2-(3-methoxyphenyl)-2-hydroxyethyl)amino)butyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester
40	5-(2-((2-(3-methoxyphenyl)-2-hydroxyethyl)amino)butyl)-1,3-benzodioxole-2,2-dicarboxylic acid, diethyl ester
41	5-(2-((2-(3-methoxyphenyl)-2-hydroxyethyl)amino)butyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dibutyl ester
42	5-(2-((2-(3-methylphenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester
43	5-(2-((2-(3-methylphenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, diethyl ester
44	5-(2-((2-(3-methylphenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dibutyl ester



TABLE A.
CONTINUED

<u>EXAMPLE</u>	<u>COMPOUND</u>
45	5-(2-((2-(3-methylphenyl)-2-hydroxyethyl)amino)butyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester
46	5-(2-((2-(3-methylphenyl)-2-hydroxyethyl)amino)butyl)-1,3-benzodioxole-2,2-dicarboxylic acid, diethyl ester
47	5-(2-((2-(3-methylphenyl)-2-hydroxyethyl)amino)butyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dibutyl ester
48	5-(2-((2-(3-dimethylaminophenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester
49	5-(2-((2-(3-dimethylaminophenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, diethyl ester
50	5-(2-((2-(3-dimethylaminophenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dibutyl ester
51	5-(2-((2-(3-dimethylaminophenyl)-2-hydroxyethyl)amino)butyl)-1,3-benzodioxole-2,2-dicarboxylic acid, diethyl ester
52	5-(2-((2-(3-dimethylaminophenyl)-2-hydroxyethyl)amino)butyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester
53	5-(2-((2-(3-dimethylaminophenyl)-2-hydroxyethyl)amino)butyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dibutyl ester
54	5-(2-((2-(3-carbomethoxyphenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester
55	5-(2-((2-(3-carbomethoxyphenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, diethyl ester



TABLE A
CONTINUED

<u>EXAMPLE</u>	<u>COMPOUND</u>
56	5-(2-((2-(3-carbomethoxyphenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dibutyl ester
57	5-(2-((2-(3-carbomethoxyphenyl)-2-hydroxyethyl)amino)butyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester
58	5-(2-((2-(3-carbomethoxyphenyl)-2-hydroxyethyl)amino)butyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dibutyl ester
59	5-(2-((2-(3-methylthiophenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester
60	5-(2-((2-(3-carbomethoxyphenyl)-2-hydroxyethyl)amino)butyl)-1,3-benzodioxole-2,2-dicarboxylic acid, diethyl ester
61	5-(2-((2-(3-methylthiophenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, diethyl ester
62	5-(2-((2-(3-methylthiophenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dibutyl ester
63	5-(2-((2-(3-methylthiophenyl)-2-hydroxyethyl)amino)butyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester
64	5-(2-((2-(3-methylthiophenyl)-2-hydroxyethyl)amino)butyl)-1,3-benzodioxole-2,2-dicarboxylic acid, diethyl ester
65	5-(2-((2-(3-methylthio-4-chlorophenyl)-2-hydroxyethyl)amino)butyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dibutyl ester
66	5-(2-((2-(3-cyanophenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester

This Sheet Has Been Replaced



TABLE A
CONTINUED

<u>EXAMPLE</u>	<u>COMPOUND</u>
67	5-(2-((2-(3-cyanophenyl)-2-hydroxyethyl) amino)propyl)-1,3-benzodioxole-2,2-di-carboxylic acid, diethyl ester
68	5-(2-((2-(3-cyanophenyl)-2-hydroxyethyl) amino)propyl)-1,3-benzodioxole-2,2-di-carboxylic acid, dibutyl ester
69	5-(2-((2-(3-cyanophenyl)-2-hydroxyethyl) amino)butyl)-1,3-benzodioxole-2,2-di-carboxylic acid, dimethyl ester
70	5-(2-((2-(3-cyanophenyl)-2-hydroxyethyl) amino)butyl)-1,3-benzodioxole-2,2-di-carboxylic acid, diethyl ester
71	5-(2-((2-(3-cyanophenyl)-2-hydroxyethyl) amino)butyl)-1,3-benzodioxole-2,2-di-carboxylic acid, dibutyl ester
72	5-(2-((2-(2-chlorophenyl)-2-hydroxyethyl) amino)propyl)-1,3-benzodioxole-2,2-di-carboxylic acid, dimethyl ester
73	5-(2-((2-(2-chlorophenyl)-2-hydroxyethyl) amino)propyl)-1,3-benzodioxole-2,2-di-carboxylic acid, diethyl ester
74	5-(2-((2-(2-bromophenyl)-2-hydroxyethyl) amino)propyl)-1,3-benzodioxole-2,2-di-carboxylic acid, dimethyl ester
75	5-(2-((2-(2-bromophenyl)-2-hydroxyethyl) amino)propyl)-1,3-benzodioxole-2,2-di-carboxylic acid, dipropyl ester
76	5-(2-((2-(2-fluorophenyl)-2-hydroxyethyl) amino)propyl)-1,3-benzodioxole-2,2-di-carboxylic acid, dimethyl ester
77	5-(2-((2-(2-fluorophenyl)-2-hydroxyethyl) amino)propyl)-1,3-benzodioxole-2,2-di-carboxylic acid, diethyl ester

THIS SHEET HAS BEEN
REPLACED



TABLE A
CONTINUED

<u>EXAMPLE</u>	<u>COMPOUND</u>
78	5-(2-((2-(2-trifluoromethylphenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester
79	5-(2-((2-(2-trifluoromethylphenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, diethyl ester
80	5-(2-((2-(2-methoxyphenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester
81	5-(2-((2-(2-methoxyphenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, diethyl ester
82	5-(2-((2-(2-methylphenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester
83	5-(2-((2-(2-methylphenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, diethyl ester
84	5-(2-((2-(2-dimethylaminophenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester
85	5-(2-((2-(2-dimethylaminophenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, diethyl ester
86	5-(2-((2-(2-methylthiophenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester
87	5-(2-((2-(2-methylthiophenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, diethyl ester




TABLE A
CONTINUED

<u>EXAMPLE</u>	<u>COMPOUND</u>
88	5-(2-((2-(2-ethoxyphenyl)-2-hydroxyethyl) amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester
89	5-(2-((2-(2-ethoxy-3-chlorophenyl)-2-hydroxyethyl) amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, diethyl ester
90	5-(2-((2-(4-chlorophenyl)-2-hydroxyethyl) amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester
91	5-(2-((2-(4-chlorophenyl)-2-hydroxyethyl) amino)butyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester
92	5-(2-((2-(4-iodophenyl)-2-hydroxyethyl) amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester
93	5-(2-((2-(4-iodophenyl)-2-hydroxyethyl) amino)butyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester
94	5-(2-((2-(4-fluorophenyl)-2-hydroxyethyl) amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester
95	5-(2-((2-(4-fluorophenyl)-2-hydroxyethyl) amino)butyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester
96	5-(2-((2-(4-trifluoromethylphenyl)-2-hydroxyethyl) amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester
97	5-(2-((2-(4-trifluoromethylphenyl)-2-hydroxyethyl) amino)butyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester



TABLE A
CONTINUED

<u>EXAMPLE</u>	<u>COMPOUND</u>
98	5-(2-((2-(4-methoxyphenyl)-2-hydroxyethyl) amino)propyl)-1,3-benzodioxole-2,2 di-carboxylic acid, dimethyl ester
99	5-(2-((2-(4-methoxyphenyl)-2-hydroxyethyl) amino)butyl)-1,3-benzodioxole-2,2 di-carboxylic acid, dimethyl ester
100	5-(2-((2-(4-ethylphenyl)-2-hydroxyethyl) amino)propyl)-1,3-benzodioxole-2,2 di-carboxylic acid, dimethyl ester
101	5-(2-((2-(4-ethylphenyl)-2-hydroxyethyl) amino)butyl)-1,3-benzodioxole-2,2 di-carboxylic acid, dimethyl ester
102	5-(2-((2-(4-dimethylaminophenyl)-2-hydroxy-ethyl) amino)propyl)-1,3-benzodioxole-2,2 dicarboxylic acid, dimethyl ester
103	5-(2-((2-(4-dimethylaminophenyl)-2-hydroxy-ethyl) amino)butyl)-1,3-benzodioxole-2,2 dicarboxylic acid, dimethyl ester
104	5-(2-((2-(4-methylthiophenyl)-2-hydroxy-ethyl) amino)propyl)-1,3-benzodioxole-2,2 dicarboxylic acid, dimethyl ester
105	5-(2-((2-(4-methylthiophenyl)-2-hydroxy-ethyl) amino)butyl)-1,3-benzodioxole-2,2 dicarboxylic acid, dimethyl ester
106	5-(2-((2-(4-cyanophenyl)-2-hydroxyethyl) amino)propyl)-1,3-benzodioxole-2,2-di-dicarboxylic acid, dimethyl ester
107	5-(2-((2-(4-cyanophenyl)-2-hydroxyethyl) amino)butyl)-1,3-benzodioxole-2,2-di-dicarboxylic acid, dimethyl ester

This Sheet Has Been Replaced



- 66 -


The above mentioned patents and publications are incorporated herein by reference.

5 Many variations of the present invention will suggest themselves to those who are skilled in the art in light of the above detailed description. All such obvious modifications are within the full intended scope of the appended claims.



THIS SHEET HAS BEEN
REPLACED

CLAIMS:

1. A compound of the formula



wherein R_1 and R_4 may be one or more groups which may be the same or different and are selected from the group consisting of hydrogen, C_1 to C_4 alkyl, C_1 to C_4 alkoxy, hydroxy, halogen, trifluoromethyl, carboxy, hydroxyalkyl, alkoxy carbonyl, C_1 to C_4 thioalkyl, sulfonyl and sulfinyl; X is a divalent radical consisting of




wherein R' is selected from the group consisting of hydrogen, C_1 to C_4 alkyl and C_1 to C_4 acyl, and Y is selected from the group consisting of carbonyl, and thiocarbonyl; R_2 and R_3 may be the same or different and are selected from the group consisting of hydrogen and C_1 to C_4 alkyl; R_5 and R_6 are selected from the group consisting of hydrogen, carboxy, alkoxy carbonyl, hydroxymethyl, $-CH_2OCH_2COOR_7$ and $-CH_2OCH_2CH_2OR_7$ where R_7 is hydrogen or C_1 to C_4 alkyl; except that R_5 and R_6 may not both be hydrogen; the asterisks denote asymmetric carbon atoms; or a pharmaceutically acceptable salt or ester thereof, the enantiomers thereof, the racemic mixtures thereof and the diastereomeric mixtures thereof.



THIS SHEET HAS BEEN
REPLACED



2. A compound as defined in Claim 1 wherein X is the divalent radical



3. A compound as defined in Claim 2 wherein R_5 and R_6 are both alkoxycarbonyl or carboxy; R_2 is hydrogen; and R_3 is C_1 to C_4 alkyl and R_1 is one or two groups.

4. The optically active compound as defined in Claim 3 which is (R,R)-5-(2-((2-(3-chlorophenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, dimethyl ester.

5. The optically active compound as defined in Claim 3 which is (R,R)-5-(2-((2-(3-chlorophenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, diethyl ester.

6. The optically active compound as defined in Claim 3 which is (R,R)-5-(2-((2-(3-chlorophenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, diisopropyl ester.

7. The optically active compound as defined in Claim 3 which is (R,R)-5-(2-((2-(3-chlorophenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, disodium salt.

8. The racemic compound as defined in Claim 3 which is (R*,S*)-(+/-)-5-(2-((2-(3-chlorophenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, diethyl ester.

9. The racemic compound as defined in Claim 3 which is (R*,S*)-(+/-)-5-(2-((2-(3-chlorophenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, diisopropyl ester.




* C L R S *

10. The racemic compound as defined in Claim 3 which is (R^*, S^*)-(+/-)-5-(2-((2-(3-chlorophenyl)-2-hydroxyethyl)amino)propyl)-1,3-benzodioxole-2,2-dicarboxylic acid, disodium salt.


11. A pharmaceutical composition for treating diabetes, hyperglycemia or obesity in humans or other mammals, comprising an effective amount of a compound as defined in Claim 1, in combination with a pharmaceutically acceptable carrier.

12. A method of treating hyperglycemia or obesity in humans or other mammals, which comprises administering to a human or other mammal an anti-obesity effective amount or an anti-hyperglycemia effective amount of a compound as defined in Claim 1.


13. A process for producing a compound of the formula:



wherein R_1 and R_4 may be one or more groups which may be the same or different and are selected from the group consisting of hydrogen, C_1 to C_4 alkyl, C_1 to C_4 alkoxy, hydroxy, halogen, trifluoromethyl, carboxy, hydroxyalkyl, alkoxycarbonyl, C_1 to C_4 thioalkyl, sulfonyl and sulfinyl; X is a divalent radical consisting of




or




wherein R' is selected from the group consisting of hydrogen, C_1 to C_4 alkyl and C_1 to C_4 acyl; and Y is selected from the group consisting of carbonyl, thiocarbonyl and methylene; R_2 and R_3 may be the same or different and are selected from the group consisting of hydrogen and C_1 to C_4 alkyl; R_5 and R_6 are selected from group consisting of hydrogen, carboxy, alkoxycarbonyl, hydroxymethyl, $-CH_2OCH_2COOR_7$ and $CH_2OCH_2CH_2OR_7$ where R_7 is selected from the group consisting of hydrogen and C_1 to C_4 alkyl; except that R_5 and R_6 may not both be hydrogen; the asterisks denote asymmetric carbon atoms; or a pharmaceutically acceptable salt or ester thereof, the enantiomers thereof, the racemic mixtures thereof and the diastereomeric mixtures thereof; said process comprising:

(a) reacting compound of the formula




with a reagent of the formula




wherein R_1 , R_2 , R_3 and R_4 are as above defined and R_8 and R_9 are C_1 to C_4 alkyl and Z is dihalomethylene, carbonyl or thiocarbonyl to produce a compound of the formula





wherein R_1 , R_2 , R_3 , R_4 and R_8 and R_9 are as above defined and; either,


(b)(i) reacting the product of step (a) with a base followed by an acid and alcohol to produce a compound of the formula



wherein R_1 , R_2 , R_3 , R_4 are as above defined and R_5 and R_6 are alkoxy carbonyl; and optionally,

(b)(ii) reacting the product of step (b)(i) with a reagent or reagents for converting the groups represented by R_5 and R_6 to carboxy, hydroxymethyl, $-\text{CH}_2\text{OCH}_2\text{COOR}_7$ and $\text{CH}_2\text{OCH}_2\text{CH}_2\text{OR}_7$ wherein R_7 is hydrogen or $\text{C}_1\text{-C}_4$ alkyl; or


(c)(i) reacting the product of step (a) with a carbonyl reducing agent to form a compound of the formula



wherein R_1, R_2, R_3, R_4 are as above defined and

(c)(ii) reacting the product of step (c)(i) with a reagent or reagents for converting the CH_2OH groups to a $-\text{CH}_2\text{OCH}_2\text{COOR}_7$ or a $-\text{CH}_2\text{OCH}_2\text{CH}_2\text{OR}_7$ group wherein R_7 is as above defined; or


(c)(iii) reacting the product of step (c)(ii) with a base to produce a compound of the formula



wherein R_{10} and R_{11} are selected from a $-\text{CH}_2\text{OCH}_2\text{COOR}_7$ group, a $-\text{CH}_2\text{OCH}_2\text{CH}_2\text{OR}_7$ group and R_7 is as above defined.


14. A method of increasing the content of lean meat in edible animals, which comprises administering to an edible animal an effective amount of the compound of Claim 1.

15. A process for producing a desired substantially pure (+) or (-) enantiomer or a derivative, salt or ester of a compound of the formula





wherein R_1 and R_4 may be one or more groups which may be the same or different and are selected from the group consisting of hydrogen, C_1 to C_4 alkyl, C_1 to C_4 alkoxy, hydroxy, halogen, trifluoromethyl, carboxy, hydroxyalkyl, alkoxycarbonyl, C_1 to C_4 thioalkyl, sulfonyl and sulfinyl; X is a divalent radical consisting of



wherein R' is hydrogen; R_2 and R_3 may be the same or different and are selected from the group consisting of hydrogen and C_1 to C_4 alkyl; R_5 and R_6 are selected from the group consisting of hydrogen, carboxy, alkoxycarbonyl, hydroxymethyl, $-\text{CH}_2\text{OCH}_2\text{COOR}_7$ and $-\text{CH}_2\text{OCH}_2\text{CH}_2\text{OR}_7$ where R_7 is hydrogen or C_1 to C_4 alkyl; except that R_5 and R_6 may not both be hydrogen; and the asterisks denote asymmetric carbon atoms; said process comprising:

(a) attaching a chiral auxiliary group at the N-9 position of a mixture of (+) and (-) enantiomers of said compound to form a new pair of diastereoisomers;

(b) separating said new pair of diastereoisomers and recovering the new (+) or (-) enantiomer; and

(c) converting the substantially pure new enantiomer into the corresponding desired substantially pure (+) or (-) enantiomer or a derivative, salt or ester of said compound.



* C L M S *