

FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5A




FIG. 5B




FIG. 6A




FIG. 6B




FIG. 7A




FIG. 7B




FIG. 8






FIG. 9A






FIG. 9B




FIG. 10A




FIG. 10B

FIG. 10C





FIG. 10D

Continuous Dynamic Filtration of
BSA-FITC from PolyBeads






FIG. 11




FIG. 12





FIG. 12







FIG. 13

b



a

FIG. 13

C





FIG. 14







FIG. 17







FIG. 19






FIG. 20A




FIG. 20B




FIG. 21






FIG. 22

FIG. 23








FIG. 25





FIG. 26A






FIG. 27A




FIG. 27B

FIG. 28




FIG. 29




FIG. 31




FIG. 33




FIG. 34




FIG. 35




FIG. 36




FIG. 37




FIG. 38





FIG. 39






FIG. 41




FIG. 42





FIG. 43






FIG. 44

FIG. 45






FIG. 46




b

Protein	Isoelectric Point (pI)	Predicted Electrophoretic Mobility Direction
Human Polyclonal IgG	6-8	Toward Cathode, Center, and Anode
Lysozyme	11	Toward Cathode


U






3 mL/min

FIG. 47A



5 mL/min

FIG. 47B



10 mL/min

FIG. 47C




FIG. 47D

FIG. 48




FIG. 49

