

Novel carboxylic acid derivatives, their preparation and use

The present invention relates to novel carboxylic acid derivatives, their preparation and use.

Endothelin is a peptide which is composed of 21 amino acids and is synthesized and released by the vascular endothelium.

Endothelin exists in three isoforms, ET-1, ET-2 and ET-3.


10 Endothelin or ET hereinafter indicates one or all isoforms of endothelin. Endothelin is a potent vasoconstrictor and has a great effect on vascular tone. It is known that this vasoconstriction is caused by the binding of endothelin to its receptor (Nature, 332, 411-415, 1988; FEBS Letters, 231, 440-444, 15 1988 and Biochem. Biophys. Res. Commun., 154, 868-875, 1988).

Increased or abnormal release of endothelin causes persistent vasoconstriction in peripheral, renal and cerebral blood vessels, which may lead to diseases. As reported in the literature, 20 elevated levels of endothelin in plasma have been found in patients with hypertension, acute myocardial infarct, pulmonary hypertension, Raynaud's syndrome, atherosclerosis and in the airways of asthmatics (Japan J. Hypertension, 12, 79 (1989), J. Vascular Med. Biology 2, 207 (1990), J. Am. Med. Association 25 264, 2868 (1990)).

Accordingly, substances which specifically inhibit the binding of endothelin to the receptor should also antagonize the various abovementioned physiological effects of endothelin and therefore 30 be valuable drugs.

It has been found (WO 94/02474) that certain carboxylic acid derivatives with the general formula Q are good inhibitors of endothelin receptors

35



40

$\text{R}^{\text{C}} = \text{COOH}$ etc.

However, this related mainly to compounds with a double bond in 45 the molecule. Besides R^{A} and R^{B} , a maximum of one hydrogen atom is permitted on the β center.


2

It has now been found, surprisingly, that this hydrogen atom can be replaced by alkyl radicals. This results in a quaternary β center with, at the same time, a large increase in the activity with regard to endothelin receptors (see Examples).

5

The invention relates to carboxylic acid derivatives of the formula I

10



where R^1 is a tetrazole [sic], nitrile [sic], COOH or a radical which can be hydrolyzed to COOH , and the other substituents have the following meanings:

- R^2 and R^3 (which can be identical or different):
- 20 phenyl or naphthyl which can be substituted by one or more of the following radicals: halogen, cyano, NO_2 , hydroxyl, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkoxy, phenoxy, C_1 - C_4 -alkylthio, amino, benzyloxy, C_1 - C_4 -alkylamino or C_1 - C_4 -dialkylamino; or
- 25 phenyl or naphthyl which are connected together in the ortho positions by a direct linkage, a methylene, ethylene or ethenylene group, or an oxygen or sulfur atom;
- 30 R^4 phenyl or naphthyl, methylenedioxyphenyl, ethylenedioxyphenyl, indanyl, indolyl, pyridyl, benzopyranyl, furanyl, benzofuranyl, isooxazolyl, isothiazolyl, 1,3,4-thiadiazolyl, pyrimidinyl, 2,3-dihydrobenzofuranyl, benzothienyl, quinolinyl, C_3 - C_7 -cycloalkyl, thienyl, oxazolyl, thiazolyl, each of which can be substituted by one or more of
- 35 the following radicals: halogen, cyano, hydroxyl, NO_2 , C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkoxy, phenoxy, C_1 - C_4 -alkylthio, amino, benzyloxy, C_1 - C_4 -alkylamino or C_1 - C_4 -dialkylamino, it being possible for the alkyl
- 40 radicals together to form a ring;
- R^5 C_1 - C_8 -alkyl, C_3 - C_6 -alkenyl, C_3 - C_6 -alkynyl or C_3 - C_8 -cycloalkyl, it being possible for each of these radicals to be substituted one or more times by: halogen, C_1 - C_4 -alkoxy,
- 45 C_1 - C_4 -alkylthio, C_1 - C_4 -alkylamino, di- C_1 - C_4 -alkylamino;


phenyl, benzyl, 1-methylnaphthyl, 2-methylnaphthyl or naphthyl, each of which can be substituted by one or more of the following radicals: halogen, cyano, hydroxyl, amino, C₁-C₄-alkyl, C₁-C₄-alkoxy, phenoxy, C₁-C₄-alkylthio,
 5 dioxomethylene [sic] or dioxoethylene [sic];

n 1 - 2.

The compounds, and the intermediates for preparing them, such as
 10 Va, may have one or more asymmetric substituted carbon atoms. Compounds of this type may be in the form of pure enantiomers or pure diastereomers or of a mixture thereof. The use of an enantiomerically pure compound as agent is preferred.

15 The invention furthermore relates to the use of the abovementioned carboxylic acid derivatives for producing drugs, in particular for producing inhibitors of endothelin receptors.

Compounds of the formula I can be prepared by initially reacting
 20 a ketone of type II with a phosphono ester of the formula III in the presence of a base to give compounds of the formula IV




Aprotic polar solvents such as DMF or THF are used as solvent.
 40

It is possible to use as base an alkali metal or alkaline earth metal hydride such as sodium hydride, potassium hydride or calcium hydride, a carbonate such as alkali metal carbonate, eg. sodium or potassium carbonate, an alkali metal or alkaline earth
 45 metal hydroxide such as sodium or potassium hydroxide, an organometallic compound such as butyllithium, an alkali metal

alcoholate such as sodium ethanolate or potassium tert-butanolate or an alkali metal amide such as lithium diisopropylamide.


The reaction is preferably carried out at a temperature in the range from 0°C to the boiling point of the solvent or solvent mixture.

The compounds of type IV can then be reacted with aromatic compounds in the presence of a catalyst to give carboxylic acid derivatives of the general formula Va




Suitable catalysts for this are strong inorganic acids and Lewis acids. Examples thereof are, inter alia, sulfuric acid, aluminum trichloride, zinc chloride or iron trichloride. When sulfuric acid is used, the free acid can be obtained directly.

Alternatively, symmetrical carboxylic acid derivatives of the formula Vb can be prepared from a β-dicarbonyl compound VI and an aromatic compound in the presence of a catalyst.



Suitable catalysts for this are strong inorganic acids and Lewis acids. Examples thereof are, inter alia, sulfuric acid, aluminum trichloride, zinc chloride or iron trichloride (see also: Gogte G.R. et al., J. Univ. Bombay, Sect. A, 27, 1958, 41).


Another possibility for preparing compounds of type Va can start from a ketone VII



5

which can be reacted with Meldrum's acid in the presence of a base such as pyridine or sodium hydride to give compounds of type VIII

5




(VIII)


10

Reaction of compounds of type VIII in diethyl ether with a Grignard reagent of the general formula IX

15




(IX)



results in compounds of type X

20



(X)

25

in which case it may be advantageous to use in addition copper salts such as copper chloride, copper bromide, copper iodide or copper cyanide and for a Lewis acid such as trimethylsilyl chloride or boron trifluoride etherate to be present.

Hydrolysis of compounds of the formula X with mineral acids such as hydrochloric acid or sulfuric acid can then afford the compounds Va ($R = OH$).

35


Further possibilities for preparing compounds Va are similar to the methods of Zimmermann H.E. et al. J. Am. Chem. Soc. 83 (1961) 1196 or Yu A.J. et al. J. Org. Chem. 23 (1958) 1004.

Compounds of the formula Va,b can be converted into the anion (or dianion for $R = H$) with a strong base such as butyllithium or lithium diisopropylamide in an inert solvent such as diethyl ether or tetrahydrofuran and under inert gas, eg. nitrogen or argon, at $-78^\circ C$ to room temperature. This anion reacts with alkylating agents of type VII at $-78^\circ C$ to room temperature.

45

6

Quenching with concentrated NH_4Cl or dilute mineral acid such as HCl results in compounds of the formula I




10

Z = halogen,
trialkylamine


Compounds of type I with $\text{R}^1 = \text{tetrazole}$ [sic] can be synthesized starting from the carboxylic acids I ($\text{R}^1 = \text{COOH}$). To do this, the
 15 carboxylic acid is reacted with thionyl chloride at room temperature to give the acid chloride, which is then reacted with aqueous ammonia solution to give the amide of the formula XII.

20




Amides of the formula XII can be reacted with oxalyl chloride or
 25 phosphorus oxychloride or trifluoroacetic anhydride in DMF or pyridine at 0°C to room temperature to give nitriles of the formula XIII

30



Reaction of nitriles of the formula XIII with sodium azide or
 35 trimethylsilyl azide in a suitable solvent such as dimethylformamide, tetrahydrofuran or 1-methyl-2-pyrrolidinone in the presence of a catalyst such as ammonium chloride (see also: Bernstein P.R. et al., Synthesis, 1987, 1133) at room temperature or elevated temperature affords the tetrazoles XIV

40




45

Compounds of the formula I can also be prepared by starting from the corresponding carboxylic acids, ie. compounds of the formula I where $R^1 = \text{COOH}$, and converting these initially in a conventional way into an activated form such as an acid halide, 5 an anhydride or imidazolidine, and then reacting the latter with an appropriate hydroxyl compound HOR. This reaction can be carried out in conventional solvents and often requires the addition of a base, in which case those mentioned above are suitable. These two steps can also be simplified, for example, by allowing the 10 carboxylic acid to act on the hydroxyl compound in the presence of a dehydrating agent such as a carbodiimide.

Compounds of the formula I can also be prepared by starting from salts of the corresponding carboxylic acids, ie. from compounds 15 of the formula I where R^1 is COR and R is OM, where M can be an alkali metal cation or the equivalent of an alkaline earth metal cation. These salts can be reacted with many compounds of the formula R-A where A is a conventional nucleofugic leaving group, for example halogen such as chlorine, bromine, iodine, or aryl- 20 or alkylsulfonyl which is unsubstituted or substituted by halogen, alkyl or haloalkyl, such as toluenesulfonyl and methylsulfonyl, or another equivalent leaving group. Compounds of the formula R-A with a reactive substituent A are known or can easily be obtained with general expert knowledge. This reaction 25 can be carried out in the usual solvents and is advantageously carried out with the addition of a base, in which case those mentioned above are suitable.


Enantiomerically pure compounds of the formula I can be obtained 30 by carrying out with racemic or diastereomeric compounds of the formula VI a classical racemate resolution with suitable enantiomerically pure bases such as brucine, strychnine, quinine, quinidine, cinchonidine, cinchonine, yohimbine, morphine, dehydroabietylamine, ephedrine (-), (+), deoxyephedrine (-), (+), 35 threo-2-amino-1-(p-nitrophenyl)-1,3-propanediol (-), (+), threo-2-(N,N-dimethylamino)-1-(p-nitrophenyl)-1,3-propanediol (+), (-) threo-2-amino-1-phenyl-1,3-propanediol (+), (-) α -methylbenzylamine (+), (-), α -(1-naphthyl)ethylamine (+), (-), α -(2-naphthyl)ethylamine (+), (-), aminomethylpinane, 40 N,N-dimethyl-1-phenylethylamine, N-methyl-1-phenylethylamine, 4-nitrophenylethylamine, pseudoephedrine, norephedrine, norpseudoephedrine, amino acid derivatives and peptide derivatives.

45 Wide variation is possible in the radical R^1 in formula I. For example, R^1 is a group



5 where R has the following meanings:

- a) a succinylimidoxy [sic] group;
- b) a 5-membered heteroaromatic radical which is linked via a
 10 nitrogen atom, such as pyrrolyl, pyrazolyl, imidazolyl and
 triazolyl, and which can carry one or two halogen atoms,
 especially fluorine and chlorine, and/or one or two of the
 following radicals:
- 15 C₁-C₄-alkyl such as methyl, ethyl, 1-propyl, 2-propyl,
 2-methyl-2-propyl, 2-methyl-1-propyl, 1-butyl, 2-butyl;
- C₁-C₄-haloalkyl, especially C₁-C₂-haloalkyl such as
 fluoromethyl, difluoromethyl, trifluoromethyl,
 20 chlorodifluoromethyl, dichlorofluoromethyl, trichloromethyl,
 1-fluoroethyl, 2-fluoroethyl, 2,2-difluoroethyl,
 2,2,2-trifluoroethyl, 2-chloro-2,2-difluoroethyl,
 2,2-dichloro-2-fluoroethyl, 2,2,2-trichloroethyl and
 pentafluoroethyl;
- 25 C₁-C₄-haloalkoxy, especially C₁-C₂-haloalkoxy such
 as difluoromethoxy, trifluoromethoxy, chlorodifluoromethoxy,
 1-fluoroethoxy, 2-fluoroethoxy, 2,2-difluoroethoxy,
 1,1,2,2-tetrafluoroethoxy, 2,2,2-trifluoroethoxy,
 30 2-chloro-1,1,2-trifluoroethoxy and pentafluoroethoxy,
 especially trifluoromethoxy;
- C₁-C₄-alkoxy such as methoxy, ethoxy, propoxy, 1-methylethoxy,
 butoxy, 1-methylpropoxy, 2-methylpropoxy, 1,1-dimethylethoxy,
 35 especially methoxy, ethoxy, 1-methylethoxy;
- C₁-C₄-alkylthio such as methylthio, ethylthio, propylthio,
 1-methylethylthio, butylthio, 1-methylpropylthio,
 2-methylpropylthio, 1,1-dimethylethylthio, especially
 40 methylthio and ethylthio;
- c) R furthermore a radical



5

where m is 0 or 1 and R⁶ and R⁷, which can be identical or different, have the following meanings:

10 hydrogen

C₁-C₈-alkyl, especially C₁-C₄-alkyl as mentioned above;

15 C₃-C₆-alkenyl such as 2-propenyl, 2-butenyl, 3-butenyl, 1-methyl-2-propenyl, 2-methyl-2-propenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 1-methyl-2-butenyl, 2-methyl-2-butenyl, 3-methyl-2-butenyl, 1-methyl-3-butenyl, 2-methyl-3-butenyl, 3-methyl-3-butenyl, 1,1-dimethyl-2-propenyl, 1,2-dimethyl-2-propenyl, 20 1-ethyl-2-propenyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, 5-hexenyl, 1-methyl-2-pentenyl, 2-methyl-2-pentenyl, 3-methyl-2-pentenyl, 4-methyl-2-pentenyl, 3-methyl-3-pentenyl, 4-methyl-3-pentenyl, 1-methyl-4-pentenyl, 2-methyl-4-pentenyl, 25 3-methyl-4-pentenyl, 4-methyl-4-pentenyl, 1,1-dimethyl-2-butenyl, 1,1-dimethyl-3-butenyl, 1,2-dimethyl-2-butenyl, 1,2-dimethyl-3-butenyl, 1,3-dimethyl-2-butenyl, 1,3-dimethyl-3-butenyl, 2,2-dimethyl-3-butenyl, 2,3-dimethyl-2-butenyl, 30 2,3-dimethyl-3-butenyl, 1-ethyl-2-butenyl, 1-ethyl-3-butenyl, 2-ethyl-2-butenyl, 2-ethyl-3-butenyl, 1,1,2-trimethyl-2-propenyl, 1-ethyl-1-methyl-2-propenyl and 1-ethyl-2-methyl-2-propenyl, especially 2-propenyl, 2-butenyl, 3-methyl-2-butenyl and 3-methyl-2-pentenyl;

35

C₃-C₆-alkynyl such as 2-propynyl, 2-butyne, 3-butyne, 1-methyl-2-propynyl, 2-pentyne, 3-pentyne, 4-pentyne, 1-methyl-3-butyne, 2-methyl-3-butyne, 1-methyl-2-butyne, 1,1-dimethyl-2-propynyl, 1-ethyl-2-propynyl, 2-hexynyl, 40 3-hexynyl, 4-hexynyl, 5-hexynyl, 1-methyl-2-pentyne, 1-methyl-3-pentyne, 1-methyl-4-pentyne, 2-methyl-3-pentyne, 2-methyl-4-pentyne, 3-methyl-4-pentyne, 4-methyl-2-pentyne, 1,1-dimethyl-2-butyne, 1,1-dimethyl-3-butyne, 1,2-dimethyl-3-butyne, 2,2-dimethyl-3-butyne, 1-ethyl-2-butyne, 1-ethyl-3-butyne, 2-ethyl-3-butyne and 1-ethyl-1-methyl-2-propynyl, preferably 2-propynyl,

45

2-butyryl, 1-methyl-2-propynyl and 1-methyl-2-butyryl, especially 2-propynyl

5 C₃-C₈-cycloalkyl, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl, cyclooctyl, it being possible for these alkyl, cycloalkyl, alkenyl and alkynyl groups each to carry one to five halogen atoms, especially fluorine or chlorine, and/or one or two of the following groups:

10 C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkoxy as mentioned above, C₃-C₆-alkenyloxy, C₃-C₆-alkenylthio, C₃-C₆-alkynyloxy, C₃-C₆-alkynylthio, with the alkenyl and alkynyl constituents in these radicals preferably
15 corresponding to the abovementioned meanings;

C₁-C₄-alkylcarbonyl such as, in particular, methylcarbonyl, ethylcarbonyl, propylcarbonyl, 1-methylethylcarbonyl, butylcarbonyl, 1-methylpropylcarbonyl,
20 2-methylpropylcarbonyl, 1,1-dimethylethylcarbonyl;

C₁-C₄-alkoxycarbonyl such as methoxycarbonyl, ethoxycarbonyl, propyloxycarbonyl, 1-methylethoxycarbonyl, butyloxycarbonyl, 1-methylpropyloxycarbonyl, 2-methylpropyloxycarbonyl,
25 1,1-dimethylethoxycarbonyl;

C₃-C₆-alkenylcarbonyl, C₃-C₆-alkynylcarbonyl, C₃-C₆-alkenyloxycarbonyl and C₃-C₆-alkynyloxycarbonyl, where the alkenyl and alkynyl radicals are preferably defined as
30 specifically stated above;


phenyl, unsubstituted or mono- or polysubstituted, eg. mono- to trisubstituted, by halogen, nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy or
35 C₁-C₄-alkylthio, such as 2-fluorophenyl, 3-chlorophenyl, 4-bromophenyl, 2-methylphenyl, 3-nitrophenyl, 4-cyanophenyl, 2-trifluoromethylphenyl, 3-methoxyphenyl, 4-trifluoroethoxyphenyl, 2-methylthiophenyl, 2,4-dichlorophenyl, 2-methoxy-3-methylphenyl,
40 2,4-dimethoxyphenyl, 2-nitro-5-cyanophenyl, 2,6-difluorophenyl;

di-C₁-C₄-alkylamino such as, in particular, dimethylamino, dipropylamino, N-propyl-N-methylamino, N-propyl-N-ethylamino,
45 diisopropylamino, N-Isopropyl-N-methylamino, N-isopropyl-N-ethylamino, N-Isopropyl-N-propylamino;

R⁶ and R⁷ furthermore phenyl which can be substituted by one or more, eg. one to three, of the following radicals: halogen, nitro, cyano, C₁-C₄-Alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy or C₁-C₄-alkylthio, as mentioned above in particular;

or R⁶ and R⁷ together form a C₄-C₇-alkylene chain which is closed to form a ring, is unsubstituted or substituted, eg. by C₁-C₄-alkyl, and which may contain a heteroatom selected from the group of oxygen, sulfur or nitrogen, such as
 -(CH₂)₄-, -(CH₂)₅-, -(CH₂)₆-, -(CH₂)₇-, -(CH₂)₂-O-(CH₂)₂-,
 -CH₂-S-(CH₂)₃-, -(CH₂)₂-O-(CH₂)₃-, -NH-(CH₂)₃-, -CH₂-NH-(CH₂)₂-,
 -CH₂-CH=CH-CH₂-, -CH=CH-(CH₂)₃-;

15 d) R furthermore a group



where k is 0, 1 and 2, p is 1, 2, 3 and 4, and R⁸ is

C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₃-C₆-alkenyl, C₃-C₆-alkynyl or unsubstituted or substituted phenyl, such as mentioned above in particular.

e) R furthermore a radical OR⁹ where R⁹ is:

hydrogen, the cation of an alkali metal such as lithium, sodium, potassium or the cation of an alkaline earth metal such as calcium, magnesium and barium, or an environmentally compatible organic ammonium ion such as tertiary C₁-C₄-alkylammonium or the ammonium ion;

C₃-C₈-cycloalkyl as mentioned above, which can carry one to three C₁-C₄-alkyl groups;

C₁-C₈-alkyl such as, in particular, methyl, ethyl, propyl, 1-methylethyl, butyl, 1-methylpropyl, 2-methylpropyl, 1,1-dimethylethyl, pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, 1,2-dimethylpropyl, 1,1-dimethylpropyl, 2,2-dimethylpropyl, 1-ethylpropyl, hexyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,3-dimethylbutyl, 1,1-dimethylbutyl, 2,2-dimethylbutyl, 3,3-dimethylbutyl, 1,1,2-trimethylpropyl, 1,2,2-trimethylpropyl, 1-ethylbutyl,

2-ethylbutyl, 1-ethyl-2-methylpropyl, which can carry one to five halogen atoms, in particular fluorine and chlorine, and/or one of the following radicals:

- 5 C₁-C₄-alkoxy, C₁-C₄-alkylthio, cyano, C₁-C₄-alkylcarbonyl, C₃-C₈-cycloalkyl, C₁-C₄-alkoxycarbonyl, phenyl, phenoxy or phenylcarbonyl, where the aromatic radicals in turn can each carry one to five halogen atoms and/or one to three of the following radicals: nitro, cyano, C₁-C₄-alkyl,
- 10 C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy and/or C₁-C₄-alkylthio, as mentioned above in particular;
- C₁-C₈-alkyl as mentioned above, which can carry one to five halogen atoms, in particular fluorine and/or chlorine, and carries one of the following radicals: a 5-membered
- 15 heteroaromatic radical containing one to three nitrogen atoms, or a 5-membered heteroaromatic radical containing one nitrogen atom and one oxygen or sulfur atom, which can carry one to four halogen atoms and/or one or two of the following
- 20 radicals:
- nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, phenyl, C₁-C₄-haloalkoxy and/or C₁-C₄-alkylthio. Particular
- 25 mention may be made of: 1-pyrazolyl, 3-methyl-1-pyrazolyl, 4-methyl-1-pyrazolyl, 3,5-dimethyl-1-pyrazolyl, 3-phenyl-1-pyrazolyl, 4-phenyl-1-pyrazolyl, 4-chloro-1-pyrazolyl, 4-bromo-1-pyrazolyl, 1-imidazolyl, 1-benzimidazolyl, 1,2,4-triazol-1-yl, 3-methyl-1,2,4-triazol-1-yl, 5-methyl-1,2,4-triazol-1-yl, 1-benzotriazolyl,
- 30 3-isopropyl-5-isoxazolyl, 3-methyl-5-isoxazolyl, 2-oxazolyl, 2-thiazolyl, 2-imidazolyl, 3-ethyl-5-isoxazolyl, 3-phenyl-5-isoxazolyl, 3-tert-butyl-5-isoxazolyl;
- C₂-C₆-alkyl which has in position 2 one of the following
- 35 radicals: C₁-C₄-alkoxyimino, C₃-C₆-alkynyloxyimino, C₃-C₆-haloalkenyloxyimino or benzyloxyimino;
- C₃-C₆-alkenyl or C₃-C₆-alkynyl, it being possible for these
- 40 groups in turn to carry one to five halogen atoms;
- R⁹ furthermore a phenyl radical which can carry one to five halogen atoms and/or one to three of the following radicals: nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy and/or C₁-C₄-alkylthio, as mentioned above in
- 45 particular;

13


- a 5-membered heteroaromatic radical which is linked via a nitrogen atom and contains one to three nitrogen atoms and which can carry one or two halogen atoms and/or one or two of the following radicals: C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, phenyl, C₁-C₄-haloalkoxy and/or C₁-C₄-alkylthio. Particular mention may be made of: 1-pyrazolyl, 3-methyl-1-pyrazolyl, 4-methyl-1-pyrazolyl, 3,5-dimethyl-1-pyrazolyl, 3-phenyl-1-pyrazolyl, 4-phenyl-1-pyrazolyl, 4-chloro-1-pyrazolyl, 4-bromo-1-pyrazolyl, 1-imidazolyl, 1-benzimidazolyl, 1,2,4-triazol-1-yl, 3-methyl-1,2,4-triazol-1-yl, 5-methyl-1,2,4-triazol-1-yl, 1-benzotriazolyl, 3,4-dichloro-1-imidazolyl;
- R⁹ furthermore a group



where R¹⁰ and R¹¹, which can be identical or different, are:

- C₁-C₈-alkyl, C₃-C₆-alkenyl, C₃-C₆-alkynyl, C₃-C₈-cycloalkyl, it being possible for these radicals to carry a C₁-C₄-alkoxy, C₁-C₄-alkylthio and/or an unsubstituted or substituted phenyl radical, as mentioned above in particular;
- phenyl, which can be substituted by one or more, eg. one to three, of the following radicals: halogen, nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy or C₁-C₄-alkylthio, where these radicals correspond in particular to those mentioned above;
- or R¹⁰ and R¹¹ together form a C₃-C₁₂-alkylene chain which can carry one to three C₁-C₄-alkyl groups and contain a heteroatom from the group of oxygen, sulfur and nitrogen, as mentioned in particular for R⁶ and R⁷.

- f) R furthermore a radical



where R¹² is:


C₁-C₄-alkyl, C₃-C₆-alkenyl, C₃-C₆-alkynyl, C₃-C₈-cycloalkyl as mentioned above in particular, it being possible for these
 5 radicals to carry a C₁-C₄-alkoxy, C₁-C₄-alkylthio and/or a phenyl radical as mentioned above;

phenyl, unsubstituted or substituted, in particular as mentioned above.

10

g) R a radical

15



20 where R¹² has the abovementioned meanings.

R¹ can furthermore be:
 tetrazole [sic] or nitrile [sic].

25 With a view to the biological effect, preferred carboxylic acid derivatives of the general formula I, both as pure enantiomers and pure diastereomers and as mixtures thereof, are those where the substituents have the following meanings:

30 R¹ tetrazole [sic], COOH or a radical which can be hydrolyzed to COOH;

R² and R³ (which can be identical or different);

35 phenyl or naphthyl which can be substituted by one or more of the following radicals: F, Cl, Br, I, cyano, NO₂, hydroxyl, methyl, ethyl, propyl, isopropyl, trifluoromethyl, 2,2,2-trifluoroethyl, methoxy, ethoxy, propoxy, isopropoxy, trifluoromethoxy, phenoxy, methylthio, ethylthio,
 40 benzyloxy, amino, methylamino, dimethylamino;

R⁴ phenyl, methylenedioxyphenyl, ethylenedioxyphenyl, indanyl, pyridyl, 2,3-dihydrobenzofuranyl, benzofuranyl, benzothienyl, 2-pyrimidinyl, 4-pyrimidinyl, 2,3-dihydrobenzothienyl, each
 45 of which can be substituted by one or more of the following radicals: F, Cl, Br, I, cyano, NO₂, methyl, ethyl, propyl, isopropyl, trifluoromethyl, methoxy, ethoxy, propoxy,

15

isopropoxy, butyloxy, tert-butyloxy, trifluoromethyloxy, phenoxy, methylthio, ethylthio, propylthio, benzyloxy, amino, methylamino, dimethylamino;

5 R⁵ methyl, ethyl, propyl, isopropyl, butyl, 2-methylpropyl, tert-butyl, pentyl, 3-methylbutyl, hexyl, 3-pentyl, 4-methylpentyl, 2-ethylbutyl, each of which can be substituted one or more times by: cyano, methoxy, ethoxy, propoxy, isopropoxy, butoxy, methylthio, ethylthio, propylthio, isopropylthio, amino, methylamino, dimethylamino;

allyl, vinyl, trifluoromethyl, 2,2,2-trifluoroethyl;

15 phenyl, benzyl, each of which can be substituted by one or more of the following radicals: F, Cl, Br, I, hydroxyl, methyl, ethyl, propyl, isopropyl, methoxy, ethoxy, propoxy, isopropoxy, methylthio, ethylthio, dioxomethylene [sic], dioxoethylene [sic];

20 n 1 - 2

25


30

35

40

45

Examples of preferred compounds are listed in the following Table:



No.	R	R ²	R ³	R ⁵	R ⁴	n
1	OH	Phenyl	Phenyl	Methyl	Phenyl	1
2	OH	Phenyl	Phenyl	Methyl	3,5-Dimethylphenyl	1
3	OH	Phenyl	Phenyl	Methyl	3,5-Dimethoxyphenyl	1
4	OH	Phenyl	Phenyl	Methyl	3,4-Methylenedioxyphenyl	1
5	OH	Phenyl	Phenyl	Methyl	2,3-Dihydro-benzofuran-5-yl	1
6	OH	Phenyl	Phenyl	Methyl	2,3-Dihydro-1H-inden-5-yl	1
7	OH	Phenyl	Phenyl	Methyl	3-Methoxy-5-methylphenyl	1
8	OH	Phenyl	Phenyl	Methyl	3,4-Ethylenedioxyphenyl	1
9	OH	Phenyl	Phenyl	Methyl	5-Methoxy-3,4-methylenedioxyphenyl	1
10	OH	Phenyl	Phenyl	Methyl	3,5-Diethylphenyl	1
11	OH	Phenyl	Phenyl	Methyl	3,4-Dimethoxyphenyl	1
12	OH	Phenyl	Phenyl	Methyl	3-Methyl-4-benzyloxyphenyl	1
13	OH	Phenyl	Phenyl	Methyl	3-Benzyloxy-5-methoxyphenyl	1
14	OH	Phenyl	Phenyl	Methyl	3-Benzyloxy-4-methoxyphenyl	1
15	OH	Phenyl	Phenyl	Methyl	3-Hydroxy-4-methoxyphenyl	1

No.	R	R ²	R ³	R ⁵	R ⁴	n
16	OH	Phenyl	Phenyl	Methyl	4-Hydroxy-3-methoxyphenyl	1
17	OH	Phenyl	Phenyl	Methyl	3,5-Bis-(trifluoromethyl)-phenyl	1
18	OH	Phenyl	Phenyl	Methyl	3-Methylphenyl	1
19	OH	Phenyl	Phenyl	Methyl	3-Methoxyphenyl	1
20	OH	Phenyl	Phenyl	Methyl	4-Benzoyloxyphenyl	1
21	OH	Phenyl	Phenyl	Methyl	4-Phenoxyphenyl	1
22	OH	Phenyl	Phenyl	Methyl	6-Methylpyridin-2-yl	1
23	OH	Phenyl	Phenyl	Methyl	6-Ethylpyridin-2-yl	1
24	OH	Phenyl	Phenyl	Methyl	6-Methoxypyridin-2-yl	1
25	OH	Phenyl	Phenyl	Methyl	4,6-Dimethylpyridin-2-yl	1
26	OH	Phenyl	Phenyl	Methyl	4,6-Diethylpyridin-2-yl	1
27	OH	Phenyl	Phenyl	Methyl	4-Methyl-6-methoxypyridin-2-yl	1
28	OH	Phenyl	Phenyl	Methyl	6,7-Dihydro-5H-cyclopenta[b]pyridin-2-yl	1
29	OH	Phenyl	Phenyl	Methyl	4,6-Dimethylpyrimidin-2-yl	1
30	OH	Phenyl	Phenyl	Methyl	4-Methoxy-6-methylpyrimidin-2-yl	1
31	OH	Phenyl	Phenyl	Methyl	4,6-Dimethoxypyrimidin-2-yl	1
32	OH	Phenyl	Phenyl	Methyl	4,6-Diethylpyrimidin-2-yl	1
33	OH	Phenyl	Phenyl	Methyl	6,7-Dihydro-5H-cyclopentapyrimidin-2-yl	1
34	OH	Phenyl	Phenyl	Methyl	Naphth-2-yl	1
35	OH	Phenyl	Phenyl	Methyl	6,7-Dihydro-5H-cyclopenta[c]pyridin-2-yl	1
36	OH	p-Methoxyphenyl	p-Methoxyphenyl	Methyl	Phenyl	1
37	OH	p-Methoxyphenyl	p-Methoxyphenyl	Methyl	3,5-Dimethylphenyl	1

No.	R	R ²	R ³	R ⁵	R ⁴	n
38	OH	p-Methoxyphenyl	p-Methoxyphenyl	Methyl	3,5-Dimethoxyphenyl	1
39	OH	p-Methoxyphenyl	p-Methoxyphenyl	Methyl	3,4-Methylenedioxyphenyl	1
40	OH	p-Methoxyphenyl	p-Methoxyphenyl	Methyl	2,3-Dihydro-benzofuran-5-yl	1
41	OH	p-Methoxyphenyl	p-Methoxyphenyl	Methyl	2,3-Dihydro-1H-inden-5-yl	1
42	OH	p-Methoxyphenyl	p-Methoxyphenyl	Methyl	3-Methoxy-5-methylphenyl	1
43	OH	p-Methoxyphenyl	p-Methoxyphenyl	Methyl	3,4-Ethylenedioxyphenyl	1
44	OH	p-Methoxyphenyl	p-Methoxyphenyl	Methyl	5-Methoxy-3,4-methylenedioxyphenyl	1
45	OH	p-Methoxyphenyl	p-Methoxyphenyl	Methyl	3,5-Diethylphenyl	1
46	OH	p-Methoxyphenyl	p-Methoxyphenyl	Methyl	3,4-Dimethoxyphenyl	1
47	OH	p-Methoxyphenyl	p-Methoxyphenyl	Methyl	3-Methyl-4-benzyloxyphenyl	1
48	OH	p-Methoxyphenyl	p-Methoxyphenyl	Methyl	3-Benzyloxy-5-methoxyphenyl	1
49	OH	p-Methoxyphenyl	p-Methoxyphenyl	Methyl	3-Benzyloxy-4-methoxyphenyl	1
50	OH	p-Methoxyphenyl	p-Methoxyphenyl	Methyl	3-Hydroxy-4-methoxyphenyl	1
51	OH	p-Methoxyphenyl	p-Methoxyphenyl	Methyl	4-Hydroxy-3-methoxyphenyl	1
52	OH	p-Methoxyphenyl	p-Methoxyphenyl	Methyl	3,5-Bis-(trifluoromethyl)-phenyl	1
53	OH	p-Methoxyphenyl	p-Methoxyphenyl	Methyl	3-Methylphenyl	1
54	OH	p-Methoxyphenyl	p-Methoxyphenyl	Methyl	3-Methoxyphenyl	1
55	OH	p-Methoxyphenyl	p-Methoxyphenyl	Methyl	4-Benzyloxyphenyl	1
56	OH	p-Methoxyphenyl	p-Methoxyphenyl	Methyl	4-Phenoxyphenyl	1
57	OH	p-Methoxyphenyl	p-Methoxyphenyl	Methyl	6-Methylpyridin-2-yl	1
58	OH	p-Methoxyphenyl	p-Methoxyphenyl	Methyl	6-Ethylpyridin-2-yl	1
59	OH	p-Methoxyphenyl	p-Methoxyphenyl	Methyl	6-Methoxypyridin-2-yl	1

No.	R	R ²	R ³	R ⁵	R ⁴	n
60	OH	p-Methoxyphenyl	p-Methoxyphenyl	Methyl	4,6-Dimethylpyridin-2-yl	1
61	OH	p-Methoxyphenyl	p-Methoxyphenyl	Methyl	4,6-Diethylpyridin-2-yl	1
62	OH	p-Methoxyphenyl	p-Methoxyphenyl	Methyl	4-Methyl-6-methoxypyridin-2-yl	1
63	OH	p-Methoxyphenyl	p-Methoxyphenyl	Methyl	6,7-Dihydro-5H-cyclopenta[b]pyridin-2-yl	1
64	OH	p-Methoxyphenyl	p-Methoxyphenyl	Methyl	4,6-Dimethylpyrimidin-2-yl	1
65	OH	p-Methoxyphenyl	p-Methoxyphenyl	Methyl	4-Methoxy-6-methylpyrimidin-2-yl	1
66	OH	p-Methoxyphenyl	p-Methoxyphenyl	Methyl	4,6-Dimethoxypyrimidin-2-yl	1
67	OH	p-Methoxyphenyl	p-Methoxyphenyl	Methyl	4,6-Diethylpyrimidin-2-yl	1
68	OH	p-Methoxyphenyl	p-Methoxyphenyl	Methyl	6,7-Dihydro-5H-cyclopentapyrimidin-2-yl	1
69	OH	p-Methoxyphenyl	p-Methoxyphenyl	Methyl	Naphth-2-yl	1
70	OH	p-Methoxyphenyl	p-Methoxyphenyl	Methyl	6,7-Dihydro-5H-cyclopenta[c]pyridin-2-yl	1
71	OH	p-Methylphenyl	p-Methylphenyl	Methyl	Phenyl	1
72	OH	p-Methylphenyl	p-Methylphenyl	Methyl	3,5-Dimethylphenyl	1
73	OH	p-Methylphenyl	p-Methylphenyl	Methyl	3,5-Dimethoxyphenyl	1
74	OH	p-Methylphenyl	p-Methylphenyl	Methyl	3,4-Methylenedioxyphenyl	1
75	OH	p-Methylphenyl	p-Methylphenyl	Methyl	2,3-Dihydro-benzofuran-5-yl	1
76	OH	p-Methylphenyl	p-Methylphenyl	Methyl	2,3-Dihydro-1H-inden-5-yl	1
77	OH	p-Methylphenyl	p-Methylphenyl	Methyl	3-Methoxy-5-methylphenyl	1
78	OH	p-Methylphenyl	p-Methylphenyl	Methyl	3,4-Ethylenedioxyphenyl	1
79	OH	p-Methylphenyl	p-Methylphenyl	Methyl	5-Methoxy-3,4-methylenedioxyphenyl	1
80	OH	p-Methylphenyl	p-Methylphenyl	Methyl	3,5-Diethylphenyl	1
81	OH	p-Methylphenyl	p-Methylphenyl	Methyl	3,4-Dimethoxyphenyl	1

No.	R	R ²	R ³	R ⁵	R ⁴	n
82	OH	p-Methylphenyl	p-Methylphenyl	Methyl	3-Methyl-4-benzyloxyphenyl	1
83	OH	p-Methylphenyl	p-Methylphenyl	Methyl	3-Benzyloxy-5-methoxyphenyl	1
84	OH	p-Methylphenyl	p-Methylphenyl	Methyl	3-Benzyloxy-4-methoxyphenyl	1
85	OH	p-Methylphenyl	p-Methylphenyl	Methyl	3-Hydroxy-4-methoxyphenyl	1
86	OH	p-Methylphenyl	p-Methylphenyl	Methyl	4-Hydroxy-3-methoxyphenyl	1
87	OH	p-Methylphenyl	p-Methylphenyl	Methyl	3,5-Bis-(trifluoromethyl)-phenyl	1
88	OH	p-Methylphenyl	p-Methylphenyl	Methyl	3-Methylphenyl	1
89	OH	p-Methylphenyl	p-Methylphenyl	Methyl	3-Methoxyphenyl	1
90	OH	p-Methylphenyl	p-Methylphenyl	Methyl	4-Benzyloxyphenyl	1
91	OH	p-Methylphenyl	p-Methylphenyl	Methyl	4-Phenoxyphenyl	1
92	OH	p-Methylphenyl	p-Methylphenyl	Methyl	6-Methylpyridin-2-yl	1
93	OH	p-Methylphenyl	p-Methylphenyl	Methyl	6-Ethylpyridin-2-yl	1
94	OH	p-Methylphenyl	p-Methylphenyl	Methyl	6-Methoxypyridin-2-yl	1
95	OH	p-Methylphenyl	p-Methylphenyl	Methyl	4,6-Dimethylpyridin-2-yl	1
96	OH	p-Methylphenyl	p-Methylphenyl	Methyl	4,6-Diethylpyridin-2-yl	1
97	OH	p-Methylphenyl	p-Methylphenyl	Methyl	4-Methyl-6-methoxypyridin-2-yl	1
98	OH	p-Methylphenyl	p-Methylphenyl	Methyl	6,7-Dihydro-5H-cyclopenta[b]pyridin-2-yl	1
99	OH	p-Methylphenyl	p-Methylphenyl	Methyl	4,6-Dimethylpyrimidin-2-yl	1
100	OH	p-Methylphenyl	p-Methylphenyl	Methyl	4-Methoxy-6-methylpyrimidin-2-yl	1
101	OH	p-Methylphenyl	p-Methylphenyl	Methyl	4,6-Dimethoxypyrimidin-2-yl	1
102	OH	p-Methylphenyl	p-Methylphenyl	Methyl	4,6-Diethylpyrimidin-2-yl	1
103	OH	p-Methylphenyl	p-Methylphenyl	Methyl	6,7-Dihydro-5H-cyclopentapyrimidin-2-yl	1

No.	R	R ²	R ³	R ⁵	R ⁴	n
104	OH	p-Methylphenyl	p-Methylphenyl	Methyl	Naphth-2-yl	1
105	OH	p-Methylphenyl	p-Methylphenyl	Methyl	6,7-Dihydro-5H-cyclopenta[c]pyridin-2-yl	1
106	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Methyl	Phenyl	1
107	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Methyl	3,5-Dimethylphenyl	1
108	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Methyl	3,5-Dimethoxyphenyl	1
109	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Methyl	3,4-Methylenedioxyphenyl	1
110	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Methyl	2,3-Dihydro-benzofuran-5-yl	1
111	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Methyl	2,3-Dihydro-1H-inden-5-yl	1
112	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Methyl	3-Methoxy-5-methylphenyl	1
113	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Methyl	3,4-Ethylenedioxyphenyl	1
114	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Methyl	5-Methoxy-3,4-methylenedioxyphenyl	1
115	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Methyl	3,5-Diethylphenyl	1
116	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Methyl	3,4-Dimethoxyphenyl	1
117	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Methyl	3-Methyl-4-benzyloxyphenyl	1
118	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Methyl	3-Benzyloxy-5-methoxyphenyl	1
119	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Methyl	3-Benzyloxy-4-methoxyphenyl	1
120	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Methyl	3-Hydroxy-4-methoxyphenyl	1
121	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Methyl	4-Hydroxy-3-methoxyphenyl	1
122	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Methyl	3,5-Bis-(trifluoromethyl)-phenyl	1
123	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Methyl	3-Methylphenyl	1
124	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Methyl	3-Methoxyphenyl	1
125	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Methyl	4-Benzyloxyphenyl	1

No.	R	R ²	R ³	R ⁵	R ⁴	n
126	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Methyl	4-Phenoxyphenyl	1
127	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Methyl	6-Methylpyridin-2-yl	1
128	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Methyl	6-Ethylpyridin-2-yl	1
129	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Methyl	6-Methoxypyridin-2-yl	1
130	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Methyl	4,6-Dimethylpyridin-2-yl	1
131	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Methyl	4,6-Diethylpyridin-2-yl	1
132	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Methyl	4-Methyl-6-methoxypyridin-2-yl	1
133	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Methyl	6,7-Dihydro-5H-cyclopenta[b]pyridin-2-yl	1
134	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Methyl	4,6-Dimethylpyrimidin-2-yl	1
135	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Methyl	4-Methoxy-6-methylpyrimidin-2-yl	1
136	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Methyl	4,6-Dimethoxypyrimidin-2-yl	1
137	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Methyl	4,6-Diethylpyrimidin-2-yl	1
138	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Methyl	6,7-Dihydro-5H-cyclopentapyrimidin-2-yl	1
139	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Methyl	Naphth-2-yl	1
140	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Methyl	6,7-Dihydro-5H-cyclopenta[c]pyridin-2-yl	1
141	OH	Phenyl	Phenyl	Ethyl	Phenyl	1
142	OH	Phenyl	Phenyl	Ethyl	3,5-Dimethylphenyl	1
143	OH	Phenyl	Phenyl	Ethyl	3,5-Dimethoxyphenyl	1
144	OH	Phenyl	Phenyl	Ethyl	3,4-Methylenedioxyphenyl	1
145	OH	Phenyl	Phenyl	Ethyl	2,3-Dihydro-benzofuran-5-yl	1
146	OH	Phenyl	Phenyl	Ethyl	2,3-Dihydro-1H-inden-5-yl	1
147	OH	Phenyl	Phenyl	Ethyl	3-Methoxy-5-methylphenyl	1

No.	R	R ²	R ³	R ⁵	R ⁴	n
148	OH	Phenyl	Phenyl	Ethyl	3,4-Ethylenedioxyphenyl	1
149	OH	Phenyl	Phenyl	Ethyl	5-Methoxy-3,4-methylenedioxyphenyl	1
150	OH	Phenyl	Phenyl	Ethyl	3,5-Diethylphenyl	1
151	OH	Phenyl	Phenyl	Ethyl	3,4-Dimethoxyphenyl	1
152	OH	Phenyl	Phenyl	Ethyl	3-Methyl-4-benzyloxyphenyl	1
153	OH	Phenyl	Phenyl	Ethyl	3-Benzyloxy-5-methoxyphenyl	1
154	OH	Phenyl	Phenyl	Ethyl	3-Benzyloxy-4-methoxyphenyl	1
155	OH	Phenyl	Phenyl	Ethyl	3-Hydroxy-4-methoxyphenyl	1
156	OH	Phenyl	Phenyl	Ethyl	4-Hydroxy-3-methoxyphenyl	1
157	OH	Phenyl	Phenyl	Ethyl	3,5-Bis-(trifluoromethyl)-phenyl	1
158	OH	Phenyl	Phenyl	Ethyl	3-Methylphenyl	1
159	OH	Phenyl	Phenyl	Ethyl	3-Methoxyphenyl	1
160	OH	Phenyl	Phenyl	Ethyl	4-Benzyloxyphenyl	1
161	OH	Phenyl	Phenyl	Ethyl	4-Phenoxyphenyl	1
162	OH	Phenyl	Phenyl	Ethyl	6-Methylpyridin-2-yl	1
163	OH	Phenyl	Phenyl	Ethyl	6-Ethylpyridin-2-yl	1
164	OH	Phenyl	Phenyl	Ethyl	6-Methoxypyridin-2-yl	1
165	OH	Phenyl	Phenyl	Ethyl	4,6-Dimethylpyridin-2-yl	1
166	OH	Phenyl	Phenyl	Ethyl	4,6-Diethylpyridin-2-yl	1
167	OH	Phenyl	Phenyl	Ethyl	4-Methyl-6-methoxypyridin-2-yl	1
168	OH	Phenyl	Phenyl	Ethyl	6,7-Dihydro-5H-cyclopenta[b]pyridin-2-yl	1
169	OH	Phenyl	Phenyl	Ethyl	4,6-Dimethylpyrimidin-2-yl	1

No.	R	R ²	R ³	R ⁵	R ⁴	n
170	OH	Phenyl	Phenyl	Ethyl	4-Methoxy-6-methylpyrimidin-2-yl	1
171	OH	Phenyl	Phenyl	Ethyl	4,6-Dimethoxypyrimidin-2-yl	1
172	OH	Phenyl	Phenyl	Ethyl	4,6-Diethylpyrimidin-2-yl	1
173	OH	Phenyl	Phenyl	Ethyl	6,7-Dihydro-5H-cyclopentapyrimidin-2-yl	1
174	OH	Phenyl	Phenyl	Ethyl	Naphth-2-yl	1
175	OH	Phenyl	Phenyl	Ethyl	6,7-Dihydro-5H-cyclopenta[c]pyridin-2-yl	1
176	OH	p-Methoxyphenyl	p-Methoxyphenyl	Ethyl	Phenyl	1
177	OH	p-Methoxyphenyl	p-Methoxyphenyl	Ethyl	3,5-Dimethylphenyl	1
178	OH	p-Methoxyphenyl	p-Methoxyphenyl	Ethyl	3,5-Dimethoxyphenyl	1
179	OH	p-Methoxyphenyl	p-Methoxyphenyl	Ethyl	3,4-Methylenedioxyphenyl	1
180	OH	p-Methoxyphenyl	p-Methoxyphenyl	Ethyl	2,3-Dihydro-benzofuran-5-yl	1
181	OH	p-Methoxyphenyl	p-Methoxyphenyl	Ethyl	2,3-Dihydro-1H-inden-5-yl	1
182	OH	p-Methoxyphenyl	p-Methoxyphenyl	Ethyl	3-Methoxy-5-methylphenyl	1
183	OH	p-Methoxyphenyl	p-Methoxyphenyl	Ethyl	3,4-Ethylenedioxyphenyl	1
184	OH	p-Methoxyphenyl	p-Methoxyphenyl	Ethyl	5-Methoxy-3,4-methylenedioxyphenyl	1
185	OH	p-Methoxyphenyl	p-Methoxyphenyl	Ethyl	3,5-Diethylphenyl	1
186	OH	p-Methoxyphenyl	p-Methoxyphenyl	Ethyl	3,4-Dimethoxyphenyl	1
187	OH	p-Methoxyphenyl	p-Methoxyphenyl	Ethyl	3-Methyl-4-benzoyloxyphenyl	1
188	OH	p-Methoxyphenyl	p-Methoxyphenyl	Ethyl	3-Benzoyloxy-5-methoxyphenyl	1
189	OH	p-Methoxyphenyl	p-Methoxyphenyl	Ethyl	3-Benzoyloxy-4-methoxyphenyl	1
190	OH	p-Methoxyphenyl	p-Methoxyphenyl	Ethyl	3-Hydroxy-4-methoxyphenyl	1
191	OH	p-Methoxyphenyl	p-Methoxyphenyl	Ethyl	4-Hydroxy-3-methoxyphenyl	1

No.	R	R ²	R ³	R ⁵	R ⁴	n
192	OH	p-Methoxyphenyl	p-Methoxyphenyl	Ethyl	3,5-Bis-(trifluoromethyl)-phenyl	1
193	OH	p-Methoxyphenyl	p-Methoxyphenyl	Ethyl	3-Methylphenyl	1
194	OH	p-Methoxyphenyl	p-Methoxyphenyl	Ethyl	3-Methoxyphenyl	1
195	OH	p-Methoxyphenyl	p-Methoxyphenyl	Ethyl	4-Benzyloxyphenyl	1
196	OH	p-Methoxyphenyl	p-Methoxyphenyl	Ethyl	4-Phenoxyphenyl	1
197	OH	p-Methoxyphenyl	p-Methoxyphenyl	Ethyl	6-Methylpyridin-2-yl	1
198	OH	p-Methoxyphenyl	p-Methoxyphenyl	Ethyl	6-Ethylpyridin-2-yl	1
199	OH	p-Methoxyphenyl	p-Methoxyphenyl	Ethyl	6-Methoxypyridin-2-yl	1
200	OH	p-Methoxyphenyl	p-Methoxyphenyl	Ethyl	4,6-Dimethylpyridin-2-yl	1
201	OH	p-Methoxyphenyl	p-Methoxyphenyl	Ethyl	4,6-Diethylpyridin-2-yl	1
202	OH	p-Methoxyphenyl	p-Methoxyphenyl	Ethyl	4-Methyl-6-methoxypyridin-2-yl	1
203	OH	p-Methoxyphenyl	p-Methoxyphenyl	Ethyl	6,7-Dihydro-5H-cyclopenta[b]pyridin-2-yl	1
204	OH	p-Methoxyphenyl	p-Methoxyphenyl	Ethyl	4,6-Dimethylpyrimidin-2-yl	1
205	OH	p-Methoxyphenyl	p-Methoxyphenyl	Ethyl	4-Methoxy-6-methylpyrimidin-2-yl	1
206	OH	p-Methoxyphenyl	p-Methoxyphenyl	Ethyl	4,6-Dimethoxypyrimidin-2-yl	1
207	OH	p-Methoxyphenyl	p-Methoxyphenyl	Ethyl	4,6-Diethylpyrimidin-2-yl	1
208	OH	p-Methoxyphenyl	p-Methoxyphenyl	Ethyl	6,7-Dihydro-5H-cyclopentapyrimidin-2-yl	1
209	OH	p-Methoxyphenyl	p-Methoxyphenyl	Ethyl	Naphth-2-yl	1
210	OH	p-Methoxyphenyl	p-Methoxyphenyl	Ethyl	6,7-Dihydro-5H-cyclopenta[c]pyridin-2-yl	1
211	OH	p-Methylphenyl	p-Methylphenyl	Ethyl	Phenyl	1
212	OH	p-Methylphenyl	p-Methylphenyl	Ethyl	3,5-Dimethylphenyl	1
213	OH	p-Methylphenyl	p-Methylphenyl	Ethyl	3,5-Dimethoxyphenyl	1

No.	R	R ²	R ³	R ⁵	R ⁴	n
214	OH	p-Methylphenyl	p-Methylphenyl	Ethyl	3,4-Methylenedioxyphenyl	1
215	OH	p-Methylphenyl	p-Methylphenyl	Ethyl	2,3-Dihydro-benzofuran-5-yl	1
216	OH	p-Methylphenyl	p-Methylphenyl	Ethyl	2,3-Dihydro-1H-inden-5-yl	1
217	OH	p-Methylphenyl	p-Methylphenyl	Ethyl	3-Methoxy-5-methylphenyl	1
218	OH	p-Methylphenyl	p-Methylphenyl	Ethyl	3,4-Ethylenedioxyphenyl	1
219	OH	p-Methylphenyl	p-Methylphenyl	Ethyl	5-Methoxy-3,4-methylenedioxyphenyl	1
220	OH	p-Methylphenyl	p-Methylphenyl	Ethyl	3,5-Diethylphenyl	1
221	OH	p-Methylphenyl	p-Methylphenyl	Ethyl	3,4-Dimethoxyphenyl	1
222	OH	p-Methylphenyl	p-Methylphenyl	Ethyl	3-Methyl-4-benzyloxyphenyl	1
223	OH	p-Methylphenyl	p-Methylphenyl	Ethyl	3-Benzyloxy-5-methoxyphenyl	1
224	OH	p-Methylphenyl	p-Methylphenyl	Ethyl	3-Benzyloxy-4-methoxyphenyl	1
225	OH	p-Methylphenyl	p-Methylphenyl	Ethyl	3-Hydroxy-4-methoxyphenyl	1
226	OH	p-Methylphenyl	p-Methylphenyl	Ethyl	4-Hydroxy-3-methoxyphenyl	1
227	OH	p-Methylphenyl	p-Methylphenyl	Ethyl	3,5-Bis-(trifluoromethyl)-phenyl	1
228	OH	p-Methylphenyl	p-Methylphenyl	Ethyl	3-Methylphenyl	1
229	OH	p-Methylphenyl	p-Methylphenyl	Ethyl	3-Methoxyphenyl	1
230	OH	p-Methylphenyl	p-Methylphenyl	Ethyl	4-Benzyloxyphenyl	1
231	OH	p-Methylphenyl	p-Methylphenyl	Ethyl	4-Phenoxyphenyl	1
232	OH	p-Methylphenyl	p-Methylphenyl	Ethyl	6-Methylpyridin-2-yl	1
233	OH	p-Methylphenyl	p-Methylphenyl	Ethyl	6-Ethylpyridin-2-yl	1
234	OH	p-Methylphenyl	p-Methylphenyl	Ethyl	6-Methoxypyridin-2-yl	1
235	OH	p-Methylphenyl	p-Methylphenyl	Ethyl	4,6-Dimethylpyridin-2-yl	1

No.	R	R ²	R ³	R ⁵	R ⁴	n
236	OH	p-Methylphenyl	p-Methylphenyl	Ethyl	4,6-Diethylpyridin-2-yl	1
237	OH	p-Methylphenyl	p-Methylphenyl	Ethyl	4-Methyl-6-methoxypyridin-2-yl	1
238	OH	p-Methylphenyl	p-Methylphenyl	Ethyl	6,7-Dihydro-5H-cyclopenta[b]pyridin-2-yl	1
239	OH	p-Methylphenyl	p-Methylphenyl	Ethyl	4,6-Dimethylpyrimidin-2-yl	1
240	OH	p-Methylphenyl	p-Methylphenyl	Ethyl	4-Methoxy-6-methylpyrimidin-2-yl	1
241	OH	p-Methylphenyl	p-Methylphenyl	Ethyl	4,6-Dimethoxypyrimidin-2-yl	1
242	OH	p-Methylphenyl	p-Methylphenyl	Ethyl	4,6-Diethylpyrimidin-2-yl	1
243	OH	p-Methylphenyl	p-Methylphenyl	Ethyl	6,7-Dihydro-5H-cyclopentapyrimidin-2-yl	1
244	OH	p-Methylphenyl	p-Methylphenyl	Ethyl	Naphth-2-yl	1
245	OH	p-Methylphenyl	p-Methylphenyl	Ethyl	6,7-Dihydro-5H-cyclopenta[c]pyridin-2-yl	1
246	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Ethyl	Phenyl	1
247	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Ethyl	3,5-Dimethylphenyl	1
248	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Ethyl	3,5-Dimethoxyphenyl	1
249	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Ethyl	3,4-Methylenedioxyphenyl	1
250	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Ethyl	2,3-Dihydro-benzofuran-5-yl	1
251	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Ethyl	2,3-Dihydro-1H-inden-5-yl	1
252	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Ethyl	3-Methoxy-5-methylphenyl	1
253	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Ethyl	3,4-Ethylenedioxyphenyl	1
254	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Ethyl	5-Methoxy-3,4-methylenedioxyphenyl	1
255	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Ethyl	3,5-Diethylphenyl	1
256	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Ethyl	3,4-Dimethoxyphenyl	1
257	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Ethyl	3-Methyl-4-benzyloxyphenyl	1

No.	R	R ²	R ³	R ⁵	R ⁴	n
258	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Ethyl	3-Benzoyloxy-5-methoxyphenyl	1
259	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Ethyl	3-Benzoyloxy-4-methoxyphenyl	1
260	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Ethyl	3-Hydroxy-4-methoxyphenyl	1
261	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Ethyl	4-Hydroxy-3-methoxyphenyl	1
262	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Ethyl	3,5-Bis-(trifluoromethyl)-phenyl	1
263	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Ethyl	3-Methylphenyl	1
264	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Ethyl	3-Methoxyphenyl	1
265	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Ethyl	4-Benzoyloxyphenyl	1
266	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Ethyl	4-Phenoxyphenyl	1
267	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Ethyl	6-Methylpyridin-2-yl	1
268	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Ethyl	6-Ethylpyridin-2-yl	1
269	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Ethyl	6-Methoxypyridin-2-yl	1
270	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Ethyl	4,6-Dimethylpyridin-2-yl	1
271	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Ethyl	4,6-Diethylpyridin-2-yl	1
272	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Ethyl	4-Methyl-6-methoxypyridin-2-yl	1
273	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Ethyl	6,7-Dihydro-5H-cyclopenta[b]pyridin-2-yl	1
274	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Ethyl	4,6-Dimethylpyrimidin-2-yl	1
275	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Ethyl	4-Methoxy-6-methylpyrimidin-2-yl	1
276	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Ethyl	4,6-Dimethoxypyrimidin-2-yl	1
277	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Ethyl	4,6-Diethylpyrimidin-2-yl	1
278	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Ethyl	6,7-Dihydro-5H-cyclopentapyrimidin-2-yl	1
279	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Ethyl	Naphth-2-yl	1

No.	R	R ²	R ³	R ⁵	R ⁴	n
280	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Ethyl	6,7-Dihydro-5H-cyclopenta[c]pyridin-2-yl	1
281	OH	Phenyl	Phenyl	Propyl	Phenyl	1
282	OH	Phenyl	Phenyl	Propyl	3,5-Dimethylphenyl	1
283	OH	Phenyl	Phenyl	Propyl	3,5-Dimethoxyphenyl	1
284	OH	Phenyl	Phenyl	Propyl	3,4-Methylenedioxyphenyl	1
285	OH	Phenyl	Phenyl	Propyl	2,3-Dihydro-benzofuran-5-yl	1
286	OH	Phenyl	Phenyl	Propyl	2,3-Dihydro-1H-inden-5-yl	1
287	OH	Phenyl	Phenyl	Propyl	3-Methoxy-5-methylphenyl	1
288	OH	Phenyl	Phenyl	Propyl	3,4-Ethylenedioxyphenyl	1
289	OH	Phenyl	Phenyl	Propyl	5-Methoxy-3,4-methylenedioxyphenyl	1
290	OH	Phenyl	Phenyl	Propyl	3,5-Diethylphenyl	1
291	OH	Phenyl	Phenyl	Propyl	3,4-Dimethoxyphenyl	1
292	OH	Phenyl	Phenyl	Propyl	3-Methyl-4-benzyloxyphenyl	1
293	OH	Phenyl	Phenyl	Propyl	3-Benzyloxy-5-methoxyphenyl	1
294	OH	Phenyl	Phenyl	Propyl	3-Benzyloxy-4-methoxyphenyl	1
295	OH	Phenyl	Phenyl	Propyl	3-Hydroxy-4-methoxyphenyl	1
296	OH	Phenyl	Phenyl	Propyl	4-Hydroxy-3-methoxyphenyl	1
297	OH	Phenyl	Phenyl	Propyl	3,5-Bis-(trifluoromethyl)-phenyl	1
298	OH	Phenyl	Phenyl	Propyl	3-Methylphenyl	1
299	OH	Phenyl	Phenyl	Propyl	3-Methoxyphenyl	1
300	OH	Phenyl	Phenyl	Propyl	4-Benzyloxyphenyl	1
301	OH	Phenyl	Phenyl	Propyl	4-Phenoxyphenyl	1

No.	R	R ²	R ³	R ⁵	R ⁴	n
302	OH	Phenyl	Phenyl	Propyl	6-Methylpyridin-2-yl	1
303	OH	Phenyl	Phenyl	Propyl	6-Ethylpyridin-2-yl	1
304	OH	Phenyl	Phenyl	Propyl	6-Methoxypyridin-2-yl	1
305	OH	Phenyl	Phenyl	Propyl	4,6-Dimethylpyridin-2-yl	1
306	OH	Phenyl	Phenyl	Propyl	4,6-Diethylpyridin-2-yl	1
307	OH	Phenyl	Phenyl	Propyl	4-Methyl-6-methoxypyridin-2-yl	1
308	OH	Phenyl	Phenyl	Propyl	6,7-Dihydro-5H-cyclopenta[b]pyridin-2-yl	1
309	OH	Phenyl	Phenyl	Propyl	4,6-Dimethylpyrimidin-2-yl	1
310	OH	Phenyl	Phenyl	Propyl	4-Methoxy-6-methylpyrimidin-2-yl	1
311	OH	Phenyl	Phenyl	Propyl	4,6-Dimethoxypyrimidin-2-yl	1
312	OH	Phenyl	Phenyl	Propyl	4,6-Diethylpyrimidin-2-yl	1
313	OH	Phenyl	Phenyl	Propyl	6,7-Dihydro-5H-cyclopentapyrimidin-2-yl	1
314	OH	Phenyl	Phenyl	Propyl	Naphth-2-yl	1
315	OH	Phenyl	Phenyl	Propyl	6,7-Dihydro-5H-cyclopenta[c]pyridin-2-yl	1
316	OH	p-Methoxyphenyl	p-Methoxyphenyl	Propyl	Phenyl	1
317	OH	p-Methoxyphenyl	p-Methoxyphenyl	Propyl	3,5-Dimethylphenyl	1
318	OH	p-Methoxyphenyl	p-Methoxyphenyl	Propyl	3,5-Dimethoxyphenyl	1
319	OH	p-Methoxyphenyl	p-Methoxyphenyl	Propyl	3,4-Methylenedioxyphenyl	1
320	OH	p-Methoxyphenyl	p-Methoxyphenyl	Propyl	2,3-Dihydro-benzofuran-5-yl	1
321	OH	p-Methoxyphenyl	p-Methoxyphenyl	Propyl	2,3-Dihydro-1H-inden-5-yl	1
322	OH	p-Methoxyphenyl	p-Methoxyphenyl	Propyl	3-Methoxy-5-methylphenyl	1
323	OH	p-Methoxyphenyl	p-Methoxyphenyl	Propyl	3,4-Ethylenedioxyphenyl	1

No.	R	R ²	R ³	R ⁵	R ⁴	n
324	OH	p-Methoxyphenyl	p-Methoxyphenyl	Propyl	5-Methoxy-3,4-methylenedioxyphenyl	1
325	OH	p-Methoxyphenyl	p-Methoxyphenyl	Propyl	3,5-Diethylphenyl	1
326	OH	p-Methoxyphenyl	p-Methoxyphenyl	Propyl	3,4-Dimethoxyphenyl	1
327	OH	p-Methoxyphenyl	p-Methoxyphenyl	Propyl	3-Methyl-4-benzyloxyphenyl	1
328	OH	p-Methoxyphenyl	p-Methoxyphenyl	Propyl	3-Benzyloxy-5-methoxyphenyl	1
329	OH	p-Methoxyphenyl	p-Methoxyphenyl	Propyl	3-Benzyloxy-4-methoxyphenyl	1
330	OH	p-Methoxyphenyl	p-Methoxyphenyl	Propyl	3-Hydroxy-4-methoxyphenyl	1
331	OH	p-Methoxyphenyl	p-Methoxyphenyl	Propyl	4-Hydroxy-3-methoxyphenyl	1
332	OH	p-Methoxyphenyl	p-Methoxyphenyl	Propyl	3,5-Bis-(trifluoromethyl)-phenyl	1
333	OH	p-Methoxyphenyl	p-Methoxyphenyl	Propyl	3-Methylphenyl	1
334	OH	p-Methoxyphenyl	p-Methoxyphenyl	Propyl	3-Methoxyphenyl	1
335	OH	p-Methoxyphenyl	p-Methoxyphenyl	Propyl	4-Benzyloxyphenyl	1
336	OH	p-Methoxyphenyl	p-Methoxyphenyl	Propyl	4-Phenoxyphenyl	1
337	OH	p-Methoxyphenyl	p-Methoxyphenyl	Propyl	6-Methylpyridin-2-yl	1
338	OH	p-Methoxyphenyl	p-Methoxyphenyl	Propyl	6-Ethylpyridin-2-yl	1
339	OH	p-Methoxyphenyl	p-Methoxyphenyl	Propyl	6-Methoxypyridin-2-yl	1
340	OH	p-Methoxyphenyl	p-Methoxyphenyl	Propyl	4,6-Dimethylpyridin-2-yl	1
341	OH	p-Methoxyphenyl	p-Methoxyphenyl	Propyl	4,6-Diethylpyridin-2-yl	1
342	OH	p-Methoxyphenyl	p-Methoxyphenyl	Propyl	4-Methyl-6-methoxypyridin-2-yl	1
343	OH	p-Methoxyphenyl	p-Methoxyphenyl	Propyl	6,7-Dihydro-5H-cyclopenta[b]pyridin-2-yl	1
344	OH	p-Methoxyphenyl	p-Methoxyphenyl	Propyl	4,6-Dimethylpyrimidin-2-yl	1
345	OH	p-Methoxyphenyl	p-Methoxyphenyl	Propyl	4-Methoxy-6-methylpyrimidin-2-yl	1

No.	R	R ²	R ³	R ⁵	R ⁴	n
346	OH	p-Methoxyphenyl	p-Methoxyphenyl	Propyl	4,6-Dimethoxypyrimidin-2-yl	1
347	OH	p-Methoxyphenyl	p-Methoxyphenyl	Propyl	4,6-Diethylpyrimidin-2-yl	1
348	OH	p-Methoxyphenyl	p-Methoxyphenyl	Propyl	6,7-Dihydro-5H-cyclopentapyrimidin-2-yl	1
349	OH	p-Methoxyphenyl	p-Methoxyphenyl	Propyl	Naphth-2-yl	1
350	OH	p-Methoxyphenyl	p-Methoxyphenyl	Propyl	6,7-Dihydro-5H-cyclopenta[c]pyridin-2-yl	1
351	OH	p-Methylphenyl	p-Methylphenyl	Propyl	Phenyl	1
352	OH	p-Methylphenyl	p-Methylphenyl	Propyl	3,5-Dimethylphenyl	1
353	OH	p-Methylphenyl	p-Methylphenyl	Propyl	3,5-Dimethoxyphenyl	1
354	OH	p-Methylphenyl	p-Methylphenyl	Propyl	3,4-Methylenedioxyphenyl	1
355	OH	p-Methylphenyl	p-Methylphenyl	Propyl	2,3-Dihydro-benzofuran-5-yl	1
356	OH	p-Methylphenyl	p-Methylphenyl	Propyl	2,3-Dihydro-1H-inden-5-yl	1
357	OH	p-Methylphenyl	p-Methylphenyl	Propyl	3-Methoxy-5-methylphenyl	1
358	OH	p-Methylphenyl	p-Methylphenyl	Propyl	3,4-Ethylenedioxyphenyl	1
359	OH	p-Methylphenyl	p-Methylphenyl	Propyl	5-Methoxy-3,4-methylenedioxyphenyl	1
360	OH	p-Methylphenyl	p-Methylphenyl	Propyl	3,5-Diethylphenyl	1
361	OH	p-Methylphenyl	p-Methylphenyl	Propyl	3,4-Dimethoxyphenyl	1
362	OH	p-Methylphenyl	p-Methylphenyl	Propyl	3-Methyl-4-benzyloxyphenyl	1
363	OH	p-Methylphenyl	p-Methylphenyl	Propyl	3-Benzyloxy-5-methoxyphenyl	1
364	OH	p-Methylphenyl	p-Methylphenyl	Propyl	3-Benzyloxy-4-methoxyphenyl	1
365	OH	p-Methylphenyl	p-Methylphenyl	Propyl	3-Hydroxy-4-methoxyphenyl	1
366	OH	p-Methylphenyl	p-Methylphenyl	Propyl	4-Hydroxy-3-methoxyphenyl	1
367	OH	p-Methylphenyl	p-Methylphenyl	Propyl	3,5-Bis-(trifluoromethyl)-phenyl	1

No.	R	R ²	R ³	R ⁵	R ⁴	n
368	OH	p-Methylphenyl	p-Methylphenyl	Propyl	3-Methylphenyl	1
369	OH	p-Methylphenyl	p-Methylphenyl	Propyl	3-Methoxyphenyl	1
370	OH	p-Methylphenyl	p-Methylphenyl	Propyl	4-Benzoyloxyphenyl	1
371	OH	p-Methylphenyl	p-Methylphenyl	Propyl	4-Phenoxyphenyl	1
372	OH	p-Methylphenyl	p-Methylphenyl	Propyl	6-Methylpyridin-2-yl	1
373	OH	p-Methylphenyl	p-Methylphenyl	Propyl	6-Ethylpyridin-2-yl	1
374	OH	p-Methylphenyl	p-Methylphenyl	Propyl	6-Methoxypyridin-2-yl	1
375	OH	p-Methylphenyl	p-Methylphenyl	Propyl	4,6-Dimethylpyridin-2-yl	1
376	OH	p-Methylphenyl	p-Methylphenyl	Propyl	4,6-Diethylpyridin-2-yl	1
377	OH	p-Methylphenyl	p-Methylphenyl	Propyl	4-Methyl-6-methoxypyridin-2-yl	1
378	OH	p-Methylphenyl	p-Methylphenyl	Propyl	6,7-Dihydro-5H-cyclopenta[b]pyridin-2-yl	1
379	OH	p-Methylphenyl	p-Methylphenyl	Propyl	4,6-Dimethylpyrimidin-2-yl	1
380	OH	p-Methylphenyl	p-Methylphenyl	Propyl	4-Methoxy-6-methylpyrimidin-2-yl	1
381	OH	p-Methylphenyl	p-Methylphenyl	Propyl	4,6-Dimethoxypyrimidin-2-yl	1
382	OH	p-Methylphenyl	p-Methylphenyl	Propyl	4,6-Diethylpyrimidin-2-yl	1
383	OH	p-Methylphenyl	p-Methylphenyl	Propyl	6,7-Dihydro-5H-cyclopentapyrimidin-2-yl	1
384	OH	p-Methylphenyl	p-Methylphenyl	Propyl	Naphth-2-yl	1
385	OH	p-Methylphenyl	p-Methylphenyl	Propyl	6,7-Dihydro-5H-cyclopenta[c]pyridin-2-yl	1
386	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Propyl	Phenyl	1
387	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Propyl	3,5-Dimethylphenyl	1
388	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Propyl	3,5-Dimethoxyphenyl	1
389	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Propyl	3,4-Methylenedioxyphenyl	1

No.	R	R ²	R ³	R ⁵	R ⁴	n
390	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Propyl	2,3-Dihydro-benzofuran-5-yl	1
391	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Propyl	2,3-Dihydro-1H-inden-5-yl	1
392	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Propyl	3-Methoxy-5-methylphenyl	1
393	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Propyl	3,4-Ethylenedioxyphenyl	1
394	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Propyl	5-Methoxy-3,4-methylenedioxyphenyl	1
395	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Propyl	3,5-Diethylphenyl	1
396	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Propyl	3,4-Dimethoxyphenyl	1
397	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Propyl	3-Methyl-4-benzyloxyphenyl	1
398	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Propyl	3-Benzyloxy-5-methoxyphenyl	1
399	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Propyl	3-Benzyloxy-4-methoxyphenyl	1
400	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Propyl	3-Hydroxy-4-methoxyphenyl	1
401	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Propyl	4-Hydroxy-3-methoxyphenyl	1
402	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Propyl	3,5-Bis-(trifluoromethyl)-phenyl	1
403	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Propyl	3-Methylphenyl	1
404	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Propyl	3-Methoxyphenyl	1
405	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Propyl	4-Benzyloxyphenyl	1
406	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Propyl	4-Phenoxyphenyl	1
407	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Propyl	6-Methylpyridin-2-yl	1
408	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Propyl	6-Ethylpyridin-2-yl	1
409	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Propyl	6-Methoxypyridin-2-yl	1
410	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Propyl	4,6-Dimethylpyridin-2-yl	1
411	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Propyl	4,6-Diethylpyridin-2-yl	1

No.	R	R ²	R ³	R ⁵	R ⁴	n
412	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Propyl	4-Methyl-6-methoxypyridin-2-yl	1
413	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Propyl	6,7-Dihydro-5H-cyclopenta[b]pyridin-2-yl	1
414	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Propyl	4,6-Dimethylpyrimidin-2-yl	1
415	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Propyl	4-Methoxy-6-methylpyrimidin-2-yl	1
416	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Propyl	4,6-Dimethoxypyrimidin-2-yl	1
417	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Propyl	4,6-Diethylpyrimidin-2-yl	1
418	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Propyl	6,7-Dihydro-5H-cyclopentapyrimidin-2-yl	1
419	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Propyl	Naphth-2-yl	1
420	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	Propyl	6,7-Dihydro-5H-cyclopenta[c]pyridin-2-yl	1
421	OH	Phenyl	Phenyl	2-Hydroxyethyl	Phenyl	1
422	OH	Phenyl	Phenyl	2-Hydroxyethyl	3,5-Dimethylphenyl	1
423	OH	Phenyl	Phenyl	2-Hydroxyethyl	3,5-Dimethoxyphenyl	1
424	OH	Phenyl	Phenyl	2-Hydroxyethyl	3,4-Methylenedioxyphenyl	1
425	OH	Phenyl	Phenyl	2-Hydroxyethyl	2,3-Dihydro-benzofuran-5-yl	1
426	OH	Phenyl	Phenyl	2-Hydroxyethyl	2,3-Dihydro-1H-inden-5-yl	1
427	OH	Phenyl	Phenyl	2-Hydroxyethyl	3-Methoxy-5-methylphenyl	1
428	OH	Phenyl	Phenyl	2-Hydroxyethyl	3,4-Ethylenedioxyphenyl	1
429	OH	Phenyl	Phenyl	2-Hydroxyethyl	5-Methoxy-3,4-methylenedioxyphenyl	1
430	OH	Phenyl	Phenyl	2-Hydroxyethyl	3,5-Diethylphenyl	1
431	OH	Phenyl	Phenyl	2-Hydroxyethyl	3,4-Dimethoxyphenyl	1
432	OH	Phenyl	Phenyl	2-Hydroxyethyl	3-Methyl-4-benzoyloxyphenyl	1
433	OH	Phenyl	Phenyl	2-Hydroxyethyl	3-Benzoyloxy-5-methoxyphenyl	1

No.	R	R ²	R ³	R ⁵	R ⁴	n
434	OH	Phenyl	Phenyl	2-Hydroxyethyl	3-Benzyloxy-4-methoxyphenyl	1
435	OH	Phenyl	Phenyl	2-Hydroxyethyl	3-Hydroxy-4-methoxyphenyl	1
436	OH	Phenyl	Phenyl	2-Hydroxyethyl	4-Hydroxy-3-methoxyphenyl	1
437	OH	Phenyl	Phenyl	2-Hydroxyethyl	3,5-Bis-(trifluoromethyl)-phenyl	1
438	OH	Phenyl	Phenyl	2-Hydroxyethyl	3-Methylphenyl	1
439	OH	Phenyl	Phenyl	2-Hydroxyethyl	3-Methoxyphenyl	1
440	OH	Phenyl	Phenyl	2-Hydroxyethyl	4-Benzyloxyphenyl	1
441	OH	Phenyl	Phenyl	2-Hydroxyethyl	4-Phenoxyphenyl	1
442	OH	Phenyl	Phenyl	2-Hydroxyethyl	6-Methylpyridin-2-yl	1
443	OH	Phenyl	Phenyl	2-Hydroxyethyl	6-Ethylpyridin-2-yl	1
444	OH	Phenyl	Phenyl	2-Hydroxyethyl	6-Methoxypyridin-2-yl	1
445	OH	Phenyl	Phenyl	2-Hydroxyethyl	4,6-Dimethylpyridin-2-yl	1
446	OH	Phenyl	Phenyl	2-Hydroxyethyl	4,6-Diethylpyridin-2-yl	1
447	OH	Phenyl	Phenyl	2-Hydroxyethyl	4-Methyl-6-methoxypyridin-2-yl	1
448	OH	Phenyl	Phenyl	2-Hydroxyethyl	6,7-Dihydro-5H-cyclopenta[b]pyridin-2-yl	1
449	OH	Phenyl	Phenyl	2-Hydroxyethyl	4,6-Dimethylpyrimidin-2-yl	1
450	OH	Phenyl	Phenyl	2-Hydroxyethyl	4-Methoxy-6-methylpyrimidin-2-yl	1
451	OH	Phenyl	Phenyl	2-Hydroxyethyl	4,6-Dimethoxypyrimidin-2-yl	1
452	OH	Phenyl	Phenyl	2-Hydroxyethyl	4,6-Diethylpyrimidin-2-yl	1
453	OH	Phenyl	Phenyl	2-Hydroxyethyl	6,7-Dihydro-5H-cyclopentapyrimidin-2-yl	1
454	OH	Phenyl	Phenyl	2-Hydroxyethyl	Naphth-2-yl	1
455	OH	Phenyl	Phenyl	2-Hydroxyethyl	6,7-Dihydro-5H-cyclopenta[c]pyridin-2-yl	1

No.	R	R ²	R ³	R ⁵	R ⁴	n
456	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Hydroxyethyl	Phenyl	1
457	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Hydroxyethyl	3,5-Dimethylphenyl	1
458	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Hydroxyethyl	3,5-Dimethoxyphenyl	1
459	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Hydroxyethyl	3,4-Methylenedioxyphenyl	1
460	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Hydroxyethyl	2,3-Dihydro-benzofuran-5-yl	1
461	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Hydroxyethyl	2,3-Dihydro-1H-inden-5-yl	1
462	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Hydroxyethyl	3-Methoxy-5-methylphenyl	1
463	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Hydroxyethyl	3,4-Ethylenedioxyphenyl	1
464	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Hydroxyethyl	5-Methoxy-3,4-methylenedioxyphenyl	1
465	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Hydroxyethyl	3,5-Diethylphenyl	1
466	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Hydroxyethyl	3,4-Dimethoxyphenyl	1
467	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Hydroxyethyl	3-Methyl-4-benzyloxyphenyl	1
468	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Hydroxyethyl	3-Benzyloxy-5-methoxyphenyl	1
469	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Hydroxyethyl	3-Benzyloxy-4-methoxyphenyl	1
470	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Hydroxyethyl	3-Hydroxy-4-methoxyphenyl	1
471	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Hydroxyethyl	4-Hydroxy-3-methoxyphenyl	1
472	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Hydroxyethyl	3,5-Bis-(trifluoromethyl)-phenyl	1
473	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Hydroxyethyl	3-Methylphenyl	1
474	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Hydroxyethyl	3-Methoxyphenyl	1
475	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Hydroxyethyl	4-Benzyloxyphenyl	1
476	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Hydroxyethyl	4-Phenoxyphenyl	1
477	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Hydroxyethyl	6-Methylpyridin-2-yl	1

No.	R	R ²	R ³	R ⁵	R ⁴	n
478	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Hydroxyethyl	6-Ethylpyridin-2-yl	1
479	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Hydroxyethyl	6-Methoxypyridin-2-yl	1
480	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Hydroxyethyl	4,6-Dimethylpyridin-2-yl	1
481	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Hydroxyethyl	4,6-Diethylpyridin-2-yl	1
482	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Hydroxyethyl	4-Methyl-6-methoxypyridin-2-yl	1
483	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Hydroxyethyl	6,7-Dihydro-5H-cyclopenta[b]pyridin-2-yl	1
484	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Hydroxyethyl	4,6-Dimethylpyrimidin-2-yl	1
485	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Hydroxyethyl	4-Methoxy-6-methylpyrimidin-2-yl	1
486	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Hydroxyethyl	4,6-Dimethoxypyrimidin-2-yl	1
487	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Hydroxyethyl	4,6-Diethylpyrimidin-2-yl	1
488	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Hydroxyethyl	6,7-Dihydro-5H-cyclopentapyrimidin-2-yl	1
489	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Hydroxyethyl	Naphth-2-yl	1
490	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Hydroxyethyl	6,7-Dihydro-5H-cyclopenta[c]pyridin-2-yl	1
491	OH	p-Methylphenyl	p-Methylphenyl	2-Hydroxyethyl	Phenyl	1
492	OH	p-Methylphenyl	p-Methylphenyl	2-Hydroxyethyl	3,5-Dimethylphenyl	1
493	OH	p-Methylphenyl	p-Methylphenyl	2-Hydroxyethyl	3,5-Dimethoxyphenyl	1
494	OH	p-Methylphenyl	p-Methylphenyl	2-Hydroxyethyl	3,4-Methylenedioxyphenyl	1
495	OH	p-Methylphenyl	p-Methylphenyl	2-Hydroxyethyl	2,3-Dihydro-benzofuran-5-yl	1
496	OH	p-Methylphenyl	p-Methylphenyl	2-Hydroxyethyl	2,3-Dihydro-1H-inden-5-yl	1
497	OH	p-Methylphenyl	p-Methylphenyl	2-Hydroxyethyl	3-Methoxy-5-methylphenyl	1
498	OH	p-Methylphenyl	p-Methylphenyl	2-Hydroxyethyl	3,4-Ethylenedioxyphenyl	1
499	OH	p-Methylphenyl	p-Methylphenyl	2-Hydroxyethyl	5-Methoxy-3,4-methylenedioxyphenyl	1

No.	R	R ²	R ³	R ⁵	R ⁴	n
500	OH	p-Methylphenyl	p-Methylphenyl	2-Hydroxyethyl	3,5-Diethylphenyl	1
501	OH	p-Methylphenyl	p-Methylphenyl	2-Hydroxyethyl	3,4-Dimethoxyphenyl	1
502	OH	p-Methylphenyl	p-Methylphenyl	2-Hydroxyethyl	3-Methyl-4-benzyloxyphenyl	1
503	OH	p-Methylphenyl	p-Methylphenyl	2-Hydroxyethyl	3-Benzyloxy-5-methoxyphenyl	1
504	OH	p-Methylphenyl	p-Methylphenyl	2-Hydroxyethyl	3-Benzyloxy-4-methoxyphenyl	1
505	OH	p-Methylphenyl	p-Methylphenyl	2-Hydroxyethyl	3-Hydroxy-4-methoxyphenyl	1
506	OH	p-Methylphenyl	p-Methylphenyl	2-Hydroxyethyl	4-Hydroxy-3-methoxyphenyl	1
507	OH	p-Methylphenyl	p-Methylphenyl	2-Hydroxyethyl	3,5-Bis-(trifluoromethyl)-phenyl	1
508	OH	p-Methylphenyl	p-Methylphenyl	2-Hydroxyethyl	3-Methylphenyl	1
509	OH	p-Methylphenyl	p-Methylphenyl	2-Hydroxyethyl	3-Methoxyphenyl	1
510	OH	p-Methylphenyl	p-Methylphenyl	2-Hydroxyethyl	4-Benzyloxyphenyl	1
511	OH	p-Methylphenyl	p-Methylphenyl	2-Hydroxyethyl	4-Phenoxyphenyl	1
512	OH	p-Methylphenyl	p-Methylphenyl	2-Hydroxyethyl	6-Methylpyridin-2-yl	1
513	OH	p-Methylphenyl	p-Methylphenyl	2-Hydroxyethyl	6-Ethylpyridin-2-yl	1
514	OH	p-Methylphenyl	p-Methylphenyl	2-Hydroxyethyl	6-Methoxypyridin-2-yl	1
515	OH	p-Methylphenyl	p-Methylphenyl	2-Hydroxyethyl	4,6-Dimethylpyridin-2-yl	1
516	OH	p-Methylphenyl	p-Methylphenyl	2-Hydroxyethyl	4,6-Diethylpyridin-2-yl	1
517	OH	p-Methylphenyl	p-Methylphenyl	2-Hydroxyethyl	4-Methyl-6-methoxypyridin-2-yl	1
518	OH	p-Methylphenyl	p-Methylphenyl	2-Hydroxyethyl	6,7-Dihydro-5H-cyclopenta[b]pyridin-2-yl	1
519	OH	p-Methylphenyl	p-Methylphenyl	2-Hydroxyethyl	4,6-Dimethylpyrimidin-2-yl	1
520	OH	p-Methylphenyl	p-Methylphenyl	2-Hydroxyethyl	4-Methoxy-6-methylpyrimidin-2-yl	1
521	OH	p-Methylphenyl	p-Methylphenyl	2-Hydroxyethyl	4,6-Dimethoxypyrimidin-2-yl	1

No.	R	R ²	R ³	R ⁵	R ⁴	n
522	OH	p-Methylphenyl	p-Methylphenyl	2-Hydroxyethyl	4,6-Diethylpyrimidin-2-yl	1
523	OH	p-Methylphenyl	p-Methylphenyl	2-Hydroxyethyl	6,7-Dihydro-5H-cyclopentapyrimidin-2-yl	1
524	OH	p-Methylphenyl	p-Methylphenyl	2-Hydroxyethyl	Naphth-2-yl	1
525	OH	p-Methylphenyl	p-Methylphenyl	2-Hydroxyethyl	6,7-Dihydro-5H-cyclopenta[c]pyridin-2-yl	1
526	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Hydroxyethyl	Phenyl	1
527	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Hydroxyethyl	3,5-Dimethylphenyl	1
528	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Hydroxyethyl	3,5-Dimethoxyphenyl	1
529	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Hydroxyethyl	3,4-Methylenedioxyphenyl	1
530	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Hydroxyethyl	2,3-Dihydro-benzofuran-5-yl	1
531	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Hydroxyethyl	2,3-Dihydro-1H-inden-5-yl	1
532	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Hydroxyethyl	3-Methoxy-5-methylphenyl	1
533	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Hydroxyethyl	3,4-Ethylenedioxyphenyl	1
534	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Hydroxyethyl	5-Methoxy-3,4-methylenedioxyphenyl	1
535	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Hydroxyethyl	3,5-Diethylphenyl	1
536	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Hydroxyethyl	3,4-Dimethoxyphenyl	1
537	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Hydroxyethyl	3-Methyl-4-benzyloxyphenyl	1
538	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Hydroxyethyl	3-Benzyloxy-5-methoxyphenyl	1
539	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Hydroxyethyl	3-Benzyloxy-4-methoxyphenyl	1
540	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Hydroxyethyl	3-Hydroxy-4-methoxyphenyl	1
541	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Hydroxyethyl	4-Hydroxy-3-methoxyphenyl	1
542	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Hydroxyethyl	3,5-Bis-(trifluoromethyl)-phenyl	1
543	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Hydroxyethyl	3-Methylphenyl	1

No.	R	R ²	R ³	R ⁵	R ⁴	n
544	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Hydroxyethyl	3-Methoxyphenyl	1
545	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Hydroxyethyl	4-Benzyloxyphenyl	1
546	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Hydroxyethyl	4-Phenoxyphenyl	1
547	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Hydroxyethyl	6-Methylpyridin-2-yl	1
548	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Hydroxyethyl	6-Ethylpyridin-2-yl	1
549	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Hydroxyethyl	6-Methoxypyridin-2-yl	1
550	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Hydroxyethyl	4,6-Dimethylpyridin-2-yl	1
551	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Hydroxyethyl	4,6-Diethylpyridin-2-yl	1
552	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Hydroxyethyl	4-Methyl-6-methoxypyridin-2-yl	1
553	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Hydroxyethyl	6,7-Dihydro-5H-cyclopenta[b]pyridin-2-yl	1
554	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Hydroxyethyl	4,6-Dimethylpyrimidin-2-yl	1
555	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Hydroxyethyl	4-Methoxy-6-methylpyrimidin-2-yl	1
556	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Hydroxyethyl	4,6-Dimethoxypyrimidin-2-yl	1
557	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Hydroxyethyl	4,6-Diethylpyrimidin-2-yl	1
558	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Hydroxyethyl	6,7-Dihydro-5H-cyclopentapyrimidin-2-yl	1
559	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Hydroxyethyl	Naphth-2-yl	1
560	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Hydroxyethyl	6,7-Dihydro-5H-cyclopenta[c]pyridin-2-yl	1
561	OH	Phenyl	Phenyl	2-Methylethyl	Phenyl	1
562	OH	Phenyl	Phenyl	2-Methylethyl	3,5-Dimethylphenyl	1
563	OH	Phenyl	Phenyl	2-Methylethyl	3,5-Dimethoxyphenyl	1
564	OH	Phenyl	Phenyl	2-Methylethyl	3,4-Methylenedioxyphenyl	1
565	OH	Phenyl	Phenyl	2-Methylethyl	2,3-Dihydro-benzofuran-5-yl	1

No.	R	R ²	R ³	R ⁵	R ⁴	n
566	OH	Phenyl	Phenyl	2-Methylethyl	2,3-Dihydro-1H-inden-5-yl	1
567	OH	Phenyl	Phenyl	2-Methylethyl	3-Methoxy-5-methylphenyl	1
568	OH	Phenyl	Phenyl	2-Methylethyl	3,4-Ethylenedioxyphenyl	1
569	OH	Phenyl	Phenyl	2-Methylethyl	5-Methoxy-3,4-methylenedioxyphenyl	1
570	OH	Phenyl	Phenyl	2-Methylethyl	3,5-Diethylphenyl	1
571	OH	Phenyl	Phenyl	2-Methylethyl	3,4-Dimethoxyphenyl	1
572	OH	Phenyl	Phenyl	2-Methylethyl	3-Methyl-4-benzyloxyphenyl	1
573	OH	Phenyl	Phenyl	2-Methylethyl	3-Benzyloxy-5-methoxyphenyl	1
574	OH	Phenyl	Phenyl	2-Methylethyl	3-Benzyloxy-4-methoxyphenyl	1
575	OH	Phenyl	Phenyl	2-Methylethyl	3-Hydroxy-4-methoxyphenyl	1
576	OH	Phenyl	Phenyl	2-Methylethyl	4-Hydroxy-3-methoxyphenyl	1
577	OH	Phenyl	Phenyl	2-Methylethyl	3,5-Bis-(trifluoromethyl)-phenyl	1
578	OH	Phenyl	Phenyl	2-Methylethyl	3-Methylphenyl	1
579	OH	Phenyl	Phenyl	2-Methylethyl	3-Methoxyphenyl	1
580	OH	Phenyl	Phenyl	2-Methylethyl	4-Benzyloxyphenyl	1
581	OH	Phenyl	Phenyl	2-Methylethyl	4-Phenoxyphenyl	1
582	OH	Phenyl	Phenyl	2-Methylethyl	6-Methylpyridin-2-yl	1
583	OH	Phenyl	Phenyl	2-Methylethyl	6-Ethylpyridin-2-yl	1
584	OH	Phenyl	Phenyl	2-Methylethyl	6-Methoxypyridin-2-yl	1
585	OH	Phenyl	Phenyl	2-Methylethyl	4,6-Dimethylpyridin-2-yl	1
586	OH	Phenyl	Phenyl	2-Methylethyl	4,6-Diethylpyridin-2-yl	1
587	OH	Phenyl	Phenyl	2-Methylethyl	4-Methyl-6-methoxypyridin-2-yl	1

No.	R	R ²	R ³	R ⁵	R ⁴	n
588	OH	Phenyl	Phenyl	2-Methylethyl	6,7-Dihydro-5H-cyclopenta[b]pyridin-2-yl	1
589	OH	Phenyl	Phenyl	2-Methylethyl	4,6-Dimethylpyrimidin-2-yl	1
590	OH	Phenyl	Phenyl	2-Methylethyl	4-Methoxy-6-methylpyrimidin-2-yl	1
591	OH	Phenyl	Phenyl	2-Methylethyl	4,6-Dimethoxypyrimidin-2-yl	1
592	OH	Phenyl	Phenyl	2-Methylethyl	4,6-Diethylpyrimidin-2-yl	1
593	OH	Phenyl	Phenyl	2-Methylethyl	6,7-Dihydro-5H-cyclopentapyrimidin-2-yl	1
594	OH	Phenyl	Phenyl	2-Methylethyl	Naphth-2-yl	1
595	OH	Phenyl	Phenyl	2-Methylethyl	6,7-Dihydro-5H-cyclopenta[c]pyridin-2-yl	1
596	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Methylethyl	Phenyl	1
597	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Methylethyl	3,5-Dimethylphenyl	1
598	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Methylethyl	3,5-Dimethoxyphenyl	1
599	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Methylethyl	3,4-Methylenedioxyphenyl	1
600	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Methylethyl	2,3-Dihydro-benzofuran-5-yl	1
601	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Methylethyl	2,3-Dihydro-1H-inden-5-yl	1
602	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Methylethyl	3-Methoxy-5-methylphenyl	1
603	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Methylethyl	3,4-Ethylenedioxyphenyl	1
604	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Methylethyl	5-Methoxy-3,4-methylenedioxyphenyl	1
605	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Methylethyl	3,5-Diethylphenyl	1
606	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Methylethyl	3,4-Dimethoxyphenyl	1
607	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Methylethyl	3-Methyl-4-benzyloxyphenyl	1
608	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Methylethyl	3-Benzyloxy-5-methoxyphenyl	1
609	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Methylethyl	3-Benzyloxy-4-methoxyphenyl	1

No.	R	R ²	R ³	R ⁵	R ⁴	n
610	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Methylethyl	3-Hydroxy-4-methoxyphenyl	1
611	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Methylethyl	4-Hydroxy-3-methoxyphenyl	1
612	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Methylethyl	3,5-Bis-(trifluoromethyl)-phenyl	1
613	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Methylethyl	3-Methylphenyl	1
614	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Methylethyl	3-Methoxyphenyl	1
615	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Methylethyl	4-Benzyloxyphenyl	1
616	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Methylethyl	4-Phenoxyphenyl	1
617	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Methylethyl	6-Methylpyridin-2-yl	1
618	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Methylethyl	6-Ethylpyridin-2-yl	1
619	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Methylethyl	6-Methoxypyridin-2-yl	1
620	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Methylethyl	4,6-Dimethylpyridin-2-yl	1
621	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Methylethyl	4,6-Diethylpyridin-2-yl	1
622	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Methylethyl	4-Methyl-6-methoxypyridin-2-yl	1
623	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Methylethyl	6,7-Dihydro-5H-cyclopenta[b]pyridin-2-yl	1
624	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Methylethyl	4,6-Dimethylpyrimidin-2-yl	1
625	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Methylethyl	4-Methoxy-6-methylpyrimidin-2-yl	1
626	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Methylethyl	4,6-Dimethoxypyrimidin-2-yl	1
627	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Methylethyl	4,6-Diethylpyrimidin-2-yl	1
628	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Methylethyl	6,7-Dihydro-5H-cyclopentapyrimidin-2-yl	1
629	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Methylethyl	Naphth-2-yl	1
630	OH	p-Methoxyphenyl	p-Methoxyphenyl	2-Methylethyl	6,7-Dihydro-5H-cyclopenta[c]pyridin-2-yl	1
631	OH	p-Methylphenyl	p-Methylphenyl	2-Methylethyl	Phenyl	1

No.	R	R ²	R ³	R ⁵	R ⁴	n
632	OH	p-Methylphenyl	p-Methylphenyl	2-Methylethyl	3,5-Dimethylphenyl	1
633	OH	p-Methylphenyl	p-Methylphenyl	2-Methylethyl	3,5-Dimethoxyphenyl	1
634	OH	p-Methylphenyl	p-Methylphenyl	2-Methylethyl	3,4-Methylenedioxyphenyl	1
635	OH	p-Methylphenyl	p-Methylphenyl	2-Methylethyl	2,3-Dihydro-benzofuran-5-yl	1
636	OH	p-Methylphenyl	p-Methylphenyl	2-Methylethyl	2,3-Dihydro-1H-inden-5-yl	1
637	OH	p-Methylphenyl	p-Methylphenyl	2-Methylethyl	3-Methoxy-5-methylphenyl	1
638	OH	p-Methylphenyl	p-Methylphenyl	2-Methylethyl	3,4-Ethylenedioxyphenyl	1
639	OH	p-Methylphenyl	p-Methylphenyl	2-Methylethyl	5-Methoxy-3,4-methylenedioxyphenyl	1
640	OH	p-Methylphenyl	p-Methylphenyl	2-Methylethyl	3,5-Diethylphenyl	1
641	OH	p-Methylphenyl	p-Methylphenyl	2-Methylethyl	3,4-Dimethoxyphenyl	1
642	OH	p-Methylphenyl	p-Methylphenyl	2-Methylethyl	3-Methyl-4-benzyloxyphenyl	1
643	OH	p-Methylphenyl	p-Methylphenyl	2-Methylethyl	3-Benzyloxy-5-methoxyphenyl	1
644	OH	p-Methylphenyl	p-Methylphenyl	2-Methylethyl	3-Benzyloxy-4-methoxyphenyl	1
645	OH	p-Methylphenyl	p-Methylphenyl	2-Methylethyl	3-Hydroxy-4-methoxyphenyl	1
646	OH	p-Methylphenyl	p-Methylphenyl	2-Methylethyl	4-Hydroxy-3-methoxyphenyl	1
647	OH	p-Methylphenyl	p-Methylphenyl	2-Methylethyl	3,5-Bis-(trifluoromethyl)-phenyl	1
648	OH	p-Methylphenyl	p-Methylphenyl	2-Methylethyl	3-Methylphenyl	1
649	OH	p-Methylphenyl	p-Methylphenyl	2-Methylethyl	3-Methoxyphenyl	1
650	OH	p-Methylphenyl	p-Methylphenyl	2-Methylethyl	4-Benzyloxyphenyl	1
651	OH	p-Methylphenyl	p-Methylphenyl	2-Methylethyl	4-Phenoxyphenyl	1
652	OH	p-Methylphenyl	p-Methylphenyl	2-Methylethyl	6-Methylpyridin-2-yl	1
653	OH	p-Methylphenyl	p-Methylphenyl	2-Methylethyl	6-Ethylpyridin-2-yl	1

No.	R	R ²	R ³	R ⁵	R ⁴	n
654	OH	p-Methylphenyl	p-Methylphenyl	2-Methylethyl	6-Methoxypyridin-2-yl	1
655	OH	p-Methylphenyl	p-Methylphenyl	2-Methylethyl	4,6-Dimethylpyridin-2-yl	1
656	OH	p-Methylphenyl	p-Methylphenyl	2-Methylethyl	4,6-Diethylpyridin-2-yl	1
657	OH	p-Methylphenyl	p-Methylphenyl	2-Methylethyl	4-Methyl-6-methoxypyridin-2-yl	1
658	OH	p-Methylphenyl	p-Methylphenyl	2-Methylethyl	6,7-Dihydro-5H-cyclopenta[b]pyridin-2-yl	1
659	OH	p-Methylphenyl	p-Methylphenyl	2-Methylethyl	4,6-Dimethylpyrimidin-2-yl	1
660	OH	p-Methylphenyl	p-Methylphenyl	2-Methylethyl	4-Methoxy-6-methylpyrimidin-2-yl	1
661	OH	p-Methylphenyl	p-Methylphenyl	2-Methylethyl	4,6-Dimethoxypyrimidin-2-yl	1
662	OH	p-Methylphenyl	p-Methylphenyl	2-Methylethyl	4,6-Diethylpyrimidin-2-yl	1
663	OH	p-Methylphenyl	p-Methylphenyl	2-Methylethyl	6,7-Dihydro-5H-cyclopentapyrimidin-2-yl	1
664	OH	p-Methylphenyl	p-Methylphenyl	2-Methylethyl	Naphth-2-yl	1
665	OH	p-Methylphenyl	p-Methylphenyl	2-Methylethyl	6,7-Dihydro-5H-cyclopenta[c]pyridin-2-yl	1
666	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Methylethyl	Phenyl	1
667	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Methylethyl	3,5-Dimethylphenyl	1
668	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Methylethyl	3,5-Dimethoxyphenyl	1
669	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Methylethyl	3,4-Methylenedioxyphenyl	1
670	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Methylethyl	2,3-Dihydro-benzofuran-5-yl	1
671	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Methylethyl	2,3-Dihydro-1H-inden-5-yl	1
672	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Methylethyl	3-Methoxy-5-methylphenyl	1
673	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Methylethyl	3,4-Ethylenedioxyphenyl	1
674	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Methylethyl	5-Methoxy-3,4-methylenedioxyphenyl	1
675	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Methylethyl	3,5-Diethylphenyl	1

No.	R	R ²	R ³	R ⁵	R ⁴	n
676	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Methylethyl	3,4-Dimethoxyphenyl	1
677	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Methylethyl	3-Methyl-4-benzyloxyphenyl	1
678	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Methylethyl	3-Benzyloxy-5-methoxyphenyl	1
679	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Methylethyl	3-Benzyloxy-4-methoxyphenyl	1
680	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Methylethyl	3-Hydroxy-4-methoxyphenyl	1
681	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Methylethyl	4-Hydroxy-3-methoxyphenyl	1
682	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Methylethyl	3,5-Bis-(trifluoromethyl)-phenyl	1
683	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Methylethyl	3-Methylphenyl	1
684	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Methylethyl	3-Methoxyphenyl	1
685	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Methylethyl	4-Benzyloxyphenyl	1
686	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Methylethyl	4-Phenoxyphenyl	1
687	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Methylethyl	6-Methylpyridin-2-yl	1
688	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Methylethyl	6-Ethylpyridin-2-yl	1
689	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Methylethyl	6-Methoxypyridin-2-yl	1
690	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Methylethyl	4,6-Dimethylpyridin-2-yl	1
691	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Methylethyl	4,6-Diethylpyridin-2-yl	1
692	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Methylethyl	4-Methyl-6-methoxypyridin-2-yl	1
693	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Methylethyl	6,7-Dihydro-5H-cyclopenta[b]pyridin-2-yl	1
694	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Methylethyl	4,6-Dimethylpyrimidin-2-yl	1
695	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Methylethyl	4-Methoxy-6-methylpyrimidin-2-yl	1
696	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Methylethyl	4,6-Dimethoxypyrimidin-2-yl	1
697	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Methylethyl	4,6-Diethylpyrimidin-2-yl	1

No.	R	R ²	R ³	R ⁵	R ⁴	n
698	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Methylethyl	6,7-Dihydro-5H-cyclopentapyrimidin-2-yl	1
699	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Methylethyl	Naphth-2-yl	1
700	OH	4-Methoxy-3-methylphenyl	4-Methoxy-3-methylphenyl	2-Methylethyl	6,7-Dihydro-5H-cyclopenta[c]pyridin-2-yl	1

The compounds of the present invention provide a novel therapeutic potential for the treatment of hypertension, pulmonary hypertension, myocardial infarct, angina pectoris, acute kidney failure, renal insufficiency, cerebral vasospasms, cerebral ischemia, subarachnoid hemorrhages, migraine, asthma, atherosclerosis, endotoxic shock, endotoxin-induced organ failure, intravascular coagulation, restenosis after angioplasty, benign prostate hyperplasia, kidney failure caused by ischemia and by intoxication, and hypertension.

The good effect of the compounds can be shown in the following tests:

15 Receptor-binding studies

Cloned human ET_A receptor-expressing CHO cells and guinea pig cerebellar membranes with > 60% ET_B receptors by comparison with ET_A receptors were employed for binding studies.

20

Membrane preparation

The ET_A receptor-expressing CHO cells were grown in F_{12} medium containing 10% fetal calf serum, 1% glutamine, 100 U/ml penicillin and 0.2% streptomycin (Gibco BRL, Gaithersburg, MD, USA). After 48 h, the cells were washed with PBS and incubated with 0.05% trypsin-containing PBS for 5 min. Then neutralization was carried out with F_{12} medium, and the cells were collected by centrifugation at 300 x g. To lyse the cells, the pellet was briefly washed with lysis buffer (5 mM tris-HCl, pH 7.4 with 10% glycerol) and then incubated at a concentration of 10^7 cells/ml of lysis buffer at 4°C for 30 min. The membranes were centrifuged at 20,000 x g for 10 min, and the pellet was stored in liquid nitrogen.

35

Guinea pig cerebella were homogenized in a Potter-Elvehjem homogenizer and obtained by differential centrifugation at 1,000 x g for 10 min and repeated centrifugation of the supernatant at 20,000 x g for 10 min.

40

Binding assays

For the ET_A and ET_B receptor binding assay, the membranes were suspended in an incubation buffer (50 mM tris-HCl, pH 7.4, with 5 mM $MnCl_2$, 40 μ g/ml bacitracin and 0.2% BSA) at a concentration of 50 μ g of protein per assay mixture and incubated in the presence and absence of test substance with 25 pM ^{125}I - ET_1 (ET_A

50

receptor assay) or 25 pM 125 -RZ₃ (ET_B receptor assay) at 25°C. The nonspecific binding was determined with 10^{-7} M ET₁. After 30 min, the free and bound radioligand were separated by filtration through GF/B glass fiber filters (Whatman, England) on a Skatron
5 cell collector (Skatron, Lier, Norway), and the filters were washed with ice-cold tris-HCl buffer, pH 7.4 with 0.2% BSA. The radioactivity collected on the filters was quantified using a Packard 2200 CA liquid scintillation counter.

10 Functional in vitro assay system for searching for endothelin receptor (subtype A) antagonists

This assay system is a functional, cell-based assay for endothelin receptors. Certain cells show, when they are
15 stimulated with endothelin 1 (ET1), an increase in the intracellular calcium concentration. This increase can be measured in intact cells which have been loaded with calcium-sensitive dyes.

20 1-Fibroblasts which were isolated from rats and in which an endogenous endothelin receptor of the A subtype had been detected were loaded with the fluorescent dye Fura 2-am as follows: after tripysinization, the cells were resuspended in buffer A (120 mM NaCl, 5 mM KCl, 1.5 mM MgCl₂, 1 mM CaCl₂, 25 mM HEPES, 10 mM
25 Glucose, pH 7.4) to a density of 2×10^6 /ml and incubated with Fura-2-am (2 μ M), Pluronic F-127 (0.04%) and DMSO (0.2%) at 37°C in the dark for 30 min. The cells were then washed twice with buffer A and resuspended at 2×10^6 /ml.

30 The fluorescence signal from 2×10^5 cells per ml with Ex/Em 380/510 was recorded continuously at 30°C. The test substances and, after an incubation time of 3 min, ET1 were added to the cells. The maximum change in the fluorescence was determined over 30 min. The response of the cells to ET1 without
35 previous addition of a test substance served as control and was set equal to 100%.

In vivo testing of ET antagonists

40 Male SD rats weighting 250 - 300 g were anesthetized with amobarbital, artificially ventilated, vagotomized and pithed. The carotid artery and the jugular vein were catheterized.

Intravenous administration of 1 μ g/kg ET1 to control animals leads
45 to a distinct rise in blood pressure which persists for a lengthy period.

51

The test compounds was [sic] injected i.v. (1 ml/kg) into the test animals 5 or 30 min before ET1 administration. To determine the ET-antagonistic properties, the increase in blood pressure in the test animals was compared with that in the control animals.

5

Endothelin-1 induced sudden death in mice

The principle of the test comprises the inhibition of the sudden heart death in mice which is caused by endothelin, probably due to constriction of the coronary vessels, by pretreatment with endothelin receptor antagonists. Intravenous injection of 10 nmol/kg endothelin in a volume of 5 ml/kg of body weight results in death of the animals within a few minutes.

15 The lethal endothelin-1 dose is checked in each case on a small group of animals. If the test substance is administered intravenously, it is usually followed after 5 min by the endothelin-1 injection which was lethal in the reference group. With other modes of administration, the times before administration are longer, where appropriate up to several hours.

The survival rate is recorded and effective doses which protect 50% of the animals for 24 h or longer against endothelin-induced heart death (ED 50) are determined.

25

Functional test for endothelin receptor antagonists on vessels

Segments of rabbit aorta with an initial tension of 2 g and a relaxation time of 1 h in Krebs-Henseleit solution at 37°C and at a pH of from 7.3 to 7.4 are initially induced to contract with K⁺. After washing out, an endothelin dose-effect plot is constructed up to the maximum.

Potential endothelin antagonists are administered to other preparations of the same vessel 15 min before starting the endothelin dose-effect plot. The effects of endothelin are calculated as a % of the K⁺ contraction. Effective endothelin antagonists result in a shift in the endothelin dose-effect plot to the right.

40

The compounds according to the invention can be administered orally or parenterally (subcutaneously, intravenously, intramuscularly, intraperitoneally) in a conventional way. Administration can also take place with vapors or sprays through the nasopharyngeal space.

The dosage depends on the age, condition and weight of the patient and on the mode of administration. As a rule, the daily dose of agent is about 0.5-50 mg/kg of body weight on oral administration and about 0.1-10 mg/kg of body weight on 5 parenteral administration.

The novel compounds can be used in conventional solid or liquid pharmaceutical forms, eg. as uncoated or (film-)coated tablets, capsules, powders, granules, suppositories, solutions, ointments, 10 creams or sprays. These are produced in a conventional way. The agents can for this purpose be processed with conventional pharmaceutical aids such as tablet binders, bulking agents, preservatives, tablet disintegrants, flow regulators, plasticizers, wetting agents, dispersants, emulsifiers, solvents, 15 release-slowing agents, antioxidants and/or propellant gases (cf. H. Sucker et al.: Pharmazeutische Technologie, Thieme-Verlag, Stuttgart, 1991). The administration forms obtained in this way normally contain from 0.1 to 90% by weight of the agent.

20 Synthetic Examples

Example 1

3,3-Bis(4-methoxyphenyl)butanoic acid

25 a) Ethyl (2E,Z)-3-(4-methoxyphenyl)-2-butenate (6.6 g, 30 mmol) were dissolved in anisole (4.9 g, 45 mmol) at 0°C, and 50 ml of 80% H₂SO₄ were cautiously added. The 2-phase mixture was vigorously stirred at room temperature for 20 h and then poured onto ice, and the product was extracted with ethyl 30 acetate. The organic phase was dried (Na₂SO₄), filtered and concentrated, the residue was taken up with ether and extracted with 2N sodium hydroxide solution, and the ether phase was discarded. The alkaline phase was adjusted to pH 2 with 2N HCl, and the product was extracted with ethyl 35 acetate. The organic phase was then dried (Na₂SO₄), filtered and concentrated, and the solid residue was stirred with diisopropyl ether. The product was filtered off with suction and dried. 5.1 g of a white powder (56%) remained. Melting point: 161-164°C

40

Further working up of the mother liquor was possible, resulting in a further 1.1 g (12%) of the acid.

45

The acid can also be prepared by the following alternative:

- b) At 0°C, 32 ml of anisole (294 mmol) were mixed with 33 ml of ethyl acetoacetate (258 mmol), 150 ml of 70% H₂SO₄ were cautiously added, and the resulting 2-phase mixture was vigorously stirred at room temperature for 72 h. The mixture was then poured onto ice and worked up further as under 1a). The residue was recrystallized from diisopropyl ether. 15.3 g (35%) of white solid remained.
- c) Similar to the preparation of 3,3-diphenylbutanoic acid (Examples 3, 4)

Example 2

- (2R,S)-3,3-Bis-(4-methoxyphenyl)-2-(3',4'-methylenedioxybenzyl)-butanoic acid

Butyllithium (13.8 ml, 22 mmol, 1.6M in hexane) was added to a solution of diisopropylamine (3.1 ml, 22 mmol) in 50 ml of dry tetrahydrofuran under nitrogen at -10°C, the mixture was stirred at -10°C for 5 min and then, at 0°C, 3,3-bis(4-methoxyphenyl)butanoic acid (3.0 g, 10 mmol) in 15 ml of absolute THF was added dropwise. After the addition was complete, the mixture was stirred at room temperature for 1 h, cooled to -20°C and, after addition of piperonyl bromide (2.6 g, 12 mmol) in 10 ml of THF, stirred at room temperature for 72 h. The mixture was then quenched with saturated NH₄Cl solution, the organic phase was separated off and the aqueous was extracted with ethyl acetate. The combined organic extracts were dried (Na₂SO₄), filtered and concentrated in a rotary evaporator. The brown residue was chromatographed on silica gel (methanol/CH₂Cl₂ 1:19), resulting in 1.3 g (30%) of product as a white foam. Melting point: 137-140°C (from diisopropyl ether)

Example 3

Ethyl 3,3-diphenylbutanoate

At 0°C, 65 g of AlCl₃ (487 mmol) were suspended in 500 ml of benzene, and 61.7 g of ethyl (2E,Z)-3-phenyl-2-butenate were slowly added. The dark red solution was stirred at room temperature for 20 h and then poured into a mixture of ice and concentrated HCl. The organic phase was separated off, and the aqueous was extracted with ethyl acetate. The combined organic phases were extracted with NaOH and then dried (Na₂SO₄), filtered and concentrated (66.8 g of dark brown oil).

54

56.5 g of this oil were distilled, resulting in 46.3 g of product as a colorless oil.

Example 4

5 3,3-Diphenylbutanoic acid

4.9 g of ethyl 3,3-diphenylbutanoate (18.3 mmol) were dissolved in 30 ml of dioxane, 36 ml of 1M KOH were added, and the mixture was stirred at 60-70°C for 6 h.

10

The dioxane was then stripped off in a rotary evaporator, and the aqueous residue was diluted with water and extracted with diethyl ether. The aqueous phase was then adjusted to pH 1 and extracted with ethyl acetate. The organic phase was dried (Na_2SO_4), filtered and concentrated. The solid residue was stirred with heptane, resulting in 2.35 g of a white powder (55%). The mother liquor was not purified further.

Example 5

20 (2R,S)-3,3-Diphenyl-2-(3',4'-methylenedioxybenzyl)butanoic acid

15 ml of butyllithium (24 mmol, 1.6M in hexane) were added dropwise to a solution of 3,3-diphenylbutanoic acid (2.4 g, 10 mmol) in 40 ml of absolute THF at -20°C, and the mixture was then stirred at from -10 to -20°C for 1 h. Then piperonyl chloride (2.2 g, 13 mmol) in 10 ml of THF was added, and the mixture was stirred at room temperature for 16 h and then quenched with saturated NH_4Cl solution. The organic phase was separated off, the aqueous phase was extracted with ethyl acetate, and then the combined organic extracts were dried (Na_2SO_4), filtered and concentrated. The residue was chromatographed on silica gel ($\text{CH}_2\text{Cl}_2/\text{MeOH}$ 19:1), resulting in 2.4 g of the required product (65%).

35 The acid was dissolved in CH_2Cl_2 and shaken with saturated sodium carbonate solution. The organic (!) phase was separated off, dried (Na_2SO_4), filtered and concentrated. 2.5 g of sodium salt of the acid were obtained.

Melting point: 308-310°C (decomposition)

40

Example 6

3,3-Bis(4-methoxy-3-methylphenyl)butanoic acid

Preparation took place as in Example 1b. However, in this case, mainly the corresponding ethyl ester was isolated so that subsequent hydrolysis was necessary (as in Example 4).

Melting point: 121-124°C

Example 7

(2R,S)-3,3-Bis(4-methoxy-3-methylphenyl)-2-(3',4'-methylenedioxybenzyl)butanoic acid

5 Preparation similar to Example 2.

3.25 ml of diisopropylamine (23 mmol), 15.6 ml of butyllithium (23 mmol, 1.5 M in hexane), 3.28 g of 3,3-bis(4-methoxy-3-methylphenyl)butanoic acid (10 mmol), 2.19 g of piperonyl chloride (13 mmol) afford 4.1 g of crude product.

10 Chromatography on silica gel (CH₂Cl₂/MeOH 19:1) afforded 1.6 g of product (35%)

Melting point: 152-153°C

Example 8

15 (2R,S)-3,3-Diphenyl-2-(3',4'-dimethoxybenzyl)butanoic acid

Preparation took place as in Example 5. 2.4 g of 3,3-diphenylbutanoic acid (10 mmol), 15.6 ml of butyllithium (23 mmol, 1.5 M in hexane), 2.2 g of 3,4-dimethoxybenzyl chloride (13 mmol) afforded 3.8 g of crude product.

Purification on silica gel (heptane/ethyl acetate 1:1) 2.1 g of product (54%)

Melting point: 141-143°C

25 Example 9

3,3-Bis(4-methoxyphenyl)pentanoic acid

Ethyl (2E,Z)-3-(4-methoxyphenyl)-2-pentenoate (7.0 g, 30 mmol) was dissolved in anisole (4.9 g, 45 mmol) at 0°C, and 50 ml of 80% H₂SO₄ were cautiously added. The 2-phase mixture was vigorously

30 stirred at room temperature for 30 h and then poured onto ice, and the product was extracted with methylene chloride. The organic phase was dried (Na₂SO₄), filtered and concentrated, the residue was taken up in ether and extracted with 2N sodium hydroxide solution, and the ether phase was discarded. The
35 alkaline phase was adjusted to pH 2 with 2N HCl, and the product was extracted with ethyl acetate. The organic phase was then dried (Na₂SO₄), filtered and concentrated, and the solid residue was stirred with heptane. The product was filtered off with suction and dried. 6.8 g of a white powder (72%) remained.

40 Melting point: 136-139°C

Example 10

(2R,S)-3,3-Bis(4-methoxyphenyl)pentanoic acid

45 29 ml of butyllithium (46 mmol, 1.6 M in hexane) were added dropwise to a solution of 3,3-bis(4-methoxyphenyl)pentanoic acid (6.2 g, 20 mmol) in 100 ml of absolute THF at -20°C, and the

56

- mixture was then stirred at room temperature for 1 h. Then, at -10°C, piperonyl chloride (4.4 g, 24 mmol) in 10 ml of THF was added, and the mixture was stirred at room temperature for 72 h and then quenched with saturated NH₄Cl solution. The organic phase
- 5 was separated off, the aqueous phase was extracted with ethyl acetate, and then the combined organic extracts were dried (Na₂SO₄), filtered and concentrated. The residue (11.2 g) was chromatographed on silica gel (CH₂Cl₂/MeOH 24:1), resulting in 3.1 g of the required product (34%).
- 10 Melting point: 84-86°C (stirred in heptane)

Example 11

Methyl 3,3-Bis(4-methoxyphenyl)hexanoate

- 15 Anisole (6.6 g, 61 mmol) was dissolved in 200 ml of dichloroethane and, at 0°C, aluminum trichloride (12.3 g, 92 mmol) was added in portions and subsequently, while stirring, methyl (2E,Z)-3-(4-methoxyphenyl)-2-hexenoate (18 g, 61 mmol) was added dropwise. The reaction mixture was stirred at 5°C for 2 h and then
- 20 at room temperature for 2 days. For workup, the mixture was poured into ice-water and extracted with CH₂Cl₂, and the combined organic phases were washed with saturated NaCl solution and dried over MgSO₄. The residue remaining after concentration was purified by chromatography on silica gel (n-heptane/7.5% ethyl acetate).
- 25 This resulted in 5.2 g (25%) of a colorless oil.
¹H-NMR (CDCl₃), δ: 0.9 (m, 3H), 1.1 and 2.2 (each m, 2H), 3.08 (s, 2H), 3.4 (s, 3H) (s, 6H), 6.8 and 7.1 (each m, 4H) ppm.

Example 12

- 30 3,3-Bis(4-methoxyphenyl)hexanoic acid

- Methyl 3,3-bis(4-methoxyphenyl)hexanoate (5.2 g, 15.2 mmol) was introduced into 20 ml of dioxane, KOH (1.05 g, 18.2 mmol) was added, and the mixture was boiled for about 1 h. It was
- 35 subsequently diluted with water and washed with ethyl acetate, and the aqueous phase was then adjusted to pH 3 with dilute HCl and extracted with ethyl acetate. The organic phase was then washed with saturated NaCl solution, dried over MgSO₄ and concentrated. Chromatography on silica gel (CH₂Cl₂/methanol 3%)
- 40 resulted in 4.1 g of a pale yellowish oil (84%).
¹H-NMR (CDCl₃), δ: 0.9 (m, 3H), 1.1 und 2.2 (each m, 2H), 3.1 (s, 3H), 3.8 (s, 6H), 6.8 and 7.1 (each m, 4H) ppm.

Example 13

45

The following compounds were prepared as in Example 5.

(2R,S)-3,3-Diphenyl-2-(methyl-2'-naphthyl)butanoic acid

Melting point: 163-166°C

FAB-MS: 380 (M⁺)

5 (2R,S)-3,3-Diphenyl-2-(3',5'-dimethylbenzyl)butanoic acid

Melting point: 141-143°C

FAB-MS: 358 (M⁺)

(2R,S)-3,3-Diphenyl-2-(4'-benzyloxy-3'-methoxybenzyl)butanoic
10 acid

Melting point: 163-166°C

FAB-MS: 466 (M⁺)

(2R,S)-3,3-Bis(4-methoxyphenyl)-2-(4'-benzyloxy-3-methoxybenzyl)-
15 butanoic acid

Melting point: 137-140°C

FAB-MS: 526 (M⁺)

(2R,S)-3,3-Diphenyl-2-(4'-hydroxy-3'-methoxybenzyl)butanoic acid
20 Melting point: 153-155°C

FAB-MS: 376 (M⁺)

(2R,S)-3,3-Bis(4-methoxyphenyl)-2-(4'-hydroxy-3-methoxybenzyl)-
butanoic acid

25 Melting point: 157-160°C

FAB-MS: 436 (M⁺)

(2R,S)-3,3-Bis(4-methoxy-3-methylphenyl)-2-(3',5'-dimethyl-
benzyl)butanoic acid

30 Melting point: 150-152°C

FAB-MS: 446 (M⁺)

(2R,S)-3,3-Bis(4-methoxyphenyl)-2-(methyl-2'-naphthyl)butanoic
acid

35 Melting point: 162-164°C

FAB-MS: 440 (M⁺)

(2R,S)-3,3-Bis(4-methoxyphenyl)-2-(3',5'-dimethylbenzyl)butanoic
acid

40 Melting point: 125-128°C

FAB-MS: 418 (M⁺)

(2R,S)-3,3-Bis(4-methoxy-3-methylphenyl)-2-(3',4'-dimethoxy-
benzyl)butanoic acid

45 Melting point: 155-157°C

FAB-MS: 478 (M⁺)

58

(2R,S)-3,3-Bis(4-methoxyphenyl)-2-(3',4'-dimethoxybenzyl)butanoic acid

Melting point: 148-150°C

FAB-MS: 450 (M⁺)

5

(2R,S)-3,3-Bis(4-methoxyphenyl)-2-(5'-methoxy-3',4'-methylene-dioxybenzyl)pentanoic acid

Melting point: 138-141°C

FAB-MS: 478 (M⁺)

10

(2R,S)-3,3-Bis(4-methoxyphenyl)-2-(5-methoxy-3',4'-methylene-dioxybenzyl)butanoic acid

Melting point: 134-136°C

FAB-MS: 464 (M⁺)

15

(2R,S)-3,3-Diphenyl-2-(5'-methoxy-3',4'-methylenedioxybenzyl)-butanoic acid

Melting point: 135-138°C

FAB-MS: 464 (M⁺)

20

(2R,S)-3,3-Bis(4-methoxyphenyl)-2-(3',4'-ethylenedioxybenzyl)-pentanoic acid

Melting point: 168-170°C

FAB-MS: 462 (M⁺)

25

(2R,S)-3,3-Bis(4-methoxyphenyl)-2-(3',4'-ethylenedioxybenzyl)-butanoic acid

Melting point: 161-163°C

FAB-MS: 448 (M⁺)

30

(2R,S)-3,3-Bis(4-methoxyphenyl)-2-(3',4'-methylenedioxybenzyl)-hexanoic acid

Melting point: 142-145°C (from n-heptane)

35 (2R,S)-3,3-Bis(4-methoxyphenyl)-2-(3',4'-ethylenedioxybenzyl)-hexanoic acid

Melting point: 163-165°C (from n-heptane/diethyl ether)




(2R,S)-3,3-Bis(4-methoxyphenyl)-2-(3',4'-methylenedioxy-5'-methoxybenzyl)hexanoic acid

40 Melting point: 180-182°C (from n-heptane/diethyl ether)

45

Example 14

The compounds prepared in Examples 2 to 10 were checked for their endothelin receptor affinity by the methods described above. A compound disclosed in WO 94/02474 was used as comparison substance. The result is reported in the following Table.


		ET _A	ET _B
10	 (WO 94/02474)	420 nM	> 6400 nM
15			
20	 (Example 2)	38 nM	2800 nM
25			
30	 (Example 10)	6 nM	1300 nM
35			
40			
45			



We claim:

A carboxylic acid derivative of the formula I

5



10

where R^1 is a tetrazole [sic], nitrile [sic], COOH or a radical which can be hydrolyzed to COOH, and the other substituents have the following meanings:

15 R^2 and R^3 (which can be identical or different):

phenyl or naphthyl which can be substituted by one or more of the following radicals: halogen, cyano, NO_2 , hydroxyl, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkoxy, 20 phenoxy, C_1 - C_4 -alkylthio, amino, benzyloxy, C_1 - C_4 -alkylamino or C_1 - C_4 -dialkylamino; or

25

phenyl or naphthyl which are connected together in the ortho positions by a direct linkage, a methylene, ethylene or ethenylene group, or an oxygen or sulfur atom;

R^4 phenyl or naphthyl, methylenedioxyphenyl, ethylenedioxyphenyl, indanyl, indolyl, pyridyl, benzopyranyl, furanyl, pyrimidinyl, benzofuranyl, isooxazolyl, 30 isothiazolyl, 1,3,4-thiadiazolyl, 2,3-dihydrobenzofuranyl, benzothienyl, quinolinyl, C_3 - C_7 -cycloalkyl, thienyl, oxazolyl, thiazolyl, each of which can be substituted by one or more of the following radicals: halogen, cyano, hydroxyl, NO_2 , C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkoxy, 35 phenoxy, C_1 - C_4 -alkylthio, amino, benzyloxy, C_1 - C_4 -alkylamino or C_1 - C_4 -dialkylamino, it being possible for the alkyl radicals together to form a ring;

R^5 C_1 - C_8 -alkyl, C_3 - C_6 -alkenyl, C_3 - C_6 -alkynyl or C_3 - C_8 -cycloalkyl, 40 it being possible for each of these radicals to be substituted one or more times by: halogen, C_1 - C_4 -alkoxy, C_1 - C_4 -alkylthio, C_1 - C_4 -alkylamino, di- C_1 - C_4 -alkylamino;

phenyl, benzyl, 1-methylnaphthyl, 2-methylnaphthyl or 45 naphthyl, each of which can be substituted by one or more of the following radicals: halogen, cyano, hydroxyl, amino,



2

C₁-C₄-alkyl, C₁-C₄-alkoxy, phenoxy, C₁-C₄-alkylthio,
dioxomethylene [sic] or dioxoethylene [sic];

n 1 - 2.

5

10

15

20

25

30

35

40

45