

5

NEPRILYSIN INHIBITORS

BACKGROUND OF THE INVENTION

FIELD OF THE INVENTION

The present invention relates to novel compounds having neprilysin-inhibition activity. The invention also relates to pharmaceutical compositions comprising such
10 compounds, processes and intermediates for preparing such compounds and methods of using such compounds to treat diseases such as hypertension, heart failure, pulmonary hypertension, and renal disease.

STATE OF THE ART


Neprilysin (neutral endopeptidase, EC 3.4.24.11) (NEP), is an endothelial
15 membrane bound Zn^{2+} metallopeptidase found in many organs and tissues, including the brain, kidneys, lungs, gastrointestinal tract, heart, and the peripheral vasculature. NEP degrades and inactivates a number of endogenous peptides, such as enkephalins, circulating bradykinin, angiotensin peptides, and natriuretic peptides, the latter of which have several effects including, for example, vasodilation and natriuresis/diuresis, as well as
20 inhibition of cardiac hypertrophy and ventricular fibrosis. Thus, NEP plays an important role in blood pressure homeostasis and cardiovascular health.

NEP inhibitors, such as thiorphan, candoxatril, and candoxatrilat, have been studied as potential therapeutics. Compounds that inhibit both NEP and angiotensin-I converting enzyme (ACE) are also known, and include omapatrilat, gempatrilat, and sampatrilat.
25 Referred to as vasopeptidase inhibitors, this latter class of compounds is described in Robl et al. (1999) *Exp. Opin. Ther. Patents* 9(12): 1665-1677.

SUMMARY

The present disclosure provides novel compounds that have been found to possess neprilysin (NEP) enzyme inhibition activity. Accordingly, compounds of the present
30 disclosure are expected to be useful and advantageous as therapeutic agents for treating conditions such as hypertension and heart failure.

One aspect of the present disclosure relates to a compound of formula I:



(I)

where:

R^1 is selected from $-OR^7$ and $-NR^8R^9$;

R^2 is H or $-P(O)(OH)_2$ or R^2 is taken together with R^7 to form $-CR^{18}R^{19}-$ or is taken
5 together with R^8 to form $-C(O)-$;

X is a $-C_{1-9}$ heteroaryl;

R^3 is absent or is selected from H; halo; $-C_{0-5}$ alkylene-OH; $-NH_2$; $-C_{1-6}$ alkyl; $-CF_3$;
 $-C_{3-7}$ cycloalkyl; $-C_{0-2}$ alkylene-O- C_{1-6} alkyl; $-C(O)R^{20}$; $-C_{0-1}$ alkylene-COOR²¹;
 $-C(O)NR^{22}R^{23}$; $-NHC(O)R^{24}$; $=O$; $-NO_2$; $-C(CH_3)=N(OH)$; phenyl optionally substituted
10 with one or two groups independently selected from halo, $-OH$, $-CF_3$, $-OCH_3$,
 $-NHC(O)CH_3$, and phenyl; naphthalenyl; pyridinyl; pyrazinyl; pyrazolyl optionally
substituted with methyl; thiophenyl optionally substituted with methyl or halo; furanyl; and
 $-CH_2$ -morpholinyl; and R^3 , when present, is attached to a carbon atom;


R^4 is absent or is selected from H; $-OH$; $-C_{1-6}$ alkyl; $-C_{1-2}$ alkylene-COOR³⁵;
15 $-CH_2OC(O)CH(R^{36})NH_2$; $-OCH_2OC(O)CH(R^{36})NH_2$; $-OCH_2OC(O)CH_3$;
 $-CH_2OP(O)(OH)_2$; $-CH_2CH(OH)CH_2OH$; $-CH[CH(CH_3)_2]-NHC(O)O-C_{1-6}$ alkyl; pyridinyl;
and phenyl or benzyl optionally substituted with one or more groups selected from halo,
 $-COOR^{35}$, $-OCH_3$, $-OCF_3$, and $-SCF_3$; and R^4 , when present, is attached to a carbon or
nitrogen atom;

20 or R^3 and R^4 are taken together to form $-phenylene-O-(CH_2)_{1-3}-$ or $-phenylene-O-$
 $CH_2-CHOH-CH_2-$;

a is 0 or 1; R^5 is selected from halo, $-CH_3$, $-CF_3$, and $-CN$;

b is 0 or an integer from 1 to 3; each R^6 is independently selected from halo, $-OH$,
 $-CH_3$, $-OCH_3$, and $-CF_3$;

25 R^7 is selected from H, $-C_{1-8}$ alkyl, $-C_{1-3}$ alkylene- C_{6-10} aryl, $-C_{1-3}$ alkylene-
 C_{1-9} heteroaryl, $-C_{3-7}$ cycloalkyl, $-[(CH_2)_2O]_{1-3}CH_3$, $-C_{1-6}$ alkylene-OC(O) R^{10} , $-C_{1-6}$ alkylene-
 $NR^{12}R^{13}$, $-C_{1-6}$ alkylene-C(O) R^{31} , $-C_{0-6}$ alkylenemorpholinyl, $-C_{1-6}$ alkylene-SO₂- C_{1-6} alkyl,



R¹⁰ is selected from -C₁₋₆alkyl, -O-C₁₋₆alkyl, -C₃₋₇cycloalkyl, -O-C₃₋₇cycloalkyl, phenyl, -O-phenyl, -NR¹²R¹³, -CH[CH(CH₃)₂]-NH₂, -CH[CH(CH₃)₂]-NHC(O)O-C₁₋₆alkyl, and -CH(NH₂)CH₂COOCH₃; and R¹² and R¹³ are independently selected from H, -C₁₋₆alkyl, and benzyl; or R¹² and R¹³ are taken together as -(CH₂)₃₋₆-, -C(O)-(CH₂)₃-, or -(CH₂)₂O(CH₂)₂-; R³¹ is selected from -O-C₁₋₆alkyl, -O-benzyl, and -NR¹²R¹³; and R³² is -C₁₋₆alkyl or -C₀₋₆alkylene-C₆₋₁₀aryl;


R⁸ is selected from H, -OH, -OC(O)R¹⁴, -CH₂COOH, -O-benzyl, -pyridyl, and -OC(S)NR¹⁵R¹⁶; R¹⁴ is selected from H, -C₁₋₆alkyl, -C₆₋₁₀aryl, -OCH₂-C₆₋₁₀aryl, -CH₂O-C₆₋₁₀aryl, and -NR¹⁵R¹⁶; and R¹⁵ and R¹⁶ are independently selected from H and -C₁₋₄alkyl;

R⁹ is selected from H, -C₁₋₆alkyl, and -C(O)-R¹⁷; and R¹⁷ is selected from H, -C₁₋₆alkyl, -C₃₋₇cycloalkyl, -C₆₋₁₀aryl, and -C₁₋₉heteroaryl;

R¹⁸ and R¹⁹ are independently selected from H, -C₁₋₆alkyl, and -O-C₃₋₇cycloalkyl, or R¹⁸ and R¹⁹ are taken together to form =O;

R²⁰ is selected from H and -C₁₋₆alkyl;

R²¹ and R³⁵ are independently selected from H, -C₁₋₆alkyl, -C₁₋₃alkylene-C₆₋₁₀aryl, -C₁₋₃alkylene-C₁₋₉heteroaryl, -C₃₋₇cycloalkyl, -[(CH₂)₂O]₁₋₃CH₃, -C₁₋₆alkylene-OC(O)R²⁵; -C₁₋₆alkylene-NR²⁷R²⁸, -C₁₋₆alkylene-C(O)R³³, -C₀₋₆alkylenemorpholinyl, -C₁₋₆alkylene-SO₂-C₁₋₆alkyl,



R²⁵ is selected from -C₁₋₆alkyl, -O-C₁₋₆alkyl, -C₃₋₇cycloalkyl, -O-C₃₋₇cycloalkyl, phenyl, -O-phenyl, -NR²⁷R²⁸, -CH[CH(CH₃)₂]-NH₂, -CH[CH(CH₃)₂]-NHC(O)O-C₁₋₆alkyl, and -CH(NH₂)CH₂COOCH₃; R²⁷ and R²⁸ are independently selected from H, -C₁₋₆alkyl, and benzyl; or R²⁷ and R²⁸ are taken together as -(CH₂)₃₋₆-, -C(O)-(CH₂)₃-, or -(CH₂)₂O(CH₂)₂-; R³³ is selected from -O-C₁₋₆alkyl, -O-benzyl, and -NR²⁷R²⁸; and R³⁴ is -C₁₋₆alkyl or -C₀₋₆alkylene-C₆₋₁₀aryl;

R²² and R²³ are independently selected from H, -C₁₋₆alkyl, -CH₂COOH, -(CH₂)₂OH; -(CH₂)₂OCH₃, -(CH₂)₂SO₂NH₂, -(CH₂)₂N(CH₃)₂, -C₀₋₁alkylene-C₃₋₇cycloalkyl, and -(CH₂)₂-imidazolyl; or R²² and R²³ are taken together to form a saturated or partially unsaturated -C₃₋₅heterocycle optionally substituted with halo, -OH, -COOH, or -CONH₂;

5 and optionally containing an oxygen atom in the ring;

R²⁴ is selected from -C₁₋₆alkyl; -C₀₋₁alkylene-O-C₁₋₆alkyl; phenyl optionally substituted with halo or -OCH₃; and -C₁₋₉heteroaryl; and

R³⁶ is selected from H, -CH(CH₃)₂, phenyl, and benzyl;

10 where each alkyl group in R¹, R³, and R⁴ is optionally substituted with 1 to 8 fluoro atoms; and;

where the methylene linker on the biphenyl is optionally substituted with one or two -C₁₋₆alkyl groups or cyclopropyl;

or a pharmaceutically acceptable salt thereof.


Another aspect of the present disclosure relates to pharmaceutical compositions
15 comprising a pharmaceutically acceptable carrier and a compound as disclosed herein. Such compositions may optionally contain other therapeutic agents. Accordingly, in yet another embodiment of the present disclosure, a pharmaceutical composition comprises a compound of the present disclosure as the first therapeutic agent, one or more secondary therapeutic agent, and a pharmaceutically acceptable carrier. Another embodiment of the
20 present disclosure relates to a combination of active agents, comprising a compound of the present disclosure and a second therapeutic agent. The compound of the present disclosure can be formulated together or separately from the additional agent(s). When formulated separately, a pharmaceutically acceptable carrier may be included with the additional agent(s). Thus, yet another embodiment of the present disclosure relates to a combination
25 of pharmaceutical compositions, the combination comprising: a first pharmaceutical composition comprising a compound of the present disclosure and a first pharmaceutically acceptable carrier; and a second pharmaceutical composition comprising a second therapeutic agent and a second pharmaceutically acceptable carrier. In another embodiment, the present disclosure relates to a kit containing such pharmaceutical
30 compositions, for example where the first and second pharmaceutical compositions are separate pharmaceutical compositions.

Compounds of the present disclosure possess NEP enzyme inhibition activity, and are therefore expected to be useful as therapeutic agents for treating patients suffering from a disease or disorder that is treated by inhibiting the NEP enzyme or by increasing the
35 levels


of its peptide substrates. Thus, one embodiment of the present disclosure relates to a method of treating patients suffering from a disease or disorder that is treated by inhibiting the NEP enzyme, comprising administering to a patient a therapeutically effective amount of a compound of the present disclosure. Another embodiment of the present disclosure relates to a method of treating hypertension, heart failure, or renal disease, comprising administering to a patient a therapeutically effective amount of a compound of the present disclosure. Still another embodiment of the present disclosure relates to a method for inhibiting a NEP enzyme in a mammal comprising administering to the mammal, a NEP enzyme-inhibiting amount of a compound of the present disclosure.

Since compounds of the present disclosure possess NEP inhibition activity, they are also useful as research tools. Accordingly, one embodiment of the present disclosure relates to a method of using a compound of the present disclosure as a research tool, the method comprising conducting a biological assay using a compound of the present disclosure. Compounds of the present disclosure can also be used to evaluate new chemical compounds. Thus another embodiment of the present disclosure relates to a method of evaluating a test compound in a biological assay, comprising: (a) conducting a biological assay with a test compound to provide a first assay value; (b) conducting the biological assay with a compound of the present disclosure to provide a second assay value; wherein step (a) is conducted either before, after or concurrently with step (b); and (c) comparing the first assay value from step (a) with the second assay value from step (b). Exemplary biological assays include a NEP enzyme inhibition assay. Still another embodiment of the present disclosure relates to a method of studying a biological system or sample comprising a NEP enzyme, the method comprising: (a) contacting the biological system or sample with a compound of the present disclosure; and (b) determining the effects caused by the compound on the biological system or sample.


Yet another embodiment of the present disclosure relates to processes and intermediates useful for preparing compounds of the present disclosure. Accordingly, another aspect of the present disclosure relates to a process of preparing compounds of formula I as disclosed herein, comprising the step of coupling a compound of formula 1 with a compound of formula 2:




to produce a compound of formula I; where P^1 is H or amino-protecting group selected from *t*-butoxycarbonyl, trityl, benzyloxycarbonyl, 9-fluorenylmethoxycarbonyl, formyl, trimethylsilyl, and *t*-butyldimethylsilyl; and where the process further comprises deprotecting the compound of formula 1 when P^1 is an amino protecting group; and where R^1 - R^6 , a, b, and X are as defined for formula I. In another aspect, there is provided a process for preparing a compound as defined herein, comprising the step of deprotecting a compound selected from:




(6),



(7),




(8), and



(9),

or a salt thereof; where R^{1P} is selected from $-O-P^3$, $-NHP^2$, and $-NH(O-P^4)$; R^{3P} is selected from $-C_{0-5}$ alkylene- $O-P^4$, $-C_{0-1}$ alkylene- $COO-P^3$, and phenyl substituted with $-O-P^4$; R^{4P} is selected from $-O-P^4$, $-C_{1-2}$ alkylene- $COO-P^3$; and phenyl or benzyl substituted with $-COO-P^3$; P^2 is an amino-protecting group selected from *t*-butoxycarbonyl, trityl, benzyloxycarbonyl, 9-fluorenylmethoxycarbonyl, formyl, trimethylsilyl, and *t*-butyldimethylsilyl; P^3 is a carboxy-protecting group selected from methyl, ethyl, *t*-butyl, benzyl, *p*-methoxybenzyl, 9-fluorenylmethyl, trimethylsilyl, *t*-butyldimethylsilyl, and diphenylmethyl; and P^4 is a hydroxyl-protecting group selected from $-C_{1-6}$ alkyl, $triC_{1-6}$ alkylsilyl, $-C_{1-6}$ alkanoyl, benzoyl, benzyl, *p*-methoxybenzyl, 9-fluorenylmethyl, and diphenylmethyl. Another embodiment of the present disclosure relates to a process of preparing a pharmaceutically acceptable salt of a compound of formula I, comprising contacting a


compound of formula I in free acid or base form with a pharmaceutically acceptable base or acid. In other embodiments, the present disclosure relates to products prepared by any of the processes described herein, as well as novel intermediates used in such process. In one embodiment of the present disclosure novel intermediates have formula 1, 6, 7, 8, or 9, or a salt thereof, as defined herein. In one aspect, there is provided an intermediate useful in the synthesis of a compound as defined herein, having formula 1:




(1),

where P^1 is H or an amino-protecting group selected from *t*-butoxycarbonyl, trityl, benzyloxycarbonyl, 9-fluorenylmethoxycarbonyl, formyl, trimethylsilyl, and *t*-butyldimethylsilyl; or a salt thereof.


In another aspect, there is provided an intermediate useful in the synthesis of a compound as defined herein, selected from:




(6),



(7),



(8), and



(9),

or a salt thereof; where R^{1P} is selected from $-O-P^3$, $-NHP^2$, and $-NH(O-P^4)$; R^{3P} is selected from $-C_{0-5}alkylene-O-P^4$, $-C_{0-1}alkylene-COO-P^3$, and phenyl substituted with $-O-P^4$; R^{4P} is selected from $-O-P^4$, $-C_{1-2}alkylene-COO-P^3$; and phenyl or benzyl substituted with $-COO-P^3$; P^2 is an amino-protecting group selected from *t*-butoxycarbonyl, trityl, benzyloxycarbonyl, 9-fluorenylmethoxycarbonyl, formyl, trimethylsilyl, and *t*-butyldimethylsilyl; P^3 is a carboxy-protecting group selected from methyl, ethyl, *t*-butyl, benzyl, *p*-methoxybenzyl, 9-fluorenylmethyl, trimethylsilyl, *t*-butyldimethylsilyl, and diphenylmethyl; and P^4 is a hydroxyl-protecting group selected from $-C_{1-6}alkyl$, $triC_{1-6}alkylsilyl$, $-C_{1-6}alkanoyl$, benzoyl, benzyl, *p*-methoxybenzyl, 9-fluorenylmethyl, and diphenylmethyl.

In another aspect, there is provided a compound as defined herein, for use in therapy.

Yet another aspect of the present disclosure relates to the use of a compound of formula I as disclosed herein or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament, especially for the manufacture of a medicament useful for treating hypertension, heart failure, or renal disease. Another embodiment of the present disclosure relates to use of a compound of the present disclosure for inhibiting a NEP enzyme in a mammal. Still another embodiment of the present disclosure relates to the use of a compound of the present disclosure as a research tool. Other embodiments of the present disclosure are disclosed herein.

DETAILED DESCRIPTION OF THE INVENTION

DEFINITIONS

When describing the compounds, compositions, methods and processes of the invention, the following terms have the following meanings unless otherwise indicated. Additionally, as used herein, the singular forms "a," "an," and "the" include the corresponding plural forms unless the context of use clearly dictates otherwise. The terms "comprising", "including," and "having" are intended to be inclusive and mean that there may be additional elements other than the listed elements. All numbers expressing quantities of ingredients, properties such as molecular weight, reaction conditions, and so forth used herein are to be understood as being modified in all instances by the term "about," unless otherwise indicated. Accordingly, the numbers set forth herein are approximations that may vary depending upon the desired properties sought to be obtained by the present invention. At least, and not as an attempt to limit the application of the doctrine of equivalents to the scope of the claims, each number should at least be construed in light of the reported significant digits and by applying ordinary rounding techniques.

The term "alkyl" means a monovalent saturated hydrocarbon group which may be linear or branched. Unless otherwise defined, such alkyl groups typically contain from 1 to 10 carbon atoms and include, for example, -C₁₋₄alkyl, -C₁₋₅alkyl, -C₂₋₅alkyl, -C₁₋₆alkyl, -C₁₋₈alkyl, and -C₁₋₁₀alkyl. Representative alkyl groups include, by way of example,
5 methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, *s*-butyl, isobutyl, *t*-butyl, *n*-pentyl, *n*-hexyl, *n*-heptyl, *n*-octyl, *n*-nonyl, *n*-decyl and the like.

When a specific number of carbon atoms is intended for a particular term used herein, the number of carbon atoms is shown preceding the term as subscript. For example, the term "-C₁₋₆alkyl" means an alkyl group having from 1 to 6 carbon atoms, and
10 the term "-C₃₋₇cycloalkyl" means a cycloalkyl group having from 3 to 7 carbon atoms, respectively, where the carbon atoms are in any acceptable configuration.

The term "alkylene" means a divalent saturated hydrocarbon group that may be linear or branched. Unless otherwise defined, such alkylene groups typically contain from 0 to 10 carbon atoms and include, for example, -C₀₋₁alkylene-, -C₀₋₆alkylene-,
15 -C₁₋₃alkylene-, and -C₁₋₆alkylene-. Representative alkylene groups include, by way of example, methylene, ethane-1,2-diyl ("ethylene"), propane-1,2-diyl, propane-1,3-diyl, butane-1,4-diyl, pentane-1,5-diyl and the like. It is understood that when the alkylene term include zero carbons such as -C₀₋₁alkylene-, such terms are intended to include the absence of carbon atoms, that is, the alkylene group is not present except for a covalent bond
20 attaching the groups separated by the alkylene term.

The term "aryl" means a monovalent aromatic hydrocarbon having a single ring (i.e., phenyl) or one or more fused rings. Fused ring systems include those that are fully unsaturated (e.g., naphthalene) as well as those that are partially unsaturated (e.g., 1,2,3,4-tetrahydronaphthalene). Unless otherwise defined, such aryl groups typically contain from
25 6 to 10 carbon ring atoms and include, for example, -C₆₋₁₀aryl. Representative aryl groups include, by way of example, phenyl and naphthalene-1-yl, naphthalene-2-yl, and the like.

The term "cycloalkyl" means a monovalent saturated carbocyclic hydrocarbon group. Unless otherwise defined, such cycloalkyl groups typically contain from 3 to 10 carbon atoms and include, for example, -C₃₋₅cycloalkyl, -C₃₋₆cycloalkyl and
30 -C₃₋₇cycloalkyl. Representative cycloalkyl groups include, by way of example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and the like.

The term "halo" means fluoro, chloro, bromo and iodo.


The term "heterocycle" is intended to include monovalent unsaturated (aromatic)

heterocycles having a single ring or two fused rings as well as monovalent saturated and partially unsaturated groups having a single ring or multiple condensed rings. The heterocycle ring can contain from 3 to 15 total ring atoms, of which 1 to 14 are ring carbon atoms, and 1 to 4 are ring heteroatoms selected from nitrogen, oxygen or sulfur. Typically, however, the heterocycle ring contains from 3 to 10 total ring atoms, of which 1 to 9 are ring carbon atoms, and 1 to 4 are ring heteroatoms. The point of attachment is at any available carbon or nitrogen ring atom. Exemplary heterocycles include, for example, -C₁₋₇heterocycle, -C₃₋₅heterocycle, -C₂₋₆heterocycle, -C₃₋₁₂heterocycle, -C₅₋₉heterocycle, -C₁₋₉heterocycle, -C₁₋₁₁heterocycle, and -C₁₋₁₄heterocycle.


10 Monovalent unsaturated heterocycles are also commonly referred to as "heteroaryl" groups. Unless otherwise defined, heteroaryl groups typically contain from 5 to 10 total ring atoms, of which 1 to 9 are ring carbon atoms, and 1 to 4 are ring heteroatoms, and include, for example, -C₁₋₉heteroaryl and -C₅₋₉heteroaryl. Representative heteroaryl groups include, by way of example, pyrrole (e.g., 3-pyrrolyl and 2H-pyrrol-3-yl), imidazole (e.g., 2-imidazolyl), furan (e.g., 2-furyl and 3-furyl), thiophene (e.g., 2-thienyl), triazole (e.g., 1,2,3-triazolyl and 1,2,4-triazolyl), pyrazole (e.g., 1H-pyrazol-3-yl), oxazole (e.g., 2-oxazolyl), isoxazole (e.g., 3-isoxazolyl), thiazole (e.g., 2-thiazolyl and 4-thiazolyl), and isothiazole (e.g., 3-isothiazolyl), pyridine (e.g., 2-pyridyl, 3-pyridyl, and 4-pyridyl), pyridylimidazole, pyridyltriazole, pyrazine, pyridazine (e.g., 3-pyridazinyl), pyrimidine (e.g., 2-pyrimidinyl), tetrazole, triazine (e.g., 1,3,5-triazinyl), indolyle (e.g., 1H-indol-2-yl, 1H-indol-4-yl and 1H-indol-5-yl), benzofuran (e.g., benzofuran-5-yl), benzothiophene (e.g., benzo/*b*/thien-2-yl and benzo/*b*/thien-5-yl), benzimidazole, benzoxazole, benzothiazole, benzotriazole, quinoline (e.g., 2-quinolyl), isoquinoline, quinazoline, quinoxaline and the like.

25 Monovalent saturated heterocycles typically contain from 3 to 10 total ring atoms, of which 2 to 9 are ring carbon atoms, and 1 to 4 are ring heteroatoms, and include, for example -C₃₋₅heterocycle. Representative monovalent saturated heterocycles include, by way of example, monovalent species of pyrrolidine, imidazolidine, pyrazolidine, piperidine, 1,4-dioxane, morpholine, thiomorpholine, piperazine, 3-pyrroline and the like.

30 In some instances, moieties may be described as being taken together to form a saturated -C₃₋₅heterocycle optionally containing an oxygen atom in the ring. Such groups include:



Monovalent partially unsaturated heterocycles typically contain from 3 to 10 total ring atoms, of which 2 to 11 are ring carbon atoms, and 1 to 3 are ring heteroatoms, and include, for example -C₃₋₅heterocycle and -C₂₋₁₂heterocycle. Representative monovalent partially unsaturated heterocycles include, by way of example, pyran, benzopyran, benzodioxole (e.g., benzo[1,3]dioxol-5-yl), tetrahydropyridazine, 2,5-dihydro-1H-pyrrole, dihydroimidazole, dihydrotriazole, dihydrooxazole, dihydroisoxazole, dihydrothiazole, dihydroisothiazole, dihydrooxadiazole, dihydrothiadiazole, tetrahydropyridazine, hexahydropyrroloquinoxaline, and dihydrooxadiazabenz[e]azulene. In some instances, moieties may be described as being taken together to form a partially unsaturated -C₃₋₅heterocycle. Such groups include:



The term "optionally substituted" means that group in question may be unsubstituted or it may be substituted one or several times, such as 1 to 3 times, or 1 to 5 times, or 1 to 8 times. For example, a phenyl group that is "optionally substituted" with halo atoms, may be unsubstituted, or it may contain 1, 2, 3, 4, or 5 halo atoms; and an alkyl group that is "optionally substituted" with fluoro atoms may be unsubstituted, or it may contain 1, 2, 3, 4, 5, 6, 7, or 8 fluoro atoms;. Similarly, a group that is "optionally substituted" with one or two -C₁₋₆alkyl groups, may be unsubstituted, or it may contain one or two -C₁₋₆alkyl groups.

As used herein, the phrase "having the formula" or "having the structure" is not intended to be limiting and is used in the same way that the term "comprising" is commonly used. For example, if one structure is depicted, it is understood that all stereoisomer and tautomer forms are encompassed, unless stated otherwise.

The term "pharmaceutically acceptable" refers to a material that is not biologically or otherwise unacceptable when used in the invention. For example, the term "pharmaceutically acceptable carrier" refers to a material that can be incorporated into a composition and administered to a patient without causing unacceptable biological effects or interacting in an unacceptable manner with other components of the composition. Such pharmaceutically acceptable materials typically have met the required standards of

toxicological and manufacturing testing, and include those materials identified as suitable inactive ingredients by the U.S. Food and Drug administration.

The term "pharmaceutically acceptable salt" means a salt prepared from a base or an acid which is acceptable for administration to a patient, such as a mammal (for example, salts having acceptable mammalian safety for a given dosage regime). However, it is understood that the salts covered by the invention are not required to be pharmaceutically acceptable salts, such as salts of intermediate compounds that are not intended for administration to a patient. Pharmaceutically acceptable salts can be derived from pharmaceutically acceptable inorganic or organic bases and from pharmaceutically acceptable inorganic or organic acids. In addition, when a compound of formula I contains both a basic moiety, such as an amine, pyridine or imidazole, and an acidic moiety such as a carboxylic acid or tetrazole, zwitterions may be formed and are included within the term "salt" as used herein. Salts derived from pharmaceutically acceptable inorganic bases include ammonium, calcium, copper, ferric, ferrous, lithium, magnesium, manganic, manganous, potassium, sodium, and zinc salts, and the like. Salts derived from pharmaceutically acceptable organic bases include salts of primary, secondary and tertiary amines, including substituted amines, cyclic amines, naturally-occurring amines and the like, such as arginine, betaine, caffeine, choline, *N,N'*-dibenzylethylenediamine, diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, *N*-ethylmorpholine, *N*-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, lysine, methylglucamine, morpholine, piperazine, piperadine, polyamine resins, procaine, purines, theobromine, triethylamine, trimethylamine, tripropylamine, tromethamine and the like. Salts derived from pharmaceutically acceptable inorganic acids include salts of boric, carbonic, hydrohalic (hydrobromic, hydrochloric, hydrofluoric or hydroiodic), nitric, phosphoric, sulfamic and sulfuric acids. Salts derived from pharmaceutically acceptable organic acids include salts of aliphatic hydroxyl acids (for example, citric, gluconic, glycolic, lactic, lactobionic, malic, and tartaric acids), aliphatic monocarboxylic acids (for example, acetic, butyric, formic, propionic and trifluoroacetic acids), amino acids (for example, aspartic and glutamic acids), aromatic carboxylic acids (for example, benzoic, *p*-chlorobenzoic, diphenylacetic, gentisic, hippuric, and triphenylacetic acids), aromatic hydroxyl acids (for example, *o*-hydroxybenzoic, *p*-hydroxybenzoic, 1-hydroxynaphthalene-2-carboxylic and 3-hydroxynaphthalene-2-carboxylic acids), ascorbic, dicarboxylic acids (for example,

fumaric, maleic, oxalic and succinic acids), glucuronic, mandelic, mucic, nicotinic, orotic, pamoic, pantothenic, sulfonic acids (for example, benzenesulfonic, camphosulfonic, edisyllic, ethanesulfonic, isethionic, methanesulfonic, naphthalenesulfonic, naphthalene-1,5-disulfonic, naphthalene-2,6-disulfonic and p-toluenesulfonic acids), xinafoic acid, and
5 the like.

As used herein, the term "prodrug" is intended to mean an inactive (or significantly less active) precursor of a drug that is converted into its active form in the body under physiological conditions, for example, by normal metabolic processes. Such compounds may not possess pharmacological activity at NEP, but may be administered orally or
10 parenterally and thereafter metabolized in the body to form compounds that are pharmacologically active at NEP. Exemplary prodrugs include esters such as C₁₋₆alkylesters and aryl-C₁₋₆alkylesters. In one embodiment, the active compound has a free carboxyl and the prodrug is an ester derivative thereof, i.e., the prodrug is an ester such as -C(O)OCH₂CH₃. Such ester prodrugs are then converted by solvolysis or under
15 physiological conditions to be the free carboxyl compound. The term is also intended to include certain protected derivatives of compounds of formula I that may be made prior to a final deprotection stage. Thus, all protected derivatives and prodrugs of compounds formula I are included within the scope of the invention.


The term "therapeutically effective amount" means an amount sufficient to effect
20 treatment when administered to a patient in need thereof, that is, the amount of drug needed to obtain the desired therapeutic effect. For example, a therapeutically effective amount for treating hypertension is an amount of compound needed to, for example, reduce, suppress, eliminate, or prevent the symptoms of hypertension, or to treat the underlying cause of hypertension. In one embodiment, a therapeutically effective amount
25 is that amount of drug needed to reduce blood pressure or the amount of drug needed to maintain normal blood pressure. On the other hand, the term "effective amount" means an amount sufficient to obtain a desired result, which may not necessarily be a therapeutic result. For example, when studying a system comprising a NEP enzyme, an "effective amount" may be the amount needed to inhibit the enzyme.

30 The term "treating" or "treatment" as used herein means the treating or treatment of a disease or medical condition (such as hypertension) in a patient, such as a mammal (particularly a human) that includes one or more of the following: (a) preventing the disease or medical condition from occurring, i.e., preventing the reoccurrence of the

disease or medical condition or prophylactic treatment of a patient that is pre-disposed to the disease or medical condition; (b) ameliorating the disease or medical condition, i.e., eliminating or causing regression of the disease or medical condition in a patient; (c) suppressing the disease or medical condition, i.e., slowing or arresting the development of the disease or medical condition in a patient; or (d) alleviating the symptoms of the disease or medical condition in a patient. For example, the term "treating hypertension" would include preventing hypertension from occurring, ameliorating hypertension, suppressing hypertension, and alleviating the symptoms of hypertension (for example, lowering blood pressure). The term "patient" is intended to include those mammals, such as humans, that are in need of treatment or disease prevention or that are presently being treated for disease prevention or treatment of a specific disease or medical condition, as well as test subjects in which compounds of the invention are being evaluated or being used in an assay, for example an animal model.

All other terms used herein are intended to have their ordinary meaning as understood by those of ordinary skill in the art to which they pertain.

In one aspect, the invention relates to compounds of formula I:



or a pharmaceutically acceptable salt thereof.


As used herein, the term "compound of the invention" includes all compounds encompassed by formula I such as the species embodied in formulas Ia-d, as well as the compounds encompassed by formulas II, and III, and species thereof. In addition, the compounds of the invention may also contain several basic or acidic groups (for example, amino or carboxyl groups) and therefore, such compounds can exist as a free base, free acid, or in various salt forms. All such salt forms are included within the scope of the invention. Furthermore, the compounds of the invention may also exist as prodrugs. Accordingly, those skilled in the art will recognize that reference to a compound herein, for example, reference to a "compound of the invention" or a "compound of formula I"

includes a compound of formula I as well as pharmaceutically acceptable salts and prodrugs of that compound unless otherwise indicated. Further, the term "or a pharmaceutically acceptable salt and/or prodrug thereof" is intended to include all permutations of salts and prodrugs, such as a pharmaceutically acceptable salt of a prodrug.


- 5 Furthermore, solvates of compounds of formula I are included within the scope of this invention.

The compounds of formula I may contain one or more chiral centers and therefore, these compounds may be prepared and used in various stereoisomeric forms. Accordingly, the invention also relates to racemic mixtures, pure stereoisomers (e.g., enantiomers and
10 diastereoisomers), stereoisomer-enriched mixtures, and the like unless otherwise indicated. When a chemical structure is depicted herein without any stereochemistry, it is understood that all possible stereoisomers are encompassed by such structure. Thus, for example, the terms "compound of formula I," "compounds of formula II," and so forth, are intended to include all possible stereoisomers of the compound. Similarly, when a particular
15 stereoisomer is shown or named herein, it will be understood by those skilled in the art that minor amounts of other stereoisomers may be present in the compositions of the invention unless otherwise indicated, provided that the utility of the composition as a whole is not eliminated by the presence of such other isomers. Individual stereoisomers may be obtained by numerous methods that are well known in the art, including chiral
20 chromatography using a suitable chiral stationary phase or support, or by chemically converting them into diastereoisomers, separating the diastereoisomers by conventional means such as chromatography or recrystallization, then regenerating the original stereoisomer.

Additionally, where applicable, all *cis-trans* or *E/Z* isomers (geometric isomers),
25 tautomeric forms and topoisomeric forms of the compounds of the invention are included within the scope of the invention unless otherwise specified. For example, if X is depicted as (R^4 being hydrogen):




it is understood that the compound may also exist in a tautomeric form such as:




30

More specifically, compounds of formula I can contain at least two chiral centers indicated by the symbols * and ** in the following formula:




In one stereoisomer, both carbon atoms identified by the * and ** symbols have the (R) configuration. This embodiment of the invention is shown in formula Ia:



In this embodiment, compounds have the (R,R) configuration at the * and ** carbon atoms or are enriched in a stereoisomeric form having the (R,R) configuration at these carbon atoms.


In another stereoisomer, both carbon atoms identified by the * and ** symbols have the (S) configuration. This embodiment of the invention is shown in formula Ib:



In this embodiment, compounds have the (S,S) configuration at the * and ** carbon atoms or are enriched in a stereoisomeric form having the (S,S) configuration at these carbon atoms.

In yet another stereoisomer, the carbon atom identified by the symbol * has the (S) configuration and the carbon atom identified by the symbol ** has the (R) configuration.


This embodiment of the invention is shown in formula Ic:



In this embodiment, compounds have the (*S,R*) configuration at the * and ** carbon atoms or are enriched in a stereoisomeric form having the (*S,R*) configuration at these carbon atoms.

In still another stereoisomer, the carbon atom identified by the symbol * has the (*R*) configuration and the carbon atom identified by the symbol ** has the (*S*) configuration.

This embodiment of the invention is shown in formula Id:



In this embodiment, compounds have the (*R,S*) configuration at the * and ** carbon atoms or are enriched in a stereoisomeric form having the (*R,S*) configuration at these carbon atoms.

Compounds of formula Ia and Ib are enantiomers and therefore, in separate aspects, this invention relates to each individual enantiomer (i.e., Ia or Ib), a racemic mixture of Ia and Ib, or an enantiomer-enriched mixture of Ia and Ib comprising predominately Ia or predominately Ib. Similarly, compounds of formula Ic and Id are enantiomers and therefore, in separate aspects, this invention relates to each individual enantiomer (i.e., Ic or Id), a racemic mixture of Ic and Id, or an enantiomer-enriched mixture of Ic and Id comprising predominately Ic or predominately Id.

In some embodiments, in order to optimize the therapeutic activity of the compounds of the invention, e.g., to treat hypertension, it may be desirable that the carbon atoms identified by the * and ** symbols have a particular (*R,R*), (*S,S*), (*S,R*), or (*R,S*)

configuration or are enriched in a stereoisomeric form having such configuration. For example, in one embodiment, the compounds of the invention have the (S,R) configuration of formula Ic or are enriched in a stereoisomeric form having the (S,R) configuration, and in another embodiment, the compounds of the invention have the (R,S) configuration of formula Id, or are enriched in a stereoisomeric form having the (R,S) configuration. In other embodiments, the compounds of the invention are present as racemic mixtures, for example as a mixture of enantiomers of formula Ia and Ib, or as a mixture of enantiomers of formula Ic and Id.


The compounds of the invention, as well as those compounds used in their synthesis, may also include isotopically-labeled compounds, that is, where one or more atoms have been enriched with atoms having an atomic mass different from the atomic mass predominately found in nature. Examples of isotopes that may be incorporated into the compounds of formula I, for example, include, but are not limited to, ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O , ^{35}S , ^{36}Cl , and ^{18}F . Of particular interest are compounds of formula I enriched in tritium or carbon-14 which can be used, for example, in tissue distribution studies; compounds of formula I enriched in deuterium especially at a site of metabolism resulting, for example, in compounds having greater metabolic stability; and compounds of formula I enriched in a positron emitting isotope, such as ^{11}C , ^{18}F , ^{15}O and ^{13}N , which can be used, for example, in Positron Emission Topography (PET) studies.

The nomenclature used herein to name the compounds of the invention is illustrated in the Examples herein. This nomenclature has been derived using the commercially available AutoNom software (MDL, San Leandro, California).

REPRESENTATIVE EMBODIMENTS

The following substituents and values are intended to provide representative examples of various aspects and embodiments of the invention. These representative values are intended to further define and illustrate such aspects and embodiments and are not intended to exclude other embodiments or to limit the scope of the invention. In this regard, the representation that a particular value or substituent is preferred is not intended in any way to exclude other values or substituents from the invention unless specifically indicated.

In one aspect, this invention relates to compounds of formula I:



(I)

R^1 is selected from $-OR^7$ and $-NR^8R^9$. The R^7 moiety is selected from:

H;

$-C_{1-8}$ alkyl, e.g., $-CH_3$, $-CH_2CH_3$, $-(CH_2)_2CH_3$, $-CH(CH_3)_2$, $-CH_2CH(CH_3)_2$,

5 $-(CH_2)_3CH_3$, $-(CH_2)_4CH_3$, $-(CH_2)_2CH(CH_3)_2$, $-(CH_2)_5CH_3$, and $-(CH_2)_6CH_3$;

$-C_{1-3}$ alkylene- C_{6-10} aryl, e.g., benzyl;

$-C_{1-3}$ alkylene- C_{1-9} heteroaryl, e.g., $-CH_2$ -pyridinyl and $-(CH_2)_2$ -pyridinyl;

$-C_{3-7}$ cycloalkyl, e.g., cyclopentyl;

$-[(CH_2)_2O]_{1-3}CH_3$, e.g., $-(CH_2)_2OCH_3$ and $-[(CH_2)_2O]_2CH_3$;

10 $-C_{1-6}$ alkylene- $OC(O)R^{10}$, e.g., $-CH_2OC(O)CH_3$, $-CH_2OC(O)CH_2CH_3$,

$-CH_2OC(O)(CH_2)_2CH_3$, $-CH_2CH(CH_3)OC(O)CH_2CH_3$, $-CH_2OC(O)OCH_3$,

$-CH_2OC(O)OCH_2CH_3$, $-CH(CH_3)OC(O)OCH_2CH_3$, $-CH(CH_3)OC(O)O-CH(CH_3)_2$,


$-CH_2CH(CH_3)OC(O)$ -cyclopentyl, $-CH_2OC(O)O$ -cyclopropyl, $-CH(CH_3)-OC(O)-O$ -

cyclohexyl, $-CH_2OC(O)O$ -cyclopentyl, $-CH_2CH(CH_3)OC(O)$ -phenyl, $-CH_2OC(O)O$ -

15 phenyl, $-CH_2OC(O)-CH[CH(CH_3)_2]-NH_2$, $-CH_2OC(O)-CH[CH(CH_3)_2]-NHC(O)OCH_3$,


and $-CH(CH_3)OC(O)-CH(NH_2)CH_2COOCH_3$;

$-C_{1-6}$ alkylene- $NR^{12}R^{13}$, e.g., $-(CH_2)_2-N(CH_3)_2$,




$-C_{1-6}$ alkylene- $C(O)R^{31}$, e.g., $-CH_2C(O)OCH_3$, $-CH_2C(O)O$ -benzyl, $-CH_2C(O)-$




20 $N(CH_3)_2$, and



$-C_{0-6}$ alkylenemorpholinyl, e.g., $-(CH_2)_2$ -morpholinyl and $-(CH_2)_3$ -morpholinyl;




-C₁₋₆alkylene-SO₂-C₁₋₆alkyl, e.g., -(CH₂)₂SO₂CH₃;



5

; and



The R¹⁰ moiety is selected from:

-C₁₋₆alkyl, e.g., -CH₃ and -CH₂CH₃;

-O-C₁₋₆alkyl, e.g., -OCH₃, -O-CH₂CH₃, and -O-CH(CH₃)₂;

10 -C₃₋₇cycloalkyl, e.g., cyclopentyl);

-O-C₃₋₇cycloalkyl, e.g., -O-cyclopropyl, -O-cyclohexyl, and -O-cyclopentyl;

phenyl;

-O-phenyl;

-NR¹²R¹³;

15 -CH[CH(CH₃)₂]-NH₂;


-CH[CH(CH₃)₂]-NHC(O)O-C₁₋₆alkyl, e.g., -CH[CH(CH₃)₂]-NHC(O)OCH₃; and

-CH(NH₂)CH₂COOCH₃.


The R¹² and R¹³ moieties are independently selected from H, -C₁₋₆alkyl (e.g., CH₃), and

benzyl. Alternately, the R¹² and R¹³ moieties can be taken together as -(CH₂)₃₋₆-, -C(O)-

20 (CH₂)₃-, or -(CH₂)₂O(CH₂)₂-, for example to form a group such as:



The R^{31} moiety is selected from $-O-C_{1-6}$ alkyl, e.g., $-OCH_3$, $-O$ -benzyl, and $-NR^{12}R^{13}$, e.g., $-N(CH_3)_2$, and



5 The R^{32} moiety is $-C_{1-6}$ alkyl (e.g., $-CH_3$ and $-C(CH_3)_3$) or $-C_{0-6}$ alkylene- C_{6-10} aryl.

The R^8 moiety is selected from:

H;

$-OH$;

$-OC(O)R^{14}$, e.g., $-OC(O)CH_3$, $-OC(O)$ -phenyl, $-OC(O)-OCH_2$ -phenyl, $-OC(O)-$

10 CH_2O -phenyl, $-OC(O)(NH_2)$, and $-OC(O)[N(CH_3)_2]$;

$-CH_2COOH$;

$-O$ -benzyl;

pyridyl; and

$-OC(S)NR^{15}R^{16}$, e.g., $-OC(S)NH_2$ and $-OC(S)N(CH_3)_2$.

15 The R^{14} moiety is selected from:

H;

$-C_{1-6}$ alkyl, e.g., $-CH_3$;

$-C_{6-10}$ aryl, e.g., phenyl;

$-OCH_2-C_{6-10}$ aryl, e.g., $-OCH_2$ -phenyl;

20 $-CH_2O-C_{6-10}$ aryl, e.g., $-CH_2O$ -phenyl; and

$-NR^{15}R^{16}$, e.g., $-NH_2$ and $N(CH_3)_2$.

The R^{15} and R^{16} moieties are independently selected from H and $-C_{1-4}$ alkyl.


The R^9 is moiety selected from H, $-C_{1-6}$ alkyl (e.g., $-CH_3$), and $-C(O)R^{17}$ (e.g., $-C(O)H$). The R^{17} moiety is selected from H, $-C_{1-6}$ alkyl (e.g., $-CH_2CH_3$), $-C_{3-7}$ cycloalkyl

25 (e.g., cyclopropyl), $-C_{6-10}$ aryl (e.g., phenyl), and $-C_{1-9}$ heteroaryl (e.g., pyridine).

In addition, each alkyl group in R^1 is optionally substituted with 1 to 8 fluoro atoms. For example, when R^1 is $-OR^7$ and R^7 is $-C_{1-8}$ alkyl, R^1 can also be a group such as $-OCH(CH_3)CF_3$, $-OCH_2CF_2CF_3$, $-OCH(CF_3)_2$, $-O(CH_2)_2CF_3$, $-OCH(CH_2F)_2$,

-OC(CF₃)₂CH₃, and -OCH(CH₃)CF₂CF₃.


In one embodiment, R¹ is selected from -OR⁷, where R⁷ is selected from H, -C₁₋₃alkyl, -C₁₋₆alkylene-OC(O)R¹⁰, -C₀₋₆alkylenemorpholinyl, -C₁₋₆alkylene-SO₂-C₁₋₆alkyl, and




where R¹⁰ is -O-C₃₋₇cycloalkyl; and -NR⁸R⁹, where R⁸ is H and R⁹ is H. In another embodiment these compounds have formula III.

In one embodiment, R¹ is selected from -OR⁷ and -NR⁸R⁹, where R⁷ is H, R⁸ is H or -OH, and R⁹ is H. In another embodiment these compounds have formula III.


In another embodiment, R¹ is -OR⁷, where R⁷ is selected from -C₁₋₃alkyl, -C₁₋₃alkylene-C₆₋₁₀aryl, -C₁₋₃alkylene-C₁₋₉heteroaryl, -C₃₋₇cycloalkyl, -[(CH₂)₂O]₁₋₃CH₃, -C₁₋₆alkylene-OC(O)R¹⁰, -C₁₋₆alkylene-NR¹²R¹³, -C₁₋₆alkylene-C(O)R³¹, -C₀₋₆alkylenemorpholinyl; -C₁₋₆alkylene-SO₂-C₁₋₆alkyl;




In yet another embodiment, R¹ is -NR⁸R⁹; where R⁸ is selected from -OC(O)R¹⁴, -CH₂COOH, -O-benzyl, pyridyl, and -OC(S)NR¹⁵R¹⁶; and R⁹ is H. In yet another embodiment, R¹ is -NR⁸R⁹, where R⁸ is H or -OH; and R⁹ is -C₁₋₆alkyl or -C(O)R¹⁷. In yet another embodiment, R¹ is -NR⁸R⁹, where R⁸ is selected from -OC(O)R¹⁴, -CH₂COOH, -O-benzyl, pyridyl, and -OC(S)NR¹⁵R¹⁶; and R⁹ is -C₁₋₆alkyl or -C(O)R¹⁷. In another embodiment these compounds have formula III. In one aspect of the invention, these compounds may find particular utility as prodrugs or as intermediates in the synthetic procedures described herein. For example, in one embodiment, R¹ is -OR⁷ and R⁷ is -C₁₋₆alkylene-OC(O)R¹⁰, such as -O-CH(CH₃)OC(O)-O-cyclohexyl:




making the compound a cilexetil ester; or R¹ is -OR⁷ and R⁷ is -C₀₋₆alkylenemorpholinyl such as -O-(CH₂)₂-morpholinyl:



making the compound a 2-morpholinoethyl or mofetil ester; or R^1 is $-OR^7$ and R^7 is




such as -O-CH₂-5-methyl-[1,3]dioxol-2-one:




making the compound a medoxomil ester.

R^2 is H or $-P(O)(OH)_2$. R^2 can also be taken together with R^7 to form $-CR^{18}R^{19}$ - or taken together with R^8 to form $-C(O)-$. R^{18} and R^{19} are independently selected from H, $-C_{1-6}$ alkyl, and $-O-C_{3-7}$ cycloalkyl, or R^{18} and R^{19} may be taken together to form $=O$. In one embodiment, R^2 is H. In another embodiment this compound has formula III.


When R^2 is taken together with R^7 to form $-CR^{18}R^{19}$ -, this embodiment can be depicted as:



and when R^{18} and R^{19} are taken together to form $=O$, this embodiment can be depicted as:



When R^2 is taken together with R^8 to form $-C(O)-$, this embodiment can be depicted as:




In one aspect of the invention, these compounds may find particular utility as prodrugs or as intermediates in the synthetic procedures described herein. In another embodiment


these compounds have formula III. Compounds where R^2 is $-P(O)(OH)_2$ may also find utility as prodrugs.

The "X" moiety is a $-C_{1-9}$ heteroaryl, and the point of attachment is at any available carbon or nitrogen ring atom. Note that in some embodiments, R^3 and/or R^4 may be absent. When present, R^3 is on any available carbon atom. When present, R^4 is on any available carbon atom or nitrogen atom. Exemplary $-C_{1-9}$ heteroaryl rings include, by way of illustration and not limitation:


pyrazole rings such as:




specific examples of which include:




imidazole rings such as:




specific examples of which include:




triazole rings, including 1,2,3-triazole such as:




as well as 1,2,4-triazole such as:




benzotriazole rings such as:




specific examples of which include:




furan rings:




specific examples of which include:




pyrrole rings:




specific examples of which include:




tetrazole rings such as:



pyrazine rings:




a specific example of which includes:




5

thiophene rings:




specific examples of which include:




10


oxazole rings:



specific examples of which include:




isoxazole rings:




15


specific examples of which include:




thiazole rings:




specific examples of which include:




isothiazole rings:




oxadiazole rings, including [1,2,4]oxadiazole such as:




10 as well as [1,2,3]oxadiazole such as:



and [1,3,4]oxadiazole:




thiadiazole rings, including [1,2,4]thiadiazole such as:




15


as well as [1,2,3]thiadiazole such as:




and [1,3,4]thiadiazole:




pyridazine rings:




5 pyridine rings:



specific examples of which include:




pyrimidine rings:




10

specific examples of which include:




pyran rings such as




15


benzimidazole rings such as:




specific examples of which include:




benzoxazole rings such as:




specific examples of which include:




benzothiazole rings such as:




specific examples of which include:




pyridylimidazole rings such as :




a specific example of which includes:



pyridyltriazole rings such as:













specific examples of which include:




In one particular embodiment, X is selected from pyrazole, imidazole, triazole,
 5 benzotriazole, furan, pyrrole, tetrazole, pyrazine, thiophene, oxazole, isoxazole, thiazole,
 isothiazole, oxadiazole, thiadiazole, pyridazine, pyridine, pyrimidine, pyran,
 benzimidazole, benzoxazole, benzothiazole, pyridylimidazole, and pyridyltriazole.

It is understood that some -C₁₋₉heteroaryl rings can exist in a tautomeric form, and
 that such tautomeric forms are part of the invention and are encompassed by the term
 10 "heteroaryl." Therefore, if a compound is depicted with a -C₁₋₉heteroaryl ring, it is
 understood that the compound can also exist in a tautomeric form and vice versa, and that
 both forms are covered by the invention.


-C ₁₋₉ heteroaryl ring	exemplary ring	exemplary tautomer(s)
pyrazole		
imidazole		
triazole		
oxazole		

-C ₁₋₉ heteroaryl ring	exemplary ring	exemplary tautomer(s)
thiazole		
isothiazole		
oxadiazole		
thiadiazole		
pyridazine		


In one particular embodiment, X is selected from pyrazole, triazole, benzotriazole, furan, tetrazole, pyrazine, thiophene, oxazole, isoxazole, thiazole, oxadiazole, pyridazine, pyridine, pyrimidine, benzoxazole, pyridylimidazole, and pyridyltriazole. In another
 5 embodiment, X is selected from pyrazole, triazole, benzotriazole, isoxazole, pyridazine, pyrimidine, and pyridyltriazole. In still another embodiment X is an isoxazole, and in one specific embodiment, have formula III:



where R¹-R⁶, a, and b are as defined for formula I. In yet another embodiment, the
 10 compounds of the invention have formula IIIa:



where R^1 , R^5 , R^6 , a , and b , are as defined for formula I. In still another embodiment, the compounds of the invention have formula IIIb:



5 where R^1 , R^5 , R^6 , a , and b , are as defined for formula I.

The R^3 moiety can be absent. When present, R^3 is attached to a carbon atom in the "X" group, and is selected from:

- H;
- halo, e.g., chloro and fluoro;
- 10 -C₀₋₅alkylene-OH, e.g., -OH, -CH₂OH, -CH(OH)CH₃, and -C(CH₃)₂-OH;
- NH₂;
- C₁₋₆alkyl, e.g., -CH₃, -(CH₂)₂CH₃, -CH(CH₃)₂, and -(CH₂)₃-CH₃;
- CF₃;
- C₃₋₇cycloalkyl, e.g., cyclopropyl and cyclohexyl;
- 15 -C₀₋₂alkylene-O-C₁₋₆alkyl, e.g., -OCH₃, -OCH₂CH₃, -CH₂-OCH₃, and -(CH₂)₂-OCH₃;
- C(O)R²⁰, e.g., -C(O)H and -C(O)CH₃;
- C₀₋₁alkylene-COOR²¹, e.g., -COOH, -CH₂-COOH, -C(O)O-CH₂CH₃, -C(O)O-(CH₂)₂OCH₃, -C(O)O-CH₂OC(O)CH₃, -CH₂-C(O)O-CH₂OC(O)CH₃, -C(O)O-CH₂OC(O)O-CH₃, -CH₂-C(O)O-CH₂OC(O)O-CH₃, -C(O)O-CH(CH₃)OC(O)O-CH₂CH₃, -C(O)O-CH(CH₃)OC(O)O-CH(CH₃)₂, -C(O)O-CH₂CH(CH₃)OC(O)-cyclopentyl, -C(O)O-CH₂OC(O)O-cyclopropyl, -C(O)O-CH(CH₃)-OC(O)-O-cyclohexyl, -C(O)O-CH₂OC(O)O-cyclopentyl, -C(O)O-CH₂CH(CH₃)OC(O)-phenyl, -C(O)O-CH₂OC(O)O-phenyl, -C(O)O-CH₂-pyridine, -C(O)O-CH₂-pyrrolidine, -C(O)O-(CH₂)₂-morpholinyl, -C(O)O-(CH₂)₃-


- morpholinyl, and $-\text{C}(\text{O})\text{O}-(\text{CH}_2)_2-\text{SO}_2-\text{CH}_3$;
- $-\text{C}(\text{O})\text{NR}^{22}\text{R}^{23}$, e.g., $-\text{C}(\text{O})\text{NH}_2$, $-\text{C}(\text{O})\text{NHCH}_3$, $-\text{C}(\text{O})\text{N}(\text{CH}_3)_2$, $-\text{C}(\text{O})\text{NH}-(\text{CH}_2)_2\text{CH}_3$, $-\text{C}(\text{O})\text{NH}-\text{CH}_2\text{COOH}$, $-\text{C}(\text{O})\text{NH}-(\text{CH}_2)_2-\text{OH}$, $-\text{C}(\text{O})\text{NH}-(\text{CH}_2)_2-\text{N}(\text{CH}_3)_2$, $-\text{C}(\text{O})\text{NH}$ -cyclopropyl, $-\text{C}(\text{O})\text{NH}-(\text{CH}_2)_2$ -imidazolyl, $-\text{C}(\text{O})\text{N}(\text{CH}_3)-\text{CH}_2\text{CH}(\text{CH}_3)_2$, and
- 5 $-\text{C}(\text{O})\text{N}(\text{CH}_3)[(\text{CH}_2)_2\text{OCH}_3]$;
- $-\text{NHC}(\text{O})\text{R}^{24}$, e.g., $-\text{NHC}(\text{O})-\text{CH}_2\text{CH}_3$, $-\text{NHC}(\text{O})-(\text{CH}_2)_3\text{CH}_3$, $-\text{NHC}(\text{O})\text{O}-\text{CH}_2\text{CH}_3$, $-\text{NHC}(\text{O})-\text{CH}_2-\text{OCH}_3$, $-\text{NHC}(\text{O})$ -2-methoxyphenyl, $-\text{NHC}(\text{O})$ -2-chlorophenyl, and $-\text{NHC}(\text{O})$ -2-pyridine;
- $=\text{O}$;
- 10 $-\text{NO}_2$;
- $-\text{C}(\text{CH}_3)=\text{N}(\text{OH})$;
- phenyl optionally substituted with one or two groups independently selected from halo, $-\text{OH}$, $-\text{CF}_3$, $-\text{OCH}_3$, $-\text{NHC}(\text{O})\text{CH}_3$, and phenyl (e.g., phenyl, 2-chlorophenyl, 2-fluorophenyl, 2-hydroxyphenyl, 2-trifluoromethylphenyl, 2-methoxyphenyl, 3-
- 15 chlorophenyl, 3-fluorophenyl, 3-methoxyphenyl, 3- $-\text{NHC}(\text{O})\text{CH}_3$ -phenyl, 4-chlorophenyl, 4-fluorophenyl, 4-methoxyphenyl, 4-biphenyl, 2,5-dichlorophenyl, 2,5-dimethoxyphenyl, 2,4-dichlorophenyl, 2-methoxy, 5-fluorophenyl, and 3,4-dichlorophenyl);
- naphthalenyl;
- pyridinyl;
- 20 pyrazinyl;
- pyrazolyl optionally substituted with methyl;
- thiophenyl optionally substituted with methyl or halo (e.g., chloro);
- furanyl; and
- $-\text{CH}_2$ -morpholinyl.
- 25 The R^{20} moiety is selected from H and $-\text{C}_{1-6}$ alkyl (e.g., $-\text{CH}_3$). The R^{21} moiety is selected from:
- H;
- $-\text{C}_{1-6}$ alkyl, e.g., $-\text{CH}_3$ and $-\text{CH}_2\text{CH}_3$;
- $-\text{C}_{1-3}$ alkylene- C_{6-10} aryl;
- 30 $-\text{C}_{1-3}$ alkylene- C_{1-9} heteroaryl, e.g., $-\text{CH}_2$ -pyridine;
- $-\text{C}_{3-7}$ cycloalkyl;
- $-\text{[(CH}_2)_2\text{O}]_{1-3}\text{CH}_3$, e.g., $-(\text{CH}_2)_2\text{OCH}_3$;
- $-\text{C}_{1-6}$ alkylene- $\text{OC}(\text{O})\text{R}^{25}$, e.g., $-\text{CH}_2\text{OC}(\text{O})\text{CH}_3$, $-\text{CH}_2\text{OC}(\text{O})\text{O}-\text{CH}_3$, $-\text{CH}_2\text{OC}(\text{O})\text{O}-$

CH₃, -CH(CH₃)OC(O)O-CH₂CH₃, -CH(CH₃)OC(O)O-CH(CH₃)₂, -CH₂CH(CH₃)OC(O)-cyclopentyl, -CH₂OC(O)O-cyclopropyl, -CH(CH₃)-OC(O)-O-cyclohexyl, -CH₂OC(O)O-cyclopentyl, -CH₂CH(CH₃)OC(O)-phenyl, and -CH₂OC(O)O-phenyl;


-C₁₋₆alkylene-NR²⁷R²⁸, e.g., -CH₂-pyrrolidine;

5 -C₁₋₆alkylene-C(O)R³³;





-C₀₋₆alkylenemorpholinyl, e.g., -(CH₂)₂-morpholinyl and -(CH₂)₃-morpholinyl:



-C₁₋₆alkylene-SO₂-C₁₋₆alkyl, e.g., -(CH₂)₂-SO₂-CH₃;



10



The R²² and R²³ moieties are independently selected from:

15

H;

-C₁₋₆alkyl, e.g., -CH₃ and -(CH₂)₂CH₃;




-CH₂COOH;

-(CH₂)₂OH;

-(CH₂)₂OCH₃;


20

-(CH₂)₂SO₂NH₂;




R^{22} and R^{23} may also be taken together to form a saturated or partially unsaturated


- 5 $-\text{C}_{3-5}$ heterocycle optionally substituted with halo, $-\text{OH}$, $-\text{COOH}$, or $-\text{CONH}_2$, and optionally containing an oxygen atom in the ring. Saturated $-\text{C}_{3-5}$ heterocycles include azetidine, pyrrolidine, piperidine and morpholine, such that exemplary R^3 groups include:



- 10 Partially unsaturated $-\text{C}_{3-5}$ heterocycles include 2,5-dihydro-1H-pyrrole, such that exemplary R^3 groups include:




The R^{24} moiety is selected from:






- 15 $-\text{C}_{0-1}\text{alkylene}-\text{O}-\text{C}_{1-6}\text{alkyl, e.g., } -\text{O}-\text{CH}_2\text{CH}_3 \text{ and } -\text{CH}_2-\text{OCH}_3;$

phenyl optionally substituted with halo or $-\text{OCH}_3$, e.g., -2chlorophenyl or -2-methoxyphenyl; and



R^{25} is selected from:

- 20 $-\text{C}_{1-6}\text{alkyl, e.g., } -\text{CH}_3, -\text{CH}_2\text{CH}_3, \text{ and } -(\text{CH}_2)_3\text{CH}_3;$



- 25 $-\text{O-phenyl;}$

-NR²⁷R²⁸;
 -CH[CH(CH₃)₂]-NH₂;
 -CH[CH(CH₃)₂]-NHC(O)O-C₁₋₆alkyl, e.g., -CH[CH(CH₃)₂]-NHC(O)OCH₃; and
 -CH(NH₂)CH₂COOCH₃.

- 5 R²⁷ and R²⁸ are independently selected from H, -C₁₋₆alkyl, and benzyl, or R²⁷ and R²⁸ are taken together as -(CH₂)₃₋₆-, -C(O)-(CH₂)₃-, or -(CH₂)₂O(CH₂)₂-; R³³ is selected from -O-C₁₋₆alkyl, -O-benzyl, and -NR²⁷R²⁸; and R³⁴ is -C₁₋₆alkyl (e.g., -CH₃ and -C(CH₃)₃) or -C₀₋₆alkylene-C₆₋₁₀aryl.


- In addition, each alkyl group in R³ is optionally substituted with 1 to 8 fluoro
 10 atoms. For example, when R³ is -C₀₋₁alkylene-COOR²¹ and R²¹ is -C₁₋₆alkyl, R³ can also be a group such as -COOCH(CH₃)CF₃, -COOCH₂CF₂CF₃, -COOCH(CF₃)₂,
 -COO(CH₂)₂CF₃, -COOCH(CH₂F)₂, -COOC(CF₃)₂CH₃, and -COOCH(CH₃)CF₂CF₃.

- In one embodiment, R³ is absent or is selected from H; halo; -C₀₋₅alkylene-OH;
 -NH₂; -C₁₋₆alkyl; -CF₃; -C₃₋₇cycloalkyl; -C₀₋₂alkylene-O-C₁₋₆alkyl; -C(O)R²⁰;
 15 -C₀₋₁alkylene-COOR²¹; -C(O)NR²²R²³; -NHC(O)R²⁴; =O; -C(CH₃)=N(OH); phenyl
 optionally substituted with one or two groups independently selected from halo, -OH, -CF₃,
 -OCH₃, -NHC(O)CH₃, and phenyl; naphthalenyl; pyridinyl; pyrazinyl; pyrazolyl
 substituted with methyl; thiophenyl substituted with methyl or halo; furanyl; and -CH₂-
 morpholinyl; R²⁰ is -C₁₋₆alkyl; R²¹ is selected from H and -C₁₋₆alkyl; R²² is selected from H
 20 and -C₁₋₆alkyl; and R²³ is selected from H, -C₁₋₆alkyl, -CH₂COOH, -(CH₂)₂OH,
 -(CH₂)₂OCH₃, -(CH₂)₂N(CH₃)₂, -C₃₋₇cycloalkyl, and -(CH₂)₂-imidazolyl; or R²² and R²³ are
 taken together to form azetidine, pyrrolidine, piperidine or morpholine, all optionally
 substituted with -OH or -CONH₂; and R²⁴ is selected from -C₁₋₆alkyl; -C₀₋₁alkylene-O-
 C₁₋₆alkyl; phenyl substituted with halo or -OCH₃; and pyridine. In other embodiments
 25 these compounds have formula III.

- In one embodiment, R³ is absent or is selected from H; halo; -C₀₋₅alkylene-OH;
 -NH₂; -C₁₋₆alkyl; -CF₃; -C₃₋₇cycloalkyl; -C₀₋₂alkylene-O-C₁₋₆alkyl; -C(O)R²⁰; -C₀₋₁alkylene-
 COOR²¹; -C(O)NR²²R²³; -NHC(O)R²⁴; =O; -NO₂; -C(CH₃)=N(OH); phenyl optionally
 substituted with one or two groups independently selected from halo, -OH, -CF₃, -OCH₃,
 30 -NHC(O)CH₃, and phenyl; naphthalenyl; pyridinyl; pyrazinyl; pyrazolyl optionally
 substituted with methyl; thiophenyl optionally substituted with methyl or halo; furanyl; and
 -CH₂-morpholinyl; and R²¹ is H. In other embodiments these compounds have formula III.

In another embodiment, R³ is -C₀₋₁alkylene-COOR²¹, and R²¹ is selected from

-C₁₋₆alkyl, -C₁₋₃alkylene-C₆₋₁₀aryl, -C₁₋₃alkylene-C₁₋₉heteroaryl, -C₃₋₇cycloalkyl,
 -[(CH₂)₂O]₁₋₃CH₃, -C₁₋₆alkylene-OC(O)R²⁵; -C₁₋₆alkylene-NR²⁷R²⁸, -C₁₋₆alkylene-C(O)R³³,
 -C₀₋₆alkylenemorpholinyl, -C₁₋₆alkylene-SO₂-C₁₋₆alkyl,



- 5 In one aspect of the invention, these compounds may find particular utility as prodrugs or as intermediates in the synthetic procedures described herein. In other embodiments these compounds have formula III.


The R⁴ moiety can be absent. When present, R⁴ is attached to a carbon or nitrogen atom in the "X" group, and is selected from:

- 10 H;
 -OH;
 -C₁₋₆alkyl, e.g., -CH₃;
 -C₁₋₂alkylene-COOR³⁵, e.g., -CH₂COOH and -(CH₂)₂-COOH;
 -CH₂OC(O)CH(R³⁶)NH₂, e.g., -CH₂OC(O)CH[CH(CH₃)₂]NH₂;
 15 -OCH₂OC(O)CH(R³⁶)NH₂, e.g., -OCH₂OC(O)CH[CH(CH₃)₂]NH₂;
 -OCH₂OC(O)CH₃;
 -CH₂OP(O)(OH)₂;
 -CH₂CH(OH)CH₂OH;
 -CH[CH(CH₃)₂]-NHC(O)O-C₁₋₆alkyl;
 20 pyridinyl; and





- phenyl or benzyl optionally substituted with one or more groups selected from halo, -COOR³⁵, -OCH₃, -OCF₃, and -SCF₃ (e.g., 4-chlorophenyl, 3-methoxyphenyl, 2,4-dichlorophenyl, 3,4-dichlorophenyl, 2-chloro, 5-fluorophenyl, 3-trifluoromethoxy, 4-chlorophenyl, 3-trifluoromethylsulfanyl, 4-chlorophenyl, 2,6-difluoro, 4-chlorophenyl, 2-chlorobenzyl, 3-chlorobenzyl, 4-chlorobenzyl, 3-carboxybenzyl, 4-carboxybenzyl, 3-methoxybenzyl, 2-chloro, 5-fluorobenzyl, 3-chloro, 5-fluorobenzyl, 2-fluoro, 4-chlorobenzyl, 3-chloro, 4-fluorobenzyl, 3-OCF₃, 4-chlorobenzyl, 3-SCF₃, 4-chlorobenzyl, 2,6-difluoro, 3-chlorobenzyl, 2,6-difluoro, 4-chlorobenzyl, and 2,3,5,6-tetrafluoro, 4-methoxy benzyl).

- 30 The R³⁵ moiety is selected from:

- H;
 -C₁₋₆alkyl, e.g., -CH₃ and -CH₂CH₃;
 -C₁₋₃alkylene-C₆₋₁₀aryl;
 -C₁₋₃alkylene-C₁₋₉heteroaryl, e.g., -CH₂-pyridine;
 5 -C₃₋₇cycloalkyl;
 -[(CH₂)₂O]₁₋₃CH₃, e.g., -(CH₂)₂OCH₃;
 -C₁₋₆alkylene-OC(O)R²⁵, e.g., -CH₂OC(O)CH₃, -CH₂OC(O)O-CH₃, -CH₂OC(O)O-CH₃, -CH(CH₃)OC(O)O-CH₂CH₃, -CH(CH₃)OC(O)O-CH(CH₃)₂, -CH₂CH(CH₃)OC(O)-cyclopentyl, -CH₂OC(O)O-cyclopropyl, -CH(CH₃)-OC(O)-O-cyclohexyl, -CH₂OC(O)O-cyclopentyl, -CH₂CH(CH₃)OC(O)-phenyl, and -CH₂OC(O)O-phenyl;
 10 -C₁₋₆alkylene-NR²⁷R²⁸, e.g., -CH₂-pyrrolidine;
 -C₁₋₆alkylene-C(O)R³³;
 -C₀₋₆alkylenemorpholinyl, e.g., -(CH₂)₂-morpholinyl and -(CH₂)₃-morpholinyl:



- 15 -C₁₋₆alkylene-SO₂-C₁₋₆alkyl, e.g., -(CH₂)₂-SO₂-CH₃;




- 20 The R²⁵, R²⁷, R²⁸, R³³, and R³⁴ moieties are defined above. The R³⁶ moiety is selected from H, -CH(CH₃)₂, phenyl, and benzyl.

In addition, each alkyl group in R⁴ is optionally substituted with 1 to 8 fluoro

atoms. For example, when R^4 is $-C_{1-2}\text{alkylene-COOR}^{35}$ and R^{35} is $-C_{1-6}\text{alkyl}$, R^4 can also be a group such as $-\text{COOCH}(\text{CH}_3)\text{CF}_3$, $-\text{COOCH}_2\text{CF}_2\text{CF}_3$, $-\text{COOCH}(\text{CF}_3)_2$, $-\text{COO}(\text{CH}_2)_2\text{CF}_3$, $-\text{COOCH}(\text{CH}_2\text{F})_2$, $-\text{COOC}(\text{CF}_3)_2\text{CH}_3$, and $-\text{COOCH}(\text{CH}_3)\text{CF}_2\text{CF}_3$.

The R^4 moiety can also be taken together with R^3 to form $-\text{phenylene-O}-(\text{CH}_2)_{1-3}-$ or $-\text{phenylene-O-CH}_2\text{-CHOH-CH}_2-$. For purposes of illustration only, these embodiments are depicted below with X being pyrazole. It is understood that other X groups can be used also.




In another particular embodiment, R^4 is absent or is selected from H; $-\text{OH}$; $-C_{1-6}\text{alkyl}$; $-C_{1-2}\text{alkylene-COOR}^{35}$; $-\text{CH}_2\text{OC(O)CH(R}^{36})\text{NH}_2$; $-\text{CH}_2\text{CH(OH)CH}_2\text{OH}$; pyridinyl; phenyl optionally substituted with one halo group; and benzyl optionally substituted with one or more groups selected from halo, $-\text{COOH}$, $-\text{OCH}_3$, $-\text{OCF}_3$, and $-\text{SCF}_3$; R^{35} is H; and R^{36} is $-\text{CH}(\text{CH}_3)_2$; or R^3 and R^4 are taken together to form $-\text{phenylene-O}-(\text{CH}_2)_{1-3}-$ or $-\text{phenylene-O-CH}_2\text{-CHOH-CH}_2-$. In other embodiments these compounds have formula III.

In one embodiment, R^4 is absent or is selected from H; $-\text{OH}$; $-C_{1-6}\text{alkyl}$; $-C_{1-2}\text{alkylene-COOR}^{35}$; $-\text{CH}_2\text{OC(O)CH(R}^{36})\text{NH}_2$; $-\text{CH}_2\text{CH(OH)CH}_2\text{OH}$; pyridinyl; and phenyl or benzyl optionally substituted with one or more groups selected from halo, $-\text{COOR}^{35}$, $-\text{OCH}_3$, $-\text{OCF}_3$, and $-\text{SCF}_3$; and R^{35} is H. In other embodiments these compounds have formula III.


In another embodiment, R^4 is selected from $-\text{OCH}_2\text{OC(O)CH}_3$; $-\text{CH}_2\text{OP(O)(OH)}_2$; $-C_{1-2}\text{alkylene-COOR}^{35}$; and phenyl or benzyl substituted with at least one $-\text{COOR}^{35}$ group; where R^{35} is selected from $-C_{1-6}\text{alkyl}$, $-C_{1-3}\text{alkylene-C}_{6-10}\text{aryl}$, $-C_{1-3}\text{alkylene-C}_{1-9}\text{heteroaryl}$,

-C₃₋₇cycloalkyl, -[(CH₂)₂O]₁₋₃CH₃, -C₁₋₆alkylene-OC(O)R²⁵; -C₁₋₆alkylene-NR²⁷R²⁸,
 -C₁₋₆alkylene-C(O)R³³, -C₀₋₆alkylenemorpholinyl, -C₁₋₆alkylene-SO₂-C₁₋₆alkyl,




In one aspect of the invention, these compounds may find particular utility as prodrugs or
 5 as intermediates in the synthetic procedures described herein. In other embodiments these compounds have formula III.

The numbering for the R⁵ and R⁶ groups is as follows:




The integer "a" is 0 or 1. The R⁵ moiety, when present, is selected from halo, -CH₃, -CF₃,
 10 and -CN. In one embodiment, a is 0. In another embodiment, a is 1, and R⁵ is halo, such as 3-chloro or 3-fluoro. The integer "b" is 0 or an integer from 1 to 3. The R⁶ moiety, when present, is independently selected from halo, -OH, -CH₃, -OCH₃, and -CF₃. In one embodiment, b is 0. In another embodiment, b is 1 and R⁶ is selected from Cl, F, -OH, -CH₃, -OCH₃, and -CF₃, such as 2'-chloro, 3'-chloro, 2'-fluoro, 3'-fluoro, 2'-hydroxy, 3'-
 15 hydroxy, 3'-methyl, 2'-methoxy, or 3'-trifluoromethyl. In another embodiment, b is 1 and R⁶ is halo, -CH₃, or -OCH₃, such as 3'-chloro, 3'-methyl, or 2'-methoxy. In one embodiment, b is 2 and R⁶ is 2'-fluoro-5'-chloro, 2',5'-dichloro, 2',5'-difluoro, 2'-methyl-5'-chloro, 3'-fluoro-5'-chloro, 3'-hydroxy-5'-chloro, 3',5'-dichloro, 3',5'-difluoro, 2'-methoxy-5'-chloro, 2'-methoxy-5'-fluoro, 2'-hydroxy-5'-fluoro, 2'-fluoro-3'-chloro, 2'-hydroxy-5'-
 20 chloro, or 2'-hydroxy-3'-chloro; and in another embodiment, b is 2 and each R⁶ is independently halo, for example, 2'-fluoro-5'-chloro and 2',5'-dichloro. In another embodiment, b is 3 and each R⁶ is independently halo or -CH₃, such as 2'-methyl-3', 5'-dichloro or 2'-fluoro-3'-methyl-5'-chloro. In yet another embodiment, a is 1 and b is 1 and R⁵ and R⁶ are independently halo, for example, 3-chloro and 3'chloro. In other
 25 embodiments these compounds have formula III. Of particular interest are compounds of


the formulas:



The methylene linker on the biphenyl is optionally substituted with one or two -C₁₋₆alkyl groups or cyclopropyl. For example, in one embodiment, the methylene linker on the biphenyl is unsubstituted; in another embodiment, the methylene linker on the biphenyl is substituted with one -C₁₋₆alkyl group (e.g., -CH₃); and in yet another embodiment, the methylene linker on the biphenyl is substituted with two -C₁₋₆alkyl groups (e.g., two -CH₃ groups); in another embodiment, the methylene linker on the biphenyl is substituted with a cyclopropyl group. These embodiments are depicted, respectively, as:



In another embodiment, R⁷ is selected from H, -C₁₋₈alkyl, -C₁₋₆alkylene-OC(O)R¹⁰, -C₀₋₆alkylenemorpholinyl, -C₁₋₆alkylene-SO₂-C₁₋₆alkyl, and




R¹⁰ is -O-C₃₋₇cycloalkyl; R⁸ is H; R⁹ is H; R² is H; R³ is absent or is selected from H; halo;

- C₀₋₅alkylene-OH; -NH₂; -C₁₋₆alkyl; -CF₃; -C₃₋₇cycloalkyl; -C₀₋₂alkylene-O-C₁₋₆alkyl; -C(O)R²⁰; -C₀₋₁alkylene-COOR²¹; -C(O)NR²²R²³; -NHC(O)R²⁴; =O; -C(CH₃)=N(OH); phenyl optionally substituted with one or two groups independently selected from halo, -OH, -CF₃, -OCH₃, -NHC(O)CH₃, and phenyl; naphthalenyl; pyridinyl; pyrazinyl; pyrazolyl substituted with methyl; thiophenyl substituted with methyl or halo; furanyl; and -CH₂-morpholinyl; R²⁰ is -C₁₋₆alkyl; R²¹ is H or -C₁₋₆alkyl; R²² is H or -C₁₋₆alkyl; and R²³ is selected from H, -C₁₋₆alkyl, -CH₂COOH, -(CH₂)₂OH, -(CH₂)₂OCH₃, -(CH₂)₂N(CH₃)₂, -C₃₋₇cycloalkyl, and -(CH₂)₂-imidazolyl; or R²² and R²³ are taken together to form azetidine, pyrrolidine, piperidine or morpholine, all optionally substituted with -OH or -CONH₂; and R²⁴ is selected from -C₁₋₆alkyl; -C₀₋₁alkylene-O-C₁₋₆alkyl; phenyl substituted with halo or -OCH₃; and pyridinyl; R⁴ is absent or is selected from H; -OH; -C₁₋₆alkyl; -C₁₋₂alkylene-COOR³⁵; -CH₂OC(O)CH(R³⁶)NH₂; -CH₂CH(OH)CH₂OH; pyridinyl; phenyl optionally substituted with one halo group; and benzyl optionally substituted with one or more groups selected from halo, -COOH, -OCH₃, -OCF₃, and -SCF₃; R³⁵ is H; and R³⁶ is -CH(CH₃)₂; or R³ and R⁴ are taken together to form -phenylene-O-(CH₂)₁₋₃- or -phenylene-O-CH₂-CHOH-CH₂-; a is 0; or a is 1 and R⁵ is 3-chloro; b is 0; or b is 1 and R⁶ is 3'-chloro, 3'-methyl, or 2'-methoxy; or b is 2 and R⁶ is 2'-fluoro-5'-chloro, 2',5'-dichloro, 2'-methyl-5'-chloro, or 3'-chloro-5'-hydroxy; and the methylene linker on the biphenyl is optionally substituted with 2 methyl groups. In one embodiment, X is selected from pyrazole, triazole, benzotriazole, furan, tetrazole, pyrazine, thiophene, oxazole, isoxazole, thiazole, oxadiazole, pyridazine, pyridine, pyrimidine, benzoxazole, pyridylimidazole, and pyridyltriazole.

- In still another embodiment, R¹ is -OR⁷; R² is H; X is selected from pyrazole, triazole, benzotriazole, isoxazole, pyridazine, pyrimidine, and pyridyltriazole; R³ is selected from H; halo; -C₀₋₅alkylene-OH; -C₁₋₆alkyl; -CF₃; -C₃₋₇cycloalkyl; -C₀₋₂alkylene-O-C₁₋₆alkyl; -C(O)R²⁰; -C₀₋₁alkylene-COOR²¹; -C(O)NR²²R²³; -NHC(O)R²⁴; =O; -C(CH₃)=N(OH); phenyl optionally substituted with one or two groups independently selected from halo, -OH, and -OCH₃; pyridinyl; pyrazinyl; and thiophenyl substituted with methyl or halo; R⁴ is selected from H; -OH; -C₁₋₆alkyl; -C₁₋₂alkylene-COOR³⁵; -CH₂CH(OH)CH₂OH; pyridinyl; and phenyl optionally substituted with one halo group; or R³ and R⁴ are taken together to form -phenylene-O-(CH₂)₁₋₃-; a is 0; or a is 1 and R⁵ is 3-chloro; b is 0; or b is 1 and R⁶ is 3'-chloro, 3'-methyl, or 2'-methoxy; or b is 2 and R⁶ is 2'-fluoro-5'-chloro, 2',5'-dichloro, 2'-methyl-5'-chloro, or 3'-chloro-5'-hydroxy; R²⁰ is


- C₁₋₆alkyl; R²¹ is H; R²² is selected from H and -C₁₋₆alkyl; and R²³ is selected from -C₁₋₆alkyl, -(CH₂)₂OCH₃, and -C₃₋₇cycloalkyl; or R²² and R²³ are taken together to form azetidine, pyrrolidine, or piperidine, all optionally substituted with -OH or -CONH₂; R²⁴ is phenyl substituted with halo or -OCH₃; R³⁵ is H; the methylene linker on the biphenyl is optionally substituted with 2 methyl groups; and R⁷ is as defined for formula I.

A particular group of compounds of formula I are those disclosed in U.S. Provisional Application No. 61/423,180, filed on December 15, 2010. This group includes compounds of formula II:



(II)


- where: R¹ is selected from -OR⁷ and -NR⁸R⁹; R⁷ is selected from H; -C₁₋₆alkyl; -C₁₋₃alkylene-C₆₋₁₀aryl; -C₁₋₃alkylene-C₁₋₉heteroaryl; -C₃₋₇cycloalkyl; -(CH₂)₂OCH₃; -C₁₋₆alkylene-OC(O)R¹⁰; -CH₂-pyridine; -CH₂-pyrrolidine; -C₀₋₆alkylenemorpholine; -C₁₋₆alkylene-SO₂-C₁₋₆alkyl;



- where R¹⁰ is selected from -C₁₋₆alkyl, -O-C₁₋₆alkyl, -C₃₋₇cycloalkyl, -O-C₃₋₇cycloalkyl, phenyl, -O-phenyl, -NR¹²R¹³, and -CH(NH₂)CH₂COOCH₃; and R¹² and R¹³ are independently selected from H, -C₁₋₆alkyl, and benzyl, or R¹² and R¹³ are taken together as -(CH₂)₃₋₆; R⁸ is selected from H; -OH; -OC(O)R¹⁴; -CH₂COOH; -O-benzyl; pyridyl; and -OC(S)NR¹⁵R¹⁶; where R¹⁴ is selected from H, -C₁₋₆alkyl, -C₆₋₁₀aryl, -OCH₂-C₆₋₁₀aryl, -CH₂O-C₆₋₁₀aryl, and -NR¹⁵R¹⁶; and R¹⁵ and R¹⁶ are independently selected from H and -C₁₋₄alkyl; R⁹ is selected from H; -C₁₋₆alkyl; and -C(O)R¹⁷; where R¹⁷ is selected from H; -C₁₋₆alkyl; -C₃₋₇cycloalkyl; -C₆₋₁₀aryl; and -C₁₋₉heteroaryl; R² is H or is taken together with R¹ to form -OCHR¹⁸R¹⁹- or -NHC(O)-; where R¹⁸ and R¹⁹ are independently selected from H, -C₁₋₆alkyl, and -O-C₃₋₇cycloalkyl, or R¹⁸ and R¹⁹ are taken together to form =O; X is a -C₁₋₉heteroaryl or a partially unsaturated -C₃₋₅heterocycle; R³ is absent or is selected from

H; halo; -C₀₋₅alkylene-OH; -NH₂; -C₁₋₆alkyl; -C₃₋₇cycloalkyl; -C₀₋₁alkylene-O-C₁₋₆alkyl; -C(O)R²⁰; -C₀₋₁alkylene-C(O)OR²¹; -C(O)NR²²R²³; -NHC(O)R²⁴; phenyl optionally substituted with one group selected from halo, -CF₃, -OCH₃, -NHC(O)CH₃, and phenyl; naphthyl; pyridine; pyrazine; pyrazole optionally substituted with methyl; thiophene optionally substituted with methyl; and furan; and R³, when present, is attached to a carbon atom; R²⁰ is selected from H and -C₁₋₆alkyl; R²¹ is selected from H; -C₁₋₆alkyl; -C₁₋₃alkylene-C₆₋₁₀aryl; -C₁₋₃alkylene-C₁₋₉heteroaryl; -C₃₋₇cycloalkyl; -(CH₂)₂OCH₃; -C₁₋₆alkylene-OC(O)R²⁵; -CH₂-pyridine; -CH₂-pyrrolidine; -C₀₋₆alkylenemorpholine; -C₁₋₆alkylene-SO₂-C₁₋₆alkyl;

10



where R²⁵ is selected from -C₁₋₆alkyl, -O-C₁₋₆alkyl, -C₃₋₇cycloalkyl, -O-C₃₋₇cycloalkyl, phenyl, -O-phenyl, -NR²⁷R²⁸, and -CH(NH₂)CH₂COOCH₃; and R²⁷ and R²⁸ are independently selected from H, -C₁₋₆alkyl, and benzyl, or R²⁷ and R²⁸ are taken together as -(CH₂)₃₋₆-; R²² and R²³ are independently selected from H; -C₁₋₆alkyl; -CH₂COOH; -(CH₂)₂OH; -(CH₂)₂OCH₃; -(CH₂)₂SO₂NH₂; -(CH₂)₂N(CH₃)₂; -C₃₋₇cycloalkyl; and -(CH₂)₂-imidazole; or R²² and R²³ are taken together to form a saturated or partially unsaturated -C₃₋₅heterocycle optionally substituted with -OH, -COOH, or -CONH₂; and optionally containing an oxygen atom in the ring; R²⁴ is selected from -C₁₋₆alkyl; -O-C₁₋₆alkyl; -CH₂-O-C₁₋₆alkyl; phenyl substituted with -OCH₃; and pyridine; R⁴ is selected from H; -C₁₋₆alkyl; phenyl or benzyl substituted with one or more groups selected from halo, -COOH, -OCH₃, -OCF₃, and -SCF₃; and R⁴ is attached to a carbon or nitrogen atom; a is 0 or 1; R⁵ is halo or -CF₃; and b is 0 or 1; R⁶ is halo; or a pharmaceutically acceptable salt thereof.

In addition, particular compounds of formula I that are of interest include those set forth in the Examples below, as well as pharmaceutically acceptable salts thereof.

GENERAL SYNTHETIC PROCEDURES

Compounds of the invention can be prepared from readily available starting materials using the following general methods, the procedures set forth in the Examples, or by using other methods, reagents, and starting materials that are known to those of ordinary skill in the art. Although the following procedures may illustrate a particular embodiment of the invention, it is understood that other embodiments of the invention can be similarly

prepared using the same or similar methods or by using other methods, reagents and starting materials known to those of ordinary skill in the art. It will also be appreciated that where typical or preferred process conditions (for example, reaction temperatures, times, mole ratios of reactants, solvents, pressures, etc.) are given, other process conditions can also be used unless otherwise stated. In some instances, reactions were conducted at room temperature and no actual temperature measurement was taken. It is understood that room temperature can be taken to mean a temperature within the range commonly associated with the ambient temperature in a laboratory environment, and will typically be in the range of about 18°C to about 30°C. In other instances, reactions were conducted at room temperature and the temperature was actually measured and recorded. While optimum reaction conditions will typically vary depending on various reaction parameters such as the particular reactants, solvents and quantities used, those of ordinary skill in the art can readily determine suitable reaction conditions using routine optimization procedures.

Additionally, as will be apparent to those skilled in the art, conventional protecting groups may be necessary or desired to prevent certain functional groups from undergoing undesired reactions. The choice of a suitable protecting group for a particular functional group as well as suitable conditions and reagents for protection and deprotection of such functional groups are well-known in the art. Protecting groups other than those illustrated in the procedures described herein may be used, if desired. For example, numerous protecting groups, and their introduction and removal, are described in T. W. Greene and G. M. Wuts, *Protecting Groups in Organic Synthesis*, Fourth Edition, Wiley, New York, 2006, and references cited therein.

Carboxy-protecting groups are suitable for preventing undesired reactions at a carboxy group, and examples include, but are not limited to, methyl, ethyl, *t*-butyl, benzyl (Bn), *p*-methoxybenzyl (PMB), 9-fluorenylmethyl (Fm), trimethylsilyl (TMS), *t*-butyldimethylsilyl (TBDMS), diphenylmethyl (benzhydryl, DPM) and the like. Amino-protecting groups are suitable for preventing undesired reactions at an amino group, and examples include, but are not limited to, *t*-butoxycarbonyl (BOC), trityl (Tr), benzyloxycarbonyl (Cbz), 9-fluorenylmethoxycarbonyl (Fmoc), formyl, trimethylsilyl (TMS), *t*-butyldimethylsilyl (TBDMS), and the like. Hydroxyl-protecting groups are suitable for preventing undesired reactions at a hydroxyl group, and examples include, but are not limited to C₁₋₆alkyls, silyl groups including triC₁₋₆alkylsilyl groups, such as trimethylsilyl (TMS), triethylsilyl (TES), and *tert*-butyldimethylsilyl (TBDMS); esters

(acyl groups) including C₁₋₆alkanoyl groups, such as formyl, acetyl, and pivaloyl, and aromatic acyl groups such as benzoyl; arylmethyl groups such as benzyl (Bn), *p*-methoxybenzyl (PMB), 9-fluorenylmethyl (Fm), and diphenylmethyl (benzhydryl, DPM); and the like.

5 Standard deprotection techniques and reagents are used to remove the protecting groups, and may vary depending upon which group is used. For example, sodium or lithium hydroxide is commonly used when the carboxy-protecting group is methyl, an acid such as TFA or HCl is commonly used when the carboxy-protecting group is ethyl or *t*-butyl, and H₂/Pd/C may be used when the carboxy-protecting group is benzyl. A BOC
10 amino-protecting group can be removed using an acidic reagent such as TFA in DCM or HCl in 1,4-dioxane, while a Cbz amino-protecting group can be removed by employing catalytic hydrogenation conditions such as H₂ (1 atm) and 10% Pd/C in an alcoholic solvent ("H₂/Pd/C"). H₂/Pd/C is commonly used when the hydroxyl-protecting group is benzyl, while NaOH is commonly used when the hydroxyl-protecting group is an acyl
15 group.

 Suitable bases for use in these schemes include, by way of illustration and not limitation, potassium carbonate, calcium carbonate, sodium carbonate, triethylamine, pyridine, 1,8-diazabicyclo-[5.4.0]undec-7-ene (DBU), *N,N*-diisopropylethylamine (DIPEA), 4-methylmorpholine, sodium hydroxide, potassium hydroxide, potassium *t*-
20 butoxide, and metal hydrides.


 Suitable inert diluents or solvents for use in these schemes include, by way of illustration and not limitation, tetrahydrofuran (THF), acetonitrile (MeCN), *N,N*-dimethylformamide (DMF), *N,N*-dimethylacetamide (DMA), dimethyl sulfoxide (DMSO), toluene, dichloromethane (DCM), chloroform (CHCl₃), carbon tetrachloride (CCl₄), 1,4-
25 dioxane, methanol, ethanol, water, and the like.

 Suitable carboxylic acid/amine coupling reagents include benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (BOP), benzotriazol-1-yloxytripyrrolidinophosphonium hexafluorophosphate (PyBOP), *N,N,N',N'*-tetramethyl-O-(7-azabenzotriazol-1-yl)uronium hexafluorophosphate (HATU), 1,3-
30 dicyclohexylcarbodiimide (DCC), *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide (EDCI), carbonyldiimidazole (CDI), 1-hydroxybenzotriazole (HOBt), and the like. Coupling reactions are conducted in an inert diluent in the presence of a base such as DIPEA, and are performed under conventional amide bond-forming conditions.

All reactions are typically conducted at a temperature within the range of about -78 C to 100°C, for example at room temperature. Reactions may be monitored by use of thin layer chromatography (TLC), high performance liquid chromatography (HPLC), and/or LCMS until completion. Reactions may be complete in minutes, or may take hours, typically from 1-2 hours and up to 48 hours. Upon completion, the resulting mixture or reaction product may be further treated in order to obtain the desired product. For example, the resulting mixture or reaction product may be subjected to one or more of the following procedures: concentrating or partitioning (for example, between EtOAc and water or between 5% THF in EtOAc and 1M phosphoric acid); extraction (for example, with EtOAc, CHCl₃, DCM, chloroform); washing (for example, with saturated aqueous NaCl, saturated aqueous NaHCO₃, Na₂CO₃ (5%), CHCl₃ or 1M NaOH); drying (for example, over MgSO₄, over Na₂SO₄, or *in vacuo*); filtering; crystallizing (for example, from EtOAc and hexanes); being concentrated (for example, *in vacuo*); and/or purification (e.g., silica gel chromatography, flash chromatography, preparative HPLC, reverse phase-HPLC, or crystallization).

Compounds of formula I, as well as their salts, can be prepared as shown in Scheme I:

Scheme I



The process comprises the step of coupling compound 1 with compound 2, where R¹-R⁶, X, a, and b are as defined for formula I, and P¹ is H or a suitable amino-protecting group, examples of which include, *t*-butoxycarbonyl, trityl, benzyloxycarbonyl, 9-fluorenylmethoxycarbonyl, formyl, trimethylsilyl, and *t*-butyldimethylsilyl. When P¹ is an amino protecting group, the process further comprises deprotecting the compound of formula 1, before or *in situ* with the coupling step.


In instances where R¹ is a group such as -OCH₃ or -OCH₂CH₃, the coupling step may be followed by a deprotection step to provide a compound of formula I where R¹ is a

group such as -OH. Thus, one method of preparing compounds of the invention involves coupling compounds 1 and 2, with an optional deprotection step to form a compound of formula I or a pharmaceutically acceptable salt thereof.

Methods of preparing compound 1 are described in the Examples. Compound 2 is generally commercially available or can be prepared using procedures that are known in the art.

Compounds of formula I, as well as their salts, can also be prepared as shown in Scheme II:

Scheme II




10

In the first step, compound 1 is coupled with compound 3 and compound 3 is coupled to compound 4, where Y and Z react *in situ* to form the R³ moiety. For example, when R³ is -C(O)NR²²R²³, Y is -COOH and Z is HNR²²R²³. Alternately, compound 3 is first coupled to compound 4, and the resulting compound is then coupled with compound 1. As with Scheme I, in instances where R¹ is a group such as -OCH₃ or -OCH₂CH₃, the coupling step may be followed by a deprotection step to provide a compound of formula I where R¹ is a group such as -OH. Thus, one method of preparing compounds of the invention involves coupling compounds 1, 2 and 3, with an optional deprotection step to form a compound of formula I or a pharmaceutically acceptable salt thereof.

20 Compounds 3 and 4 are generally commercially available or can be prepared using procedures that are known in the art.

Compounds of formula I, as well as their salts, can also be prepared as shown in Scheme III: {Based on Example 9}


Scheme III




Again, as with Schemes I and II, this is a standard coupling reaction between a compound of formula I, where R^1 is $-OH$ and compound 5, to yield a compound of formula I, where

5 R^1 is $-NR^8R^9$.

Certain intermediates described herein are believed to be novel and accordingly, such compounds are provided as further aspects of the invention including, for example, the compounds of formula 1 or a salt thereof:




10 where P^1 is H or an amino-protecting group selected from *t*-butoxycarbonyl, trityl, benzyloxycarbonyl, 9-fluorenylmethoxycarbonyl, formyl, trimethylsilyl, and *t*-butyldimethylsilyl; and R^1 , R^2 , R^5 , R^6 , a and b are as defined for formula I. Another intermediate of the invention has formula 6 or a salt thereof:




15 where R^{1p} is selected from $-O-P^3$, $-NHP^2$, and $-NH(O-P^4)$; where P^2 is an amino-protecting group selected from *t*-butoxycarbonyl, trityl, benzyloxycarbonyl, 9-

fluorenylmethoxycarbonyl, formyl, trimethylsilyl, and *t*-butyldimethylsilyl; P³ is a carboxy-protecting group selected from methyl, ethyl, *t*-butyl, benzyl, *p*-methoxybenzyl, 9-fluorenylmethyl, trimethylsilyl, *t*-butyldimethylsilyl, and diphenylmethyl; P⁴ is a hydroxyl-protecting group selected from -C₁₋₆alkyl, triC₁₋₆alkylsilyl, -C₁₋₆alkanoyl, benzoyl, benzyl, *p*-methoxybenzyl, 9-fluorenylmethyl, and diphenylmethyl; and R², R³, R⁴, R⁵, R⁶, a, b, and X are as defined for formula I. Another intermediate of the invention has formula 7 or a salt thereof:




where R^{3P} is selected from -C₀₋₅alkylene-O-P⁴, -C₀₋₁alkylene-COO-P³, and phenyl substituted with -O-P⁴; P³ is a carboxy-protecting group selected from methyl, ethyl, *t*-butyl, benzyl, *p*-methoxybenzyl, 9-fluorenylmethyl, trimethylsilyl, *t*-butyldimethylsilyl, and diphenylmethyl; P⁴ is a hydroxyl-protecting group selected from -C₁₋₆alkyl, triC₁₋₆alkylsilyl, -C₁₋₆alkanoyl, benzoyl, benzyl, *p*-methoxybenzyl, 9-fluorenylmethyl, and diphenylmethyl; and R¹, R², R⁴, R⁵, R⁶, a, b, and X are as defined for formula I. Still another intermediate of the invention has formula 8 or a salt thereof:



where R^{4P} is selected from -O-P⁴, -C₁₋₂alkylene-COO-P³; and phenyl or benzyl substituted with -COO-P³; P³ is a carboxy-protecting group selected from methyl, ethyl, *t*-butyl, benzyl, *p*-methoxybenzyl, 9-fluorenylmethyl, trimethylsilyl, *t*-butyldimethylsilyl, and diphenylmethyl; P⁴ is a hydroxyl-protecting group selected from -C₁₋₆alkyl, triC₁₋₆alkylsilyl, -C₁₋₆alkanoyl, benzoyl, benzyl, *p*-methoxybenzyl, 9-fluorenylmethyl, and diphenylmethyl; and R¹, R², R³, R⁵, R⁶, a, b, and X are as defined for formula I. Yet

another intermediate of the invention has formula 9 or a salt thereof:



(9),

where R^{3p} is selected from $-C_{0-5}alkylene-O-P^4$, $-C_{0-1}alkylene-COO-P^3$, and phenyl substituted with $-O-P^4$; R^{4p} is selected from $-O-P^4$, $-C_{1-2}alkylene-COO-P^3$; and phenyl or benzyl substituted with $-COO-P^3$; P^3 is a carboxy-protecting group selected from methyl, ethyl, *t*-butyl, benzyl, *p*-methoxybenzyl, 9-fluorenylmethyl, trimethylsilyl, *t*-butyldimethylsilyl, and diphenylmethyl; P^4 is a hydroxyl-protecting group selected from $-C_{1-6}alkyl$, $triC_{1-6}alkylsilyl$, $-C_{1-6}alkanoyl$, benzoyl, benzyl, *p*-methoxybenzyl, 9-fluorenylmethyl, and diphenylmethyl; and R^1 , R^2 , R^5 , R^6 , a , b , and X are as defined for formula I. Thus, another method of preparing compounds of the invention involves deprotecting a compound of formula 1, 6, 7, 8, 9, or a salt thereof.

Further details regarding specific reaction conditions and other procedures for preparing representative compounds of the invention or intermediates thereof are described in the Examples set forth below.

UTILITY

Compounds of the invention possess neprilysin (NEP) inhibition activity, that is, the compounds are able to inhibit enzyme-catalytic activity. In another embodiment, the compounds do not exhibit significant inhibitory activity of the angiotensin-converting enzyme. One measure of the ability of a compound to inhibit NEP activity is the inhibition constant (pK_i). The pK_i value is the negative logarithm to base 10 of the dissociation constant (K_i), which is typically reported in molar units. Compounds of the invention of particular interest are those having a pK_i at NEP greater than or equal to 6.0, particularly those having a pK_i greater than or equal to 7.0, and even more particularly those having a pK_i greater than or equal to 8.0. In one embodiment, compounds of interest have a pK_i in the range of 6.0-6.9; in another embodiment, compounds of interest have a pK_i in the range of 7.0-7.9; in yet another embodiment, compounds of interest have a pK_i in the range of 8.0-8.9; and in still another embodiment, compounds of interest have a pK_i in the range of

greater than or equal to 9.0. Such values can be determined by techniques that are well known in the art, as well as in the assays described herein.

Another measure of the ability of a compound to inhibit NEP activity is the apparent inhibition constant (IC_{50}), which is the molar concentration of compound that results in half-maximal inhibition of substrate conversion by the NEP enzyme. The pIC_{50} value is the negative logarithm to base 10 of the IC_{50} . Compounds of the invention that are of particular interest, include those that exhibit a pIC_{50} for NEP greater than or equal to about 5.0. Compounds of interest also include those having a pIC_{50} for NEP $\geq$ about 6.0 or a pIC_{50} for NEP $\geq$ about 7.0. In another embodiment, compounds of interest have a pIC_{50} for NEP within the range of about 7.0-11.0; and in another embodiment, within the range of about 8.0-11.0, such as within the range of about 8.0-10.0.

It is noted that in some cases, compounds of the invention may possess weak NEP inhibition activity. In such cases, those of skill in the art will recognize that these compounds still have utility as research tools.

Exemplary assays to determine properties of compounds of the invention, such as the NEP inhibiting activity, are described in the Examples and include by way of illustration and not limitation, assays that measure NEP inhibition (described in Assay 1). Useful secondary assays include assays to measure ACE inhibition (also described in Assay 1) and aminopeptidase P (APP) inhibition (described in Sulpizio et al. (2005) *JPET* 315:1306-1313). A pharmacodynamic assay to assess the *in vivo* inhibitory potencies for ACE and NEP in anesthetized rats is described in Assay 2 (see also Seymour et al. (1985) *Hypertension* 7(Suppl 1):I-35-I-42 and Wagle et al. (1992) *Can. J. Physiol. Pharmacol.* 70:1525-1528), where ACE inhibition is measured as the percent inhibition of the angiotensin I pressor response and NEP inhibition is measured as increased urinary cyclic guanosine 3', 5'-monophosphate (cGMP) output.

There are many *in vivo* assays that can be used to ascertain further utilities of the compounds of the invention. The conscious spontaneously hypertensive rat (SHR) model is a renin dependent hypertension model, and is described in Assay 3. See also Intengan et al. (1999) *Circulation* 100(22):2267-2275 and Badyal et al. (2003) *Indian Journal of Pharmacology* 35:349-362. The conscious desoxycorticosterone acetate-salt (DOCA-salt) rat model is a volume dependent hypertension model that is useful for measuring NEP activity, and is described in Assay 4. See also Trapani et al. (1989) *J. Cardiovasc. Pharmacol.* 14:419-424, Intengan et al. (1999) *Hypertension* 34(4):907-913, and Badyal et

al. (2003) *supra*). The DOCA-salt model is particularly useful for evaluating the ability of a test compound to reduce blood pressure as well as to measure a test compound's ability to prevent or delay a rise in blood pressure. The Dahl salt-sensitive (DSS) hypertensive rat model is a model of hypertension that is sensitive to dietary salt (NaCl), and is described in
5 Assay 5. See also Rapp (1982) *Hypertension* 4:753-763. The rat monocrotaline model of pulmonary arterial hypertension described, for example, in Kato et al. (2008) *J. Cardiovasc. Pharmacol.* 51(1):18-23, is a reliable predictor of clinical efficacy for the treatment of pulmonary arterial hypertension. Heart failure animal models include the DSS rat model for heart failure and the aorto-caval fistula model (AV shunt), the latter of which
10 is described, for example, in Norling et al. (1996) *J. Amer. Soc. Nephrol.* 7:1038-1044. Other animal models, such as the hot plate, tail-flick and formalin tests, can be used to measure the analgesic properties of compounds of the invention, as well as the spinal nerve ligation (SNL) model of neuropathic pain. See, for example, Malmberg et al. (1999) *Current Protocols in Neuroscience* 8.9.1-8.9.15.

15 Compounds of the invention are expected to inhibit the NEP enzyme in any of the assays listed above, or assays of a similar nature. Thus, the aforementioned assays are useful in determining the therapeutic utility of compounds of the invention, for example, their utility as antihypertensive agents or antidiarrheal agents. Other properties and utilities of compounds of the invention can be demonstrated using other *in vitro* and *in vivo* assays
20 well-known to those skilled in the art. Compounds of formula I may be active drugs as well as prodrugs. Thus, when discussing the activity of compounds of the invention, it is understood that any such prodrugs may not exhibit the expected activity in an assay, but are expected to exhibit the desired activity once metabolized.

Compounds of the invention are expected to be useful for the treatment and/or
25 prevention of medical conditions responsive to NEP inhibition. Thus it is expected that patients suffering from a disease or disorder that is treated by inhibiting the NEP enzyme or by increasing the levels of its peptide substrates, can be treated by administering a therapeutically effective amount of a compound of the invention. For example, by inhibiting NEP, the compounds are expected to potentiate the biological effects of
30 endogenous peptides that are metabolized by NEP, such as the natriuretic peptides, bombesin, bradykinins, calcitonin, endothelins, enkephalins, neurotensin, substance P and vasoactive intestinal peptide. Thus, these compounds are expected to have other physiological actions, for example, on the renal, central nervous, reproductive and


gastrointestinal systems.

In one embodiment of the invention, patients suffering from a disease or disorder that is treated by inhibiting the NEP enzyme, are treated by administering a compound of the invention that is in its active form, i.e., a compound of formula I where R^1 is selected from $-OR^7$ and $-NR^8R^9$, R^7 is H, R^8 is H or $-OH$, R^9 is H, and R^2-R^6 , a, b, and X are as defined for formula I.

In another embodiment, patients are treated by administering a compound that is metabolized *in vitro* to form a compound of formula I where R^1 is selected from $-OR^7$ and $-NR^8R^9$, R^7 is H, R^8 is H or $-OH$, R^9 is H, and R^2-R^6 , a, b, and X are as defined for formula I. In an exemplary embodiment, patients are treated by administering a compound that is metabolized *in vitro* to form a compound of formula III where R^1 is $-OR^7$ and R^7 is H.

In another embodiment, patients are treated by administering a compound of the invention that is in its prodrug form at the R^1 group, i.e., a compound of formula I where:

R^1 is $-OR^7$; and R^7 is selected from $-C_{1-8}alkyl$, $-C_{1-3}alkylene-C_{6-10}aryl$, $-C_{1-3}alkylene-C_{1-9}heteroaryl$, $-C_{3-7}cycloalkyl$, $-[(CH_2)_2O]_{1-3}CH_3$, $-C_{1-6}alkylene-OC(O)R^{10}$, $-C_{1-6}alkylene-NR^{12}R^{13}$, $-C_{1-6}alkylene-C(O)R^{31}$, $-C_{0-6}alkylenemorpholinyl$, $-C_{1-6}alkylene-SO_2-C_{1-6}alkyl$,



R^1 is $-NR^8R^9$; R^8 is selected from $-OC(O)R^{14}$, $-CH_2COOH$, $-O-benzyl$, $pyridyl$, and $-OC(S)NR^{15}R^{16}$; and R^9 is H; or

R^1 is $-NR^8R^9$; R^8 is selected from $-OC(O)R^{14}$, $-CH_2COOH$, $-O-benzyl$, $pyridyl$, and $-OC(S)NR^{15}R^{16}$; and R^9 is $-C_{1-6}alkyl$ or $-C(O)R^{17}$;

R^1 is $-NR^8R^9$; R^8 is selected from H or $-OH$; and R^9 is selected from $-C_{1-6}alkyl$, and $-C(O)R^{17}$;

R^1 is $-OR^7$ and R^2 is taken together with R^7 to form $-CR^{18}R^{19}$;-; or

R^1 is $-NR^8R^9$ and R^2 is taken together with R^8 to form $-C(O)$ -;

and R^{10} , $R^{12}-R^{17}$, R^{31} , R^{32} , R^3-R^6 , a, b, and X are as defined for formula I. In an exemplary embodiment, patients are treated by administering a compound of the invention that is in its prodrug form at the R^1 group and has formula III.

30

Cardiovascular Diseases

By potentiating the effects of vasoactive peptides like the natriuretic peptides and bradykinin, compounds of the invention are expected to find utility in treating and/or preventing medical conditions such as cardiovascular diseases. See, for example, Roques et al. (1993) *Pharmacol. Rev.* 45:87-146 and Dempsey et al. (2009) *Amer. J. of Pathology* 174(3):782-796. Cardiovascular diseases of particular interest include hypertension and heart failure. Hypertension includes, by way of illustration and not limitation: primary hypertension, which is also referred to as essential hypertension or idiopathic hypertension; secondary hypertension; hypertension with accompanying renal disease; severe hypertension with or without accompanying renal disease; pulmonary hypertension, including pulmonary arterial hypertension; and resistant hypertension. Heart failure includes, by way of illustration and not limitation: congestive heart failure; acute heart failure; chronic heart failure, for example with reduced left ventricular ejection fraction (also referred to as systolic heart failure) or with preserved left ventricular ejection fraction (also referred to as diastolic heart failure); and acute and chronic decompensated heart failure, with or without accompanying renal disease. Thus, one embodiment of the invention relates to a method for treating hypertension, particularly primary hypertension or pulmonary arterial hypertension, comprising administering to a patient a therapeutically effective amount of a compound of the invention.

For treatment of primary hypertension, the therapeutically effective amount is typically the amount that is sufficient to lower the patient's blood pressure. This would include both mild-to-moderate hypertension and severe hypertension. When used to treat hypertension, the compound may be administered in combination with other therapeutic agents such as aldosterone antagonists, aldosterone synthase inhibitors, angiotensin-converting enzyme inhibitors and dual-acting angiotensin-converting enzyme/neprilysin inhibitors, angiotensin-converting enzyme 2 (ACE2) activators and stimulators, angiotensin-II vaccines, anti-diabetic agents, anti-lipid agents, anti-thrombotic agents, AT₁ receptor antagonists and dual-acting AT₁ receptor antagonist/neprilysin inhibitors, β_1 -adrenergic receptor antagonists, dual-acting β -adrenergic receptor antagonist/ α_1 -receptor antagonists, calcium channel blockers, diuretics, endothelin receptor antagonists, endothelin converting enzyme inhibitors, neprilysin inhibitors, natriuretic peptides and their analogs, natriuretic peptide clearance receptor antagonists, nitric oxide donors, non-steroidal anti-inflammatory agents, phosphodiesterase inhibitors (specifically PDE-V inhibitors), prostaglandin receptor agonists, renin inhibitors, soluble guanylate cyclase

stimulators and activators, and combinations thereof. In one particular embodiment of the invention, a compound of the invention is combined with an AT₁ receptor antagonist, a calcium channel blocker, a diuretic, or a combination thereof, and used to treat primary hypertension. In another particular embodiment of the invention, a compound of the invention is combined with an AT₁ receptor antagonist, and used to treat hypertension with
5 accompanying renal disease. When used to treat resistant hypertension, the compound may be administered in combination with other therapeutic agents such as aldosterone synthase inhibitors.

For treatment of pulmonary arterial hypertension, the therapeutically effective
10 amount is typically the amount that is sufficient to lower the pulmonary vascular resistance. Other goals of therapy are to improve a patient's exercise capacity. For example, in a clinical setting, the therapeutically effective amount can be the amount that improves a patient's ability to walk comfortably for a period of 6 minutes (covering a distance of approximately 20-40 meters). When used to treat pulmonary arterial
15 hypertension the compound may be administered in combination with other therapeutic agents such as α -adrenergic receptor antagonists, β_1 -adrenergic receptor antagonists, β_2 -adrenergic receptor agonists, angiotensin-converting enzyme inhibitors, anticoagulants, calcium channel blockers, diuretics, endothelin receptor antagonists, PDE-V inhibitors, prostaglandin analogs, selective serotonin reuptake inhibitors, and combinations thereof.
20 In one particular embodiment of the invention, a compound of the invention is combined with a PDE-V inhibitor or a selective serotonin reuptake inhibitor and used to treat pulmonary arterial hypertension.

Another embodiment of the invention relates to a method for treating heart failure, in particular congestive heart failure (including both systolic and diastolic congestive heart
25 failure), comprising administering to a patient a therapeutically effective amount of a compound of the invention. Typically, the therapeutically effective amount is the amount that is sufficient to lower blood pressure and/or improve renal functions. In a clinical setting, the therapeutically effective amount can be the amount that is sufficient to improve cardiac hemodynamics, like for instance reduction in wedge pressure, right atrial pressure,
30 filling pressure, and vascular resistance. In one embodiment, the compound is administered as an intravenous dosage form. When used to treat heart failure, the compound may be administered in combination with other therapeutic agents such as adenosine receptor antagonists, advanced glycation end product breakers, aldosterone

antagonists, AT₁ receptor antagonists, β_1 -adrenergic receptor antagonists, dual-acting β -adrenergic receptor antagonist/ α_1 -receptor antagonists, chymase inhibitors, digoxin, diuretics, endothelin converting enzyme (ECE) inhibitors, endothelin receptor antagonists, natriuretic peptides and their analogs, natriuretic peptide clearance receptor antagonists, nitric oxide donors, prostaglandin analogs, PDE-V inhibitors, soluble guanylate cyclase activators and stimulators, and vasopressin receptor antagonists. In one particular embodiment of the invention, a compound of the invention is combined with an aldosterone antagonist, a β_1 -adrenergic receptor antagonist, an AT₁ receptor antagonist, or a diuretic, and used to treat congestive heart failure.

10 *Diarrhea*

As NEP inhibitors, compounds of the invention are expected to inhibit the degradation of endogenous enkephalins and thus such compounds may also find utility for the treatment of diarrhea, including infectious and secretory/watery diarrhea. See, for example, Baumer et al. (1992) *Gut* 33:753-758; Farthing (2006) *Digestive Diseases* 24:47-58; and Marçais-Collado (1987) *Eur. J. Pharmacol.* 144(2):125-132. When used to treat diarrhea, compounds of the invention may be combined with one or more additional antidiarrheal agents.

Renal Diseases

By potentiating the effects of vasoactive peptides like the natriuretic peptides and bradykinin, compounds of the invention are expected to enhance renal function (see Chen et al. (1999) *Circulation* 100:2443-2448; Lipkin et al. (1997) *Kidney Int.* 52:792-801; and Dussaule et al. (1993) *Clin. Sci.* 84:31-39) and find utility in treating and/or preventing renal diseases. Renal diseases of particular interest include diabetic nephropathy, chronic kidney disease, proteinuria, and particularly acute kidney injury or acute renal failure (see Sharkovska et al. (2011) *Clin. Lab.* 57:507-515 and Newaz et al. (2010) *Renal Failure* 32:384-390). When used to treat renal disease, the compound may be administered in combination with other therapeutic agents such as angiotensin-converting enzyme inhibitors, AT₁ receptor antagonists, and diuretics.

Preventative Therapy

By potentiating the effects of the natriuretic peptides, compounds of the invention are also expected to be useful in preventative therapy, due to the antihypertrophic and antifibrotic effects of the natriuretic peptides (see Potter et al. (2009) *Handbook of Experimental Pharmacology* 191:341-366), for example in preventing the progression of

cardiac insufficiency after myocardial infarction, preventing arterial restenosis after angioplasty, preventing thickening of blood vessel walls after vascular operations, preventing atherosclerosis, and preventing diabetic angiopathy.

Glaucoma

5 By potentiating the effects of the natriuretic peptides, compounds of the invention are expected to be useful to treat glaucoma. See, for example, Diestelhorst et al. (1989) *International Ophthalmology* 12:99-101. When used to treat glaucoma, compounds of the invention may be combined with one or more additional antiglaucoma agents.

Pain Relief

10 As NEP inhibitors, compounds of the invention are expected to inhibit the degradation of endogenous enkephalins and thus such compounds may also find utility as analgesics. See, for example, Roques et al. (1980) *Nature* 288:286-288 and Thanawala et al. (2008) *Current Drug Targets* 9:887-894. When used to treat pain, the compounds of the invention may be combined with one or more additional antinociceptive drugs such as
15 aminopeptidase N or dipeptidyl peptidase III inhibitors, non-steroidal anti-inflammatory agents, monoamine reuptake inhibitors, muscle relaxants, NMDA receptor antagonists, opioid receptor agonists, 5-HT_{1D} serotonin receptor agonists, and tricyclic antidepressants.

Other Utilities

Due to their NEP inhibition properties, compounds of the invention are also
20 expected to be useful as antitussive agents, as well as find utility in the treatment of portal hypertension associated with liver cirrhosis (see Sansoe et al. (2005) *J. Hepatol.* 43:791-798), cancer (see Vesely (2005) *J. Investigative Med.* 53:360-365), depression (see Noble et al. (2007) *Exp. Opin. Ther. Targets* 11:145-159), menstrual disorders, preterm labor, pre-eclampsia, endometriosis, reproductive disorders (for example, male and female
25 infertility, polycystic ovarian syndrome, implantation failure), and male and female sexual dysfunction, including male erectile dysfunction and female sexual arousal disorder. More specifically, the compounds of the invention are expected to be useful in treating female sexual dysfunction (see Pryde et al. (2006) *J. Med. Chem.* 49:4409-4424), which is often defined as a female patient's difficulty or inability to find satisfaction in sexual expression.
30 This covers a variety of diverse female sexual disorders including, by way of illustration and not limitation, hypoactive sexual desire disorder, sexual arousal disorder, orgasmic disorder and sexual pain disorder. When used to treat such disorders, especially female sexual dysfunction, compounds of the invention may be combined with one or more of the

following secondary agents: PDE-V inhibitors, dopamine agonists, estrogen receptor agonists and/or antagonists, androgens, and estrogens. Due to their NEP inhibition properties, compounds of the invention are also expected to have anti-inflammatory properties, and are expected to have utility as such, particularly when used in combination
5 with statins.

Recent studies suggest that NEP plays a role in regulating nerve function in insulin-deficient diabetes and diet induced obesity. Coppey et al. (2011) *Neuropharmacology* 60:259-266. Therefore, due to their NEP inhibition properties, compounds of the invention are also expected to be useful in providing protection from nerve impairment caused by
10 diabetes or diet induced obesity.

The amount of the compound of the invention administered per dose or the total amount administered per day may be predetermined or it may be determined on an individual patient basis by taking into consideration numerous factors, including the nature and severity of the patient's condition, the condition being treated, the age, weight, and
15 general health of the patient, the tolerance of the patient to the active agent, the route of administration, pharmacological considerations such as the activity, efficacy, pharmacokinetics and toxicology profiles of the compound and any secondary agents being administered, and the like. Treatment of a patient suffering from a disease or medical condition (such as hypertension) can begin with a predetermined dosage or a dosage
20 determined by the treating physician, and will continue for a period of time necessary to prevent, ameliorate, suppress, or alleviate the symptoms of the disease or medical condition. Patients undergoing such treatment will typically be monitored on a routine basis to determine the effectiveness of therapy. For example, in treating hypertension, blood pressure measurements may be used to determine the effectiveness of treatment.
25 Similar indicators for other diseases and conditions described herein, are well known and are readily available to the treating physician. Continuous monitoring by the physician will insure that the optimal amount of the compound of the invention will be administered at any given time, as well as facilitating the determination of the duration of treatment. This is of particular value when secondary agents are also being administered, as their selection,
30 dosage, and duration of therapy may also require adjustment. In this way, the treatment regimen and dosing schedule can be adjusted over the course of therapy so that the lowest amount of active agent that exhibits the desired effectiveness is administered and, further, that administration is continued only so long as is necessary to successfully treat the

disease or medical condition.

Research Tools

Since compounds of the invention possess NEP enzyme inhibition activity, such compounds are also useful as research tools for investigating or studying biological systems or samples having a NEP enzyme, for example to study diseases where the NEP enzyme or its peptide substrates plays a role. Any suitable biological system or sample having a NEP enzyme may be employed in such studies which may be conducted either *in vitro* or *in vivo*. Representative biological systems or samples suitable for such studies include, but are not limited to, cells, cellular extracts, plasma membranes, tissue samples, isolated organs, mammals (such as mice, rats, guinea pigs, rabbits, dogs, pigs, humans, and so forth), and the like, with mammals being of particular interest. In one particular embodiment of the invention, NEP enzyme activity in a mammal is inhibited by administering a NEP-inhibiting amount of a compound of the invention. Compounds of the invention can also be used as research tools by conducting biological assays using such compounds.

When used as a research tool, a biological system or sample comprising a NEP enzyme is typically contacted with a NEP enzyme-inhibiting amount of a compound of the invention. After the biological system or sample is exposed to the compound, the effects of inhibiting the NEP enzyme are determined using conventional procedures and equipment, such as by measuring receptor binding in a binding assay or measuring ligand-mediated changes in a functional assay. Exposure encompasses contacting cells or tissue with the compound, administering the compound to a mammal, for example by *i.p.*, *p.o.*, *i.v.*, *s.c.*, or inhaled administration, and so forth. This determining step can involve measuring a response (a quantitative analysis) or can involve making an observation (a qualitative analysis). Measuring a response involves, for example, determining the effects of the compound on the biological system or sample using conventional procedures and equipment, such as enzyme activity assays and measuring enzyme substrate or product mediated changes in functional assays. The assay results can be used to determine the activity level as well as the amount of compound necessary to achieve the desired result, that is, a NEP enzyme-inhibiting amount. Typically, the determining step will involve determining the effects of inhibiting the NEP enzyme.

Additionally, compounds of the invention can be used as research tools for evaluating other chemical compounds, and thus are also useful in screening assays to

discover, for example, new compounds having NEP-inhibiting activity. In this manner, a compound of the invention is used as a standard in an assay to allow comparison of the results obtained with a test compound and with compounds of the invention to identify those test compounds that have about equal or superior activity, if any. For example, pK_i data for a test compound or a group of test compounds is compared to the pK_i data for a compound of the invention to identify those test compounds that have the desired properties, for example, test compounds having a pK_i value about equal or superior to a compound of the invention, if any. This aspect of the invention includes, as separate embodiments, both the generation of comparison data (using the appropriate assays) and the analysis of test data to identify test compounds of interest. Thus, a test compound can be evaluated in a biological assay, by a method comprising the steps of: (a) conducting a biological assay with a test compound to provide a first assay value; (b) conducting the biological assay with a compound of the invention to provide a second assay value; wherein step (a) is conducted either before, after or concurrently with step (b); and (c) comparing the first assay value from step (a) with the second assay value from step (b). Exemplary biological assays include a NEP enzyme inhibition assay.

PHARMACEUTICAL COMPOSITIONS AND FORMULATIONS

Compounds of the invention are typically administered to a patient in the form of a pharmaceutical composition or formulation. Such pharmaceutical compositions may be administered to the patient by any acceptable route of administration including, but not limited to, oral, rectal, vaginal, nasal, inhaled, topical (including transdermal), ocular, and parenteral modes of administration. Further, the compounds of the invention may be administered, for example orally, in multiple doses per day (for example, two, three, or four times daily), in a single daily dose or a single weekly dose. It will be understood that any form of the compounds of the invention, (that is, free base, free acid, pharmaceutically acceptable salt, solvate, etc.) that is suitable for the particular mode of administration can be used in the pharmaceutical compositions discussed herein.

Accordingly, in one embodiment, the invention relates to a pharmaceutical composition comprising a pharmaceutically acceptable carrier and a compound of the invention. The compositions may contain other therapeutic and/or formulating agents if desired. When discussing compositions, the "compound of the invention" may also be referred to herein as the "active agent," to distinguish it from other components of the formulation, such as the carrier. Thus, it is understood that the term "active agent" includes

compounds of formula I as well as pharmaceutically acceptable salts, solvates and prodrugs of that compound.

The pharmaceutical compositions of the invention typically contain a therapeutically effective amount of a compound of the invention. Those skilled in the art will recognize, however, that a pharmaceutical composition may contain more than a therapeutically effective amount, such as in bulk compositions, or less than a therapeutically effective amount, that is, individual unit doses designed for multiple administration to achieve a therapeutically effective amount. Typically, the composition will contain from about 0.01-95 wt% of active agent, including, from about 0.01-30 wt%, such as from about 0.01-10 wt%, with the actual amount depending upon the formulation itself, the route of administration, the frequency of dosing, and so forth. In one embodiment, a composition suitable for an oral dosage form, for example, may contain about 5-70 wt%, or from about 10-60 wt% of active agent.

Any conventional carrier or excipient may be used in the pharmaceutical compositions of the invention. The choice of a particular carrier or excipient, or combinations of carriers or excipients, will depend on the mode of administration being used to treat a particular patient or type of medical condition or disease state. In this regard, the preparation of a suitable composition for a particular mode of administration is well within the scope of those skilled in the pharmaceutical arts. Additionally, carriers or excipients used in such compositions are commercially available. By way of further illustration, conventional formulation techniques are described in *Remington: The Science and Practice of Pharmacy*, 20th Edition, Lippincott Williams & White, Baltimore, Maryland (2000); and H. C. Ansel et al., *Pharmaceutical Dosage Forms and Drug Delivery Systems*, 7th Edition, Lippincott Williams & White, Baltimore, Maryland (1999).

Representative examples of materials which can serve as pharmaceutically acceptable carriers include, but are not limited to, the following: sugars, such as lactose, glucose and sucrose; starches, such as corn starch and potato starch; cellulose, such as microcrystalline cellulose, and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; powdered tragacanth; malt; gelatin; talc; excipients, such as cocoa butter and suppository waxes; oils, such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; glycols, such as propylene glycol; polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol; esters, such as ethyl oleate and ethyl laurate; agar; buffering agents, such as magnesium hydroxide and

aluminum hydroxide; alginic acid; pyrogen-free water; isotonic saline; Ringer's solution; ethyl alcohol; phosphate buffer solutions; compressed propellant gases, such as chlorofluorocarbons and hydrofluorocarbons; and other non-toxic compatible substances employed in pharmaceutical compositions.

5 Pharmaceutical compositions are typically prepared by thoroughly and intimately mixing or blending the active agent with a pharmaceutically acceptable carrier and one or more optional ingredients. The resulting uniformly blended mixture may then be shaped or loaded into tablets, capsules, pills, canisters, cartridges, dispensers and the like using conventional procedures and equipment.

10 In one embodiment, the pharmaceutical compositions are suitable for oral administration. Suitable compositions for oral administration may be in the form of capsules, tablets, pills, lozenges, cachets, dragees, powders, granules; solutions or suspensions in an aqueous or non-aqueous liquid; oil-in-water or water-in-oil liquid emulsions; elixirs or syrups; and the like; each containing a predetermined amount of the
15 active agent.

 When intended for oral administration in a solid dosage form (capsules, tablets, pills and the like), the composition will typically comprise the active agent and one or more pharmaceutically acceptable carriers, such as sodium citrate or dicalcium phosphate. Solid dosage forms may also comprise: fillers or extenders, such as starches, microcrystalline
20 cellulose, lactose, sucrose, glucose, mannitol, and/or silicic acid; binders, such as carboxymethylcellulose, alginates, gelatin, polyvinyl pyrrolidone, sucrose and/or acacia; humectants, such as glycerol; disintegrating agents, such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and/or sodium carbonate; solution retarding agents, such as paraffin; absorption accelerators, such as quaternary ammonium
25 compounds; wetting agents, such as cetyl alcohol and/or glycerol monostearate; absorbents, such as kaolin and/or bentonite clay; lubricants, such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and/or mixtures thereof; coloring agents; and buffering agents.

 Release agents, wetting agents, coating agents, sweetening, flavoring and
30 perfuming agents, preservatives and antioxidants may also be present in the pharmaceutical compositions. Exemplary coating agents for tablets, capsules, pills and like, include those used for enteric coatings, such as cellulose acetate phthalate, polyvinyl acetate phthalate, hydroxypropyl methylcellulose phthalate, methacrylic acid-methacrylic acid ester

copolymers, cellulose acetate trimellitate, carboxymethyl ethyl cellulose, hydroxypropyl methyl cellulose acetate succinate, and the like. Examples of pharmaceutically acceptable antioxidants include: water-soluble antioxidants, such as ascorbic acid, cysteine hydrochloride, sodium bisulfate, sodium metabisulfate sodium sulfite and the like; oil-soluble antioxidants, such as ascorbyl palmitate, butylated hydroxyanisole, butylated hydroxytoluene, lecithin, propyl gallate, alpha-tocopherol, and the like; and metal-chelating agents, such as citric acid, ethylenediamine tetraacetic acid, sorbitol, tartaric acid, phosphoric acid, and the like.

Compositions may also be formulated to provide slow or controlled release of the active agent using, by way of example, hydroxypropyl methyl cellulose in varying proportions or other polymer matrices, liposomes and/or microspheres. In addition, the pharmaceutical compositions of the invention may contain opacifying agents and may be formulated so that they release the active agent only, or preferentially, in a certain portion of the gastrointestinal tract, optionally, in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes. The active agent can also be in micro-encapsulated form, optionally with one or more of the above-described excipients.

Suitable liquid dosage forms for oral administration include, by way of illustration, pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. Liquid dosage forms typically comprise the active agent and an inert diluent, such as, for example, water or other solvents, solubilizing agents and emulsifiers, such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, oils (for example, cottonseed, groundnut, corn, germ, olive, castor and sesame oils), glycerol, tetrahydrofuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. Suspensions may contain suspending agents such as, for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminium metahydroxide, bentonite, agar-agar and tragacanth, and mixtures thereof.

When intended for oral administration, the pharmaceutical compositions of the invention may be packaged in a unit dosage form. The term "unit dosage form" refers to a physically discrete unit suitable for dosing a patient, that is, each unit containing a predetermined quantity of the active agent calculated to produce the desired therapeutic effect either alone or in combination with one or more additional units. For example, such

unit dosage forms may be capsules, tablets, pills, and the like.

In another embodiment, the compositions of the invention are suitable for inhaled administration, and will typically be in the form of an aerosol or a powder. Such compositions are generally administered using well-known delivery devices, such as a nebulizer, dry powder, or metered-dose inhaler. Nebulizer devices produce a stream of high velocity air that causes the composition to spray as a mist that is carried into a patient's respiratory tract. An exemplary nebulizer formulation comprises the active agent dissolved in a carrier to form a solution, or micronized and combined with a carrier to form a suspension of micronized particles of respirable size. Dry powder inhalers administer the active agent as a free-flowing powder that is dispersed in a patient's air-stream during inspiration. An exemplary dry powder formulation comprises the active agent dry-blended with an excipient such as lactose, starch, mannitol, dextrose, polylactic acid, polylactide-co-glycolide, and combinations thereof. Metered-dose inhalers discharge a measured amount of the active agent using compressed propellant gas. An exemplary metered-dose formulation comprises a solution or suspension of the active agent in a liquefied propellant, such as a chlorofluorocarbon or hydrofluoroalkane. Optional components of such formulations include co-solvents, such as ethanol or pentane, and surfactants, such as sorbitan trioleate, oleic acid, lecithin, glycerin, and sodium lauryl sulfate. Such compositions are typically prepared by adding chilled or pressurized hydrofluoroalkane to a suitable container containing the active agent, ethanol (if present) and the surfactant (if present). To prepare a suspension, the active agent is micronized and then combined with the propellant. Alternatively, a suspension formulation can be prepared by spray drying a coating of surfactant on micronized particles of the active agent. The formulation is then loaded into an aerosol canister, which forms a portion of the inhaler.

Compounds of the invention can also be administered parenterally (for example, by subcutaneous, intravenous, intramuscular, or intraperitoneal injection). For such administration, the active agent is provided in a sterile solution, suspension, or emulsion. Exemplary solvents for preparing such formulations include water, saline, low molecular weight alcohols such as propylene glycol, polyethylene glycol, oils, gelatin, fatty acid esters such as ethyl oleate, and the like. Parenteral formulations may also contain one or more anti-oxidants, solubilizers, stabilizers, preservatives, wetting agents, emulsifiers, and dispersing agents. Surfactants, additional stabilizing agents or pH-adjusting agents (acids, bases or buffers) and anti-oxidants are particularly useful to provide stability to the

formulation, for example, to minimize or avoid hydrolysis of ester and amide linkages that may be present in the compound. These formulations may be rendered sterile by use of a sterile injectable medium, a sterilizing agent, filtration, irradiation, or heat. In one particular embodiment, the parenteral formulation comprises an aqueous cyclodextrin solution as the pharmaceutically acceptable carrier. Suitable cyclodextrins include cyclic molecules containing six or more α -D-glucopyranose units linked at the 1,4 positions by a linkages as in amylose, β -cyclodextrin or cycloheptaamylose. Exemplary cyclodextrins include cyclodextrin derivatives such as hydroxypropyl and sulfobutyl ether cyclodextrins such as hydroxypropyl- β -cyclodextrin and sulfobutyl ether β -cyclodextrin. Exemplary buffers for such formulations include carboxylic acid-based buffers such as citrate, lactate and maleate buffer solutions.

Compounds of the invention can also be administered transdermally using known transdermal delivery systems and excipients. For example, the compound can be admixed with permeation enhancers, such as propylene glycol, polyethylene glycol monolaurate, azacycloalkan-2-ones and the like, and incorporated into a patch or similar delivery system. Additional excipients including gelling agents, emulsifiers and buffers, may be used in such transdermal compositions if desired.

Secondary Agents

The compounds of the invention may be useful as the sole treatment of a disease or may be combined with one or more additional therapeutic agents in order to obtain the desired therapeutic effect. Thus, in one embodiment, pharmaceutical compositions of the invention contain other drugs that are co-administered with a compound of the invention. For example, the composition may further comprise one or more drugs (also referred to as "secondary agents(s)"). Such therapeutic agents are well known in the art, and include adenosine receptor antagonists, α -adrenergic receptor antagonists, β_1 -adrenergic receptor antagonists, β_2 -adrenergic receptor agonists, dual-acting β -adrenergic receptor antagonist/ α_1 -receptor antagonists, advanced glycation end product breakers, aldosterone antagonists, aldosterone synthase inhibitors, aminopeptidase N inhibitors, androgens, angiotensin-converting enzyme inhibitors and dual-acting angiotensin-converting enzyme/neprilysin inhibitors, angiotensin-converting enzyme 2 activators and stimulators, angiotensin-II vaccines, anticoagulants, anti-diabetic agents, antidiarrheal agents, anti-glaucoma agents, anti-lipid agents, antinociceptive agents, anti-thrombotic agents, AT_1 receptor antagonists and dual-acting AT_1 receptor antagonist/neprilysin inhibitors and

multifunctional angiotensin receptor blockers, bradykinin receptor antagonists, calcium channel blockers, chymase inhibitors, digoxin, diuretics, dopamine agonists, endothelin converting enzyme inhibitors, endothelin receptor antagonists, HMG-CoA reductase inhibitors, estrogens, estrogen receptor agonists and/or antagonists, monoamine reuptake
5 inhibitors, muscle relaxants, natriuretic peptides and their analogs, natriuretic peptide clearance receptor antagonists, neprilysin inhibitors, nitric oxide donors, non-steroidal anti-inflammatory agents, N-methyl d-aspartate receptor antagonists, opioid receptor agonists, phosphodiesterase inhibitors, prostaglandin analogs, prostaglandin receptor agonists, renin inhibitors, selective serotonin reuptake inhibitors, sodium channel blocker, soluble
10 guanylate cyclase stimulators and activators, tricyclic antidepressants, vasopressin receptor antagonists, and combinations thereof. Specific examples of these agents are detailed herein.

Accordingly, in yet another aspect of the invention, a pharmaceutical composition comprises a compound of the invention, a second active agent, and a pharmaceutically
15 acceptable carrier. Third, fourth etc. active agents may also be included in the composition. In combination therapy, the amount of compound of the invention that is administered, as well as the amount of secondary agents, may be less than the amount typically administered in monotherapy.

Compounds of the invention may be physically mixed with the second active agent
20 to form a composition containing both agents; or each agent may be present in separate and distinct compositions which are administered to the patient simultaneously or at separate times. For example, a compound of the invention can be combined with a second active agent using conventional procedures and equipment to form a combination of active agents comprising a compound of the invention and a second active agent. Additionally, the
25 active agents may be combined with a pharmaceutically acceptable carrier to form a pharmaceutical composition comprising a compound of the invention, a second active agent and a pharmaceutically acceptable carrier. In this embodiment, the components of the composition are typically mixed or blended to create a physical mixture. The physical mixture is then administered in a therapeutically effective amount using any of the routes
30 described herein.

Alternatively, the active agents may remain separate and distinct before administration to the patient. In this embodiment, the agents are not physically mixed together before administration but are administered simultaneously or at separate times as

separate compositions. Such compositions can be packaged separately or may be packaged together in a kit. When administered at separate times, the secondary agent will typically be administered less than 24 hours after administration of the compound of the invention, ranging anywhere from concurrent with administration of the compound of the invention to
5 about 24 hours post-dose. This is also referred to as sequential administration. Thus, a compound of the invention can be orally administered simultaneously or sequentially with another active agent using two tablets, with one tablet for each active agent, where sequential may mean being administered immediately after administration of the compound of the invention or at some predetermined time later (for example, one hour
10 later or three hours later). It is also contemplated that the secondary agent may be administered more than 24 hours after administration of the compound of the invention. Alternatively, the combination may be administered by different routes of administration, that is, one orally and the other by inhalation.

In one embodiment, the kit comprises a first dosage form comprising a compound
15 of the invention and at least one additional dosage form comprising one or more of the secondary agents set forth herein, in quantities sufficient to carry out the methods of the invention. The first dosage form and the second (or third, etc.) dosage form together comprise a therapeutically effective amount of active agents for the treatment or prevention of a disease or medical condition in a patient.

20 Secondary agent(s), when included, are present in a therapeutically effective amount such that they are typically administered in an amount that produces a therapeutically beneficial effect when co-administered with a compound of the invention. The secondary agent can be in the form of a pharmaceutically acceptable salt, solvate, optically pure stereoisomer, and so forth. The secondary agent may also be in the form of
25 a prodrug, for example, a compound having a carboxylic acid group that has been esterified. Thus, secondary agents listed herein are intended to include all such forms, and are commercially available or can be prepared using conventional procedures and reagents.

In one embodiment, compounds of the invention are administered in combination with an adenosine receptor antagonist, representative examples of which include, but are
30 not limited to, naxifylline, rolofylline, SLV-320, theophylline, and tonapofylline.

In one embodiment, compounds of the invention are administered in combination with an α -adrenergic receptor antagonist, representative examples of which include, but are not limited to, doxazosin, prazosin, tamsulosin, and terazosin.

Compounds of the invention may also be administered in combination with a β_1 -adrenergic receptor antagonist (" β_1 -blockers"). Representative β_1 -blockers include, but are not limited to, acebutolol, alprenolol, amosulalol, arotinolol, atenolol, befunolol, betaxolol, bevantolol, bisoprolol, bopindolol, bucindolol, bucumolol, bufetolol, bufuralol, bunitrolol, 5 bupranolol, bubridine, butofilolol, carazolol, carteolol, carvedilol, celiprolol, cetamolol, cloranolol, dilevalol, epanolol, esmolol, indenolol, labetolol, levobunolol, mepindolol, metipranolol, metoprolol such as metoprolol succinate and metoprolol tartrate, moprolool, nadolol, nadoxolol, nebivalol, nipradilol, oxprenolol, penbutolol, perbutolol, pindolol, practolol, pronethalol, propranolol, sotalol, sufinalol, talindol, tertatolol, tilisolol, timolol, 10 toliprolol, xibenolol, and combinations thereof. In one particular embodiment, the β_1 -antagonist is selected from atenolol, bisoprolol, metoprolol, propranolol, sotalol, and combinations thereof. Typically, the β_1 -blocker will be administered in an amount sufficient to provide from about 2-900 mg per dose.

In one embodiment, compounds of the invention are administered in combination 15 with a β_2 -adrenergic receptor agonist, representative examples of which include, but are not limited to, albuterol, bitolterol, fenoterol, formoterol, indacaterol, isocetharine, levalbuterol, metaproterenol, pirbuterol, salbutamol, salmefamol, salmeterol, terbutaline, vilanterol, and the like. Typically, the β_2 -adrenergic receptor agonist will be administered in an amount sufficient to provide from about 0.05-500 μ g per dose.

20 In one embodiment, compounds of the invention are administered in combination with an advanced glycation end product (AGE) breaker, examples of which include, by way of illustration and not limitation, alagebrium (or ALT-711), and TRC4149.

In another embodiment, compounds of the invention are administered in combination with an aldosterone antagonist, representative examples of which include, but 25 are not limited to, eplerenone, spironolactone, and combinations thereof. Typically, the aldosterone antagonist will be administered in an amount sufficient to provide from about 5-300 mg per day.

In one embodiment, compounds of the invention are administered in combination with an aminopeptidase N or dipeptidyl peptidase III inhibitor, examples of which include, 30 by way of illustration and not limitation, bestatin and PC18 (2-amino-4-methylsulfonyl butane thiol, methionine thiol).

Compounds of the invention can also be administered in combination with an angiotensin-converting enzyme (ACE) inhibitor. Representative ACE inhibitors include,

but are not limited to, accupril, alacepril, benazepril, benazeprilat, captopril, ceranapril, cilazapril, delapril, enalapril, enalaprilat, fosinopril, fosinoprilat, imidapril, lisinopril, moexipril, monopril, moveltopril, pentopril, perindopril, quinapril, quinaprilat, ramipril, ramiprilat, saralasin acetate, spirapril, temocapril, trandolapril, zofenopril, and

- 5 combinations thereof. In a particular embodiment, the ACE inhibitor is selected from: benazepril, captopril, enalapril, lisinopril, ramipril, and combinations thereof. Typically, the ACE inhibitor will be administered in an amount sufficient to provide from about 1-150 mg per day.

- In another embodiment, compounds of the invention are administered in
 10 combination with a dual-acting angiotensin-converting enzyme/neprilysin (ACE/NEP) inhibitor, examples of which include, but are not limited to: AVE-0848 ((4*S*,7*S*,12*bR*)-7-[3-methyl-2(*S*)-sulfanylbutyramido]-6-oxo-1,2,3,4,6,7,8,12*b*-octahydropyrido[2,1-*a*][2]-benzazepine-4-carboxylic acid); AVE-7688 (ilepatril) and its parent compound; BMS-182657 (2-[2-oxo-3(*S*)-[3-phenyl-2(*S*)-sulfanylpropionamido]-2,3,4,5-tetrahydro-1*H*-1-
 15 benzazepin-1-yl]acetic acid); CGS-26303 ([N-[2-(biphenyl-4-yl)-1(*S*)-(1*H*-tetrazol-5-yl)ethyl]amino]methylphosphonic acid); CGS-35601 (N-[1-[4-methyl-2(*S*)-sulfanylpentanamido]cyclopentylcarbonyl]-L-tryptophan); fasidotril; fasidotrilate; enalaprilat; ER-32935 ((3*R*,6*S*,9*aR*)-6-[3(*S*)-methyl-2(*S*)-sulfanylpentanamido]-5-oxoperhydrothiazolo[3,2-*a*]azepine-3-carboxylic acid); gempatrilat; MDL-101264
 20 ((4*S*,7*S*,12*bR*)-7-[2(*S*)-(2-morpholinoacetylthio)-3-phenylpropionamido]-6-oxo-1,2,3,4,6,7,8,12*b*-octahydropyrido[2,1-*a*][2]benzazepine-4-carboxylic acid); MDL-101287 ([4*S*-[4*α*,7*α*(*R**),12*bβ*]]-7-[2-(carboxymethyl)-3-phenylpropionamido]-6-oxo-1,2,3,4,6,7,8,12*b*-octahydropyrido[2,1-*a*][2]benzazepine-4-carboxylic acid); omapatrilat; RB-105 (N-[2(*S*)-(mercaptomethyl)-3(*R*)-phenylbutyl]-L-alanine); sampatrilat; SA-898
 25 ((2*R*,4*R*)-N-[2-(2-hydroxyphenyl)-3-(3-mercaptopropionyl)thiazolidin-4-ylcarbonyl]-L-phenylalanine); Sch-50690 (N-[1(*S*)-carboxy-2-[N2-(methanesulfonyl)-L-lysylamino]ethyl]-L-valyl-L-tyrosine); and combinations thereof, may also be included. In one particular embodiment, the ACE/NEP inhibitor is selected from: AVE-7688, enalaprilat, fasidotril, fasidotrilate, omapatrilat, sampatrilat, and combinations thereof.

- 30 In one embodiment, compounds of the invention are administered in combination with an angiotensin-converting enzyme 2 (ACE2) activator or stimulator.

In one embodiment, compounds of the invention are administered in combination with an angiotensin-II vaccine, examples of which include, but are not limited to

ATR12181 and CYT006-AngQb.

In one embodiment, compounds of the invention are administered in combination with an anticoagulant, representative examples of which include, but are not limited to: coumarins such as warfarin; heparin; and direct thrombin inhibitors such as argatroban,
5 bivalirudin, dabigatran, and lepirudin.

In yet another embodiment, compounds of the invention are administered in combination with an anti-diabetic agent. Representative anti-diabetic agents include injectable drugs as well as orally effective drugs, and combinations thereof. Examples of injectable drugs include, but are not limited to, insulin and insulin derivatives. Examples
10 of orally effective drugs include, but are not limited to: biguanides such as metformin; glucagon antagonists; α -glucosidase inhibitors such as acarbose and miglitol; dipeptidyl peptidase IV inhibitors (DPP-IV inhibitors) such as alogliptin, denagliptin, linagliptin, saxagliptin, sitagliptin, and vildagliptin; meglitinides such as repaglinide; oxadiazolidinediones; sulfonylureas such as chlorpropamide, glimepiride, glipizide,
15 glyburide, and tolazamide; thiazolidinediones such as pioglitazone and rosiglitazone; and combinations thereof.

In another embodiment, compounds of the invention are administered in combination with antidiarrheal treatments. Representative treatment options include, but are not limited to, oral rehydration solutions (ORS), loperamide, diphenoxylate, and
20 bismuth subsalicylate.

In yet another embodiment, a compound of the invention is administered in combination with an anti-glaucoma agent. Representative anti-glaucoma agents include, but are not limited to: α -adrenergic agonists such as brimonidine; β_1 -adrenergic receptor antagonists; topical β_1 -blockers such as betaxolol, levobunolol, and timolol; carbonic
25 anhydrase inhibitors such as acetazolamide, brinzolamide, or dorzolamide; cholinergic agonists such as cevimeline and DMXB-anabaseine; epinephrine compounds; miotics such as pilocarpine; and prostaglandin analogs.

In yet another embodiment, compounds of the invention are administered in combination with an anti-lipid agent. Representative anti-lipid agents include, but are not
30 limited to: cholesteryl ester transfer protein inhibitors (CETPs) such as anacetrapib, dalcetrapib, and torcetrapib; statins such as atorvastatin, fluvastatin, lovastatin, pravastatin, rosuvastatin and simvastatin; and combinations thereof.

In one embodiment, compounds of the invention are administered in combination

with an anti-thrombotic agent. Representative anti-thrombotic agents include, but are not limited to: aspirin; anti-platelet agents such as clopidogrel, prasugrel, and ticlopidine; heparin, and combinations thereof.

In one embodiment, compounds of the invention are administered in combination
5 with an AT₁ receptor antagonist, also known as angiotensin II type I receptor blockers (ARBs). Representative ARBs include, but are not limited to, abitesartan, azilsartan (e.g., azilsartan medoxomil), benzylosartan, candesartan, candesartan cilexetil, elisartan, embusartan, enoltasartan, eprosartan, EXP3174, fonsartan, forasartan, glycylosartan, irbesartan, isoteoline, losartan, medoximil, milfasartan, olmesartan (e.g., olmesartan
10 medoxomil), opomisartan, prazosartan, ripisartan, saprisartan, saralasin, sarmesin, TAK-591, tasosartan, telmisartan, valsartan, zolasartan, and combinations thereof. In a particular embodiment, the ARB is selected from azilsartan medoxomil, candesartan cilexetil, eprosartan, irbesartan, losartan, olmesartan medoxomil, irbesartan, saprisartan, tasosartan, telmisartan, valsartan, and combinations thereof. Exemplary salts and/or prodrugs include
15 candesartan cilexetil, eprosartan mesylate, losartan potassium salt, and olmesartan medoxomil. Typically, the ARB will be administered in an amount sufficient to provide from about 4-600 mg per dose, with exemplary daily dosages ranging from 20-320 mg per day.

Compounds of the invention may also be administered in combination with a dual-
20 acting agent, such as an AT₁ receptor antagonist/neprilysin inhibitor (ARB/NEP) inhibitor, examples of which include, but are not limited to, compounds described in U.S. Publication Nos. 2008/0269305 and 2009/0023228, both to Allegretti et al. filed on April 23, 2008, such as the compound, 4'-{2-ethoxy-4-ethyl-5-[(S)-2-mercapto-4-methylpentanoylamino)-methyl]imidazol-1-ylmethyl}-3'-fluorobiphenyl-2-carboxylic acid.

25 Compounds of the invention may also be administered in combination with multifunctional angiotensin receptor blockers as described in Kurtz & Klein (2009) *Hypertension Research* 32:826-834.

In one embodiment, compounds of the invention are administered in combination with a bradykinin receptor antagonist, for example, icatibant (HOE-140). It is expected
30 that this combination therapy may present the advantage of preventing angioedema or other unwanted consequences of elevated bradykinin levels.

In one embodiment, compounds of the invention are administered in combination with a calcium channel blocker. Representative calcium channel blockers include, but are

not limited to, amlodipine, anipamil, aranipine, barnidipine, bencyclane, benidipine, bepridil, clentiazem, cilnidipine, cinnarizine, diltiazem, efonidipine, elgodipine, etafenone, felodipine, fendiline, flunarizine, gallopamil, isradipine, lacidipine, lercanidipine, lidoflazine, lomerizine, manidipine, mibefradil, nicardipine, nifedipine, niguldipine, 5 niludipine, nilvadipine, nimodipine, nisoldipine, nitrendipine, nivaldipine, perhexiline, prenylamine, ryosidine, semotiadil, terodiline, tiapamil, verapamil, and combinations thereof. In a particular embodiment, the calcium channel blocker is selected from amlodipine, bepridil, diltiazem, felodipine, isradipine, lacidipine, nicardipine, nifedipine, niguldipine, niludipine, nimodipine, nisoldipine, ryosidine, verapamil, and combinations 10 thereof. Typically, the calcium channel blocker will be administered in an amount sufficient to provide from about 2-500 mg per dose.

In one embodiment, compounds of the invention are administered in combination with a chymase inhibitor, such as TPC-806 and 2-(5-formylamino-6-oxo-2-phenyl-1,6-dihydropyrimidine-1-yl)-N-[{3,4-dioxo-1-phenyl-7-(2-pyridyloxy)}-2-heptyl]acetamide 15 (NK3201).

In one embodiment, compounds of the invention are administered in combination with a diuretic. Representative diuretics include, but are not limited to: carbonic anhydrase inhibitors such as acetazolamide and dichlorphenamide; loop diuretics, which include sulfonamide derivatives such as acetazolamide, ambuside, azosernide, bumetanide, 20 butazolamide, chloraminophenamide, clofenamide, clopamide, clorexolone, disulfamide, ethoxolamide, furosemide, mefruside, methazolamide, pirtanide, torsemide, tripamide, and xipamide, as well as non-sulfonamide diuretics such as ethacrynic acid and other phenoxyacetic acid compounds such as tienilic acid, indacrinone and quincarbate; osmotic diuretics such as mannitol; potassium-sparing diuretics, which include aldosterone 25 antagonists such as spironolactone, and Na⁺ channel inhibitors such as amiloride and triamterene; thiazide and thiazide-like diuretics such as althiazide, bendroflumethiazide, benzylhydrochlorothiazide, benzthiazide, buthiazide, chlorthalidone, chlorothiazide, cyclopenthiazide, cyclothiazide, epithiazide, ethiazide, fenquizone, flumethiazide, hydrochlorothiazide, hydroflumethiazide, indapamide, methylclothiazide, meticrane, 30 metolazone, paraflutizide, polythiazide, quinethazone, teclothiazide, and trichloromethiazide; and combinations thereof. In a particular embodiment, the diuretic is selected from amiloride, bumetanide, chlorothiazide, chlorthalidone, dichlorphenamide, ethacrynic acid, furosemide, hydrochlorothiazide, hydroflumethiazide, indapamide,

methylclothiazide, metolazone, torsemide, triamterene, and combinations thereof. The diuretic will be administered in an amount sufficient to provide from about 5-50 mg per day, more typically 6-25 mg per day, with common dosages being 6.25 mg, 12.5 mg or 25 mg per day.

- 5 Compounds of the invention may also be administered in combination with an endothelin converting enzyme (ECE) inhibitor, examples of which include, but are not limited to, phosphoramidon, CGS 26303, and combinations thereof.

 In a particular embodiment, compounds of the invention are administered in combination with an endothelin receptor antagonist. Representative endothelin receptor
10 antagonists include, but are not limited to: selective endothelin receptor antagonists that affect endothelin A receptors, such as avosentan, ambrisentan, atrasentan, BQ-123, clazosentan, darusentan, sitaxentan, and zibotentan; and dual endothelin receptor antagonists that affect both endothelin A and B receptors, such as bosentan, macitentan, tezosentan).

- 15 In yet another embodiment, a compound of the invention is administered in combination with one or more HMG-CoA reductase inhibitors, which are also known as statins. Representative statins include, but are not limited to, atorvastatin, fluvastatin, lovastatin, pitavastatin, pravastatin, rosuvastatin and simvastatin.

 In one embodiment, compounds of the invention are administered in combination
20 with a monoamine reuptake inhibitor, examples of which include, by way of illustration and not limitation, norepinephrine reuptake inhibitors such as atomoxetine, bupropion and the bupropion metabolite hydroxybupropion, maprotiline, reboxetine, and viloxazine; selective serotonin reuptake inhibitors (SSRIs) such as citalopram and the citalopram metabolite desmethylcitalopram, dapoxetine, escitalopram (e.g., escitalopram oxalate),
25 fluoxetine and the fluoxetine desmethyl metabolite norfluoxetine, fluvoxamine (e.g., fluvoxamine maleate), paroxetine, sertraline and the sertraline metabolite demethylsertraline; dual serotonin-norepinephrine reuptake inhibitors (SNRIs) such as bicifadine, duloxetine, milnacipran, nefazodone, and venlafaxine; and combinations thereof.

- 30 In another embodiment, compounds of the invention are administered in combination with a muscle relaxant, examples of which include, but are not limited to: carisoprodol, chlorzoxazone, cyclobenzaprine, diflunisal, metaxalone, methocarbamol, and combinations thereof.

In one embodiment, compounds of the invention are administered in combination with a natriuretic peptide or analog, examples of which include but are not limited to: carperitide, CD-NP (Nile Therapeutics), CU-NP, nesiritide, PL-3994 (Palatin Technologies, Inc.), ularitide, cenderitide, and compounds described in Ogawa et al (2004) *J. Biol. Chem.* 279:28625-31. These compounds are also referred to as natriuretic peptide receptor-A (NPR-A) agonists. In another embodiment, compounds of the invention are administered in combination with a natriuretic peptide clearance receptor (NPR-C) antagonist such as SC-46542, cANF (4-23), and AP-811 (Veale (2000) *Bioorg Med Chem Lett* 10:1949-52). For example, AP-811 has shown synergy when combined with the NEP inhibitor, thiorphan (Wegner (1995) *Clin. Exper. Hypert.* 17:861-876).

In another embodiment, compounds of the invention are administered in combination with a neprilysin (NEP) inhibitor. Representative NEP inhibitors include, but are not limited to: AHU-377; candoxatril; candoxatrilat; dexecadotril ((+)-N-[2(R)-(acetylthiomethyl)-3-phenylpropionyl]glycine benzyl ester); CGS-24128 (3-[3-(biphenyl-4-yl)-2-(phosphonomethylamino)propionamido]propionic acid); CGS-24592 ((S)-3-[3-(biphenyl-4-yl)-2-(phosphonomethylamino)propionamido]propionic acid); CGS-25155 (N-[9(R)-(acetylthiomethyl)-10-oxo-1-azacyclodecan-2(S)-ylcarbonyl]-4(R)-hydroxy-L-proline benzyl ester); 3-(1-carbamoylcyclohexyl)propionic acid derivatives described in WO 2006/027680 to Hepworth et al. (Pfizer Inc.); JMV-390-1 (2(R)-benzyl-3-(N-hydroxycarbamoyl)propionyl-L-isoleucyl-L-leucine); ecadotril; phosphoramidon; retrothiorphan; RU-42827 (2-(mercaptomethyl)-N-(4-pyridinyl)benzenepropionamide); RU-44004 (N-(4-morpholinyl)-3-phenyl-2-(sulfanylmethyl)propionamide); SCH-32615 ((S)-N-[N-(1-carboxy-2-phenylethyl)-L-phenylalanyl]-β-alanine) and its prodrug SCH-34826 ((S)-N-[N-[1-[(2,2-dimethyl-1,3-dioxolan-4-yl)methoxy]carbonyl]-2-phenylethyl]-L-phenylalanyl]-β-alanine); sialorphan; SCH-42495 (N-[2(S)-(acetylsulfanylmethyl)-3-(2-methylphenyl)propionyl]-L-methionine ethyl ester); spinorphin; SQ-28132 (N-[2-(mercaptomethyl)-1-oxo-3-phenylpropyl]leucine); SQ-28603 (N-[2-(mercaptomethyl)-1-oxo-3-phenylpropyl]-β-alanine); SQ-29072 (7-[[2-(mercaptomethyl)-1-oxo-3-phenylpropyl]amino]heptanoic acid); thiorphan and its prodrug racecadotril; UK-69578 (cis-4-[[[1-[2-carboxy-3-(2-methoxyethoxy)propyl]cyclopentyl]carbonyl]amino]cyclohexanecarboxylic acid); UK-447,841 (2-{1-[3-(4-chlorophenyl)propylcarbamoyl]-cyclopentylmethyl}-4-methoxybutyric acid); UK-505,749 ((R)-2-methyl-3-{1-[3-(2-methylbenzothiazol-6-yl)propylcarbamoyl]cyclopentyl}propionic acid); 5-biphenyl-4-yl-4-

(3-carboxypropionylamino)-2-methylpentanoic acid and 5-biphenyl-4-yl-4-(3-carboxypropionylamino)-2-methylpentanoic acid ethyl ester (WO 2007/056546); daglutril [(3*S*,2'*R*)-3-{1-[2'-(ethoxycarbonyl)-4'-phenylbutyl]-cyclopentan-1-carbonylamino}-2,3,4,5-tetrahydro-2-oxo-1*H*-1-benzazepine-1-acetic acid] described in

5 WO 2007/106708 to Khder et al. (Novartis AG); and combinations thereof. In a particular embodiment, the NEP inhibitor is selected from AHU-377, candoxatril, candoxatrilat, CGS-24128, phosphoramidon, SCH-32615, SCH-34826, SQ-28603, thiorphan, and combinations thereof. In a particular embodiment, the NEP inhibitor is a compound such as daglutril, which also has activity both as an inhibitor of the endothelin converting

10 enzyme (ECE) and of NEP. Other dual acting ECE/NEP compounds can also be used. The NEP inhibitor will be administered in an amount sufficient to provide from about 20-800 mg per day, with typical daily dosages ranging from 50-700 mg per day, more commonly 100-600 or 100-300 mg per day.

In one embodiment, compounds of the invention are administered in combination

15 with a nitric oxide donor, examples of which include, but are not limited to nicorandil; organic nitrates such as pentaerythritol tetranitrate; and sydnonimines such as linsidomine and molsidomine.

In yet another embodiment, compounds of the invention are administered in combination with a non-steroidal anti-inflammatory agent (NSAID). Representative

20 NSAIDs include, but are not limited to: acetaminophen, acetylsalicylic acid, alclofenac, alminoprofen, amfenac, amiprilose, amoxicillin, anilofenac, apazone, azapropazone, benorilate, benoxaprofen, bezpiperylon, broperamol, bucloxic acid, carprofen, clidanac, diclofenac, diflunisal, diflunolone, enolicam, etodolac, etoricoxib, fenbufen, fenclofenac, fenclozic acid, fenoprofen, fentiazac, feprazole, flufenamic acid, flufenisal, fluprofen,

25 flurbiprofen, furofenac, ibufenac, ibuprofen, indomethacin, indoprofen, isoxepac, isoxicam, ketoprofen, ketorolac, lofenizole, lornoxicam, meclofenamate, meclofenamic acid, mefenamic acid, meloxicam, mesalamine, miroprofen, mofebutazone, nabumetone, naproxen, niflumic acid, oxaprozin, oxipinac, oxyphenbutazone, phenylbutazone, piroxicam, piroprofen, pranoprofen, salsalate, sudoxicam, sulfasalazine, sulindac, suprofen,

30 tenoxicam, tiopinac, tiaprofenic acid, tioprofen, tolafenamic acid, tolmetin, triflumidate, zidometacin, zomepirac, and combinations thereof. In a particular embodiment, the NSAID is selected from etodolac, flurbiprofen, ibuprofen, indomethacin, ketoprofen, ketorolac, meloxicam, naproxen, oxaprozin, piroxicam, and combinations thereof.

In one embodiment, compounds of the invention are administered in combination with an N-methyl d-aspartate (NMDA) receptor antagonist, examples of which include, by way of illustration and not limitation, including amantadine, dextromethorphan, dextropropoxyphene, ketamine, ketobemidone, memantine, methadone, and so forth.

5 In still another embodiment, compounds of the invention are administered in combination with an opioid receptor agonist (also referred to as opioid analgesics). Representative opioid receptor agonists include, but are not limited to: buprenorphine, butorphanol, codeine, dihydrocodeine, fentanyl, hydrocodone, hydromorphone, levallorphan, levorphanol, meperidine, methadone, morphine, nalbuphine, nalmefene, 10 nalorphine, naloxone, naltrexone, nalorphine, oxycodone, oxymorphone, pentazocine, propoxyphene, tramadol, and combinations thereof. In certain embodiments, the opioid receptor agonist is selected from codeine, dihydrocodeine, hydrocodone, hydromorphone, morphine, oxycodone, oxymorphone, tramadol, and combinations thereof.

In a particular embodiment, compounds of the invention are administered in 15 combination with a phosphodiesterase (PDE) inhibitor, particularly a PDE-V inhibitor. Representative PDE-V inhibitors include, but are not limited to, avanafil, lodenafil, mirodenafil, sildenafil (Revatio[®]), tadalafil (Adcirca[®]), vardenafil (Levitra[®]), and udenafil.

In another embodiment, compounds of the invention are administered in combination with a prostaglandin analog (also referred to as prostanoids or prostacyclin 20 analogs). Representative prostaglandin analogs include, but are not limited to, beraprost sodium, bimatoprost, epoprostenol, iloprost, latanoprost, tafluprost, travoprost, and treprostinil, with bimatoprost, latanoprost, and tafluprost being of particular interest.

In yet another embodiment, compounds of the invention are administered in combination with a prostaglandin receptor agonist, examples of which include, but are not 25 limited to, bimatoprost, latanoprost, travoprost, and so forth.

Compounds of the invention may also be administered in combination with a renin inhibitor, examples of which include, but are not limited to, aliskiren, enalkiren, remikiren, and combinations thereof.

In another embodiment, compounds of the invention are administered in 30 combination with a selective serotonin reuptake inhibitor (SSRI). Representative SSRIs include, but are not limited to: citalopram and the citalopram metabolite desmethylcitalopram, dapoxetine, escitalopram (e.g., escitalopram oxalate), fluoxetine and the fluoxetine desmethyl metabolite norfluoxetine, fluvoxamine (e.g., fluvoxamine

maleate), paroxetine, sertraline and the sertraline metabolite demethylsertraline, and combinations thereof.

In one embodiment, compounds of the invention are administered in combination with a 5-HT_{1D} serotonin receptor agonist, examples of which include, by way of illustration and not limitation, triptans such as almotriptan, avitriptan, eletriptan, frovatriptan, naratriptan rizatriptan, sumatriptan, and zolmitriptan.

In one embodiment, compounds of the invention are administered in combination with a sodium channel blocker, examples of which include, by way of illustration and not limitation, carbamazepine, fosphenytoin, lamotrigine, lidocaine, mexiletine, oxcarbazepine, phenytoin, and combinations thereof.

In one embodiment, compounds of the invention are administered in combination with a soluble guanylate cyclase stimulator or activator, examples of which include, but are not limited to ataciguat, riociguat, and combinations thereof.

In one embodiment, compounds of the invention are administered in combination with a tricyclic antidepressant (TCA), examples of which include, by way of illustration and not limitation, amitriptyline, amitriptylinoxide, butriptyline, clomipramine, demexiptiline, desipramine, dibenzepin, dimetacrine, dosulepin, doxepin, imipramine, imipraminoxide, lofepramine, melitracen, metapramine, nitroxazepine, nortriptyline, noxiptiline, pipofezine, propizepine, protriptyline, quinupramine, and combinations thereof.

In one embodiment, compounds of the invention are administered in combination with a vasopressin receptor antagonist, examples of which include, by way of illustration and not limitation, conivaptan and tolvaptan.

Combined secondary therapeutic agents may also be helpful in further combination therapy with compounds of the invention. For example, compounds of the invention can be combined with a diuretic and an ARB, or a calcium channel blocker and an ARB, or a diuretic and an ACE inhibitor, or a calcium channel blocker and a statin. Specific examples include, a combination of the ACE inhibitor enalapril (in the maleate salt form) and the diuretic hydrochlorothiazide, which is sold under the mark Vaseretic[®], or a combination of the calcium channel blocker amlodipine (in the besylate salt form) and the ARB olmesartan (in the medoxomil prodrug form), or a combination of a calcium channel blocker and a statin, all may also be used with the compounds of the invention. Other therapeutic agents such as α_2 -adrenergic receptor agonists and vasopressin receptor

antagonists may also be helpful in combination therapy. Exemplary α_2 -adrenergic receptor agonists include clonidine, dexmedetomidine, and guanfacine.

The following formulations illustrate representative pharmaceutical compositions of the invention.

5 *Exemplary Hard Gelatin Capsules For Oral Administration*

A compound of the invention (50 g), 440 g spray-dried lactose and 10 g magnesium stearate are thoroughly blended. The resulting composition is then loaded into hard gelatin capsules (500 mg of composition per capsule). Alternately, a compound of the invention (20 mg) is thoroughly blended with starch (89 mg), microcrystalline cellulose (89 mg) and
10 magnesium stearate (2 mg). The mixture is then passed through a No. 45 mesh U.S. sieve and loaded into a hard gelatin capsule (200 mg of composition per capsule).

Alternately, a compound of the invention (30 g), a secondary agent (20 g), 440 g spray-dried lactose and 10 g magnesium stearate are thoroughly blended, and processed as described above.

15 *Exemplary Gelatin Capsule Formulation For Oral Administration*

A compound of the invention (100 mg) is thoroughly blended with polyoxyethylene sorbitan monooleate (50 mg) and starch powder (250 mg). The mixture is then loaded into a gelatin capsule (400 mg of composition per capsule). Alternately, a compound of the invention (70 mg) and a secondary agent (30 mg) are thoroughly blended with
20 polyoxyethylene sorbitan monooleate (50 mg) and starch powder (250 mg), and the resulting mixture loaded into a gelatin capsule (400 mg of composition per capsule).

Alternately, a compound of the invention (40 mg) is thoroughly blended with microcrystalline cellulose (Avicel PH 103; 259.2 mg) and magnesium stearate (0.8 mg). The mixture is then loaded into a gelatin capsule (Size #1, White, Opaque) (300 mg of
25 composition per capsule).

Exemplary Tablet Formulation For Oral Administration

A compound of the invention (10 mg), starch (45 mg) and microcrystalline cellulose (35 mg) are passed through a No. 20 mesh U.S. sieve and mixed thoroughly. The granules so produced are dried at 50-60 °C and passed through a No. 16 mesh U.S. sieve.
30 A solution of polyvinylpyrrolidone (4 mg as a 10 % solution in sterile water) is mixed with sodium carboxymethyl starch (4.5 mg), magnesium stearate (0.5 mg), and talc (1 mg), and this mixture is then passed through a No. 16 mesh U.S. sieve. The sodium carboxymethyl starch, magnesium stearate and talc are then added to the granules. After mixing, the

mixture is compressed on a tablet machine to afford a tablet weighing 100 mg.

Alternately, a compound of the invention (250 mg) is thoroughly blended with microcrystalline cellulose (400 mg), silicon dioxide fumed (10 mg), and stearic acid (5 mg). The mixture is then compressed to form tablets (665 mg of composition per tablet).

5 Alternately, a compound of the invention (400 mg) is thoroughly blended with cornstarch (50 mg), croscarmellose sodium (25 mg), lactose (120 mg), and magnesium stearate (5 mg). The mixture is then compressed to form a single-scored tablet (600 mg of composition per tablet).

Alternately, a compound of the invention (100 mg) is thoroughly blended with
10 cornstarch (100 mg) with an aqueous solution of gelatin (20 mg). The mixture is dried and ground to a fine powder. Microcrystalline cellulose (50 mg) and magnesium stearate (5 mg) are then admixed with the gelatin formulation, granulated and the resulting mixture compressed to form tablets (100 mg of the compound of the invention per tablet).

Exemplary Suspension Formulation For Oral Administration

15 The following ingredients are mixed to form a suspension containing 100 mg of the compound of the invention per 10 mL of suspension:

Ingredients	Amount
Compound of the invention	1.0 g
Fumaric acid	0.5 g
Sodium chloride	2.0 g
Methyl paraben	0.15 g
Propyl paraben	0.05 g
Granulated sugar	25.5 g
Sorbitol (70% solution)	12.85 g
Veegum [®] K (magnesium aluminum silicate)	1.0 g
Flavoring	0.035 mL
Colorings	0.5 mg
Distilled water	q.s. to 100 mL

Exemplary Liquid Formulation For Oral Administration

A suitable liquid formulation is one with a carboxylic acid-based buffer such as
20 citrate, lactate and maleate buffer solutions. For example, a compound of the invention (which may be pre-mixed with DMSO) is blended with a 100 mM ammonium citrate buffer and the pH adjusted to pH 5, or is blended with a 100 mM citric acid solution and the pH adjusted to pH 2. Such solutions may also include a solubilizing excipient such as a cyclodextrin, for example the solution may include 10 wt% hydroxypropyl- β -cyclodextrin.

25 Other suitable formulations include a 5% NaHCO₃ solution, with or without

cyclodextrin.

Exemplary Injectable Formulation For Administration By Injection

A compound of the invention (0.2 g) is blended with 0.4 M sodium acetate buffer solution (2.0 mL). The pH of the resulting solution is adjusted to pH 4 using 0.5 N aqueous hydrochloric acid or 0.5 N aqueous sodium hydroxide, as necessary, and then sufficient water for injection is added to provide a total volume of 20 mL. The mixture is then filtered through a sterile filter (0.22 micron) to provide a sterile solution suitable for administration by injection.

Exemplary Compositions For Administration By Inhalation

10 A compound of the invention (0.2 mg) is micronized and then blended with lactose (25 mg). This blended mixture is then loaded into a gelatin inhalation cartridge. The contents of the cartridge are administered using a dry powder inhaler, for example.

Alternately, a micronized compound of the invention (10 g) is dispersed in a solution prepared by dissolving lecithin (0.2 g) in demineralized water (200 mL). The resulting suspension is spray dried and then micronized to form a micronized composition comprising particles having a mean diameter less than about 1.5 μm . The micronized composition is then loaded into metered-dose inhaler cartridges containing pressurized 1,1,1,2-tetrafluoroethane in an amount sufficient to provide about 10 μg to about 500 μg of the compound of the invention per dose when administered by the inhaler.

20 Alternately, a compound of the invention (25 mg) is dissolved in citrate buffered (pH 5) isotonic saline (125 mL). The mixture is stirred and sonicated until the compound is dissolved. The pH of the solution is checked and adjusted, if necessary, to pH 5 by slowly adding aqueous 1 N NaOH. The solution is administered using a nebulizer device that provides about 10 μg to about 500 μg of the compound of the invention per dose.

25 EXAMPLES

The following Preparations and Examples are provided to illustrate specific embodiments of the invention. These specific embodiments, however, are not intended to limit the scope of the invention in any way unless specifically indicated.

The following abbreviations have the following meanings unless otherwise indicated and any other abbreviations used herein and not defined have their standard, generally accepted meaning:

AcOH	acetic acid
(Boc) ₂ O	di- <i>t</i> -butyl dicarbonate

	DCC	dicyclohexylcarbodiimide
	DCM	dichloromethane or methylene chloride
	DIBAL	diisobutylaluminum hydride
	DIPEA	<i>N,N</i> -diisopropylethylamine
5	DMAP	4-dimethylaminopyridine
	DMF	<i>N,N</i> -dimethylformamide
	DMSO	dimethyl sulfoxide
	Dnp	2,4-dinitrophenyl
	EDCI	<i>N</i> -(3-dimethylaminopropyl)- <i>N'</i> -ethylcarbodiimide
10	Et ₃ N	triethylamine
	Et ₂ O	diethyl ether
	EtOAc	ethyl acetate
	EtOH	ethanol
15	HATU	<i>N,N,N',N'</i> -tetramethyl- <i>O</i> -(7-azabenzotriazol-1-yl)uronium hexafluorophosphate
	HCTU	(2-(6-chloro-1H-benzotriazole-1-yl)-1,1,3,3-tetramethylaminium hexafluorophosphate)
	HEPES	4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid
	HOAt	1-hydroxy-7-azabenzotriazole
20	LiHMDS	lithium hexamethyl disilazide
	Mca	(7-methoxycoumarin-4-yl)acyl
	MeCN	acetonitrile
	MeOH	methanol
	Pd(PPh ₃) ₄	tetrakis(triphenylphosphine)palladium(0)
25	SilicaCat [®] DPP-Pd	silica based diphenylphosphine palladium (II) catalyst
	SilicaCat [®] Pd(0)	silica based palladium (0) catalyst
	TFA	trifluoroacetic acid
	THF	tetrahydrofuran

Unless noted otherwise, all materials, such as reagents, starting materials and
 30 solvents, were purchased from commercial suppliers (such as Sigma-Aldrich, Fluka
 Riedel-de Haën, and the like) and were used without further purification.

Reactions were run under nitrogen atmosphere, unless noted otherwise. The
 progress of reactions were monitored by thin layer chromatography (TLC), analytical high

performance liquid chromatography (anal. HPLC), and mass spectrometry, the details of which are given in specific examples. Solvents used in analytical HPLC were as follows: solvent A was 98% H₂O/2% MeCN /1.0 mL/L TFA; solvent B was 90% MeCN/10% H₂O/1.0 mL/L TFA.

Reactions were worked up as described specifically in each preparation for example; commonly reaction mixtures were purified by extraction and other purification methods such as temperature-, and solvent-dependent crystallization, and precipitation. In addition, reaction mixtures were routinely purified by preparative HPLC, typically using Microsorb C18 and Microsorb BDS column packings and conventional eluents. Progress of reactions was typically measured by liquid chromatography mass spectrometry (LCMS). Characterization of isomers were done by Nuclear Overhauser effect spectroscopy (NOE). Characterization of reaction products was routinely carried out by mass and ¹H-NMR spectrometry. For NMR measurement, samples were dissolved in deuterated solvent (CD₃OD, CDCl₃, or DMSO-*d*₆), and ¹H-NMR spectra were acquired with a Varian Gemini 2000 instrument (400 MHz) under standard observation conditions. Mass spectrometric identification of compounds was typically conducted using an electrospray ionization method (ESMS) with an Applied Biosystems (Foster City, CA) model API 150 EX instrument or an Agilent (Palo Alto, CA) model 1200 LC/MSD instrument.

Preparation 1

Oxodiperoxymolybdenum(pyridine)(hexamethylphosphorictriamide)




Molybdenum oxide (MoO₃; 30 g, 0.2 mol) and 30% hydrogen peroxide (150 mL) were combined, with stirring. The reaction vessel was placed in an oil bath equilibrated at 40°C and heated until the internal temperature reached 35°C. The heating bath was then removed and replaced by a water bath to control the mildly exothermic reaction so that an internal temperature of 35-40°C was maintained. After the initial exothermic period (~ 30 minutes), the reaction vessel was returned to the 40°C oil bath and stirred for a total of 3.5 hours to form a yellow solution with a small amount of suspended white solid. After cooling to 20°C, the solution was filtered and the resulting yellow filtrate was cooled to 10°C (ice bath with stirring) and hexamethylphosphoric triamide ((Me₂N)₃PO; HMPA; 37.3 g, 0.2 mol) was added dropwise over 5 minutes, resulting in the formation of a yellow crystalline precipitate. Stirring was continued for a total of 15 minutes at 10°C, and the

product was filtered and pressed dry. After 30 minutes under vacuum, the filter cake was combined with MeOH (20 mL) and stirred at 40°C. Additional MeOH was slowly added until the solids dissolved. The saturated solution was cooled in the refrigerator, yielding a yellow solid (appeared as needles). The solid mass was physically broken, filtered and washed with cold MeOH (20-30 mL) to yield oxodiperoxymolybdenum(aqua) (hexamethylphosphoric triamide) ($\text{MoO}_5 \cdot \text{H}_2\text{O} \cdot \text{HMPA}$, 46-50 g).

$\text{MoO}_5 \cdot \text{H}_2\text{O} \cdot \text{HMPA}$ was dried over phosphorus oxide in a vacuum desiccator, shielded from the light, for 24 hours at 0.2 mm Hg to yield a somewhat hygroscopic yellow solid, $\text{MoO}_5 \cdot \text{HMPA}$. $\text{MoO}_5 \cdot \text{HMPA}$ (36.0 g, 0.1 mol) was dissolved in THF (150 mL) and the solution was filtered to remove any precipitate. The filtrate was then stirred at 20°C while dry pyridine (8.0 g, 0.1 mol) was added over 10 minutes. The crystalline, yellow product was collected, washed with dry THF (25 mL) and anhydrous ether (200 mL) and dried in a vacuum desiccator (1 hour, 0.2 mm Hg) to yield the title compound, oxodiperoxymolybdenum(pyridine)(hexamethylphosphoric triamide) ($\text{MoO}_5 \cdot \text{Py} \cdot \text{HMPA}$) as a finely divided yellow solid (36-38 g).


Preparation 2

(S)-2-Biphenyl-4-ylmethyl-5-oxopyrrolidine-1-carboxylic Acid *t*-Butyl Ester




To a stirred solution of (*R*)-3-biphenyl-4-yl-2-*t*-butoxycarbonylamino-2-oxopropanoic acid (50 g, 146.5 mmol), Meldrum's acid (23.3 g, 161.1 mmol) and DMAP (27.8 g, 227 mmol) in anhydrous DCM (500 mL) was added a solution of DCC (33.3 g, 161.1 mmol) in anhydrous DCM (200 mL) over 1 hour at -5°C under nitrogen. The mixture was stirred at -5°C for 8 hours, then refrigerated overnight, during which tiny crystals of dicyclohexylurea precipitated. After filtration the mixture was washed with 5% KHSO_4 (4 x 200 mL), saturated aqueous NaCl (1 x 200 mL) and dried over MgSO_4 overnight. The resulting solution was evaporated to give the crude compound 1 (68 g) as a light yellow

solid). LC-MS: $[M^+Na]$:490, $[2M^+Na]$:957.




To a solution of crude compound 1 (68 g, 146.5 mmol) in anhydrous DCM (1000 mL) was added AcOH (96.8 g, 1.6 mol) at -5°C under nitrogen. The resulting mixture was stirred at -5°C for 0.5 hours, then NaBH_4 (13.9 g, 366 mmol) was added in small portions over 2 hours. After stirring for another hour at -5°C , saturated aqueous NaCl (300 mL) was added. The organic layer was washed with saturated aqueous NaCl (2 x 300 mL) then water (2 x 300 mL), dried over MgSO_4 , filtered, and evaporated to give the crude compound 2, which was further purified by chromatography (hexanes:EtOAc=5:1) to give purified compound 2 (46 g) as a light yellow solid. LC-MS: $[M^+Na]$:476, $[2M^+Na]$:929.




A stirred solution of purified compound 2 (46 g, 101 mmol) in anhydrous toluene (300 mL) was heated to reflux under nitrogen for 3 hours. After evaporation of the solvent, the residue was purified by chromatography (hexanes:EtOAc=10:1) to yield the title compound (27 g) as a light yellow solid.

LC-MS: $[M^+Na]$:374, $[2M^+Na]$:725 ; ^1H NMR (300 MHz, CDCl_3): δ 7.64-7.62 (m, 4H), 7.51-7.46 (m, 2H), 7.42-7.39 (m, 1H), 7.39-7.30 (m, 2H), 4.50-4.43 (m, 1H), 3.27-3.89 (m, 1H), 2.88-2.80 (m, 1H), 2.48-2.42 (m, 2H), 2.09-1.88 (m, 2H), 1.66 (s, 9H).

Preparation 3

(2R,4R)-4-Amino-5-biphenyl-4-yl-2-hydroxypentanoic Acid Ethyl Ester and (2R,4S)-4-Amino-5-biphenyl-4-yl-2-hydroxypentanoic Acid Ethyl Ester

- 5 Step 1: To a stirred solution of (*S*)-2-biphenyl-4-ylmethyl-5-oxopyrrolidine-1-carboxylic acid *t*-butyl ester (4.4 g, 12.4 mmol) in anhydrous THF (70 mL) was added a solution of 1 M LiHMDS in THF (28 mL) over 15 minutes at -65°C under nitrogen. After stirring for 3 hours at -65°C, oxodiperoxymolybdenum(pyridine)(hexamethylphosphorictriamide) (9 g, 18.6 mmol) was added. The mixture was stirred for another 2 hours
- 10 at -35°C, then saturated aqueous Na₂S₂O₃ (60 mL) was added. The organic layer was collected and washed with saturated aqueous NH₄Cl (60 mL x 3) and saturated aqueous NaCl (60 mL x 2), then dried over Na₂SO₄, and the solvent was removed under reduced pressure to yield the crude product which was further purified by chromatography (hexanes:EtOAc = 5:1) to yield compound 1 as a white solid (1.8 g). LC-MS:
- 15 [2M+Na]:757.




- Step 2: To a solution of compound 1 (1.8 g, 5.0 mmol) in anhydrous DCM (50 mL) was added DMAP (122 mg, 1 mmol) and Et₃N (1.5 g, 14.9 mmol) at 0°C under nitrogen. After stirring for 0.5 hour at 0°C, benzyl chloride (1.0 g, 7.4 mmol) was added over 15
- 20 minutes. The mixture was stirred for an additional 2 hours at 0°C, then saturated aqueous NaHCO₃ (50 mL) was added. The organic layer was collected and washed with saturated

aqueous NaHCO_3 (50 mL x 2) and saturated aqueous NaCl (50 mL x 1), then dried over Na_2SO_4 . The solids were filtered out and the filtrate was concentrated to yield the crude product which was further purified by chromatography (hexanes:EtOAc = 4:1) to yield compound 2A (471 mg) and compound 2B (883 mg) as white solids. LC-MS:

5 [M+Na]:494; [2M+Na]:965.

Compound 2A: ^1H NMR (300 MHz, CDCl_3): δ (ppm) = 8.02 (m, 2H), 7.57-7.25 (m, 12H), 5.42 (m, 1H), 4.50 (m, 1H), 3.26-3.21 (m, 1H), 2.90 (m, 1H), 2.58 (m, 1H), 2.15-2.05 (m, 1H), 1.62 (m, 9H)


Compound 2B: ^1H NMR (300 MHz, CDCl_3): δ (ppm) = 8.06 (m, 2H), 7.58-7.18 (m, 12H), 5.53-5.41 (m, 1H), 4.39 (m, 1H), 3.57-3.54 (m, 1H), 2.87-2.80 (m, 1H), 2.48-2.44 (m, 1H), 1.98 (m, 1H), 1.63 (m, 9H).



Step 3: To a stirred solution of compound 2A (471 mg, 1 mmol) in anhydrous EtOH (10 mL) was added anhydrous K_2CO_3 (691 mg, 5 mmol) at room temperature under nitrogen. After stirring for 20 hours at room temperature, the solids were filtered out. To the filtrate was added water (30 mL), DCM (30 mL) and saturated aqueous NaCl (5 mL). The aqueous layer was separated and extracted with DCM (30 mL x 3). The combined organic layers were washed with saturated aqueous NaCl (50 mL), dried over Na_2SO_4 , and concentrated to yield the crude product which was further purified by chromatography (hexanes:EtOAc = 6:1) to yield compound 3 as a white solid (275 mg). LC-MS:

15
20


[M+Na]:436, [2M+Na]:849.




Step 4: To EtOH (5 mL) was added acetyl chloride (685 mg) at -30°C . After stirring for 1 hour at -30°C , a solution of compound 3 (275 mg, 665 μmol) in anhydrous

EtOH (5 mL) was added. The mixture was heated to 25°C and stirred for 3 hours at 25°C. After evaporation of the solvent, the residue was washed with cold anhydrous Et₂O (10 mL) to yield (2*R*,4*R*)-4-amino-5-biphenyl-4-yl-2-hydroxypentanoic acid ethyl ester as a white solid HCl salt (207 mg). LC-MS: [M+H]:314, [2M+Na]:649.

- 5 ¹H NMR (300 MHz, CDCl₃): δ (ppm) = 7.99 (m, 3H), 7.66-7.64 (m, 4H), 7.48-7.35 (m, 5H), 6.08 (m, 1H), 4.21 (m, 1H), 4.09-4.05 (m, 2H), 3.52 (m, 1H), 2.97-2.95 (m, 2H), 1.89-1.87 (m, 2H), 1.19-1.14 (m, 3H).



- Step 5: To a stirred solution of compound 2B (883 mg, 1.9 mmol) in anhydrous EtOH (15 mL) was added anhydrous K₂CO₃ (1293 mg, 9.4 mmol) at room temperature under nitrogen. After stirring for 20 hours at room temperature, the solids were filtered out. To the filtrate was added water (30 mL), DCM (30 mL) and saturated aqueous NaCl (5 mL). The aqueous layer was separated and extracted with DCM (30 mL x 3). The combined organic layers were washed with saturated aqueous NaCl (50 mL), dried over Na₂SO₄, and concentrated to yield the crude product which was further purified by chromatography (hexanes:EtOAc = 6:1) to yield compound 4 as a white solid (524 mg). LC-MS: [M+Na]:436, [2M+Na]:849.
- 10
15




- Step 6: To EtOH (8 mL) was added acetyl chloride (1300 mg) at -30°C. After stirring for 1 hour at -30°C, a solution of compound 4 (524 mg, 1.3 mmol) in anhydrous EtOH (8 mL) was added. The mixture was heated to 25°C and stirred for 3 hours at 25°C. After evaporation of the solvent, the residue was washed with cold anhydrous Et₂O (10 mL) to yield (2*R*,4*S*)-4-amino-5-biphenyl-4-yl-2-hydroxypentanoic acid ethyl ester as a white solid HCl salt (395 mg). LC-MS: [M+H]:314, [2M+Na]:649.
- 20


^1H NMR (300 MHz, CDCl_3): δ (ppm) = 8.14 (m, 3H), 7.66-7.62 (m, 4H), 7.47-7.31 (m, 5H), 5.87-5.85 (m, 1H), 4.34 (m, 1H), 4.08-4.00 (m, 2H), 3.48 (m, 1H), 3.09 (m, 1H), 2.85-2.81 (m, 1H), 1.88 (m, 1H), 1.76 (m, 1H), 1.15-1.10 (m, 3H).

Alternate Synthesis of

5 *(2R,4R)-4-Amino-5-biphenyl-4-yl-2-hydroxypentanoic Acid Ethyl Ester*



A solution of [(*S*)-2-Biphenyl-4-yl-1-(2,2-dimethyl-4,6-dioxo-[1,3]dioxan-5-ylmethyl)ethyl]-carbamic acid *t*-butyl ester (143 g, 320 mmol) in anhydrous toluene (1 L) was heated to reflux under nitrogen overnight. The solvent was removed under reduced
 10 pressure to give (*S*)-2-biphenyl-4-ylmethyl-5-oxopyrrolidine-1-carboxylic acid *t*-butyl ester, which was directly used without further purification and added to a solution of 3N HCl in EtOAc (1.2 L). The resulting mixture was stirred for 3 hours at room temperature. The solvent was removed in *vacuo*. The residue was re-crystallized in EtOAc (300 mL) to yield a first batch of the deprotected compound (56 g). The mother liquid was subject to
 15 chromatography column (eluted by EtOAc and hexanes from 1:1 to 100% EtOAc) to yield a second batch of the deprotected compound (8 g). To a suspension of the deprotected compound (64 g, 250 mmol) in anhydrous THF (500 mL) was added dropwise BuLi (100 mL, 2.5 M in hexanes) at -78°C . After stirring for 0.5 hour, pivaloyl chloride (34 g, 0.28 mol) was added dropwise. The mixture was stirred for 1 hour at -78°C . The reaction was
 20 then quenched with saturated aqueous NH_4Cl , and the mixture extracted with EtOAc. The extract was dried over MgSO_4 and concentrated to yield (*S*)-5-biphenyl-4-ylmethyl-1-(2,2-dimethylpropionyl)pyrrolidin-2-one (85 g) as white solid.




(*S*)-5-Biphenyl-4-ylmethyl-1-(2,2-dimethylpropionyl)pyrrolidin-2-one (40 g, 120 mmol) was dissolved in anhydrous THF (400 mL) and stirred at -78°C under nitrogen. To this was added 1.5 eq of a 2.0M THF solution of sodium bis(trimethylsilyl)amide dropwise over 5 minutes. The light yellow mixture was stirred for 20 minutes under nitrogen at -78°C, followed by the slow dropwise addition of oxaziridine (53 g, 180 mmol) as a 200 mL solution in THF. The mixture was stirred for 0.5 hour. The reaction was quenched with saturated aqueous NH₄Cl, and the mixture was extracted with EtOAc (1 L). The extract was washed with 1N HCl (1 L), dried over MgSO₄, and concentrated to a 500 mL volume. The precipitated white solid was filtered, the filtrate was concentrated to remove solvent after the addition of silica gel (200 g). The residue was put on a column (8×80 cm) of silica gel (900 g), which was previously packed in hexanes. Elution was carried out initially with DCM:hexanes (1:1). Once the oxaziridine and imine were completely collected, the column was eluted with DCM to obtain (*3R,5R*)-5-Biphenyl-4-ylmethyl-1-(2,2-dimethylpropionyl)-3-hydroxypyrrolidin-2-one (21 g, 98% purity) as yellow oil.

(*3R,5R*)-5-Biphenyl-4-ylmethyl-1-(2,2-dimethylpropionyl)-3-hydroxypyrrolidin-2-one (56 g, 0.156 mol) was dissolved in EtOH (700 mL) and 12N HCl (700 mL). The mixture was heated to 90~95°C for 20 hours. The mixture was concentrated on an 80°C water bath under reduced pressure. EtOH (100 mL) was added to the residue, and the resulting mixture was filtered to yield a yellow solid. This solid was suspended in 3N HCl/EtOH (800 mL). The mixture was refluxed for 3 hours. The solution was concentrated to a reduced volume (~200 mL volume), and ether (200 mL) was added. The resulting slightly yellow solid was filtered and dried under reduced pressure to yield (*2R,4R*)-4-amino-5-biphenyl-4-yl-2-hydroxypentanoic acid ethyl ester (43 g).

Preparation 4

(*2R,4R*)-4-Amino-5-biphenyl-4-yl-2-hydroxypentanoic Acid Butyl Ester




1-Butanol (40 mL, 400 mmol) was cooled in an ice bath and acetyl chloride was added dropwise until a concentration of 4N HCl was achieved. The mixture was warmed

to room temperature. (2*R*,4*R*)-4-amino-5-biphenyl-4-yl-2-hydroxypentanoic acid ethyl ester (HCl salt; 250 mg, 710 μ mol) was added to the acidified butanol (15 mL) and the mixture was capped and heated to 80°C for 3 hours. The resulting product was dried under vacuum to yield the title compound as a white/orange flaky solid HCl salt (220 mg).

5

EXAMPLE 1



A. (2*R*,4*R*)-4-[(5-Acetyl-2H-pyrazole-3-carbonyl)amino]-5-biphenyl-4-yl-2-hydroxypentanoic Acid Ethyl Ester ($R^7 = -CH_2CH_3$)

(2*R*,4*R*)-4-Amino-5-biphenyl-4-yl-2-hydroxypentanoic acid ethyl ester (HCl salt; 500 mg, 1 mmol, 1.0 eq.), 5-acetyl-2H-pyrazole-3-carboxylic acid (330.4 mg, 2.1 mmol, 1.5 eq.), and HATU (820 mg, 2.1 mmol, 1.5 eq.) were combined in DMF (5 mL), and the resulting mixture was stirred for 2 minutes. DIPEA (750 μ L) was added and the mixture was stirred for 1 hour. The mixture was dried under vacuum, and the product was purified using reverse phase chromatography (10-70% MeCN/H₂O; 0.05% TFA over 70 minutes) to yield the title compound (300 mg, purity 98%) as a TFA salt. MS m/z [$M+H$]⁺ calc'd for C₂₅H₂₇N₃O₅, 450.20; found 450.2.

B. (2*R*,4*R*)-4-[(5-Acetyl-2H-pyrazole-3-carbonyl)amino]-5-biphenyl-4-yl-2-hydroxypentanoic Acid ($R^7 = H$)

A portion (200 mg) of the purified ethyl ester (compound A) was dissolved in EtOH (~10 mL), and 10N NaOH was added to make the solution basic (~200 μ L). The mixture was monitored over 1 hour until final deprotection was complete. The mixture was acidified with an equal volume (~200 μ L) of AcOH and dried under vacuum. The product was purified using reverse phase chromatography to yield the title compound as a TFA salt (160 mg; purity 100%). MS m/z [$M+H$]⁺ calc'd for C₂₃H₂₃N₃O₅, 422.16; found 422.2.

C. (2R,4R)-4-[(5-Acetyl-2H-pyrazole-3-carbonyl)amino]-5-biphenyl-4-yl-2-hydroxypentanoic Acid Butyl Ester ($R^7 = -(CH_2)_3CH_3$)

(2R,4R)-4-Amino-5-biphenyl-4-yl-2-hydroxypentanoic acid butyl ester (HCl salt; 500 mg, 1 mmol, 1.0 eq.), 3-acetyl-1H-pyrazole-5-carboxylic acid (330.4 mg, 2.1 mmol, 1.5 eq.), and HATU (820 mg, 2.1 mmol, 1.5 eq.) were combined in DMF (5 mL), and the resulting mixture was stirred for 2 minutes. DIPEA (750 μ L) was added and the mixture was stirred for 1 hour. The mixture was dried under vacuum, and the product was purified using reverse phase chromatography to yield the title compound as a TFA salt (45 mg; purity 95%). MS m/z $[M+H]^+$ calc'd for $C_{27}H_{31}N_3O_5$, 478.23; found 478.4.

10 D. (2R,4R)-4-[(5-Acetyl-2H-pyrazole-3-carbonyl)amino]-5-biphenyl-4-yl-2-hydroxypentanoic Acid 2-methoxyethyl Ester ($R^7 = -(CH_2)_2OCH_3$)

A saturated HCl solution in 2-methoxyethanol was prepared by adding acetyl chloride (~400 μ L) dropwise to 10 mL of the anhydrous alcohol. To this solution was added (2R,4R)-4-[(5-acetyl-2H-pyrazole-3-carbonyl)amino]-5-biphenyl-4-yl-2-hydroxypentanoic acid ethyl ester (21.3 mg, 47.4 μ mol) and the resulting mixture was heated to 60°C for 1-2 hours. The material was dried under vacuum. The product was then purified using reverse phase chromatography to yield the title compound as a TFA salt (10 mg; purity 100%) MS m/z $[M+H]^+$ calc'd for $C_{26}H_{29}N_3O_6$, 480.21; found 480.4.

20 E. (2R,4R)-4-[(5-Acetyl-2H-pyrazole-3-carbonyl)amino]-5-biphenyl-4-yl-2-hydroxypentanoic Acid 5-methyl-2-oxo-[1,3]dioxol-4-ylmethyl Ester ($R^7 = -CH_2-5-methyl-[1,3]dioxol-2-one$)

(2R,4R)-4-[(5-Acetyl-2H-pyrazole-3-carbonyl)amino]-5-biphenyl-4-yl-2-hydroxypentanoic acid (20 mg, 50 μ mol) was dissolved in dry DMF (2 mL). Dicesium carbonate (50 mg, 0.2 mmol) was added, and the resulting mixture was stirred at 0°C for 1 hour. 4-Chloromethyl-5-methyl-1,3-dioxol-2-one (20 mg, 0.1 mmol) was added and the resulting mixture was stirred at room temperature overnight. EtOAc (20 mL) was added and the mixture was washed with saturated aqueous NaCl (100 mL) then dried under vacuum. The material was purified using reverse phase chromatography to yield the title compound as a TFA salt (5 mg, purity 100%). MS m/z $[M+H]^+$ calc'd for $C_{28}H_{27}N_3O_8$, 534.18; found 534.4.

F. (2R,4R)-4-[(5-Acetyl-2H-pyrazole-3-carbonyl)amino]-5-biphenyl-4-yl-2-hydroxypentanoic Acid 2-Morpholin-4-yl-ethyl Ester ($R^7 = -(CH_2)_2-morpholine$)
(2R,4R)-4-[(5-Acetyl-2H-pyrazole-3-carbonyl)amino]-5-biphenyl-4-yl-2-


hydroxypentanoic acid (50 mg, 120 μ mol, 1.1 eq.) and HATU (42 mg, 110 μ mol, 1.0 eq.) were combined in DMF (5 mL) and the resulting mixture was stirred for 2 minutes. 4-Morpholineethanol (73 mg, 560 μ mol, 5.0 eq.) was added, followed by the addition of DIPEA. The resulting mixture was stirred for 1 hour, then the material was dried under vacuum. The product was then purified using reverse phase chromatography to yield the title compound as a TFA salt (approximately 4 mg, purity 95%). MS m/z $[M+H]^+$ calc'd for $C_{29}H_{34}N_4O_6$, 535.25; found 535.4.

G. (2R,4R)-4-[(5-Acetyl-2H-pyrazole-3-carbonyl)amino]-5-biphenyl-4-yl-2-hydroxypentanoic acid 2-Methanesulfonylethyl Ester ($R^7 = -(CH_2)_2-SO_2-CH_3$)

2-(Methylsulfonyl)ethanol (690 mg, 5.6 mmol) was warmed to 40°C and 0.5eq of acetyl chloride was added with stirring for 5 minutes. (2R,4R)-4-[(5-Acetyl-2H-pyrazole-3-carbonyl)amino]-5-biphenyl-4-yl-2-hydroxypentanoic acid ethyl ester (50 mg, 110 μ mol) was added and the reaction vessel was capped and heated to 70°C for 2 hours. The product was then purified using reverse phase chromatography to yield the title compound as a TFA salt (approximately 28 mg, purity 95%). MS m/z $[M+H]^+$ calc'd for $C_{26}H_{29}N_3O_7S$, 528.17; found 528.4.


Preparation 5

Acetoxy(diethoxyphosphoryl)acetic Acid Ethyl Ester




Ethyl 2-oxoacetate(50%) (74 g, 724.8 mmol, 1.0 eq.) was added dropwise with stirring at 0°C to a solution of diethyl hydrogen phosphite (50 g, 362.1 mmol, 1.0 eq.) in toluene (100 mL), under nitrogen. Et_3N (110 g, 1.1 mol, 3.0 eq.) was added dropwise with stirring at 0°C. The resulting solution was stirred for 1 hour at room temperature. To the mixture was added acetic anhydride (37 g, 362.4 mmol, 1.0 eq.) dropwise with stirring at 0°C. The resulting solution was stirred overnight at room temperature. The pH value of the solution was adjusted to 6 with 2N HCl. The resulting solution was extracted with DCM (3x150 mL) and the organic layers were combined, dried over Na_2SO_4 , and concentrated under vacuum. The residue was loaded onto a silica gel column with EtOAc:hexanes (1:2~1:5) to yield the title compound (52 g) as a light yellow liquid.


Preparation 6

[(*R*)-2-Biphenyl-4-yl-1-(methoxymethylcarbamoyl)ethyl]carbamic Acid *t*-Butyl Ester

Step 1: *N*-*t*-Boc-D-tyrosine (80.0 g, 284.4 mmol, 1.0 eq.), *O*, *N*-dimethyl-
 5 hydroxylamine hydrochloride (33.3 g, 341.4 mmol, 1.2 eq.), 1-ethyl-3-((3-dimethylamino)-
 propyl)carbodiimide hydrochloride (81.8 g, 426.7 mmol, 1.5 eq.), hydroxybenzotriazole
 (57.6 g, 426.4 mmol, 1.5 eq.), and DCM (dried) (800 mL) were combined with stirring
 under nitrogen. The resulting mixture was cooled to 0°C and Et₃N (44.0 g, 434.8 mmol,
 1.53 eq.) was added dropwise with stirring over 40 minutes. The resulting solution was
 10 stirred overnight at room temperature, then washed with saturated aqueous NaHCO₃
 solution (2x300 mL). The organic layer was separated and the aqueous layer was re-
 extracted with DCM (200 mL). The organic phase was combined, dried over Na₂SO₄, and
 concentrated under vacuum to yield crude compound 1 (100 g) as a yellow solid.




Step 2: Compound 1 (100.0 g, crude, 308.3 mmol, 1.0 eq.), pyridine (61.0 g, 771.2
 mmol, 2.5 eq.), and DCM (dried) (1000 mL) were combined with stirring under nitrogen.
 The resulting mixture was cooled to -20°C and then trifluoromethanesulfonic anhydride
 (104.0 g, 368.6 mmol, 1.2 eq.) was added dropwise with stirring at -20°C over 30 minutes.
 The resulting solution was then stirred for 30 minutes at -20~-15°C. The reaction was then
 20 quenched by the addition of water (250 mL). The resulting mixture was washed with a
 0.5N NaOH solution (2x300 mL) and a 15% citric acid solution (2x300 mL). The basic
 aqueous layer was re-extracted with DCM (200 mL), and the organic phase was combined,
 dried over Na₂SO₄, and concentrated under vacuum. The residue was loaded onto a silica
 gel column with EtOAc:hexanes (0~1:5) to yield compound 2 (97.7 g) as pale-yellow oil.



Step 3: Phenylboronic acid (53.0 g, 434.7 mmol, 2.03 eq.), potassium carbonate (45.0 g, 325.6 mmol, 1.52 eq.), tetrakis(triphenylphosphine)palladium(0) (12.5 g, 10.8 mmol, 0.05 eq.), and toluene (distilled) (800 mL) were combined with stirring under nitrogen. The resulting suspension was degassed and heated to 80°C before adding a 200 mL toluene solution of compound 2 (97.7g, 214.1 mmol, 1.0 eq.). The thick suspension was stirred overnight at 80°C. The reaction progress was monitored by LCMS, which indicated an incomplete reaction. After the crude product/starting material mixture was recovered by chromatography, additional phenylboronic acid (21.2 g), potassium carbonate (18.0g), tetrakis(triphenylphosphine)palladium(0) (5.0 g) was added and the resulting mixture suspended in toluene (distilled) (700 mL). The resulting mixture was heated to 80°C and stirred overnight. Upon completion, the reaction was quenched by the addition of water (300 mL). The mixture was extracted with EtOAc (2x300 mL) and the organic layers were combined. The organic phase was washed with saturated aqueous NaCl (1x300 mL), dried over Na₂SO₄, and concentrated under vacuum. The residue was loaded onto a silica gel column with EtOAc:hexanes (0~1:3) to yield the title compound (50.0 g) as a white solid.

Preparation 7

[(R)-2-Biphenyl-4-yl-1-formylethyl]carbamic Acid *t*-Butyl Ester




20


[(R)-2-Biphenyl-4-yl-1-(methoxymethylcarbamoyl)ethyl]carbamic acid *t*-butyl ester (7.5 g, 19.5 mmol, 1.0 eq.) was combined with THF (distilled) (100 mL) under nitrogen. Lithium aluminum hydride (750 mg, 19.8 mmol, 1.0 eq.) was added to the resulting stirred solution in several batches at -5~0°C over a period of 30 minutes. The

resulting solution was stirred for 30 minutes at -5-0°C. The reaction was then quenched by the addition of a solution of KHSO₄ (6.6 g) in water (35 mL). The resulting solution was combined with a 1N HCl (75 mL) solution, and stirred for five minutes. EtOAc (100 mL) was added and the resulting solution was extracted with EtOAc (3x100 mL) and the organic layers were combined to yield the title compound (6.3 g) as a yellow oil.

Preparation 8


(R)-4-Amino-5-biphenyl-4-yl-2-hydroxypentanoic Acid Ethyl Ester

Step 1: A solution of acetoxy(diethoxyphosphoryl)acetic acid ethyl ester (15.6 g, 55.3 mmol, 1.2 equiv) in THF (dried) (150 mL), under nitrogen, was cooled to -78°C. LiHMDS (1M in THF) (55.3 mL, 1.2 eq.) was added dropwise with stirring at -78°C. After stirring for 30 minutes at that temperature, a solution of ((R)-2-biphenyl-4-yl-1-formylethyl)carbamic acid *t*-butyl ester (15.0 g, crude, 1.0 eq.) in THF (dried) (30 mL) was added dropwise over 15 minutes. Stirring was continued for 1.5 hours at -78°C before the mixture was poured into a cold solution with water (200 mL) and EtOAc (200 mL). The organic layer was repeatedly separated and the aqueous layer was reextracted with EtOAc (2x100 mL). The combined organic layers were dried over Na₂SO₄, filtered, and evaporated, and the residue was purified by flash chromatography (EtOAc/hexanes=0~1:10) to give compound 1 (10.5 g) as a white solid.




Step 2: A stirred solution of compound 1 (10.5 g, 23.2 mmol, 1.0 eq.) in EtOH (anhydrous) (100 mL) was combined with palladium carbon (1.0 g), under nitrogen. The mixture was purged four times with hydrogen and then hydrogen was bubbled over 2 hours at room temperature. The palladium carbon was filtered out, and the filtrate was

concentrated under vacuum to yield crude compound 2 (10.0 g) as a pale-yellow oil, which was used without further purification.




Step 3: Compound 2 (10.0 g, 22.0 mmol, 1.0 eq.) in EtOH (anhydrous) (100 mL) was combined with potassium carbonate (6.1 g, 44.1 mmol, 2.0 eq.) and the resulting solution was stirred for 2 hours at room temperature. The solids were filtered out and the filtrate was concentrated under vacuum. The residue was loaded onto a silica gel column (EtOAc/hexanes=0~1:5) to yield compound 3 (6.0 g) as a white solid.



Step 4: Compound 3 (6.0 g, 14.5 mmol, 1.0 eq.) was dissolved in DCM (dried) (120 mL), and HCl was bubbled into the mixture over 5~6 hours at room temperature. Solid precipitate was observed. The mixture was concentrated to half volume then filtered. The solids were collected and washed with cold EtOAc, and dried over reduced pressure to yield the title compound (4.2 g) as an off-white solid HCl salt. LC-MS (ES, m/z): 314 [M-HCl+H]⁺.

¹H NMR (300 MHz, DMSO): δ (ppm) = 8.07 (s, 1.9H), 7.96 (s, 1.2H), 7.65-7.69 (m, 4.0H), 7.45-7.50 (m, 2.0H), 7.33-7.39 (m, 3.0H), 6.05-6.07 (m, 0.63H), 5.88-5.90 (m, 0.88H), 4.32-4.38 (m, 0.80H), 4.18-4.31 (m, 0.51H), 4.05-4.11 (m, 2H), 3.50 (s, 1H), 2.75-3.05 (m, 2.8H), 1.83-1.94 (m, 1H), 1.71-1.82 (m, 1H), 1.10-1.20 (m, 3.3H).

EXAMPLE 2



- A. 5-((1*R*,3*S*)-1-Biphenyl-4-ylmethyl-3-ethoxycarbonyl-3-hydroxypropylcarbamoyl)-2H-pyrazole-3-carboxylic Acid ($R^7 = -CH_2CH_3$)
- 5 B. 5-((1*R*,3*R*)-1-Biphenyl-4-ylmethyl-3-ethoxycarbonyl-3-hydroxypropylcarbamoyl)-2H-pyrazole-3-carboxylic Acid ($R^7 = -CH_2CH_3$)
- C. 5-((1*R*,3*S*)-1-Biphenyl-4-ylmethyl-3-carboxy-3-hydroxypropylcarbamoyl)-2H-pyrazole-3-carboxylic Acid ($R^7 = H$)
- 10 D. 5-((1*R*,3*R*)-1-Biphenyl-4-ylmethyl-3-carboxy-3-hydroxypropylcarbamoyl)-2H-pyrazole-3-carboxylic Acid (D; $R^7 = H$)

(*R*)-4-Amino-5-biphenyl-4-yl-2-hydroxypentanoic acid ethyl ester (500 mg, 1.6 mmol, 1.0 eq.), 3,5-pyrazoledicarboxylic acid (250 mg, 1.6 mmol, 1.6 eq.), and HATU (610 mg, 1.6 mmol, 1.0 eq.) were combined in DMF (5 mL), and the resulting mixture was stirred for 2 minutes. DIPEA (1 mL) was added and the mixture was stirred for 1 hour.

- 15 The mixture was dried under vacuum, and the product was purified using reverse phase chromatography (10-70% MeCN/H₂O; 0.05% TFA over 70 minutes) to yield 5-((*R*)-1-biphenyl-4-ylmethyl-3-ethoxycarbonyl-3-hydroxypropylcarbamoyl)-2H-pyrazole-3-carboxylic acid as a TFA salt.


- 20 This material was separated by preparative HPLC, purified using reverse phase chromatography and dried under vacuum to yield compounds A and B as TFA salts. Compounds A and B were each dissolved in EtOH (~100 mL), and 10N NaOH was added to make the solutions basic (~200 μ L). Each mixture was monitored over 1 hour until final deprotection was complete. Each mixture was acidified with an equal volume (~200 μ L) of AcOH and dried under vacuum. The products were purified using reverse phase
- 25 chromatography to yield compound C (52 mg; purity 100%; MS m/z [M+H]⁺ calc'd for C₂₂H₂₁N₃O₆, 424.14; found 424.2) and compound D (65 mg; purity 100%; MS m/z [M+H]⁺ calc'd for C₂₂H₂₁N₃O₆, 424.14; found 424.2) as TFA salts.

E. 5-((1*R*,3*R*)-1-Biphenyl-4-ylmethyl-3-butoxycarbonyl-3-hydroxy-propylcarbamoyl)-1H-pyrazole-3-carboxylic Acid ($R^7 = -(CH_2)_3CH_3$)

(2*R*,4*R*)-4-Amino-5-biphenyl-4-yl-2-hydroxypentanoic acid butyl ester (HCl salt; 108 mg, 286 μ mol, 1.0 eq.), 3,5-pyrazoledicarboxylic acid (66.9 mg, 429 μ mol, 1.5 eq.), and HATU (160 mg, 430 μ mol, 1.5 eq.) were combined in DMF (5 mL), and the resulting mixture was stirred for 2 minutes. DIPEA (150 μ L) was added and the mixture was stirred for 1 hour. The mixture was dried under vacuum, and the product was purified using reverse phase chromatography to yield the title compound as a TFA salt (60 mg; purity 98%). MS m/z $[M+H]^+$ calc'd for $C_{26}H_{29}N_3O_6$, 480.21; found 480.4).

10

EXAMPLE 3



A. (2*S*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-((5-[(2-methoxyethyl)methylcarbamoyl]-2H-pyrazole-3-carbonyl)amino)pentanoic Acid Ethyl Ester ($R^7 = -CH_2CH_3$)

B. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-((5-[(2-methoxyethyl)methylcarbamoyl]-2H-pyrazole-3-carbonyl)amino)pentanoic Acid Ethyl Ester ($R^7 = -CH_2CH_3$)

15

C. (2*S*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-((5-[(2-methoxyethyl)methylcarbamoyl]-2H-pyrazole-3-carbonyl)amino)pentanoic Acid ($R^7 = H$)

D. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-((5-[(2-methoxyethyl)methylcarbamoyl]-2H-pyrazole-3-carbonyl)amino)pentanoic Acid ($R^7 = H$)

20

(*R*)-4-Amino-5-biphenyl-4-yl-2-hydroxypentanoic acid ethyl ester (900 mg, 3.0 mmol, 1.0 eq.), 3,5-pyrazoledicarboxylic acid (448 mg, 2.9 mmol, 1.0 eq.), N-(2-methoxyethyl)methylamine (256 mg, 2.9 mmol, 1.0 eq.), and HATU (1090 mg, 2.9 mmol, 1.0 eq.) were combined in DMF (5 mL), and the resulting mixture was stirred for 2 minutes. DIPEA (1 mL) was added and the mixture was stirred for 1 hour. The mixture was dried under vacuum, and the product was purified using reverse phase chromatography (10-70% MeCN/H₂O; 0.05% TFA over 70 minutes) to yield (*R*)-5-biphenyl-4-yl-2-


25

hydroxy-4-({5-[(2-methoxyethyl)methylcarbamoyl]-2H-pyrazole-3-carbonyl}amino)-pentanoic acid ethyl ester (25 mg) as a TFA salt.

This material was separated by preparative HPLC, purified using reverse phase chromatography and dried under vacuum to yield compounds A and B as TFA salts.

- 5 Compounds A and B were each dissolved in EtOH (~10 mL), and 10N NaOH was added to make the solutions basic (~200 μ L). Each mixture was monitored over 1 hour until final deprotection was complete. Each mixture was acidified with an equal volume (~200 μ L) of AcOH and dried under vacuum. The products were purified using reverse phase chromatography to yield compounds C (20.6 mg; purity 100%; MS m/z $[M+H]^+$ calc'd for $C_{26}H_{30}N_4O_6$, 495.22; found 495.22) and D (8.3 mg; purity 100%; MS m/z $[M+H]^+$ calc'd for $C_{26}H_{30}N_4O_6$, 495.22; found 495.22) as TFA salts.

EXAMPLE 4




- A. (2R,4R)-4-{{5-[(Azetidine-1-carbonyl)-2H-pyrazole-3-carbonyl]amino}-5-biphenyl-4-yl}-2-hydroxypentanoic Acid Ethyl Ester ($R^7 = -CH_2CH_3$)

- (2R,4R)-4-Amino-5-biphenyl-4-yl-2-hydroxypentanoic acid ethyl ester (HCl salt; 100 mg, 286 μ mol, 1.0 eq.), 3,5-pyrazoledicarboxylic acid (66.9 mg, 429 μ mol, 1.5 eq.), and HATU (160 mg, 430 μ mol, 1.5 eq.) were combined in DMF (5 mL), and the resulting mixture was stirred for 2 minutes. DIPEA (150 μ L) was added and the mixture was stirred for 1 hour. Azetidine hydrochloride (40.0 mg, 429 μ mol, 1.5 eq.) was added with a second equivalent of HATU, and the mixture was stirred briefly. Additional DIPEA (2.0 eq.) was added and the mixture was stirred. The mixture was dried under vacuum then purified using reverse phase chromatography (10-70% MeCN/H₂O; 0.05% TFA over 70 minutes) to yield the title compound as a TFA salt (20 mg; purity 95%). MS m/z $[M+H]^+$ calc'd for $C_{27}H_{30}N_4O_5$, 491.22; found 491.4.

B. (2*R*,4*R*)-4-[[5-(Azetidine-1-carbonyl)-2*H*-pyrazole-3-carbonyl]amino]-5-biphenyl-4-yl-2-hydroxy-pentanoic acid ($R^7 = H$)

(2*R*,4*R*)-4-Amino-5-biphenyl-4-yl-2-hydroxypentanoic acid ethyl ester (HCl salt; 50 mg, 0.1 mmol, 1.0 eq.), 3,5-pyrazoledicarboxylic acid monohydrate (29.9 mg, 173 μ mol, 1.2 eq.), and HATU (82 mg, 210 μ mol, 1.5 eq.) were combined in DMF (5 mL), and the resulting mixture was stirred for 2 minutes. DIPEA (75 μ L) was added and the mixture was stirred for 1 hour. Azetidine hydrochloride (20.0 mg, 214 μ mol, 1.5 eq.) was added with a second equivalent of HATU, and the mixture was stirred briefly. The mixture was dried under vacuum to yield the crude ester. The crude ester was dissolved in EtOH (~10 mL), and 10N NaOH was added to make the solution basic (~200 μ L). The mixture was monitored over 1 hour until final deprotection was complete. The mixture was acidified with an equal volume (~200 μ L) of AcOH and dried under vacuum. The product was purified using reverse phase chromatography to yield the title compound as a TFA salt (17 mg; purity 95%). MS m/z $[M+H]^+$ calc'd for $C_{25}H_{26}N_4O_5$, 463.19; found 463.4.

EXAMPLE 5




A. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[[5-(pyrrolidine-1-carbonyl)-2*H*-pyrazole-3-carbonyl]amino]pentanoic Acid Ethyl Ester ($R^7 = -CH_2CH_3$)

(2*R*,4*R*)-4-Amino-5-biphenyl-4-yl-2-hydroxypentanoic acid ethyl ester (HCl salt; 85 mg, 240 μ mol, 1.0 eq.), 3,5-pyrazoledicarboxylic acid monohydrate (40 mg, 0.2 mmol, 1.0 eq.), and HATU (90 mg, 0.2 mmol, 1.0 eq.) were combined in DMF (5 mL), and the resulting mixture was stirred for 2 minutes. DIPEA (150 μ L) was added and the mixture was stirred for 1 hour. A second equivalent of HATU was added, followed by pyrrolidine (20.3 μ L, 243 μ mol, 1.0 eq.) (1N in THF) and a second equivalent of DIPEA. Upon completion of the reaction, the mixture was dried under vacuum then purified using reverse phase chromatography (10-70% MeCN/H₂O; 0.05% TFA over 70 minutes) to yield the title compound as a TFA salt (65 mg; purity 98%). MS m/z $[M+H]^+$ calc'd for $C_{28}H_{32}N_4O_5$, 505.24; found 505.4.

B. (2R,4R)-5-Biphenyl-4-yl-2-hydroxy-4-[[5-(pyrrolidine-1-carbonyl)-2H-pyrazole-3-carbonyl]-amino]pentanoic Acid ($R^7 = H$)

(2R,4R)-4-Amino-5-biphenyl-4-yl-2-hydroxypentanoic acid ethyl ester (HCl salt; 50 mg, 0.1 mmol, 1.0 eq.), 3,5-pyrazoledicarboxylic acid monohydrate (20.0 mg, 0.1 mmol, 1.0 eq.), and HATU (50 mg, 0.1 mmol, 1.0 eq.) were combined in DMF (5 mL), and the resulting mixture was stirred for 2 minutes. DIPEA (75 μ L) was added and the mixture was stirred for 1 hour. A second equivalent of HATU was added, followed by pyrrolidine (20.0 mg, 280 μ mol, 2.0 eq.) and a second equivalent of DIPEA. The mixture was dried under vacuum to yield the crude ester. The crude ester was dissolved in EtOH (~10 mL), and 10N NaOH was added to make the solution basic (~200 μ L). The mixture was monitored over 1 hour until final deprotection was complete. The mixture was acidified with an equal volume (~200 μ L) of AcOH and dried under vacuum. The product was purified using reverse phase chromatography to yield the title compound as a TFA salt (25 mg; purity 95%). MS m/z $[M+H]^+$ calc'd for $C_{26}H_{28}N_4O_5$, 477.21; found 477.0.

EXAMPLE 6




A. (2R,4R)-5-Biphenyl-4-yl-2-hydroxy-4-[(2-hydroxypyrimidine-5-carbonyl)amino]pentanoic Acid Ethyl Ester ($R^7 = -CH_2CH_3$)

B. (2R,4R)-5-Biphenyl-4-yl-2-hydroxy-4-[(2-hydroxypyrimidine-5-carbonyl)amino]pentanoic Acid ($R^7 = H$)

(2R,4R)-4-Amino-5-biphenyl-4-yl-2-hydroxypentanoic acid ethyl ester (HCl salt; 200 mg, 572 μ mol, 1.0 eq.) and 2-hydroxypyrimidine-5-carboxylic acid (88.1 mg, 629 μ mol) were suspended in DMF (5.0 mL). HATU (239 mg, 629 μ mol) was added followed by DIPEA (299 μ L), and the resulting mixture was stirred at room temperature until the reaction was complete (~2 hours). The mixture was divided into two equal portions and both solutions were concentrated. One portion was purified by preparative HPLC (10-70% MeCN/ H_2O) to produce compound A as a TFA salt (59.2 mg). MS m/z $[M+H]^+$ calc'd for

$C_{24}H_{25}N_3O_5$, 436.18; found 436.4. The remaining portion was dissolved in THF (2.0 mL). LiOH monohydrate (120 mg, 2.9 mmol) in water (2.0 mL) was added and the mixture was stirred at room temperature for ~ 1 hour. The reaction was quenched by addition of AcOH and the solution was concentrated. The crude product was purified by preparative HPLC (10-70% MeCN/H₂O) to yield compound B as a TFA salt (27.8 mg). MS m/z $[M+H]^+$ calc'd for $C_{22}H_{21}N_3O_5$, 408.15; found 408.4.

EXAMPLE 7



A. (2R,4R)-5-Biphenyl-4-yl-2-hydroxy-4-[(3-hydroxyisoxazole-5-carbonyl)amino]pentanoic Acid Ethyl Ester ($R^7 = -CH_2CH_3$)

3-Hydroxyisoxazole-5-carboxylic acid (81.2 mg, 629 μ mol, 1.1 eq.) was dissolved in DMF (5.0 mL). DIPEA (299 μ L) and HOAt (85.6 mg, 629 μ mol, 1.1 eq.) were added, followed by EDCI (111 μ L, 1.1 eq.). The solution was stirred for 5 minutes. (2R,4R)-4-Amino-5-biphenyl-4-yl-2-hydroxypentanoic acid ethyl ester (HCl salt; 200 mg, 572 μ mol, 1.0 eq.) was then added and the resulting mixture were stirred at room temperature overnight. The mixture was concentrated, and the crude product was purified by preparative HPLC (10-70% MeCN/H₂O) to yield the title compound as a TFA salt (50.8 mg, 98% purity). MS m/z $[M+H]^+$ calc'd for $C_{23}H_{24}N_2O_6$, 425.16; found 425.4.

B. (2R,4R)-5-Biphenyl-4-yl-2-hydroxy-4-[(3-hydroxyisoxazole-5-carbonyl)amino]pentanoic Acid ($R^7 = H$)

(2R,4R)-4-Amino-5-biphenyl-4-yl-2-hydroxypentanoic acid ethyl ester (HCl salt; 200 mg, 572 μ mol, 1.0 eq.) and 3-hydroxyisoxazole-5-carboxylic acid (81.2 mg, 629 μ mol) were suspended in DMF (5.0 mL). HATU (239 mg, 629 μ mol) was added followed by DIPEA (299 μ L), and the resulting mixture was stirred at room temperature overnight. The mixture was concentrated and the crude compound was used without purification. The crude material was dissolved in THF (3.0 mL). LiOH monohydrate (240 mg, 5.7 mmol) in water (3.0 mL) was added and the mixture was stirred at room temperature for ~ 1 hour.

The mixture was concentrated and the crude product was purified by preparative HPLC (10-70% MeCN/H₂O) to yield the title compound as a TFA salt (36.6 mg, 97% purity). MS *m/z* [M+H]⁺ calc'd for C₂₁H₂₀N₂O₆, 397.13; found 397.2.


- C. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(3-hydroxyisoxazole-5-carbonyl)amino]pentanoic acid 5-methyl-2-oxo-[1,3]dioxol-4-ylmethyl Ester (R⁷ = -CH₂-5-methyl-[1,3]dioxol-2-one)
- (2*R*,4*R*)-4-Amino-5-biphenyl-4-yl-2-hydroxy-pentanoic acid ethyl ester; hydrochloride (1.6 g, 4.6 mmol, 1.0 eq.) was dissolved in DCM (100 mL) and cooled to 0°C. Di-*t*-butyldicarbonate (1.2 g, 5.4 mmol, 1.2 eq.) was added as a solid followed by DIPEA (1.6 mL, 9.1 mmol). The resulting mixture was stirred overnight at room temperature. The product was purified by flash chromatography (0-75% EtOAc/hexanes) then dissolved in EtOH and sufficient equivalents of 10N NaOH to make the solution basic (pH = 14). The reaction was monitored until deprotection was complete (~1 hour). EtOAc (200 mL) was added and the mixture was washed with 1N HCl (100 mL) followed by saturated aqueous NaCl (100 mL). The organic layer was retained and dried over anhydrous MgSO₄ for 10 minutes prior to filtration and evacuation to dryness, to yield (2*R*,4*R*)-5-biphenyl-4-yl-4-*t*-butoxycarbonylamino-2-hydroxypentanoic acid (1.2 g).
- (2*R*,4*R*)-5-Biphenyl-4-yl-4-*t*-butoxycarbonylamino-2-hydroxypentanoic acid (500 mg, 1.0 mmol, 1.0 eq.) was dissolved in dry DMF (10.0 mL). The solution was cooled to 0°C and dicesium carbonate (465 mg, 1.4 mmol, 1.1 eq.) was added. The mixture was stirred for 30 minutes and 4-chloromethyl-5-methyl-1,3-dioxol-2-one (212 mg, 1.4 mmol, 1.1 eq.) was added. The resulting mixture was stirred for an additional 1 hour at 0°C, then warmed to room temperature while stirring. The mixture was further stirred overnight. EtOAc (20 mL) was added, followed by washing with saturated aqueous NaCl (100 mL). The product was dried under vacuum and purified using reverse phase chromatography (10-70% MeCN/H₂O; 0.05% TFA over 70 minutes) to yield (2*R*,4*R*)-5-biphenyl-4-yl-4-*t*-butoxycarbonylamino-2-hydroxypentanoic acid 5-methyl-2-oxo-[1,3]dioxol-4-ylmethyl ester. This material was dissolved in 4N HCL in dioxane and stirred for 2 hours at room temperature. The material was dried under vacuum then azeotroped with toluene to yield (2*R*,4*R*)-4-amino-5-biphenyl-4-yl-2-hydroxypentanoic acid 5-methyl-2-oxo-[1,3]dioxol-4-ylmethyl ester, which was used in the next step without further purification.

(2*R*,4*R*)-4-amino-5-biphenyl-4-yl-2-hydroxypentanoic acid 5-methyl-2-oxo-[1,3]dioxol-4-ylmethyl ester (100 mg, 0.3 mmol, 1.0 eq.) and 3-hydroxyisoxazole-5-

carboxylic acid (50.5 mg, 391 μ mol, 1.5 eq.) were combined in DMF (5 mL) and the resulting mixture was stirred for 2 minutes. DIPEA was added and the resulting mixture was stirred for 1 hour. The material was dried under vacuum then purified using reverse phase chromatography to yield the title compound (40 mg, 98% purity). MS m/z $[M+H]^+$

5 calc'd for $C_{26}H_{24}N_2O_9$, 509.15; found 509.4.

EXAMPLE 8



A. (2R,4R)-5-Biphenyl-4-yl-2-hydroxy-4-([5-(1-hydroxy-1-methylethyl)-1H-pyrazole-3-carbonyl]amino}pentanoic acid ethyl ester ($R^7 = -CH_2CH_3$)

10 (2R,4R)-4-Amino-5-biphenyl-4-yl-2-hydroxypentanoic acid ethyl ester (HCl salt; 100 mg, 0.3 mmol, 1.0 eq.), 5-(1-hydroxy-1-methylethyl)-1H-pyrazole-3-carboxylic acid (73 mg, 429 μ mmol, 1.5 eq.), and HATU (160 mg, 430 μ mol, 1.5 eq.) were combined in DMF (5 mL), and the resulting mixture was stirred for 2 minutes. DIPEA (149 μ L) was added and the mixture was stirred for 1 hour. Upon completion of the reaction, the mixture

15 was dried under vacuum then purified using reverse phase chromatography (10-70% MeCN/ H_2O ; 0.05% TFA over 70 minutes) to yield the title compound as a TFA salt (50 mg). MS m/z $[M+H]^+$ calc'd for $C_{26}H_{31}N_3O_5$, 466.23; found 466.4.

B. (2R,4R)-5-Biphenyl-4-yl-2-hydroxy-4-([5-(1-hydroxy-1-methylethyl)-1H-pyrazole-3-carbonyl]amino}pentanoic acid ($R^7 = H$)

20 (2R,4R)-4-Amino-5-biphenyl-4-yl-2-hydroxypentanoic acid ethyl ester (HCl salt; 60 mg, 0.2 mmol, 1.0 eq.), 5-(1-hydroxy-1-methylethyl)-1H-pyrazole-3-carboxylic acid (43.8 mg, 257 μ mol, 150 eq.), and HATU (98 mg, 260 μ mol, 1.5 eq.) were combined in DMF (5 mL), and the resulting mixture was stirred for 2 minutes. DIPEA (89.6 μ L) was added and the mixture was stirred for 1 hour. The mixture was dried under vacuum to


25 yield the crude ester. The crude ester was dissolved in EtOH (~10 mL), and 10N NaOH was added to make the solution basic (~200 μ L). The mixture was monitored over 1 hour until final deprotection was complete. The mixture was acidified with an equal volume (~200 μ L) of AcOH and dried under vacuum. The product was purified using reverse

phase chromatography to yield the title compound as a TFA salt (30 mg). MS m/z $[M+H]^+$ calc'd for $C_{24}H_{27}N_3O_5$, 438.20; found 438.4.

EXAMPLE 9


2-Hydroxypyrimidine-5-carboxylic acid ((1*R*,3*R*)-1-biphenyl-4-ylmethyl-3-carbamoyl-3-hydroxypropyl)amide

5




(2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(2-hydroxypyrimidine-5-carbonyl)amino]pentanoic acid (14.5 mg, 35.6 μ mol, 1.0 eq.) and HATU (13.5 mg, 35.6 μ mol, 1.0 eq.) were combined in DMF (5 mL), and the resulting mixture was stirred for 2 minutes. 0.5 M ammonia in dioxane was added, followed by DIPEA (12.4 μ L). The resulting mixture was stirred for 1 hour then dried under vacuum. The product was purified using reverse phase chromatography (10-70% MeCN/ H_2O ; 0.05% TFA over 70 minutes) to yield the title compound (2 mg). MS m/z $[M+H]^+$ calc'd for $C_{22}H_{22}N_4O_4$, 407.16; found 407.4.

15 Following the procedures described in the examples herein, and substituting the appropriate starting materials and reagents, the compounds in Examples 10 to 15, having the following formula, were also prepared:



EXAMPLE 10


20 These were prepared as the parent compound or as a TFA salt.



Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
1	-OCH ₂ CH ₃	H	H	C ₂₂ H ₂₄ N ₄ O ₄	409.18	409.4
2	-OH	H	H	C ₂₀ H ₂₀ N ₄ O ₄	381.15	381.4

1. (*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(3H-[1,2,3]triazole-4-carbonyl)amino]pentanoic acid ethyl ester (TFA salt)


5 2. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(3H-[1,2,3]triazole-4-carbonyl)amino]pentanoic acid (TFA salt)



Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
3	-OH	-OH	H	C ₂₀ H ₂₀ N ₄ O ₅	397.14	397.0
4	-OCH ₂ CH ₃	-OH	H	C ₂₂ H ₂₄ N ₄ O ₅	425.17	425.2

3. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(1-hydroxy-1H-[1,2,3]triazole-4-carbonyl)-amino]-pentanoic acid

10 4. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(1-hydroxy-1H-[1,2,3]triazole-4-carbonyl)-amino]-pentanoic acid ethyl ester (TFA salt)



Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
5	-OCH ₂ CH ₃	H	H	C ₂₂ H ₂₄ N ₄ O ₄	409.18	409.4
6	-OH	H	H	C ₂₀ H ₂₀ N ₄ O ₄	381.15	381.4
7	-OH	-CH ₃	4-chlorophenyl	C ₂₇ H ₂₅ ClN ₄ O ₄	505.16	506.2
8	-OCH ₂ CH ₃	-OH	H	C ₂₂ H ₂₄ N ₄ O ₅	425.17	425.2
9	-OH	-OH	H	C ₂₀ H ₂₀ N ₄ O ₅	397.14	397.2


15 5. (*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(1H-[1,2,4]triazole-3-carbonyl)amino]pentanoic acid ethyl ester (TFA salt)

6. (*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(1H-[1,2,4]triazole-3-carbonyl)amino]pentanoic acid (TFA salt)

7. (*R*)-5-Biphenyl-4-yl-4-{[1-(4-chlorophenyl)-5-methyl-1H-[1,2,4]triazole-3-carbonyl]-amino}-2-hydroxypentanoic acid (TFA salt)

5 8. (*2R,4R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(5-hydroxy-1H-[1,2,4]triazole-3-carbonyl)amino]-pentanoic acid ethyl ester (TFA salt)


9. (*2R,4R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(5-hydroxy-1H-[1,2,4]triazole-3-carbonyl)amino]-pentanoic acid (TFA salt)



Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
10	-OH	2-pyridine	H	C ₂₅ H ₂₃ N ₅ O ₄	458.18	458.2

10

10. (*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(5-pyridin-2-yl-4H-[1,2,4]triazole-3-carbonyl)amino]pentanoic acid (TFA salt)




Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
11	-OH	-CH ₃	H	C ₂₁ H ₂₂ N ₄ O ₄	395.16	395.2
12	-OH	-Cl	H	C ₂₀ H ₁₉ ClN ₄ O ₄	415.11	415.4

15 11. (*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(5-methyl-2H-[1,2,4]triazole-3-carbonyl)amino]pentanoic acid (TFA salt)

12. (*2R,4R*)-5-Biphenyl-4-yl-4-[(5-chloro-2H-[1,2,4]triazole-3-carbonyl)amino]-2-hydroxypentanoic acid


EXAMPLE 11

20 These were all prepared as TFA salts.




Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
1	-OH	H	3-carboxybenzyl	C ₃₂ H ₂₈ N ₄ O ₆	565.20	565.6
2	-OH	H	4-carboxybenzyl	C ₃₂ H ₂₈ N ₄ O ₆	565.20	565.6
3	-OH	H	H	C ₂₄ H ₂₂ N ₄ O ₄	431.16	431.2
4	-OH	F	H	C ₂₄ H ₂₁ FN ₄ O ₄	449.15	449.0
5	-OCH ₂ CH ₃	Cl	H	C ₂₆ H ₂₅ ClN ₄ O ₄	493.16	493.2

1. 3-[5-((*R*)-1-Biphenyl-4-ylmethyl-3-carboxy-3-hydroxypropylcarbamoyl)-benzotriazol-1-ylmethyl]-benzoic acid
- 5 2. 4-[5-((*IR*,*3S*)-1-Biphenyl-4-ylmethyl-3-carboxy-3-hydroxypropylcarbamoyl)-benzotriazol-1-ylmethyl]-benzoic acid
3. (*R*)-4-[(1*H*-Benzotriazole-5-carbonyl)amino]-5-biphenyl-4-yl-2-hydroxypentanoic acid
4. (*R*)-5-Biphenyl-4-yl-4-[(7-fluoro-1*H*-benzotriazole-5-carbonyl)amino]-2-hydroxy-
- 10 pentanoic acid
5. (*R*)-5-Biphenyl-4-yl-4-[(7-chloro-1*H*-benzotriazole-5-carbonyl)amino]-2-hydroxy-pentanoic acid ethyl ester




Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
6	-OH	-CH ₃	H	C ₂₅ H ₂₄ N ₄ O ₄	445.18	445.2

- 15 6. (*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(7-methyl-3*H*-benzotriazole-5-carbonyl)amino]pentanoic acid




Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
7	-OH	Cl	H	C ₂₄ H ₂₁ ClN ₄ O ₄	465.13	465.0

7. (*R*)-5-Biphenyl-4-yl-4-[(4-chloro-1H-benzotriazole-5-carbonyl)amino]-2-hydroxy-pentanoic acid




Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
8	-OH	-CH ₃	H	C ₂₆ H ₂₄ N ₂ O ₅	445.17	445.4

8. (*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(2-methyl-benzooxazole-6-carbonyl)-amino]-pentanoic acid



Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
9	-OH	H	H	C ₂₄ H ₂₂ N ₄ O ₄	431.16	431.2


9. (*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(3H-imidazo[4,5-b]pyridine-6-carbonyl)-amino]-pentanoic acid



Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
10	-OCH ₂ CH ₃	H	H	C ₂₅ H ₂₅ N ₅ O ₄	460.19	460.4
11	-OH	H	H	C ₂₃ H ₂₁ N ₅ O ₄	432.16	432.2

10. (*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(1*H*-[1,2,3]triazolo[4,5-*b*]pyridine-6-carbonyl)-amino]-pentanoic acid ethyl ester

11. (*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(1*H*-[1,2,3]triazolo[4,5-*b*]pyridine-6-carbonyl)-amino]-pentanoic acid



5

-XR³R⁴ =

Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
12	-OCH ₂ CH ₃	H	H	C ₂₅ H ₂₅ N ₅ O ₄	460.19	460.4
13	-OH	H	H	C ₂₃ H ₂₁ N ₅ O ₄	432.16	432.4
14	-OCH ₂ CH ₃	H	H	C ₂₅ H ₂₅ N ₅ O ₄	460.19	460.4
15	-OH	H	H	C ₂₃ H ₂₁ N ₅ O ₄	432.16	432.4

12. (*2S,4R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(3*H*-[1,2,3]triazolo[4,5-*b*]pyridine-6-carbonyl)-amino]-pentanoic acid ethyl ester

13. (*2S,4R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(3*H*-[1,2,3]triazolo[4,5-*b*]pyridine-6-carbonyl)-amino]-pentanoic acid


14. (*2R,4R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(3*H*-[1,2,3]triazolo[4,5-*b*]pyridine-6-carbonyl)-amino]-pentanoic acid ethyl ester

15. (*2R,4R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(3*H*-[1,2,3]triazolo[4,5-*b*]pyridine-6-carbonyl)-amino]-pentanoic acid

15

EXAMPLE 12


These were prepared as the parent compound or as a TFA salt.



Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
1	-OH	-COOH	H	C ₂₃ H ₂₁ NO ₇	424.13	424.0


1. 5-((*R*)-1-Biphenyl-4-ylmethyl-3-carboxy-3-hydroxypropylcarbamoyl)furan-2-carboxylic acid (TFA salt)

20













Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
2	-OH	-COOH	H	C ₂₃ H ₂₁ NO ₆ S	440.11	440.0

2. 5-((*R*)-1-Biphenyl-4-ylmethyl-3-carboxy-3-hydroxypropylcarbamoyl)thiophene-2-carboxylic acid




5








Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
3	-OH	-OCH ₃	H	C ₂₂ H ₂₂ N ₂ O ₆	411.15	411.2
4	-OH		H	C ₂₇ H ₂₃ ClN ₂ O ₅	491.13	491.2
5	-OH		H	C ₂₉ H ₂₇ N ₃ O ₇	530.19	530.4
6	-OH		H	C ₂₇ H ₂₃ ClN ₂ O ₅	491.13	491.4
7	-OH		H	C ₂₇ H ₃₀ N ₂ O ₅	463.22	463.4
8	-OH		H	C ₂₇ H ₂₃ FN ₂ O ₅	475.16	475.0
9	-OH		H	C ₂₇ H ₂₃ FN ₂ O ₅	475.16	475.0


Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
10	-OH		H	C ₂₅ H ₂₄ N ₄ O ₅	461.17	461.2
11	-OH		H	C ₂₇ H ₂₂ Cl ₂ N ₂ O ₅	525.09	525.4
12	-OH		H	C ₂₇ H ₂₂ Cl ₂ N ₂ O ₅	525.09	525.4
13	-OH		H	C ₂₈ H ₂₆ N ₂ O ₆	487.18	487.2
14	-OH	-COOH	H	C ₂₂ H ₂₀ N ₂ O ₇	425.13	425.2

3. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(3-methoxyisoxazole-5-carbonyl)amino]-pentanoic acid
4. (2*R*,4*R*)-5-Biphenyl-4-yl-4-[[3-(4-chlorophenyl)-isoxazole-5-carbonyl]-amino]-2-
- 5 hydroxy-pentanoic acid (TFA salt)
5. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[[3-(2-methoxy-benzoylamino)-isoxazole-5-carbonyl]-amino]-pentanoic acid (TFA salt)
6. (2*R*,4*R*)-5-Biphenyl-4-yl-4-[[3-(2-chlorophenyl)-isoxazole-5-carbonyl]-amino]-2-
- hydroxy-pentanoic acid (TFA salt)
- 10 7. (2*R*,4*R*)-5-Biphenyl-4-yl-4-[(3-cyclohexyl-isoxazole-5-carbonyl)-amino]-2-
- hydroxy-pentanoic acid (TFA salt)
8. (2*R*,4*R*)-5-Biphenyl-4-yl-4-[[3-(3-fluorophenyl)-isoxazole-5-carbonyl]-amino]-2-
- hydroxy-pentanoic acid
9. (2*R*,4*R*)-5-Biphenyl-4-yl-4-[[3-(2-fluorophenyl)-isoxazole-5-carbonyl]-amino]-2-
- 15 hydroxy-pentanoic acid
10. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[[3-(1-methyl-1*H*-pyrazol-4-yl)-isoxazole-5-carbonyl]-amino]-pentanoic acid (TFA salt)
11. (2*R*,4*R*)-5-Biphenyl-4-yl-4-[[3-(2,5-dichlorophenyl)-isoxazole-5-carbonyl]-amino]-2-hydroxy-pentanoic acid


12. (2*R*,4*R*)-5-Biphenyl-4-yl-4-{{3-(3,4-dichlorophenyl)-isoxazole-5-carbonyl]-amino}-2-hydroxy-pentanoic acid
13. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-{{3-(2-methoxy-phenyl)-isoxazole-5-carbonyl]-amino}-pentanoic acid
- 5 14. 5-((1*R*,3*R*)-1-Biphenyl-4-ylmethyl-3-carboxy-3-hydroxy-propylcarbamoyl)-isoxazole-3-carboxylic acid



Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
15	-OH		H	C ₂₈ H ₂₆ N ₂ O ₆	487.18	487.4
16	-OH		H	C ₂₇ H ₂₃ ClN ₂ O ₅	491.13	491.4
17	-OH		H	C ₂₇ H ₂₃ ClN ₂ O ₅	491.13	491.2
18	-OH		H	C ₂₈ H ₂₆ N ₂ O ₆	487.18	487.2
19	-OH		H	C ₂₆ H ₂₉ N ₃ O ₆	480.21	480.2
20	-OH		H	C ₂₈ H ₂₆ N ₂ O ₆	487.18	487.2
21	-OH		H	C ₂₇ H ₂₄ N ₂ O ₆	473.16	473.0


Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
22	-OH		H	C ₂₈ H ₂₄ ClN ₃ O ₆	534.14	535.0

15. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4- {[5-(2-methoxy-phenyl)-isoxazole-3-carbonyl]-amino}-pentanoic acid (TFA salt)
16. (2*R*,4*R*)-5-Biphenyl-4-yl-4- {[5-(2-chlorophenyl)-isoxazole-3-carbonyl]-amino}-2-hydroxy-pentanoic acid (TFA salt)
17. (2*R*,4*R*)-5-Biphenyl-4-yl-4- {[5-(4-chlorophenyl)-isoxazole-3-carbonyl]-amino}-2-hydroxy-pentanoic acid (TFA salt)
18. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4- {[5-(3-methoxy-phenyl)-isoxazole-3-carbonyl]-amino}-pentanoic acid (TFA salt)
19. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4- [(5-morpholin-4-ylmethyl-isoxazole-3-carbonyl)-amino]-pentanoic acid (TFA salt)
20. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4- {[5-(4-methoxy-phenyl)-isoxazole-3-carbonyl]-amino}-pentanoic acid (TFA salt)
21. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4- {[5-(2-hydroxy-phenyl)-isoxazole-3-carbonyl]-amino}-pentanoic acid (TFA salt)
22. (2*R*,4*R*)-5-Biphenyl-4-yl-4- {[5-(2-chlorobenzoylamino)-isoxazole-3-carbonyl]-amino}-2-hydroxy-pentanoic acid (TFA salt)




Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
23	-OH	phenyl	H	C ₂₇ H ₂₄ N ₂ O ₅	457.17	457.4

23. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4- [(2-phenyl-oxazole-5-carbonyl)-amino]-pentanoic acid (TFA salt)




Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
24	-OH	-NHC(O)CH ₃	H	C ₂₃ H ₂₃ N ₃ O ₅ S	454.14	454.2

24. (2*R*,4*R*)-4-[(2-Acetylamino-thiazole-5-carbonyl)-amino]-5-biphenyl-4-yl-2-hydroxy-pentanoic acid (TFA salt)




Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
25	-OH		H	C ₂₉ H ₂₇ N ₃ O ₆ S	546.16	546.2

25. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-{[2-(2-methoxy-benzoylamino)-thiazole-4-carbonyl]-amino}-pentanoic acid (TFA salt)



Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
26	-OH	2-chlorophenyl	absent	C ₂₆ H ₂₂ ClN ₃ O ₅	492.12	492.4

26. (2*R*,4*R*)-5-Biphenyl-4-yl-4-{[3-(2-chlorophenyl)-[1,2,4]oxadiazole-5-carbonyl]-amino}-2-hydroxy-pentanoic acid



Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found


Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
27	-OH	=O	H	C ₂₀ H ₁₉ N ₃ O ₆	398.13	398.2

27. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(5-oxo-4,5-dihydro-[1,2,4]oxadiazole-3-carbonyl)-amino]-pentanoic acid

5


EXAMPLE 13

These were prepared as the parent compound or as a TFA salt.



Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
1	-OH	-COOH	H	C ₂₃ H ₂₁ N ₃ O ₆	436.14	436.2


10 1. 5-((*R*)-1-Biphenyl-4-ylmethyl-3-carboxy-3-hydroxypropylcarbamoyl)-pyrazine-2-carboxylic acid (TFA salt)



Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
2	-OCH ₂ CH ₃	H	H	C ₂₄ H ₂₅ N ₃ O ₄	420.18	420.4


2. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(pyrimidine-5-carbonyl)-amino]-pentanoic

15 acid ethyl ester (TFA salt)




Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
3	-OH	-COOH	H	C ₂₃ H ₂₁ N ₃ O ₆	436.14	436.2
4	-OCH ₂ CH ₃	-OH	H	C ₂₄ H ₂₅ N ₃ O ₅	436.18	436.4
5	-OH	-OH	H	C ₂₂ H ₂₁ N ₃ O ₅	408.15	408.2

3. 6-((*R*)-1-Biphenyl-4-ylmethyl-3-carboxy-3-hydroxypropylcarbamoyl)-pyridazine-3-carboxylic acid (TFA salt)
4. (*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(6-hydroxypyridazine-3-carbonyl)-amino]-pentanoic acid ethyl ester
5. (*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(6-hydroxypyridazine-3-carbonyl)-amino]-pentanoic acid (TFA salt)




Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
6	-OH	H	H	C ₂₂ H ₂₃ N ₃ O ₅	410.16	410.0

6. (*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(6-oxo-1,4,5,6-tetrahydro-pyridazine-3-carbonyl)-amino]-pentanoic acid



Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
7	-OH	-COOH	H	C ₂₄ H ₂₂ N ₂ O ₆	435.15	435.2

7. 6-((*R*)-1-Biphenyl-4-ylmethyl-3-carboxy-3-hydroxypropylcarbamoyl)nicotinic acid




Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
8	-OH	-COOH	H	C ₂₄ H ₂₂ N ₂ O ₆	435.15	435.2

8. 5-((*R*)-1-Biphenyl-4-ylmethyl-3-carboxy-3-hydroxypropylcarbamoyl)-pyridine-2-carboxylic acid

EXAMPLE 14

This was prepared as the parent compound.




Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
1	-OH	absent	H	C ₁₉ H ₁₉ N ₅ O ₄	382.14	382.2

1. (*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(1H-tetrazole-5-carbonyl)amino]pentanoic acid





5


EXAMPLE 15

These were prepared as TFA salts.



Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
1	-OH	H	H	C ₂₁ H ₂₁ N ₃ O ₄	380.15	380.0
2	-OH	-CH ₂ COOH	H	C ₂₃ H ₂₃ N ₃ O ₆	438.16	438.4
3	-OH	-CH ₂ OH	H	C ₂₂ H ₂₃ N ₃ O ₅	410.16	410.4
4	-OH	-CH ₂ OCH ₃	H	C ₂₃ H ₂₅ N ₃ O ₅	424.18	424.4
5	-OH	-COOH	-CH ₃	C ₂₃ H ₂₃ N ₃ O ₆	438.16	438.4
6	-OH	-COOH	3-SCF ₃ , 4-Cl-benzyl	C ₃₀ H ₂₅ ClF ₃ N ₃ O ₆ S	648.11	648.0
7	-OH	-COOH	3-Cl-benzyl	C ₂₉ H ₂₆ ClN ₃ O ₆	548.15	548.2
8	-OH	-COOH	2,6-diF, 4-Cl-benzyl	C ₂₉ H ₂₄ ClF ₂ N ₃ O ₆	584.13	584.0
9	-OH	-COOH	2-Cl-benzyl	C ₂₉ H ₂₆ ClN ₃ O ₆	548.15	548.2
10	-OH	-COOH	2-Cl, 5-F-benzyl	C ₂₉ H ₂₅ ClFN ₃ O ₆	566.14	566.0
11	-OH	-COOH	3-Cl, 5-F-benzyl	C ₂₉ H ₂₅ ClFN ₃ O ₆	566.14	566.0
12	-OH	-COOH	3-OCF ₃ , 4-Cl-benzyl	C ₃₀ H ₂₅ ClF ₃ N ₃ O ₇	632.13	632.0
13	-OH	-COOH	2-F, 4-Cl-benzyl	C ₂₉ H ₂₅ ClFN ₃ O ₆	566.14	566.2
14	-OH	-COOH	3-Cl, 4-F-benzyl	C ₂₉ H ₂₅ ClFN ₃ O ₆	566.14	566.2
15	-OH	-COOH	4-Cl-benzyl	C ₂₉ H ₂₆ ClN ₃ O ₆	548.15	548.2


Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
16	-OH	-COOH	2,6-diF, 3-Cl- benzyl	C ₂₉ H ₂₄ ClF ₂ N ₃ O ₆	584.13	584.0
17	-OH	-COOH	3- methoxy- benzyl	C ₃₀ H ₂₉ N ₃ O ₇	544.20	544.2
18	-OCH ₂ -CH ₃	-C(O)O- CH ₂ CH ₃	H	C ₂₆ H ₂₉ N ₃ O ₆	480.21	480.2
19	-OH	-CH ₃	H	C ₂₂ H ₂₃ N ₃ O ₄	394.17	394.2
20	-OH		H	C ₂₆ H ₂₄ N ₄ O ₄	457.18	457.4
21	-OH		H	C ₂₆ H ₂₄ N ₄ O ₄	457.18	457.2
22	-OH		H	C ₂₅ H ₂₃ N ₅ O ₄	458.18	458.2
23	-OH	1-methyl-1H- pyrazole	H	C ₂₅ H ₂₅ N ₅ O ₄	460.19	460.2
24	-OH	2-Cl-phenyl	H	C ₂₇ H ₂₄ ClN ₃ O ₄	490.15	490.0
25	-OH	3-Cl-phenyl	H	C ₂₇ H ₂₄ ClN ₃ O ₄	490.15	490.4
26	-OH	2,4-diCl- phenyl	H	C ₂₇ H ₂₃ Cl ₂ N ₃ O ₄	524.11	524.0
27	-OH	2,5-diCl- phenyl	H	C ₂₇ H ₂₃ Cl ₂ N ₃ O ₄	524.11	524.0
28	-OH	2-hydroxy- phenyl	H	C ₂₇ H ₂₅ N ₃ O ₅	472.18	472.6
29	-OCH ₂ -CH ₃	2-hydroxy- phenyl	H	C ₂₉ H ₂₉ N ₃ O ₅	500.21	500.4
30	-OH	2-hydroxy- phenyl	-CH ₂ - CHOH- CH ₂ OH	C ₃₀ H ₃₁ N ₃ O ₇	546.22	546.4
31	-OH	3-NHC(O)- CH ₃ -phenyl	H	C ₂₉ H ₂₈ N ₄ O ₅	513.21	513.2
32	-OH		H	C ₂₉ H ₂₈ N ₄ O ₆	529.20	530.4
33	-OH	phenyl	H	C ₂₇ H ₂₅ N ₃ O ₄	456.18	456.2
34	-OH	phenyl	-CH ₃	C ₂₈ H ₂₇ N ₃ O ₄	470.20	470.2
35	-OH	phenyl	-CH ₂ - COOH	C ₂₉ H ₂₇ N ₃ O ₆	514.19	514.6
36	-OH	phenyl	-(CH ₂) ₂ - COOH	C ₃₀ H ₂₉ N ₃ O ₆	528.21	528.2
37	-OH	furan	H	C ₂₅ H ₂₃ N ₃ O ₅	446.16	446.2
38	-OH	naphthyl	H	C ₃₁ H ₂₇ N ₃ O ₄	506.20	506.2
39	-OH	4-biphenyl	H	C ₃₃ H ₂₉ N ₃ O ₄	532.22	532.2
40	-OH	4-Cl-phenyl	H	C ₂₇ H ₂₄ ClN ₃ O ₄	490.15	490.2




Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
41		-C(CH ₃) ₂ OH	H	C ₂₉ H ₃₁ N ₃ O ₈	550.21	550.4









1. (R)-5-Biphenyl-4-yl-2-hydroxy-4-[(1H-pyrazole-3-carbonyl)amino]pentanoic acid
2. (2R,4R)-5-Biphenyl-4-yl-4-[(5-carboxymethyl-1H-pyrazole-3-carbonyl)amino]-2-hydroxy-pentanoic acid
- 5 3. (2R,4R)-5-Biphenyl-4-yl-2-hydroxy-4-[(5-hydroxymethyl-1H-pyrazole-3-carbonyl)-amino]-pentanoic acid
4. (2R,4R)-5-Biphenyl-4-yl-2-hydroxy-4-[(5-methoxymethyl-1H-pyrazole-3-carbonyl)-amino]-pentanoic acid
5. 5-((1R,3R)-1-Biphenyl-4-ylmethyl-3-carboxy-3-hydroxypropylcarbamoyl)-2-methyl-2H-pyrazole-3-carboxylic acid
- 10 6. 5-((R)-1-Biphenyl-4-ylmethyl-3-carboxy-3-hydroxypropylcarbamoyl)-2-(4-chloro-3-trifluoromethylsulfanyl-benzyl)-2H-pyrazole-3-carboxylic acid
7. 5-((R)-1-Biphenyl-4-ylmethyl-3-carboxy-3-hydroxypropylcarbamoyl)-2-(3-chlorobenzyl)-2H-pyrazole-3-carboxylic acid
- 15 8. 5-((R)-1-Biphenyl-4-ylmethyl-3-carboxy-3-hydroxy-propylcarbamoyl)-2-(4-chloro-2,6-difluorobenzyl)-2H-pyrazole-3-carboxylic acid
9. 5-((R)-1-Biphenyl-4-ylmethyl-3-carboxy-3-hydroxy-propylcarbamoyl)-2-(2-chlorobenzyl)-2H-pyrazole-3-carboxylic acid
10. 5-((R)-1-Biphenyl-4-ylmethyl-3-carboxy-3-hydroxy-propylcarbamoyl)-2-(2-chloro-20 5-fluorobenzyl)-2H-pyrazole-3-carboxylic acid
11. 5-((R)-1-Biphenyl-4-ylmethyl-3-carboxy-3-hydroxy-propylcarbamoyl)-2-(3-chloro-5-fluorobenzyl)-2H-pyrazole-3-carboxylic acid
12. 5-((R)-1-Biphenyl-4-ylmethyl-3-carboxy-3-hydroxy-propylcarbamoyl)-2-(4-chloro-3-trifluoromethoxybenzyl)-2H-pyrazole-3-carboxylic acid
- 25 13. 5-((R)-1-Biphenyl-4-ylmethyl-3-carboxy-3-hydroxy-propylcarbamoyl)-2-(4-chloro-2-fluorobenzyl)-2H-pyrazole-3-carboxylic acid
14. 5-((R)-1-Biphenyl-4-ylmethyl-3-carboxy-3-hydroxy-propylcarbamoyl)-2-(3-chloro-4-fluorobenzyl)-2H-pyrazole-3-carboxylic acid

15. 5-((*R*)-1-Biphenyl-4-ylmethyl-3-carboxy-3-hydroxy-propylcarbamoyl)-2-(4-chlorobenzyl)-2H-pyrazole-3-carboxylic acid
16. 5-((*R*)-1-Biphenyl-4-ylmethyl-3-carboxy-3-hydroxy-propylcarbamoyl)-2-(3-chloro-2,6-difluorobenzyl)-2H-pyrazole-3-carboxylic acid
- 5 17. 5-((*R*)-1-Biphenyl-4-ylmethyl-3-carboxy-3-hydroxy-propylcarbamoyl)-2-(3-methoxy-benzyl)-2H-pyrazole-3-carboxylic acid
18. 5-((*R*)-1-Biphenyl-4-ylmethyl-3-ethoxycarbonyl-3-hydroxy-propylcarbamoyl)-2H-pyrazole-3-carboxylic acid ethyl ester
19. (*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(5-methyl-1H-pyrazole-3-
- 10 carbonyl)amino]pentanoic acid
20. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(5-pyridin-4-yl-1H-pyrazole-3-carbonyl)-amino]-pentanoic acid
21. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(5-pyridin-3-yl-1H-pyrazole-3-carbonyl)-amino]-pentanoic acid
- 15 22. (*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(5-pyrazin-2-yl-1H-pyrazole-3-carbonyl)amino]-pentanoic acid
23. (*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(1'-methyl-2H,1'H-[3,4']bipyrazolyl-5-carbonyl)amino]-pentanoic acid
24. (*R*)-5-Biphenyl-4-yl-4-{[5-(2-chlorophenyl)-1H-pyrazole-3-carbonyl]amino}-2-
- 20 hydroxy-pentanoic acid
25. (2*R*,4*R*)-5-Biphenyl-4-yl-4-{[5-(3-chlorophenyl)-1H-pyrazole-3-carbonyl]-amino}-2-hydroxy-pentanoic acid
26. (2*R*,4*R*)-5-Biphenyl-4-yl-4-{[5-(2,4-dichlorophenyl)-1H-pyrazole-3-carbonyl]-amino}-2-hydroxy-pentanoic acid
- 25 27. (2*R*,4*R*)-5-Biphenyl-4-yl-4-{[5-(2,5-dichlorophenyl)-1H-pyrazole-3-carbonyl]-amino}-2-hydroxy-pentanoic acid
28. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-{[5-(2-hydroxy-phenyl)-1H-pyrazole-3-carbonyl]-amino}-pentanoic acid
29. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-{[5-(2-hydroxy-phenyl)-1H-pyrazole-3-
- 30 carbonyl]-amino}-pentanoic acid ethyl ester
30. (2*R*,4*R*)-5-Biphenyl-4-yl-4-{[1-(2,3-dihydroxy-propyl)-5-(2-hydroxy-phenyl)-1H-pyrazole-3-carbonyl]-amino}-2-hydroxy-pentanoic acid










31. (R)-4- {[5-(3-Acetylamino-phenyl)-1H-pyrazole-3-carbonyl]amino}-5-biphenyl-4-yl-2-hydroxypentanoic acid
32. (2R,4R)-5-Biphenyl-4-yl-2-hydroxy-4- {[5-(2-methoxy-benzoylamino)-1H-pyrazole-3-carbonyl]-amino}-pentanoic acid
- 5 33. (2R,4R)-5-Biphenyl-4-yl-2-hydroxy-4-[(5-phenyl-1H-pyrazole-3-carbonyl)-amino]-pentanoic acid
34. (R)-5-Biphenyl-4-yl-2-hydroxy-4-[(1-methyl-5-phenyl-1H-pyrazole-3-carbonyl)amino]-pentanoic acid
35. (2R,4R)-5-Biphenyl-4-yl-4-[(1-carboxymethyl-5-phenyl-1H-pyrazole-3-carbonyl)-amino]-2-hydroxy-pentanoic acid
- 10 36. (2R,4R)-5-Biphenyl-4-yl-4- {[1-(2-carboxy-ethyl)-5-phenyl-1H-pyrazole-3-carbonyl]-amino}-2-hydroxy-pentanoic acid
37. (R)-5-Biphenyl-4-yl-4-[(5-furan-2-yl-1H-pyrazole-3-carbonyl)amino]-2-hydroxy-pentanoic acid
- 15 38. (R)-5-Biphenyl-4-yl-2-hydroxy-4-[(5-naphthalen-2-yl-1H-pyrazole-3-carbonyl)amino]-pentanoic acid
39. (R)-5-Biphenyl-4-yl-4-[(5-biphenyl-4-yl-1H-pyrazole-3-carbonyl)amino]-2-hydroxy-pentanoic acid
40. (R)-5-Biphenyl-4-yl-4- {[5-(4-chlorophenyl)-1H-pyrazole-3-carbonyl]amino}-2-hydroxy-pentanoic acid
- 20 41. (2R,4R)-5-Biphenyl-4-yl-2-hydroxy-4- {[5-(1-hydroxy-1-methylethyl)-1H-pyrazole-3-carbonyl]amino}pentanoic acid 5-methyl-2-oxo-[1,3]dioxol-4-ylmethyl ester



Ex.	R ¹	R ³	R ⁴	Formula	MS m/z: [M+H] ⁺	
					calcd	found
42	-OH		H	C ₂₆ H ₂₈ N ₄ O ₆	493.20	493.2
43	-OCH ₂ -CH ₃		H	C ₂₉ H ₃₃ N ₅ O ₆	548.24	548.2
44	-OH		H	C ₂₇ H ₂₉ N ₅ O ₆	520.21	520.2

Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
45	-OCH ₂ -CH ₃		H	C ₂₉ H ₃₃ N ₅ O ₆	548.24	548.2
46	-OH		H	C ₂₇ H ₂₉ N ₅ O ₆	520.21	520.2
47	-OCH ₂ -CH ₃		H	C ₂₉ H ₃₄ N ₄ O ₆	535.25	535.2
48	-OH		H	C ₂₇ H ₃₀ N ₄ O ₆	507.22	507.2
49	-OCH ₂ -CH ₃		H	C ₂₉ H ₃₄ N ₄ O ₆	535.25	535.2
50	-OH		H	C ₂₇ H ₃₀ N ₄ O ₆	507.22	507.2
51	-OCH ₂ -CH ₃		H	C ₂₈ H ₃₂ N ₄ O ₆	521.23	521.4
52	-OCH ₂ -CH ₃	-C(O)NH-cyclopropyl	H	C ₂₇ H ₃₀ N ₄ O ₅	491.22	491.2
53	-OH		H	C ₂₆ H ₂₈ N ₄ O ₆	493.20	493.2
54	-OCH ₂ -CH ₃	-C(O)NH-cyclopropyl	H	C ₂₇ H ₃₀ N ₄ O ₅	491.22	491.2
55	-OH	-C(O)NH-cyclopropyl	H	C ₂₅ H ₂₆ N ₄ O ₅	463.19	463.2
56	-OH	-C(O)NH-CH ₂ COOH	H	C ₂₄ H ₂₄ N ₄ O ₇	481.16	481.2
57	-OH	-C(O)NH-cyclopropyl	H	C ₂₅ H ₂₆ N ₄ O ₅	463.19	463.2
58	-OCH ₂ -CH ₃	-C(O)NH ₂	H	C ₂₄ H ₂₆ N ₄ O ₅	451.19	451.2
59	-OH	-C(O)NH ₂	H	C ₂₂ H ₂₂ N ₄ O ₅	423.16	424.0
60	-OH	-C(O)-NHCH ₃	H	C ₂₃ H ₂₄ N ₄ O ₅	437.17	437.4
61	-OH	-C(O)NH-(CH ₂) ₂ CH ₃	H	C ₂₅ H ₂₈ N ₄ O ₅	465.21	465.2
62	-OCH ₂ -CH ₃	-C(O)-N(CH ₃) ₂	H	C ₂₆ H ₃₀ N ₄ O ₅	479.22	479.2
63	-OH	-C(O)-N(CH ₃) ₂	H	C ₂₄ H ₂₆ N ₄ O ₅	451.19	451.2
64	-OCH ₂ -CH ₃	-C(O)-N(CH ₃)-CH ₂ -CH(CH ₃) ₂	H	C ₂₉ H ₃₆ N ₄ O ₅	521.27	521.2
65	-OH	-C(O)-N(CH ₃)-CH ₂ -CH(CH ₃) ₂	H	C ₂₇ H ₃₂ N ₄ O ₅	493.24	493.2

Ex.	R ¹	R ³	R ⁴	Formula	MS m/z: [M+H] ⁺	
					calcd	found
66	-OCH ₂ -CH ₃	-C(O)-N(CH ₃)-CH ₂ -CH(CH ₃) ₂	H	C ₂₉ H ₃₆ N ₄ O ₅	521.27	521.2
67	-OH	-C(O)-N(CH ₃)-CH ₂ -CH(CH ₃) ₂	H	C ₂₇ H ₃₂ N ₄ O ₅	493.24	493.4
68	-OCH ₂ -CH ₃	-C(O)NH-(CH ₂) ₂ -imidazole	H	C ₂₉ H ₃₂ N ₆ O ₅	545.24	545.2
69	-OH	-C(O)NH-(CH ₂) ₂ -imidazole	H	C ₂₇ H ₂₈ N ₆ O ₅	517.21	517.2
70	-OH	-C(O)-NH-(CH ₂) ₂ -OH	H	C ₂₄ H ₂₆ N ₄ O ₆	467.19	467.2
71	-OH	-C(O)NH-(CH ₂) ₂ -N(CH ₃) ₂	H	C ₂₆ H ₃₁ N ₅ O ₅	494.23	494.2
72	-OH	cyclopropyl	H	C ₂₄ H ₂₅ N ₃ O ₄	420.18	420.2
73	-OH	-CH(CH ₃) ₂	H	C ₂₄ H ₂₇ N ₃ O ₄	422.20	422.2
74	-OH	-(CH ₂) ₂ -CH ₃	H	C ₂₄ H ₂₇ N ₃ O ₄	422.20	422.2
75	-OH	-(CH ₂) ₃ -CH ₃	H	C ₂₅ H ₂₉ N ₃ O ₄	436.22	436.2
76	-OH	phenyl	H	C ₂₇ H ₂₅ N ₃ O ₄	456.18	456.2
77	-OH	phenyl	-(CH ₂) ₂ -COOH	C ₃₀ H ₂₉ N ₃ O ₆	528.21	528.4
78	-OH	2-hydroxy-phenyl	-CH ₂ -CHOH-CH ₂ OH	C ₃₀ H ₃₁ N ₃ O ₇	546.22	546.2
79	-OH	2-methoxy-phenyl	H	C ₂₈ H ₂₇ N ₃ O ₅	486.20	486.6
80	-OH	2,5-dimethoxy-phenyl	H	C ₂₉ H ₂₉ N ₃ O ₆	516.21	516.2
81	-OH	2-CF ₃ -phenyl	H	C ₂₈ H ₂₄ F ₃ N ₃ O ₄	524.17	524.2
82	-OH	4-F-phenyl	H	C ₂₇ H ₂₄ FN ₃ O ₄	474.18	474.2
83	-OH	4-Cl-phenyl	H	C ₂₇ H ₂₄ ClN ₃ O ₄	490.15	490.4
84	-OH	2-Cl-phenyl	H	C ₂₇ H ₂₄ ClN ₃ O ₄	490.15	490.4
85	-OCH ₂ -CH ₃	2-Cl-phenyl	H	C ₂₉ H ₂₈ ClN ₃ O ₄	518.18	518.4
86	-OH	2-methoxy,5-fluorophenyl	H	C ₂₈ H ₂₆ FN ₃ O ₅	504.19	504.2
87	-OCH-(CH ₃) ₂	2-methoxy,5-fluorophenyl	H	C ₃₁ H ₃₂ FN ₃ O ₅	546.23	546.2
88	-OCH ₂ -CH ₃	2-methoxy,5-fluoro-phenyl	H	C ₃₀ H ₃₀ FN ₃ O ₅	532.22	532.4
89	-OH	2-methyl-thiophene	H	C ₂₆ H ₂₅ N ₃ O ₄ S	476.16	476.0
90	-OH	2-chloro-thiophene	H	C ₂₅ H ₂₂ ClN ₃ O ₄ S	496.10	497.0
91	-OH	-NHC(O)-CH ₂ CH ₃	H	C ₂₄ H ₂₆ N ₄ O ₅	451.19	451.2
92	-OH	-NHC(O)-(CH ₂) ₃ CH ₃	H	C ₂₆ H ₃₀ N ₄ O ₅	479.22	479.2
93	-OH	-NHC(O)-O-CH ₂ CH ₃	H	C ₂₄ H ₂₆ N ₄ O ₆	467.19	467.2

Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
94	-OH	-NHC(O)-CH ₂ -OCH ₃	H	C ₂₄ H ₂₆ N ₄ O ₆	467.19	467.2
95	-OH	-NHC(O)-2-methoxy-phenyl	H	C ₂₉ H ₂₈ N ₄ O ₆	529.20	529.2
96	-OH		H	C ₂₈ H ₂₅ ClN ₄ O ₅	533.15	533.2
97	-OCH ₂ -CH ₃		H	C ₃₀ H ₂₉ ClN ₄ O ₅	561.18	561.2
98	-OCH ₂ -CH-(CH ₃) ₂		H	C ₃₂ H ₃₃ ClN ₄ O ₅	589.21	589.2
99	-O-CH-(CH ₃) ₂		H	C ₃₁ H ₃₁ ClN ₄ O ₅	575.20	575.2
100	-OH	-NHC(O)-2-pyridine	H	C ₂₇ H ₂₅ N ₅ O ₅	500.19	500.2
101	-OH	-NH ₂	H	C ₂₁ H ₂₂ N ₄ O ₄	395.16	395.0
102	-OH		H	C ₂₆ H ₂₄ N ₄ O ₄	457.18	457.6
103	-OCH ₂ -CH ₃		H	C ₂₈ H ₂₈ N ₄ O ₄	485.21	485.6
104	-OCH ₂ -CH-(CH ₃) ₂		H	C ₃₀ H ₃₂ N ₄ O ₄	513.24	513.6
105			H	C ₃₁ H ₂₈ N ₄ O ₇	569.20	569.4
106	-OH	pyrazine	H	C ₂₅ H ₂₃ N ₅ O ₄	458.18	458.2

Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
107	-O-CH-(CH ₃)O C(O)O-cyclohexyl	-C(O)CH ₃	H	C ₃₂ H ₃₇ N ₃ O ₈	592.26	592.6
108	-NH ₂	-C(O)CH ₃	H	C ₂₃ H ₂₄ N ₄ O ₄	421.18	421.2
109	-OH	-CH(OH)-CH ₃	H	C ₂₃ H ₂₅ N ₃ O ₅	424.18	424.2
110	-OH	-CH(OH)-CH ₃	H	C ₂₃ H ₂₅ N ₃ O ₅	424.18	424.4
111	-OH	-C(CH ₃) =N(OH)	H	C ₂₃ H ₂₄ N ₄ O ₅	437.17	437.4
112	-OH	-(CH ₂) ₃ -CH ₃	H	C ₂₅ H ₂₉ N ₃ O ₄	436.22	436.4


42. (*R*)-5-Biphenyl-4-yl-2-hydroxy-4-{{5-(3-hydroxypyrrolidine-1-carbonyl)-2H-pyrazole-3-carbonyl]-amino}-pentanoic acid
43. (*2S,4R*)-5-Biphenyl-4-yl-4-{{5-(3-carbamoylpyrrolidine-1-carbonyl)-2H-pyrazole-3-carbonyl]-amino}-2-hydroxy-pentanoic acid ethyl ester
44. (*2S,4R*)-5-Biphenyl-4-yl-4-{{5-(3-carbamoylpyrrolidine-1-carbonyl)-2H-pyrazole-3-carbonyl]-amino}-2-hydroxy-pentanoic acid
45. (*2R,4R*)-5-Biphenyl-4-yl-4-{{5-(3-carbamoylpyrrolidine-1-carbonyl)-2H-pyrazole-3-carbonyl]-amino}-2-hydroxy-pentanoic acid ethyl ester
46. (*2R,4R*)-5-Biphenyl-4-yl-4-{{5-(3-carbamoylpyrrolidine-1-carbonyl)-2H-pyrazole-3-carbonyl]-amino}-2-hydroxy-pentanoic acid
47. (*2R,4R*)-5-Biphenyl-4-yl-2-hydroxy-4-{{5-(4-hydroxypiperidine-1-carbonyl)-2H-pyrazole-3-carbonyl]-amino}-pentanoic acid ethyl ester
48. (*2R,4R*)-5-Biphenyl-4-yl-2-hydroxy-4-{{5-(4-hydroxypiperidine-1-carbonyl)-2H-pyrazole-3-carbonyl]-amino}-pentanoic acid
49. (*2S,4R*)-5-Biphenyl-4-yl-2-hydroxy-4-{{5-(4-hydroxypiperidine-1-carbonyl)-2H-pyrazole-3-carbonyl]amino}-pentanoic acid ethyl ester
50. (*2S,4R*)-5-Biphenyl-4-yl-2-hydroxy-4-{{5-(4-hydroxypiperidine-1-carbonyl)-2H-pyrazole-3-carbonyl]-amino}-pentanoic acid
51. (*R*)-5-Biphenyl-4-yl-2-hydroxy-4-{{5-(morpholine-4-carbonyl)-2H-pyrazole-3-carbonyl]-amino}-pentanoic acid ethyl ester
52. (*2S,4R*)-5-Biphenyl-4-yl-4-[(5-cyclopropylcarbamoyl-2H-pyrazole-3-carbonyl)-amino]-2-hydroxypentanoic acid ethyl ester
53. (*R*)-5-Biphenyl-4-yl-2-hydroxy-4-{{5-(morpholine-4-carbonyl)-2H-pyrazole-3-carbonyl]amino}pentanoic acid

54. (2*R*,4*R*)-5-Biphenyl-4-yl-4-[(5-cyclopropylcarbamoyl-2H-pyrazole-3-carbonyl)-amino]-2-hydroxypentanoic acid ethyl ester
55. (2*S*,4*R*)-5-Biphenyl-4-yl-4-[(5-cyclopropylcarbamoyl-2H-pyrazole-3-carbonyl)-amino]-2-hydroxypentanoic acid
- 5 56. (*R*)-5-Biphenyl-4-yl-4-{[5-(carboxymethyl-carbamoyl)-2H-pyrazole-3-carbonyl]-amino}-2-hydroxy-pentanoic acid
57. (2*R*,4*R*)-5-Biphenyl-4-yl-4-[(5-cyclopropylcarbamoyl-2H-pyrazole-3-carbonyl)-amino]-2-hydroxypentanoic acid
58. (*R*)-5-Biphenyl-4-yl-4-[(5-carbamoyl-2H-pyrazole-3-carbonyl)-amino]-2-hydroxy-
10 pentanoic acid ethyl ester
59. (*R*)-5-Biphenyl-4-yl-4-[(5-carbamoyl-2H-pyrazole-3-carbonyl)-amino]-2-hydroxy-pentanoic acid
60. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(5-methylcarbamoyl-2H-pyrazole-3-carbonyl)-amino]pentanoic acid
- 15 61. (*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(5-propylcarbamoyl-2H-pyrazole-3-carbonyl)-amino]pentanoic acid
62. (2*R*,4*R*)-5-Biphenyl-4-yl-4-[(5-dimethylcarbamoyl-2H-pyrazole-3-carbonyl)-amino]-2-hydroxy-pentanoic acid ethyl ester
63. (2*R*,4*R*)-5-Biphenyl-4-yl-4-[(5-dimethylcarbamoyl-2H-pyrazole-3-carbonyl)-
20 amino]-2-hydroxy-pentanoic acid
64. (2*S*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-{[5-(isobutyl-methyl-carbamoyl)-2H-pyrazole-3-carbonyl]-amino}-pentanoic acid ethyl ester
65. (2*S*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-{[5-(isobutylmethyl-carbamoyl)-2H-pyrazole-3-carbonyl]amino}-pentanoic acid
- 25 66. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-{[5-(isobutylmethyl-carbamoyl)-2H-pyrazole-3-carbonyl]amino}-pentanoic acid ethyl ester
67. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-{[5-(isobutylmethyl-carbamoyl)-2H-pyrazole-3-carbonyl]amino}-pentanoic acid
68. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-({5-[2-(1H-imidazol-4-yl)-ethylcarbamoyl]-
30 2H-pyrazole-3-carbonyl}amino)pentanoic acid ethyl ester
69. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-({5-[2-(3H-imidazol-4-yl)-ethylcarbamoyl]-2H-pyrazole-3-carbonyl}amino)pentanoic acid

70. (R)-5-Biphenyl-4-yl-2-hydroxy-4- {[5-(2-hydroxy-ethylcarbamoyl)-2H-pyrazole-3-carbonyl]-amino} -pentanoic acid
71. (R)-5-Biphenyl-4-yl-4- {[5-(2-dimethylamino-ethylcarbamoyl)-2H-pyrazole-3-carbonyl]-amino} -2-hydroxy-pentanoic acid
- 5 72. (R)-5-Biphenyl-4-yl-4- [(5-cyclopropyl-2H-pyrazole-3-carbonyl)-amino]-2-hydroxy-pentanoic acid
73. (R)-5-Biphenyl-4-yl-2-hydroxy-4- [(5-isopropyl-2H-pyrazole-3-carbonyl)amino]-pentanoic acid
74. (R)-5-Biphenyl-4-yl-2-hydroxy-4- [(5-propyl-2H-pyrazole-3-carbonyl)amino]-
10 pentanoic acid
75. (R)-5-Biphenyl-4-yl-4- [(5-butyl-2H-pyrazole-3-carbonyl)-amino]-2-hydroxy-pentanoic acid
76. (R)-5-Biphenyl-4-yl-2-hydroxy-4- [(5-phenyl-2H-pyrazole-3-carbonyl)-amino]-pentanoic acid
- 15 77. (2R,4R)-5-Biphenyl-4-yl-4- {[2-(2-carboxy-ethyl)-5-phenyl-2H-pyrazole-3-carbonyl]-amino} -2-hydroxy-pentanoic acid
78. (2R,4R)-5-Biphenyl-4-yl-4- {[2-(2,3-dihydroxy-propyl)-5-(2-hydroxy-phenyl)-2H-pyrazole-3-carbonyl]-amino} -2-hydroxy-pentanoic acid
79. (2R,4R)-5-Biphenyl-4-yl-2-hydroxy-4- {[5-(2-methoxy-phenyl)-2H-pyrazole-3-
20 carbonyl]-amino} -pentanoic acid
80. (2R,4R)-5-Biphenyl-4-yl-4- {[5-(2,5-dimethoxy-phenyl)-2H-pyrazole-3-carbonyl]-amino} -2-hydroxy-pentanoic acid
81. (R)-5-Biphenyl-4-yl-2-hydroxy-4- {[5-(2-trifluoromethyl-phenyl)-2H-pyrazole-3-carbonyl]-amino} -pentanoic acid
- 25 82. (R)-5-Biphenyl-4-yl-4- {[5-(4-fluorophenyl)-2H-pyrazole-3-carbonyl]amino} -2-hydroxy-pentanoic acid
83. (2R,4R)-5-Biphenyl-4-yl-4- {[5-(4-chlorophenyl)-2H-pyrazole-3-carbonyl]-
84. (2R,4R)-5-Biphenyl-4-yl-4- {[5-(2-chlorophenyl)-2H-pyrazole-3-carbonyl]amino} -2-hydroxy-pentanoic acid
- 30 85. (2R,4R)-5-Biphenyl-4-yl-4- {[5-(2-chlorophenyl)-2H-pyrazole-3-carbonyl]-amino} -2-hydroxy-pentanoic acid ethyl ester
86. (2R,4R)-5-Biphenyl-4-yl-4- {[5-(5-fluoro-2-methoxy-phenyl)-2H-pyrazole-3-carbonyl]-amino} -2-hydroxy-pentanoic acid

87. (2*R*,4*R*)-5-Biphenyl-4-yl-4-[[5-(5-fluoro-2-methoxy-phenyl)-2H-pyrazole-3-carbonyl]-amino}-2-hydroxy-pentanoic acid isopropyl ester (HCL salt)
88. (2*R*,4*R*)-5-Biphenyl-4-yl-4-[[5-(5-fluoro-2-methoxy-phenyl)-2H-pyrazole-3-carbonyl]-amino}-2-hydroxy-pentanoic acid ethyl ester
- 5 89. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[[5-(5-methylthiophen-2-yl)-2H-pyrazole-3-carbonyl]-amino}-pentanoic acid
90. (2*R*,4*R*)-5-Biphenyl-4-yl-4-[[5-(5-chlorothiophen-2-yl)-2H-pyrazole-3-carbonyl]-amino}-2-hydroxy-pentanoic acid
91. (*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(5-propionylamino-2H-pyrazole-3-carbonyl)-amino]-pentanoic acid
- 10 92. (*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(5-pentanoylamino-2H-pyrazole-3-carbonyl)-amino]-pentanoic acid
93. (*R*)-5-Biphenyl-4-yl-4-[(5-ethoxycarbonylamino-2H-pyrazole-3-carbonyl)-amino]-2-hydroxy-pentanoic acid
- 15 94. (*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[[5-(2-methoxy-acetylamino)-2H-pyrazole-3-carbonyl]-amino}-pentanoic acid
95. (*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[[5-(2-methoxy-benzoylamino)-2H-pyrazole-3-carbonyl]-amino}-pentanoic acid
96. (2*R*,4*R*)-5-Biphenyl-4-yl-4-[[5-(2-chlorobenzoylamino)-2H-pyrazole-3-carbonyl]-amino}-2-hydroxy-pentanoic acid
- 20 97. (2*R*,4*R*)-5-Biphenyl-4-yl-4-[[5-(2-chlorobenzoylamino)-2H-pyrazole-3-carbonyl]-amino}-2-hydroxy-pentanoic acid ethyl ester
98. (2*R*,4*R*)-5-Biphenyl-4-yl-4-[[5-(2-chlorobenzoylamino)-2H-pyrazole-3-carbonyl]-amino}-2-hydroxy-pentanoic acid isobutyl ester
- 25 99. (2*R*,4*R*)-5-Biphenyl-4-yl-4-[[5-(2-chlorobenzoylamino)-2H-pyrazole-3-carbonyl]-amino}-2-hydroxy-pentanoic acid isopropyl ester
100. (*R*)-5-Biphenyl-4-yl-2-hydroxy-4-({5-[(pyridine-2-carbonyl)-amino]-2H-pyrazole-3-carbonyl}-amino)-pentanoic acid
101. (*R*)-4-[(5-Amino-2H-pyrazole-3-carbonyl)-amino]-5-biphenyl-4-yl-2-hydroxy-pentanoic acid
- 30 102. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(5-pyridin-2-yl-2H-pyrazole-3-carbonyl)-amino]-pentanoic acid
103. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(5-pyridin-2-yl-2H-pyrazole-3-carbonyl)-

- amino]-pentanoic acid ethyl ester
104. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(5-pyridin-2-yl-2*H*-pyrazole-3-carbonyl)-amino]-pentanoic acid isobutyl ester
105. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(5-pyridin-2-yl-2*H*-pyrazole-3-carbonyl)-amino]-pentanoic acid 5-methyl-2-oxo-[1,3]dioxol-4-ylmethyl ester
106. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(5-pyrazin-2-yl-2*H*-pyrazole-3-carbonyl)-amino]pentanoic acid
107. (2*R*,4*R*)-4-[(5-Acetyl-2*H*-pyrazole-3-carbonyl)amino]-5-biphenyl-4-yl-2-hydroxy-pentanoic acid 1-cyclohexyloxycarbonyloxyethyl ester
108. 5-Acetyl-2*H*-pyrazole-3-carboxylic acid ((1*R*,3*R*)-1-biphenyl-4-ylmethyl-3-carbamoyl-3-hydroxypropyl)amide
109. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-{[5-((*S*)-1-hydroxyethyl)-2*H*-pyrazole-3-carbonyl]amino}pentanoic acid
110. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-{[5-((*R*)-1-hydroxyethyl)-2*H*-pyrazole-3-carbonyl]amino}pentanoic acid
111. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(5-{1-[(*E*)-hydroxyimino]-ethyl}-2*H*-pyrazole-3-carbonyl)-amino]-pentanoic acid
112. (2*R*,4*R*)-5-Biphenyl-4-yl-4-[(5-butyl-2*H*-pyrazole-3-carbonyl)-amino]-2-hydroxy-pentanoic acid




Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
113	-OH	H	H	C ₂₁ H ₂₁ N ₃ O ₄	380.15	380.2

113. (*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(1*H*-pyrazole-4-carbonyl)amino]pentanoic acid

EXAMPLE 16

Following the procedures described in the examples herein, and substituting the appropriate starting materials and reagents, compounds having the following formula were prepared as TFA salts.




Ex.	-XR ³ R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
			calcd	found
1		C ₂₉ H ₂₇ N ₃ O ₅	498.20	498.8
2		C ₂₉ H ₂₇ N ₃ O ₅	498.20	498.6
3		C ₃₀ H ₂₉ N ₃ O ₅	512.21	512.6
4		C ₃₀ H ₂₉ N ₃ O ₆	528.21	528.4
5		C ₃₀ H ₂₉ N ₃ O ₆	528.21	528.4
6		C ₂₈ H ₂₅ N ₃ O ₅	484.18	484.6

- 5 1. (2*R*,4*R*)-5-Biphenyl-4-yl-4-[(4,5-dihydro-6-oxa-3,3a-diaza-benzo[e]azulene-2-carbonyl)-amino]-2-hydroxy-pentanoic acid

2. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(8-oxa-11,14-diazatricyclo[9.2.1.0^{2,7}]^{2,4,6,12}-pentaene-12-carbonyl)-amino]-pentanoic acid
3. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(10-oxa-5,6-diazatricyclo[9.4.0.0^{2,6}]^{2,4,11,13}-pentaene-4-carbonyl)-amino]-pentanoic acid
- 5 4. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(*S*)-8-hydroxy-10-oxa-5,6-diazatricyclo[9.4.0.0^{2,6}]^{2,4,11,13}-pentaene-4-carbonyl)-amino]-pentanoic acid
5. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(*R*)-8-hydroxy-10-oxa-5,6-diazatricyclo[9.4.0.0^{2,6}]^{2,4,11,13}-pentaene-4-carbonyl)-amino]-pentanoic acid
6. (2*R*,4*R*)-5-Biphenyl-4-yl-2-hydroxy-4-[(5-oxa-3,3a-diazacyclopenta[*a*]naphthalene-2-carbonyl)-amino]-pentanoic acid


Preparation 9

(*S*)-2-(4-Bromobenzyl)-5-oxopyrrolidine-1-carboxylic Acid *t*-Butyl Ester

To a solution of (*R*)-2-amino-3-(4-bromophenyl)propionic acid (50 g, 0.2 mol) in MeCN (700 mL) was added a solution of NaOH (16.4 g, 0.4 mol) in water (700 mL) at -5°C. After stirring for 10 minutes, a solution of (Boc)₂O (44.7 g, 0.2 mol) in MeCN (100 mL) was added. The mixture was warmed to room temperature and stirred overnight. After the evaporation of the MeCN, the residue was diluted with DCM (800 mL) and acidified with 1 M HCl to pH = 2 at -5°C. The aqueous was extracted with DCM (3x200 mL). The combined organic layers were washed with saturated aqueous NaCl (500 mL), dried over Na₂SO₄ and concentrated to yield compound 1 (66.5 g) as a white solid. LC-MS: 366 (M+Na), 709 (2M+Na).

To a solution of compound 1 (66.5 g, 193 μmol), Meldrum's acid (33.4 g, 232 mmol) and DMAP (37.7 g, 309 mmol) in anhydrous DCM (600 mL), was added dropwise a solution of DCC (47.9 g, 232 mmol) in anhydrous DCM (200 mL) over 1 hour at -5°C under nitrogen. The mixture was stirred at -5°C for 8 hours, then refrigerated overnight. Crystals of dicyclohexylurea were observed. The mixture was filtered, washed with 5% KHSO₄ (5x200 mL), saturated aqueous NaCl (200 mL) and dried over anhydrous MgSO₄ under refrigeration overnight. The solution was then evaporated to yield crude compound

2 (91 g) as a light yellow solid. LC-MS: 492(M+Na), 961(2M+Na).




To a solution of crude compound 2 (91 g, 193 mmol) in anhydrous DCM (1 L) was added AcOH (127.5 g, 2.1 mol) at -5°C under nitrogen. The mixture was stirred at -5°C for 30 minutes, then NaBH₄ (18.3 g, 483 mmol) was added in small portions over 1 hour. After stirring for another 1 hour at -5°C, saturated aqueous NaCl (500 mL) was added. The organic layer was washed with saturated aqueous NaCl (2x300 mL) and water (2x300 mL), dried over MgSO₄, filtered, and concentrated to yield the crude product, which was further purified by washing with Et₂O to yield compound 3 (68 g) as a light yellow solid. LC-MS: 478 (M+Na), 933 (2M+Na).

A solution of compound 3 (68 g, 149 mmol) in anhydrous toluene (500 mL) was refluxed under nitrogen for 3 hours. After evaporation of the solvent, the residue was purified by chromatography (hexanes:EtOAc=10:1) to yield the title compound (38 g) as a light yellow oil. LC-MS: 376 (M+Na), 729 (2M+Na).

Preparation 10

(2R,4R)-4-Amino-5-(4-bromophenyl)-2-hydroxypentanoic Acid Ethyl Ester




To a solution of (S)-2-(4-bromobenzyl)-5-oxopyrrolidine-1-carboxylic acid *t*-butyl ester (38 g, 107 mmol) in anhydrous DCM (250 mL) was added TFA (20 mL, 0.27 mol) at -5°C under nitrogen. The mixture was warmed to room temperature and stirred overnight. After evaporation of the solvent, the residue was diluted with EtOAc (300 mL) and washed with saturated aqueous NaHCO₃ (3x200 mL), water (200 mL), saturated aqueous NaCl (250 mL), dried over Na₂SO₄ and concentrated to yield crude compound 1 (24 g) as a light yellow solid. LC-MS: 254 (M+H).

To a solution of NaH (8.6 g, 250 mmol) in anhydrous THF (200 mL) was added

dropwise a solution of compound 1 (24 g, 94 mmol) in anhydrous THF (200 mL) over 30 minutes at 0°C under nitrogen. The mixture was warmed to room temperature and stirred for 2 hours. After cooling to 0°C, pivaloyl chloride (18 g, 150 mmol) was added dropwise over 30 minutes. The mixture was warmed to room temperature and stirred overnight.

- 5 The reaction was quenched with saturated aqueous NH₄Cl (300 mL) and extracted with EtOAc (3x200 mL). The combined organic layers were washed with saturated aqueous NaCl (300 mL), dried over MgSO₄, filtered and concentrated to yield the crude product, which was further purified by chromatography (hexanes:EtOAc=25:1) to yield compound 2 (18 g) as a light yellow solid. LC-MS: 360(M+Na).



10

To a solution of compound 2 (18 g, 53 mmol) in anhydrous THF (250 mL) was added dropwise HMDSNa (47.7 mL, 96 mmol) over 30 minutes at -78°C under nitrogen. After stirring at -78°C for 90 minutes, a solution of (+)-(8,8-

- 15 dichlorocamphorylsulfonyl)oxaziridine (31.6 g, 106 mmol) was added dropwise over 30 minutes. After stirring at -78°C for 2 hours, the reaction was quenched with saturated aqueous NH₄Cl (400 mL) and extracted with EtOAc (3x300 mL). The combined organic layers were washed with saturated aqueous NaCl (300 mL), dried over MgSO₄, filtered, and concentrated to give the crude product which was further purified by chromatography (hexanes:EtOAc=15:1) to yield compound 3 (8.9 g) as a light yellow solid. LC-MS:

20 376(M+Na).

A solution of compound 3 (8.9 g, 25 mmol) in concentrated HCl (81 mL, 81 mmol) was heated at 100°C for 16 hours. The mixture was then concentrated to yield the crude product which was further purified by washing with Et₂O to yield compound 4 (7 g) as a light yellow solid HCl salt. LC-MS: 323 (M+H).

- 25 A solution of compound 4 (7 g, 22 mmol) in EtOH (10 mL) was combined with 8M HCl in EtOH (120 mL, 960 mmol) at room temperature. The mixture was heated at 50°C for 16 hours, then concentrated. The crude product was further purified by washing with Et₂O to yield the title compound (6 g) as a light yellow solid HCl salt. LC-MS: 352 (M+H).

Preparation 11


1-Hydroxy-1H-1,2,3-triazole-4-carboxylic Acid

1-Hydroxy-1H-[1,2,3]triazole-4-carboxylic acid ethyl ester (2.0 g, 13 mmol) and
 5 EtOH (25 mL) were combined with a pre-dissolved solution of lithium hydroxide monohydrate (1.6 g, 38.2 mmol) and water (10 mL), and stirred at room temperature for 4 hours. The mixture was partially concentrated and acidified with HCl to cause precipitation. The solid was filtered and dried under vacuum to yield the title compound (1.3 g).

10

Preparation 12

(2R,4R)-5-(4-Bromo-phenyl)-2-hydroxy-4-[(1-hydroxy-1H-1,2,3-triazole-4-carbonyl)-amino]-pentanoic Acid Ethyl Ester




1-Hydroxy-1H-1,2,3-triazole-4-carboxylic acid (163 mg, 1.3 mmol), HCTU (523
 15 mg, 1.3 mmol), and DMF were combined and stirred for 5 minutes at room temperature. DIPEA (661 μ L, 3.8 mmol) and (2R,4R)-4-amino-5-(4-bromophenyl)-2-hydroxypentanoic acid ethyl ester (4 mg, 1.3 mmol) were added and the resulting mixture was stirred for 10 minutes. The reaction mixture was evaporated under reduced pressure and purified (C18 column; 20-70% MeCN in water with 0.05% TFA) to yield the title compound (330 mg).

20


EXAMPLE 17

A. (2R,4R)-5-(3'-Chlorobiphenyl-4-yl)-2-hydroxy-4-[(1-hydroxy-1H-[1,2,3]triazole-4-carbonyl)-amino]pentanoic Acid



1-Hydroxy-1H-1,2,3-triazole-4-carboxylic acid (17.6 mg, 136 μ mol) and HCTU (56.4 mg, 136 μ mol) were combined in DMF and stirred for 5 minutes at room temperature. DIPEA (71.2 μ l, 409 μ mol) and (2*R*,4*R*)-4-amino-5-(4-bromophenyl)-2-hydroxypentanoic acid ethyl ester (43 mg, 140 μ mol) were added and the resulting mixture was stirred for 10 minutes. The reaction mixture was evaporated under reduced pressure and purified (C18 reverse phase column). The purified material was combined with 3-chlorophenylboronic acid, pinacol ester (48.1 mg, 202 μ mol), K_2CO_3 (41.8 mg, 302 μ mol), EtOH (1 ml) and water (0.3 ml). Oxygen was removed (high vacuum) and SilicaCat[®]Pd(0) (0.09 mmol/g loading; 112 mg, 10.1 μ mol) was quickly added under nitrogen. The mixture was microwaved at 100°C for 20 minutes, then concentrated. The material was redissolved in AcOH and purified by preparative HPLC to yield the title compound (12 mg; purity 95%). MS m/z $[M+H]^+$ calc'd for $C_{20}H_{19}ClN_4O_5$, 431.10; found 431.4.

B. (2*R*,4*R*)-5-(3'-Chlorobiphenyl-4-yl)-2-hydroxy-4-[(1-hydroxy-1H-[1,2,3]triazole-4-carbonyl)-amino]pentanoic Acid Ethyl Ester





(2*R*,4*R*)-5-(4-Bromo-phenyl)-2-hydroxy-4-[(1-hydroxy-1H-1,2,3-triazole-4-carbonyl)-amino]-pentanoic acid ethyl ester (120 mg, 281 μ mol) and 3-chlorophenylboronic acid, pinacol ester (120 mg, 506 μ mol) were combined with K_2CO_3 (116 mg, 842 μ mol), EtOH (3 mL), and water (0.8 mL). SilicaCat[®]DPP-Pd (280 μ mol/g loading; 1 mg, 28 μ mol) was then added and the mixture was heated at 90°C and the reaction monitored by LC/MS. The reaction was stopped after 2 hours, and the mixture was filtered, concentrated and purified (C18 reverse phase column; 30-70% MeCN in

water with 0.05% TFA). EtOH (5.0 mL, 86 mmol) and 4 M HCl in dioxane (1.5 mL, 6.0 mmol) was added and the resulting mixture was stirred at room temperature for 40 minutes, then concentrated and purified (C18 reverse phase column; 30-90% MeCN in water with 0.05% TFA) to yield the title compound (20 mg; purity 95%). MS m/z $[M+H]^+$ calc'd for $C_{22}H_{23}ClN_4O_5$, 459.14; found 459.4.


EXAMPLE 18

Following the procedures described in the examples herein, and substituting the appropriate starting materials and reagents, compounds having the following formula were prepared as the parent compound or as a TFA salt.




Ex.	R ¹	R ³	R ⁴	Formula	MS m/z : $[M+H]^+$	
					calcd	found
1	-OCH ₂ CH ₃	H	H	C ₂₂ H ₂₃ ClN ₄ O ₄	443.14	443.4

1. (2*R*,4*R*)-5-(3'-Chlorobiphenyl-4-yl)-2-hydroxy-4-[(3*H*-[1,2,3]triazole-4-carbonyl)-amino]-pentanoic acid ethyl ester (TFA salt)



Ex.	R ¹	R ³	R ⁴	Formula	MS m/z : $[M+H]^+$	
					calcd	found
2	-OH	H	H	C ₂₀ H ₁₉ ClN ₄ O ₄	415.11	415.4


2. (2*R*,4*R*)-5-(3'-Chlorobiphenyl-4-yl)-2-hydroxy-4-[(1*H*-[1,2,3]triazole-4-carbonyl)-amino]-pentanoic acid (TFA salt)



Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
3	-OH	H	-OH	C ₂₄ H ₂₁ ClN ₄ O ₅	481.12	481.2
4	-OCH ₂ CH ₃	H	-OH	C ₂₆ H ₂₅ ClN ₄ O ₅	509.15	509.2

3. (2*R*,4*R*)-5-(3'-Chlorobiphenyl-4-yl)-2-hydroxy-4-[(3-hydroxy-3*H*-benzotriazole-5-carbonyl)-amino]-pentanoic acid


5 4. (2*R*,4*R*)-5-(3'-Chlorobiphenyl-4-yl)-2-hydroxy-4-[(3-hydroxy-3*H*-benzotriazole-5-carbonyl)-amino]-pentanoic acid ethyl ester



Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
5	-OCH ₂ CH(CH ₃) ₂	H	-OH	C ₂₈ H ₂₉ ClN ₄ O ₅	537.18	537.2
6	-OCH(CH ₃) ₂	H	-OH	C ₂₇ H ₂₇ ClN ₄ O ₅	523.17	523.2


5. (2*R*,4*R*)-5-(3'-Chlorobiphenyl-4-yl)-2-hydroxy-4-[(3-hydroxy-3*H*-benzotriazole-5-carbonyl)-amino]-pentanoic acid isobutyl ester

6. (2*R*,4*R*)-5-(3'-Chlorobiphenyl-4-yl)-2-hydroxy-4-[(3-hydroxy-3*H*-benzotriazole-5-carbonyl)-amino]-pentanoic acid isopropyl ester




Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
7	-OH	-C(O)CH ₃	H	C ₂₃ H ₂₁ ClN ₂ O ₆	457.11	457.2
8	-OH	-OCH ₃	H	C ₂₂ H ₂₁ ClN ₂ O ₆	445.11	445.2
9	-OH	Cl	H	C ₂₁ H ₁₈ Cl ₂ N ₂ O ₅	449.06	450.3
10	-OH	-CH(CH ₃) ₂	H	C ₂₄ H ₂₅ ClN ₂ O ₅	457.15	457.4
11	-OH	-(CH ₂) ₂ CH ₃	H	C ₂₄ H ₂₅ ClN ₂ O ₅	457.15	457.4
12	-OH	-CH ₂ -CH(CH ₃) ₂	H	C ₂₅ H ₂₇ ClN ₂ O ₅	471.16	471.4
13	-OH	-C(OH)(CH ₃) ₂	H	C ₂₄ H ₂₅ ClN ₂ O ₆	473.14	472.8
14	-OH	-C(CH ₃) ₃	H	C ₂₅ H ₂₇ ClN ₂ O ₅	471.16	470.8

7. (2*R*,4*R*)-4-[(3-Acetyl-isoxazole-5-carbonyl)-amino]-5-(3'-chlorobiphenyl-4-yl)-2-hydroxy-pentanoic acid
8. (2*R*,4*R*)-5-(3'-Chlorobiphenyl-4-yl)-2-hydroxy-4-[(3-methoxy-isoxazole-5-carbonyl)-amino]-pentanoic acid
9. (2*R*,4*R*)-5-(3'-Chloro-biphenyl-4-yl)-4-[(3-chloro-isoxazole-5-carbonyl)-amino]-2-hydroxy-pentanoic acid
10. (2*R*,4*R*)-5-(3'-Chloro-biphenyl-4-yl)-2-hydroxy-4-[(3-isopropyl-isoxazole-5-carbonyl)-amino]-pentanoic acid
11. (2*R*,4*R*)-5-(3'-Chloro-biphenyl-4-yl)-2-hydroxy-4-[(3-propyl-isoxazole-5-carbonyl)-amino]-pentanoic acid
12. (2*R*,4*R*)-5-(3'-Chloro-biphenyl-4-yl)-2-hydroxy-4-[(3-isobutyl-isoxazole-5-carbonyl)-amino]-pentanoic acid
13. (2*R*,4*R*)-5-(3'-Chloro-biphenyl-4-yl)-2-hydroxy-4-[[3-(1-hydroxy-1-methyl-ethyl)-isoxazole-5-carbonyl]-amino]-pentanoic acid
14. (2*R*,4*R*)-4-[(3-tert-Butyl-isoxazole-5-carbonyl)-amino]-5-(3'-chloro-biphenyl-4-yl)-2-hydroxy-pentanoic acid

















Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
15	-OH	2-chlorophenyl	H	C ₂₇ H ₂₂ Cl ₂ N ₂ O ₅	525.09	525.4
16	-OH	2-hydroxyphenyl	H	C ₂₇ H ₂₃ ClN ₂ O ₆	507.12	507.2

15. (2*R*,4*R*)-5-(3'-Chlorobiphenyl-4-yl)-4-[[5-(2-chlorophenyl)-isoxazole-3-carbonyl]-amino]-2-hydroxy-pentanoic acid
16. (2*R*,4*R*)-5-(3'-Chlorobiphenyl-4-yl)-2-hydroxy-4-[[5-(2-hydroxyphenyl)-isoxazole-3-carbonyl]-amino]-pentanoic acid




Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found


Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
17	-OH	2-hydroxy-phenyl	H	C ₂₇ H ₂₄ ClN ₃ O ₅	506.14	506.0
18	-OCH ₂ CH ₃	2-hydroxy-phenyl	H	C ₂₉ H ₂₈ ClN ₃ O ₅	534.17	534.2
19	-OH		H	C ₂₅ H ₂₂ ClN ₅ O ₄	492.14	492.2
20	-OCH ₂ CH ₃		H	C ₂₇ H ₂₆ ClN ₅ O ₄	520.17	520.2
21	-OCH ₂ -CH(CH ₃)		H	C ₂₉ H ₃₀ ClN ₅ O ₄	548.20	548.2
22			H	C ₃₀ H ₂₆ ClN ₅ O ₇	604.15	604.1
23	-OH		H	C ₂₆ H ₂₃ ClN ₄ O ₄	491.14	491.2
24	-OCH ₂ CH ₃		H	C ₂₈ H ₂₇ ClN ₄ O ₄	519.17	519.2
25	-OCH ₂ -CH(CH ₃)		H	C ₃₀ H ₃₁ ClN ₄ O ₄	547.20	547.1
26	-OCH(CH ₃)		H	C ₂₉ H ₂₉ ClN ₄ O ₄	533.19	533.2
27	-OH		H	C ₂₈ H ₂₄ Cl ₂ N ₄ O ₅	567.11	567.1


Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
28	-OCH ₂ CH ₃		H	C ₃₀ H ₂₈ Cl ₂ N ₄ O ₅	595.14	595.1
29	-OCH ₂ -CH(CH ₃)		H	C ₃₂ H ₃₂ Cl ₂ N ₄ O ₅	623.18	623.2
30	-OCH(CH ₃)		H	C ₃₁ H ₃₀ Cl ₂ N ₄ O ₅	609.16	609.2
31	-OH		H	C ₂₉ H ₂₇ ClN ₄ O ₆	563.16	563.2
32	-OH	-CH(OH)-CH ₃	H	C ₂₃ H ₂₄ ClN ₃ O ₅	458.14	458.4
33	-OCH ₂ CH ₃			C ₂₈ H ₂₇ ClN ₄ O ₄	519.17	519.4

17. (2*R*,4*R*)-5-(3'-Chlorobiphenyl-4-yl)-2-hydroxy-4-[[5-(2-hydroxy-phenyl)-2H-pyrazole-3-carbonyl]-amino]-pentanoic acid (TFA salt)
18. (2*R*,4*R*)-5-(3'-Chlorobiphenyl-4-yl)-2-hydroxy-4-[[5-(2-hydroxy-phenyl)-2H-pyrazole-3-carbonyl]-amino]-pentanoic acid ethyl ester (TFA salt)
- 5 19. (2*R*,4*R*)-5-(3'-Chlorobiphenyl-4-yl)-2-hydroxy-4-[(5-pyrazin-2-yl-2H-pyrazole-3-carbonyl)-amino]-pentanoic acid (TFA salt)
20. (2*R*,4*R*)-5-(3'-Chloro-biphenyl-4-yl)-2-hydroxy-4-[(5-pyrazin-2-yl-2H-pyrazole-3-carbonyl)-amino]-pentanoic acid ethyl ester
- 10 21. (2*R*,4*R*)-5-(3'-Chloro-biphenyl-4-yl)-2-hydroxy-4-[(5-pyrazin-2-yl-2H-pyrazole-3-carbonyl)-amino]-pentanoic acid isobutyl ester
22. (2*S*,4*S*)-5-(3'-Chloro-biphenyl-4-yl)-2-hydroxy-4-[(5-pyrazin-2-yl-2H-pyrazole-3-


- carbonyl)-amino]-pentanoic acid 5-methyl-2-oxo-[1,3]dioxol-4-ylmethyl ester (TFA salt)
23. (2*R*,4*R*)-5-(3'-Chlorobiphenyl-4-yl)-2-hydroxy-4-[(5-pyridin-2-yl-2H-pyrazole-3-carbonyl)-amino]-pentanoic acid (TFA salt)
24. (2*R*,4*R*)-5-(3'-Chloro-biphenyl-4-yl)-2-hydroxy-4-[(5-pyridin-2-yl-2H-pyrazole-3-carbonyl)-amino]-pentanoic acid ethyl ester
25. (2*R*,4*R*)-5-(3'-Chloro-biphenyl-4-yl)-2-hydroxy-4-[(5-pyridin-2-yl-2H-pyrazole-3-carbonyl)-amino]-pentanoic acid isobutyl ester
26. (2*R*,4*R*)-5-(3'-Chloro-biphenyl-4-yl)-2-hydroxy-4-[(5-pyridin-2-yl-2H-pyrazole-3-carbonyl)-amino]-pentanoic acid isopropyl ester
27. (2*R*,4*R*)-4-{[5-(2-Chlorobenzoylamino)-2H-pyrazole-3-carbonyl]-amino}-5-(3'-chlorobiphenyl-4-yl)-2-hydroxy-pentanoic acid (TFA salt)
28. (2*R*,4*R*)-4-{[5-(2-Chloro-benzoylamino)-2H-pyrazole-3-carbonyl]-amino}-5-(3'-chloro-biphenyl-4-yl)-2-hydroxy-pentanoic acid ethyl ester (TFA salt)
29. (2*R*,4*R*)-4-{[5-(2-Chloro-benzoylamino)-2H-pyrazole-3-carbonyl]-amino}-5-(3'-chloro-biphenyl-4-yl)-2-hydroxy-pentanoic acid isobutyl ester (TFA salt)
30. (2*R*,4*R*)-4-{[5-(2-Chloro-benzoylamino)-2H-pyrazole-3-carbonyl]-amino}-5-(3'-chloro-biphenyl-4-yl)-2-hydroxy-pentanoic acid isopropyl ester (TFA salt)
31. (2*R*,4*R*)-5-(3'-Chloro-biphenyl-4-yl)-2-hydroxy-4-{[5-(2-methoxy-benzoylamino)-2H-pyrazole-3-carbonyl]-amino}-pentanoic acid
32. (2*R*,4*R*)-5-(3'-Chloro-biphenyl-4-yl)-2-hydroxy-4-{[5-(1-hydroxy-ethyl)-2H-pyrazole-3-carbonyl]-amino}-pentanoic acid
33. (2*R*,4*R*)-5-(3'-Chloro-biphenyl-4-yl)-2-hydroxy-4-[(5-pyridin-3-yl-2H-pyrazole-3-carbonyl)-amino]-pentanoic acid ethyl ester (TFA salt)



Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
34	-OH		H	C ₂₆ H ₂₃ ClN ₄ O ₄	491.14	491.2
35	-OH	-OH	-CH ₃	C ₂₂ H ₂₂ ClN ₃ O ₅	444.12	444.4
36	-OCH ₂ - CH(CH ₃)	-OH	-CH ₃	C ₂₆ H ₃₀ ClN ₃ O ₅	500.19	500.0

Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
37	-OH	-OH		C ₂₆ H ₂₃ ClN ₄ O ₅	507.14	506.8
38	-OH	-OH	2-fluoro-phenyl	C ₂₇ H ₂₃ ClFN ₃ O ₅	524.13	523.8
39	-OH	-OH	3-chloro-phenyl	C ₂₇ H ₂₃ Cl ₂ N ₃ O ₅	540.10	539.6
40	-OH	-OCH ₂ -CH ₃	H	C ₂₃ H ₂₄ ClN ₃ O ₅	458.14	458.4

34. (2*R*,4*R*)-5-(3'-Chlorobiphenyl-4-yl)-2-hydroxy-4-[(5-pyridin-3-yl-1*H*-pyrazole-3-carbonyl)-amino]-pentanoic acid (TFA salt)
35. (2*R*,4*R*)-5-(3'-Chloro-biphenyl-4-yl)-2-hydroxy-4-[(5-hydroxy-1-methyl-1*H*-pyrazole-3-carbonyl)-amino]-pentanoic acid
36. (2*R*,4*R*)-5-(3'-Chloro-biphenyl-4-yl)-2-hydroxy-4-[(5-hydroxy-1-methyl-1*H*-pyrazole-3-carbonyl)-amino]-pentanoic acid isobutyl ester
37. (2*R*,4*R*)-5-(3'-Chloro-biphenyl-4-yl)-2-hydroxy-4-[(5-hydroxy-1-pyridin-2-yl-1*H*-pyrazole-3-carbonyl)-amino]-pentanoic acid (TFA salt)
38. (2*R*,4*R*)-5-(3'-Chloro-biphenyl-4-yl)-4-{[1-(2-fluoro-phenyl)-5-hydroxy-1*H*-pyrazole-3-carbonyl]-amino}-2-hydroxy-pentanoic acid
39. (2*R*,4*S*)-5-(3'-Chloro-biphenyl-4-yl)-4-{[1-(3-chloro-phenyl)-5-hydroxy-1*H*-pyrazole-3-carbonyl]-amino}-2-hydroxy-pentanoic acid
40. (2*R*,4*R*)-5-(3'-Chloro-biphenyl-4-yl)-4-[(5-ethoxy-1*H*-pyrazole-3-carbonyl)-amino]-2-hydroxy-pentanoic acid



Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
41	-OH	=O	phenyl	C ₂₇ H ₂₄ ClN ₃ O ₅	506.14	506.2
42	-OH	=O	3-chlorophenyl	C ₂₆ H ₂₂ Cl ₂ N ₄ O ₅	541.10	540.6
43	-OH	=O	2-chlorophenyl	C ₂₆ H ₂₂ Cl ₂ N ₄ O ₅	541.10	540.6
44	-OH	=O	4-fluorophenyl	C ₂₆ H ₂₂ ClFN ₄ O ₅	525.13	524.8


41. (2*R*,4*R*)-5-(3'-Chlorobiphenyl-4-yl)-2-hydroxy-4-[(5-oxo-1-phenyl-2,5-dihydro-1*H*-pyrazole-3-carbonyl)-amino]-pentanoic acid (TFA salt)
42. (2*R*,4*R*)-5-(3'-Chloro-biphenyl-4-yl)-4-{[1-(3-chloro-phenyl)-5-oxo-4,5-dihydro-

1H-[1,2,4]triazole-3-carbonyl]-amino}-2-hydroxy-pentanoic acid

43. (2*R*,4*R*)-5-(3'-Chloro-biphenyl-4-yl)-4-{[1-(2-chloro-phenyl)-5-oxo-4,5-dihydro-1H-[1,2,4]triazole-3-carbonyl]-amino}-2-hydroxy-pentanoic acid


44. (2*R*,4*R*)-5-(3'-Chloro-biphenyl-4-yl)-4-{[1-(4-fluoro-phenyl)-5-oxo-4,5-dihydro-

5 1H-[1,2,4]triazole-3-carbonyl]-amino}-2-hydroxy-pentanoic acid



Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
45	-OH	-CF ₃	H	C ₂₁ H ₁₈ ClF ₃ N ₄ O ₄	483.10	483.0


10 45. (2*R*,4*R*)-5-(3'-Chlorobiphenyl-4-yl)-2-hydroxy-4-[(5-trifluoromethyl-4H-[1,2,4]triazole-3-carbonyl)-amino]-pentanoic acid (TFA salt)



Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
46	-OH	Cl	H	C ₂₀ H ₁₈ Cl ₂ N ₄ O ₄	449.07	449.0


46. (2*R*,4*R*)-5-(3'-Chlorobiphenyl-4-yl)-4-[(5-chloro-1H-[1,2,4]triazole-3-carbonyl)-

15 amino]-2-hydroxy-pentanoic acid (TFA salt)



Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
47	-OH	=O	phenyl	C ₂₆ H ₂₃ ClN ₄ O ₅	507.14	507.2


47. (2*R*,4*R*)-5-(3'-Chlorobiphenyl-4-yl)-2-hydroxy-4-[(5-oxo-1-phenyl-4,5-dihydro-1H-[1,2,4]triazole-3-carbonyl)-amino]-pentanoic acid (TFA salt)



Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
48	-OH	=O	benzyl	C ₂₉ H ₂₆ ClN ₃ O ₅	532.16	532.2
49	-OH	=O	phenyl	C ₂₈ H ₂₄ ClN ₃ O ₅	518.14	518.2

48. (2*R*,4*R*)-4-[(1-Benzyl-6-oxo-1,6-dihydro-pyridazine-3-carbonyl)-amino]-5-(3'-chlorobiphenyl-4-yl)-2-hydroxy-pentanoic acid (TFA salt)

5 49. (2*R*,4*R*)-5-(3'-Chlorobiphenyl-4-yl)-2-hydroxy-4-[(6-oxo-1-phenyl-1,6-dihydro-pyridazine-3-carbonyl)-amino]-pentanoic acid (TFA salt)



Ex.	R ¹	R ³	R ⁴	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
50	-OH	-OH	H	C ₂₂ H ₂₀ ClN ₃ O ₅	442.11	442.0


50. (2*R*,4*R*)-5-(3'-Chlorobiphenyl-4-yl)-2-hydroxy-4-[(6-hydroxy-pyridazine-3-carbonyl)-amino]-pentanoic acid (TFA salt)


10

EXAMPLE 19

Following the procedures described in the examples herein, and substituting the appropriate starting materials and reagents, compounds having the following formula were prepared as the parent compound.

15






Ex.	R ¹	R ³	R ⁴	b	R ⁶	Formula	MS <i>m/z</i> : [M+H] ⁺	
							calcd	found
1	-OH	H	-OH	2	2',5'-diCl	C ₂₀ H ₁₈ Cl ₂ N ₄ O ₅	465.07	465.2
2	-OCH ₂ -CH(CH ₃) ₂	H	-OH	2	2',5'-diCl	C ₂₄ H ₂₆ Cl ₂ N ₄ O ₅	521.13	521.2
3	-OCH-(CH ₃) ₂	H	-OH	2	2',5'-diCl	C ₂₃ H ₂₄ Cl ₂ N ₄ O ₅	507.11	507.2
4	-O-CH ₂ CH ₃	H	-OH	2	2',5'-diCl	C ₂₂ H ₂₂ Cl ₂ N ₄ O ₅	493.10	493.2
5	-O-(CH ₂) ₆ -CH ₃	H	-OH	2	2',5'-diCl	C ₂₇ H ₃₂ Cl ₂ N ₄ O ₅	563.18	563.2
6	-OH	H	-OH	2	2'-F, 5'-Cl	C ₂₀ H ₁₈ ClFN ₄ O ₅	449.10	449.2
7	-OCH ₂ -CH(CH ₃) ₂	H	-OH	2	2'-F, 5'-Cl	C ₂₄ H ₂₆ ClFN ₄ O ₅	505.16	505.2
8	-OCH-(CH ₃) ₂	H	-OH	2	2'-F, 5'-Cl	C ₂₃ H ₂₄ ClFN ₄ O ₅	491.14	491.4
9	-O-CH ₂ CH ₃	H	-OH	2	2'-F, 5'-Cl	C ₂₂ H ₂₂ ClFN ₄ O ₅	477.13	477.0
10	-O-(CH ₂) ₆ -CH ₃	H	-OH	2	2'-F, 5'-Cl	C ₂₇ H ₃₂ ClFN ₄ O ₅	547.20	547.2
11	-OCH-(CH ₃) ₂	H	-O-CH ₂ O-C(O)-CH-(CH ₃) ₂ -NH ₂	2	2'-F, 5'-Cl	C ₂₉ H ₃₅ ClFN ₅ O ₇	620.22	620.0
12	-OH	H	-OH	2	2'-CH ₃ , 5'-Cl	C ₂₁ H ₂₁ ClN ₄ O ₅	445.12	445.4
13	-OCH ₂ -CH(CH ₃) ₂	H	-OH	2	2'-CH ₃ , 5'-Cl	C ₂₅ H ₂₉ ClN ₄ O ₅	501.18	501.4
14	-OCH-(CH ₃) ₂	H	-OH	2	2'-CH ₃ , 5'-Cl	C ₂₄ H ₂₇ ClN ₄ O ₅	487.17	487.4
15	-O-CH ₂ CH ₃	H	-OH	2	2'-CH ₃ , 5'-Cl	C ₂₃ H ₂₅ ClN ₄ O ₅	473.15	473.2
16	-O-(CH ₂) ₆ -CH ₃	H	-OH	2	2'-CH ₃ , 5'-Cl	C ₂₈ H ₃₅ ClN ₄ O ₅	543.23	543.4
17	-OH	H	-OH	1	2'-OCH ₃	C ₂₁ H ₂₂ N ₄ O ₆	427.15	427.2

1. (2*R*,4*R*)-5-(2',5'-Dichlorobiphenyl-4-yl)-2-hydroxy-4-[(1-hydroxy-1H-

- [1,2,3]triazole-4-carbonyl)-amino]-pentanoic acid
2. (2*R*,4*R*)-5-(2',5'-Dichloro-biphenyl-4-yl)-2-hydroxy-4-[(1-hydroxy-1*H*-[1,2,3]triazole-4-carbonyl)-amino]-pentanoic acid isobutyl ester
3. (2*R*,4*R*)-5-(2',5'-Dichloro-biphenyl-4-yl)-2-hydroxy-4-[(1-hydroxy-1*H*-[1,2,3]triazole-4-carbonyl)-amino]-pentanoic acid isopropyl ester
- 5 4. (2*R*,4*R*)-5-(2',5'-Dichloro-biphenyl-4-yl)-2-hydroxy-4-[(1-hydroxy-1*H*-[1,2,3]triazole-4-carbonyl)-amino]-pentanoic acid ethyl ester
5. (2*R*,4*R*)-5-(2',5'-Dichloro-biphenyl-4-yl)-2-hydroxy-4-[(1-hydroxy-1*H*-[1,2,3]triazole-4-carbonyl)-amino]-pentanoic acid heptyl ester
- 10 6. (2*R*,4*R*)-5-(5'-Chloro-2'-fluorobiphenyl-4-yl)-2-hydroxy-4-[(1-hydroxy-1*H*-[1,2,3]triazole-4-carbonyl)-amino]-pentanoic acid
7. (2*R*,4*R*)-5-(5'-Chloro-2'-fluorobiphenyl-4-yl)-2-hydroxy-4-[(1-hydroxy-1*H*-[1,2,3]triazole-4-carbonyl)-amino]-pentanoic acid isobutyl ester
8. (2*R*,4*R*)-5-(5'-Chloro-2'-fluorobiphenyl-4-yl)-2-hydroxy-4-[(1-hydroxy-1*H*-[1,2,3]triazole-4-carbonyl)-amino]-pentanoic acid isopropyl ester
- 15 9. (2*R*,4*R*)-5-(5'-Chloro-2'-fluorobiphenyl-4-yl)-2-hydroxy-4-[(1-hydroxy-1*H*-[1,2,3]triazole-4-carbonyl)-amino]-pentanoic acid ethyl ester
10. (2*R*,4*R*)-5-(5'-Chloro-2'-fluorobiphenyl-4-yl)-2-hydroxy-4-[(1-hydroxy-1*H*-[1,2,3]triazole-4-carbonyl)-amino]-pentanoic acid heptyl ester
- 20 11. (2*R*,4*R*)-4-[[1-(*S*)-2-Amino-3-methyl-butyryloxymethoxy]-1*H*-[1,2,3]triazole-4-carbonyl]-amino}-5-(5'-chloro-2'-fluoro-biphenyl-4-yl)-2-hydroxy-pentanoic acid isopropyl ester
12. (2*R*,4*R*)-5-(5'-Chloro-2'-methyl-biphenyl-4-yl)-2-hydroxy-4-[(1-hydroxy-1*H*-[1,2,3]triazole-4-carbonyl)-amino]-pentanoic acid
- 25 13. (2*R*,4*R*)-5-(5'-Chloro-2'-methyl-biphenyl-4-yl)-2-hydroxy-4-[(1-hydroxy-1*H*-[1,2,3]triazole-4-carbonyl)-amino]-pentanoic acid isobutyl ester
14. (2*R*,4*R*)-5-(5'-Chloro-2'-methyl-biphenyl-4-yl)-2-hydroxy-4-[(1-hydroxy-1*H*-[1,2,3]triazole-4-carbonyl)-amino]-pentanoic acid isopropyl ester
15. (2*R*,4*R*)-5-(5'-Chloro-2'-methyl-biphenyl-4-yl)-2-hydroxy-4-[(1-hydroxy-1*H*-[1,2,3]triazole-4-carbonyl)-amino]-pentanoic acid ethyl ester
- 30 16. (2*R*,4*R*)-5-(5'-Chloro-2'-methyl-biphenyl-4-yl)-2-hydroxy-4-[(1-hydroxy-1*H*-[1,2,3]triazole-4-carbonyl)-amino]-pentanoic acid heptyl ester
17. (2*R*,4*R*)-2-Hydroxy-4-[(1-hydroxy-1*H*-[1,2,3]triazole-4-carbonyl)-amino]-5-(2'-


methoxy-biphenyl-4-yl)-pentanoic acid

5



Ex.	R ¹	R ³	R ⁴	b	R ⁶	Formula	MS <i>m/z</i> : [M+H] ⁺	
							calcd	found
18	-OH	H	-OCH ₂ O-C(O)-CH ₃	2	2'-F, 5'-Cl	C ₂₃ H ₂₂ ClFN ₄ O ₇	521.12	521.1


18. (2*R*,4*R*)-4-[(3-Acetoxymethoxy-3*H*-[1,2,3]triazole-4-carbonyl)-amino]-5-(5'-chloro-2'-fluoro-biphenyl-4-yl)-2-hydroxy-pentanoic acid



10

Ex.	R ¹	R ³	R ⁴	b	R ⁶	Formula	MS <i>m/z</i> : [M+H] ⁺	
							calcd	found
19	-OH	-OCH ₃	H	2	2'-F, 5'-Cl	C ₂₂ H ₂₀ ClFN ₂ O ₆	463.10	463.2


19. (2*R*,4*R*)-5-(5'-Chloro-2'-fluoro-biphenyl-4-yl)-2-hydroxy-4-[(3-methoxy-isoxazole-5-carbonyl)-amino]-pentanoic acid



Ex.	R ¹	R ³	R ⁴	b	R ⁶	Formula	MS <i>m/z</i> : [M+H] ⁺	
							calcd	found
20	-OH	=O	phenyl	2	2'-F, 5'-Cl	C ₂₆ H ₂₂ ClFN ₄ O ₅	525.13	525.2

15

20. (2*R*,4*R*)-5-(5'-Chloro-2'-fluoro-biphenyl-4-yl)-2-hydroxy-4-[(5-oxo-1-phenyl-4,5-dihydro-1*H*-[1,2,4]triazole-3-carbonyl)-amino]-pentanoic acid





Ex.	R ¹	R ⁴	b	R ⁶	Formula	MS <i>m/z</i> : [M+H] ⁺	
						calcd	found
21	-OH	-OCH ₂ O- C(O)-CH ₃	2	2',5'- diCl	C ₂₇ H ₂₄ Cl ₂ N ₄ O ₇	587.10	587.1
22	-OCH ₂ CH- (CH ₃) ₂	-OCH ₂ O- C(O)CH- (CH ₃) ₂ NH ₂	1	3'-Cl	C ₃₄ H ₄₀ ClN ₅ O ₇	666.26	666.0

21. (2*R*,4*R*)-4-[(3-Acetoxymethoxy-3*H*-benzotriazole-5-carbonyl)-amino]-5-(2',5'-dichloro-biphenyl-4-yl)-2-hydroxy-pentanoic acid
22. (2*R*,4*R*)-4-{[3-((*S*)-2-Amino-3-methyl-butyryloxymethoxy)-3*H*-benzotriazole-5-carbonyl]-amino}-5-(3'-chloro-biphenyl-4-yl)-2-hydroxy-pentanoic acid isobutyl ester

10

EXAMPLE 20


Following the procedures described in the examples herein, and substituting the appropriate starting materials and reagents, compounds having the following formula were prepared as the parent compound or as a TFA salt.



15


Ex.	R ³	R ⁴	R ⁵	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
1	H	H	Cl	C ₂₀ H ₁₉ ClN ₄ O ₄	415.11	415.4

1. (2*R*,4*R*)-5-(3-Chlorobiphenyl-4-yl)-2-hydroxy-4-[(1*H*-[1,2,3]triazole-4-carbonyl)-amino]-pentanoic acid (TFA salt)



Ex.	R ³	R ⁴	R ⁵	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
1	H	H	Cl	C ₂₀ H ₁₉ ClN ₄ O ₄	415.11	415.2


- 5 2. (2*R*,4*R*)-5-(3-Chlorobiphenyl-4-yl)-2-hydroxy-4-[(3*H*-[1,2,3]triazole-4-carbonyl)-amino]-pentanoic acid




Ex.	R ³	R ⁴	R ⁵	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
3	=O	3-chlorophenyl	Cl	C ₂₆ H ₂₂ Cl ₂ N ₄ O ₅	541.10	541.2
4	=O	phenyl	Cl	C ₂₆ H ₂₃ ClN ₄ O ₅	507.14	507.2

3. (2*R*,4*R*)-5-(3-Chloro-biphenyl-4-yl)-4-{[1-(3-chloro-phenyl)-5-oxo-4,5-dihydro-1*H*-[1,2,4]triazole-3-carbonyl]-amino}-2-hydroxy-pentanoic acid

4. (2*R*,4*R*)-5-(3-Chloro-biphenyl-4-yl)-2-hydroxy-4-[(5-oxo-1-phenyl-4,5-dihydro-1*H*-[1,2,4]triazole-3-carbonyl)-amino]-pentanoic acid




Ex.	R ³	R ⁴	R ⁵	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
5	-C(O)N(CH ₃) ₂	H	Cl	C ₂₄ H ₂₅ ClN ₄ O ₅	485.15	486.2
6	-C(O)N(CH ₃)-[(CH ₂) ₂ OCH ₃]	H	Cl	C ₂₆ H ₂₉ ClN ₄ O ₆	529.18	529.2
7	-C(O)-CH ₃	H	Cl	C ₂₃ H ₂₂ ClN ₃ O ₅	456.12	456.2
8		H	Cl	C ₂₆ H ₂₃ ClN ₄ O ₄	491.14	491.2


Ex.	R ³	R ⁴	R ⁵	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
9		H	Cl	C ₂₆ H ₂₃ ClN ₄ O ₄	491.14	493.2

5. (2*R*,4*R*)-5-(3-Chloro-biphenyl-4-yl)-4-[(5-dimethylcarbamoyl-2H-pyrazole-3-carbonyl)-amino]-2-hydroxy-pentanoic acid
6. (2*R*,4*R*)-5-(3-Chloro-biphenyl-4-yl)-2-hydroxy-4-({5-[(2-methoxy-ethyl)-methyl-carbamoyl]-2H-pyrazole-3-carbonyl}-amino)-pentanoic acid
- 5 7. (2*R*,4*R*)-4-[(5-Acetyl-2H-pyrazole-3-carbonyl)-amino]-5-(3-chloro-biphenyl-4-yl)-2-hydroxy-pentanoic acid
8. (2*R*,4*R*)-5-(3-Chloro-biphenyl-4-yl)-2-hydroxy-4-[(5-pyridin-3-yl-2H-pyrazole-3-carbonyl)-amino]-pentanoic acid (TFA salt)
- 10 9. (2*R*,4*R*)-5-(3-Chloro-biphenyl-4-yl)-2-hydroxy-4-[(5-pyridin-2-yl-2H-pyrazole-3-carbonyl)-amino]-pentanoic acid (TFA salt)

EXAMPLE 21


- Following the procedures described in the examples herein, and substituting the appropriate starting materials and reagents, compounds having the following formula were
- 15 prepared as the parent compound.





Ex.	R ³	R ⁴	R ⁵	R ⁶	Formula	MS <i>m/z</i> : [M+H] ⁺	
						calcd	found
1	H	H	Cl	Cl	C ₂₀ H ₁₈ Cl ₂ N ₄ O ₄	449.07	449.0

1. (2*R*,4*R*)-5-(3,3'-Dichlorobiphenyl-4-yl)-2-hydroxy-4-[(3H-[1,2,3]triazole-4-carbonyl)-amino]-pentanoic acid




5

Ex.	R ³	R ⁴	R ⁵	R ⁶	Formula	MS <i>m/z</i> : [M+H] ⁺	
						calcd	found
2	-CH ₂ OCH ₃	H	Cl	Cl	C ₂₃ H ₂₃ Cl ₂ N ₃ O ₅	492.10	492.2

2. (2*R*,4*R*)-5-(3,3'-Dichloro-biphenyl-4-yl)-2-hydroxy-4-[(5-methoxymethyl-2H-pyrazole-3-carbonyl)-amino]-pentanoic acid

Preparation 13


- 10 (2*R*,4*R*)-5-(4-Bromophenyl)-2-hydroxy-4-[(3-hydroxy-3H-benzotriazole-5-carbonyl)-amino]-pentanoic Acid Ethyl Ester



- 15 HCTU (837 mg, 2.0 mmol) was added to a solution of 1-hydroxy-1H-1,2,3-benzotriazole-6-carboxylic acid (362 mg, 2.0 mmol) and the mixture was stirred at room temperature for 10 minutes. DIPEA (529 μ L, 3.0 mmol) and (2*R*,4*R*)-4-amino-5-(4-bromophenyl)-2-hydroxy-pentanoic acid ethyl ester (320 mg, 1.0 mmol) were then added and the resulting mixture was stirred at room temperature for 15 minutes. The mixture was concentrated *in vacuo* and the resulting residue was purified (C18 column, 55g, 20-70% MeCN in water with 5% TFA) to yield the title compound as a white solid 365 mg).

EXAMPLE 22

(2R,4R)-5-(5'-Chloro-2'-fluorobiphenyl-4-yl)-2-hydroxy-4-[(3-hydroxy-3H-benzotriazole-5-carbonyl)-amino]-pentanoic Acid




5 SilicaCat[®]DPP-Pd (0.3 mmol/g loading; 37.4 mg, 10 μ mol) was added to a solution of (2R,4R)-5-(4-bromophenyl)-2-hydroxy-4-[(3-hydroxy-3H-benzotriazole-5-carbonyl)-amino]-pentanoic acid ethyl ester (50.0 mg, 105 μ mol), the phenyl boronic acid (157 μ mol, 1.5 eq.) and K₂CO₃ (43.4 mg, 314 μ mol) in EtOH (500 μ L) and water (150 μ L), and the reaction mixture was heated at 100°C for 10 minutes. The mixture was allowed to cool to

10 room temperature, after which 1.0 M of LiOH in water (838 μ L, 838 μ mol) was added and the mixture was stirred for 30 minutes. The mixture was filtered, the filtrate was concentrated *in vacuo* and purified by preparative HPLC to yield the title compound (17.9 mg; purity 98%). MS *m/z* [M+H]⁺ calc'd for C₂₄H₂₀ClFN₄O₅, 499.11; found 499.2.

15

EXAMPLE 23

Following the procedures described in the examples herein, and substituting the appropriate starting materials and reagents, compounds having the following formula were prepared as the parent compound or as a TFA salt.



20


Ex.	R ¹	a	R ⁶	Formula	MS <i>m/z</i> : [M+H] ⁺	
					calcd	found
1	-OH	1	3'-CH ₃	C ₂₅ H ₂₄ N ₄ O ₅	461.17	461.2
2	-OH	2	2'-CH ₃ , 5'-Cl	C ₂₅ H ₂₄ ClN ₄ O ₅	496.14	495.2
3	-OH	1	2'-OCH ₃	C ₂₅ H ₂₄ N ₄ O ₆	477.17	477.2
4	-OH	2	2',5'-diCl	C ₂₄ H ₂₀ Cl ₂ N ₄ O ₅	515.08	515.0
5	-OCH ₂ CH ₃	2	2',5'-diCl	C ₂₆ H ₂₄ Cl ₂ N ₄ O ₅	543.11	543.2
6	-OCH ₂ CH(CH ₃) ₂	2	2',5'-diCl	C ₂₈ H ₂₈ Cl ₂ N ₄ O ₅	571.14	571.2
7	-OCH(CH ₃) ₂	2	2',5'-diCl	C ₂₇ H ₂₆ Cl ₂ N ₄ O ₅	557.13	557.1
8	-OH	2	3'-Cl, 5'-OH	C ₂₄ H ₂₁ ClN ₄ O ₆	497.12	497.2

1. (2*S*,4*S*)-2-Hydroxy-4-[(3-hydroxy-3H-benzotriazole-5-carbonyl)-amino]-5-(3'-methyl-biphenyl-4-yl)-pentanoic acid
2. (2*S*,4*S*)-5-(5'-Chloro-2'-methyl-biphenyl-4-yl)-2-hydroxy-4-[(3-hydroxy-1H-benzotriazole-5-carbonyl)-amino]-pentanoic acid
3. (2*S*,4*S*)-2-Hydroxy-4-[(3-hydroxy-3H-benzotriazole-5-carbonyl)-amino]-5-(2'-methoxy-biphenyl-4-yl)-pentanoic acid
4. (2*S*,4*S*)-5-(2',5'-Dichlorobiphenyl-4-yl)-2-hydroxy-4-[(3-hydroxy-3H-benzotriazole-5-carbonyl)-amino]-pentanoic acid
- 10 5. (2*R*,4*R*)-5-(2',5'-Dichlorobiphenyl-4-yl)-2-hydroxy-4-[(3-hydroxy-3H-benzotriazole-5-carbonyl)-amino]-pentanoic acid ethyl ester
6. (2*R*,4*R*)-5-(2',5'-Dichlorobiphenyl-4-yl)-2-hydroxy-4-[(3-hydroxy-3H-benzotriazole-5-carbonyl)-amino]-pentanoic acid isobutyl ester
7. (2*R*,4*R*)-5-(2',5'-Dichlorobiphenyl-4-yl)-2-hydroxy-4-[(3-hydroxy-3H-benzotriazole-5-carbonyl)-amino]-pentanoic acid isopropyl ester (TFA salt)
- 15 8. (2*S*,4*S*)-5-(3'-Chloro-5'-hydroxy-biphenyl-4-yl)-2-hydroxy-4-[(3-hydroxy-3H-benzotriazole-5-carbonyl)-amino]-pentanoic acid

Preparation 14

(*S*)-2-(1-Biphenyl-4-yl-1-methylethyl)-5-oxo-pyrrolidine-1-carboxylic Acid *t*-Butyl Ester


20



To a solution of 2-(4-bromophenyl)acetonitrile (130.0 g, 0.7 mol) and iodomethane (103.9 mL, 1.7 mol) in THF (1.0 L) was added NaH (60% dispersion in mineral oil, 66.7 g, 1.7 mol) in small portions at 10°C. After completion of the addition, the mixture was stirred at 10°C for another 2 hours. The mixture was poured into ice water (2.0 L) and

extracted with EtOAc (1.5 L). The organic layer was washed with saturated aqueous NaCl, dried over anhydrous MgSO_4 and concentrated to yield compound 1 (175 g, containing mineral oil) as a yellow oil, which was used directly without further purification. ^1H NMR (CDCl_3 , 300 MHz) δ 7.52 (d, J = 9.0 Hz, 2H), 7.38 (d, J = 9.0 Hz, 2H), 1.72 (s, 6H).

5 To a solution of compound 1 (175 g, containing mineral oil) in DCM (1.0 L) was added DIBAL (1.0 M solution in DCM, 700 mL, 0.70 mol) dropwise at -78°C . The reaction mixture was stirred at -78°C for 1.5 hours and then quenched carefully with 3.0 N HCl (1.0 L). The resulting mixture was stirred at room temperature overnight and the organic layer was washed with saturated aqueous NaCl, dried over anhydrous Na_2SO_4 and
 10 concentrated to yield compound 2 (180 g) as a yellow oil, which was used directly without further purification. ^1H NMR (CDCl_3 , 300 MHz) δ 9.48 (s, 1H), 7.53 (d, J = 11.0 Hz, 2H), 7.17 (d, J = 11.0 Hz, 2H), 1.46 (s, 6H).




To an aqueous solution of NaCN (32.7 g in 1.0 L of H_2O , 0.7 mol) were added
 15 $(\text{NH}_4)_2\text{CO}_3$ (380 g, 4.0 mol) and compound 2 (180 g). The reaction mixture was refluxed overnight and then concentrated under reduced pressure at 75°C . Water (350 mL) was added to the residue and the mixture was concentrated again. The residue was suspended in petroleum ether (700 mL) and water (250 mL) and the resulting mixture was stirred at room temperature for 15 minutes. The precipitate was collected by filtration and dried to
 20 yield compound 3 (150 g) as a white solid. ^1H NMR ($\text{DMSO}-d_6$, 300 MHz) δ 10.39 (s, 1H), 8.05 (s, 1H), 7.48 (d, J = 9.0 Hz, 2H), 7.28 (d, J = 9.0 Hz, 2H), 4.17 (s, 1H), 1.42 (s, 3H), 1.34 (s, 3H).

A suspension of compound 3 (150 g, 0.51 mol) in 6.0 N NaOH (400 mL) and ethane-1,2-diol (300 mL) was stirred at 120°C for 38 hours. The mixture was cooled to
 25 room temperature and neutralized with an HCl solution. The precipitate was collected by filtration and dried to yield compound 4 (250 g, containing NaCl salt) as a white solid. ^1H NMR ($\text{DMSO}-d_6$, 300 MHz) δ 7.35 (d, J = 9.0 Hz, 2H), 7.17 (d, J = 9.0 Hz, 2H), 3.22 (s, 1H), 1.16 (s, 3H), 1.15 (s, 3H).

To a suspension of compound 4 (250 g, containing NaCl salt) in MeOH (1.0 L) was
 30 added thionyl chloride (72.0 mL, 1.0 mol) dropwise at 5°C . The mixture was refluxed

overnight and the solvent was removed under reduced pressure. The residue was partitioned between DCM (1.0 L) and saturated aqueous NaHCO₃ (1.5 L). The organic layer was washed with saturated aqueous NaCl, dried over anhydrous Na₂SO₄ and concentrated to yield the corresponding methyl ester (90.0 g). 2-Phenylacetyl chloride (48.6 g, 0.32 mol) was added dropwise to a solution of the ester (90.0 g) and Et₃N (56.5 mL, 0.41 mol) in DCM (1.0 L) at 0°C and the mixture was stirred at 0°C for 30 minutes. The mixture was washed with 1.0 N HCl (500 mL) and saturated aqueous NaCl, respectively. The organic layer was dried over anhydrous Na₂SO₄ and concentrated to yield compound 5 (120 g). ¹H NMR (CDCl₃, 300 MHz) δ 7.32 (m, 5H), 7.18 (m, 2H), 6.95 (m, 2H), 5.68 (br s, 1H), 4.76 (d, J = 9.0 Hz, 1H), 3.57 (s, 3H), 3.53 (d, J = 5.0 Hz, 2H), 1.30 (s, 3H), 1.25 (s, 3H).




To a solution of compound 5 (120 g, 0.30 mol) in MeOH (500 mL) was added 4.0 N NaOH (200 mL). The mixture was stirred at room temperature for 4 hours and then the pH was adjusted to pH=1 with 3.0 N HCl. The resulting mixture was extracted with EtOAc (2x300mL). The combined extracts were washed with saturated aqueous NaCl, dried over anhydrous Na₂SO₄, and concentrated under reduced pressure. The residue was recrystallized from EtOAc/hexanes to yield compound 6 (82.0 g). ¹H NMR (DMSO-d₆, 300 MHz) δ 7.41 (d, J = 6.0 Hz, 2H), 7.22 (m, 5H), 6.99 (d, J = 6.0 Hz, 2H), 4.65 (d, J = 9.0 Hz, 1H), 3.52 (d, J = 14.0 Hz, 1H), 3.36 (d, J = 14.0 Hz, 1H), 1.34 (s, 3H), 1.30 (s, 3H).

A suspension of compound 6 (82.0 g, 0.21 mol) in distilled water (3.0 L) was adjusted to pH=8.5 with 3.0 N LiOH and a clear solution was formed. Immobilized Penicillinase (20.0 g) was added and the resulting mixture was stirred at 37°C for 60 hours. The mixture was filtered and the filtrate was adjusted to pH=1 with 3.0 N HCl and extracted with EtOAc. The combined extracts were washed with saturated aqueous NaCl, dried over anhydrous Na₂SO₄ and concentrated to yield compound 7 (59.0 g, 80% ee, containing 2-phenylacetic acid).


A suspension of compound 7 (59.0 g, containing 2-phenylacetic acid) in 6.0 N HCl (500 mL) was refluxed overnight. The mixture was washed with EtOAc (300 mL) and the

aqueous phase was concentrated under reduced pressure to yield the corresponding amino acid as its hydrochloride salt. The salt was dissolved in water (300 mL) and the solution was adjusted to pH=11. A solution of (Boc)₂O (33.0 g, 0.2 mol) in acetone (200 mL) was added and the mixture was stirred at room temperature for 2 hours. The mixture was washed with hexanes (200 mL) and the aqueous phase was adjusted to pH=2. The resulting mixture was extracted with EtOAc (2x300 mL). The combined extracts were washed with saturated aqueous NaCl, dried over anhydrous Na₂SO₄ and concentrated to yield compound 8 (37.0 g) as a white solid. ¹H NMR (CDCl₃, 300 MHz) δ 9.48 (br s, 1H), 7.46 (d, J = 7.0 Hz, 2H), 7.26 (d, J = 7.0 Hz, 2H), 5.02 (br s, 1H), 4.56 (d, J = 9.0 Hz, 1H), 1.39 (s, 9H).



A mixture of compound 8 (37.0 g, 0.1 mol) in dioxane (200 mL) and 1.0 N K₂CO₃ (200 mL) was degassed for 30 minutes with nitrogen, followed by the addition of phenylboronic acid (13.4 g, 0.1 mol) and Pd(PPh₃)₄ (1.6 g, 1.4 mmol). The mixture was heated at 75°C for 8 hours and then cooled to room temperature. The mixture was washed with EtOAc/hexanes (150 mL, 1:1) and the aqueous phase was adjusted to pH=2 and extracted with EtOAc (2x300 mL). The combined extracts were washed with saturated aqueous NaCl, dried over anhydrous Na₂SO₄ and concentrated to yield compound 9 (31.0 g, 84% yield) as a white solid.

A solution of compound 9 (31.0 g, 84 mmol), Meldrum's acid (13.3 g, 92 mmol) and DMAP (15.4 g, 0.13 mol) in DCM (400 mL) was cooled to -5°C and a solution of DCC (19.0 g, 92 mmol) in DCM (200 mL) was added over 1 hour. The mixture was stirred at -5°C overnight. The precipitate was filtered off and the filtrate was washed with 1.0 N HCl (2x700 mL) and saturated aqueous NaCl, respectively. After the organic layer containing compound 10 was dried over anhydrous MgSO₄, it was used directly for the next step without concentration.




A solution of compound 10 in DCM (600 mL) was cooled to -5°C and AcOH (45.0 mL) was added. Then NaBH_4 (7.0 g, 0.2 mol) was added in small portions over 30 minutes and the mixture was stirred at -5°C for 3 hours. Water (50.0 mL) was added dropwise followed by addition of saturated aqueous NaCl (450 mL). The organic layer was washed with water (2x300 mL) and saturated aqueous NaHCO_3 (2x300 mL), dried over anhydrous MgSO_4 and concentrated to yield compound 11 (32.0 g, 75% ee) as an off-white solid. After recrystallization from EtOH, chirally pure compound 11 (13.0 g) was obtained. ^1H NMR (CDCl_3 , 300 MHz) δ 7.61 (m, 10H), 4.46 (br s, 1H), 4.26 (m, 1H), 3.72 (br s, 1H), 2.23 (m, 1H), 1.79 (s, 3H), 1.76 (s, 3H), 1.48 (s, 6H), 1.39 (s, 9H).

A solution of compound 11 (13.0 g, 27.0 mmol) in toluene (100.0 mL) was refluxed for 3 hours. After evaporation of the solvent, the residue was recrystallized from hexanes/EtOAc (3:1) to yield the title compound (8.0 g) as a white solid.

Preparation 15


(2R,4S)-4-Amino-5-biphenyl-4-yl-2-hydroxy-5-methylhexanoic Acid Ethyl Ester



A mixture of (*S*)-2-(1-biphenyl-4-yl-1-methylethyl)-5-oxo-pyrrolidine-1-carboxylic acid *t*-butyl ester (14.0 g, 36.9 mmol, racemic) in a 3.0 N HCl-EtOAc solution (150 mL) was stirred at room temperature for 3 hours. The solvent was removed under reduced pressure to yield compound 1 (10.0 g) as a white solid.

To a solution of compound 1 (10.0 g, 35.8 mmol) in THF (80.0 mL) was added BuLi (2.5 M in hexanes, 15.0 mL) dropwise at -78°C . After the mixture was stirred for 30 minutes pivaloyl chloride (4.8 mL, 39.4 mmol) was added dropwise. The mixture was stirred at -78°C for 1 hour and then quenched with saturated aqueous NH_4Cl . The resulting mixture was extracted with EtOAc and the combined extracts were washed with saturated

aqueous NaCl, dried over anhydrous MgSO_4 and concentrated. The residue was purified by flash column chromatography on silica gel to yield compound 2 (9.0 g) as a white solid.




To a solution of compound 2 (9.0 g, 24.7 mmol) in THF (50.0 mL) was added sodium bis(trimethylsilyl)amide (2.0 M in THF, 18.5 mL, 37.0 mmol) dropwise at -78°C . The mixture was stirred for 20 minutes and a solution of oxaziridine derivative (10.8 g, 37.0 mmol) in THF (30.0 mL) was added dropwise. The mixture was stirred at -78°C for 30 minutes and then quenched with saturated aqueous NH_4Cl . The resulting mixture was extracted with EtOAc (1.0 L) and the extract was washed with 1.0 N HCl and saturated aqueous NaCl, dried over anhydrous MgSO_4 and evaporated to remove most of the solvent. The precipitate was filtered off and the filtrate was concentrated. The residue was purified by flash column chromatography on silica gel (DCM:hexanes= 1:1 to DCM) to yield compound 3 (4.3 g, racemic). This racemate was subjected to chiral AD-column chromatography to afford chirally pure compound 3 (1.4 g). ^1H NMR (DMSO- d_6 , 300 MHz) δ 7.63 (m, 4H), 7.49 (m, 4H), 4.83 (d, 1H), 3.29 (m, 1H), 2.31 (m, 2H), 1.40 (s, 3H), 1.36 (s, 3H), 1.28 (s, 9H). LC-MS (ESI): m/z 380.1 $[\text{M}+\text{H}]^+$.

A solution of compound 3 (1.7 g, 160 mmol) in EtOH (15.0 mL) and 12.0 N HCl (15.0 mL) was heated at $90\sim 95^\circ\text{C}$ for 20 hours. The solvent was removed and the residue was treated with a 3.0 N HCl-EtOH solution (25.0 mL) under reflux for another 3 hours. After removal of the solvent, the residue was purified by preparative HPLC to yield the title compound (0.6 g) as a foamy solid HCl salt. ^1H NMR (DMSO- d_6 , 300 MHz) δ 7.88 (br s, 3H), 7.68 (m, 4H), 7.49 (m, 4H), 7.35 (m, 1H), 6.11 (br s, 1H), 4.11 (br s, 1H), 4.05 (q, 2H), 3.61 (br s, 1H), 1.67 (m, 2H), 1.40 (s, 3H), 1.36 (s, 3H), 1.09 (t, 3H). LC-MS (ESI): m/z 342.1 $[\text{M}+\text{H}]^+$.

EXAMPLE 24


5-[(S)-2-Biphenyl-4-yl-1-((R)-2-carboxy-2-hydroxy-ethyl)-2-methyl-propylcarbamoyl]-
1H-pyrazole-3-carboxylic Acid



- 5 (2*R*,4*S*)-4-Amino-5-biphenyl-4-yl-2-hydroxy-5-methyl-hexanoic acid ethyl ester (70 mg, 0.2 mmol), 3,5-pyrazoledicarboxylic acid (32 mg, 0.2 mmol), and HCTU (85 mg, 0.2 mmol) were combined in DMF (5 mL) and stirred for 2 minutes. DIPEA (79 mg, 0.6 mmol) was added and the resulting mixture was stirred for 1 hour at 50°C. The reaction mixture was evaporated under reduced pressure. The crude material was dissolved in
- 10 EtOH, and add enough equivalents of 10N NaOH were added to make the mixture basic. The reaction was monitor closely over 1 hour until final deprotection is complete. The mixture was then re-acidified with an equal volume of acetic acid and evaporated under reduced pressure. The product was purified using reverse phase chromatography (10-70% MeCN gradient) to yield the title compound as a TFA salt (14 mg; purity 90%). MS *m/z*
- 15 $[M+H]^+$ calc'd for $C_{24}H_{25}N_3O_6$, 452.17; found 452.2.

EXAMPLE 25

- Following the procedures described in the examples herein, and substituting the appropriate starting materials and reagents, compounds having the following formula were
- 20 prepared as the parent compound or as a TFA salt.



Ex.	$-XR^3R^4$	Formula	MS m/z : $[M+H]^+$	
			calcd	found
1		$C_{22}H_{24}N_4O_4$	409.18	409.4
2		$C_{22}H_{24}N_4O_5$	425.17	425.2
3		$C_{23}H_{24}N_2O_6$	425.16	425.2

1. (2*R*,4*S*)-5-Biphenyl-4-yl-2-hydroxy-5-methyl-4-[(1*H*-[1,2,3]triazole-4-carbonyl)-amino]-hexanoic acid (TFA salt)

2. (2*R*,4*S*)-5-Biphenyl-4-yl-2-hydroxy-4-[(1-hydroxy-1*H*-[1,2,3]triazole-4-carbonyl)-amino]-5-methyl-hexanoic acid (TFA salt)

3. (2*R*,4*S*)-5-Biphenyl-4-yl-2-hydroxy-4-[(3-hydroxy-isoxazole-5-carbonyl)-amino]-5-methyl-hexanoic acid

ASSAY 1

In vitro Assays for the Quantitation of Inhibitor Potencies

10 at Human and Rat NEP, and Human ACE

The inhibitory activities of compounds at human and rat neprilysin (EC 3.4.24.11; NEP) and human angiotensin converting enzyme (ACE) were determined using *in vitro* assays as described below.

Extraction of NEP Activity from Rat Kidneys

15 Rat NEP was prepared from the kidneys of adult Sprague Dawley rats. Whole kidneys were washed in cold phosphate buffered saline (PBS) and brought up in ice-cold lysis buffer (1% Triton X-114, 150 mM NaCl, 50 mM tris(hydroxymethyl) aminomethane (Tris) pH 7.5; Bordier (1981) *J. Biol. Chem.* 256: 1604-1607) in a ratio of 5 mL of buffer for every gram of kidney. Samples were homogenized on ice using a polytron hand held
20 tissue grinder. Homogenates were centrifuged at 1000 x g in a swinging bucket rotor for 5 minutes at 3°C. The pellet was resuspended in 20 mL of ice cold lysis buffer and incubated on ice for 30 minutes. Samples (15-20 mL) were then layered onto 25 mL of ice-cold cushion buffer (6% w/v sucrose, 50 mM pH 7.5 Tris, 150 mM NaCl, 0.06%, Triton X-114), heated to 37°C for 3-5 minutes and centrifuged at 1000 x g in a swinging
25 bucket rotor at room temperature for 3 minutes. The two upper layers were aspirated off, leaving a viscous oily precipitate containing the enriched membrane fraction. Glycerol

was added to a concentration of 50% and samples were stored at -20°C. Protein concentrations were quantitated using a BCA detection system with bovine serum albumin (BSA) as a standard.

Enzyme Inhibition Assays

5 Recombinant human NEP and recombinant human ACE were obtained commercially (R&D Systems, Minneapolis, MN, catalog numbers 1182-ZN and 929-ZN, respectively). The fluorogenic peptide substrate Mca-D-Arg-Arg-Leu-Dap-(Dnp)-OH (Medeiros et al. (1997) *Braz. J. Med. Biol. Res.* 30:1157-62; Anaspec, San Jose, CA) and Abz-Phe-Arg-Lys(Dnp)-Pro-OH (Araujo et al. (2000) *Biochemistry* 39:8519-8525; 10 Bachem, Torrance, CA) were used in the NEP and ACE assays respectively.

The assays were performed in 384-well white opaque plates at 37°C using the fluorogenic peptide substrates at a concentration of 10 µM in Assay Buffer (NEP: 50 mM HEPES, pH 7.5, 100 mM NaCl, 0.01% polyethylene glycol sorbitan monolaurate (Tween-20), 10 µM ZnSO₄; ACE: 50 mM HEPES, pH 7.5, 100 mM NaCl, 0.01% Tween-20, 1 µM 15 ZnSO₄). The respective enzymes were used at concentrations that resulted in quantitative proteolysis of 1 µM of substrate after 20 minutes at 37°C.

Test compounds were assayed over the range of concentrations from 10 µM to 20 pM. Test compounds were added to the enzymes and incubated for 30 minute at 37°C prior to initiating the reaction by the addition of substrate. Reactions were terminated after 20 minutes of incubation at 37°C by the addition of glacial acetic acid to a final 20 concentration of 3.6% (v/v).

Plates were read on a fluorometer with excitation and emission wavelengths set to 320 nm and 405 nm, respectively. Inhibition constants were obtained by nonlinear regression of the data using the equation (GraphPad Software, Inc., San Diego, CA):

$$25 \quad v = v_0 / [1 + (I / K')]$$

where v is the reaction rate, v_0 is the uninhibited reaction rate, I is the inhibitor concentration and K' is the apparent inhibition constant.

Compounds of the invention were tested in this assay and found to have pK_i values at human NEP as follows. In general, either the prodrug compounds did not inhibit the 30 enzyme in this *in vitro* assay, or the prodrugs were not tested (n.d.) since activity would not be expected.

Ex.	pK _i
I-A	n.d.
I-B	≥9.0

Ex.	pK _i
15-64	n.d.
15-65	7.0-7.9

Ex.	pK _i
1-C	n.d.
1-D	n.d.
1-E	n.d.
1-F	n.d.
1-G	n.d.
2-A	n.d.
2-B	n.d.
2-C	≥9.0
2-D	≥9.0
2-E	n.d.
3-A	n.d.
3-B	n.d.
3-C	8.0-8.9
3-D	≥9.0
4-A	n.d.
4-B	≥9.0
5-A	n.d.
5-B	≥9.0
6-A	n.d.
6-B	≥9.0
7-A	n.d.
7-B	≥9.0
7-C	n.d.
8-A	n.d.
8-B	≥9.0
9	7.0-7.9
10-1	n.d.
10-2	8.0-8.9
10-3	≥9.0
10-4	n.d.
10-5	n.d.
10-6	7.0-7.9
10-7	7.0-7.9
10-8	n.d.
10-9	8.0-8.9
10-10	8.0-8.9
10-11	7.0-7.9
10-12	8.0-8.9
11-1	8.0-8.9
11-2	7.0-7.9
11-3	8.0-8.9
11-4	8.0-8.9
11-5	n.d.
11-6	8.0-8.9
11-7	8.0-8.9
11-8	7.0-7.9

Ex.	pK _i
15-66	n.d.
15-67	≥9.0
15-68	n.d.
15-69	7.0-7.9
15-70	8.0-8.9
15-71	8.0-8.9
15-72	7.0-7.9
15-73	7.0-7.9
15-74	7.0-7.9
15-75	8.0-8.9
15-76	8.0-8.9
15-77	7.0-7.9
15-77	8.0-8.9
15-79	≥9.0
15-80	≥9.0
15-81	7.0-7.9
15-82	8.0-8.9
15-83	≥9.0
15-84	≥9.0
15-85	n.d.
15-86	≥9.0
15-87	n.d.
15-88	n.d.
15-89	≥9.0
15-90	≥9.0
15-91	7.0-7.9
15-92	7.0-7.9
15-93	7.0-7.9
15-94	7.0-7.9
15-95	8.0-8.9
15-96	≥9.0
15-97	n.d.
15-98	n.d.
15-99	n.d.
15-100	8.0-8.9
15-101	7.0-7.9
15-102	≥9.0
15-103	n.d.
15-104	n.d.
15-105	n.d.
15-106	8.0-8.9
15-107	n.d.
15-108	7.0-7.9
15-109	8.0-8.9
15-110	8.0-8.9
15-111	≥9.0

Ex.	pK _i
11-9	7.0-7.9
11-10	n.d.
11-11	8.0-8.9
11-12	n.d.
11-13	≥9.0
11-14	n.d.
11-15	7.0-7.9
12-1	8.0-8.9
12-2	8.0-8.9
12-3	8.0-8.9
12-4	≥9.0
12-5	≥9.0
12-6	≥9.0
12-7	≥9.0
12-8	≥9.0
12-9	≥9.0
12-10	8.0-8.9
12-11	≥9.0
12-12	8.0-8.9
12-13	≥9.0
12-14	7.0-7.9
12-15	8.0-8.9
12-16	≥9.0
12-17	8.0-8.9
12-18	8.0-8.9
12-19	7.0-7.9
12-20	8.0-8.9
12-21	≥9.0
12-22	≥9.0
12-23	8.0-8.9
12-24	7.0-7.9
12-25	7.0-7.9
12-26	8.0-8.9
12-27	≥9.0
13-1	8.0-8.9
13-2	n.d.
13-3	7.0-7.9
13-4	n.d.
13-5	7.0-7.9
13-6	7.0-7.9
13-7	7.0-7.9
13-8	7.0-7.9
14	8.0-8.9
15-1	7.0-7.9
15-2	8.0-8.9
15-3	8.0-8.9

Ex.	pK _i
15-112	8.0-8.9
15-113	7.0-7.9
16-1	≥9.0
16-2	7.0-7.9
16-3	8.0-8.9
16-4	8.0-8.9
16-5	7.0-7.9
16-6	8.0-8.9
17-A	≥9.0
17-B	n.d.
18-1	n.d.
18-2	≥9.0
18-3	≥9.0
18-4	n.d.
18-5	n.d.
18-6	n.d.
18-7	≥9.0
18-8	≥9.0
18-9	≥9.0
18-10	≥9.0
18-11	≥9.0
18-12	≥9.0
18-13	≥9.0
18-14	≥9.0
18-15	≥9.0
18-16	≥9.0
18-17	≥9.0
18-18	n.d.
18-19	≥9.0
18-20	n.d.
18-21	n.d.
18-22	n.d.
18-23	≥9.0
18-24	n.d.
18-25	n.d.
18-26	n.d.
18-27	≥9.0
18-28	n.d.
18-29	n.d.
18-30	n.d.
18-31	≥9.0
18-32	≥9.0
18-33	n.d.
18-34	≥9.0
18-35	≥9.0
18-36	n.d.

Ex.	pK _i
15-4	8.0-8.9
15-5	≥9.0
15-6	7.0-7.9
15-7	8.0-8.9
15-8	8.0-8.9
15-9	8.0-8.9
15-10	8.0-8.9
15-11	8.0-8.9
15-12	7.0-7.9
15-13	8.0-8.9
15-14	8.0-8.9
15-15	8.0-8.9
15-16	8.0-8.9
15-17	7.0-7.9
15-18	n.d.
15-19	7.0-7.9
15-20	8.0-8.9
15-21	≥9.0
15-22	8.0-8.9
15-23	7.0-7.9
15-24	8.0-8.9
15-25	≥9.0
15-26	≥9.0
15-27	≥9.0
15-28	≥9.0
15-29	n.d.
15-30	≥9.0
15-31	8.0-8.9
15-32	8.0-8.9
15-33	≥9.0
15-34	8.0-8.9
15-35	≥9.0
15-36	8.0-8.9
15-37	8.0-8.9
15-38	8.0-8.9
15-39	7.0-7.9
15-40	8.0-8.9
15-41	n.d.
15-42	≥9.0
15-43	n.d.
15-44	8.0-8.9
15-45	n.d.
15-46	≥9.0
15-47	n.d.
15-48	≥9.0
15-49	n.d.

Ex.	pK _i
18-37	≥9.0
18-38	≥9.0
18-39	≥9.0
18-40	≥9.0
18-41	≥9.0
18-42	≥9.0
18-43	≥9.0
18-44	≥9.0
18-45	≥9.0
18-46	≥9.0
18-47	≥9.0
18-48	8.0-8.9
18-49	8.0-8.9
18-50	≥9.0
19-1	≥9.0
19-2	n.d.
19-3	n.d.
19-4	n.d.
19-5	n.d.
19-6	≥9.0
19-7	n.d.
19-8	n.d.
19-9	n.d.
19-10	n.d.
19-11	n.d.
19-12	≥9.0
19-13	n.d.
19-14	n.d.
19-15	n.d.
19-16	n.d.
19-17	≥9.0
19-18	≥9.0
19-19	≥9.0
19-20	≥9.0
19-21	≥9.0
19-22	n.d.
20-1	≥9.0
20-2	≥9.0
20-3	≥9.0
20-4	≥9.0
20-5	≥9.0
20-6	≥9.0
20-7	≥9.0
20-8	≥9.0
20-9	8.0-8.9
21-1	≥9.0

Ex.	pK _i
15-50	7.0-7.9
15-51	n.d.
15-52	n.d.
15-53	8.0-8.9
15-54	n.d.
15-55	7.0-7.9
15-56	8.0-8.9
15-57	≥9.0
15-58	n.d.
15-59	8.0-8.9
15-60	≥9.0
15-61	≥9.0
15-62	n.d.
15-63	≥9.0

Ex.	pK _i
21-2	≥9.0
22	≥9.0
23-1	≥9.0
23-2	≥9.0
23-3	≥9.0
23-4	≥9.0
23-5	n.d.
23-6	n.d.
23-7	n.d.
23-8	≥9.0
24	≥9.0
25-1	7.0-7.9
25-2	8.0-8.9
25-3	≥9.0

n.d. = not determined

ASSAY 2

5 Pharmacodynamic (PD) assay for ACE and NEP Activity in Anesthetized Rats

Male, Sprague Dawley, normotensive rats are anesthetized with 120 mg/kg (i.p.) of inactin. Once anesthetized, the jugular vein, carotid artery (PE 50 tubing) and bladder (flared PE 50 tubing) catheters are cannulated and a tracheotomy is performed (Teflon Needle, size 14 gauge) to facilitate spontaneous respiration. The animals are then allowed a
10 60 minute stabilization period and kept continuously infused with 5 mL/kg/h of saline (0.9%) throughout, to keep them hydrated and ensure urine production. Body temperature is maintained throughout the experiment by use of a heating pad. At the end of the 60 minute stabilization period, the animals are dosed intravenously (i.v.) with two doses of AngI (1.0 µg/kg, for ACE inhibitor activity) at 15 minutes apart. At 15 minutes post-
15 second dose of AngI, the animals are treated with vehicle or test compound. Five minutes later, the animals are additionally treated with a bolus i.v. injection of atrial natriuretic peptide (ANP; 30 µg/kg). Urine collection (into pre-weighted eppendorf tubes) is started immediately after the ANP treatment and continued for 60 minutes. At 30 and 60 minutes into urine collection, the animals are re-challenged with AngI. Blood pressure
20 measurements are done using the Notocord system (Kalamazoo, MI). Urine samples are frozen at -20 °C until used for the cGMP assay. Urine cGMP concentrations are determined by Enzyme Immuno Assay using a commercial kit (Assay Designs, Ann Arbor, Michigan, Cat. No. 901-013). Urine volume is determined gravimetrically. Urinary cGMP output is calculated as the product of urine output and urine cGMP concentration. ACE

inhibition is assessed by quantifying the % inhibition of pressor response to AngI. NEP inhibition is assessed by quantifying the potentiation of ANP-induced elevation in urinary cGMP output.

ASSAY 3

5 *In Vivo* Evaluation of Antihypertensive Effects in the Conscious SHR Model of Hypertension

Spontaneously hypertensive rats (SHR, 14-20 weeks of age) are allowed a minimum of 48 hours acclimation upon arrival at the testing site with free access to food and water. For blood pressure recording, these animals are surgically implanted with
10 small rodent radiotransmitters (telemetry unit; DSI Models TA11PA-C40 or C50-PXT, Data Science Inc., USA). The tip of the catheter connected to the transmitter is inserted into the descending aorta above the iliac bifurcation and secured in place with tissue adhesive. The transmitter is kept intraperitoneally and secured to the abdominal wall while closing of the abdominal incision with a non-absorbable suture. The outer skin is
15 closed with suture and staples. The animals are allowed to recover with appropriate post operative care. On the day of the experiment, the animals in their cages are placed on top of the telemetry receiver units to acclimate to the testing environment and baseline recording. After at least of 2 hours baseline measurement is taken, the animals are then dosed with vehicle or test compound and followed out to 24 hours post-dose blood
20 pressure measurement. Data is recorded continuously for the duration of the study using Notocord software (Kalamazoo, MI) and stored as electronic digital signals. Parameters measured are blood pressure (systolic, diastolic and mean arterial pressure) and heart rate.

ASSAY 4

25 *In Vivo* Evaluation of Antihypertensive Effects in the Conscious DOCA-Salt Rat Model of Hypertension

CD rats (male, adult, 200-300 grams, Charles River Laboratory, USA) are allowed a minimum of 48 hours acclimation upon arrival at the testing site before they are placed on a high salt diet. One week after the start of the high salt diet (8% in food or 1% NaCl in drinking water), a deoxycorticosterone acetate (DOCA) pellet (100 mg, 90 days release
30 time, Innovative Research of America, Sarasota, FL) is implanted subcutaneously and unilateral nephrectomy is performed. At this time, the animals are also surgically implanted with small rodent radiotransmitters for blood pressure measurement (see Assay 3 for details). The animals are allowed to recover with appropriate post operative care.

Study design, data recording, and parameters measured is similar to that described for Assay 3.

ASSAY 5

In Vivo Evaluation of Antihypertensive Effects

in the Conscious Dahl/SS Rat Model of Hypertension


Male, Dahl salt sensitive rats (Dahl/SS, 6-7 weeks of age from Charles River Laboratory, USA) are allowed at least 48 hours of acclimation upon arrival at the testing site before they were placed on a 8% NaCl high salt diet (TD.92012, Harlan, USA) then surgically implanted with small rodent radiotransmitters for blood pressure measurement (see Assay 3 for details). The animals are allowed to recover with appropriate post operative care. At approximately 4 to 5 weeks from the start of high salt diet, these animals are expected to become hypertensive. Once the hypertension level is confirmed, these animals are used for the study while continued with the high salt diet to maintain their hypertension level. Study design, data recording, and parameters measured is similar to that described in Assay 3.

While the present invention has been described with reference to specific aspects or embodiments thereof, it will be understood by those of ordinary skilled in the art that various changes can be made or equivalents can be substituted without departing from the true spirit and scope of the invention. Additionally, to the extent permitted by applicable patent statutes and regulations, all publications, patents and patent applications cited herein are hereby incorporated by reference in their entirety to the same extent as if each document had been individually incorporated by reference herein.

CLAIMS

WHAT IS CLAIMED IS:

1. A compound of formula I:



5 where:

R¹ is selected from -OR⁷ and -NR⁸R⁹;R² is H or -P(O)(OH)₂ or R² is taken together with R⁷ to form -CR¹⁸R¹⁹- or is taken together with R⁸ to form -C(O)-;X is a -C₁₋₉heteroaryl;

10 R³ is absent or is selected from H; halo; -C₀₋₅alkylene-OH; -NH₂; -C₁₋₆alkyl; -CF₃; -C₃₋₇cycloalkyl; -C₀₋₂alkylene-O-C₁₋₆alkyl; -C(O)R²⁰; -C₀₋₁alkylene-COOR²¹; -C(O)NR²²R²³; -NHC(O)R²⁴; =O; -NO₂; -C(CH₃)=N(OH); phenyl optionally substituted with one or two groups independently selected from halo, -OH, -CF₃, -OCH₃, -NHC(O)CH₃, and phenyl; naphthalenyl; pyridinyl; pyrazinyl; pyrazolyl optionally substituted with methyl; thiophenyl optionally substituted with methyl or halo; furanyl; and

15 -CH₂-morpholinyl; and R³, when present, is attached to a carbon atom;

R⁴ is absent or is selected from H; -OH; -C₁₋₆alkyl; -C₁₋₂alkylene-COOR³⁵;

-CH₂OC(O)CH(R³⁶)NH₂; -OCH₂OC(O)CH(R³⁶)NH₂; -OCH₂OC(O)CH₃;

-CH₂OP(O)(OH)₂; -CH₂CH(OH)CH₂OH; -CH[CH(CH₃)₂]-NHC(O)O-C₁₋₆alkyl; pyridinyl;


20 and phenyl or benzyl optionally substituted with one or more groups selected from halo, -COOR³⁵, -OCH₃, -OCF₃, and -SCF₃; and R⁴, when present, is attached to a carbon or nitrogen atom;

or R³ and R⁴ are taken together to form -phenylene-O-(CH₂)₁₋₃- or -phenylene-O-CH₂-CHOH-CH₂-;

25 a is 0 or 1; R⁵ is selected from halo, -CH₃, -CF₃, and -CN;

b is 0 or an integer from 1 to 3; each R⁶ is independently selected from halo, -OH, -CH₃, -OCH₃, and -CF₃;

R^7 is selected from H, $-C_{1-8}$ alkyl, $-C_{1-3}$ alkylene- C_{6-10} aryl, $-C_{1-3}$ alkylene- C_{1-9} heteroaryl, $-C_{3-7}$ cycloalkyl, $-[(CH_2)_2O]_{1-3}CH_3$, $-C_{1-6}$ alkylene- $OC(O)R^{10}$, $-C_{1-6}$ alkylene- $NR^{12}R^{13}$, $-C_{1-6}$ alkylene- $C(O)R^{31}$, $-C_{0-6}$ alkylenemorpholinyl, $-C_{1-6}$ alkylene- SO_2-C_{1-6} alkyl,



- 5 R^{10} is selected from $-C_{1-6}$ alkyl, $-O-C_{1-6}$ alkyl, $-C_{3-7}$ cycloalkyl, $-O-C_{3-7}$ cycloalkyl, phenyl, $-O$ -phenyl, $-NR^{12}R^{13}$, $-CH[CH(CH_3)_2]-NH_2$, $-CH[CH(CH_3)_2]-NHC(O)O-C_{1-6}$ alkyl, and $-CH(NH_2)CH_2COOCH_3$; and R^{12} and R^{13} are independently selected from H, $-C_{1-6}$ alkyl, and benzyl; or R^{12} and R^{13} are taken together as $-(CH_2)_{3-6}-$, $-C(O)-(CH_2)_3-$, or $-(CH_2)_2O(CH_2)_2-$; R^{31} is selected from $-O-C_{1-6}$ alkyl, $-O$ -benzyl, and $-NR^{12}R^{13}$; and R^{32} is
- 10 $-C_{1-6}$ alkyl or $-C_{0-6}$ alkylene- C_{6-10} aryl;


R^8 is selected from H, $-OH$, $-OC(O)R^{14}$, $-CH_2COOH$, $-O$ -benzyl, $-pyridyl$, and $-OC(S)NR^{15}R^{16}$; R^{14} is selected from H, $-C_{1-6}$ alkyl, $-C_{6-10}$ aryl, $-OCH_2-C_{6-10}$ aryl, $-CH_2O-C_{6-10}$ aryl, and $-NR^{15}R^{16}$; and R^{15} and R^{16} are independently selected from H and $-C_{1-4}$ alkyl;

- 15 R^9 is selected from H, $-C_{1-6}$ alkyl, and $-C(O)-R^{17}$; and R^{17} is selected from H, $-C_{1-6}$ alkyl, $-C_{3-7}$ cycloalkyl, $-C_{6-10}$ aryl, and $-C_{1-9}$ heteroaryl;

R^{18} and R^{19} are independently selected from H, $-C_{1-6}$ alkyl, and $-O-C_{3-7}$ cycloalkyl, or R^{18} and R^{19} are taken together to form $=O$;

R^{20} is selected from H and $-C_{1-6}$ alkyl;

- 20 R^{21} and R^{35} are independently selected from H, $-C_{1-6}$ alkyl, $-C_{1-3}$ alkylene- C_{6-10} aryl, $-C_{1-3}$ alkylene- C_{1-9} heteroaryl, $-C_{3-7}$ cycloalkyl, $-[(CH_2)_2O]_{1-3}CH_3$, $-C_{1-6}$ alkylene- $OC(O)R^{25}$, $-C_{1-6}$ alkylene- $NR^{27}R^{28}$, $-C_{1-6}$ alkylene- $C(O)R^{33}$, $-C_{0-6}$ alkylenemorpholinyl, $-C_{1-6}$ alkylene- SO_2-C_{1-6} alkyl,



- 25 R^{25} is selected from $-C_{1-6}$ alkyl, $-O-C_{1-6}$ alkyl, $-C_{3-7}$ cycloalkyl, $-O-C_{3-7}$ cycloalkyl, phenyl, $-O$ -phenyl, $-NR^{27}R^{28}$, $-CH[CH(CH_3)_2]-NH_2$, $-CH[CH(CH_3)_2]-NHC(O)O-C_{1-6}$ alkyl, and $-CH(NH_2)CH_2COOCH_3$; R^{27} and R^{28} are independently selected from H, $-C_{1-6}$ alkyl, and

benzyl; or R^{27} and R^{28} are taken together as $-(CH_2)_{3-6}-$, $-C(O)-(CH_2)_3-$, or $-(CH_2)_2O(CH_2)_2-$; R^{33} is selected from $-O-C_{1-6}alkyl$, $-O-benzyl$, and $-NR^{27}R^{28}$; and R^{34} is $-C_{1-6}alkyl$ or $-C_{0-6}alkylene-C_{6-10}aryl$;

R^{22} and R^{23} are independently selected from H, $-C_{1-6}alkyl$, $-CH_2COOH$, $-(CH_2)_2OH$,
 5 $-(CH_2)_2OCH_3$, $-(CH_2)_2SO_2NH_2$, $-(CH_2)_2N(CH_3)_2$, $-C_{0-1}alkylene-C_{3-7}cycloalkyl$, and $-(CH_2)_2-imidazolyl$; or R^{22} and R^{23} are taken together to form a saturated or partially unsaturated $-C_{3-5}heterocycle$ optionally substituted with halo, $-OH$, $-COOH$, or $-CONH_2$; and optionally containing an oxygen atom in the ring;

R^{24} is selected from $-C_{1-6}alkyl$; $-C_{0-1}alkylene-O-C_{1-6}alkyl$; phenyl optionally
 10 substituted with halo or $-OCH_3$; and $-C_{1-9}heteroaryl$; and

R^{36} is selected from H, $-CH(CH_3)_2$, phenyl, and benzyl;


where each alkyl group in R^1 , R^3 , and R^4 is optionally substituted with 1 to 8 fluoro atoms; and;

where the methylene linker on the biphenyl is optionally substituted with one or
 15 two $-C_{1-6}alkyl$ groups or cyclopropyl;
 or a pharmaceutically acceptable salt thereof.

2. The compound of Claim 1, where X is selected from pyrazole, imidazole, triazole, benzotriazole, furan, pyrrole, tetrazole, pyrazine, thiophene, oxazole, isoxazole, thiazole, isothiazole, oxadiazole, thiadiazole, pyridazine, pyridine, pyrimidine, pyran,
 20 benzimidazole, benzoxazole, benzothiazole, pyridylimidazole, and pyridyltriazole.

3. The compound of Claim 2, where X is selected from pyrazole, triazole, benzotriazole, furan, tetrazole, pyrazine, thiophene, oxazole, isoxazole, thiazole, oxadiazole, pyridazine, pyridine, pyrimidine, benzoxazole, pyridylimidazole, and pyridyltriazole.

25 4. The compound of Claim 3, having formula III:




(III)

5. The compound as claimed in any one of Claims 1 to 4, where R^1 is selected from

-OR⁷ and -NR⁸R⁹, R⁷ is H, R⁸ is H or -OH, and R⁹ is H.

6. The compound as claimed in any one of Claims 1 to 4, where:

- R¹ is -OR⁷; and R⁷ is selected from -C₁₋₈alkyl, -C₁₋₃alkylene-C₆₋₁₀aryl, -C₁₋₃alkylene-C₁₋₉heteroaryl, -C₃₋₇cycloalkyl, -[(CH₂)₂O]₁₋₃CH₃, -C₁₋₆alkylene-OC(O)R¹⁰,
 5 -C₁₋₆alkylene-NR¹²R¹³, -C₁₋₆alkylene-C(O)R³¹, -C₀₋₆alkylenemorpholinyl, -C₁₋₆alkylene-SO₂-C₁₋₆alkyl,



R¹ is -NR⁸R⁹; R⁸ is selected from -OC(O)R¹⁴, -CH₂COOH, -O-benzyl, pyridyl, and -OC(S)NR¹⁵R¹⁶; and R⁹ is H; or


- 10 R¹ is -NR⁸R⁹; R⁸ is selected from -OC(O)R¹⁴, -CH₂COOH, -O-benzyl, pyridyl, and -OC(S)NR¹⁵R¹⁶; and R⁹ is -C₁₋₆alkyl or -C(O)R¹⁷;

R¹ is -NR⁸R⁹; R⁸ is selected from H and -OH; and R⁹ is selected from -C₁₋₆alkyl, and -C(O)R¹⁷;

R¹ is -OR⁷ and R² is taken together with R⁷ to form -CR¹⁸R¹⁹-; or

- 15 R¹ is -NR⁸R⁹ and R² is taken together with R⁸ to form -C(O)-.

7. The compound as claimed in any one of Claims 1 to 4, where R¹ is -OR⁷, and R⁷ is selected from H, -C₁₋₈alkyl, -C₁₋₆alkylene-OC(O)R¹⁰, -C₀₋₆alkylenemorpholinyl, -C₁₋₆alkylene-SO₂-C₁₋₆alkyl, and



- 20 R¹⁰ is -O-C₃₋₇cycloalkyl; and R³² is -CH₃.


8. The compound as claimed in any one of Claims 1 to 4, where R¹ is -NR⁸R⁹, where R⁸ is H and R⁹ is H.

9. The compound as claimed in any one of Claims 1 to 8, where R² is H.

10. The compound as claimed in any one of Claims 1 to 9, where R³ is absent or is
 25 selected from H; halo; -C₀₋₅alkylene-OH; -NH₂; -C₁₋₆alkyl; -CF₃; -C₃₋₇cycloalkyl; -C₀₋₂alkylene-O-C₁₋₆alkyl; -C(O)R²⁰; -C₀₋₁alkylene-COOR²¹; -C(O)NR²²R²³; -NHC(O)R²⁴, =O; -NO₂; -C(CH₃)=N(OH); phenyl optionally substituted with one or two groups

independently selected from halo, -OH, -CF₃, -OCH₃, -NHC(O)CH₃, and phenyl; naphthalenyl; pyridinyl; pyrazinyl; pyrazolyl optionally substituted with methyl; thiophenyl optionally substituted with methyl or halo; furanyl; and -CH₂-morpholinyl; and R²¹ is H.

- 5 11. The compound as claimed in any one of Claims 1 to 9, where R³ is -C₀₋₁alkylene-COOR²¹; and R²¹ is selected from -C₁₋₆alkyl, -C₁₋₃alkylenc-C₆₋₁₀aryl, -C₁₋₃alkylenc-C₁₋₉heteroaryl, -C₃₋₇cycloalkyl, -[(CH₂)₂O]₁₋₃CH₃, -C₁₋₆alkylene-OC(O)R²⁵, -C₁₋₆alkylene-NR²⁷R²⁸, -C₁₋₆alkylene-C(O)R³³, -C₀₋₆alkylenemorpholinyl, -C₁₋₆alkylene-SO₂-C₁₋₆alkyl,




12. The compound as claimed in any one of Claims 1 to 9, where R³ is absent or is selected from H; halo; -C₀₋₅alkylene-OH; -NH₂; -C₁₋₆alkyl; -CF₃; -C₃₋₇cycloalkyl; -C₀₋₂alkylene-O-C₁₋₆alkyl; -C(O)R²⁰; -C₀₋₁alkylene-COOR²¹; -C(O)NR²²R²³; -NHC(O)R²⁴; =O; -C(CH₃)=N(OH); phenyl optionally substituted with one or two groups independently selected from halo, -OH, -CF₃, -OCH₃, -NHC(O)CH₃, and phenyl; naphthalenyl; pyridinyl; pyrazinyl; pyrazolyl substituted with methyl; thiophenyl substituted with methyl or halo; furanyl; and -CH₂-morpholinyl; R²⁰ is -C₁₋₆alkyl; R²¹ is selected from H and -C₁₋₆alkyl; R²² is selected from H and -C₁₋₆alkyl; and R²³ is selected from H, -C₁₋₆alkyl, -CH₂COOH, -(CH₂)₂OH, -(CH₂)₂OCH₃, -(CH₂)₂N(CH₃)₂, -C₃₋₇cycloalkyl, and -(CH₂)₂-imidazolyl; or
15 R²² and R²³ are taken together to form azetidine, pyrrolidine, piperidine or morpholine, all optionally substituted with -OH or -CONH₂; and R²⁴ is selected from -C₁₋₆alkyl; -C₀₋₁alkylene-O-C₁₋₆alkyl; phenyl substituted with halo or -OCH₃; and pyridine.


13. The compound as claimed in any one of Claims 1 to 12, where R⁴ is absent or is selected from H; -OH; -C₁₋₆alkyl; -C₁₋₂alkylene-COOR³⁵; -CH₂OC(O)CH(R³⁶)NH₂; -CH₂CH(OH)CH₂OH; pyridinyl; and phenyl or benzyl optionally substituted with one or
25 more groups selected from halo, -COOR³⁵, -OCH₃, -OCF₃, and -SCF₃; and R³⁵ is H.

14. The compound as claimed in any one of Claims 1 to 12, where R⁴ is selected from -OCH₂OC(O)CH₃; -CH₂OP(O)(OH)₂; -C₁₋₂alkylene-COOR³⁵; and phenyl or benzyl substituted with at least one -COOR³⁵ group; where R³⁵ is selected from -C₁₋₆alkyl,
30 -C₁₋₃alkylene-C₆₋₁₀aryl, -C₁₋₃alkylene-C₁₋₉heteroaryl, -C₃₋₇cycloalkyl, -[(CH₂)₂O]₁₋₃CH₃,

-C₁₋₆alkylene-OC(O)R²⁵, -C₁₋₆alkylene-NR²⁷R²⁸, -C₁₋₆alkylene-C(O)R³³,
 -C₀₋₆alkylenemorpholinyl, -C₁₋₆alkylene-SO₂-C₁₋₆alkyl,



15. The compound as claimed in any one of Claims 1 to 12, where R⁴ is absent or is
 5 selected from H; -OH; -C₁₋₆alkyl; -C₁₋₂alkylene-COOR³⁵; -CH₂OC(O)CH(R³⁶)NH₂;
 -CH₂CH(OH)CH₂OH; pyridinyl; phenyl optionally substituted with one halo group; and
 benzyl optionally substituted with one or more groups selected from halo, -COOH, -OCH₃,
 -OCF₃, and -SCF₃; R³⁵ is H; and R³⁶ is -CH(CH₃)₂; or R³ and R⁴ are taken together to form
 -phenylene-O-(CH₂)₁₋₃- or -phenylene-O-CH₂-CHOH-CH₂-.
- 10 16. The compound as claimed in any one of Claims 1 to 15, where a is 0; or a is 1 and
 R⁵ is 3-chloro.
17. The compound as claimed in any one of Claims 1 to 16, where b is 0; or b is 1 and
 R⁶ is 3'-chloro, 3'-methyl, or 2'-methoxy; or b is 2 and R⁶ is 2'-fluoro-5'-chloro, 2',5'-
 dichloro, 2'-methyl-5'-chloro, or 3'-chloro-5'-hydroxy.
- 15 18. The compound as claimed in any one of Claims 1 to 17, where the methylene linker
 on the biphenyl is substituted with 2 methyl groups.
19. The compound of Claim 1, where
 R¹ is -OR⁷; R⁷ is selected from H, -C₁₋₈alkyl, -C₁₋₆alkylene-OC(O)R¹⁰,
 -C₀₋₆alkylenemorpholinyl, -C₁₋₆alkylene-SO₂-C₁₋₆alkyl, and



- 20 R¹⁰ is -O-C₃₋₇cycloalkyl; R³² is -CH₃; R⁸ is H; R⁹ is H;
 R² is H;

- X is selected from pyrazole, triazole, benzotriazole, furan, tetrazole, pyrazine,
 thiophene, oxazole, isoxazole, thiazole, oxadiazole, pyridazine, pyridine, pyrimidine,
 25 benzoxazole, pyridylimidazole, and pyridyltriazole;
 R³ is absent or is selected from H; halo; -C₀₋₅alkylene-OH; -NH₂; -C₁₋₆alkyl; -CF₃;
 -C₃₋₇cycloalkyl; -C₀₋₂alkylene-O-C₁₋₆alkyl; -C(O)R²⁰; -C₀₋₁alkylene-COOR²¹;

- C(O)NR²²R²³; -NHC(O)R²⁴; =O; -C(CH₃)=N(OH); phenyl optionally substituted with one or two groups independently selected from halo, -OH, -CF₃, -OCH₃, -NHC(O)CH₃, and phenyl; naphthalenyl; pyridinyl; pyrazinyl; pyrazolyl substituted with methyl; thiophenyl substituted with methyl or halo; furanyl; and -CH₂-morpholinyl;
- 5 R⁴ is absent or is selected from H; -OH; -C₁₋₆alkyl; -C₁₋₂alkylene-COOR³⁵; -CH₂OC(O)CH(R³⁶)NH₂; -CH₂CH(OH)CH₂OH; pyridinyl; phenyl optionally substituted with one halo group; and benzyl optionally substituted with one or more groups selected from halo, -COOH, -OCH₃, -OCF₃, and -SCF₃;
- or R³ and R⁴ are taken together to form -phenylene-O-(CH₂)₁₋₃- or -phenylenc-O-
- 10 CH₂-CHOH-CH₂-;
- a is 0; or a is 1 and R⁵ is 3-chloro;
- b is 0; or b is 1 and R⁶ is 3'-chloro, 3'-methyl, or 2'-methoxy; or b is 2 and R⁶ is 2'-fluoro-5'-chloro, 2',5'-dichloro, 2'-methyl-5'-chloro, or 3'-chloro-5'-hydroxy;
- R²⁰ is -C₁₋₆alkyl;
- 15 R²¹ is selected from H and -C₁₋₆alkyl;
- R²² is selected from H and -C₁₋₆alkyl; and R²³ is selected from H, -C₁₋₆alkyl, -CH₂COOH, -(CH₂)₂OH, -(CH₂)₂OCH₃, -(CH₂)₂N(CH₃)₂, -C₃₋₇cycloalkyl, and -(CH₂)₂-imidazolyl; or R²² and R²³ are taken together to form azetidine, pyrrolidine, piperidine or morpholine, all optionally substituted with -OH or -CONH₂;
- 20 R²⁴ is selected from -C₁₋₆alkyl; -O-C₁₋₆alkyl; -CH₂-O-C₁₋₆alkyl; phenyl substituted with halo or -OCH₃; and pyridinyl;
- R³⁵ is H;
- R³⁶ is -CH(CH₃)₂; and
- the methylene linker on the biphenyl is optionally substituted with 2 methyl groups.
- 25 20. The compound of Claim 1, where
- R¹ is -OR⁷;
- R² is H;
- X is selected from pyrazole, triazole, benzotriazole, isoxazole, pyridazine, pyrimidine, and pyridyltriazole;
- 30 R³ is selected from H; halo; -C₀₋₅alkylene-OH; -C₁₋₆alkyl; -CF₃; -C₃₋₇cycloalkyl; -C₀₋₂alkylene-O-C₁₋₆alkyl; -C(O)R²⁰; -C₀₋₁alkylene-COOR²¹; -C(O)NR²²R²³; -NHC(O)R²⁴; =O; -C(CH₃)=N(OH); phenyl optionally substituted with one or two groups independently selected from halo, -OH, and -OCH₃; pyridinyl; pyrazinyl; and thiophenyl substituted with

methyli or halo;

R⁴ is selected from H; -OH; -C₁₋₆alkyl; -C₁₋₂alkylene-COOR³⁵;

-CH₂CH(OH)CH₂OH; pyridinyl; and phenyl optionally substituted with one halo group; or

R^3 and R^4 are taken together to form -phenylene-O-(CH₂)₁₋₃-;

5 a is 0; or a is 1 and R⁵ is 3-chloro;

b is 0; or b is 1 and R⁶ is 3'-chloro, 3'-methyl, or 2'-methoxy; or b is 2 and R⁶ is 2'-fluoro-5'-chloro, 2',5'-dichloro, 2'-methyl-5'-chloro, or 3'-chloro-5'-hydroxy;

R²⁰ is -C₁₋₆alkyl;

 R^{21} is $\mathbb{H}$;


10 R²² is selected from H and -C₁₋₆alkyl; and R²³ is selected from -C₁₋₆alkyl, -(CH₂)₂OCH₃, and -C₃₋₇cycloalkyl; or R²² and R²³ are taken together to form azetidine, pyrrolidine, or piperidine, all optionally substituted with -OH or -CONH₂;

R^{24} is phenyl substituted with halo or $-OCH_3$;

 R^{35} is H; and


15 the methylene linker on the biphenyl is optionally substituted with 2 methyl groups.

21. A process for preparing a compound as claimed in any one of Claims 1 to 20, comprising the step of coupling a compound of formula 1 with a compound of formula 2:



to produce a compound of formula I; where P¹ is H or an amino-protecting group selected from *t*-butoxycarbonyl, trityl, benzyloxycarbonyl, 9-fluorenylmethoxycarbonyl, formyl, 20 trimethylsilyl, and *t*-butyldimethylsilyl; and where the process further comprises deprotecting the compound of formula I when P¹ is an amino protecting group.


22. An intermediate useful in the synthesis of a compound as claimed in any one of Claims 1 to 20, having formula 1:




(1),

where P^1 is H or an amino-protecting group selected from *t*-butoxycarbonyl, trityl, benzyloxycarbonyl, 9-fluorenylmethoxycarbonyl, formyl, trimethylsilyl, and *t*-butyldimethylsilyl; or a salt thereof.


23. A process for preparing a compound as claimed in any one of Claims 1 to 20, comprising the step of deprotecting a compound selected from:




(6),



(7),



(8), and




(9),

- or a salt thereof; where R^{1P} is selected from $-O-P^3$, $-NHP^2$, and $-NH(O-P^4)$; R^{3P} is selected from $-C_{0-5}alkylene-O-P^4$, $-C_{0-1}alkylene-COO-P^3$, and phenyl substituted with $-O-P^4$; R^{4P} is selected from $-O-P^4$, $-C_{1-2}alkylene-COO-P^3$; and phenyl or benzyl substituted with $-COO-P^3$; P^2 is an amino-protecting group selected from *t*-butoxycarbonyl, trityl, benzyloxycarbonyl, 9-fluorenylmethoxycarbonyl, formyl, trimethylsilyl, and *t*-butyldimethylsilyl; P^3 is a carboxy-protecting group selected from methyl, ethyl, *t*-butyl,

benzyl, *p*-methoxybenzyl, 9-fluorenylmethyl, trimethylsilyl, *t*-butyldimethylsilyl, and diphenylmethyl; and P⁴ is a hydroxyl-protecting group selected from -C₁₋₆alkyl, triC₁₋₆alkylsilyl, -C₁₋₆alkanoyl, benzoyl, benzyl, *p*-methoxybenzyl, 9-fluorenylmethyl, and diphenylmethyl.

- 5 24. An intermediate useful in the synthesis of a compound as claimed in any one of Claims 1 to 20, selected from:



- or a salt thereof; where R^{1P} is selected from -O-P³, -NHP², and -NH(O-P⁴); R^{3P} is selected from -C₀₋₅alkylene-O-P⁴, -C₀₋₁alkylene-COO-P³, and phenyl substituted with -O-P⁴; R^{4P} is selected from -O-P⁴, -C₁₋₂alkylene-COO-P³; and phenyl or benzyl substituted with -COO-P³; P² is an amino-protecting group selected from *t*-butoxycarbonyl, trityl, benzyloxycarbonyl, 9-fluorenylmethoxycarbonyl, formyl, trimethylsilyl, and *t*-butyldimethylsilyl; P³ is a carboxy-protecting group selected from methyl, ethyl, *t*-butyl, benzyl, *p*-methoxybenzyl, 9-fluorenylmethyl, trimethylsilyl, *t*-butyldimethylsilyl, and diphenylmethyl; and P⁴ is a hydroxyl-protecting group selected from -C₁₋₆alkyl, triC₁₋₆alkylsilyl, -C₁₋₆alkanoyl, benzoyl, benzyl, *p*-methoxybenzyl, 9-fluorenylmethyl, and diphenylmethyl.

25. A pharmaceutical composition comprising a compound as claimed in any one of Claims 1 to 20 and a pharmaceutically acceptable carrier.

26. The pharmaceutical composition of Claim 25, further comprising a therapeutic agent selected from adenosine receptor antagonists, α -adrenergic receptor antagonists, β_1 -adrenergic receptor antagonists, β_2 -adrenergic receptor agonists, dual-acting β -adrenergic receptor antagonist/ α_1 -receptor antagonists, advanced glycation end product breakers, aldosterone antagonists, aldosterone synthase inhibitors, aminopeptidase N inhibitors, androgens, angiotensin-converting enzyme inhibitors and dual-acting angiotensin-converting enzyme/neprilysin inhibitors, angiotensin-converting enzyme 2 activators and stimulators, angiotensin-II vaccines, anticoagulants, anti-diabetic agents, antidiarrheal agents, anti-glaucoma agents, anti-lipid agents, antinociceptive agents, anti-thrombotic agents, AT_1 receptor antagonists and dual-acting AT_1 receptor antagonist/neprilysin inhibitors and multifunctional angiotensin receptor blockers, bradykinin receptor antagonists, calcium channel blockers, chymase inhibitors, digoxin, diuretics, dopamine agonists, endothelin converting enzyme inhibitors, endothelin receptor antagonists, HMG-CoA reductase inhibitors, estrogens, estrogen receptor agonists and/or antagonists, monoamine reuptake inhibitors, muscle relaxants, natriuretic peptides and their analogs, natriuretic peptide clearance receptor antagonists, neprilysin inhibitors, nitric oxide donors, non-steroidal anti-inflammatory agents, N-methyl d-aspartate receptor antagonists, opioid receptor agonists, phosphodiesterase inhibitors, prostaglandin analogs, prostaglandin receptor agonists, renin inhibitors, selective serotonin reuptake inhibitors, sodium channel blocker, soluble guanylate cyclase stimulators and activators, tricyclic antidepressants, vasopressin receptor antagonists, and combinations thereof.
27. The pharmaceutical composition of Claim 26, wherein the therapeutic agent is an AT_1 receptor antagonist.
28. A compound as claimed in any one of Claims 1 to 20, for use in therapy.
29. A compound as claimed in Claim 28, for use in treating hypertension, heart failure, or renal disease.
30. The use of a compound as claimed in any one of Claims 1 to 20, for the manufacture of a medicament for treating hypertension, heart failure, or renal disease.