
Abstract

2-Heterocyclicethylamine derivatives and their use as pharmaceutical

Compounds of formula I,



I

wherein

A represents C=C, S or O;

Q represents a 5- or 6-membered unsaturated heterocyclic ring having a nitrogen atom in the position adjacent to the point of attachment, 0-3 further heteroatoms selected from N, O and S, and substituents R^6 and R^7;

R^1 represents H or C_{1-6} alkyl;

R^2 represents H, C_{1-6} alkyl, C_{3-6} alkenyl, C_{3-6} alkynyl, C_{3-6} cycloalkyl or NH_2CH_2CO-;

in addition, R^1 and R^2 taken together may form a C_{3-5} alkylene chain;

R^3 represents H, C_{1-6} alkyl, C_{3-6} alkenyl, C_{3-6} alkynyl;

R^4 and R^5 independently represent H, OH, C_{1-6} alkoxy, C_{1-6} alkyl, halogen, trifluoromethyl or NR^8R^9;

R^6 and R^7 independently represent H, OH, C_{1-6} alkoxy, C_{1-6} alkyl, halogen, trifluoromethyl, C_{1-6} hydroxyalkyl, amidino, $CONH_2$ or NR^8R^9;

in addition, R^6 and R^7 may independently represent O when substituted on N;

R^8 and R^9 independently represent H or C_{1-6} alkyl;

with various provisos;

and pharmaceutically acceptable derivatives thereof;

are useful as pharmaceuticals, in particular in the treatment of neurological disorders.