with a compound or salt of the present invention, thereby inhibiting the replication of HCV virus in the cells.

In addition, the present invention features methods of using the compounds or salts of the present invention, or compositions comprising the same, to treat HCV infection. The methods comprise administering a compound or salt of the present invention, or a pharmaceutical composition comprising the same, to a patient in need thereof, thereby reducing the blood or tissue level of HCV virus in the patient.

The present invention also features use of the compounds or salts of the present invention for the manufacture of medicaments for the treatment of HCV infection.

Furthermore, the present invention features processes of making the compounds or salts of the invention.

The present invention also features a compound, which is dimethyl (2S,2'S)-1,1'-((2S,2'S)-2,2'-(4,4'-((2S,5S)-1-(4-*tert*-butylphenyl)pyrrolidine-2,5-diyl)bis(4,1-phenylene))bis(azanediyl)bis(oxomethylene)bis(pyrrolidine-2,1-diyl))bis(3-methyl-1-oxobutane-2,1-diyl)dicarbamate.


The present invention also features a pharmaceutical composition comprising the compound as defined herein or a pharmaceutically acceptable salt thereof.

The present invention also features the use of the compound as defined herein, or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for treating a patient infected with HCV.

Other features, objects, and advantages of the present invention are apparent in the detailed description that follows. It should be understood, however, that the detailed description, while indicating preferred embodiments of the invention, are given by way of illustration only, not limitation. Various changes and modifications within the scope of the invention will become apparent to those skilled in the art from the detailed description.

DETAILED DESCRIPTION

The present invention features compounds having Formula I, and pharmaceutically acceptable salts thereof,



I

wherein:

X is C_3-C_{12}carbocycle or 3- to 12-membered heterocycle, and is optionally substituted with one or more R_A;

L_1 and L_2 are each independently selected from bond; or C_1-C_6alkylene, C_2-C_6alkenylene or C_2-C_6alkynylene, each of which is independently optionally substituted at each occurrence with one or more R_L;

L_3 is bond or $-L_S-K-L_S'-$, wherein K is selected from bond, $-O-$, $-S-$, $-N(R_B)-$, $-C(O)-$, $-S(O)_2-$, $-S(O)-$, $-OS(O)-$, $-OS(O)_2-$, $-S(O)_2O-$, $-S(O)O-$, $-C(O)O-$, $-OC(O)-$, $-OC(O)O-$, $-C(O)N(R_B)-$, $-N(R_B)C(O)-$, $-N(R_B)C(O)O-$, $-OC(O)N(R_B)-$, $-N(R_B)S(O)-$, $-N(R_B)S(O)_2-$, $-S(O)N(R_B)-$, $-S(O)_2N(R_B)-$, $-C(O)N(R_B)C(O)-$, $-N(R_B)C(O)N(R_B')-$, $-N(R_B)SO_2N(R_B')-$, or $-N(R_B)S(O)N(R_B')-$;