

(51) International Patent Classification:
C07D 499/87 (2006.01)(21) International Application Number:
PCT/EP2015/060733(22) International Filing Date:
14 May 2015 (14.05.2015)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
1408649.0 15 May 2014 (15.05.2014) GB(71) Applicant: ALLECRA THERAPEUTICS SAS [FR/FR];
13 rue Village Neuf, F-68300 Saint Louise (FR).

(72) Inventors: FAINI, Andrea; Via Cascella, 6, I-20138 Milano (IT). FORZATTI, Marco; Via Molise, 15, I-20900 Monza (IT). FOGLIATO, Giovanni; Via Mazzini, 22, I-24030 Barzana (BG) (IT). BIONDI, Stefano; Via Copernico, 22, I-20016 Pero (MI) (IT).

(74) Agents: GILANI, Anwar et al.; Venner Shipley LLP, Byron House, Cambridge Business Park, Cowley Road, Cambridge, Cambridgeshire CB4 0WZ (GB).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).


Published:

— without international search report and to be republished upon receipt of that report (Rule 48.2(g))


(54) Title: METHOD



Formula (IIIa)



(IIa)



(VIII)

(57) Abstract: A method of forming a compound of formula (IIIa); wherein: R¹ in each occurrence is independently selected from H, halogen, amino, C₁₋₅ alkyl, C₁₋₅ alkenyl and C₁₋₅ alkynyl; R² in each occurrence is independently selected from H, halogen, amino, C₁₋₅ alkyl, C₁₋₅ alkenyl and C₁₋₅ alkynyl; R⁴ is C₁₋₅ alkyl; and R⁶ is a C₁₋₅ fluoroalkyl; and PG is a protecting group, the method comprising the step of reacting a compound of formula (IIa) with a compound of formula (VIII).