

CHLORIN DERIVATIVE USEFUL IN PHOTODYNAMIC THERAPY AND DIAGNOSIS**Technical field**

5 The present invention relates to chlorin e4 sodium and a process for its preparation. Chlorin e4 sodium is suitable for use in photodynamic therapy, cytoluminescent therapy and photodynamic diagnosis, for example for treating or detecting a tumour. The present invention also relates to a pharmaceutical composition comprising chlorin e4 sodium, to the use of chlorin e4 sodium in the manufacture
10 of a phototherapeutic or photodiagnostic agent, and to a method of photodynamic therapy, cytoluminescent therapy or photodynamic diagnosis, for example for treating or detecting a tumour.

Background art

15

Porphyrins and their derivatives are known photosensitive chemical compounds, which can absorb light photons and emit them at higher wavelengths. There are many applications for such unique properties and PDT (photodynamic therapy) is one of them.

20

During light activation, a molecule goes from the ground state (S_0) to a singlet excited state (S_1), ($S_0 \rightarrow S_1$ transition), and then falls back down to the ground state (S_0), ($S_1 \rightarrow S_0$ transition), emitting light at higher wavelengths in the form of fluorescence. There is another energy state known as the triplet state (T_1). The
25 triplet state (T_1) cannot be efficiently populated by direct excitation. However, intersystem crossing $S_1 \rightarrow T_1$ occurs isoenergetically followed by deactivation to T_1 . The $S_1 \rightarrow T_1$ change is formally spin-forbidden, but it is a downhill process and in some molecules intersystem crossing occurs so efficiently that fluorescence is quenched. Such compounds are useful for generating excited triplets by excitation
30 transfer (a phenomenon known as sensitization) in molecules where the intersystem crossing is minimal.

Figure 1 shows a modified Jablonski diagram for a typical photosensitizer, outlining the principle photophysical processes of interest.

At the triplet state (T_1) molecules become very unstable and can react with
5 molecular oxygen ($^3O_2$), splitting the latter into two singlet oxygens ($^1O_2$). Singlet oxygen ($^1O_2$) is known as a “scavenger” and can destroy abnormal biological cells and remove free radicals.


Such unique properties make many photosensitive molecules good photosensitizers
10 for PDT.

Presently, there are two generations of photosensitizers for PDT. The first generation comprises hem porphyrins (blood derivatives), and the second for the most part are chlorophyll derivatives. The later compounds are known as chlorins
15 and bacteriochlorins.

Chlorins show photophysical properties similar to those of the porphyrin systems, but have an enhanced, red-shifted Q band (670nm) which makes chlorin-containing systems better candidates for PDT. *Chlorophyll a*, the green photosynthetic pigment,
20 is one prototype of the chlorin class of natural products. It or its derivatives can be extracted from certain *Spirulina Platensis* species without any contamination with *chlorophyll b*, thus avoiding a tedious chromatographic separation.


A water-soluble chlorophyll derivative, known as tri-sodium copper(II) chlorin e6,
25 has been used in the treatment of various human ailments with no evidence of toxicity, skin sensitization or other serious side effects.


Chlorin e4 was studied recently and was shown to display good photosensitive activity. It was indicated that chlorin e4 has a protective effect against
30 indomethacin-induced gastric lesions in rats and TAA- or CCl_4 -induced acute liver injuries in mice. It was therefore suggested that chlorin e4 may be a promising new drug candidate for anti-gastrelcosis and liver injury protection. WO 2009/040411 suggests the use of a chlorin e4 zinc complex in photodynamic therapy.



However, there is an ongoing need for better photosensitizers. There is a need for compounds that have a high singlet oxygen quantum yield and for compounds that have a strong photosensitizing ability, preferably in organic and aqueous media. There is also a need for compounds that have a high fluorescence quantum yield.

Summary of the inventionIn one aspect, the present invention refers to chlorin e4 sodium, comprising less than 10% of other stereoisomers. In another aspect, the present invention refers to chlorin e4 sodium, comprising chlorin e4 and sodium in a ratio of 1:2 and comprising less than 10% of other stereoisomers. In another aspect, the present invention refers to chlorin e4 sodium, having the structure:






and comprising less than 10% of other stereoisomers. In another aspect, the present invention refers to a process of preparing chlorin e4 sodium comprising less than 10% of other stereoisomers, comprising the step of reacting chlorin e4 with a sodium compound. In another aspect, the present invention refers to the chlorin e4 sodium of the present invention, for use in the treatment of atherosclerosis; multiple sclerosis; diabetes; diabetic retinopathy; arthritis; rheumatoid arthritis; a fungal, viral, chlamydial, bacterial, nanobacterial or parasitic infectious disease; HIV; Aids; infection with sars virus, Asian (chicken) flu virus, herpes simplex or herpes zoster; hepatitis; viral hepatitis; a cardiovascular disease; coronary artery stenosis; carotid artery stenosis; intermittent claudication; a dermatological condition; acne; psoriasis; a disease characterised by benign or malignant cellular hyperproliferation or by areas of neovascularisation; a benign or malignant tumour; early cancer; cervical dysplasia; soft tissue sarcoma; a germ cell tumour; retinoblastoma; age-related macular degeneration; lymphoma; Hodgkin's lymphoma; head and neck cancer; oral or mouth cancer; or cancer of the blood, prostate, cervix, uterus, vaginal or other female adnexa, breast, naso-pharynx, trachea, larynx, bronchi, bronchioles, lung, hollow organs, esophagus, stomach, bile duct, intestine, colon, colorectum, rectum, bladder, ureter, kidney, liver, gall bladder, spleen, brain, lymphatic system, bones, skin or pancreas. In still another aspect, the present invention refers to a use of chlorin e4 sodium of the present invention in the manufacture of a medicament for the treatment of a benign or malignant tumour;

early cancer; cervical dysplasia; soft tissue sarcoma; a germ cell tumour; retinoblastoma; age-related macular degeneration; lymphoma; Hodgkin's lymphoma; head and neck cancer; oral or mouth cancer; or cancer of the blood, prostate, cervix, uterus, vaginal or other female adnexa, breast, naso-pharynx, trachea, larynx, bronchi, bronchioles, lung, hollow organs, esophagus, stomach, bile duct, intestine, colon, colorectum, rectum, bladder, ureter, kidney, liver, gall bladder, spleen, brain, lymphatic system, bones, skin or pancreas.

Further described herein, in a first example there is provided chlorin e4 sodium. Preferably the chlorin e4 sodium comprises chlorin e4 and sodium in a ratio of 1:2.


10 Preferably the chlorin e4 sodium has the structure:



15

20


Preferably the chlorin e4 sodium has the structure:




The skilled person will appreciate that the negative charges on the chlorin structures
5 shown in the above representations are delocalized over the conjugated system.

Chlorin e4 sodium has two chiral centres. The chlorin e4 sodium described herein
is preferably substantially enantiomerically pure, which means that the chlorin e4
sodium comprises less than 10% of other stereoisomers, preferably less than 5%,
10 preferably less than 3%, preferably less than 2%, preferably less than 1%.

A second example provides a process of preparing chlorin e4 sodium, comprising
the step of reacting chlorin e4 with a sodium compound. Preferably the chlorin e4
sodium obtained comprises chlorin e4 and sodium in a ratio of 1:2. Preferably the
15 chlorin e4 sodium obtained has the structure:



Preferably the chlorin e4 sodium obtained has the structure:



- 5 The skilled person will appreciate that the negative charges on the chlorin structures shown in the above representations are delocalized over the conjugated system.

Preferably the sodium compound is sodium hydroxide, sodium carbonate, a carboxylic acid sodium salt (such as sodium stearate or sodium acetate), a sodium
 10 halide (such as sodium fluoride) or sodium hydride. Preferably the sodium compound is sodium hydroxide, a carboxylic acid sodium salt (such as sodium stearate or sodium acetate), a sodium halide (such as sodium fluoride) or sodium

hydride. Preferably the sodium compound is sodium hydroxide or sodium carbonate. More preferably the sodium compound is sodium hydroxide.

5 Optionally chlorin e4 is reacted with the sodium compound in the presence of a base. Preferably the base is an amine base, preferably an aromatic amine base such as thiamine pyrophosphate or DMAP. If used, the base is preferably thiamine pyrophosphate.

10 Preferably chlorin e4, the sodium compound and the optional base are used in a molar ratio of 0.15-1 : 45-60 : 0-12.

15 Preferably chlorin e4 is reacted with the sodium compound in the presence of a solvent. Preferably the solvent is selected from an alcohol (such as methanol, ethanol, n-propanol or isopropanol), water, DMF, chloroform, DMSO, toluene or a mixture thereof. Preferably the solvent is selected from methanol, ethanol, water or a mixture thereof. Preferably the solvent is selected from methanol, water or a mixture thereof. Preferably the solvent is a mixture of methanol and water.

20 Preferably the reaction is carried out at a temperature of 60°C or less. Preferably the reaction is carried out at a temperature in the range of 25-60°C.

Preferably the reaction is carried out over 1 to 12 hours, preferably over 3 to 8 hours.

25 The chlorin e4 sodium obtained may be purified by filtering, for example through a 45µm Millipore filter. Alternatively or additionally, the chlorin e4 sodium obtained may be purified by chromatography, for example by flash chromatography on a silica gel column. The chlorin e4 sodium may be isolated by lyophilisation.

30 A third example provides chlorin e4 sodium preparable by a process according to the second example. The third example also provides chlorin e4 sodium when prepared by a process according to the second example.

A fourth example provides a pharmaceutical composition comprising chlorin e4 sodium according to the first or third example and a pharmaceutically acceptable carrier or diluent.

- 5 Preferably the chlorin e4 sodium according to the first or third example is suitable for use in medicine.

Preferably the chlorin e4 sodium according to the first or third example and the pharmaceutical composition according to the fourth example are suitable for use in
10 photodynamic therapy or cytoluminescent therapy.

Preferably the chlorin e4 sodium according to first or third example and the pharmaceutical composition according to the fourth example are suitable for the treatment of atherosclerosis; multiple sclerosis; diabetes; diabetic retinopathy;
15 arthritis; rheumatoid arthritis; a fungal, viral, chlamydial, bacterial, nanobacterial or parasitic infectious disease; HIV; Aids; infection with sars virus, Asian (chicken) flu virus, herpes simplex or herpes zoster; hepatitis; viral hepatitis; a cardiovascular disease; coronary artery stenosis; carotid artery stenosis; intermittent claudication; a dermatological condition; acne; psoriasis; a disease characterised by benign or
20 malignant cellular hyperproliferation or by areas of neovascularisation; a benign or malignant tumour; early cancer; cervical dysplasia; soft tissue sarcoma; a germ cell tumour; retinoblastoma; age-related macular degeneration; Hodgkin's lymphoma; or cancer of the blood, prostate, cervix, uterus, vaginal or other female adnexa, breast, naso-pharynx, trachea, larynx, bronchi, bronchioles, lung, hollow organs, esophagus,
25 stomach, bile duct, intestine, colon, colorectum, rectum, bladder, ureter, kidney, liver, gall bladder, spleen, brain, lymphatic system or bones.

Preferably the chlorin e4 sodium according to the first or third example and the pharmaceutical composition according to the fourth

example are suitable for the treatment of a disease characterised by benign or malignant cellular hyperproliferation or by areas of neovascularisation.

5 Preferably the chlorin e4 sodium according to the first or third example and the pharmaceutical composition according to the fourth example are suitable for the treatment of a benign or malignant tumour.

10 Preferably the chlorin e4 sodium according to the first or third example and the pharmaceutical composition according to the fourth example are suitable for the treatment of early cancer; cervical dysplasia; soft tissue sarcoma; a germ cell tumour; retinoblastoma; age-related macular degeneration; lymphoma; Hodgkin's lymphoma; head and neck cancer; oral or mouth cancer; or cancer of the blood, prostate, cervix, uterus, vaginal or other female adnexa, breast, naso-pharynx, trachea, larynx, bronchi, bronchioles, lung, hollow organs, esophagus, stomach, bile duct, intestine, colon, colorectum, rectum, bladder, ureter, kidney, liver, gall bladder, spleen, brain, lymphatic system, bones, skin or pancreas. Preferably they are suitable for the treatment of early cancer; cervical dysplasia; soft tissue sarcoma; a germ cell tumour; retinoblastoma; age-related macular degeneration; Hodgkin's lymphoma; or cancer of the blood, prostate, cervix, uterus, vaginal or other female adnexa, breast, naso-pharynx, trachea, larynx, bronchi, bronchioles, lung, hollow organs, esophagus, stomach, bile duct, intestine, colon, colorectum, rectum, bladder, ureter, kidney, liver, gall bladder, spleen, brain, lymphatic system or bones. Preferably they are suitable for the treatment of lymphoma; Hodgkin's lymphoma; oral or mouth cancer; or cancer of the prostate, cervix, vaginal or other female adnexa, breast, naso-pharynx, lung, esophagus, kidney, liver, brain, lymphatic system, bones, skin or pancreas. Preferably they are suitable for the treatment of lymphoma; oral or mouth cancer; or cancer of the prostate, cervix, vaginal or other female adnexa, breast, naso-pharynx, lung, esophagus, bones or skin.

30 Preferably the chlorin e4 sodium according to the first or third example and the pharmaceutical composition according to the fourth example are suitable for use in photodynamic diagnosis.

Preferably the chlorin e4 sodium according to the first or third example and the pharmaceutical composition according to the fourth example are suitable for the detection of atherosclerosis; multiple sclerosis; diabetes; diabetic retinopathy; arthritis; rheumatoid arthritis; a fungal, viral, chlamydial, bacterial, nanobacterial or parasitic infectious disease; HIV; Aids; infection with sars virus, Asian (chicken) flu virus, herpes simplex or herpes zoster; hepatitis; viral hepatitis; a cardiovascular disease; coronary artery stenosis; carotid artery stenosis; intermittent claudication; a dermatological condition; acne; psoriasis; a disease characterised by benign or malignant cellular hyperproliferation or by areas of neovascularisation; a benign or malignant tumour; early cancer; cervical dysplasia; soft tissue sarcoma; a germ cell tumour; retinoblastoma; age-related macular degeneration; Hodgkin's lymphoma; or cancer of the blood, prostate, cervix, uterus, vaginal or other female adnexa, breast, naso-pharynx, trachea, larynx, bronchi, bronchioles, lung, hollow organs, esophagus, stomach, bile duct, intestine, colon, colorectum, rectum, bladder, ureter, kidney, liver, gall bladder, spleen, brain, lymphatic system or bones.

Preferably the chlorin e4 sodium according to the first or third example and the pharmaceutical composition according to the fourth example are suitable for the detection of an area that is affected by benign or malignant cellular hyperproliferation or by neovascularisation.

Preferably the chlorin e4 sodium according to the first or third example and the pharmaceutical composition according to the fourth example are suitable for the detection of a benign or malignant tumour.

Preferably the chlorin e4 sodium according to the first or third example and the pharmaceutical composition according to the fourth example are suitable for the detection of early cancer; cervical dysplasia; soft tissue sarcoma; a germ cell tumour; retinoblastoma; age-related macular degeneration; lymphoma; Hodgkin's lymphoma; head and neck cancer; oral or mouth cancer; or cancer of the blood, prostate, cervix, uterus, vaginal or other

female adnexa, breast, naso-pharynx, trachea, larynx, bronchi, bronchioles, lung, hollow organs, esophagus, stomach, bile duct, intestine, colon, colorectum, rectum, bladder, ureter, kidney, liver, gall bladder, spleen, brain, lymphatic system, bones, skin or pancreas. Preferably they are suitable for the detection of early cancer; cervical dysplasia; soft tissue sarcoma; a germ cell tumour; retinoblastoma; age-related macular degeneration; Hodgkin's lymphoma; or cancer of the blood, prostate, cervix, uterus, vaginal or other female adnexa, breast, naso-pharynx, trachea, larynx, bronchi, bronchioles, lung, hollow organs, esophagus, stomach, bile duct, intestine, colon, colorectum, rectum, bladder, ureter, kidney, liver, gall bladder, spleen, brain, lymphatic system or bones. Preferably they are suitable for the detection of lymphoma; Hodgkin's lymphoma; oral or mouth cancer; or cancer of the prostate, cervix, vaginal or other female adnexa, breast, naso-pharynx, lung, esophagus, kidney, liver, brain, lymphatic system, bones, skin or pancreas. Preferably they are suitable for the detection of lymphoma; oral or mouth cancer; or cancer of the prostate, cervix, vaginal or other female adnexa, breast, naso-pharynx, lung, esophagus, bones or skin.

Preferably the chlorin e4 sodium according to the first or third example and the pharmaceutical composition according to the fourth example are suitable for the fluorescent or phosphorescent detection of the diseases listed above, preferably for the fluorescent or phosphorescent detection and quantification of the said diseases.

Preferably the chlorin e4 sodium according to the first or third example and the pharmaceutical composition according to the fourth example are adapted for administration simultaneous with or prior to administration of irradiation or sound, preferably for administration prior to administration of irradiation.

If the chlorin e4 sodium according to the first or third example or the pharmaceutical composition according to the fourth example are for use in photodynamic therapy or cytoluminescent therapy, then they are preferably adapted for administration 5 to 100 hours before the

irradiation, preferably 6 to 72 hours before the irradiation, preferably 24 to 48 hours before the irradiation.

If the chlorin e4 sodium according to the first or third example or the
5 pharmaceutical composition according to the fourth example are for use in
photodynamic diagnosis, then they are preferably adapted for administration 3 to 60
hours before the irradiation, preferably 8 to 40 hours before the irradiation.

Preferably the irradiation used in the photodynamic therapy, cytoluminescent
10 therapy or photodynamic diagnosis is electromagnetic radiation with a wavelength
in the range of from 500nm to 1000nm, preferably from 600nm to 790nm. The
electromagnetic radiation may be administered for about 5-60 minutes, preferably
for about 15-20 minutes, at about 0.1-5W, preferably at about 1W. In one example,
two sources of electromagnetic radiation are used (for example a laser light and an
15 LED light), one source with a wavelength in the range of from 600nm to 690nm,
preferably about 660nm, the other source with a wavelength in the range of from
700nm to 790nm, preferably about 760nm. In another example, two sources of
electromagnetic radiation are used (for example a laser light and an LED light), both
sources adapted to provide irradiation with a wavelength in the range of from
20 600nm to 690nm, preferably about 660nm, and irradiation with a wavelength in the
range of from 700nm to 790nm, preferably about 760nm. In another example, the
irradiation may be provided by a prostate, anal, vaginal, mouth and nasal device for
insertion into a body cavity. In another example, the irradiation may be provided by
interstitial light activation, for example, using a fine needle to insert an optical fibre
25 laser into the lung, liver, lymph nodes or breast. In another example, the
irradiation may be provided by endoscopic light activation, for example, for
delivering light to the lung, stomach, colon, bladder or neck.

The pharmaceutical composition according to the fourth example may be in a form
30 suitable for oral, parental (including intravenous,

subcutaneous, intramuscular, intradermal, intratracheal, intraperitoneal, intraarticular, intraabdominal, intracranial and epidural), transdermal, airway (aerosol), rectal, vaginal or topical (including buccal, mucosal and sublingual) administration. The pharmaceutical composition may also be in a form suitable for
5 administration by enema or for administration by injection into a tumour. Preferably the pharmaceutical composition is in a form suitable for oral, parental or airway administration, preferably in a form suitable for oral or parental administration, preferably in a form suitable for oral administration.

10 In one preferred example, the pharmaceutical composition is in a form suitable for oral administration. Preferably the pharmaceutical composition is provided in the form of a tablet, capsule, hard or soft gelatine capsule, caplet, troche or lozenge, as a powder or granules, or as an aqueous solution, suspension or dispersion. More preferably the pharmaceutical composition is provided in the form of an aqueous
15 solution, suspension or dispersion for oral administration, or alternatively in the form of a freeze-dried powder which can be mixed with water before administration to provide an aqueous solution, suspension or dispersion for oral administration. Preferably the pharmaceutical composition is in a form suitable for providing 0.01 to 10 mg/kg/day of the chlorin e4 sodium, preferably 0.1 to 5 mg/kg/day,
20 preferably about 3 mg/kg/day.

In another preferred example, the pharmaceutical composition is in a form suitable for parental administration. Preferably the pharmaceutical composition is in a form suitable for intravenous administration. Preferably the pharmaceutical composition
25 is provided in the form of an aqueous solution for parental administration, or alternatively in the form of a freeze-dried powder which can be mixed with water before administration to provide an aqueous solution for parental administration. Preferably the pharmaceutical composition is an aqueous solution or suspension having a pH of from 6 to 8.5.

30

In another preferred example, the pharmaceutical composition is in a form suitable for airway administration. Preferably the pharmaceutical composition is provided in the form of an aqueous solution, suspension or dispersion for airway

administration, or alternatively in the form of a freeze-dried powder which can be mixed with water before administration to provide an aqueous solution, suspension or dispersion for airway administration. Preferably the pharmaceutical composition is in a form suitable for providing 0.01 to 10 mg/kg/day of the chlorin e4 sodium, preferably 0.1 to 5 mg/kg/day, preferably about 3 mg/kg/day.

A fifth example provides use of chlorin e4 sodium according to the first or third example in the manufacture of a medicament for the treatment of atherosclerosis; multiple sclerosis; diabetes; diabetic retinopathy; arthritis; rheumatoid arthritis; a fungal, viral, chlamydial, bacterial, nanobacterial or parasitic infectious disease; HIV; Aids; infection with sars virus, Asian (chicken) flu virus, herpes simplex or herpes zoster; hepatitis; viral hepatitis; a cardiovascular disease; coronary artery stenosis; carotid artery stenosis; intermittent claudication; a dermatological condition; acne; psoriasis; a disease characterised by benign or malignant cellular hyperproliferation or by areas of neovascularisation; a benign or malignant tumour; early cancer; cervical dysplasia; soft tissue sarcoma; a germ cell tumour; retinoblastoma; age-related macular degeneration; Hodgkin's lymphoma; or cancer of the blood, prostate, cervix, uterus, vaginal or other female adnexa, breast, naso-pharynx, trachea, larynx, bronchi, bronchioles, lung, hollow organs, esophagus, stomach, bile duct, intestine, colon, colorectum, rectum, bladder, ureter, kidney, liver, gall bladder, spleen, brain, lymphatic system or bones.

The fifth example also provides use of chlorin e4 sodium according to the first or third example in the manufacture of a phototherapeutic agent for use in photodynamic therapy or cytoluminescent therapy. Preferably the phototherapeutic agent is suitable for the treatment of atherosclerosis; multiple sclerosis; diabetes; diabetic retinopathy; arthritis; rheumatoid arthritis; a fungal, viral, chlamydial, bacterial, nanobacterial or parasitic infectious disease; HIV; Aids; infection with sars virus, Asian (chicken) flu virus, herpes simplex or herpes zoster; hepatitis; viral hepatitis; a cardiovascular disease; coronary artery stenosis; carotid artery stenosis; intermittent claudication; a dermatological condition; acne; psoriasis; a disease characterised by benign or malignant cellular hyperproliferation or by areas of neovascularisation; a benign or

malignant tumour; early cancer; cervical dysplasia; soft tissue sarcoma; a germ cell tumour; retinoblastoma; age-related macular degeneration; Hodgkin's lymphoma; or cancer of the blood, prostate, cervix, uterus, vaginal or other female adnexa, breast, naso-pharynx, trachea, larynx, bronchi, bronchioles, lung, hollow organs, esophagus, stomach, bile duct, intestine, colon, colorectum, rectum, bladder, ureter, kidney, liver, gall bladder, spleen, brain, lymphatic system or bones.

Preferably the medicament or the phototherapeutic agent of the fifth example is suitable for the treatment of a disease characterised by benign or malignant cellular hyperproliferation or by areas of neovascularisation.

Preferably the medicament or the phototherapeutic agent of the fifth example is suitable for the treatment of a benign or malignant tumour.

Preferably the medicament or the phototherapeutic agent of the fifth example is suitable for the treatment of early cancer; cervical dysplasia; soft tissue sarcoma; a germ cell tumour; retinoblastoma; age-related macular degeneration; lymphoma; Hodgkin's lymphoma; head and neck cancer; oral or mouth cancer; or cancer of the blood, prostate, cervix, uterus, vaginal or other female adnexa, breast, naso-pharynx, trachea, larynx, bronchi, bronchioles, lung, hollow organs, esophagus, stomach, bile duct, intestine, colon, colorectum, rectum, bladder, ureter, kidney, liver, gall bladder, spleen, brain, lymphatic system, bones, skin or pancreas. Preferably it is suitable for the treatment of early cancer; cervical dysplasia; soft tissue sarcoma; a germ cell tumour; retinoblastoma; age-related macular degeneration; Hodgkin's lymphoma; or cancer of the blood, prostate, cervix, uterus, vaginal or other female adnexa, breast, naso-pharynx, trachea, larynx, bronchi, bronchioles, lung, hollow organs, esophagus, stomach, bile duct, intestine, colon, colorectum, rectum, bladder, ureter, kidney, liver, gall bladder, spleen, brain, lymphatic system or bones. Preferably it is suitable for the treatment of lymphoma; Hodgkin's lymphoma; oral or mouth cancer; or cancer of the prostate, cervix, vaginal or other female adnexa, breast, naso-pharynx, lung, esophagus, kidney, liver, brain, lymphatic system, bones, skin or pancreas. Preferably it is suitable for the

treatment of lymphoma; oral or mouth cancer; or cancer of the prostate, cervix, vaginal or other female adnexa, breast, naso-pharynx, lung, esophagus, bones or skin.

- 5 The fifth example also provides use of chlorin e4 sodium according to the first or third example in the manufacture of a photodiagnostic agent for use in photodynamic diagnosis.

10 Preferably the photodiagnostic agent of the fifth example is suitable for the detection of atherosclerosis; multiple sclerosis; diabetes; diabetic retinopathy; arthritis; rheumatoid arthritis; a fungal, viral, chlamydial, bacterial, nanobacterial or parasitic infectious disease; HIV; Aids; infection with sars virus, Asian (chicken) flu virus, herpes simplex or herpes zoster; hepatitis; viral hepatitis; a cardiovascular disease; coronary artery stenosis; carotid artery stenosis; intermittent claudication; a
15 dermatological condition; acne; psoriasis; a disease characterised by benign or malignant cellular hyperproliferation or by areas of neovascularisation; a benign or malignant tumour; early cancer; cervical dysplasia; soft tissue sarcoma; a germ cell tumour; retinoblastoma; age-related macular degeneration; Hodgkin's lymphoma; or cancer of the blood, prostate, cervix, uterus, vaginal or other female adnexa, breast,
20 naso-pharynx, trachea, larynx, bronchi, bronchioles, lung, hollow organs, esophagus, stomach, bile duct, intestine, colon, colorectum, rectum, bladder, ureter, kidney, liver, gall bladder, spleen, brain, lymphatic system or bones.

25 Preferably the photodiagnostic agent of the fifth example is suitable for the detection of an area that is affected by benign or malignant cellular hyperproliferation or by neovascularisation.

Preferably the photodiagnostic agent of the fifth example is suitable for the detection of a benign or malignant tumour.

30

Preferably the photodiagnostic agent of the fifth example is suitable for the detection of early cancer; cervical dysplasia; soft tissue sarcoma; a germ cell tumour; retinoblastoma; age-related macular degeneration; lymphoma;

Hodgkin's lymphoma; head and neck cancer; oral or mouth cancer; or cancer of the blood, prostate, cervix, uterus, vaginal or other female adnexa, breast, naso-pharynx, trachea, larynx, bronchi, bronchioles, lung, hollow organs, esophagus, stomach, bile duct, intestine, colon, colorectum, rectum, bladder, ureter, kidney, liver, gall bladder, spleen, brain, lymphatic system, bones, skin or pancreas. Preferably it is
5 suitable for the detection of early cancer; cervical dysplasia; soft tissue sarcoma; a germ cell tumour; retinoblastoma; age-related macular degeneration; Hodgkin's lymphoma; or cancer of the blood, prostate, cervix, uterus, vaginal or other female adnexa, breast, naso-pharynx, trachea, larynx, bronchi, bronchioles, lung, hollow
10 organs, esophagus, stomach, bile duct, intestine, colon, colorectum, rectum, bladder, ureter, kidney, liver, gall bladder, spleen, brain, lymphatic system or bones. Preferably it is suitable for the detection of lymphoma; Hodgkin's lymphoma; oral or mouth cancer; or cancer of the prostate, cervix, vaginal or other female adnexa, breast, naso-pharynx, lung, esophagus, kidney, liver, brain, lymphatic system, bones,
15 skin or pancreas. Preferably it is suitable for the detection of lymphoma; oral or mouth cancer; or cancer of the prostate, cervix, vaginal or other female adnexa, breast, naso-pharynx, lung, esophagus, bones or skin.

Preferably the photodiagnostic agent of the fifth example is suitable for the
20 fluorescent or phosphorescent detection of the said diseases, preferably the fluorescent or phosphorescent detection and quantification of the said diseases.

Preferably the medicament, the phototherapeutic agent or the photodiagnostic agent is adapted for administration simultaneous with or prior to administration of
25 irradiation or sound, preferably for administration prior to administration of irradiation.

If the medicament or the phototherapeutic agent is for use in photodynamic therapy or cytoluminescent therapy, then it is preferably adapted for administration
30 hours before the irradiation, preferably 6 to 72 hours before the irradiation, preferably 24 to 48 hours before the irradiation.

If the photodiagnostic agent is for use in photodynamic diagnosis, then it is preferably adapted for administration 3 to 60 hours before the irradiation, preferably 8 to 40 hours before the irradiation.

- 5 Preferably the irradiation used in the photodynamic therapy, cytoluminescent therapy or photodynamic diagnosis is electromagnetic radiation with a wavelength in the range of from 500nm to 1000nm, preferably from 600nm to 790nm. The electromagnetic radiation may be administered for about 5-60 minutes, preferably for about 15-20 minutes, at about 0.1-5W, preferably at about 1W. In one example,
- 10 two sources of electromagnetic radiation are used (for example a laser light and an LED light), one source with a wavelength in the range of from 600nm to 690nm, preferably about 660nm, the other source with a wavelength in the range of from 700nm to 790nm, preferably about 760nm. In another example, two sources of electromagnetic radiation are used (for example a laser light and an LED light), both
- 15 sources adapted to provide irradiation with a wavelength in the range of from 600nm to 690nm, preferably about 660nm, and irradiation with a wavelength in the range of from 700nm to 790nm, preferably about 760nm. In another example, the irradiation may be provided by a prostate, anal, vaginal, mouth and nasal device for insertion into a body cavity. In another example, the irradiation may be provided by
- 20 interstitial light activation, for example, using a fine needle to insert an optical fibre laser into the lung, liver, lymph nodes or breast. In another example, the irradiation may be provided by endoscopic light activation, for example, for delivering light to the lung, stomach, colon, bladder or neck.
- 25 A sixth example provides a method of treating atherosclerosis; multiple sclerosis; diabetes; diabetic retinopathy; arthritis; rheumatoid arthritis; a fungal, viral, chlamydial, bacterial, nanobacterial or parasitic infectious disease; HIV; Aids; infection with sars virus, Asian (chicken) flu virus, herpes simplex or herpes zoster; hepatitis; viral hepatitis; a cardiovascular disease; coronary artery stenosis; carotid
- 30 artery stenosis; intermittent claudication; a dermatological condition; acne; psoriasis; a disease characterised by benign or

malignant cellular hyperproliferation or by areas of neovascularisation; a benign or malignant tumour; early cancer; cervical dysplasia; soft tissue sarcoma; a germ cell tumour; retinoblastoma; age-related macular degeneration; Hodgkin's lymphoma; or cancer of the blood, prostate, cervix, uterus, vaginal or other female adnexa, breast,
5 naso-pharynx, trachea, larynx, bronchi, bronchioles, lung, hollow organs, esophagus, stomach, bile duct, intestine, colon, colorectum, rectum, bladder, ureter, kidney, liver, gall bladder, spleen, brain, lymphatic system or bones; the method comprising administering a therapeutically effective amount of chlorin e4 sodium according to the first or third example to a human or animal in need thereof.

10

The sixth example also provides a method of photodynamic therapy or cytoluminescent therapy of a human or animal disease, the method comprising administering a therapeutically effective amount of chlorin e4 sodium according to the first or third example to a human or animal in need thereof. Preferably the
15 human or animal disease is atherosclerosis; multiple sclerosis; diabetes; diabetic retinopathy; arthritis; rheumatoid arthritis; a fungal, viral, chlamydial, bacterial, nanobacterial or parasitic infectious disease; HIV; Aids; infection with sars virus, Asian (chicken) flu virus, herpes simplex or herpes zoster; hepatitis; viral hepatitis; a cardiovascular disease; coronary artery stenosis; carotid artery stenosis; intermittent
20 claudication; a dermatological condition; acne; psoriasis; a disease characterised by benign or malignant cellular hyperproliferation or by areas of neovascularisation; a benign or malignant tumour; early cancer; cervical dysplasia; soft tissue sarcoma; a germ cell tumour; retinoblastoma; age-related macular degeneration; Hodgkin's lymphoma; or cancer of the blood, prostate, cervix, uterus, vaginal or other female
25 adnexa, breast, naso-pharynx, trachea, larynx, bronchi, bronchioles, lung, hollow organs, esophagus, stomach, bile duct, intestine, colon, colorectum, rectum, bladder, ureter, kidney, liver, gall bladder, spleen, brain, lymphatic system or bones.

Preferably the method of the sixth example is a method of treating benign or
30 malignant cellular hyperproliferation or areas of neovascularisation.

Preferably the method of the sixth example is a method of treating a benign or malignant tumour.

Preferably the method of the sixth example is a method of treating early cancer;
5 cervical dysplasia; soft tissue sarcoma; a germ cell tumour; retinoblastoma; age-related macular degeneration; lymphoma; Hodgkin's lymphoma; head and neck cancer; oral or mouth cancer; or cancer of the blood, prostate, cervix, uterus, vaginal or other female adnexa, breast, naso-pharynx, trachea, larynx, bronchi, bronchioles, lung, hollow organs, esophagus, stomach, bile duct, intestine, colon,
10 colorectum, rectum, bladder, ureter, kidney, liver, gall bladder, spleen, brain, lymphatic system, bones, skin or pancreas. Preferably the method is a method of treating early cancer; cervical dysplasia; soft tissue sarcoma; a germ cell tumour; retinoblastoma; age-related macular degeneration; Hodgkin's lymphoma; or cancer of the blood, prostate, cervix, uterus, vaginal or other female adnexa, breast, naso-pharynx, trachea, larynx, bronchi, bronchioles, lung, hollow organs, esophagus,
15 stomach, bile duct, intestine, colon, colorectum, rectum, bladder, ureter, kidney, liver, gall bladder, spleen, brain, lymphatic system or bones. Preferably the method is a method of treating lymphoma; Hodgkin's lymphoma; oral or mouth cancer; or cancer of the prostate, cervix, vaginal or other female adnexa, breast, naso-pharynx,
20 lung, esophagus, kidney, liver, brain, lymphatic system, bones, skin or pancreas. Preferably the method is a method of treating lymphoma; oral or mouth cancer; or cancer of the prostate, cervix, vaginal or other female adnexa, breast, naso-pharynx, lung, esophagus, bones or skin.

25 The sixth example also provides a method of photodynamic diagnosis of a human or animal disease, the method comprising administering a diagnostically effective amount of chlorin e4 sodium according to the first or third example to a human or animal. Preferably the human or animal disease is atherosclerosis; multiple sclerosis; diabetes; diabetic retinopathy; arthritis; rheumatoid arthritis; a fungal,
30 viral, chlamydial, bacterial, nanobacterial or parasitic infectious disease; HIV; Aids; infection with sars virus, Asian (chicken) flu virus, herpes simplex or herpes zoster; hepatitis; viral hepatitis; a cardiovascular

disease; coronary artery stenosis; carotid artery stenosis; intermittent claudication; a dermatological condition; acne; psoriasis; a disease characterised by benign or malignant cellular hyperproliferation or by areas of neovascularisation; a benign or malignant tumour; early cancer; cervical dysplasia; soft tissue sarcoma; a germ cell tumour; retinoblastoma; age-related macular degeneration; Hodgkin's lymphoma; or cancer of the blood, prostate, cervix, uterus, vaginal or other female adnexa, breast, naso-pharynx, trachea, larynx, bronchi, bronchioles, lung, hollow organs, esophagus, stomach, bile duct, intestine, colon, colorectum, rectum, bladder, ureter, kidney, liver, gall bladder, spleen, brain, lymphatic system or bones. Preferably the human or animal disease is characterised by benign or malignant cellular hyperproliferation or by areas of neovascularisation. Preferably the human or animal disease is a benign or malignant tumour. Preferably the human or animal disease is early cancer; cervical dysplasia; soft tissue sarcoma; a germ cell tumour; retinoblastoma; age-related macular degeneration; lymphoma; Hodgkin's lymphoma; head and neck cancer; oral or mouth cancer; or cancer of the blood, prostate, cervix, uterus, vaginal or other female adnexa, breast, naso-pharynx, trachea, larynx, bronchi, bronchioles, lung, hollow organs, esophagus, stomach, bile duct, intestine, colon, colorectum, rectum, bladder, ureter, kidney, liver, gall bladder, spleen, brain, lymphatic system, bones, skin or pancreas. Preferably the human or animal disease is early cancer; cervical dysplasia; soft tissue sarcoma; a germ cell tumour; retinoblastoma; age-related macular degeneration; Hodgkin's lymphoma; or cancer of the blood, prostate, cervix, uterus, vaginal or other female adnexa, breast, naso-pharynx, trachea, larynx, bronchi, bronchioles, lung, hollow organs, esophagus, stomach, bile duct, intestine, colon, colorectum, rectum, bladder, ureter, kidney, liver, gall bladder, spleen, brain, lymphatic system or bones. Preferably the human or animal disease is lymphoma; Hodgkin's lymphoma; oral or mouth cancer; or cancer of the prostate, cervix, vaginal or other female adnexa, breast, naso-pharynx, lung, esophagus, kidney, liver, brain, lymphatic system, bones, skin or pancreas. Preferably the human or animal disease is lymphoma; oral or mouth cancer; or cancer of the prostate, cervix, vaginal or other female adnexa, breast, naso-pharynx, lung, esophagus, bones or skin. Preferably the method of photodynamic diagnosis is suitable for the fluorescent or phosphorescent detection of the said diseases,

preferably for the fluorescent or phosphorescent detection and quantification of the said diseases.

In any of the methods of the sixth example, the human or animal is preferably
5 further subjected to irradiation or sound simultaneous with or after the
administration of the chlorin e4 sodium. Preferably the human or animal is
subjected to irradiation after the administration of the chlorin e4 sodium.

If the method is a method of photodynamic therapy or cytoluminescent therapy,
10 then the human or animal is preferably subjected to irradiation 5 to 100 hours after
administration of the chlorin e4 sodium, preferably 6 to 72 hours after
administration, preferably 24 to 48 hours after administration.

If the method is a method of photodynamic diagnosis, then the human or animal is
15 preferably subjected to irradiation 3 to 60 hours after administration of the chlorin
e4 sodium, preferably 8 to 40 hours after administration.

Preferably the irradiation is electromagnetic radiation with a wavelength in the range
of from 500nm to 1000nm, preferably from 600nm to 790nm. The electromagnetic
20 radiation may be administered for about 5-60 minutes, preferably for about 15-20
minutes, at about 0.1-5W, preferably at about 1W. In one example, two sources of
electromagnetic radiation are used (for example a laser light and an LED light), one
source with a wavelength in the range of from 600nm to 690nm, preferably about
660nm, the other source with a wavelength in the range of from 700nm to 790nm,
25 preferably about 760nm. In another example, two sources of electromagnetic radiation are
used (for example a laser light and an LED light), both sources adapted to provide
irradiation with a wavelength in the range of from 600nm to 690nm, preferably about
660nm, and irradiation with a wavelength in the range of from 700nm to 790nm, preferably
about 760nm. In another example, the irradiation may be provided by a prostate, anal,
30 vaginal, mouth and nasal device for insertion into a body cavity. In another example, the
irradiation may be provided by interstitial light activation, for example, using a

fine needle to insert an optical fibre laser into the lung, liver, lymph nodes or breast. In another example, the irradiation may be provided by endoscopic light activation, for example, for delivering light to the lung, stomach, colon, bladder or neck.

5 Preferably the human or animal is a human.

Brief description of the drawings

Figure 1 shows a modified Jablonski diagram for a typical photosensitizer, outlining
10 the principle photophysical processes of interest.

Figure 2 depicts schematically the metalation and demetalation of a porphyrin ring.

Figure 3 depicts schematically the reaction of 1,3-diphenylisobenzofuran (DPIBF)
15 with two singlet oxygens ($^1\text{O}_2$).

Figure 4 shows the absorption spectrum of a chlorin e4 sodium solution (about 10 μM) in ethanol.

20 Figure 5 shows the absorption spectrum of a chlorin e4 sodium solution (about 10 μM) in water.

Figure 6 shows the absorption spectrum of a chlorin e4 sodium solution (about 10 μM) in water with 0.2 M detergent sodium dodecyl sulphate (SDS).
25

Figure 7 shows the fluorescence spectrum of a chlorin e4 sodium solution (about 10 μM) in ethanol. Excitation wavelength was 403 nm; the absorbance at this wavelength was 0.10; cuvette size 1cm.

30 Figure 8 shows the fluorescence spectrum of a chlorin e4 sodium solution (about 10 μM) in water. Excitation wavelength was 403 nm; the absorbance at this wavelength was 0.13; cuvette size 1 cm.

Figure 9 shows the fluorescence spectrum of a chlorin e4 sodium solution (about 10 μ M) in water with 0.2 M detergent sodium dodecyl sulphate (SDS). Excitation wavelength was 403 nm; the absorbance at this wavelength was 0.13; cuvette size 1
5 cm.

Figure 10 shows the fluorescence excitation spectrum of a chlorin e4 sodium solution (about 10 μ M) in ethanol. Fluorescence was measured at 710 nm; the absorbance at 403 nm was 0.15 in 1 cm; the optical path was 0.5 cm.

10

Figure 11 shows the absorption spectra of a solution of DPIBF and chlorin e4 sodium in ethanol before and 14 minutes after irradiation by red light (660 nm).

Figure 12 shows kinetic traces (1) and the spectrum (2) of singlet oxygen
15 phosphorescence after pulsed laser excitation of chlorin e4 sodium in ethanol.

Figure 13 shows kinetic traces (1) and the spectrum (2) of singlet oxygen phosphorescence after pulsed laser excitation of chlorin e4 sodium in water.

20 Figure 14 shows kinetic traces of singlet oxygen phosphorescence after pulsed laser excitation of chlorin e4 sodium in water with 0.2 M detergent sodium dodecyl sulphate (SDS).

Detailed description of the invention

25

Metal insertion into the central part of a porphyrin or chlorin ring is known as metalation and the general chemical process is depicted in Figure 2. Porphyrins and chlorins can form complexes with a wide variety of metals (M). These metal complexes not only protect the inner nitrogen atoms from electrophilic reagents and
30 strong bases, but also have a pronounced effect on the reactivity of the macrocycles. A common geometry of these complexes is octahedral, with the metal ions occupying the centre of the N₄ porphyrin plane and the metal ligands being in *trans* positions (Figure 2).

Chlorin e4 has nine side chains, two of which carry a carboxylic acid moiety. These carboxylic acid moieties can form salts with a wide variety of metals (M).

5 Experimental details

Example – general synthetic details

Method 1: 0.15-1mmol of chlorin e4 is dissolved in 60-100ml of methanol. To this
10 solution, 15-20ml of sodium hydroxide (3M water solution) is added. The mixture
is stirred until the synthesis is complete after about 6 hours. Then the mixture is
filtered through a 45µm Millipore filter. The product is purified by flash
chromatography with a methanol mobile phase. Then the solution is distilled and
lyophilized. In a preferred embodiment of the present invention, the reaction is
15 carried out at a temperature in the range of 25-60°C.

Method 2: 0.15-1mmol of chlorin e4 is dissolved in 60-100ml of methanol. To this
solution, 15-20ml of sodium hydroxide (3M water solution) is added. Separately, 5-
12mmol of thiamine pyrophosphate is dissolved in 40-60ml of water, and the latter
20 solution is added to the previous one. The mixture is stirred until the synthesis is
complete after about 6 hours. Then the mixture is filtered through a 45µm
Millipore filter. The product is purified by flash chromatography with a methanol
mobile phase. Then the solution is distilled and lyophilized. In a preferred
embodiment of the present invention, the reaction is carried out at a temperature in
25 the range of 25-60°C.

Example – synthesis of chlorin e4 sodium

Method 1: 0.5mmol (276mg) of chlorin e4 was dissolved in 60ml of methanol, and
30 the solution was stirred slowly for 10 minutes at 40°C. 15ml of sodium hydroxide
(3M water solution) was added, and stirring was continued for a further 6 hours at
60°C. Then the reaction mixture was filtered through a 45µm Millipore filter. The
product was purified by flash chromatography on a dry silica gel column using a

methanol mobile phase. Then the solution was reduced in volume on a rotary evaporator and the product was isolated as a dry powder by lyophilization in 85% yield.

5 The chlorin e4 sodium product was characterised by ^1H and ^{13}C NMR.

^1H NMR (D_2O) (ppm): 8.82 (1H), 8.64 (1H), 6.52 (1H), 6.35 (1H), 5.25 (1H), 5.12 (1H), 4.68 (2H), 4.09 (3H), 3.52 (3H), 2.87 (1H), 2.71 (2H), 2.59 (3H), 2.44 (2H), 2.06 (1H), 1.82 (3H), 0.99 (6H).

10

^{13}C NMR (D_2O) (ppm): 185.7 (CO_2H), 180.8 (CO_2H), 171.6 ($\text{C}_{\text{quat}}=$), 170.7 ($\text{C}_{\text{quat}}=$), 148.0 ($\text{C}_{\text{quat}}=$), 144.7 ($\text{C}_{\text{quat}}=$), 139.0 ($\text{C}_{\text{quat}}=$), 137.5 ($\text{C}_{\text{quat}}=$), 136.5 ($\text{C}_{\text{quat}}=$), 136.2 ($\text{C}_{\text{quat}}=$), 135.3 ($\text{C}_{\text{quat}}=$), 133.9 ($\text{C}_{\text{quat}}=$), 133.6 ($\text{C}_{\text{quat}}=$), 132.5 ($\text{C}_{\text{quat}}=$), 131.2 ($\text{C}_{\text{quat}}=$), 129.5 ($\text{CH}=\text{}$), 121.6 ($\text{CH}_2=\text{}$), 121.4 ($\text{C}_{\text{quat}}=$), 109.4 ($\text{C}_{\text{quat}}=$), 101.2 ($\text{CH}=\text{}$), 97.4 (CH=), 96.2 (CH=), 56.8 (CH), 50.9 (CH), 38.0 (CH_2), 34.5 (CH_2), 26.1 (Me), 21.7 (Me), 20.1 (CH_2), 18.7 (Me), 14.0 (Me), 12.6 (Me), 10.7 (Me).

15

Method 2: 0.5mmol (276mg) of chlorin e4 was dissolved in 60ml of methanol, and the solution was stirred slowly for 10 minutes at 40°C. 15ml of sodium hydroxide (3M water solution) was added, and stirring was continued for a further 20 minutes. Separately, 6mmol (378mg) of thiamine pyrophosphate was dissolved in 50ml of water, and the solution was stirred at 60°C. The thiamine pyrophosphate solution was added to the chlorin e4 solution, and the reaction mixture was stirred slowly for 6 hours at 60°C. Then the reaction mixture was filtered through a 45 μm Millipore filter. The product was purified by flash chromatography on a dry silica gel column using a methanol mobile phase. Then the solution was reduced in volume on a rotary evaporator and the product was isolated as a dry powder by lyophilization in 85% yield.

25

30 The chlorin e4 sodium product was characterised by ^1H and ^{13}C NMR. The same results as for method 1 were obtained.

Example – *in vitro* studies

The following experiments were carried out to study the absorption and fluorescence spectra of chlorin e4 sodium, the quantum yields of chlorin e4 sodium
 5 fluorescence and the photosensitizing ability of chlorin e4 sodium in three systems – solutions in ethanol, pure water and water containing detergent sodium dodecyl sulfate (SDS).

Absorption and fluorescence measurements

10

Absorption spectra were recorded using computerized spectrophotometers Hitachi 3400 (Japan) and SF-2000 (Russia). Fluorescence spectra were measured with a computerized spectrofluorimeter Perkin-Elmer MPF-44B. 10 mm wide quartz cuvettes were used for absorption and fluorescence measurements. Fluorescence
 15 was detected at a right angle to the excitation beam. The spectral bandwidth of the monochromator was 3 nm. To avoid fluorescence re-absorption by the pigment molecules, diluted chlorin e4 sodium solutions were used with absorbance at 403 nm which did not exceed 0.2 cm in the 1 cm cuvette. The fluorescence quantum yields (Φ_F) were determined using as a reference, solutions of TPPS (m-tetrakis(p-sulfonatophenyl)porphine) in ethanol. According to the literature, the most likely
 20 fluorescence quantum yield of TPPS in air-saturated water and ethanol is equal to 0.08 (K. Kalyanasundaram and M. Neumann-Spallart, Photophysical and redox properties of water soluble porphyrins in aqueous media, J. Phys. Chem., 1982, 86, 5163-5169; and A.T. Gradyushko, A.N. Sevchenko, K.N. Solovyov and M.P. Tsvirko, Energetics of photophysical processes in chlorophyll-like molecules, Photochem. Photobiol., 1970, 11, 387-400). The fluorescence of the samples was
 25 excited by the monochromatic light corresponding to the maximum of the Soret bands: 415 nm for TPPS in ethanol and 413 nm for TPPS in water, and 403 nm for chlorin e4 sodium in water and ethanol. The monochromator bandwidth was 3 nm.
 30 The fluorescence quantum yields (Φ_F) were calculated using the following equation:

$$\Phi_F = \Phi_{TPP} \cdot \frac{S_c \cdot (1 - 10^{-D_{TPP}}) / (1 - 10^{-D_c})}{S_{TPP}}$$

where Φ_{TPP} is the quantum yield of TPPS fluorescence, S_e is the square under the fluorescence spectrum of the chlorin e4 sodium sample, S_{TPP} is the square under the fluorescence spectrum of TPPS, and D_e and D_{TPP} are absorbance (optical density) of the chlorin e4 sodium and TPPS respectively at the excitation wavelengths. The
5 obtained results were collated using relative units of intensity of the excited light at the excitation wavelengths. The intensities of the monochromatic light were measured using a calibrated silicon detector from Optical Power Meter System ThorLabs (Karlsfeld, Germany).

10 *Detection of singlet oxygen phosphorescence*

Singlet oxygen phosphorescence was detected using measurement of infrared phosphorescence at 1270 nm corresponding to the transition of oxygen molecules from the singlet to the triplet state. Phosphorescence was recorded at room
15 temperature with the laser phosphorescence spectrometer described at A.A. Krasnovsky Jr., Photodynamic action and singlet oxygen, Biofizika, 2004, 49, 305-321; and A.A. Krasnovsky Jr., Singlet oxygen and primary mechanisms of photodynamic therapy and photodynamic diseases, In: Photodynamic therapy at the cellular level, ed. A.B. Uzdensky, Research Singpost, Kerala, India, 2007, 17-62. For
20 excitation, nanosecond pulses of 511 nm radiation from a copper-vapour laser (Troitzk, Physics Institute of the Russian Academy of Science) were employed. Laser radiation passed additionally through a green light filter SZS-22 (4 mm), which transmitted laser light and cut off IR laser radiation caused by heating of the laser tube. Pulse duration was 20 ns, pulse energy varied from 18 to 40 μJ and the
25 pulse repetition rate was 12 kHz. The laser light was collimated on the surface of a 1 cm rectangular quartz cuvette. The diameter of the illuminated spot on the surface of the cell was 5 mm. For kinetic measurements another spectrometer was used without a monochromator. In this case, phosphorescence resulting from the right angle to the excitation beam was focused on the photocathode of a cooled S-1
30 photomultiplier (FEU-112) through a cut-off IR light filter and an interference filter with the transmission maximum at 1270 nm. The phosphorescence spectra were measured using a MS-80 high throughput grid monochromator. In this case, phosphorescence was focused on the entry slit of this monochromator. The

bandwidth of the monochromator corresponded to a 20 nm spectral interval. The monochromatic light was focused on the photocathode of a cooled S-1 photomultiplier (FEU-83). The photomultiplier signal was registered using time-correlated single photon counting and averaged over many laser pulses. In most experiments, one-channel on time was equal to 81 ns. When the pigment bleaching reached 10%, the solution was replaced by a fresh one and measurement was continued. As chlorin e4 sodium bleached rather rapidly, the solutions were changed after each 60-180 s irradiation. Reasonable phosphorescent curves were obtained after 10-20 minutes signal accumulation and therefore samples were changed many times. As described previously (D.A. Butorina, A.A. Krasnovsky Jr. and A.V. Priezzev, Investigation of the kinetic parameters of singlet molecular oxygen in aqueous porphyrin solutions, Influence of detergents and the quencher sodium azide, Biofizika, 2003, 48, 189-196; A.A. Krasnovsky Jr., Luminescence and photochemical studies of singlet oxygen photonics, J. Photochem. Photobiol. A: Chem., 2008, 196, 210-218; and N. Kuznetsova, D. Makarov, O. Yuzhakova, A. Strizhakov, Y. Roumbal, L. Ulanova, A. Krasnovsky and O. Kaliya, Photophysical properties and photodynamic activity of octacationic oxotitanium(IV) phthalocyanines, Photochemical Photobiological Sciences, 2009, 8, 1724-1733), for computer analysis the following two-exponential equation was used:

$$I(t) = I_0 \cdot [\exp(-k_{decay} \cdot t) - \exp(-k_{rise} \cdot t)] + n$$

where k_{decay} and k_{rise} are the rate constants of the decay and rise phases of the phosphorescence curves after a laser pulse, n is the off noise of the photomultiplier, I_0 is the zero-time phosphorescence intensity, and $I(t)$ is the phosphorescence intensity at time t . To reduce contribution of the luminescence of the pigment in the quartz cuvette, which was observed shortly after laser pulses, readings were taken starting from 400 ns after laser pulses in ethanol and 1 μ s in water. In addition, phosphorescence and kinetic traces were measured also from the control chlorin e4 sodium solutions containing high concentration of sodium azide, which completely quenched singlet oxygen. This procedure revealed a strong rapidly decaying component of delayed luminescence in the chlorin e4 sodium solutions, which was not so easily quenched by sodium azide. The luminescent curves obtained from the control solutions were approximated by exponentials and

subtracted from the experimental kinetic traces obtained for the chlorin e4 sodium solutions where sodium azide was not added.

For measurements, Origin 6.0 software was used. This software operates using the
5 least-square approximation method. Origin allowed variation of the parameters I_0 ,
 k_{decay} , k_{rise} and n in order to obtain their values corresponding to the minimum
coefficient x^2 . In chlorin e4 sodium solutions, measurements always yielded k_{decay} ,
which was somewhat shorter than in the reference solutions of TPPS in the same
solvents. In the solvents used in this study, the lifetime of singlet oxygen, which is
10 responsible for k_{decay} , is independent of the pigments. Therefore this result indicates
that "the azide control" does not allow to fully eliminate the rapidly decaying
component from the overall decay curves. It is possible that luminescence
responsible for this component is partially quenched by sodium azide. In order to
eliminate the error caused by this effect, for final measurements of phosphorescent
15 curves in chlorin e4 sodium solutions, fixed k_{decay} obtained in the TPPS solutions
was used. In this case as Origin indicated, the correlation coefficient R^2 was not
less than 98% and error of k_{rise} determination was less than $\pm 5\%$ in all solvents.
The quantum yields (Φ_Δ) of singlet oxygen photogeneration by chlorin e4 sodium
were determined from comparison of the overall singlet oxygen phosphorescence
20 intensities in the chlorin e4 sodium solutions and the reference solutions of TPPS in
the same solvents. According to the literature, the most probable values of
quantum yields of singlet oxygen photogeneration by TPPS are 0.6 in water and 0.7
in ethanol and detergent containing water (F. Wilkinson, W.P. Helman and A.B.
Ross, Quantum yields of the photosensitized formation of the lowest electronically
25 excited singlet states of molecular oxygen, J. Phys. Chem. Ref. Data, 1993, 22(2),
113-262; and C. Tanielian, C. Wolf and M. Esch, Singlet oxygen production in
water: Aggregation and charge transfer effects, J. Phys. Chem., 1996, 100, 6555-
6560). The overall phosphorescence intensity was calculated by integrating the
phosphorescence kinetic traces. The absorbance of the pigment solutions at 511
30 nm was usually about 0.1-0.2 in 5 mm cuvette. This corresponded to a pigment
concentration equal to about 20 μM .

Detection of singlet oxygen using chemical conjugates

The second method of singlet oxygen detection was based on photosensitized oxygenation of the singlet oxygen conjugate 1,3-diphenylisobenzofuran (DPIBF) (Acros Organics, United Kingdom). DPIBF chemically binds to two singlet oxygens ($^1\text{O}_2$) (Figure 3). This method was applied to chlorin e4 sodium solutions in ethanol. The major absorption band of DPIBF is located at 414-415 nm. The molar absorption coefficient in this maximum, ϵ_{DPIBF} , is known to be $23500 \text{ M}^{-1}\text{cm}^{-1}$. Tested solutions were irradiated by the monochromatic light from a xenon lamp using the excitation system of the spectrofluorimeter Perkin-Elmer MPF-44B, the spectral width of the monochromator bandwidth corresponding to 5 nm. Radiant power was measured using the calibrated detector ThorLabs PM-100 (Germany). The singlet oxygen quantum yields (Φ_{Δ}) were calculated according to the equation:

$$\Phi_{\Delta} = \frac{\eta_p \cdot \frac{\Delta D_{\text{DPIBF}}}{\epsilon_{\text{DPIBF}} \cdot l \cdot t_{\text{irr}}} \cdot V \cdot N_A}{\frac{I_{\text{irr}}}{h\nu(1-T)}}$$

where ΔD_{DPIBF} is the reduction in the optical density of DPIBF at 414 nm during the irradiation time, ϵ_{DPIBF} is the molar absorption coefficient of DPIBF at 414 nm, l is the optical path equal to 1 cm, V is the volume of tested solutions (usually 1.2-1.5 ml), N_A is the Avogadro constant, I_{irr} is the radiant power (in W) of the monochromatic light from the excitation system of the Perkin-Elmer spectrofluorimeter that includes the xenon lamp and the excitation monochromator with the bandwidth corresponding to the 5 nm wavelength interval, $h\nu$ is the energy of one photon in J, $(1-T)$ is the absorption factor of the tested solutions, t_{irr} is the irradiation time in seconds, and η_p is the fraction of $^1\text{O}_2$ molecules which is captured by DPIBF in the studied solutions. This parameter was calculated using known kinetic equations for singlet oxygen consumption by chemical conjugates and the rate constant of singlet oxygen reaction with DPIBF equal to $10^9 \text{ M}^{-1}\text{s}^{-1}$ (see A.A. Krasnovsky Jr., Y.V. Roumbal and A.A. Strizhakov, Rates of $^1\text{O}_2$ production upon direct excitation of molecular oxygen by 1270 nm laser radiation in air-saturated alcohols and micellar aqueous dispersions, Chem. Phys. Lett., 2008, 458, 195-199; and references therein).

Results

Absorption spectra of chlorin e4 sodium in ethanol, pure water and water
 5 containing detergent sodium dodecyl sulfate (SDS) are shown in Figures 4-6. The
 major absorption bands are listed in Table 1. The strongest are the Soret band at
 403 nm and the red band at 654-662 nm. Analysis of the absorption spectra
 suggests that chlorin e4 sodium was mostly monomeric in all solvents. In water
 having no detergent, the red absorption maximum was slightly shifted to the shorter
 10 wavelength region (from 662 to 654 nm) and a weak shoulder at about 665-670 nm
 was seen. In addition, the observed absorbance in the green region was increased
 compared to ethanol and detergent containing solutions. These data indicate that
 the solution in water having no detergent may contain aggregated (dimeric and
 oligomeric) molecules of chlorin e4 sodium, which disaggregate in the presence of
 15 the detergent.

Solvents	Absorption maxima (nm $\pm$ 1 nm)	Fluorescence maxim (nm $\pm$ 1 nm)
Ethanol	403, 504, 606, 662	666, 720
Water	402, 504, 602, 654	657, 710
Water + 0.2 M SDS	403, 504, 608, 662	665, 720

Table 1 – Positions of the major absorption and
 fluorescence bands of chlorin e4 sodium solutions

20 Fluorescence measurements supported this assumption (Figures 7-9). The
 fluorescence spectra were rather similar in ethanol and SDS containing water.
 Figure 10 shows the excitation spectrum of chlorin e4 sodium fluorescence in
 ethanol. It is seen that it practically coincides with the absorption spectrum of this
 pigment. This observation indicates that fluorescence belongs to chlorin e4 sodium.
 25 In water without SDS, the fluorescence maximum was shifted to shorter
 wavelengths and a shoulder at about 710 nm became markedly stronger than in
 ethanol. The fluorescence quantum yield was found to be highest in ethanol,
 slightly less in SDS containing water and the smallest in water without SDS. These
 data indicate that aggregation of chlorin e4 sodium occurs in water. However, the
 30 chlorin e4 sodium fluorescence quantum yield was still high in water without SDS,

showing that an appreciable amount of monomeric chlorin e4 sodium molecules are present in aqueous solutions without detergent. These experiments show that chlorin e4 sodium is a highly fluorescent pigment and emits a large quantity of photons. Hence chlorin e4 sodium can be used efficiently as a fluorescent pigment
5 for photodynamic diagnosis.

Singlet oxygen generation was studied in ethanol solutions using both chemical singlet oxygen conjugate DPIBF and measurement of chlorin e4 sodium photosensitized phosphorescence of singlet oxygen at 1270 nm. Figure 11
10 illustrates the experiment with DPIBF. The absorption maximum of DPIBF lies at 420 nm. It is seen from Figure 11 that irradiation of an ethanol solution containing both chlorin e4 sodium and DPIBF by red light (660 nm), which is absorbed by chlorin e4 sodium but not by DPIBF, caused bleaching of the DPIBF maximum at 420 nm. This effect results from reaction of DPIBF with singlet oxygen, which is
15 formed upon irradiation of chlorin e4 sodium. The quantum yield of singlet oxygen generation obtained from this experiment was shown to be 0.68 ± 0.05 in ethanol. Kinetic traces of chlorin e4 sodium photosensitized phosphorescence of singlet oxygen after laser pulses are shown in Figures 12-14. Table 2 summarizes kinetic parameters of the phosphorescent curves. The rise time reflects the rate of singlet
20 oxygen generation by triplet pigment molecules. It corresponds with the lifetime of the pigment triplet state in air-saturated pigment solution. In ethanol, the rise time was equal to 300 ns. The decay time corresponds with the singlet oxygen lifetime in ethanol (13.5 μ s). The singlet oxygen quantum yield in ethanol determined using these kinetic curves was equal to 0.63 ± 0.05 , which is close to that obtained using
25 DPIBF. Thus both methods provided similar results. Table 2 indicates that chlorin e4 sodium is a potent photosensitizer of singlet oxygen formation in alcoholic as well as aqueous systems, though in water without SDS the photosensitizing activity was lower than in ethanol. This effect may have been due to partial aggregation of pigment molecules, which is known to lead to the loss of photosensitizing activity.
30 These experiments show that chlorin e4 sodium has strong photosensitizing ability in organic solvent (alcohol) and in aqueous media, with a high singlet oxygen quantum yield. Hence chlorin e4 sodium can be used efficiently in photodynamic therapy.

Solvents	Rise time, μ s	Decay time, μ s
Ethanol	0.3 ± 0.3	13.5 ± 0.3
Water	1.8 ± 0.4	3.3 ± 0.3
Water + 0.2 M SDS	1.2 ± 0.3	3.85 ± 0.3

Table 2 – Kinetic parameters of photosensitized singlet oxygen phosphorescence in air saturated chlorin e4 sodium solutions

5 Conclusion

Chlorin e4 sodium shows strong fluorescence and strong photosensitizing ability in organic solvent (alcohol) and in aqueous media. In water, the chlorin e4 sodium photosensitizing activity decreases due to aggregation of pigment molecules, but is
 10 still high. Addition of detergent causes disaggregation and an increase of both fluorescence and photosensitizing ability. This shows that chlorin e4 sodium is active in both hydrophilic and aqueous media. Therefore this pigment is promising for application in photodynamic medicine.

15 Example – *in vivo* studies

The following cancer patients were treated successfully with chlorin e4 sodium. Most of the patients were late stage metastatic cancer patients who had previously unsuccessfully used traditional treatment therapies. Most of the patients were in
 20 poor health prior to treatment with chlorin e4 sodium.

Example 1

A 70 year old female patient with invasive ductile breast cancer and lymph node
 25 metastases in the right arm pit, was treated successfully. Prior to treatment a large cancer mass in the right breast was observed and a small cancer mass in the right arm pit. The treatment consisted of three courses. For each course, five doses of 62.5-187.5mg chlorin e4 sodium were administered orally over the course of eight days. 24-48 hours after each oral dose of chlorin e4 sodium, the patient was treated
 30 with laser and LED light having a wavelength of 660nm and 760nm.

Following treatment, doppler ultrasound analysis showed that the large cancer mass in the breast showed almost no vascularity, no blood vessels leading to it. The cancer mass had not reduced in size, however, the outline of the mass was solid and smooth, indicating an inactive, non-growing tumour. The smaller lymph node cancer mass also showed almost no vascularity and it had reduced in size from 1.9cm to 1cm.

Subsequently the patient was treated with two further courses. Again, for each course, five doses of 62.5-187.5mg chlorin e4 sodium were administered orally over the course of eight days. 24-48 hours after each oral dose of chlorin e4 sodium, the patient was treated with laser and LED light having a wavelength of 660nm and 760nm. Additionally an interstitial light treatment using a fine needle was performed on the breast with an optical fibre laser having a wavelength of 660nm and 760nm.

Doppler ultrasound analysis obtained after the fourth treatment course, showed that the large cancer mass in the breast had reduced in size by about 25%. PET/CT scans obtained after the fifth and final treatment course, showed a complete response in the breast and lymph nodes. After each treatment course, an inflammatory response was observed in the breast tumour site, which is typical as the immune system attacks the tumour. Towards the end of the second treatment course, the patient experienced a burning sensation in the tumour site of her breast. The patient is currently alive and well, two years and three months after starting her treatment and five months after completion of her treatment.

Example 2

A 62 year old male patient with naso pharyngeal carcinoma with extension to a lymph node in the left neck, was treated successfully. The treatment consisted of three courses. For each course, five doses of 62.5-187.5mg chlorin e4 sodium were administered orally over the course of eight days. 24-48 hours after each oral dose

of chlorin e4 sodium, the patient was treated with laser and LED light having a wavelength of 660nm and 760nm.

Immediately following treatment, a biopsy of the treated lymph node of the neck
5 showed a benign or non-cancerous lymph node, no cancer cells were detected. This was confirmed by another biopsy ten months after treatment. A comparison of CT scans, obtained one month before and ten months after treatment, showed no spread of tumours and that the size of the tumours was reduced.

10 Subsequently the patient was treated with a further course. Again, for the course, five doses of 62.5-187.5mg chlorin e4 sodium were administered orally over the course of eight days. 24-48 hours after each oral dose of chlorin e4 sodium, the patient was treated with laser and LED light having a wavelength of 660nm and 760nm.

15

Ultrasound images and biopsies obtained after the final treatment course, showed a complete response. The patient suffered no side effects, but an inflammatory response was observed after each treatment, which is typical as the immune system attacks the tumour. He is currently alive and well, two years and six months after
20 starting his treatment and five months after completion of his treatment.

Example 3

A 71 year old male patient with advanced metastatic kidney disease with metastases
25 in the gall bladder, liver and lung, was treated successfully. The treatment consisted of three courses. For each course, five doses of 62.5-187.5mg chlorin e4 sodium were administered orally over the course of eight days. 24-48 hours after each oral dose of chlorin e4 sodium, the patient was treated with laser and LED light having a wavelength of 660nm and 760nm.

30

PET/CT scans, obtained just before treatment and two months, three months, five months and 15 months after treatment, showed that throughout the patient's body

some tumours disappeared altogether and the remaining tumours reduced in size. No further metastasis was observed.

Subsequently the patient was treated with a further two courses. Again, for each course, five doses of 62.5-187.5mg chlorin e4 sodium were administered orally over the course of eight days. 24-48 hours after each oral dose of chlorin e4 sodium, the patient was treated with laser and LED light having a wavelength of 660nm and 760nm. Additionally an interstitial light treatment using a fine needle was performed on the liver and lung with an optical fibre laser having a wavelength of 660nm and 760nm.

PET/CT scans again showed a good treatment response. The patient suffered no side effects from the treatment. Unfortunately the patient has recently died. Prior to the five treatment courses, he was an advanced stage cancer patient with a life expectancy of only six months, but he managed to live an active life for two years and four months.

Example 4

A 20 year old female patient with stage IVB Hodgkin's lymphoma with metastases in the liver, lymphatic system, spleen, lung and bones, was treated successfully. The treatment consisted of three courses. For each course, five doses of 62.5-187.5mg chlorin e4 sodium were administered orally over the course of eight days. 24-48 hours after each oral dose of chlorin e4 sodium, the patient was treated with laser and LED light having a wavelength of 660nm and 760nm.

CT scans showed that in the areas of the patient's body that were treated with laser and LED light so far, some tumours disappeared altogether and some tumours reduced in size.

Subsequently the patient was treated with a further two courses. Again, for each course, five doses of 62.5-187.5mg chlorin e4 sodium were administered orally over the course of eight days. 24-48 hours after each oral dose of chlorin e4 sodium, the

patient was treated with laser and LED light having a wavelength of 660nm and 760nm.

PET/CT scans again showed a good treatment response. Unfortunately the patient
5 has recently died. Prior to the five treatment courses, she was in an extremely poor condition, suffering from extreme weight loss and lethargy, too weak to walk, and requiring constant pain relief. Before her treatment, her life expectancy was only two weeks, but she managed to live for two years and two months, at least part of it active.

10

Example 5

A 82 year old male patient with lung cancer was treated successfully. The treatment consisted of one course of five doses of 62.5-187.5mg chlorin e4 sodium
15 administered orally over the course of eight days. 24-48 hours after each oral dose of chlorin e4 sodium, the patient was treated with laser and LED light having a wavelength of 660nm and 760nm.

Three months after treatment, the patient was coughing up blood and tumour
20 tissue. A CT scan showed tumour breakdown.

Example 6

A 46 year old male patient with lung cancer was treated successfully. The treatment
25 consisted of three courses. For each course, five doses of 62.5-187.5mg chlorin e4 sodium were administered orally over the course of eight days. 24-48 hours after each oral dose of chlorin e4 sodium, the patient was treated with laser and LED light having a wavelength of 660nm and 760nm.

30 A CT scan, obtained immediately after the last treatment course, showed complete recovery from lung cancer.

Example 7

A 54 year old female patient with advanced pancreatic cancer with metastases in the liver, was treated successfully. Prior to treatment the patient was in poor health
5 generally and suffering from extreme weight loss, pain and other typical symptoms of advanced pancreatic cancer. The treatment consisted of three courses. For each course, five doses of 62.5-187.5mg chlorin e4 sodium were administered orally over the course of eight days. About 24 hours after each oral dose of chlorin e4 sodium, the patient was treated with laser and LED light having a wavelength of 660nm and
10 760nm.

PET/CT scans of the pancreas, obtained before and after treatment, showed improvements and a good treatment response in the pancreatic area. The patient had no photosensitive reaction and suffered less pain. Unfortunately bowel
15 blocking and malnutrition caused the patient's death.

Example 8

A 67 year old female patient with non-small cell lung cancer with metastases in the lymph nodes and bones, was treated successfully. The treatment consisted of two
20 courses. For each course, five doses of 62.5-187.5mg chlorin e4 sodium were administered orally over the course of eight days. About 24 hours after each oral dose of chlorin e4 sodium, the patient was treated with laser and LED light having a wavelength of 660nm and 760nm.

PET/CT scans of the lungs, obtained before and after treatment, showed a remarkable response to treatment, with many tumours disappearing and the remaining tumours showing no or low metabolic activity after treatment. The patient suffered no side effects and her health improved. She is currently alive and
25
30 well, 10 months after starting her treatment and seven months after completion of her treatment.

Example 9

A 54 year old female patient with breast cancer in the right breast was treated successfully. The treatment consisted of three courses. For each course, five doses
5 of 62.5-187.5mg chlorin e4 sodium were administered orally over the course of eight days. 24-48 hours after each oral dose of chlorin e4 sodium, the patient was treated with laser and LED light having a wavelength of 660nm and 760nm.

Mammography, ultrasound and PET/CT scans of the breasts, obtained before and
10 after treatment, showed a complete response. The patient suffered no side effects. She is currently alive and well, one year and seven months after starting her treatment and 10 months after completion of her treatment.

Example 10

15

A 33 year old male patient with naso pharyngeal carcinoma with metastases in the lymph nodes of the left neck and in the lung, was treated successfully. The treatment consisted of five courses. For each course, five doses of 62.5-187.5mg chlorin e4 sodium were administered orally over the course of eight days. 24-48
20 hours after each oral dose of chlorin e4 sodium, the patient was treated with laser and LED light having a wavelength of 660nm and 760nm. Additionally an interstitial light treatment using a fine needle was performed on the left lung to a tumour that was large in size with an optical fibre laser having a wavelength of 660nm and 760nm.

25

PET/CT scans showed a very good treatment response. The patient is currently alive and well, one year and five months after starting his treatment and two months after completion of his treatment.

30 Example 11

A 44 year old female patient with lung cancer with metastases in the lymph nodes and sacrum bone, was treated successfully. The patient had an operation to remove

the primary tumour of the lung and completed four cycles of chemotherapy. Subsequent PET/CT scans showed tumours in the lymph nodes of the neck, the lymph nodes in the left arm pit and the sacrum bone. The treatment consisted of five courses. For each course, five doses of 62.5-187.5mg chlorin e4 sodium were administered orally over the course of eight days. 24-48 hours after each oral dose of chlorin e4 sodium, the patient was treated with laser and LED light having a wavelength of 660nm and 760nm. Additionally, for two of the treatment courses, an interstitial light treatment using a fine needle was performed on the sacrum bone with an optical fibre laser having a wavelength of 660nm and 760nm. Additionally, mesothelium bone growth therapy was administered for the last two treatment courses.

PET/CT scans showed a complete response, with no active tumours in the patient's body. The patient is currently alive and well, 10 months after starting her treatment and four months after completion of her treatment.

Example 12

A 52 year old male patient with brain cancer with metastases in the lung and lymph nodes, was treated successfully. An MRI scan showed a large tumour mass in the brain. The patient had an operation to debulk the tumour pressing against his brain, but he did not have any chemotherapy or radiotherapy. Subsequent PET/CT scans showed active tumours in the brain with metastases in both lungs and the lymph nodes in the right cervical and right supraclavicular regions. The patient experienced occasional epileptic seizures and severe headaches. The treatment consisted of two courses. For each course, five doses of 62.5-187.5mg chlorin e4 sodium were administered orally over the course of eight days. 24-48 hours after each oral dose of chlorin e4 sodium, the patient was treated with laser and LED light having a wavelength of 660nm and 760nm.

30

PET/CT scans showed a complete response in the lungs and no metabolic activity in the area of the brain tumour. The lymph nodes were not yet clear, and the patient will undergo a third treatment course.

Example – clinical study with patients suffering from advanced non-small cell lung cancer

- 5 Through pathology or cytology diagnosis, 66 advanced non-small cell lung cancer patients were selected and randomly divided into two groups. Group A had 32 patients who were treated with chlorin e4 sodium. Group B had 34 patients who were treated with both chemotherapy and radiotherapy.
- 10 For group A, chlorin e4 sodium was administered at a dosage of 2 mg per kilogram of body weight, with two thirds administered by inhalation and one third administered orally. After the irradiation target(s) had been determined, an optical fibre for laser treatment purposes was guided into the lung tumor(s) via an interstitial procedure. Irradiation light intensity was 200 J/cm^2 , irradiation power
- 15 was 1,000 mW, and irradiation lasted for eight to ten minutes. Laser irradiation into the tumor was two to five centimeters in depth. Irradiation was either performed phase by phase (sub-paragraph treatment) based on tumor size or performed repeatedly.
- 20 For group B, for the radiotherapy, a 15 MV X-ray radiation therapy irradiation field covered primary lung tumor lesion(s) and mediastinal lymphatic drainage area, with a dosage of 65-70 Gy, administered on every first day in a week of radiotherapy. For the chemotherapy, 20 mg cisplatin was administered by intravenous application.
- 25 One and two years survival rates for groups A and B were 93.75%, 70.60% and 68.75, 32.35% respectively, the difference was statistically significant ($P < 0.05$). The combined complete and partial remission rates for groups A and B were 56.20% and 21.30%, the rate of group A was significantly higher than that of group B, the difference was statistically significant ($P < 0.05$).

Example – study with mice with subcutaneous S180 sarcoma

In this experiment the effects of photodynamic therapy mediated by three different photosensitizers, namely Duteroporphyrin, chlorin e4 sodium and 5-ALA, on mouse
5 S180 sarcoma were observed. Duteroporphyrin and 5-ALA are porphyrin derivatives.

40 Kunming mice with well grown subcutaneous S180 sarcoma were randomly divided into four groups: (A) Duteroporphyrin group (10mg/kg), (B) chlorin e4
10 sodium group (20mg/kg), (C) 5-ALA group (100mg/kg) and (D) blank control group. The mice in groups A and B were subjected to irradiation 8-10 hours after tail vein injection of the photosensitizers, while the mice in group C were subjected to irradiation 3 hours after tail vein injection of the photosensitizer. A laser with a wavelength of 630nm was used for groups A and C, while a dual-frequency laser
15 with wavelengths of 660nm and 760nm was used for group B. The local tumour sites were irradiated vertically, with the diameter of the laser spots being 1.2-1.5cm. The whole tumour and 3-5mm of normal tissue around the tumour were covered by the laser. The lasers were applied to groups A, B and C for 20 minutes, with a power density of 150mW/cm² and an energy density of 180J/cm². The mice in
20 control group D were not subjected to irradiation.

After the treatments, tumour changes in appearance were observed. Black eschars were seen in groups A and B after treatments, while the eschars observed in group C were thinner and brownish red in colour. Skins and fur were fully recovered in all
25 three treatment groups after the eschars had fallen off; group C was the fastest in skin repairing.

From the tumour growth curves it could be seen that there was no obvious difference in sizes of tumours between the treatment groups A, B and C and the
30 control group D six days after the treatments. This was possibly caused by the oedema and eschars caused by the treatment which occurred at the irradiation sites. Observation of the tumours' sizes 21 days after the treatments indicated that the mean volume differences between all treatment groups A, B and C and the control

group D were statistically significant ($P < 0.05$), the mean volume being statistically significantly smaller in treatment groups A, B and C than the control group D. The mean volume difference between group B and group C was also statistically significant ($P < 0.05$), the mean volume being statistically significantly smaller in
5 group B than group C.

Moreover, all treatment groups showed statistically significant tumour inhibition compared to the control group D, and group B showed a statistically significantly better therapeutic effect than group C ($P < 0.05$). The tumour inhibition rates of
10 groups A, B and C were respectively 62.5%, 75.4% and 37.2%.

Based on the analysis of apoptotic and necrotic tumour cells by FCM 24 hours after treatments, the proportions of apoptotic and necrotic tumour cells were a lot higher and living cells were obviously fewer in all treatment groups A, B and C than in the
15 control group D, indicating that all treatment groups showed clear effects in killing cancer cells.

The pathological sections of the tumour tissues were observed via HE staining. The pathological changes of tumour tissues in the treatment groups A, B and C 24 hours
20 after the treatment included: a wide range of tumour necrosis, tumour vascular disruption and neutrophil infiltration.


The mechanisms of killing tumour cells by photodynamic therapy include directly killing tumour cells, damaging tumour vessels and stimulating patients' immune
25 systems. These factors may benefit each other and they are all long-term tumour-controlling mechanisms.

To sum up, this study verified that photodynamic therapy with Duteroporphyrin, chlorin e4 sodium and 5-ALA can inhibit the growth of S180 sarcoma in mice.
30 Chlorin e4 sodium had a similar or slightly better therapeutic effect than Duteroporphyrin, and a statistically significantly better therapeutic effect than 5-ALA.

It will be understood that the present invention has been described above by way of example only. The examples are not intended to limit the scope of the invention. Various modifications and embodiments can be made without departing from the scope and spirit of the invention, which is defined by the following claims only.


Claims

1. Chlorin e4 sodium, comprising less than 10% of other stereoisomers.
- 5 2. Chlorin e4 sodium, comprising chlorin e4 and sodium in a ratio of 1:2 and comprising less than 10% of other stereoisomers.
3. Chlorin e4 sodium, having the structure:



10 and comprising less than 10% of other stereoisomers.

4. Chlorin e4 sodium, having the structure:




and comprising less than 10% of other stereoisomers.


5. A process of preparing chlorin e4 sodium comprising less than 10% of other stereoisomers, comprising the step of reacting chlorin e4 with a sodium compound.

6. The process of claim 5, wherein the chlorin e4 sodium obtained comprises chlorin e4 and sodium in a ratio of 1:2.

7. The process of claim 5 or 6, wherein the chlorin e4 sodium obtained has the structure:



8. The process of any one of claims 5 to 7, wherein the chlorin e4 sodium obtained has the structure:



9. The process of any one of claims 5 to 8, wherein the sodium compound is sodium hydroxide.
- 5 10. The process of any one of claims 5 to 9, wherein chlorin e4 is reacted with the sodium compound in the presence of a base.
11. The process of claim 10, wherein the base is an amine base.
- 10 12. The process of claim 11, wherein the amine base is thiamine pyrophosphate.
13. The process of any one of claims 5 to 12, wherein chlorin e4, the sodium compound and the optional base are used in a molar ratio of 0.15-1 : 45-60 : 0-12.
- 15 14. Chlorin e4 sodium preparable by a process of any one of claims 5 to 13.
15. Chlorin e4 sodium when prepared by a process of any one of claims 5 to 13.
16. The chlorin e4 sodium of any one of claims 1-4 or 14-15, for use in
20 medicine.
17. The chlorin e4 sodium of any one of claims 1-4 or 14-16, for use in photodynamic therapy or cytoluminescent therapy.
- 25 18. The chlorin e4 sodium of any one of claims 1-4 or 14-17, for use in the treatment of atherosclerosis; multiple sclerosis; diabetes; diabetic retinopathy; arthritis; rheumatoid arthritis; a fungal, viral, chlamydial, bacterial, nanobacterial or parasitic infectious disease; HIV; Aids; infection with sars virus, Asian (chicken) flu virus, herpes simplex or herpes zoster; hepatitis; viral hepatitis; a cardiovascular
30 disease; coronary artery stenosis; carotid artery stenosis; intermittent claudication; a dermatological condition; acne; psoriasis; a disease characterised by benign or malignant cellular hyperproliferation or by areas of neovascularisation; a benign or malignant tumour; early cancer; cervical dysplasia; soft tissue sarcoma; a germ cell

tumour; retinoblastoma; age-related macular degeneration; lymphoma; Hodgkin's lymphoma; head and neck cancer; oral or mouth cancer; or cancer of the blood, prostate, cervix, uterus, vaginal or other female adnexa, breast, naso-pharynx, trachea, larynx, bronchi, bronchioles, lung, hollow organs, esophagus, stomach, bile duct, intestine, colon, colorectum, rectum, bladder, ureter, kidney, liver, gall bladder, spleen, brain, lymphatic system, bones, skin or pancreas.

19. The chlorin e4 sodium of any one of claims 1-4 or 14-18, for use in the treatment of a disease characterised by benign or malignant cellular hyperproliferation or by areas of neovascularisation.

20. The chlorin e4 sodium of any one of claims 1-4 or 14-19, for use in the treatment of a benign or malignant tumour.

21. The chlorin e4 sodium of any one of claims 1-4 or 14-20, for use in the treatment of early cancer; cervical dysplasia; soft tissue sarcoma; a germ cell tumour; retinoblastoma; age-related macular degeneration; lymphoma; Hodgkin's lymphoma; head and neck cancer; oral or mouth cancer; or cancer of the blood, prostate, cervix, uterus, vaginal or other female adnexa, breast, naso-pharynx, trachea, larynx, bronchi, bronchioles, lung, hollow organs, esophagus, stomach, bile duct, intestine, colon, colorectum, rectum, bladder, ureter, kidney, liver, gall bladder, spleen, brain, lymphatic system, bones, skin or pancreas.

22. The chlorin e4 sodium of any one of claims 1-4 or 14-16, for use in photodynamic diagnosis.

23. The chlorin e4 sodium of any one of claims 16 to 22, wherein the chlorin e4 sodium is adapted for administration prior to administration of irradiation.

24. The chlorin e4 sodium of claim 23, wherein the irradiation is electromagnetic radiation with a wavelength in the range of from 500nm to 1000nm.

25. A pharmaceutical composition comprising chlorin e4 sodium of any one of claims 1-4 or 14-24 and a pharmaceutically acceptable carrier or diluent.
26. The pharmaceutical composition of claim 25, for use in photodynamic
5 therapy or cytoluminescent therapy.
27. The pharmaceutical composition of claim 25 or 26, for use in the treatment of atherosclerosis; multiple sclerosis; diabetes; diabetic retinopathy; arthritis; rheumatoid arthritis; a fungal, viral, chlamydial, bacterial, nanobacterial or parasitic
10 infectious disease; HIV; Aids; infection with sars virus, Asian (chicken) flu virus, herpes simplex or herpes zoster; hepatitis; viral hepatitis; a cardiovascular disease; coronary artery stenosis; carotid artery stenosis; intermittent claudication; a dermatological condition; acne; psoriasis; a disease characterised by benign or malignant cellular hyperproliferation or by areas of neovascularisation; a benign or
15 malignant tumour; early cancer; cervical dysplasia; soft tissue sarcoma; a germ cell tumour; retinoblastoma; age-related macular degeneration; lymphoma; Hodgkin's lymphoma; head and neck cancer; oral or mouth cancer; or cancer of the blood, prostate, cervix, uterus, vaginal or other female adnexa, breast, naso-pharynx, trachea, larynx, bronchi, bronchioles, lung, hollow organs, esophagus, stomach, bile
20 duct, intestine, colon, colorectum, rectum, bladder, ureter, kidney, liver, gall bladder, spleen, brain, lymphatic system, bones, skin or pancreas.
28. The pharmaceutical composition of any one of claims 25 to 27, for use in the treatment of a disease characterised by benign or malignant cellular
25 hyperproliferation or by areas of neovascularisation.
29. The pharmaceutical composition of any one of claims 25 to 28, for use in the treatment of a benign or malignant tumour.
30. The pharmaceutical composition of any one of claims 25 to 29, for use in the
30 treatment of early cancer; cervical dysplasia; soft tissue sarcoma; a germ cell tumour; retinoblastoma; age-related macular degeneration; lymphoma; Hodgkin's lymphoma; head and neck cancer; oral or mouth cancer; or cancer of the blood, prostate,

cervix, uterus, vaginal or other female adnexa, breast, naso-pharynx, trachea, larynx, bronchi, bronchioles, lung, hollow organs, esophagus, stomach, bile duct, intestine, colon, colorectum, rectum, bladder, ureter, kidney, liver, gall bladder, spleen, brain, lymphatic system, bones, skin or pancreas.

5

31. The pharmaceutical composition of claim 25, for use in photodynamic diagnosis.

32. The pharmaceutical composition of any one of claims 25 to 31, wherein the
10 pharmaceutical composition is adapted for administration prior to administration of irradiation.

33. The pharmaceutical composition claim 32, wherein the irradiation is electromagnetic radiation with a wavelength in the range of from 500nm to 1000nm.

15

34. The pharmaceutical composition of any one of claims 25 to 33, wherein the pharmaceutical composition is in a form suitable for oral, parenteral (including intravenous, subcutaneous, intramuscular, intradermal, intratracheal, intraperitoneal, intraarticular, intraabdominal, intracranial and epidural), transdermal, airway
20 (aerosol), rectal, vaginal or topical (including buccal, mucosal and sublingual) administration.

35. The pharmaceutical composition of claim 34, wherein the pharmaceutical composition is in a form suitable for oral or parenteral administration.

25

36. Use of chlorin e4 sodium of any one of claims 1-4 or 14-24 in the manufacture of a medicament for the treatment of a benign or malignant tumour; early cancer; cervical dysplasia; soft tissue sarcoma; a germ cell tumour; retinoblastoma; age-related macular degeneration; lymphoma; Hodgkin's lymphoma;
30 head and neck cancer; oral or mouth cancer; or cancer of the blood, prostate, cervix, uterus, vaginal or other female adnexa, breast, naso-pharynx, trachea, larynx, bronchi, bronchioles, lung, hollow organs, esophagus, stomach, bile duct, intestine,

colon, colorectum, rectum, bladder, ureter, kidney, liver, gall bladder, spleen, brain, lymphatic system, bones, skin or pancreas.

37. Use of chlorin e4 sodium of any one of claims 1-4 or 14-24 in the
5 manufacture of a phototherapeutic agent for use in photodynamic therapy or cytoluminescent therapy.

38. The use of claim 37, wherein the phototherapeutic agent is for the treatment
of a benign or malignant tumour; early cancer; cervical dysplasia; soft tissue
10 sarcoma; a germ cell tumour; retinoblastoma; age-related macular degeneration; lymphoma; Hodgkin's lymphoma; head and neck cancer; oral or mouth cancer; or cancer of the blood, prostate, cervix, uterus, vaginal or other female adnexa, breast, naso-pharynx, trachea, larynx, bronchi, bronchioles, lung, hollow organs, esophagus, stomach, bile duct, intestine, colon, colorectum, rectum, bladder, ureter, kidney,
15 liver, gall bladder, spleen, brain, lymphatic system, bones, skin or pancreas.

39. The use of any one of claims 36 to 38, wherein the medicament or the phototherapeutic agent is for the treatment of a benign or malignant tumour.

40. The use of any one of claims 36 to 39, wherein the medicament or the
20 phototherapeutic agent is for the treatment of early cancer; cervical dysplasia; soft tissue sarcoma; a germ cell tumour; retinoblastoma; age-related macular degeneration; lymphoma; Hodgkin's lymphoma; head and neck cancer; oral or mouth cancer; or cancer of the blood, prostate, cervix, uterus, vaginal or other
25 female adnexa, breast, naso-pharynx, trachea, larynx, bronchi, bronchioles, lung, hollow organs, esophagus, stomach, bile duct, intestine, colon, colorectum, rectum, bladder, ureter, kidney, liver, gall bladder, spleen, brain, lymphatic system, bones, skin or pancreas.

41. Use of chlorin e4 sodium of any one of claims 1-4 or 14-24 in the
30 manufacture of a photodiagnostic agent for use in photodynamic diagnosis.

42. The use of any one of claims 36 to 41, wherein the medicament, the phototherapeutic agent or the photodiagnostic agent is adapted for administration prior to administration of irradiation.

5 43. The use of claim 42, wherein the irradiation is electromagnetic radiation with a wavelength in the range of from 500nm to 1000nm.