



159159

COMPOUNDS FOR TREATMENT OF INFLAMMATION, DIABETES AND RELATED DISORDERS

CROSS REFERENCE TO RELATED APPLICATION

[0001] This application claims priority from U.S. Provisional Application No. 60/334,818, filed November 29, 2001, and is incorporated herein, in its entirety, by reference.

FIELD OF THE INVENTION

[0002] The invention is directed to compounds, for example, heterocyclic derivatives of acyl urea, thiourea, carbamate and thiocarbamate compounds, that provide a variety of useful pharmacological effects. The compounds are useful, for example, in lowering blood glucose levels in hyperglycemic disorders, such as diabetes mellitus, and for treating related disorders, such as obesity and hyperlipidemia. Furthermore, these compounds are useful for treatment of disorders associated with insulin resistance, such as polycystic ovary syndrome, and for the treatment of inflammation, inflammatory and immunological diseases, particularly those mediated by pro-inflammatory cytokines (such as TNF-alpha, IL-1 beta and IL-6), type 4 and type 3 phosphodiesterase (PDE4 and PDE3, respectively), p44/42 mitogen-activated protein (MAP) kinase, cyclooxygenase-2 (COX-2) and/or inducible nitric oxide synthase (iNOS).

BACKGROUND

[0003] The causes of diabetes mellitus are not yet known, although both genetics and environment seem to be major factors. Type 1 diabetes, also known as insulin-dependent diabetes mellitus (IDDM), is an autoimmune disease in which the responsible autoantigen is still unidentified. Since their insulin-producing pancreatic cells are destroyed, Type 1 diabetics need to take insulin parenterally to survive. On the other hand, type 2 diabetes, also called non-insulin-dependent diabetes mellitus (NIDDM), the more common form, is a metabolic disorder resulting from the body's inability to make a sufficient amount of insulin or to properly use the insulin that is produced. Impaired insulin secretion and insulin resistance are considered the major



G00002

defects; however, the precise genetic factors involved in the mechanism remain unknown.

[0004] Other than insulin administered parenterally and as shown in Table 1, there are generally four major classes of oral hypoglycemic agents currently used in the treatment of diabetes mellitus:

Class	Approved Drugs	Mechanisms of Action	Limitations
Sulfonylurea	four (1st generation) and two (2nd generation)	stimulates pancreas to release more insulin	hypoglycemia; may increase cardiovascular risk; contra-indicated in liver and renal dysfunction; hyperinsulinemia
Biguanide	metformin	reduces glucose production by liver; improves insulin sensitivity	lactic acidosis; GI side effects
Alpha-glucosidase inhibitor	acarbose	reduces glucose absorption by gut	GI side effects; requires frequent postprandial dosing
Thiazolidinedione	troglitazone (withdrawn) rosiglitazone pioglitazone	stimulates nuclear PPAR-gamma receptor; reduces insulin resistance	edema; contra-indicated in heart failure; long onset of action; weight gain; frequent liver function testing

TABLE 1

[0005] As is shown in the above table, each of the current agents available for use in treatment of diabetes mellitus has several disadvantages. Accordingly, there is a need for the identification and development of new agents, particularly, water

soluble agents which can be orally administered, for use in the treatment of diabetes mellitus and other hyperglycemic disorders.

[0006] Moreover, while the thiazolidinedione class has gained more widespread use in recent years as insulin sensitizers to combat "insulin resistance", a condition in which the patient becomes less responsive to the effects of insulin, there is a need for frequent liver testing for patients using these compounds. In fact, the known thiazolidinediones are not effective for a significant portion of the patient population. In addition, the first drug in this class to be approved by the FDA, troglitazone, was withdrawn from the market due to problems of liver toxicity. Thus, there is a continuing need for nontoxic, more widely effective insulin sensitizers.

[0007] As indicated above, the invention is also directed to the treatment of immunological diseases or inflammation, in particular, such diseases as are mediated by cytokines, COX-2 and iNOS. The principal elements of the immune system are macrophages or antigen-presenting cells, T cells and B cells. Macrophages are important mediators of inflammation and also provide the necessary "help" for T cell stimulation and proliferation. For example, macrophages make the cytokines IL-1, IL-12 and TNF-alpha, all of which are potent pro-inflammatory molecules. Cytokine production may lead to the secretion of other cytokines, altered cellular function, cell division or differentiation. In addition, activation of macrophages results in the induction of enzymes, such as COX-2 and iNOS, and in the production of free radicals capable of damaging normal cells. Many factors activate macrophages, including bacterial products, superantigens and interferon gamma. It is believed that phosphotyrosine kinases and other cellular kinases are involved in the activation process. Since macrophages are sentinel to the development of an immune response, agents that modify their function, specifically their cytokine secretion profile, are likely to determine the direction and potency of the immune response.

[0008] Inflammation is the body's normal response to injury or infection. However, in inflammatory diseases such as rheumatoid arthritis, pathologic inflammatory processes can lead to morbidity and mortality. The cytokine tumor necrosis factor-alpha (TNF-alpha) plays a central role in the inflammatory response

and has been targeted as a point of intervention in inflammatory disease. TNF-alpha is a polypeptide hormone released by activated macrophages and other cells. At low concentrations, TNF-alpha participates in the protective inflammatory response by activating leukocytes and promoting their migration to extravascular sites of inflammation (Moser et al., *J Clin Invest*, 83:444-55, 1989). At higher concentrations, TNF-alpha can act as a potent pyrogen and induce the production of other pro-inflammatory cytokines (Haworth et al., *Eur J Immunol*, 21:2575-79, 1991; Brennan et al., *Lancet*, 2:244-7, 1989). TNF-alpha also stimulates the synthesis of acute-phase proteins. In rheumatoid arthritis, a chronic and progressive inflammatory disease affecting about 1% of the adult U.S. population, TNF-alpha mediates the cytokine cascade that leads to joint damage and destruction (Arend et al., *Arthritis Rheum*, 38:151-60, 1995). Inhibitors of TNF-alpha, including soluble TNF receptors (etanercept) (Goldenberg, *Clin Ther*, 21:75-87, 1999) and anti-TNF-alpha antibody (infliximab) (Luong et al., *Ann Pharmacother*, 34:743-60, 2000), the contents of each of which are incorporated herein by reference, have recently been approved by the U.S. Food and Drug Administration (FDA) as agents for the treatment of rheumatoid arthritis.

[0009] Elevated levels of TNF-alpha have also been implicated in many other disorders and disease conditions, including cachexia (Fong et al., *Am J Physiol*, 256:R659-65, 1989), septic shock syndrome (Tracey et al., *Proc Soc Exp Biol Med*, 200:233-9, 1992), osteoarthritis (Venn et al., *Arthritis Rheum*, 36:819-26, 1993), inflammatory bowel disease such as Crohn's disease and ulcerative colitis (Murch et al., *Gut*, 32:913-7, 1991), Behcet's disease (Akoglu et al., *J Rheumatol*, 17:1107-8, 1990), Kawasaki disease (Matsubara et al., *Clin Immunol Immunopathol*, 56:29-36, 1990), cerebral malaria (Grau et al., *N Engl J Med*, 320:1586-91, 1989), adult respiratory distress syndrome (Millar et al., *Lancet* 2:712-4, 1989), asbestosis and silicosis (Bissonnette et al., *Inflammation*, 13:329-39, 1989), pulmonary sarcoidosis (Baughman et al., *J Lab Clin Med*, 115:36-42, 1990), asthma (Shah et al., *Clin Exp Allergy*, 25:1038-44, 1995), AIDS (Dezube et al., *J Acquir Immune Defic Syndr*, 5:1099-104, 1992), meningitis (Waage et al., *Lancet*, 1:355-7, 1987), psoriasis (Oh et al., *J Am Acad Dermatol*, 42:829-30, 2000), spondyloarthritides such as ankylosing spondylitis (Braun et al., *Curr Opin Rheumatol* 13:245-9, 2001; Marzo-Ortega et al., *Arthritis Rheum* 44:2112-7, 2001), graft versus host reaction (Nestel et

al., J Exp Med, 175:405-13, 1992), multiple sclerosis (Sharief et al., N Engl J Med, 325:467-72, 1991), systemic lupus erythematosus (Maury et al., Int J Tissue React, 11:189-93, 1989), diabetes (Hotamisligil et al., Science, 259:87-91, 1993) and atherosclerosis (Bruunsgaard et al., Clin Exp Immunol, 121:255-60, 2000), the contents of each of which are incorporated herein by reference. It can be seen from the references cited above that inhibitors of TNF-alpha are potentially useful in the treatment of a wide variety of diseases.

[00010] Interleukin-6 (IL-6) is another pro-inflammatory cytokine that exhibits pleiotropy and redundancy of action. IL-6 participates in the immune response, inflammation and hematopoiesis. It is a potent inducer of the hepatic acute phase response and is a powerful stimulator of the hypothalamic-pituitary-adrenal axis that is under negative control by glucocorticoids. IL-6 promotes the secretion of growth hormone but inhibits release of thyroid stimulating hormone. Elevated levels of IL-6 are seen in several inflammatory diseases, and inhibition of the IL-6 cytokine subfamily has been suggested as a strategy to improve therapy for rheumatoid arthritis (Carroll et al., Inflamm Res, 47:1-7, 1998). In addition, IL-6 has been implicated in the progression of atherosclerosis and the pathogenesis of coronary heart disease (Yudkin et al., Atherosclerosis, 148:209-14, 1999). Overproduction of IL-6 is also seen in steroid withdrawal syndrome, conditions related to deregulated vasopressin secretion, and osteoporosis associated with increased bone resorption, such as in cases of hyperparathyroidism and sex-steroid deficiency (Papanicolaou et al., Ann Intern Med, 128:127-37, 1998). Since excessive production of IL-6 is implicated in several disease states, it is highly desirable to develop compounds that inhibit IL-6 secretion.

[00011] The cytokine IL-1 beta also participates in the inflammatory response. It stimulates thymocyte proliferation, fibroblast growth factor activity, and the release of prostaglandin from synovial cells. Elevated or unregulated levels of the cytokine IL-1 beta have been associated with a number of inflammatory diseases and other disease states, including but not limited to adult respiratory distress syndrome (Meduri et al, Chest 107:1062-73, 1995), allergy (Hastie et al, Cytokine 8:730-8, 1996), Alzheimer's disease (O'Barr et al, J Neuroimmunol 109:87-94, 2000), anorexia (Laye et al, Am J Physiol Regul Integr Comp Physiol 279:R93-8, 2000),

asthma (Sousa et al, Thorax 52:407-10, 1997), atherosclerosis (Dewberry et al, Arterioscler Thromb Vasc Biol 20:2394-400, 2000), brain tumors (Ilyin et al, Mol Chem Neuropathol 33:125-37, 1998), cachexia (Nakatani et al, Res Commun Mol Pathol Pharmacol 102:241-9, 1998), carcinoma (Ikemoto et al, Anticancer Res 20:317-21, 2000), chronic arthritis (van den Berg et al, Clin Exp Rheumatol 17:S105-14, 1999), chronic fatigue syndrome (Cannon et al, J Clin Immunol 17:253-61, 1997), CNS trauma (Herx et al, J Immunol 165:2232-9, 2000), epilepsy (De Simoni et al, Eur J Neurosci 12:2623-33, 2000), fibrotic lung diseases (Pan et al, Pathol Int 46:91-9, 1996), fulminant hepatic failure (Sekiyama et al, Clin Exp Immunol 98:71-7, 1994), gingivitis (Biesbrock et al, Monogr Oral Sci 17:20-31, 2000), glomerulonephritis (Kluth et al, J Nephrol 12:66-75, 1999), Guillain-Barre syndrome (Zhu et al, Clin Immunol Immunopathol 84:85-94, 1997), heat hyperalgesia (Oprea et al, J Neurosci 20:6289-93, 2000), hemorrhage and endotoxemia (Parsey et al, J Immunol 160:1007-13, 1998), inflammatory bowel disease (Olson et al, J Pediatr Gastroenterol Nutr 16:241-6, 1993), leukemia (Estrov et al, Leuk Lymphoma 24:379-91, 1997), leukemic arthritis (Rudwaleit et al, Arthritis Rheum 41:1695-700, 1998), systemic lupus erythematosus (Mao et al, Autoimmunity 24:71-9, 1996), multiple sclerosis (Martin et al, J Neuroimmunol 61:241-5, 1995), osteoarthritis (Hernvann et al, Osteoarthritis Cartilage 4:139-42, 1996), osteoporosis (Zheng et al, Maturitas 26:63-71, 1997), Parkinson's disease (Bessler et al, Biomed Pharmacother 53:141-5, 1999), POEMS syndrome (Gherardi et al, Blood 83:2587-93, 1994), pre-term labor (Dudley, J Reprod Immunol 36:93-109, 1997), psoriasis (Bonifati et al, J Biol Regul Homeost Agents 11:133-6, 1997), reperfusion injury (Clark et al, J Surg Res 58:675-81, 1995), rheumatoid arthritis (Seitz et al, J Rheumatol 23:1512-6, 1996), septic shock (van Deuren et al, Blood 90:1101-8, 1997), systemic vasculitis (Brooks et al, Clin Exp Immunol 106:273-9, 1996), temporal mandibular joint disease (Nordahl et al, Eur J Oral Sci 106:559-63, 1998), tuberculosis (Tsao et al, Tuber Lung Dis 79:279-85, 1999), viral rhinitis (Roseler et al, Eur Arch Otorhinolaryngol Suppl 1:S61-3, 1995), the contents of each of which are incorporated herein by reference, and pain and/or inflammation resulting from strain, sprain, trauma, surgery, infection or other disease processes. Since overproduction of IL-1 beta is associated with numerous disease conditions, it is desirable to develop compounds that inhibit the production or activity of IL-1 beta.

[00012] Cyclooxygenase is an enzyme that catalyzes a rate-determining step in the biosynthesis of prostaglandins, which are important mediators of inflammation and pain. The enzyme occurs as at least two distinct isomers, cyclooxygenase-1 (COX-1) and cyclooxygenase-2 (COX-2). The COX-1 isomer is constitutively expressed in the gastric mucosa, platelets and other cells and is involved in the maintenance of homeostasis in mammals, including protecting the integrity of the digestive tract. The COX-2 isomer, on the other hand, is not constitutively expressed but rather is induced by various agents, such as cytokines, mitogens, hormones and growth factors. In particular, COX-2 is induced during the inflammatory response (DeWitt DL, *Biochim Biophys Acta*, 1083:121-34, 1991; Seibert et al., *Receptor*, 4:17-23, 1994.). Aspirin and other conventional non-steroid anti-inflammatory drugs (NSAIDs) are non-selective inhibitors of both COX-1 and COX-2. They can be effective in reducing inflammatory pain and swelling, but since they hamper the protective action of COX-1, they produce undesirable side effects of gastrointestinal pathology. Therefore, agents that selectively inhibit COX-2 but not COX-1 are preferable for treatment of inflammatory diseases. Recently, a diarylpyrazole sulfonamide (celecoxib) that selectively inhibits COX-2 has been approved by the FDA for use in the treatment of osteoarthritis and adult rheumatoid arthritis (Luong et al., *Ann Pharmacother*, 34:743-60, 2000; Penning et al., *J Med Chem*, 40:1347-65, 1997). Another selective COX-2 inhibitor, rofecoxib, has been approved by the FDA for treatment of osteoarthritis, acute pain and primary dysmenorrhea (Scott and Lamb, *Drugs*, 58:499-505, 1999; Morrison et al., *Obstet Gynecol*, 94:504-8, 1999; Saag et al, *Arch Fam Med*, 9:1124-34, 2000). COX-2 is also expressed in many cancers and precancerous lesions, and there is mounting evidence that selective COX-2 inhibitors may be useful for treating and preventing colorectal, breast and other cancers (Taketo MM, *J Natl Cancer Inst*, 90:1609-20, 1998; Fournier et al., *J Cell Biochem Suppl*, 34:97-102, 2000; Masferrer et al., *Cancer Res*, 60:1306-11, 2000), the contents of each of which are incorporated herein by reference. In 1999 celecoxib was approved by the FDA as an adjunct to usual care for patients with familial adenomatous polyposis, a condition which, left untreated, generally leads to colorectal cancer.

[00013] Production of nitric oxide by iNOS has been associated with both beneficial and detrimental effects in inflammation, inflammatory diseases and related

disorders. For example, deleterious effects have been implicated in the pathogenesis of abdominal aortic aneurysms (Johanning et al, *J Vasc Surg* 33:579-86, 2001), acute endotoxemia (Henningsson et al, *Am J Physiol Cell Physiol* 280:C1242-54, 2001), amyotrophic lateral sclerosis (Sasaki et al, *Neurosci Lett* 291:44-8, 2000), atherosclerosis (Behr-Roussel et al, *Circulation* 102:1033-8, 2000), bladder cancer (Wolf et al, *Virchows Arch* 437:662-6, 2000), colon cancer (Watanabe et al, *Biofactors* 12:129-33, 2000), cystitis (Alfieri et al, *Naunyn Schmiedeberg's Arch Pharmacol* 363:353-7, 2001), HIV-1 encephalitis (Zhao et al, *J Neuroimmunol* 115:182-91, 2001), inflammatory bowel disease (Singer et al, *Gastroenterology* 111:871-85, 1996), multiple sclerosis (Pozza et al, *Brain Res* 855:39-46, 2000), osteoarthritis (Pelletier et al, *Osteoarthritis Cartilage* 7:416-8, 1999), osteoporosis (Armour et al, *J Bone Miner Res* 14:2137-42, 1999), portal hypertension (Schroeder et al, *Dig Dis Sci* 45:2405-10, 2000), pulmonary edema in endotoxin shock (Lee et al, *Clin Exp Pharmacol Physiol* 28:315-20, 2001), rheumatoid arthritis (van't Hof et al, *Rheumatology (Oxford)* 39:1004-8, 2000), sepsis (Nishina et al, *Anesth Analg* 92:959-66, 2001), severe burn/smoke inhalation injury (Soejima et al, *Am J Respir Crit Care Med* 163:745-52, 2001), and ulcerative colitis (Ikeda et al, *Am J Gastroenterol* 92:1339-41, 1997), the contents of each of which are incorporated herein by reference. Since the production of nitric oxide by iNOS has been implicated in the pathogenesis of inflammatory and related immunological diseases, it is desirable to develop compounds that inhibit iNOS activity or expression.

[00014] Phosphodiesterases (PDEs) are responsible for the hydrolysis of intracellular cyclic adenosine and guanosine monophosphate (cAMP and cGMP), which converts these second messengers into their inactive forms. There are 11 major families of PDEs, designated PDE1 to PDE11. Type 4 phosphodiesterase (PDE4) is found in airway smooth muscle cells and in immune and inflammatory cells. PDE4 activity has been associated with a wide variety of inflammatory and autoimmune diseases, and PDE4 inhibitors have been studied as potential therapeutic agents for such diseases as asthma, chronic obstructive pulmonary disease, rheumatoid arthritis, multiple sclerosis and type 2 diabetes (Burnouf and Pruniaux, *Current Pharm Des*, 8:1255-96, 2002; Dal Piaz and Giovannoni, *Eur J Med Chem*, 35:463-80, 2000). Type 3 phosphodiesterase (PDE3) is localized in platelets and cardiac and vascular smooth muscle cells. Inhibitors of PDE3 have

been proposed as possible drugs for the treatment of acute respiratory distress syndrome (Schermyly et al, J Pharmacol Exp Ther, 292:512-20, 2000), cancer (Shimizu et al, Anticancer Drugs, 13:875-80, 2002; Murata et al, Anticancer Drugs, 12:79-83, 2001), cardiomyopathy (Alharethi and Movsesian, Expert Opin Investig Drugs, 11:1529-36, 2002), congestive heart failure (Movsesian, J Am Coll Cardiol, 34:318-24, 1999), erectile dysfunction (Kuthe et al, Curr Opin Investig Drugs, 3:1489-95, 2002), and T-cell-mediated autoimmune disorders (Bielekova et al, J Immunol 164:1117-24, 2000), the contents of each of which are incorporated herein by reference.





[00015] Activation of lymphocyte and macrophage immune response to pathogens involve complex intracellular signaling pathways involving a cascade of various phosphorylating enzymes, kinases that ultimately regulate cytokine production and cell apoptosis. Key kinases include p44/42 MAP kinase (also known as ERK1/ERK2), P38 MAP kinase, MEK, and IRAK/NFkB. While different processes utilize different aspects of the pathway, the bacterial coat-derived protein LPS has been shown to activate multiple mitogen-activated protein kinases, including the extracellular signal-regulated receptor kinases ERK1 and ERK2. LPS-induced TNF-alpha production by human monocytes involves activation of ERK1/ERK2 (van der Bruggen et al, Infect Immun, 67:3824-9, 1999). As TNF-alpha is a key mediator of autoimmune disease, blocking the ERK pathway has potential for the treatment of inflammatory and immunological diseases such as lupus (Yi et al, J Immunol, 165:6627-34, 2000), rheumatoid arthritis (Neff et al, Cell Microbiol, 3:703-12, 2001; Schett et al, Arthritis Rheum, 43:2501-12, 2000), psoriasis (van der Bruggen et al, Infect Immun, 67:3824-9, 1999) and destruction of pancreatic islet beta cells in Type I diabetes (Pavlovic et al, Eur Cytokine Netw 11:267-74, 2000), the contents of each of which are incorporated herein by reference.







[00016] It will be appreciated from the foregoing that, while there have been extensive prior efforts to provide compounds for inhibiting, for example, TNF-alpha, IL-1 beta, IL-6, COX-2, PDE4 or other agents considered responsible for inflammation or inflammatory diseases, e.g. arthritis, there still remains a need for new and improved compounds for effectively treating or inhibiting such diseases. A principal object of the invention is to provide compounds which are effective for such


treatments as well as for the treatment of, for example, diabetes, coronary heart disease, insulin resistance and related disorders.

SUMMARY OF THE INVENTION

[00017] The invention is directed to compounds, for example, heterocyclic derivatives of acyl urea, thiourea, carbamate and thiocarbamate compounds, for providing a variety of useful pharmacological effects. The compounds are useful, for example, in lowering blood glucose levels in hyperglycemic disorders, such as diabetes mellitus, and for treating related disorders, such as obesity and hyperlipidemia. Furthermore, these compounds are useful for treatment of disorders associated with insulin resistance, such as polycystic ovary syndrome, and for the treatment of inflammation and immunological diseases, particularly those mediated by pro-inflammatory cytokines (such as TNF-alpha, IL-1 beta and IL-6), type 4 phosphodiesterase (PDE4), type 3 phosphodiesterase (PDE3), p44/42 mitogen activated protein (MAP) kinase, cyclooxygenase-2 (COX-2) and/or inducible nitric oxide synthase (iNOS). In particular, the invention discloses compounds of the Formulas I-XIII as well as the pharmaceutically acceptable salts and solvates thereof:







[00018] wherein the stereocenters marked with an asterisk (*) may be R- or S-; the bond represented by a dashed line plus a solid line may be a double bond or a single bond, and when the bond is a double bond it may be in the E or Z configuration, and when the bond is a single bond the resulting stereocenters may have the R- or S- configuration; and

[00019] R₁, R₂, R₃, R₄, R₅, R₆ and R₇ are each independently selected from the group consisting of

H; optionally substituted C₁-C₂₀ linear or branched alkyl including chloroalkyl or fluoroalkyl; optionally substituted C₂-C₂₀ linear or branched alkenyl; optionally substituted C₆-C₂₀ aryl, linear or branched alkylaryl, linear or branched alkenylaryl; COOR where R is H, optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl or optionally substituted C₆-C₁₀ aryl, sodium, potassium or other pharmaceutically acceptable counter-ion such as calcium, magnesium, ammonium, tromethamine and the like; CONR'R'', where R' and R'' are independently H, optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl or optionally substituted C₆-C₁₀ aryl or where NR'R'' represents a cyclic moiety such as morpholine, piperidine, piperazine and the like; optionally substituted C₁-C₆

amidoalkyl; NH₂; C₁-C₂₀ alkylamino, bis(alkylamino), cycloalkylamino or cyclic amino; OH; optionally substituted C₁-C₂₀ alkoxy including trifluoromethoxy and the like; optionally substituted C₁-C₂₀ alkanoyl; optionally substituted C₁-C₂₀ acyloxy; halo; optionally substituted C₁-C₂₀ alkylcarboxylamino; cyano; nitro; SO₂NR'''R'''' where R''' and R'''' are independently H, C₁-C₂₀ alkyl or aryl; SO₂R''' where R''' is H, C₁-C₂₀ alkyl or aryl; SO₃R''' where R''' is H, C₁-C₂₀ alkyl or aryl; and C₄-C₈ heterocycles such as tetrazolyl, imidazolyl, pyrrolyl, pyridyl, indolyl and the like; and wherein when individual aromatic rings possess adjacent substituents, these substituents may be joined to form a ring such as a methylenedioxy or ethylenedioxy group, and the like, including lactones and lactams;

[00020] R₈ and R₉ are each independently selected from the group consisting of

H; optionally substituted C₁-C₂₀ linear or branched alkyl; optionally substituted C₂-C₂₀ linear or branched alkenyl; optionally substituted C₆-C₁₀ aryl or heteroaryl; COOR where R is H, optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl or optionally substituted C₆-C₁₀ aryl, sodium, potassium or other pharmaceutically acceptable counter-ion such as calcium, magnesium, ammonium, tromethamine and the like; CONR'R'', where R' and R'' are independently H, alkoxy, optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl, optionally substituted C₃-C₁₀ cycloalkyl or cycloalkenyl or optionally substituted C₆-C₁₀ aryl or heteroaryl, preferably 2-, 3- or 4-pyridyl; or where NR'R'' represents a cyclic moiety such as morpholine, piperidine, hydroxypiperidine, imidazole, piperazine, methylpiperazine and the like; NH₂; C₁-C₂₀ alkylamino, bis(alkylamino), cycloalkylamino or cyclic amino; OH; C₁-C₂₀ alkoxy; C₁-C₂₀ alkanoyl; C₁-C₂₀ acyloxy; halo; C₁-C₂₀ alkylcarboxylamino; cyano; nitro; SO₂NR'''R'''' where R''' and R'''' are independently H, C₁-C₂₀ alkyl or aryl; SO₂R''' where R''' is H, C₁-C₂₀ alkyl or aryl; SO₃R''' where R''' is H, C₁-C₂₀ alkyl or aryl; and tetrazolyl; and wherein R₈ and R₉ together may be joined to form a C₄-C₈ heterocyclic ring, including lactone or lactam;

[00021] R₁₀ and R₁₁ are each independently selected from the group consisting of

H; optionally substituted C₁-C₂₀ linear or branched alkyl; optionally substituted C₂-C₂₀ linear or branched alkenyl; optionally substituted C₆-C₁₀ aryl or heteroaryl; COOR where R is H, optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl or optionally substituted C₆-C₁₀ aryl, sodium, potassium or other pharmaceutically acceptable counter-ion such as calcium, magnesium, ammonium, tromethamine and the like; CONR'R'', where R' and R'' are independently H, optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl or optionally substituted C₆-C₁₀ aryl or where NR'R'' represents a cyclic moiety such as morpholine, piperidine, piperazine and the like; NH₂; C₁-C₂₀ alkylamino, bis(alkylamino), cycloalkylamino or cyclic amino; OH; C₁-C₂₀ alkoxy; C₁-C₂₀ alkanoyl; C₁-C₂₀ acyloxy; halo; C₁-C₂₀ alkylcarboxylamino; cyano; nitro; SO₂NR'''R'''' where R''' and R'''' are independently H, C₁-C₂₀ alkyl or aryl; SO₂R''' where R''' is H, C₁-C₂₀ alkyl or aryl; SO₃R''' where R''' is H, C₁-C₂₀ alkyl or aryl; and tetrazolyl; and wherein R₁₀ and R₁₁ together may be joined to form a C₄-C₈ heterocyclic ring, including lactone or lactam;

[00022] R₁₂, R₁₃, R₁₈, R₁₉ and R₂₀ are each independently selected from the group consisting of

H; optionally substituted C₁-C₂₀ linear or branched alkyl; optionally substituted C₂-C₂₀ linear or branched alkenyl; optionally substituted C₆-C₁₀ aryl or heteroaryl; COOR where R is optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl or optionally substituted C₆-C₁₀ aryl, sodium, potassium or other pharmaceutically acceptable counter-ion such as calcium, magnesium, ammonium, tromethamine and the like; CONR'R'', where R' and R'' are independently H, optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl or optionally substituted C₆-C₁₀ aryl or where NR'R'' represents a cyclic moiety such as morpholine, piperidine, piperazine and the like; C₁-C₂₀ alkanoyl; C₁-C₂₀ alkylamido; C₆-C₂₀ aroyl or heteroaroyl; SO₂R''' where R''' is H, C₁-C₂₀ alkyl or aryl; morpholinocarbonylmethyl; piperazinocarbonylmethyl; and piperadinocarbonylmethyl;

[00023] R_{12} and R_{13} may be absent, or R_{12} and R_{13} together may be an optionally substituted heterocyclic ring, preferably morpholine, piperidine, piperazine, and N-methyl piperidine.

[00024] R_{14} is selected from the group consisting of

H; optionally substituted C_1 - C_{20} linear or branched alkyl including chloroalkyl and fluoroalkyl; optionally substituted C_2 - C_{20} linear or branched alkenyl; optionally substituted C_6 - C_{10} aryl or heteroaryl; COOR where R is H, optionally substituted C_1 - C_{20} alkyl, optionally substituted C_2 - C_{20} alkenyl or optionally substituted C_6 - C_{10} aryl, sodium, potassium or other pharmaceutically acceptable counter-ion such as calcium, magnesium, ammonium, tromethamine and the like; CONR'R", where R' and R" are independently H, optionally substituted C_1 - C_{20} alkyl, optionally substituted C_2 - C_{20} alkenyl or optionally substituted C_6 - C_{10} aryl or where NR'R" represents a cyclic moiety such as morpholine, piperidine, piperazine and the like; cyano; and tetrazolyl;

[00025] R_{15} , R_{16} , and R_{17} are each independently selected from the group consisting of

H; optionally substituted C_1 - C_{20} linear or branched alkyl including chloroalkyl and fluoroalkyl; optionally substituted C_2 - C_{20} linear or branched alkenyl; optionally substituted C_6 - C_{10} aryl or heteroaryl; COOR where R is H, optionally substituted C_1 - C_{20} alkyl, optionally substituted C_2 - C_{20} alkenyl or optionally substituted C_6 - C_{10} aryl, sodium, potassium or other pharmaceutically acceptable counter-ion such as calcium, magnesium, ammonium, tromethamine and the like; CONR'R", where R' and R" are independently H, optionally substituted C_1 - C_{20} alkyl, optionally substituted C_2 - C_{20} alkenyl or optionally substituted C_6 - C_{10} aryl or where NR'R" represents a cyclic moiety such as morpholine, piperidine, piperazine and the like; NH_2 ; C_1 - C_{20} alkylamino, bis(alkylamino), cycloalkylamino or cyclic amino; OH; C_1 - C_{20} alkoxy; C_1 - C_{20} alkanoyl; C_1 - C_{20} acyloxy; halo; C_1 - C_{20} alkylcarboxylamino; cyano; nitro; $SO_2NR'''R'''$ where R''' and R''' are independently H, C_1 - C_{20} alkyl or aryl; SO_2R''' where R''' is H, C_1 - C_{20} alkyl or aryl; SO_3R''' where R''' is H, C_1 - C_{20} alkyl or aryl; and tetrazolyl;

[00026] X is independently selected from the group consisting of O; N; S; S=O; SO₂; or NR^{''''}, where R^{''''} may be H or optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl, optionally substituted C₁-C₂₀ acyl, optionally substituted C₁-C₂₀ acyloxy and optionally substituted C₁-C₂₀ alkoxy carbonyl;

[00027] Y is independently O, S or NH;

[00028] Z is OR_a where R_a is selected from the group consisting of H; optionally substituted C₁-C₂₀ linear or branched alkyl including chloroalkyl or fluoroalkyl and the like; optionally substituted C₂-C₂₀ linear or branched alkenyl; optionally substituted C₆-C₁₀ aryl or heteroaryl; optionally substituted C₆-C₂₀ aroyl or heteroaroyl; optionally substituted C₁-C₂₀ alkanoyl; and SO₂R^{'''} where R^{'''} is H, C₁-C₂₀ alkyl or aryl;
or

[00029] Z is NR_bR_c where R_b and R_c are independently selected from the group consisting of H; optionally substituted C₁-C₂₀ linear or branched alkyl including chloroalkyl or fluoroalkyl and the like; optionally substituted C₂-C₂₀ linear or branched alkenyl; optionally substituted C₆-C₁₀ aryl or heteroaryl; optionally substituted C₃-C₁₀ cycloalkyl or cycloalkenyl; COOZ₁ where Z₁ is optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl or optionally substituted C₆-C₁₀ aryl; optionally substituted C₆-C₂₀ aroyl or heteroaroyl; optionally substituted C₁-C₂₀ alkanoyl; and SO₂R^{'''} where R^{'''} is H, C₁-C₂₀ alkyl or aryl; and wherein R_b and R_c together may be joined to form a 3-6 membered ring such as aziridine, morpholine, piperidine, piperazine and the like;
or

[00030] Z is CR_dR_eR_f where R_d, R_e and R_f are each independently selected from the group consisting of

H; optionally substituted C₁-C₂₀ linear or branched alkyl including chloroalkyl or fluoroalkyl and the like; optionally substituted C₂-C₂₀ linear or branched

alkenyl; optionally substituted C₆-C₁₀ aryl or heteroaryl; optionally substituted C₃-C₁₀ cycloalkyl or cycloalkenyl; COOR where R is H, optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl or optionally substituted C₆-C₁₀ aryl, sodium, potassium or other pharmaceutically acceptable counter-ion such as calcium, magnesium, ammonium, tromethamine and the like; NH₂; C₁-C₂₀ alkylamino, bis(alkylamino); cycloalkylamino or cyclic amino; OH; optionally substituted C₁-C₂₀ alkoxy including trifluoromethoxy and the like; optionally substituted C₁-C₂₀ alkanoyl; optionally substituted C₁-C₂₀ acyloxy; optionally substituted C₆-C₂₀ aroyl or heteroaroyl; halo; cyano; nitro; optionally substituted C₁-C₂₀ alkylcarboxylamino; SO₂NR'''R''' where R''' and R''' are independently H, C₁-C₂₀ alkyl or aryl; SO₂R''' where R''' is H, C₁-C₂₀ alkyl or aryl; and SO₃R''' where R''' is H, C₁-C₂₀ alkyl or aryl; and wherein R_d and R_e together may be joined to form a 3-6 membered ring such as aziridine, morpholine, piperidine, piperazine and the like; and the resulting stereocenter may have the R- or S- configuration; or the grouping C(=Y)Z may represent hydrogen or R₁₂ or may be absent.

[00031] Q is OR_a where R_a is selected from the group consisting of H; optionally substituted C₁-C₂₀ linear or branched alkyl including chloroalkyl or fluoroalkyl and the like; optionally substituted C₂-C₂₀ linear or branched alkenyl; optionally substituted C₆-C₁₀ aryl or heteroaryl; optionally substituted C₆-C₂₀ aroyl or heteroaroyl; optionally substituted C₁-C₂₀ alkanoyl; and SO₂R''' where R''' is H, C₁-C₂₀ alkyl or aryl; or

[00032] Q is NR_bR_c where R_b and R_c are independently selected from the group consisting of H; optionally substituted C₁-C₂₀ linear or branched alkyl including chloroalkyl or fluoroalkyl and the like; optionally substituted C₂-C₂₀ linear or branched alkenyl; optionally substituted C₆-C₁₀ aryl or heteroaryl; optionally substituted C₃-C₁₀ cycloalkyl or cycloalkenyl; COOZ₁ where Z₁ is optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl or optionally substituted C₆-C₁₀ aryl; optionally substituted C₆-C₂₀ aroyl or heteroaroyl; optionally substituted C₁-C₂₀ alkanoyl; and SO₂R''' where R''' is H, C₁-C₂₀ alkyl or aryl;

and wherein R_b and R_c together may be joined to form a 3-6 membered ring such as aziridine, morpholine, piperidine, piperazine and the like;
or

[00033] Q is SR_g , SOR_g or SO_2R_g where R_g is selected from the group consisting of

H; optionally substituted C_1 - C_{20} linear or branched alkyl including chloroalkyl or fluoroalkyl and the like; optionally substituted C_2 - C_{20} linear or branched alkenyl; optionally substituted C_1 - C_{20} acyl; optionally substituted C_1 - C_{20} alkoxycarbonyl; C_2 - C_{20} alkoxy; optionally substituted C_6 - C_{10} aryl or heteroaryl; and optionally substituted C_6 - C_{10} aroyl or heteroaroyl.

[00034] Group A is optionally substituted C_2 - C_{20} linear or branched alkenyl; optionally substituted C_6 - C_{20} aryl, linear or branched alkylaryl, linear or branched alkenylaryl; optionally substituted heteroaryls like pyridine, indole, morpholine, piperidine, piperazine, tetrazolyl and the like; COR where R is optionally substituted C_1 - C_{20} linear or branched alkyl; optionally substituted C_2 - C_{20} linear or branched alkenyl; optionally substituted C_6 - C_{20} aryl, linear or branched alkylaryl, linear or branched alkenylaryl; optionally substituted heteroaryls like pyridine, indole, morpholine, piperidine, piperazine, tetrazolyl and the like;

[00035] Group B is OH, C_1 - C_{20} alkoxy; SO_2R where R may be H or linear or branched C_1 - C_{20} alkyl.

[00036] Group Het represents a heterocyclic ring which is pyridyl, indolyl, tetrazolyl, imidazolyl, morphonyl, piperidinyl, piperazinyl, thiophenyl or the like.

[00037] These compounds are useful for treating diabetes and other diseases linked to insulin resistance, such as coronary artery disease and peripheral vascular disease, and also for treating or inhibiting inflammation or inflammatory diseases such as inflammatory arthritides and collagen vascular

diseases, which are caused by, for example, cytokines or inducible enzymes such as TNF-alpha, IL-1, IL-6, iNOS and/or COX-2. The compounds are also useful for treating or preventing other diseases mediated by cytokines, iNOS and/or COX-2, such as cancer.

[00038] Another aspect of the invention is a method of treating diabetes and related diseases comprising the step of administering to a subject suffering from a diabetic or related condition a therapeutically effective amount of a compound of Formulas I - XIII. Additionally, the invention provides a method of treating inflammation or inflammatory diseases or diseases mediated by cytokines, iNOS, PDE4, PDE3, p44/42 MAP kinase and/or COX-2 by administering to a subject in need of such treatment an effective amount of a compound according to Formulas I - XIII. Further, pharmaceutical compositions containing a therapeutically effective amount of one or more compounds according to Formulas I - XIII together with a pharmaceutically or physiologically acceptable carrier, for use in the treatments contemplated herein, are also provided.

BRIEF DESCRIPTION OF THE DRAWINGS

[00039] FIG. 1 shows a graph of the dose-dependent increase in glucose uptake in 3T3-L1 adipocytes treated with varying concentrations of a compound according to the invention.

[00040] FIG. 2 shows a graph of the enhancement of glucose uptake in 3T3-L1 adipocytes treated with a compound according to the invention in addition to varying concentrations of insulin.

[00041] FIG. 3 shows a graph of the lowering of blood glucose levels in ob/ob mice treated with a compound according to the invention.

[00042] FIGS. 4A and 4B show graphs of the lowering of serum triglycerides and free fatty acid levels, respectively, in ob/ob mice treated with a compound according to the invention.

[00043] FIG. 5 shows a graph of the inhibition of LPS-induced TNF-alpha production in mouse RAW264.7 cells treated with varying concentrations of a compound according to the invention.

[00044] FIG. 6 shows a graph of the inhibition of LPS-induced IL-1 beta production in mouse RAW264.7 cells treated with varying concentrations of a compound according to the invention.

[00045] FIG. 7 shows a graph of the inhibition of LPS-induced IL-6 production in mouse RAW264.7 cells treated with varying concentrations of a compound according to the invention.

[00046] FIG. 8 shows photos of Western blots demonstrating the inhibition of LPS-induced iNOS and COX-2 production in mouse RAW264.7 cells treated with varying concentrations of a compound according to the invention.

[00047] FIG. 9 shows a graph of median clinical scores over time demonstrating improvement of collagen induced arthritis in mice using varying concentrations of a compound according to the invention.

DETAILED DESCRIPTION OF THE INVENTION

[00048] The invention is based on the discovery that the compounds described herein are useful in the treatment of diabetes and other diseases linked to insulin resistance, such as coronary artery disease and peripheral vascular disease, and also for the treatment or inhibition of inflammation or inflammatory diseases such as inflammatory arthritides and collagen vascular diseases, which are caused by, for example, cytokines or inducible enzymes such as TNF-alpha, IL-1, IL-6, PDE4, PDE3, p44/42 MAP kinase, iNOS and/or COX-2.

Definitions

[00049] As utilized herein, the following terms, unless otherwise indicated, shall be understood to have the following meanings:

[00050] "Alkyl", alone or in combination, means a straight-chain or branched-chain alkyl radical containing preferably 1-20 carbon atoms, more preferably 1-10 carbon atoms, and most preferably 1-6 carbon atoms. Exemplary alkyl radicals include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, pentyl, isopentyl, neopentyl, iso-amyl, hexyl and the like.

[00051] "Alkenyl", alone or in combination, means a straight-chain or branched-chain hydrocarbon radical having one or more double bonds, preferably 1-2 double bonds and more preferably one double bond, and containing preferably 2-20 carbon atoms, more preferably 2-10 carbon atoms, and still more preferably 2-6 carbon atoms. Exemplary alkenyl radicals include ethenyl, propenyl, 2-methylpropenyl, n-butenyl, isobutenyl, and include groups containing multiple sites of unsaturation such as 1,3-butadiene and 1,4-butadienyl and the like.

[00052] "Alkoxy", alone or in combination, means a radical of the type "R-O-" wherein R can be hydrogen, linear or branched alkyl, or linear or branched alkenyl as previously defined and "O" is an oxygen atom. Exemplary alkoxy radicals include methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, iso-butoxy, sec-butoxy, tert-butoxy and the like.

[00053] "Alkoxycarbonyl", alone or in combination, means a radical of the type "R-O-C(O)-" wherein "R-O-" is an alkoxy radical as previously defined and "C(O)-" is a carbonyl radical. Exemplary alkoxycarbonyl groups include methoxycarbonyl and ethoxycarbonyl.

[00054] "Alkylcarboxylamino" means a group RCON(R)- where R can be independently hydrogen, linear or branched alkyl, or linear or branched alkenyl as previously defined.

[00055] "Alkanoyl", alone or in combination, means a radical of the type "R-C(O)-" wherein "R" is an alkyl radical as previously defined and "-C(O)-" is a carbonyl radical. Exemplary alkanoyl radicals include acetyl, trifluoroacetyl, hydroxyacetyl, propionyl, butyryl, valeryl, 4-methylvaleryl and the like.

[00056] "Halo" or "halogen", alone or in combination, means chloro, bromo, fluoro or iodo radicals.

[00057] "Aryl", alone or in combination, means an aromatic carbocyclic radical containing about 6 to about 10 carbon atoms, which is optionally substituted with one or more substituents selected from alkyl, alkoxy, halogen, hydroxy, amino, azido, nitro, cyano, haloalkyl, carboxy, alkoxycarbonyl, cycloalkyl, alkanoylamino, amido, amidino, alkoxycarbonylamino, N-alkylamidino, alkylamino, dialkylamino, aminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, N-alkylamido, N,N-dialkylamido, aralkoxycarbonylamino, alkylthio, alkylsulfinyl, alkylsulfonyl, oxo and the like. Exemplary aryl radicals include phenyl, o-tolyl, 4-methoxyphenyl, 2-(tert-butoxy)phenyl, 3-methyl-4-methoxyphenyl, 2-fluorophenyl, 2-chlorophenyl, 3-nitrophenyl, 3-aminophenyl, 3-acetamidophenyl, 2-amino-3-(aminomethyl)phenyl, 6-methyl-2-aminophenyl, 2-amino-3-methylphenyl, 4,6-dimethyl-2-aminophenyl, 4-hydroxyphenyl, 3-methyl-4-hydroxyphenyl, 4-(2-methoxyphenyl)phenyl, 2-amino-1-naphthyl, 2-naphthyl, 1-methyl-3-amino-2-naphthyl, 2,3-diamino-1-naphthyl, 4,8-dimethoxy-2-naphthyl and the like.


[00058] "Acyloxy" or "Acylamino" group means an oxygen or amino group, respectively, bonded to an acyl group (RCO) where R can be hydrogen, linear or branched alkyl, or linear or branched alkenyl.


[00059] "Alkylamido" means the group RN(H)CO- where R can be hydrogen, linear or branched alkyl, or linear or branched alkenyl, as previously defined.

[00060] The reference to "optionally substituted" in the definition of the compounds throughout this disclosure is intended to include any substituent which does not negatively affect the activity of the compounds. Typical substitution includes, for example, lower (C₁-C₆) alkyl; halogen such as fluoro, chloro and bromo; nitro; amino; lower alkylamino; carboxylate, lower alkyl carboxylate, hydroxy, lower alkoxy, sulfonamide, cyano, or the like.

[00061] The invention is directed to compounds, for example, heterocyclic derivatives of acyl urea, thiourea, carbamate and thiocarbamate compounds, that

provide a variety of useful pharmacological effects. The compounds are useful, for example, in lowering blood glucose levels in hyperglycemic disorders, such as diabetes mellitus, and for treating related disorders, such as obesity and hyperlipidemia. Furthermore, these compounds are useful for treatment of disorders associated with insulin resistance, such as polycystic ovary syndrome, and for the treatment of inflammation, inflammatory and immunological diseases, particularly those mediated by pro-inflammatory cytokines (such as TNF-alpha, IL-1 beta and IL-6), type 4 phosphodiesterase (PDE4), type 3 phosphodiesterase (PDE3), p44/42 mitogen activated protein (MAP) kinase, cyclooxygenase-2 (COX-2) and/or inducible nitric oxide synthase (iNOS). In particular, the invention discloses compounds of the Formulas I-XIII as well as the pharmaceutically acceptable salts and solvates thereof:





[00062] wherein the stereocenters marked with an asterisk (*) may be R- or S-; the bond represented by a dashed line plus a solid line may be a double bond or a single bond, and when the bond is a double bond it may be in the E or Z configuration, and when the bond is a single bond the resulting stereocenters may have the R- or S- configuration; and

[00063] R₁, R₂, R₃, R₄, R₅, R₆ and R₇ are each independently selected from the group consisting of

H; optionally substituted C₁-C₂₀ linear or branched alkyl including chloroalkyl or fluoroalkyl; optionally substituted C₂-C₂₀ linear or branched alkenyl; optionally substituted C₆-C₂₀ aryl, linear or branched alkylaryl, linear or branched alkenylaryl; COOR where R is H, optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl or optionally substituted C₆-C₁₀ aryl, sodium, potassium or other pharmaceutically acceptable counter-ion such as calcium, magnesium, ammonium, tromethamine and the like; CONR'R", where R' and R" are independently H, optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl or optionally substituted C₆-C₁₀ aryl or where NR'R" represents a cyclic moiety such as morpholine, piperidine, piperazine and the like; optionally substituted C₁-C₆ amidoalkyl; NH₂; C₁-C₂₀ alkylamino, bis(alkylamino), cycloalkylamino or cyclic amino; OH; optionally substituted C₁-C₂₀ alkoxy including trifluoromethoxy and the like; optionally substituted C₁-C₂₀ alkanoyl; optionally substituted C₁-C₂₀ acyloxy; halo; optionally substituted C₁-C₂₀ alkylcarboxylamino; cyano; nitro; SO₂NR'''R''' where R''' and R''' are independently H, C₁-C₂₀ alkyl or aryl; SO₂R''' where R''' is H, C₁-C₂₀ alkyl or aryl; SO₃R''' where R''' is H, C₁-C₂₀ alkyl or aryl; and C₄-C₈ heterocycles such as tetrazolyl, imidazolyl, pyrrolyl, pyridyl, indolyl and the like; or when individual aromatic rings possess adjacent substituents, these substituents may be joined to form a ring such as a methylenedioxy or ethylenedioxy group, and the like, including lactones and lactams;

[00064] R₈ and R₉ are each independently selected from the group consisting of

H; optionally substituted C₁-C₂₀ linear or branched alkyl; optionally substituted C₂-C₂₀ linear or branched alkenyl; optionally substituted C₆-C₁₀ aryl or heteroaryl; COOR where R is H, optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl or optionally substituted C₆-C₁₀ aryl, sodium, potassium or other pharmaceutically acceptable counter-ion such as calcium, magnesium, ammonium, tromethamine and the like; CONR'R", where R' and

R" are independently H, alkoxy, optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl, optionally substituted C₃-C₁₀ cycloalkyl or cycloalkenyl or optionally substituted C₆-C₁₀ aryl or heteroaryl, preferably 2-, 3- or 4-pyridyl or where NR'R" represents a cyclic moiety such as morpholine, piperidine, hydroxypiperidine, imidazole, piperazine, methylpiperazine and the like; NH₂; C₁-C₂₀ alkylamino, bis(alkylamino), cycloalkylamino or cyclic amino; OH; C₁-C₂₀ alkoxy; C₁-C₂₀ alkanoyl; C₁-C₂₀ acyloxy; halo; C₁-C₂₀ alkylcarboxylamino; cyano; nitro; SO₂NR'''R'''' where R''' and R'''' are independently H, C₁-C₂₀ alkyl or aryl; SO₂R''' where R''' is H, C₁-C₂₀ alkyl or aryl; SO₃R''' where R''' is H, C₁-C₂₀ alkyl or aryl; and tetrazolyl; wherein R₈ and R₉ together may be joined to form a C₄-C₈ heterocyclic ring, including lactone or lactam;

[00065] R₁₀ and R₁₁ are each independently selected from the group consisting of

H; optionally substituted C₁-C₂₀ linear or branched alkyl; optionally substituted C₂-C₂₀ linear or branched alkenyl; optionally substituted C₆-C₁₀ aryl or heteroaryl; COOR where R is H, optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl or optionally substituted C₆-C₁₀ aryl, sodium, potassium or other pharmaceutically acceptable counter-ion such as calcium, magnesium, ammonium, tromethamine and the like; CONR'R", where R' and R" are independently H, optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl or optionally substituted C₆-C₁₀ aryl or where NR'R" represents a cyclic moiety such as morpholine, piperidine, piperazine and the like; NH₂; C₁-C₂₀ alkylamino, bis(alkylamino), cycloalkylamino or cyclic amino; OH; C₁-C₂₀ alkoxy; C₁-C₂₀ alkanoyl; C₁-C₂₀ acyloxy; halo; C₁-C₂₀ alkylcarboxylamino; cyano; nitro; SO₂NR'''R'''' where R''' and R'''' are independently H, C₁-C₂₀ alkyl or aryl; SO₂R''' where R''' is H, C₁-C₂₀ alkyl or aryl; SO₃R''' where R''' is H, C₁-C₂₀ alkyl or aryl; and tetrazolyl; wherein R₁₀ and R₁₁ together may be joined to form a C₄-C₈ heterocyclic ring, including lactone or lactam;

[00066] R₁₂, R₁₃, R₁₈, R₁₉ and R₂₀ are each independently selected from the group consisting of

H; optionally substituted C₁-C₂₀ linear or branched alkyl; optionally substituted C₂-C₂₀ linear or branched alkenyl; optionally substituted C₆-C₁₀ aryl or heteroaryl; COOR where R is optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl or optionally substituted C₆-C₁₀ aryl, sodium, potassium or other pharmaceutically acceptable counter-ion such as calcium, magnesium, ammonium, tromethamine and the like; CONR'R'', where R' and R'' are independently H, optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl or optionally substituted C₆-C₁₀ aryl or where NR'R'' represents a cyclic moiety such as morpholine, piperidine, piperazine and the like; C₁-C₂₀ alkanoyl; C₁-C₂₀ alkylamido; C₆-C₂₀ aroyl or heteroaroyl; SO₂R''' where R''' is H, C₁-C₂₀ alkyl or aryl; morpholinocarbonylmethyl; piperazinocarbonylmethyl; and piperidinocarbonylmethyl;

[00067] R₁₂ and R₁₃ may be absent, or R₁₂ and R₁₃ together may be an optionally substituted heterocyclic ring, preferably morpholine, piperidine, piperazine, and N-methyl piperidine.

[00068] R₁₄ is selected from the group consisting of
H; optionally substituted C₁-C₂₀ linear or branched alkyl including chloroalkyl and fluoroalkyl; optionally substituted C₂-C₂₀ linear or branched alkenyl; optionally substituted C₆-C₁₀ aryl or heteroaryl; COOR where R is H, optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl or optionally substituted C₆-C₁₀ aryl, sodium, potassium or other pharmaceutically acceptable counter-ion such as calcium, magnesium, ammonium, tromethamine and the like; CONR'R'', where R' and R'' are independently H, optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl or optionally substituted C₆-C₁₀ aryl or where NR'R'' represents a cyclic moiety such as morpholine, piperidine, piperazine and the like; cyano; and tetrazolyl;

[00069] R₁₅, R₁₆, and R₁₇ are each independently selected from the group consisting of

H; optionally substituted C₁-C₂₀ linear or branched alkyl including chloroalkyl and fluoroalkyl; optionally substituted C₂-C₂₀ linear or branched alkenyl;

optionally substituted C₆-C₁₀ aryl or heteroaryl; COOR where R is H, optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl or optionally substituted C₆-C₁₀ aryl, sodium, potassium or other pharmaceutically acceptable counter-ion such as calcium, magnesium, ammonium, tromethamine and the like; CONR'R", where R' and R" are independently H, optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl or optionally substituted C₆-C₁₀ aryl or where NR'R" represents a cyclic moiety such as morpholine, piperidine, piperazine and the like; NH₂; C₁-C₂₀ alkylamino, bis(alkylamino), cycloalkylamino or cyclic amino; OH; C₁-C₂₀ alkoxy; C₁-C₂₀ alkanoyl; C₁-C₂₀ acyloxy; halo; C₁-C₂₀ alkylcarboxylamino; cyano; nitro; SO₂NR'''R''' where R''' and R''' are independently H, C₁-C₂₀ alkyl or aryl; SO₂R''' where R''' is H, C₁-C₂₀ alkyl or aryl; SO₃R''' where R''' is H, C₁-C₂₀ alkyl or aryl; and tetrazolyl;

[00070] X is independently selected from the group consisting of O; N; S; S=O; SO₂; or NR''''', where R''''' may be H or optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl, optionally substituted C₁-C₂₀ acyl, optionally substituted C₁-C₂₀ acyloxy and optionally substituted C₁-C₂₀ alkoxycarbonyl;

[00071] Y is independently O, S or NH;

[00072] Z is OR_a where R_a is selected from the group consisting of H; optionally substituted C₁-C₂₀ linear or branched alkyl including chloroalkyl or fluoroalkyl and the like; optionally substituted C₂-C₂₀ linear or branched alkenyl; optionally substituted C₆-C₁₀ aryl or heteroaryl; optionally substituted C₆-C₂₀ aroyl or heteroaroyl; optionally substituted C₁-C₂₀ alkanoyl; and SO₂R''' where R''' is H, C₁-C₂₀ alkyl or aryl;
or

[00073] Z is NR_bR_c where R_b and R_c are independently selected from the group consisting of

H; optionally substituted C₁-C₂₀ linear or branched alkyl including chloroalkyl or fluoroalkyl and the like; optionally substituted C₂-C₂₀ linear or branched

alkenyl; optionally substituted C₆-C₁₀ aryl or heteroaryl; COOZ₁ where Z₁ is optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl or optionally substituted C₆-C₁₀ aryl; optionally substituted C₆-C₂₀ aroyl or heteroaroyl; optionally substituted C₁-C₂₀ alkanoyl; and SO₂R''' where R''' is H, C₁-C₂₀ alkyl or aryl; and wherein R_b and R_c together may be joined to form a 3-6 membered ring such as aziridine, morpholine, piperidine, piperazine and the like;
or

[00074] Z is CR_dR_eR_f where R_d, R_e and R_f are each independently selected from the group consisting of

H; optionally substituted C₁-C₂₀ linear or branched alkyl including chloroalkyl or fluoroalkyl and the like; optionally substituted C₂-C₂₀ linear or branched alkenyl; optionally substituted C₆-C₁₀ aryl or heteroaryl; COOR where R is H, optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl or optionally substituted C₆-C₁₀ aryl, sodium, potassium or other pharmaceutically acceptable counter-ion such as calcium, magnesium, ammonium, tromethamine and the like; NH₂; C₁-C₂₀ alkylamino, bis(alkylamino); cycloalkylamino or cyclic amino; OH; optionally substituted C₁-C₂₀ alkoxy including trifluoromethoxy and the like; optionally substituted C₁-C₂₀ alkanoyl; optionally substituted C₁-C₂₀ acyloxy; optionally substituted C₆-C₂₀ aroyl or heteroaroyl; halo; cyano; nitro; optionally substituted C₁-C₂₀ alkylcarboxylamino; SO₂NR'''R''' where R''' and R''' are independently H, C₁-C₂₀ alkyl or aryl; SO₂R''' where R''' is H, C₁-C₂₀ alkyl or aryl; and SO₃R''' where R''' is H, C₁-C₂₀ alkyl or aryl; and wherein R_d and R_e together may be joined to form a 3-6 membered ring such as aziridine, morpholine, piperidine, piperazine and the like; and the resulting stereocenter may have the R- or S-configuration; or
the grouping -C(=Y)Z may represent hydrogen or R₁₂ or may be absent.

[00075] Q is OR_a where R_a is selected from the group consisting of

H; optionally substituted C₁-C₂₀ linear or branched alkyl including chloroalkyl or fluoroalkyl and the like; optionally substituted C₂-C₂₀ linear or branched alkenyl; optionally substituted C₆-C₁₀ aryl or heteroaryl; optionally substituted

C₆-C₂₀ aroyl or heteroaroyl; optionally substituted C₁-C₂₀ alkanoyl; and SO₂R''' where R''' is H, C₁-C₂₀ alkyl or aryl;
or

[00076] Q is NR_bR_c where R_b and R_c are independently selected from the group consisting of

H; optionally substituted C₁-C₂₀ linear or branched alkyl including chloroalkyl or fluoroalkyl and the like; optionally substituted C₂-C₂₀ linear or branched alkenyl; optionally substituted C₆-C₁₀ aryl or heteroaryl; optionally substituted C₃-C₁₀ cycloalkyl or cycloalkenyl; COOZ₁ where Z₁ is optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl or optionally substituted C₆-C₁₀ aryl; optionally substituted C₆-C₂₀ aroyl or heteroaroyl; optionally substituted C₁-C₂₀ alkanoyl; and SO₂R''' where R''' is H, C₁-C₂₀ alkyl or aryl; and wherein R_b and R_c together may be joined to form a 3-6 membered ring such as aziridine, morpholine, piperidine, piperazine and the like;
or

[00077] Q is SR_g, SOR_g or SO₂R_g where R_g is selected from the group consisting of

H; optionally substituted C₁-C₂₀ linear or branched alkyl including chloroalkyl or fluoroalkyl and the like; optionally substituted C₂-C₂₀ linear or branched alkenyl; optionally substituted C₁-C₂₀ acyl; optionally substituted C₁-C₂₀ alkoxycarbonyl; C₂-C₂₀ alkoxy; optionally substituted C₆-C₁₀ aryl or heteroaryl; and optionally substituted C₆-C₁₀ aroyl or heteroaroyl.

[00078] Group A is optionally substituted C₂-C₂₀ linear or branched alkenyl; optionally substituted C₆-C₂₀ aryl, linear or branched alkylaryl, linear or branched alkenylaryl; optionally substituted heteroaryls like pyridine, indole, morpholine, piperidine, piperazine, tetrazolyl and the like; COR where R is optionally substituted C₁-C₂₀ linear or branched alkyl; optionally substituted C₂-C₂₀ linear or branched alkenyl; optionally substituted C₆-C₂₀ aryl, linear or branched alkylaryl, linear or branched alkenylaryl; optionally substituted heteroaryls like pyridine, indole, morpholine, piperidine, piperazine, tetrazolyl and the like;

[00079] Group B is OH, C₁-C₂₀ alkoxy; SO₂R where R may be H or linear or branched C₁-C₂₀ alkyl.

[00080] Group Het represents a heterocyclic ring which is pyridyl, indolyl, tetrazolyl, imidazolyl, morphonyl, piperidinyl, piperazinyl, thiophenyl or the like.

[00081] These compounds are useful for treating diabetes and other diseases linked to insulin resistance, such as coronary artery disease and peripheral vascular disease, and also for treating or inhibiting inflammation or inflammatory diseases such as inflammatory arthritides and collagen vascular diseases, which are caused by, for example, cytokines or inducible enzymes such as TNF-alpha, IL-1, IL-6, PDE4, PDE3, p44/42 MAP kinase, iNOS and/or COX-2. The compounds are also useful for treating or preventing other diseases mediated by cytokines, PDE4, PDE3, p44/42 MAP kinase, iNOS and/or COX-2, such as cancer.

[00082] As indicated above, the compounds of the invention include bonds, designated in Formulas I - XIII with a dashed line plus a solid line, that may be either a double bond or a single bond. When such a bond is a double bond, it may have either the E or Z configuration. On the other hand, when such a bond is a single bond, the resulting stereocenters may be in the R- and/or S- configurations. Likewise, compounds of the invention with other stereocenters, designated in Formulas I - XIII with an asterisk, may be R- and/or S- stereoisomers. The invention contemplates racemic mixtures of such stereoisomers as well as the individual, separated stereoisomers. The individual stereoisomers may be obtained by the use of an optically active resolving agent. Alternatively, a desired enantiomer may be obtained by stereospecific synthesis using an optically pure starting material of known configuration.

[00083] Generally, R- or S- refers to the configuration of the stereoisomers. The determination of whether the configuration is R- (rectus)

or S- (sinister) is based on the priority of the atoms in a compound. Similarly, E- or Z- configuration is used when describing compounds with double bonds and wherein the determination is based on the priority of the atom on each carbon of a double bond.

[00084] The following compounds are representative of the preferred compounds according to Formula I:

[00085] 3-(3,5-Dimethoxyphenyl)-2-{4-[4-(3-oxo-3-ureidopropyl)-phenoxy]-phenyl}-acrylic acid methyl ester **(1)**;

[00086] 3-(3,5-Dimethoxyphenyl)-2-{4-[4-(3-oxo-3-ureidopropyl)-phenoxy]-phenyl}-acrylic acid **(6)**;

[00087] 3-(3,5-Dimethoxyphenyl)-2-{4-[4-(3-ethoxycarbonylamino-3-oxo-propyl)-phenoxy]-phenyl}-acrylic acid methyl ester **(8)**;

[00088] 2-{4-[4-(3-Benzoyloxycarbonylamino-3-oxo-propyl)-phenoxy]-phenyl}-3-(3,5-dimethoxyphenyl)-acrylic acid methyl ester **(9)**;

[00089] 3-(3,5-Dimethoxyphenyl)-2-{4-[4-(3-oxo-3-ureidopropyl)-phenoxy]-phenyl}-propionic acid **(10)**;

[00090] 3-(3,5-Dimethoxyphenyl)-2-{4-[4-(3-oxo-3-ureidopropenyl)-phenoxy]-phenyl}-acrylic acid **(11)**;

[00091] 3-(3,5-Dimethoxyphenyl)-2-{4-[4-(3-oxo-3-ureidopropyl)-phenoxy]-phenyl}-acrylic acid ethyl ester **(12)**;

[00092] 3-(3,5-Dimethoxyphenyl)-*N,N*-dimethyl-2-{4-[4-(3-oxo-3-ureidopropyl)-phenoxy]-phenyl}-acrylamide **(13)**;

[00093] 2-(4-{4-[3-(3-Cyclohexylureido)-3-oxopropyl]-phenoxy}-phenyl)-3-(3,5-dimethoxyphenyl)-acrylic acid **(14)**.

[00094] The following are preferred compounds according to Formula II:
[3-(4-Phenoxyphenyl)-propionyl]-urea **(15)**;

[00095] {3-[4-(4-Methoxyphenoxy)-phenyl]-acryloyl}-urea **(16)**.

[00096] The following are preferred compounds according to Formula III:
2-{4-[4-(3-Acetylureidomethyl)-phenoxy]-phenyl}-3-(3,5-dimethoxyphenyl)-acrylic acid methyl ester **(17)**;

[00097] 2-{4-[4-(3-Acetylthioureidomethyl)-phenoxy]-phenyl}-3-(3,5-dimethoxyphenyl)-acrylic acid **(18)**.

- [00098]** The following are preferred compounds according to Formula IV:
1-Acetyl-3-[4-(4-methoxyphenoxy)-benzyl]-urea (**24**);
- [00099]** Acetyl-3-[4-(3,4-dimethoxyphenoxy)-benzyl]-urea (**25**).
- [000100]** The following are more preferred compounds for their anti-inflammatory properties:
- [000101]** 3-(3,5-Dimethoxyphenyl)-*N,N*-dimethyl-2-{4-[4-(3-oxo-3-ureidopropyl)-phenoxy]-phenyl}-acrylamide (**13**);
- [000102]** 2-{4-[4-(2-Carbamoylethyl)-phenoxy]-phenyl}-3-(3,5-dimethoxyphenyl)-*N,N*-dimethylacrylamide (**31**);
- [000103]** 3-(4-{4-[2-(3,5-Dimethoxyphenyl)-1-dimethylcarbamoylvinyl]-phenoxy}-phenyl)-propionic acid ethyl ester (**37**);
- [000104]** *N*-{4-[2-(3,5-Dimethoxyphenyl)-1-dimethylcarbamoylvinyl]-phenyl}-3-hydroxybenzamide (**44**);
- [000105]** 3-(3,5-Dimethoxyphenyl)-2-(4-hydroxyphenyl)-*N,N*-dimethylacrylamide (**49**);
- [000106]** [3-(4-{4-[2-(3,5-Dimethoxyphenyl)-1-(piperidine-1-carbonyl)-vinyl]-phenoxy}-phenyl)-propionyl]-urea (**51**);
- [000107]** 2-{4-[4-(3-Acetylamino-3-oxopropyl)-phenoxy]-phenyl}-3-(4-fluorophenyl)-*N,N*-dimethylacrylamide (**56**);
- [000108]** 2-(4-{4-[2-(3,5-Dimethoxyphenyl)-1-dimethylcarbamoylvinyl]-phenoxy}-benzyl)-malonic acid (**58**);
- [000109]** 2-(4-{4-[2-(3,5-Dimethoxyphenyl)-1-dimethylcarbamoylvinyl]-phenoxy}-benzyl)-malonamide (**59**);
- [000110]** 3-(3,5-Dimethoxyphenyl)-*N,N*-dimethyl-2-[4-(pyridin-2-yloxy)-phenyl]-acrylamide (**66**);
- [000111]** *N*-{4-[2-(3,5-Dimethoxyphenyl)-1-dimethylcarbamoyl-vinyl]-phenyl}-benzamide (**67**) ;
- [000112]** 2-{4-[4-(1-Dimethylcarbamoyl-2-pyridin-3-yl-vinyl)-phenoxy]-benzyl}-malonamide (**71**);
- [000113]** 3-{4-[4-(2-Benzo[1,3]dioxol-5-yl-1-dimethylcarbamoyl-vinyl)-phenoxy]-phenyl}-propionic acid ethyl ester (**69**);
- [000114]** 3-Benzo[1,3]dioxol-5-yl-2-{4-[4-(2-carbamoylethyl)-phenoxy]-phenyl}-*N,N*-dimethyl-acrylamide (**72**);

- [000115]** N,N-Dimethyl-2-{4-[4-(3-oxo-3-ureidopropyl)-phenoxy]-phenyl}-3-pyridin-3-yl-acrylamide (**73**);
- [000116]** The following are more preferred compounds for their antidiabetic properties:
- [000117]** 3-(3,5-Dimethoxyphenyl)-2-{4-[4-(3-ethoxycarbonylamino-3-oxopropyl)-phenoxy]-phenyl}-acrylic acid methyl ester (**8**);
- [000118]** (4-{4-[2-(3,5-Dimethoxyphenyl)-1-dimethylcarbamoyl-vinyl]-phenoxy}-benzyl)-carbamic acid methyl ester (**29**);
- [000119]** 2-{4-[4-(2-Carbamoylethyl)-phenoxy]-phenyl}-3-(3,5-dimethoxyphenyl)-N,N-dimethylacrylamide (**31**);
- [000120]** 3-(3,5-Dimethoxyphenyl)-N,N-dimethyl-2-{4-[4-(3-morpholin-4-yl-3-oxopropyl)-phenoxy]-phenyl}-acrylamide (**40**);
- [000121]** [3-(4-{4-[2-(3,5-Dimethoxyphenyl)-1-(piperidine-1-carbonyl)-vinyl]-phenoxy}-phenyl)-propionyl]-urea (**51**);
- [000122]** 2-{4-[4-(3-Acetylamino-3-oxopropyl)-phenoxy]-phenyl}-3-(4-fluorophenyl)-N,N-dimethylacrylamide (**56**);
- [000123]** 3-(3,5-Dimethoxyphenyl)-2-{4-[4-(3-oxo-3-ureidopropyl)-phenoxy]-phenyl}-N-pyridin-4-ylacrylamide (**60**);
- [000124]** N-(4-Chlorophenyl)-3-(3,5-dimethoxyphenyl)-2-{4-[4-(3-oxo-3-ureidopropyl)-phenoxy]-phenyl}-acrylamide (**61**);
- [000125]** 3-(3,5-Dimethoxyphenyl)-N,N-dimethyl-2-(4-{4-[2-(2-morpholin-4-yl-2-oxoethylcarbamoyl)-ethyl]-phenoxy}-phenyl)-acrylamide (**63**);
- [000126]** 3-(3,5-Dimethoxyphenyl)-N,N-dimethyl-2-(4-{4-[3-(4-methylpiperazin-1-yl)-3-oxopropyl]-phenoxy}-phenyl)-acrylamide (**64**).

[000127] However, it will be appreciated that the invention also contemplates the provision and use of other compounds according to Formulas I – XIII.

[000128] The compounds according to the present invention may be combined with a physiologically acceptable carrier or vehicle to provide a pharmaceutical composition, such as, lyophilized powder in the form of tablet or capsule with various fillers and binders. Similarly, the compounds may be coadministered with other agents. Co-administration shall mean the administration of at least two agents to a

subject so as to provide the beneficial effects of the combination of both agents. For example, the agents may be administered simultaneously or sequentially over a period of time. The effective dosage of a compound in the composition can be widely varied as selected by those of ordinary skill in the art and may be empirically determined. Moreover, the compounds of the present invention can be used alone or in combination with one or more additional agents depending on the indication and the desired therapeutic effect. For example, in the case of diabetes, insulin resistance and associated conditions or complications, including obesity and hyperlipidemia, such additional agent(s) may be selected from the group consisting of: insulin or an insulin mimetic, a sulfonylurea (such as acetohexamide, chlorpropamide, glimepiride, glipizide, glyburide, tolbutamide and the like) or other insulin secretagogue (such as nateglinide, repaglinide and the like), a thiazolidinedione (such as pioglitazone, rosiglitazone and the like) or other peroxisome proliferator-activated receptor (PPAR)-gamma agonist, a fibrate (such as bezafibrate, clofibrate, fenofibrate, gemfibrozol and the like) or other PPAR-alpha agonist, a PPAR-delta agonist, a biguanide (such as metformin), a statin (such as fluvastatin, lovastatin, pravastatin, simvastatin and the like) or other hydroxymethylglutaryl (HMG) CoA reductase inhibitor, an alpha-glucosidase inhibitor (such as acarbose, miglitol, voglibose and the like), a bile acid-binding resin (such as cholestyramine, celestipol and the like), a high density lipoprotein (HDL)-lowering agent such as apolipoprotein A-I (apoA1), niacin and the like, probucol and nicotinic acid. In the case of inflammation, inflammatory diseases, autoimmune disease and other such cytokine mediated disorders, the additional agent(s) may be selected from the group consisting of: a nonsteroidal anti-inflammatory drug (NSAID) (such as diclofenac, diflunisal, ibuprofen, naproxen and the like), a cyclooxygenase-2 inhibitor (such as celecoxib, rofecoxib and the like), a corticosteroid (such as prednisone, methylprednisone and the like) or other immunosuppressive agent (such as methotrexate, leflunomide, cyclophosphamide, azathioprine and the like), a disease-modifying antirheumatic drug (DMARD) (such as injectable gold, penicillamine, hydroxychloroquine, sulfasalazine and the like), a TNF-alpha inhibitor (such as etanercept, infliximab and the like), other cytokine inhibitor (such as soluble cytokine receptor, anti-cytokine antibody and the like), other immune modulating agent (such as cyclosporin, tacrolimus, rapamycin and the like) and a narcotic agent (such as hydrocodone, morphine, codeine, tramadol and the like). The combination therapy

contemplated by the invention includes, for example, administration of the inventive compound and additional agent(s) in a single pharmaceutical formulation as well as administration of the inventive compound and additional agent(s) in separate pharmaceutical formulations.

[000129] Another aspect of the invention is a method of treating diabetes and related diseases comprising the step of administering to a subject suffering from a diabetic or related condition a therapeutically effective amount of a compound of Formulas I -XIII. Additionally, the invention provides a method of treating inflammation or inflammatory diseases or diseases mediated by cytokines, PDE4, PDE3, p44/42 MAP kinase, iNOS and/or COX-2 by administering to a subject in need of such treatment an effective amount of a compound according to Formulas I - XIII. Further, pharmaceutical compositions containing a therapeutically effective amount of one or more compounds according to Formulas I – XIII together with a pharmaceutically or physiologically acceptable carrier, for use in the treatments contemplated herein, are also provided.


[000130] The compounds of the invention are useful for the treatment of diabetes, characterized by the presence of elevated blood glucose levels, that is, hyperglycemic disorders such as diabetes mellitus, including both type 1 and 2 diabetes, as well as other hyperglycemic related disorders such as obesity, increased cholesterol, hyperlipidemia such as hypertriglyceridemia, kidney related disorders and the like. The compounds are also useful for the treatment of disorders linked to insulin resistance and/or hyperinsulinemia, which include, in addition to diabetes, hyperandrogenic conditions such as polycystic ovary syndrome (Ibanez et al., J. Clin Endocrinol Metab, 85:3526-30, 2000; Taylor A.E., Obstet Gynecol Clin North Am, 27:583-95, 2000), coronary artery disease such as atherosclerosis and vascular restenosis, and peripheral vascular disease. Additionally, the compounds of the present invention are also useful for the treatment of inflammation and immunological diseases that include those mediated by signaling pathways linked to pro-inflammatory cytokines, such as rheumatoid arthritis, ankylosing spondylitis, multiple sclerosis, inflammatory bowel disease, psoriasis, and contact and atopic dermatitis.

[000131] By "treatment", it is meant that the compounds of the invention are administered in an amount which is at least sufficient to, for example, reduce the blood glucose level in a patient suffering from a hyperglycemic disorder or to inhibit or prevent the development of pro-inflammatory cytokine or like responses in a patient suffering from inflammatory or immunological disease. In the case of diabetes, the compound is usually administered in the amount sufficient to reduce the blood glucose level, free fatty acid level, triglyceride level and/or the like level sufficient to improve or alleviate the symptoms and/or reduce the risk of complications associated with elevated levels of these parameters. A variety of subjects may be treated with the present compounds to reduce blood glucose levels such as livestock, wild or rare animals, pets, as well as humans. The compounds may be administered to a subject suffering from hyperglycemic disorder using any convenient administration technique, including intravenous, intradermal, intramuscular, subcutaneous, oral and the like. However, oral daily dosage is preferred. The dosage delivered to the host will necessarily depend upon the route by which the compound is delivered, but generally ranges from about 0.1 to about 500 mg/kg human body weight or typically from about 0.1 to about 50 mg/kg human body weight. Generally similar types of administration and dosages are also contemplated when the compounds of the invention are used to treat inflammatory or immunological disease.

[000132] The compounds of this invention may be used in formulations using acceptable pharmaceutical vehicles for enteral, or parenteral, administration, such as, for example, water, alcohol, gelatin, gum arabic, lactose, amylase, magnesium stearate, talc, vegetable oils, polyalkylene glycol, and the like. The compounds can be formulated in solid form, e.g., as tablets, capsules, drages and suppositories, or in the liquid form, e.g., solutions, suspensions and emulsions. The preparations may also be delivered transdermally or by topical application.


[000133] The syntheses of representative compounds according to the present invention are illustrated in Schemes I and II. Further examples illustrating the syntheses of additional compounds according to the present invention are also given below.

SCHEME I



[000134] Scheme 1 details the synthesis of compounds 1-6. Scheme 2 details the synthesis of 17. It is to be understood that the Schemes 1 and 2 are representative schemes and are not intended to be limited to the compounds disclosed.

SCHEME II



[000135] The following examples are provided to further illustrate the present invention and are not intended to limit the invention in any way.

EXAMPLE 1

Synthesis of 3-(3,5-dimethoxyphenyl)-2-{4-[4-(3-oxo-3-ureidopropyl)-phenoxy]-phenyl}-acrylic acid methyl ester (1) [see Scheme I]

[000136] Step 1: Synthesis of 3-(3,5-dimethoxyphenyl)-2-(4-hydroxyphenyl)-acrylic acid (2). To a mixture of 3,5-dimethoxybenzaldehyde (120 g, 0.72 mol) and *p*-hydroxyphenyl acetic acid (110 g, 0.72 mol) was added acetic anhydride (240 mL) and triethylamine (161 mL, 1.6 equiv.). This non-homogeneous mixture on heating becomes homogeneous at ~70°C. After being stirred at 130°C for 4 hr, the mixture was cooled to room temperature. HCl (15%, 500 mL) was added to the reaction mixture slowly in 30min keeping temperature below 5-10°C. The solid was dissolved

in 3N aqueous NaOH (1.2 L) and stirred for 0.5 hr. The filtrate was acidified, maintaining a temperature at 25-30°C, with conc. HCl (~700 mL) to pH 1. The precipitated product was filtered and washed with water to give crude product (~300 g, wet cake). The crude product was dissolved by heating in ethanol and recrystallized by adding equal volume of water. The product was dried overnight in a vacuum oven at 40°C. Yield: 161 g, 74%. Analysis: ¹HNMR (DMSO-d₆): δ 12.48 (br, 1H), 9.42 (s, 1H), 7.59 (s, 1H), 6.95 (d, J=8.0 Hz, 2H), 6.76 (d, J=8.0 Hz, 2H), 6.35 (t, J=2.2 Hz, 1H), 6.27 (d, J=2.2 Hz, 2H), 3.56 (s, 6H).

[000137] (b) Step 2: Synthesis of 3-(3,5-dimethoxyphenyl)-2-[4-(4-formylphenoxy)-phenyl]-acrylic acid (3). **2** (64.0 g, 0.21 mol) was dissolved in 320 mL anhydrous DMSO under nitrogen, and potassium *tert*-butoxide (48.0 g, 0.43 mol) was added in lots. When the solution became homogenous, *p*-fluorobenzaldehyde (27 mL, 0.22 mol) was added and the mixture was heated at 100°C for 5 hr. After cooling to room temperature, the solution was poured into 1 L water and extracted with ether (2 x 500 mL). The aqueous phase was acidified with 5% HCl to ~ pH 4 and the precipitated product was collected by suction filtration. The wet filter cake was dissolved in a minimum of boiling acetone and recrystallized with addition of water. After chilling to 4°C for 3 hr, the solid was collected by vacuum filtration. The product was dried overnight at 40°C in a vacuum oven. Yield: 62 g, 73%. Analysis: ¹HNMR (DMSO-d₆): δ 12.87 (s, 1H), 9.94 (s, 1H), 7.95 (d, J=8.2 Hz, 2H), 7.72 (s, 1H), 7.27 (d, J=8.0 Hz, 2H), 7.19 (d, J=8.0 Hz, 2H), 7.15 (d, J=8.2 Hz, 2H), 6.42 (t, J=1.6 Hz, 1H), 6.29 (d, J=2.0 Hz, 2H), 3.60 (s, 6H).

[000138] (c) Step 3: Synthesis of 3-(3,5-dimethoxyphenyl)-2-[4-[4-(2-ethoxycarbonyl-vinyl)-phenoxy]-phenyl]-acrylic acid (4). Triethylphosphonoacetate (7.14 mL, 36 mmol) was added to a suspension of NaH (60% in mineral oil, 2.64 g, 66 mmol) in anhydrous THF (100 mL) at 0°C under argon, and the mixture was stirred for 15 min. A solution of aldehyde **3**, (12.12 g, 30 mmol) in THF (100 mL) was added and the mixture was stirred for 1h. The mixture was quenched with saturated aqueous ammonium chloride solution (5 mL), diluted with ethyl acetate (300 mL) and acidified with 5% aqueous HCl to pH 1. The ethyl acetate layer was separated, and the aqueous layer was extracted with ethyl acetate (100 mL). The combined

organic layers were washed with brine, dried over anhydrous MgSO_4 , filtered and concentrated. The crude product was purified by recrystallization from a mixture of chloroform/methanol. The compound was suspended in hot methanol (200 mL) and a minimum volume (~30–40 mL) of chloroform was added to yield **4**. Yield: 12.39 g, 87.1%. Analysis: ^1H NMR ($\text{DMSO}-d_6$): δ 7.77 (d, $J=8.4$ Hz, 2H), 7.69 (s, 1H), 7.65 (d, $J=16$ Hz, 2H), 7.23 (d, 8.8 Hz, 2H), 7.11 (d, $J=8.8$ Hz, 2H), 7.01 (d, $J=8.4$ Hz, 2H), 6.57 (d, $J=16$ Hz, 2H), 6.41 (t, $J=2$ Hz, 1H), 6.28 (d, $J=1.6$ Hz, 2H), 4.18 (q, $J=7.2$ Hz, 2H), 3.59 (s, 6H), 1.26 (t, $J=7.2$ Hz, 3H).

[000139] (d) Step 4: Synthesis of 3-(3,5-dimethoxyphenyl)-2-{4-[4-(2-ethoxycarbonyl-ethyl)-phenoxy]-phenyl}-acrylic acid (**5**). To a suspension of Raney Ni (10.0 g, Raney 2800 nickel in water active catalyst) in ethanol-dioxane (2:1, 50 mL) was added a solution of **4** (13.0 g, 27.4 mmol) in a mixture of ethanol-dioxane (2:1, 400 mL), and the resulting mixture was stirred vigorously for 15 hr under hydrogen at atmospheric pressure. Completion of the reaction was monitored by HPLC (time varies with the speed of stirring). Catalyst was filtered through a bed of Celite® diatomaceous earth, the bed was washed with ethanol-dioxane (2:1, 200 mL), and solvent was evaporated. The solid obtained was dissolved in hot toluene (150 mL) and cooled at 4°C overnight. Solid separated was filtered and washed with ice-cold toluene (50 mL) and dried at 55°C for 6 hr. Yield: 11.61 g, 90.5%. Analysis: ^1H NMR ($\text{DMSO}-d_6$): δ 12.75 (s, 1H), 7.68 (s, 1H), 7.26 (d, $J=8.4$ Hz, 2H), 7.17 (d, $J=8.4$ Hz, 2H), 6.99 (d, $J=8.4$ Hz, 2H), 6.94 (d, $J=8.4$ Hz, 2H), 6.39 (t, $J=2.0$ Hz, 1H), 6.27 (d, $J=1.6$ Hz, 2H), 4.06 (q, $J=7.2$ Hz, 2H), 3.57 (s, 6H), 2.84 (t, $J=8$ Hz, 2H), 2.60 (t, $J=8$ Hz, 2H), 1.15 (t, $J=8$ Hz, 3H).

[000140] (e) Step 5: Synthesis of 3-(3,5-dimethoxyphenyl)-2-{4-[4-(3-oxo-3-ureido-propyl)-phenoxy]-phenyl}-acrylic acid (**6**). To a solution of sodium ethoxide in ethanol (21% w/w, 65 mL) under argon was added ethyl acetate (3.12 mL), then refluxed for 20 min. Urea (18 g, 0.3 mol) was dissolved in the above-mentioned sodium ethoxide in ethanol solution at 75°C . To this solution was added **5** (13 g, 0.027 mol) in one lot. After all dissolved, the resulting mixture was stirred at 75°C for another 5 min, cooled quickly in 15 min to $15\text{--}20^\circ\text{C}$, TFA (13 mL) added, and then adjusted to pH 4–5 with 5% HCl. After stirring at room temperature for 1 hr, the

mixture was slowly added to water (520 mL). The solid separated was filtered and refluxed in 10% isopropanol in ethyl acetate (150 mL) for 20 min. The mixture was allowed to cool to room temperature, then incubated overnight at 4°C. The mixture was filtered and solid was dried. Yield: 8.5 g. Analysis: ¹HNMR (DMSO-d₆): δ12.35 (br, 1H), 10.20 (s, 1H), 7.75 (br, 1H), 7.68 (s, 1H), 7.26 (d, J=8.4 Hz, 2H), 7.17 (d, J=8.4 Hz, 2H), 6.99 (d, J=8.4 Hz, 2H), 6.94 (d, J=8.4 Hz, 2H), 6.39 (t, J=2.4 Hz, 1H), 6.27 (d, J=2.4 Hz, 2H), 3.57 (s, 6H), 2.81 (t, J=7.2 Hz, 2H), 2.54 (t, J=7.2 Hz, 2H).


[000141] (f) Step 6: Synthesis of 3-(3,5-dimethoxyphenyl)-2-{4-[4-(3-oxo-3-ureido-propyl)-phenoxy]-phenyl}-acrylic acid methyl ester (1). To a stirred solution of **6** (5 g, 0.01 mol) in dry DMF (35 mL) under argon was added K₂CO₃ (1.38 g, 0.01 mol). To this, dimethyl sulfate (3.8 g, 0.03 mol) was added and stirred at room temperature for 30 min. The reaction mixture was acidified with 5% aqueous HCl and extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate and evaporated. The oily residue was dissolved in hexane/ethyl acetate (2:3, 30 mL) with stirring, and incubated overnight at 4°C for crystallization. The solid was collected by vacuum filtration and dried. Yield: 3.3 g, 65%. Analysis: ¹HNMR (DMSO-d₆): δ10.17 (br, 1H), 7.72 (br, 2H), 7.72 (s, 1H), 7.25 (d, J=8.4 Hz, 2H), 7.18 (d, J=6.8 Hz, 2H), 7.21 (s overlapped, 1H), 7.01 (d, J=6.8 Hz, 2H), 6.96 (d, J=8.4 Hz, 2H), 6.41 (t, J=2.2 Hz, 1H), 6.28 (d, J=2.2 Hz, 2H), 3.73 (s, 3H), 3.57 (s, 6H), 2.84 (t, J=7.2 Hz, 2H), 2.61 (t, J=7.2 Hz, 2H).

EXAMPLE 2

Synthesis of 3-(3,5-dimethoxyphenyl)-2-{4-[4-(3-ethoxycarbonylamino-3-oxo-propyl)-phenoxy]-phenyl}-acrylic acid methyl ester (**8**)

[000142] 2-{4-[4-(2-Carbamoyl-ethyl)-phenoxy]-phenyl}-3-(3,5-dimethoxyphenyl)-acrylic acid methyl ester (**7**) was obtained as a byproduct in the synthesis of 3-(3,5-dimethoxy-phenyl)-2-{4-[4-(2,4-dioxothiazolidin-5-ylmethyl)-phenoxy]-phenyl}-acrylic acid methyl ester, performed essentially as shown in PCT/US99/09982 (WO 99/58127). **7** (460 mg, 1.0 mmol) was taken up in dry THF (6 mL) and cooled to -78°C. To this solution, lithium diisopropyl amide (LDA) (2M, 0.55 mL, 1.1 mmol) was added and stirred for 10 min. Ethyl chloroformate (0.11 mL, 1.2 mmol) was added

and stirred overnight at room temperature. The reaction was quenched with saturated aqueous ammonium chloride solution and ethyl acetate (50 mL) was added. The organic layer was washed with brine (2 x 20 mL), dried on anhydrous magnesium sulfate and evaporated under reduced pressure. The crude product was purified by silica gel chromatography and eluted with hexane-ethyl acetate (7:3). Yield: 264 mg, 49.8%.




[000143] Analysis: ^1H NMR (DMSO- d_6): δ 10.52 (s, 1H), 7.70 (s, 1H), 7.24 (d, $J=8.4$ Hz, 2H), 7.17 (d, $J=8.4$ Hz, 2H), 6.99 (d, $J=8.4$ Hz, 2H), 6.94 (d, $J=8.4$ Hz, 2H), 6.40 (t, $J=2.1$ Hz, 1H), 6.27 (d, $J=2.1$ Hz, 2H), 4.07 (q, $J=7.2$ Hz, 2H), 3.70 (s, 3H), 3.56 (s, 6H), 2.76 (m, 4H), 1.19 (t, $J=7.2$ Hz, 3H).

EXAMPLE 3

Synthesis of 2-[4-[4-(3-benzoyloxycarbonylamino-3-oxo-propyl)-phenoxy]-phenyl]-3-(3,5-dimethoxyphenyl)-acrylic acid methyl ester (**9**)

[000144] **7** (1.38, 3.0 mmol) prepared as in Example 2 was taken up in dry THF (20 mL) and cooled to -78°C . To this solution, LDA (2M, 1.8 mL, 3.6 mmol) was added and stirred for 10 min. Benzyl chloroformate (0.67 g, 3.9 mmol) was added and stirred overnight at room temperature. The reaction was quenched with saturated aqueous ammonium chloride solution, and ethyl acetate (150 mL) was added. The organic layer was washed with brine (2 x 25 mL), dried on anhydrous magnesium sulfate and evaporated under reduced pressure. The crude product was purified by silica gel chromatography and eluted with hexane-ethyl acetate (7:3). Yield: 0.68g, 37.3%.




[000145] Analysis: ^1H NMR ($\text{DMSO}-d_6$): δ 10.65 (s, 1H), 7.72 (s, 1H), 7.38-7.39 (m, 5H), 7.25 (d, $J=8.4$ Hz, 2H), 7.18 (d, $J=8.4$ Hz, 2H), 7.00 (d, $J=8.4$ Hz, 2H), 6.94 (d, $J=8.4$ Hz, 2H), 6.41 (t, $J=2.0$ Hz, 1H), 6.28 (d, $J=2.0$ Hz, 2H), 5.12 (s, 2H), 3.72 (s, 3H), 3.57 (s, 6H), 2.79 (m, 4H).

EXAMPLE 4

Synthesis of 3-(3,5-dimethoxyphenyl)-2-{4-[4-(3-oxo-3-ureidopropyl)-phenoxy]-phenyl}-propionic acid (**10**)

[000146] 3-(3,5-Dimethoxyphenyl)-2-{4-[4-(2-ethoxycarbonylvinyl)-phenoxy]-phenyl}-acrylic acid (**4**, 2.37 g, 5.0 mmol) was dissolved in a mixture of ethanol-dioxane (2:1, 150 mL), and palladium charcoal (10%, 500 mg) was added. The mixture was stirred under hydrogen for 15 hr. Catalyst was then removed by filtration, and solvent was evaporated under reduced pressure to yield 3-(3,5-dimethoxyphenyl)-2-{4-[4-(2-ethoxycarbonylvinyl)-phenoxy]-phenyl}-propionic acid (**18**) quantitatively. Urea (0.21 g, 3.58 mmol) was dissolved in sodium ethoxide (2.7 M, 2.2 mL, 5.92 mmol) at 80°C under argon, and to this a solution of **18** (1.13 g, 2.37 mmol) in anhydrous ethanol (15 mL) was added and heated at this temperature for 13 hr. Ethanol was evaporated under reduced pressure, water (20 mL) was added, acidified to pH 1 by 5% aqueous HCl and extracted with ethyl acetate (50 mL). The organic layer was washed with water (2 x 25 mL), brine (2 x 20 mL), dried over anhydrous magnesium sulfate and evaporated. The crude product was purified by silica gel chromatography and eluted with hexane-ethyl acetate (3:7) containing acetic acid (1%), followed by recrystallization from ethanol. Yield: 256 mg, 22.8%.




[000147] Analysis: ^1H NMR ($\text{DMSO}-d_6$): δ 12.37 (s, 1H), 10.17 (s, 1H), 7.74 (br, 1H), 7.31 (d, $J=9.2$ Hz, 2H), 7.21 (d, $J=9.2$ Hz, 2H), 6.91 (d, $J=8.4$ Hz, 2H), 6.90 (d, $J=8.4$ Hz, 2H), 6.33 (d, $J=2.0$ Hz, 2H), 6.29 (t, $J=2.0$ Hz, 1H), 3.83 (t, $J=8.0$ Hz, 1H), 3.68 (s, 6H), 3.19 (dd, $J=14.4$ & 8.4 Hz, 1H), 2.88-2.80 (m, 3H), 2.59 (t, $J=8.0$ Hz, 2H).

EXAMPLE 5

Synthesis of 3-(3,5-dimethoxyphenyl)-2-{4-[4-(3-oxo-3-ureidopropenyl)-phenoxy]-phenyl}-acrylic acid (**11**)

[000148] Urea (0.21 g, 3.58 mmol) was dissolved in sodium ethoxide (2.7 M, 2.2 mL, 5.92 mmol) at 80°C under argon, and to this a solution of **4** (1.14 g, 2.37 mmol) in anhydrous ethanol (15 mL) was added and heated at this temperature for 13 hr. Ethanol was evaporated under reduced pressure, water (20 mL) was added, acidified to pH 1 by 5% aqueous HCl and extracted with ethyl acetate (50 mL). The organic layer was washed with water (2 x 25 mL), brine (2 x 20 mL), dried over anhydrous magnesium sulfate and evaporated. The crude product was purified by silica gel chromatography and eluted with hexane-ethyl acetate (3:7) containing acetic acid (1%), followed by recrystallization from ethanol. Yield: 167 mg, 14.4%.




[000149] Analysis: ^1H NMR (DMSO- d_6): δ 12.51 (br, 1H), 10.30 (s, 1H), 7.92 (br, 1H), 7.77 (d, $J=9.2$ Hz, 2H), 7.68 (s, 1H), 7.65 (d, $J=16.0$ Hz, 1H), 7.30 (br, 1H), 7.22 (d, $J=8.8$ Hz, 2H), 7.10 (d, $J=8.8$ Hz, 2H), 7.03 (d, $J=9.2$ Hz, 2H), 6.73 (d, $J=16.0$ Hz, 1H), 6.40 (t, $J=2.0$ Hz, 1H), 6.28 (d, $J=2$ Hz, 2H), 3.59 (s, 6H).

EXAMPLE 6

Synthesis of 3-(3,5-dimethoxyphenyl)-2-{4-[4-(3-oxo-3-ureidopropyl)-phenoxy]-phenyl}-acrylic acid ethyl ester (**12**)

[000150] To a stirred solution of **6** (0.40 g, 0.81 mmol) in dry DMSO (3 mL) was added K_2CO_3 (0.14 g, 0.98 mmol). To this, diethyl sulfate (0.115 g, 0.91 mmol) was added and stirred at room temperature for 30 min. The reaction mixture was poured into water (30 mL) and extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate and evaporated. The crude product was purified by column chromatography over silica gel and eluted with hexanes-ethyl acetate (3:1). Yield: 0.39 g, 92.2%.




[000151] Analysis: ^1H NMR (DMSO- d_6): δ 10.17 (s, 1H), 7.74 (br, 1H), 7.70 (s, 1H), 7.25 (d, $J=8.4$ Hz, 2H), 7.24 (overlapped, 1H), 7.18 (d, $J=8.4$ Hz, 2H), 7.00 (d,

J=8.4 Hz, 2H), 6.95 (d, J=8.4 Hz, 2H), 6.41 (t, J=1.6 Hz, 1H), 6.28 (d, J=1.6 Hz, 2H), 4.19 (q, J=8.0 Hz, 2H), 3.57 (s, 6H), 2.83 (t, J=7.2 Hz, 2H), 2.60 (t, J=7.2 Hz, 2H), 1.25 (t, J=8.0 Hz, 3H).

EXAMPLE 7

Synthesis of 3-(3,5-dimethoxyphenyl)-*N,N*-dimethyl-2-{4-[4-(3-oxo-3-ureido-propyl)-phenoxy]-phenyl}-acrylamide (**13**)

[000152] To a stirred solution of **6** (1.68 g, 3.43 mmol) in dry DMF (30 mL) was added carbonyldiimidazole (1.1 g, 6.86 mmol), and the reaction mixture was heated to 60°C for 1 hr. The reaction mixture was cooled to 0°C and a solution of dimethylamine in THF (2 M, 8.6 mL, 17.2 mmol) was added and stirred for 18 hr. The reaction mixture was diluted with water (100 mL) and extracted with ethyl acetate (100 mL). The organic phase was then rinsed sequentially with 10% citric acid (2 X 50mL), water (2 X 50mL), and brine (20 mL), then dried over anhydrous magnesium sulfate and evaporated. The crude product was purified by silica gel chromatography using hexane-ethyl acetate (3:7) containing 1% acetic acid. Yield: 1.77 g, 100%.




[000153] Analysis: ¹HNMR (DMSO-d₆): δ10.17 (br, 1H), 7.74 (br, 1H), 7.27 (d, J=9.2 Hz, 2H), 7.23 (d, J=8.8 Hz, 2H), 7.23 (br, 1H), 6.79 (d, J=9.2 Hz, 2H), 6.93 (d, J=8.8 Hz, 2H), 6.56 (s, 1H), 6.34 (t, J=2 Hz, 1H), 6.29 (s, 1H), 6.28 (s, 1H), 3.58 (s, 6H), 3.05 (br, 3H), 2.90 (br, 3H), 2.82 (t, J=7.2 Hz, J=8.0 Hz, 2H), 2.59 (t, J=8.0 Hz, J=7.2 Hz, 2H).

EXAMPLE 8

Synthesis of 2-(4-{4-[3-(3-cyclohexylureido)-3-oxopropyl]-phenoxy}-phenyl)-3-(3,5-dimethoxyphenyl)-acrylic acid (**14**)

[000154] Cyclohexylurea (1.3 g, 9 mmol) was dissolved in sodium ethoxide in ethanol (21% w/w, 3 mL) at 75°C. To this solution **5** was added (0.5 g, 1.1 mmol) in one lot. The resulting mixture was stirred at 75°C for 5 min, then cooled quickly to 40-50°C. TFA (0.5 mL) was added and then 5% aqueous HCl (1N, 0.6 mL). After stirring at room temperature for 1 hr, the mixture was left overnight at 4°C. The solid separated was filtered and refluxed in ethyl acetate (4 mL) for 20 min. The mixture was allowed to cool to room temperature, filtered and the crude product was purified by silica gel chromatography using hexane-ethyl acetate (1:1). Yield: 0.27 g, 45%.




[000155] Analysis: ^1H NMR (DMSO- d_6): δ 12.74 (s, 1H), 10.30 (s, 1H), 8.32 (br, 1H), 7.67 (s, 1H), 7.24 (d, J =8.8 Hz, 2H), 7.16 (d, J =8.8 Hz, 2H), 6.90 (d, J =8.4 Hz, 2H), 6.94 (d, J =8.4 Hz, 2H), 6.34 (t, J =2.4 Hz, 1H), 6.27 (d, J =2.4 Hz, 2H), 3.58 (s, 6H), 2.83 (t, J =7.6 Hz, 2H), 2.59 (t, J =7.6 Hz, 2H), 1.78 (m, 2H), 1.61 (m, 2H), 1.51 (m, 1H), 1.32-1.16 (m, 5H).

EXAMPLE 9

Synthesis of [3-(4-phenoxyphenyl)-propionyl]-urea (**15**)

[000156] 4-Phenoxy-benzaldehyde was reacted with triethyl phosphonoacetate to yield 3-(4-phenoxyphenyl)-acrylic acid ethyl ester, which was then reduced with H_2 using palladium-on-carbon catalyst to yield 3-(4-phenoxyphenyl)-propionic acid methyl ester (**19**). Urea (1.20 g, 19.99 mmol) was dissolved in sodium ethoxide (2 M, 6.7 mL, 13.4 mmol) at 80°C under argon, and to this a solution of **19** (1.71 g,

6.67 mmol) in anhydrous ethanol (8 mL) was added and heated at this temperature for 1 hr. Ethanol was evaporated under reduced pressure, water (20 mL) was added, acidified to pH 1 by 5% aqueous HCl and extracted with ethyl acetate (50 mL). The organic layer was washed with water (2 x 25 mL), brine (2 x 20 mL), dried over anhydrous magnesium sulfate and evaporated. The crude product was purified by silica gel chromatography and eluted with hexane-ethyl acetate (1:1) containing acetic acid (1%) followed by recrystallization from ethanol. Yield: 113 mg, 5.6%.



[000157] Analysis: ^1H NMR (DMSO- d_6): δ 10.18 (s, 1H), 7.74 (br, 1H), 7.38 (d, $J=7.6$ Hz, 1H), 7.36 (d, $J=7.6$ Hz, 1H), 7.22 (d, $J=8.8$ Hz, 2H), 7.17 (t, $J=7.2$ Hz, 1H), 6.97 (d, $J=7.2$ Hz, 2H), 6.93 (d, $J=8.8$ Hz, 2H), 2.82 (t, $J=7.2$ Hz, 2H), 2.59 (t, $J=7.2$ Hz, 2H).

EXAMPLE 10

Synthesis of 2-{4-[4-(3-acetyluroidomethyl)-phenoxy]-phenyl}-3-(3,5-dimethoxyphenyl)-acrylic acid methyl ester (**17**) [see Scheme II]

[000158] Step 1: Synthesis of 3-(3,5-dimethoxyphenyl)-2-[4-(4-hydroxymethyl)-phenoxy]-phenyl]-acrylic acid methyl ester (**22**). 3-(3,5-Dimethoxy-phenyl)-2-[4-(4-formylphenoxy)-phenyl]-acrylic acid methyl ester (**21**) was first prepared by converting the corresponding free acid (**3**) to the methyl ester by addition of DMF, K_2CO_3 and dimethyl sulfate in a manner analogous to Example 1(f) above. Sodium borohydride (0.125 g, 3.3 mmol) was added to a suspension of **21** (1.26 g, 3 mmol) in ethanol (20 mL) and stirred at room temperature for 1 hr. The reaction was quenched with 5% aqueous HCl, and ethanol was evaporated under reduced pressure. Residue was taken up in ethyl acetate (50 mL) and washed with brine (2 x

20 mL), dried over anhydrous magnesium sulfate and evaporated. The crude product was purified by silica gel chromatography and eluted with hexanes-ethyl acetate (1:1). Yield: 1.14 g, 95.0%. Analysis: ^1H NMR (DMSO- d_6): δ 7.72 (s, 1H), 7.36 (d, J =8.8 Hz, 2H), 7.19 (d, J =8.8 Hz, 2H), 7.01 (d, J =8.4 Hz, 2H), 6.99 (d, J =8.4 Hz, 2H), 6.41 (t, J =2.4 Hz, 1H), 6.28 (d, J =2.4 Hz, 2H), 5.18 (t, J =6.4 Hz, 1H), 4.49 (d, J =4.8 Hz, 2H), 3.72 (s, 3H), 3.57 (s, 6H).

[000159] (b) Step 2: Synthesis of 2-[4-(4-bromomethylphenoxy)-phenyl]-3-(3,5-dimethoxyphenyl)-acrylic acid methyl ester (23). To a stirred solution of **22** (1.05 g, 2.5 mmol) in dichloromethane (10 mL) at 10°C, PBr_3 (1 M, 3.75 mL) was added and stirred for 1 hr. The reaction was quenched with saturated aqueous sodium bicarbonate solution. The organic layer was washed with water (20 mL), brine (2 x 30 mL), dried over anhydrous magnesium sulfate and evaporated. The crude product was purified by silica gel chromatography and eluted with hexanes-ethyl acetate (4:1). Yield: 0.85 g, 70.4%. Analysis: ^1H NMR (DMSO- d_6): δ 7.73 (s, 1H), 7.49 (d, J =8.4 Hz, 2H), 7.22 (d, J =8.4 Hz, 2H), 7.07 (d, J =8.4 Hz, 2H), 7.00 (d, J =8.4 Hz, 2H), 6.42 (t, J =2.4 Hz, 1H), 6.28 (d, J =2.4 Hz, 2H), 4.74 (s, 2H), 3.73 (s, 3H), 3.58 (s, 6H).

[000160] (c) Synthesis of 2-[4-[4-(3-acetylureidomethyl)-phenoxy]-phenyl]-3-(3,5-dimethoxyphenyl)-acrylic acid methyl ester (17). To a stirred suspension of sodium hydride (60% in oil, 0.11 g, 2.8 mmol) in dimethylformamide (2 mL), *N*-acylurea (0.11 g, 1.12 mmol) was added and stirred at room temperature for 30 min. A solution of **23** (0.54 g, 1.12 mmol) in dimethylformamide (3 mL) was added and heated overnight at 80°C. The reaction was quenched with water and extracted with ethyl acetate (3 x 30 mL). The combined organic layer was washed with brine (2 x 25 mL), dried over anhydrous magnesium sulfate and evaporated. The crude product was purified by silica gel column chromatography and eluted with hexanes-ethyl acetate (3:7) containing 1% acetic acid. Yield: 0.16 g, 28.4%. Analysis: ^1H NMR (DMSO- d_6): δ 8.34 (t, J =5.6 Hz, 1H), 7.72 (s, 1H), 7.29 (d, J =8.4 Hz, 1H), 7.19 (d, J =8.4 Hz, 2H), 7.02 (d, J =8.4 Hz, 2H), 6.99 (d, J =8.4 Hz, 2H), 6.42 (t, J =8.4 Hz, 1H), 6.28 (d, J =2.4 Hz, 2H), 4.24 (d, J =5.2 Hz), 3.73 (s, 3H), 3.57 (s, 6H), 1.87 (s, 3H).

General Procedure for Conversion of Carboxylic Acids to Amides

[000161] A mixture of carboxylic acid (1.1 mmol) and carbonyldiimidazole (1.3 mmol) in DMF (20 mL) was heated at 60°C for 30 min. After the reaction mixture was cooled to room temperature, a solution of amine (2M, 1 mL, 2.0 mmol) was added and stirred for 18 hr. To the reaction mixture water (100 mL) was added and extracted with ethyl acetate (3 x 60 mL). The organic phase was washed with 10% citric acid (20 mL), water (2 x 50 mL), and brine (50 mL), then dried over anhydrous magnesium sulfate and removed the solvent. The crude product was purified by silica gel chromatography.

EXAMPLE 11


Synthesis of N,N-dimethyl-2-[4-[4-(3-oxo-3-ureidopropyl)-phenoxy]-phenyl]-acetamide (**26**)

[000162] Urea (0.78 g, 13 mmol) and 3-[4-(4-carboxymethylphenoxy)-phenyl]-propionic acid ethyl ester, **24** (0.5 g, 1.5 mmol) were dissolved in sodium ethoxide in ethanol (2M, 6.5 mL, 13 mmol) at 80°C under argon, and the reaction mixture was heated at this temperature for 1h. The reaction was then quenched by TFA (0.5 mL) after cooling to 5°C. Water (40 mL) was added to the reaction mixture. The crude product was filtered and purified by silica gel chromatography and eluted with hexane-ethyl acetate (1:1) containing acetic acid (1%) followed by recrystallization from toluene yielded **25** (0.28 g, 54%).

[000163] Analysis: ¹HNMR (DMSO-d₆): δ 12.28 (br, 1H), 7.73 (br, 1H), 7.24 (d, J=8.8 Hz, 2H), 7.23, (br, 1H), 7.21 (d, J=8.8 Hz, 2H), 6.93, (d, J=8.8 Hz, 2H), 6.92 (d, J=8.8 Hz, 2H), 3.54 (s, 2H), 2.81 (t, J=7.2 Hz, 2H), 2.58 (t, J=7.2 Hz, 2H).

[000164] Following the general procedure for conversion of carboxylic acids to amides mentioned above and using dimethyl amine as amine, **25** was converted to **26** in 97% yield.

[000165] Analysis: $^1\text{H NMR}$ ($\text{DMSO}-d_6$): δ 10.17 (s, 1H), 7.73 (s, 1H), 7.22 (s, 1H), 7.21 (d, $J=8.0$ Hz, 2H), 7.19 (d, $J=8.0$ Hz, 2H), 6.92 (d, $J=8.0$ Hz, 2H), 6.90 (d, $J=8.0$ Hz, 2H), 3.65 (s, 2H), 3.00 (s, 3H), 2.81 (t, $J=8.0$ Hz, 2H), 2.58 (t, $J=8.0$ Hz, 2H).



EXAMPLE 12

Synthesis of (4-{4-[2-(3,5-dimethoxyphenyl)-1-dimethylcarbamoyl-vinyl]-phenoxy}-benzyl)-carbamic acid methyl ester (**29**)


[000166] Reaction of 3-(3,5-dimethoxyphenyl)-2-{4-[4-(2,4-dioxothiazolidin-3-ylmethyl)-phenoxy]-phenyl}-acrylic acid, **27**, (0.4 g, 0.77 mmol) with 5% LiOH (2mL) in methanol (19 mL) was carried out at room temperature for 18 h.

The reaction mixture was acidified to pH 3 by 5% aqueous HCl and extracted with ethyl acetate (2 x 50 mL). The organic layer was washed with water (2 x 50 mL), brine (2 x 20 mL), dried over anhydrous magnesium sulfate and evaporated. The crude product was purified by silica gel chromatography and eluted with hexane-ethyl acetate (1:1) containing acetic acid (1%). Yield (**28**): 0.31g, 83%.

Analysis: $^1\text{H NMR}$ ($\text{DMSO}-d_6$): δ 12.75 (br, 1H), 7.68 (t, $J=4.6$ Hz, 1H), 7.67 (s, 1H), 7.28 (d, $J=8.8$ Hz, 2H), 7.17 (d, $J=8.8$ Hz, 2H), 7.01 (d, $J=8.8$ Hz, 2H), 6.97 (d, $J=8.8$ Hz, 2H), 6.39 (t, $J=2.8$ Hz, 1H), 6.27 (d, $J=2.4$ Hz, 2H), 4.17 (d, $J=6.4$ Hz, 2H), 3.58 (s, 6H), 3.55 (s, 3H).

[000167] Following the general procedure for conversion of carboxylic acids to amides mentioned above and using dimethyl amine as amine, **28** was converted to **29** in 96% yield.

[000168] Analysis: $^1\text{H NMR}$ (DMSO-d_6): δ 7.68 (t, $J=4.6$ Hz, 1H), 7.28 (d, $J=8.8$ Hz, 2H), 7.27 (d, $J=8.8$ Hz, 2H), 6.98 (d, $J=8.8$ Hz, 2H), 6.96 (d, $J=8.8$ Hz, 2H), 6.57 (s, 1H), 6.35 (t, $J=2.8$ Hz, 1H), 6.28 (d, $J=2.4$ Hz, 2H), 4.16 (d, $J=6.4$ Hz, 2H), 3.59 (s, 6H), 3.55 (s, 3H), 3.05 (br, 3H), 2.91 (br, 3H).



EXAMPLE 13


Synthesis of 2-{4-[4-(2-carbamoyl-1H-imidazol-1-yl)phenoxy]phenyl}-3-(3,5-dimethoxyphenyl)-N,N-dimethylacrylamide (**31**)

[000169] Urea (0.78 g, 13 mmol) and 3-(3,5-dimethoxyphenyl)-2-{4-[4-(2-ethoxycarbonyl-1H-imidazol-1-yl)phenoxy]phenyl}acrylic acid **5** (0.45 g, 1.5 mmol) were dissolved in sodium ethoxide in ethanol (2M, 6.5 mL, 13 mmol) at 80°C under argon, and the reaction mixture was heated at this temperature for 5 h. The reaction was then quenched by TFA (0.5 mL) after cooling to 5°C. Water (40 mL) was added to the reaction mixture. The crude product was filtered and purified by silica gel chromatography and eluted with hexane-ethyl acetate (1:1) containing acetic acid (1%). Yield (**30**): 0.39 g, 93%.

[000170] Analysis: $^1\text{H NMR}$ (DMSO-d_6): δ 12.73 (br, 1H), 7.68 (s, 1H), 7.29 (br, 1H), 7.24 (d, $J=8.8$ Hz, 2H), 7.65 (d, $J=8.8$ Hz, 2H), 6.99 (d, $J=8.8$ Hz, 2H), 6.92 (d, $J=8.8$ Hz, 2H), 6.78 (br, 1H), 6.39 (t, $J=2.4$ Hz, 1H), 6.27 (d, $J=2$ Hz, 2H), 3.57 (s, 6H), 2.79 (t, $J=8.0$ Hz, 2H), 2.35 (t, $J=8.0$ Hz, 2H).

[000171] Following the general procedure for conversion of carboxylic acids to amides mentioned above and using dimethyl amine as amine, **30** was converted to **31** in 98% yield.

[000172] Analysis: $^1\text{H NMR}$ (DMSO-d_6): δ 7.30 (br, 1H), 7.28 (d, $J=8.8$ Hz, 2H), 7.23 (d, $J=8.8$ Hz, 2H), 6.95 (d, $J=8.8$ Hz, 2H), 6.92 (d, $J=8.8$ Hz, 2H), 6.79 (br, 1H), 6.56 (s, 1H), 6.34 (t, $J=2.4$ Hz, 1H), 6.28 (d, $J=2$ Hz, 2H), 3.58 (s, 6H), 3.05 (br, 3H), 2.90 (br, 3H), 2.77 (t, $J=8.0$ Hz, 2H), 2.34 (t, $J=8.0$ Hz, 2H).



EXAMPLE 14


Synthesis of 2-[4-(4-acetylaminophenoxy)-phenyl]-3-(3,5-dimethoxyphenyl)-N,N-dimethylacrylamide (**34**)

[000173] Compound **2** was reacted with 1-fluoro-4-nitrobenzene in the presence of NaH in DMF to give 3-(3,5-dimethoxyphenyl)-2-[4-(4-nitrophenoxy)-phenyl]-acrylic acid (**32**). Reduction of **32** (10 g, 24 mmol) with zinc dust (15 g, 230 mmol) in acetic acid (100 mL) was accomplished at 120°C for 15 h, the mixture was cooled to room temperature. Water (250 mL) was slowly added to the reaction mixture. The precipitated product was filtered and washed with water (70 mL) to give crude product. The product was recrystallized from toluene. Yield (**33**): 9.7 g, 94%.

[000174] Analysis: $^1\text{H NMR}$ (DMSO-d_6): δ 12.35 (br, 1H), 9.96 (s, 1H), 7.67 (s, 1H), 7.60 (d, $J=8.8$ Hz, 2H), 7.15 (d, $J=8.8$ Hz, 2H), 6.97 (d, $J=8.8$ Hz, 2H), 6.96 (d, $J=8.8$ Hz, 2H), 6.34 (t, $J=2.8$ Hz, 1H), 6.28 (d, $J=2.4$ Hz, 2H), 3.58 (s, 6H), 2.03 (s, 3H).

[000175] Following the general procedure for conversion of carboxylic acids to amides mentioned above and using dimethylamine as amine, **33** was converted to **34** in 98% yield.

[000176] Analysis: $^1\text{H NMR}$ ($\text{DMSO}-d_6$): δ 9.96 (s, 1H), 7.60 (d, $J=8.8$ Hz, 2H), 7.25 (d, $J=8.8$ Hz, 2H), 6.97 (d, $J=8.8$ Hz, 2H), 6.93 (d, $J=8.8$ Hz, 2H), 6.55 (s, 1H), 6.34 (t, $J=2.8$ Hz, 1H), 6.28 (d, $J=2.4$ Hz, 2H), 3.58 (s, 6H), 3.04 (br, 3H), 2.90 (br, 3H), 2.03 (s, 3H).



EXAMPLE 15


Synthesis of 3-(3,5-dimethoxyphenyl)-2-[4-(4-methanesulfonylphenoxy)-phenyl]-N,N-dimethylacrylamide (**36**)

[000177] Compound **2** (3 g, 10 mmol) was dissolved in anhydrous DMF (70 mL) under nitrogen, and potassium carbonate (1.4 g, 10 mol) was added in lots. When the solution became homogeneous, 4-fluorophenyl methyl sulfone (1.74 g, 10 mmol) was added and the mixture was heated at 150°C for 2 h. After cooling to room temperature, the solution was poured into water (150 mL). The mixture was acidified with 5% HCl to ~ pH 4 and the solidified product was collected by suction filtration. The crude product was recrystallized with toluene. Yield(**35**): 4.3 g, 96%.

[000178] Analysis: $^1\text{H NMR}$ ($\text{DMSO}-d_6$): δ 12.72 (br, 1H), 7.94 (d, $J=8.8$ Hz, 2H), 7.72 (s, 1H), 7.80 (d, $J=8.4$ Hz, 2H), 7.18 (d, $J=8.8$ Hz, 2H), 7.17 (d, $J=8.4$ Hz, 2H), 6.42 (t, $J=2.8$ Hz, 1H), 6.28 (d, $J=2.4$ Hz, 2H), 3.59 (s, 6H), 3.21 (s, 3H).

[000179] Following the general procedure for conversion of carboxylic acids to amides mentioned above and using dimethylamine as amine, **35** was converted to **36** in 96% yield.

[000180] Analysis: $^1\text{H NMR}$ (DMSO-d_6): δ 7.93 (d, $J=8.8$ Hz, 2H), 7.38 (d, $J=8.4$ Hz, 2H), 7.17 (d, $J=8.8$ Hz, 2H), 7.16 (d, $J=8.4$ Hz, 2H), 6.62 (s, 1H), 6.36 (t, $J=2.8$ Hz, 1H), 6.29 (d, $J=2.4$ Hz, 2H), 3.59 (s, 6H), 3.20 (s, 3H), 3.08 (br, 3H), 2.92 (br, 3H).




EXAMPLE 16

Synthesis of 3-(4-{4-[2-(3,5-dimethoxyphenyl)-1-dimethylcarbamoylvinyl]-phenoxy}-phenyl)-propionic acid ethyl ester (**37**)

[000181] Following the general procedure for conversion of carboxylic acids to amides mentioned above and using dimethyl amine as amine, **5** was converted to **37** in 97% yield.

[000182] Analysis: $^1\text{H NMR}$ (DMSO-d_6): δ 7.28 (d, $J=8.8$ Hz, 2H), 7.23 (d, $J=8.8$ Hz, 2H), 6.95 (d, $J=8.8$ Hz, 2H), 6.92 (d, $J=8.8$ Hz, 2H), 6.56 (s, 1H), 6.34 (t, $J=2.4$ Hz, 1H), 6.28 (d, $J=2$ Hz, 2H), 4.04 (q, $J=6.8$ Hz, 2H), 3.58 (s, 6H), 3.05 (br, 3H), 2.90 (br, 3H), 2.84 (t, $J=8.4$ Hz, 2H), 2.61 (t, $J=8.4$ Hz, 2H), 1.15 (t, $J=6.4$ Hz, 3H).




EXAMPLE 17

Synthesis of 2-[4-[4-(N-ureido-2-carbamoylethyl)-phenoxy]-phenyl]-3-(3,5-dimethoxyphenyl)-N,N- dimethylacrylamide (**39**)

[000183] Hydrolysis of **13** with 1N NaOH yielded **38**. The 1,1-carbonyl-diimidazole (CDI) derivative was made by the general procedure for conversion of carboxylic acids to amides mentioned above. The CDI intermediate of **38** was converted to **39** by reacting this with semicarbazide in 73% yield.

[000184] Analysis: ^1H NMR (DMSO- d_6): δ 9.48 (br, 1H), 7.72 (br, 1H), 7.28 (d, $J=8.8$ Hz, 2H), 7.25 (d, $J=8.8$ Hz, 2H), 6.95 (d, $J=8.8$ Hz, 2H), 6.92 (d, $J=8.8$ Hz, 2H), 6.56 (s, 1H), 6.34 (t, $J=2.4$ Hz, 1H), 6.28 (d, $J=2$ Hz, 2H), 5.86 (s, 2H), 3.58 (s, 6H), 3.05 (br, 3H), 2.90 (br, 3H), 2.77 (t, $J=8.0$ Hz, 2H), 2.39 (t, $J=8.0$ Hz, 2H).




EXAMPLE 18

Synthesis of 3-(3,5-dimethoxyphenyl)-N,N-dimethyl-2-{4-[4-(3-morpholin-4-yl-3-oxopropyl)-phenoxy]-phenyl}-acrylamide (**40**)

[000185] The CDI intermediate of **38** was converted to **40** by reacting it with morpholine in 94% yield.

[000186] Analysis: ^1H NMR (DMSO- d_6): δ 7.27 (d, $J=8.8$ Hz, 2H), 7.26 (d, $J=8.8$ Hz, 2H), 6.95 (d, $J=8.8$ Hz, 2H), 6.92 (d, $J=8.8$ Hz, 2H), 6.56 (s, 1H), 6.34 (t, $J=2.4$ Hz, 1H), 6.28 (d, $J=2$ Hz, 2H), 3.58 (s, 6H), 3.49 (m, 4H), 3.41 (m, 4H), 3.05 (br, 3H), 2.90 (br, 3H), 2.77 (t, $J=8.0$ Hz, 2H), 2.39 (t, $J=8.0$ Hz, 2H).




EXAMPLE 19

Synthesis of 2-(4-{4-[2-(3,5-dimethoxyphenyl)-1-dimethylcarbamoylvinyl]-phenoxy}-benzyl)-malonic acid dimethyl ester (**43**)

[000187] Condensation of **3** with malonic acid dimethyl ester in the presence of sodium hydride as base resulted in **41**, which on reduction with zinc/acetic acid yielded **42**. Conversion of **42** to **43** was accomplished by the general procedure for conversion of carboxylic acids to amides mentioned above in 94% yield.

Analysis: ^1H NMR (DMSO- d_6): δ 7.29 (d, $J=8.8$ Hz, 2H), 7.23 (d, $J=8.8$ Hz, 2H), 6.96 (d, $J=8.8$ Hz, 2H), 6.92 (d, $J=8.8$ Hz, 2H), 6.57 (s, 1H), 6.34 (t, $J=2.4$ Hz, 1H), 6.28 (d, $J=2$ Hz, 2H), 3.87 (t, $J=8$ Hz, 1H), 3.61 (s, 6H), 3.58 (s, 6H), 3.08 (d, $J=7.6$ Hz, 2H), 3.05 (br, 3H), 2.91 (br, 3H).




EXAMPLE 20

Synthesis of N-{4-[2-(3,5-dimethoxyphenyl)-1-dimethylcarbamoylvinyl]-phenyl}-3-hydroxybenzamide (**44**)

[000188] A mixture of 2-(4-aminophenyl)-3-(3,5-dimethoxyphenyl)-N,N-dimethylacrylamide, **43**, (0.59 g, 1.5 mmol), benzotriazol-1-yloxytris-(dimethylamino)-phosphonium hexafluorophosphate (BOP, 0.88g, 2.0 mmol), 3-hydroxybenzoic acid (0.28g, 2.0 mmol), triethylamine (0.2 g, 2.0 mmol) in DMF (8.0 mL) was stirred for 3h at room temperature. The reaction mixture was poured in water (50mL) and solid separated was filtered, dried and purity was checked by HPLC (97.6%).

[000189] Analysis: $^1\text{H NMR}$ (DMSO-d_6): δ 10.29 (s, 1H), 9.81 (s, 1H), 7.79 (d, $J=6.8\text{Hz}$, 2H), 7.43 (d, $J=8.0\text{Hz}$, 1H), 7.37 (t, $J=7.6\text{Hz}$, 2H), 7.29 (d, $J=8.4\text{Hz}$, 2H), 7.02 (m, 1H), 6.60 (s, 1H), 6.40 (t, $J=2.0\text{Hz}$, 1H), 6.36 (d, $J=2.0\text{Hz}$, 2H), 3.63 (s, 6H), 3.08 (brs, 3H), 2.96 (brs, 3H).



EXAMPLE 21


Synthesis of N,N-dimethyl-2-{4-[4-(3-oxo-3-ureidopropenyl)-phenoxy]-phenyl}-3-pyridin-3-ylacrylamide (**47**)

[000190] Synthesis of **45** from 3-pyridinecarboxaldehyde was performed following Scheme I. Urea (0.78 g, 13 mmol) and 2-{4-[4-(2-ethoxycarbonyl-vinyl)-phenoxy]-phenyl}-3-pyridin-3-ylacrylic acid, **45** (0.5 g, 1.2 mmol) was dissolved in sodium ethoxide in ethanol (2M, 6.5 mL, 13 mmol) at 80°C under argon, and the reaction mixture was heated at this temperature for 1 h. The reaction was then quenched by TFA (0.5 mL) after cooling to 5°C. Water (40 mL) was added to the reaction mixture. The crude product was filtered and purified by silica gel chromatography and eluted with hexanes-ethyl acetate (1:1) containing acetic acid (1%) followed by recrystallization from toluene. Yield (**46**): 0.33 g, 63%.

[000191] Analysis: $^1\text{H NMR}$ (DMSO- d_6): δ 12.78 (br, 1H), 10.29 (s, 1H), 8.42 (dd, $J=4.8, 1.6$ Hz, 1H), 8.35 (d, $J=2.4$ Hz, 1H), 7.92 (br, 1H), 7.66 (d, $J=16$ Hz, 1H), 7.64 (d, $J=8.8$ Hz, 2H), 7.36 (tt, $J=8.4, 1.6$ Hz, 1H), 7.30 (br, 1H), 7.28 (m, 1H), 7.23 (d, $J=8.8$ Hz, 2H), 7.11 (d, $J=8.8$ Hz, 2H), 7.09 (d, $J=8.8$ Hz, 2H), 6.73 (d, $J=16$ Hz, 1H).

[000192] Following the general procedure for conversion of carboxylic acids to amides mentioned above, **46** was converted to **47**.

[000193] Analysis: $^1\text{H NMR}$ (DMSO- d_6): δ 10.30 (s, 1H), 8.39 (dd, $J=4.8, 1.6$ Hz, 1H), 8.34 (d, $J=2.4$ Hz, 1H), 7.92 (br, 1H), 7.66 (d, $J=16$ Hz, 1H), 7.64 (d, $J=8.8$ Hz, 2H), 7.45 (tt, $J=8.4, 1.6$ Hz, 1H), 7.32 (br, 1H), 7.29 (d, $J=8.8$ Hz, 2H), 7.26 (m, 1H), 7.11 (d, $J=8.8$ Hz, 2H), 7.05 (d, $J=8.8$ Hz, 2H), 6.73 (d, $J=16$ Hz, 1H), 6.70 (s, 1H), 3.07 (br, 3H), 2.93 (br, 3H).




EXAMPLE 22

Synthesis of 3-(3,5-dimethoxyphenyl)-2-(4-hydroxyphenyl)-N,N-dimethylacrylamide (49)

[000194] Following the general procedure for conversion of carboxylic acids to amides mentioned above and using dimethyl amine as amine, **2** was converted to **49**.

[000195] Analysis: ^1H NMR (DMSO- d_6): δ 9.59 (s, 1H), 7.07 (d, $J=8.8$, 2H), 6.73 (d, $J=8.8$ Hz, 2H), 6.43 (s, 1H), 6.23 (t, $J=2.4$ Hz, 1H), 6.29 (d, $J=2.4$ Hz, 2H), 3.57 (s, 6H), 2.99 (brs, 3H), 2.89 (brs, 3H).




EXAMPLE 23

Synthesis of [3-(4-{4-[2-(3,5-dimethoxyphenyl)-1-(piperidine-1-carbonyl)-vinyl]-phenoxy}-phenyl)-propionyl]-urea (51)

[000196] Following the general procedure for conversion of carboxylic acids to amides mentioned above and using piperidine as amine, **6** was converted to **51**.

[000197] Analysis: ^1H NMR (DMSO- d_6): δ 10.16 (s, 1H), 7.73 (brs, 1H), 7.26 (d, $J=8.8$ Hz, 2H), 7.23 (d, $J=8.8$ Hz, 2H), 6.98 (d, $J=8.8$ Hz, 2H), 6.93 (d, $J=8.8$ Hz, 2H), 6.55 (s, 1H), 6.34 (t, $J=2.4$ Hz, 1H), 6.29 (d, $J=2.4$ Hz, 2H), 3.58 (s, 6H), 3.50 (br, 4H), 2.82 (t, $J=7.6$ Hz, 2H), 2.59 (t, $J=7.6$ Hz, 2H), 1.58 (br, 2H) 1.40-1.45 (br, 4H).




EXAMPLE 24

Synthesis of 3-(3,5-dimethoxyphenyl)-N,N-diethyl-2-{4-[4-(3-oxo-3-ureidopropyl)phenoxy]phenyl}-acrylamide (**53**)

[000198] Following the general procedure for conversion of carboxylic acids to amides mentioned above and using diethylamine as amine, **6** was converted to **53**.

[000199] Analysis: ^1H NMR (DMSO- d_6): δ 10.17 (s, 1H), 7.70 (brs, 1H), 7.26 (overlapped d, $J=8.8\text{Hz}$, 2H), 7.23 (overlapped d, $J=8.8\text{Hz}$, 2H), 6.97 (d, $J=8.8\text{Hz}$, 2H), 6.92 (d, $J=8.8\text{Hz}$, 2H), 6.54 (s, 1H), 6.34 (t, $J=2.0\text{Hz}$, 1H), 6.29 (d, $J=2.0\text{Hz}$, 2H), 3.32-3.37 (br, 4H), 3.59 (s, 6H), 2.82 (t, $J=7.6\text{Hz}$, 2H), 2.59 (t, $J=7.6\text{Hz}$, 2H), 1.03 (br, 3H), 0.92 (br, 3H).



EXAMPLE 25


Synthesis of 2-{4-[4-(3-acetylamino-3-oxopropyl)-phenoxy]-phenyl}-3-(4-fluorophenyl)-N,N-dimethylacrylamide (**56**)

[000200] To a solution of {4-[4-(2-carbamoylethyl)-phenoxy]-phenyl}-acetic acid, **54**, (0.45g, 1.5mmol) in acetic anhydride (15 mL) was added 4-fluorobenzaldehyde (0.17 mL, 1.6 mmol) and potassium acetate (0.17g, 1.8 mmol) and refluxed overnight. Reaction mixture was poured in water (50 mL) and extracted with ethyl acetate (2 x 50mL). The crude product was purified by silica gel chromatography to yield **55**.

[000201] Analysis: $^1\text{H NMR}$ ($\text{DMSO}-d_6$): δ 12.50 (br, 1H), 10.64 (s, 1H), 7.74 (s, 1H), 7.27 (d, $J=8.4\text{Hz}$, 2H), 7.10-7.15 (m, 6H), 6.99 (d, $J=8.4\text{Hz}$, 2H), 6.97 (d, $J=8.4\text{Hz}$, 2H), 2.81 (d, $J=6.8\text{Hz}$, 2H), 2.76 (d, $J=6.8\text{Hz}$, 2H), 2.15 (s, 3H).

[000202] Following the general procedure for conversion of carboxylic acids to amides mentioned above and using dimethylamine as amine, **55** was converted to **56**.

[000203] Analysis: $^1\text{H NMR}$ ($\text{DMSO}-d_6$): δ 10.62 (s, 1H), 7.26 (d, $J=8.4\text{Hz}$, 2H), 7.22 (d, $J=8.4\text{Hz}$, 2H), 7.15 (d, $J=8.4\text{Hz}$, 2H), 7.05 (d, $J=8.4\text{Hz}$, 2H), 6.97 (d, $J=8.0\text{Hz}$, 2H), 6.94 (d, $J=8.0\text{Hz}$, 2H), 6.63 (s, 1H), 2.81 (d, $J=6.8\text{Hz}$, 2H), 2.76 (d, $J=6.8\text{Hz}$, 2H), 2.15 (s, 3H).




EXAMPLE 26

Synthesis of 2-(4-{4-[2-(3,5-dimethoxyphenyl)-1-dimethylcarbamoylvinyl]-phenoxy}-benzyl)-malonic acid (**58**) and 2-(4-{4-[2-(3,5-dimethoxyphenyl)-1-dimethylcarbamoylvinyl]-phenoxy}-benzyl)-malonamide (**59**)

[000204] To a solution of 2-(4-{4-[2-(3,5-dimethoxyphenyl)-1-dimethylcarbamoylvinyl]-phenoxy}-benzyl)-malonic acid dimethyl ester, **43** (0.40g, 0.73mmol) in DMF (6 mL) and ethanol (10 mL), ammonium hydroxide (20 mL, 28%) and 1N NaOH (0.36 mL, 0.36 mmol) was added and stirred overnight at room temperature. Solvent was evaporated and the crude product was purified by silica gel chromatography to yield **58** and **59**.

Analysis: ^1H NMR (DMSO- d_6 + D_2O) of **58**: δ 7.20 (d, $J=8.4\text{Hz}$, 2H), 7.17 (d, $J=8.4\text{Hz}$, 2H), 6.90 (d, $J=8.4\text{Hz}$, 2H), 6.81 (d, $J=8.4\text{Hz}$, 2H), 6.51 (s, 1H), 6.29 (t, $J=2.0\text{Hz}$, 1H), 6.21 (d, $J=2.0\text{Hz}$, 2H), 3.53 (s, 6H), 3.13 (br, 1H), 3.01 (brs, 3H), 2.92 (br, 2H), 2.86 (brs, 3H).

[000205] Analysis: ^1H NMR (DMSO- d_6) of **59**: δ 7.28 (d, $J=8.8\text{ Hz}$, 2H), 7.26 (br, 2H), 7.22 (d, $J=8.8\text{ Hz}$, 2H), 7.03 (br, 2H), 6.97 (d, $J=8.8\text{ Hz}$, 2H), 6.90 (d, $J=8.8\text{ Hz}$, 2H), 6.56 (s, 1H), 6.34 (t, $J=2.4\text{ Hz}$, 1H), 6.28 (d, $J=2\text{ Hz}$, 2H), 3.58 (s, 6H), 3.29 (t, $J=8\text{ Hz}$, 1H), 3.05 (br, 3H), 2.95 (d, $J=7.6\text{ Hz}$, 2H), 2.91 (br, 3H).




EXAMPLE 27

Synthesis of 3-(3,5-dimethoxyphenyl)-2-{4-[4-(3-oxo-3-ureidopropyl)-phenoxy]-phenyl}-N-pyridin-4-ylacrylamide (**60**)

[000206] Following the general procedure for conversion of carboxylic acids to amides mentioned above and using 4-aminopyridine as amine, **6** was converted to **60**.

[000207] Analysis: ^1H NMR (DMSO- d_6): δ 10.17 (s, 1H), 8.24 (brs, 1H), 7.71 (br, 2H), 7.53 (d, $J=8.8\text{Hz}$, 2H), 7.44 (s, 1H), 7.25 (d, $J=8.4\text{Hz}$, 2H), 7.22 (br, 1H), 7.03 (d, $J=9.2\text{Hz}$, 2H), 7.99 (d, $J=8.4\text{Hz}$, 2H), 6.47 (d, $J=2.4\text{Hz}$, 2H), 6.43 (t, $J=2.4\text{Hz}$, 2H), 3.65 (s, 6H), 2.83 (t, $J=7.6\text{Hz}$, 2H), 2.60 (t, $J=7.6\text{Hz}$, 2H).




EXAMPLE 28

Synthesis of N-(4-chlorophenyl)-3-(3,5-dimethoxyphenyl)-2-{4-[4-(3-oxo-3-ureidopropyl)-phenoxy]-phenyl}-acrylamide (**61**)

[000208] Following the general procedure for conversion of carboxylic acids to amides mentioned above and using 4-chloroaniline as amine, **6** was converted to **61**.

[000209] Analysis: ^1H NMR (DMSO- d_6): δ 10.16 (s, 1H), 8.24 (brs, 1H), 7.65 (brs, 1H), 7.53 (d, $J=8.8\text{Hz}$, 2H), 7.44 (s, 1H), 7.25 (d, $J=8.8\text{Hz}$, 2H), 7.22 (br, 1H), 7.03 (d, $J=8.8\text{Hz}$, 2H), 7.00 (d, $J=8.8\text{Hz}$, 2H), 6.47 (d, $J=2.4\text{Hz}$, 2H), 6.43 (d, $J=2.4\text{Hz}$, 1H), 3.66 (s, 6H), 2.83 (t, $J=8.0\text{Hz}$, 2H), 2.60 (t, $J=8.0\text{Hz}$, 2H).




61

EXAMPLE 29

Synthesis of 3-(3,5-dimethoxyphenyl)-N,N-dimethyl-2-(4-{4-[2-(2-morpholin-4-yl-2-oxoethylcarbamoyl)-ethyl]-phenoxy}-phenyl)-acrylamide (**63**)

[000210] Following the general procedure for conversion of carboxylic acids to amides mentioned above and using 2-amino-1-morpholin-4-yl-ethanone as amine, 3-(4-{4-[2-(3,5-dimethoxyphenyl)-1-dimethylcarbamoylvinyl]-phenoxy}-phenyl)-propionic acid, **38**, was converted to **63**.

[000211] Analysis: ^1H NMR (DMSO- d_6): δ 7.99 (t, $J=5.6\text{Hz}$, 1H), 7.27 (d, $J=8.8\text{Hz}$, 2H), 7.24 (d, $J=8.8\text{Hz}$, 2H), 6.97 (d, $J=8.8\text{Hz}$, 2H), 6.92 (d, $J=8.8\text{Hz}$, 2H), 6.56 (s, 1H), 6.34 (t, $J=2.0\text{Hz}$, 1H), 6.28 (d, $J=2.0\text{Hz}$, 2H), 3.93 (d, $J=5.6\text{Hz}$, 2H), 3.56 (s, 6H), 3.52-3.56 (m, 4H), 3.40-3.42 (m, 4H), 3.05 (brs, 3H), 2.91 (brs, 3H), 2.80 (t, $J=7.6\text{Hz}$, 2H), 2.46 (t, $J=7.6\text{Hz}$, 2H).




EXAMPLE 30

Synthesis of 3-(3,5-dimethoxyphenyl)-N,N-dimethyl-2-(4-{4-[3-(4-methylpiperazin-1-yl)-3-oxopropyl]-phenoxy}-phenyl)-acrylamide (**64**)

[000212] Following the general procedure for conversion of carboxylic acids to amides mentioned above and using 4-methylpiperazine as amine, 3-(4-{4-[2-(3,5-dimethoxyphenyl)-1-dimethylcarbamoylvinyl]-phenoxy}-phenyl)-propionic acid, **38**, was converted to **64**.

[000213] Analysis: ^1H NMR (DMSO- d_6): δ 7.28 (d, $J=2.8\text{Hz}$, 2H), 7.25 (d, $J=2.8\text{Hz}$, 2H), 6.96 (d, $J=8.8\text{Hz}$, 2H), 6.92 (d, $J=8.6\text{Hz}$, 2H), 6.56 (s, 1H), 6.34 (t, $J=2.0\text{Hz}$, 1H), 6.28 (d, $J=2.0\text{Hz}$, 2H), 6.19 (s, 6H), 3.40 (dt, $J=18.0$ and 4.8Hz), 3.04 (brs, 3H), 2.90 (brs, 3H), 2.79 (t, $J=8.0$, 2H), 2.60 (t, $J=8.0\text{Hz}$, 2H), 2.20 (t, $J=5.2\text{Hz}$, 2H), 2.14 (s, 3H).




EXAMPLE 31

Synthesis of 3-(3,5-dimethoxyphenyl)-N,N-dimethyl-2-[4-(pyridin-2-yloxy)-phenyl]-acrylamide (**66**)

[000214] A solution of 3-(3,5-dimethoxyphenyl)-2-(4-hydroxyphenyl)-acrylic acid, **2**, (0.6 g, 2.0 mmol), 2-fluoropyridine (0.19 g, 2.0 mmol) in dimethyl acetamide (4.0 mL) was heated in presence of potassium carbonate (0.28 g, 2.0 mmol) at 175°C for 2 h, and then quenched with water (25 mL), neutralized with dilute HCl and extracted with ethyl acetate (2 x 50mL). Organic layer was dried and evaporated. The crude product was purified by silica gel chromatography to yield **65** (0.15 g, 19.9%).

[000215] A mixture of 3-(3,5-dimethoxyphenyl)- 2-[4-(pyridin-2-yloxy)-phenyl]-acrylic acid, **65**, (0.11g, 0.3 mmol), benzotriazol-1-yloxytris-(dimethylamino)-phosphonium hexafluorophosphate (BOP, 0.15 g, 0.35 mmol), dimethylamine in THF (2M, 0.5 mL, 1.0 mmol), triethylamine (0.035 g, 035 mmol) in DMF (6.0 mL) was stirred for 3 h at room temperature. The reaction mixture was poured in water (50.0 mL) and extracted with ethyl acetate (2 x 50 mL). Solvent was evaporated under reduced pressure and residue was purified by silica gel chromatography to yield **66**.

[000216] Analysis: ¹HNMR (DMSO-d₆): δ 8.14 (m, 1H), 7.88 (m, 1H), 7.33 (d, J=8.8 Hz, 2H), 7.14 (m, 3H), 7.05 (d, J=8.4Hz, 2H), 6.59 (s, 1H), 6.34 (t, J=2.0Hz, 1H), 6.31 (d, J=2.0Hz, 2H), 3.58(s, 6H), 3.10 (brs, 3H), 2.92 (brs, 3H).



EXAMPLE 32

Measurement of Increased Glucose Uptake in 3T3-L1 Adipocytes Treated With a Compound of the Present Invention

[000217] The effect of treatment with **1** on glucose uptake was measured in 3T3-L1 differentiated adipocytes. The assay was conducted essentially according to the method of Tafuri SR, *Endocrinology*, 137, 4706-4712 (1996). The adipocytes were incubated with different concentrations of the test compound for 48 hours in Dulbecco's modified Eagle's medium (DMEM) containing 10% fetal bovine serum (FBS), then washed and incubated in glucose-free, serum-free medium for 60 minutes at 37°C. Then ¹⁴C-deoxyglucose was added and the cells were incubated for 30 minutes at room temperature. After washing, the cells were lysed (0.1% SDS) and the radioactivity was measured to determine the amount of glucose uptake. Glucose uptake was calculated as a percentage of the basal level seen in cells not

treated with drug. As shown in FIG. 1, treatment with **1** resulted in a dose-dependent increase in glucose uptake.

EXAMPLE 33

Measurement of Enhanced Glucose Uptake in 3T3-L1 Adipocytes Treated With Insulin in Combination with a Compound of the Present Invention

[000218] The ability of **1** to enhance insulin-stimulated glucose uptake was assessed in 3T3-L1 adipocytes essentially as described above in Example 32. Adipocytes were incubated with either vehicle (0.1% DMSO) or test compound (5 μ M **1**) for 48 hours in DMEM plus 10% FBS. The cells were then serum-starved, incubated for 30 minutes with different concentrations of insulin, and then glucose uptake was carried out for 10 minutes at room temperature. When compared to treatment with vehicle, treatment with **1** enhanced the stimulation of glucose uptake by insulin (see Figure 2).

EXAMPLE 34

Measurement of the Glucose-Lowering Effect in ob/ob Mice Treated With a Compound of the Present Invention

[000219] The glucose-lowering effect of **1** was measured in ob/ob mice, an animal model for type 2 diabetes. At the onset of diabetes, seven-week-old male ob/ob mice received daily oral doses of either vehicle (0.5% CMC) or **1** (10 mg/kg) by gavage for seven days. Blood glucose levels were measured on day 0 (24 hours prior to administration of the first dose), day 1 (immediately prior to the first dose), and on days 2, 4, 6 and 8 (24 hours following administration of the prior dose). Significant decreases in blood glucose levels were recorded on day 6 (36% decrease, $p < 0.05$) and day 8 (23% decrease, $p < 0.05$) in the drug-treated versus the vehicle-treated animals (see Figure 3).

EXAMPLE 35

Measurement of the Lipid-Lowering Effects in ob/ob Mice Treated With a Compound of the Present Invention

[000220] The lipid-lowering effects of **1** also were measured in ob/ob mice following one week of treatment. In the experiment described above in Example 34, the concentrations of serum triglycerides and free fatty acids were determined on day 8. Significant decreases were observed in the levels of serum triglycerides (49% lower, $p < 0.05$) and free fatty acids (19% lower, $p < 0.05$) in the drug-treated versus the vehicle treated mice (see Figure 4).

EXAMPLE 36

Measurement of the Inhibition of LPS-induced TNF-alpha Production in RAW264.7 Cells Treated With a Compound of the Present Invention

[000221] The ability of **1** to inhibit LPS-induced TNF-alpha production was assessed in the mouse macrophage cell line RAW264.7. The RAW cells were preincubated with either 1 μ M dexamethasone (Dex) or 10, 30 or 100 μ M **1** for 1 hour at 37°C in RPMI-1640 containing 10% FBS. After 1 hour LPS (0.1 μ g/ml) was added and cells were incubated an additional 6 hours. Cell supernatant was then collected, aliquoted and frozen at -70°C, and an aliquot used to determine the concentration of TNF-alpha by ELISA. As shown in Figure 5, treatment with **1** significantly inhibited the LPS-induced production of TNF-alpha. The inhibitory effect approached that seen with dexamethasone.

EXAMPLE 37

Measurement of the Inhibition of LPS-induced IL-1 Beta Production in RAW264.7 Cells Treated With a Compound of the Present Invention

[000222] The ability of **1** to inhibit LPS-induced IL-1 beta production was also examined in RAW264.7 cells. The RAW cells were preincubated with either 1 μ M dexamethasone (Dex) or 10, 30 or 100 μ M **1** for 1 hour at 37°C in RPMI-1640 containing 10% FBS. After 1 hour LPS (0.1 μ g/ml) was added and cells were incubated an additional 6 hours. Cell supernatant was then collected, aliquoted and

frozen at -70°C, and an aliquot used to determine the concentration of IL-1 beta by ELISA. As shown in Figure 6, treatment with **1** significantly inhibited the LPS-induced production of IL-1 beta. The inhibition seen with **1** was of the same approximate magnitude as that seen with dexamethasone.

EXAMPLE 38

Measurement of the Inhibition of LPS-induced IL-6 Production in RAW264.7 Cells Treated With a Compound of the Present Invention

[000223] The ability of **1** to inhibit LPS-induced IL-6 production was also measured in RAW264.7 cells. The RAW cells were preincubated with either 1 µM dexamethasone (Dex) or 10, 30 or 100 µM **1** for 1 hour at 37°C in RPMI-1640 containing 10% FBS. After 1 hour LPS (0.1 µg/ml) was added and cells were incubated an additional 6 hours. Cell supernatant was then collected, aliquoted and frozen at -70°C, and an aliquot used to determine the concentration of IL-6 by ELISA. As shown in Figure 7, treatment with **1** significantly inhibited the LPS-induced production of IL-6. The inhibitory effect was greater than that observed with dexamethasone.

EXAMPLE 39

Measurement of the Inhibition of LPS-induced iNOS and COX-2 Production in RAW264.7 Cells Treated With a Compound of the Present Invention

[000224] The ability of **1** to inhibit LPS-induced production of iNOS and COX-2 was also measured in RAW264.7 cells. The RAW cells were preincubated with either 1 µM dexamethasone (Dex) or 10, 30 or 100 µM **1** (Test Cpd) or other reference compound (Ref Cpd A or Ref Cpd B) for 1 hour at 37°C in RPMI-1640 containing 10% FBS. After 1 hour LPS (0.1 µg/ml) was added and cells were incubated an additional 6 hours. Cells receiving no drug treatment, incubated with or without LPS, served as controls. Cells were lysed and 25 µg/well of total protein was electrophoresed on 4-20% Tris-glycine SDS gels. Proteins were transferred to nitrocellulose membrane, and the resulting blot was probed with antibody to iNOS, then stripped and reprobed with antibody to COX-2, and then stripped and reprobed with antibody to COX-1. As shown in Figure 8, treatment with **1** exhibited dose-

dependent inhibition of LPS-induced iNOS production. In addition, treatment with 1 selectively inhibited production of COX-2 but not COX-1 in LPS-stimulated cells.

EXAMPLE 40

Inhibition of LPS-induced TNF-alpha Release by Human Monocytes With Compounds of the Present Invention

[000225] Frozen human elutriated monocytes (Advanced Biotechnologies Incorporated) were thawed and each 1-ml vial mixed with ~12 ml of 10% FBS complete medium (10% heat-inactivated fetal bovine serum in RPMI 1640 medium supplemented with 100 U/ml penicillin, 100 µg/ml streptomycin and 50 µM 2-mercaptoethanol). Cells were centrifuged at 800 rpm for 10 min at room temperature using a Beckman GS-6 centrifuge with GH-3.8 rotor, and the cell pellets were resuspended in 20% FBS complete medium (20% heat-inactivated FBS in RPMI 1640 medium supplemented with 100 U/ml penicillin, 100 µg/ml streptomycin and 50 µM 2-mercaptoethanol) and centrifuged again at 800 rpm for 10 min at room temperature. Cell pellets were resuspended in 20% FBS complete medium, and the cell suspensions were pooled and passed through a 70-micron cell strainer to remove any aggregates or clumps. The cell suspension was adjusted to 2.5×10^6 cells/ml in 20% FBS complete medium. Cell suspension (160 µl, 4×10^5 cells) was added into each well of a 96-well tissue-culture treated polystyrene plate and incubated at 37°C for 1 - 2.5 h. Cells were pretreated with vehicle (DMSO) or test compound (0.3, 1.0, 3.0, 10 or 30 µM) in 20% FBS complete medium for 1 h at 37°C. After pretreatment, lipopolysaccharides (LPS) from *Salmonella typhimurium* in 20% FBS complete medium were added to the cells. The final concentrations were 0.1% DMSO and 10 ng/ml LPS in a final volume of 200 µl/well. The cells were incubated for 20 h at 37°C, and then the supernatants were harvested and aliquots of the supernatants frozen at -80°C for subsequent analysis. Cells on the plates were assayed for cell viability using the MTS/PMS assay (Cory AH et al., Cancer Commun 3:207-212, 1991). The concentration of TNF-alpha in the cell supernatants was determined using quantitative sandwich enzyme immunoassay for human TNF-alpha (R&D Systems). The mean percent inhibition of TNF-alpha release relative to vehicle was calculated for each concentration of test compound from multiple

determinations. As shown in Table 2, the compounds of the invention caused significant inhibition of LPS-induced TNF-alpha release by human monocytes.

TABLE 2

Test Compound	Percent Inhibition of TNF-alpha Release				
	0.3 μ M	1.0 μ M	3.0 μ M	10 μ M	30 μ M
49	-	-	14%	47%	54%
31	-	51%	73%	83%	86%
37	-	17%	38%	65%	78%
13	15%	40%	70%	78%	78%
51	-	-	25%	57%	*
56	1%	-	6%	-	54%
66	27%	-	53%	-	84%
67	40%	-	62%	-	89%
44	32%	-	67%	-	91%
71	20%	-	47%	-	65%
69	1%	-	22%	-	50%
58	6%	-	13%	-	53%
59	27%	-	69%	-	80%

* Cell viability < 70%

EXAMPLE 41

Stimulation of Glucose Uptake in 3T3-L1 Adipocytes With Compounds of the Present Invention

[000226] Differentiation of mouse 3T3-L1 adipocytes was carried out using the method of Greenberg AS, et al., J Biol Chem 276:45456-61, 2001. Briefly, 3T3-L1 fibroblasts were differentiated to adipocytes by incubation in DMEM containing 10% FBS, 72 μ g/ml porcine insulin, 0.5 mM 3-isobutylmethylxanthine (IBMX) and 400 ng/ml dexamethasone for 2 x 48 h at 37°C. Differentiated cells were maintained in media containing 10% FBS without insulin, IBMX or dexamethasone until they were used for experiments. The effect of treatment with compounds of the invention on glucose uptake by differentiated adipocytes was assessed essentially according to the method of Tafuri SR, Endocrinology 137:4706-12, 1996. Adipocytes were incubated with vehicle (0.1 % DMSO) or test compound (0.1, 1.0 or 10 μ M) for 48 h in DMEM containing 10% FBS, then washed and incubated in high-glucose, serum-free medium for 3 h at 37°C. Cells were then washed, incubated for 20 min in glucose-free, serum-free medium containing 100 nM insulin, then supplemented with 2.5 μ Ci/ml ¹⁴C-deoxyglucose in 0.1 mM cold deoxyglucose and further incubated for

10 min at room temperature. After washing, cells were lysed with 0.5% SDS and the radioactivity was measured in a scintillation counter to determine the amount of glucose uptake. The mean percent stimulation of glucose uptake relative to vehicle (set at 100%) was calculated for each concentration of test compound from triplicate determinations. As shown in Table 3, the compounds of the invention caused significant stimulation of glucose uptake in differentiated adipocytes.

TABLE 3

Test Compound	Percent Stimulation of Glucose Uptake		
	0.1 μM	1.0 μM	10 μM
31	107%	119%	161%
8	115%	132%	171%
60	93%	120%	229%
61	93%	120%	229%
51	93%	107%	136%
29	106%	124%	120%
40	126%	117%	126%
63	107%	112%	139%
64	108%	113%	127%
56	83%	100%	126%

EXAMPLE 42

Inhibition of PDE4 and PDE3 Activity With a Compound of the Present Invention

[000227] Compound **13** was examined for its ability to inhibit the activity of PDE4 and PDE3 enzymes. PDE4 partially purified from human U-937 promonocytic cells and PDE3 partially purified from human platelets were used. Test compound (1, 10 or 30 μ M) or vehicle (0.1% DMSO) was incubated with 0.2 μ g PDE4 enzyme or 1 μ g PDE3 enzyme and 1 μ M cAMP containing 0.01 μ g [3 H]cAMP in Tris buffer pH 7.5 for 20 min at 30°C. The reaction was terminated by boiling for 2 min and the resulting AMP was converted to adenosine by addition of 10 mg/ml snake venom nucleotidase and further incubation at 30°C for 10 min. Unhydrolyzed cAMP was bound to AGI-X2 resin, and remaining [3 H]adenosine in the aqueous phase was quantitated by scintillation counting. The mean percent inhibition of PDE4 or PDE3 activity was calculated from duplicate determinations (Table 4). Compound **13** exhibited significant inhibition of both PDE4 ($IC_{50} < 1 \mu$ M) and PDE3 ($IC_{50} = 13.6 \mu$ M) enzyme activities.

TABLE 4

Enzyme Assay	Percent Inhibition of Enzyme Activity		
	1 μ M	10 μ M	30 μ M
PDE4	85%	98%	102%
PDE3	20%	52%	55%

EXAMPLE 43

Inhibition of LPS-induced Phosphorylation of p44/42 MAP Kinase With a Compound of the Present Invention

[000228] Compound 13 was examined for its ability to inhibit LPS-induced and LPS/IFN-gamma induced phosphorylation of p44/42 MAP kinase. RAW 264.7 gamma NO(-) cells were seeded at 1×10^6 /well (2 ml per well) in 6-well tissue culture plates one day prior to the experiment. To start the experiment, cells were washed 2X with RPMI 1640 medium, 0.5% FBS, 100 U/ml penicillin, 100 μ g/ml streptomycin, 1mM sodium pyruvate, and then pretreated with vehicle (0.1% DMSO) or test compound (10 or 30 μ M) at 37°C for 1 h. After pretreatment, cells were incubated in RPMI 1640 medium, 10% FBS, 100 U/ml penicillin, 100 μ g/ml streptomycin, 1mM sodium pyruvate, containing 10 ng/ml LPS or LPS (10 ng/ml)/IFN-gamma (10 U/ml) at 37°C for 15 min. Cells were then washed 2X with cold PBS (pH 7.4) and lysed in 20 mM Tris-HCl (pH 7.5), 150 mM NaCl, 1 mM Na₂EDTA, 1 mM EGTA, 1% Triton, 2.5 mM sodium pyrophosphate, 1 mM beta-glycerophosphate, 1 mM Na₃VO₄, 1 μ g/ml leupeptin, 1 mM PMSF on ice for 10 min. Lysed cells were collected and centrifuged at $\sim 20,800 \times g$ for 10 min at 4°C. Supernatants (lysates) were collected, aliquoted, and stored frozen at -80°C until use. Lysates (29 μ g of total proteins per sample) were subjected to SDS-polyacrylamide (4-20%) gel electrophoresis, and the separated proteins were transferred to nitrocellulose membranes. Membranes were blocked with 5% non-fat dry milk, 10 mM Tris-HCl (pH 8.0), 150 mM NaCl, 0.1% Tween®-20 at room temperature for 1 h and then were blotted with mAb against phospho-p44/42 MAP kinase (Thr 202/Tyr 204) at room temperature for 1 h. The membranes were then washed and incubated with a horseradish peroxidase-linked anti-mouse secondary antibody at room temperature for 1 h. The signals were detected using ECL Western blotting detection reagents. The results showed that

compound **13** inhibited LPS-induced phosphorylation of p44/42 MAP kinase at 30 μ M but not 10 μ M, while the compound inhibited LPS/IFN- γ induced phosphorylation of p44/42 in a dose-dependent manner at 30 μ M and 10 μ M (data not shown).

EXAMPLE 44

Inhibition of Anti-CD3/Anti-CD28 Stimulated Lymphocyte Proliferation With a Compound of the Present Invention

[000229] Compound **13** was examined for its ability to inhibit anti-CD3/anti-CD28 stimulated lymphocyte proliferation. Binding of antigen, or antibodies, to CD3/CD28 triggers activation and proliferation of T-lymphocytes, which are key steps involved in mounting an immune response (Abbas, Lichtman and Pober, Cellular and Molecular Immunology, 3rd edition, W.B. Saunders Company, Philadelphia, 1997). Mesenteric lymph nodes were collected from BALB/c mice (female, ~8 weeks old), and the cells were isolated in PBS (pH 7.4) and mixed with culture medium (RPMI 1640 medium, 10% FBS, 100 U/ml penicillin, 100 μ g/ml streptomycin, 50 μ M 2-mercaptoethanol). The cell suspension was centrifuged at 900 rpm for 10 min at room temperature using a Beckman GS-6 centrifuge with GH-3.8 rotor. After centrifugation, cell pellets were resuspended in culture medium and centrifuged again at 900 rpm for 10 min at room temperature. Cell pellets were resuspended in culture medium and cells were counted. 2×10^5 lymph node cells per well were added into a 96-well cell culture plate. For the treatment (n=4), vehicle (DMSO) or test compound was added into cells. Cells were treated with purified hamster anti-mouse CD3 ϵ (2 μ g/ml) and anti-mouse CD28 (0.2 μ g/ml) monoclonal antibodies or with culture medium. The final concentrations were 0.1 % DMSO and 10 μ M test compound in a final volume of 200 μ l per well. Cells were incubated at 37°C for 67 h, and then cells on plates were centrifuged at 900 rpm for 10 min at room temperature using a Beckman GS-6 Centrifuge with GH-3.8 rotor. 150 μ l of supernatant from each well was subsequently harvested and frozen at -80°C for further analysis (ELISA). For the cells on the plate, 150 μ l of culture medium was added into each well to replace the harvested supernatants and 40 μ l of MTS/PMS solution was added into each well. After further incubation at 37°C for 140 min, the plate was read at 505 nm in a microplate spectrophotometer. The O.D. values (after subtracting the mean O.D. of blank wells)

were used to compare the proliferation of treated cells. As shown in Table 5, 10 μ M of compound **13** caused about 50% inhibition of the proliferation of mouse mesenteric lymph node cells stimulated by anti-CD3/anti-CD28 monoclonal antibodies. Inhibition of CD3/CD28 mediated T-cell proliferation demonstrates compound **13** is able to block an immunologically-relevant cellular response, probably via interactions with a step in the signal transduction cascade. This indicates that compound **13** has immunomodulatory activity, which may be useful for the treatment of immunoproliferative disorders.

TABLE 5

Treatment	O.D. (Mean $\pm$ SD)
DMSO	0.020 $\pm$ 0.006
DMSO + anti-CD3/anti-CD28 mAbs	1.372 $\pm$ 0.125
Test compound + anti-CD3/anti-CD28 mAbs	0.578 $\pm$ 0.012

EXAMPLE 45

Improvement of Collagen Induced Arthritis in Mice Using a Compound of the Present Invention

[000230] Collagen-induced arthritis (CIA) was induced in 45 DBA/1J mice using immunization with chicken collagen Type II. The induction schedule was as follows: on Day 0, 100 μ g/100 μ l in Complete Freund's Adjuvant (CFA) intradermally; on Day 21, 100 μ g/100 μ l in Incomplete Freund's Adjuvant subcutaneously; on Day 31, 100 μ g/100 μ l in CFA subcutaneously; all given at the base of the tail. On Day 35 animals received lipopolysaccharides (detoxified from E. coli serotype O111:B4; 40 μ g/mL) intraperitoneally. When signs of arthritis appeared, mice were assigned into four treatment groups: vehicle control (0.5% carboxymethylcellulose (CMC)); compound **31** (40 mg/kg suspension in CMC); compound **31** (100 mg/kg in CMC); positive control (dexamethasone; 5 mg/kg). The animals were dosed per oral by gavage, twice daily for 14 days, at a dose volume of 250 μ l per mouse per dose. The study was scored blindly to the different treatment groups. Mice were weighed and arthritis was scored three times a week. Arthritis was scored as a count of affected limbs and digits, evaluated as: erythema and swelling of tarsal, the ankle to the metatarsal joints, up to restriction of movement and deformity of the joints.

Plasma was collected from the animals 4 hours following the final dose, for measurement of circulating drug levels. At termination, animals were euthanized and hind limbs removed for histopathologic examination, hind limbs were collected in formalin. Decalcified tissue was sectioned longitudinally, parallel to the bones, and hematoxylin and eosin stained sections were scored using a standard rheumatoid arthritis scoring system by a veterinary histopathologist who was blinded to the treatment groups. Animals in all groups had moderate arthritis prior to the start of dosing (Day 0) as shown in Figure 9. The vehicle group exhibited an increase in severity over the course of the study with a tendency to plateau from about Day 10. The low dose of compound **31** had no apparent effect on the animals compared with vehicle controls. The high dose prevented the increase in severity, to about the same extent as dexamethasone. Histology showed that the vehicle group and the low-dose compound **31** group had marked chronic inflammation of synovium with pannus formation and destruction of bone and cartilage, while in the dexamethasone group the joints were within normal limits. At high dose of compound **31** there was a reduction in incidence and severity of pannus formation, inflammation cell infiltration and bone/cartilage damage. Thus a dose-dependent effect of compound **31** was observed on both the soft tissue and bone and cartilage, consistent with a disease-modifying activity of the compound in this model.


[000231] It will be evident from the above that the compounds according to the present invention not only lower blood glucose level, triglyceride level and free fatty acid level in diabetic conditions, but also inhibit TNF-alpha, IL-6, IL-1 beta, COX-2 and iNOS production in inflammation, as well as inhibit PDE4 and PDE3 activity, phosphorylation of p44/42 MAP kinase and lymphocyte proliferation. The properties demonstrated above indicate that the compounds of the invention should be useful in the treatment of disorders associated with insulin resistance, hyperglycemia, hyperlipidemia, coronary artery disease and peripheral vascular disease and for the treatment of inflammation, inflammatory diseases, immunological diseases, proliferative diseases and cancer, especially those mediated by cytokines, cyclooxygenase, phosphodiesterase and/or MAP kinase.

EXAMPLE 46

Synthesis of N-{4-[2-(3,5-dimethoxyphenyl)-1-dimethylcarbamoylvinyl]-phenyl}-benzamide (67)

[000232] A mixture of 2-(4-aminophenyl)-3-(3,5-dimethoxyphenyl)-N,N-dimethylacrylamide, **43**, (0.49g, 1.2 mmol) and benzoyl chloride (0.26 g, 1.8 mmol) in anhydrous benzene (18.0 mL) was heated at 90°C for 2 h. Solvent was evaporated and crude product was purified by silica gel chromatography.

[000233] Analysis: ¹HNMR (DMSO-d₆): δ 10.33 (s, 1H), 7.96 (d, J=8.8Hz, 2H), 7.76 (d, J=8.8Hz, 1H), 7.51-7.62 (m, 3H), 7.26 (d, J=9.2Hz, 2H), 6.55 (s, 1H), 6.35 (t, J=2.0Hz, 1H), 6.31 (d, J=2.0Hz, 2H), 3.58 (s, 6H), 3.03 (brs, 3H), 2.91 (brs, 3H).




EXAMPLE 47

Synthesis of 3-{4-[4-(2-benzo[1,3]dioxol-5-yl)-1-dimethylcarbamoylvinyl]-phenoxy]-phenyl}-propionic acid ethyl ester (69)

[000234] A mixture of 3-{4-[4-(2-ethoxycarbonylethyl)-phenoxy]-phenyl}-2-oxopropionic acid, **24** (1.0 g, 3.0 mmol), piperonal (0.67 g, 0.45 mmol), triethylamine (5.12 mL) and acetic anhydride (5 mL) was heated at 80°C for 3 h. Reaction mixture was poured in water (50 mL). Solid separated was filtered and boiled in toluene, cooled and filtered. Crude solid was purified by silica gel chromatography to yield **68**, 0.35g (yield, 25.0%).

[000235] A mixture of 4-benzo[1,3]dioxol-5-yl-3-{4-[4-(2-ethoxycarbonylethyl)-phenoxy]-phenyl}-2-oxobut-3-enoic acid, **68**, (0.08 g, 0.17 mmol), benzotriazol-1-yloxytris-(dimethylamino)-phosphonium hexafluorophosphate (BOP, 0.09 g, 0.21 mmol), triethylamine (36 μL, 0.25 mmol), dimethylamine in THF (2M, 0.25 mL, 0.5

mmol) in DMF (2.0 mL) was stirred for 10 min at room temperature. Reaction mixture was poured in water (20 mL). Solid separated was filtered and boiled in toluene, cooled and filtered. Crude solid was purified by silica gel chromatography to yield **69**. Analysis: ^1H NMR (DMSO- d_6): δ 7.24 (d, $J=8.8\text{Hz}$, 4H), 6.95 (overlapped d, $J=8.8\text{Hz}$, 4H), 6.80 (d, $J=8.0\text{Hz}$, 1H), 6.68 (d, $J=8.0\text{Hz}$, 1H), 6.54 (s, 1H), 6.51 (s, 1H), 5.96 (s, 2H), 4.05 (q, $J=4.0\text{Hz}$, 2H), 3.05 (brs, 3H), 2.85 (brs, 3H), 2.80 (t, $J=6.0\text{Hz}$, 2H), 2.60 (t, $J=6.0\text{Hz}$, 2H) and 1.15 (t, $J=4.0\text{Hz}$, 3H).




EXAMPLE 48

Synthesis of 2-[4-[4-(1-dimethylcarbamoyl-2-pyridin-3-ylvinyl)-phenoxy]-benzyl]-malonamide (**71**)

[000236] To a solution of 2-[4-[4-(1-dimethylcarbamoyl-2-pyridin-3-ylvinyl)-phenoxy]-benzyl]-malonic acid dimethyl ester, **70** (0.49 g, 1.0 mmol), in DMF (5 mL), ammonium hydroxide (28% in water, 12 mL) was added and stirred overnight at room temperature. Reaction mixture was poured in water (30 mL) and extracted with chloroform (5 x 25 mL). The organic layer was dried on anhydrous magnesium sulfate and evaporated. The crude product was purified by silica gel chromatography to yield **71**, 0.23g (yield, 24.5%).


[000237] Analysis: ^1H NMR ($\text{CDCl}_3 + \text{CD}_3\text{OD}$): δ 8.32 (m, 2H), 7.40 (m, 1H), 7.18 (overlapped d, $J=8.0\text{Hz}$, 2H), 7.20 (overlapped d, $J=8.0\text{Hz}$, 2H), 7.12 (m, 1H), 6.92 (d, $J=8.0\text{Hz}$, 2H), 6.84 (d, $J=8.0\text{Hz}$, 2H), 6.60 (s, 1H), 3.22 (d, $J=12.0\text{Hz}$), 3.12 (brd, $J=12.0\text{Hz}$), 2.98 (brs, 3H), 2.96 (brs, 3H).










[000238] It will be appreciated that various modifications may be made in the invention as described above and as defined in the following claims wherein:

What is claimed is:

1. A compound having a structure selected from Formulas I-XIII:





wherein the stereocenters marked with an asterisk (*) are R- or S-; the bond represented by a dashed line plus a solid line is a double bond or a single bond; and wherein

R₁, R₂, R₃, R₄, R₅, R₆ and R₇ are each independently selected from the group consisting of

H; optionally substituted C₁-C₂₀ linear or branched alkyl, chloroalkyl or fluoroalkyl; optionally substituted C₂-C₂₀ linear or branched alkenyl; optionally substituted C₆-C₂₀ aryl, linear or branched alkylaryl, linear or branched alkenylaryl; COOR where R is H, optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl or optionally substituted C₆-C₁₀ aryl, sodium, potassium, calcium, magnesium, ammonium, tromethamine; CONR'R'', where R' and R'' are independently H, optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl or optionally substituted C₆-C₁₀ aryl or where NR'R'' represents a cyclic moiety selected from morpholine, piperidine, piperazine; optionally substituted C₁-C₆ amidoalkyl; NH₂; C₁-C₂₀ alkylamino, bis(alkylamino), cycloalkylamino or cyclic amino; OH; optionally substituted C₁-C₂₀ alkoxy, optionally substituted C₁-C₂₀ alkanoyl; optionally substituted C₁-C₂₀ acyloxy; halo; optionally substituted C₁-C₂₀ alkylcarboxylamino; cyano; nitro; SO₂NR'''R'''' where R''' and R'''' are independently H, C₁-C₂₀ alkyl or aryl; SO₂R''' where R''' is H, C₁-C₂₀ alkyl or aryl; SO₃R''' where R''' is H, C₁-C₂₀ alkyl or aryl; and C₄-C₈ heterocycles selected from tetrazolyl, imidazolyl, pyrrolyl, pyridyl and indolyl;

R₈ and R₉ are each independently selected from the group consisting of

H; optionally substituted C₁-C₂₀ linear or branched alkyl; optionally substituted C₂-C₂₀ linear or branched alkenyl; optionally substituted C₆-C₁₀ aryl or heteroaryl; COOR where R is H, optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl or optionally substituted C₆-C₁₀ aryl, sodium, potassium, calcium, magnesium, ammonium, or tromethamine; CONR'R'', where R' and R'' are independently H, alkoxy, optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl, optionally substituted C₃-C₁₀ cycloalkyl or cycloalkenyl or optionally substituted C₆-C₁₀ aryl or heteroaryl, or where NR'R'' represents a cyclic moiety selected from morpholine, piperidine, hydroxypiperidine, imidazole, piperazine, or methylpiperazine; NH₂; C₁-C₂₀ alkylamino, bis(alkylamino), cycloalkylamino or cyclic amino; OH; C₁-C₂₀ alkoxy; C₁-C₂₀ alkanoyl; C₁-C₂₀ acyloxy; halo; C₁-C₂₀ alkylcarboxylamino;

cyano; nitro; $\text{SO}_2\text{NR}'''\text{R}''''$ where R''' and R'''' are independently H, $\text{C}_1\text{-C}_{20}$ alkyl or aryl; $\text{SO}_2\text{R}'''$ where R''' is H, $\text{C}_1\text{-C}_{20}$ alkyl or aryl; $\text{SO}_3\text{R}'''$ where R''' is H, $\text{C}_1\text{-C}_{20}$ alkyl or aryl; and tetrazolyl;

R_{10} and R_{11} are each independently selected from the group consisting of

H; optionally substituted $\text{C}_1\text{-C}_{20}$ linear or branched alkyl; optionally substituted $\text{C}_2\text{-C}_{20}$ linear or branched alkenyl; optionally substituted $\text{C}_6\text{-C}_{10}$ aryl or heteroaryl; COOR where R is H, optionally substituted $\text{C}_1\text{-C}_{20}$ alkyl, optionally substituted $\text{C}_2\text{-C}_{20}$ alkenyl or optionally substituted $\text{C}_6\text{-C}_{10}$ aryl, sodium, potassium, calcium, magnesium, ammonium, or tromethamine; $\text{CONR}'\text{R}''$, where R' and R'' are independently H, optionally substituted $\text{C}_1\text{-C}_{20}$ alkyl, optionally substituted $\text{C}_2\text{-C}_{20}$ alkenyl or optionally substituted $\text{C}_6\text{-C}_{10}$ aryl or where $\text{NR}'\text{R}''$ represents a cyclic moiety selected from morpholine, piperidine, or piperazine; NH_2 ; $\text{C}_1\text{-C}_{20}$ alkylamino, bis(alkylamino), cycloalkylamino or cyclic amino; OH; $\text{C}_1\text{-C}_{20}$ alkoxy; $\text{C}_1\text{-C}_{20}$ alkanoyl; $\text{C}_1\text{-C}_{20}$ acyloxy; halo; $\text{C}_1\text{-C}_{20}$ alkylcarboxylamino; cyano; nitro; $\text{SO}_2\text{NR}'''\text{R}''''$ where R''' and R'''' are independently H, $\text{C}_1\text{-C}_{20}$ alkyl or aryl; $\text{SO}_2\text{R}'''$ where R''' is H, $\text{C}_1\text{-C}_{20}$ alkyl or aryl; $\text{SO}_3\text{R}'''$ where R''' is H, $\text{C}_1\text{-C}_{20}$ alkyl or aryl; and tetrazolyl;

R_{12} , R_{13} , R_{18} , R_{19} and R_{20} are each independently selected from the group consisting of

H; optionally substituted $\text{C}_1\text{-C}_{20}$ linear or branched alkyl; optionally substituted $\text{C}_2\text{-C}_{20}$ linear or branched alkenyl; optionally substituted $\text{C}_6\text{-C}_{10}$ aryl or heteroaryl; COOR where R is optionally substituted $\text{C}_1\text{-C}_{20}$ alkyl, optionally substituted $\text{C}_2\text{-C}_{20}$ alkenyl or optionally substituted $\text{C}_6\text{-C}_{10}$ aryl, sodium, potassium, calcium, magnesium, ammonium, or tromethamine; $\text{CONR}'\text{R}''$, where R' and R'' are independently H, optionally substituted $\text{C}_1\text{-C}_{20}$ alkyl, optionally substituted $\text{C}_2\text{-C}_{20}$ alkenyl or optionally substituted $\text{C}_6\text{-C}_{10}$ aryl or where $\text{NR}'\text{R}''$ represents a cyclic moiety selected from morpholine, piperidine and piperazine; $\text{C}_1\text{-C}_{20}$ alkanoyl; $\text{C}_1\text{-C}_{20}$ alkylamido; $\text{C}_6\text{-C}_{20}$ aroyl or heteroaroyl; $\text{SO}_2\text{R}'''$ where R''' is H, $\text{C}_1\text{-C}_{20}$ alkyl or aryl; morpholinocarbonylmethyl; piperazinocarbonylmethyl; and piperidinocarbonylmethyl;

R₁₂ and R₁₃ may be absent, or R₁₂ and R₁₃ together may be an optionally substituted heterocyclic ring selected from morpholine, piperidine, piperazine, and N-methylpiperidine;

R₁₄ is selected from the group consisting of

H; optionally substituted C₁-C₂₀ linear or branched alkyl including chloroalkyl and fluoroalkyl; optionally substituted C₂-C₂₀ linear or branched alkenyl; optionally substituted C₆-C₁₀ aryl or heteroaryl; COOR where R is H, optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl or optionally substituted C₆-C₁₀ aryl, sodium, potassium, calcium, magnesium, ammonium, or tromethamine; CONR'R'', where R' and R'' are independently H, optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl or optionally substituted C₆-C₁₀ aryl or where NR'R'' represents a cyclic moiety selected from morpholine, piperidine and piperazine; cyano; and tetrazolyl;

R₁₅, R₁₆ and R₁₇ are each independently selected from the group consisting of

H; optionally substituted C₁-C₂₀ linear or branched alkyl including chloroalkyl and fluoroalkyl; optionally substituted C₂-C₂₀ linear or branched alkenyl; optionally substituted C₆-C₁₀ aryl or heteroaryl; COOR where R is H, optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl or optionally substituted C₆-C₁₀ aryl, sodium, potassium, calcium, magnesium, ammonium, and tromethamine; CONR'R'', where R' and R'' are independently H, optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl or optionally substituted C₆-C₁₀ aryl or where NR'R'' represents a cyclic moiety selected from morpholine, piperidine, and piperazine; NH₂; C₁-C₂₀ alkylamino, bis(alkylamino), cycloalkylamino or cyclic amino; OH; C₁-C₂₀ alkoxy; C₁-C₂₀ alkanoyl; C₁-C₂₀ acyloxy; halo; C₁-C₂₀ alkylcarboxylamino; cyano; nitro; SO₂NR'''R'''' where R''' and R'''' are independently H, C₁-C₂₀ alkyl or aryl; SO₂R''' where R''' is H, C₁-C₂₀ alkyl or aryl; SO₃R''' where R''' is H, C₁-C₂₀ alkyl or aryl; and tetrazolyl;

X is independently selected from

O; N; S; S=O; SO₂; or NR''''', where R''''' may be H or optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl, optionally substituted C₁-

C₂₀ acyl, optionally substituted C₁-C₂₀ acyloxy and optionally substituted C₁-C₂₀ alkoxycarbonyl;

Y is independently selected from O, S or NH;

Z is selected from the group consisting of

OR_a, wherein R_a is selected from the group consisting of H; optionally substituted C₁-C₂₀ linear or branched alkyl, chloroalkyl or fluoroalkyl, optionally substituted C₂-C₂₀ linear or branched alkenyl; optionally substituted C₆-C₁₀ aryl or heteroaryl; optionally substituted C₆-C₂₀ aroyl or heteroaroyl; optionally substituted C₁-C₂₀ alkanoyl; and SO₂R''' where R''' is H, C₁-C₂₀ alkyl or aryl;

NR_bR_c, wherein R_b and R_c are independently selected from the group consisting of H; optionally substituted C₁-C₂₀ linear or branched alkyl, chloroalkyl or fluoroalkyl; optionally substituted C₂-C₂₀ linear or branched alkenyl; optionally substituted C₆-C₁₀ aryl or heteroaryl; optionally substituted C₃-C₁₀ cycloalkyl or cycloalkenyl; COOZ₁ where Z₁ is optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl or optionally substituted C₆-C₁₀ aryl; optionally substituted C₆-C₂₀ aroyl or heteroaroyl; optionally substituted C₁-C₂₀ alkanoyl; and SO₂R''' where R''' is H, C₁-C₂₀ alkyl or aryl; and wherein R_b and R_c together may be joined to form a 3-6 membered ring selected from aziridine, morpholine, piperidine and piperazine; and

CR_dR_eR_f, wherein R_d, R_e and R_f are each independently selected from the group consisting of H; optionally substituted C₁-C₂₀ linear or branched alkyl, chloroalkyl or fluoroalkyl, optionally substituted C₂-C₂₀ linear or branched alkenyl; optionally substituted C₆-C₁₀ aryl or heteroaryl; optionally substituted C₃-C₁₀ cycloalkyl or cycloalkenyl; COOR where R is H, optionally substituted C₁-C₂₀ alkyl, optionally substituted C₂-C₂₀ alkenyl or optionally substituted C₆-C₁₀ aryl, sodium, potassium, calcium, magnesium, ammonium, or tromethamine; NH₂; C₁-C₂₀ alkylamino, bis(alkylamino); cycloalkylamino or cyclic amino; OH; optionally substituted C₁-C₂₀ alkoxy, trifluoromethoxy; optionally substituted C₁-C₂₀ alkanoyl; optionally substituted C₁-C₂₀ acyloxy; optionally substituted C₆-C₂₀ aroyl or heteroaroyl; halo; cyano; nitro; optionally substituted C₁-C₂₀ alkylcarboxylamino; SO₂NR'''R''' where R''' and R''' are independently H, C₁-C₂₀ alkyl or aryl; SO₂R''' where R''' is H, C₁-C₂₀ alkyl or aryl; and SO₃R''' where R''' is H, C₁-C₂₀ alkyl or aryl; or

the grouping $-C(=Y)Z$ may represent hydrogen or R_{12} or may be absent;

Q is selected from the group consisting of

OR_a where R_a is selected from the group consisting of H; optionally substituted C_1-C_{20} linear or branched alkyl, chloroalkyl or fluoroalkyl; optionally substituted C_2-C_{20} linear or branched alkenyl; optionally substituted C_6-C_{10} aryl or heteroaryl; optionally substituted C_6-C_{20} aroyl or heteroaroyl; optionally substituted C_1-C_{20} alkanoyl; and SO_2R''' where R''' is H, C_1-C_{20} alkyl or aryl; and

NR_bR_c where R_b and R_c are independently selected from the group consisting of H; optionally substituted C_1-C_{20} linear or branched alkyl, chloroalkyl or fluoroalkyl; optionally substituted C_2-C_{20} linear or branched alkenyl; optionally substituted C_6-C_{10} aryl or heteroaryl; optionally substituted C_3-C_{10} cycloalkyl or cycloalkenyl; $COOZ_1$ where Z_1 is optionally substituted C_1-C_{20} alkyl, optionally substituted C_2-C_{20} alkenyl or optionally substituted C_6-C_{10} aryl; optionally substituted C_6-C_{20} aroyl or heteroaroyl; optionally substituted C_1-C_{20} alkanoyl; and SO_2R''' where R''' is H, C_1-C_{20} alkyl or aryl; and wherein R_b and R_c together may be joined to form a 3-6 membered ring such as aziridine, morpholine, piperidine, piperazine and the like; and

SR_g , SOR_g or SO_2R_g where R_g is selected from the group consisting of H; optionally substituted C_1-C_{20} linear or branched alkyl, chloroalkyl or fluoroalkyl; optionally substituted C_2-C_{20} linear or branched alkenyl; optionally substituted C_1-C_{20} acyl; optionally substituted C_1-C_{20} alkoxycarbonyl; C_2-C_{20} alkoxy; optionally substituted C_6-C_{10} aryl or heteroaryl; and optionally substituted C_6-C_{10} aroyl or heteroaroyl.

Group A is optionally substituted C_2-C_{20} linear or branched alkenyl; optionally substituted C_6-C_{20} aryl, linear or branched alkylaryl, linear or branched alkenylaryl; optionally substituted heteroaryl selected from pyridine, indole, morpholine, piperidine, tetrazolyl and piperazine; COR where R is optionally substituted C_1-C_{20} linear or branched alkyl; optionally substituted C_2-C_{20} linear or branched alkenyl; optionally substituted C_6-C_{20} aryl, linear or branched alkylaryl, linear or branched alkenylaryl;

optionally substituted heteroaryl selected from pyridine, indole, morpholine, piperidine, piperazine, and tetrazolyl;

Group B is OH, C₁-C₂₀ alkoxy; SO₂R where R may be H or linear or branched C₁-C₂₀ alkyl;

Group Het represents a heterocyclic ring selected from pyridyl, indolyl, tetrazolyl, imidazolyl, morphonyl, piperidinyl, piperazinyl and thiophenyl.

2. The compound of Claim 1 wherein the bond represented by a dashed line plus a solid line is a double bond and is in the E or Z configuration.
3. The compound of Claim 1 wherein the bond represented by a dashed line plus a solid line is a single bond wherein the resulting stereocenters have the R- or S- configuration.
4. The compound of Claim 1 wherein at least one aromatic ring possesses more than one adjacent substituent, and wherein the substituents are joined to form a ring.
5. The compound of Claim 1 wherein R₈ and R₉ together are joined to form a C₄-C₈ heterocyclic ring.
6. The compound of Claim 1 wherein R₁₀ and R₁₁ together are joined to form a C₄-C₈ heterocyclic ring.
7. The compound of Claim 1 wherein R_d and R_e together are joined to form a 3-6 membered ring and wherein the resulting stereocenter has an R- or S- configuration.
8. The compound of Claim 1 which is of Formula I.


9. The compound of Claim 8 selected from the group consisting of:
 - 3-(3,5-Dimethoxyphenyl)-2-{4-[4-(3-oxo-3-ureidopropyl)-phenoxy]-phenyl}-acrylic acid methyl ester;
 - 3-(3,5-Dimethoxyphenyl)-2-{4-[4-(3-oxo-3-ureido-propyl)-phenoxy]-phenyl}-acrylic acid;
 - 3-(3,5-Dimethoxyphenyl)-2-{4-[4-(3-ethoxycarbonylamino-3-oxo-propyl)-phenoxy]-phenyl}-acrylic acid methyl ester;
 - 2-{4-[4-(3-Benzoyloxycarbonylamino-3-oxopropyl)-phenoxy]-phenyl}-3-(3,5-dimethoxyphenyl)-acrylic acid methyl ester;
 - 3-(3,5-Dimethoxyphenyl)-2-{4-[4-(3-oxo-3-ureidopropyl)-phenoxy]-phenyl}-propionic acid;
 - 3-(3,5-Dimethoxyphenyl)-2-{4-[4-(3-oxo-3-ureidopropenyl)-phenoxy]-phenyl}-acrylic acid;
 - 3-(3,5-Dimethoxyphenyl)-2-{4-[4-(3-oxo-3-ureidopropyl)-phenoxy]-phenyl}-acrylic acid ethyl ester;
 - 3-(3,5-Dimethoxyphenyl)-*N,N*-dimethyl-2-{4-[4-(3-oxo-3-ureido-propyl)-phenoxy]-phenyl}-acrylamide; and
 - 2-(4-{4-[3-(3-Cyclohexylureido)-3-oxopropyl]-phenoxy}-phenyl)-3-(3,5-dimethoxyphenyl)-acrylic acid.
10. The compound of Claim 1 which is of Formula II.
11. The compound of Claim 10 selected from the group consisting of
 - [3-(4-Phenoxyphenyl)-propionyl]-urea; and
 - {3-[4-(4-Methoxyphenoxy)-phenyl]-acryloyl}-urea.
12. The compound of Claim 1 which is of Formula III.
13. The compound of Claim 12 selected from the group consisting of
 - 2-{4-[4-(3-Acetylureidomethyl)-phenoxy]-phenyl}-3-(3,5-dimethoxy-phenyl)-acrylic acid methyl ester; and
 - 2-{4-[4-(3-Acetylthioureidomethyl)-phenoxy]-phenyl}-3-(3,5-dimethoxy-phenyl)-acrylic acid.

14. The compound of Claim 1 which is of formula IV.
15. The compound of Claim 14 selected from the group consisting of he
1-Acetyl-3-[4-(4-methoxyphenoxy)-benzyl]-urea; and
1-Acetyl-3-[4-(3,4-dimethoxyphenoxy)-benzyl]-urea.
16. A composition comprising the compound of Claim 1 and a
pharmaceutically acceptable carrier.
17. A method of treatment of diabetes comprising the co-administration of the
compound of Claim 1 and an agent selected from the group consisting of:
insulin or an insulin mimetic,
a sulfonylurea or other insulin secretagogue,
a thiazolidinedione,
a fibrate or other PPAR-alpha agonist,
a PPAR-delta agonist,
a biguanide,
a statin or other hydroxymethylglutaryl (HMG) CoA reductase
inhibitor,
an alpha-glucosidase inhibitor,
a bile acid-binding resin,
apoA1,
niacin,
probucol,
and nicotinic acid.
18. A method of treatment of inflammatory or immunological disease
comprising the co-administration of the compound of Claim 1 and an agent
selected from the group consisting of:
a nonsteroidal anti-inflammatory drug (NSAID),
a cyclooxygenase-2 inhibitor,
a corticosteroid or other immunosuppressive agent,
a disease-modifying antirheumatic drug (DMARD),
a TNF-alpha inhibitor,

other cytokine inhibitor,
other immune modulating agent,
and a narcotic agent.


19. A method of treatment of inflammatory or immunological disease comprising the step of administering to a subject in need of such treatment a therapeutically effective amount of the compound of Claim 1.
20. The method of claim 19 wherein the compound is administered in combination with an agent selected from the group consisting of NSAID, cyclooxygenase-2 inhibitor, immunosuppressive agent, DMARD, TNF-alpha inhibitor, cytokine inhibitor, immune modulating agent and a narcotic agent.
21. The method of Claim 20 wherein the agent is corticosteroid or methotrexate.
22. A method of treatment for diabetes comprising the step of administering to a subject suffering from a diabetic condition a therapeutically effective amount of the compound of Claim 1.
23. The method of claim 20 wherein the compound is administered in combination with an agent selected from the group consisting of: insulin, insulin mimetic, insulin secretagogue, PPAR-gamma agonist, PPAR-alpha agonist, a PPAR-delta agonist, biguanide, HMG CoA reductase inhibitor, alpha-glucosidase inhibitor, bile acid-binding resin, apoA1, niacin, probucol and nicotinic acid.
24. The method of Claim 21 wherein the agent is selected from the group consisting of sulfonylurea, thiazolidinedione, fibrate, and statin.
25. The method of claim 22, wherein the agent is sulfonylurea.

26. A method of inhibiting the activity of TNF-alpha, IL-1, IL-6, PDE4, PDE3, p44/42 MAP kinase, iNOS or COX-2 comprising administering to a host in need of such inhibition a therapeutically effective amount of the compound of Claim 1.
27. A method of inhibiting the undesired action of cytokine, phosphodiesterase, MAP kinase or cyclooxygenase comprising administering to a host in need of such treatment a therapeutically effective amount of the compound of Claim 1.
28. A method of treating hyperlipidemia comprising administering to a host in need of such treatment a therapeutically effective amount of the compound of Claim 1.
29. A method of treating coronary heart disease comprising administering to a host in need of such treatment a therapeutically effective amount of the compound of Claim 1.
30. A method of treating cancer or proliferative disease comprising administering to a host in need of such treatment a therapeutically effective amount of the compound of Claim 1.
31. The method of Claim 19 wherein the inflammatory or immunological disease is selected from the group consisting of: rheumatoid arthritis, osteoarthritis, ankylosing spondylitis, psoriasis, psoriatic arthritis, asthma, acute respiratory distress syndrome, chronic obstructive pulmonary disease, and multiple sclerosis.
32. A compound of the formula



wherein the compound is 3-(3,5-Dimethoxyphenyl)-2-{4-[4-(3-oxo-3-ureidopropyl)-phenoxy]-phenyl}-acrylic acid methyl ester.

33. A compound of the formula



wherein the compound is 3-(3,5-Dimethoxyphenyl)-*N,N*-dimethyl-2-{4-[4-(3-oxo-3-ureidopropyl)-phenoxy]-phenyl}-acrylamide.

34. An anti-inflammatory composition comprising an effective amount of a compound selected from the group consisting of:

3-(3,5-Dimethoxyphenyl)-*N,N*-dimethyl-2-{4-[4-(3-oxo-3-ureidopropyl)-phenoxy]-phenyl}-acrylamide (**13**);

2-{4-[4-(2-Carbamoylethyl)-phenoxy]-phenyl}-3-(3,5-dimethoxyphenyl)-N,N-dimethylacrylamide (31);

3-(4-{4-[2-(3,5-Dimethoxyphenyl)-1-dimethylcarbamoylvinyl]-phenoxy}-phenyl)-propionic acid ethyl ester (37);

N-{4-[2-(3,5-Dimethoxyphenyl)-1-dimethylcarbamoylvinyl]-phenyl}-3-hydroxybenzamide (44);

3-(3,5-Dimethoxyphenyl)-2-(4-hydroxyphenyl)-N,N-dimethylacrylamide (49);

[3-(4-{4-[2-(3,5-Dimethoxyphenyl)-1-(piperidine-1-carbonyl)-vinyl]-phenoxy}-phenyl)-propionyl]-urea (51);

2-{4-[4-(3-Acetylamino-3-oxopropyl)-phenoxy]-phenyl}-3-(4-fluorophenyl)-N,N-dimethylacrylamide (56);

2-(4-{4-[2-(3,5-Dimethoxyphenyl)-1-dimethylcarbamoylvinyl]-phenoxy}-benzyl)-malonic acid (58);

2-(4-{4-[2-(3,5-Dimethoxyphenyl)-1-dimethylcarbamoylvinyl]-phenoxy}-benzyl)-malonamide (59);

3-(3,5-Dimethoxyphenyl)-N,N-dimethyl-2-[4-(pyridin-2-yloxy)-phenyl]-acrylamide (66);

N,N-dimethyl-2-{4-[4-(3-oxo-3-ureidopropenyl)-phenoxy]-phenyl}-3-pyridin-3-ylacrylamide (47);

N-{4-[2-(3,5-Dimethoxyphenyl)-1-dimethylcarbamoyl-vinyl]-phenyl}-benzamide (67) ;

2-{4-[4-(1-Dimethylcarbamoyl-2-pyridin-3-yl-vinyl)-phenoxy]-benzyl}-malonamide (71);

3-{4-[4-(2-Benzo[1,3]dioxol-5-yl-1-dimethylcarbamoyl-vinyl)-phenoxy]-phenyl}-propionic acid ethyl ester (69);

3-Benzo[1,3]dioxol-5-yl-2-{4-[4-(2-carbamoylethyl)-phenoxy]-phenyl}-N,N-dimethyl-acrylamide (72);

N,N-Dimethyl-2-{4-[4-(3-oxo-3-ureidopropyl)-phenoxy]-phenyl}-3-pyridin-3-yl-acrylamide (73)

35. An antidiabetic composition comprising an effective amount of a compound selected from the group consisting of:
- 3-(3,5-Dimethoxyphenyl)-2-{4-[4-(3-ethoxycarbonylamino-3-oxo-propyl)-phenoxy]-phenyl}-acrylic acid methyl ester (8);

(4-{4-[2-(3,5-Dimethoxyphenyl)-1-dimethylcarbamoyl-vinyl]-phenoxy}-benzyl)-
carbamic acid methyl ester (**29**);

2-{4-[4-(2-Carbamoylethyl)-phenoxy]-phenyl}-3-(3,5-dimethoxyphenyl)-N,N-
dimethylacrylamide (**31**);

3-(3,5-Dimethoxyphenyl)-N,N-dimethyl-2-{4-[4-(3-morpholin-4-yl-3-oxopropyl)-
phenoxy]-phenyl}-acrylamide (**40**);

[3-(4-{4-[2-(3,5-Dimethoxyphenyl)-1-(piperidine-1-carbonyl)-vinyl]-phenoxy}-
phenyl)-propionyl]-urea (**51**);

2-{4-[4-(3-Acetyl-amino-3-oxopropyl)-phenoxy]-phenyl}-3-(4-fluorophenyl)-N,N-
dimethylacrylamide (**56**);

3-(3,5-Dimethoxyphenyl)-2-{4-[4-(3-oxo-3-ureidopropyl)- phenoxy]-phenyl}-N-
pyridin-4-ylacrylamide (**60**);

N-(4-Chlorophenyl)-3-(3,5-dimethoxyphenyl)-2-{4-[4-(3-oxo-3-ureidopropyl)-
phenoxy]-phenyl}-acrylamide (**61**);

3-(3,5-Dimethoxyphenyl)-N,N-dimethyl-2-(4-{4-[2-(2-morpholin-4-yl-2-
oxoethylcarbamoyl)-ethyl]-phenoxy}-phenyl)-acrylamide (**63**);

3-(3,5-Dimethoxyphenyl)-N,N-dimethyl-2-(4-{4-[3-(4-methylpiperazin-1-yl)-3-
oxopropyl]-phenoxy}-phenyl)-acrylamide (**64**).

36. 2-{4-[4-(2-Carbamoylethyl)-phenoxy]-phenyl}-3-(3,5-dimethoxyphenyl)-N,N-
dimethylacrylamide (**31**).