

RAD51 INHIBITORS

RELATED APPLICATION

This application claims priority to, and the benefit of U.S. Provisional Application No. 62/816,998, filed on March 12, 2019, the entire contents of which are incorporated by reference.


BACKGROUND

RAD51 is a member of the RAD51 family which promotes the repair of DNA double strand breaks (DSB). RAD51 protein is highly conserved in most eukaryotes, from yeast to humans. The human RAD51 is a 339-amino acid protein that plays a major role in DNA replication and repair by homologous recombination (HR). RAD51 catalyzes strand transfer between a broken sequence and undamaged homologous template to allow re-synthesis of the damaged region.

Studies have demonstrated sensitization to certain DNA damaging therapies associated with defects in proteins that promote HR DNA repair. This sensitization is particularly dramatic for DNA cross-linking chemotherapeutic drugs and ionizing radiation. It has been shown that HR can be partially inhibited in order to sensitize cells to DNA damaging therapies. For example, inhibition of XRCC3 (a RAD51 paralog protein) using a synthetic peptide sensitized Chinese Hamster Ovary (CHO) cells to cisplatin and inhibited the formation of sub-nuclear RAD51 foci in response to DNA damage. Researchers have inhibited the expression of the RAD51 protein itself or blocked its function by over-expressing a dominant negative BRC peptide fragment derived from BRCA2. In view of the connection between increased sensitivity to DNA damaging therapies and defects in HR DNA repair-related proteins, there is a need for compounds that inhibit RAD51. The present application addresses the need.

SUMMARY

The present application relates to a compound of Formula I:



or a pharmaceutically acceptable salt or solvate thereof, wherein the definition of each variable is provided herein below.

The present application also relates to a pharmaceutical composition comprising a compound disclosed herein or a pharmaceutically acceptable salt or solvate thereof and a
5 pharmaceutically acceptable carrier or diluent.

The present application further relates to a method of treating a disease or disorder in which RAD51 plays a role, for example, cancer, an autoimmune disease, an immune deficiency, or a neurodegenerative disease. The method comprises administering to a subject in need thereof an effective amount of a compound disclosed herein or a pharmaceutically
10 acceptable salt or solvate thereof or a pharmaceutical composition disclosed herein.

The present application further relates to use of a compound disclosed herein, or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition disclosed herein, in the manufacture of a medicament for the treatment of a disease or disorder in which RAD51 plays a role, for example, a cancer, an autoimmune disease, an
15 immune deficiency, or a neurodegenerative disease.

The present application further relates to a compound disclosed herein, or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition disclosed herein, for use in treating a disease or disorder in which RAD51 plays a role, for example, a cancer, an autoimmune disease, an immune deficiency, or a neurodegenerative
20 disease.

The present application further relates to use of a compound disclosed herein, or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition disclosed herein, in treating a disease or disorder in which RAD51 plays a role, for example, a cancer, an autoimmune disease, an immune deficiency, or a neurodegenerative disease.

25 The present application provides compounds and compositions with an improved efficacy and safety profile relative to known RAD51 inhibitors. The present application also provides agents with novel mechanisms of action toward RAD51 in the treatment of various types of diseases. Ultimately the present application provides a novel therapeutic strategy for the treatment of diseases and disorders associated with RAD51.


30 The details of the application are set forth in the accompanying description below. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present application, illustrative methods and materials are now described. Other features, objects, and advantages of the application will be apparent from the

description and from the claims. In the specification and the appended claims, the singular forms also include the plural unless the context clearly dictates otherwise. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this application belongs. All patents and publications cited in this specification are incorporated herein by reference in their entireties.

DETAILED DESCRIPTION

Compounds of the Application

In one aspect, the present application relates to a compound of Formula I:



or a pharmaceutically acceptable salt or solvate thereof, wherein:

the thiazolyl ring is optionally substituted with F or Cl;

the ring Cy is C₃-C₇ cycloalkyl, bridged C₆-C₁₂ cycloalkyl, or saturated heterocyclyl comprising one or two 3- to 7-membered rings and 1-3 heteroatoms selected from N, O, and S, wherein the cycloalkyl or heterocyclyl moiety is optionally substituted with one or more groups selected from halogen, OH, CN, NH₂, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy, and C₁-C₄ haloalkoxy;

X₁ is NR₈ or O, or, when X₁ is bonded to a nitrogen atom in the ring Cy, X₁ is absent;

X₂ is NR₈ or O;

R₁ is H or C₁-C₆ alkyl optionally substituted with halogen, OH, C₁-C₆ alkoxy, or C₆-C₁₀ aryloxy;

R₂ is H or C₁-C₆ alkyl optionally substituted with halogen, OH, C₁-C₆ alkoxy, or C₆-C₁₀ aryloxy;

or R₁ and R₂, together with the nitrogen atom to which they are attached, form a heterocyclyl comprising one or two 3- to 7-membered rings and 1-3 heteroatoms selected from N, O, and S;


R₃ is C₁-C₆ alkyl optionally substituted with one or more groups selected from halogen, OH, and CN, phenyl, CH₂-phenyl, C₃-C₇ cycloalkyl, CH₂-(C₃-C₇) cycloalkyl, heterocyclyl, or CH₂-heterocyclyl, wherein the heterocyclyl comprises one or two 3- to 7-membered rings and 1-3 heteroatoms selected from N, O, and S, wherein the cycloalkyl, phenyl, or heterocyclyl moiety is optionally substituted with one or more groups selected from halogen, OH, CN, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy, and C₁-C₄ haloalkoxy;

A₁, A₂, A₃, and A₄ are each independently N or C(R₄);

each R₄ is independently H, halogen, CN, OH, N(R₆')₂, C₁-C₆ alkoxy, C(=O)N(R₆)₂, C(=O)OR₆, C(=O)R₆, Q-T, C₆-C₁₀ aryl, or heteroaryl comprising one or two 5- or 6-membered rings and 1-3 heteroatoms selected from N, O, and S, wherein the aryl or heteroaryl moiety is optionally substituted with one or more R₉;

each Q is independently C₁-C₄ alkylene or O-(C₁-C₄) alkylene wherein the oxygen

atom is bonded to the ring



each T is independently C₁-C₄ alkoxy, OH, N(R₆)₂, N(R₅)C(=O)R₆, N(R₅)C(=O)OR₆, C(=O)N(R₆)₂, C(=O)OR₆, C(=O)R₆, C₆-C₁₀ aryl, or heteroaryl comprising one or two 5- or 6-membered rings and 1-3 heteroatoms selected from N, O, and S, wherein the aryl or heteroaryl moiety is optionally substituted with one or more R₉;

each R₅ is independently H or C₁-C₄ alkyl;

each R₆' is independently H, C₁-C₆ alkyl optionally substituted with one or more R₇, C₁-C₆ haloalkyl, C₃-C₇ cycloalkyl, heterocyclyl comprising one or two 3- to 7-membered rings and 1-3 heteroatoms selected from N, O, and S, C₆-C₁₀ aryl, or heteroaryl comprising one or two 5- or 6-membered rings and 1-3 heteroatoms selected from N, O, and S, wherein the cycloalkyl, heterocyclyl, aryl, or heteroaryl moiety is optionally substituted with one or more R₉, wherein at least one R₆' is not H;

or two R₆' together with the atoms to which they are attached form a 3- to 10-membered heterocyclyl comprising 1-3 heteroatoms selected from N, O, and S, wherein the heterocyclyl is optionally substituted with one or more R₉;

each R₆ is independently H, C₁-C₆ alkyl optionally substituted with one or more R₇, C₁-C₆ haloalkyl, C₆-C₁₀ aryl, or heteroaryl comprising one or two 5- or 6-membered rings and 1-3 heteroatoms selected from N, O, and S, wherein the aryl or heteroaryl moiety is optionally substituted with one or more R₉;


or two R₆ together with the atoms to which they are attached form a 3- to 10-membered heterocyclyl comprising 1-3 heteroatoms selected from N, O, and S, wherein the heterocyclyl is optionally substituted with one or more R₉;

each R₇ is independently N(R₈)₂, OR₈, C₆-C₁₀ aryl, or heteroaryl comprising one or two 5- or 6-membered rings and 1-3 heteroatoms selected from N, O, and S;


each R₈ is independently H or C₁-C₆ alkyl; and

each R₉ is independently oxo, halogen, OH, CN, NH₂, N(C₁-C₄ alkyl)₂, C₁-C₆ alkyl, N(C₁-C₄ alkyl)₂, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, or C₁-C₆ haloalkoxy, wherein the C₁-C₆ alkyl is optionally substituted with one or more oxo, OH, O(C₁-C₄ alkyl), CN, NH₂, NH(C₁-C₄ alkyl), or N(C₁-C₄ alkyl)₂.

In one aspect, the present application relates to a compound of Formula I:



or a pharmaceutically acceptable salt or solvate thereof, wherein:

the thiazolyl ring  is optionally substituted with F or Cl;

the ring Cy is C₃-C₇ cycloalkyl, bridged C₆-C₁₂ cycloalkyl, or saturated heterocyclyl comprising one or two 3- to 7-membered rings and 1-3 heteroatoms selected from N, O, and S, wherein the cycloalkyl or heterocyclyl moiety is optionally substituted with one or more groups selected from halogen, OH, CN, NH₂, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy, and C₁-C₄ haloalkoxy;

X₁ is NR₈ or O, or, when X₁ is bonded to a nitrogen atom in the ring Cy, X₁ is absent; X₂ is NR₈ or O;

R₁ is H or C₁-C₆ alkyl optionally substituted with halogen, OH, C₁-C₆ alkoxy, or C₆-C₁₀ aryloxy;

R₂ is H or C₁-C₆ alkyl optionally substituted with halogen, OH, C₁-C₆ alkoxy, or C₆-C₁₀ aryloxy;

or R₁ and R₂, together with the nitrogen atom to which they are attached, form a 3-7 membered heterocyclic ring comprising 1 or 2 heteroatoms selected from N, O, and S;


R₃ is C₁-C₆ alkyl optionally substituted with one or more groups selected from halogen, OH, and CN, phenyl, CH₂-phenyl, C₃-C₇ cycloalkyl, CH₂-(C₃-C₇) cycloalkyl, heterocyclyl, or CH₂-heterocyclyl, wherein the heterocyclyl comprises one 3- to 7-membered ring and 1 or 2 heteroatoms selected from N, O, and S, wherein the cycloalkyl, phenyl, or heterocyclyl moiety is optionally substituted with one or more groups selected from halogen, OH, CN, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy, and C₁-C₄ haloalkoxy;

A₁, A₂, A₃, and A₄ are each independently N or C(R₄);

each R₄ is independently H, halogen, CN, OH, N(R₆')₂, C₁-C₄ alkoxy, C(=O)N(R₆)₂, C(=O)OR₆, C(=O)R₆, Q-T, C₆-C₁₀ aryl, or heteroaryl comprising one or two 5- or 6-membered rings and 1-3 heteroatoms selected from N, O, and S, wherein the aryl or heteroaryl moiety is optionally substituted with one or more R₉;

each Q is independently C₁-C₄ alkylene or O-(C₁-C₄) alkylene wherein the oxygen

atom is bonded to the ring



each T is independently C₁-C₄ alkoxy, NH₂, NH(C₁-C₄ alkyl), N(C₁-C₄ alkyl)₂, N(R₅)C(=O)R₆, N(R₅)C(=O)OR₆, C(=O)N(R₆)₂, C(=O)OR₆, or C(=O)R₆;

each R₅ is independently H or C₁-C₄ alkyl;

each R₆' is independently H, C₁-C₆ alkyl optionally substituted with one or more R₇, C₁-C₆ haloalkyl, C₆-C₁₀ aryl, or heteroaryl comprising one or two 5- or 6-membered rings and 1-3 heteroatoms selected from N, O, and S, wherein the aryl or heteroaryl moiety is optionally substituted with one or more R₉, wherein at least one R₆' is not H;


each R₆ is independently H, C₁-C₆ alkyl optionally substituted with one or more R₇, C₁-C₆ haloalkyl, C₆-C₁₀ aryl, or heteroaryl comprising one or two 5- or 6-membered rings and 1-3 heteroatoms selected from N, O, and S, wherein the aryl or heteroaryl moiety is optionally substituted with one or more R₉;

each R₇ is independently N(R₈)₂ or OR₈;

each R₈ is independently H or C₁-C₆ alkyl; and


each R₉ is independently OH, CN, NH₂, N(C₁-C₄ alkyl)₂, C₁-C₆ alkyl optionally substituted with one or more groups selected from OH, CN, NH₂, and N(C₁-C₄ alkyl)₂, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, or C₁-C₆ haloalkoxy.

In some embodiments, the compound of Formula I is of Formula Ia or Ib:





or a pharmaceutically acceptable salt or solvate thereof.

In some embodiments, the compound of Formula I is of Formula Ic:




5 or a pharmaceutically acceptable salt or solvate thereof.

In some embodiments, the compound of Formula I is of Formula Id, Ie, If, or Ig:




or a pharmaceutically acceptable salt or solvate thereof.

10 In some embodiments, the compound of Formula I is of Formula IIa or IIb:




or a pharmaceutically acceptable salt or solvate thereof.

In some embodiments, the compound of Formula I is of Formula IIc:




or a pharmaceutically acceptable salt or solvate thereof.


In some embodiments, the compound of Formula I is of Formula IId, IIe, IIIf, or IIg:




or a pharmaceutically acceptable salt or solvate thereof.

For a compound of any of Formula I, Ia, Ib, Ic, Id, Ie, If, Ig, IIa, IIb, IIc, IId, IIe, IIIf, or IIg, where applicable:

(A1) In some embodiments, the thiazolyl ring  is unsubstituted.

(A2) In some embodiments, the thiazolyl ring  is substituted with F.

(A3) In some embodiments, the thiazolyl ring  is substituted with Cl.

(B1) In some embodiments, the ring Cy is C₃-C₇ cycloalkyl selected from cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and cycloheptyl, each of which is optionally substituted with one or more groups selected from halogen (*e.g.*, F, Cl, Br, or I), OH, CN, NH₂, C₁-C₄ alkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, or *t*-butyl), C₁-C₄ haloalkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, or *t*-

butyl, each of which is substituted with one or more halogen (*e.g.*, F, Cl, Br, or I)), C₁-C₄ alkoxy (*e.g.*, methoxy, ethoxy, propoxy, or butoxy), and C₁-C₄ haloalkoxy (*e.g.*, methoxy, ethoxy, propoxy, or butoxy, each of which is substituted with one or more halogen (*e.g.*, F, Cl, Br, or I)). In some embodiments, the ring Cy is cyclohexyl, optionally substituted as
 5 described herein.

(B2) In some embodiments, the ring Cy is bridged C₆-C₁₂ cycloalkyl selected from bicyclo[2.1.1]hexyl, bicyclo[2.2.1]heptyl, bicyclo[2.2.2]octyl, bicyclo[3.2.1]octyl, bicyclo[4.3.1]decyl, bicyclo[3.3.1]nonyl, bornyl, bornenyl, norbornyl, norbornenyl, 6,6-dimethylbicyclo [3.1.1]heptyl, and adamantyl, each of which is optionally substituted with
 10 one or more groups selected from halogen (*e.g.*, F, Cl, Br, or I), OH, CN, NH₂, C₁-C₄ alkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, or *t*-butyl), C₁-C₄ haloalkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, or *t*-butyl, each of which is substituted with one or more halogen (*e.g.*, F, Cl, Br, or I)), C₁-C₄ alkoxy (*e.g.*, methoxy, ethoxy, propoxy, or butoxy), and C₁-C₄ haloalkoxy (*e.g.*, methoxy, ethoxy, propoxy, or
 15 butoxy, each of which is substituted with one or more halogen (*e.g.*, F, Cl, Br, or I)).

(B3) In some embodiments, the ring Cy is saturated heterocyclyl selected from aziridinyl, oxiranyl, thiiranyl, diaziridinyl, oxaziridinyl, dioxiranyl, azetidiny, oxetanyl, thietanyl, diazetidinyl, dioxetanyl, dithietanyl, pyrrolidinyl, tetrahydrofuranyl, tetrahydrothiophenyl, imidazolidinyl, pyrazolidinyl, oxazolidinyl, isoxazolidinyl,
 20 thiazolidinyl, isothiazolidinyl, dioxolanyl, dithiolanyl, piperidinyl, tetrahydropyranyl, thianyl, piperazinyl, morpholinyl, thiomorpholinyl, dioxanyl, dithianyl, hexahydro-1,3,5-triazinyl, trioxanyl, trithianyl, azepanyl, oxepanyl, thiepanyl, diazepanyl, diazaspiro[4.4]nonyl, diazaspiro[3.5]nonyl, hexahydropyridazinyl, hexahydropyrimidinyl, tetrahydrothiopyranyl, thiomorpholinyl, tropanyl, valerolactanyl, azanorbornyl, quinuclidinyl, isoquinuclidinyl,
 25 azabicyclo[2.2.1]heptanyl, 2-azabicyclo[3.2.1]octanyl, azabicyclo[3.2.1]octanyl, azabicyclo[3.2.2]nonanyl, azabicyclo[3.3.0]nonanyl, azabicyclo [3.3.1]nonanyl, diazabicyclo[2.2.1]heptanyl, diazabicyclo[3.2.1]octanyl, octahydropyrrolo[3,4-b]pyrrolyl, and octahydropyrrolo[3,4-c]pyrrolyl, each of which is optionally substituted with one or more groups selected from halogen (*e.g.*, F, Cl, Br, or I), OH, CN, NH₂, C₁-C₄ alkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, or *t*-butyl), C₁-C₄ haloalkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, or *t*-butyl, each of which is substituted with
 30 one or more halogen (*e.g.*, F, Cl, Br, or I)), C₁-C₄ alkoxy (*e.g.*, methoxy, ethoxy, propoxy, or

butoxy), and C₁-C₄ haloalkoxy (*e.g.*, methoxy, ethoxy, propoxy, or butoxy, each of which is substituted with one or more halogen (*e.g.*, F, Cl, Br, or I)).

(C1) In some embodiments, X₁ is NR₈ and X₂ is NR₈. In a further embodiment, each R₈ is H. In another further embodiment, one R₈ is H, and the other R₈ is C₁-C₆ alkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl, pentyl (straight or branched), or hexyl (straight or branched)). In another further embodiment, each R₈ is independently C₁-C₆ alkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl, pentyl (straight or branched), or hexyl (straight or branched)).

(C2) In some embodiments, X₁ is NR₈ and X₂ is O. In a further embodiment, R₈ is H. In another further embodiment, R₈ is C₁-C₆ alkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl, pentyl (straight or branched), or hexyl (straight or branched)).

(C3) In some embodiments, X₁ is O and X₂ is NR₈. In a further embodiment, R₈ is H. In another further embodiment, R₈ is C₁-C₆ alkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl, pentyl (straight or branched), or hexyl (straight or branched)).

(C4) In some embodiments, X₁ is O and X₂ is O.

(C5) In some embodiments, X₁ is absent and X₂ is NR₈. In a further embodiment, R₈ is H. In another further embodiment, R₈ is C₁-C₆ alkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl, pentyl (straight or branched), or hexyl (straight or branched)).

(C6) In some embodiments, X₁ is absent and X₂ is O.

(D1) In some embodiments, R₁ is H.

(D2) In some embodiments, R₁ is C₁-C₆ alkyl selected from methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl, pentyl (straight or branched), and hexyl (straight or branched), each of which is optionally substituted with one or more groups selected from halogen (*e.g.*, F, Cl, Br, or I), OH, C₁-C₆ alkoxy (*e.g.*, methoxy, ethoxy, propoxy, butoxy, pentoxy, or hexyloxy), and C₆-C₁₀ aryloxy (*e.g.*, phenoxy). In some embodiments, R₁ is C₁-C₄ alkyl selected from methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, and *t*-butyl, each of which is optionally substituted with one or more groups selected from halogen (*e.g.*, F, Cl, Br, or I), OH, C₁-C₆ alkoxy (*e.g.*, methoxy, ethoxy, propoxy, butoxy, pentoxy, or hexyloxy), and C₆-C₁₀ aryloxy (*e.g.*, phenoxy). In some embodiments, R₁ is C₁-C₄ alkyl selected from methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, and *t*-butyl, each of which is optionally substituted with one or more groups selected from halogen (*e.g.*, F, Cl,

Br, or I) and OH. In some embodiments, R₁ is unsubstituted C₁-C₆ alkyl. In some embodiments, R₁ is unsubstituted C₁-C₄ alkyl.

(E1) In some embodiments, R₂ is H.

(E2) In some embodiments, R₂ is C₁-C₆ alkyl selected from methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl, pentyl (straight or branched), and hexyl (straight or branched), each of which is optionally substituted with one or more groups selected from halogen (*e.g.*, F, Cl, Br, or I), OH, C₁-C₆ alkoxy (*e.g.*, methoxy, ethoxy, propoxy, butoxy, pentoxy, or hexyloxy), and C₆-C₁₀ aryloxy (*e.g.*, phenoxy). In some embodiments, R₂ is C₁-C₄ alkyl selected from methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, and *t*-butyl, each of which is optionally substituted with one or more groups selected from halogen (*e.g.*, F, Cl, Br, or I), OH, C₁-C₆ alkoxy (*e.g.*, methoxy, ethoxy, propoxy, butoxy, pentoxy, or hexyloxy), and C₆-C₁₀ aryloxy (*e.g.*, phenoxy). In some embodiments, R₂ is C₁-C₄ alkyl selected from methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, and *t*-butyl, each of which is optionally substituted with one or more groups selected from halogen (*e.g.*, F, Cl, Br, or I) and OH. In some embodiments, R₂ is unsubstituted C₁-C₆ alkyl. In some embodiments, R₂ is unsubstituted C₁-C₄ alkyl.

(DE) In some embodiments, R₁ and R₂, together with the nitrogen atom to which they are attached, form heterocyclic ring selected from aziridinyl, oxiranyl, thiiranyl, diaziridinyl, oxaziridinyl, dioxiranyl, azetidiny, oxetanyl, thietanyl, diazetidinyl, dioxetanyl, dithietanyl, pyrrolidinyl, tetrahydrofuranyl, tetrahydrothiophenyl, imidazolidinyl, pyrazolidinyl, oxazolidinyl, isoxazolidinyl, thiazolidinyl, isothiazolidinyl, dioxolanyl, dithiolanyl, piperidinyl, tetrahydropyranyl, thianyl, piperazinyl, morpholinyl, thiomorpholinyl, dioxanyl, dithianyl, hexahydro-1,3,5-triazinyl, trioxanyl, trithianyl, azepanyl, oxepanyl, thiepanyl, diazepanyl, hexahydropyridazinyl, hexahydropyrimidinyl, tetrahydrothiopyranyl, and thiomorpholinyl. In some embodiments, R₁ and R₂, together with the nitrogen atom to which they are attached, form azetidiny or pyrrolidinyl.

(F1) In some embodiments, R₃ is C₁-C₆ alkyl selected from methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl, pentyl (straight or branched), and hexyl (straight or branched), each of which is optionally substituted with one or more groups selected from halogen (*e.g.*, F, Cl, Br, or I), OH, and CN. In some embodiments, R₃ is C₁-C₄ alkyl selected from methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, and *t*-butyl, each of which is optionally substituted with one or more groups selected from halogen (*e.g.*, F, Cl, Br, or I), OH, and CN. In some embodiments, R₃ is unsubstituted C₁-C₆ alkyl. In some embodiments,

R₃ is unsubstituted C₁-C₄ alkyl. In some embodiments, R₃ is unsubstituted methyl. In some embodiments, R₃ is unsubstituted *i*-propyl. In some embodiments, R₃ is unsubstituted *t*-butyl.

(F2) In some embodiments, R₃ is phenyl or CH₂-phenyl, wherein the phenyl moiety is optionally substituted with one or more groups selected from halogen (*e.g.*, F, Cl, Br, or I), OH, CN, C₁-C₄ alkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, or *t*-butyl), C₁-C₄ haloalkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, or *t*-butyl, each of which is substituted with one or more halogen (*e.g.*, F, Cl, Br, or I)), C₁-C₄ alkoxy (*e.g.*, methoxy, ethoxy, propoxy, or butoxy), and C₁-C₄ haloalkoxy (*e.g.*, methoxy, ethoxy, propoxy, or butoxy, each of which is substituted with one or more halogen (*e.g.*, F, Cl, Br, or I)).

(F3) In some embodiments, R₃ is C₃-C₇ cycloalkyl or CH₂-(C₃-C₇) cycloalkyl, wherein the cycloalkyl moiety is selected from cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and cycloheptyl, each of which is optionally substituted with one or more groups selected from halogen (*e.g.*, F, Cl, Br, or I), OH, CN, C₁-C₄ alkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, or *t*-butyl), C₁-C₄ haloalkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, or *t*-butyl, each of which is substituted with one or more halogen (*e.g.*, F, Cl, Br, or I)), C₁-C₄ alkoxy (*e.g.*, methoxy, ethoxy, propoxy, or butoxy), and C₁-C₄ haloalkoxy (*e.g.*, methoxy, ethoxy, propoxy, or butoxy, each of which is substituted with one or more halogen (*e.g.*, F, Cl, Br, or I)). In some embodiments, R₃ is cyclopropyl or cyclobutyl, or CH₂-cyclopropyl or CH₂-cyclobutyl, each of which is optionally substituted as described herein. In some embodiments, R₃ is cyclopropyl optionally substituted as described herein.

(F4) In some embodiments, R₃ is heterocyclyl or CH₂-heterocyclyl, wherein the heterocyclyl moiety is selected from aziridinyl, oxiranyl, thiiranyl, diaziridinyl, oxaziridinyl, dioxiranyl, azetidiny, oxetanyl, thietanyl, diazetidinyl, dioxetanyl, dithietanyl, pyrrolidinyl, tetrahydrofuranyl, tetrahydrothiophenyl, imidazolidinyl, pyrazolidinyl, oxazolidinyl, isoxazolidinyl, thiazolidinyl, isothiazolidinyl, dioxolanyl, dithiolanyl, piperidinyl, tetrahydropyranyl, thianyl, piperazinyl, morpholinyl, thiomorpholinyl, dioxanyl, dithianyl, hexahydro-1,3,5-triazinyl, trioxanyl, trithianyl, azepanyl, oxepanyl, thiepanyl, diazepanyl, hexahydropyridazinyl, hexahydropyrimidinyl, tetrahydrothiopyranyl, and thiomorpholinyl, each of which is optionally substituted with one or more groups selected from halogen (*e.g.*, F, Cl, Br, or I), OH, CN, C₁-C₄ alkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, or *t*-butyl), C₁-C₄ haloalkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-

butyl, or *t*-butyl, each of which is substituted with one or more halogen (*e.g.*, F, Cl, Br, or I)), C₁-C₄ alkoxy (*e.g.*, methoxy, ethoxy, propoxy, or butoxy), and C₁-C₄ haloalkoxy (*e.g.*, methoxy, ethoxy, propoxy, or butoxy, each of which is substituted with one or more halogen (*e.g.*, F, Cl, Br, or I)). In some embodiments, R₃ is unsubstituted oxetanyl.

5 (G1) In some embodiments, A₁, A₂, A₃, and A₄ are each C(R₄).

(G2) In some embodiments, one, two, or three of A₁, A₂, A₃, and A₄ are N, and the remaining of A₁, A₂, A₃, and A₄ are C(R₄).

(G3) In some embodiments, one of A₁, A₂, A₃, and A₄ is N, and the remaining of A₁, A₂, A₃, and A₄ are C(R₄).

10 (G3a) In some embodiments, A₁ is N, and A₂, A₃, and A₄ are C(R₄).

(G3b) In some embodiments, A₂ is N, and A₁, A₃, and A₄ are C(R₄).

(G3c) In some embodiments, A₃ is N, and A₁, A₂, and A₄ are C(R₄).

(G3d) In some embodiments, A₄ is N, and A₁, A₂, and A₃ are C(R₄).

(G4) In some embodiments, two of A₁, A₂, A₃, and A₄ are N, and the remaining of A₁, A₂, A₃, and A₄ are C(R₄).

(G4a) In some embodiments, A₁ and A₂ are N, and A₃ and A₄ are C(R₄).

(G4b) In some embodiments, A₁ and A₃ are N, and A₂ and A₄ are C(R₄).

(G4c) In some embodiments, A₁ and A₄ are N, and A₂ and A₃ are C(R₄).

(G4d) In some embodiments, A₂ and A₃ are N, and A₁ and A₄ are C(R₄).

20 (G4e) In some embodiments, A₂ and A₄ are N, and A₁ and A₃ are C(R₄).

(G4f) In some embodiments, A₃ and A₄ are N, and A₁ and A₂ are C(R₄).

(G5) In some embodiments, three of A₁, A₂, A₃, and A₄ are N, and the remaining of A₁, A₂, A₃, and A₄ is C(R₄).

(G5a) In some embodiments, A₂, A₃, and A₄ are N, and A₁ is C(R₄).

25 (G5b) In some embodiments, A₁, A₃, and A₄ are N, and A₂ is C(R₄).

(G5c) In some embodiments, A₁, A₂, and A₄ are N, and A₃ is C(R₄).

(G5d) In some embodiments, A₁, A₂, and A₃ are N, and A₄ is C(R₄).

(G6) In some embodiments, A₁, A₂, A₃, and A₄ are each N.

(G7) In some embodiments, at most two of A₁, A₂, A₃, and A₄ are N.

30 (H1) In some embodiments, each R₄ is H.

(H2) In some embodiments, at least one R₄ is halogen, CN, OH, N(R₆')₂, C₁-C₆ alkoxy, C(=O)N(R₆)₂, C(=O)OR₆, C(=O)R₆, Q-T, C₆-C₁₀ aryl, or heteroaryl comprising one

or two 5- or 6-membered rings and 1-3 heteroatoms selected from N, O, and S, wherein the aryl or heteroaryl moiety is optionally substituted with one or more R₉.

(H2a) In some embodiments, at least one R₄ is halogen (*e.g.*, F, Cl, Br, or I), CN, OH, N(R₆')₂, C₁-C₆ alkoxy (*e.g.*, methoxy, ethoxy, propoxy, butoxy, pentoxy, or hexoxy),

5 C(=O)N(R₆)₂, C(=O)OR₆, C(=O)R₆, or Q-T.

(H2b) In some embodiments, at least one R₄ is CN, OH, N(R₆')₂, C₁-C₆ alkoxy (*e.g.*, methoxy, ethoxy, propoxy, butoxy, pentoxy, or hexoxy), C(=O)N(R₆)₂, C(=O)OR₆, C(=O)R₆, or Q-T.

(H2c) In some embodiments, at least one R₄ is C(=O)N(R₆)₂, C(=O)OR₆, C(=O)R₆, or
10 Q-T.

(H2d) In some embodiments, at least one R₄ is Q-T.

(H2e) In some embodiments, at least one R₄ is N(R₆')₂, and one of R₆' is H, and the other R₆' is C₁-C₆ alkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl, pentyl (straight or branched), or hexyl (straight or branched), each of which is
15 optionally substituted with one or more R₇), C₁-C₆ haloalkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl, pentyl (straight or branched), or hexyl (straight or branched), each of which is substituted with one or more halogen (*e.g.*, F, Cl, Br, or I)), C₃-C₇ cycloalkyl (*e.g.*, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, or cycloheptyl), heterocyclyl comprising one or two 3- to 7-membered rings and 1-3 heteroatoms selected
20 from N, O, and S (*e.g.*, aziridinyl, oxiranyl, thiiranyl, diaziridinyl, oxaziridinyl, dioxiranyl, azetidiny, oxetanyl, thietanyl, diazetidinyl, dioxetanyl, dithietanyl, pyrrolidinyl, tetrahydrofuranyl, tetrahydrothiophenyl, imidazolidinyl, pyrazolidinyl, oxazolidinyl, isoxazolidinyl, thiazolidinyl, isothiazolidinyl, dioxolanyl, dithiolanyl, piperidinyl, tetrahydropyranyl, thianyl, piperazinyl, morpholinyl, thiomorpholinyl, dioxanyl, dithianyl,
25 hexahydro-1,3,5-triazinyl, trioxanyl, trithianyl, azepanyl, oxepanyl, thiepanyl, diazepanyl, hexahydropyridazinyl, hexahydropyrimidinyl, tetrahydrothiopyranyl, or thiomorpholinyl), C₆-C₁₀ aryl (*e.g.*, phenyl), or heteroaryl comprising one or two 5- or 6-membered rings and 1-3 heteroatoms selected from N, O, and S (*e.g.*, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, pyrrolyl, pyrazolyl, imidazolyl, thiazolyl, oxazolyl, isooxazolyl, isothiazolyl, thiadiazolyl,
30 oxadiazolyl, thiophenyl, furanyl, isoquinolinyl, indolyl, pyrazolopyridinyl, indazolyl, benzofuranyl, benzoxazolyl, benzothiazolyl, or benzotriazolyl), wherein the cycloalkyl, heterocyclyl, aryl, or heteroaryl moiety is optionally substituted with one or more R₉.

(H2e-1) In some embodiments, at least one R_4 is $N(R_6')_2$, and one of R_6' is H, and the other R_6' is C_1 - C_6 alkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl, pentyl (straight or branched), or hexyl (straight or branched), each of which is optionally substituted with one or more R_7), or C_1 - C_6 haloalkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl, pentyl (straight or branched), or hexyl (straight or branched), each of which is substituted with one or more halogen (*e.g.*, F, Cl, Br, or I)).

(H2e-2) In some embodiments, at least one R_4 is $N(R_6')_2$, and one of R_6' is H, and the other R_6' is C_1 - C_6 alkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl, pentyl (straight or branched), or hexyl (straight or branched), each of which is optionally substituted with one or more R_7).

(H2e-3) In some embodiments, at least one R_4 is $N(R_6')_2$, and one of R_6' is H, and the other R_6' is C_1 - C_6 haloalkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl, pentyl (straight or branched), or hexyl (straight or branched), each of which is substituted with one or more halogen (*e.g.*, F, Cl, Br, or I)).

(H2e-4) In some embodiments, at least one R_4 is $N(R_6')_2$, and one of R_6' is H, and the other R_6' is C_3 - C_7 cycloalkyl (*e.g.*, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, or cycloheptyl), wherein the cycloalkyl moiety is optionally substituted with one or more R_9 .

(H2e-5) In some embodiments, at least one R_4 is $N(R_6')_2$, and one of R_6' is H, and the other R_6' is heterocyclyl comprising one or two 3- to 7-membered rings and 1-3 heteroatoms selected from N, O, and S (*e.g.*, aziridinyl, oxiranyl, thiranyl, diaziridinyl, oxaziridinyl, dioxiranyl, azetidiny, oxetanyl, thietanyl, diazetidinyl, dioxetanyl, dithietanyl, pyrrolidinyl, tetrahydrofuranyl, tetrahydrothiophenyl, imidazolidinyl, pyrazolidinyl, oxazolidinyl, isoxazolidinyl, thiazolidinyl, isothiazolidinyl, dioxolanyl, dithiolanyl, piperidinyl, tetrahydropyranyl, thianyl, piperazinyl, morpholinyl, thiomorpholinyl, dioxanyl, dithianyl, hexahydro-1,3,5-triazinyl, trioxanyl, trithianyl, azepanyl, oxepanyl, thiepanyl, diazepanyl, hexahydropyridazinyl, hexahydropyrimidinyl, tetrahydrothiopyranyl, or thiomorpholinyl), wherein the heterocyclyl moiety is optionally substituted with one or more R_9 . In some embodiments, at least one R_4 is $N(R_6')_2$, and one of R_6' is H, and the other R_6' is heterocyclyl comprising one 4- to 6-membered ring and 1-3 heteroatoms selected from N, O, and S, as described herein, wherein the heterocyclyl moiety is optionally substituted with one or more R_9 . In some embodiments, at least one R_4 is $N(R_6')_2$, and one of R_6' is H, and the other R_6' is heterocyclyl comprising one 4- to 5-membered ring and 1-3 heteroatoms selected from N, O, and S, as described herein, wherein the heterocyclyl moiety is optionally substituted with one

or more R₉. In some embodiments, at least one R₄ is N(R₆')₂, and one of R₆' is H, and the other R₆' is heterocyclyl comprising one 4-membered ring and 1-2 heteroatoms selected from N, O, and S, as described herein, wherein the heterocyclyl moiety is optionally substituted with one or more R₉. In some embodiments, at least one R₄ is N(R₆')₂, and one of R₆' is H, and the other R₆' is heterocyclyl comprising one 4-membered ring and 1 heteroatom selected from N, O, and S, as described herein, wherein the heterocyclyl moiety is optionally substituted with one or more R₉.

(H2e-6) In some embodiments, at least one R₄ is N(R₆')₂, and one of R₆' is H, and the other R₆' is C₆-C₁₀ aryl (*e.g.*, phenyl), wherein the aryl moiety is optionally substituted with one or more R₉.

(H2e-7) In some embodiments, at least one R₄ is N(R₆')₂, and one of R₆' is H, and the other R₆' is heteroaryl comprising one or two 5- or 6-membered rings and 1-3 heteroatoms selected from N, O, and S (*e.g.*, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, pyrrolyl, pyrazolyl, imidazolyl, thiazolyl, oxazolyl, isooxazolyl, isothiazolyl, thiadiazolyl, oxadiazolyl, thiophenyl, furanyl, isoquinolinyl, indolyl, pyrazolopyridinyl, indazolyl, benzofuranyl, benzoxazolyl, benzothiazolyl, or benzotriazolyl), wherein the heteroaryl moiety is optionally substituted with one or more R₉. In some embodiments, at least one R₄ is N(R₆')₂, and one of R₆' is H, and the other R₆' is heteroaryl comprising one 5- or 6-membered ring and 1-3 heteroatoms selected from N, O, and S, as described herein, wherein the heteroaryl moiety is optionally substituted with one or more R₉. In some embodiments, at least one R₄ is N(R₆')₂, and one of R₆' is H, and the other R₆' is heteroaryl comprising one 5-membered ring and 1-3 heteroatoms selected from N, O, and S, as described herein, wherein the heteroaryl moiety is optionally substituted with one or more R₉. In some embodiments, at least one R₄ is N(R₆')₂, and one of R₆' is H, and the other R₆' is heteroaryl comprising one 6-membered ring and 1-3 heteroatoms selected from N, O, and S, as described herein, wherein the heteroaryl moiety is optionally substituted with one or more R₉.

(H2e-8) In some embodiments, at least one R₄ is N(R₆')₂, and the two R₆' are each independently C₁-C₆ alkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl, pentyl (straight or branched), or hexyl (straight or branched), each of which is optionally substituted with one or more R₇), or C₁-C₆ haloalkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl, pentyl (straight or branched), or hexyl (straight or branched), each of which is substituted with one or more halogen (*e.g.*, F, Cl, Br, or I)).

(H2e-9) In some embodiments, at least one R_4 is $N(R_6')_2$, and the two R_6' together with the atoms to which they are attached form a 3- to 10-membered heterocyclyl comprising 1-3 heteroatoms selected from N, O, and S (*e.g.*, aziridinyl, oxiranyl, thiiranyl, diaziridinyl, oxaziridinyl, dioxiranyl, azetidiny, oxetanyl, thietanyl, diazetidinyl, dioxetanyl, dithietanyl, pyrrolidinyl, tetrahydrofuranyl, tetrahydrothiophenyl, imidazolidinyl, pyrazolidinyl, oxazolidinyl, isoxazolidinyl, thiazolidinyl, isothiazolidinyl, dioxolanyl, dithiolanyl, piperidinyl, tetrahydropyranyl, thianyl, piperazinyl, morpholinyl, thiomorpholinyl, dioxanyl, dithianyl, hexahydro-1,3,5-triazinyl, trioxanyl, trithianyl, azepanyl, oxepanyl, thiepanyl, diazepanyl, hexahydropyridazinyl, hexahydropyrimidinyl, tetrahydrothiopyranyl, or thiomorpholinyl), wherein the heterocyclyl is optionally substituted with one or more R_9 . In some embodiments, at least one R_4 is $N(R_6')_2$, and the two R_6' together with the atoms to which they are attached form a 3- to 6-membered heterocyclyl comprising 1-3 heteroatoms selected from N, O, and S, as described herein, and is optionally substituted with one or more R_9 . In some embodiments, at least one R_4 is $N(R_6')_2$, and the two R_6' together with the atoms to which they are attached form a 4- to 6-membered heterocyclyl comprising 1-3 heteroatoms selected from N, O, and S, as described herein, and is optionally substituted with one or more R_9 . In some embodiments, at least one R_4 is $N(R_6')_2$, and the two R_6' together with the atoms to which they are attached form azetidiny, piperidinyl, morpholinyl, or thiomorpholino, each of which is optionally substituted with one or more R_9 .

(H2e-10) In some embodiments, at least one R_4 is $N(R_6')_2$, and one of R_6' is C_1 - C_6 alkyl or C_1 - C_6 haloalkyl, as described herein, and the other R_6' is cycloalkyl, heterocyclyl comprising one or two 3- to 7-membered rings and 1-3 heteroatoms selected from N, O, and S, C_6 - C_{10} aryl, or heteroaryl comprising one or two 5- or 6-membered rings and 1-3 heteroatoms selected from N, O, and S, as described herein, wherein the cycloalkyl, heterocyclyl, aryl, or heteroaryl moiety is optionally substituted with one or more R_9 .

(H2f) In some embodiments, at least one R_4 is C_6 - C_{10} aryl (*e.g.*, phenyl) or heteroaryl comprising one or two 5- or 6-membered rings and 1-3 heteroatoms selected from N, O, and S (*e.g.*, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, pyrrolyl, pyrazolyl, imidazolyl, thiazolyl, oxazolyl, isooxazolyl, isothiazolyl, thiadiazolyl, oxadiazolyl, thiophenyl, furanyl, isoquinolinyl, indolyl, pyrazolopyridinyl, indazolyl, benzofuranyl, benzoxazolyl, benzothiazolyl, or benzotriazolyl), wherein the aryl or heteroaryl moiety is optionally substituted with one or more R_9 .

(H2f-1) In some embodiments, at least one R₄ is C₆-C₁₀ aryl (*e.g.*, phenyl), wherein the aryl moiety is optionally substituted with one or more R₉.

(H2f-2) In some embodiments, at least one R₄ is heteroaryl comprising one or two 5- or 6-membered rings and 1-3 heteroatoms selected from N, O, and S (*e.g.*, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, pyrrolyl, pyrazolyl, imidazolyl, thiazolyl, oxazolyl, isooxazolyl, isothiazolyl, thiadiazolyl, oxadiazolyl, thiophenyl, furanyl, isoquinolinyl, indolyl, pyrazolopyridinyl, indazolyl, benzofuranyl, benzoxazolyl, benzothiazolyl, or benzotriazolyl), wherein the heteroaryl moiety is optionally substituted with one or more R₉. In some embodiments, at least one R₄ is heteroaryl comprising one 5- or 6-membered ring and 1-3 heteroatoms selected from N, O, and S, as described herein, wherein the heteroaryl moiety is optionally substituted with one or more R₉. In some embodiments, at least one R₄ is heteroaryl comprising one 5-membered ring and 1-3 heteroatoms selected from N, O, and S, as described herein, wherein the heteroaryl moiety is optionally substituted with one or more R₉. In some embodiments, at least one R₄ is heteroaryl comprising one 6-membered ring and 1-3 heteroatoms selected from N, O, and S, as described herein, wherein the heteroaryl moiety is optionally substituted with one or more R₉.

(H3) In some embodiments, only one R₄ is halogen, CN, OH, N(R₆')₂, C₁-C₆ alkoxy, C(=O)N(R₆)₂, C(=O)OR₆, C(=O)R₆, Q-T, C₆-C₁₀ aryl, or heteroaryl comprising one or two 5- or 6-membered rings and 1-3 heteroatoms selected from N, O, and S, wherein the aryl or heteroaryl moiety is optionally substituted with one or more R₉.

(H3a) In some embodiments, only one R₄ is halogen (*e.g.*, F, Cl, Br, or I), CN, OH, N(R₆')₂, C₁-C₆ alkoxy (*e.g.*, methoxy, ethoxy, propoxy, butoxy, pentoxy, or hexoxy), C(=O)N(R₆)₂, C(=O)OR₆, C(=O)R₆, or Q-T.

(H3b) In some embodiments, only one R₄ is CN, OH, N(R₆')₂, C₁-C₆ alkoxy (*e.g.*, methoxy, ethoxy, propoxy, butoxy, pentoxy, or hexoxy), C(=O)N(R₆)₂, C(=O)OR₆, C(=O)R₆, or Q-T.

(H3c) In some embodiments, only one R₄ is C(=O)N(R₆)₂, C(=O)OR₆, C(=O)R₆, or Q-T.

(H3d) In some embodiments, only one R₄ is Q-T.

(H3e) In some embodiments, only one R₄ is N(R₆')₂, and one of R₆' is H, and the other R₆' is C₁-C₆ alkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl, pentyl (straight or branched), or hexyl (straight or branched), each of which is optionally substituted with one or more R₇), C₁-C₆ haloalkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-

propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl, pentyl (straight or branched), or hexyl (straight or branched), each of which is substituted with one or more halogen (*e.g.*, F, Cl, Br, or I)), C₃-C₇ cycloalkyl, heterocyclyl comprising one or two 3- to 7-membered rings and 1-3 heteroatoms selected from N, O, and S (*e.g.*, aziridinyl, oxiranyl, thiiranyl, diaziridinyl, oxaziridinyl, dioxiranyl, azetidiny, oxetanyl, thietanyl, diazetidinyl, dioxetanyl, dithietanyl, pyrrolidinyl, tetrahydrofuranyl, tetrahydrothiophenyl, imidazolidinyl, pyrazolidinyl, oxazolidinyl, isoxazolidinyl, thiazolidinyl, isothiazolidinyl, dioxolanyl, dithiolanyl, piperidinyl, tetrahydropyranyl, thianyl, piperazinyl, morpholinyl, thiomorpholinyl, dioxanyl, dithianyl, hexahydro-1,3,5-triazinyl, trioxanyl, trithianyl, azepanyl, oxepanyl, thiepanyl, diazepanyl, hexahydropyridazinyl, hexahydropyrimidinyl, tetrahydrothiopyranyl, or thiomorpholinyl), C₆-C₁₀ aryl (*e.g.*, phenyl), or heteroaryl comprising one or two 5- or 6-membered rings and 1-3 heteroatoms selected from N, O, and S (*e.g.*, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, pyrrolyl, pyrazolyl, imidazolyl, thiazolyl, oxazolyl, isooxazolyl, isothiazolyl, thiadiazolyl, oxadiazolyl, thiophenyl, furanyl, isoquinolinyl, indolyl, pyrazolopyridinyl, indazolyl, benzofuranyl, benzoxazolyl, benzothiazolyl, or benzotriazolyl), wherein the cycloalkyl, heterocyclyl, aryl, or heteroaryl moiety is optionally substituted with one or more R₉.

(H3e-1) In some embodiments, only one R₄ is N(R₆')₂, and one of R₆' is H, and the other R₆' is C₁-C₆ alkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl, pentyl (straight or branched), or hexyl (straight or branched), each of which is optionally substituted with one or more R₇), or C₁-C₆ haloalkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl, pentyl (straight or branched), or hexyl (straight or branched), each of which is substituted with one or more halogen (*e.g.*, F, Cl, Br, or I)).

(H3e-2) In some embodiments, only one R₄ is N(R₆')₂, and one of R₆' is H, and the other R₆' is C₁-C₆ alkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl, pentyl (straight or branched), or hexyl (straight or branched), each of which is optionally substituted with one or more R₇).

(H3e-3) In some embodiments, only one R₄ is N(R₆')₂, and one of R₆' is H, and the other R₆' is C₁-C₆ haloalkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl, pentyl (straight or branched), or hexyl (straight or branched), each of which is substituted with one or more halogen (*e.g.*, F, Cl, Br, or I)).

(H3e-4) In some embodiments, only one R₄ is N(R₆')₂, and one of R₆' is H, and the other R₆' is C₃-C₇ cycloalkyl (*e.g.*, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, or cycloheptyl), wherein the cycloalkyl moiety is optionally substituted with one or more R₉.

(H3e-5) In some embodiments, only one R_4 is $N(R_6')_2$, and one of R_6' is H, and the other R_6' is heterocyclyl comprising one or two 3- to 7-membered rings and 1-3 heteroatoms selected from N, O, and S (*e.g.*, aziridinyl, oxiranyl, thiranyl, diaziridinyl, oxaziridinyl, dioxiranyl, azetidiny, oxetanyl, thietanyl, diazetidinyl, dioxetanyl, dithietanyl, pyrrolidinyl, tetrahydrofuranyl, tetrahydrothiophenyl, imidazolidinyl, pyrazolidinyl, oxazolidinyl, isoxazolidinyl, thiazolidinyl, isothiazolidinyl, dioxolanyl, dithiolanyl, piperidinyl, tetrahydropyranyl, thianyl, piperazinyl, morpholinyl, thiomorpholinyl, dioxanyl, dithianyl, hexahydro-1,3,5-triazinyl, trioxanyl, trithianyl, azepanyl, oxepanyl, thiepanyl, diazepanyl, hexahydropyridazinyl, hexahydropyrimidinyl, tetrahydrothiopyranyl, or thiomorpholinyl), wherein the heterocyclyl moiety is optionally substituted with one or more R_9 . In some embodiments, only one R_4 is $N(R_6')_2$, and one of R_6' is H, and the other R_6' is heterocyclyl comprising one 4- to 6-membered ring and 1-3 heteroatoms selected from N, O, and S, as described herein, wherein the heterocyclyl moiety is optionally substituted with one or more R_9 . In some embodiments, only one R_4 is $N(R_6')_2$, and one of R_6' is H, and the other R_6' is heterocyclyl comprising one 4- to 5-membered ring and 1-3 heteroatoms selected from N, O, and S, as described herein, wherein the heterocyclyl moiety is optionally substituted with one or more R_9 . In some embodiments, only one R_4 is $N(R_6')_2$, and one of R_6' is H, and the other R_6' is heterocyclyl comprising one 4-membered ring and 1-2 heteroatoms selected from N, O, and S, as described herein, wherein the heterocyclyl moiety is optionally substituted with one or more R_9 . In some embodiments, only one R_4 is $N(R_6')_2$, and one of R_6' is H, and the other R_6' is heterocyclyl comprising one 4-membered ring and 1 heteroatom selected from N, O, and S, as described herein, wherein the heterocyclyl moiety is optionally substituted with one or more R_9 .

(H3e-6) In some embodiments, only one R_4 is $N(R_6')_2$, and one of R_6' is H, and the other R_6' is C_6 - C_{10} aryl (*e.g.*, phenyl), wherein the aryl moiety is optionally substituted with one or more R_9 .

(H3e-7) In some embodiments, only one R_4 is $N(R_6')_2$, and one of R_6' is H, and the other R_6' is heteroaryl comprising one or two 5- or 6-membered rings and 1-3 heteroatoms selected from N, O, and S (*e.g.*, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, pyrrolyl, pyrazolyl, imidazolyl, thiazolyl, oxazolyl, isooxazolyl, isothiazolyl, thiadiazolyl, oxadiazolyl, thiophenyl, furanyl, isoquinolinyl, indolyl, pyrazolopyridinyl, indazolyl, benzofuranyl, benzoxazolyl, benzothiazolyl, or benzotriazolyl), wherein the heteroaryl moiety is optionally substituted with one or more R_9 . In some embodiments, only one R_4 is $N(R_6')_2$, and one of

R₆' is H, and the other R₆' is heteroaryl comprising one 5- or 6-membered ring and 1-3 heteroatoms selected from N, O, and S, as described herein, wherein the heteroaryl moiety is optionally substituted with one or more R₉. In some embodiments, only one R₄ is N(R₆')₂, and one of R₆' is H, and the other R₆' is heteroaryl comprising one 5-membered ring and 1-3 heteroatoms selected from N, O, and S, as described herein, wherein the heteroaryl moiety is optionally substituted with one or more R₉. In some embodiments, only one R₄ is N(R₆')₂, and one of R₆' is H, and the other R₆' is heteroaryl comprising one 6-membered ring and 1-3 heteroatoms selected from N, O, and S, as described herein, wherein the heteroaryl moiety is optionally substituted with one or more R₉.

10 (H3e-8) In some embodiments, only one R₄ is N(R₆')₂, and the two R₆' are each independently C₁-C₆ alkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl, pentyl (straight or branched), or hexyl (straight or branched), each of which is optionally substituted with one or more R₇), or C₁-C₆ haloalkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl, pentyl (straight or branched), or hexyl (straight or branched), each of which is substituted with one or more halogen (*e.g.*, F, Cl, Br, or I)).

(H3e-9) In some embodiments, only one R₄ is N(R₆')₂, and the two R₆' together with the atoms to which they are attached form a 3- to 10-membered heterocyclyl comprising 1-3 heteroatoms selected from N, O, and S (*e.g.*, aziridinyl, oxiranyl, thiiranyl, diaziridinyl, oxaziridinyl, dioxiranyl, azetidiny, oxetanyl, thietanyl, diazetidinyl, dioxetanyl, dithietanyl, pyrrolidinyl, tetrahydrofuranyl, tetrahydrothiophenyl, imidazolidinyl, pyrazolidinyl, oxazolidinyl, isoxazolidinyl, thiazolidinyl, isothiazolidinyl, dioxolanyl, dithiolanyl, piperidinyl, tetrahydropyranyl, thianyl, piperazinyl, morpholinyl, thiomorpholinyl, dioxanyl, dithianyl, hexahydro-1,3,5-triazinyl, trioxanyl, trithianyl, azepanyl, oxepanyl, thiepanyl, diazepanyl, hexahydropyridazinyl, hexahydropyrimidinyl, tetrahydrothiopyranyl, or thiomorpholinyl), wherein the heterocyclyl is optionally substituted with one or more R₉. In some embodiments, only one R₄ is N(R₆')₂, and the two R₆' together with the atoms to which they are attached form a 3- to 6-membered heterocyclyl comprising 1-3 heteroatoms selected from N, O, and S, as described herein, and is optionally substituted with one or more R₉. In some embodiments, only one R₄ is N(R₆')₂, and the two R₆' together with the atoms to which they are attached form a 4- to 6-membered heterocyclyl comprising 1-3 heteroatoms selected from N, O, and S, as described herein, and is optionally substituted with one or more R₉. In some embodiments, only one R₄ is N(R₆')₂, and the two R₆' together with the atoms to which

they are attached form azetidiny, piperidiny, morpholiny, or thiomorpholino, each of which is optionally substituted with one or more R₉.

(H3e-10) In some embodiments, only one R₄ is N(R₆')₂, and one of R₆' is C₁-C₆ alkyl or C₁-C₆ haloalkyl, as described herein, and the other R₆' is cycloalkyl, heterocyclyl comprising one or two 3- to 7-membered rings and 1-3 heteroatoms selected from N, O, and S, C₆-C₁₀ aryl, or heteroaryl comprising one or two 5- or 6-membered rings and 1-3 heteroatoms selected from N, O, and S, as described herein, wherein the cycloalkyl, heterocyclyl, aryl, or heteroaryl moiety is optionally substituted with one or more R₉.

(H3f) In some embodiments, only one R₄ is C₆-C₁₀ aryl (*e.g.*, phenyl) or heteroaryl comprising one or two 5- or 6-membered rings and 1-3 heteroatoms selected from N, O, and S (*e.g.*, pyridiny, pyraziny, pyrimidiny, pyridaziny, pyrroly, pyrazoly, imidazoly, thiazoly, oxazoly, isooxazoly, isothiazoly, thiadiazoly, oxadiazoly, thiopheny, furany, isoquinolinyl, indoly, pyrazolopyridiny, indazoly, benzofurany, benzoxazoly, benzothiazoly, or benzotriazoly), wherein the aryl or heteroaryl moiety is optionally substituted with one or more R₉.

(H3f-1) In some embodiments, only one R₄ is C₆-C₁₀ aryl (*e.g.*, phenyl), wherein the aryl moiety is optionally substituted with one or more R₉.

(H3f-2) In some embodiments, only one R₄ is heteroaryl comprising one or two 5- or 6-membered rings and 1-3 heteroatoms selected from N, O, and S (*e.g.*, pyridiny, pyraziny, pyrimidiny, pyridaziny, pyrroly, pyrazoly, imidazoly, thiazoly, oxazoly, isooxazoly, isothiazoly, thiadiazoly, oxadiazoly, thiopheny, furany, isoquinolinyl, indoly, pyrazolopyridiny, indazoly, benzofurany, benzoxazoly, benzothiazoly, or benzotriazoly), wherein the heteroaryl moiety is optionally substituted with one or more R₉. In some embodiments, only one R₄ is heteroaryl comprising one 5- or 6-membered ring and 1-3 heteroatoms selected from N, O, and S, as described herein, wherein the heteroaryl moiety is optionally substituted with one or more R₉. In some embodiments, only one R₄ is heteroaryl comprising one 5-membered ring and 1-3 heteroatoms selected from N, O, and S, as described herein, wherein the heteroaryl moiety is optionally substituted with one or more R₉. In some embodiments, only one R₄ is heteroaryl comprising one 6-membered ring and 1-3 heteroatoms selected from N, O, and S, as described herein, wherein the heteroaryl moiety is optionally substituted with one or more R₉.

(H3g) In some embodiments, at least one R₄ is C₁-C₆ alkoxy (*e.g.*, methoxy, ethoxy, propoxy, butoxy, pentoxy, or hexoxy).

(H3g-1) In some embodiments, only one R₄ is C₁-C₆ alkoxy (*e.g.*, methoxy, ethoxy, propoxy, butoxy, pentoxy, or hexoxy).

(I1) In some embodiments, at least one Q is C₁-C₄ alkylene (*e.g.*, methylene, ethylene, propylene, or butylene).

5 (I2) In some embodiments, at least one Q is O-(C₁-C₄) alkylene (*e.g.*, O-methylene, O-ethylene, O-propylene, or O-butylene).

(J1) In some embodiments, at least one T is C₁-C₄ alkoxy (*e.g.*, methoxy, ethoxy, propoxy, or butoxy), OH, N(R₆)₂, N(R₅)C(=O)R₆, N(R₅)C(=O)OR₆, C(=O)N(R₆)₂, C(=O)OR₆, C(=O)R₆, C₆-C₁₀ aryl, or heteroaryl comprising one or two 5- or 6-membered
10 rings and 1-3 heteroatoms selected from N, O, and S, wherein the aryl or heteroaryl moiety is optionally substituted with one or more R₉.

(J2) In some embodiments, at least one T is C₁-C₄ alkoxy (*e.g.*, methoxy, ethoxy, propoxy, or butoxy).

(J3) In some embodiments, at least one T is NH₂, NH(R₆), or N(R₆)₂.

15 (J3-1) In some embodiments, at least one T is NH₂, NH(R₆), or N(R₆)₂, wherein R₆ is C₁-C₄ alkyl moiety is selected from methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, and *t*-butyl.

(J4) In some embodiments, at least one T is N(R₅)C(=O)R₆ or N(R₅)C(=O)OR₆.

(J5) In some embodiments, at least one T is C(=O)N(R₆)₂, C(=O)OR₆, or C(=O)R₆.

20 (J6) In some embodiments, at least one T is OH.

(J7) In some embodiments, at least one T is C₆-C₁₀ aryl or heteroaryl comprising one or two 5- or 6-membered rings and 1-3 heteroatoms selected from N, O, and S, wherein the aryl or heteroaryl moiety is optionally substituted with one or more R₉.

25 (J7-1) In some embodiments, at least one T is C₆-C₁₀ aryl (*e.g.*, phenyl), optionally substituted with one or more R₉.

(J7-2) In some embodiments, at least one T is heteroaryl comprising one or two 5- or 6-membered rings and 1-3 heteroatoms selected from N, O, and S (*e.g.*, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, pyrrolyl, pyrazolyl, imidazolyl, thiazolyl, oxazolyl, isooxazolyl, isothiazolyl, thiadiazolyl, oxadiazolyl, thiophenyl, furanyl, isoquinolinyl, indolyl,
30 pyrazolopyridinyl, indazolyl, benzofuranyl, benzoxazolyl, benzothiazolyl, or benzotriazolyl), optionally substituted with one or more R₉. In some embodiments, at least one T is heteroaryl comprising one 5- or 6-membered ring and 1-3 heteroatoms selected from N, O, and S, as described herein, wherein the heteroaryl moiety is optionally substituted with one or more R₉.

In some embodiments, at least one T is heteroaryl comprising one 5-membered ring and 1-3 heteroatoms selected from N, O, and S, as described herein, wherein the heteroaryl moiety is optionally substituted with one or more R₉. In some embodiments, at least one T is heteroaryl comprising one 6-membered ring and 1-3 heteroatoms selected from N, O, and S, as

5 described herein, wherein the heteroaryl moiety is optionally substituted with one or more R₉.

(K1) In some embodiments, each R₅ is H.

(K2) In some embodiments, at least one R₅ is C₁-C₄ alkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, or *t*-butyl).

(L1) In some embodiments, each R₆ is H.

10 (L2) In some embodiments, at least one R₆ is C₁-C₆ alkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl, pentyl (straight or branched), or hexyl (straight or branched), each of which is optionally substituted with one or more R₇), C₁-C₆ haloalkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl, pentyl (straight or branched), or hexyl (straight or branched), each of which is substituted with one
15 or more halogen (*e.g.*, F, Cl, Br, or I)), C₆-C₁₀ aryl (*e.g.*, phenyl), or heteroaryl comprising one or two 5- or 6-membered rings and 1-3 heteroatoms selected from N, O, and S (*e.g.*, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, pyrrolyl, pyrazolyl, imidazolyl, thiazolyl, oxazolyl, isooxazolyl, isothiazolyl, thiadiazolyl, oxadiazolyl, thiophenyl, furanyl, isoquinolinyl, indolyl, pyrazolopyridinyl, indazolyl, benzofuranyl, benzoxazolyl,
20 benzothiazolyl, or benzotriazolyl), wherein the aryl or heteroaryl moiety is optionally substituted with one or more R₉.

(L2a) In some embodiments, at least one R₆ is C₁-C₆ alkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl, pentyl (straight or branched), or hexyl (straight or branched), each of which is optionally substituted with one or more R₇), or C₁-C₆
25 haloalkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl, pentyl (straight or branched), or hexyl (straight or branched), each of which is substituted with one or more halogen (*e.g.*, F, Cl, Br, or I)).

(L2b) In some embodiments, at least one R₆ is C₁-C₆ alkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl, pentyl (straight or branched), or hexyl
30 (straight or branched), each of which is optionally substituted with one or more R₇).

(L2c) In some embodiments, at least one R₆ is C₆-C₁₀ aryl (*e.g.*, phenyl), wherein the aryl moiety is optionally substituted with one or more R₉.

(L2d) In some embodiments, at least one R₆ is heteroaryl comprising one or two 5- or 6-membered rings and 1-3 heteroatoms selected from N, O, and S (*e.g.*, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, pyrrolyl, pyrazolyl, imidazolyl, thiazolyl, oxazolyl, isooxazolyl, isothiazolyl, thiadiazolyl, oxadiazolyl, thiophenyl, furanyl, isoquinolinyl, indolyl, pyrazolopyridinyl, indazolyl, benzofuranyl, benzoxazolyl, benzothiazolyl, or benzotriazolyl), wherein the heteroaryl moiety is optionally substituted with one or more R₉. In some embodiments, at least one R₆ is heteroaryl comprising one 5- or 6-membered ring and 1-3 heteroatoms selected from N, O, and S, as described herein, wherein the heteroaryl moiety is optionally substituted with one or more R₉. In some embodiments, at least one R₆ is heteroaryl comprising one 5-membered ring and 1-3 heteroatoms selected from N, O, and S, as described herein, wherein the heteroaryl moiety is optionally substituted with one or more R₉. In some embodiments, at least one R₆ is heteroaryl comprising one 6-membered ring and 1-3 heteroatoms selected from N, O, and S, as described herein, wherein the heteroaryl moiety is optionally substituted with one or more R₉.

(L2e) In some embodiments, two R₆ together with the atoms to which they are attached form a 3- to 10-membered heterocyclyl comprising 1-3 heteroatoms selected from N, O, and S (*e.g.*, aziridinyl, oxiranyl, thiiranyl, diaziridinyl, oxaziridinyl, dioxiranyl, azetidiny, oxetanyl, thietanyl, diazetidinyl, dioxetanyl, dithietanyl, pyrrolidinyl, tetrahydrofuranyl, tetrahydrothiophenyl, imidazolidinyl, pyrazolidinyl, oxazolidinyl, isoxazolidinyl, thiazolidinyl, isothiazolidinyl, dioxolanyl, dithiolanyl, piperidinyl, tetrahydropyranyl, thianyl, piperazinyl, morpholinyl, thiomorpholinyl, dioxanyl, dithianyl, hexahydro-1,3,5-triazinyl, trioxanyl, trithianyl, azepanyl, oxepanyl, thiepanyl, diazepanyl, hexahydropyridazinyl, hexahydropyrimidinyl, tetrahydrothiopyranyl, or thiomorpholinyl), wherein the heterocyclyl is optionally substituted with one or more R₉. In some embodiments, two R₆ together with the atoms to which they are attached form a 3- to 6-membered heterocyclyl comprising 1-3 heteroatoms selected from N, O, and S, as described herein, and is optionally substituted with one or more R₉. In some embodiments, two R₆ together with the atoms to which they are attached form a 4- to 6-membered heterocyclyl comprising 1-3 heteroatoms selected from N, O, and S, as described herein, and is optionally substituted with one or more R₉.

(M1a) In some embodiments, each R₇ is independently N(R₈)₂, and each R₈ is H.

(M1b) In some embodiments, each R_7 is independently $N(R_8)_2$, and one R_8 is H, and the other R_8 is C_1 - C_6 alkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl, pentyl (straight or branched), or hexyl (straight or branched)).

(M1c) In some embodiments, each R_7 is independently $N(R_8)_2$, and each R_8 is
 5 independently C_1 - C_6 alkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl, pentyl (straight or branched), or hexyl (straight or branched)).

(M2a) In some embodiments, each R_7 is independently OR_8 , and R_8 is H.

(M2b) In some embodiments, each R_7 is independently OR_8 , and R_8 is C_1 - C_6 alkyl
 10 (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl, pentyl (straight or branched), or hexyl (straight or branched)).

(M2c) In some embodiments, each R_7 is independently C_6 - C_{10} aryl (*e.g.*, phenyl) or heteroaryl comprising one or two 5- or 6-membered rings and 1-3 heteroatoms selected from N, O, and S (*e.g.*, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, pyrrolyl, pyrazolyl, imidazolyl, thiazolyl, oxazolyl, isooxazolyl, isothiazolyl, thiadiazolyl, oxadiazolyl,
 15 thiophenyl, furanyl, isoquinolinyl, indolyl, pyrazolopyridinyl, indazolyl, benzofuranyl, benzoxazolyl, benzothiazolyl, or benzotriazolyl).

(M2c-1) In some embodiments, each R_7 is independently C_6 - C_{10} aryl (*e.g.*, phenyl).

(M2c-2) In some embodiments, each R_7 is independently heteroaryl comprising one or two 5- or 6-membered rings and 1-3 heteroatoms selected from N, O, and S (*e.g.*,
 20 pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, pyrrolyl, pyrazolyl, imidazolyl, thiazolyl, oxazolyl, isooxazolyl, isothiazolyl, thiadiazolyl, oxadiazolyl, thiophenyl, furanyl, isoquinolinyl, indolyl, pyrazolopyridinyl, indazolyl, benzofuranyl, benzoxazolyl, benzothiazolyl, or benzotriazolyl).


(N1) In some embodiments, each R_9 is independently oxo, halogen (*e.g.*, F, Cl, Br, or
 25 I), OH, CN, NH_2 , $N(C_1-C_4 \text{ alkyl})_2$ (*e.g.*, dimethylamino, methylethylamino, diethylamino, methylpropylamino, ethylpropylamino, dipropylamino, methylbutylamino, ethylbutylamino, propylbutylamino, or dibutylamino), C_1 - C_6 alkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl, pentyl (straight or branched), or hexyl (straight or branched)), C_1 - C_6 haloalkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl,
 30 pentyl (straight or branched), or hexyl (straight or branched), each of which is substituted with one or more halogen (*e.g.*, F, Cl, Br, or I)), C_1 - C_6 alkoxy (*e.g.*, methoxy, ethoxy, propoxy, butoxy, pentoxy, or hexyloxy), or C_1 - C_6 haloalkoxy (*e.g.*, methoxy, ethoxy, propoxy, butoxy, pentoxy, or hexyloxy, each of which is substituted with one or more


halogen (*e.g.*, F, Cl, Br, or I)), wherein the C₁-C₆ alkyl is optionally substituted with one or more groups selected from oxo, OH, O(C₁-C₄ alkyl), CN, NH₂, and N(C₁-C₄ alkyl)₂ (*e.g.*, dimethylamino, methylethylamino, diethylamino, methylpropylamino, ethylpropylamino, dipropylamino, methylbutylamino, ethylbutylamino, propylbutylamino, or dibutylamino).


- 5 (N2) In some embodiments, each R₉ is independently OH, CN, NH₂, N(C₁-C₄ alkyl)₂ (*e.g.*, dimethylamino, methylethylamino, diethylamino, methylpropylamino, ethylpropylamino, dipropylamino, methylbutylamino, ethylbutylamino, propylbutylamino, or dibutylamino), C₁-C₆ alkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl, pentyl (straight or branched), or hexyl (straight or branched)), C₁-C₆ haloalkyl (*e.g.*, methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl, pentyl (straight or branched), or hexyl (straight or branched)), each of which is substituted with one or more halogen (*e.g.*, F, Cl, Br, or I)), C₁-C₆ alkoxy (*e.g.*, methoxy, ethoxy, propoxy, butoxy, pentoxy, or hexyloxy), or C₁-C₆ haloalkoxy (*e.g.*, methoxy, ethoxy, propoxy, butoxy, pentoxy, or hexyloxy, each of which is substituted with one or more halogen (*e.g.*, F, Cl, Br, or I)), wherein the C₁-C₆ alkyl is optionally substituted with one or more groups selected from OH, CN, NH₂, and N(C₁-C₄ alkyl)₂ (*e.g.*, dimethylamino, methylethylamino, diethylamino, methylpropylamino, ethylpropylamino, dipropylamino, methylbutylamino, ethylbutylamino, propylbutylamino, or dibutylamino).


- For a compound of any of Formula I, Ia, Ib, Ic, Id, Ie, If, Ig, IIa, IIb, IIc, IId, IIe, IIg, or IIg, where applicable, each of the substituent groups illustrated herein for any of Cy, X₁, X₂, A₁, A₂, A₃, A₄, R₁, R₂, R₃, R₄, R₅, R₆, R₆', R₇, R₈, R₉, Q, and T, can be combined with any of the substituent groups illustrated herein for one or more of the remainder of Cy, X₁, X₂, A₁, A₂, A₃, A₄, R₁, R₂, R₃, R₄, R₅, R₆, R₆', R₇, R₈, R₉, Q, and T.

For example:


- 25 (O1) The thiazolyl ring  is as illustrated in any of (A1)-(A3), and the Cy ring is as illustrated in (B1).

- (O2) The thiazolyl ring  is as illustrated in any of (A1)-(A3), and the Cy ring is as illustrated in (B2).


- (O3) The thiazolyl ring  is as illustrated in any of (A1)-(A3), and the Cy ring is as illustrated in (B3).




(P1) The thiazolyl ring is as illustrated in any of (A1)-(A3), and A₁, A₂, A₃, and A₄ are each as illustrated in (G1).




(P2) The thiazolyl ring is as illustrated in any of (A1)-(A3), and A₁, A₂, A₃, and A₄ are each as illustrated in (G2).




5 (P3) The thiazolyl ring is as illustrated in any of (A1)-(A3), and A₁, A₂, A₃, and A₄ are each as illustrated in any of (G3)-(G3d).




(P4) The thiazolyl ring is as illustrated in any of (A1)-(A3), and A₁, A₂, A₃, and A₄ are each as illustrated in any of (G4)-(G4f).



10 (P5) The thiazolyl ring is as illustrated in any of (A1)-(A3), and A₁, A₂, A₃, and A₄ are each as illustrated in any of (G5)-(G5d).



(P6) The thiazolyl ring is as illustrated in any of (A1)-(A3), and A₁, A₂, A₃, and A₄ are each as illustrated in any of (G6).



(P7) The thiazolyl ring is as illustrated in any of (A1)-(A3), and A₁, A₂, A₃, and A₄ are each as illustrated in any of (G7).

15 (Q1a) The Cy ring is as illustrated in (B1), and A₁, A₂, A₃, and A₄ are each as illustrated in (G1).

(Q1b) The Cy ring is as illustrated in (B1), and A₁, A₂, A₃, and A₄ are each as illustrated in (G2).

20 (Q1c) The Cy ring is as illustrated in (B1), and A₁, A₂, A₃, and A₄ are each as illustrated in any of (G3)-(G3d).

(Q1d) The Cy ring is as illustrated in (B1), and A₁, A₂, A₃, and A₄ are each as illustrated in any of (G4)-(G4f).

(Q1e) The Cy ring is as illustrated in (B1), and A₁, A₂, A₃, and A₄ are each as illustrated in any of (G5)-(G5d).

(Q1f) The Cy ring is as illustrated in (B1), and A₁, A₂, A₃, and A₄ are each as illustrated in any of (G6).

(Q1g) The Cy ring is as illustrated in (B1), and A₁, A₂, A₃, and A₄ are each as illustrated in any of (G7).

5 (Q2a) The Cy ring is as illustrated in (B2), and A₁, A₂, A₃, and A₄ are each as illustrated in (G1).

(Q2b) The Cy ring is as illustrated in (B2), and A₁, A₂, A₃, and A₄ are each as illustrated in (G2).

(Q2c) The Cy ring is as illustrated in (B2), and A₁, A₂, A₃, and A₄ are each as
10 illustrated in any of (G3)-(G3d).

(Q2d) The Cy ring is as illustrated in (B2), and A₁, A₂, A₃, and A₄ are each as illustrated in any of (G4)-(G4f).

(Q2e) The Cy ring is as illustrated in (B2), and A₁, A₂, A₃, and A₄ are each as illustrated in any of (G5)-(G5d).

15 (Q2f) The Cy ring is as illustrated in (B2), and A₁, A₂, A₃, and A₄ are each as illustrated in any of (G6).

(Q2g) The Cy ring is as illustrated in (B2), and A₁, A₂, A₃, and A₄ are each as illustrated in any of (G7).

(Q3a) The Cy ring is as illustrated in (B3), and A₁, A₂, A₃, and A₄ are each as
20 illustrated in (G1).

(Q3b) The Cy ring is as illustrated in (B3), and A₁, A₂, A₃, and A₄ are each as illustrated in (G2).


(Q3c) The Cy ring is as illustrated in (B3), and A₁, A₂, A₃, and A₄ are each as illustrated in any of (G3)-(G3d).


25 (Q3d) The Cy ring is as illustrated in (B3), and A₁, A₂, A₃, and A₄ are each as illustrated in any of (G4)-(G4f).


(Q3e) The Cy ring is as illustrated in (B3), and A₁, A₂, A₃, and A₄ are each as illustrated in any of (G5)-(G5d).


(Q3f) The Cy ring is as illustrated in (B3), and A₁, A₂, A₃, and A₄ are each as
30 illustrated in any of (G6).


(Q3g) The Cy ring is as illustrated in (B3), and A₁, A₂, A₃, and A₄ are each as illustrated in any of (G7).


(Q4a) The thiazolyl ring  and the Cy ring are illustrated as in any of (O1)-(O3), and A₁, A₂, A₃, and A₄ are each as illustrated in (G1).


(Q4b) The thiazolyl ring  and the Cy ring are illustrated as in any of (O1)-(O3), and A₁, A₂, A₃, and A₄ are each as illustrated in (G2).


5 (Q4c) The thiazolyl ring  and the Cy ring are illustrated as in any of (O1)-(O3), and A₁, A₂, A₃, and A₄ are each as illustrated in any of (G3)-(G3d).


(Q4d) The thiazolyl ring  and the Cy ring are illustrated as in any of (O1)-(O3), and A₁, A₂, A₃, and A₄ are each as illustrated in any of (G4)-(G4f).


10 (Q4e) The thiazolyl ring  and the Cy ring are illustrated as in any of (O1)-(O3), and A₁, A₂, A₃, and A₄ are each as illustrated in any of (G5)-(G5d).


(Q4f) The thiazolyl ring  and the Cy ring are illustrated as in any of (O1)-(O3), and A₁, A₂, A₃, and A₄ are each as illustrated in any of (G6).


(Q4g) The thiazolyl ring  and the Cy ring are illustrated as in any of (O1)-(O3), and A₁, A₂, A₃, and A₄ are each as illustrated in any of (G7).


15 (R1) The thiazolyl ring , the Cy ring, and/or A₁, A₂, A₃, and A₄ are each as illustrated, as applicable, in any or (A1)-(A3), (B1)-(B3), (G1)-(G7), and (O1)-(Q4g), R₄ is as illustrated in (H1).


20 (R2) The thiazolyl ring , the Cy ring, and/or A₁, A₂, A₃, and A₄ are each as illustrated, as applicable, in any or (A1)-(A3), (B1)-(B3), (G1)-(G7), and (O1)-(Q4g), R₄ is as illustrated in (H2).


(R3) The thiazolyl ring , the Cy ring, and/or A₁, A₂, A₃, and A₄ are each as illustrated, as applicable, in any or (A1)-(A3), (B1)-(B3), (G1)-(G7), and (O1)-(Q4g), R₄ is as illustrated in (H2a).


(R4) The thiazolyl ring , the Cy ring, and/or A₁, A₂, A₃, and A₄ are each as illustrated, as applicable, in any or (A1)-(A3), (B1)-(B3), (G1)-(G7), and (O1)-(Q4g), R₄ is as illustrated in (H2b).


(R5) The thiazolyl ring , the Cy ring, and/or A₁, A₂, A₃, and A₄ are each as illustrated, as applicable, in any or (A1)-(A3), (B1)-(B3), (G1)-(G7), and (O1)-(Q4g), R₄ is as illustrated in (H2c).


(R6) The thiazolyl ring , the Cy ring, and/or A₁, A₂, A₃, and A₄ are each as illustrated, as applicable, in any or (A1)-(A3), (B1)-(B3), (G1)-(G7), and (O1)-(Q4g), R₄ is as illustrated in (H2d).


(R7) The thiazolyl ring , the Cy ring, and/or A₁, A₂, A₃, and A₄ are each as illustrated, as applicable, in any or (A1)-(A3), (B1)-(B3), (G1)-(G7), and (O1)-(Q4g), R₄ is as illustrated in any of (H2e)-(H2e-10).


(R8) The thiazolyl ring , the Cy ring, and/or A₁, A₂, A₃, and A₄ are each as illustrated, as applicable, in any or (A1)-(A3), (B1)-(B3), (G1)-(G7), and (O1)-(Q4g), R₄ is as illustrated in (H2f)-(H2f-2).


(R9) The thiazolyl ring , the Cy ring, and/or A₁, A₂, A₃, and A₄ are each as illustrated, as applicable, in any or (A1)-(A3), (B1)-(B3), (G1)-(G7), and (O1)-(Q4g), R₄ is as illustrated in (H3).


(R10) The thiazolyl ring , the Cy ring, and/or A₁, A₂, A₃, and A₄ are each as illustrated, as applicable, in any or (A1)-(A3), (B1)-(B3), (G1)-(G7), and (O1)-(Q4g), R₄ is as illustrated in (H3a).


(R11) The thiazolyl ring , the Cy ring, and/or A₁, A₂, A₃, and A₄ are each as illustrated, as applicable, in any or (A1)-(A3), (B1)-(B3), (G1)-(G7), and (O1)-(Q4g), R₄ is as illustrated in (H3b).

5 (R12) The thiazolyl ring , the Cy ring, and/or A₁, A₂, A₃, and A₄ are each as illustrated, as applicable, in any or (A1)-(A3), (B1)-(B3), (G1)-(G7), and (O1)-(Q4g), R₄ is as illustrated in (H3c).

(R13) The thiazolyl ring , the Cy ring, and/or A₁, A₂, A₃, and A₄ are each as illustrated, as applicable, in any or (A1)-(A3), (B1)-(B3), (G1)-(G7), and (O1)-(Q4g), R₄ is as illustrated in (H3d).

10 (R14) The thiazolyl ring , the Cy ring, and/or A₁, A₂, A₃, and A₄ are each as illustrated, as applicable, in any or (A1)-(A3), (B1)-(B3), (G1)-(G7), and (O1)-(Q4g), R₄ is as illustrated in any of (H3e)-(H3e-10).

15 (R15) The thiazolyl ring , the Cy ring, and/or A₁, A₂, A₃, and A₄ are each as illustrated, as applicable, in any or (A1)-(A3), (B1)-(B3), (G1)-(G7), and (O1)-(Q4g), R₄ is as illustrated in (H3f-1)-(H3f-2).






(R16) The thiazolyl ring , the Cy ring, and/or A₁, A₂, A₃, and A₄ are each as illustrated, as applicable, in any or (A1)-(A3), (B1)-(B3), (G1)-(G7), and (O1)-(Q4g), R₄ is as illustrated in (H3g)-(H3g-1).

Non-limiting illustrative compounds of the application are listed in Table 1.

Table 1

Cmpd No.	Structure	Chemical Name
1		isopropyl (<i>trans</i> -4-(5-(4-(aminomethyl)-2-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
2		isopropyl (<i>trans</i> -4-(5-(4-(acetamidomethyl)-2-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
3		isopropyl (<i>trans</i> -4-(5-(2-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl)-4-(pyrazolo[1,5- <i>a</i>]pyridin-3-yl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
4		isopropyl (<i>trans</i> -4-(5-(2-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl)-4-(1-(2,2,2-trifluoroethyl)-1 <i>H</i> -pyrazol-4-yl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
5		isopropyl (<i>trans</i> -4-(5-(2-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl)-4-(pyridin-3-yl)phenyl)thiazol-2-yl)cyclohexyl)carbamate






Cmpd No.	Structure	Chemical Name
6		isopropyl (<i>trans</i> -4-(5-(2-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl)-4-(1 <i>H</i> -pyrazol-3-yl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
7		isopropyl (<i>trans</i> -4-(5-(2-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl)-4-(6-hydroxy pyridin-3-yl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
8		isopropyl (<i>trans</i> -4-(5-(2-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl)-4-(2,4-dimethoxy pyrimidin-5-yl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
9		isopropyl (<i>trans</i> -4-(5-(2-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl)-4-(2 <i>H</i> -indazol-6-yl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
10		isopropyl (<i>trans</i> -4-(5-(4-(2-aminopyrimidin-5-yl)-2-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl) phenyl)thiazol-2-yl)cyclohexyl)carbamate

Cmpd No.	Structure	Chemical Name
11		isopropyl (<i>trans</i> -4-(5-(4-(benzofuran-5-yl)-2-(<i>N</i> -(<i>tert</i> -butyl) sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
12		isopropyl (<i>trans</i> -4-(5-(2-(<i>N</i> -(<i>tert</i> -butyl) sulfamoyl)-4-(pyrimidin-5-yl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
13		isopropyl (<i>trans</i> -4-(5-(2-(<i>N</i> -(<i>tert</i> -butyl) sulfamoyl)-4-(3,6-dimethoxypyridazin-4-yl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
14		isopropyl (<i>trans</i> -4-(5-(2-(<i>N</i> -(<i>tert</i> -butyl) sulfamoyl)-4-(2-methylthiazol-5-yl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
15		isopropyl (<i>trans</i> -4-(5-(4-(benzo[<i>d</i>]thiazol-6-yl)-2-(<i>N</i> -(<i>tert</i> -butyl) sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate






Cmpd No.	Structure	Chemical Name
16		isopropyl (<i>trans</i> -4-(5-(2-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl)-4-((dimethylamino)methyl)pyridin-3-yl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
17		isopropyl (<i>trans</i> -4-(5-(2-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl)-4-(5-methyl-1 <i>H</i> -pyrazol-4-yl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
18		isopropyl (<i>trans</i> -4-(5-(2-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl)-4-(6-cyano pyridin-3-yl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
19		isopropyl (<i>trans</i> -4-(5-(4-(1 <i>H</i> -benzo[<i>d</i>][1,2,3]triazol-6-yl)-2-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
20		isopropyl (<i>trans</i> -4-(5-(2-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl)-4-(6-(hydroxymethyl)pyridin-3-yl)phenyl)thiazol-2-yl)cyclohexyl)carbamate

Cmpd No.	Structure	Chemical Name
21		isopropyl (<i>trans</i> -4-(5-(2-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl)-4-(3-methyl-1 <i>H</i> -indol-6-yl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
22		isopropyl (<i>trans</i> -4-(5-(2-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl)-4-(2-(trifluoromethyl)pyridin-4-yl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
23		isopropyl (<i>trans</i> -4-(5-(2-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl)-4-hydroxyphenyl)thiazol-2-yl)cyclohexyl)carbamate
24		isopropyl (<i>trans</i> -4-(5-(4-hydroxy-2-sulfamoylphenyl)thiazol-2-yl)cyclohexyl)carbamate
25		isopropyl <i>N</i> -[<i>trans</i> -4-[5-[4-[2-(<i>tert</i> -butoxycarbonylamino)ethoxy]-2-(<i>tert</i> -butyl)sulfamoyl]phenyl]thiazol-2-yl]cyclohexyl)carbamate

Cmpd No.	Structure	Chemical Name
26		isopropyl (<i>trans</i> -4-(5-(4-(2-aminoethoxy)-2-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
27		isopropyl (<i>trans</i> -4-(5-(2-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl)-4-isopropoxyphenyl)thiazol-2-yl)cyclohexyl)carbamate
28		isopropyl (<i>trans</i> -4-(5-(2-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl)-4-methoxyphenyl)thiazol-2-yl)cyclohexyl)carbamate
29		<i>tert</i> -butyl (2-(3-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl)-4-(2-(<i>trans</i> -4-((iso propoxycarbonyl)amino)cyclohexyl)thiazol-5-yl)phenoxy)ethyl)(methyl)carbamate
30		isopropyl (<i>trans</i> -4-(5-(2-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl)-4-(2-(dimethylamino)ethoxy)phenyl)thiazol-2-yl)cyclohexyl)carbamate

Cmpd No.	Structure	Chemical Name
31		isopropyl (<i>trans</i> -4-(5-(2-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl)-4-(2-methoxyethoxy)phenyl)thiazol-2-yl)cyclohexyl)carbamate
32		isopropyl (<i>trans</i> -4-(5-(2-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl)-4-(2-(methylamino)ethoxy)phenyl)thiazol-2-yl)cyclohexyl)carbamate
33		isopropyl (<i>trans</i> -4-(5-(2-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl)-4-(isobutoxymethyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
34		oxetan-3-yl (<i>trans</i> -4-(5-(2-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl)-4-(isobutoxymethyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
35		isopropyl (<i>trans</i> -4-(5-(4-(dimethylamino)-2-(<i>N</i> -ethylsulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate






Cmpd No.	Structure	Chemical Name
36		isopropyl (<i>trans</i> -4-(5-(4-cyano-2-(<i>N</i> -ethylsulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
37		isopropyl (<i>trans</i> -4-(5-(3-(<i>N</i> -ethylsulfamoyl)pyridin-4-yl)thiazol-2-yl)cyclohexyl)carbamate
38		isopropyl (<i>trans</i> -4-(5-(4-((1 <i>H</i> -imidazol-2-yl)amino)-2-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
39		oxetan-3-yl (<i>trans</i> -4-(5-(4-((1 <i>H</i> -imidazol-2-yl)amino)-2-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
40		isopropyl (<i>trans</i> -4-(5-(2-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl)-4-(2-(isopropylamino)-2-oxoethyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate

Cmpd No.	Structure	Chemical Name
41		oxetan-3-yl (<i>trans</i> -4-(5-(2-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl)-4-(2-(isopropylamino)-2-oxoethyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
42		isopropyl (<i>trans</i> -4-(5-(2-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl)-4-(oxazol-2-ylamino)phenyl)thiazol-2-yl)cyclohexyl)carbamate
43		oxetan-3-yl (<i>trans</i> -4-(5-(2-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl)-4-(oxazol-2-ylamino)phenyl)thiazol-2-yl)cyclohexyl)carbamate
44		isopropyl (<i>trans</i> -4-(5-(2-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl)-4-(2-sulfamoylphenyl)thiazol-2-yl)cyclohexyl)carbamate
45		isopropyl (<i>trans</i> -4-(5-(2-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl)-4-(2-ethylsulfamoylphenyl)thiazol-2-yl)cyclohexyl)carbamate






Cmpd No.	Structure	Chemical Name
46		isopropyl (<i>trans</i> -4-(5-(2-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
47		isopropyl (<i>trans</i> -4-(5-(2-(azetidin-1-ylsulfonyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
48		3-(<i>N</i> -ethylsulfamoyl)-4-(2-(<i>trans</i> -4-((isopropoxycarbonyl)amino)cyclohexyl)thiazol-5-yl)benzoic acid
49		isopropyl (<i>trans</i> -4-(5-(4-(4-carbamoyl-2-(<i>N</i> -ethylsulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
50		isopropyl (<i>trans</i> -4-(5-(2-(<i>N</i> -ethylsulfamoyl)-4-(isopropylcarbamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
51		isopropyl (<i>trans</i> -4-(5-(2-(<i>N</i> -ethylsulfamoyl)-4-(isobutoxymethyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate

Cmpd No.	Structure	Chemical Name
52		oxetan-3-yl (<i>trans</i> -4-(5-(2-(<i>N</i> -ethylsulfamoyl)-4-(isobutoxymethyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
53		isopropyl (<i>trans</i> -4-(5-(2-(azetidin-1-ylsulfonyl)-4-(dimethylamino)phenyl)thiazol-2-yl)cyclohexyl)carbamate
54		isopropyl (<i>trans</i> -4-(5-(2-(azetidin-1-ylsulfonyl)-4-cyanophenyl)thiazol-2-yl)cyclohexyl)carbamate
55		isopropyl (<i>trans</i> -4-(5-(3-(azetidin-1-ylsulfonyl)pyridin-4-yl)thiazol-2-yl)cyclohexyl)carbamate
56		isopropyl (<i>trans</i> -4-(5-(4-((1 <i>H</i> -imidazol-2-yl)amino)-2-(<i>N</i> -ethylsulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate






Cmpd No.	Structure	Chemical Name
57		oxetan-3-yl (<i>trans</i> -4-(5-(4-((1 <i>H</i> -imidazol-2-yl)amino)-2-(<i>N</i> -ethylsulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
58		isopropyl (<i>trans</i> -4-(5-(2-(<i>N</i> -ethylsulfamoyl)-4-(2-(isopropylamino)-2-oxoethyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
59		oxetan-3-yl (<i>trans</i> -4-(5-(2-(<i>N</i> -ethylsulfamoyl)-4-(2-(isopropylamino)-2-oxoethyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
60		isopropyl (<i>trans</i> -4-(5-(2-(<i>N</i> -ethylsulfamoyl)-4-(oxazol-2-ylamino)phenyl)thiazol-2-yl)cyclohexyl)carbamate
61		oxetan-3-yl (<i>trans</i> -4-(5-(2-(<i>N</i> -ethylsulfamoyl)-4-(oxazol-2-ylamino)phenyl)thiazol-2-yl)cyclohexyl)carbamate






Cmpd No.	Structure	Chemical Name
62		oxetan-3-yl (<i>trans</i> -4-(5-(2-sulfamoylphenyl)thiazol-2-yl)cyclohexyl)carbamate
63		oxetan-3-yl (<i>trans</i> -4-(5-(2-(<i>N</i> -ethylsulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
64		oxetan-3-yl (<i>trans</i> -4-(5-(2-(<i>N</i> -(<i>tert</i> -butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
65		oxetan-3-yl (<i>trans</i> -4-(5-(2-(azetidin-1-ylsulfonyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
66		3-(<i>N</i> -ethylsulfamoyl)-4-(2-(<i>trans</i> -4-(((oxetan-3-yl)oxy)carbonyl)amino)cyclohexyl)thiazol-5-yl)benzoic acid

Cmpd No.	Structure	Chemical Name
67		oxetan-3-yl (<i>trans</i> -4-(5-(4-carbamoyl-2-(<i>N</i> -ethylsulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
68		oxetan-3-yl (<i>trans</i> -4-(5-(2-(<i>N</i> -ethylsulfamoyl)-4-(isopropylcarbamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate
69		isopropyl (<i>trans</i> - <i>N</i> -(4-(5-(2-(ethylsulfamoyl)-4-((3-isopropyl-oxetan-3-yl)amino)phenyl)thiazol-2-yl)cyclohexyl))carbamate
70		isopropyl <i>trans</i> - <i>N</i> -[4-[5-[2-(ethylsulfamoyl)-4-[(3-isobutyl-oxetan-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
71		methyl <i>trans</i> -3-(ethylsulfamoyl)-4-[2-[4-(isopropoxycarbonylamino)cyclohexyl]thiazol-5-yl]benzoate






Cmpd No.	Structure	Chemical Name
72		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-(hydroxymethyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate
73		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(1 <i>H</i> -imidazol-2-ylamino)methyl]phenyl]thiazol-2-yl]cyclohexyl]carbamate
74		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(oxazol-2-ylamino)methyl]phenyl]thiazol-2-yl]cyclohexyl]carbamate
75		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-(1 <i>H</i> -imidazol-2-yl)phenyl]thiazol-2-yl]cyclohexyl]carbamate
76		methyl <i>trans</i> -2-[3-(ethylsulfamoyl)-4-[2-[4-(isopropoxycarbonylamino)cyclohexyl]thiazol-5-yl]phenyl]acetate






Cmpd No.	Structure	Chemical Name
77		<i>trans</i> -2-[3-(ethylsulfamoyl)-4-[2-[4-(isopropoxycarbonylamino)cyclohexyl]thiazol-5-yl]phenyl]acetic acid
78		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-(1H-imidazol-2-ylmethyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate
79		isopropyl <i>trans</i> -N-[4-[5-[4-[2-(benzylamino)-2-oxo-ethyl]-2-(ethyl-sulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate
80		isopropyl <i>trans</i> -N-[4-[5-[4-[2-(isopropyl(methyl)amino)-2-oxo-ethyl]-2-(ethyl-sulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate
81		isopropyl <i>trans</i> -N-[4-[5-[4-[2-(benzyl(methyl)amino)-2-oxo-ethyl]-2-(ethyl-sulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate
82		isopropyl <i>trans</i> -N-[4-[5-[4-[2-oxo-2-(pyrrolidin-1-yl)ethyl]-2-(ethyl-sulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate

Cmpd No.	Structure	Chemical Name
83		isopropyl <i>trans</i> -N-[4-[5-[4-[2-oxo-2-(piperidin-1-yl)ethyl]-2-(ethyl-sulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate
84		isopropyl <i>trans</i> -N-[4-[5-[4-[2-oxo-2-(morpholin-4-yl)ethyl]-2-(ethyl-sulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate
85		isopropyl <i>trans</i> -N-[4-[5-[4-[2-oxo-2-(3-hydroxyazetidin-1-yl)ethyl]-2-(ethyl-sulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate
86		isopropyl <i>trans</i> -N-[4-[5-[4-[2-amino-2-oxo-ethyl]-2-(ethyl-sulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate
87		isopropyl <i>trans</i> -N-[4-[5-[4-[2-oxo-2-(azetidin-1-yl)ethyl]-2-(ethyl-sulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate






Cmpd No.	Structure	Chemical Name
88		isopropyl <i>trans</i> -N-[4-[5-(4-amino-2-pyrrolidin-1-ylsulfonyl-phenyl)thiazol-2-yl]cyclohexyl]carbamate
89		isopropyl <i>trans</i> -N-[4-[5-(4-(methylamino)-2-pyrrolidin-1-ylsulfonyl-phenyl)thiazol-2-yl]cyclohexyl]carbamate
90		isopropyl <i>trans</i> -N-[4-[5-[4-(1 <i>H</i> -imidazol-2-ylamino)-2-sulfamoyl-phenyl]thiazol-2-yl]cyclohexyl]carbamate
91		isopropyl <i>trans</i> -N-[4-[5-[2-(dimethylsulfamoyl)-4-(1 <i>H</i> -imidazol-2-ylamino)phenyl]thiazol-2-yl]cyclohexyl]carbamate
92		isopropyl <i>trans</i> -N-[4-[5-[4-(1 <i>H</i> -imidazol-2-ylamino)-2-pyrrolidin-1-ylsulfonyl-phenyl]thiazol-2-yl]cyclohexyl]carbamate






Cmpd No.	Structure	Chemical Name
93		isopropyl <i>trans</i> -N-[4-[5-(2-methylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate
94		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-(1H-imidazol-2-yl)carbamoyl]phenyl]thiazol-2-yl]cyclohexyl]carbamate
95		isopropyl <i>trans</i> -N-[4-[5-[4-pyrrolidin-1-ylcarbonyl-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate
96		oxetan-3-yl 4-(5-(2-(N-ethylsulfamoyl)-4-(2-(isopropylamino)-2-oxoethyl)phenyl)thiazol-2-yl)piperazine-1-carboxylate
97		isopropyl <i>trans</i> -N-[4-[5-[4-[2-(benzylamino)-2-oxo-ethyl]-2-(<i>tert</i> -butylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate
98		oxetan-3-yl <i>trans</i> -N-[4-[5-[4-[2-(benzylamino)-2-oxo-ethyl]-2-(<i>tert</i> -butylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate

Cmpd No.	Structure	Chemical Name
99		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-isopentyloxy-phenyl]thiazol-2-yl]cyclohexyl]carbamate
100		oxetan-3-yl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-isopentyloxy-phenyl]thiazol-2-yl]cyclohexyl]carbamate
101		isopropyl <i>trans</i> -N-[4-[5-[2-(<i>tert</i> -butylsulfamoyl)-4-isopentyloxy-phenyl]thiazol-2-yl]cyclohexyl]carbamate
102		oxetan-3-yl <i>trans</i> -N-[4-[5-[2-(<i>tert</i> -butylsulfamoyl)-4-isopentyloxy-phenyl]thiazol-2-yl]cyclohexyl]carbamate
103		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-(oxazol-2-ylmethyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate

Cmpd No.	Structure	Chemical Name
104		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-oxazol-2-yl]-phenyl]thiazol-2-yl]cyclohexyl]carbamate
105		isopropyl <i>trans</i> -N-[4-[5-[2-(1-methylimidazol-2-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
106		isopropyl <i>cis</i> -N-[4-[5-[2-(1-methylimidazol-2-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
107		isopropyl <i>trans</i> -N-[4-[5-[2-(1-isobutylimidazol-2-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
108		cyclopropyl <i>trans</i> -N-[4-[5-[4-amino-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate

Cmpd No.	Structure	Chemical Name
109		cyclopropyl <i>trans</i> - <i>N</i> -[4-[5-[2-(ethylsulfamoyl)-4-(1 <i>H</i> -imidazol-2-ylamino)phenyl]thiazol-2-yl]cyclohexyl]carbamate
110		isopropyl <i>cis</i> - <i>N</i> -[4-[5-[2-(<i>tert</i> -butylsulfamoyl)-4-(1 <i>H</i> -imidazol-2-ylamino)phenyl]thiazol-2-yl]cyclohexyl]carbamate
111		isopropyl <i>trans</i> - <i>N</i> -[4-[5-[2-(ethylsulfamoyl)-4-[(4-isopropyl-2-pyridyl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
112		isopropyl <i>trans</i> - <i>N</i> -[4-[5-[2-(ethylsulfamoyl)-4-[(6-isopropyl-2-pyridyl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
113		isopropyl <i>trans</i> - <i>N</i> -[4-[5-[2-(ethylsulfamoyl)-4-[(5-isopropyl-2-pyridyl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate

Cmpd No.	Structure	Chemical Name
114		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[3-isopropoxyazetidin-1-yl]phenyl]thiazol-2-yl]cyclohexyl]carbamate
115		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(5-isopropyl-1,3,4-thiadiazol-2-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
116		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(3-methyloxetan-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
117		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[methyl(1-methyl-1 <i>H</i> -imidazol-2-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
118		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(1,3,4-oxadiazol-2-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate

Cmpd No.	Structure	Chemical Name
119		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(1,3,4-thiadiazol-2-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
120		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(1H-pyrazol-4-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
121		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(1H-pyrazol-5-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
122		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(pyrazin-2-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
123		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(pyridazin-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate






Cmpd No.	Structure	Chemical Name
124		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(pyrimidin-4-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
125		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(pyridin-4-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
126		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(pyridin-2-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
127		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(pyrimidin-2-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
128		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(pyridin-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate







Cmpd No.	Structure	Chemical Name
129		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(1-isopropyl-1 <i>H</i> -pyrazol-5-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
130		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[4-hydroxypiperidin-1-yl]phenyl]thiazol-2-yl]cyclohexyl]carbamate
131		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(2-hydroxyethyl)(methyl))amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
132		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(2-methoxyethyl)(methyl))amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
133		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[1,1-dioxidothiomorpholino]phenyl]thiazol-2-yl]cyclohexyl]carbamate

Cmpd No.	Structure	Chemical Name
134		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[morpholino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
135		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[piperidin-1-yl]phenyl]thiazol-2-yl]cyclohexyl]carbamate
136		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(2-isopropylpyrimidin-4-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
137		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(oxetan-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
138		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(3-isopropylimidazol-4-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate

Cmpd No.	Structure	Chemical Name
139		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(1-isopropyl-1 <i>H</i> -imidazol-4-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
140		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(6-isopropylpyrazin-2-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
141		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(5-methyl-1 <i>H</i> -imidazol-2-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
142		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(5-isopropyl-1 <i>H</i> -imidazol-2-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
143		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(6-isopropylpyridazin-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
144		isopropyl <i>trans</i> -N-[4-[5-[4-(azetidin-3-ylamino)-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate


Cmpd No.	Structure	Chemical Name
145		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(1-isobutyl)azetidin-3-yl]amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
146		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[[1-(2-methylpropanoyl)azetidin-3-yl]amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
147		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(3-methylazetidin-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
148		isopropyl <i>trans</i> -N-[4-[5-[2-(azetidin-1-ylsulfonyl)-4-(1H-imidazol-2-ylamino)phenyl]thiazol-2-yl]cyclohexyl]carbamate
149		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-(2H-triazol-4-ylamino)phenyl]thiazol-2-yl]cyclohexyl]carbamate
150		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[1H-imidazol-2-yl(methyl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate

Cmpd No.	Structure	Chemical Name
151		isopropyl <i>trans</i> -N-[4-[5-[2-(<i>tert</i> -butylsulfamoyl)-4-(4 <i>H</i> -1,2,4-triazol-3-ylamino)phenyl]thiazol-2-yl]cyclohexyl]carbamate
152		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-(4 <i>H</i> -1,2,4-triazol-3-ylamino)phenyl]thiazol-2-yl]cyclohexyl]carbamate
153		isopropyl <i>trans</i> -[4-[5-[4-(oxazol-2-ylamino)-2-pyrrolidin-1-ylsulfonyl-phenyl]thiazol-2-yl]cyclohexyl]carbamate
154		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(1-isopropylimidazol-2-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
155		isopropyl <i>trans</i> -N-[4-[5-[4-[(2-chloropyrimidin-4-yl)amino]-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate

Cmpd No.	Structure	Chemical Name
156		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(2-hydroxypyrimidin-4-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
157		isopropyl <i>trans</i> -N-[4-[5-[4-[(6-ethylpyridazin-3-yl)amino]-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate
158		isopropyl <i>trans</i> -N-[4-[5-[4-[(5-ethylpyridazin-3-yl)amino]-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate
159		isopropyl <i>trans</i> -N-[4-[5-[4-[(6-chloropyridazin-3-yl)amino]-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate
160		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(6-oxo-1H-pyridazin-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
161		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(5-methylpyridazin-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate

Cmpd No.	Structure	Chemical Name
162		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(6-methylpyridazin-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
163		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(5-isopropylpyrazin-2-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
164		isopropyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(5-isopropylpyridazin-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
165		methyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(3-methyloxetan-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
166		methyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(1 <i>H</i> -pyrazol-4-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
167		methyl <i>trans</i> -N-[4-[5-[2-(ethylsulfamoyl)-4-[(1 <i>H</i> -pyrazol-5-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate

Cmpd No.	Structure	Chemical Name
168		isopropyl <i>trans</i> - <i>N</i> -[4-[5-[2-(ethylsulfamoyl)-4-[(3 <i>R</i>)-1-isobutyl-2-oxo-azetidin-3-yl]amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
169		isopropyl <i>trans</i> - <i>N</i> -[4-[5-[2-(ethylsulfamoyl)-4-[(3 <i>S</i>)-1-isobutyl-2-oxo-azetidin-3-yl]amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
170		<i>tert</i> -butyl <i>trans</i> - <i>N</i> -[4-[5-[2-(ethylsulfamoyl)-4-[(2-isopropylpyrazol-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
171		isopropyl <i>trans</i> - <i>N</i> -[4-[5-[2-(ethylsulfamoyl)-4-[(6-isopropylpyrimidin-4-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
172		isopropyl <i>trans</i> - <i>N</i> -[4-[5-[2-(ethylsulfamoyl)-4-[(1-methyl-6-oxo-pyridazin-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate
173		isopropyl <i>trans</i> - <i>N</i> -[4-[5-[2-(<i>tert</i> -butylsulfamoyl)-4-[(6-chloropyridazin-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate

Cmpd No.	Structure	Chemical Name
174		isopropyl <i>trans</i> -N-[4-[5-[2-(<i>tert</i> -butylsulfamoyl)-4-(pyridazin-3-ylamino)phenyl]thiazol-2-yl]cyclohexyl]carbamate

In some embodiments, a compound of the present application (*e.g.*, a compound of any of the formulae or any individual compounds disclosed herein) is a pharmaceutically acceptable salt. In another embodiment, a compound of the present application (*e.g.*, a compound of any of the formulae or any individual compounds disclosed herein) is a solvate. In another embodiment, a compound of the present application (*e.g.*, a compound of any of the formulae or any individual compounds disclosed herein) is a hydrate.

The compounds of the present application may form salts which are also within the scope of this application. Reference to a compound of the Formula herein is understood to include reference to salts thereof, unless otherwise indicated.

Representative "pharmaceutically acceptable salts" include, *e.g.*, water-soluble and water-insoluble salts, such as the acetate, amsonate (4,4-diaminostilbene-2,2-disulfonate), benzenesulfonate, benzonate, bicarbonate, bisulfate, bitartrate, borate, bromide, butyrate, calcium, calcium edetate, camsylate, carbonate, chloride, citrate, clavulariate, dihydrochloride, edetate, edisylate, estolate, esylate, fumerate, fiunarate, gluceptate, gluconate, glutamate, glycolylarsanilate, hexafluorophosphate, hexylresorcinate, hydrabamine, hydrobromide, hydrochloride, hydroxynaphthoate, iodide, isothionate, lactate, lactobionate, laurate, magnesium, malate, maleate, mandelate, mesylate, methylbromide, methylnitrate, methylsulfate, mucate, napsylate, nitrate, N-methylglucamine ammonium salt, 3-hydroxy-2-naphthoate, oleate, oxalate, palmitate, pamoate (1,1-methene-bis-2-hydroxy-3-naphthoate, einbonate), pantothenate, phosphate/diphosphate, picrate, polygalacturonate, propionate, p-toluenesulfonate, salicylate, stearate, subacetate, succinate, sulfate, sulfosalicylate, suramate, tannate, tartrate, teoclate, tosylate, triethiodide, and valerate salts.

"Solvate" means solvent addition forms that contain either stoichiometric or non stoichiometric amounts of solvent. Some compounds or salts have a tendency to trap a fixed

molar ratio of solvent molecules in the crystalline solid state, thus forming a solvate. If the solvent is water, the solvate formed is a hydrate; and if the solvent is alcohol, the solvate formed is an alcoholate. Hydrates are formed by the combination of one or more molecules of water with one molecule of the substance in which the water retains its molecular state as

5 H₂O.

Compounds having one or more chiral centers can exist in various stereoisomeric forms. Stereoisomers are compounds that differ only in their spatial arrangement. Stereoisomers include all diastereomeric, enantiomeric, and epimeric forms as well as racemates and mixtures thereof.

10 The term “geometric isomer” refers to cyclic compounds having at least two substituents, wherein the two substituents are both on the same side of the ring (*cis*) or wherein the substituents are each on opposite sides of the ring (*trans*). When a disclosed compound is named or depicted by structure without indicating stereochemistry, it is understood that the name or the structure encompasses one or more of the possible

15 stereoisomers, or geometric isomers, or a mixture of the encompassed stereoisomers or geometric isomers. When a geometric isomer is depicted by name or structure, it is to be understood that the named or depicted isomer exists to a greater degree than another isomer, that is that the geometric isomeric purity of the named or depicted geometric isomer is greater than 50%, such as at least 60%, 70%, 80%, 90%, 99%, or 99.9% pure by weight. Geometric

20 isomeric purity is determined by dividing the weight of the named or depicted geometric isomer in the mixture by the total weight of all of the geometric isomers in the mixture.

“Chiral isomer” means a compound with at least one chiral center. Compounds with more than one chiral center may exist either as an individual diastereomer or as a mixture of diastereomers, termed “diastereomeric mixture”. When one chiral center is present, a

25 stereoisomer may be characterized by the absolute configuration (R or S) of that chiral center. Absolute configuration refers to the arrangement in space of the substituents attached to the chiral center. The substituents attached to the chiral center under consideration are ranked in accordance with the *Sequence Rule* of Cahn, Ingold and Prelog. (Cahn *et al.*, *Angew. Chem. Inter. Edit.* 1966, 5, 385; errata 511; Cahn *et al.*, *Angew. Chem.* 1966, 78, 413; Cahn and

30 Ingold, *J. Chem. Soc.* 1951 (London), 612; Cahn *et al.*, *Experientia* 1956, 12, 81; Cahn, *J. Chem. Educ.* 1964, 41, 116).

In some embodiments of the application, the compounds of the application are diastereomers. In some embodiments, the compounds are the syn diastereomer. In other embodiments, the compounds are the anti diastereomer.

Racemic mixture means 50% of one enantiomer and 50% of its corresponding enantiomer. When a compound with one chiral center is named or depicted without indicating the stereochemistry of the chiral center, it is understood that the name or structure encompasses both possible enantiomeric forms (*e.g.*, both enantiomerically-pure, enantiomerically-enriched or racemic) of the compound. When a compound with two or more chiral centers is named or depicted without indicating the stereochemistry of the chiral centers, it is understood that the name or structure encompasses all possible diastereomeric forms (*e.g.*, diastereomerically pure, diastereomerically enriched and equimolar mixtures of one or more diastereomers (*e.g.*, racemic mixtures) of the compound.

Enantiomeric and diastereomeric mixtures can be resolved into their component enantiomers or stereoisomers by well-known methods, such as chiral-phase gas chromatography, chiral-phase high performance liquid chromatography, crystallizing the compound as a chiral salt complex, or crystallizing the compound in a chiral solvent. Enantiomers and diastereomers also can be obtained from diastereomerically- or enantiomerically-pure intermediates, reagents, and catalysts by well-known asymmetric synthetic methods.

When a compound is designated by a name or structure that indicates a single enantiomer, unless indicated otherwise, the compound is at least 60%, 70%, 80%, 90%, 99% or 99.9% optically pure (also referred to as “enantiomerically pure”). Optical purity is the weight in the mixture of the named or depicted enantiomer divided by the total weight in the mixture of both enantiomers.

When the stereochemistry of a disclosed compound is named or depicted by structure, and the named or depicted structure encompasses more than one stereoisomer (*e.g.*, as in a diastereomeric pair), it is to be understood that one of the encompassed stereoisomers or any mixture of the encompassed stereoisomers is included. It is to be further understood that the stereoisomeric purity of the named or depicted stereoisomers is at least 60%, 70%, 80%, 90%, 99% or 99.9% by weight. The stereoisomeric purity in this case is determined by dividing the total weight in the mixture of the stereoisomers encompassed by the name or structure by the total weight in the mixture of all of the stereoisomers.

It is also possible that the compounds of the application may exist in different tautomeric forms, and all such forms are embraced within the scope of the application. “Tautomer” is one of two or more structural isomers that exist in equilibrium and is readily converted from one isomeric form to another. This conversion results in the formal migration of a hydrogen atom accompanied by a switch of adjacent conjugated double bonds. Tautomers exist as a mixture of a tautomeric set in solution. In solid form, usually one tautomer predominates. In solutions where tautomerization is possible, a chemical equilibrium of the tautomers will be reached. The exact ratio of the tautomers depends on several factors, including temperature, solvent and pH. The concept of tautomers that are interconvertible by tautomerizations is called tautomerism. Of the various types of tautomerism that are possible, two are commonly observed. In keto-enol tautomerism a simultaneous shift of electrons and a hydrogen atom occurs. Ring-chain tautomerism arises as a result of the aldehyde group (-CHO) in a sugar chain molecule reacting with one of the hydroxy groups (-OH) in the same molecule to give it a cyclic (ring-shaped) form as exhibited by glucose. Common tautomeric pairs are: ketone-enol, amide-nitrile, lactam-lactim, amide-imidic acid tautomerism in heterocyclic rings (*e.g.*, in nucleobases such as guanine, thymine and cytosine), amine-enamine and enamine-imine.

The application also comprehends isotopically-labeled compounds, which are identical to those recited in the each of the formulae described herein, but for the fact that one or more atoms are replaced by an atom having an atomic mass or mass number different from the atomic mass or mass number most commonly found in nature. Examples of isotopes that can be incorporated into compounds of the application include isotopes of hydrogen, carbon, nitrogen, fluorine, such as ^3H , ^{11}C , ^{14}C , ^2H and ^{18}F .

Compounds of the application that contain the aforementioned isotopes and/or other isotopes of other atoms are within the scope of the present application. Isotopically-labeled compounds of the present application, for example those into which radioactive isotopes such as ^3H , ^{14}C are incorporated, are useful in drug and/or substrate tissue distribution assays. Tritiated, *i.e.*, ^3H , and carbon-14, *i.e.*, ^{14}C , isotopes are useful for their ease of preparation and detectability. ^{11}C and ^{18}F isotopes are useful in PET (positron emission tomography). PET is useful in brain imaging. Further, substitution with heavier isotopes such as deuterium, *i.e.*, ^2H , can afford certain therapeutic advantages resulting from greater metabolic stability, for example, increased *in vivo* half-life or reduced dosage requirements and, hence, may be preferred in some circumstances. Isotopically labeled compounds of the application, can

generally be prepared by carrying out the procedures disclosed in the Schemes and/or in the Examples described herein, by substituting a readily available isotopically labeled reagent for a non-isotopically labeled reagent. In some embodiments, the compounds of the application are not isotopically labelled.

5

Methods for Preparing the Compounds

The compounds of the present application may be made by a variety of methods, including standard chemistry. Suitable synthetic routes are depicted in the Schemes given below.

10

The compounds of Formula I may be prepared by methods known in the art of organic synthesis as set forth in part by the following synthetic schemes. In the scheme described below, it is well understood that protecting groups for sensitive or reactive groups are employed where necessary in accordance with general principles of chemistry. Protecting groups are manipulated according to standard methods of organic synthesis (T. W. Greene and P. G. M. Wuts, "Protective Groups in Organic Synthesis", Third edition, Wiley, New York 1999). These groups are removed at a convenient stage of the compound synthesis using methods that are readily apparent to those skilled in the art. The selection processes, as well as the reaction conditions and order of their execution, shall be consistent with the preparation of the compounds of the present application.

15

20

Those skilled in the art will recognize if a stereocenter exists in the compounds of Formula I. Accordingly, the present application includes both possible stereoisomers (unless specified in the synthesis) and includes not only racemic compound but the individual enantiomers and/or diastereomers as well. When a compound is desired as a single enantiomer or diastereomer, it may be obtained by stereospecific synthesis or by resolution of the final product or any convenient intermediate. Resolution of the final product, an intermediate, or a starting material may be affected by any suitable method known in the art. See, for example, "Stereochemistry of Organic Compounds" by E. L. Eliel, S. H. Wilen, and L. N. Mander (Wiley-Interscience, 1994).


25

30

The compounds described herein may be made from commercially available starting materials or synthesized using known organic, inorganic, and/or enzymatic processes. The compounds of the present application can be prepared in a number of ways well known to those skilled in the art of organic synthesis. By way of example, the compounds of the present application can be synthesized using the methods described below, together with


synthetic methods known in the art of synthetic organic chemistry, or variations thereon as appreciated by those skilled in the art. Preferred methods include but are not limited to those methods described below. The compounds of the present application (*i.e.*, a compound of Formula I) can be synthesized by following the steps outlined in General Schemes and/or General Methods below. Starting materials are either commercially available or made by known procedures in the reported literature or as illustrated.

General Scheme 1: Synthesis of Compounds VII where Cy is connected to the thiazole via a carbon-carbon bond



- 10 A carboxylic acid I is converted to the amide II using HBTU and ammonium chloride. The resulting amide is reacted with Lawesson's reagent in THF to provide the thioamide III. Cyclization with 2-bromo-1,1-diethoxyethane under acidic conditions gives the thiazole IV which is then brominated using *N*-bromosuccinimide. The bromothiazole V is then reacted under Suzuki coupling conditions with the boronate VI to give the Rad51 inhibitor VII or an
- 15 intermediate which is further transformed using conventional chemistry protocols.

General Scheme 2: Synthesis of Compounds XI where Cy is connected to the thiazole via a carbon-nitrogen bond




2,5-dibromothiazole (VIII) is reacted with a cycloalkylamine or heterocyclic amine (IX) in DMF to give the bromothiazole X. A Suzuki coupling with the boronate VI provides the Rad51 inhibitor XI or an intermediate which is further transformed using conventional chemistry protocols.

The following General Methods further illustrate the preparation of the compounds of the present application. Although certain reagents/materials and conditions are described in the General Methods, it is understood that equivalent reagents/materials and conditions may also be used.

General Method	Reaction Name
A	Amide coupling reaction
B	Borate formation
C	Suzuki Reaction A
D	Suzuki Reaction B
E	Deprotection of tert-butylsulfonamide
F	Buchwald Reaction
G	Sulfamide formation
H	Reductive amination
I	Deprotection of SEM group
J	Hydrolysis reaction
K	Substitution Reaction
L	Sandmeyer reaction

10


General Method A



To a solution of carboxylic acid (1.0 *eq.*) and DIEA (15.0 *eq.*) in DMF may be added NH₄Cl (10.0 *eq.*) and HATU (1.3 *eq.*). The mixture may be stirred at 25°C for 12 h,


then poured into H₂O and extracted with EtOAc (3x). The combined organic layers may be washed with brine, dried over Na₂SO₄, filtered, and concentrated to yield the amide product.

General Method B



- 5 A mixture of Boc-protected amine (1.0 eq.), B₂Pin₂ (3.0 eq.), KOAc (3.0 eq.) and Pd(dppf)Cl₂.CH₂Cl₂ (0.1 eq.) in dioxane may be degassed and purged with N₂ (3x) and stirred at 80°C for 12 h. The reaction mixture may be concentrated and the residue may be purified by prep-TLC (SiO₂, Petroleum ether/Ethyl acetate=3/1) to yield the borate product.

General Method C




10

A mixture of borate (1.0 eq.), thiazole (1.2 eq.), Na₂CO₃ (3.0 eq.), Pd(PPh₃)₄ (0.1 eq.) and KF (3.0 eq.) in EtOH/H₂O/Tol. may be degassed and purged with N₂ (3x) and then stirred at 80°C for 12 h. The reaction mixture may be concentrated and the residue may be purified by prep-TLC (SiO₂, Petroleum ether/Ethyl acetate=2/1) to yield the Suzuki Reaction


15 A product.

General Method D




A mixture of brominated thiazole moiety (1.0 *eq.*), 3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrazolo[1,5-*a*]pyridine (1.2 *eq.*), KOAc (3.0 *eq.*) and Pd(AmPhos)Cl₂ (0.1 *eq.*) in 1:0.25 EtOH/H₂O may be degassed and purged with N₂ (3x), and then stirred at 80°C for 12 h. The reaction mixture may be concentrated and the residue may be purified by prep-HPLC (column: Welch Ultimate AQ-C18 150*30mm*5um; mobile phase: [water(0.1%TFA)-ACN]; B%: 50%-80%, 12min) to yield the Suzuki Reaction B product.

General Method E




To a solution of *N*-(3-isopropoxyloxetan-3-yl)-2-methyl-propane-2-sulfonamide (1 *eq.*) in DCM may be added HCl/EtOAc (4 M, 4.4 *eq.*). The mixture may be stirred at 0°C for 5 min, then the reaction mixture may be diluted with MTBE then the solid may be formed. The residue may be concentrated under reduced pressure to remove solvent at 0°C. Then the residue may be triturated with MTBE:Petroleum ether = 1:1 and filtered, the filter cake may be collected to yield 3-isopropoxyloxetan-3-amine which may be used into the next step without further purification.


General Method F




A mixture of 3-isopropoxyloxetan-3-amine (2 *eq.*), brominated thiazole moiety (1 *eq.*), K₂CO₃ (3 *eq.*) and BrettPhos Pd G3 (0.1 *eq.*) in *t*-BuOH may be stirred at 110°C for 12 h under Ar atmosphere. The reaction mixture may be concentrated under reduced pressure, then the residue may be purified by prep-TLC (SiO₂, petroleum ether: ethyl acetate = 2:1) and purified by prep-HPLC (column: YMC-Actus Triart C18 100*30mm*5um; mobile phase: [water(0.1%TFA)-ACN]; B%: 25%-60%, 10min) to yield the Buchwald Reaction product.

General Method G

- To a solution of TEA (3 *eq.*) in DCM may be added alkylamine (1.1 *eq.*) at -20°C. Then the methyl 4-bromo-3-chlorosulfonylbenzoate (1.0 *eq.*) may be added batchwise. The mixture may be stirred at -20°C for 5 min, then the reaction mixture may be quenched with HCl (1M, pH=2) and extracted with DCM, the organic layers may be dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield the sulfamide formation product.

General Method H


- To a mixture of 1-(2-trimethylsilylethoxymethyl)imidazol-2-amine (1.5 *eq.*) and formyl-phenyl-thiazol moiety (1 *eq.*) in DCM may be added Ti(i-PrO)₄ (2 *eq.*) and Et₃N (0.1 *eq.*), then the mixture may be stirred at 25°C for 16 h under N₂ atmosphere. The mixture may be added NaBH₄ (1.5 *eq.*) at 0°C, and stirred at 25°C for 2 h. The reaction mixture may be quenched with H₂O at 25°C and concentrated under reduced pressure. The residue may be purified by prep-TLC (SiO₂, petroleum ether: ethyl acetate=0:1) to yield the reductive amination product.

General Method I

- A mixture of SEM protected conjugated imidazole (1 *eq.*) in TFA and DCM may be stirred at 20°C for 1 h. The reaction mixture may be concentrated under reduced pressure to remove DCM at 25°C. The residue may be diluted with EtOH and stirred at 60°C for 3 h. The

mixture may be concentrated under reduced pressure. The residue may be purified by prep-HPLC (TFA condition). (column: Welch Ultimate AQ-C18 150*30mm*5um;mobile phase: [water(0.1%TFA)-ACN];B%: 18%-48%,12min) to yield the deprotection of SEM group product.


5 General Method J



To a solution of ester (1 *eq.*) in 2:1 MeOH:H₂O, may be added LiOH (3 *eq.*). The mixture may be stirred at 50°C for 2 h. The mixture may be concentrated under reduced pressure. The residue may be adjusted pH = 2 with HCl (6M) and extracted with EtOAc (2x).


- 10 The combined organic layers may be dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield the hydrolysis reaction product.

General Method K



- 15 To a solution of amine (1 *eq.*) in n-BuOH may be added TsOH (3 *eq.*) and 2-bromo-1H-imidazole (5 *eq.*). The mixture may be stirred at 140°C for 12 h. The reaction mixture may be concentrated under reduced pressure and purified by prep-HPLC (column: Waters Xbridge BEH C18 100*30mm*10um;mobile phase: [water(0.1%TFA)-ACN];B%: 30%-90%, 10 min) to yield the substitution reaction product.

General Method L



To a solution of amine (1 *eq.*) in MeCN may be added CuBr₂ (0.4 *eq.*) and tert-butyl nitrite (1.5 *eq.*). The mixture may be stirred at 60°C for 1 h and concentrated. The residue may be diluted with H₂O and extracted with EtOAc (3x). The combined organic layers may be dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue may be purified by prep-TLC (SiO₂, petroleum ether: ethyl acetate = 1:2) to yield the Sandmeyer reaction product.

Definitions

The articles "a" and "an" are used in this application to refer to one or more than one (*i.e.*, at least one) of the grammatical object of the article. By way of example, "an element" means one element or more than one element.

The term "and/or" is used in this application to mean either "and" or "or" unless indicated otherwise.

The term "halo" or "halogen" as used herein includes fluoro, chloro, bromo, and iodo.

The term "alkyl," as used herein, refers to saturated, straight or branched-chain hydrocarbon radicals containing, in some embodiments, between one and six carbon atoms. Examples of C₁-C₈ alkyl radicals include, but are not limited to, methyl, ethyl, propyl, isopropyl, *n*-butyl, *tert*-butyl, neopentyl, *n*-hexyl, *n*-heptyl, and *n*-octyl radicals. Examples of C₁-C₆ alkyl radicals include, but are not limited to, methyl, ethyl, propyl, isopropyl, *n*-butyl, *tert*-butyl, neopentyl, and *n*-hexyl radicals.

The term "alkoxy" means an alkyl radical attached through an oxygen linking atom, represented by -O-alkyl. For example, "(C₁-C₄) alkoxy" includes methoxy, ethoxy, propoxy, and butoxy.

The terms "haloalkyl" and "haloalkoxy" means alkyl or alkoxy, as the case may be, substituted with one or more halogen atoms.

An "alkylene group" is a saturated aliphatic branched or straight-chain divalent hydrocarbon radical. Unless otherwise specified, an alkylene group typically has 1-6 carbon atoms, *e.g.*, (C₁-C₆) alkylene.

The term “cycloalkyl” means a monocyclic saturated hydrocarbon ring system. For example, C₃-C₇ cycloalkyl includes cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl, cycloheptyl.

5 A bridged cycloalkyl means a bicyclic hydrocarbon ring system in which the two rings share at least three adjacent ring carbon atoms. For example, a bridged cycloalkyl has 6-12 ring carbon atoms. Examples include, but are not limited to, bicyclo[2.1.1]hexyl, bicyclo[2.2.1]heptyl, bicyclo[2.2.2]octyl, bicyclo[3.2.1]octyl, bicyclo[4.3.1]decyl, bicyclo[3.3.1]nonyl, bornyl, bornenyl, norbornyl, norbornenyl, 6,6-dimethylbicyclo[3.1.1]heptyl, and adamantyl.

10 The terms “heterocyclyl”, “heterocyclic ring”, and “heterocyclic group”, are used interchangeably herein, and means saturated or unsaturated non-aromatic 4-10 membered ring radical containing from 1 to 4 ring heteroatoms, which may be the same or different, selected from N, O, or S. It can be monocyclic, bicyclic or tricyclic (*e.g.*, a fused or bridged bicyclic or tricyclic ring). Examples of include, but are not limited to, azetidiny, morpholinyl, thiomorpholinyl, pyrrolidinonyl, pyrrolidinyl, piperidinyl, piperazinyl, 15 hydantoinyl, valerolactamyl, oxiranyl, oxetanyl, dihydroimidazole, dihydrofuranyl, dihydropyranyl, dihydropyridinyl, dihydropyrimidinyl, dihydrothienyl, dihydrothiophenyl, dihydrothiopyranyl, tetrahydroimidazole, tetrahydrofuranyl, tetrahydropyranyl, tetrahydrothienyl, tetrahydropyridinyl, tetrahydropyrimidinyl, tetrahydrothiophenyl, and tetrahydrothiopyranyl. A heterocyclic ring optionally contains one or more double bonds 20 and/or is optionally fused with one or more aromatic rings (for example, tetrahydronaphthyridine, indolinone, dihydropyrrolotriazole, imidazopyrimidine, quinolinone, dioxaspirodecane). Examples of 3-7 membered monocyclic heterocyclic ring include, but are not limited to, azetidiny, morpholinyl, thiomorpholinyl, pyrrolidinonyl, 25 pyrrolidinyl, piperidinyl, piperazinyl, hydantoinyl, valerolactamyl, oxiranyl, oxetanyl, dihydroimidazole, dihydrofuranyl, dihydropyranyl, dihydropyridinyl, dihydropyrimidinyl, dihydrothienyl, dihydrothiophenyl, dihydrothiopyranyl, tetrahydroimidazole, tetrahydrofuranyl, tetrahydropyranyl, tetrahydrothienyl, tetrahydropyridinyl, tetrahydropyrimidinyl, tetrahydrothiophenyl, and tetrahydrothiopyranyl.

30 A bridged heterocyclyl means a bicyclic ring system containing from 1 to 4 ring heteroatoms in which the two rings share at least three adjacent ring atoms. For example, a bridged heterocyclyl has 6-12 ring atoms. Examples include, but are not limited to, azanorbornyl, quinuclidinyl, isoquinuclidinyl, tropanyl, azabicyclo[3.2.1]octanyl,

azabicyclo[2.2.1]heptanyl, 2-azabicyclo[3.2.1]octanyl, azabicyclo[3.2.1]octanyl, azabicyclo[3.2.2]nonanyl, azabicyclo[3.3.0]nonanyl, and azabicyclo [3.3.1]nonanyl.

The terms “heteroaryl”, “heteroaromatic”, “heteroaryl ring”, “heteroaryl group”, “heteroaromatic ring”, and “heteroaromatic group”, are used interchangeably herein.

- 5 “Heteroaryl”, when used alone or as part of a larger moiety as in “heteroaralkyl” or “heteroarylalkoxy”, refers to aromatic ring groups having five to ten ring atoms selected from carbon and at least one (typically 1 to 4, more typically 1 or 2) heteroatoms (*e.g.*, oxygen, nitrogen, or sulfur). “Heteroaryl” includes monocyclic rings and polycyclic rings in which a monocyclic heteroaromatic ring is fused to one or more other aromatic or
- 10 heteroaromatic rings. “Heteroaryl” includes monocyclic and bicyclic ring systems.

- “Monocyclic 5-6 membered heteroaromatic ring (or heteroaryl)” means a monocyclic heteroaromatic ring having five or six ring atoms selected from carbon and at least one (typically 1 to 3, more typically 1 or 2) heteroatoms (*e.g.*, oxygen, nitrogen, or sulfur). Examples of monocyclic 5-6 membered heteroaromatic ring groups include furanyl (*e.g.*, 2-
- 15 furanyl, 3-furanyl), imidazolyl (*e.g.*, N-imidazolyl, 2-imidazolyl, 4-imidazolyl, 5-imidazolyl), isoxazolyl (*e.g.*, 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl), oxadiazolyl (*e.g.*, 2-oxadiazolyl, 5-oxadiazolyl), oxazolyl (*e.g.*, 2-oxazolyl, 4-oxazolyl, 5-oxazolyl), pyrazolyl (*e.g.*, 3-pyrazolyl, 4-pyrazolyl), pyrrolyl (*e.g.*, 1-pyrrolyl, 2-pyrrolyl, 3-pyrrolyl), pyridyl (*e.g.*, 2-pyridyl, 3-pyridyl, 4-pyridyl), pyrimidinyl (*e.g.*, 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl),
- 20 pyridazinyl (*e.g.*, 3-pyridazinyl), thiazolyl (*e.g.*, 2-thiazolyl, 4-thiazolyl, 5-thiazolyl), isothiazolyl, triazolyl (*e.g.*, 2-triazolyl, 5-triazolyl), tetrazolyl (*e.g.*, tetrazolyl), and thienyl (*e.g.*, 2-thienyl, 3-thienyl).

- If a group is described as being “substituted,” a non-hydrogen substituent replaces a hydrogen atom on a carbon or nitrogen. Thus, for example, a substituted alkyl is an alkyl
- 25 wherein at least one non-hydrogen substituent is in the place of a hydrogen atom on the alkyl substituent. To illustrate, monofluoroalkyl is alkyl substituted with a fluoro substituent, and difluoroalkyl is alkyl substituted with two fluoro substituents. It should be recognized that if there is more than one substitution on a substituent, each non-hydrogen substituent can be identical or different (unless otherwise stated). As used herein, many moieties (*e.g.*, alkyl, cycloalkyl, or a heterocyclic ring) are referred to as being either “substituted” or “optionally
- 30 substituted”. It will be appreciated that the phrase “optionally substituted” is used interchangeably with the phrase “substituted or unsubstituted.” When a moiety is modified by one of these terms, unless otherwise noted, it denotes that any portion of the moiety that is

known to one skilled in the art as being available for substitution can be substituted, which includes one or more substituents. If more than one substituent is present, then each substituent is independently selected. Such means for substitution are well-known in the art and/or taught by the instant application. The optional substituents can be any substituents that are suitable to attach to the moiety. A person of ordinary skill in the art will recognize that the compounds and definitions provided do not include impermissible substituent patterns (*e.g.*, methyl substituted with 5 different groups, and the like). Such impermissible substitution patterns are clearly recognized by a person of ordinary skill in the art. When a group is described as being optionally substituted by “one or more” substituents, it denotes that the group is optionally substituted by one, two, three, four, five or six substituents. In some embodiments, a group is optionally substituted by 1-3 substituents. In some embodiments, a group is optionally substituted by 1-2 substituents. In some embodiments, a group is optionally substituted by one substituent.

Suitable substituents are those which do not have a significant adverse effect on the ability of the compound to inhibit RAD51. Where suitable substituents are not specifically enumerated, exemplary substituents include, but are not limited to, halo, CN, alkyl, alkoxy, halomethyl, halomethoxy, (C₁-C₅)alkyl, halo(C₁-C₅)alkyl, (C₁-C₅)alkoxy, NO₂, OR^c, NR^aR^b, S(O)_iR^a, NR^aS(O)_iR^b, S(O)_iNR^aR^b, C(=O)OR^a, OC(=O)OR^a, C(=S)OR^a, O(C=S)R^a, C(=O)NR^aR^b, NR^aC(=O)R^b, C(=S)NR^aR^b, NR^aC(=S)R^b, NR^a(C=O)OR^b, O(C=O)NR^aR^b, NR^a(C=S)OR^b, O(C=S)NR^aR^b, NR^a(C=O)NR^aR^b, NR^a(C=S)NR^aR^b, C(=S)R^a, C(=O)R^a, (C₃-C₆) cycloalkyl, monocyclic heteroaryl, and phenyl, wherein the (C₃-C₆) cycloalkyl, monocyclic heteroaryl, and phenyl substituents are optionally and independently substituted, for example, with CH₃, halomethyl, halo, methoxy, or halomethoxy. Each R^a and each R^b are independently H or (C₁-C₆) alkyl, wherein the (C₁-C₆) alkyl group represented by R^a or R^b is optionally substituted, for example, with hydroxyl or (C₁-C₃) alkoxy; R^c is H, halo(C₁-C₆) alkyl, or (C₁-C₆) alkyl, wherein the (C₁-C₆) alkyl group represented by R^c is optionally substituted, for example with hydroxyl or (C₁-C₃) alkoxy; and i is 0, 1, or 2. =O is also a suitable substituent for alkyl, cycloalkyl, and a heterocyclic ring.

Suitable substituents may also include: -F, -Cl, -Br, -I, -OH, protected hydroxy, -NO₂, -CN, -NH₂, protected amino, -NH-C₁-C₁₂-alkyl, -NH-C₂-C₁₂-alkenyl, -NH-C₂-C₁₂-alkenyl, -NH-C₃-C₁₂-cycloalkyl, -NH-aryl, -NH-heteroaryl, -NH-heterocycloalkyl, -dialkylamino, -diaryl amino, -diheteroaryl amino, -O-C₁-C₁₂-alkyl, -O-C₂-C₁₂-alkenyl, -O-C₂-C₁₂-alkenyl,

- O-C₃-C₁₂-cycloalkyl, -O-aryl, -O-heteroaryl, -O-heterocycloalkyl, -C(O)-C₁-C₁₂-alkyl, -C(O)-C₂-C₁₂-alkenyl, -C(O)-C₂-C₁₂-alkenyl, -C(O)-C₃-C₁₂-cycloalkyl, -C(O)-aryl, -C(O)-heteroaryl, -C(O)-heterocycloalkyl, -CONH₂, -CONH-C₁-C₁₂-alkyl, -CONH-C₂-C₁₂-alkenyl, -CONH-C₂-C₁₂-alkenyl, -CONH-C₃-C₁₂-cycloalkyl, -CONH-aryl, -CONH-heteroaryl, -CONH-heterocycloalkyl, -OCO₂-C₁-C₁₂-alkyl, -OCO₂-C₂-C₁₂-alkenyl, -OCO₂-C₂-C₁₂-alkenyl, -OCO₂-C₃-C₁₂-cycloalkyl, -OCO₂-aryl, -OCO₂-heteroaryl, -OCO₂-heterocycloalkyl, -OCONH₂, -OCONH-C₁-C₁₂-alkyl, -OCONH-C₂-C₁₂-alkenyl, -OCONH-C₂-C₁₂-alkenyl, -OCONH-C₃-C₁₂-cycloalkyl, -OCONH-aryl, -OCONH-heteroaryl, -OCONH-heterocycloalkyl, -NHC(O)-C₁-C₁₂-alkyl, -NHC(O)-C₂-C₁₂-alkenyl, -NHC(O)-C₂-C₁₂-alkenyl, -NHC(O)-C₃-C₁₂-cycloalkyl, -NHC(O)-aryl, -NHC(O)-heteroaryl, -NHC(O)-heterocycloalkyl, -NHCO₂-C₁-C₁₂-alkyl, -NHCO₂-C₂-C₁₂-alkenyl, -NHCO₂-C₂-C₁₂-alkenyl, -NHCO₂-C₃-C₁₂-cycloalkyl, -NHCO₂-aryl, -NHCO₂-heteroaryl, -NHCO₂-heterocycloalkyl, -NHC(O)NH₂, -NHC(O)NH-C₁-C₁₂-alkyl, -NHC(O)NH-C₂-C₁₂-alkenyl, -NHC(O)NH-C₂-C₁₂-alkenyl, -NHC(O)NH-C₃-C₁₂-cycloalkyl, -NHC(O)NH-aryl, -NHC(O)NH-heteroaryl, -NHC(O)NH-heterocycloalkyl, -NHC(S)NH₂, -NHC(S)NH-C₁-C₁₂-alkyl, -NHC(S)NH-C₂-C₁₂-alkenyl, -NHC(S)NH-C₂-C₁₂-alkenyl, -NHC(S)NH-C₃-C₁₂-cycloalkyl, -NHC(S)NH-aryl, -NHC(S)NH-heteroaryl, -NHC(S)NH-heterocycloalkyl, -NHC(NH)NH₂, -NHC(NH)NH-C₁-C₁₂-alkyl, -NHC(NH)NH-C₂-C₁₂-alkenyl, -NHC(NH)NH-C₂-C₁₂-alkenyl, -NHC(NH)NH-C₃-C₁₂-cycloalkyl, -NHC(NH)NH-aryl, -NHC(NH)NH-heteroaryl, -NHC(NH)NH-heterocycloalkyl, -NHC(NH)-C₁-C₁₂-alkyl, -NHC(NH)-C₂-C₁₂-alkenyl, -NHC(NH)-C₂-C₁₂-alkenyl, -NHC(NH)-C₃-C₁₂-cycloalkyl, -NHC(NH)-aryl, -NHC(NH)-heteroaryl, -NHC(NH)-heterocycloalkyl, -C(NH)NH-C₁-C₁₂-alkyl, -C(NH)NH-C₂-C₁₂-alkenyl, -C(NH)NH-C₂-C₁₂-alkenyl, -C(NH)NH-C₃-C₁₂-cycloalkyl, -C(NH)NH-aryl, -C(NH)NH-heteroaryl, -C(NH)NH-heterocycloalkyl, -S(O)-C₁-C₁₂-alkyl, -S(O)-C₂-C₁₂-alkenyl, -S(O)-C₂-C₁₂-alkenyl, -S(O)-C₃-C₁₂-cycloalkyl, -S(O)-aryl, -S(O)-heteroaryl, -S(O)-heterocycloalkyl -SO₂NH₂, -SO₂NH-C₁-C₁₂-alkyl, -SO₂NH-C₂-C₁₂-alkenyl, -SO₂NH-C₂-C₁₂-alkenyl, -SO₂NH-C₃-C₁₂-cycloalkyl, -SO₂NH-aryl, -SO₂NH-heteroaryl, -SO₂NH-heterocycloalkyl, -NHSO₂-C₁-C₁₂-alkyl, -NHSO₂-C₂-C₁₂-alkenyl, -NHSO₂-C₂-C₁₂-alkenyl, -NHSO₂-C₃-C₁₂-cycloalkyl, -NHSO₂-aryl, -NHSO₂-heteroaryl, -NHSO₂-heterocycloalkyl, -CH₂NH₂, -CH₂SO₂CH₃, -aryl, -arylalkyl, -heteroaryl, -heteroarylalkyl, -heterocycloalkyl, -C₃-C₁₂-cycloalkyl, polyalkoxyalkyl, polyalkoxy, -methoxymethoxy, -methoxyethoxy, -SH,

-S-C₁-C₁₂-alkyl, -S-C₂-C₁₂-alkenyl, -S-C₂-C₁₂-alkenyl, -S-C₃-C₁₂-cycloalkyl, -S-aryl, -S-heteroaryl, -S-heterocycloalkyl, or methylthiomethyl.

A "patient" or "subject" is a mammal, *e.g.*, a human, mouse, rat, guinea pig, dog, cat, horse, cow, pig, or non-human primate, such as a monkey, chimpanzee, baboon or rhesus.

5 An "effective amount" or "therapeutically effective amount" when used in connection with a compound or pharmaceutical composition is an amount effective for treating or preventing a disease in a subject as described herein.

The term "treating" with regard to a subject, refers to improving at least one symptom of the subject's disorder. Treating includes curing, improving, or at least partially
10 ameliorating the disorder.

The compounds of the present application, or a pharmaceutically acceptable salt or solvate thereof, can also be used to prevent a disease, condition or disorder. As used herein, "preventing" or "prevent" describes reducing or eliminating the onset of the symptoms or complications of the disease, condition or disorder.

15 The term "disorder" is used in this application to mean, and is used interchangeably with, the terms disease, condition, or illness, unless otherwise indicated.

As used herein, the term diseases or disorders in which RAD51 plays a role means any disease or other deleterious condition in which RAD51 is known to play a role. Accordingly, another embodiment of the present application relates to treating or lessening
20 the severity of one or more diseases in which RAD51 is known to play a role.

Pharmaceutical Compositions

The compounds disclosed therein are RAD51 inhibitors. The pharmaceutical composition of the present application comprises one or more RAD51 inhibitors, or a
25 pharmaceutically acceptable salt or solvate thereof, and a pharmaceutically acceptable carrier or diluent.

A "pharmaceutical composition" is a formulation containing the compound of the present application in a form suitable for administration to a subject. In some embodiments, the pharmaceutical composition is in bulk or in unit dosage form. The unit dosage form is any
30 of a variety of forms, including, for example, a capsule, an IV bag, a tablet, a single pump on an aerosol inhaler or a vial. The quantity of active ingredient (*e.g.*, a formulation of the disclosed compound or a pharmaceutically acceptable salt or solvate thereof) in a unit dose of composition is an effective amount and is varied according to the particular treatment

involved. One skilled in the art will appreciate that it is sometimes necessary to make routine variations to the dosage depending on the age and condition of the patient. The dosage will also depend on the route of administration. A variety of routes are contemplated, including oral, pulmonary, rectal, parenteral, transdermal, subcutaneous, intravenous, intramuscular, intraperitoneal, inhalational, buccal, sublingual, intrapleural, intrathecal, intranasal, and the like. Dosage forms for the topical or transdermal administration of a compound of this application include powders, sprays, ointments, pastes, creams, lotions, gels, solutions, patches and inhalants. In some embodiments, the active compound is mixed under sterile conditions with a pharmaceutically acceptable carrier, and with any preservatives, buffers or propellants that are required.

As used herein, the phrase "pharmaceutically acceptable" refers to those compounds, materials, compositions, carriers, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

"Pharmaceutically acceptable carrier" and "pharmaceutically acceptable diluent" refer to a substance that aids the formulation and/or administration of an active agent to and/or absorption by a subject and can be included in the compositions of the present application without causing a significant adverse toxicological effect on the subject. Non-limiting examples of pharmaceutically acceptable carriers and/or diluents include water, NaCl, normal saline solutions, lactated Ringer's, normal sucrose, normal glucose, binders, fillers, disintegrants, lubricants, coatings, sweeteners, flavors, salt solutions (such as Ringer's solution), alcohols, oils, gelatins, carbohydrates such as lactose, amylose or starch, fatty acid esters, hydroxymethylcellulose, polyvinyl pyrrolidone, and colors, and the like. Such preparations can be sterilized and, if desired, mixed with auxiliary agents such as lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, coloring, and/or aromatic substances and the like that do not deleteriously react with or interfere with the activity of the compounds provided herein. One of ordinary skill in the art will recognize that other pharmaceutical excipients are suitable for use with disclosed compounds.

The term "carrier", as used in this application, encompasses carriers, excipients, and diluents and means a material, composition or vehicle, such as a liquid or solid filler, diluent, excipient, solvent or encapsulating material, involved in carrying or transporting a

pharmaceutical agent from one organ, or portion of the body, to another organ, or portion of the body of a subject.

The pharmaceutical compositions of the present teachings optionally include one or more pharmaceutically acceptable carriers and/or diluents therefor, such as lactose, starch, cellulose and dextrose. Other excipients, such as flavoring agents; sweeteners; and preservatives, such as methyl, ethyl, propyl and butyl parabens, can also be included. More complete listings of suitable excipients can be found in the Handbook of Pharmaceutical Excipients (5th Ed., Pharmaceutical Press (2005)). A person skilled in the art would know how to prepare formulations suitable for various types of administration routes. Conventional procedures and ingredients for the selection and preparation of suitable formulations are described, for example, in Remington's Pharmaceutical Sciences (2003 - 20th edition) and in The United States Pharmacopeia: The National Formulary (USP 24 NF19) published in 1999. The carriers, diluents and/or excipients are "acceptable" in the sense of being compatible with the other ingredients of the pharmaceutical composition and not deleterious to the recipient thereof.

Pharmaceutical compositions of the application are formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, *e.g.*, intravenous, intradermal, subcutaneous, oral (*e.g.*, inhalation), transdermal (topical), and transmucosal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates, and agents for the adjustment of tonicity such as sodium chloride or dextrose. The pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

A compound or pharmaceutical composition of the application can be administered to a subject in many of the well-known methods currently used for chemotherapeutic treatment. For example, for treatment of cancers, a compound of the application may be injected directly into tumors, injected into the blood stream or body cavities or taken orally or applied through the skin with patches. The dose chosen should be sufficient to constitute effective treatment

but not as high as to cause unacceptable side effects. The state of the disease condition (*e.g.*, cancer, precancer, and the like) and the health of the patient should preferably be closely monitored during and for a reasonable period after treatment.

The term “therapeutically effective amount”, as used herein, refers to an amount of a pharmaceutical agent to treat, ameliorate, or prevent an identified disease or condition, or to exhibit a detectable therapeutic or inhibitory effect. The effect can be detected by any assay method known in the art. The precise effective amount for a subject will depend upon the subject’s body weight, size, and health; the nature and extent of the condition; and the therapeutic or combination of therapeutics selected for administration. Therapeutically effective amounts for a given situation can be determined by routine experimentation that is within the skill and judgment of the clinician.

For any compound, the therapeutically effective amount can be estimated initially either in cell culture assays, *e.g.*, of neoplastic cells, or in animal models, usually rats, mice, rabbits, dogs, or pigs. The animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans. Therapeutic/prophylactic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or experimental animals, *e.g.*, ED₅₀ (the dose therapeutically effective in 50% of the population) and LD₅₀ (the dose lethal to 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index, and it can be expressed as the ratio, LD₅₀/ED₅₀. Pharmaceutical compositions that exhibit large therapeutic indices are preferred. The dosage may vary within this range depending upon the dosage form employed, sensitivity of the patient, and the route of administration.

Dosage and administration are adjusted to provide sufficient levels of the active agent(s) or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, general health of the subject, age, weight, and gender of the subject, diet, time and frequency of administration, drug combination(s), reaction sensitivities, and tolerance/response to therapy. Long-acting pharmaceutical compositions may be administered every 3 to 4 days, every week, or once every two weeks depending on half-life and clearance rate of the particular formulation.

The pharmaceutical compositions containing active compound (*i.e.*, a compound of Formula (I)) of the present application may be manufactured in a manner that is generally known, *e.g.*, by means of conventional mixing, dissolving, granulating, dragee-making,

levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes. Pharmaceutical compositions may be formulated in a conventional manner using one or more pharmaceutically acceptable carriers comprising excipients and/or auxiliaries that facilitate processing of the active compound into preparations that can be used pharmaceutically. Of course, the appropriate formulation is dependent upon the route of administration chosen.

Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EL™ (BASF, Parsippany, N.J.) or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringeability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions can be prepared by incorporating the active compound in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle that contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, methods of preparation are vacuum drying and freeze-drying that yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

Oral compositions generally include an inert diluent or an edible pharmaceutically acceptable carrier. They can be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash, wherein the compound in the fluid carrier is applied orally and swished and expectorated or swallowed. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

For administration by inhalation, the compound is delivered in the form of an aerosol spray from pressured container or dispenser, which contains a suitable propellant, *e.g.*, a gas such as carbon dioxide, or a nebulizer.

Systemic administration can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the active compound is formulated into ointments, salves, gels, or creams as generally known in the art.

The active compound can be prepared with pharmaceutically acceptable carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to infected cells with monoclonal antibodies to viral antigens) can also be used as pharmaceutically acceptable carriers. These

can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Pat. No. 4,522,811.

It is especially advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used
5 herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the application are dictated by and directly dependent on the unique characteristics of the active compound and the particular therapeutic
10 effect to be achieved.

In therapeutic applications, the dosages of the pharmaceutical compositions used in accordance with the application vary depending on the agent, the age, weight, and clinical condition of the recipient patient, and the experience and judgment of the clinician or practitioner administering the therapy, among other factors affecting the selected dosage.
15 Generally, the dose should be sufficient to result in slowing, and preferably regressing, the growth of the tumors and also preferably causing complete regression of the cancer. Dosages can range from about 0.01 mg/kg per day to about 5000 mg/kg per day. An effective amount of a pharmaceutical agent is that which provides an objectively identifiable improvement as noted by the clinician or other qualified observer. For example, regression of a tumor in a
20 subject may be measured with reference to the diameter of a tumor. Decrease in the diameter of a tumor indicates regression. Regression is also indicated by failure of tumors to reoccur after treatment has stopped. As used herein, the term "dosage effective manner" refers to amount of an active compound to produce the desired biological effect in a subject or cell.

The pharmaceutical compositions can be included in a container, pack, or dispenser
25 together with instructions for administration.

The dosage regimen utilizing the compound is selected in accordance with a variety of factors including type, species, age, weight, sex and medical condition of the patient; the severity of the condition to be treated; the route of administration; the renal and hepatic function of the patient; and the particular compound or pharmaceutically acceptable salt or
30 solvate thereof employed. An ordinarily skilled physician or veterinarian can readily determine and prescribe the effective amount of the drug required to prevent, counter or arrest the progress of the condition.

Techniques for formulation and administration of the disclosed compound of the application can be found in *Remington: the Science and Practice of Pharmacy*, 19th edition, Mack Publishing Co., Easton, PA (1995). In an embodiment, the compound described herein, and the pharmaceutically acceptable salts or solvates thereof, are used in pharmaceutical
5 preparations in combination with a pharmaceutically acceptable carrier or diluent. Suitable pharmaceutically acceptable carriers include inert solid fillers or diluents and sterile aqueous or organic solutions. The compound or pharmaceutically acceptable salts or solvates thereof will be present in such pharmaceutical compositions in amounts sufficient to provide the desired dosage amount in the range described herein.

10 All percentages and ratios used herein, unless otherwise indicated, are by weight. Other features and advantages of the present application are apparent from the different examples. The provided examples illustrate different components and methodology useful in practicing the present application. The examples do not limit the claimed application. Based on the present application the skilled artisan can identify and employ other components and
15 methodology useful for practicing the present application.

Methods of Using the Compounds

The present application provides a method of treating a subject with a disease or disorder which can be ameliorated by inhibition of RAD51, by administering to the subject
20 an effective amount of one or more disclosed compounds, or a pharmaceutically acceptable salt or solvate thereof, or the corresponding pharmaceutical composition. Diseases which can be ameliorated by inhibition of RAD51 include treating cancer, autoimmune disease, immune deficiency, or neurodegenerative disease.

The present application further relates to a method of treating a disease or disorder in
25 which RAD51 plays a role, for example, cancer, an autoimmune disease, an immune deficiency, or a neurodegenerative disease. The method comprises administering to a subject in need thereof an effective amount of a compound disclosed herein or a pharmaceutically acceptable salt or solvate thereof or a pharmaceutical composition disclosed herein.

In one aspect, described herein is a method of treating cancer, autoimmune disease,
30 immune deficiency, or neurodegenerative disease, the method comprising administering a therapeutically effective dose of a composition as described herein, *e.g.*, a composition comprising a compound of the present application, to a subject in need of treatment for cancer, autoimmune disease, immune deficiency, or neurodegenerative disease.

In some embodiments, described herein is a method of treating cancer comprising administering a therapeutically effective dose of a composition as described herein, *e.g.*, a composition comprising a compound of the present application, to a subject in need of treatment for cancer.

5 In some embodiments, described herein is a method of treating autoimmune disease comprising administering a therapeutically effective dose of a composition as described herein, *e.g.*, a composition comprising a compound of the present application, to a subject in need of treatment for autoimmune disease.

10 In some embodiments, described herein is a method of treating immune deficiency comprising administering a therapeutically effective dose of a composition as described herein, *e.g.*, a composition comprising a compound of the present application, to a subject in need of treatment for immune deficiency.

15 In some embodiments, described herein is a method of treating neurodegenerative disease comprising administering a therapeutically effective dose of a composition as described herein, *e.g.*, a composition comprising a compound of the present application, to a subject in need of treatment for neurodegenerative disease.

20 The present application further relates to use of a compound disclosed herein, or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition disclosed herein, in the manufacture of a medicament for the treatment of a disease or disorder in which RAD51 plays a role, for example, a cancer, an autoimmune disease, an immune deficiency, or a neurodegenerative disease.

25 The present application provides use of a compound disclosed herein, or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition disclosed herein, in the manufacture of a medicament for the treatment of a disease or disorder which can be ameliorated by inhibition of RAD51.

 In one aspect, described herein is use of a compound disclosed herein, or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition disclosed herein, in the manufacture of a medicament for the treatment of a cancer, autoimmune disease, immune deficiency, or neurodegenerative disease.

30 In one aspect, described herein is use of a compound disclosed herein, or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition disclosed herein, in the manufacture of a medicament for the treatment of a cancer.

In one aspect, described herein is use of a compound disclosed herein, or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition disclosed herein, in the manufacture of a medicament for the treatment of an autoimmune disease.

5 In one aspect, described herein is use of a compound disclosed herein, or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition disclosed herein, in the manufacture of a medicament for the treatment of an immune deficiency.

10 In one aspect, described herein is use of a compound disclosed herein, or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition disclosed herein, in the manufacture of a medicament for the treatment of a neurodegenerative disease.

The present application further relates to a compound disclosed herein, or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition disclosed herein, for use in treating a disease or disorder in which RAD51 plays a role, for example, a cancer, an autoimmune disease, an immune deficiency, or a neurodegenerative disease.

15 The present application provides a compound disclosed herein, or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition disclosed herein, for use in treating a disease or disorder which can be ameliorated by inhibition of RAD51.

20 In one aspect, described herein is a compound disclosed herein, or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition disclosed herein, for use in treating a cancer, autoimmune disease, immune deficiency, or neurodegenerative disease.

25 In one aspect, described herein is a compound disclosed herein, or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition disclosed herein, for use in treating a cancer.

In one aspect, described herein is a compound disclosed herein, or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition disclosed herein, for use in treating an autoimmune disease.

30 In one aspect, described herein is a compound disclosed herein, or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition disclosed herein, for use in treating an immune deficiency.

In one aspect, described herein is a compound disclosed herein, or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition disclosed herein, for use in treating a neurodegenerative disease.

5 The present application further relates to use of a compound disclosed herein, or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition disclosed herein, in treating a disease or disorder in which RAD51 plays a role, for example, a cancer, an autoimmune disease, an immune deficiency, or a neurodegenerative disease.

10 The present application provides use of a compound disclosed herein, or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition disclosed herein, in treating a disease or disorder which can be ameliorated by inhibition of RAD51.

15 In one aspect, described herein is use of a compound disclosed herein, or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition disclosed herein, in treating a cancer, autoimmune disease, immune deficiency, or neurodegenerative disease.

In one aspect, described herein is use of a compound disclosed herein, or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition disclosed herein, in treating a cancer.

20 In one aspect, described herein is use of a compound disclosed herein, or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition disclosed herein, in treating an autoimmune disease.

In one aspect, described herein is use of a compound disclosed herein, or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition disclosed herein, in treating an immune deficiency.

25 In one aspect, described herein is use of a compound disclosed herein, or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition disclosed herein, in treating a neurodegenerative disease.

30 In some embodiments, the subject can be a subject determined to have an increased level of DNA damage occurring in one or more cell types relative to a reference level. As used herein, "DNA damage" refers to breaks, nicks, and mutations of the DNA present in a cell. In some embodiments, the DNA damage can comprise one or more of single-strand breaks (*e.g.*, "nicks"), double strand breaks (DSBs), and mutations. In some embodiments, the DNA damage can be one or more DSBs. As used herein, "mutation" refers to a change or

difference in the genetic material of a cell as compared to a reference wildtype cell, *e.g.*, a deletion, an insertion, a SNP, a gene rearrangement, and/or the introduction of an exogenous gene or sequence.

In some embodiments, the subject can be determined to have an increased level of DNA damage if the subject is determined to have an increased level and/or activity of a DNA damage process or DNA editing enzyme. As used herein, "DNA damage process" refers to any activity or process in a cell which causes one or more types of DNA damage to occur.

In some embodiments, an increased level of DNA damage can be an increased level of mutations, *e.g.*, by determining the overall mutation status in all or a portion of the genome of a cell. An overall mutation status at least 2% greater, *e.g.*, 2% greater or more, 3% greater or more, 5% greater or more, 10% greater or more, or 20% greater or more than the overall mutation status in a reference cell can be indicative of an increased, elevated, and/or significant level of a DNA editing enzyme activity. In some embodiments, the level of hyper mutations can be determined. In some embodiments, the overall mutation status in the whole genome or a portion thereof can be determined using FISH, whole genome sequencing, high throughput sequencing, exome sequencing, hybridization, and/or PCR. In some embodiments the activity of a DNA editing enzyme can be measured by determining the level of hypermutations in the specific target genes including, but not limited to IGH, BCL6, MYC, BCL1 1A, CD93, PIM1 and/or PAX5. In some embodiments the DNA editing enzyme is AID.

In some embodiments, a level of mutation in specific target genes including IGH, BCL6, MYC, BCL1 1A, CD93, PIM1 and/or PAX5 which is at least 2% greater, *e.g.*, 2% greater or more, 3% greater or more, 5% greater or more, 10% greater or more, or 20% greater or more than the level of mutation in IGH, BCL6, MYC, BCL1 1A, CD93, PIM1 and/or PAX5 in a reference cell can be indicative of an increased, elevated, and/or significant level of AID activity.

In some embodiments, an increased level of DNA damage can be an increased level of double strand breaks (DSBs). The level of DSBs can be determined, by way of non-limiting example, by karyotyping, by γ -H2AX foci formation, and/or by using FISH analysis to detect DNA double strand breaks, *e.g.*, DNA breakage detection fish (DBD-FISH) (Volpi and Bridger, BioTechniques, Vol. 45, No. 4, October 2008, pp. 385-409).

In some embodiments, an increased level of DNA damage can be an increased level of single strand breaks. The level of single-strand breaks in DNA can be determined, by way of non-limiting example, by COMET assays, FISH, or the use of single-strand break-specific

probes. Detection of DNA breaks, both single and double -stranded is known in the art and described further, at, *e.g.*, Kumari *et al.*, EXCLI Journal 2009 7:44-62 and Motalleb *et al.*, Research Journal of Applied Sciences, Engineering and Technology 2012 4: 1888-1894; each of which is incorporated by reference herein in its entirety.

5 In some embodiments, an increased level of activity of a DNA damage process can comprise an increased level and/or activity of a DNA editing enzyme. In some embodiments, the technology described herein is directed to treating cells having an active DNA editing enzyme with a compound of the present application. In some embodiments, the technology described herein is directed to treating cells having an increased level and/or activity of a
10 DNA editing enzyme with a compound of the present application. As used herein, "DNA editing enzyme" refers to an enzyme which normally catalyzes the mutation, exchange or excision of DNA segments, particularly enzymes which can generate or promote the generation of point mutations, DNA single strand breaks, DNA double-strand breaks or protein-DNA adducts. A DNA editing enzyme, as referred to herein, is not necessarily site-
15 specific in its action. Similarly, it is not necessarily cell specific. In some embodiments, the cell is a B cell expressing a detectable amount of such an enzyme.

Non-limiting examples of DNA editing enzymes include, but are not limited to Recombination Activating Gene 1 (RAG1; NCBI Gene ID: 5896), Recombination Activating Gene 1 (RAG2; NCBI Gene ID: 5897), Sporulation-specific protein 11 (SPO1 1; NCBI Gene
20 ID: 23626), APOBEC family members a Type 1 Topoisomerase; a Type 2 Topoisomerase; and/or AID. In some embodiments, the DNA editing enzyme can be AID.

In some embodiments, the DNA editing enzyme can be a member of the APOBEC (apolipoprotein B mRNA editing enzyme, catalytic polypeptide -like) family. As used herein "APOBEC family" refers to a family of cytidine deaminase enzymes having an N-terminal
25 zinc-dependent cytidine deaminase catalytic domain comprising and a C-terminal pseudocatalytic domain. Non-limiting examples of APOBEC family members include AID, APOBEC 1 (*e.g.*, NCBI Gene ID: 339), APOBEC2 (*e.g.*, NCBI Gene ID: 10930), APOBEC3A (*e.g.*, NCBI Gene ID: 200315), APOBEC3B (*e.g.*, NCBI Gene ID: 9582), APOBEC3C (*e.g.*, NCBI Gene ID: 27350), APOBEC3E (*e.g.*, NCBI Gene ID: 140564),
30 APOBEC3F (*e.g.*, NCBI Gene ID:200316), APOBEC3G (*e.g.*, NCBI Gene ID: 60489), APOBEC3H (*e.g.*, NCBI Gene ID: 164668), and APOBEC4(*e.g.*, NCBI Gene ID: 403314).

In some embodiments, the DNA editing enzyme can be a Type 1 topoisomerase. In some embodiments, the DNA editing enzyme can be a Type 2 topoisomerase.

Topoisomerases generate breaks in DNA to help uncoil or relax the strand. Type II topoisomerases hydrolyze ATP to generate DSB cuts, while Type I topoisomerases generate single-stranded breaks. Non-limiting examples of Type II topoisomerases can include topoisomerase II (*e.g.*, NCBI Gene ID: 7153 and 7155). Non-limiting examples of Type I
5 topoisomerases can include topoisomerase I (*e.g.*, NCBI Gene ID: 7150).

Embodiments of the technology described herein are based on the discovery that the compounds described herein can inhibit DNA repair mechanisms, *e.g.*, homologous recombination repair. Activation-induced cytidine deaminase (AID, or AICDA, also known as ARP2, CDA2 or HIGM2), a DNA-editing enzyme that is a member of the apolipoprotein
10 B mRNA editing enzymes, catalytic polypeptide -like (APOBEC), will cause widespread genomic breaks and cell death in cells with diminished homologous recombination ability (*e.g.*, cells with diminished DNA double strand break repair abilities). Accordingly, provided herein is a method of causing cell death comprising detecting increased expression of a DNA-
15 editing enzyme (*e.g.*, AID) in a cell and thereafter contacting the cell with a compound of the present application; thereby resulting in cell death. Accordingly, provided herein is a method of causing cell death comprising increasing expression of a DNA-editing enzyme (*e.g.*, AID) in a cell and thereafter contacting the cell with a compound of the present application;
20 thereby resulting in cell death. Accordingly, provided herein is a method of causing cell death comprising administering to a cell a therapeutically effective amount of a DNA editing enzyme (*e.g.*, AID) and thereafter contacting the cell with a compound of the present application; thereby resulting in cell death.

AID, encoded by the AICDA gene (NCBI Gene ID: 57379), is required for proper B-cell function and is most prominently expressed in centroblast B-cells. The protein is involved in somatic hypermutation, gene conversion, and class-switch recombination of
25 immunoglobulin genes. AID is normally expressed almost exclusively in antigen-activated germinal center B-cells, where it initiates immunoglobulin isotype class switching (Manis *et al.*, 2002, Trends Immunol, 23, 31-39; Chaudhuri and Alt, Nat Rev Immunol, 2004, 4, 541-552; Longerich *et al.*, Curr Opin Immunol, 2006, 18, 164-174; Chaudhuri *et al.*, Adv Immunol 2007, 94, 157-214). AID is required for somatic hypermutation and
30 immunoglobulin class switching in activated B cells. AID expression is regulated by CD40 ligand, B-cell receptor, IL4R, or Toll-like receptor stimulation (Crouch *et al.*, J Exp Med 2007 204: 1145-1156; Muramatsu *et al.*, J Biol Chem 1999 274: 18470-6). After activation, AID is transiently upregulated, induces point mutations or DNA double strand breaks in a

sequence nonspecific manner within immunoglobulin genes, and is then downregulated (Longerich *et al.*, Curr Opin Immunol, 2006, 18, 164-176; Chaudhuri *et al.*, Adv Immunol 2007, 94, 157-214). Overall, AID is active in only a tiny population of normal cells (antigen-activated B-cells) at any given time. The genomic rearrangements and mutations controlled by AID lead to the development of antigen-recognition diversity, receptor editing and lymphoid effector function required for functional adaptive immunity (Mills, *et al.*, Immunol Rev 2003 194:77-95). It has been reported that AID has off-target point mutation activities (Liu, M. *et al.*, Nature 2008, 451, 841-845; Liu and Schatz, Trends Immunol. 2009, 30, 173-181; Perez-Duran *et al.*, Carcinogenesis. 2007, 28(12):2427-33). Robbiani *et al.* has reported off-target activities of AID in B- cells, especially c-myc/IgH translocations (Robbiani *et al.*, Mol Cell 2009, 36(4):631-41). AID expression accelerates the rate of tumor development in Bcl6 transgenic mice (Pasqualucci *et al.*, 2008, Nat. Genet. 40, 108-112). However, deregulated AID does not necessarily cause malignancy or translocation-associated cancer on its own in B cells (Muto *et al.*, 2006, Proc. Natl. Acad. Sci. USA 103, 2752-2757; Okazaki *et al.*, 2003, J. Exp. Med. 197, 1173-1181; Shen *et al.*, 2008, Mol. Immunol. 45, 1883-1892). In addition, despite its role in promoting c-myc/IgH translocation, AID is not required for the development of plasmacytosis or plasmacytoma in IL-6 transgenic or pristane-treated mice, respectively (Kovalchuk *et al.*, 2007, J. Exp. Med. 204, 2989-3001; Ramiro *et al.*, 2004, J. Exp. Med. 200, 1103-1110). However, most human B cell lymphoma-associated translocations do not involve c-myc, and many do not involve Ig genes (Kuppers, 2005, Oncogene 20, 5580-5594).

Overexpression of AID has been reported in chronic lymphocytic leukemia (CLL) (Hancer *et al.*, Leuk Lymphoma. 2011 Jan;52(1):79-84; Heintel *et al.*, Leukemia. 2004 Apr; 18(4):756-62). Further, AID expression has been shown to be correlated with blast crisis B lineage leukemia and therapy resistance in myeloid leukemia and to be associated with generally poor prognosis in chronic B lymphocytic leukemia (Mao *et al.*, Br J Dermatol 2001, 145: 117-122; Chaudhuri *et al.*, Nature 2004, 430:992-8). Further expression of AID in tumor cells from a variety of cancers has been reported including but not limited to lung, breast, gastric, colon, intestinal, liver cancer and choriangiocarcinoma (Greeve *et al.*, Blood 2003, 1010, 3574-3580; Feldhahn *et al.*, J Exp Med 2007, 204, 1157-1166; Kotani *et al.*, PNAS USA 2007, 104, 1616-1620; Engels *et al.*, 2008, Appl Immunohistochem Mol Morphol 16, 521-529; Klemm *et al.*, 2009, Cancer Cell 6, 232-245; Palacios *et al.*, 2010, Blood 115(22), 4488-4496; Leuenberger *et al.*, 2009, Mod Pathol 32, 177-186; Gruber *et al.*,

2010, *Cancer Res* 70, 7411-7420; inflammatory cancer (Marusawa 2008, *Int J Biochem Cell Biol.* 40, 399-402); follicular lymphoma (Hardianti *et al.*, 2004, *Leukemia* 18, 826-831; Shikata *et al.*, 2012, *Cancer Sci.* 103(3):415-21); thyroid cancer (Qiu *et al.*, 2012, *Mod Pathol* 25(1),36-45); breast cancer (Borchert *et al.*, 2011, *BMC Cancer* 11:347); Marusawa, *et al.*,
 5 2011, *Adv Immunol* 111 : 109-41; Zhang *et al.*, 2012, *Hum Pathol* 43(3):423-34; Komori *et al.*, 2008, *Hepatology* 47(3):888-896; Hockley 2010, *Leukemia* 24(5): 1084-6; adult T-cell leukemia (Nakamura *et al.*, 2011, *Br J Dermatol.* 165(2):437-9). All of the references in the foregoing paragraph are incorporated by reference herein in their entireties.

Elevated levels of AID have been reported in arthritis (Xu *et al.*, *Scand. J. Immunol.*
 10 2009, 296, 2033-6) and in the MRL/Fas(lpr/lpr) mouse lupus model (White *et al.*, 2011, *Autoimmunity* 44(8), 585-98). All of the references in the foregoing paragraph are incorporated by reference herein in their entireties.

When DSB repair is inhibited, the extent of the DSBs generated by AID is much higher than previously suspected and the extent of genomic damage is so severe as to result in
 15 cell death. Accordingly, In some embodiments of the technology described herein, there is provided a method of treatment comprising; (a) selecting a subject having cells that express elevated levels of activation-induced cytidine deaminase (AID); and (b) administering a therapeutically effective amount of an inhibitor of double strand break repair (*e.g.*, a compound of the present application) to the subject; wherein an elevated level of AID is a
 20 level of AID that is higher than the level of AID in cells of the same type from a healthy individual. In some embodiments, the cells expressing elevated levels of AID are B cells. In some embodiments, the B cell expressing elevated levels of AID is a cancerous B cells or a B cell associated with autoimmune disease. In some embodiments, the subject can be a human subject.

25 Methods provided herein treat cancers, autoimmune disorders, immune deficiencies, or neurodegenerative disorders by inhibiting DNA double strand break repair. This inhibition proves lethal to cells expressing AID, as AID generates widespread genomic breaks, and the treatment with a double strand break repair inhibitor prevents the repair of these lesions which are being generated by the cell itself. This results in cell death in the subject which is
 30 specific to the cells expressing AID, *e.g.*, cancerous B cells and/or autoimmune cells. Accordingly, as described herein, In some embodiments there is a provided a treatment paradigm that selectively induces self-destruction of certain diseased cells, while reducing the unintended side effects in healthy tissues.

In some embodiments, an increased level and/or activity of a DNA editing enzyme can be an increased level of DNA editing enzyme mRNA. mRNA levels can be assessed using, *e.g.*, biochemical and molecular biology techniques such as Northern blotting or other hybridization assays, nuclease protection assay, reverse transcription (quantitative RT-PCR) techniques, RNA-Seq, high throughput sequencing and the like. Such assays are well known to those in the art. In some embodiments, nuclear "run-on" (or "run-off") transcription assays are used (see *e.g.*, Methods in Molecular Biology, Volume: 49, Sep-27-1995, Page Range: 229-238). Arrays can also be used; arrays, and methods of analyzing mRNA using such arrays have been described previously, *e.g.*, in EP0834575, EP0834576, W096/31622, U.S. Pat. No. 5,837,832 or WO98/30883. WO97/10365 provides methods for monitoring of expression levels of a multiplicity of genes using high density oligonucleotide arrays.

In some embodiments, a subject can be determined to have an increased level of DNA damage occurring in one or more cell types relative to a reference level if the subject has been exposed to an agent that is known to cause such DNA damage. Non-limiting examples of such agents can include a viral infection with a DNA integrating virus (*e.g.*, adeno-associated virus, retrovirus, human T-lymphotropic virus, HIV-1, oncovirus, hepatitis virus, hepatitis B virus); DNA damaging chemicals (*e.g.*, acetaldehyde, polycyclic aromatic hydrocarbons, benzenes, nitrosamines, tobacco smoke, aflatoxin, and the like); DNA damaging chemotherapeutic agents (*e.g.*, bleomycin, mitomycin, nitrogen mustards (*e.g.*, mechlorethamine, cyclophosphamide, melphalan, chlorambucil, ifosfamide and busulfan), nitrosoureas (*e.g.*, N-Nitroso-N-methylurea (MNU), carmustine (BCNU), lomustine (CCNU) and semustine (MeCCNU), fotemustine and streptozotocin), tetrazines (*e.g.*, dacarbazine, mitozolomide and temozolomide), aziridines (*e.g.*, thiotepa, mytomycin and diaziquone (AZQ)), cisplatins (*e.g.*, cisplatin, carboplatin and oxaliplatin) procarbazine and hexamethylmelamine); and ionizing or ultraviolet radiation. Exposure to such agents can be the result of an accident, infection and/or environmental exposure or the result of a therapeutic administration of such agents.

In some embodiments, the increased level of DNA damage can be occurring in a cell type affected by the cancer, autoimmune disease, immunodeficiency, and/or neurodegenerative disease. In some embodiments, the subject is determined to have an increased level of DNA damage occurring in a cell selected from the group consisting of: a cancer cell; an immune system cell; or a nervous system cell.

In some embodiments, the DNA editing enzyme can be AID. In some embodiments, the level of AID can be the level of AID in a blood cell. In some embodiments, the level of AID can be the level of AID in a B cell.

In some embodiments, an increased level of AID can be a detectable level of AID, *e.g.*, as described below herein.

In some embodiments, the subject can be a human subject.

In some embodiments, methods provided herein treat cancers and/or autoimmune disorders by inhibiting DNA double strand break repair. This inhibition proves lethal to cells expressing AID, as AID generates widespread genomic breaks, and the treatment with a double strand break repair inhibitor prevents the repair of these lesions which are being generated by the cell itself. This results in cell death in the subject which is specific to the cells expressing AID, *e.g.*, cancerous B cells and/or autoimmune cells. Accordingly, as described herein, In some embodiments there is a provided a treatment paradigm that selectively induces self-destruction of certain diseased cells, while reducing the unintended side effects in healthy tissues.

In some embodiments, methods of detecting cancers in patients with increased levels of DNA damage or increased levels of DNA editing enzymes are disclosed in WO2016/094897, incorporated herein by reference.

In some embodiments, the cancer to be treated is a type with high expression of a DNA editing enzyme. In some embodiments, the cancer to be treated is a B-cell neoplasm.

Another embodiment is a method of treating a cancer by administering to the subject an effective amount of one or more disclosed compounds, or a pharmaceutically acceptable salt or solvate thereof, or the corresponding pharmaceutical composition. In one aspect, the cancer is selected from the group consisting of lymphoma, leukemia, and a plasma cell neoplasm. In another aspect, the cancer selected from the group consisting of carcinoma and sarcoma.

In some embodiments, the cancer to be treated is a lymphoma. Lymphomas which can be treated by the disclosed methods include Non-Hodgkin's lymphoma; Burkitt's lymphoma; small lymphocytic lymphoma; lymphoplasmacytic lymphoma; MALT lymphoma; follicular lymphoma; diffuse large B-cell lymphoma; mantle cell lymphoma; and T-cell lymphoma.

Lymphoma is a malignancy in the lymphatic cells of the immune system (*e.g.*, B cells, T cells, or natural killer (NK) cells). Lymphomas often originate in the lymph nodes and present as solid tumors. They can metastasize to other organs such as the brain, bone, or skin.

Extranodal sites are often located in the abdomen. Lymphomas are closely related to the lymphoid leukemia and in some cases a particular form of cancer is categorized as both a lymphoma and a leukemia.

Leukemias which can be treated by the disclosed methods include acute
5 lymphoblastic leukemia (ALL); Burkitt's leukemia; B-cell leukemia; B-cell acute lymphoblastic leukemia; chronic lymphocytic leukemia (CLL); acute myelogenous leukemia (AML); chronic myelogenous leukemia (CML); and T-cell acute lymphoblastic leukemia (T-ALL).

In some embodiments, the cancer to be treated is B-cell neoplasms, B-cell leukemia,
10 B-cell acute lymphoblastic leukemia, chronic lymphocytic leukemia, chronic myelogenous leukemia, Burkitt's leukemia, acute myelogenous leukemia and/or T-ALL. The maturation of B cells most typically ceases or substantially decreases when the foreign antigen has been neutralized. Occasionally, however, proliferation of a particular B cell will continue unabated; such proliferation can result in a cancer referred to as "B-cell lymphoma" or a "B-
15 cell leukemia." In some embodiments the cancer to be treated is chronic lymphocytic leukemia (CLL) or chronic myelogenous leukemia (CML).

In some embodiments, the cancer to be treated is a plasma cell neoplasm. Examples for plasma cell neoplasms include multiple myeloma; plasma cell myeloma; plasma cell leukemia and plasmacytoma.

20 Carcinomas which can be treated by the disclosed methods include colon cancer; liver cancer; gastric cancer; intestinal cancer; esophageal cancer; breast cancer; ovarian cancer; head and neck cancer; lung cancer; and thyroid cancer.

Sarcomas which can be treated by the disclosed methods include soft tissue sarcoma and bone sarcoma.

25 Any cancer characterized by high levels of DNA damage and/or DNA editing enzyme expression can be treated with a compound as described herein, *e.g.*, a compound of the present application. For example, sarcomas, epithelial cell cancer (carcinomas), colon cancer, gastric cancer, intestinal cancer, liver cancer, hepatocellular cancer, breast cancer, thyroid cancer, esophageal cancer, lung cancer, brain cancer, head and neck cancer, melanoma, renal
30 cancer, prostate cancer, hemangioma, rhabdomyosarcoma, chondrosarcoma, osteosarcoma, fibrosarcoma and cholangiocarcinoma may be characterized by high levels of a DNA editing enzyme expression, *e.g.*, AID. In some embodiments the cancer to be treated is colon cancer,

liver cancer, gastric cancer, intestinal cancer, breast cancer, lung cancer, thyroid cancer and/or cholangiocarcinoma.

In some embodiments, cancers that can be treated by the disclosed methods include cancer of the bladder, blood, bone, bone marrow, brain, breast, colon, esophagus, 5 gastrointestinal, gum, head, kidney, liver, lung, nasopharynx, neck, ovary, prostate, skin, stomach, testis, tongue, or uterus. In addition, the cancer may specifically be of the following histological type, though it is not limited to these: neoplasm, malignant; carcinoma; carcinoma, undifferentiated; giant and spindle cell carcinoma; sarcomas; small cell carcinoma; papillary carcinoma; squamous cell carcinoma; lymphoepithelial carcinoma; basal 10 cell carcinoma; pilomatrix carcinoma; transitional cell carcinoma; papillary transitional cell carcinoma; adenocarcinoma; gastrinoma, malignant; cholangiocarcinoma; hepatocellular carcinoma; combined hepatocellular carcinoma and cholangiocarcinoma; trabecular adenocarcinoma; adenoid cystic carcinoma; adenocarcinoma in adenomatous polyp; adenocarcinoma, familial polyposis coli; solid carcinoma; carcinoid tumor, malignant; 15 branchiolo-alveolar adenocarcinoma; papillary adenocarcinoma; chromophobe carcinoma; acidophil carcinoma; oxyphilic adenocarcinoma; basophil carcinoma; clear cell adenocarcinoma; granular cell carcinoma; follicular adenocarcinoma; papillary and follicular adenocarcinoma; nonencapsulating sclerosing carcinoma; adrenal cortical carcinoma; endometroid carcinoma; skin appendage carcinoma; apocrine adenocarcinoma; sebaceous 20 adenocarcinoma; ceruminous adenocarcinoma; mucoepidermoid carcinoma; cystadenocarcinoma; papillary cystadenocarcinoma; papillary serous cystadenocarcinoma; mucinous cystadenocarcinoma; mucinous adenocarcinoma; signet ring cell carcinoma; infiltrating duct carcinoma; medullary carcinoma; lobular carcinoma; inflammatory carcinoma; paget's disease, mammary; acinar cell carcinoma; adenosquamous carcinoma; 25 adenocarcinoma w/squamous metaplasia; thymoma, malignant; ovarian stromal tumor, malignant; thecoma, malignant; granulosa cell tumor, malignant; androblastoma, malignant; sertoli cell carcinoma; leydig cell tumor, malignant; lipid cell tumor, malignant; paraganglioma, malignant; extra-mammary paraganglioma, malignant; pheochromocytoma; glomangiosarcoma; malignant melanoma; amelanotic melanoma; superficial spreading 30 melanoma; malig melanoma in giant pigmented nevus; epithelioid cell melanoma; blue nevus, malignant; sarcoma; fibrosarcoma; fibrous histiocytoma, malignant; myxosarcoma; liposarcoma; leiomyosarcoma; rhabdomyosarcoma; embryonal rhabdomyosarcoma; alveolar rhabdomyosarcoma; stromal sarcoma; mixed tumor, malignant; mullerian mixed tumor;

nephroblastoma; hepatoblastoma; carcinosarcoma; mesenchymoma, malignant; brenner tumor, malignant; phyllodes tumor, malignant; synovial sarcoma; mesothelioma, malignant; dysgerminoma; embryonal carcinoma; teratoma, malignant; struma ovarii, malignant; choriocarcinoma; mesonephroma, malignant; hemangiosarcoma; hemangioendothelioma, malignant; Kaposi's sarcoma; hemangiopericytoma, malignant; lymphangiosarcoma; osteosarcoma; juxtacortical osteosarcoma; chondrosarcoma; chondroblastoma, malignant; mesenchymal chondrosarcoma; giant cell tumor of bone; Ewing's sarcoma; odontogenic tumor, malignant; ameloblastic odontosarcoma; ameloblastoma, malignant; ameloblastic fibrosarcoma; pinealoma, malignant; chordoma; glioma, malignant; ependymoma; astrocytoma; protoplasmic astrocytoma; fibrillary astrocytoma; astroblastoma; glioblastoma; oligodendroglioma; oligodendroblastoma; primitive neuroectodermal; cerebellar sarcoma; ganglioneuroblastoma; neuroblastoma; retinoblastoma; olfactory neurogenic tumor; meningioma, malignant; neurofibrosarcoma; neurilemmoma, malignant; granular cell tumor, malignant; malignant lymphoma; Hodgkin's disease; hodgkin's; paragranuloma; malignant lymphoma, small lymphocytic; malignant lymphoma, large cell, diffuse; malignant lymphoma, follicular; mycosis fungoides; other specified non-Hodgkin's lymphomas; malignant histiocytosis; multiple myeloma; mast cell sarcoma; immunoproliferative small intestinal disease; leukemia; lymphoid leukemia; plasma cell leukemia; erythroleukemia; lymphosarcoma cell leukemia; myeloid leukemia; basophilic leukemia; eosinophilic leukemia; monocytic leukemia; mast cell leukemia; megakaryoblastic leukemia; myeloid sarcoma; and hairy cell leukemia.

In some embodiments, for the disclosed method, the cancer is characterized by mutations in the *mutS* homologues (*e.g.*, MSH2, MSH3, and MSH6), *mutL* homologues (*e.g.*, MLH1), or mismatch repair endonuclease PMS2. Mutations are changes in the genetic code. They include point mutations and frameshift mutations. In a point mutation, one nucleotide is swapped out for another. Therefore, the mutation occurs at a single point or location within the DNA strand. Frameshift mutations are due to either insertions or deletions of nucleotides. This causes the entire DNA strand to elongate or to shrink in size. Thus, frameshift mutations may alter all of the codons that occur after the deletion or insertion. The mutations referred to herein include, but are not limited to, insertions, deletions, duplications, inversions, or other recognized point mutations. It has now been found that RAD51 inhibitors are particularly effective in treating cancers with mutations in MSH (*e.g.*, *MSH6*), MLH, or PMS2.

MutS Homolog 2 (*MSH2*) is a protein that in humans is encoded by the *MSH2* gene, which is located on chromosome 2. *MSH2* is a tumor suppressor gene and more specifically a caretaker gene that codes for a DNA mismatch repair (MMR) protein, *MSH2*, which forms a heterodimer with *MSH6* to make the human MutS α mismatch repair complex. It also

5 dimerizes with *MSH3* to form the MutS β DNA repair complex. *MSH2* is involved in many different forms of DNA repair, including transcription-coupled repair, homologous recombination, and base excision repair. Examples of the mutations in *MSH2* include, but are not limited to, g.47630253_47630254del, g.47702411_47702421del, g.47709913_47709915inv, g.47635629_47635634del, g.47637227_47637236dup,

10 g.47639550_47639561del, g.(?_47630206)_(47710367_?)del, g.(?_47630206)_(47643569_47656880)del, g.47630263_47643568del, g.(?_47630206)_(47657081_47672686)del, g.47630263_47657080del, g.(?_47630206)_(47672797_47690169)del, g.47630263_47672796del, g.(?_47630206)_(47672797_47690169)del, g.(?_47630206)_(47693948_47698103)del,

15 g.47630263_47693947del, g.(?_47630206)_(47698202_47702163)del, g.(?_47630206)_(47630542_47635539)del, g.(?_47630206)_(47708011_47709917)del, g.(?_47630206)_(47635695_47637232)del, g.(?_47630206)_(47635695_47637232)del, g.(?_47630206)_(47637512_47639552)del, g.(?_47630206)_(47639700_47641407)del, g.(?_47630206)_(47641558_47643434)del, g.47618487_47650860delins(155),

20 g.47628578_47638433del, g.47595033_47662777del, g.47583175_47667707del, g.47625602_47636880del, g.47554933_47699909del, g.47629508_47649552del, g.47629375_47651274del, g.(?_47630206)_(47630542_47635539)del, g.(?_47630206)_(47635695_47637232)del, g.47643509_47643510del, g.47643529_47643530dup, g.47656746_47657199dup, g.47656661_47663325del,

25 g.(47643569_47656880)_(47710367_?)del, g.(47643569_47656880)_(47710367_?)del, g.47656881_47657080del, g.(47643569_47656880)_(47657081_47672686)del, g.(47643569_47656880)_(47657081_47672686)del, g.(47643569_47656880)_(47657081_47672686)del, g.(47643569_47656880)_(47657081_47672686)dup,

30 g.(47643569_47656880)_(47657081_47672686)dup, g.(47643569_47656880)_(47672797_47690169)del, g.(47643569_47656880)_(47693948_47698103)del, g.47656881_47693947del, g.(47643569_47656880)_(47702410_47703505)del, g.47656881_47656882ins(173),

- g.47656901_47656902insA, g.47656903del, g.47656912del, g.47630440del, g.47656923del,
g.47656931_47656932dup, g.47656943del, g.47656943_47656949delinsCCCAGA,
g.47656948dup, g.47656996dup, g.47657000_47657001dup, g.47630449del,
g.47657007dup, g.47657008del, g.47657020_47657023dup, g.47657025_47657026del,
5 g.47657026dup, g.47657030_47657031del, g.47657047_47657050del, g.47657053del,
g.47657053_47657057del, g.47657064del, g.47657073dup, g.47657312_47676594del,
g.47668611_47674615del, g.47672116_47675123del, g.47666463_47677632del,
g.47666403_47677572del, g.(47657081_47672686)_(47710367_?)del,
g.(47657081_47672686)_(47710367_?)inv,
10 g.47671507_47675022delinsCATTCTCTTTGAAAA, g.47657278_47676557del,
g.47672687_47672796del, g.(47657081_47672686)_(47672797_47690169)del,
g.(47657081_47672686)_(47672797_47690169)del,
g.(47657081_47672686)_(47693948_47698103)del,
g.(47657081_47672686)_(47698202_47702163)del,
15 g.(47657081_47672686)_(47708011_47709917)del, g.47672691dup, g.47672697dup,
g.47672721_47672744delins47672748_47672771inv, g.47672728_47672729del,
g.47672731dup, g.47672750_47672751insGG, g.47672755_47672758del,
g.47672762_47672763del, g.47630466_47630494del, g.47686194_47697740del,
g.(47672797_47690169)_(47710367_?)del,
20 g.(47672797_47690169)_(47690294_47693796)del,
g.(47672797_47690169)_(47693948_47698103)del, g.47690170_47693947del,
g.(47672797_47690169)_(47693948_47698103)del,
g.(47672797_47690169)_(47693948_47698103)dup,
g.(47672797_47690169)_(47705659_47707834)del, g.47690173del, g.47690191del,
25 g.47690216_47690217dup, g.47690227del, g.47690227dup, g.47690228_47690232del,
g.47690230_47690231del, g.47690240del, g.47690240_47690243del, g.47630475del,
g.47630475_47630476del, g.47690259_47690260delinsCT, g.47690277dup, g.47690280del,
g.47690283dup, g.(47690294_47693796)_(47702410_47703505)del,
g.47630484_47630485insG, g.47693838_47693839del, g.47693862del, g.47693864del,
30 g.47693873del, g.47693880dup, g.47693913del, g.47693924_47693925dup, g.47630493del,
g.47697730_47706125del, g.(47693948_47698103)_(47710367_?)del,
g.(47693948_47698103)_(47698202_47702163)del,
g.(47693948_47698103)_(47705659_47707834)del, g.47698107del, g.47698109del,

- g.47698109_47698110insA, g.47630496del, g.47698118del, g.47698125del, g.47698129dup,
g.47698138_47698139del, g.47698142_47698146del, g.47698144dup,
g.47698147_47698148del, g.47698147_47698148dup, g.47698147_47698148insT,
g.47698159del, g.47698162del, g.47698506_47703472del, g.47701803_47708848del,
5 g.(47698202_47702163)_(47710367_?)del,
g.(47698202_47702163)_(47702410_47703505)del,
g.(47698202_47702163)_(47703711_47705410)del,
g.(47698202_47702163)_(47705659_47707834)del, g.47702164del,
g.47702175_47702176insA, g.47702183_47702186del, g.47702185_47702186insCT,
10 g.47702190_47702192del, g.47702191dup, g.47702192_47702193del, g.47702213del,
g.47702231del, g.47702242dup, g.47702257del, g.47702262_47702263dup,
g.47630516_47630517dup, g.47630517del, g.47630517dup, g.47702289_47702290inv,
g.47702293_47702296del, g.47702301dup, g.47702315del, g.47702315del,
g.47702328_47702329del, g.47630522dup, g.47702339del, g.47702371_47702374dup,
15 g.47702384_47702385del, g.47702386_47702389del, g.47702388del,
g.47702388_47702389del, g.47702390del, g.47702390_47702391del,
g.47702400_47702401del, g.47703506_47703710del, g.47703506_47708010del,
g.47703510del, g.47703515del, g.47703521_47703522del, g.47703535_47703536del,
g.47703546_47703547del, g.47703548_47703611dup, g.47630534del, g.47703571dup,
20 g.47703574_47703581del, g.47703585dup, g.47630350del, g.47632107_47668733del,
g.47703613del, g.(47630542_47635539)_(47643569_47656880)del,
g.(47630542_47635539)_(47643569_47656880)inv,
g.(47630542_47635539)_(47657081_47672686)del, g.47635540_47657080del,
g.(47630542_47635539)_(47672797_47690169)del,
25 g.(47630542_47635539)_(47690294_47693796)del,
g.(47630542_47635539)_(47705659_47707834)del, g.47635540_47635694del,
g.(47630542_47635539)_(47635695_47637232)del,
g.(47630542_47635539)_(47635695_47637232)del,
g.(47630542_47635539)_(47637512_47639552)del, g.47703635dup, g.47703641dup,
30 g.47635542_47635549del, g.47703660_47703663del, g.47703667dup, g.47630351dup,
g.47703704del, g.47703826_47707938del,
g.(47703711_47705410)_(47705659_47707834)del, g.47705428_47705431del,
g.47705437_47705438insA, g.47635551_47635552del, g.47705440_47705441del,

- g.47705461del, g.47705490del, g.47705494del, g.47705495del, g.47635557_47635558del,
g.47705505del, g.47705535dup, g.47705547del, g.47705560_47705561dup, g.47705561dup,
g.47705562dup, g.47705588del, g.47705608_47705609del, g.47705618dup, g.47705627dup,
g.47635571_47635601delins(217), g.(47705659_47707834)_(47710367_?)del,
5 g.(47705659_47707834)_(47708011_47709917)del, g.47707842_47707843del,
g.47707861del, g.47707861_47707874dup, g.47707878_47707884del,
g.47707878_47707884del, g.47707883del, g.47707895_47707905del, g.47707897del,
g.47707901_47707902del, g.47707905_47707906del, g.47707921del, g.47635583dup,
g.47635583_47635584del, g.47707969_47707973del, g.47707996_47707997ins(115),
10 g.47708009_47708010del, g.(47708011_47709917)_(47710367_?)del,
g.47635591_47635592del, g.47635597_47635618dup, g.47635606_47635607del,
g.47630359dup, g.47635672del, g.47635675_47635678del, g.47630364dup, g.47635680dup,
g.47636862_47639040del, g.47636781_47638831del, g.47636753_47638155del,
g.47636552_47638597del, g.(47635695_47637232)_(47643569_47656880)del,
15 g.(47635695_47637232)_(47643569_47656880)del,
g.(47635695_47637232)_(47657081_47672686)del,
g.(47635695_47637232)_(47672797_47690169)del,
g.(47635695_47637232)_(47698202_47702163)del,
g.(47635695_47637232)_(47637512_47639552)del,
20 g.(47635695_47637232)_(47641558_47643434)del, g.47637234del,
g.47637246_47637247del, g.47637253_47637254del, g.47637254_47637255del,
g.47637254_47637255del, g.47637265del, g.47637274del, g.47637282del, g.47637320del,
g.47637372_47637375del, g.47637377_47637449dup, g.47637379del, g.47637384del,
g.47637394_47637395del, g.47637396_47637397del, g.47637417del,
25 g.47637427_47637435del, g.47637437_47637439del, g.47637453del, g.47637458dup,
g.47637479_47637482dup, g.47637482dup, g.47637504_47637505del,
g.47637508_47637511del, g.47638050_47653430del, g.47638302_47648462del,
g.47638478_47648643del, g.(47637512_47639552)_(47710367_?)del,
g.(47637512_47639552)_(47643569_47656880)del, g.47639553_47643568del,
30 g.(47637512_47639552)_(47657081_47672686)del,
g.(47637512_47639552)_(47657081_47672686)del,
g.(47637512_47639552)_(47672797_47690169)del,
g.(47637512_47639552)_(47639700_47641407)del,

- g.(47637512_47639552)_(47641558_47643434)del, g.47639557_47639561del,
 g.47639582_47639586delinsTAAT, g.47639583_47639584del, g.47639594del,
 g.47639594dup, g.47639598del, g.47639603_47639604del, g.47639611_47639612del,
 g.47639612del, g.47639618_47639621del, g.47639624_47639628delinsTTA,
 5 g.47630401dup, g.47639632dup, g.47639638_47639641dup, g.47639638_47639641dup,
 g.47639639del, g.47639639del, g.47639642dup, g.47630403_47630404insC, g.47639653del,
 g.47639666del, g.47639666_47639669del, g.47639668del, g.47639670_47639673delinsTT,
 g.47639674_47639675dup, g.47639695_47639696del, g.47639707_47642985del,
 g.47641402_47642007del, g.(47639700_47641407)_(47643569_47656880)del,
 10 g.47641408_47643568del, g.(47639700_47641407)_(47657081_47672686)del,
 g.(47639700_47641407)_(47672797_47690169)del,
 g.(47639700_47641407)_(47641558_47643434)del,
 g.(47639700_47641407)_(47641558_47643434)del, g.47641410del,
 g.47641425_47641426del, g.47641426_47641429del, g.47630412del, g.47641451del,
 15 g.47641454dup, g.47641455dup, g.47641469del, g.47641478del, g.47641488_47641491del,
 g.47641496_47641497del, g.47641503del, g.47641513_47641514dup,
 g.47641530_47641537dup, g.47642509_47655432del,
 g.(47641558_47643434)_(47643569_47656880)del,
 g.(47641558_47643434)_(47693948_47698103)del, g.47630424_47630433del,
 20 g.47643450dup, g.47643462_47643463del, g.47643462_47643463ins(4),
 g.47643464_47643465insNC_000022.10:35788169_35788352, g.47643465dup.

MutS Homolog 3 (*MSH3*) is a human homologue of the bacterial mismatch repair protein MutS that participates in the mismatch repair (MMR) system. MSH3 typically forms the heterodimer MutS β with MSH2 in order to correct long insertion/deletion loops and base-
 25 base mispairs in microsatellites during DNA synthesis. Deficient capacity for MMR is found in approximately 15% of colorectal cancers, and somatic mutations in the *MSH3* gene can be found in nearly 50% of MMR-deficient colorectal cancers. Examples of the mutations in *MSH3* include, but are not limited to, g.79970809del.

MSH6 encodes MutS homologue 6 (MSH6), a member of the Mutator S (MutS)
 30 family of proteins that are involved in DNA mismatch repair (MMR). The MSH6 protein forms a heterodimer with MutS homologue 2 (MSH2) in both human and yeast. Human MSH2/6 recognizes single base-base mismatches and short insertion/deletion loops. Upon recognition of a mismatch, MSH2/6 complex binds and exchanges ADP for ATP, resulting in

a conformational change to the complex that precedes base pair dissolution, base excision, and repair.

MSH6 mutations include frameshift and/or nonsense mutations and can result in non-functional *MSH6* and loss of protein expression. Examples include a frameshift mutation at
5 *MSH6* amino acid residue 290 and a compounding missense T1189I.

Inactivating *MSH6* mutations can be detected in cancers by routine diagnostics methods. These methods include, but are not limited to, obtaining cancer cells and other diagnostic indicators such as peripheral blood mononuclear cells (PBMCs), PBMC subpopulations, circulating blasts (CD34+ cells), circulating tumor cells and circulating
10 exosomes/cancer cells by biopsy and blood tests and by obtaining lymphatic or other bodily fluids. It is then determined from the cancer cells or other diagnostic indicators whether the cancer exhibits an inactivating *MSH6* mutation is by methodology known in the art, for example, direct DNA sequencing and multiplex ligation dependent probe amplification, RNA sequencing (RNA-Seq), microarray, quantitative PCR, or NanoString™ gene expression
15 panels, or *MSH6* protein by immunohistochemistry, flow cytometry, immunocytochemistry or Western blot. Methods for identifying inactivating *MSH6* mutations are disclosed in Houlleberghs H, Goverde A, Lusseveld J, Dekker M, Bruno MJ, *et al.* (2017) Suspected Lynch syndrome associated *MSH6* variants: A functional assay to determine their pathogenicity. PLOS Genetics 13(5):
20 e1006765. <https://doi.org/10.1371/journal.pgen.1006765>.

Examples of the mutations in *MSH6* include, but are not limited to,
g.48032846_48032849del, g.48032846_48032849del, g.48032846_48032849del,
g.48033337_48033342del, g.48033420_48033422del, g.(?_48010221)_(48034092)del,
g.(?_48010221)_(48018263_48023032)del, g.47998510_48020183del,
25 g.48007276_48020272del, g.48026207del, g.48026223del, g.48026223del,
g.48026257_48026261del, g.48026261_48026265del, g.48026312_48026313del,
g.48026398del, g.48026543_48026544dup, g.48026693dup, g.48026702del, g.48026712del,
g.48026718dup, g.48026736_48026737delinsAG, g.48026736_48026737delinsG,
g.48026750_48026751del, g.48026754_48026757del, g.48026756_48026759del,
30 g.48026759_48026760del, g.48026906del, g.48026928_48026931del, g.48026941dup,
g.48026991del, g.48027023_48027024del, g.48027079del, g.48027079_48027082dup,
g.48027167_48027168del, g.48027172_48027173dup, g.48027178_48027185del,
g.48027184_48027185del, g.48027272_48027275del, g.48027470_48027471del,

- g.48027501_48027502del, g.48027501_48027502delTG, g.48027657dup,
 g.48027691_48027694del, g.48027733_48027736dup, g.48027794_48027796delinsC,
 g.48027841_48027842del, g.48027887del, g.48027890dup, g.48027973_48027980del,
 g.48028067del, g.48028098del, g.48028106del, g.48028175_48028176del,
 5 g.48028241_48028242del, g.48028241_48028242delTT, g.48028272_48028284dup,
 g.48028277_48028278del, g.48030558_48030559del, g.48030126_48032394del,
 g.48030568del, g.48030581_48030584del, g.48030584_48030585dup, g.48030607del,
 g.48030645_48030646insT, g.48030647del, g.48030647dup, g.48030649dup,
 g.48030654_48030660del, g.48030659dup, g.48030697_48030698del, g.48030698del,
 10 g.48030706del, g.48030710dup, g.48030727_48030728insC, g.48030765_48030829del,
 c.3438+797_3438+798insTATins1839_3439-428,
 c.3438+797_3438+798insTATins1839_3439-428, g.48032121_48032122del,
 g.48032123_48032124del, g.48032124dup, g.48032126_48032129del,
 g.48032129_48032130insA, g.48032129_48032132dup,
 15 g.(48032167_48032756)_(48034092_?)del, g.48032809_48032812del, g.48032835dup,
 g.48032846_48032849del, g.48033374_48033402dup, g.48033395_48033398del,
 g.48033421_48033433del, g.48033425_48033428dup, g.48033453_48033454insA,
 g.48033494_48033523del, g.48033495_48033496del, g.48033593dup,
 g.48033610_48033613dup, g.48033629_48033635del, g.48033636_48033639dup,
 20 g.48033676_48033682del, g.48033707dup, g.48033709_48033716dup,
 g.48033721_48033724dup, g.48033727_48033730dup, g.48033728_48033746dup,
 g.(48033742_48033743)_(48033742_48033743)ins(32), g.48033746dup,
 g.48033748_48033751del, g.48033758_48033768del, g.48033773_48033774insATCA,
 g.48033773_48033776dup, g.48033785_48033789dup, g.48033887_48033910inv,
 25 g.(48018263_48023032)_(48032167_48032756)del,
 g.(48018263_48023032)_(48023203_48025749)del, g.48023097_48023098del,
 g.48025773dup, g.48025832del, g.48025860_48025861insT, g.48025884_48025885del,
 g.48025967dup.

- MutL homolog 1, colon cancer, nonpolyposis type 2 (E. coli) is a protein that in
 30 humans is encoded by the MLH1 gene located on Chromosome 3. It is a gene commonly
 associated with hereditary nonpolyposis colorectal cancer.

Examples of the mutations in *MSH6* include, but are not limited to,
 g.37089113_37089115del, g.37089175del, g.37090379_37090393del,

- g.37038201_37038202del, g.37042531_37042542del, g.37053339_37053355del,
g.37053354del, g.37053590_37053591insT, g.37034841_37092337del,
g.(?_37034841)_37092337?)del, g.(?_37034841)_37061955_37067127del,
g.(?_37034841)_37035155_37038109del, g.(?_37034841)_37035155_37038109del,
5 g.(?_37034841)_37070424_37081676del, g.(?_37034841)_37083823_37089009del,
g.37034841_37083822del, g.(?_37034841)_37038201_37042445del,
g.(?_37034841)_37042545_37045891del, g.37034841_37042544del,
g.(?_37034841)_37042545_37045891del, g.(?_37034841)_37042545_37045891del,
g.(?_37034841)_37045966_37048481del, g.(?_37034841)_37050397_37053310del,
10 g.(?_37034841)_37059091_37061800del, g.37034658_37038806del,
g.36961079_37138741del, g.37061923del, g.37061927del, g.37061933del, g.37061939del,
g.37061942dup, g.37035140_37035141del, g.37070417del, g.37070417_37070418insT,
g.37070419dup, g.37070422_37070423insT, g.37080355_37083368del,
g.(37070424_37081676)_37092337?)del,
15 g.(37070424_37081676)_37081786_37083758del,
g.(37070424_37081676)_37083823_37089009del, g.37038148_37038151del,
g.37038149del, g.37038149dup, g.37081690_37081691del, g.37081691_37081692del,
g.37081706_37081708del, g.37081710_37081711del, g.37035053_37035066del,
g.37038154del, g.37038154_37038157del, g.37081738_37081739del, g.37081740del,
20 g.37081753dup, g.37081757_37081761dup, g.37081782_37081783insAAGT,
g.37081787_37081793delinsATTT, g.(37081786_37083758)_37083823_37089009del,
g.(37081786_37083758)_37089175_37090007del, g.37083759del, g.37083780dup,
g.37083781_37083784del, g.37083781_37083784delCTCA, g.37083808_37083809del,
g.37083816del, g.37086069_37089606del, g.37084092_37089247del,
25 g.37084590_37089786del, g.(37083823_37089009)_37092337?)del,
g.(37083823_37089009)_37089175_37090007del, g.37089010_37089174del,
g.(37083823_37089009)_37090509_37091976del, g.37089023del,
g.37089026_37089027del, g.37089027del, g.37089036del, g.37089036dup, g.37038168dup,
g.37089042del, g.37089047del, g.37089050_37089053del, g.37089056_37089057del,
30 g.37089061_37089062del, g.37089078_37089096del, g.37089090dup, g.37089099dup,
g.37089107_37089110dup, g.37089109_37089110del, g.37089130_37089132del,
g.37089130_37089132delAAG, g.37089131delinsTTCTT, g.37089133del, g.37089133delG,
g.37089144del, g.37089155del, g.37089155_37089161del, g.37089158_37089161del,

- g.37089162_37089166del, g.37089171del,
 g.(37089175_37090007)_(37090101_37090394)del, g.37035056_37035072del,
 g.37090013del, g.37090015dup, g.37038183_37038184del, g.37090024_37090037dup,
 g.37090025_37090053dup, g.37090027dup, g.37038184dup, g.37090031_37090032insT,
 5 g.37090041del, g.37090057del, g.37090064_37090067del, g.37038188del, g.37090082del,
 g.37090086_37090087del, g.37090087_37090088del, g.37090097_37090101delinsC,
 g.37090099del, g.37038191dup, g.(37090101_37090394)_(37092337_?)del,
 g.37035057_37035073del, g.37090405dup, g.37090411_37090415del, g.37090414del,
 g.37038194del, g.37038198del, g.37090472_37090478del, g.37039445_37059613dup,
 10 g.37039760_37052440del, g.37090481_37090482del, g.37090483_37090484del,
 g.37090483_37092045del, g.37040732_37043185delinsACATAGTA,
 g.37042445_37042446del, g.(37038201_37042445)_(37042545_37045891)del,
 g.(37038201_37042445)_(37048555_37050304)del,
 g.(37038201_37042445)_(37050397_37053310)del,
 15 g.(37038201_37042445)_(37053591_37055922)del, g.37090497_37090498del,
 g.37090497_37090498delTC, g.37090504_37090507del,
 g.(37090509_37091976)_(37092337_?)del, g.(37090509_37091976)_(37092337_?)dup,
 g.37091977_37091978del, g.37091978_37091987del, g.37042448_37042451del,
 g.37091984_37091990del, g.37042451_37042453del, g.37092020_37092021del,
 20 g.37092022_37092068dup, g.37092027_37092028del, g.37092027_37092028dup,
 g.37092030dup, g.37092052_37092055del, g.37092054_37092055del,
 g.37092068_37092071dup, g.37092091dup, g.37092094_37092097delins(30),
 g.37092096_37092106del, g.37092097del, g.37092125_37092126delAA,
 g.37092125_37092126del, g.37092139_37092142dup, g.37092142dup, g.37035060dup,
 25 g.37042469_37042470del, g.37042470del, g.37042482dup, g.37042485del, g.37042499del,
 g.37042546dup, g.37044472_37046589del, g.37045648_37049941del,
 g.37045095_37054651del, g.37045072_37046861del,
 g.(37042545_37045891)_(37045966_37048481)del,
 g.(37042545_37045891)_(37092337_?)del,
 30 g.(37042545_37045891)_(37048555_37050304)del,
 g.(37042545_37045891)_(37050397_37053310)del, g.37045892_37050396del,
 g.37035069del, g.37045926del, g.37045931del, g.37045939_37045940dup,
 g.37045957_37045958del, g.37045963del, g.37035075del, g.37048067_37049287del,

- g.(37045966_37048481)_(37048555_37050304)del,
g.(37045966_37048481)_(37050397_37053310)del, g.37048483del,
g.37048483_37048503delinsT, g.37048486_37048487delinsGTT, g.37048489del,
g.37048490del, g.37035076_37035077insCCCA, g.37035077_37035078dup,
5 g.37048505_37048508del, g.37048521del, g.37048529dup, g.37035082dup,
g.37049873_37052281del, g.37049839_37052249del, g.37049800_37052209del,
g.37049640_37050445del, g.37050305_37050396del,
g.(37048555_37050304)_(37050397_37053310)del, g.37050305_37050396del,
g.37050319_37050320del, g.37050339del, g.37050348del, g.37050353_37050354del,
10 g.37050354dup, g.37050364del, g.37050375_37050376insGA, g.37035090del,
g.37050382_37050383delinsAT, g.37050382_37050383delinsCT, g.37050390_37050396del,
g.37052950_37060990del, g.(37050397_37053310)_(37067499_37070274)dup,
g.(37050397_37053310)_(37053591_37055922)del,
g.(37050397_37053310)_(37056036_37058996)del, g.37053353del,
15 g.37053510_37053511del, g.37035099del, g.37053545_37053546insT, g.37053562del,
g.37053578del, g.37053578dup, g.37053585del, g.37053586_37053589del, g.37053591del,
g.37053590_37053591delinsAT, g.37055920_37055921del, g.37055914_37055938del,
g.(37053591_37055922)_(37070424_37081676)del,
g.(37053591_37055922)_(37083823_37089009)del,
20 g.(37053591_37055922)_(37059091_37061800)del, g.37035105del, g.37055928dup,
g.37035106_37035116del, g.37055938del, g.37035108del, g.37055972_37055975del,
g.37055976_37055979del, g.37035111del, g.37055990dup, g.37035114del, g.37035116del,
g.37056036del, g.37056037dup, g.37058993_37059001del,
g.(37056036_37058996)_(37070424_37081676)del,
25 g.(37056036_37058996)_(37059091_37061800)del, g.37058997_37059000del,
g.37059014_37059017del, g.37059017_37059021del, g.37059027_37059030dup,
g.37035122del, g.37059062_37059063insT, g.37059065_37059066del, g.37059066del,
g.37059066dup, g.37059072_37059073del, g.37059072_37059073dup,
g.37059090_37059093del, g.37061595_37061913del, g.37061308_37066756del,
30 g.37061207_37063077del, g.(37059091_37061800)_(37092337_?)del,
g.(37059091_37061800)_(37061955_37067127)del, g.37061801_37061954del,
g.(37059091_37061800)_(37083823_37089009)del, g.37061803dup, g.37061804del,
g.37061817del, g.37061837_37061838dup, g.37061844del, g.37061851dup, g.37061855dup,

g.37061870del, g.37061904_37061906del, g.37061910del, g.37035047del,
g.[37049179_37051317delinsTG;37051667_37054327delinsCA].

Human *PMS2* related genes are located at bands 7p12, 7p13, 7q11, and 7q22. Exons 1 through 5 of these homologues share high degree of identity to human *PMS2*. The product of this gene is involved in DNA mismatch repair. The protein forms a heterodimer with *MLH1* and this complex interacts with *MSH2* bound to mismatched bases. Defects in this gene are associated with hereditary nonpolyposis colorectal cancer, with Turcot syndrome, and are a cause of supratentorial primitive neuroectodermal tumors.

Examples of the mutations in *PMS2* include, but are not limited to,

- 10 g.(?_6012870)_(6048737_?)del, g.6012870_6048737del,
g.(6027252_6029430)_(6048737_?)del, g.(6045663_6048627)_(6048737_?)del,
g.6029554del, g.6029499dup, g.6029495_6029496del, g.6029462_6029463delinsTAAA,
g.5992485_6028601del, g.(6018328_6022454)_(6027252_6029430)del,
g.(6013174_6017218)_(6027252_6029430)del, g.6027226_6027227ins(20), g.6027175del,
15 g.6027090dup, g.6036705_6044207delinsCG, g.6026666dup, g.6026628del, g.6043671del,
g.6026565dup, g.6026565dupT, g.6018315_6018316del, g.6018306_6018310del,
g.6018306_6018310delAGTTA, g.6043633_6043634dup, g.6018256_6018259del,
g.6015623_6017501del, g.6016429_6017479del, g.6017300_6017303del,
g.6045579_6045674delinsATTT, g.(6043690_6045522)_(6045663_6048627)del,
20 g.(?_6012870)_(6042268_6043320)del, g.(6035265_6036956)_(6042268_6043320)del,
g.6038283_6039384del, g.6038901del, g.6038851dup,
g.(6035265_6036956)_(6037055_6038738)del,
g.6037019_6037024delinsCTTCACACACA, g.6036980del, g.6036958dup,
g.6035323_6035324insJN866832.1, g.(6022623_6026389)_(6035265_6036956)del,
25 g.(6031689_6035164)_(6035265_6036956)del, g.6035204_6035207del,
g.6035205_6035206del, g.(?_6012870)_(6031689_6035164)del,
g.(6027252_6029430)_(6031689_6035164)del,
g.(6029587_6031603)_(6031689_6035164)del, g.6028725_6029882del,
g.(?_6012870)_(6029587_6031603)del.

- 30 The present application provides a method of treating patients with Lynch syndrome to reduce the likelihood of developing or treating cancers derived from Lynch syndrome patients, by administering to the subject an effective amount of one or more disclosed

compounds, or a pharmaceutically acceptable salt or solvate thereof, or the corresponding pharmaceutical composition.

Lynch syndrome is a hereditary disorder caused by a mutation in a mismatch repair gene in which affected individuals have a higher than normal chance of developing colorectal cancer, endometrial cancer, and various other types of aggressive cancers, often at a young age – also called hereditary nonpolyposis colon cancer (HNPCC).

The mutations of specific mismatch repair (MMR) genes including but not limited to MLH1, MSH2, MSH6, PMS2, and EPCAM-TACSTD1 deletions are responsible for Lynch syndrome. These genes work in repairing mistakes made when DNA is copied in preparation for cell division. The defects in the genes disallow repair of DNA mistakes and as cells divide, errors stack and uncontrollable cell growth may result in cancer.

Those with Lynch syndrome carry up to an 85% risk of contracting colon cancer as well as a higher than average risk for endometrial cancer, stomach cancer, pancreatic cancer, kidney/ureter tract cancer, hepatobiliary tract cancer, gastric tract cancer, prostate cancer, ovarian cancer, gallbladder duct cancer, brain cancer, small intestine cancer, breast cancer, and skin cancer.

In some embodiments for the disclosed method, the method is a method of treating cancer derived from Lynch syndrome, selected from the group consisting of colon cancer, endometrial cancer, stomach cancer, pancreatic cancer, kidney/ureter tract cancer, hepatobiliary tract cancer, gastric tract cancer, prostate cancer, ovarian cancer, gallbladder duct cancer, brain cancer, small intestine cancer, breast cancer, and skin cancer.

In some embodiments, the method is a method of treating autoimmune disease. Exemplary autoimmune diseases include lupus erythematosus; Wiskott-Aldrich syndrome; autoimmune lymphoproliferative syndrome; myasthenia gravis; rheumatoid arthritis (RA); lupus nephritis; multiple sclerosis; systemic lupus erythematosus; discoid lupus; subacute cutaneous lupus erythematosus; cutaneous lupus erythematosus including chilblain lupus erythematosus; chronic arthritis; Sjogren's syndrome; inflammatory chronic rhinosinusitis; colitis; celiac disease; inflammatory bowel disease; Barrett's esophagus; inflammatory gastritis; autoimmune nephritis; autoimmune vasculitis; autoimmune hepatitis; autoimmune carditis; autoimmune encephalitis; autoimmune diabetes; autoimmune diabetes nephritis; psoriasis; Graft-versus-host disease (GvHD); and autoimmune mediated hematological disease.

In one aspect of this embodiment, the method is a method of treating immune deficiency selected from the group consisting of Autoimmune Lymphoproliferative Syndrome (ALPS), Autoimmune polyglandular syndrome type 1 (APS-1), BENTA Disease, Caspase Eight Deficiency State (CEDS), Chronic Granulomatous Disease (CGD), Common
 5 Variable Immunodeficiency (CVID), Congenital Neutropenia Syndromes, CTLA4 Deficiency, DOCK8 Deficiency, GATA2 Deficiency, Glycosylation Disorders With Immunodeficiency, hyper-immunoglobulin E syndrome (HIES), Hyper-Immunoglobulin M (Hyper-IgM) Syndromes, Leukocyte adhesion deficiency (LAD), LRBA deficiency, PI3 Kinase disease, PLCG2-associated antibody deficiency and immune dysregulation (PLAID),
 10 severe combined immunodeficiency (SCID), STAT3 gain-of-function disease, Warts, Hypogammaglobulinemia, Infections, and Myelokathexis Syndrome (WHIMS), X-Linked Agammaglobulinemia (XLA), X-Linked Lymphoproliferative Disease (XLP), and XMEN Disease.

As used herein, the term "immune deficiency" refers to a condition in which a portion
 15 or some portions of cell components constituting an immune system are defective or dysfunction, so that a normal immune mechanism is damaged. In other words, "immune deficiency" means a condition under which: congenital immunity and/or acquired immunity are suppressed and/or decreased. In some embodiments, the immune -deficiency subject is an immunocompromised subject. Non-limiting examples of immune deficiencies can include
 20 AIDS, hypogammaglobulinemia, agammaglobulinemia, granulocyte deficiency, chronic granulomatous disease, asplenia, SCID, complement deficiency, and/or sickle cell anemia.

In another aspect of this embodiment, the method is a method of treating a neurodegenerative disorder selected from the group consisting of multiple sclerosis, Parkinson's disease (PD), Alzheimer's disease (AD), Dentatorubropallidoluysian atrophy
 25 (DRPLA), Huntington's Disease (HD), Spinocerebellar ataxia Type 1 (SCA1), Spinocerebellar ataxia Type 2 (SCA2), Spinocerebellar ataxia Type 3 (SCA3), Spinocerebellar ataxia 6 (SCA6), Spinocerebellar ataxia Type 7 (SCA7), Spinocerebellar ataxia Type 8 (SCA8), Spinocerebellar ataxia Type 12 (SCA12), Spinocerebellar ataxia Type 17 (SCA17), Spinobulbar Muscular Ataxia/Kennedy Disease (SBMA), Fargile X syndrome
 30 (FRAXA), Fragile XE mental retardation (FRAXE), and Myotonic dystrophy (DM).

A "subject" is a mammal, preferably a human, but can also be an animal in need of veterinary treatment, *e.g.*, companion animals (*e.g.*, dogs, cats, and the like), farm animals

(*e.g.*, cows, sheep, pigs, horses, and the like) and laboratory animals (*e.g.*, rats, mice, guinea pigs, and the like).

In some embodiments, the methods disclosed herein further comprise co-administering an effective amount of a DNA repair inhibitor, a DNA damage response (DDR) inhibitor, a DNA damaging agent or an immunomodulatory agent to the subject being treated for cancer, in addition to an effective amount of a disclosed RAD51 inhibitor.

The term “DNA repair inhibitor” refers to any agent that targets components/processes which a cell uses to repair mutations or changes in DNA and restore the DNA to its original state and prevents the repair of DNA. Examples of DNA repair inhibitors include: RPA inhibitors, APE1 inhibitors, DNA ligase inhibitors, DNA polymerase inhibitors, Parp inhibitors etc.

The term “DNA damage response inhibitor” refers to any agent that targets components/processes involved in detecting DNA lesions, signaling the presence of DNA damage, and/or promote the repair of DNA damage. Examples of DNA damage response inhibitors include checkpoint inhibitors, ATM and ATR inhibitors, DNA-PK inhibitors, etc.

The term “DNA damaging agent” refers to any agent that directly or indirectly damages DNA. The DNA damaging agents is selected from the group consisting of: exposure to a DNA damaging chemical; exposure to a chemotherapeutic agent; exposure to a radiochemotherapy, and exposure to ionizing or ultraviolet radiation. Specific examples of DNA-damaging chemotherapeutic agents include alkylating agents, nitrosoureas, anti-metabolites, plant alkaloids, plant extracts and radioisotopes. Specific examples of the chemotherapeutic agents also include DNA-damaging drugs, for example, 5-fluorouracil (5-FU), capecitabine, S-1 (Tegafur, 5-chloro-2,4-dihydroxypyridine and oxonic acid), 5-ethynyluracil, arabinosyl cytosine (ara-C), 5-azacytidine (5-AC), 2',2'-difluoro-2'-deoxycytidine (dFdC), purine antimetabolites (mercaptopurine, azathiopurine, thioguanine), gemcitabine hydrochlorine (Gemzar), pentostatin, allopurinol, 2-fluoro-arabinosyl-adenine (2F-ara-A), hydroxyurea, sulfur mustard (bischloroethylsulfide), mechlorethamine, melphalan, chlorambucil, cyclophosphamide, ifosfamide, thiotepa, AZQ, mitomycin C, dianhydrogalactitol, dibromoducitol, alkyl sulfonate (busulfan), nitrosoureas (BCNU, CCNU, 4-methyl CCNU or ACNU), procarbazine, decarbazine, rebeccamycin, anthracyclins such as doxorubicin (adriamycin; ADR), daunorubicin (Cerubicine), idarubicin (Idamycin) and epirubicin (Ellence), anthracyclin analogs such as mitoxantrone, actinomycin D, non-intercalating topoisomerase inhibitors such as epipodophyllotoxins (etoposide or VP16,

teniposide or VM-26), podophylotoxin, bleomycin (Bleo), pepleomycin, compounds that form adducts with nucleic acid including platinum derivatives, *e.g.*, cisplatin (CDDP), trans analog of cisplatin, carboplatin, iproplatin, tetraplatin and oxaliplatin, as well as camptothecin, topotecan, irinotecan (CPT-11), and SN-38. Specific examples of nucleic acid
5 damaging treatments include radiation *e.g.*, ultraviolet (UV), infrared (IR), or .alpha.-, .beta.-, or .gamma.-radiation, as well as environmental shock, *e.g.*, hyperthermia.

“Immunomodulatory agent” means an agent that modulates an immune response to an antigen but is not the antigen or derived from the antigen. “Modulate”, as used herein, refers to inducing, enhancing, suppressing, directing, or redirecting an immune response. Such
10 agents include immunostimulatory agents, such as adjuvants, that stimulate (or boost) an immune response to an antigen but is not an antigen or derived from an antigen. There are several distinct types of immunomodulatory agents, which include, but are not limited to, Toll-like Receptor (TLR) agonists and Toll-like Receptor (TLR) antagonists. Such agents also include immunosuppressants. The immunomodulatory agent is selected from the group
15 consisting of immune checkpoint modulators, Toll-like receptor (TLR) agonists, cell-based therapies, cytokines and cancer vaccines.

In some embodiments, the subject is determined to have an increased level and/or activity of a DNA damage process or DNA editing enzyme. In one aspect of this embodiment, the DNA editing enzyme is selected from the group consisting of activation
20 induced cytidine deaminase (AID or AICDA), APOBEC2, APOBEC3A, APOBEC3C, APOBEC3D, APOBEC3F, APOBEC3G, APOBEC3H, APOBEC4, a Type 1 Topoisomerase, a Type 2 Topoisomerase, Recombination Activating Gene 1 (RAG 1), and Recombination Activating Gene 2 (RAG2).

In some embodiments, blood cells obtained from the subject have been determined to
25 have a detectable level of activation-induced cytidine deaminase (AID).

In some embodiments, B cells obtained from the subject have been determined to have a detectable level of activation-induced cytidine deaminase (AID).

In some embodiments, the detectable level of activation-induced cytidine deaminase (AID) is statistically significantly higher than the level of AID expressed in unactivated B-
30 cells or normal non-immune cells from a healthy subject.

The precise amount of compound administered to provide an “effective amount” to the subject will depend on the mode of administration, the type, and severity of the disease, and on the characteristics of the subject, such as general health, age, sex, body weight, and

tolerance to drugs. The skilled artisan will be able to determine appropriate dosages depending on these and other factors. When administered in combination with other therapeutic agents, *e.g.*, when administered in combination with an anti-cancer agent, an “effective amount” of any additional therapeutic agent(s) will depend on the type of drug used. Suitable dosages are known for approved therapeutic agents and can be adjusted by the skilled artisan according to the condition of the subject, the type of condition(s) being treated and the amount of a compound of the application being used by following, for example, dosages reported in the literature and recommended in the *Physician’s Desk Reference* (57th ed., 2003).

The term “effective amount” means an amount when administered to the subject which results in beneficial or desired results, including clinical results, *e.g.*, inhibits, suppresses or reduces the symptoms of the condition being treated in the subject as compared to a control. For example, a therapeutically effective amount can be given in unit dosage form (*e.g.*, 0.1 mg to about 50 g per day, alternatively from 1 mg to about 5 grams per day).

The terms “administer”, “administering”, “administration”, and the like, as used herein, refer to methods that may be used to enable delivery of compositions to the desired site of biological action. These methods include, but are not limited to, intraarticular (in the joints), intravenous, intramuscular, intratumoral, intradermal, intraperitoneal, subcutaneous, orally, topically, intrathecally, inhalationally, transdermally, rectally, and the like.

Administration techniques that can be employed with the agents and methods described herein are found in *e.g.*, Goodman and Gilman, *The Pharmacological Basis of Therapeutics*, current ed.; Pergamon; and Remington’s, *Pharmaceutical Sciences* (current edition), Mack Publishing Co., Easton, Pa.

In addition, the disclosed RAD51 inhibitors can be co-administered with other therapeutic agents. As used herein, the terms “co-administration”, “administered in combination with”, and their grammatical equivalents, are meant to encompass administration of two or more therapeutic agents to a single subject, and are intended to include treatment regimens in which the agents are administered by the same or different route of administration or at the same or different times. In some embodiments the one or more compounds described herein will be co-administered with other agents. These terms encompass administration of two or more agents to the subject so that both agents and/or their metabolites are present in the subject at the same time. They include simultaneous administration in separate compositions, administration at different times in separate

compositions, and/or administration in a composition in which both agents are present. Thus, in some embodiments, the compounds described herein and the other agent(s) are administered in a single composition. In some embodiments, the compounds described herein and the other agent(s) are admixed in the composition.

5 The particular mode of administration and the dosage regimen will be selected by the attending clinician, taking into account the particulars of the case (*e.g.*, the subject, the disease, the disease state involved, the particular treatment). Treatment can involve daily or multi-daily or less than daily (such as weekly or monthly *etc.*) doses over a period of a few days to months, or even years. However, a person of ordinary skill in the art would
10 immediately recognize appropriate and/or equivalent doses looking at dosages of approved compositions for treating a RAD51 mediated disease using the disclosed RAD51 inhibitors for guidance.

 The compounds or the corresponding pharmaceutical compositions taught herein can be administered to a patient in a variety of forms depending on the selected route of
15 administration, as will be understood by those skilled in the art. The compounds of the present teachings may be administered, for example, by oral, parenteral, buccal, sublingual, nasal, rectal, patch, pump or transdermal administration and the pharmaceutical compositions formulated accordingly. Parenteral administration includes intravenous, intraperitoneal, subcutaneous, intramuscular, transepithelial, nasal, intrapulmonary, intrathecal, rectal and
20 topical modes of administration. Parenteral administration can be by continuous infusion over a selected period of time.

 The pharmaceutical composition of the application is formulated to be compatible with its intended route of administration. In an embodiment, the composition is formulated in accordance with routine procedures as a pharmaceutical composition adapted for intravenous,
25 subcutaneous, intramuscular, oral, intranasal, or topical administration to human beings. In preferred embodiments, the pharmaceutical composition is formulated for intravenous administration.

 Typically, for oral therapeutic administration, a compound of the present teachings may be incorporated with excipient and used in the form of ingestible tablets, buccal tablets,
30 troches, capsules, elixirs, suspensions, syrups, wafers, and the like.

 Typically for parenteral administration, solutions of a compound of the present teachings can generally be prepared in water suitably mixed with a surfactant such as hydroxypropylcellulose. Dispersions can also be prepared in glycerol, liquid polyethylene

glycols, DMSO and mixtures thereof with or without alcohol, and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.


Typically, for injectable use, sterile aqueous solutions or dispersion of, and sterile
 5 powders of, a compound described herein for the extemporaneous preparation of sterile injectable solutions or dispersions are appropriate.

EXAMPLES


Abbreviations

10	Ac	acetyl
	ACN	acetonitrile
	aq	aqueous
	Bn	benzyl
	Boc	<i>tert</i> -butoxycarbonyl
15	br.	Broad
	CAN	cerium ammonium nitrate
	d	doublet (when in reference to an ¹ H NMR spectra)
	DCM	dichloromethane
	DIEA(DIPEA)	diisopropylethylamine
20	DMA	dimethylacetamide
	DMAP	4-dimethylaminopyridine
	DMF	<i>N,N</i> -dimethylformamide
	DMSO	dimethylsulfoxide
	dppf	1,1'-bis(diphenylphosphino) ferrocene
25	eq	equivalent
	EtOAc	ethyl acetate
	h	hour
	HBTU	<i>N,N,N',N'</i> -tetramethyl-O-(1 <i>H</i> -benzotriazol-1-yl)uronium hexafluorophosphate
30	HPLC	high performance liquid chromatography
	LC-MS	liquid chromatography coupled to mass spectrometry
	m	multiplet (when in reference to an ¹ H NMR spectra)
	MS ESI	mass spectra, electrospray ionization
	NBS	<i>N</i> -bromosuccinimide
35	NMR	nuclear magnetic resonance
	prep	preparative
	Py	pyridine
	s	singlet (when in reference to an ¹ H NMR spectra)
	sat	saturated
40	SFC	supercritical fluid chromatography
	t	triplet (when in reference to an ¹ H NMR spectra)
	TEA	triethylamine
	TFA	trifluoroacetic acid
	THF	tetrahydrofuran
45	TLC	thin layer chromatography
	Tol	toluene

Example 1. Preparation of isopropyl (*trans*-4-(5-(4-(aminomethyl)-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (Compound 1)




5 a) Synthesis of 4-bromo-3-(*tert*-butylsulfamoyl)benzamide (**General Method A**)



To a solution of 4-bromo-3-(*tert*-butylsulfamoyl)benzoic acid (1.9 g, 5.65 mmol, 1.0 *eq.*) and DIEA (84.77 mmol, 14.77 mL, 15.0 *eq.*) in DMF (50 mL) was added NH₄Cl (3.02 g, 56.51 mmol, 10.0 *eq.*) and HATU (2.79 g, 7.35 mmol, 1.3 *eq.*). The mixture was stirred at 25°C for 12 h, then poured into H₂O (200 mL) and extracted with EtOAc (30 mL x 3). The combined organic layers were washed with brine (100 mL), dried over Na₂SO₄, filtered and concentrated to give 4-bromo-3-(*tert*-butylsulfamoyl)benzamide (2.2 g, crude) as a yellow gum. [M+H] = 335.0/337.0.


b) Synthesis of 5-(aminomethyl)-2-bromo-*N*-*tert*-butyl-benzene sulfonamide



To a solution of 4-bromo-3-(*tert*-butylsulfamoyl)benzamide (1 g, 2.98 mmol, 1.0 *eq.*) in THF (30 mL) was added BH₃-Me₂S (10 M in THF, 1.49 mL, 5 *eq.*). The mixture was stirred at 50°C for 2 h. The reaction mixture was quenched by 1N HCl (2 mL) at 0°C, stirred at 50°C for 1 hr, then concentrated under reduced pressure to give 5-(aminomethyl)-2-bromo-


N-*tert*-butyl-benzenesulfonamide (1.5 g, crude, HCl salt) as a white solid. $[M+H]^+ = 321.1/323.1$.

c) Synthesis of *tert*-butyl *N*-[[4-bromo-3-(*tert*-butylsulfamoyl) phenyl]methyl]carbamate




To a solution of 5-(aminomethyl)-2-bromo-*N*-*tert*-butyl-benzenesulfonamide (HCl salt, 1.5 g, 4.67 mmol, 1.0 *eq.*) and Boc_2O (1.22 g, 5.60 mmol, 1.2 *eq.*) in MeOH (10 mL) and H_2O (10 mL) was added Na_2CO_3 (990 mg, 9.34 mmol, 2.0 *eq.*). The mixture was stirred at 25°C for 12 h under N_2 atmosphere and then concentrated. The residue was purified by prep-HPLC to give *tert*-butyl *N*-[[4-bromo-3-(*tert*-butylsulfamoyl) phenyl]methyl]carbamate (190 mg, 451 μmol , 10% yield) as a white solid. ^1H NMR (400MHz, CDCl_3) δ = 7.99 (d, $J=1.3$ Hz, 1H), 7.59 (d, $J=8.2$ Hz, 1H), 7.27 - 7.22 (m, 1H), 4.25 (br d, $J=5.7$ Hz, 2H), 1.38 (s, 9H), 1.14 (s, 9H). $[M+\text{Na}]^+ = 443.1/445.1$.

d) Synthesis of *tert*-butyl *N*-[[3-(*tert*-butylsulfamoyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]methyl]carbamate (**General Method B**)




A mixture of *tert*-butyl *N*-[[4-bromo-3-(*tert*-butylsulfamoyl)phenyl]methyl]carbamate (180 mg, 427.2 μmol , 1.0 *eq.*), B_2Pin_2 (326 mg, 1.28 mmol, 3.0 *eq.*), KOAc (126 mg, 1.28 mmol, 3.0 *eq.*) and $\text{Pd}(\text{dppf})\text{Cl}_2 \cdot \text{CH}_2\text{Cl}_2$ (35 mg, 42.7 μmol , 0.1 *eq.*) in dioxane (5 mL) was degassed and purged with N_2 for 3 times and stirred at 80°C for 12 h. The reaction mixture was concentrated and the residue was purified by prep-TLC (SiO_2 , Petroleum ether/Ethyl acetate=3/1) to give *tert*-butyl *N*-[[3-(*tert*-butylsulfamoyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]methyl]carbamate (20 mg, 42.7 μmol , 10% yield) as a yellow gum. ESI $[M+H]^+ = 469.2$.

e) Synthesis of isopropyl *N*-[*trans*-4-[5-[4-[(*tert*-butoxycarbonylamino) methyl]-2-(*tert*-butylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl] carbamate (**General Method C**)




- A mixture of *tert*-butyl *N*-[[3-(*tert*-butylsulfamoyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]methyl]carbamate (20 mg, 42.7 μ mol, 1.0 *eq.*), isopropyl (*trans*-4-(5-bromothiazol-2-yl)cyclohexyl)carbamate (18 mg, 51.2 μ mol, 1.2 *eq.*), Na₂CO₃ (14 mg, 128 μ mol, 3.0 *eq.*), Pd(PPh₃)₄ (4.9 mg, 4.3 μ mol, 0.1 *eq.*) and KF (7.4 mg, 128 μ mol, 3.0 *eq.*) in EtOH (1 mL)/H₂O (0.3 mL)/Tol. (1 mL) was degassed and purged with N₂ for 3 times and then stirred at 80°C for 12 h. The reaction mixture was concentrated and the residue was purified by prep-TLC (SiO₂, Petroleum ether/Ethyl acetate=2/1) to give isopropyl *N*-[*trans*-4-[5-[4-[(*tert*-butoxycarbonylamino)methyl]-2-(*tert*-butylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (20 mg, crude) as a yellow gum. ESI [M+H] = 609.3.
- f) Synthesis of isopropyl (*trans*-4-(5-(4-(aminomethyl)-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (**Compound 1**)




- A solution of isopropyl *N*-[*trans*-4-[5-[4-[(*tert*-butoxycarbonylamino)methyl]-2-(*tert*-butylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (20 mg, crude) in HCl/MeOH (4 M, 1 mL) was stirred at 25°C for 30 mins. The reaction mixture was concentrated and the residue was purified by prep-HPLC (column: Nano-Micro UniSil 5-100 C18 ULTRA 100*250mm 5 μ m; mobile phase: [water(0.1%TFA)-ACN]; B%: 30%-60%, 10min) to give isopropyl (*trans*-4-(5-(4-(aminomethyl)-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (1.03 mg, 1.47 μ mol, 89% purity, TFA salt) as a pale yellow solid.
- ¹H NMR (400MHz, methanol-*d*₄) δ = 8.32 (d, J=1.7 Hz, 1H), 7.81 (s, 1H), 7.75 - 7.70 (m, 1H), 7.60 (d, J=7.8 Hz, 1H), 4.86 - 4.82 (m, 1H), 4.28 (s, 2H), 3.52 - 3.43 (m, 1H), 3.11 - 3.00 (m, 1H), 2.26 (br d, J=13.7 Hz, 2H), 2.09 (br d, J=11.1 Hz, 2H), 1.79 - 1.68 (m, 2H), 1.50 - 1.37 (m, 2H), 1.25 (br d, J=6.1 Hz, 6H), 1.09 (s, 9H). ESI [M+H] = 509.2.

- Example 2. Preparation of isopropyl (*trans*-4-(5-(4-(acetamidomethyl)-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (Compound 2)**




a) Synthesis of *N*-[[4-bromo-3-(*tert*-butylsulfamoyl)phenyl]methyl]acetamide




To a solution of 5-(aminomethyl)-2-bromo-*N*-*tert*-butylbenzenesulfonamide (0.4 g, 1.25 mmol, 1.0 *eq.*) in DCM (5 mL) were added DIEA (6.25 mmol, 1.1 mL, 5.0 *eq.*), DMAP (15.2 mg, 125 μmol, 0.1 *eq.*) and Ac₂O (153 mg, 1.5 mmol, 1.2 *eq.*). The mixture was stirred at 25°C for 1 hr, then poured into H₂O (20 mL) and extracted with EtOAc (20 mL x 3). The combined organic layers were dried over Na₂SO₄, filtered and concentrated to give *N*-[[4-bromo-3-(*tert*-butylsulfamoyl)phenyl]methyl]acetamide (320 mg, crude) as a yellow oil. ESI [M+H] = 363.1/365.1

b) Synthesis of *N*-[[3-(*tert*-butylsulfamoyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]methyl]acetamide




A mixture of *N*-[[4-bromo-3-(*tert*-butylsulfamoyl)phenyl]methyl]acetamide (0.3 g, 826 μmol, 1.0 *eq.*), B₂Pin₂ (629 mg, 2.5 mmol, 3.0 *eq.*), KOAc (243 mg, 2.5 mmol, 3 *eq.*) and Pd(dppf)Cl₂.CH₂Cl₂ (67.5 mg, 82.6 μmol, 0.1 *eq.*) in dioxane (5 mL) was degassed and purged with N₂ for 3 times and then stirred at 80°C for 12 h. The reaction mixture was concentrated and the residue was purified by column chromatography (SiO₂, Petroleum ether/Ethyl acetate = 10/1 to 0:1) to give *N*-[[3-(*tert*-butylsulfamoyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]methyl]acetamide (70 mg, 171 μmol, 21% yield) as a yellow gum. ESI [M+H] = 411.3.

c) Synthesis of isopropyl (*trans*-4-(5-(4-(acetamidomethyl)-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (**Compound 2**)




A mixture of *N*-[[3-(*tert*-butylsulfamoyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]methyl]acetamide (66 mg, 161 μ mol, 0.93 *eq.*), isopropyl *N*-[*trans*-4-(5-bromothiazol-2-yl)cyclohexyl]carbamate (60 mg, 173 μ mol, 1.0 *eq.*), Na₂CO₃ (54.9 mg, 518 μ mol, 3.0 *eq.*), KF (30.1 mg, 518 μ mol, 3.0 *eq.*) and Pd(PPh₃)₄ (19.9 mg, 17.3 μ mol, 0.1 *eq.*) in H₂O (0.3 mL)/EtOH (1.0 mL)/Tol. (1.0 mL) was degassed and purged with N₂ for 3 times. The mixture was stirred at 80°C for 12 h under N₂ atmosphere and then concentrated. The residue was purified by prep-TLC (SiO₂, Petroleum ether/Ethyl acetate=1/1) to give isopropyl (*trans*-4-(5-(4-(acetamidomethyl)-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (16.99 mg, 30.85 μ mol, 18% yield, 100% purity) as a white solid. ¹H NMR (400MHz, methanol-*d*₄) δ = 8.10 (d, *J*=1.6 Hz, 1H), 7.81 (s, 1H), 7.57 (dd, *J*=1.6, 7.9 Hz, 1H), 7.47 (d, *J*=7.8 Hz, 1H), 4.88 - 4.81 (m, 1H), 4.48 (s, 2H), 3.52 - 3.43 (m, 1H), 3.07 (br t, *J*=12.2 Hz, 1H), 2.26 (br d, *J*=12.1 Hz, 2H), 2.10 (br d, *J*=9.9 Hz, 2H), 2.04 (s, 3H), 1.80 - 1.66 (m, 2H), 1.50 - 1.38 (m, 2H), 1.24 (d, *J*=6.1 Hz, 6H), 1.11 (s, 9H). ESI [M+H] = 551.2.

Example 3. Preparation of isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-(pyrazolo[1,5-*a*]pyridin-3-yl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (Compound 3**)**




a) Synthesis of isopropyl (*trans*-4-(5-(4-bromo-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate



To a solution of *tert*-butyl nitrite (2.2 g, 21.2 mmol, 1.5 *eq.*) in MeCN (20 mL) was added CuBr₂ (3.16 g, 14.2 mmol, 1.0 *eq.*) and isopropyl (*trans*-4-(5-(4-amino-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (7.0 g, 14.2 mmol, 1.0 *eq.*). The mixture was stirred at 60°C for 1 hr, then poured into 1N NaHCO₃ solution (100 mL) at 25°C and extracted with EtOAc (50 mL x 3). The combined organic layers dried over Na₂SO₄, filtered and concentrated. The residue was purified by column chromatography (SiO₂, Petroleum ether/Ethyl acetate = 10:1 to 5:1 to 0:1) to give isopropyl (*trans*-4-(5-(4-bromo-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl)thiazol-2-yl) cyclohexyl)carbamate (4.2 g, 7.52 mmol, 53% yield) as a white solid. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.29 (d, *J* = 2.1 Hz, 1H), 7.79 (s, 2H), 7.42 (d, *J* = 8.2 Hz, 1H), 4.84 (br d, *J* = 6.4 Hz, 1H), 3.53 - 3.43 (m, 1H), 3.03 (s, 1H), 2.25 (br d, *J* = 12.5 Hz, 2H), 2.13 - 2.06 (m, 2H), 1.77 - 1.64 (m, 2H), 1.43 (br dd, *J* = 2.4, 12.6 Hz, 2H), 1.25 (dd, *J* = 2.0, 6.7 Hz, 6H), 1.13 - 1.08 (m, 9H). ESI [M+H] = 558.1/560.1.

b) Synthesis of isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-(pyrazolo[1,5-*a*]pyridin-3-yl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (**Compound 3**; General Method D)



A mixture of isopropyl (*trans*-4-(5-(4-bromo-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (40 mg, 71.6 μmol, 1.0 *eq.*), 3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrazolo[1,5-*a*]pyridine (21 mg, 86 μmol, 1.2 *eq.*), KOAc (21.08 mg, 214.84 μmol, 3.0 *eq.*) and Pd(AmPhos)Cl₂ (5.07 mg, 7.16 μmol, 0.1 *eq.*) in EtOH (1 mL)/H₂O (0.25 mL) was degassed and purged with N₂ for 3 times, and then stirred at 80°C for 12 h. The reaction mixture was concentrated and the residue was purified by prep-HPLC (column: Welch Ultimate AQ-C18 150*30mm*5μm; mobile phase: [water(0.1%TFA)-ACN]; B%: 50%-80%, 12min) to give isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-(pyrazolo[1,5-*a*]pyridin-3-yl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (22.64 mg, 37.62 μmol, 52.53% yield, 99% purity) as a white solid. ¹H NMR (400MHz, DMSO-*d*₆) δ = 8.82 (d,

J=7.0 Hz, 1H), 8.52 (s, 1H), 8.34 (d, J=1.8 Hz, 1H), 8.03 (d, J=9.0 Hz, 1H), 7.98 - 7.94 (m, 1H), 7.77 (s, 1H), 7.56 (d, J=7.9 Hz, 1H), 7.50 - 7.43 (m, 1H), 7.17 (s, 1H), 7.09 - 7.00 (m, 2H), 4.76 (td, J=6.2, 12.5 Hz, 1H), 2.95 (tt, J=3.6, 11.9 Hz, 1H), 2.16 (br d, J=11.6 Hz, 2H), 1.94 (br d, J=10.4 Hz, 2H), 1.68 - 1.55 (m, 2H), 1.43 - 1.31 (m, 2H), 1.18 (d, J=6.2 Hz, 6H), 1.08 (s, 9H). ESI [M+H] = 596.2.

Example 4. Preparation of isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-(1-(2,2,2-trifluoroethyl)-1*H*-pyrazol-4-yl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (Compound 4)

From isopropyl (*trans*-4-(5-(4-bromo-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate, using the corresponding boric acid or boronate and under the same reaction conditions as for **Compound 3**. ¹H NMR (400 MHz, DMSO-*d*₆) δ = 8.43 (s, 1H), 8.19 (d, J = 1.8 Hz, 1H), 8.12 (s, 1H), 7.84 (dd, J = 1.8, 7.9 Hz, 1H), 7.70 (s, 1H), 7.45 (d, J = 7.9 Hz, 1H), 7.07 - 6.89 (m, 2H), 5.17 (q, J = 9.0 Hz, 2H), 4.71 (td, J = 6.2, 12.3 Hz, 1H), 3.36 - 3.21 (m, 1H), 2.89 (ddd, J = 3.3, 8.5, 11.7 Hz, 1H), 2.11 (br d, J = 11.7 Hz, 2H), 1.89 (br d, J = 10.4 Hz, 2H), 1.55 (dq, J = 2.5, 12.7 Hz, 2H), 1.37 - 1.25 (m, 2H), 1.13 (d, J = 6.2 Hz, 6H), 1.01 (s, 9H). ESI [M+H] = 628.2.

Example 5. Preparation of isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-(pyridin-3-yl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (Compound 5)

From isopropyl (*trans*-4-(5-(4-bromo-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate, using the corresponding boric acid or boronate and under the same reaction conditions as for **Compound 3**. ¹H NMR (400 MHz, DMSO-*d*₆) δ = 9.21 (s, 1H), 8.84 (br d, J = 5.1 Hz, 1H), 8.67 (br d, J = 7.9 Hz, 1H), 8.41 (s, 1H), 8.07 (br d, J = 7.9 Hz, 1H), 8.00 - 7.89 (m, 1H), 7.77 (s, 1H), 7.65 (d, J = 7.9 Hz, 1H), 7.10 (s, 1H), 6.98 (br d, J = 7.1 Hz, 1H), 4.71 (td, J = 6.0, 12.2 Hz, 1H), 3.37 - 3.23 (m, 1H), 2.92 (br t, J = 11.8 Hz, 1H), 2.12 (br d, J = 12.1 Hz, 2H), 1.90 (br d, J = 11.0 Hz, 2H), 1.63 - 1.49 (m, 2H), 1.38 - 1.27 (m, 2H), 1.13 (d, J = 6.2 Hz, 6H), 1.00 (s, 9H). ESI [M+H] = 557.2.

Example 6. Preparation of isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-(1*H*-pyrazol-3-yl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (Compound 6)

From isopropyl (*trans*-4-(5-(4-bromo-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate, using the corresponding boric acid or boronate and under the same reaction conditions as for **Compound 3**. ¹H NMR (400 MHz, DMSO-*d*₆) δ = 8.52 (d, J=1.54 Hz, 1H), 7.99 (dd, J=1.76, 7.94 Hz, 1H), 7.81 (d, J=1.98 Hz, 1H), 7.73 (s, 1H), 7.49 (d, J=7.94 Hz, 1H), 7.07 (s, 1H), 7.00 (br d, J=7.50 Hz, 1H), 6.81 (d, J=2.21 Hz, 1H), 4.73 (td,

J=6.17, 12.35 Hz, 1H), 3.31 (br d, J=7.94 Hz, 1H), 2.85-2.98 (m, 1H), 2.13 (br d, J=12.35 Hz, 2H), 1.91 (br d, J=9.92 Hz, 2H), 1.50-1.67 (m, 2H), 1.25-1.43 (m, 2H), 1.15 (d, J=6.39 Hz, 6H), 1.04 (s, 9H). ESI [M+H] = 546.2.

Example 7. Preparation of isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-(6-hydroxy pyridin-3-yl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (Compound 7)

From isopropyl (*trans*-4-(5-(4-bromo-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl) thiazol-2-yl)cyclohexyl)carbamate, using the corresponding boric acid or boronate and under the same reaction conditions as for **Compound 3**. ¹H NMR (400MHz, DMSO-*d*₆) δ = 8.17 (d, J=1.8 Hz, 1H), 7.93 - 7.81 (m, 3H), 7.74 (br s, 1H), 7.50 (d, J=8.1 Hz, 1H), 7.11 (br s, 1H), 7.03 (br d, J=7.6 Hz, 1H), 6.50 (d, J=9.4 Hz, 1H), 4.75 (td, J=6.1, 12.4 Hz, 1H), 3.40 - 3.24 (m, 1H), 2.95 (br d, J=9.9 Hz, 1H), 2.15 (br d, J=12.2 Hz, 2H), 1.93 (br d, J=10.0 Hz, 2H), 1.68 - 1.52 (m, 2H), 1.41 - 1.30 (m, 2H), 1.17 (d, J=6.2 Hz, 6H), 1.05 (s, 9H). ESI [M+H] = 573.2.

Example 8. Preparation of isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-(2,4-dimethoxy pyrimidin-5-yl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (Compound 8)

From isopropyl (*trans*-4-(5-(4-bromo-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl) thiazol-2-yl)cyclohexyl)carbamate, using the corresponding boric acid or boronate and under the same reaction conditions as for **Compound 3**. ¹H NMR (400MHz, DMSO-*d*₆) δ = 8.50 (s, 1H), 8.25 (d, J=1.8 Hz, 1H), 7.86 - 7.72 (m, 2H), 7.56 (d, J=8.1 Hz, 1H), 7.21 (s, 1H), 7.05 (br d, J=7.8 Hz, 1H), 4.75 (td, J=6.1, 12.4 Hz, 1H), 3.98 (d, J=4.0 Hz, 6H), 3.42 - 3.26 (m, 1H), 2.94 (tt, J=3.5, 11.9 Hz, 1H), 2.16 (br d, J=12.0 Hz, 2H), 1.93 (br d, J=10.0 Hz, 2H), 1.67 - 1.52 (m, 2H), 1.44 - 1.27 (m, 2H), 1.17 (d, J=6.2 Hz, 6H), 1.13 - 1.03 (m, 9H). ESI [M+H] = 618.3.

Example 9. Preparation of isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-(2H-indazol-6-yl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (Compound 9)

From isopropyl (*trans*-4-(5-(4-bromo-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl) thiazol-2-yl)cyclohexyl)carbamate, using the corresponding boric acid or boronate and under the same reaction conditions as for **Compound 3**. ¹H NMR (400MHz, DMSO-*d*₆) δ = 8.39 (d, J=1.7 Hz, 1H), 8.15 (s, 1H), 8.02 (dd, J=1.8, 8.0 Hz, 1H), 7.93 (d, J=8.3 Hz, 1H), 7.84 (s, 1H), 7.79 (s, 1H), 7.60 (d, J=7.9 Hz, 1H), 7.49 (d, J=8.4 Hz, 1H), 7.21 (s, 1H), 7.03 (br d, J=7.6 Hz, 1H), 4.76 (td, J=6.2, 12.4 Hz, 1H), 3.34 (br d, J=7.3 Hz, 1H), 3.01 - 2.90 (m, 1H), 2.17 (br d, J=12.1 Hz, 2H), 1.94 (br d, J=10.4 Hz, 2H), 1.67 - 1.53 (m, 2H), 1.43 - 1.30 (m, 2H), 1.18 (d, J=6.2 Hz, 6H), 1.09 (s, 9H). ESI [M+H] = 596.3.

Example 10. Preparation of isopropyl (*trans*-4-(5-(4-(2-aminopyrimidin-5-yl)-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (Compound 10)

From isopropyl (*trans*-4-(5-(4-bromo-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate, using the corresponding boric acid or boronate and under the same reaction conditions as for **Compound 3**. ¹H NMR (400MHz, DMSO-*d*₆) δ = 8.66 (s, 2H), 8.23 (d, J=1.7 Hz, 1H), 7.90 (dd, J=1.8, 8.0 Hz, 1H), 7.78 - 7.69 (m, 1H), 7.53 (d, J=7.9 Hz, 1H), 7.10 (s, 1H), 7.05 - 6.97 (m, 3H), 4.75 (td, J=6.2, 12.4 Hz, 1H), 3.42 - 3.26 (br s, 1H), 2.94 (tt, J=3.5, 11.8 Hz, 1H), 2.15 (br d, J=12.0 Hz, 2H), 1.93 (br d, J=10.3 Hz, 2H), 1.67 - 1.51 (m, 2H), 1.43 - 1.27 (m, 2H), 1.17 (d, J=6.2 Hz, 6H), 1.06 (s, 9H). ESI [M+H] = 573.2.

Example 11. Preparation of isopropyl (*trans*-4-(5-(4-(benzofuran-5-yl)-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (Compound 11)

From isopropyl (*trans*-4-(5-(4-bromo-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate, using the corresponding boric acid or boronate and under the same reaction conditions as for **Compound 3**. ¹H NMR (400 MHz, DMSO-*d*₆) δ = 8.32 (d, J = 2.0 Hz, 1H), 8.06 (d, J = 2.2 Hz, 1H), 8.01 (d, J = 1.5 Hz, 1H), 7.95 (dd, J = 2.0, 7.9 Hz, 1H), 7.77 - 7.71 (m, 2H), 7.68 - 7.64 (m, 1H), 7.56 (d, J = 7.9 Hz, 1H), 7.12 (s, 1H), 7.07 - 7.04 (m, 1H), 7.03 - 6.98 (m, 1H), 4.78 - 4.68 (m, 1H), 3.27 (br s, 1H), 2.98 - 2.86 (m, 1H), 2.21 - 2.09 (m, 2H), 1.91 (br dd, J = 2.0, 12.3 Hz, 2H), 1.56 (br d, J = 2.2 Hz, 2H), 1.40 - 1.27 (m, 2H), 1.15 (d, J = 6.2 Hz, 6H), 1.05 (s, 9H). ESI [M+H] = 596.2.

Example 12. Preparation of isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-(pyrimidin-5-yl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (Compound 12)

From isopropyl (*trans*-4-(5-(4-bromo-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate, using the corresponding boric acid or boronate and under the same reaction conditions as for **Compound 3**. ¹H NMR (400 MHz, DMSO-*d*₆) δ = 9.23 (s, 1H), 9.18 (s, 2H), 8.37 (d, J = 2.0 Hz, 1H), 8.05 (dd, J = 1.9, 8.0 Hz, 1H), 7.75 (s, 1H), 7.63 (d, J = 7.9 Hz, 1H), 7.13 (s, 1H), 7.01 (br d, J = 7.7 Hz, 1H), 4.71 (s, 1H), 3.31 (br d, J=7.72 Hz, 1H), 2.97 - 2.86 (m, 1H), 2.12 (br d, J = 11.7 Hz, 2H), 1.89 (br d, J = 10.1 Hz, 2H), 1.56 (dq, J = 2.9, 12.8 Hz, 2H), 1.36 - 1.25 (m, 2H), 1.13 (d, J = 6.4 Hz, 6H), 1.02 (s, 9H). ESI [M+H] = 558.2.

Example 13. Preparation of isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-(3,6-dimethoxypyridazin-4-yl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (Compound 13)

From isopropyl (*trans*-4-(5-(4-bromo-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate, using the corresponding boric acid or boronate and under the same

reaction conditions as for **Compound 3**. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.33 (d, J=1.76 Hz, 1H), 7.87 (dd, J=1.76, 7.94 Hz, 1H), 7.76 (s, 1H), 7.59 (d, J=7.94 Hz, 1H), 7.35 (s, 1H), 7.22 (s, 1H), 6.93-7.05 (m, 1H), 4.73 (td, J=6.17, 12.35 Hz, 1H), 3.99 (d, J=2.20 Hz, 6H), 3.31 (br d, J=7.72 Hz, 1H), 2.81-3.01 (m, 1H), 2.13 (br d, J=12.57 Hz, 2H), 1.84-1.99 (m, 2H), 1.49-1.64 (m, 2H), 1.24-1.43 (m, 2H), 1.15 (d, J=6.17 Hz, 6H), 1.08 (s, 9H). ESI [M+H] = 618.2.

Example 14. Preparation of isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-(2-methylthiazol-5-yl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (Compound 14)

From isopropyl (*trans*-4-(5-(4-bromo-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate, using the corresponding boric acid or boronate and under the same reaction conditions as for **Compound 3**. ¹H NMR (400 MHz, DMSO-*d*₆) δ = 8.24 - 8.15 (m, 2H), 7.93 (dd, J = 2.0, 8.1 Hz, 1H), 7.77 (s, 1H), 7.55 (d, J = 8.1 Hz, 1H), 7.27 (s, 1H), 7.03 (br d, J = 7.7 Hz, 1H), 4.75 (td, J = 6.2, 12.4 Hz, 1H), 3.39 - 3.25 (m, 1H), 2.94 (tt, J = 3.4, 11.8 Hz, 1H), 2.72 (s, 3H), 2.15 (br d, J = 11.9 Hz, 2H), 1.93 (br d, J = 10.5 Hz, 2H), 1.59 (dq, J = 2.9, 12.7 Hz, 2H), 1.41 - 1.29 (m, 2H), 1.17 (d, J = 6.2 Hz, 6H), 1.07 (s, 9H). ESI [M+H] = 577.2.

Example 15. Preparation of isopropyl (*trans*-4-(5-(4-(benzo[*d*]thiazol-6-yl)-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (Compound 15)

From isopropyl (*trans*-4-(5-(4-bromo-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate, using the corresponding boric acid or boronate and under the same reaction conditions as for **Compound 3**. ¹H NMR (400 MHz, DMSO-*d*₆) δ = 9.47 (s, 1H), 8.63 (d, J = 1.6 Hz, 1H), 8.42 (d, J = 2.0 Hz, 1H), 8.24 (d, J = 8.6 Hz, 1H), 8.05 (dd, J = 2.0, 8.1 Hz, 1H), 7.92 (dd, J = 1.8, 8.6 Hz, 1H), 7.79 (s, 1H), 7.63 (d, J = 8.1 Hz, 1H), 7.14 (s, 1H), 7.09 - 6.95 (m, 1H), 4.76 (td, J = 6.3, 12.4 Hz, 1H), 3.34 (br d, J = 4.6 Hz, 1H), 3.02 - 2.89 (m, 1H), 2.17 (br d, J = 11.9 Hz, 2H), 1.94 (br d, J = 10.4 Hz, 2H), 1.68 - 1.54 (m, 2H), 1.42 - 1.30 (m, 2H), 1.18 (d, J = 6.2 Hz, 6H), 1.07 (s, 9H). ESI [M+H] = 613.2.

Example 16. Preparation of isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-(5-((dimethylamino)methyl)pyridin-3-yl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (Compound 16)

From isopropyl (*trans*-4-(5-(4-bromo-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate, using the corresponding boric acid or boronate and under the same reaction conditions as for **Compound 3**. ¹H NMR (400MHz, DMSO-*d*₆) δ = 9.08 (d, J=2.1 Hz, 1H), 8.75 (d, J=1.7 Hz, 1H), 8.42 (d, J=1.8 Hz, 1H), 8.36 (s, 1H), 8.07 (dd, J=2.0, 7.9 Hz,

1H), 7.80 (s, 1H), 7.68 (d, J=7.9 Hz, 1H), 7.14 - 6.96 (m, 2H), 4.75 (td, J=6.1, 12.4 Hz, 1H), 4.46 (br s, 2H), 3.45 - 3.25 (m, 1H), 3.07 - 2.89 (m, 1H), 2.83 (br s, 6H), 2.17 (br d, J=11.7 Hz, 2H), 1.94 (br d, J=10.3 Hz, 2H), 1.69 - 1.52 (m, 2H), 1.46 - 1.27 (m, 2H), 1.17 (d, J=6.2 Hz, 6H), 1.03 (s, 9H). ESI [M+H] = 614.3.

5 **Example 17. Preparation of isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-(5-methyl-1*H*-pyrazol-4-yl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (Compound 17)**

From isopropyl (*trans*-4-(5-(4-bromo-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate, using the corresponding boric acid or boronate and under the same reaction conditions as for **Compound 3**. ¹H NMR (400 MHz, DMSO-*d*₆) δ = 8.12 (d, J = 1.8
10 Hz, 1H), 7.93 (s, 1H), 7.76 - 7.69 (m, 2H), 7.48 (d, J = 7.9 Hz, 1H), 7.15 - 7.00 (m, 2H), 4.75 (td, J = 6.2, 12.5 Hz, 1H), 3.40 - 3.26 (m, 1H), 2.93 (tt, J = 3.4, 11.8 Hz, 1H), 2.43 (s, 3H), 2.15 (br d, J = 12.0 Hz, 2H), 1.93 (br d, J = 10.4 Hz, 2H), 1.59 (dq, J = 2.8, 12.7 Hz, 2H), 1.41 - 1.29 (m, 2H), 1.17 (d, J = 6.2 Hz, 6H), 1.07 (s, 9H). ESI [M+H] = 560.2.

15 **Example 18. Preparation of isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-(6-cyano pyridin-3-yl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (Compound 18)**

From isopropyl (*trans*-4-(5-(4-bromo-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate, using the corresponding boric acid or boronate and under the same reaction conditions as for **Compound 3**. ¹H NMR (400 MHz, methanol-*d*₄) δ = 9.08 (d, J = 1.5 Hz, 1H), 8.49 (d, J = 2.2 Hz, 1H), 8.36 - 8.32 (m, 1H), 8.04 - 7.99 (m, 2H), 7.85 - 7.83
20 (m, 1H), 7.67 (d, J = 7.9 Hz, 1H), 4.81 - 4.75 (m, 1H), 3.48 - 3.44 (m, 1H), 3.07 - 3.03 (m, 1H), 2.27 - 2.23 (m, 2H), 2.10 - 2.05 (m, 2H), 1.75 - 1.69 (m, 2H), 1.46 - 1.40 (m, 2H), 1.25 - 1.19 (m, 6H), 1.10 (s, 9H). ESI [M+H] = 582.2.

Example 19. Preparation of isopropyl (*trans*-4-(5-(4-(1*H*-benzo[*d*][1,2,3]triazol-6-yl)-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (Compound 19)

25 From isopropyl (*trans*-4-(5-(4-bromo-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate, using the corresponding boric acid or boronate and under the same reaction conditions as for **Compound 3**. ¹H NMR (400 MHz, DMSO-*d*₆) δ = 8.38 (d, J = 2.0 Hz, 1H), 8.23 (br s, 1H), 8.08 - 8.01 (m, 2H), 7.82 - 7.74 (m, 2H), 7.59 (d, J = 7.9 Hz, 1H), 7.18 (s, 1H), 7.00 (br d, J = 7.5 Hz, 1H), 4.73 (td, J = 6.2, 12.3 Hz, 1H), 3.31 (td, J = 3.7, 7.3
30 Hz, 1H), 2.96 - 2.89 (m, 1H), 2.14 (br d, J = 11.9 Hz, 2H), 1.91 (br d, J = 9.9 Hz, 2H), 1.58 (dq, J = 2.8, 12.8 Hz, 2H), 1.39 - 1.29 (m, 2H), 1.15 (d, J = 6.2 Hz, 6H), 1.06 (s, 9H). ESI [M+H] = 597.2.

Example 20. Preparation of isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-(6-(hydroxymethyl)pyridin-3-yl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (Compound 20)

From isopropyl (*trans*-4-(5-(4-bromo-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate, using the corresponding boric acid or boronate and under the same reaction conditions as for **Compound 3**. ¹H NMR (400 MHz, DMSO-*d*₆) δ = 8.92 (s, 1H), 8.36 (d, J = 1.6 Hz, 2H), 8.08 - 8.01 (m, 1H), 7.79 (s, 1H), 7.72 (d, J = 7.8 Hz, 1H), 7.64 (d, J = 8.1 Hz, 1H), 7.19 (s, 1H), 7.04 (br d, J = 8.2 Hz, 1H), 4.78 - 4.73 (m, 1H), 4.70 (s, 2H), 3.39 - 3.29 (m, 1H), 3.01 - 2.89 (m, 1H), 2.16 (br d, J = 12.7 Hz, 2H), 1.97 - 1.90 (m, 2H), 1.60 (dd, J = 2.6, 12.4 Hz, 2H), 1.42 - 1.31 (m, 2H), 1.18 (d, J = 6.2 Hz, 6H), 1.06 (s, 9H). ESI [M+H] = 587.2.


Example 21. Preparation of isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-(3-methyl-1*H*-indol-6-yl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (Compound 21)

From isopropyl (*trans*-4-(5-(4-bromo-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate, using the corresponding boric acid or boronate and under the same reaction conditions as for **Compound 3**. ¹H NMR (400 MHz, DMSO-*d*₆) δ = 8.35 (d, J = 2.0 Hz, 1H), 7.95 (dd, J = 2.0, 8.1 Hz, 1H), 7.77 (s, 1H), 7.68 (d, J = 0.9 Hz, 1H), 7.63 (d, J = 8.3 Hz, 1H), 7.55 (d, J = 8.1 Hz, 1H), 7.38 (dd, J = 1.5, 8.3 Hz, 1H), 7.22 (d, J = 0.9 Hz, 1H), 7.17 (s, 1H), 7.04 (br d, J = 7.8 Hz, 1H), 4.76 (s, 1H), 3.39 - 3.28 (m, 1H), 2.99 - 2.90 (m, 1H), 2.30 (d, J = 0.6 Hz, 3H), 2.17 (br d, J = 11.5 Hz, 2H), 1.94 (br d, J = 10.1 Hz, 2H), 1.60 (dq, J = 2.8, 12.7 Hz, 2H), 1.42 - 1.28 (m, 2H), 1.18 (d, J = 6.2 Hz, 6H), 1.09 (s, 9H). ESI [M+H] = 609.3.

Example 22. Preparation of isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-(2-(trifluoromethyl)pyridin-4-yl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (Compound 22)

From isopropyl (*trans*-4-(5-(4-bromo-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate, using the corresponding boric acid or boronate and under the same reaction conditions as for **Compound 3**. ¹H NMR (400 MHz, DMSO-*d*₆) δ = 8.91 (d, J = 5.0 Hz, 1H), 8.49 (d, J = 1.8 Hz, 1H), 8.27 (s, 1H), 8.21 (dd, J = 1.8, 8.1 Hz, 1H), 8.13 (d, J = 4.9 Hz, 1H), 7.81 (s, 1H), 7.69 (d, J = 8.1 Hz, 1H), 7.24 (s, 1H), 7.03 (br d, J = 7.8 Hz, 1H), 4.80 - 4.71 (m, 1H), 3.40 - 3.27 (m, 1H), 2.90 (s, 1H), 2.17 (br d, J = 11.7 Hz, 2H), 1.98 - 1.88 (m, 2H), 1.67 - 1.54 (m, 2H), 1.41 - 1.29 (m, 2H), 1.17 (d, J = 6.2 Hz, 6H), 1.06 (s, 9H). ESI [M+H] = 625.2.


Example 23. Preparation of isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-hydroxyphenyl)thiazol-2-yl)cyclohexyl)carbamate (Compound 23) and isopropyl (*trans*-4-(5-(4-hydroxy-2-sulfamoylphenyl)thiazol-2-yl)cyclohexyl)carbamate (Compound 24)



- 5 Isopropyl (*trans*-4-(5-(4-amino-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (1 g, 2.02 mmol, 1.0 *eq.*) was dissolved in H₂SO₄ (30%, aq., 12 mL), cooled to 0°C and added a solution of NaNO₂ (279 mg, 4.04 mmol, 2.0 *eq.*) in H₂O (2 mL). The mixture was stirred at 0°C for 1 hr, then added H₂O (28 mL) and the mixture was stirred at 65°C for 11 h. The reaction mixture was concentrated and the residue was purified by
- 10 prep-HPLC (column: Xtimate C18 10μ 250 mm *50mm; mobile phase: [water(0.1%TFA)-ACN]; B%: 30%-60%, 20min) to give isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-hydroxyphenyl)thiazol-2-yl)cyclohexyl)carbamate (**Compound 23**) (0.3 g, 581 μmol, 28.74% yield, 96% purity) as a yellow solid and isopropyl (*trans*-4-(5-(4-hydroxy-2-sulfamoylphenyl)thiazol-2-yl)cyclohexyl)carbamate (**Compound 24**) (0.3 g, 669 μmol, 33%
- 15 yield, 98% purity) as a yellow solid.


- ¹H NMR (400MHz, methanol-*d*₄) δ = 7.75 (s, 1H), 7.60 (d, J=2.6 Hz, 1H), 7.32 (d, J=8.4 Hz, 1H), 7.03 (dd, J=2.6, 8.4 Hz, 1H), 4.87 - 4.81 (m, 1H), 3.47 (tt, J=3.8, 11.6 Hz, 1H), 3.06 (tt, J=3.5, 12.1 Hz, 1H), 2.30 - 2.21 (m, 2H), 2.14 - 2.03 (m, 2H), 1.72 (dq, J=3.1, 12.8 Hz, 2H), 1.43 (dq, J=3.2, 12.6 Hz, 2H), 1.24 (br d, J=6.1 Hz, 6H), 1.13 (s, 9H). ESI [M+H] = 496.2.
- 20 ¹H NMR (400MHz, methanol-*d*₄) δ = 7.77 (s, 1H), 7.58 (d, J=2.6 Hz, 1H), 7.33 (d, J=8.3 Hz, 1H), 7.03 (dd, J=2.6, 8.3 Hz, 1H), 4.87 - 4.82 (m, 1H), 3.47 (tt, J=4.0, 11.6 Hz, 1H), 3.06 (tt, J=3.3, 12.0 Hz, 1H), 2.29 - 2.18 (m, 2H), 2.15 - 2.00 (m, 2H), 1.72 (dq, J=2.9, 12.8 Hz, 2H), 1.43 (dq, J=3.4, 12.6 Hz, 2H), 1.24 (br d, J=6.1 Hz, 6H). ESI [M+H] = 440.1.

- Example 24. Preparation of isopropyl *N*-[*trans*-4-[5-[4-[2-(*tert*-butoxycarbonylamino)ethoxy]-2-(*tert*-butylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 25)**
- 25



To a solution of isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-hydroxyphenyl)thiazol-2-yl)cyclohexyl)carbamate (50 mg, 101 μ mol, 1.0 *eq.*) in DMF (1 mL) was added K_2CO_3 (42 mg, 303 μ mol, 3.0 *eq.*) and *tert*-butyl *N*-(2-bromoethoxy) carbamate (27 mg, 121 μ mol, 1.2 *eq.*). The mixture was stirred at 50°C for 12 h, then diluted with sat. aq. LiCl (10 mL) and extracted with EtOAc (20 mL x 2). The combined organic layers were washed with sat. aq. LiCl (10 mL x 2), dried over Na_2SO_4 , filtered and concentrated. The residue was purified by prep-HPLC (column: Ximate C18 150*25mm*5 μ m; mobile phase: [water(10mM NH_4HCO_3)-ACN]; B%: 50%-80%, 8min) to give isopropyl *N*-[*trans*-4-[5-[4-[2-(*tert*-butoxycarbonylamino)ethoxy]-2-(*tert*-butylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (4.98 mg, 7.72 μ mol, 8% yield, 99% purity) as a white solid. 1H NMR (400MHz, methanol-*d*₄) δ = 7.76 - 7.68 (m, 2H), 7.41 (d, *J*=8.6 Hz, 1H), 7.21 (dd, *J*=2.7, 8.6 Hz, 1H), 4.86 - 4.81 (m, 1H), 4.13 (t, *J*=5.4 Hz, 2H), 3.49 (t, *J*=5.4 Hz, 2H), 3.44 (br s, 1H), 3.09 - 2.95 (m, 1H), 2.25 (br d, *J*=12.5 Hz, 2H), 2.09 (br d, *J*=10.9 Hz, 2H), 1.79 - 1.62 (m, 2H), 1.47 (s, 9H), 1.45 - 1.36 (m, 2H), 1.25 (br d, *J*=6.1 Hz, 6H), 1.12 (s, 9H). ESI [$M+H$] = 639.3.


Example 25. Preparation of isopropyl (*trans*-4-(5-(4-(2-aminoethoxy)-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (Compound 26)



A solution of isopropyl *N*-[*trans*-4-[5-[4-[2-(*tert*-butoxycarbonylamino)ethoxy]-2-(*tert*-butylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (75 mg, 117.4 μ mol) in HCl/MeOH (4 M, 1 mL) was stirred at 25°C for 0.5 hr. The mixture was then concentrated and purified by prep-HPLC (column: Nano-Micro UniSil 5-100 C18 ULTRA 100X250mm 5 μ m; mobile phase: [water(0.1%TFA)-ACN]; B%: 25%-45%, 10min) to give isopropyl (*trans*-4-(5-(4-(2-aminoethoxy)-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (10.21 mg, 15.64 μ mol, 13% yield, 100% purity, TFA salt) as


a yellow solid. ^1H NMR (400MHz, methanol- d_4) δ = 7.69 (d, J =2.7 Hz, 1H), 7.60 (s, 1H), 7.35 (d, J =8.4 Hz, 1H), 7.17 (dd, J =2.7, 8.6 Hz, 1H), 4.74 - 4.68 (m, 1H), 4.24 (t, J =5.0 Hz, 2H), 3.40 - 3.35 (m, 1H), 3.32 (br t, J =4.8 Hz, 2H), 2.91 (tt, J =3.5, 12.0 Hz, 1H), 2.13 (br d, J =12.0 Hz, 2H), 1.97 (br d, J =9.9 Hz, 2H), 1.59 (dq, J =2.8, 12.8 Hz, 2H), 1.31 (dq, J =3.2, 12.6 Hz, 2H), 1.13 (br d, J =6.1 Hz, 6H), 0.98 (s, 9H). ESI $[\text{M}+\text{H}]^+ = 539.2$.

Example 26. Preparation of isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-isopropoxy phenyl)thiazol-2-yl)cyclohexyl)carbamate (Compound 27)



To a solution of isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-hydroxyphenyl)thiazol-2-yl)cyclohexyl)carbamate (30 mg, 60.5 μmol , 1.0 *eq.*) in DMF (5 mL) was added K_2CO_3 (16.7 mg, 121 μmol , 2 *eq.*) and 2-iodopropane (20.6 mg, 121 μmol , 2 *eq.*). The mixture was stirred at 80°C for 12 h, then concentrated and purified by prep-HPLC to give isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-isopropoxyphenyl)thiazol-2-yl)cyclohexyl)carbamate (18 mg, 33.19 μmol , 55% yield, 99% purity) as a white solid. ^1H NMR (400 MHz, methanol- d_4) δ = 7.69 (s, 1H), 7.62 (d, J = 2.6 Hz, 1H), 7.38 - 7.35 (m, 1H), 7.14 (dd, J = 2.6, 8.6 Hz, 1H), 4.84 - 4.78 (m, 1H), 4.70 (td, J = 6.0, 12.1 Hz, 1H), 3.45 (tt, J = 3.9, 11.6 Hz, 1H), 3.04 - 2.97 (m, 1H), 2.26 - 2.18 (m, 2H), 2.10 - 2.03 (m, 2H), 1.69 (dq, J = 3.0, 12.8 Hz, 2H), 1.46 - 1.38 (m, 2H), 1.36 (d, J = 6.0 Hz, 6H), 1.22 (br d, J = 6.2 Hz, 6H), 1.09 (s, 9H). ESI $[\text{M}+\text{H}]^+ = 538.2$.


Example 27. Preparation of isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-methoxy phenyl)thiazol-2-yl)cyclohexyl)carbamate (Compound 28)



To a solution of isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-hydroxy phenyl)thiazol-2-yl)cyclohexyl)carbamate (30 mg, 60.5 μmol , 1.0 *eq.*) in DMF (4 mL) was added NaHCO_3 (10 mg, 121 μmol , 2 *eq.*) and MeI (10.3 mg, 72.6 μmol , 1.2 *eq.*). The mixture was stirred at 60°C for 12 h, then concentrated and purified by prep-HPLC (column: Welch


Ultimate AQ-C18 150*30mm*5um;mobile phase: [water(0.1%TFA)-ACN];B%: 40%-70%,12min) to give isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-methoxyphenyl)thiazol-2-yl)cyclohexyl)carbamate (2.34 mg, 4.26 umol, 7% yield, 93% purity) as a pale yellow solid. ¹H NMR (400MHz, methanol-*d*₄) δ = 7.75 - 7.62 (m, 2H), 7.42 (d, J=8.4 Hz, 1H), 7.20 (dd, J=2.8, 8.5 Hz, 1H), 4.84 (br d, J=5.5 Hz, 1H), 3.92 (s, 3H), 3.53 - 3.38 (m, 1H), 3.11 - 2.97 (m, 1H), 2.25 (br d, J=13.1 Hz, 2H), 2.09 (br d, J=10.4 Hz, 2H), 1.78 - 1.64 (m, 2H), 1.49 - 1.37 (m, 2H), 1.31 - 1.21 (m, 6H), 1.11 (s, 9H). ESI [M+H] =510.2.

Example 28. Preparation of *tert*-butyl (2-(3-(*N*-(*tert*-butyl)sulfamoyl)-4-(2-(*trans*-4-((isopropoxycarbonyl)amino)cyclohexyl)thiazol-5-yl)phenoxy)ethyl)(methyl) carbamate (Compound 29)



A mixture of isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-hydroxy phenyl)thiazol-2-yl)cyclohexyl)carbamate (140 mg, 282 umol, 1.0 eq.), *tert*-butyl *N*-(2-hydroxyethyl)-*N*-methyl-carbamate (74 mg, 424 umol, 1.5 eq.) and PPh₃ (148 mg, 565 umol, 2.0 eq.) in anhydrous THF (2 mL) was degassed and purged with N₂ for 3 times, then DIAD (86 mg, 424 umol, 1.5 eq.) was added and the mixture was stirred at 30°C for 12 h under N₂ atmosphere. The reaction mixture was concentrated and the residue was purified prep-HPLC (column: Waters Xbridge 150*25 5u;mobile phase: [water(10mM NH₄HCO₃)-ACN];B%: 40%-70%,10min) to give *tert*-butyl (2-(3-(*N*-(*tert*-butyl)sulfamoyl)-4-(2-(*trans*-4-((isopropoxycarbonyl)amino)cyclohexyl)thiazol-5-yl)phenoxy)ethyl)(methyl)carbamate (7.58 mg, 98% purity) as a pale yellow solid. ¹H NMR (400MHz, methanol-*d*₄) δ = 7.70 (br s, 2H), 7.42 (br d, J=7.9 Hz, 1H), 7.23 (br s, 1H), 4.87 - 4.77 (m, 1H), 4.25 (br s, 2H), 3.69 (br s, 2H), 3.47 (br s, 1H), 3.00 (br s, 4H), 2.25 (br d, J=11.0 Hz, 2H), 2.09 (br d, J=10.5 Hz, 2H), 1.71 (br d, J=12.1 Hz, 2H), 1.54 - 1.35 (m, 11H), 1.25 (br d, J=5.0 Hz, 6H), 1.11 (br s, 9H). ESI [M+H] =653.3.

Example 29. Preparation of isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-(2-(dimethyl amino)ethoxy)phenyl)thiazol-2-yl)cyclohexyl)carbamate (Compound 30)




To a solution of isopropyl (*trans*-4-(5-(4-(2-aminoethoxy)-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (13 mg, 24.1 μ mol, 1.0 *eq.*) in MeOH (1 mL) was added AcOH (one drop), HCHO (3.6 mg, 120.7 μ mol, 5.0 *eq.*) followed by

5 NaBH₃CN (4.5 mg, 72.4 μ mol, 3.0 *eq.*) after 0.5 hr. The mixture was stirred at 25°C for 12 h and then concentrated. The residue was purified by prep-HPLC (column: Welch Ultimate AQ-C18 150*30mm*5 μ m; mobile phase: [water(0.1%TFA)-ACN]; B%: 25%-55%, 12min) to give isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-(2-(dimethylamino)ethoxy)phenyl)thiazol-2-yl)cyclohexyl) carbamate (20.6 mg, 99% purity, TFA salt) as a white solid. ¹H

10 NMR (400MHz, methanol-*d*₄) δ = 7.81 (br s, 1H), 7.73 (s, 1H), 7.49 (br d, *J*=8.3 Hz, 1H), 7.31 (br d, *J*=7.0 Hz, 1H), 4.86 (br d, *J*=5.7 Hz, 1H), 4.49 (br s, 2H), 3.68 (br s, 2H), 3.55 - 3.41 (m, 1H), 3.03 (s, 7H), 2.25 (br d, *J*=11.6 Hz, 2H), 2.09 (br d, *J*=10.9 Hz, 2H), 1.71 (q, *J*=12.0 Hz, 2H), 1.50 - 1.36 (m, 2H), 1.24 (br d, *J*=5.4 Hz, 6H), 1.10 (s, 9H). ESI [M+H] = 567.2.

15 **Example 30. Preparation of isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-(2-methoxyethoxy)phenyl)thiazol-2-yl)cyclohexyl)carbamate (Compound 31)**




A mixture of isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-hydroxyphenyl)thiazol-2-yl)cyclohexyl)carbamate (30 mg, 60.5 μ mol, 1.0 *eq.*), 1-bromo-2-methoxyethane (6.7 mg, 48.4 μ mol, 0.8 *eq.*) and K₂CO₃ (16.73 mg, 121.05 μ mol, 2 *eq.*) in DMF (2 mL) was

20 stirred at 30°C for 12 h under N₂ atmosphere. The mixture was concentrated and purified by prep-HPLC (column: Welch Ultimate AQ-C18 150X30mmX5 μ m; mobile phase: [water(0.1%TFA)-ACN]; B%: 40%-70%, 12min) to give isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-(2-methoxyethoxy)phenyl)thiazol-2-yl)cyclohexyl)carbamate (19.84 mg,

25 35.70 μ mol, 59% yield, 99% purity) as pale yellow solid. ¹H NMR (400 MHz, methanol-*d*₄) δ = 7.79 (s, 1H), 7.72 (d, *J* = 2.7 Hz, 1H), 7.42 (s, 1H), 7.26 - 7.20 (m, 1H), 4.87 - 4.74 (m,


1H), 4.25 (dd, J = 3.7, 5.3 Hz, 2H), 3.81 (dd, J = 3.7, 5.3 Hz, 2H), 3.54 - 3.43 (m, 4H), 3.12 - 3.02 (m, 1H), 2.31 - 2.19 (m, 2H), 2.13 - 2.07 (m, 2H), 1.73 (br dd, J = 2.8, 12.7 Hz, 2H), 1.50 - 1.36 (m, 2H), 1.24 (br d, J = 6.2 Hz, 6H), 1.13 (s, 9H). ESI [M+H] = 554.3.

Example 31. Preparation of isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-(2-(methyl amino)ethoxy)phenyl)thiazol-2-yl)cyclohexyl)carbamate (Compound 32)




A solution of *tert*-butyl (2-(3-(*N*-(*tert*-butyl)sulfamoyl)-4-(2-(*trans*-4-((isopropoxy carbonyl)amino)cyclohexyl)thiazol-5-yl)phenoxy)ethyl)(methyl)carbamate (0.1 g, 153 μ mol, 1.0 *eq.*) in HCl/MeOH (4 M, 5 mL) was stirred at 20°C for 30 mins. The reaction mixture was concentrated and purified by prep-HPLC (column: Waters Xbridge 150*25 μ m; mobile phase: [water(10mM NH₄HCO₃)-ACN]; B%: 20%-50%, 10min) to give isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-(2-(methylamino)ethoxy)phenyl) thiazol-2-yl)cyclohexyl)carbamate (6.84 mg, 12.25 μ mol, 8% yield, 99% purity) as a yellow solid. ¹H NMR (400MHz, methanol-*d*₄) δ = 7.78 - 7.69 (m, 2H), 7.43 (d, J=8.4 Hz, 1H), 7.23 (dd, J=2.5, 8.5 Hz, 1H), 4.86 - 4.80 (m, 1H), 4.22 (br t, J=4.8 Hz, 2H), 3.55 - 3.41 (m, 1H), 3.11 - 2.95 (m, 3H), 2.50 (br s, 3H), 2.25 (br d, J=12.2 Hz, 2H), 2.09 (br d, J=10.8 Hz, 2H), 1.79 - 1.63 (m, 2H), 1.49 - 1.35 (m, 2H), 1.24 (br d, J=6.1 Hz, 6H), 1.11 (s, 9H). ESI [M+H] = 553.2.

Example 32: Preparation of oxetan-3-yl *trans*-N-[4-[5-[2-(*tert*-butylsulfamoyl)-4-(isobutoxymethyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 34)




a) Synthesis of 4-bromo-3-(*tert*-butylsulfamoyl)benzoic acid



5

To a solution of 2-methylpropan-2-amine (2.9 g, 40.1 mmol, 3.0 *eq.*) in DCM (80 mL) was added 4-bromo-3-chlorosulfonylbenzoic acid (4.0 g, 13.4 mmol, 1 *eq.*) with THF (20 mL) at -10°C, then the mixture was stirred at 15°C for 2 h. The reaction mixture was concentrated under reduced pressure, the residue was diluted with HCl (20 mL) and extracted with EtOAc 150 mL (50 mL X 3). The combined organic layers were dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue which was triturated with petroleum ether yielding 4-bromo-3-(*tert*-butylsulfamoyl)benzoic acid (4.5 g, 13.4 mmol, 100 % yield) as a white solid. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.70 (d, J=1.98 Hz, 1H), 8.03 (dd, J=8.16, 1.98 Hz, 1H) 7.63 (d, J=8.38 Hz, 1H) 1.19 (s, 9H).


15 b) Synthesis of 2-bromo-N-*tert*-butyl-5-(hydroxymethyl)benzenesulfonamide



To a solution of 4-bromo-3-(*tert*-butylsulfamoyl)benzoic acid (3.5 g, 10.4 mmol, 1 *eq.*) in THF (20 mL) was added BH₃-Me₂S (10 M, 12.49 mL, 12 *eq.*) at 0°C and stirred at 15°C for 12 h. The reaction mixture was quenched by addition MeOH (15 mL) at 15°C,


then the reaction mixture was concentrated under reduced pressure. The crude product was purified by reversed-phase HPLC (TFA condition) to yield 2-bromo-*N*-*tert*-butyl-5-(hydroxymethyl)benzenesulfonamide (3g) as a white solid. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.17 (d, *J*=2.08 Hz, 1H), 7.76 (d, *J*=8.07 Hz, 1H), 7.42 - 7.52 (m, 1H), 4.65 (s, 2H), 1.21 (s, 9H). ESI[M-H] = 319.9/321.9.

c) Synthesis of 2-bromo-5-(bromomethyl)-*N*-*tert*-butyl-benzenesulfonamide




To a solution of 2-bromo-*N*-*tert*-butyl-5-(hydroxymethyl)benzenesulfonamide (2.9 g, 9 mmol, 1 *eq.*) in DCM (50 mL) was added CBr₄ (4.6 g, 13.0 mmol, 1.5 *eq.*) and PPh₃ (3.6 g, 13.0 mmol, 1.5 *eq.*) and stirred at 15°C for 2 h. The reaction mixture was diluted with H₂O (15 mL) and extracted with DCM 150 mL (50 mL X 3). The combined organic layers were dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by column chromatography (SiO₂, petroleum ether: ethyl acetate=1:0 to 8:1 to 5:1 to 2:1 to 1:1) to yield 2-bromo-5-(bromomethyl)-*N*-*tert*-butyl-benzenesulfonamide (5.1 g, 13.0 mmol, quantitative yield) as a white solid. ESI[M-H] = 383.8.

d) Synthesis of 2-bromo-*N*-*tert*-butyl-5-(isobutoxymethyl)benzene sulfonamide




To a solution of 2-methylpropan-1-ol (3.3 g, 45.0 mmol, 5 *eq.*) in DMF (15 mL) was added NaH (1.1 g, 27.0 mmol, 60% purity, 3.0 *eq.*) and 2-bromo-5-(bromomethyl)-*N*-*tert*-butyl-benzenesulfonamide (3.5 g, 9.0 mmol, 1.0 *eq.*). The mixture was stirred at 25°C for 2 h. The reaction mixture was poured into NaCl (sat.aq.) (15 mL) at 25°C, extracted with EtOAc (30 mL), The combined organic layers were washed with NaCl 60 mL (20 mL X 3), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by prep-HPLC (column: Welch Xtimate C18 250*50mm*10um;mobile phase: [water(0.1%TFA)-ACN];B%: 45%-75%,20min) to yield 2-bromo-*N*-*tert*-butyl-5-(isobutoxymethyl)benzenesulfonamide (820 mg, 2.2 mmol, 24% yield) as a white solid. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.12 (d, *J*=1.98 Hz, 1 H), 7.74 (d, *J*=8.16 Hz, 1 H), 7.41 (dd, *J*=8.16, 2.21 Hz, 1 H), 3.25 - 3.33 (m, 4 H), 1.85 - 1.95 (m, 1 H), 1.18 (s, 9 H), 0.93 (d, *J*=6.84 Hz, 6 H).

- e) Synthesis of *N-tert*-butyl-5-(isobutoxymethyl)-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzenesulfonamide



- From 2-bromo-*N-tert*-butyl-5-(isobutoxymethyl)benzenesulfonamide, using General Method B. ESI[M+H] = 426.2.

- f) Synthesis of oxetan-3-yl *trans*-*N*-[4-[5-[2-(*tert*-butylsulfamoyl)-4-(isobutoxymethyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 34**)




- From *N-tert*-butyl-5-(isobutoxymethyl)-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzenesulfonamide, using General Method C. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.15 (d, J=1.32 Hz, 1H), 7.75 (s, 1H), 7.57 (dd, J=7.72, 1.32 Hz, 1H), 7.45 (d, J=7.72 Hz, 1H), 5.31 - 5.40 (m, 1H), 4.87 (s, 1H), 4.84 (br s, 1H), 4.60 (s, 4H), 3.39 - 3.50 (m, 1H), 3.29 (d, J=1.98 Hz, 2H), 2.98 - 3.06 (m, 1H), 2.24 (br d, J=12.57 Hz, 2H), 2.02 - 2.11 (m, 2H), 1.92 (dt, J=13.40, 6.64 Hz, 1H), 1.64 - 1.76 (m, 2H), 1.38 - 1.49 (m, 2H), 1.08 (s, 9H), 0.95 (d, J=6.84 Hz, 6H). ESI [M+H] = 580.3.


Example 33. Preparation of isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-(isobutoxymethyl)phenyl)thiazol-2-yl)cyclohexyl]carbamate (Compound 33**)**

- Following the same protocol and under the same reaction conditions as for Compound 34, Compound 33 was prepared. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.06 (s, 1H), 7.67 (s, 1H), 7.48 (br d, J=7.58 Hz, 1H), 7.36 (d, J=7.70 Hz, 1H), 4.51 (s, 2H), 3.30 - 3.39 (m, 1H), 3.20 (br s, 3H), 2.92 (tt, J=11.83, 3.03 Hz, 1H), 2.14 (br d, J=12.47 Hz, 2H), 1.98 (br d, J=11.00 Hz, 2H), 1.75 - 1.89 (m, 1H), 1.55 - 1.67 (m, 2H), 1.32 (qd, J=12.59, 2.69 Hz, 2H), 1.13 (br d, J=5.99 Hz, 6H), 1.00 (s, 9H), 0.86 (d, J=6.72 Hz, 6H). ESI [M+H] = 566.3.

- Example 34. Preparation of isopropyl *trans*-*N*-[4-[5-[4-cyano-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 36**)**




a) Synthesis of 2-bromo-*N*-ethyl-5-nitro-benzenesulfonamide



From 1-(2-bromo-5-nitro-phenyl)sulfonyl chloride and ethylamine, using General


- 5 Method G. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.82 (d, *J* = 2.9 Hz, 1H), 8.30 (dd, *J* = 2.6, 8.6 Hz, 1H), 8.07 (d, *J* = 8.6 Hz, 1H), 3.01 (q, *J* = 7.3 Hz, 2H), 1.08 (t, *J* = 7.3 Hz, 3H).

b) Synthesis of 5-amino-2-bromo-*N*-ethyl-benzenesulfonamide




- To a solution of 2-bromo-*N*-ethyl-5-nitro-benzenesulfonamide (100.0 g, 323.5 mmol, 10 1.0 *eq.*) in EtOH (600 mL), H₂O (200 mL) and THF (600 mL) was added Fe (90.3 g, 1.6 mol, 5.0 *eq.*) and NH₄Cl (51.9 g, 970.4 mmol, 3.0 *eq.*). The mixture was stirred at 80°C for 2 h and filtered. Then concentrated under reduced pressure, the residue diluted with H₂O (60 mL) and extracted with EtOAc 1200 mL (400 mL X 3). The combined organic layers were dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by 15 column chromatography (SiO₂, petroleum ether: ethyl acetate = 10:1 to 0:1) to yield 5-amino-2-bromo-*N*-ethyl-benzenesulfonamide (82.0 g, 293.7 mmol, 91% yield) as a yellow solid. ¹H NMR (400MHz, methanol-*d*₄) δ = 7.44 - 7.37 (m, 2H), 6.74 (dd, *J*=2.9, 8.5 Hz, 1H), 2.95 (q, *J*=7.2 Hz, 2H), 1.08 (t, *J*=7.3 Hz, 3H). ESI [M+H] = 279.0/281.0.

- c) Synthesis of 5-amino-*N*-ethyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzenesulfonamide 20




From 5-amino-2-bromo-*N*-ethyl-benzenesulfonamide, using General Method B. ¹H NMR (400 MHz, methanol-*d*₄) δ = 7.97 (s, 1H), 7.49 (br d, *J*=7.94 Hz, 2H), 2.98 (s, 2H), 1.35 (s, 12H), 1.00-1.08 (m, 3H). ESI [M+H] = 327.3.

- 5 d) Synthesis of isopropyl *trans*-*N*-[4-[5-[4-amino-2-(ethylsulfamoyl)phenyl] thiazol-2-yl]cyclohexyl]carbamate




From 5-amino-*N*-ethyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzene sulfonamide, using General Method C. ESI [M+H] = 467.3.

- 10 e) Synthesis of isopropyl *trans*-*N*-[4-[5-[4-bromo-2-(ethylsulfamoyl)phenyl] thiazol-2-yl]cyclohexyl]carbamate




From isopropyl *trans*-*N*-[4-[5-[4-amino-2-(ethylsulfamoyl)phenyl] thiazol-2-yl] cyclohexyl]carbamate, using General Method L. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.21 (d, *J*=1.47 Hz, 1 H), 7.78 - 7.86 (m, 2 H), 7.44 (d, *J*=8.19 Hz, 1 H), 4.85 (br d, *J*=5.87 Hz, 1 H), 3.48 (ddd, *J*=11.43, 7.70, 3.97 Hz, 1 H), 3.00 - 3.10 (m, 1 H), 2.89 (q, *J*=7.21 Hz, 2 H), 2.22 - 2.30 (m, 2 H), 2.10 (br d, *J*=10.64 Hz, 2 H), 1.72 (qd, *J*=12.76, 2.57 Hz, 2 H), 1.43 (qd, *J*=12.55, 2.93 Hz, 2 H), 1.25 (br d, *J*=5.99 Hz, 6 H), 1.04 (t, *J*=7.21 Hz, 3 H). ESI [M+H] = 530.1/532.1.

- 20 f) Synthesis of isopropyl *trans*-*N*-[4-[5-[4-cyano-2-(ethylsulfamoyl) phenyl] thiazol-2-yl]cyclohexyl]carbamate (**Compound 36**)




A solution of isopropyl *trans*-N-[4-[5-[4-bromo-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (60 mg, 113 μ mol, 1 *eq.*), Zn(CN)₂ (27 mg, 226 μ mol, 2 *eq.*), Pd(PPh₃)₄ (13 mg, 11 μ mol, 0.1 *eq.*) in NMP (2 mL) was stirred for 1 h under microwave at 170°C. Then diluted with H₂O (10 mL), extracted with EtOAc 40 mL (20 mL X 2). The combined organic layers were dried over Na₂SO₄, filtered and concentrated to give a residue. The residue was purified by prep-TLC (SiO₂, petroleum ether: ethyl acetate = 0:1) and prep-HPLC (column: Waters Xbridge BEH C18 100*25mm*5 μ m; mobile phase: [water(10mM NH₄HCO₃)-ACN]; B%: 20%-50%, 10min.) to yield isopropyl *trans*-N-[4-[5-[4-cyano-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (810 μ g, 1 μ mol, 1.3% yield, 86% purity) as a yellow solid. ¹H NMR (400MHz, methanol-*d*₄) δ = 8.40 (s, 1H), 8.01 (d, J=8.1 Hz, 1H), 7.87 (s, 1H), 7.76 - 7.65 (m, 1H), 4.83 (br s, 1H), 3.55 - 3.39 (m, 1H), 3.14 - 2.99 (m, 1H), 2.95 - 2.85 (m, 2H), 2.27 (br d, J=12.6 Hz, 2H), 2.10 (br d, J=11.7 Hz, 2H), 1.79 - 1.67 (m, 2H), 1.50 - 1.37 (m, 2H), 1.25 (br d, J=6.1 Hz, 6H), 1.11 - 0.98 (m, 3H). ESI [M+H]⁺ = 477.1.

Example 35. Preparation of isopropyl *trans*-N-[4-[5-[2-(*tert*-butylsulfamoyl)-4-[2-(isopropylamino)-2-oxo-ethyl]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 40)




a) Synthesis of 2-[4-bromo-3-(*tert*-butylsulfamoyl)phenyl]-N-isopropyl-acetamide




From 2-[4-bromo-3-(*tert*-butylsulfamoyl)phenyl]-*N*-isopropyl-acetic acid and isopropylamine, using General Method A. ^1H NMR (400MHz, methanol- d_4) δ = 7.97 (s, 1H), 7.71 (d, J =7.9 Hz, 1H), 7.37 (dd, J =2.1, 8.0 Hz, 1H), 4.00 - 3.86 (m, 1H), 3.50 (s, 2H), 1.18 (s, 9H), 1.13 (d, J =6.6 Hz, 6H). ESI[M+H] = 391.0/393.0.

- 5 b) Synthesis of 2-[3-(*tert*-butylsulfamoyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]-*N*-isopropyl-acetamide



- From 2-[4-bromo-3-(*tert*-butylsulfamoyl)phenyl]-*N*-isopropyl-acetamide, using General Method D. ^1H NMR (400MHz, methanol- d_4) δ = 7.92 (d, J =1.1 Hz, 1H), 7.66 (d, J =7.7 Hz, 1H), 7.55 - 7.39 (m, 1H), 3.98 - 3.89 (m, 1H), 3.58 - 3.51 (m, 2H), 1.40 (s, 9H), 1.25 - 1.16 (m, 12H), 1.15 - 1.10 (m, 6H). ESI[M+H] = 439.2.

- c) Synthesis of isopropyl *trans*-*N*-[4-[5-[2-(*tert*-butylsulfamoyl)-4-[2-(isopropylamino)-2-oxo-ethyl]phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 40**)




- 15 From 2-[3-(*tert*-butylsulfamoyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]-*N*-isopropyl-acetamide, using General Method C. ^1H NMR (400MHz, methanol- d_4) δ = 8.11 (br s, 1H), 7.77 (s, 1H), 7.54 (br d, J =7.3 Hz, 1H), 7.42 (br d, J =7.5 Hz, 1H), 4.82 - 4.74 (m, 1H), 4.01 - 3.87 (m, 1H), 3.57 (br s, 2H), 3.45 (br s, 1H), 3.08 - 2.95 (m, 1H), 2.23 (br d, J =12.3 Hz, 2H), 2.06 (br d, J =10.8 Hz, 2H), 1.78 - 1.61 (m, 2H), 1.49 - 1.33 (m, 2H), 1.21 (br d, J =4.9 Hz, 6H), 1.14 (br d, J =6.4 Hz, 6H), 1.08 (s, 9H). ESI [M+H] = 579.3.

Example 36. Preparation of oxetan-3-yl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-(2-(isopropylamino)-2-oxoethyl)phenyl)thiazol-2-yl)cyclohexyl]carbamate (Compound 41**)**


- Following the same protocol and under the same reaction conditions as for Compound 40, Compound 41 was prepared. ^1H NMR (400MHz, DMSO- d_6) δ = 8.05 (br d, J =7.5 Hz, 1H), 7.97 (s, 1H), 7.67 (s, 1H), 7.49 - 7.41 (m, 2H), 7.40 - 7.34 (m, 1H), 6.96 (s, 1H), 5.26 (quin, J =5.7 Hz, 1H), 4.73 (t, J =6.8 Hz, 2H), 4.53 - 4.37 (m, 2H), 3.78 (qd, J =6.8, 13.6 Hz, 1H), 3.47 (s, 2H), 3.29 (s, 1H), 2.98 - 2.84 (m, 1H), 2.13 (br d, J =12.1 Hz, 2H), 1.91 (br d,

J=10.1 Hz, 2H), 1.66 - 1.46 (m, 2H), 1.42 - 1.23 (m, 2H), 1.03 (d, J=6.6 Hz, 6H), 1.01 (s, 9H). ESI [M+H] = 593.3.

Example 37. Preparation of isopropyl *trans*-N-[4-[5-(2-sulfamoylphenyl)thiazol-2-yl]cyclohexyl]carbamate (Compound 44)




a) Synthesis of isopropyl *trans*-N-[4-[5-(2-aminophenyl)thiazol-2-yl]cyclohexyl]carbamate



From 2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline, using General Method C. ¹H NMR (400 MHz, methanol-*d*₄) δ = 7.46 - 7.27 (m, 1H), 7.21 - 7.06 (m, 2H), 6.83 (d, J = 8.2 Hz, 1H), 6.71 (t, J = 7.5 Hz, 1H), 4.82 (br d, J = 6.4 Hz, 1H), 3.52 - 3.38 (m, 1H), 2.98 (tt, J = 3.5, 12.1 Hz, 1H), 2.26 - 2.16 (m, 2H), 2.06 (br d, J = 10.6 Hz, 2H), 1.68 (dq, J = 2.8, 12.9 Hz, 2H), 1.40 (dq, J = 3.3, 12.6 Hz, 2H), 1.24 - 1.18 (m, 6H). ESI [M+H] = 360.1.

b) Synthesis of isopropyl *trans*-N-[4-[5-(2-chlorosulfonylphenyl)thiazol-2-yl]cyclohexyl]carbamate

15




To a solution of isopropyl *trans*-N-[4-[5-(2-aminophenyl)thiazol-2-yl]cyclohexyl]carbamate (1.0 g, 2.8 mmol, 1.0 *eq.*) in H₂O (1 mL) and AcOH (5 mL) was added conc. HCl (5 mL) at 0°C followed by a solution of NaNO₂ (960 mg, 14 mmol, 5 *eq.*) in H₂O (1 mL), the mixture was stirred at 0°C for 2.5 h (mixture 1). To another mixture of CuCl₂ (374 mg, 3 mmol, 1 *eq.*) in H₂O (4 mL) was added a solution of SO₂ in AcOH (20 mL, 2 M) and stirred at 25°C for 1 h (mixture 2). To the mixture 2 was added the mixture 1 at 0°C. The mixture was

20

stirred at 0°C for 0.5 h. The reaction mixture was quenched with H₂O (20 mL) at 0°C and filtered to yield crude product isopropyl *trans-N*-[4-[5-(2-chlorosulfonylphenyl)thiazol-2-yl]cyclohexyl]carbamate (1.2 g, crude) as a pale yellow solid, without any purification. ESI [M+H] = 443.0.

- 5 c) Synthesis of isopropyl *trans-N*-[4-[5-(2-sulfamoylphenyl)thiazol-2-yl]cyclohexyl]carbamate (**Compound 44**)



- To a solution of TEA (137 mg, 1 mmol, 2 *eq.*), NH₃.H₂O (395 mg, 3 mmol, 30% purity, 5 *eq.*) in THF (2 mL) was added a solution of isopropyl *trans-N*-[4-[5-(2-chlorosulfonylphenyl)thiazol-2-yl]cyclohexyl]carbamate (300 mg, 677 μmol, 1 *eq.*) in THF (2 mL) at 0°C. The mixture was stirred at 20°C for 1 h and concentrated. The residue was purified by prep-HPLC (column: Welch Ultimate AQ-C18 150*30mm*5μm; mobile phase: [water(0.1%TFA)-ACN]; B%: 35%-65%, 12min) to yield isopropyl *trans-N*-[4-[5-(2-sulfamoylphenyl)thiazol-2-yl]cyclohexyl]carbamate (92 mg, 216 μmol, 32% yield, 99% purity) as a yellow solid. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.22 - 8.12 (m, 1H), 7.87 - 7.79 (m, 1H), 7.70 - 7.58 (m, 2H), 7.56 - 7.50 (m, 1H), 4.87 - 4.79 (m, 1H), 3.55 - 3.40 (m, 1H), 3.12 - 2.98 (m, 1H), 2.32 - 2.18 (m, 2H), 2.14 - 2.01 (m, 2H), 1.79 - 1.65 (m, 2H), 1.50 - 1.35 (m, 2H), 1.25 (br d, J = 6.2 Hz, 6H). ESI [M+H]=424.1.

- Example 38. Preparation of isopropyl (*trans*-4-(5-(2-(*N*-ethylsulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (Compound 45)**

- Following the same protocol and under the same reaction conditions as for Compound 44, Compound 45 was prepared. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.06 (dd, J = 1.0, 7.8 Hz, 1H), 7.72 (s, 1H), 7.68 - 7.55 (m, 2H), 7.49 (dd, J = 1.0, 7.4 Hz, 1H), 4.84 - 4.76 (m, 1H), 3.45 (ddd, J = 4.1, 7.6, 11.5 Hz, 1H), 3.00 (tt, J = 3.4, 12.1 Hz, 1H), 2.85 (q, J = 7.3 Hz, 2H), 2.31 - 2.17 (m, 2H), 2.12 - 1.98 (m, 2H), 1.69 (dq, J = 2.9, 12.8 Hz, 2H), 1.41 (dq, J = 3.1, 12.6 Hz, 2H), 1.22 (br d, J = 6.2 Hz, 6H), 1.00 (t, J = 7.2 Hz, 3H). ESI [M+H]=452.1.

Example 39. Preparation of isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (Compound 46)


- Following the same protocol and under the same reaction conditions as for Compound 44, Compound 46 was prepared. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.15 (dd, J = 1.3, 7.7

Hz, 1H), 7.76 (s, 1H), 7.68 - 7.53 (m, 2H), 7.48 (dd, J = 1.3, 7.5 Hz, 1H), 4.86 - 4.76 (m, 1H), 3.45 (tt, J = 3.8, 11.5 Hz, 1H), 3.02 (tt, J = 3.5, 12.0 Hz, 1H), 2.30 - 2.17 (m, 2H), 2.12 - 2.01 (m, 2H), 1.70 (dq, J = 2.6, 12.9 Hz, 2H), 1.41 (dq, J = 3.1, 12.6 Hz, 2H), 1.22 (br d, J = 6.2 Hz, 6H), 1.07 (s, 9H). ESI [M+H]⁺=480.2.


5 **Example 40. Preparation of isopropyl *trans*-N-[4-[5-(2-methylsulfamoylphenyl)thiazol-2-yl]cyclohexyl]carbamate (Compound 93)**

Following the same protocol and under the same reaction conditions as for Compound 44, Compound 93 was prepared. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.18 - 7.99 (m, 1H), 7.80 - 7.77 (m, 1H), 7.71 - 7.61 (m, 2H), 7.54 (dd, J = 1.2, 7.2 Hz, 1H), 4.87 - 4.80 (m, 1H), 3.54 - 3.43 (m, 1H), 3.11 - 2.99 (m, 1H), 2.30 - 2.23 (m, 2H), 2.13 - 2.04 (m, 2H), 1.80 - 1.65 (m, 2H), 1.51 - 1.37 (m, 2H), 1.25 (br d, J = 6.1 Hz, 6H). ESI [M+H]⁺=438.1.

Example 41. Preparation of oxetan-3-yl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-(isobutoxymethyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 52)




15 a) **Synthesis of 2-bromo-N-ethyl-5-(hydroxymethyl)benzenesulfonamide**



To a solution of 4-bromo-3-(ethylsulfamoyl)benzoic acid (1.8 g, 5.8 mmol, 1.0 eq.) in THF (40 mL) was added BH₃·Me₂S (10.0 M, 5.8 mL, 10.0 eq.) at 0°C and stirred at 25°C for 12 h. The reaction mixture was quenched by MeOH 50 mL at 0°C, and was stirred at 50°C for 1 h. Then the reaction mixture was concentrated under reduced pressure. The crude product was purified by reversed-phase HPLC (0.1% TFA condition) to yield 2-bromo-


N-ethyl-5-(hydroxymethyl)benzenesulfonamide (1.2 g, 4.2 mmol, 71% yield) as a yellow oil. ESI [M-H] = 291.8/293.8.

b) Synthesis of [4-bromo-3-(ethylsulfamoyl)phenyl]methyl methanesulfonate




To a solution of 2-bromo-*N*-ethyl-5-(hydroxymethyl)benzenesulfonamide (1.1 g, 3.8 mmol, 1.0 *eq*) in DCM (15 mL) was added TEA (770 mg, 8 mmol, 2 *eq*) and methanesulfonyl chloride (0.15 g, 1.31 mmol, 3.44e-1 *eq.*) at 0°C. The mixture was stirred at 25°C for 2 h. The reaction mixture was concentrated under reduced pressure. The residue was diluted with H₂O 20 mL and extracted with DCM 60mL (20 mL X 3). The combined organic layers were dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO₂, petroleum ether: ethyl acetate=1:0 to 0:1) to yield [4-bromo-3-(ethylsulfamoyl)phenyl]methyl methanesulfonate (600 mg, 1 mmol, 42% yield) as a pale yellow solid. ESI [M-H] = 369.8/371.8.

c) Synthesis of 2-bromo-*N*-ethyl-5-(isobutoxymethyl)benzenesulfonamide




To a solution of 2-methylpropan-1-ol (1 g, 13 mmol, 8 *eq*) in DMF (10 mL) was added NaH (58 mg, 1.5 mmol, 60% purity, 0.9 *eq*) at 0°C. The mixture was stirred at 25°C for 30 min. To the mixture was added a solution of [4-bromo-3-(ethylsulfamoyl)phenyl]methyl methanesulfonate (600 mg, 1.61 mmol, 1 *eq*) in DMF (10 mL), then stirred at 25°C for 1 h. The reaction was quenched with sat.aq. NH₄Cl 50 mL at 0°C, extracted with EtOAc 150 mL (50 mL X 3). The combined organic layers were washed with sat.aq. NaCl 45 mL (15 mL X 3), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO₂, petroleum ether: ethyl acetate=1:0 to 0:1) to yield 2-bromo-*N*-ethyl-5-(isobutoxymethyl)benzenesulfonamide (260 mg, 742.28 umol, 46% yield) as a pale yellow oil. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.07 (d, J = 1.8 Hz, 1H), 7.76 (d, J = 7.9 Hz, 1H), 7.44 (dd, J = 2.0, 8.2 Hz, 1H), 4.53 (s, 2H), 3.29 (br s, 2H), 2.94 (q, J = 7.3 Hz, 2H), 1.90 (td, J = 6.6, 13.2 Hz, 1H), 1.05 (t, J = 7.3 Hz, 3H), 0.94 (d, J = 6.8 Hz, 6H). ESI [M-H] = 347.9/349.9.

d) Synthesis of *N*-ethyl-5-(isobutoxymethyl)-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzenesulfonamide



From 2-bromo-*N*-ethyl-5-(isobutoxymethyl)benzenesulfonamide, using General Method B. ¹H NMR (400 MHz, methanol-*d*₄) δ = 7.76 (d, *J* = 19.6 Hz, 1H), 7.58 - 7.41 (m, 2H), 4.48 (d, *J* = 3.5 Hz, 2H), 3.18 (s, 2H), 2.89 - 2.72 (m, 2H), 1.87 - 1.72 (m, 1H), 1.22 - 1.05 (m, 12H), 0.94 (td, *J* = 7.3, 9.1 Hz, 3H), 0.84 (d, *J* = 6.7 Hz, 6H). ESI [M+H] = 398.2.

e) Synthesis of oxetan-3-yl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-(isobutoxymethyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 52**)




From *N*-ethyl-5-(isobutoxymethyl)-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzenesulfonamide, using General Method C. ¹H NMR (400 MHz, DMSO-*d*₆) δ = 7.90 (s, 1H), 7.69 (s, 1H), 7.55 (br d, *J* = 7.9 Hz, 1H), 7.49 - 7.38 (m, 3H), 5.26 (quin, *J* = 5.7 Hz, 1H), 4.73 (t, *J* = 6.8 Hz, 2H), 4.56 (s, 2H), 4.48 - 4.39 (m, 2H), 3.29 (br s, 1H), 3.23 (d, *J* = 6.6 Hz, 2H), 2.92 (tt, *J* = 3.3, 11.8 Hz, 1H), 2.77 (br d, *J* = 6.8 Hz, 2H), 2.13 (br d, *J* = 11.9 Hz, 2H), 1.97 - 1.88 (m, 2H), 1.87 - 1.78 (m, 1H), 1.63 - 1.48 (m, 2H), 1.43 - 1.28 (m, 2H), 0.93 (t, *J* = 7.3 Hz, 3H), 0.88 (d, *J* = 6.8 Hz, 6H). ESI [M+H] = 552.2.


Example 42. Preparation of isopropyl (*trans*-4-(5-(2-(*N*-ethylsulfamoyl)-4-(isobutoxymethyl)phenyl)thiazol-2-yl)cyclohexyl]carbamate (Compound 51**)**

Following the same protocol and under the same reaction conditions as for Compound 52, Compound 51 was prepared. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.08 (s, 1H), 7.77 (s, 1H), 7.62 (d, *J* = 7.8 Hz, 1H), 7.50 (d, *J* = 7.8 Hz, 1H), 4.84 (br s, 1H), 4.63 (s, 2H), 3.47 (tdd, *J* = 3.9, 7.7, 11.4 Hz, 1H), 3.33 (s, 2H), 3.04 (tt, *J* = 3.3, 12.0 Hz, 1H), 2.89 (q, *J* = 7.2 Hz, 2H), 2.26 (br d, *J* = 12.5 Hz, 2H), 2.13 - 2.05 (m, 2H), 2.01 - 1.88 (m, 1H), 1.72 (dq, *J* = 2.4, 12.8 Hz, 2H), 1.50 - 1.37 (m, 2H), 1.24 (br d, *J* = 6.0 Hz, 6H), 1.04 (t, *J* = 7.2 Hz, 3H), 0.98 (d, *J* = 6.7 Hz, 6H). ESI [M+H] = 538.3.

Example 43. Preparation of isopropyl *trans*-*N*-[4-[5-[2-(azetidin-1-ylsulfonyl)-4-cyano-phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 54**)**




a) Synthesis of 1-(2-bromo-5-nitro-phenyl)sulfonylazetidine



From 1-(2-bromo-5-nitro-phenyl)sulfonyl chloride and azetidine, using General


5 Method G.

b) Synthesis of 3-(azetidin-1-ylsulfonyl)-4-bromo-aniline




To a solution of 1-(2-bromo-5-nitro-phenyl)sulfonylazetidine (5.2 g, 16.1 mmol, 1.0 *eq.*) in EtOH (30 mL), H₂O (10 mL) and THF (30 mL) was added Fe (4.5 g, 80.8 mmol, 5.0 *eq.*) and NH₄Cl (2.6 g, 48.5 mmol, 3.0 *eq.*). The mixture was stirred at 80°C for 12 h. The reaction mixture was filtered. The filtrate was concentrated and the residue was diluted with H₂O (30 mL) and extracted with EtOAc 90 mL (30 mL X 3). The combined organic layers were dried over Na₂SO₄, filtered and concentrated. The residue was purified by column chromatography (SiO₂, petroleum ether: ethyl acetate = 10:1 to 1:10) to yield 3-(azetidin-1-ylsulfonyl)-4-bromo-aniline (2.8 g, 9.6 mmol, 60% yield) as a white solid. ESI [M+H] = 291.0/293.0.

c) Synthesis of 3-(azetidin-1-ylsulfonyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline



A mixture of 4,4,5,5-tetramethyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1,3,2-dioxaborolane (11.3 g, 44.7 mmol, 5.0 *eq.*), 3-(azetidin-1-ylsulfonyl)-4-bromo-aniline (2.6 g, 8.9 mmol, 1.0 *eq.*), K₃PO₄ (9.5 g, 44.7 mmol, 5.0 *eq.*), Pd(PPh₃)₄ (1.0 g, 893.0 μmol, 0.1 *eq.*) in dioxane (120 mL) was stirred at 90°C for 3 h under N₂ atmosphere. The reaction mixture was concentrated and diluted with H₂O (30 mL) and extracted with EtOAc 90 mL (30 mL X 3). The combined organic layers were dried over Na₂SO₄, filtered and concentrated. The residue was purified by column chromatography (SiO₂, petroleum ether: ethyl acetate = 100:1 to 1:100) to yield crude 3-(azetidin-1-ylsulfonyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (3.8 g, crude) as a dark gum. ¹H NMR (400 MHz, methanol-*d*₄) δ = 7.26 (d, *J* = 7.9 Hz, 1H), 7.08 (d, *J* = 2.0 Hz, 1H), 6.91 (dd, *J* = 2.1, 8.0 Hz, 1H), 3.82 (t, *J* = 7.7 Hz, 4H), 2.05 (quin, *J* = 7.8 Hz, 2H), 1.33 (s, 12H). ESI [M+H] = 339.2.

d) Synthesis of isopropyl *trans*-N-[4-[5-[4-amino-2-(azetidin-1-ylsulfonyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate




15

From 3-(azetidin-1-ylsulfonyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline, using General Method C. ¹H NMR (400 MHz, methanol-*d*₄) δ = 7.59 (s, 1H), 7.36 (d, *J* = 2.3 Hz, 1H), 7.20 (d, *J* = 8.3 Hz, 1H), 6.89 (dd, *J* = 2.4, 8.3 Hz, 1H), 4.86 - 4.77 (m, 1H), 3.69 (t, *J* = 7.6 Hz, 4H), 3.53 - 3.42 (m, 1H), 3.07 - 2.93 (m, 1H), 2.28 - 2.17 (m, 2H), 2.16 - 2.11 (m, 2H), 2.08 (br d, *J* = 7.8 Hz, 2H), 1.79 - 1.63 (m, 2H), 1.50 - 1.35 (m, 2H), 1.24 (br d, *J* = 6.1 Hz, 6H). ESI [M+H] = 479.2.


20

e) Synthesis of isopropyl *trans*-N-[4-[5-[2-(azetidin-1-ylsulfonyl)-4-bromophenyl]thiazol-2-yl]cyclohexyl]carbamate (**General Method L**)




To a solution of isopropyl *trans*-*N*-[4-[5-[4-amino-2-(azetidin-1-ylsulfonyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (200 mg, 418 μ mol, 1 *eq.*) in MeCN (6 mL) was added CuBr₂ (37 mg, 167 μ mol, 0.4 *eq.*) and tert-butyl nitrite (64 mg, 627 μ mol, 1.5 *eq.*). The mixture was stirred at 60°C for 1 h and concentrated. The residue was diluted with H₂O (5 mL) and extracted with EtOAc 60 mL (20 mL X 3). The combined organic layers were dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (SiO₂, petroleum ether: ethyl acetate = 1:2) to yield isopropyl *trans*-*N*-[4-[5-[2-(azetidin-1-ylsulfonyl)-4-bromo-phenyl]thiazol-2-yl]cyclohexyl]carbamate (120 mg, 221 μ mol, 53% yield) as a white solid. ESI [M+H] = 542.0/544.0.

- 10 f) Synthesis of isopropyl *trans*-*N*-[4-[5-[2-(azetidin-1-ylsulfonyl)-4-cyano-phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 54**)




- To a solution of isopropyl *trans*-*N*-[4-[5-[2-(azetidin-1-ylsulfonyl)-4-bromo-phenyl]thiazol-2-yl]cyclohexyl]carbamate (60 mg, 110 μ mol, 1 *eq.*) in NMP (2 mL) was added Zn(CN)₂ (26 mg, 221 μ mol, 2 *eq.*) and Pd(PPh₃)₄ (13 mg, 11 μ mol, 0.1 *eq.*). The mixture was stirred at 140°C under microwave for 1 h and filtered. The filtrate was purified by prep-HPLC (column: Nano-Micro UniSil 5-100 C18 ULTRA 100*250mm 5 μ m; mobile phase: [water(0.1%TFA)-ACN]; B%: 45%-60%, 10min) to yield isopropyl *trans*-*N*-[4-[5-[2-(azetidin-1-ylsulfonyl)-4-cyano-phenyl]thiazol-2-yl]cyclohexyl] carbamate (15 mg, 31 μ mol, 28% yield, 100% purity) as a yellow solid. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.42 (s, 1H), 8.04 (br d, J=7.83 Hz, 1H), 7.85 (s, 1H), 7.75 (d, J=7.95 Hz, 1H), 4.77 - 4.86 (m, 1H), 3.75 (br t, J=7.64 Hz, 4H), 3.44 - 3.53 (m, 1H), 3.03 - 3.16 (m, 1H), 2.26 (br d, J=12.59 Hz, 2H), 2.14 - 2.21 (m, 2H), 2.10 (br d, J=11.98 Hz, 2H), 1.73 (br d, J=12.59 Hz, 2H), 1.38 - 1.50 (m, 2H), 1.25 (br d, J=5.87 Hz, 6H). ESI [M+H] = 489.1.

- 25 **Example 44. Preparation of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[2-(isopropylamino)-2-oxo-ethyl]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 58)**




a) Synthesis of 2-(4-bromo-3-chlorosulfonyl-phenyl)acetate




5 HSO₃Cl (200 mL) was added slowly to a flask containing methyl 2-(4-bromo-phenyl)acetate (20 g, 87 mmol, 1 *eq.*) was stirred at 0°C for 5 h. The reaction mixture was quenched with ice-water (1 L) and extracted with DCM 2 L (1 L X 2). The combined organic layers were washed with H₂O (500 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield 2-(4-bromo-3-chlorosulfonyl-phenyl)acetate (30 g, crude) as a yellow oil.

10 b) Synthesis of methyl 2-[4-bromo-3-(ethylsulfamoyl)phenyl]acetate



From 2-(4-bromo-3-chlorosulfonyl-phenyl)acetate, using General Method G. ESI[M-H] = 333.8/335.8.


c) Synthesis of 2-[4-bromo-3-(ethylsulfamoyl)phenyl]acetic acid



15


From methyl 2-[4-bromo-3-(ethylsulfamoyl)phenyl]acetate, using General Method J. ESI[M-H] = 319.9/321.9.

d) Synthesis of 2-[4-bromo-3-(ethylsulfamoyl)phenyl]-N-isopropyl-acetamide




From 2-[4-bromo-3-(ethylsulfamoyl)phenyl]acetic acid and isopropylamine, using General Method A. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.03 (d, *J*=2.21 Hz, 1H), 7.73 (d, *J*=8.16 Hz, 1H), 7.39 (dd, *J*=2.20, 7.94 Hz, 1H), 3.94 (spt, *J*=6.58 Hz, 1H), 3.51 (s, 2H), 2.88-2.98 (m, 2H), 1.13 (d, *J*=6.62 Hz, 6H), 1.04 (t, *J*=7.28 Hz, 3H). ESI [M+H] = 363.0/365.0.

e) Synthesis of 2-[3-(ethylsulfamoyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]-N-isopropyl-acetamide



From 2-[4-bromo-3-(ethylsulfamoyl)phenyl]-N-isopropyl-acetamide, using General Method B. ESI [M+H] = 411.1.

f) Synthesis of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[2-(isopropyl-amino)-2-oxo-ethyl]phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 58**)



From 2-[3-(ethylsulfamoyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]-N-isopropyl-acetamide, using General Method C. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.01 (d, *J*=1.54 Hz, 1H), 7.71 (s, 1H), 7.55 (dd, *J*=7.72, 1.76 Hz, 1H), 7.43 (d, *J*=7.94 Hz, 1H), 4.76 - 4.84 (m, 1H), 3.96 (dt, *J*=13.12, 6.67 Hz, 1H), 3.59 (s, 2H), 3.38 - 3.50 (m, 1H), 2.94 - 3.06 (m, 1H), 2.85 (q, *J*=7.20 Hz, 2H), 2.19 - 2.28 (m, 2H), 2.06 (br d, *J*=10.14 Hz, 2H), 1.63 - 1.75 (m, 2H), 1.34 - 1.46 (m, 2H), 1.22 (br d, *J*=6.17 Hz, 6H), 1.15 (d, *J*=6.62 Hz, 6H), 1.00 (t, *J*=7.17 Hz, 3H). ESI [M+H] = 551.2.

Example 45. Preparation of oxetan-3-yl (*trans*-4-(5-(2-(N-ethylsulfamoyl)-4-(2-(isopropylamino)-2-oxoethyl)phenyl)thiazol-2-yl)cyclohexyl]carbamate (Compound 59**)**


Following the same protocol and under the same reaction conditions as for Compound 58, Compound 59 was prepared. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.01 (s, 1H), 7.71 (s, 1H), 7.55 (br d, *J*=7.72 Hz, 1H), 7.43 (d, *J*=7.72 Hz, 1H), 5.29 - 5.41 (m, 1H), 4.87 (br s, 2H), 4.59 (br t, *J*=6.06 Hz, 2H), 3.96 (dt, *J*=13.23, 6.62 Hz, 1H), 3.59 (s, 2H), 3.39 - 3.51 (m, 1H),

2.96 - 3.07 (m, 1H), 2.85 (q, J=7.28 Hz, 2H), 2.24 (br d, J=12.35 Hz, 2H), 1.99 - 2.13 (m, 2H), 1.62 - 1.75 (m, 2H), 1.36 - 1.49 (m, 2H), 1.15 (d, J=6.62 Hz, 6H), 1.00 (t, J=7.28 Hz, 3H). ESI [M+H] = 564.9.

Example 46. Preparation of oxetan-3-yl 4-(5-(2-(N-ethylsulfamoyl)-4-(2-(isopropylamino)-2-oxoethyl)phenyl)thiazol-2-yl)piperazine-1-carboxylate (Compound 96)


Following the same protocol and under the same reaction conditions as for Compound 58, Compound 96 was prepared. ¹H NMR (400 MHz, methanol-*d*₄) δ = 7.99 (d, J=1.54 Hz, 1H), 7.52 (dd, J=7.83, 1.65 Hz, 1H), 7.42 - 7.46 (m, 1H), 7.28 (s, 1H), 5.36 - 5.43 (m, 1H), 4.88 (t, J=7.17 Hz, 2H), 4.65 (dd, J=7.83, 5.40 Hz, 2H), 3.95 (dt, J=13.18, 6.53 Hz, 1H), 3.59 - 3.75 (m, 4H), 3.51 - 3.58 (m, 6H), 2.86 (q, J=7.28 Hz, 2H), 1.15 (d, J=6.62 Hz, 6H), 1.01 (t, J=7.17 Hz, 3H). ESI [M+H] = 551.9.

Example 47. Preparation of *trans*-3-(ethylsulfamoyl)-4-[2-[4-(oxetan-3-yloxy-carbonylamino)cyclohexyl]thiazol-5-yl]benzoic acid (Compound 66)



15


a) Synthesis of 4-bromo-3-(ethylsulfamoyl)benzoic acid



20


From 4-bromo-3-(chlorosulfonyl)benzoic acid, using General Method G. ¹H NMR (400 MHz, DMSO-*d*₆) δ = 8.48 (s, 1H), 7.87 (d, J = 8.2 Hz, 1H), 7.72 (d, J = 8.2 Hz, 1H), 2.92 (quin, J = 6.8 Hz, 2H), 1.00 (t, J = 7.1 Hz, 3H). ESI [M-H] = 305.9/307.9.

b) Synthesis of 3-(ethylsulfamoyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoic acid




From 4-bromo-3-(ethylsulfamoyl)benzoic acid, using General Method B. ESI [M-H]
= 354.0.

- c) Synthesis of *trans*-3-(ethylsulfamoyl)-4-[2-[4-(oxetan-3-yloxycarbonyl)-
5 amino)cyclohexyl]thiazol-5-yl]benzoic acid (**Compound 66**)



- From 3-(ethylsulfamoyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoic acid
and *trans*-oxetan-3-yl-(4-(5-bromothiazol-2-yl)cyclohexyl)carbamate, using General Method
D. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.63 (d, J = 1.5 Hz, 1H), 8.18 (dd, J = 1.5, 7.9 Hz,
10 1H), 7.80 (s, 1H), 7.55 (d, J = 7.7 Hz, 1H), 5.35 (br t, J = 5.6 Hz, 1H), 4.86 - 4.81 (m, 2H),
4.62 - 4.56 (m, 2H), 3.51 - 3.38 (m, 1H), 3.10 - 2.98 (m, 1H), 2.91 (q, J = 7.3 Hz, 2H), 2.25
(br d, J = 12.6 Hz, 2H), 2.12 - 2.02 (m, 2H), 1.74 - 1.63 (m, 2H), 1.47 - 1.40 (m, 2H), 1.04 (t,
J = 7.3 Hz, 3H). ESI [M+H] = 510.1.


- Example 48. Preparation of *trans*-3-(ethylsulfamoyl)-4-[2-[4-(isopropoxyloxycarbonyl)-
15 amino)cyclohexyl]thiazol-5-yl]benzoic acid (Compound 48)**



- From 3-(ethylsulfamoyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoic acid
and *trans*-isopropyl-(4-(5-bromothiazol-2-yl)cyclohexyl)carbamate, using General Method
D. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.57 (s, 1H), 8.14 (br d, J = 7.9 Hz, 1H), 7.77 (br s,
20 1H), 7.54 (br d, J = 7.7 Hz, 1H), 4.74 (br s, 1H), 3.36 (br t, J = 11.5 Hz, 1H), 3.03 - 2.89 (m,
1H), 2.80 (q, J = 7.2 Hz, 2H), 2.16 (br d, J = 13.0 Hz, 2H), 1.98 (br d, J = 11.0 Hz, 2H), 1.69 -


1.54 (m, 2H), 1.38 - 1.28 (m, 2H), 1.22 - 1.08 (m, 6H), 0.94 (t, $J = 7.3$ Hz, 3H). ESI $[M+H] = 496.1$.

Example 49. Preparation of isopropyl (*trans*-*N*-(4-(5-(2-(ethylsulfamoyl)-4-((3-isopropyl-oxetan-3-yl)amino)phenyl)thiazol-2-yl)cyclohexyl))carbamate (Compound 69)




69

a) Synthesis of 2-methyl-*N*-(oxetan-3-ylidene)propane-2-sulfinamide




To a solution of oxetan-3-one (5.0 g, 69.4 mmol, 1.0 *eq.*) and 2-methylpropane-2-sulfinamide (10.1 g, 83.3 mmol, 1.2 *eq.*) in DCM (60 mL) was added tetraisopropoxy-titanium (39.4 g, 138.8 mmol, 2.0 *eq.*). The mixture was stirred at 40°C for 12 h. Then the reaction was quenched with sat. aq. NaHCO_3 (30 mL) and filtered. The filtrate was diluted with H_2O 30 mL and extracted with DCM 60 mL (20 mL X 3). The combined organic layers were dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified by column chromatography (SiO_2 , petroleum ether: ethyl acetate = 10:1 to 0:1) to yield 2-methyl-*N*-(oxetan-3-ylidene)propane-2-sulfinamide (6.0 g, 34.2 mmol, 49.3% yield) as yellow oil. ^1H NMR (400 MHz, CDCl_3) $\delta = 5.81 - 5.66$ (m, 1H), 5.64 - 5.53 (m, 1H), 5.46 - 5.30 (m, 2H), 1.20 (s, 9H).

b) Synthesis of *N*-(3-isopropyl-oxetan-3-yl)-2-methyl-propane-2-sulfinamide




To a solution of 2-methyl-*N*-(oxetan-3-ylidene)propane-2-sulfinamide (5.7 g, 32.4 mmol, 1.0 *eq.*) in THF (100 mL) was added isopropylolithium (1.6 M, 40.5 mL, 2.0 *eq.*) (1.6 M in hexane) under N₂ atmosphere at -78°C. The mixture was stirred at -78°C for 45 min. The reaction mixture was quenched with THF: AcOH = 4:1 (50 mL) at -78°C, then was diluted with H₂O 20 mL and extracted with EtOAc 150 mL (50 mL X 3). The combined organic layers were washed with sat. aq. NaCl 30 mL and dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by prep-HPLC (column: Welch Xtimate C18 250*50mm*10um; mobile phase: [water (0.05% ammonia hydroxide v/v)-ACN]; B%: 2%-40%, 20min) to yield *N*-(3-isopropoxyloxetan-3-yl)-2-methyl-propane-2-sulfinamide (300 mg, 1.4 mmol, 4.2% yield) as a white solid. ¹H NMR (400 MHz, CDCl₃) δ = 4.86 (d, J = 6.8 Hz, 1H), 4.77 (d, J = 6.8 Hz, 1H), 4.55 (dd, J = 6.8, 11.9 Hz, 2H), 3.40 (s, 1H), 2.32 (td, J = 6.8, 13.7 Hz, 1H), 1.26 (s, 9H), 1.09 - 0.91 (m, 6H). ESI [M+H] = 220.2.

c) Synthesis of 3-isopropoxyloxetan-3-amine (**General Method E**)



To a solution of *N*-(3-isopropoxyloxetan-3-yl)-2-methyl-propane-2-sulfinamide (100 mg, 456 μmol, 1 *eq.*) in DCM (1 mL) was added HCl/EtOAc (4 M, 500 μL, 4.4 *eq.*). The mixture was stirred at 0°C for 5 min, then the reaction mixture was diluted with MTBE 3 mL then the solid was formed. The residue was concentrated under reduced pressure to remove solvent at 0°C. Then the residue was triturated with MTBE: Petroleum ether = 1:1 (3 mL) and filtered, the filter cake was collected to yield 3-isopropoxyloxetan-3-amine (20 mg, crude) which was used into the next step without further purification. ¹H NMR (400 MHz, DMSO-*d*₆) δ = 8.61 (br s, 2H), 4.58 - 4.46 (m, 4H), 2.20 - 2.10 (m, 1H), 0.94 (d, J = 6.8 Hz, 6H). ESI [M+H] = 116.1.


d) Synthesis of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[(3-isopropoxyloxetan-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 69**; **General Method F**)




A mixture of 3-isopropoxyloxetan-3-amine (13 mg, 113 μmol, 2 *eq.*), isopropyl *N*-[4-[5-[4-bromo-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (30 mg, 56 μmol, 1

eq.), K₂CO₃ (23 mg, 169 μ mol, 3 eq.) and BrettPhos Pd G3 (5 mg, 5 μ mol, 0.1 eq.) in *t*-BuOH (2 mL) was stirred at 110°C for 12 h under Ar atmosphere. The reaction mixture was concentrated under reduced pressure, then the residue was purified by prep-TLC(SiO₂, petroleum ether: ethyl acetate = 2:1) and purified by prep-HPLC (column: YMC-Actus Triart C18 100*30mm*5 μ m; mobile phase: [water(0.1%TFA)-ACN]; B%: 25%-60%, 10min) to yield isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[(3-isopropoxyloxetan-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl] carbamate (1 mg, 1.9 μ mol, 3.3% yield, 89.8% purity) as a yellow gum. ¹H NMR (400 MHz, DMSO-*d*₆) δ = 7.53 (s, 1H), 7.34 - 7.22 (m, 1H), 7.11 (d, *J* = 8.4 Hz, 1H), 7.05 - 6.90 (m, 3H), 6.48 (dd, *J* = 1.8, 7.9 Hz, 1H), 4.77 - 4.69 (m, 1H), 4.64 (d, *J* = 6.0 Hz, 2H), 4.49 (d, *J* = 6.6 Hz, 2H), 3.26 - 3.23 (m, 1H), 2.86 - 2.79 (m, 1H), 2.79 - 2.70 (m, 2H), 2.28 - 2.20 (m, 1H), 2.13 - 2.06 (m, 2H), 1.93 - 1.86 (m, 2H), 1.57 - 1.45 (m, 2H), 1.37 - 1.28 (m, 2H), 1.13 (d, *J* = 6.2 Hz, 6H), 1.01 - 0.90 (m, 9H). ESI [M+H] = 565.1.

Example 50. Preparation of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[(3-isobutyloxetan-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 70)




a) Synthesis of *N*-(3-isobutyloxetan-3-yl)-2-methyl-propane-2-sulfonamide



To a solution of 2-methyl-*N*-(oxetan-3-ylidene)propane-2-sulfonamide (7.0 g, 40.0 mmol, 1.0 eq.) in THF (4 mL) was added isobutyryllithium (1.6 M, 50.0 mL, 2.0 eq.) (1M in THF) dropwise at -78°C under N₂ atmosphere. The mixture was stirred at -78°C for 45 min. Then the reaction mixture was quenched by additional THF: AcOH = 4:1 (50 mL) at -78°C and the reaction was diluted with H₂O 20 mL and extracted with EtOAc 150 mL (50 mL X 3). The combined organic layers were washed with sat.aq. NaCl (3 mL), and dried over Na₂SO₄, filtered and concentrated under reduced pressure. Then the residue was purified by


prep-HPLC (column: Welch Xtimate C18 250*50mm*10um; mobile phase: [water (0.05% ammonia hydroxide v/v)-ACN]; B%: 5%-48%, 20min) to yield *N*-(3-isobutyloxyetan-3-yl)-2-methyl-propane-2-sulfonamide (130 mg, 557 μ mol, 1.4% yield) was obtained as a white solid. ^1H NMR (400 MHz, CDCl_3) δ = 4.69 (t, J = 7.0 Hz, 2H), 4.57 (d, J = 6.9 Hz, 1H), 4.51 (d, J = 6.6 Hz, 1H), 3.56 (s, 1H), 1.95 (t, J = 6.9 Hz, 1H), 1.86 - 1.73 (m, 2H), 1.19 (s, 9H), 0.84 (d, J = 6.4 Hz, 6H). ESI $[\text{M}+\text{H}] = 234.1$.

b) Synthesis of 3-isobutyloxyetan-3-amine




From *N*-(3-isobutyloxyetan-3-yl)-2-methyl-propane-2-sulfonamide, using the conditions of General Method E. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ = 4.59 - 4.55 (m, 2H), 4.54 - 4.50 (m, 2H), 1.84 - 1.74 (m, 3H), 0.92 (t, J = 6.3 Hz, 2H), 0.90 - 0.86 (m, 6H).

c) Synthesis of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[(3-isobutyloxyetan-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 70**)




From isopropyl *N*-[4-[5-[4-bromo-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate and 3-isobutyloxyetan-3-amine, using General Method F. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ = 7.69 - 7.54 (m, 1H), 7.37 (t, J = 5.6 Hz, 1H), 7.22 (d, J = 8.4 Hz, 1H), 7.17 - 7.11 (m, 1H), 7.08 (br d, J = 7.3 Hz, 1H), 7.03 (d, J = 1.8 Hz, 1H), 6.58 (dd, J = 2.1, 8.7 Hz, 1H), 4.87 - 4.73 (m, 1H), 4.63 (s, 4H), 3.41 - 3.35 (m, 1H), 3.01 - 2.90 (m, 1H), 2.88 - 2.79 (m, 2H), 2.24 - 2.12 (m, 2H), 2.06 (br d, J = 7.1 Hz, 2H), 2.01 - 1.90 (m, 2H), 1.67 - 1.52 (m, 3H), 1.46 - 1.32 (m, 2H), 1.23 (d, J = 6.2 Hz, 6H), 1.03 (t, J = 7.2 Hz, 3H), 0.89 (d, J = 6.6 Hz, 6H). ESI $[\text{M}+\text{H}] = 579.3$.

Example 51. Preparation of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[(1*H*-imidazol-2-ylamino)methyl]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 73**)**




a) Synthesis of 4-bromo-3-(chlorosulfonyl)benzoic acid



- 5 A solution of 4-bromobenzoic acid (20.0 g, 99.5 mmol, 1.0 *eq.*) in HSO₃Cl (120 mL) was stirred at 100°C for 12 h, then the reaction mixture was poured into ice water (600 mL) slowly. The mixture was stirred for 0.5 h, then filtrated to yield 4-bromo-3-chlorosulfonylbenzoic acid (23.4 g, crude) as a pale-yellow solid used directly without any purification.


b) Synthesis of methyl 4-bromo-3-chlorosulfonylbenzoate



- 10 A solution of 4-bromo-3-chlorosulfonylbenzoic acid (23.4 g, 78.0 mmol, 1.0 *eq.*) in SOCl₂ (92.7 g, 780.0 mmol, 10.0 *eq.*) was stirred at 80°C for 3 h. Then the mixture was concentrated under reduced pressure. The residue was diluted in Tol (50 mL) and MeOH (3.3 g, 101.4 mmol, 1.3 *eq.*). The mixture was stirred at 110°C for 2 h. Then concentrated under reduced pressure. The residue was purified by column chromatography (SiO₂, petroleum
15 ether : ethyl acetate = 1:0 to 0:1) to yield methyl 4-bromo-3-chlorosulfonylbenzoate (10.5 g,


33.2 mmol, 42.5% yield) as a white solid. ^1H NMR (400 MHz, methanol- d_4) δ = 8.75 (d, J = 2.0 Hz, 1H), 8.27 (dd, J = 2.1, 8.3 Hz, 1H), 8.16 (d, J = 8.3 Hz, 1H), 4.02 - 3.98 (m, 3H).

c) Synthesis of methyl 4-bromo-3-(ethylsulfonyl)benzoate (**General Method G**)




To a solution of TEA (7.8 g, 76.5 mmol, 3 *eq.*) in DCM (140 mL) was added ethylamine (1.3 g, 28.1 mmol, 1.1 *eq.*) at -20°C . Then the methyl 4-bromo-3-chlorosulfonyl-benzoate (8.0 g, 25.5 mmol, 1.0 *eq.*) was added batchwise. The mixture was stirred at -20°C for 5 min, then the reaction mixture was quenched with HCl(1M, 50 mL, pH=2) and extracted with DCM 150 mL, the organic layers were dried over Na_2SO_4 , filtered and concentrated under reduced pressure to yield methyl 4-bromo-3-(ethylsulfonyl)benzoate (8.0 g, 24.8 mmol, 97.3% yield) as a pale-yellow solid. ^1H NMR (400 MHz, methanol- d_4) δ = 8.63 (d, J = 1.8 Hz, 1H), 8.10 - 8.00 (m, 1H), 7.93 (d, J = 8.2 Hz, 1H), 3.94 (s, 3H), 2.96 (q, J = 7.3 Hz, 2H), 1.05 (t, J = 7.2 Hz, 3H).

d) Synthesis of methyl 3-(ethylsulfonyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate



From methyl 4-bromo-3-(ethylsulfonyl)benzoate, using General Method B. ^1H NMR (400 MHz, methanol- d_4) δ = 8.51 - 8.39 (m, 1H), 8.20 (dd, J = 1.5, 7.7 Hz, 1H), 7.74 (d, J = 7.7 Hz, 1H), 3.95 (s, 3H), 2.96 - 2.88 (m, 2H), 1.40 (s, 12H), 1.05 - 1.00 (m, 3H). ESI $[\text{M}+\text{H}] = 370.1$.

e) Synthesis of methyl *trans*-3-(ethylsulfonyl)-4-[2-[4-(isopropoxycarbonyl)amino)cyclohexyl]thiazol-5-yl]benzoate (**Compound 71**)




From methyl 3-(ethylsulfonyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate, using General Method D. ^1H NMR (400 MHz, methanol- d_4) δ = 8.68 (d, J = 1.6

Hz, 1H), 8.24 (dd, $J = 1.8, 8.0$ Hz, 1H), 7.85 (s, 1H), 7.66 (d, $J = 8.0$ Hz, 1H), 4.84 (br s, 1H), 3.99 (s, 3H), 3.48 (tt, $J = 3.8, 11.6$ Hz, 1H), 3.05 (ddd, $J = 3.4, 8.7, 11.9$ Hz, 1H), 2.91 (q, $J = 7.2$ Hz, 2H), 2.27 (br d, $J = 12.1$ Hz, 2H), 2.14 - 2.05 (m, 2H), 1.73 (dq, $J = 3.0, 12.8$ Hz, 2H), 1.44 (dq, $J = 3.2, 12.6$ Hz, 2H), 1.25 (br d, $J = 6.1$ Hz, 6H), 1.05 (t, $J = 7.3$ Hz, 3H). ESI


5 [M+H] = 510.2.

f) Synthesis of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-(hydroxymethyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 72**)



To a solution of *trans*-methyl 3-(ethylsulfamoyl)-4-[2-[4-(isopropoxycarbonylamino)-cyclohexyl]thiazol-5-yl]benzoate (500 mg, 981 μ mol, 1 *eq.*) in THF (6 mL) was added LiAlH₄ (74 mg, 2 mmol, 2 *eq.*), then stirred at 0°C for 2 h. The reaction was quenched with H₂O 10 mL and extracted with EtOAc 30 mL (10 mL X 3). The combined organic layers were dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by column chromatography (SiO₂, Petroleum ether: Ethyl acetate = 1:0 to 0:1) to yield isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-(hydroxymethyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (210 mg, 436 μ mol, 44.4% yield, 98.6% purity) as a white solid. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.11 (s, 1H), 7.81 (s, 1H), 7.65 (d, $J = 7.1$ Hz, 1H), 7.51 (d, $J = 7.8$ Hz, 1H), 4.85 - 4.80 (m, 1H), 4.75 (s, 2H), 3.48 (ddd, $J = 3.6, 7.7, 11.6$ Hz, 1H), 3.14 - 3.01 (m, 1H), 2.90 (q, $J = 7.3$ Hz, 2H), 2.28 (br d, $J = 12.0$ Hz, 2H), 2.15 - 2.07 (m, 2H), 1.80 - 1.66 (m, 2H), 1.52 - 1.38 (m, 2H), 1.25 (br d, $J = 6.0$ Hz, 6H), 1.04 (t, $J = 7.2$ Hz, 3H). ESI [M+H] = 482.1.


g) Synthesis of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-formyl-phenyl]thiazol-2-yl]cyclohexyl]carbamate



To a solution of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-(hydroxymethyl)-phenyl]thiazol-2-yl]cyclohexyl]carbamate (100 mg, 207 μ mol, 1 *eq.*) in CHCl₃ (1


mL)/MeOH (1 mL) was added MnO₂ (361 mg, 4 mmol, 20 *eq.*). The mixture was stirred at 60°C for 2 h. The reaction mixture was filtered, and the filtrate was concentrated under reduced pressure. The residue was purified by prep-TLC (SiO₂, petroleum ether: ethyl acetate = 1:2) to yield isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-formyl-phenyl]thiazol-2-yl]cyclohexyl]carbamate (30 mg, 62.55 μmol, 30.13% yield) as a white solid. ESI [M+H] = 480.2.

h) Synthesis of isopropyl-*trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[[[1-(2-trimethylsilylethoxymethyl)imidazol-2-yl]amino]methyl]phenyl]thiazol-2-yl]cyclohexyl]carbamate (**General Method H**)



To a mixture of 1-(2-trimethylsilylethoxymethyl)imidazol-2-amine (20 mg, 94 μmol, 1.5 *eq.*) and isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-formyl-phenyl]thiazol-2-yl]cyclohexyl]carbamate (30 mg, 62 μmol, 1 *eq.*) in DCM (1 mL) was added Ti(i-PrO)₄ (35 mg, 125 μmol, 37 μL, 2 *eq.*) and Et₃N (633 μg, 6 μmol, 0.1 *eq.*), then the mixture was stirred at 25°C for 16 h under N₂ atmosphere. The mixture was added NaBH₄ (3.6 mg, 93.9 μmol, 1.5 *eq.*) at 0°C, and stirred at 25°C for 2 h. The reaction mixture was quenched with H₂O 10 mL at 25°C and concentrated under reduced pressure. The residue was purified by prep-TLC (SiO₂, petroleum ether: ethyl acetate=0:1) to yield isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[[[1-(2-trimethylsilylethoxy-methyl)imidazol-2-yl]amino]methyl]phenyl]thiazol-2-yl]cyclohexyl]carbamate (50 mg, crude) as a yellow oil. ESI [M+H] = 677.3.


20 i) Synthesis of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[(1*H*-imidazol-2-ylamino)methyl]phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 73; General Method I**)



A mixture of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[[[1-(2-trimethylsilylethoxymethyl)imidazol-2-yl]amino]methyl]phenyl]thiazol-2-yl]cyclohexyl]carbamate (50


mg, 74 μ mol, 1 *eq.*) in TFA (1 mL) and DCM (1 mL) was stirred at 20°C for 1 h. The reaction mixture was concentrated under reduced pressure to remove DCM at 25°C. The residue was diluted with EtOH (1 mL) and stirred at 60°C for 3 h. The mixture was concentrated under reduced pressure. The residue was purified by prep-HPLC (TFA condition). (column: Welch Ultimate AQ-C18 150*30mm*5 μ m; mobile phase: [water(0.1%TFA)-ACN]; B%: 18%-48%, 12min) to yield isopropyl *trans-N*-[4-[5-[2-(ethylsulfamoyl)-4-[(1*H*-imidazol-2-ylamino)methyl]phenyl]thiazol-2-yl]cyclohexyl]carbamate (9 mg, 13 μ mol, 18% yield, 100% purity, TFA) as a pale yellow solid. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.08 (d, *J* = 1.1 Hz, 1H), 7.74 (s, 1H), 7.64 (dd, *J* = 1.4, 7.8 Hz, 1H), 7.52 (d, *J* = 7.9 Hz, 1H), 6.88 (s, 2H), 4.85 - 4.74 (m, 1H), 4.65 (s, 2H), 3.51 - 3.37 (m, 1H), 3.08 - 2.96 (m, 1H), 2.81 (q, *J* = 7.3 Hz, 2H), 2.33 - 2.18 (m, 2H), 2.07 (br d, *J* = 10.4 Hz, 2H), 1.69 (dq, *J* = 3.0, 12.8 Hz, 2H), 1.41 (dq, *J* = 3.1, 12.6 Hz, 2H), 1.22 (br d, *J* = 6.2 Hz, 6H), 0.97 (t, *J* = 7.3 Hz, 3H). ESI [M+H] = 547.1.

Example 52. Preparation of isopropyl *trans-N*-[4-[5-[2-(ethylsulfamoyl)-4-[(oxazol-2-ylamino) methyl]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 74)



From isopropyl *trans-N*-[4-[5-[2-(ethylsulfamoyl)-4-formyl-phenyl]thiazol-2-yl]cyclohexyl]carbamate and 2-amino oxazole, using General Method H. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.04 (s, 1H), 7.72 (s, 1H), 7.64 - 7.55 (m, 1H), 7.46 (d, *J* = 7.7 Hz, 1H), 7.32 (d, *J* = 1.1 Hz, 1H), 6.76 (d, *J* = 0.9 Hz, 1H), 4.60 (br s, 1H), 4.56 (s, 2H), 3.53 - 3.39 (m, 1H), 3.06 - 2.94 (m, 1H), 2.84 (q, *J* = 7.3 Hz, 2H), 2.29 - 2.18 (m, 2H), 2.07 (br d, *J* = 9.9 Hz, 2H), 1.77 - 1.60 (m, 2H), 1.50 - 1.36 (m, 2H), 1.23 (br d, *J* = 6.2 Hz, 6H), 1.00 (t, *J* = 7.2 Hz, 3H). ESI [M+H] = 548.3.

Example 53. Preparation of isopropyl *trans-N*-[4-[5-[2-(ethylsulfamoyl)-4-(1*H*-imidazol-2-yl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 75)



To a solution of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-formyl-phenyl]thiazol-2-yl]cyclohexyl]carbamate (30 mg, 62 μ mol, 1 eq.) in EtOH (1 mL) was added oxaldehyde (27 mg, 187 μ mol, 24 μ L, 3 eq.) (40% aq.) and $\text{NH}_3 \cdot \text{H}_2\text{O}$ (146 mg, 1.3 mmol, 30% purity, 20 eq.) at 0°C. The mixture was stirred at 25°C for 2 h, and concentrated under reduced pressure.

5 The residue was purified by prep-HPLC (column: Welch Ultimate AQ-C18


150*30mm*5 μ m; mobile phase: [water(0.1%TFA)-ACN]; B%: 30%-60%, 12min) to yield

isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-(1*H*-imidazol-2-yl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (16 mg, 25 μ mol, 40% yield, 99.3% purity, TFA) as a yellow solid.


¹H NMR (400 MHz, methanol-*d*₄) δ = 8.70 (d, *J* = 2.0 Hz, 1H), 8.15 (dd, *J* = 2.0, 8.2 Hz, 1H),
 10 7.86 (s, 1H), 7.78 (d, *J* = 8.2 Hz, 1H), 7.72 (s, 2H), 4.83 - 4.78 (m, 1H), 3.45 (tdd, *J* = 4.0,
 7.6, 11.7 Hz, 1H), 3.04 (tt, *J* = 3.5, 12.0 Hz, 1H), 2.87 (q, *J* = 7.3 Hz, 2H), 2.26 (br d, *J* = 12.1
 Hz, 2H), 2.12 - 2.03 (m, 2H), 1.77 - 1.65 (m, 2H), 1.48 - 1.36 (m, 2H), 1.22 (br d, *J* = 6.2 Hz,
 6H), 1.05 - 0.92 (m, 3H). ESI [M+H] = 518.0.

Example 54. Preparation of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-(1*H*-imidazol-

15 **2-ylmethyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 78)**




a) Synthesis of methyl 2-[3-(ethylsulfamoyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]acetate




From methyl 2-(4-bromo-3-(*N*-ethylsulfamoyl)phenyl)acetate, using General Method B. ^1H NMR (400MHz, $\text{DMSO}-d_6$) δ = 7.97 - 7.83 (m, 1H), 7.70 (s, 1H), 7.51 - 7.44 (m, 2H), 3.81 (s, 2H), 3.64 - 3.58 (m, 3H), 2.88 - 2.75 (m, 2H), 1.29 (s, 12H), 0.93 (t, $J=7.2$ Hz, 3H). ESI $[\text{M}+\text{H}] = 384.1$.

- 5 b) Synthesis of methyl *trans*-2-[3-(ethylsulfamoyl)-4-[2-[4-(isopropoxycarbonylamino)cyclohexyl]thiazol-5-yl]phenyl]acetate (**Compound 76**)



- From methyl 2-[3-(ethylsulfamoyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]acetate and isopropyl (*trans*-4-(5-bromothiazol-2-yl)cyclohexyl)carbamate, using General Method D. ^1H NMR (400MHz, methanol- d_4) δ = 8.02 (d, $J=1.4$ Hz, 1H), 7.75 (s, 1H), 7.58 (dd, $J=1.6, 7.9$ Hz, 1H), 7.47 (d, $J=7.9$ Hz, 1H), 4.82 (br s, 1H), 3.84 (s, 2H), 3.74 (s, 3H), 3.47 (tt, $J=3.8, 11.5$ Hz, 1H), 3.35 - 3.35 (m, 1H), 2.89 (q, $J=7.3$ Hz, 2H), 2.31 - 2.21 (m, 2H), 2.14 - 2.03 (m, 2H), 1.72 (dq, $J=2.8, 12.9$ Hz, 2H), 1.43 (dq, $J=3.1, 12.5$ Hz, 2H), 1.24 (br d, $J=6.1$ Hz, 6H), 1.04 (t, $J=7.3$ Hz, 3H). ESI $[\text{M}+\text{H}] = 524.2$.


- 15 c) Synthesis of *trans*-2-[3-(ethylsulfamoyl)-4-[2-[4-(isopropoxycarbonylamino)cyclohexyl]thiazol-5-yl]phenyl]acetic acid (**Compound 77; General Method J**)



- To a solution of methyl *trans*-2-[3-(ethylsulfamoyl)-4-[2-[4-(isopropoxycarbonylamino) cyclohexyl]thiazol-5-yl]phenyl]acetate (800 mg, 1.5 mmol, 1 *eq.*) in MeOH (8 mL), H_2O (4 mL) was added LiOH (110 mg, 4.5 mmol, 3 *eq.*). The mixture was stirred at 50°C for 2 h. The mixture was concentrated under reduced pressure. The residue was adjusted pH = 2 with HCl(6M) and extracted with EtOAc 40 mL (20 mL X 2). The combined organic layers were dried over Na_2SO_4 , filtered and concentrated under reduced pressure to yield *trans*-2-[3-(ethylsulfamoyl)-4-[2-[4-(isopropoxycarbonyl-amino) cyclohexyl]thiazol-5-yl]phenyl]acetic acid (750 mg, crude) as a yellow oil. ^1H NMR (400MHz, methanol- d_4) δ = 8.00 (s, 1H), 7.74 (s, 1H), 7.56 (br d, $J=7.9$ Hz, 1H), 7.45 (d,


J=7.7 Hz, 1H), 4.89 - 4.88 (m, 1H), 3.76 (s, 2H), 3.45 (br t, J=11.0 Hz, 1H), 3.01 (br t, J=12.2 Hz, 1H), 2.87 (q, J=7.2 Hz, 2H), 2.24 (br d, J=12.1 Hz, 2H), 2.07 (br d, J=10.4 Hz, 2H), 1.77 - 1.62 (m, 2H), 1.48 - 1.33 (m, 2H), 1.22 (br d, J=6.0 Hz, 6H), 1.01 (t, J=7.3 Hz, 3H). ESI [M+H] = 510.2.

- 5 d) Synthesis of isopropyl *trans*-N-[4-[5-[4-[2-(2,2-dimethoxyethylamino)-2-oxo-ethyl]-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate




From *trans*-2-[3-(ethylsulfamoyl)-4-[2-[4-(isopropoxycarbonylamino)cyclohexyl]thiazol-5-yl]phenyl]acetic acid and 2,2-dimethoxyethanamine, using General Method A. ESI [M+H] = 597.3.

- 10 e) Synthesis of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-(1*H*-imidazol-2-ylmethyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 78**)



A solution of isopropyl *trans*-N-[4-[5-[4-[2-(2,2-dimethoxyethylamino)-2-oxo-ethyl]-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (60 mg, 100 μ mol, 1 *eq.*) and NH₄OAc (16 mg, 201 μ mol, 2 *eq.*) in AcOH (2 mL) was heated at 100°C for 30 min under microwave. The mixture was concentrated, then the residue was purification by prep-TLC (petroleum ether: ethyl acetate = 0:1) to yield isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-(1*H*-imidazol-2-ylmethyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (2 mg, 3.4 μ mol, 3.4% yield, 91.2% purity) as pale yellow solid. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.02 (s, 1H), 7.73 (s, 1H), 7.47-7.61 (m, 4H), 4.67-4.77 (m, 1H), 4.51 (s, 2H), 3.44 (br s, 1H), 3.01 (s, 1H), 2.75-2.88 (m, 2H), 2.24 (br d, J=14.99 Hz, 2H), 2.02 (br d, J=18.52 Hz, 2H), 1.60-1.82 (m, 2H), 1.33-1.51 (m, 2H), 1.22 (br d, J=6.17 Hz, 6H), 0.92-1.05 (m, 3H). ESI [M+H] = 532.2.

- 25 **Example 55. Preparation of isopropyl *trans*-N-[4-[5-[4-[2-(benzylamino)-2-oxo-ethyl]-2-(ethyl-sulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 79)**



From *trans*-2-[3-(ethylsulfamoyl)-4-[2-[4-(isopropoxycarbonylamino)cyclohexyl]thiazol-5-yl]phenyl]acetic acid and benzylamine, using General Method A. ¹H NMR (400MHz, methanol-*d*₄) δ = 8.04 (s, 1H), 7.79 (s, 1H), 7.58 (br d, *J*=7.5 Hz, 1H), 7.45 (d, *J*=7.7 Hz, 1H), 7.34 - 7.18 (m, 5H), 4.87 - 4.77 (m, 1H), 4.38 (s, 2H), 3.69 (s, 2H), 3.45 (br t, *J*=11.8 Hz, 1H), 3.11 - 2.97 (m, 1H), 2.84 (q, *J*=7.3 Hz, 2H), 2.24 (br d, *J*=12.8 Hz, 2H), 2.07 (br d, *J*=10.4 Hz, 2H), 1.78 - 1.62 (m, 2H), 1.50 - 1.32 (m, 2H), 1.22 (br d, *J*=6.2 Hz, 6H), 1.00 (t, *J*=7.3 Hz, 3H). ESI [M+H] = 599.2

Example 56. Preparation of isopropyl *trans*-N-[4-[5-[4-[2-(isopropyl(methyl)amino)-2-oxo-ethyl]-2-(ethyl-sulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 80)

Following the same General Method A, Compound 80 was prepared. ¹H NMR (400MHz, methanol-*d*₄) δ = 7.96 (br d, *J*=3.1 Hz, 1H), 7.80 (s, 1H), 7.57 - 7.50 (m, 1H), 7.49 - 7.44 (m, 1H), 4.84 - 4.75 (m, 1H), 4.27 (td, *J*=6.7, 13.1 Hz, 1H), 3.95 (s, 1H), 3.89 (s, 1H), 3.45 (br t, *J*=11.7 Hz, 1H), 3.05 (br t, *J*=12.0 Hz, 1H), 2.93 (s, 2H), 2.86 (q, *J*=7.3 Hz, 2H), 2.81 (s, 1H), 2.25 (br d, *J*=12.6 Hz, 2H), 2.07 (br d, *J*=11.2 Hz, 2H), 1.78 - 1.63 (m, 2H), 1.48 - 1.34 (m, 2H), 1.22 (br d, *J*=6.2 Hz, 6H), 1.17 (d, *J*=6.6 Hz, 3H), 1.13 (d, *J*=6.8 Hz, 3H), 1.01 (t, *J*=7.2 Hz, 3H). ESI [M+H] = 565.3.

Example 57. Preparation of isopropyl *trans*-N-[4-[5-[4-[2-(benzyl(methyl)amino)-2-oxo-ethyl]-2-(ethyl-sulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 81)

Following the same General Method A, Compound 81 was prepared. ¹H NMR (400MHz, methanol-*d*₄) δ = 8.01 (s, 0.5H), 7.93 (s, 0.5H), 7.82 - 7.76 (m, 1H), 7.59 - 7.53 (m, 1H), 7.47 (br d, *J*=7.3 Hz, 1H), 7.37 - 7.29 (m, 2H), 7.28 - 7.22 (m, 2H), 7.15 (br d, *J*=7.3 Hz, 1H), 4.83 - 4.77 (m, 1H), 4.72 (s, 1H), 4.62 (s, 1H), 4.03 - 3.93 (m, 2H), 3.45 (br t, *J*=11.6 Hz, 1H), 3.10 - 3.00 (m, 3H), 2.96 (s, 1H), 2.90 - 2.77 (m, 2H), 2.24 (br d, *J*=12.3 Hz, 2H), 2.07 (br d, *J*=11.7 Hz, 2H), 1.71 (q, *J*=12.3 Hz, 2H), 1.41 (q, *J*=11.8 Hz, 2H), 1.22 (br d, *J*=5.7 Hz, 6H), 1.00 (t, *J*=7.1 Hz, 3H). ESI [M+H] = 613.2.

Example 58. Preparation of isopropyl *trans*-N-[4-[5-[4-[2-oxo-2-(pyrrolidin-1-yl)ethyl]-2-(ethyl-sulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 82)

Following the same General Method A, Compound 82 was prepared. ¹H NMR (400MHz, methanol-*d*₄) δ = 7.97 (s, 1H), 7.79 (s, 1H), 7.56 - 7.52 (m, 1H), 7.48 - 7.44 (m,

1H), 4.83 (br s, 1H), 3.85 (s, 2H), 3.58 (t, J=6.7 Hz, 2H), 3.45 (br t, J=6.8 Hz, 3H), 3.05 (br t, J=12.1 Hz, 1H), 2.87 (q, J=7.3 Hz, 2H), 2.24 (br d, J=12.6 Hz, 2H), 2.12 - 2.03 (m, 2H), 2.03 - 1.96 (m, 2H), 1.94 - 1.86 (m, 2H), 1.80 - 1.63 (m, 2H), 1.49 - 1.33 (m, 2H), 1.22 (br d, J=6.0 Hz, 6H), 1.02 (t, J=7.3 Hz, 3H). ESI [M+H] = 563.2.

5 **Example 59. Preparation of isopropyl *trans*-N-[4-[5-[4-[2-oxo-2-(piperidin-1-yl)ethyl]-2-(ethyl-sulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 83)**

Following the same General Method A, Compound 83 was prepared. ¹H NMR (400MHz, methanol-*d*₄) δ = 7.96 (s, 1 H), 7.73 (s, 1 H), 7.50 - 7.56 (m, 1 H), 7.41 - 7.47 (m, 1 H), 4.76 - 4.84 (m, 1 H), 3.91 (s, 2 H), 3.56 (dt, J=14.17, 5.49 Hz, 4 H), 3.40 - 3.49 (m, 1 H), 2.97 - 3.06 (m, 1 H), 2.85 (q, J=7.28 Hz, 2 H), 2.24 (br d, J=12.57 Hz, 2 H), 2.02 - 2.11 (m, 2 H), 1.61 - 1.75 (m, 4 H), 1.46 - 1.59 (m, 4 H), 1.35 - 1.45 (m, 2 H), 1.22 (br d, J=5.95 Hz, 6 H), 1.01 (t, J=7.28 Hz, 3 H). ESI [M+H] = 577.3.

Example 60. Preparation of isopropyl *trans*-N-[4-[5-[4-[2-oxo-2-(morpholin-4-yl)ethyl]-2-(ethyl-sulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 84)

15 Following the same General Method A, Compound 84 was prepared. ¹H NMR (400MHz, methanol-*d*₄) δ = 7.96 (s, 1 H), 7.79 (s, 1 H), 7.51 - 7.57 (m, 1 H), 7.43 - 7.49 (m, 1 H), 4.80 - 4.86 (m, 1 H), 3.93 (s, 2 H), 3.57 - 3.70 (m, 8 H), 3.40 - 3.51 (m, 1 H), 2.98 - 3.09 (m, 1 H), 2.86 (q, J=7.13 Hz, 2 H), 2.21 - 2.29 (m, 2 H), 2.07 (br d, J=11.69 Hz, 2 H), 1.65 - 1.77 (m, 2 H), 1.35 - 1.48 (m, 2 H), 1.17 - 1.26 (m, 6 H), 1.01 (t, J=7.17 Hz, 3 H). ESI [M+H] = 579.3.

Example 61. Preparation of isopropyl *trans*-N-[4-[5-[4-[2-oxo-2-(3-hydroxyazetidin-1-yl)ethyl]-2-(ethyl-sulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 85)

25 Following the same General Method A, Compound 85 was prepared. ¹H NMR (400MHz, methanol-*d*₄) δ = 8.00 (d, J=1.5 Hz, 1H), 7.90 (s, 1H), 7.62 - 7.56 (m, 1H), 7.54 - 7.48 (m, 1H), 4.79 - 4.75 (m, 1H), 4.62 (tt, J=4.3, 6.7 Hz, 1H), 4.57 - 4.48 (m, 1H), 4.25 (dd, J=7.9, 10.6 Hz, 1H), 4.10 - 4.04 (m, 1H), 3.80 (dd, J=4.2, 10.6 Hz, 1H), 3.73 - 3.64 (m, 2H), 3.49 (tt, J=3.9, 11.6 Hz, 1H), 3.19 - 3.09 (m, 1H), 2.92 (q, J=7.2 Hz, 2H), 2.29 (br d, J=12.5 Hz, 2H), 2.11 (br d, J=10.0 Hz, 2H), 1.75 (dq, J=3.1, 12.8 Hz, 2H), 1.52 - 1.38 (m, 2H), 1.25 (br d, J=6.1 Hz, 6H), 1.06 (t, J=7.3 Hz, 3H). ESI [M+H] = 565.2.

30 **Example 62. Preparation of isopropyl *trans*-N-[4-[5-[4-[2-amino-2-oxo-ethyl]-2-(ethyl-sulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 86)**


Following the same General Method A, Compound 86 was prepared. ¹H NMR (400MHz, methanol-*d*₄) δ = 8.05 (d, J=1.6 Hz, 1H), 7.80 (s, 1H), 7.64 - 7.58 (m, 1H), 7.48 (d,

J=7.9 Hz, 1H), 4.88 - 4.87 (m, 1H), 3.68 (s, 2H), 3.52 - 3.42 (m, 1H), 3.17 - 3.00 (m, 1H), 2.90 (q, J=7.3 Hz, 2H), 2.27 (br d, J=12.1 Hz, 2H), 2.10 (br d, J=10.3 Hz, 2H), 1.80 - 1.66 (m, 2H), 1.50 - 1.38 (m, 2H), 1.25 (br d, J=6.1 Hz, 6H), 1.04 (t, J=7.2 Hz, 3H). ESI [M+H] = 509.2.


5 **Example 63. Preparation of isopropyl *trans*-N-[4-[5-[4-[2-oxo-2-(azetidin-1-yl)ethyl]-2-(ethyl-sulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 87)**

Following the same General Method A, Compound 87 was prepared. ¹H NMR (400MHz, methanol-*d*₄) δ = 7.99 (d, J=1.6 Hz, 1H), 7.81 (s, 1H), 7.57 (dd, J=1.8, 7.9 Hz, 1H), 7.52 - 7.45 (m, 1H), 4.89 - 4.88 (m, 1H), 4.34 (t, J=7.7 Hz, 2H), 4.06 (t, J=7.8 Hz, 2H),
 10 3.65 (s, 2H), 3.48 (tt, J=3.9, 11.6 Hz, 1H), 3.08 (tt, J=3.4, 12.0 Hz, 1H), 2.90 (q, J=7.3 Hz, 2H), 2.41 - 2.32 (m, 2H), 2.30 - 2.23 (m, 2H), 2.15 - 2.06 (m, 2H), 1.73 (dq, J=2.9, 12.9 Hz, 2H), 1.51 - 1.37 (m, 2H), 1.25 (br d, J=6.1 Hz, 6H), 1.05 (t, J=7.2 Hz, 3H). ESI [M+H] = 549.2.

15 **Example 64. Preparation of isopropyl *trans*-N-[4-[5-(4-(methylamino)-2-pyrrolidin-1-ylsulfonyl-phenyl)thiazol-2-yl]cyclohexyl]carbamate (Compound 89)**




a) **Synthesis of 1-(2-bromo-5-nitrophenyl)sulfonylpyrrolidine**




From 2-bromo-5-nitrobenzenesulfonyl chloride General Method G. ^1H NMR (400 MHz, methanol- d_4) δ = 8.81 (d, J =2.69 Hz, 1 H), 8.33 (dd, J =8.68, 2.69 Hz, 1 H), 8.13 (d, J =8.68 Hz, 1 H), 3.44 - 3.49 (m, 4 H), 1.97 (dt, J =6.48, 3.48 Hz, 4 H).

b) Synthesis of 4-bromo-3-pyrrolidin-1-ylsulfonyl-aniline




A mixture of 1-(2-bromo-5-nitro-phenyl)sulfonylpyrrolidine (3.0 g, 9.0 mmol, 1.0 *eq.*), NH_4Cl (1.4 g, 27.0 mmol, 3.0 *eq.*), Fe (2.5 g, 45.0 mmol, 5.0 *eq.*) in THF (30 mL), EtOH (30 mL) and H_2O (10 mL) was stirred at 80°C for 4 h. The reaction mixture was filtered, and the filtrate was concentrated. The residue was diluted with H_2O (10 mL) and extracted with EtOAc 150 mL (50 mL X 3). The combined organic layers dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified by column chromatography (SiO_2 , Petroleum ether: Ethyl acetate = 10:1 to 1:1 to 0:1) to yield 4-bromo-3-pyrrolidin-1-ylsulfonyl-aniline (2.3 g, 7.5 mmol, 84% yield) as a white solid. ESI $[\text{M}+\text{H}] = 304.9/306.9$.

c) Synthesis of 3-pyrrolidin-1-ylsulfonyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline



From 4-bromo-3-pyrrolidin-1-ylsulfonyl-aniline, using General Method B. ^1H NMR (400MHz, $\text{DMSO}-d_6$) δ = 7.13 - 7.04 (m, 1H), 6.96 (s, 1H), 6.80 - 6.70 (m, 1H), 3.16 (br t, J =6.3 Hz, 4H), 1.67 - 1.62 (m, 4H), 1.25 (s, 12H). ESI $[\text{M}+\text{H}] = 353.1$.


d) Synthesis of isopropyl *trans*-*N*-[4-[5-(4-amino-2-pyrrolidin-1-ylsulfonyl-phenyl)thiazol-2-yl]cyclohexyl]carbamate (**Compound 88**)



From 3-pyrrolidin-1-ylsulfonyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline, using General Method D. ^1H NMR (400MHz, methanol- d_4) δ = 7.78 - 7.66 (m,


1H), 7.41 (br s, 1H), 7.24 (br d, J=8.2 Hz, 1H), 7.00 - 6.90 (m, 1H), 4.87 - 4.77 (m, 1H), 3.45 (br t, J=11.7 Hz, 1H), 3.12 - 3.05 (m, 1H), 3.01 (br t, J=6.1 Hz, 4H), 2.20 (br d, J=12.1 Hz, 2H), 2.07 (br d, J=10.6 Hz, 2H), 1.86 - 1.75 (m, 4H), 1.74 - 1.61 (m, 2H), 1.48 - 1.34 (m, 2H), 1.22 (br d, J=6.0 Hz, 6H). ESI [M+H]⁺=493.2.

- 5 e) Synthesis of isopropyl *trans*-N-[4-[5-(4-(methylamino)-2-pyrrolidin-1-ylsulfonyl-phenyl)thiazol-2-yl]cyclohexyl]carbamate (**Compound 89**)




- A mixture of isopropyl *trans*-N-[4-[5-(4-amino-2-pyrrolidin-1-ylsulfonyl-phenyl)thiazol-2-yl]cyclohexyl]carbamate (170 mg, 345 μmol, 1 *eq.*), TFA (4 mg, 35 μmol, 0.1 *eq.*) in diethoxymethoxyethane (2 mL) was stirred at 100°C for 12 h. The mixture was concentrated followed by addition of EtOH (1 mL) and NaBH₄ (39 mg, 1 mmol, 3 *eq.*) at 0°C. The mixture was stirred at 80°C for 2 h and quenched by H₂O (2 mL) and extracted with EtOAc 10 mL (5 mL X 2). The combined organic layers were washed with H₂O (5 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (column: Welch Ultimate AQ-C18 150*30mm*5μm; mobile phase: [water(0.1%TFA)-ACN]; B%: 40%-70%, 12min) to yield isopropyl *trans*-N-[4-[5-(4-(methylamino)-2-pyrrolidin-1-ylsulfonyl-phenyl)thiazol-2-yl]cyclohexyl]carbamate (34 mg, 64 μmol, 19% yield, 92% purity) as a yellow solid. ¹H NMR (400MHz, methanol-*d*₄) δ = 7.67 (s, 1H), 7.29 - 7.19 (m, 2H), 6.80 (dd, J=2.2, 8.4 Hz, 1H), 4.83 (br s, 1H), 3.45 (br t, J=11.6 Hz, 1H), 3.09 - 3.02 (m, 1H), 2.99 (br t, J=6.4 Hz, 4H), 2.83 (s, 3H), 2.20 (br d, J=12.1 Hz, 2H), 2.07 (br d, J=11.2 Hz, 2H), 1.78 (br t, J=6.5 Hz, 4H), 1.75 - 1.61 (m, 2H), 1.47 - 1.34 (m, 2H), 1.22 (br d, J=6.0 Hz, 6H). ESI [M+H]⁺ = 507.2.

- Example 65. Preparation of isopropyl *trans*-N-[4-[5-[4-(1*H*-imidazol-2-ylamino)-2-sulfamoyl-phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 90; General Method K)**




To a solution of isopropyl *trans-N*-[4-[5-[4-amino-2-(tert-butylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (200 mg, 404 μ mol, 1 *eq.*) in *n*-BuOH (4 mL) was added TsOH (209 mg, 1 mmol, 3 *eq.*) and 2-bromo-1H-imidazole (297 mg, 2 mmol, 5 *eq.*). The mixture was stirred at 140°C for 12 h. The reaction mixture was concentrated under reduced pressure and purified by prep-HPLC (column: Waters Xbridge BEH C18 100*30mm*10 μ m; mobile phase: [water(0.1%TFA)-ACN]; B%: 30%-90%, 10 min) to yield isopropyl *trans-N*-[4-[5-[4-(1H-imidazol-2-ylamino)-2-sulfamoyl-phenyl]thiazol-2-yl]cyclohexyl]carbamate (12 mg, 20 μ mol, 40% yield, 100% purity, TFA) as a yellow solid. ¹H NMR (400MHz, methanol-*d*₄) δ = 7.97 (d, J=2.0 Hz, 1H), 7.80 - 7.74 (m, 1H), 7.58 - 7.52 (m, 1H), 7.51 - 7.45 (m, 1H), 7.10 (s, 2H), 4.83 - 4.76 (m, 1H), 3.45 (br t, J=11.6 Hz, 1H), 3.09 - 2.96 (m, 1H), 2.24 (br d, J=12.8 Hz, 2H), 2.11 - 1.96 (m, 2H), 1.83 - 1.64 (m, 2H), 1.47 - 1.32 (m, 2H), 1.22 (br d, J=6.0 Hz, 6H). ESI [M+H] = 505.1.

Example 66. Preparation of isopropyl *trans-N*-[4-[5-[2-(dimethylsulfamoyl)-4-(1H-imidazol-2-ylamino)phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 91)




a) Synthesis of 2-bromo-*N,N*-dimethyl-5-nitro-benzenesulfonamide



From 2-bromo-5-nitrobenzenesulfonyl chloride and dimethylamine, using General Method G. ESI [M+H] = 308.9/310.9.


b) Synthesis of 5-amino-2-bromo-*N,N*-dimethyl-benzenesulfonamide



To a solution of 2-bromo-*N,N*-dimethyl-5-nitro-benzenesulfonamide (4.6 g, 14.8 mmol, 1.0 *eq.*) in H₂O (20 mL), EtOH (60 mL) and THF (60 mL) was added Fe (4.1 g, 74.1


mmol, 5.0 *eq.*) and NH₄Cl (2.4 g, 44.5 mmol, 3.0 *eq.*). The mixture was stirred at 80°C for 2 h then filtered, and the filtrate was concentrated under reduced pressure. The residue was diluted with H₂O (20 mL) and extracted with EtOAc 90 mL (30 mL X 3). The combined organic layers were dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by column chromatography (SiO₂, Petroleum ether: Ethyl acetate = 1:0 to 0:1) to yield 5-amino-2-bromo-*N,N*-dimethyl-benzenesulfonamide (3.9 g, 14.0 mmol, 94 % yield) as a pale yellow solid. ¹H NMR (400 MHz, methanol-*d*₄) δ = 7.43 (d, J = 8.6 Hz, 1H), 7.35 (d, J = 2.8 Hz, 1H), 6.75 (dd, J = 2.8, 8.6 Hz, 1H), 2.86 (s, 6H). ESI [M+H] = 278.9/281.0.

- 10 c) Synthesis of 2-bromo-5-(1*H*-imidazol-2-ylamino)-*N,N*-dimethyl-benzene sulfonamide




- From 5-amino-2-bromo-*N,N*-dimethyl-benzenesulfonamide, using General Method K. ¹H NMR (400 MHz, methanol-*d*₄) δ = 7.97 - 7.84 (m, 2H), 7.41 (dd, J = 2.8, 8.5 Hz, 1H), 7.12 (s, 2H), 2.93 (s, 6H). ESI [M+H] = 345.0/347.0.

- 15 d) Synthesis of 5-(1*H*-imidazol-2-ylamino)-*N,N*-dimethyl-2-(4,4,5,5-tetra methyl-1,3,2-dioxaborolan-2-yl)benzenesulfonamide




- A solution of 2-bromo-5-(1*H*-imidazol-2-ylamino)-*N,N*-dimethyl-benzenesulfonamide (380 mg, 1 mmol, 1 *eq.*), 4,4,5,5-tetramethyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1,3,2-dioxaborolane (1.4 g, 5.5 mmol, 5.0 *eq.*), [2-(2-aminophenyl)phenyl]-chloro-palladium;bis(1-adamantyl)-butyl-phosphane (74 mg, 110 umol, 0.1 *eq.*), DIEA (711 mg, 6 mmol, 5 *eq.*) in MeOH (10 mL) was stirred at 50°C for 12 h under N₂ atmosphere. The reaction mixture was concentrated and purified by column chromatography (SiO₂, petroleum ether: ethyl acetate = 1:0 to THF) to yield 5-(1*H*-imidazol-2-ylamino)-*N,N*-dimethyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzenesulfonamide (300 mg, crude) as a brown oil. ¹H NMR (400 MHz, methanol-*d*₄) δ = 7.53 - 7.40 (m, 1H), 7.39 - 7.31 (m, 1H), 7.29 - 7.14 (m, 1H), 6.92 (s, 1H), 6.82 (d, J = 2.4 Hz, 1H), 3.72 (td, J = 6.6, 13.2 Hz, 3H), 3.22 (q, J = 7.5 Hz, 3H), 1.38 - 1.36 (m, 12H). ESI [M+H] = 393.1.

e) Synthesis of isopropyl *trans*-*N*-[4-[5-[2-(dimethylsulfamoyl)-4-(1*H*-imidazol-2-ylamino)phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 91**)




From 5-(1*H*-imidazol-2-ylamino)-*N,N*-dimethyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzenesulfonamide, using General Method D. ¹H NMR (400 MHz, methanol-*d*₄) δ = 7.90 (d, *J* = 2.3 Hz, 1H), 7.72 (s, 1H), 7.62 - 7.51 (m, 2H), 7.14 (s, 2H), 4.84 - 4.78 (m, 1H), 3.47 (tt, *J* = 3.9, 11.6 Hz, 1H), 3.05 (tt, *J* = 3.5, 12.0 Hz, 1H), 2.65 - 2.55 (m, 6H), 2.32 - 2.18 (m, 2H), 2.14 - 2.03 (m, 2H), 1.71 (dq, *J* = 2.9, 12.8 Hz, 2H), 1.44 (dq, *J* = 3.3, 12.5 Hz, 2H), 1.25 (br d, *J* = 6.3 Hz, 6H). ESI [*M*+*H*] = 533.2.

10 **Example 67. Preparation of isopropyl *trans*-*N*-[4-[5-[4-(1*H*-imidazol-2-ylamino)-2-pyrrolidin-1-ylsulfonyl-phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 92**)**




a) Synthesis of *N*-(4-bromo-3-(pyrrolidin-1-ylsulfonyl-phenyl)-1*H*-imidazol-2-amine




15 From 4-bromo-3-(pyrrolidin-1-ylsulfonyl)aniline, using General Method K. ¹H NMR (400 MHz, methanol-*d*₄) δ = 7.82 - 7.92 (m, 2 H), 7.38 (dd, *J*=8.49, 2.76 Hz, 1 H), 7.06 - 7.11 (m, 2 H), 3.36 - 3.48 (m, 4 H), 1.93 (dt, *J*=6.45, 3.50 Hz, 4 H). ESI [*M*+*H*] = 370.9/372.9.

b) Synthesis of *N*-[3-(pyrrolidin-1-ylsulfonyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]-1*H*-imidazol-2-amine




A mixture of 4,4,5,5-tetramethyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1,3,2-dioxaborolane (654 mg, 3 mmol, 5 *eq.*), *N*-(4-bromo-3-pyrrolidin-1-ylsulfonyl-phenyl)-1*H*-imidazol-2-amine (250 mg, 515 μ mol, 1 *eq.*, TFA), [2-(2-amino-phenyl)phenyl]-chloro-
 5 palladium; bis(1-adamantyl)-butyl-phosphane (34 mg, 52 μ mol, 0.1 *eq.*), DIEA (333 mg, 3 mmol, 5 *eq.*) in MeOH (15 mL) was degassed and purged with N₂ for 3 times, and then the mixture was stirred at 50°C for 12 h under N₂ atmosphere. The reaction mixture was concentrated and purified by column chromatography (SiO₂, petroleum ether: ethyl acetate = 1:0 to 5:1 to 1:1 to 0:1) to yield *N*-[3-pyrrolidin-1-ylsulfonyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]-1*H*-imidazol-2-amine (190 mg, 454 μ mol, 88% yield) as a white solid. ESI [M+H] = 419.2.

c) Synthesis of isopropyl *trans*-*N*-[4-[5-[4-(1*H*-imidazol-2-ylamino)-2-pyrrolidin-1-ylsulfonyl-phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 92**)




From *N*-[3-pyrrolidin-1-ylsulfonyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]-1*H*-imidazol-2-amine, using General Method D. ¹H NMR (400 MHz, methanol-*d*₄) δ = 7.91 (d, *J*=1.75 Hz, 1 H), 7.76 (br s, 1 H), 7.50 - 7.59 (m, 2 H), 7.13 (s, 2 H), 4.82 (br d, *J*=6.00 Hz, 1 H), 3.46 (br t, *J*=11.57 Hz, 1 H), 3.04 (br s, 1 H), 2.97 (br t, *J*=6.25 Hz, 4 H), 2.21 (br d, *J*=12.63 Hz, 2 H), 2.07 (br d, *J*=10.88 Hz, 2 H), 1.78 (br t, *J*=6.25 Hz, 4 H), 1.69
 20 (br d, *J*=12.01 Hz, 2 H), 1.38 - 1.48 (m, 2 H), 1.23 (br d, *J*=6.13 Hz, 6 H). ESI [M+H] = 559.2.

Example 68. Preparation of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-(1*H*-imidazol-2-ylcarbamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 94)




a) Synthesis of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[[1-(2-trimethylethoxymethyl)-1*H*-imidazol-2-yl]carbamoyl]phenyl]thiazol-2-yl]cyclohexyl] carbamate



5 From *trans*-3-(ethylsulfamoyl)-4-[2-[4-(isopropoxyloxycarbonyl-amino) cyclohexyl]thiazol-5-yl]benzoic acid (**Compound 48**) and 1-((2-(trimethylsilyl)ethoxy)methyl)-1*H*-imidazol-2-amine, using General Method A. ESI [M-H] = 691.3.

b) Synthesis of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-(1*H*-imidazol-2-yl)carbamoyl]phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 94**)




10

From isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[[1-(2-trimethylethoxymethyl)-1*H*-imidazol-2-yl]carbamoyl]phenyl]thiazol-2-yl]cyclohexyl] carbamate, using General Method I. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.78 (d, J=1.63 Hz, 1 H), 8.28 (dd, J=8.00, 1.63 Hz, 1 H), 7.89 (s, 1 H), 7.75 (d, J=7.88 Hz, 1 H), 7.29 (s, 2 H), 4.77 - 4.84 (m, 1 H), 3.42 - 3.53 (m, 1 H), 3.07 (tt, J=11.99, 3.39 Hz, 1 H), 2.90 (q, J=7.21 Hz, 2 H), 2.29 (br d, J=12.76 Hz, 2 H), 2.05 - 2.14 (m, 2 H), 1.74 (qd, J=12.88, 3.00 Hz, 2 H), 1.45 (qd, J=12.51, 3.00 Hz, 2 H), 1.20 - 1.31 (m, 6 H), 1.03 (t, J=7.19 Hz, 3 H). ESI [M+H] = 561.2.

15

Example 69. Preparation of isopropyl *trans*-N-[4-[5-[4-carbamoyl-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 49)



From *trans*-3-(ethylsulfamoyl)-4-[2-[4-(isopropylloxycarbonyl-amino)cyclohexyl]thiazol-5-yl]benzoic acid (**Compound 48**) and ammonium chloride, using General Method A. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.56 (d, *J* = 1.5 Hz, 1H), 8.09 (dd, *J* = 1.8, 7.9 Hz, 1H), 7.85 (s, 1H), 7.61 (d, *J* = 7.9 Hz, 1H), 4.85 - 4.74 (m, 1H), 3.52 - 3.40 (m, 1H), 3.05 (tt, *J* = 3.5, 11.9 Hz, 1H), 2.89 (q, *J* = 7.2 Hz, 2H), 2.32 - 2.20 (m, 2H), 2.11 - 2.04 (m, 2H), 1.71 (dq, *J* = 2.8, 13.0 Hz, 2H), 1.48 - 1.36 (m, 2H), 1.22 (br d, *J* = 6.2 Hz, 6H), 1.02 (t, *J* = 7.3 Hz, 3H). ESI [M+H] = 495.1.

Example 70. Preparation of isopropyl (*trans*-4-(5-(2-(*N*-ethylsulfamoyl)-4-


(isopropylcarbamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (Compound 50)

Following the same protocol and under the same reaction conditions as for Compound 94, Compound 50 was prepared. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.52 (s, 1H), 8.05 (br d, *J* = 7.5 Hz, 1H), 7.83 (s, 1H), 7.61 (d, *J* = 7.9 Hz, 1H), 4.88 - 4.80 (m, 1H), 4.25 (quin, *J* = 6.6 Hz, 1H), 3.48 (ddd, *J* = 4.2, 7.7, 11.4 Hz, 1H), 3.12 - 2.99 (m, 1H), 2.91 (q, *J* = 7.1 Hz, 2H), 2.27 (br d, *J* = 12.5 Hz, 2H), 2.17 - 2.02 (m, 2H), 1.84 - 1.64 (m, 2H), 1.49 - 1.40 (m, 2H), 1.30 (d, *J* = 6.6 Hz, 6H), 1.27 - 1.16 (m, 6H), 1.04 (t, *J* = 7.2 Hz, 3H). ESI [M+H] = 537.2.

Example 71. Preparation of isopropyl *trans*-N-[4-[5-[4-pyrrolidin-1-ylcarbonyl-2-(ethylsulfamoyl)phenyl] thiazol-2-yl]cyclohexyl]carbamate (Compound 95)

Following the same protocol and under the same reaction conditions as for Compound 94, Compound 95 was prepared. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.20 (d, *J* = 1.5 Hz, 1H), 7.85 - 7.76 (m, 2H), 7.60 (d, *J* = 7.9 Hz, 1H), 4.84 - 4.76 (m, 1H), 3.63 (t, *J* = 6.8 Hz, 2H), 3.51 (t, *J* = 6.5 Hz, 2H), 3.48 - 3.40 (m, 1H), 3.03 (tt, *J* = 3.5, 12.0 Hz, 1H), 2.88 (q, *J* = 7.3 Hz, 2H), 2.30 - 2.19 (m, 2H), 2.07 (br dd, *J* = 3.2, 13.3 Hz, 2H), 2.03 - 1.90 (m, 4H), 1.71 (dq, *J* = 2.9, 12.9 Hz, 2H), 1.47 - 1.36 (m, 2H), 1.22 (br d, *J* = 6.2 Hz, 6H), 1.02 (t, *J* = 7.3 Hz, 3H). ESI [M+H] = 549.3.

Example 72. Preparation of oxetan-3-yl *trans*-N-[4-[5-[4-carbamoyl-2-(ethylsulfamoyl)phenyl] thiazol-2-yl]cyclohexyl]carbamate (Compound 67)




From *trans*-3-(ethylsulfamoyl)-4-[2-[4-(oxetan-3-yloxycarbonyl-amino)cyclohexyl]thiazol-5-yl]benzoic acid (**Compound 66**) and ammonium chloride, using General Method A. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.56 (d, *J* = 1.8 Hz, 1H), 8.08 (dd, *J* = 1.8, 7.9 Hz, 1H), 7.81 (s, 1H), 7.60 (d, *J* = 7.9 Hz, 1H), 5.35 (quin, *J* = 5.7 Hz, 1H), 4.84 (br s, 2H), 4.65 - 4.54 (m, 2H), 3.45 (ddd, *J* = 3.7, 7.7, 11.5 Hz, 1H), 3.09 - 2.97 (m, 1H), 2.89 (q, *J* = 7.3 Hz, 2H), 2.29 - 2.19 (m, 2H), 2.08 (br d, *J* = 10.4 Hz, 2H), 1.71 (dq, *J* = 3.0, 12.8 Hz, 2H), 1.50 - 1.36 (m, 2H), 1.02 (t, *J* = 7.2 Hz, 3H). ESI [*M*+*H*] = 509.2.


Example 73. Preparation of oxetan-3-yl (*trans*-4-(5-(2-(*N*-ethylsulfamoyl)-4-(isopropylcarbamoyl)phenyl)thiazol-2-yl)cyclohexyl)carbamate (Compound 68**)**

Following the same protocol and under the same reaction conditions as for Compound 67, Compound 68 was prepared. ¹H NMR (400 MHz, DMSO-*d*₆) δ = 8.53 (d, *J* = 7.7 Hz, 1H), 8.39 (d, *J* = 1.8 Hz, 1H), 8.07 (dd, *J* = 1.7, 8.0 Hz, 1H), 7.76 (s, 1H), 7.59 (d, *J* = 7.9 Hz, 1H), 7.53 (br s, 1H), 7.43 (d, *J* = 7.5 Hz, 1H), 5.26 (quin, *J* = 5.7 Hz, 1H), 4.73 (t, *J* = 6.9 Hz, 2H), 4.46 - 4.40 (m, 2H), 4.15 - 4.03 (m, 1H), 3.29 (s, 1H), 2.94 (tt, *J* = 3.6, 11.9 Hz, 1H), 2.77 (q, *J* = 7.1 Hz, 2H), 2.14 (br d, *J* = 11.7 Hz, 2H), 1.91 (br d, *J* = 9.7 Hz, 2H), 1.62 - 1.51 (m, 2H), 1.41 - 1.30 (m, 2H), 1.16 (d, *J* = 6.6 Hz, 6H), 0.93 (t, *J* = 7.3 Hz, 3H). ESI [*M*+*H*] = 551.2.

Example 74. Preparation of isopropyl *trans*-*N*-[4-[5-[4-[2-(benzylamino)-2-oxo-ethyl]-2-(*tert*-butylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 97**)**




a) Synthesis of methyl 2-[4-bromo-3-(*tert*-butylsulfamoyl)phenyl]acetate



To a solution of 2-methylpropan-2-amine (2.6 g, 36.6 mmol, 3 *eq.*) in DCM (60 mL) was added DMAP (149 mg, 1 mmol, 0.1 *eq.*), and methyl 2-(4-bromo-3-chlorosulfonylphenyl)acetate (4.0 g, 12.2 mmol, 1 *eq.*) at 0°C. The mixture was stirred at 25°C for 2 h, then diluted with H₂O (100 mL) and extracted with DCM 400 mL (200 mL X 2). The combined organic layers were dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was slurried in the solution (petroleum ether : ethyl acetate=20:1, 10 mL) for 1 h. Then it was filtered to yield methyl 2-[4-bromo-3-(*tert*-butylsulfamoyl)phenyl]acetate (2.2 g, 6.0 mmol, 49.4 % yield) as a yellow solid. ¹H NMR (400MHz, methanol-*d*₄) δ = 8.16 - 8.00 (m, 1H), 7.80 - 7.62 (m, 1H), 7.36 (dd, J=2.0, 8.2 Hz, 1H), 3.73 (s, 2H), 3.69 - 3.64 (m, 3H), 1.21 - 1.15 (m, 9H). ESI[M-H] = 361.9/363.9.


b) Synthesis of 2-[4-bromo-3-(*tert*-butylsulfamoyl)phenyl]acetic acid



15


From methyl 2-[4-bromo-3-(*tert*-butylsulfamoyl)phenyl]acetate, using General Method J.

c) Synthesis of *N*-benzyl-2-[4-bromo-3-(*tert*-butylsulfamoyl)phenyl] acetamide




From 2-[4-bromo-3-(*tert*-butylsulfamoyl)phenyl]acetic acid and benzylamine, using General Method A. ¹H NMR (400MHz, methanol-*d*₄) δ = 8.11 (d, *J*=2.0 Hz, 1H), 7.76 - 7.65 (m, 1H), 7.44 - 7.33 (m, 2H), 7.32 - 7.18 (m, 4H), 4.36 (s, 2H), 3.59 (s, 2H), 1.21 - 1.10 (m, 9H). ESI [M+H] = 439.0/441.0.

d) Synthesis of *N*-benzyl-2-[3-(*tert*-butylsulfamoyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]acetamide



From *N*-benzyl-2-[4-bromo-3-(*tert*-butylsulfamoyl)phenyl] acetamide, using General Method B. ESI[M+H] = 487.2.

e) Synthesis of isopropyl *trans*-*N*-[4-[5-[4-[2-(benzylamino)-2-oxo-ethyl]-2-(*tert*-butylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 97**)




From *N*-benzyl-2-[3-(*tert*-butylsulfamoyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]acetamide, using General Method C. ¹H NMR (400MHz, methanol-*d*₄) δ = 8.14 (d, *J*=1.5 Hz, 1H), 7.79 (s, 1H), 7.56 (dd, *J*=1.7, 7.8 Hz, 1H), 7.43 (d, *J*=7.7 Hz, 1H), 7.34 - 7.19 (m, 5H), 4.84 (br s, 1H), 4.38 (s, 2H), 3.67 (s, 2H), 3.51 - 3.39 (m, 1H), 3.03 (tt, *J*=3.4, 12.0 Hz, 1H), 2.23 (br d, *J*=12.6 Hz, 2H), 2.07 (br d, *J*=10.6 Hz, 2H), 1.76 - 1.60 (m, 2H), 1.41 (dq, *J*=3.2, 12.5 Hz, 2H), 1.22 (br d, *J*=6.2 Hz, 6H), 1.13 - 0.99 (m, 9H). ESI [M+H] = 627.3.

20 **Example 75. Preparation of oxetan-3-yl *trans*-*N*-[4-[5-[4-[2-(benzylamino)-2-oxo-ethyl]-2-(*tert*-butylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 98)**


Following the same protocol and under the same reaction conditions as for Compound 97, Compound 98 was prepared. ¹H NMR (400MHz, DMSO-*d*₆) δ = 8.66 (t, *J*=6.0 Hz, 1H), 8.01 (s, 1H), 7.68 (s, 1H), 7.50 (br d, *J*=7.9 Hz, 1H), 7.45 - 7.36 (m, 2H), 7.33 - 7.26 (m, 2H), 7.25 - 7.18 (m, 3H), 6.94 (s, 1H), 5.26 (quin, *J*=5.7 Hz, 1H), 4.73 (t, *J*=6.9 Hz, 2H), 4.52 - 4.39 (m, 2H), 4.26 (d, *J*=5.7 Hz, 2H), 3.60 (s, 2H), 3.30 - 3.20 (m, 1H), 2.99 - 2.82 (m, 1H),

2.12 (br d, $J=11.5$ Hz, 2H), 1.90 (br d, $J=10.4$ Hz, 2H), 1.65 - 1.48 (m, 2H), 1.42 - 1.27 (m, 2H), 1.00 (s, 9H). ESI $[M+H]^+ = 641.3$.

Example 76. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-isopentyloxy-phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 99)




a) Synthesis of 2-bromo-N-ethyl-5-hydroxy-benzenesulfonamide



To a solution of 5-amino-2-bromo-N-ethyl-benzenesulfonamide (5 g, 17.91 mmol, 1 eq) in H₂O (200 mL) was added trifluoroborane; hydrofluoride (7.8 g, 35.8 mmol, 200 mL, 40% purity). Then NaNO₂ (1.4 g, 19.7 mmol, 1.1 eq) in H₂O (140 mL) was added batch at 0°C for 30 mins. Then, sat.aq. CuSO₄ (416.0 g, 2.6 mol, 145.52 eq) and CuO (1.42 g, 17.91 mmol, 225.44 uL, 1 eq) was added batchwise. The mixture was stirred at 20°C for 2 h. The reaction mixture was extracted with DCM 150 mL (50 mL X 3). The combined organic layers were dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude product was purified by reversed-phase HPLC (0.1% TFA condition) to yield 2-bromo-N-ethyl-5-hydroxy-benzenesulfonamide (2.2 g) as a yellow gum. ESI $[M+H]^+ = 279.9/281.9$.


b) Synthesis of 2-bromo-N-ethyl-5-isopentyloxy-benzenesulfonamide



A mixture of 2-bromo-N-ethyl-5-hydroxy-benzenesulfonamide (700 mg, 2.5 mmol, 1 eq), 1-bromo-3-methyl-butane (1.1 g, 7.5 mol, 3.0 eq), K₂CO₃ (1.0 g, 7.5 mmol, 3.0 eq)


in DMF (7 mL) was stirred at 60°C for 2 h under N₂ atmosphere. The reaction mixture was quenched by H₂O 70 mL and extracted with EtOAc 120 mL (40 mL X 3). The combined organic layers were dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by column chromatography (SiO₂, petroleum ether: ethyl acetate=50:1 to 0:1) to yield 2-bromo-*N*-ethyl-5-isopentyloxy-benzenesulfonamide (920mg) as colorless oil. ¹H NMR (400 MHz, methanol-*d*₄) δ = 7.65 (d, J = 8.8 Hz, 1H), 7.59 (d, J = 2.9 Hz, 1H), 7.16 - 7.15 (m, 1H), 4.06 (t, J = 6.5 Hz, 2H), 2.94 (q, J = 7.2 Hz, 2H), 1.84 (td, J = 6.7, 13.5 Hz, 1H), 1.74 - 1.61 (m, 2H), 1.06 (t, J = 7.2 Hz, 3H), 0.98 (d, J = 6.6 Hz, 6H). ESI [M+H] = 349.9/352.0.

- 10 c) Synthesis of *N*-ethyl-5-isopentyloxy-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzenesulfonamide



- From 2-bromo-*N*-ethyl-5-isopentyloxy-benzenesulfonamide, using General Method B. ¹H NMR (400 MHz, methanol-*d*₄) δ = 7.64 (d, J = 8.2 Hz, 1H), 7.42 (d, J = 2.4 Hz, 1H), 7.14 (dd, J = 2.5, 8.3 Hz, 1H), 4.09 (t, J = 6.5 Hz, 2H), 2.90 (q, J = 7.3 Hz, 2H), 1.70 (q, J = 6.6 Hz, 2H), 1.39 (s, 1H), 1.38 (s, 12H), 1.03 (t, J = 7.2 Hz, 3H), 0.98 (d, J = 6.6 Hz, 6H). ESI [M+H] = 398.1.

- d) Synthesis of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-isopentyloxy-phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 99**)




- From *N*-ethyl-5-isopentyloxy-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzenesulfonamide, using General Method C. ¹H NMR (400 MHz, DMSO-*d*₆) δ = 7.60 (s, 1H), 7.46 (t, J = 5.6 Hz, 1H), 7.43 - 7.36 (m, 2H), 7.19 (dd, J = 2.6, 8.6 Hz, 1H), 7.01 (br d, J = 7.7 Hz, 1H), 4.72 (td, J = 6.3, 12.4 Hz, 1H), 4.08 (t, J = 6.5 Hz, 2H), 3.36 - 3.22 (m, 1H), 2.93 - 2.84 (m, 1H), 2.82 - 2.72 (m, 2H), 2.11 (br d, J = 11.9 Hz, 2H), 1.89 (br d, J = 10.4 Hz, 2H), 1.82 - 1.71 (m, 1H), 1.63 (q, J = 6.5 Hz, 2H), 1.54 (dq, J = 2.8, 12.8 Hz, 2H), 1.38 - 1.25 (m, 2H), 1.14 (d, J = 6.2 Hz, 6H), 0.98 - 0.93 (m, 3H), 0.92 (d, J = 6.6 Hz, 6H). ESI [M+H] = 538.2.


Example 77. Preparation of oxetan-3-yl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-isopentyloxy-phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 100)

Following the same protocol and under the same reaction conditions as for Compound 99, Compound 100 was prepared. ¹H NMR (400 MHz, DMSO-*d*₆) δ = 7.60 (s, 1H), 7.55 - 7.29 (m, 4H), 7.19 (dd, J = 2.5, 8.5 Hz, 1H), 5.25 (quin, J = 5.7 Hz, 1H), 4.73 (t, J = 6.8 Hz, 2H), 4.49 - 4.37 (m, 2H), 4.08 (t, J = 6.5 Hz, 2H), 3.30 - 3.25 (m, 1H), 2.90 (tt, J = 3.3, 11.8 Hz, 1H), 2.78 (q, J = 7.3 Hz, 2H), 2.12 (br d, J = 11.9 Hz, 2H), 1.90 (br d, J = 10.6 Hz, 2H), 1.83 - 1.70 (m, 1H), 1.67 - 1.59 (m, 2H), 1.59 - 1.46 (m, 2H), 1.39 - 1.29 (m, 2H), 0.98 - 0.93 (m, 3H), 0.92 (d, J = 6.4 Hz, 6H). ESI [M+H] = 552.2.

Example 78. Preparation of isopropyl *trans*-N-[4-[5-[2-(*tert*-butylsulfamoyl)-4-isopentyloxy-phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 101)




a) Synthesis of 2-bromo-*N*-*tert*-butyl-5-hydroxy-benzenesulfonamide




To isopropyl 5-amino-2-bromo-*N*-*tert*-butyl-benzenesulfonamide (10.0 g, 32.6 mmol, 1.0 *eq.*) dissolved in H₂SO₄ (120 mL, 30% aq.) was added a solution of NaNO₂ (2.3 g, 1.1 mol, 34.5 *eq.*) in H₂O (20 mL) at 0°C and the mixture was stirred at 0°C for 30 min. Then was added H₂O (280 mL) to the mixture and stirred at 65°C for 11 h. The mixture was extracted with EtOAc (300 mL X 3). The combined organic layers were dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by reversed-phase chromatography to yield 2-bromo-*N*-*tert*-butyl-5-hydroxy-benzene sulfonamide (1.4 g, 4.5 mmol, 14% yield) was obtained as a red solid. ESI [M-H] = 305.8/307.8.

b) Synthesis of 2-bromo-*N*-*tert*-butyl-5-isopentyloxy-benzenesulfonamide




To a solution of 2-bromo-*N*-*tert*-butyl-5-hydroxy-benzenesulfonamide (1.5 g, 4.7 mmol, 1.0 *eq.*) in DMF (30 mL) was added K₂CO₃ (2.0 g, 14.1 mmol, 3.0 *eq.*) and 1-bromo-3-methyl-butane (1.1 g, 7.1 mmol, 1.5 *eq.*). The reaction mixture was stirred at 100°C for 12 h. The mixture was poured into water (100 mL) and the solid filtrated, yielding 2-bromo-*N*-*tert*-butyl-5-isopentyloxy-benzenesulfonamide (1.4 g, crude) as a yellow solid. ¹H NMR (400 MHz, methanol-*d*₄) δ = 7.61 - 7.70 (m, 2 H), 7.04 (dd, *J*=8.68, 2.69 Hz, 1 H), 4.08 (t, *J*=6.54 Hz, 2 H), 1.80 - 1.92 (m, 1 H), 1.71 (q, *J*=6.48 Hz, 2 H), 1.14 - 1.26 (m, 9 H), 0.99 (s, 6 H).

c) Synthesis of *N*-*tert*-butyl-5-isopentyloxy-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzenesulfonamide



From 2-bromo-*N*-*tert*-butyl-5-isopentyloxy-benzenesulfonamide, using General Method B. ¹H NMR (400MHz, CDCl₃) δ = 7.66 (d, *J*=8.2 Hz, 1H), 7.50 (d, *J*=2.4 Hz, 1H), 6.92 - 6.89 (m, 1H), 3.98 - 3.94 (m, 2H), 1.64 - 1.59 (m, 3H), 1.32 (s, 12H), 1.14 (s, 9H), 0.91 - 0.88 (m, 6H). ESI [M+H] = 426.2.

15 d) Synthesis of isopropyl *trans*-*N*-[4-[5-[2-(*tert*-butylsulfamoyl)-4-isopentyloxy-phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 101**)




From *N*-*tert*-butyl-5-isopentyloxy-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzenesulfonamide, using General Method C. ¹H NMR (400MHz, methanol-*d*₄) δ = 7.68 - 7.53 (m, 2H), 7.30 (d, *J*=8.4 Hz, 1H), 7.08 (dd, *J*=2.6, 8.4 Hz, 1H), 4.77 - 4.69 (m, 1H), 4.03 (t, *J*=6.5 Hz, 2H), 3.36 (ddd, *J*=3.9, 7.8, 11.5 Hz, 1H), 2.92 (br t, *J*=12.1 Hz, 1H), 2.14 (br d, *J*=12.2 Hz, 2H), 1.98 (br d, *J*=10.6 Hz, 2H), 1.83 - 1.71 (m, 1H), 1.67 - 1.54 (m, 4H), 1.38 - 1.24 (m, 2H), 1.14 (br d, *J*=6.1 Hz, 6H), 1.01 (s, 9H), 0.91 (d, *J*=6.6 Hz, 6H). ESI [M+H] = 566.2.


25 **Example 79. Preparation of oxetan-3-yl *trans*-*N*-[4-[5-[2-(*tert*-butylsulfamoyl)-4-isopentyloxy-phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 102)**

Following the same protocol and under the same reaction conditions as for Compound 101, Compound 102 was prepared. ¹H NMR (400MHz, methanol-*d*₄) δ = 7.73 - 7.65 (m, 2H), 7.40 (d, J=8.4 Hz, 1H), 7.18 (dd, J=2.6, 8.5 Hz, 1H), 5.45 - 5.32 (m, 1H), 4.87 (br s, 2H), 4.66 - 4.59 (m, 2H), 4.14 (t, J=6.5 Hz, 2H), 3.52 - 3.40 (m, 1H), 3.03 (br t, J=12.0 Hz, 1H), 2.25 (br d, J=12.3 Hz, 2H), 2.09 (br d, J=11.6 Hz, 2H), 1.89 (quind, J=6.7, 13.4 Hz, 1H), 1.79 - 1.64 (m, 4H), 1.52 - 1.39 (m, 2H), 1.11 (s, 9H), 1.02 (d, J=6.6 Hz, 6H). ESI [M+H] = 580.2.

Example 80. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-(oxazol-2-ylmethyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 103)




- 10 a) Synthesis of *trans*-2-[2-(4-aminocyclohexyl)thiazol-5-yl]-N-ethyl-5-(oxazol-2-ylmethyl)benzenesulfonamide




- 15 A mixture of isopropyl *trans*-N-[4-[5-[4-[2-(2,2-dimethoxyethylamino)-2-oxo-ethyl]-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (50 mg, 83 μ mol, 1 *eq.*) in PPA (1 mL) was stirred at 70°C for 1 h then the mixture warmed to 120°C for 3 h. The reaction mixture was diluted with MeOH (1 mL). The residue was purified by prep-HPLC (column: Phenomenex Luna C18 100*30mm*5 μ m; mobile phase: [water(0.1%TFA)-ACN]; B%: 10%-35%, 12min) to yield *trans*-2-[2-(4-amino-cyclohexyl)thiazol-5-yl]-N-ethyl-5-(oxazol-2-ylmethyl)benzenesulfonamide (5 mg, 9 μ mol, 11% yield, TFA) as a white solid. ESI [M+H] = 447.2.

- 20 b) Synthesis of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-(oxazol-2-ylmethyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 103**)




To a solution of *trans*-2-[2-(4-aminocyclohexyl)thiazol-5-yl]-*N*-ethyl-5-(oxazol-2-ylmethyl)benzenesulfonamide (5 mg, 9 μ mol, 1 *eq.*, TFA) in dioxane (1 mL) was added sat. aq. Na_2CO_3 (0.3 mL) and isopropyl chloroformate (2 mg, 17 μ mol, 2 *eq.*). The mixture was stirred at 25°C for 2 h. The reaction mixture was concentrated under reduced pressure and purified by prep-HPLC (column: Welch Ultimate AQ-C18 150*30mm*5 μ m; mobile phase: [water(0.1%TFA)-ACN]; B%: 33%-63%, 12min) to yield isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-(oxazol-2-ylmethyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (720 μ g, 1 μ mol, 14 % yield, 95 % purity) as a yellow gum. ^1H NMR (400 MHz, methanol- d_4) δ = 7.89 (d, J = 1.3 Hz, 1H), 7.78 (s, 1H), 7.63 (s, 1H), 7.48 (dd, J = 1.4, 7.9 Hz, 1H), 7.38 (d, J = 7.9 Hz, 1H), 7.04 (s, 1H), 4.54 (br s, 1H), 4.21 (s, 2H), 3.40 - 3.34 (m, 1H), 2.97 - 2.87 (m, 1H), 2.75 (q, J = 7.3 Hz, 2H), 2.18 - 2.08 (m, 2H), 2.02 - 1.92 (m, 2H), 1.66 - 1.55 (m, 2H), 1.38 - 1.28 (m, 2H), 1.13 (br d, J = 6.1 Hz, 6H), 0.91 (t, J = 7.3 Hz, 3H). ESI $[\text{M}+\text{H}] = 533.0$.

Example 81. Preparation of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-oxazol-2-yl-phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 104)




a) Synthesis of isopropyl *trans*-*N*-[4-[5-[4-(2,2-dimethoxyethylcarbamoyl)-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate




From *trans*-3-(ethylsulfamoyl)-4-[2-[4-(isopropylloxycarbonyl-amino) cyclohexyl]thiazol-5-yl]benzoic acid (**Compound 48**), using General Method A. ¹H NMR (400MHz, methanol-*d*₄) δ = 8.54 (s, 1H), 8.06 (br d, *J*=7.9 Hz, 1H), 7.84 (s, 1H), 7.63 (br d, *J*=7.8 Hz, 1H), 5.00 - 4.96 (m, 1H), 4.61 (br t, *J*=5.5 Hz, 1H), 3.55 (br d, *J*=5.3 Hz, 2H), 3.49 (br d, *J*=11.0 Hz, 1H), 3.44 (s, 6H), 3.37 (br d, *J*=5.3 Hz, 1H), 2.95 - 2.89 (m, 2H), 2.35 - 2.23 (m, 2H), 2.10 (br d, *J*=10.8 Hz, 2H), 1.79 - 1.67 (m, 2H), 1.53 - 1.43 (m, 2H), 1.25 (br d, *J*=5.9 Hz, 6H), 1.10 - 0.98 (m, 3H). ESI [M+H] = 583.2.

b) Synthesis of *trans*-2-[2-(4-aminocyclohexyl)thiazol-5-yl]-N-ethyl-5-oxazol-2-yl-benzenesulfonamide



A solution of isopropyl *trans*-N-[4-[5-[4-(2,2-dimethoxyethylcarbamoyl)-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (50 mg, 85 μ mol, 1 *eq.*) in PPA (2 mL) was stirred at 70°C for 1 h then warmed to 120°C stirred for 2 h. The reaction was concentrated under reduced pressure. The residue was purified by prep-HPLC (column: Nano-micro Kromasil C18 80*25mm 3 μ m; mobile phase: [water(0.1%TFA)-ACN]; B%: 15%-45%, 7min) to yield *trans*-2-[2-(4-aminocyclohexyl)thiazol-5-yl]-N-ethyl-5-oxazol-2-yl-benzenesulfonamide (17 mg, 31 μ mol, 36 % yield, TFA) as brown oil. ESI [M+H] = 433.2.


c) Synthesis of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-oxazol-2-yl-phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 104**)



To a solution of *trans*-2-[2-(4-aminocyclohexyl)thiazol-5-yl]-N-ethyl-5-oxazol-2-yl-benzenesulfonamide (15 mg, 27 μ mol, 1 *eq.*, TFA) in dioxane (2 mL) was added sat. aq. Na₂CO₃ (0.3 mL) and isopropyl chloroformate (26 mg, 213 μ mol, 8 *eq.*). The mixture was stirred at 25°C for 1 h. The reaction was concentrated and purified by prep-HPLC (column: Nano-micro Kromasil C18 80*25mm 3 μ m; mobile phase: [water(0.1%TFA)-ACN]; B%: 45%-62%, 7min) to yield isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-oxazol-2-yl-

phenyl]thiazol-2-yl]cyclohexyl]carbamate (2 mg, 4 μ mol, 11% yield, 94% purity) as a yellow solid. ^1H NMR (400MHz, methanol- d_4) δ = 8.72 (d, J =1.5 Hz, 1H), 8.27 (dd, J =1.8, 8.0 Hz, 1H), 8.10 (s, 1H), 7.86 (s, 1H), 7.69 (d, J =8.0 Hz, 1H), 7.41 (s, 1H), 4.96 - 4.91 (m, 1H), 3.53 - 3.43 (m, 1H), 3.10 - 3.02 (m, 1H), 2.95 (q, J =7.3 Hz, 2H), 2.28 (br d, J =12.1 Hz, 2H), 2.14 - 2.04 (m, 2H), 1.80 - 1.67 (m, 2H), 1.44 (dq, J =2.9, 12.6 Hz, 2H), 1.25 (br d, J =6.1 Hz, 6H), 1.07 (t, J =7.2 Hz, 3H). ESI $[\text{M}+\text{H}] = 519.0$.


Example 82. Preparation of compound isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[(1-methylimidazol-2-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 105) and isopropyl *cis*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[(1-methylimidazol-2-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 106)




From isopropyl *trans*-*N*-[4-[5-[4-amino-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate and 2-bromo-1-methyl-1*H*-imidazole, using General Method K. **(Compound 105)**, ^1H NMR (400 MHz, methanol- d_4) δ = 7.74 (d, J = 2.4 Hz, 1H), 7.63 - 7.54 (m, 1H), 7.40 - 7.23 (m, 2H), 7.01 (d, J = 2.0 Hz, 1H), 6.92 (d, J = 1.8 Hz, 1H), 4.72 (br d, J = 6.2 Hz, 1H), 3.61 - 3.51 (m, 3H), 3.42 - 3.30 (m, 1H), 2.96 - 2.87 (m, 1H), 2.77 (q, J = 7.2 Hz, 2H), 2.18 - 2.09 (m, 2H), 1.97 (br d, J = 9.9 Hz, 2H), 1.59 (dq, J = 2.9, 12.9 Hz, 2H), 1.37 - 1.24 (m, 2H), 1.13 (br d, J = 6.1 Hz, 6H), 0.91 (t, J = 7.2 Hz, 3H). ESI $[\text{M}+\text{H}] = 547.2$.

(Compound 106), ^1H NMR (400 MHz, methanol- d_4) δ = 7.73 (d, J = 2.3 Hz, 1H), 7.69 (s, 1H), 7.37 - 7.31 (m, 1H), 7.25 (dd, J = 2.4, 8.4 Hz, 1H), 6.97 (d, J = 1.2 Hz, 1H), 6.86 (d, J = 1.3 Hz, 1H), 4.84 (br s, 1H), 3.77 (br d, J = 4.9 Hz, 1H), 3.60 (s, 3H), 3.17 (br dd, J = 4.2, 8.6 Hz, 1H), 2.91 (q, J = 7.2 Hz, 2H), 2.09 - 1.94 (m, 4H), 1.91 - 1.71 (m, 4H), 1.25 (d, J = 6.2 Hz, 6H), 1.05 (t, J = 7.2 Hz, 3H). ESI $[\text{M}+\text{H}] = 547.2$.

Example 83. Preparation of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[(1-isobutylimidazol-2-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 107)




a) Synthesis of 2-bromo-1-isobutyl-1H-imidazole



To a mixture of 2-bromo-1H-imidazole (300 mg, 2 mmol, 1 *eq.*), NaH (245 mg, 6 mmol, 60% purity, 3 *eq.*) in THF (3 mL) was added 1-bromo-2-methylpropane (838.55 mg, 6.12 mmol, 665.52 uL, 3 *eq.*) and stirred at 25°C for 6 h under N₂ atmosphere. The reaction mixture was quenched by addition H₂O 10 mL, extracted with EtOAc 60 mL (20mL X 3). The combined organic layers were washed with sat.aq. NaCl 60 mL (12 mL X 5), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by prep-HPLC (Nano-micro Kromasil C18 100*30mm 5um; mobile phase: [water(0.1%TFA)-ACN];B%:3%-35%,7min) to yield 2-bromo-1-isobutyl-1H-imidazole (400 mg, 2 mmol, 97% yield) as yellow oil. ¹H NMR (400 MHz, CDCl₃) δ = 7.37 (d, J=1.76 Hz, 1H), 7.10 (d, J=1.76 Hz, 1H), 3.84 (d, J=7.50 Hz, 2H), 2.05-2.27 (m, 1H), 0.98 (d, J=6.84 Hz, 6H). ESI [M+H] = 203.0/205.0.


b) Synthesis of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(1-isobutyl-imidazol-2-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 107**)




From isopropyl *trans*-N-[4-[5-[4-amino-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate and 2-bromo-1-isobutyl-1H-imidazole, using General Method K. ¹H NMR (400 MHz, methanol-*d*₄) δ = 7.63-7.70 (m, 2H), 7.31 (d, J=8.38 Hz, 1H), 7.18 (dd, J=2.43, 8.38 Hz, 1H), 7.00 (d, J=1.54 Hz, 1H), 6.88 (d, J=1.54 Hz, 1H), 4.83 (br d, J=6.39 Hz, 1H), 3.69-3.79 (m, 2H), 3.45 (br t, J=11.80 Hz, 1H), 2.93-3.05 (m, 1H), 2.86 (q, J=7.28

Hz, 2H), 2.23 (br d, $J=12.79$ Hz, 2H), 2.01-2.12 (m, 3H), 1.63-1.75 (m, 2H), 1.35-1.47 (m, 2H), 1.23 (br d, $J=6.17$ Hz, 6H), 1.01 (t, $J=7.28$ Hz, 3H), 0.90 (d, $J=6.62$ Hz, 6H). ESI $[M+H]^+ = 589.3$.

Example 84. Preparation of cyclopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-(1*H*-imidazol-2-ylamino)phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 109)




a) Synthesis of cyclopropyl *trans*-*N*-[4-(5-bromothiazol-2-yl)cyclohexyl] carbamate




To a solution of cyclopropanol (293 mg, 5 mmol, 5 eq) in DCE (3 mL) was added DIEA (651 mg, 5 mmol, 5 eq) at 0°C. The mixture was stirred at 0°C for 10 min, followed by the addition of triphosgene (598 mg, 2 mmol, 2 eq) at 0°C. The mixture was stirred at 25°C for 30 min (mixture 1). To a solution of *trans*-4-(5-bromothiazol-2-yl)cyclohexanamine (300 mg, 1 mmol, 1 eq, HCl) in DIEA (3 mL) was added mixture 1 and stirred at 25°C for 2 h. The reaction mixture was quenched by addition sat. aq. Na_2CO_3 5 mL, and extracted with dichloromethane 30 mL (10 mL X 3). The combined organic layers were dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified prep-TLC (SiO_2 , petroleum ether: ethyl acetate=2:1) to yield cyclopropyl *trans*-*N*-[4-(5-bromothiazol-2-yl)cyclohexyl]carbamate (140 mg, 406 μmol , 40% yield) as a yellow solid. ^1H NMR (400MHz, methanol- d_4) δ = 7.60 (s, 1H), 3.95 (br d, $J=3.3$ Hz, 1H), 3.43 (br s, 1H), 3.04 - 2.81 (m, 1H), 2.22 - 1.89 (m, 4H), 1.72 - 1.49 (m, 2H), 1.46 - 1.11 (m, 4H), 0.65 - 0.61 (m, 2H). ESI $[M+H]^+ = 345.1/347.0$.

b) Synthesis of cyclopropyl *trans*-*N*-[4-[5-[4-amino-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 108)




From cyclopropyl *trans*-*N*-[4-(5-bromothiazol-2-yl)cyclohexyl]carbamate and 5-amino-*N*-ethyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzenesulfonamide, using General Method C. ¹H NMR (400MHz, methanol-*d*₄) δ = 7.74 (s, 1H), 7.40 (d, *J*=2.4 Hz, 1H), 7.21 (d, *J*=8.2 Hz, 1H), 6.91 (dd, *J*=2.4, 8.4 Hz, 1H), 3.96 (br d, *J*=3.3 Hz, 1H), 3.47 (br t, *J*=11.5 Hz, 1H), 3.06 (br t, *J*=12.2 Hz, 1H), 2.87 (q, *J*=7.3 Hz, 2H), 2.23 (br d, *J*=12.8 Hz, 2H), 2.13 - 1.95 (m, 2H), 1.83 - 1.59 (m, 2H), 1.41 (dq, *J*=3.2, 12.6 Hz, 2H), 1.02 (t, *J*=7.2 Hz, 3H), 0.63 (br d, *J*=5.1 Hz, 4H). ESI [M+H] = 465.1.

c) Synthesis of cyclopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-(1*H*-imidazol-2-ylamino)phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 109**)



From cyclopropyl *trans*-*N*-[4-[5-[4-amino-2-(ethylsulfamoyl) phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 108**) and 2-bromo-1*H*-imidazole, using General Method K. ¹H NMR (400MHz, methanol-*d*₄) δ = 7.90 (d, *J*=1.5 Hz, 1H), 7.72 (s, 1H), 7.53 - 7.40 (m, 2H), 7.05 (s, 2H), 3.97 (br d, *J*=3.5 Hz, 1H), 3.47 (br s, 1H), 3.01 (br t, *J*=11.9 Hz, 1H), 2.87 (q, *J*=7.3 Hz, 2H), 2.25 (br d, *J*=12.8 Hz, 2H), 2.08 (br d, *J*=10.1 Hz, 2H), 1.85 - 1.63 (m, 2H), 1.50 - 1.33 (m, 2H), 1.01 (t, *J*=7.2 Hz, 3H), 0.64 (br d, *J*=4.6 Hz, 4H). ESI [M+H] = 531.1.

Example 85. Preparation of isopropyl *trans*-*N*-[4-[5-[2-(*tert*-butylsulfamoyl)-4-(1*H*-imidazol-2-ylamino)phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 38) and isopropyl *cis*-*N*-[4-[5-[2-(*tert*-butylsulfamoyl)-4-(1*H*-imidazol-2-ylamino)phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 110)




From isopropyl *trans*-*N*-[4-[5-[4-amino-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate and 2-bromo-1*H*-imidazole, using General Method K.


(**Compound 38**), ¹H NMR (400MHz, methanol-*d*₄) δ = 7.95 (t, *J* = 1.3 Hz, 1H), 7.73 (s, 1H), 7.47 (d, *J* = 1.3 Hz, 2H), 7.04 (s, 2H), 4.85 - 4.79 (m, 1H), 3.55 - 3.37 (m, 1H), 3.01 (tt, *J* = 3.5, 12.0 Hz, 1H), 2.32 - 2.16 (m, 2H), 2.13 - 1.99 (m, 2H), 1.69 (dq, *J* = 3.1, 12.9 Hz, 2H), 1.41 (dq, *J* = 3.2, 12.5 Hz, 2H), 1.22 (br d, *J* = 6.2 Hz, 6H), 1.09 (s, 9H). ESI [M+H] = 561.2.

(**Compound 110**), ¹H NMR (400MHz, methanol-*d*₄) δ = 7.96 (d, *J* = 2.2 Hz, 1H), 7.76 (s, 1H), 7.59 - 7.42 (m, 2H), 7.08 (s, 2H), 4.83 - 4.75 (m, 1H), 3.82 - 3.66 (m, 1H), 3.23 - 3.08 (m, 1H), 2.08 - 1.92 (m, 4H), 1.89 - 1.68 (m, 4H), 1.23 (d, *J* = 6.2 Hz, 6H), 1.10 (s, 9H). ESI [M+H] = 561.2.

Example 86. Preparation of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[(4-isopropenyl-2-pyridyl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 111)




15 a) Synthesis of 4-isopropenylpyridin-2-amine




A mixture of 2-isopropenyl-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (2.53 g, 15 mmol, 1.3 *eq.*), 4-bromopyridin-2-amine (2 g, 12 mmol, 1.0 *eq.*), Cs₂CO₃ (11.3 g, 35 mmol, 3.0 *eq.*) and Pd(dppf)Cl₂ (846 mg, 1 mmol, 0.1 *eq.*) in H₂O (50 mL) and dioxane (100 mL) was stirred at 90°C for 4 h under N₂ atmosphere. The reaction was concentrated, and then purified by column chromatography (SiO₂, petroleum ether: ethyl acetate=10:1 to 1:1) to yield 4-isopropenylpyridin-2-amine (1.5 g, 11 mmol, 97% yield) as a brown solid. ESI [M+H]⁺ = 135.1.

b) Synthesis of 4-isopropylpyridin-2-amine



A mixture of 4-isopropenylpyridin-2-amine (1.5 g, 11 mmol, 1 *eq.*), Pd/C (500 mg, 10% purity) in EtOAc (300 mL) was degassed and purged with H₂ (30 psi) for 3 times, and then the mixture was stirred at 25°C for 12 h under H₂ atmosphere. The reaction mixture was filtered, and the filtrate was concentrated. The residue was purified by prep-HPLC (Neutral condition: column: Waters Xbridge Prep OBD C18 150*40mm*10um; mobile phase: [water(10mM NH₄HCO₃)-ACN]; B%: 15%-45%, 8min) to yield 4-isopropylpyridin-2-amine (800 mg, 6 mmol, 53% yield) as yellow oil. ¹H NMR (400MHz, methanol-*d*₄) δ = 7.78 (d, J=5.5 Hz, 1H), 6.56 - 6.44 (m, 2H), 2.82 - 2.71 (m, 1H), 1.23 (d, J=6.9 Hz, 6H). ESI [M+H]⁺ = 137.0.

c) Synthesis of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(4-isopropyl-2-pyridyl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 111**)



From isopropyl *trans*-N-[4-[5-[4-bromo-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate and 4-isopropylpyridin-2-amine, using General Method F. ¹H NMR (400MHz, methanol-*d*₄) δ = 8.16 (d, J=2.3 Hz, 1H), 7.97 (d, J=6.9 Hz, 1H), 7.79 (s, 1H), 7.72 (dd, J=2.2, 8.3 Hz, 1H), 7.60 (d, J=8.1 Hz, 1H), 7.11 - 7.08 (m, 2H), 4.85 - 4.85 (m, 1H), 3.48

(br t, J=11.5 Hz, 1H), 3.10 - 2.98 (m, 2H), 2.91 (q, J=7.2 Hz, 2H), 2.28 (br d, J=12.6 Hz, 2H), 2.10 (br d, J=10.8 Hz, 2H), 1.80 - 1.67 (m, 2H), 1.51 - 1.38 (m, 2H), 1.33 (d, J=6.9 Hz, 6H), 1.25 (br d, J=6.1 Hz, 6H), 1.05 (t, J=7.3 Hz, 3H). ESI [M+H] = 586.3.

Example 87. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(6-isopropyl-2-pyridyl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 112)

Following the same protocol and under the same reaction conditions as for Compound 111, Compound 112 was prepared. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.46 (d, J = 2.0 Hz, 1H), 7.85 - 7.77 (m, 2H), 7.76 (s, 1H), 7.48 (d, J = 8.3 Hz, 1H), 6.91 (d, J = 7.9 Hz, 2H), 4.82 (br s, 1H), 3.54 - 3.43 (m, 1H), 3.13 - 3.00 (m, 2H), 2.91 (q, J = 7.2 Hz, 2H), 2.32 - 2.23 (m, 2H), 2.10 (br dd, J = 2.9, 12.8 Hz, 2H), 1.80 - 1.67 (m, 2H), 1.50 - 1.41 (m, 2H), 1.38 (d, J = 7.0 Hz, 6H), 1.25 (br d, J = 6.1 Hz, 6H), 1.04 (t, J = 7.2 Hz, 3H). ESI [M+H] = 586.1.

Example 88. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(5-isopropyl-2-pyridyl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 113)

Following the same protocol and under the same reaction conditions as for Compound 111, Compound 113 was prepared. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.05 - 7.99 (m, 1H), 7.96 - 7.89 (m, 1H), 7.79 (s, 1H), 7.67 (s, 1H), 7.62 - 7.55 (m, 1H), 7.47 (d, J = 8.2 Hz, 1H), 7.21 - 7.07 (m, 1H), 4.74 - 4.70 (m, 1H), 3.36 (tt, J = 3.9, 11.6 Hz, 1H), 2.98 - 2.84 (m, 2H), 2.80 (q, J = 7.2 Hz, 2H), 2.16 (br d, J = 12.1 Hz, 2H), 1.98 (br d, J = 10.3 Hz, 2H), 1.61 (dq, J = 2.3, 12.7 Hz, 2H), 1.38 - 1.25 (m, 2H), 1.20 (d, J = 7.0 Hz, 6H), 1.13 (br d, J = 6.2 Hz, 6H), 0.93 (t, J = 7.2 Hz, 3H). ESI [M+H] = 586.1.

Example 89. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[3-isopropoxyazetidin-1-yl]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 114)

Following the same protocol and under the same reaction conditions as for Compound 111, Compound 114 was prepared. ¹H NMR (400 MHz, methanol-*d*₄) δ = 7.62 (s, 1H), 7.28 (d, J=8.38 Hz, 1H), 7.08 (d, J=2.38 Hz, 1H), 6.68 (dd, J=8.32, 2.44 Hz, 1H), 4.72 - 4.80 (m, 1H), 4.57 - 4.63 (m, 1H), 4.25 (t, J=7.25 Hz, 2H), 3.71 - 3.79 (m, 3H), 3.40 - 3.50 (m, 1H), 2.93 - 3.05 (m, 1H), 2.87 (q, J=7.21 Hz, 2H), 2.19 - 2.30 (m, 2H), 2.02 - 2.14 (m, 2H), 1.62 - 1.78 (m, 2H), 1.37 - 1.48 (m, 2H), 1.24 (br d, J=6.13 Hz, 6H), 1.20 (d, J=6.13 Hz, 6H), 1.03 (t, J=7.25 Hz, 3H). ESI [M+H] = 565.3.

Example 90. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(5-isopropyl-1,3,4-thiadiazol-2-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 115)

Following the same protocol and under the same reaction conditions as for Compound 111, Compound 115 was prepared. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.46 (d, J=2.25 Hz,

1H), 7.85 (dd, J=8.38, 2.38 Hz, 1H), 7.77 (br s, 1H), 7.47 (d, J=8.38 Hz, 1H), 4.84 (br s, 1H), 3.42 - 3.53 (m, 1H), 3.35 - 3.41 (m, 1H), 3.04 (br s, 1H), 2.97 (q, J=7.25 Hz, 2H), 2.26 (br d, J=12.63 Hz, 2H), 2.09 (br d, J=10.51 Hz, 2H), 1.72 (q, J=11.30 Hz, 2H), 1.43 (d, J=6.88 Hz, 8H), 1.25 (br d, J=6.00 Hz, 6H), 1.09 (t, J=7.25 Hz, 3H). ESI [M+H] = 593.0..

5 **Example 91. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(3-methyloxetan-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 116)**

Following the same protocol and under the same reaction conditions as for Compound 111, Compound 116 was prepared. ¹H NMR (400MHz, DMSO-*d*₆) δ = 7.56 (s, 1H), 7.31 (br s, 1H), 7.23 - 7.15 (m, 1H), 7.02 (br d, J=7.7 Hz, 1H), 6.98 - 6.90 (m, 2H), 6.52 (dd, J=2.2, 8.3 Hz, 1H), 4.80 - 4.70 (m, 1H), 4.63 (br d, J=5.9 Hz, 2H), 4.51 (br d, J=5.7 Hz, 2H), 3.31- 3.25 (m, 1H), 2.93 - 2.82 (m, 1H), 2.81 - 2.77 (m, 2H), 2.18 - 2.07 (m, 2H), 1.91 (br d, J=10.6 Hz, 2H), 1.58 (s, 3H), 1.57 - 1.47 (m, 2H), 1.40 - 1.26 (m, 2H), 1.17 (br d, J=6.1 Hz, 6H), 0.98 (t, J=7.2 Hz, 3H). ESI [M+H] = 537.2.

15 **Example 92. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[methyl(1-methyl-1H-imidazol-2-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 117)**

Following the same protocol and under the same reaction conditions as for Compound 111, Compound 117 was prepared. ¹H NMR (400 MHz, methanol-*d*₄) δ = 7.78 - 7.67 (m, 1H), 7.61 - 7.48 (m, 4H), 7.28 - 7.13 (m, 1H), 4.86 - 4.79 (m, 1H), 3.61 (s, 3H), 3.58 - 3.55 (m, 3H), 3.52 - 3.40 (m, 1H), 3.08 - 3.00 (m, 1H), 2.85 (q, J = 7.2 Hz, 2H), 2.30 - 2.22 (m, 2H), 2.14 - 2.06 (m, 2H), 1.78 - 1.65 (m, 2H), 1.43 (dq, J = 3.2, 12.7 Hz, 2H), 1.25 (br d, J = 6.1 Hz, 6H), 1.04 - 0.99 (m, 3H). ESI [M+H] = 561.3.

25 **Example 93. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(1,3,4-oxadiazol-2-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 118)**

Following the same protocol and under the same reaction conditions as for Compound 111, Compound 118 was prepared. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.49 (d, J=2.38 Hz, 1H), 8.31 (s, 1H), 7.93 (dd, J=8.38, 2.38 Hz, 1H), 7.82 (s, 1H), 7.66 (d, J=8.25 Hz, 1H), 4.79 - 4.84 (m, 1H), 3.42 - 3.53 (m, 1H), 3.06 (tt, J=12.04, 3.35 Hz, 1H), 2.96 (q, J=7.21 Hz, 2H), 2.20 - 2.32 (m, 2H), 2.04 - 2.14 (m, 2H), 1.65 - 1.80 (m, 2H), 1.34 - 1.51 (m, 2H), 1.25 (br d, J=6.13 Hz, 6H), 1.02 - 1.12 (m, 3H). ESI [M+H] = 535.2.

30 **Example 94. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(1,3,4-thiadiazol-2-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 119)**

Following the same protocol and under the same reaction conditions as for Compound 111, Compound 119 was prepared. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.90 (s, 1H), 8.51 (d, J = 2.5 Hz, 1H), 7.91 (dd, J = 2.4, 8.4 Hz, 1H), 7.79 (s, 1H), 7.50 (d, J = 8.4 Hz, 1H), 4.84 (br s, 1H), 3.48 (tt, J = 3.8, 11.6 Hz, 1H), 3.12 - 3.03 (m, 1H), 2.99 (q, J = 7.2 Hz, 2H), 2.32 - 2.22 (m, 2H), 2.14 - 2.05 (m, 2H), 1.73 (dq, J = 3.0, 12.9 Hz, 2H), 1.44 (dq, J = 3.1, 12.6 Hz, 2H), 1.25 (br d, J = 6.1 Hz, 6H), 1.10 (t, J = 7.2 Hz, 3H). ESI [M+H] = 551.2.

Example 95. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(1H-pyrazol-4-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 120)

Following the same protocol and under the same reaction conditions as for Compound 111, Compound 120 was prepared. ¹H NMR (400 MHz, methanol-*d*₄) δ = 7.68 (s, 1H), 7.61 (s, 2H), 7.45 (d, J = 2.4 Hz, 1H), 7.24 (d, J = 8.4 Hz, 1H), 6.97 (dd, J = 2.4, 8.4 Hz, 1H), 4.84 - 4.75 (m, 1H), 3.51 - 3.39 (m, 1H), 3.01 (tt, J = 3.4, 12.1 Hz, 1H), 2.86 (q, J = 7.2 Hz, 2H), 2.27 - 2.17 (m, 2H), 2.11 - 2.03 (m, 2H), 1.68 (dq, J = 3.0, 12.8 Hz, 2H), 1.40 (dq, J = 3.3, 12.6 Hz, 2H), 1.22 (br d, J = 6.2 Hz, 6H), 1.01 (t, J = 7.2 Hz, 3H). ESI [M+H] = 533.2.

Example 96. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(1H-pyrazol-5-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 121)

Following the same protocol and under the same reaction conditions as for Compound 111, Compound 121 was prepared. ¹H NMR (400 MHz, methanol-*d*₄) δ = 7.97 (d, J = 2.4 Hz, 1H), 7.67 (s, 1H), 7.57 (d, J = 2.4 Hz, 1H), 7.41 (dd, J = 2.4, 8.4 Hz, 1H), 7.28 (d, J = 8.4 Hz, 1H), 6.01 (d, J = 2.4 Hz, 1H), 4.82 - 4.74 (m, 1H), 3.45 (tt, J = 3.8, 11.5 Hz, 1H), 3.05 - 2.95 (m, 1H), 2.91 (q, J = 7.1 Hz, 2H), 2.27 - 2.18 (m, 2H), 2.10 - 2.01 (m, 2H), 1.68 (dq, J = 3.0, 12.8 Hz, 2H), 1.40 (dq, J = 3.1, 12.6 Hz, 2H), 1.22 (br d, J = 6.2 Hz, 6H), 1.04 (t, J = 7.3 Hz, 3H). ESI [M+H] = 533.2.

Example 97. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(pyrazin-2-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 122)

Following the same protocol and under the same reaction conditions as for Compound 111, Compound 122 was prepared. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.59 (d, J = 2.2 Hz, 1H), 8.26 - 8.16 (m, 2H), 7.99 - 7.90 (m, 2H), 7.72 (s, 1H), 7.41 (d, J = 8.4 Hz, 1H), 4.82 (br s, 1H), 3.52 - 3.40 (m, 1H), 3.06 - 2.98 (m, 1H), 2.95 (q, J = 7.1 Hz, 2H), 2.29 - 2.18 (m, 2H), 2.11 - 2.01 (m, 2H), 1.70 (dq, J = 3.2, 12.8 Hz, 2H), 1.41 (dq, J = 3.4, 12.5 Hz, 2H), 1.22 (br d, J = 6.2 Hz, 6H), 1.07 (t, J = 7.3 Hz, 3H). ESI [M+H] = 545.2.

Example 98. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(pyridazin-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 123)

Following the same protocol and under the same reaction conditions as for Compound 111, Compound 123 was prepared. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.95 (d, J=4.25 Hz, 1H) 8.56 (d, J=2.38 Hz, 1H) 7.92 - 8.04 (m, 2H) 7.72 - 7.82 (m, 2H) 7.55 (d, J=8.38 Hz, 1H) 4.76 - 4.89 (m, 1H) 3.40 - 3.54 (m, 1H) 3.06 (tt, J=12.10, 3.35 Hz, 1H) 2.92 (q, J=7.25 Hz, 2H) 2.22 - 2.35 (m, 2H) 2.00 - 2.13 (m, 2H) 1.73 (qd, J=12.78, 2.94 Hz, 2H) 1.44 (qd, J=12.57, 3.06 Hz, 2H) 1.25 (br d, J=6.13 Hz, 6H) 1.05 (t, J=7.25 Hz, 3H). ESI [M+H] = 545.2.

Example 99. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(pyrimidin-4-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 124)

Following the same protocol and under the same reaction conditions as for Compound 111, Compound 124 was prepared. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.96 (s, 1H), 8.55 (d, J=1.88 Hz, 1H) 8.40 (dd, J=7.13, 1.13 Hz, 1H), 8.02 (dd, J=8.32, 2.19 Hz, 1H), 7.81 (s, 1H) 7.60 (d, J=8.38 Hz, 1H), 7.15 (d, J=7.13 Hz, 1H), 4.76 - 4.90 (m, 1H), 3.42 - 3.54 (m, 1H), 3.01 - 3.12 (m, 1H), 2.92 (q, J=7.25 Hz, 2H), 2.21 - 2.32 (m, 2H), 2.04 - 2.14 (m, 2H), 1.73 (qd, J=12.84, 2.88 Hz, 2H), 1.44 (qd, J=12.57, 3.06 Hz, 2H), 1.25 (br d, J=6.13 Hz, 6H), 1.05 (t, J=7.25 Hz, 3H). ESI [M+H] = 545.2.

Example 100. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(pyridin-4-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 125)

Following the same protocol and under the same reaction conditions as for Compound 111, Compound 125 was prepared. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.28 (d, J = 7.3 Hz, 2H), 8.02 (d, J = 2.0 Hz, 1H), 7.79 (s, 1H), 7.69 - 7.57 (m, 2H), 7.27 (d, J = 7.3 Hz, 2H), 4.85 - 4.77 (m, 1H), 3.45 (tt, J = 3.8, 11.5 Hz, 1H), 3.03 (tt, J = 3.3, 12.0 Hz, 1H), 2.88 (q, J = 7.1 Hz, 2H), 2.29 - 2.20 (m, 2H), 2.11 - 2.01 (m, 2H), 1.70 (dq, J = 3.0, 12.8 Hz, 2H), 1.41 (dq, J = 3.1, 12.6 Hz, 2H), 1.22 (br d, J = 6.2 Hz, 6H), 1.01 (t, J = 7.3 Hz, 3H). ESI [M+H] = 544.2.

Example 101. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(pyridin-2-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 126)

Following the same protocol and under the same reaction conditions as for Compound 111, Compound 126 was prepared. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.10 (d, J = 2.3 Hz, 1H), 7.98 (dd, J = 1.1, 5.9 Hz, 1H), 7.89 (ddd, J = 1.8, 7.2, 8.8 Hz, 1H), 7.70 - 7.60 (m, 2H), 7.46 (d, J = 8.3 Hz, 1H), 7.11 (d, J = 8.8 Hz, 1H), 7.02 - 6.93 (m, 1H), 4.74 - 4.67 (m, 1H), 3.36 (tt, J = 3.9, 11.6 Hz, 1H), 2.94 (tt, J = 3.5, 12.1 Hz, 1H), 2.81 (q, J = 7.2 Hz, 2H), 2.20 - 2.11 (m, 2H), 2.02 - 1.94 (m, 2H), 1.61 (dq, J = 3.1, 12.9 Hz, 2H), 1.38 - 1.26 (m, 2H), 1.13 (d, J = 6.1 Hz, 6H), 0.94 (t, J = 7.2 Hz, 3H). ESI [M+H] = 544.2.

Example 102. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(pyrimidin-2-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 127)

Following the same protocol and under the same reaction conditions as for Compound 111, Compound 127 was prepared. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.66 (d, J = 2.3 Hz, 1H), 8.53 (d, J = 4.8 Hz, 2H), 7.97 (dd, J = 2.3, 8.4 Hz, 1H), 7.79 (s, 1H), 7.43 (d, J = 8.4 Hz, 1H), 6.91 (t, J = 4.9 Hz, 1H), 4.87 - 4.80 (m, 1H), 3.48 (tt, J = 3.8, 11.5 Hz, 1H), 3.11 - 3.02 (m, 1H), 2.99 (q, J = 7.3 Hz, 2H), 2.31 - 2.21 (m, 2H), 2.14 - 2.05 (m, 2H), 1.72 (dq, J = 2.9, 12.8 Hz, 2H), 1.43 (dq, J = 3.2, 12.6 Hz, 2H), 1.25 (br d, J = 6.1 Hz, 6H), 1.10 (t, J = 7.3 Hz, 3H). ESI [M+H] = 545.2.

Example 103. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(pyridin-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 128)

Following the same protocol and under the same reaction conditions as for Compound 111, Compound 128 was prepared. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.54 (d, J=2.63 Hz, 1H), 8.30 (d, J=5.50 Hz, 1H), 8.22 - 8.27 (m, 1H), 7.90 - 7.96 (m, 2H), 7.78 (s, 1H), 7.51 - 7.59 (m, 2H), 4.82 - 4.88 (m, 1H), 3.42 - 3.52 (m, 1H), 3.06 (tt, J=12.04, 3.53 Hz, 1H), 2.90 (q, J=7.17 Hz, 2H), 2.20 - 2.34 (m, 2H), 2.03 - 2.17 (m, 2H), 1.74 - 1.75 (m, 2H), 1.38 - 1.51 (m, 2H), 1.25 (d, J=6.13 Hz, 6H), 1.04 (t, J=7.25 Hz, 3H). ESI [M+H] = 544.2.

Example 104. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(1-

isopropyl-1H-pyrazol-5-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 129)

Following the same protocol and under the same reaction conditions as for Compound 111, Compound 129 was prepared. ¹H NMR (400 MHz, methanol-*d*₄) δ = 7.74 (s, 1H), 7.61 (d, J=1.75 Hz, 1H), 7.49 (d, J=2.50 Hz, 1H), 7.34 (d, J=8.38 Hz, 1H), 7.04 (dd, J=8.32, 2.44 Hz, 1H), 6.18 (d, J=2.00 Hz, 1H), 4.77 - 4.87 (m, 1H), 4.62 (dt, J=13.35, 6.64 Hz, 1H), 3.40 - 3.53 (m, 1H), 3.06 (tt, J=12.07, 3.38 Hz, 1H), 2.87 (q, J=7.25 Hz, 2H), 2.19 - 2.31 (m, 2H), 2.02 - 2.17 (m, 2H), 1.72 (qd, J=12.82, 2.81 Hz, 2H), 1.46 (d, J=6.63 Hz, 8H), 1.24 (br d, J=6.13 Hz, 6H), 1.02 (t, J=7.25 Hz, 3H). ESI [M+H] = 575.2.

Example 105. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(4-

hydroxypiperidin-1-yl]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 130)

Following the same protocol and under the same reaction conditions as for Compound 111, Compound 130 was prepared. ¹H NMR (400 MHz, methanol-*d*₄) δ = 7.79 (s, 1H), 7.61

(d, J = 2.4 Hz, 1H), 7.35 (d, J = 8.6 Hz, 1H), 7.22 (dd, J = 2.4, 8.6 Hz, 1H), 4.81 (br s, 1H), 3.85 (tt, J = 4.0, 8.5 Hz, 1H), 3.80 - 3.65 (m, 2H), 3.45 (tdd, J = 3.8, 7.5, 11.2 Hz, 1H), 3.20 - 3.00 (m, 3H), 2.87 (q, J = 7.2 Hz, 2H), 2.24 (br d, J = 12.6 Hz, 2H), 2.08 (br d, J = 10.4 Hz, 2H), 1.99 (br dd, J = 3.6, 9.4 Hz, 2H), 1.79 - 1.56 (m, 4H), 1.47 - 1.34 (m, 2H), 1.22 (br d, J = 6.2 Hz, 6H), 1.02 (t, J = 7.2 Hz, 3H). ESI [M+H] = 551.2.

Example 106. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(2-hydroxyethyl) (methyl))amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 131)

Following the same protocol and under the same reaction conditions as for Compound 111, Compound 131 was prepared. ¹H NMR (400 MHz, methanol-*d*₄) δ = 7.68 (s, 1H), 7.38 (d, J = 2.9 Hz, 1H), 7.27 (d, J = 8.6 Hz, 1H), 6.95 (dd, J = 2.9, 8.6 Hz, 1H), 4.82 - 4.78 (m, 1H), 3.80 - 3.70 (m, 2H), 3.67 - 3.54 (m, 2H), 3.51 - 3.40 (m, 1H), 3.09 (s, 3H), 3.02 (tt, J = 3.4, 12.0 Hz, 1H), 2.87 (q, J = 7.1 Hz, 2H), 2.31 - 2.15 (m, 2H), 2.10 - 2.00 (m, 2H), 1.69 (dq, J = 2.9, 12.9 Hz, 2H), 1.41 (dq, J = 3.2, 12.6 Hz, 2H), 1.22 (br d, J = 6.2 Hz, 6H), 1.02 (t, J = 7.3 Hz, 3H). ESI [M+H] = 525.2.

Example 107. Preparation of isopropyl (*trans*-4-(5-(2-(N-ethylsulfamoyl)-4-(oxazol-2-ylamino)phenyl)thiazol-2-yl)cyclohexyl]carbamate (Compound 60)

Following the same protocol and under the same reaction conditions as for Compound 111, Compound 60 was prepared. ¹H NMR (400MHz, methanol-*d*₄) δ = 8.37 (d, J=2.3 Hz, 1H), 7.89 - 7.73 (m, 2H), 7.54 (d, J=0.9 Hz, 1H), 7.46 (d, J=8.4 Hz, 1H), 7.03 (s, 1H), 4.85 (br s, 1H), 3.54 - 3.45 (m, 1H), 3.15 - 3.02 (m, 1H), 2.96 (q, J=7.3 Hz, 2H), 2.27 (br d, J=12.1 Hz, 2H), 2.10 (br d, J=10.6 Hz, 2H), 1.81 - 1.67 (m, 2H), 1.49 - 1.36 (m, 2H), 1.25 (br d, J=6.1 Hz, 6H), 1.09 (t, J=7.2 Hz, 3H). ESI [M+H] = 534.2.

Example 108. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(2-methoxyethyl) (methyl))amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 132)

Following the same protocol and under the same reaction conditions as for Compound 111, Compound 132 was prepared. ¹H NMR (400MHz, methanol-*d*₄) δ = 7.68 (s, 1H), 7.40 (d, J=2.6 Hz, 1H), 7.29 (d, J=8.6 Hz, 1H), 6.97 (dd, J=2.7, 8.6 Hz, 1H), 3.71 - 3.57 (m, 4H), 3.47 (br t, J=11.6 Hz, 1H), 3.37 (s, 3H), 3.09 (s, 3H), 3.06 - 2.97 (m, 1H), 2.94 - 2.85 (m, 2H), 2.25 (br d, J=12.6 Hz, 2H), 2.14 - 2.03 (m, 2H), 1.78 - 1.64 (m, 2H), 1.49 - 1.35 (m, 2H), 1.25 (br d, J=6.0 Hz, 6H), 1.08 - 1.01 (m, 3H). ESI [M+H] = 539.2.

Example 109. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[1,1-dioxidothiomorpholino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 133)

Following the same protocol and under the same reaction conditions as for Compound 111, Compound 133 was prepared. ¹H NMR (400MHz, methanol-*d*₄) δ = 7.73 (s, 1H), 7.67 - 7.63 (m, 1H), 7.41 (d, J=8.4 Hz, 1H), 7.29 (dd, J=2.8, 8.6 Hz, 1H), 4.87 - 4.81 (m, 1H), 4.02 (br s, 4H), 3.53 - 3.42 (m, 1H), 3.20 (br s, 4H), 3.05 (ddd, J=3.5, 8.7, 12.1 Hz, 1H), 2.88 (q, J=7.2 Hz, 2H), 2.26 (br d, J=12.5 Hz, 2H), 2.09 (br d, J=11.0 Hz, 2H), 1.79 - 1.65 (m, 2H), 1.50 - 1.38 (m, 2H), 1.25 (br d, J=6.1 Hz, 6H), 1.04 (t, J=7.3 Hz, 3H). ESI [M+H] = 585.2.

Example 110. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[morpholino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 134)

Following the same protocol and under the same reaction conditions as for Compound 111, Compound 134 was prepared. ¹H NMR (400MHz, methanol-*d*₄) δ = 7.74 (br s, 1H), 7.57 (br s, 1H), 7.34 (br d, J=7.9 Hz, 1H), 7.17 (br d, J=7.3 Hz, 1H), 4.82 - 4.74 (m, 1H), 3.85 (br s, 4H), 3.45 (br s, 1H), 3.28 (br s, 4H), 3.01 (br s, 1H), 2.85 (q, J=7.2 Hz, 2H), 2.24 (br d, J=12.6 Hz, 2H), 2.07 (br d, J=10.6 Hz, 2H), 1.68 (br s, 2H), 1.49 - 1.34 (m, 2H), 1.22 (br d, J=6.0 Hz, 6H), 1.01 (t, J=7.3 Hz, 3H). ESI [M+H] = 537.2.

Example 111. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[piperidin-1-yl]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 135)

Following the same protocol and under the same reaction conditions as for Compound 111, Compound 135 was prepared. ¹H NMR (400MHz, methanol-*d*₄) δ = 7.76 (s, 1H), 7.66 (br s, 1H), 7.45 - 7.33 (m, 1H), 7.26 (br d, J=7.1 Hz, 1H), 4.86 - 4.79 (m, 1H), 3.53 - 3.46 (m, 1H), 3.41 (br s, 4H), 3.07 (br t, J=11.4 Hz, 1H), 2.88 (q, J=7.1 Hz, 2H), 2.26 (br d, J=12.7 Hz, 2H), 2.10 (br d, J=11.0 Hz, 2H), 1.86 - 1.63 (m, 8H), 1.50 - 1.36 (m, 2H), 1.24 (br d, J=5.7 Hz, 6H), 1.04 (br t, J=7.2 Hz, 3H). ESI [M+H] = 535.2.

Example 112. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(2-isopropylpyrimidin-4-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 136)


Following the same protocol and under the same reaction conditions as for Compound 111, Compound 136 was prepared. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.83 (br s, 1H), 8.30 (d, J=7.21 Hz, 1H), 7.90 (dd, J=2.20, 8.31 Hz, 1H), 7.77-7.83 (m, 1H), 7.60 (d, J=8.31 Hz, 1H), 7.01 (d, J=7.09 Hz, 1H), 4.81-4.85 (m, 1H), 3.48 (tt, J=3.88, 11.58 Hz, 1H), 3.21 (spt, J=6.83 Hz, 1H), 3.05 (tt, J=3.50, 12.03 Hz, 1H), 2.87 (q, J=7.21 Hz, 2H), 2.28 (br d, J=12.35 Hz, 2H), 2.10 (br d, J=9.90 Hz, 2H), 1.73 (dq, J=2.93, 12.92 Hz, 2H), 1.47 (d, J=6.85

Hz, 6H), 1.29-1.45 (m, 2H), 1.25 (br d, $J=6.11$ Hz, 6H), 1.01 (t, $J=7.21$ Hz, 3H). ESI $[M+H]^+$ = 587.1.

Example 113. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(oxetan-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 137)

5 Following the same protocol and under the same reaction conditions as for Compound 111, Compound 137 was prepared. ^1H NMR (400MHz, $\text{DMSO-}d_6$) δ = 7.53 (s, 1H), 7.25 (t, $J=5.7$ Hz, 1H), 7.19 - 7.11 (m, 2H), 7.04 (d, $J=2.4$ Hz, 1H), 6.99 (br d, $J=7.7$ Hz, 1H), 6.62 (dd, $J=2.4, 8.4$ Hz, 1H), 4.85 (t, $J=6.5$ Hz, 2H), 4.72 (td, $J=6.2, 12.5$ Hz, 1H), 4.57 (sxt, $J=6.4$ Hz, 1H), 4.45 - 4.38 (m, 2H), 3.30 - 3.22 (m, 1H), 2.90 - 2.80 (m, 1H), 2.80 - 2.72 (m, 2H),
 10 2.10 (br d, $J=11.2$ Hz, 2H), 1.89 (br d, $J=10.1$ Hz, 2H), 1.61 - 1.44 (m, 2H), 1.39 - 1.21 (m, 2H), 1.14 (d, $J=6.4$ Hz, 6H), 0.95 (t, $J=7.2$ Hz, 3H). ESI $[M+H]^+$ = 523.2.

Example 114. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(3-isopropylimidazol-4-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 138)




a) Synthesis of 1-isopropyl-5-nitro-1H-imidazole



20 To a solution of 5-nitro-1H-imidazole (15 g, 133 mmol, 1.0 *eq.*) in DMF (50 mL) was added K_2CO_3 (55 g, 398 mmol, 3.0 *eq.*) and 2-iodopropane (24.81 g, 146 mmol, 1.1 *eq.*). The mixture was stirred at 65°C for 12 h. The reaction was filtered and filtrate was concentrated under reduced pressure. The residue was purified by prep-HPLC (neutral condition; column: Phenomenex luna C18 250*80mm*10 um; mobile phase: [water(0.1%TFA)-ACN]; B%: 5%-30%, 20min) to yield 1-isopropyl-5-nitro-1H-imidazole (1.7 g, 11 mmol, 8% yield) as


a yellow solid. ^1H NMR (400MHz, CDCl_3) δ = 8.19 (s, 1H), 8.11 (s, 1H), 5.33 (br d, J =6.8 Hz, 1H), 1.63 (br d, J =6.6 Hz, 6H). ESI $[\text{M}+\text{H}] = 156.1$.

b) Synthesis of 3-isopropylimidazol-4-amine




- 5 A mixture of 1-isopropyl-5-nitro-1H-imidazole (200 mg, 1 mmol, 1.0 *eq.*), Fe (360 mg, 6 mmol, 5 *eq.*), NH_4Cl (207 mg, 4 mmol, 3.0 *eq.*) in THF (10 mL), EtOH (10 mL) and H_2O (3 mL) was stirred at 80°C for 2 h under N_2 atmosphere. The reaction was filtered, then the filter was concentrated under reduced pressure. The residue was diluted with H_2O 20 mL and extracted with EtOAc 90 mL (30 mL X 3). The combined organic
- 10 layers dried over Na_2SO_4 , filtered and concentrated under reduced pressure to yield 3-isopropylimidazol-4-amine (150 mg, crude) as a black oil. ESI $[\text{M}+\text{H}] = 126.1$.

c) Synthesis of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(3-isopropylimidazol-4-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 138**)




- 15 From isopropyl *trans*-N-[4-[5-[4-bromo-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate, using General Method F. ^1H NMR (400 MHz, methanol- d_4) δ = 9.12 (s, 1H), 7.69 (s, 1H), 7.59 (s, 1H), 7.48 (d, J =2.38 Hz, 1H), 7.38 (d, J =8.38 Hz, 1H), 7.05 (dd, J =8.25, 2.38 Hz, 1H), 4.78 - 4.83 (m, 1H), 4.61 (dt, J =13.41, 6.61 Hz, 1H), 3.46 (qd, J =7.61, 3.56 Hz, 1H), 3.02 (ddd, J =11.88, 8.69, 3.44 Hz, 1H), 2.84 (q, J =7.21 Hz, 2H), 2.25 (br d, J =12.26 Hz, 2H), 2.09 (br d, J =10.38 Hz, 2H), 1.65 - 1.77 (m, 2H), 1.58 (d, J =6.63 Hz, 6H),
- 20 1.39 - 1.48 (m, 2H), 1.24 (br d, J =6.00 Hz, 6H), 1.01 (t, J =7.19 Hz, 3H). ESI $[\text{M}+\text{H}] = 575.1$.


Example 115. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(1-isopropyl-1H-imidazol-4-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 139**)**



a) Synthesis of 1-isopropylimidazol-4-amine




To a solution of 1-isopropyl-4-nitroimidazole (1.0 g, 6.5 mmol, 1.0 *eq.*) in EtOAc (50 mL) was added Pd/C (0.1 g, 10% purity) and the mixture was stirred under H₂ (15 psi) at 25°C for 3 h, and then filtered and concentrated under reduced pressure. The residue was purified by prep-HPLC (TFA condition). column: Phenomenex luna C18 250*50mm*10 um; mobile phase: [water(0.1%TFA)-ACN]; B%: 1%-10%, 10min to yield 1-isopropylimidazol-4-amine (390 mg, 3.12 mmol, 48% yield) as a yellow oil. ¹H NMR (400 MHz, CDCl₃) δ = 7.91 - 7.77 (m, 1H), 6.34 - 6.23 (m, 1H), 4.42 - 4.30 (m, 1H), 1.53 (d, J = 6.7 Hz, 6H). ESI [M+H] = 126.1.


b) Synthesis of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(1-isopropyl-1H-imidazol-4-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 139**)

From isopropyl *trans*-N-[4-[5-[4-bromo-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate and 1-isopropylimidazol-4-amine, using General Method F. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.91 - 8.86 (m, 1H), 7.70 (s, 1H), 7.63 - 7.59 (m, 2H), 7.42 - 7.37 (m, 1H), 7.18 - 7.12 (m, 1H), 4.87 - 4.81 (m, 1H), 4.73 - 4.62 (m, 1H), 3.54 - 3.43 (m, 1H), 3.03 (tt, J = 3.5, 12.0 Hz, 1H), 2.86 (q, J = 7.2 Hz, 2H), 2.31 - 2.21 (m, 2H), 2.13 - 2.04 (m, 2H), 1.77 - 1.67 (m, 2H), 1.65 - 1.61 (m, 6H), 1.49 - 1.39 (m, 2H), 1.25 (br d, J = 6.1 Hz, 6H), 1.07 - 0.98 (m, 3H). ESI [M+H] = 575.3.

Example 116. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(6-isopropylpyrazin-2-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 140)




a) Synthesis of 6-isopropenylpyrazin-2-amine



A mixture of 2-isopropenyl-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (1.0 g, 6.0 mmol, 1.3 *eq.*), 6-bromopyrazin-2-amine (820 mg, 5 mmol, 1.0 *eq.*), Cs₂CO₃ (4.6 g, 14 mmol, 3.0 *eq.*) and Pd(dppf)Cl₂ (345 mg, 471 μmol, 0.1 *eq.*) in dioxane (20 mL) and H₂O (10 mL) was stirred at 80°C for 12 h under N₂ atmosphere. The reaction mixture was concentrated under reduced pressure. The residue was diluted with H₂O 20 mL and extracted with EtOAc 90 mL (30 mL X 3). The combined organic layers dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by column chromatography (SiO₂, petroleum ether: ethyl acetate=10:1 to 5:1 to 3:1 to 1:1 to 0:1) to yield 6-isopropenylpyrazin-2-amine (730 mg, crude) as a white solid. ¹H NMR (400 MHz, methanol-*d*₄) δ = 7.91 (s, 1 H), 7.79 (s, 1 H), 5.87 (s, 1 H), 5.29 (s, 1 H), 2.14 (s, 3 H). ESI [M+H] = 136.1.


b) Synthesis of 6-isopropylpyrazin-2-amine



To a solution of 6-isopropenylpyrazin-2-amine (700 mg, 5 mmol, 1 *eq.*) in EtOAc (20 mL) was added AcOH (31 mg, 518 μmol, 0.1 *eq.*) and Pd/C (100 mg, 10% purity). The mixture was stirred under H₂ (15 psi) at 30°C for 12 h. The mixture was filtered, the filter liquor was concentrated under reduced pressure. The residue was purified by prep-HPLC (column: Phenomenex Luna C18 200*40mm*10μm; mobile phase: [water(10mM


NH_4HCO_3)-ACN];B%: 1%-30%,10min) to yield 6-isopropylpyrazin-2-amine (300 mg) as a white solid. ^1H NMR (400 MHz, methanol- d_4) δ = 7.73 (s, 1 H) 7.63 (s, 1 H) 2.82 - 2.92 (m, 1 H) 1.27 (d, J=6.88 Hz, 6 H). ESI $[\text{M}+\text{H}] = 138.2$.

- c) Synthesis of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[(6-isopropylpyrazin-2-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 140**)




- From isopropyl *trans*-*N*-[4-[5-[4-bromo-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate and 6-isopropylpyrazin-2-amine, using General Method F. ^1H NMR (400 MHz, methanol- d_4) δ = 8.84 (d, J=2.25 Hz, 1H), 8.03 (s, 1H), 7.94 (dd, J=8.38, 2.25 Hz, 1H), 7.90 (s, 1H), 7.72 (s, 1H) 7.43 (d, J=8.38 Hz, 1H), 4.81 - 4.85 (m, 1H), 3.41 - 3.55 (m, 1H), 2.99 - 3.13 (m, 2H), 2.90 (q, J=7.13 Hz, 2H), 2.27 (br d, J=12.26 Hz, 2H), 2.10 (br d, J=10.51 Hz, 2H), 1.66 - 1.77 (m, 2H), 1.42 - 1.51 (m, 2H), 1.39 (d, J=6.88 Hz, 6H), 1.25 (br d, J=6.13 Hz, 6H), 1.03 (t, J=7.19 Hz, 3H). ESI $[\text{M}+\text{H}] = 587.1$.

- Example 117. Preparation of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[(5-methyl-1*H*-imidazol-2-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 141**)**




- a) Synthesis of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-guanidino-phenyl]thiazol-2-yl]cyclohexyl]carbamate



To a solution of isopropyl *trans*-*N*-[4-[5-[4-amino-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (0.5 g, 1 mmol, 1 *eq.*) in HCl (10 mL) was added cyanamide (180 mg, 2 mmol, 2 *eq.*). The mixture was stirred at 60°C for 6 h, and then filtered. The residue was purified by prep-HPLC (TFA condition). column: Phenomenex luna C18 250*50mm*10 um;mobile phase: [water(0.1%TFA)-ACN];B%: 20%-50%,10min to yield isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-guanidino-phenyl]thiazol-2-yl] cyclohexyl]carbamate (0.31 g, 497.85 umol, 46% yield, TFA salt) as a white solid. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.04 - 7.94 (m, 1H), 7.82 - 7.74 (m, 1H), 7.64 - 7.55 (m, 2H), 4.84 - 4.75 (m, 1H), 3.55 - 3.40 (m, 1H), 3.14 - 3.02 (m, 1H), 2.94 - 2.83 (m, 2H), 2.33 - 2.23 (m, 2H), 2.12 - 2.04 (m, 2H), 1.80 - 1.64 (m, 2H), 1.56 - 1.38 (m, 2H), 1.25 (br d, *J* = 6.1 Hz, 6H), 1.08 - 0.98 (m, 3H). ESI [M+H] = 509.1.

b) Synthesis of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[(5-methyl-1*H*-imidazol-2-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 141**)



15

To a solution of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-guanidino-phenyl]thiazol-2-yl]cyclohexyl]carbamate (136 mg, 219 umol, 1.5 *eq.*, TFA), TEA (44 mg, 438 umol, 3 *eq.*) in DMF (3 mL) was added 1-bromopropan-2-one (20 mg, 146 umol, 1 *eq.*). The mixture was stirred at 25°C for 5 h. The residue was purified by prep-HPLC (TFA condition). column: Phenomenex Luna C18 100*30mm*5um;mobile phase: [water(0.1%TFA)-ACN];B%: 10%-40%,12min to yield isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[(5-methyl-1*H*-imidazol-2-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (98% purity, TFA salt) (15 mg) as a pale yellow solid. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.22 - 8.08 (m, 1H), 7.86 - 7.70 (m, 3H), 6.89 - 6.75 (m, 1H), 5.01 - 4.90 (m, 1H), 3.53 - 3.41 (m, 1H), 3.11 - 3.01 (m, 1H), 2.92 (q, *J* = 7.1 Hz, 2H), 2.34 - 2.20 (m, 5H), 2.10 (br d, *J* = 10.6 Hz, 2H), 1.80

25


- 1.64 (m, 2H), 1.51 - 1.36 (m, 2H), 1.25 (br d, J = 6.0 Hz, 6H), 1.13 - 0.97 (m, 3H). ESI [M+H] = 547.1.

Example 118. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(5-isopropyl-1*H*-imidazol-2-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate

5 (Compound 142)


Following the same protocol and under the same reaction conditions as for Compound 141, Compound 142 was prepared. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.26 - 8.14 (m, 1H), 7.85 - 7.80 (m, 2H), 7.77 - 7.73 (m, 1H), 6.93 - 6.85 (m, 1H), 4.85 - 4.81 (m, 1H), 3.53 - 3.39 (m, 1H), 3.14 - 2.98 (m, 1H), 2.97 - 2.85 (m, 3H), 2.33 - 2.24 (m, 2H), 2.13 - 2.05 (m, 2H), 1.80 - 1.63 (m, 2H), 1.51 - 1.37 (m, 2H), 1.33 (d, J = 6.8 Hz, 6H), 1.25 (br d, J = 6.1 Hz, 6H), 1.10 - 0.99 (m, 3H). ESI [M+H] = 575.1.

Example 119. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(6-isopropylpyridazin-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 143)



15


a) Synthesis of 6-isopropenylpyridazin-3-amine




A mixture of 2-isopropenyl-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (1.16 g, 7 mmol, 1.2 *eq.*), 6-bromopyridazin-3-amine (1 g, 6 mmol, 1 *eq.*), Cs₂CO₃ (5.6 g, 17 mmol, 3 *eq.*), Pd(dppf)Cl₂ (421 mg, 575 μmol, 0.1 *eq.*) in dioxane (20 mL)/H₂O (10 mL) was stirred at 80°C for 12 h under N₂ atmosphere, and then concentrated under reduced pressure to give a residue which was purified by column chromatography (SiO₂, petroleum ether : ethyl acetate=10:1 to 0:1) to yield 6-isopropenylpyridazin-3-amine (550 mg, 4 mmol, 71% yield) as a pale yellow solid. ESI [M+H] = 136.1.

20

b) Synthesis of 6-isopropylpyridazin-3-amine




To a solution of 6-isopropenylpyridazin-3-amine (540 mg, 4 mmol, 1 *eq.*) and AcOH (24 mg, 400 μ mol, 0.1 *eq.*) in EtOAc (30 mL) was added Pd/C (50 mg, 10% purity) and the mixture was stirred under H₂ (15 psi) at 25°C for 12 h. The reaction mixture was filtered and concentrated under reduced pressure to give a residue which was purified by prep-HPLC (neutral condition). column: Waters Xbridge Prep OBD C18 150*40mm*10 μ m; mobile phase: [water(10mM NH₄HCO₃)-ACN]; B%: 1%-15%, 8min to yield 6-isopropylpyridazin-3-amine (300 mg, 2 mmol, 55% yield) as a white solid. ¹H NMR (400 MHz, CDCl₃) δ = 7.18 - 7.09 (m, 1H), 6.78 - 6.69 (m, 1H), 4.91 - 4.66 (m, 2H), 3.27 - 3.10 (m, 1H), 1.30 (d, J = 7.0 Hz, 6H). ESI [M+H] = 138.2.


c) Synthesis of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(6-isopropylpyridazin-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 143**)

From isopropyl *trans*-N-[4-[5-[4-bromo-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate and 6-isopropylpyridazin-3-amine, using General Method F. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.63 - 8.49 (m, 1H), 7.99 - 7.94 (m, 1H), 7.93 - 7.88 (m, 1H), 7.75 (s, 1H), 7.62 - 7.55 (m, 1H), 7.50 (d, J = 8.3 Hz, 1H), 4.86 - 4.82 (m, 1H), 3.48 (tt, J = 3.8, 11.5 Hz, 1H), 3.31 - 3.25 (m, 1H), 3.10 - 2.98 (m, 1H), 2.91 (q, J = 7.2 Hz, 2H), 2.31 - 2.23 (m, 2H), 2.14 - 2.06 (m, 2H), 1.79 - 1.66 (m, 2H), 1.50 - 1.43 (m, 2H), 1.42 (d, J = 7.0 Hz, 6H), 1.25 (br d, J = 6.2 Hz, 6H), 1.07 - 0.98 (m, 3H). ESI [M+H] = 587.1.

Example 120. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[[1-(2-methylpropanoyl)azetidin-3-yl]amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 146)




a) Synthesis of *tert*-butyl *trans*-3-[3-(ethylsulfamoyl)-4-[2-[4-(isopropoxycarbonylamino)cyclohexyl]thiazol-5-yl]anilino]azetidine-1-carboxylate



5 From isopropyl *trans*-*N*-[4-[5-[4-bromo-2-(ethylsulfamoyl)phenyl]thiazol-2-yl] cyclohexyl]carbamate and *tert*-butyl 3-aminoazetidine-1-carboxylate, using General Method F. ESI $[\text{M}+\text{H}] = 622.2$.

b) Synthesis of isopropyl *trans*-*N*-[4-[5-[4-(azetidin-3-ylamino)-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 144**)




10

A mixture of *tert*-butyl *trans*-3-[3-(ethylsulfamoyl)-4-[2-[4-(isopropoxycarbonylamino)cyclohexyl]thiazol-5-yl]anilino]azetidine-1-carboxylate (100 mg, 161 μmol , 1 *eq.*) in HCl/MeOH (2 mL) (4 M) was stirred at 25°C for 30 min under N_2 atmosphere. The reaction mixture was concentrated under reduced pressure to yield isopropyl *trans*-*N*-[4-[5-


[4-(azetidin-3-ylamino)-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (89.76 mg, crude, HCl salt) as a yellow oil. ¹H NMR (400MHz, methanol-*d*₄) δ = 7.73 (s, 1H), 7.30 (dd, J=2.8, 5.0 Hz, 2H), 6.90 - 6.71 (m, 1H), 4.88 - 4.77 (m, 1H), 4.71 - 4.56 (m, 1H), 4.46 (br t, J=8.9 Hz, 2H), 4.15 - 3.97 (m, 2H), 3.47 (br t, J=11.6 Hz, 1H), 3.07 (br t, J=12.0 Hz, 1H), 2.87 (q, J=7.2 Hz, 2H), 2.25 (br d, J=12.2 Hz, 2H), 2.09 (br d, J=10.9 Hz, 2H), 1.78 - 1.59 (m, 2H), 1.52 - 1.35 (m, 2H), 1.24 (br d, J=5.9 Hz, 6H), 1.03 (br t, J=7.2 Hz, 3H). ESI [M+H] = 522.2.

c) Synthesis of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[(1-isobutyl-azetidin-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 145**)




To a solution of isopropyl *trans*-*N*-[4-[5-[4-(azetidin-3-ylamino)-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (89 mg, 159 μ mol, 1.0 *eq.*, HCl salt) in DCE (2 mL) was added AcOH (1 mg, 16 μ mol, 0.1 *eq.*), 2-methylpropanal (11.50 mg, 159 μ mol, 1.0 *eq.*), NaBH(OAc)₃ (101 mg, 478 μ mol, 3.0 *eq.*) at 0°C. The mixture was stirred at 15 25°C for 2 h. The reaction mixture was concentrated under reduced pressure. The residue was purified by prep-HPLC (neutral condition; column: Waters Xbridge BEH C18 100*30mm*10 μ m; mobile phase: [water(10mM NH₄HCO₃)-ACN]; B%: 45%-75%, 10min) to yield isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[(1-isobutylazetidin-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (80 mg, 138 μ mol, 86.8% yield) as a 20 yellow solid. ¹H NMR (400MHz, methanol-*d*₄) δ = 7.62 (s, 1H), 7.33 - 7.22 (m, 2H), 6.86 - 6.71 (m, 1H), 4.80 (br d, J=6.0 Hz, 1H), 4.77 - 4.65 (m, 1H), 4.55 (br d, J=7.1 Hz, 1H), 4.44 (br s, 1H), 4.25 (br s, 1H), 3.99 (br d, J=9.5 Hz, 1H), 3.51 - 3.35 (m, 1H), 3.24 - 3.09 (m, 2H), 3.06 - 2.92 (m, 1H), 2.82 (q, J=7.1 Hz, 2H), 2.22 (br d, J=11.9 Hz, 2H), 2.06 (br d, J=10.4 Hz, 2H), 2.02 - 1.91 (m, 1H), 1.78 - 1.59 (m, 2H), 1.49 - 1.31 (m, 2H), 1.22 (d, J=6.2 Hz, 25 6H), 1.02 (d, J=6.6 Hz, 6H), 0.99 (t, J=7.3 Hz, 3H). ESI [M+H] = 578.2.

d) Synthesis of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[[1-(2-methylpropanoyl)azetidin-3-yl]amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 146**)




To a solution of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(1-isobutylazetidin-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (30 mg, 52 μ mol, 1.0 eq.) in DCM (2 mL) was added KMnO₄ (25 mg, 156 μ mol, 3.0 eq.) and benzyltriethylammonium chloride (35 mg, 156 μ mol, 3 eq.). The mixture was stirred at 0°C for 0.5 h. The reaction was quenched with sat. aq. Na₂SO₃ 20 mL and extracted with DCM 20 mL (10 mL X 2). The combined organic layers were dried with Na₂SO₄ and concentrated under reduced pressure. The residue was purified by prep-HPLC (TFA condition; column: Phenomenex Luna C18 150*30mm*5 μ m; mobile phase: [water(0.1%TFA)-ACN]; B%: 45%-75%, 10min) to yield isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(1-(2-methylpropanoyl)azetidin-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (6.1 mg, 9.6 μ mol, 18% yield, 92.8% purity) as a pale yellow solid. ¹H NMR (400MHz, methanol-*d*₄) δ = 7.64 (s, 1H), 7.24 (dd, J=2.9, 5.3 Hz, 2H), 6.76 (dd, J=2.4, 8.4 Hz, 1H), 4.83 - 4.75 (m, 1H), 4.70 - 4.59 (m, 1H), 4.42 - 4.29 (m, 2H), 4.10 - 3.96 (m, 1H), 3.88 - 3.73 (m, 1H), 3.51 - 3.35 (m, 1H), 2.99 (br t, J=12.0 Hz, 1H), 2.85 (q, J=7.3 Hz, 2H), 2.57 (td, J=6.6, 13.5 Hz, 1H), 2.22 (br d, J=12.8 Hz, 2H), 2.06 (br d, J=10.8 Hz, 2H), 1.76 - 1.60 (m, 2H), 1.47 - 1.33 (m, 2H), 1.22 (br d, J=6.0 Hz, 6H), 1.08 (dd, J=5.1, 6.6 Hz, 6H), 1.01 (t, J=7.3 Hz, 3H). ESI [M+H] = 592.1.

Example 121. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(3-methylazetidin-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 147)




20

a) Synthesis of *tert*-butyl *trans*-3-[3-(ethylsulfamoyl)-4-[2-[4-(isopropoxycarbonylamino)cyclohexyl]thiazol-5-yl]anilino]-3-methyl-azetidine-1-carboxylate




From isopropyl *trans*-*N*-[4-[5-[4-bromo-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate and *tert*-butyl 3-amino-3-methylazetidine-1-carboxylate, using General Method F. ESI [M+H] = 636.2.

- 5 b) Synthesis of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[(3-methylazetidin-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 147**)




- A solution of *tert*-butyl *trans*-3-[3-(ethylsulfamoyl)-4-[2-[4-(isopropoxycarbonylamino)cyclohexyl]thiazol-5-yl]anilino]-3-methylazetidine-1-carboxylate (150 mg, 236 μ mol, 1 *eq.*) in HCl/MeOH (2 mL, 4M) was stirred at 25°C for 2 h. The reaction was concentrated. The residue was purified by prep-HPLC (TFA condition: column: Phenomenex luna C18 250*50mm*10 μ m; mobile phase: [water(0.1%TFA)-ACN]; B%: 15%-45%, 10min) to yield isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[(3-methylazetidin-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (6.7 mg, 10 μ mol, 4.3% yield, 99.4% purity, TFA) as a pale yellow solid. ¹H NMR (400MHz, methanol-*d*₄) δ = 7.70 - 7.66 (m, 1H), 7.30 (d, J=8.4 Hz, 1H), 7.22 (d, J=2.5 Hz, 1H), 6.71 (dd, J=2.4, 8.3 Hz, 1H), 4.85 (br s, 1H), 4.25 - 4.14 (m, 4H), 3.47 (ddd, J=4.1, 7.7, 11.5 Hz, 1H), 3.04 (br t, J=12.0 Hz, 1H), 2.86 (q, J=7.1 Hz, 2H), 2.25 (br d, J=12.3 Hz, 2H), 2.13 - 2.03 (m, 2H), 1.76 - 1.66 (m, 5H), 1.43 (dq, J=3.0, 12.5 Hz, 2H), 1.28 - 1.20 (m, 6H), 1.03 (t, J=7.2 Hz, 3H). ESI [M+H] = 536.2.

- 20 **Example 122. Preparation of isopropyl *trans*-*N*-[4-[5-[2-(azetidin-1-ylsulfonyl)-4-(1*H*-imidazol-2-ylamino)phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 148)**




a) Synthesis of isopropyl *trans*-*N*-[4-[5-[2-(azetidin-1-ylsulfonyl)-4-[[1-(2-trimethylsilylethoxymethyl)-1*H*-imidazol-2-yl]amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate



From isopropyl *trans*-*N*-[4-[5-[2-(azetidin-1-ylsulfonyl)-4-bromo-phenyl]thiazol-2-yl]cyclohexyl]carbamate and 1-((2-(trimethylsilyl)ethoxy)methyl)-1*H*-imidazol-2-amine, using General Method F. ESI [M+H] = 675.3.

b) Synthesis of isopropyl *trans*-*N*-[4-[5-[2-(azetidin-1-ylsulfonyl)-4-(1*H*-imidazol-2-ylamino)phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 148**)


10



From isopropyl *trans*-*N*-[4-[5-[2-(azetidin-1-ylsulfonyl)-4-[[1-(2-trimethylsilylethoxy-methyl)-1*H*-imidazol-2-yl]amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate, using General Method I. ¹H NMR (400 MHz, methanol-*d*₄) δ =

15 7.89 (d, *J* = 2.2 Hz, 1H), 7.71 (s, 1H), 7.62 - 7.48 (m, 2H), 7.11 (s, 2H), 4.84 - 4.76 (m, 1H), 3.68 (t, *J* = 7.7 Hz, 4H), 3.54 - 3.38 (m, 1H), 3.04 (tt, *J* = 3.5, 12.1 Hz, 1H), 2.27 - 2.17 (m, 2H), 2.17 - 2.09 (m, 2H), 2.07 (br d, *J* = 10.8 Hz, 2H), 1.70 (dq, *J* = 2.9, 12.8 Hz, 2H), 1.41 (dq, *J* = 3.1, 12.6 Hz, 2H), 1.22 (br d, *J* = 6.2 Hz, 6H). ESI [M+H] = 545.2.

Example 123. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-(2*H*-triazol-4-ylamino)phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 149)



a) Synthesis of trimethyl-[2-[(4-nitrotriazol-2-yl)methoxy]ethyl]silane




5

To a solution of 4-nitro-2*H*-triazole (1 g, 8.7 mmol, 1 *eq.*) in THF (20 mL) was added SEM-Cl (1.75 g, 10.5 mmol, 1.2 *eq.*) and NaH (385 mg, 9.6 mmol, 60% in oil, 1.1 *eq.*) at 0°C. The mixture was stirred at 25°C for 12 h. The reaction mixture was quenched by sat. aq. NH₄Cl (10 mL) at 0°C, and then extracted with ethyl acetate (60 mL) (20 mL X 3).

10 The combined organic layers were dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue, which was purified by MPLC (SiO₂, petroleum ether: ethyl acetate = 50:1 to 10:1) to yield trimethyl-[2-[(4-nitrotriazol-2-yl)methoxy]ethyl]silane (0.76 g, 3.1 mmol, 35% yield) as colorless oil. ¹H NMR (400 MHz, CDCl₃) δ = 8.30 - 8.21 (m, 1H), 5.67 - 5.56 (m, 2H), 3.79 - 3.68 (m, 2H), 1.01 - 0.92 (m, 2H), 0.05 - 0.05 (m, 9H).

15 b) Synthesis of 2-(2-(trimethylsilylethoxymethyl)triazol-4-amine




To a solution of trimethyl-[2-[(4-nitrotriazol-2-yl)methoxy]ethyl]silane (0.76 g, 3.1 mmol, 1 *eq.*) in ethyl acetate (20 mL) was added Pd/C (0.05 g, 10% purity). The mixture was stirred under H₂ (15 psi) at 25°C for 12 h. The reaction mixture was filtered and concentrated under reduced pressure to give a residue, which was purified by prep-HPLC (neutral condition: column: Welch Xtimate C18 150*30mm*5um; mobile phase: [water(10mM

20

NH_4HCO_3) - ACN]; B%: 20%-50%, 8min) to yield 2-(2-trimethylsilyl-ethoxymethyl)triazol-4-amine (0.3 g, 1.4 mmol, 45% yield) as a white solid. ^1H NMR (400 MHz, CDCl_3) δ = 7.17 - 7.03 (m, 1H), 5.55 - 5.43 (m, 2H), 3.86 - 3.73 (m, 2H), 3.68 - 3.54 (m, 2H), 1.00 - 0.82 (m, 2H), 0.00 (s, 9H). ESI $[\text{M}+\text{H}] = 215.1$.


- 5 c) Synthesis of isopropyl *trans-N*-[4-[5-[2-(ethylsulfamoyl)-4-[[2-(2-trimethylsilylethoxymethyl)triazol-4-yl]amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate



From isopropyl *trans-N*-[4-[5-[4-bromo-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate and 2-(2-trimethylsilyl-ethoxymethyl)triazol-4-amine, using General


- 10 Method F. ESI $[\text{M}+\text{H}] = 664.3$.

- d) Synthesis of isopropyl *trans-N*-[4-[5-[2-(ethylsulfamoyl)-4-(2*H*-triazol-4-ylamino)phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 149**)




- 15 From isopropyl *trans-N*-[4-[5-[2-(ethylsulfamoyl)-4-[[2-(2-trimethylsilylethoxy-methyl)triazol-4-yl]amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate, using General Method I. ^1H NMR (400 MHz, methanol- d_4) δ = 8.15 - 8.03 (m, 1H), 7.80 - 7.71 (m, 1H), 7.58 - 7.50 (m, 1H), 7.49 - 7.44 (m, 1H), 7.41 - 7.33 (m, 1H), 4.80 - 4.71 (m, 1H), 3.54 - 3.42 (m, 1H), 3.12 - 3.03 (m, 1H), 2.99 - 2.90 (m, 2H), 2.33 - 2.23 (m, 2H), 2.16 - 2.03 (m, 2H), 1.81 - 1.66 (m, 2H), 1.52 - 1.36 (m, 2H), 1.34 - 1.20 (m, 6H), 1.12 - 1.05 (m, 3H). ESI $[\text{M}+\text{H}] = 534.2$.

- 20 **Example 124. Preparation of isopropyl *trans-N*-[4-[5-[2-(ethylsulfamoyl)-4-[1*H*-imidazol-2-yl(methyl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 150)**




a) Synthesis of trimethyl-[2-[(2-nitroimidazol-1-yl)methoxy]ethyl]silane




A mixture of 2-nitro-1*H*-imidazole (9 g, 79.5 mmol, 1 *eq.*), NaH (9.55 g, 238.7 mmol, 60% purity, 3 *eq.*) in THF (150 mL) was degassed and purged with N₂ for 3 times, and then the mixture was stirred at 0°C for 30 min under N₂ atmosphere, then to the mixture was added 2-(chloromethoxy)ethyl-trimethyl-silane (15.92 g, 95.5 mmol, 16.9 mL, 1.2 *eq.*) at 0°C. The mixture was stirred at 25°C for 2 h. The reaction mixture was quenched by pouring into sat. aq. NH₄Cl 50 mL at 0°C and then diluted with H₂O (50 mL) and extracted with ethyl acetate 450 mL (150 mL X 3). The combined organic layers were dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue, which was purified by column chromatography (SiO₂, petroleum ether: ethyl acetate = 10:1 to 2:1) to yield trimethyl-[2-[(2-nitroimidazol-1-yl)methoxy] ethyl]silane (13 g, 53.4 mmol, 67% yield) as a white solid. ¹H NMR (400 MHz, CDCl₃) δ = 7.34 (d, J=0.88 Hz, 1H), 7.18 (d, J=0.88 Hz, 1H), 5.78 (s, 2H) 3.59 - 3.72 (m, 2H), 0.87 - 1.04 (m, 2H), 0.04 - 0.02 (m, 9H).

b) Synthesis of 1-(2-trimethylsilylethoxymethyl)imidazol-2-amine




A mixture of trimethyl-[2-[(2-nitroimidazol-1-yl)methoxy]ethyl]silane (4 g, 16.44 mmol, 1 eq.), Pd/C (200 mg, 10% purity), in ethyl acetate (20 mL) was degassed and purged with N₂ for 3 times, and then the mixture was stirred at 25°C for 3 h under H₂ atmosphere. The reaction was filtered and the filtrate was concentrated under reduced pressure to give a residue which was purified by prep-HPLC (column: Welch Xtimate C18 250*50mm*10um; mobile phase: [water(10mM NH₄HCO₃)-ACN]; B%: 20%-50%, 23min) to yield 1-(2-trimethylsilylethoxymethyl)imidazol-2-amine (3.1 g, 14.5 mmol, 88% yield) as a white solid. ¹H NMR (400 MHz, methanol-*d*₄) δ = 6.67 (d, J=1.63 Hz, 1H), 6.48 (d, J=1.75 Hz, 1H), 5.13 (s, 2H), 3.46 - 3.61 (m, 2H), 0.84 - 0.96 (m, 2H), 0.00 (s, 9H). ESI [M+H] = 214.2.

c) Synthesis of *N*-methyl-1-(2-trimethylsilylethoxymethyl)imidazol-2-amine




A solution of 1-(2-trimethylsilylethoxymethyl)imidazol-2-amine (2 g, 9.3 mmol, 1 eq.) in diethoxymethoxyethane (20 mL) was stirred at 100°C for 3 h. The reaction mixture was concentrated under reduced pressure. The residue was dissolved in EtOH (20 mL) and NaBH₄ (1.06 g, 28.1 mmol, 3 eq.) was added to the solution in portions at 0°C and stirred at 80°C for 2 h. The reaction mixture was quenched with H₂O (10 mL) at 30°C, the reaction mixture was concentrated under reduced pressure to remove solvent, and then the mixture was diluted with H₂O (20 mL) and extracted with ethyl acetate 90 mL (30 mL X 3). The combined organic layers dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue which was purified by prep-HPLC (column: Agela DuraShell C18 250*25mm*10um; mobile phase: [water(10mM NH₄HCO₃)-ACN]; B%: 5%-33%, 22min) to yield *N*-methyl-1-(2-trimethylsilylethoxy-methyl)imidazol-2-amine (470 mg, 2.0 mmol, 22% yield) as a pale yellow oil. ¹H NMR (400 MHz, methanol-*d*₄) δ = 6.69 (d, J=1.71 Hz, 1H), 6.44 - 6.57 (m, 1H), 5.10 (s, 2H), 3.52 - 3.63 (m, 2H), 2.89 (s, 3H), 0.88 - 0.95 (m, 2H), 0.00 (s, 9H). ESI [M+H] = 228.0.

d) Synthesis of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[methyl-1-(2-trimethylsilylethoxymethyl)imidazol-2-yl]amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate




From isopropyl *trans*-*N*-[4-[5-[4-bromo-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate and *N*-methyl-1-(2-trimethylsilylethoxy-methyl)imidazol-2-amine, using General Method F. ESI [M+H] = 677.3.

- 5 e) Synthesis of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[1*H*-imidazol-2-yl(methyl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 150**)




- From isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[methyl-[1-(2-trimethylsilylethoxymethyl)imidazol-2-yl]amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate, using General
- 10 Method I. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.11 (d, *J*=2.25 Hz, 1H), 7.80 (s, 1H), 7.65 - 7.76 (m, 2H), 7.06 (s, 2H), 4.75 (br s, 1H), 3.60 (s, 3H), 3.48 (tt, *J*=11.58, 3.99 Hz, 1H), 3.05 (tt, *J*=12.05, 3.52 Hz, 1H), 2.90 (q, *J*=7.25 Hz, 2H), 2.22 - 2.34 (m, 2H), 2.02 - 2.15 (m, 2H), 1.73 (qd, *J*=12.84, 3.00 Hz, 2H), 1.37 - 1.53 (m, 2H), 1.25 (br d, *J*=6.25 Hz, 6H), 1.00 - 1.08 (m, 3H). ESI [M+H] = 547.2.

- 15 **Example 125. Preparation of isopropyl *trans*-*N*-[4-[5-[2-(*tert*-butylsulfamoyl)-4-(4*H*-1,2,4-triazol-3-ylamino)phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 151)**



a) Synthesis of 2-(4H-1,2,4-triazol-3-yl)isoindoline-1,3-dione



A mixture of 1H-1,2,4-triazol-3-amine (8.4 g, 99.9 mmol, 1 *eq*) and isobenzofuran-1,3-dione (14.80 g, 99.9 mmol, 1 *eq*) in dioxane (200 mL) the mixture was stirred at 120°C for 10 h under N₂ atmosphere. The mixture was cooled to 25°C, and then filtered to yield 2-(4H-1,2,4-triazol-3-yl)isoindoline-1,3-dione (14 g, 65.3 mmol, 65.4% yield) as a white solid. ESI [M-H]⁺=212.9.


b) Synthesis of 2-[4-(2-trimethylsilylethoxymethyl)-1,2,4-triazol-3-yl] isoindoline-1,3-dione



To a mixture of 2-(chloromethoxy)ethyl-trimethyl-silane (5.84 g, 35.0 mmol, 1.5 *eq*), 2-(4H-1,2,4-triazol-3-yl)isoindoline-1,3-dione (5 g, 23.3 mmol, 1 *eq*) was added NaH (1.40 g, 35.0 mmol, 60% in oil, 1.5 *eq*) in DMF (50 mL), and then the mixture was stirred at 50°C for 5 h under N₂ atmosphere. The reaction mixture was quenched with H₂O (20 mL) and extracted with ethyl acetate 180 mL (60 mL X 3). The combined organic layers were washed with sat.aq. NaCl 100 mL (20 mL X 5), dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield 2-[4-(2-


trimethylsilylethoxymethyl)-1,2,4-triazol-3-yl]isoindoline-1,3-dione (9 g, crude) as yellow oil. ESI [M+H] = 345.2.

c) Synthesis of 4-(2-trimethylsilylethoxymethyl)-1,2,4-triazol-3-amine




To a solution of 2-[4-(2-trimethylsilylethoxymethyl)-1,2,4-triazol-3-yl]isoindoline-1,3-dione (8 g, 23.2 mmol, 1 *eq*) in THF (40 mL) and MeOH (40 mL) was added NH₂-NH₂·H₂O (1.74 g, 34.8 mmol, 98% purity, 1.5 *eq*), and then the mixture was stirred at 25°C for 10 h under N₂ atmosphere. The mixture was concentrated, and purified by column chromatography (SiO₂, petroleum ether: ethyl acetate = 1:1 to 0:1) to yield 4-(2-trimethylsilylethoxymethyl)-1,2,4-triazol-3-amine (2 g, 9.3 mmol, 40% yield) as white solid. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.11-8.25 (m, 1H), 5.25 (br s, 2H), 3.51-3.69 (m, 2H), 0.81-1.00 (m, 2H), -0.10-0.10 (m, 9H). ESI [M+H] = 215.1.

d) Synthesis of isopropyl *trans*-*N*-[4-[5-[4-bromo-2-(*tert*-butylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate




From isopropyl (*trans*-4-(5-(4-amino-2-(*N*-(*tert*-butyl)sulfamoyl)phenyl)thiazol-2-yl)cyclohexyl]carbamate, using General Method L. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.26 (d, *J* = 2.2 Hz, 1H), 7.83 - 7.76 (m, 2H), 7.41 (d, *J* = 8.2 Hz, 1H), 4.85 - 4.78 (m, 1H), 3.52 - 3.39 (m, 1H), 3.03 (tt, *J* = 3.6, 12.0 Hz, 1H), 2.28 - 2.18 (m, 2H), 2.12 - 2.02 (m, 2H), 1.70 (dq, *J* = 2.6, 12.9 Hz, 2H), 1.46 - 1.34 (m, 2H), 1.22 (br d, *J* = 6.2 Hz, 6H), 1.09 (s, 9H).

e) Synthesis of isopropyl *trans*-*N*-[4-[5-[2-(*tert*-butylsulfamoyl)-4-[[4-(2-trimethylsilylethoxymethyl)-1,2,4-triazol-3-yl]amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate



A mixture of isopropyl *trans*-N-[4-[5-[4-bromo-2-(*tert*-butylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (150 mg, 268.5 μ mol, 1 *eq.*), 4-(2-trimethylsilylethoxymethyl)-1,2,4-triazol-3-amine (86 mg, 402.8 μ mol, 1.5 *eq.*), Cs₂CO₃ (175 mg, 537.1 μ mol, 2 *eq.*), Brettphos (14 mg, 26.8 μ mol, 0.1 *eq.*) and Pd₂(dba)₃ (24 mg, 26.8 μ mol, 0.1 *eq.*) in *t*-BuOH (1.5 mL) and toluene (1.5 mL) was stirred at 100°C for 12 h under Ar atmosphere. The reaction mixture was concentrated, and purified by prep-TLC (SiO₂, petroleum ether: ethyl acetate = 1: 1) to yield isopropyl *trans*-N-[4-[5-[2-(*tert*-butylsulfamoyl)-4-[[4-(2-trimethylsilylethoxymethyl)-1,2,4-triazol-3-yl]amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (205 mg, crude) as a yellow solid. ESI [M+H] = 692.3.


f) Synthesis of isopropyl *trans*-N-[4-[5-[2-(*tert*-butylsulfamoyl)-4-(4*H*-1,2,4-triazol-3-ylamino)phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 151**)




Isopropyl *trans*-N-[4-[5-[2-(*tert*-butylsulfamoyl)-4-[[4-(2-trimethylsilylethoxymethyl)-1,2,4-triazol-3-yl]amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (80 mg, 115.6 μ mol, 1 *eq.*) and TsOH.H₂O (32 mg, 173.4 μ mol, 1.5 *eq.*) were taken up into a microwave tube with MeOH (3 mL). The sealed tube was heated at 100°C for 0.5 h under microwave. The reaction mixture was concentrated under reduced pressure and purified by prep-HPLC (TFA condition: column: Nano-micro Kromasil C18 80*25mm 3 μ m; mobile phase: [water(0.1%TFA)-ACN]; B%: 24%-58%, 7min) to yield isopropyl *trans*-N-[4-[5-[2-(*tert*-butylsulfamoyl)-4-(4*H*-1,2,4-triazol-3-ylamino)phenyl]thiazol-2-yl]cyclohexyl]carbamate (7 mg, 13.3 μ mol, 12% yield, 100% purity) as a pale yellow solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ = 9.74 (s, 1H), 8.40 (d, *J* = 2.4 Hz, 1H), 8.30 (br s, 1H), 7.68 (dd, *J* = 2.3, 8.5 Hz, 1H), 7.62 (s, 1H), 7.30 (d, *J* = 8.4 Hz, 1H), 7.00 (br d, *J* = 7.7 Hz, 1H), 6.79 (s, 1H), 4.72 (td, *J* =

6.2, 12.5 Hz, 1H), 3.30 (td, $J = 3.6, 7.4$ Hz, 1H), 2.87 (tt, $J = 3.4, 11.9$ Hz, 1H), 2.11 (br d, $J = 11.7$ Hz, 2H), 1.90 (br d, $J = 10.4$ Hz, 2H), 1.55 (dq, $J = 2.6, 12.7$ Hz, 2H), 1.37 - 1.25 (m, 2H), 1.14 (d, $J = 6.4$ Hz, 6H), 1.07 (s, 9H). ESI $[M+H] = 562.3$.

Example 126. Preparation of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-(4*H*-1,2,4-triazol-3-ylamino)phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 152)




a) Synthesis of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[[4-(2-trimethylsilylethoxymethyl)-1,2,4-triazol-3-yl]amino]phenyl]thiazol-2-yl]cyclohexyl] carbamate




A mixture of isopropyl *trans*-*N*-[4-[5-[4-bromo-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (0.13 g, 245.0 μmol , 1 *eq.*), 4-(2-trimethylsilylethoxy-methyl)-1,2,4-triazol-3-amine (105 mg, 490.1 μmol , 2 *eq.*), $\text{Pd}_2(\text{dba})_3$ (22 mg, 24.5 μmol , 0.1 *eq.*), Cs_2CO_3 (159 mg, 490.1 μmol , 2 *eq.*) and dicyclohexyl-[3,6-dimethoxy-2-(2,4,6-triisopropylphenyl)phenyl]phosphane (13 mg, 24.5 μmol , 0.1 *eq.*) in toluene (2 mL) and t-BuOH (2 mL) was stirred at 100°C for 12 h under N_2 atmosphere. The reaction mixture was concentrated under reduced pressure to give a residue which was purified by prep-TLC (SiO_2 , petroleum ether: ethyl acetate = 0:1) to yield isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[[4-(2-trimethylsilylethoxymethyl)-1,2,4-triazol-3-yl]amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (120 mg, 180.7 μmol , 74% yield) as a yellow solid. ESI $[M+H] = 664.3$.

b) Synthesis of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-(4*H*-1,2,4-triazol-3-ylamino)phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 152**)


**152**

A mixture of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[[4-(2-trimethylsilyl)ethoxymethyl]-1,2,4-triazol-3-yl]amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (95 mg, 143.0 μmol , 1 *eq.*) and $\text{TsOH}\cdot\text{H}_2\text{O}$ (40 mg, 214.6 μmol , 1.5 *eq.*) were taken up into a microwave tube with MeOH (2 mL). The sealed tube was heated at 100°C for 0.5 h under microwave. The reaction mixture was concentrated under reduced pressure to give a residue, which was purified by prep-HPLC (TFA condition: column: Phenomenex Luna C18 100*30mm*5 μm ; mobile phase: [water(0.1%TFA)-ACN]; B%: 15%-50%, 12min) to yield isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-(4*H*-1,2,4-triazol-3-ylamino)phenyl]thiazol-2-yl]cyclohexyl]carbamate (38 mg, 72.9 μmol , 51% yield) as a yellow gum. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ = 9.94 - 9.78 (m, 1H), 8.44 - 8.33 (m, 2H), 7.85 - 7.80 (m, 1H), 7.75 - 7.71 (m, 1H), 7.45 - 7.40 (m, 1H), 7.38 - 7.33 (m, 1H), 7.15 - 7.06 (m, 1H), 4.84 (td, J = 6.2, 12.4 Hz, 1H), 3.45 - 3.35 (m, 1H), 3.04 - 2.90 (m, 3H), 2.27 - 2.19 (m, 2H), 2.05 - 1.97 (m, 2H), 1.71 - 1.60 (m, 2H), 1.50 - 1.36 (m, 2H), 1.26 (d, J = 6.3 Hz, 6H), 1.12 - 1.04 (m, 3H). ESI $[\text{M}+\text{H}] = 534.3$.

Example 127. Preparation of isopropyl *trans*-[4-[5-[4-(oxazol-2-ylamino)-2-pyrrolidin-1-ylsulfonyl-phenyl]thiazol-2-yl]cyclohexyl]carbamate (153)




- 20 a) Synthesis of isopropyl *trans*-*N*-[4-[5-[4-(oxazol-2-ylamino)-2-pyrrolidin-1-ylsulfonyl-phenyl]thiazol-2-yl]cyclohexyl]carbamate



From isopropyl *trans*-*N*-[4-[5-(4-amino-2-pyrrolidin-1-ylsulfonyl-phenyl)thiazol-2-yl]cyclohexyl]carbamate, using General Method L. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.23 (d, *J*=2.00 Hz, 1H), 7.86 (dd, *J*=8.19, 2.06 Hz, 1H), 7.76 (s, 1H), 7.46 (d, *J*=8.13 Hz, 1H), 4.80 - 4.86 (m, 1H), 3.47 (tt, *J*=11.44, 3.63 Hz, 1H), 3.02 - 3.09 (m, 1H), 2.97 (br t, *J*=6.63 Hz, 4H), 2.19 - 2.25 (m, 2H), 2.09 (br d, *J*=10.26 Hz, 2H), 1.77 - 1.82 (m, 4H), 1.64 - 1.75 (m, 2H), 1.41 - 1.46 (m, 2H), 1.24 (br d, *J*=6.00 Hz, 6H). ESI [M+H] = 556.1/558.1.

b) Synthesis of isopropyl *trans*-[4-[5-[4-(oxazol-2-ylamino)-2-pyrrolidin-1-ylsulfonyl-phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 153**)



10

From isopropyl *trans*-*N*-[4-[5-(4-bromo-2-pyrrolidin-1-ylsulfonyl-phenyl)thiazol-2-yl]cyclohexyl]carbamate and 2-amino oxazole, using General Method F. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.27 (d, *J*=2.38 Hz, 1H), 7.68 (dd, *J*=8.38, 2.38 Hz, 1H), 7.58 (s, 1H), 7.41 (d, *J*=0.88 Hz, 1H), 7.34 (d, *J*=8.38 Hz, 1H), 6.89 (d, *J*=0.75 Hz, 1H), 4.72 (br s, 1H), 3.30 - 3.41 (m, 1H), 3.00 (s, 4H), 2.85 - 2.94 (m, 1H), 2.05 - 2.16 (m, 2H), 1.92 - 2.01 (m, 2H), 1.76 (dt, *J*=6.47, 3.46 Hz, 4H), 1.59 (qd, *J*=12.86, 2.94 Hz, 2H), 1.31 (qd, *J*=12.55, 3.13 Hz, 2H), 1.13 (br d, *J*=6.13 Hz, 6H). ESI [M+H] = 560.1.

15


Example 128. Preparation of isopropyl (*trans*-4-(5-(2-(*N*-(*tert*-butyl)sulfamoyl)-4-(oxazol-2-ylamino)phenyl)thiazol-2-yl)cyclohexyl]carbamate (Compound 42**)**

20

Following the same protocol and under the same reaction conditions as for Compound 153, Compound 42 was prepared. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.44 (d, *J* = 2.4 Hz, 1H), 7.81 (dd, *J* = 2.5, 8.4 Hz, 1H), 7.73 (s, 1H), 7.52 (d, *J* = 1.0 Hz, 1H), 7.42 (d, *J* = 8.4 Hz, 1H), 7.00 (d, *J* = 1.0 Hz, 1H), 4.84 - 4.80 (m, 1H), 3.48 (tt, *J* = 3.8, 11.7 Hz, 1H), 3.02 (tt, *J* =

3.5, 12.1 Hz, 1H), 2.29 - 2.21 (m, 2H), 2.12 - 2.05 (m, 2H), 1.72 (dq, $J = 3.0, 12.9$ Hz, 2H), 1.49 - 1.38 (m, 2H), 1.25 (br d, $J = 6.1$ Hz, 6H), 1.16 (s, 9H). ESI $[M+H] = 562.2$.

Example 129. Preparation of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-(1*H*-imidazol-2-ylamino)phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 56)




5

From isopropyl *trans*-*N*-[4-[5-[4-amino-2-(ethylsulfamoyl)phenyl]thiazol-2-yl] cyclohexyl]carbamate and 2-bromo-1*H*-imidazole, using General Method K. ^1H NMR (400 MHz, methanol- d_4) $\delta = 7.90$ (d, $J = 2.4$ Hz, 1H), 7.78 - 7.69 (m, 1H), 7.59 - 7.44 (m, 2H), 7.12 (s, 2H), 4.85 - 4.76 (m, 1H), 3.52 - 3.38 (m, 1H), 3.02 (tt, $J = 3.4, 12.0$ Hz, 1H), 2.93 - 2.80 (m, 2H), 2.34 - 2.14 (m, 2H), 2.13 - 2.02 (m, 2H), 1.77 - 1.63 (m, 2H), 1.41 (dq, $J = 3.1, 12.6$ Hz, 2H), 1.22 (br d, $J = 6.0$ Hz, 6H), 1.05 - 0.97 (m, 3H). ESI $[M+H] = 533.2$.


10

Example 130. Preparation of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[(1-isopropylimidazol-2-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 154)



15

a) Synthesis of 2-bromo-1-isopropyl-imidazole




20

To a mixture of 2-bromo-1*H*-imidazole (300 mg, 2 mol, 1 *eq.*), NaH (245 mg, 6 mmol, 60% in oil, 3 *eq.*) in THF (3 mL) was added 2-bromopropane (753 mg, 6 mmol, 3 *eq.*). The mixture was stirred at 25°C for 6 h under N_2 atmosphere. The reaction mixture was quenched by H_2O (10 mL) and extracted with EtOAc 60 mL (20 mL X 3). The combined organic layers were washed with sat. aq. NaCl 60 mL (12 mL X 5), dried over Na_2SO_4 ,


filtered and concentrated under reduced pressure. The residue was purified by prep-HPLC (column: Nano-micro Kromasil C18 100*30mm 5um; mobile phase: [water(0.1%TFA)-ACN]; B%: 1%-25%, 10min) to yield 2-bromo-1-isopropyl-imidazole (300 mg, 2 mmol, 78% yield) as yellow oil. ¹H NMR (400 MHz, CDCl₃) δ = 7.36 (d, J=1.76 Hz, 1H), 7.17 (s, 1H), 4.59 (spt, J=6.73 Hz, 1H), 1.50 (d, J=6.62 Hz, 6H). ESI [M+H] = 188.9/190.9.

b) Synthesis of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(1-isopropylimidazol-2-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 154**)




From isopropyl *trans*-N-[4-[5-[4-amino-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate and 2-bromo-1-isopropyl-imidazole, using General Method K. ¹H NMR (methanol-*d*₄, 400 MHz) δ = 7.87 (d, 1H, J=2.4 Hz), 7.7-7.8 (m, 1H), 7.5-7.6 (m, 1H), 7.4-7.5 (m, 2H), 7.23 (d, 1H, J=2.4 Hz), 4.63 (quin, 1H, J=6.8 Hz), 3.45 (br t, 1H, J=11.6 Hz), 3.1-3.2 (m, 1H), 3.0-3.1 (m, 1H), 2.8-2.9 (m, 2H), 2.24 (br d, 2H, J=12.6 Hz), 2.0-2.1 (m, 2H), 1.6-1.8 (m, 2H), 1.55 (d, 6H, J=6.6 Hz), 1.3-1.5 (m, 2H), 1.22 (br d, 6H, J=6.0 Hz), 0.9-1.0 (m, 3H). ESI [M+H] = 575.2.

Example 131. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(2-hydroxypyrimidin-4-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 156)




a) Synthesis of isopropyl *trans*-N-[4-[5-[4-[(2-chloropyrimidin-4-yl)amino]-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 155**)



To a solution of isopropyl *trans*-N-[4-[5-[4-amino-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (200 mg, 429 μ mol, 1.0 eq.) in i-PrOH (2 mL) was added 2,4-dichloropyrimidine (127 mg, 857 μ mol, 2.0 eq.) and HCl (12 M, one drop) and the mixture was stirred at 25°C for 1 h. The reaction mixture was then concentrated under reduced pressure to yield isopropyl *trans*-N-[4-[5-[4-[(2-chloropyrimidin-4-yl)amino]-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (350 mg, crude) as a yellow oil. 40 mg of crude was purified by prep-HPLC (column: Phenomenex Luna C18 150*30mm*5 μ m; mobile phase: [water(0.1%TFA)-ACN]; B%: 45%-65%, 10min) to give pure compound CYT-2056 (2.93 mg, 86.4% purity) as a gray solid. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.50 (d, J=2.20 Hz, 1H), 8.20 (d, J=5.87 Hz, 1H), 7.90-8.00 (m, 1H), 7.77 (s, 1H), 7.48-7.52 (m, 1H), 6.80 (d, J=5.99 Hz, 1H), 4.82-4.84 (m, 1H), 3.48 (ddd, J=3.55, 7.83, 11.74 Hz, 1H), 3.03 (q, J=7.30 Hz, 3H), 2.27 (br d, J=12.23 Hz, 2H), 2.05-2.15 (m, 2H), 1.73 (dq, J=3.12, 12.82 Hz, 2H), 1.44 (dq, J=3.36, 12.53 Hz, 2H), 1.25 (br d, J=6.11 Hz, 6H), 1.10 (t, J=7.21 Hz, 3H). ESI [M+H]⁺ = 579.2.


b) Synthesis of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(2-hydroxypyrimidin-4-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 156**)




A mixture of isopropyl *trans*-N-[4-[5-[4-[(2-chloropyrimidin-4-yl)amino]-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (250 mg, 432 μ mol, 1.0 eq.) in formic acid (2.5 mL) was stirred at 70°C for 12 h. The reaction mixture was concentrated under reduced pressure and then purified by prep-HPLC (column: Phenomenex Luna C18 200*40mm*10 μ m; mobile phase: [water(0.2%FA)-ACN]; B%: 1%-30%, 8min) to yield isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(2-hydroxypyrimidin-4-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (7 mg, 13 μ mol, 3% yield, 97% purity) as a white solid. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.82 (s, 1H), 7.94 (dd, J=2.14, 8.38

Hz, 1H), 7.74 (s, 1H), 7.59 (d, J=7.09 Hz, 1H), 7.48 (d, J=8.31 Hz, 1H), 6.09 (d, J=7.09 Hz, 1H), 4.76-4.83 (m, 1H), 3.42-3.55 (m, 1H), 2.93-3.08 (m, 3H), 2.26 (br d, J=12.23 Hz, 2H), 2.10 (br d, J=10.39 Hz, 2H), 1.65-1.79 (m, 2H), 1.37-1.51 (m, 2H), 1.25 (br d, J=6.11 Hz, 6H), 1.13 (t, J=7.21 Hz, 3H). ESI [M+H]⁺ = 561.2.

5 **Example 132. Preparation of isopropyl *trans*-N-[4-[5-[4-[(6-ethylpyridazin-3-yl) amino]-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 157)**




a) **Synthesis of 6-vinylpyridazin-3-amine**



10 A mixture of 4,4,5,5-tetramethyl-2-vinyl-1,3,2-dioxaborolane (1.1 g, 6.9 mmol, 1.2 *eq.*), 6-bromopyridazin-3-amine (1 g, 5.8 mmol, 1.0 *eq.*), Cs₂CO₃ (5.6 g, 17.3 mmol, 3.0 *eq.*) and Pd(dppf)Cl₂ (420 mg, 575 μmol, 0.1 *eq.*) in dioxane (20 mL)/H₂O (10 mL) was stirred at 80°C for 3 h under N₂ atmosphere. The reaction mixture was concentrated under reduced pressure and then purified by column chromatography (SiO₂, petroleum ether: ethyl acetate=10:1 to 0:1) to yield 6-vinylpyridazin-3-amine (0.3 g, 2.5 mmol, 43% yield) as a pale yellow solid. ESI [M+H]⁺ = 122.0.


b) **Synthesis of 6-ethylpyridazin-3-amine**



20 To a solution of 6-vinylpyridazin-3-amine (0.3 g, 2.5 mmol, 1.0 *eq.*) and AcOH (15 mg, 248 μmol, 0.1 *eq.*) in EtOAc (20 mL) was added Pd/C (100 mg, 10% purity) under N₂. The suspension was degassed under vacuum and purged with H₂ for 3 times and then stirred under H₂ (15 psi) at 25°C for 2 h. The reaction mixture was filtered, the filtrate was concentrated under reduced pressure and then purified by prep-HPLC (column: Phenomenex


Gemini-NX C18 75*30mm*3um; mobile phase: [water(10mM NH₄HCO₃)-ACN]; B%: 1%-15%, 10min) to yield 6-ethylpyridazin-3-amine (220 mg, 1.8 mmol, 72% yield) as a white solid. ¹H NMR (400 MHz, methanol-*d*₄) δ = 7.30 - 7.24 (m, 1H), 6.96 - 6.86 (m, 1H), 2.80 - 2.69 (m, 2H), 1.33 - 1.19 (m, 3H). ESI [M+H]⁺ = 124.2.

- 5 c) Synthesis of isopropyl *trans*-*N*-[4-[5-[4-[(6-ethylpyridazin-3-yl) amino]-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 157**)




- From isopropyl *trans*-*N*-[4-[5-[4-bromo-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl] carbamate and 6-ethylpyridazin-3-amine, using General Method F. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.61 - 8.53 (m, 1H), 8.11 - 8.02 (m, 1H), 7.95 - 7.88 (m, 1H), 7.87 - 7.79 (m, 1H), 7.79 - 7.74 (m, 1H), 7.56 - 7.49 (m, 1H), 4.86 (br d, *J* = 6.8 Hz, 1H), 3.53 - 3.41 (m, 1H), 3.12 - 3.00 (m, 3H), 2.94 - 2.84 (m, 2H), 2.32 - 2.19 (m, 2H), 2.15 - 1.94 (m, 2H), 1.78 - 1.57 (m, 2H), 1.53 - 1.37 (m, 5H), 1.29 - 1.18 (m, 6H), 1.09 - 0.92 (m, 3H). ESI [M+H]⁺ = 573.2.

- 15 **Example 133. Preparation of isopropyl *trans*-*N*-[4-[5-[4-[(5-ethylpyridazin-3-yl) amino]-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 158**)**




- a) Synthesis of 3,6-dichloro-4-ethyl-pyridazine




AgNO_3 (570 mg, 3.4 mmol, 0.1 *eq.*), propionic acid (3.0 g, 40.3 mmol, 1.2 *eq.*) and TFA (0.1 mL) were added in that order into a solution of 3,6-dichloropyridazine (5.0 g, 34.0 mmol, 1.0 *eq.*) in H_2O (50 mL). The reaction solution was heated to 70°C , then slowly added $(\text{NH}_4)_2\text{S}_2\text{O}_8$ (15.3 g, 68 mmol, 2.0 *eq.*). The reaction was stirred at 70°C for 12 h, then extracted with dichloromethane (90 mL, $30\text{ mL} \times 3$). The combined organic layers were dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO_2 , petroleum ether: ethyl acetate=10:1 to 5:1) to yield 3,6-dichloro-4-ethyl-pyridazine (3.1 g, 17.5 mmol, 52% yield) as a yellow oil which was used directly in the next step.

b) Synthesis of 6-chloro-*N*-[(2,4-dimethoxyphenyl)methyl]-5-ethyl-pyridazin-3-amine




To a solution of 3,6-dichloro-4-ethyl-pyridazine (3.1 g, 17.5 mmol, 1 *eq.*) in $n\text{-BuOH}$ (50 mL) was added (2,4-dimethoxyphenyl)methanamine (4.4 g, 26.3 mmol, 1.5 *eq.*) and K_2CO_3 (4.8 g, 35.0 mmol, 2.0 *eq.*). The mixture was stirred at 100°C for 12 h, and then concentrated under reduced pressure and purified by column chromatography (SiO_2 , petroleum ether: ethyl acetate=3:1 to 1:1) to yield 6-chloro-*N*-[(2,4-dimethoxyphenyl)methyl]-5-ethyl-pyridazin-3-amine (700 mg, 2 mmol, 13% yield) as a pale yellow oil.

c) Synthesis of *N*-[(2,4-dimethoxyphenyl)methyl]-5-ethyl-pyridazin-3-amine




A mixture of 6-chloro-*N*-[(2,4-dimethoxyphenyl)methyl]-5-ethyl-pyridazin-3-amine (600 mg, 2.0 mmol, 1.0 *eq.*), Pd/C (10 mg, 10% purity), TEA (20 mg, 195 μmol , 0.1 *eq.*) and HCOONH_4 (3.7 g, 58.5 mmol, 30 *eq.*) in EtOH (40 mL) was stirred at 80°C for 1 h under N_2 atmosphere. The reaction mixture was filtered, and the filtrate was concentrated under reduced pressure to remove solvent. The residue was diluted with H_2O (10 mL), extracted with EtOAc (60 mL, $20\text{ mL} \times 3$), dried over Na_2SO_4 , filtered and concentrated under reduced pressure to yield *N*-[(2,4-dimethoxy-phenyl)methyl]-5-ethyl-pyridazin-3-amine (700 mg, crude) as a yellow oil. ESI $[\text{M}+\text{H}] = 274.1$.

d) Synthesis of 5-ethylpyridazin-3-amine




A mixture of *N*-[(2,4-dimethoxyphenyl)methyl]-5-ethylpyridazin-3-amine (600 mg, 2 mmol, 1.0 *eq.*) in DCM (6 mL)/TFA (2 mL) was stirred at 25°C for 1 h. The reaction mixture was concentrated under reduced pressure and then purified by prep-HPLC (column: Phenomenex luna C18 250*50mm*10 um; mobile phase: [water(0.1%TFA)-ACN]; B%: 1%-20%, 10min) to yield 5-ethylpyridazin-3-amine (250 mg, 2 mmol, 92% yield) as a white solid. ¹H NMR (400MHz, methanol-*d*₄) δ = 8.34 (d, *J*=1.4 Hz, 1H), 7.26 (s, 1H), 2.77 (q, *J*=7.4 Hz, 2H), 1.32 (t, *J*=7.5 Hz, 3H). ESI [M+H]⁺ = 124.0.

10 e) Synthesis of isopropyl *trans*-*N*-[4-[5-[4-[(5-ethylpyridazin-3-yl)amino]-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 158**)




From isopropyl *trans*-*N*-[4-[5-[4-bromo-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl] carbamate and 5-ethylpyridazin-3-amine, using General Method F. ¹H NMR (400MHz, methanol-*d*₄) δ = 8.94 (s, 1H), 8.52 (d, *J*=2.4 Hz, 1H), 7.95 (dd, *J*=2.4, 8.4 Hz, 1H), 7.78 (s, 1H), 7.61 (s, 1H), 7.55 (d, *J*=8.4 Hz, 1H), 4.87 - 4.82 (m, 1H), 3.48 (tt, *J*=4.0, 11.6 Hz, 1H), 3.05 (tt, *J*=3.5, 12.0 Hz, 1H), 2.97 - 2.83 (m, 4H), 2.27 (br d, *J*=12.3 Hz, 2H), 2.10 (br d, *J*=10.3 Hz, 2H), 1.73 (dq, *J*=3.0, 12.8 Hz, 2H), 1.51 - 1.34 (m, 5H), 1.25 (br d, *J*=6.1 Hz, 6H), 1.05 (t, *J*=7.2 Hz, 3H). ESI [M+H]⁺ = 573.2.

20 **Example 134. Preparation of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[(6-oxo-1*H*-pyridazin-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 160)**




a) Synthesis of isopropyl *trans*-N-[4-[5-[4-[(6-chloropyridazin-3-yl)amino]-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 159**)



- 5 A mixture of isopropyl *trans*-N-[4-[5-[4-amino-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (500 mg, 1 mmol, 1.0 *eq.*), 3,6-dichloropyridazine (160 mg, 1 mmol, 1.0 *eq.*) in *i*-PrOH (5 mL) was stirred at 90°C for 12 h. The reaction mixture was concentrated under reduced pressure to remove solvent. The residue was triturated with EtOAc (10 mL) to yield isopropyl *trans*-N-[4-[5-[4-[(6-chloropyridazin-3-yl)amino]-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (200 mg, 345 μ mol, 32% yield) a yellow solid. ¹H NMR (400MHz, methanol-*d*₄) δ = 8.55 (d, *J*=2.4 Hz, 1H), 8.04 - 7.94 (m, 1H), 7.77 (s, 1H), 7.56 (d, *J*=9.5 Hz, 1H), 7.45 (d, *J*=8.6 Hz, 1H), 7.23 (d, *J*=9.3 Hz, 1H), 4.82 - 4.77 (m, 1H), 3.45 (br t, *J*=11.7 Hz, 1H), 3.09 - 3.00 (m, 1H), 2.95 (q, *J*=7.2 Hz, 2H), 2.24 (br d, *J*=13.5 Hz, 2H), 2.07 (br d, *J*=11.2 Hz, 2H), 1.78 - 1.60 (m, 2H), 1.48 - 1.35 (m, 2H), 1.22 (br d, *J*=6.2 Hz, 6H), 1.13 - 0.99 (m, 3H). ESI [M+H] = 579.2.


b) Synthesis of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(6-oxo-1H-pyridazin-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 160**)



A mixture of isopropyl *trans*-*N*-[4-[5-[4-[(6-chloropyridazin-3-yl)amino]-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (30 mg, 52 μ mol, 1 *eq.*) in AcOH (1 mL) was stirred at 100°C for 2 h and then concentrated under reduced pressure to remove AcOH. The residue was purified by SFC (column: (s,s) WHELK-O1

- 5 (250mm*30mm,5 μ m);mobile phase: [0.1%NH₃H₂O IPA];B%: 50%-50%,min), then by prep-HPLC (column: Phenomenex Luna C18 150*30mm*5 μ m;mobile phase: [water(0.1%TFA)-ACN];B%: 30%-60%,10min) to yield isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[(6-oxo-1*H*-pyridazin-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl] carbamate (1 mg, 2 μ mol, 3% yield, 97% purity) as a pale yellow solid. ¹H NMR (400MHz, methanol-*d*₄) δ = 8.38 (d, J=2.4
- 10 Hz, 1H), 7.82 (dd, J=2.5, 8.5 Hz, 1H), 7.72 - 7.65 (m, 1H), 7.37 (d, J=8.6 Hz, 1H), 7.28 (d, J=9.7 Hz, 1H), 6.95 (d, J=9.9 Hz, 1H), 4.83 - 4.73 (m, 1H), 3.45 (br t, J=11.6 Hz, 1H), 3.05 - 2.94 (m, 1H), 2.89 (q, J=7.4 Hz, 2H), 2.23 (br d, J=13.2 Hz, 2H), 2.07 (br d, J=10.6 Hz, 2H), 1.76 - 1.62 (m, 2H), 1.47 - 1.35 (m, 2H), 1.29 - 1.20 (m, 6H), 1.06 - 0.98 (m, 3H). ESI [M+H] = 561.2.

15 **Example 135. Preparation of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[(5-methylpyridazin-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 161)**




- From isopropyl *trans*-*N*-[4-[5-[4-bromo-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl] carbamate and 5-methylpyridazin-3-amine, using General Method F. ¹H NMR
- 20 (400 MHz, methanol-*d*₄) δ = 8.86 (br s, 1H), 8.46 (d, J = 2.2 Hz, 1H), 7.89 (dd, J = 2.2, 8.4 Hz, 1H), 7.76 (s, 1H), 7.59 - 7.49 (m, 2H), 4.84 - 4.78 (m, 1H), 3.45 (tt, J = 3.9, 11.6 Hz, 1H), 3.03 (tt, J = 3.3, 12.0 Hz, 1H), 2.89 (q, J = 7.3 Hz, 2H), 2.51 (s, 3H), 2.29 - 2.19 (m, 2H), 2.13 - 2.03 (m, 2H), 1.70 (dq, J = 3.0, 12.8 Hz, 2H), 1.47 - 1.33 (m, 2H), 1.22 (br d, J = 6.0 Hz, 6H), 1.02 (t, J = 7.3 Hz, 3H). ESI [M+H] = 559.3.

25 **Example 136. Preparation of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[(6-methylpyridazin-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 162)**


- Following the same protocol and under the same reaction conditions as for Compound 161, Compound 162 was prepared. ¹H NMR (400MHz, DMSO-*d*₆) δ = 10.08 (br s, 1H), 8.40 (d, J=2.3 Hz, 1H), 8.00 (dd, J=2.3, 8.5 Hz, 1H), 7.75 - 7.65 (m, 2H), 7.52 - 7.29 (m, 3H), 7.02
- 30 (br d, J=8.6 Hz, 1H), 3.34 - 3.30 (m, 1H), 3.34 - 3.30 (m, 1H), 2.99 - 2.78 (m, 3H), 2.58 (s,

3H), 2.60 - 2.56 (m, 2H), 2.22 - 2.10 (m, 2H), 1.67 - 1.50 (m, 2H), 1.45 - 1.27 (m, 2H), 1.17 (d, J=6.3 Hz, 6H), 0.97 (t, J=7.2 Hz, 3H). ESI [M+H] = 559.2.

Example 137. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(5-isopropylpyrazin-2-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 163)




a) Synthesis of 5-isopropenylpyrazin-2-amine



A mixture of 5-bromopyrazin-2-amine (1.0 g, 5.8 mmol, 1.0 *eq.*), 2-isopropenyl-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (1.3 g, 7.5 mmol, 1.3 *eq.*), Cs₂CO₃ (5.6 g, 17.2 mmol, 3.0 *eq.*) and Pd(dppf)Cl₂ (421 mg, 575 μmol, 0.1 *eq.*) in dioxane (20 mL)/H₂O (10 mL) was stirred at 80°C for 12 h under N₂ atmosphere. The reaction mixture was concentrated under reduced pressure to remove solvent, then the residue was diluted with H₂O (10 mL) and extracted with EtOAc (90 mL, 30 mL X 3). The combined organic layer was dried over Na₂SO₄, filtered and concentrated. The residue was purified by column chromatography (SiO₂, petroleum ether: ethyl acetate=10:1 to 5:1 to 1:1 to 0:1) to yield 5-isopropenylpyrazin-2-amine (750 mg, 97% yield) as a white solid. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.06 (s, 1 H), 7.90 (d, J=1.10 Hz, 1 H), 5.63 (d, J=0.66 Hz, 1 H), 5.00 - 5.13 (m, 1 H), 2.11 (s, 3 H). ESI [M+H] = 136.1.


b) Synthesis of 5-isopropylpyrazin-2-amine



To a solution of 5-isopropenylpyrazin-2-amine (490 mg, 4 mmol, 1.0 *eq.*) in EtOAc (20 mL) was added AcOH (22 mg, 363 μmol, 0.1 *eq.*) and Pd/C (100 mg, 10% purity). The


suspension was degassed under vacuum and purged with H₂ for 3 times. The mixture was stirred under H₂ (15 psi) at 25°C for 4 h, then filtered and concentrated under reduced pressure to yield 5-isopropylpyrazin-2-amine (500 mg, crude) as a pale yellow oil. ¹H NMR (400 MHz, methanol-*d*₄) δ = 7.87 (d, J=1.32 Hz, 1 H), 7.78 (d, J=1.10 Hz, 1 H), 2.91 (dt, J=13.89, 6.95 Hz, 1 H), 1.23 (d, J=7.06 Hz, 6 H). ESI [M+H]⁺=138.1.

c) Synthesis of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[(5-isopropylpyrazin-2-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 163**)




From isopropyl *trans*-*N*-[4-[5-[4-bromo-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl] carbamate and 5-isopropylpyrazin-2-amine, using General Method F. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.58 (d, J=2.38 Hz, 1 H), 8.16 (dd, J=8.88, 1.25 Hz, 2 H), 7.94 (dd, J=8.38, 2.38 Hz, 1 H), 7.74 (s, 1 H), 7.42 (d, J=8.38 Hz, 1 H), 4.78 - 4.86 (m, 1 H), 3.43 - 3.52 (m, 1 H), 3.00 - 3.10 (m, 2 H), 2.97 (q, J=7.21 Hz, 2 H), 2.22 - 2.30 (m, 2 H), 2.06 - 2.14 (m, 2 H), 1.72 (qd, J=12.84, 3.00 Hz, 2 H), 1.43 (qd, J=12.57, 3.31 Hz, 2 H), 1.33 (d, J=6.88 Hz, 6 H), 1.25 (br d, J=6.13 Hz, 6 H), 1.09 (t, J=7.25 Hz, 3 H). ESI [M+H]⁺=587.2.

Example 138. Preparation of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[(5-isopropylpyridazin-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 164)



a) Synthesis of 3,6-dichloro-4-isopropyl-pyridazine




AgNO_3 (57 mg, 336 μmol , 0.1 *eq.*), 2-methylpropanoic acid (355 mg, 4 mmol, 1.2 *eq.*) and TFA (0.1 mL) were added (in order) to a solution of 3,6-dichloropyridazine (500 mg, 3 mmol, 1.0 *eq.*) in H_2O (5 mL). The reaction solution was heated to 70°C , then slowly

5 added $(\text{NH}_4)_2\text{S}_2\text{O}_8$ (1.5 g, 6.7 mmol, 2.0 *eq.*). The reaction was stirred for 12 h at 70°C , then extracted with dichloromethane (90 mL, 30 mL X 3). The combined organic layers were dried over Na_2SO_4 , filtered and concentrated under reduced pressure to yield 3,6-dichloro-4-isopropylpyridazine (650 mg, crude) as a yellow oil. ESI $[\text{M}+\text{H}] = 191.0/193.0$.

b) Synthesis of 6-chloro-*N*-[(2,4-dimethoxyphenyl)methyl]-5-isopropylpyridazin-3-amine


10 amine



To a solution of (2,4-dimethoxyphenyl)methanamine (853 mg, 4.5 mmol, 1.5 *eq.*) and 3,6-dichloro-4-isopropylpyridazine (650 mg, 3 mmol, 1.0 *eq.*) in *n*-BuOH (30 mL) was added K_2CO_3 (1.4 g, 10.2 mmol, 3.0 *eq.*) and the mixture was stirred at 100°C for

15 12 h. The reaction mixture was concentrated and purified by column chromatography (SiO_2 , petroleum ether: ethyl acetate=10:1 to 0:1) to yield 6-chloro-*N*-[(2,4-dimethoxyphenyl)methyl]-5-isopropylpyridazin-3-amine (320 mg, 994 μmol , 29 % yield) as a light yellow oil which was used directly in the next step.

c) Synthesis of *N*-[(2,4-dimethoxyphenyl)methyl]-5-isopropylpyridazin-3-amine




To a solution of 6-chloro-*N*-[(2,4-dimethoxyphenyl)methyl]-5-isopropylpyridazin-3-amine (300 mg, 932 μmol , 1.0 *eq.*) and Pd/C (300 mg, 10% purity) in EtOH (10 mL) was added TEA (573 mg, 6 mmol, 6.0 *eq.*) and HCOONH_4 (1.8 g, 28.0 mmol, 30.0 *eq.*). The mixture was stirred at 80°C for 1 h, then filtered and concentrated. The residue was diluted

25 with H_2O (10 mL) and extracted with EtOAc (45 mL, 15 mL X 3). The combined organic layers were dried over Na_2SO_4 , filtered and concentrated under reduced pressure to yield *N*-


[(2,4-dimethoxyphenyl)methyl]-5-isopropyl-pyridazin-3-amine (250 mg, crude) as a yellow oil. ESI [M+H]⁺ = 287.9.

d) Synthesis of 5-isopropylpyridazin-3-amine




5 A solution of *N*-[(2,4-dimethoxyphenyl)methyl]-5-isopropyl-pyridazin-3-amine (230 mg, 800 μmol, 1.0 *eq.*) in DCM (20 mL) and TFA (7 mL) was stirred at 25°C for 1 h. The mixture was concentrated, diluted with H₂O (20 mL) and extracted with EtOAc (60 mL, 20 mL X 3). The combined organic layers were dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (column: Nano-micro Kromasil C18 100*40mm 10μm; mobile phase: [water(10mM NH₄HCO₃)-ACN]; B%: 1%-20%, 8min) to yield 5-isopropylpyridazin-3-amine (30 mg, 218 μmol, 27% yield) as a white solid. ¹H NMR (400MHz, methanol-*d*₄) δ = 8.42 (s, 1H), 7.25 (s, 1H), 3.04 (spt, J=6.9 Hz, 1H), 1.33 (d, J=6.8 Hz, 6H). ESI [M+H]⁺ = 138.0.

e) Synthesis of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[(5-isopropylpyridazin-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 164**)




From isopropyl *trans*-*N*-[4-[5-[4-bromo-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl] carbamate and 5-isopropylpyridazin-3-amine, using General Method F. ¹H NMR (400MHz, DMSO-*d*₆) δ = 9.90 (s, 1H), 8.84 (s, 1H), 8.40 (d, J=2.1 Hz, 1H), 8.07 (dd, J=2.1, 8.4 Hz, 1H), 7.69 (s, 1H), 7.47 (d, J=8.4 Hz, 1H), 7.39 (t, J=5.5 Hz, 1H), 7.11 (s, 1H), 7.02 (br d, J=7.3 Hz, 1H), 4.75 (td, J=6.2, 12.5 Hz, 1H), 3.32 (br s, 1H), 3.00 - 2.90 (m, 2H), 2.89 - 2.80 (m, 2H), 2.16 (br d, J=11.6 Hz, 2H), 1.93 (br d, J=10.0 Hz, 2H), 1.67 - 1.53 (m, 2H), 1.42 - 1.29 (m, 2H), 1.26 (d, J=6.9 Hz, 6H), 1.18 (d, J=6.3 Hz, 6H), 0.99 (t, J=7.2 Hz, 3H). ESI [M+H]⁺ = 587.3.

25 **Example 139. Preparation of methyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[(3-methyl-oxetan-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 165)**



a) Synthesis of methyl *trans*-*N*-[4-[5-[4-bromo-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate




To a solution of methyl *trans*-*N*-[4-[5-[4-amino-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (2.5 g, 5.7 mmol, 1.0 *eq.*) in MeCN (50 mL) was added CuBr (981 mg, 7 mmol, 1.2 *eq.*), [(1*S*,4*R*)-7,7-dimethyl-2-oxo-norbornan-1-yl]methanesulfonic acid (4.0 g, 17.1 mmol, 3.0 *eq.*), TBAB (11.0 g, 34.2 mmol, 6.0 *eq.*) and NaNO₂ (1.2 g, 17.1 mmol, 3 *eq.*). The reaction mixture was stirred at 25°C for 1 h, then quenched with

with Na₂CO₃ sat.aq. (50 mL), and concentrated under reduced pressure to remove MeCN. The residue was diluted with H₂O (50 mL) and extracted with EtOAc (300 mL, 100 mL X 3), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by reversed-phase HPLC(0.1% TFA condition) to yield methyl *trans*-*N*-[4-[5-[4-bromo-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (1.4 g, 2.8 mmol, 48.9 %

yield) as a yellow solid. ¹H NMR (400MHz, methanol-*d*₄) δ = 8.21 (d, J=1.6 Hz, 1H), 7.93 - 7.71 (m, 2H), 7.43 (d, J=8.1 Hz, 1H), 3.65 (s, 3H), 3.48 (br t, J=11.6 Hz, 1H), 3.04 (br s, 1H), 2.89 (q, J=7.1 Hz, 2H), 2.26 (br d, J=12.5 Hz, 2H), 2.09 (br d, J=11.0 Hz, 2H), 1.72 (q, J=11.9 Hz, 2H), 1.51 - 1.37 (m, 2H), 1.04 (t, J=7.2 Hz, 3H). ESI [M+H]⁺ = 502.1/504.1.

b) Synthesis of methyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[(3-methyl-oxetan-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 165**)



From methyl *trans*-N-[4-[5-[4-bromo-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl] carbamate and 3-methyloxetan-3-amine, using General Method F. ¹H NMR (400MHz, methanol-*d*₄) δ = 7.60 (s, 1H), 7.20 (d, J=8.4 Hz, 1H), 7.11 (d, J=2.4 Hz, 1H), 6.60 (dd, J=2.5, 8.3 Hz, 1H), 4.79 (d, J=6.2 Hz, 2H), 4.59 (d, J=6.0 Hz, 2H), 3.62 (s, 3H), 3.50 - 3.39 (m, 1H), 3.03 - 2.92 (m, 1H), 2.85 (q, J=7.1 Hz, 2H), 2.21 (br d, J=12.8 Hz, 2H), 2.06 (br d, J=10.8 Hz, 2H), 1.74 - 1.60 (m, 5H), 1.40 (dq, J=3.1, 12.5 Hz, 2H), 1.01 (t, J=7.3 Hz, 3H). ESI [M+H] = 509.2.


Example 140. Preparation of methyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(1H-pyrazol-4-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 166)

Following the same protocol and under the same reaction conditions as for Compound 165, Compound 166 was prepared. ¹H NMR (400MHz, methanol-*d*₄) δ = 7.71 (s, 1H), 7.64 (s, 2H), 7.47 (d, J=2.4 Hz, 1H), 7.27 (d, J=8.4 Hz, 1H), 7.00 (dd, J=2.5, 8.4 Hz, 1H), 3.65 (s, 3H), 3.48 (ddd, J=4.0, 7.5, 11.5 Hz, 1H), 3.10 - 2.99 (m, 1H), 2.88 (q, J=7.3 Hz, 2H), 2.25 (br d, J=13.0 Hz, 2H), 2.09 (br d, J=10.5 Hz, 2H), 1.78 - 1.65 (m, 2H), 1.50 - 1.37 (m, 2H), 1.03 (t, J=7.3 Hz, 3H). ESI [M+H] = 505.2.


Example 141. Preparation of methyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(1H-pyrazol-5-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 167)

Following the same protocol and under the same reaction conditions as for Compound 165, Compound 167 was prepared. ¹H NMR (400MHz, methanol-*d*₄) δ = 8.00 (d, J=1.5 Hz, 1H), 7.87 (s, 1H), 7.75 - 7.62 (m, 1H), 7.52 - 7.42 (m, 1H), 7.36 (d, J=8.3 Hz, 1H), 6.11 (br s, 1H), 3.65 (s, 3H), 3.56 - 3.42 (m, 1H), 3.13 (br t, J=11.8 Hz, 1H), 2.95 (q, J=7.2 Hz, 2H), 2.27 (br d, J=12.3 Hz, 2H), 2.10 (br d, J=10.8 Hz, 2H), 1.82 - 1.65 (m, 2H), 1.54 - 1.36 (m, 2H), 1.07 (t, J=7.2 Hz, 3H). ESI [M+H] = 505.2.

Example 142. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(3*R*)-1-isobutyl-2-oxo-azetidin-3-yl]amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 168)




a) Synthesis of benzyl *N*-[(1*R*)-1-(hydroxymethyl)-2-(4-methoxyanilino)-2-oxo-ethyl]carbamate



- 5 To a solution of 4-methoxyaniline (25.7 g, 208.7 mmol, 1.0 *eq.*) in THF (300 mL) and DCM (100 mL) was added (2*R*)-2-(benzyloxycarbonylamino)-3-hydroxypropanoic acid (10.0 g, 41.7 mmol, 0.2 *eq.*) and EDCI (8.0 g, 41.7 mmol, 0.2 *eq.*) at 0°C for 30 min. The mixture was stirred at 25°C for 12 h and concentrated under reduced pressure. The residue was purified by column chromatography (SiO₂, petroleum ether: ethyl acetate = 10:1 to 0:1)
- 10 to yield benzyl *N*-[(1*R*)-1-(hydroxymethyl)-2-(4-methoxy-anilino)-2-oxo-ethyl]carbamate (8.0 g, 23.2 mmol, 11% yield) as a white solid. ¹H NMR (400 MHz, methanol-*d*₄) δ = 6.73 (d, *J* = 1.1 Hz, 9H), 4.86 - 4.84 (m, 2H), 4.39 - 4.31 (m, 1H), 3.86 (br d, *J* = 5.4 Hz, 2H), 3.80 - 3.75 (m, 3H). ESI [M+H] = 345.0.


b) Synthesis of benzyl *N*-[(3*R*)-1-(4-methoxyphenyl)-2-oxo-azetidin-3-yl]carbamate



15


To a solution of benzyl *N*-[(1*R*)-1-(hydroxymethyl)-2-(4-methoxyanilino)-2-oxo-ethyl]carbamate (7.0 g, 20.3 mmol, 1.0 *eq.*) in DMF (200 mL) was added 1-imidazol-1-ylsulfonylimidazole (6.0 g, 30.5 mmol, 1.5 *eq.*) at 0°C under N₂ atmosphere. The resulting mixture was stirred at 0°C for additional 0.5 h and then cooled to -20°C. Under vigorous stirring, NaH (1.2 g, 30.5 mmol, 60% purity, 1.5 *eq.*) was added portionwise. The resulting suspension was vigorously stirred at -20°C for additional 1 h. The reaction mixture was quenched with water (200 mL). The collected precipitate was filtered, washed with additional water (100 mL) and then dried under reduced pressure to yield benzyl *N*-[(3*R*)-1-(4-methoxyphenyl)-2-oxo-azetidin-3-yl]carbamate (4.5 g, 13.8 mmol, 68% yield) as a pale yellow solid. ¹H NMR (400 MHz, CDCl₃) δ = 7.32 - 7.08 (m, 7H), 6.83 - 6.72 (m, 2H), 5.70 - 5.60 (m, 1H), 5.10 - 5.01 (m, 2H), 4.96 - 4.87 (m, 1H), 3.89 - 3.78 (m, 1H), 3.70 (s, 3H), 3.53 - 3.43 (m, 1H). ESI [M+H] = 327.1.

c) Synthesis of benzyl *N*-[(3*R*)-2-oxoazetidin-3-yl]carbamate



To a solution of benzyl *N*-[(3*R*)-1-(4-methoxyphenyl)-2-oxo-azetidin-3-yl]carbamate (2.3 g, 7.1 mmol, 1.0 *eq.*) in MeCN (110 mL)/H₂O (88 mL) was added CAN (11.6 g, 21.0 mmol, 3.0 *eq.*) at 0°C. The reaction mixture was stirred at 0°C for 15 min, then quenched by addition of sat. aq. NaHCO₃ (110 mL) at 0°C. The mixture was filtered, and the filtrate was extracted with EtOAc (120 mL, 40 mL X 3). The combined organic layers dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by prep-HPLC (column: Phenomenex luna C18 250*50mm*10 um; mobile phase: [water(0.1%TFA)-ACN]; B%: 10%-40%, 10min) to yield benzyl *N*-[(3*R*)-2-oxoazetidin-3-yl]carbamate (890 mg, 4 mmol, 57% yield) as a purple solid. ¹H NMR (400 MHz, CDCl₃) δ = 7.45 - 7.33 (m, 5H), 6.08 - 5.93 (m, 1H), 5.68 - 5.52 (m, 1H), 5.25 - 5.08 (m, 2H), 4.97 - 4.82 (m, 1H), 3.76 - 3.57 (m, 1H), 3.46 - 3.34 (m, 1H). ESI [M+H] = 221.1.


d) Synthesis of (3*R*)-3-aminoazetidin-2-one



To a solution of benzyl *N*-[(3*R*)-2-oxoazetidin-3-yl]carbamate (890 mg, 4 mmol, 1 *eq.*) in THF (50 mL) was added Pd(OH)₂ (50 mg, 10% purity) and Pd/C (50 mg, 10% purity). The suspension was stirred under H₂ (50 psi) at 50°C for 4 h, then filtered and concentrated under reduced pressure to yield (3*R*)-3-aminoazetidin-2-one (500 mg, crude) as a purple oil.


5 ESI [2M+H] = 173.1.

e) Synthesis of (3*R*)-3-(dibenzylamino)azetidin-2-one



To a solution of (3*R*)-3-aminoazetidin-2-one (500 mg, 6 mmol, 1.0 *eq.*), benzaldehyde (555 mg, 5 mmol, 0.9 *eq.*) in MeOH (20 mL) was added AcOH (34 mg, 580 μmol, 0.1 *eq.*) and NaBH₃CN (1.1 g, 17.4 mmol, 3 *eq.*). The mixture was stirred at 25°C for 12 h and then quenched with NH₃·H₂O (5 mL). The mixture was concentrated under reduced pressure and then purified by column chromatography (SiO₂, Petroleum ether: Ethyl acetate=10:1 to 1:1) to yield (3*R*)-3-(dibenzylamino)azetidin-2-one (620 mg, crude) as a yellow oil. ¹H NMR (400 MHz, CDCl₃) δ = 7.44 - 7.29 (m, 10H), 6.58 - 6.42 (m, 1H), 4.44 - 4.31 (m, 2H), 4.26 (br d, J = 2.5 Hz, 1H), 4.11 (br d, J = 13.1 Hz, 2H), 3.28 (br d, J = 5.4 Hz, 1H), 3.11 - 3.04 (m, 1H). ESI [M+H] = 267.1.


f) Synthesis of (3*R*)-3-(dibenzylamino)-1-isobutylazetidin-2-one



To a solution of (3*R*)-3-(dibenzylamino)azetidin-2-one (600 mg, 2 mmol, 1 *eq.*) in DMF (2 mL) was added NaH (135 mg, 3 mmol, 60% in oil, 1.5 *eq.*) and 1-bromo-2-methylpropane (926 mg, 7 mmol, 3 *eq.*). The mixture was stirred at 25°C for 1 h and then quenched with NH₄Cl (10 mL) at 0°C. The mixture was diluted with H₂O (20 mL) and extracted with EtOAc (60 mL, 20 mL X 3). The combined organic layers were washed with brine (30 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by prep-HPLC (column: Nano-micro Kromasil C18 100*40mm 10μm; mobile phase: [water(0.1%TFA)-ACN]; B%: 40%-67%, 8min) to yield (3*R*)-3-(dibenzylamino)-1-isobutylazetidin-2-one (220 mg, 682 μmol, 30% yield) as a yellow oil. ¹H NMR (400 MHz, CDCl₃) δ = 7.53 - 7.23 (m, 10H), 4.54 - 4.37 (m, 2H), 4.34 - 4.21 (m, 1H),


4.13 (br d, $J = 13.0$ Hz, 2H), 3.39 - 3.32 (m, 1H), 3.12 - 3.02 (m, 1H), 2.99 - 2.87 (m, 2H), 1.79 - 1.68 (m, 1H), 0.90 - 0.76 (m, 6H). ESI $[M+H]^+ = 323.1$.

g) Synthesis of (3*R*)-3-amino-1-isobutyl-azetidin-2-one



5 To a solution of (*R*)-3-(dibenzylamino)-1-isobutylazetidin-2-one (220 mg, 682 μ mol, 1 *eq.*) in THF (50 mL) and *i*-PrOH (50 mL) was added $Pd(OH)_2$ (20 mg, 10%) and Pd/C (20 mg, 10%). The suspension was stirred under H_2 (50 psi) at 50°C for 12 h. The reaction mixture was filtered and concentrated under reduced pressure to yield (3*R*)-3-amino-1-isobutyl-azetidin-2-one (210 mg, crude) as a black gum. ESI $[2M+H]^+ = 285.2$.

10 h) Synthesis of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[(3*R*)-1-iso-butyl-2-oxo-azetidin-3-yl]amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 168**)




From isopropyl *trans*-*N*-[4-[5-[4-bromo-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl] carbamate and (3*R*)-3-amino-1-isobutyl-azetidin-2-one, using General Method F. 1H NMR (400 MHz, methanol- d_4) $\delta = 7.75$ - 7.63 (m, 1H), 7.44 - 7.36 (m, 1H), 7.33 - 7.23 (m, 1H), 6.98 - 6.87 (m, 1H), 4.91 - 4.87 (m, 1H), 3.81 - 3.75 (m, 1H), 3.53 - 3.42 (m, 1H), 3.38 - 3.36 (m, 1H), 3.29 (dd, $J = 2.0, 5.7$ Hz, 1H), 3.18 - 3.08 (m, 2H), 3.07 - 2.96 (m, 1H), 2.94 - 2.83 (m, 2H), 2.25 (br d, $J = 12.5$ Hz, 2H), 2.14 - 2.05 (m, 2H), 2.01 - 1.90 (m, 1H), 1.79 - 1.60 (m, 2H), 1.49 - 1.36 (m, 2H), 1.25 (br d, $J = 6.1$ Hz, 6H), 1.04 (t, $J = 7.2$ Hz, 3H), 0.99 (dd, $J = 3.8, 6.7$ Hz, 6H). ESI $[M+H]^+ = 592.2$.

Example 143. Preparation of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[(3*S*)-1-isobutyl-2-oxo-azetidin-3-yl]amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 169**)**


Following the same protocol and under the same reaction conditions as for Compound 168, Compound 169 was prepared. 1H NMR (400 MHz, methanol- d_4) $\delta = 7.66$ (s, 1H), 7.39 (d, $J = 2.5$ Hz, 1H), 7.26 (d, $J = 8.4$ Hz, 1H), 6.93 (dd, $J = 2.4, 8.4$ Hz, 1H), 4.81 (br s, 1H), 3.79 (t, $J = 5.2$ Hz, 1H), 3.52 - 3.42 (m, 1H), 3.37 (br s, 1H), 3.29 (dd, $J = 2.0, 5.6$ Hz, 1H), 3.18 - 3.07 (m, 2H), 3.07 - 2.97 (m, 1H), 2.89 (q, $J = 7.2$ Hz, 2H), 2.25 (br d, $J = 12.3$ Hz,

2H), 2.09 (br d, $J = 10.3$ Hz, 2H), 2.01 - 1.91 (m, 1H), 1.76 - 1.63 (m, 2H), 1.48 - 1.37 (m, 2H), 1.25 (br d, $J = 6.0$ Hz, 6H), 1.08 - 0.96 (m, 9H). ESI $[M+H] = 592.3$.

Example 144. Preparation of *tert*-butyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[(2-isopropylpyrazol-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 170)




a) Synthesis of *tert*-butyl *trans*-*N*-[4-[5-[4-amino-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate



A mixture of 5-amino-*N*-ethyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzenesulfonamide (2.8 g, 8.5 mmol, 1.4 eq), *tert*-butyl *trans*-*N*-[4-(5-bromo-thiazol-2-yl)cyclohexyl]carbamate (2.2 g, 6.1 mmol, 1 eq), KF (530 mg, 9 mmol, 1.5 eq), Na₂CO₃ (1.9 g, 18.3 mmol, 3 eq) and Pd(PPh₃)₄ (703 mg, 609 μmol, 0.1 eq) in toluene (30 mL), EtOH (30 mL) and H₂O (10 mL) was stirred at 80°C for 2 h under N₂ atmosphere. The reaction mixture was concentrated under reduced pressure, then diluted with H₂O (20 mL) and extracted with EtOAc (90 mL, 30 mL X 3), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue which was purified by reversed-phase HPLC (0.1% TFA condition) to yield *tert*-butyl *trans*-*N*-[4-[5-[4-amino-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (2.9 g, 6.0 mmol, 99% yield) as a brown solid. ¹H NMR (400MHz, methanol-*d*₄) $\delta = 7.64$ (s, 1H), 7.60 (br s, 1H), 7.56 (d, $J=3.0$ Hz, 1H), 7.47 (d, $J=3.4$ Hz, 1H), 3.46 - 3.39 (m, 1H), 3.37 (s, 2H), 3.02 (tt, $J=3.5, 12.1$ Hz, 1H), 2.19 (br d, $J=13.5$ Hz, 2H),

2.05 (br d, $J=10.1$ Hz, 2H), 1.66 (dt, $J=10.1, 12.9$ Hz, 2H), 1.46 (s, 9H), 1.45 - 1.34 (m, 3H), 1.23 - 0.89 (m, 2H). ESI $[M+H]^+ = 481.2$.


b) Synthesis of *tert*-butyl *trans*-*N*-[4-[5-[4-bromo-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate



To a solution of *tert*-butyl *trans*-*N*-[4-[5-[4-amino-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (1.0 g, 2.1 mmol, 1.0 *eq.*) in MeCN (30 mL) was added TBAB (4.0 g, 12.5 mmol, 6.0 *eq.*), CuBr (358 mg, 2.5 mmol, 1.2 *eq.*), [(1*S*,4*R*)-7,7-dimethyl-2-oxo-norbornan-1-yl]methanesulfonic acid (1.5 g, 6.0 mmol, 3.0 *eq.*) and NaNO₂ (430 mg, 6 mmol, 3 *eq.*). The mixture was stirred at 25°C for 1 h, then quenched with sat. aq. Na₂CO₃ (20 mL). The mixture was concentrated, diluted with H₂O (20 mL) and extracted with EtOAc (150 mL, 50 mL X 3), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue which was purified by column chromatography (SiO₂, petroleum ether: ethyl acetate = 3:1 to 1:1) to yield *tert*-butyl *trans*-*N*-[4-[5-[4-bromo-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl] carbamate (500 mg, 918 μ mol, 44% yield) as a yellow oil. ¹H NMR (400MHz, methanol-*d*₄) δ = 8.18 (d, $J=2.0$ Hz, 1H), 7.80 (dd, $J=2.0, 8.2$ Hz, 1H), 7.74 (s, 1H), 7.40 (d, $J=8.2$ Hz, 1H), 3.39 (br t, $J=11.7$ Hz, 1H), 2.99 (tt, $J=3.5, 12.0$ Hz, 1H), 2.85 (q, $J=7.3$ Hz, 2H), 2.23 (br d, $J=12.8$ Hz, 2H), 2.10 - 1.99 (m, 2H), 1.75 - 1.58 (m, 2H), 1.44 (s, 9H), 1.41 - 1.32 (m, 2H), 1.01 (t, $J=7.3$ Hz, 3H). ESI $[M+H]^+ = 544.0/546.0$.

10
15
20

c) Synthesis of *tert*-butyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[(2-isopropyl-pyrazol-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 170**)




From *tert*-butyl *trans*-*N*-[4-[5-[4-bromo-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate and 3-amino-2-isopropylpyrazole, using General Method F. ¹H NMR (400MHz, methanol-*d*₄) δ = 7.77 (s, 1H), 7.63 (d, $J=1.8$ Hz, 1H), 7.50 (d, $J=2.5$ Hz, 1H), 7.35


25

(d, $J=8.4$ Hz, 1H), 7.05 (dd, $J=2.5, 8.4$ Hz, 1H), 6.19 (d, $J=2.0$ Hz, 1H), 4.63 (spt, $J=6.6$ Hz, 1H), 3.42 (br t, $J=11.4$ Hz, 1H), 3.12 - 3.01 (m, 1H), 2.87 (q, $J=7.2$ Hz, 2H), 2.26 (br d, $J=12.9$ Hz, 2H), 2.08 (br d, $J=10.5$ Hz, 2H), 1.79 - 1.65 (m, 2H), 1.48 - 1.44 (m, 15H), 1.44 - 1.35 (m, 2H), 1.02 (t, $J=7.3$ Hz, 3H). ESI $[M+H]^+ = 589.3$.

5 **Example 145. Preparation of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(6-isopropylpyrimidin-4-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 171)**



a) Synthesis of 6-isopropenylpyrimidin-4-amine




10

A mixture of 2-isopropenyl-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (1.2 g, 7.1 mmol, 1.3 *eq.*), 6-bromopyrimidin-4-amine (950 mg, 5 mmol, 1 *eq.*), Cs_2CO_3 (5.3 g, 16.4 mmol, 3 *eq.*) and $\text{Pd}(\text{dppf})\text{Cl}_2$ (399 mg, 546 μmol , 0.1 *eq.*) in H_2O (15 mL)/dioxane (30 mL) was stirred at 90°C for 12 h under N_2 atmosphere. The reaction mixture was concentrated under reduced pressure to remove solvent. The residue was diluted with H_2O (20 mL) and extracted with EtOAc (60 mL, 20 mL X 3). The combined organic layers were dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give a residue which was purified by column chromatography (SiO_2 , petroleum ether: ethyl acetate=1:0 to 0:1) to yield 6-isopropenylpyrimidin-4-amine (330 mg, 2 mmol, 45% yield) as a pale yellow solid. ^1H NMR (400 MHz, methanol- d_4) δ = 8.19 (d, $J = 0.9$ Hz, 1H), 6.51 (d, $J = 1.0$ Hz, 1H), 5.84 (d, $J = 0.8$ Hz, 1H), 5.25 (t, $J = 1.5$ Hz, 1H), 2.00 (s, 3H). ESI $[M+H]^+ = 136.0$.

15


20

b) Synthesis of 6-isopropylpyrimidin-4-amine



To a solution of 6-isopropenylpyrimidin-4-amine (330 mg, 2 mmol, 1 *eq.*) in EtOAc (20 mL) was added Pd/C (50 mg, 10% purity) and the suspension was stirred under H₂ (15 psi) at 25°C for 2 h. The reaction mixture was filtered, and the filtrate was concentrated under reduced pressure to yield 6-isopropylpyrimidin-4-amine (320 mg, 2 mmol, 96% yield) as a white solid. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.15 (d, *J* = 0.8 Hz, 1H), 6.31 (d, *J* = 0.6 Hz, 1H), 2.67 (td, *J* = 6.9, 13.9 Hz, 1H), 1.13 (d, *J* = 7.0 Hz, 6H). ESI [M+H]⁺ = 138.0.

c) Synthesis of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[(6-isopropylpyrimidin-4-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 171**)




10

From isopropyl *trans*-*N*-[4-[5-[4-bromo-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate and 6-isopropylpyrimidin-4-amine, using General Method F. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.82 (s, 1H), 8.52 (d, *J* = 2.2 Hz, 1H), 7.97 (dd, *J* = 2.2, 8.4 Hz, 1H), 7.76 (s, 1H), 7.55 (d, *J* = 8.4 Hz, 1H), 6.91 (s, 1H), 4.83 - 4.77 (m, 1H), 3.50 - 3.39 (m, 1H), 3.12 - 2.96 (m, 2H), 2.89 (q, *J* = 7.2 Hz, 2H), 2.31 - 2.16 (m, 2H), 2.13 - 2.00 (m, 2H), 1.76 - 1.63 (m, 2H), 1.38 (d, *J* = 6.8 Hz, 8H), 1.22 (br d, *J* = 6.2 Hz, 6H), 1.02 (t, *J* = 7.3 Hz, 3H). ESI [M+H]⁺ = 587.3.


15

Example 146. Preparation of isopropyl *trans*-*N*-[4-[5-[2-(ethylsulfamoyl)-4-[(1-methyl-6-oxo-pyridazin-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 172)




20

a) Synthesis of 6-imino-2-methyl-1*H*-pyridazin-3-one




To a solution of 6-imino-1,2-dihydropyridazin-3-one (250 mg, 2 mmol, 1 *eq.*) in EtOH (5 mL) was added NaOH (180 mg, 5 mmol, 2 *eq.*) and MeI (351 mg, 2 mmol, 1 *eq.*). The mixture was stirred at 80°C for 2 h and then quenched with NH₃·H₂O (0.1 mL) at 25°C. The mixture was purified by prep-TLC (SiO₂, dichloromethane : methanol = 8:1) and further purified by prep-HPLC (column: Waters Xbridge BEH C18 100*25mm*5um; mobile phase: [water(10mM NH₄HCO₃)-ACN]; B%: 1%-12%, 10min) to yield 6-imino-2-methyl-1H-pyridazin-3-one (120 mg, 959 umol, 43% yield) as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ = 6.91 (d, J = 9.7 Hz, 1H), 6.71 (d, J = 9.5 Hz, 1H), 5.73 (s, 2H), 3.39 (s, 3H). ESI [2M+H] = 251.1.

b) Synthesis of isopropyl *trans*-N-[4-[5-[2-(ethylsulfamoyl)-4-[(1-methyl-6-oxo-pyridazin-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (**Compound 172**)




From isopropyl *trans*-N-[4-[5-[4-bromo-2-(ethylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate and 6-imino-2-methyl-1H-pyridazin-3-one, using General Method F. ¹H NMR (400 MHz, methanol-*d*₄) δ = 8.47 (d, J = 2.4 Hz, 1H), 7.80 (dd, J = 2.3, 8.3 Hz, 1H), 7.72 (s, 1H), 7.40 (d, J = 8.4 Hz, 1H), 7.24 (d, J = 9.7 Hz, 1H), 6.95 (d, J = 9.7 Hz, 1H), 4.82 - 4.77 (m, 1H), 3.74 (s, 3H), 3.54 - 3.38 (m, 1H), 3.09 - 2.97 (m, 1H), 2.95 - 2.84 (m, 2H), 2.30 - 2.18 (m, 2H), 2.11 - 2.01 (m, 2H), 1.77 - 1.62 (m, 2H), 1.49 - 1.34 (m, 2H), 1.22 (br d, J = 6.2 Hz, 6H), 1.03 (t, J = 7.2 Hz, 3H). ESI [M+H] = 575.3.

Example 147. Preparation of isopropyl *trans*-N-[4-[5-[2-(*tert*-butylsulfamoyl)-4-[(6-chloro pyridazin-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 173)



To a solution of isopropyl *trans*-N-[4-[5-[4-amino-2-(*tert*-butylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (20 mg, 40 μ mol, 1 eq) in *i*-PrOH (2 mL) was added 3,6-dichloropyridazine (18 mg, 121 μ mol, 3 eq) and the mixture was stirred at 90°C for 1 h. The reaction mixture was concentrated under reduced pressure and purified by prep-HPLC (column: Welch Xtimate C18 150*25mm*5 μ m; mobile phase: [water(0.04% HCl)-ACN]; B%: 50%-80%, 10 min) to yield isopropyl *trans*-N-[4-[5-[2-(*tert*-butylsulfamoyl)-4-[(6-chloropyridazin-3-yl)amino]phenyl]thiazol-2-yl]cyclohexyl]carbamate (14 mg, 23 μ mol, 57 % yield, 96 % purity) as a white solid. ^1H NMR (400 MHz, methanol- d_4) δ = 8.65 - 8.54 (m, 1H), 8.12 - 8.03 (m, 1H), 7.89 - 7.81 (m, 1H), 7.60 - 7.54 (m, 1H), 7.50 - 7.43 (m, 1H), 7.27 - 7.18 (m, 1H), 4.84 - 4.80 (m, 1H), 3.53 - 3.41 (m, 1H), 3.15 - 3.05 (m, 1H), 2.30 - 2.22 (m, 2H), 2.14 - 2.04 (m, 2H), 1.80 - 1.65 (m, 2H), 1.50 - 1.35 (m, 2H), 1.22 (br d, J = 6.2 Hz, 6H), 1.17 - 1.15 (m, 9H). ESI $[\text{M}+\text{H}]^+ = 607.2/609.2$.

Example 148. Preparation of isopropyl *trans*-N-[4-[5-[2-(*tert*-butylsulfamoyl)-4-(pyridazin-3-ylamino)phenyl]thiazol-2-yl]cyclohexyl]carbamate (Compound 174)



A mixture of 3-bromopyridazine (13 mg, 81 μ mol, 2 eq), isopropyl *trans*-N-[4-[5-[4-amino-2-(*tert*-butylsulfamoyl)phenyl]thiazol-2-yl]cyclohexyl]carbamate (20 mg, 40 μ mol, 1 eq), $\text{Pd}_2(\text{dba})_3$ (4 mg, 4 μ mol, 0.1 eq), Cs_2CO_3 (40 mg, 121 μ mol, 3 eq) and Xantphos (2 mg, 4 μ mol, 0.1 eq) in dioxane (2 mL) was stirred at 110°C for 12 h under N_2 atmosphere. The reaction mixture was concentrated and purified by prep-HPLC (column: Nano-micro Kromasil C18 100*40mm 3 μ m; mobile phase: [water (0.1% TFA)-ACN]; B%: 30%-60%, 8 min) to yield isopropyl *trans*-N-[4-[5-[2-(*tert*-butylsulfamoyl)-4-(pyridazin-3-ylamino)phenyl]thiazol-2-yl]cyclohexyl]carbamate (4 mg, 6 μ mol, 15% yield, 100% purity) as a yellow solid. ^1H NMR (400 MHz, methanol- d_4) δ = 8.89 (br d, J = 4.50 Hz, 1 H),

8.61 (d, J=2.13 Hz, 1 H), 8.00 (dd, J=8.38, 2.00 Hz, 1 H), 7.90 (br dd, J=9.07, 4.82 Hz, 1 H), 7.79 (s, 1 H), 7.63 (br d, J=9.13 Hz, 1 H), 7.51 (d, J=8.38 Hz, 1 H), 4.78 (br s, 1 H), 3.44 - 3.52 (m, 1 H), 3.05 (ddd, J=11.94, 8.69, 3.50 Hz, 1 H), 2.26 (br d, J=11.88 Hz, 2 H), 2.10 (br d, J=10.13 Hz, 2 H), 1.67 - 1.79 (m, 2 H), 1.38 - 1.49 (m, 2 H), 1.25 (br d, J=6.00 Hz, 6 H), 1.13 (s, 9 H). ESI [M+H]⁺ = 573.2

Example 149. Compound Primary Screening

Primary screening was a phenotypic screen that utilized the synthetic lethal interaction between AID and RAD51 to identify compounds that were both potent and on target. AID expressing cells are dependent upon RAD51 for survival; inhibiting RAD51 in AID positive cells results in a cytotoxic effect. Based on such an effect, compounds that were potent in AID positive cells and were significantly less potent in AID negative cells were identified.

Materials and Supplies

Plastic ware and consumables needed for the experiment include: cell culture media; Evaporation Buffer media; 100% DMSO; 96 well U-bottom sterile culture plates; 250 mL bottle; 1.5 mL Opaque amber epi tubes; Epi Tube rack; 300 mL reservoirs; 25 mL reservoir; 25 mL serological pipette tips; 5 mL serological pipette tips P1000 Pipette Tips; and P200 Pipette Tips.

Equipment needed for this experiment include: Viaflo 384 liquid handler; Eppendorf serological pipette; Eppendorf P1000 Pipette; and Eppendorf P200 Pipette.

Daudi Cell Culture is also needed for this experiment.

Lastly, compounds (*e.g.*, the compounds of this application) to be tested are needed.

Procedure

All steps were performed in a sterile environment inside the Biosafety cabinet.

A 96 well u-bottom plate was prepared by writing the experiment number, plate number, date and initials in the top right corner of the plate lid. With a sterile 300 ml reservoir, and 25 ml serological pipette, evaporation buffer media was pipetted into reservoir in 25 ml increments. Using the liquid handler, 150 ul of evaporation buffer media was pipetted from reservoir into rows A and H, and Columns 1 and 12 of the 96 well u-bottom plate. Cell cultures were counted to obtain the density of cells per ml, and the culture viability. The cell density information was used to obtain 1,000,000 cells from culture using a 5mL serological pipette into an epi tube. The cell density information from the culture was used to calculate the number of cells and volume of media needed for the assay to seed 1250

cells in 130 μ L of media per available culture well in the 96 well u-bottom plate. Rows B through F were used for cells (50 wells in total), with row G left for an empty media control. The calculation was overestimated by 10 mL to account for the dead volume in the 300 mL reservoir. Once the media volume was calculated, the appropriate volume of media was

5 pipetted in 25 mL increments into the 250 mL bottle using a 25 mL serological pipette. The 250 mL bottle was capped tightly, and placed into a 37°C water bath for 2 minutes. While the culture media was warming, 10 mL of fresh media was pipetted from the 500 mL culture media bottle into a sterile 25 mL reservoir. Using the Eppendorf multichannel pipette, 130 μ L of media was pipetted from the 25 mL reservoir into row G of the 96 well u-bottom plate.

10 Once the 250 mL bottle of media was warmed, the volume of culture needed was pipetted into the bottle, and mixed gently with a 25 mL serological pipette as to not create bubbles, and then the contents of the bottle were pipetted into a new 300 mL reservoir. Using the liquid handler, 130 μ L of culture was pipetted from the 300 mL reservoir into rows B through F of the 96 well u-bottom plate. Once the culture was added, the plate was placed into a 37 °C

15 incubator until the compound master plate was prepared for use.

Two 96 well u-bottom plates were prepared by writing the master plate name in the upper right corner of the plate lid. Labeling one DMSO master and the other Media Master. The compounds of interest were obtained from the laboratory freezer, and placed into a 25 well storage box with a lid, and set the box aside. The compounds were vortexed after

20 thawing but before use. Using an automatic multichannel pipette, 20 μ L of 100% DMSO was pipetted into wells B3-B11 through G3-G11 of the DMSO master plate. For each compound on the master plate, 50 μ L of the compound were pipetted in the appropriate well of row 2 (reference plate map to determine appropriate well). A serial dilution was prepared beginning by aspirating 20 μ L from row 2 and mixing with row 3, repeating until row 11 was reached.

25 Using the liquid handler, 194 μ L of Daudi media was dispensed into wells B2-B11 through G2-G11 of the Media master plate. Using the liquid handler, 6 μ L from the DMSO master plate was aspirated and dispensed into the media master plate, mixing 100 μ L twice.

Compounds from master plate were then added to the culture plate. The culture plates were removed from the incubator, and set inside the biosafety cabinet. Using a liquid handler,

30 20 μ L from wells B2 to B11 through G2 to G11 of master plate were aspirated, and dispensed into wells B2 to B11 through G2 to G11 of culture plate. This set was continued with each culture plate. Once the culture plates acquired their 20 μ L of compound dilutions, they were placed back into the incubator, until their reads on Day 7 of experiment.

Cell death was measured on Day 7 of the experiment using Cell-Titer Glo and a Promega Plate reader.

Percent cell death and EC₅₀ values were calculated by comparing the cell viability of the compound treated wells to the non-treated wells. Normalized RLU values were obtained by subtracting the media well values from each of the wells in the same column, and then dividing that value by the DMSO treated cells values. The percent kill was then calculated by subtracting the normalized RLU value from 1 and multiplying by 100. The average normalized percent kill value and standard error of the mean was then calculated. The kill values were then inputted into Prism with the corresponding standard errors. In Prism a non-linear regression line was plotted with the data points using a semi-log scale, and the EC₅₀ value was calculated. For compounds that showed good potency in the Daudi cell line, the assay was repeated using WI-38 cells (AID negative).

The biological activity of the compounds of the present application measured above is listed below.

Table 2.

Cmpd No.	Cellular activity in AID + (Daudi)
1	D
2	D
3	B
4	B
5	C
6	C
7	C
8	B
9	C
10	B
11	C
12	D
13	B
14	C
15	C
16	C
17	C
18	C
19	D
20	C
21	C
22	C
23	B
24	B
25	C

26	C
27	C
28	D
29	C
30	D
31	C
32	D
33	B
34	C
36	C
38	A
40	A
41	B
42	A
44	C
45	C
46	C
48	C
49	B
50	B
51	B
52	B
54	C
56	A
58	B
59	B
60	A
66	C
67	C
68	B
69	A
70	B
71	C
72	C
73	C
74	B
75	C
76	C
77	C
78	C
79	A
80	B
81	B
82	B
83	B
84	B
85	B
86	B

87	B
88	B
89	B
90	B
91	B
92	B
93	C
94	A
95	B
96	D
97	A
98	A
99	B
100	B
101	C
102	C
103	B
104	B
105	A
106	A
107	A
108	A
109	B
110	B
111	B
112	A
113	A
114	B
115	A
116	A
117	B
118	B
119	A
120	A
121	A
122	A
123	A
124	A
125	A
126	A
127	B
128	A
129	A
130	B
131	A
132	B
133	C
134	B

135	B
136	A
137	A
138	B
139	A
140	B
141	C
142	C
143	A
144	B
145	B
146	A
147	C
148	A
149	A
150	B
151	A
152	A
153	A
154	A
155	B
156	B
157	A
158	B
159	A
160	B
161	A
162	A
163	B
164	B
165	C
166	C
167	B
168	B
169	A
170	B
171	B
172	B
173	B
174	B

EC₅₀: A: $\leq 0.1 \mu\text{M}$, B: $0.1 \mu\text{M} - 1 \mu\text{M}$, C: $1 \mu\text{M} - 10 \mu\text{M}$, D: $> 10 \mu\text{M}$

EQUIVALENTS


The details of one or more embodiments of the disclosure are set forth in the accompanying description above. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present disclosure, the preferred methods and materials are now described. Other features, objects, and advantages of the disclosure will be apparent from the description and from the claims. In the specification and the appended claims, the singular forms include plural referents unless the context clearly dictates otherwise. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure belongs. All patents and publications cited in this specification are incorporated by reference.

The foregoing description has been presented only for the purposes of illustration and is not intended to limit the disclosure to the precise form disclosed, but by the claims appended hereto.

15

CLAIMS


1. A compound of Formula I:



or a pharmaceutically acceptable salt or solvate thereof, wherein:

- 5 the ring Cy is C₃-C₇ cycloalkyl, bridged C₆-C₁₂ cycloalkyl, or saturated heterocyclyl comprising one or two 3- to 7-membered rings and 1-3 heteroatoms selected from N, O, and S, wherein the cycloalkyl or heterocyclyl moiety is optionally substituted with one or more groups selected from halogen, OH, CN, NH₂, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy, and C₁-C₄ haloalkoxy;
- 10 X₁ is NR₈ or O, or, when X₁ is bonded to a nitrogen atom in the ring Cy, X₁ is absent; X₂ is NR₈ or O;
- R₁ is H or C₁-C₆ alkyl optionally substituted with halogen, OH, C₁-C₆ alkoxy, or C₆-C₁₀ aryloxy;
- R₂ is H or C₁-C₆ alkyl optionally substituted with halogen, OH, C₁-C₆ alkoxy, or C₆-C₁₀ aryloxy;
- 15 or R₁ and R₂, together with the nitrogen atom to which they are attached, form a heterocyclyl comprising one or two 3- to 7-membered rings and 1-3 heteroatoms selected from N, O, and S;
- R₃ is C₁-C₆ alkyl optionally substituted with one or more groups selected from halogen, OH, and CN, phenyl, CH₂-phenyl, C₃-C₇ cycloalkyl, CH₂-(C₃-C₇) cycloalkyl, heterocyclyl, or CH₂-heterocyclyl, wherein the heterocyclyl comprises one or two 3- to 7-membered rings and 1-3 heteroatoms selected from N, O, and S, wherein the cycloalkyl, phenyl, or heterocyclyl moiety is optionally substituted with one or more groups selected from halogen, OH, CN, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy, and C₁-C₄ haloalkoxy;
- 20 A₁, A₂, A₃, and A₄ are each independently N or C(R₄);
- each R₄ is independently H, halogen, CN, OH, N(R₆')₂, C₁-C₆ alkoxy, C(=O)N(R₆)₂, C(=O)OR₆, C(=O)R₆, Q-T, C₆-C₁₀ aryl, or heteroaryl comprising one or two 5- or 6-membered rings and 1-3 heteroatoms selected from N, O, and S, wherein the aryl or heteroaryl moiety is optionally substituted with one or more R₉;
- 25

each Q is independently C₁-C₄ alkylene or O-(C₁-C₄) alkylene wherein the oxygen



each T is independently C₁-C₄ alkoxy, OH, N(R₆)₂, N(R₅)C(=O)R₆, N(R₅)C(=O)OR₆, C(=O)N(R₆)₂, C(=O)OR₆, C(=O)R₆, C₆-C₁₀ aryl, or heteroaryl comprising one or two 5- or 6-
5 membered rings and 1-3 heteroatoms selected from N, O, and S, wherein the aryl or heteroaryl moiety is optionally substituted with one or more R₉;

each R₅ is independently H or C₁-C₄ alkyl;

each R₆' is independently H, C₁-C₆ alkyl optionally substituted with one or more R₇, C₁-C₆ haloalkyl, C₃-C₇ cycloalkyl, heterocyclyl comprising one or two 3- to 7-membered
10 rings and 1-3 heteroatoms selected from N, O, and S, C₆-C₁₀ aryl, or heteroaryl comprising one or two 5- or 6-membered rings and 1-3 heteroatoms selected from N, O, and S, wherein the cycloalkyl, heterocyclyl, aryl, or heteroaryl moiety is optionally substituted with one or more R₉, wherein at least one R₆' is not H;

or two R₆' together with the atoms to which they are attached form a 3- to 10-
15 membered heterocyclyl comprising 1-3 heteroatoms selected from N, O, and S, wherein the heterocyclyl is optionally substituted with one or more R₉;


each R₆ is independently H, C₁-C₆ alkyl optionally substituted with one or more R₇, C₁-C₆ haloalkyl, C₆-C₁₀ aryl, or heteroaryl comprising one or two 5- or 6-membered rings and
20 1-3 heteroatoms selected from N, O, and S, wherein the aryl or heteroaryl moiety is optionally substituted with one or more R₉;


or two R₆ together with the atoms to which they are attached form a 3- to 10-
membered heterocyclyl comprising 1-3 heteroatoms selected from N, O, and S, wherein the heterocyclyl is optionally substituted with one or more R₉;

each R₇ is independently N(R₈)₂, OR₈, C₆-C₁₀ aryl, or heteroaryl comprising one or
25 two 5- or 6-membered rings and 1-3 heteroatoms selected from N, O, and S;

each R₈ is independently H or C₁-C₆ alkyl; and

each R₉ is independently oxo, halogen, OH, CN, NH₂, N(C₁-C₄ alkyl)₂, C₁-C₆ alkyl, N(C₁-C₄ alkyl)₂, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, or C₁-C₆ haloalkoxy, wherein the C₁-C₆ alkyl
30 alkyl), or N(C₁-C₄ alkyl)₂.

2. The compound of claim 1, wherein the thiazolyl ring  is unsubstituted.

3. The compound of claim 1, wherein the thiazolyl ring  is substituted with F or Cl.

5

4. The compound of any one of claims 1-3, wherein the ring Cy is C₃-C₇ cycloalkyl optionally substituted with one or more groups selected from halogen, OH, CN, NH₂, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy, and C₁-C₄ haloalkoxy.

10 5. The compound of any one of claims 1-4, wherein the ring Cy is cyclohexyl optionally substituted with one or more groups selected from halogen, OH, CN, NH₂, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy, and C₁-C₄ haloalkoxy.

15 6. The compound of any one of claims 1-3, wherein the ring Cy is bridged C₆-C₁₂ cycloalkyl optionally substituted with one or more groups selected from halogen, OH, CN, NH₂, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy, and C₁-C₄ haloalkoxy.

20 7. The compound of any one of claims 1-3, wherein the ring Cy is saturated heterocyclyl comprising one or two 3- to 7-membered rings and 1-3 heteroatoms selected from N, O, and S, optionally substituted with one or more groups selected from halogen, OH, CN, NH₂, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy, and C₁-C₄ haloalkoxy.

8. The compound of any one of claims 1-7, wherein A₁, A₂, A₃, and A₄ are each C(R₄).

25 9. The compound of any one of claims 1-7, wherein one, two, or three of A₁, A₂, A₃, and A₄ are N.

10. The compound of any one of claims 1-7, wherein at most two of A₁, A₂, A₃, and A₄ are N.

30

11. The compound of any one of claims 1-7, wherein one of A₁, A₂, A₃, and A₄ is N.

12. The compound of any one of claims 1-7, wherein two of A₁, A₂, A₃, and A₄ is N.
13. The compound of any one of claims 1-12, wherein X₁ is NR₈ and X₂ is NR₈.
- 5 14. The compound of any one of claims 1-12, wherein X₁ is NR₈ and X₂ is O.
15. The compound of any one of claims 1-12 and 14, wherein X₁ is NH and X₂ is O.
- 10 16. The compound of any one of claims 1-12, wherein X₁ is O and X₂ is NR₈.
17. The compound of any one of claims 1-12, wherein X₁ is O and X₂ is O.
18. The compound of any one of claims 1-12, wherein X₁ is absent and X₂ is NR₈.
- 15 19. The compound of any one of claims 1-12, wherein X₁ is absent and X₂ is O.
20. The compound of any one of claims 1-19, wherein R₃ is C₁-C₆ alkyl optionally substituted with one or more groups selected from halogen, OH, and CN.
- 20 21. The compound of any one of claims 1-20, wherein R₃ is *i*-propyl.
22. The compound of any one of claims 1-19, wherein R₃ is phenyl or CH₂-phenyl, wherein the phenyl moiety is optionally substituted with one or more groups selected from
- 25 halogen, OH, CN, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy, and C₁-C₄ haloalkoxy.
23. The compound of any one of claims 1-19, wherein R₃ is C₃-C₇ cycloalkyl or CH₂-(C₃-C₇) cycloalkyl, wherein the cycloalkyl moiety is optionally substituted with one or more groups selected from halogen, OH, CN, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy, and C₁-
- 30 C₄ haloalkoxy.
24. The compound of any one of claims 1-19, wherein R₃ is heterocyclyl or CH₂-heterocyclyl, wherein the heterocyclyl moiety is optionally substituted with one or more

groups selected from halogen, OH, CN, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy, and C₁-C₄ haloalkoxy.

25. The compound of any one of claims 1-19 and 24, wherein R₃ is oxetanyl.
- 5 26. The compound of any one of claims 1-25, wherein R₁ is H.
27. The compound of any one of claims 1-25, wherein R₁ is C₁-C₆ alkyl optionally substituted with halogen, OH, C₁-C₆ alkoxy, or C₆-C₁₀ aryloxy.
- 10 28. The compound of any one of claims 1-25 and 27, wherein R₁ is ethyl or *t*-butyl.
29. The compound of any one of claims 1-28, wherein R₂ is H.
- 15 30. The compound of any one of claims 1-28, wherein R₂ is C₁-C₆ alkyl optionally substituted with halogen, OH, C₁-C₆ alkoxy, or C₆-C₁₀ aryloxy.
31. The compound of any one of claims 1-28, wherein R₁ and R₂, together with the nitrogen atom to which they are attached, form a heterocyclyl.
- 20 32. The compound of any one of claims 1-31, wherein each R₄ is H.
33. The compound of any one of claims 1-31, wherein at least one R₄ is halogen, CN, OH, N(R₆')₂, C₁-C₄ alkoxy, C(=O)N(R₆)₂, C(=O)OR₆, C(=O)R₆, Q-T, C₆-C₁₀ aryl, or
- 25 heteroaryl comprising one or two 5- or 6-membered rings and 1-3 heteroatoms selected from N, O, and S, wherein the aryl or heteroaryl moiety is optionally substituted with one or more R₉.
34. The compound of any one of claims 1-31, wherein at least one R₄ is halogen, CN, OH, N(R₆')₂, C₁-C₄ alkoxy, C(=O)N(R₆)₂, C(=O)OR₆, C(=O)R₆, or Q-T.
- 30 35. The compound of any one of claims 1-31, wherein at least one R₄ is C(=O)N(R₆)₂, C(=O)OR₆, C(=O)R₆, or Q-T.


36. The compound of any one of claims 1-31, wherein at least one R₄ is Q-T.
37. The compound of any one of claims 1-31, wherein at least one R₄ is N(R₆')₂.
- 5 38. The compound of any one of claims 1-31, 33, 34, and 37, wherein one of R₆' is H, and the other R₆' is C₁-C₆ alkyl optionally substituted with one or more R₇, C₁-C₆ haloalkyl, C₃-C₇ cycloalkyl, heterocyclyl comprising one or two 3- to 7-membered rings and 1-3 heteroatoms selected from N, O, and S, C₆-C₁₀ aryl, or heteroaryl comprising one or two 5- or
- 10 6-membered rings and 1-3 heteroatoms selected from N, O, and S, wherein the aryl or heteroaryl moiety is optionally substituted with one or more R₉.
39. The compound of any one of claims 1-31, 33, 34, 37, and 38, wherein one of R₆' is H, and the other R₆' is C₁-C₆ alkyl optionally substituted with one or more R₇ or C₁-C₆
- 15 haloalkyl.
40. The compound of any one of claims 1-31, 33, 34, 37, and 38, wherein one of R₆' is H, and the other R₆' is C₃-C₇ cycloalkyl or heterocyclyl optionally substituted with one or more R₉.
- 20 41. The compound of any one of claims 1-31, 33, 34, 37, and 38, wherein one of R₆' is H, and the other R₆' is C₆-C₁₀ aryl or heteroaryl optionally substituted with one or more R₉.
42. The compound of any one of claims 1-31, 33, 34, and 37, wherein two R₆' are each
- 25 C₁-C₆ alkyl optionally substituted with one or more R₇ or C₁-C₆ haloalkyl.
43. The compound of any one of claims 1-31 and 33, wherein at least one R₄ is C₆-C₁₀ aryl optionally substituted with one or more R₉.
- 30 44. The compound of any one of claims 1-31 and 33, wherein at least one R₄ is heteroaryl optionally substituted with one or more R₉.

45. The compound of any one of claims 1-31 and 33-36, wherein at least one Q is C₁-C₄ alkylene.
46. The compound of any one of claims 1-31 and 33-36, wherein at least one Q is C₁-C₄ O-alkylene.
47. The compound of any one of claims 1-31, 33-36, 45, and 46, wherein at least one T is C₁-C₄ alkoxy.
48. The compound of any one of claims 1-31, 33-36, 45, and 46, wherein at least one T is NH₂, NH(C₁-C₄ alkyl), or N(C₁-C₄ alkyl)₂.
49. The compound of any one of claims 1-31, 33-36, 45, and 46, wherein at least one T is N(R₅)C(=O)R₆ or N(R₅)C(=O)OR₆.
50. The compound of any one of claims 1-31, 33-36, 45, and 46, wherein at least one T is C(=O)N(R₆)₂, C(=O)OR₆, or C(=O)R₆.
51. The compound of any one of claims 1-31, 33-36, 45, 46, 49, and 50, wherein each R₆ is H.
52. The compound of any one of claims 1-31, 33-36, 45, 46, 49, and 50, wherein at least one R₆ is C₁-C₆ alkyl optionally substituted with one or more R₇, C₁-C₆ haloalkyl, C₆-C₁₀ aryl, or heteroaryl comprising one or two 5- or 6-membered rings and 1-3 heteroatoms selected from N, O, and S, wherein the aryl or heteroaryl moiety is optionally substituted with one or more R₉.
53. The compound of any one of claims 1-31, 33-36, 45, 46, 49, 50, and 52, wherein at least one R₆ is C₁-C₆ alkyl optionally substituted with one or more R₇ or C₁-C₆ haloalkyl.
54. The compound of any one of claims 1-31, 33-36, 45, 46, 49, 50, and 52, wherein at least one R₆ is C₆-C₁₀ aryl or heteroaryl, wherein the aryl or heteroaryl moiety is optionally substituted with one or more R₉.

55. The compound of any one of claims 1-31, 33-36, 45, 46, 49, 50, 52, and 54, wherein at least one R₆ is C₆-C₁₀ aryl optionally substituted with one or more R₉.


56. The compound of any one of claims 1-31, 33-36, 45, 46, 49, 50, 52, and 54, wherein at least one R₆ is heteroaryl optionally substituted with one or more R₉.

57. The compound of claim 1, of Formula Ia or Ib:




or a pharmaceutically acceptable salt or solvate thereof.


58. The compound of claim 1, of Formula Ic:



or a pharmaceutically acceptable salt or solvate thereof.

59. The compound of claim 1, of Formula Id, Ie, If, or Ig:





or a pharmaceutically acceptable salt or solvate thereof.

60. A compound selected from Table 1 or a pharmaceutically acceptable salt or solvate thereof.

61. A pharmaceutical composition comprising the compound of any one of claims 1-60 or a pharmaceutically acceptable salt or solvate thereof, and a pharmaceutically acceptable carrier or diluent.

10

62. A method of treating cancer, an autoimmune disease, an immune deficiency, or a neurodegenerative disease, comprising administering to a subject in need thereof an effective amount of the compound of any one of claims 1-60 or a pharmaceutically acceptable salt or solvate thereof.

15

63. The method of claim 62, wherein the cancer is selected from the group consisting of lymphoma, leukemia, and a plasma cell neoplasm.

64. The method of claim 63, wherein the cancer is lymphoma.

20

65. The method of claim 64, wherein the lymphoma is a B-cell lymphoma.

66. The method of claim 65, wherein the B-cell lymphoma is diffuse large B-cell lymphoma.

25

67. The method of claim 64, wherein the lymphoma is mantle cell lymphoma.

68. Use of the compound of any one of claims 1-60 or a pharmaceutically acceptable salt or solvate thereof in the manufacture of a medicament for the treatment of cancer, an autoimmune disease, an immune deficiency, or a neurodegenerative disease.
- 5 69. A compound of any one of claims 1-60 or a pharmaceutically acceptable salt or solvate thereof for treating cancer, an autoimmune disease, an immune deficiency, or a neurodegenerative disease.