



- (51) **International Patent Classification:**
C11D 1/83 (2006.01) *C11D 1/66* (2006.01)
- (21) **International Application Number:**
PCT/EP2014/054911
- (22) **International Filing Date:**
13 March 2014 (13.03.2014)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
13160613.9 22 March 2013 (22.03.2013) EP
- (71) **Applicant:** BASF SE [DE/DE]; 67056 Ludwigshafen (DE).
- (72) **Inventors:** BAUER, Frederic; Im Linsenbusch 25, 67146 Deidesheim (DE). ESKUCHEN, Rainer; Alt Langenfeld 42, 40764 Langenfeld (DE). TROPSCH, Jürgen; Im oberen Berg 81, 67354 Römerberg (DE). SÜLING, Carsten; Albrecht-Dürer-Ring 20c, 67227 Frankenthal (DE).
- (81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM,


AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).


Published:

— with international search report (Art. 21(3))

- (54) **Title:** MIXTURES OF ALKYL POLYGLYCOSIDES, THEIR PREPARATION, AND USES



(I)



- (57) **Abstract:** The current invention is directed towards mixtures, comprising (A) in the range of from 15 to 85 % by weight of at least one compound of the general formula (I) (B) in the range of from 85 to 15 % by weight of at least one compound of the general formula (II), wherein the integers are defined as follows: R¹ is C₃-C₄-alkyl, linear or branched, R² is C₅-C₆-alkyl, linear or branched, G¹, G² are different or identical and selected from monosaccharides with 4 to 6 carbon atoms, x, y are numbers in the range of from 1.1 to 4, R³ is C₃-C₉-alkyl, linear or branched, the percentages referring to the total mixture.

