

METHOD OF DETECTING AND QUANTIFYING GENOMIC AND GENE EXPRESSION ALTERATIONS USING RNA

FIELD OF INVENTION

[0001] The present invention relates to the detection and quantification of nucleic acid. In particular, the present invention relates to the detection and quantification of RNA.

BACKGROUND

[0002] Circulating biomarkers are promising tools used for cancer detection, prognostication and prediction of cancer treatment response. These circulating biomarkers typically include DNA samples such as cell-free DNA (cfDNA) and circulating-tumor cells. It is known that various RNA molecules are also potential biomarkers for the diagnosis and prognosis of various diseases such as cancer and are useful for early cancer diagnosis, tumor progression monitor, and prediction of therapy response. It is also known that cancer cells release cell-free RNA (cfRNA) into the body circulation. These cancer-associated cfRNA, also known as circulating-tumor RNA (ctRNA) can be found in the serum and blood plasma of cancer patients. While both cfDNA and cfRNA are promising cancer biomarkers, the measurement of cfDNA is traditionally preferred due to its stability in biological fluids. Despite the discovery of RNA in blood plasma and serum over 20 years ago, there is still a general perception that extracellular RNA in blood is extremely unstable and highly fragmented, given the relative instability of RNA compared to DNA, which in itself is unstable when fragmented in the blood, due to the presence of a high concentration of ribonucleases in the blood circulation. Multiple studies have documented the presence of tumor-specific circulating RNA (ctRNA) in serum and plasma in cancer patients. Current non-oncology clinical applications of cfRNA include the measurement of maternal and fetal cfRNA transcripts to monitor longitudinal phenotypic changes in both the mother and the fetus and to assess fetal gestational age. It is known that in blood circulation, cfRNA occur in free form, bound to proteins or lipids, or as exosomes protected in various types of membrane-derived microvesicles, making them highly stable. It is plausible that plasma cfRNA is a mixture of RNA protected by RNA binding proteins and RNA contained within extracellular vesicles. The wide availability of cfRNA in plasma, serum and many other bodily fluids and their paradoxical stability makes them potential candidates for the development of biomarkers for rapid, sensitive and inexpensive diagnostics. Furthermore, detection of ctRNA provides the same mutational information as ctDNA, but additionally, it can also provide quantitative information about the expression levels of target

genes of interest, and can potentially increase the sensitivity of detection of variants with low allelic frequencies due to the overexpression of tumor-specific transcripts. Lastly, the expression of various ctRNA species is dysregulated due to uncontrolled cell proliferation, rendering it a potential valuable tool for cancer detection. At present, the most common
5 technique for detection of cfRNA is using quantitative Real-time Polymerase Chain Reaction (qRT-PCR). However, methods involving qRT-PCR are often limited by their sensitivity when assaying low input samples. NGS may be more well-suited due the ability to detect novel cfRNAs and differentiate RNA isoforms. With hybridization-based library preparation methods, sequence-specific biases due to enzymatic ligation during library construction step
10 leads to biased representation of transcripts, particularly during analysis of small RNA. Targeted NGS assays such as hybridization capture or amplicon sequencing may also allow for sensitive quantification of cfRNAs (as opposed to whole transcriptome analysis with low conversion efficiency).

[0003] Many cancer genes exhibit genomic alterations, and these genomic alteration events
15 have been discovered in a wide variety of tumors. Targeted DNA-based next generation sequencing techniques specifically designed to detect rearrangements in kinases can effectively detect oncogenic kinase fusions with high confidence. However, there are technical limitations to the ability of such DNA-based assays to detect certain genomic alterations, such as gene fusions. DNA-based assays can only identify fusions in genes where the genomic
20 rearrangements occur in typically short introns effectively covered in the panel. Some clinically important fusions arise from rearrangements in very long introns, the complete coverage of which would significantly compromise coverage of the remainder of the genes on the panel. Hence, there are gaps in the coverage of certain introns resulting in blind spots in the detection of potential rearrangement breakpoint. Fusion detection using DNA does not provide direct
25 evidence that the rearrangement produces a fusion expressed at the mRNA level, a particular problem for rearrangements that appear non-canonical at the genomic DNA level. In fact, in one study in lung cancer tissue samples, it was shown that by using RNA sequencing, alterations were detected in 14% (36/254) cases which were otherwise negative for clinically actionable mutations by DNA sequencing. For example, gene fusion events involving
30 Neurotrophic Receptor Tyrosine Kinase (NTRK) gene (NTRK1/2/3) and neuregulin-1 (NRG1) gene cannot be effectively covered in a targeted DNA sequencing panel without compromising on the cost of sequencing and the coverage of the remaining genes in the sequencing panel.

[0004] Apart from detecting genomic alterations events, the ability to accurately quantify genomic expression of relevant cancer biomarkers non-invasively is important for predicting

the response to cancer therapies and making the appropriate treatment decision. For example, the gene expression level of Programmed death-ligand 1 (PD-L1) is a predictive cancer biomarker used to identify cancer patients with a greater likelihood of responding to immunotherapy. PD-L1 is also a potential predictive biomarker to measure the sensitivity of tumors to immune checkpoint blockade drug inhibitors such as anti-PD-1 inhibitors (pembrolizumab and nivolumab), anti-cytotoxic T-lymphocyte-associated protein 4 (CTLA-4) inhibitors, (ipilimumab and tremelimumab) and anti- programmed death protein 1 (PD-1) (atezolizumab, durvalumab and avelumab). Other genetic biomarkers that are useful for predicting the likelihood of responding to immune checkpoint inhibitor therapy include T cell immunoglobulin and mucin domain-containing protein 3 (TIM-3), Lymphocyte Activating 3 (LAG-3) and cytotoxic T-lymphocyte-associated protein 4 (CTLA-4). The ability to quantify the expression of these target biomarkers longitudinally and non-invasively can be very useful for monitoring treatment response and making treatment decisions.

[0005] Conventional assays routinely detect for genomic alterations at the DNA level, limiting the scope of detection to DNA genomic alterations such as mutations and quantifying genomic copy number changes.

[0006] Thus, there is a need to provide a method for sensitive detection and quantification of genomic alteration events and expression of genes associated with disease (such as cancer) that overcomes, or at least ameliorates, one or more of the disadvantages described above. There is a need to provide a method to simultaneously detect genomic alterations such as structural rearrangements, and gene expression using an alternative sample input such as RNA (such as circulating cell-free RNA (cfRNA)).

SUMMARY

[0007] In one aspect, the present disclosure refers to a method of detecting genomic alteration and/or detecting gene expression and/or quantifying a level of gene expression using RNA in a biological sample, comprising:

- (a) extracting RNA from the biological sample and converting the RNA to complementary DNA (cDNA);
- (b) performing a plurality of multiplexed PCR reactions on the converted cDNA using
 - (I) a plurality of forward and reverse primer pairs specific to a plurality of **target genes that are capable of undergoing genomic alteration**.

wherein each forward primer of the plurality of forward and reverse primer pairs specific to the plurality of target genes that are capable of undergoing genomic alteration is complementary to a sequence located about 50 base pairs upstream of an exonic junction of each target gene that is capable of undergoing genomic alteration,

wherein each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of target genes that are capable of undergoing genomic alteration is complementary to a sequence located about 50 base pairs downstream of an exonic junction of each target gene that is capable of undergoing genomic alteration,

wherein each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of target genes that are capable of undergoing genomic alteration comprises a barcode sequence on its 5' end, wherein the barcode sequence of each reverse primer corresponding to each target gene that is capable of undergoing genomic alteration is different, and/or

(II) a plurality of forward and reverse primer pairs specific to a plurality of control housekeeping genes, wherein:

(i) each forward primer of the plurality of forward and reverse primer pairs specific to the plurality of control housekeeping genes is complementary to a sequence spanning an exon-exon junction of each control housekeeping gene,

wherein each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of control housekeeping genes is complementary to a sequence about 100 base pairs downstream of the sequence spanning the exon-exon junction of each control housekeeping gene,

wherein each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of control housekeeping genes comprises a barcode sequence on its 5' end, wherein the barcode sequence of each reverse primer corresponding to each control housekeeping gene is different;

(ii) each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of control housekeeping genes is complementary

to a sequence spanning an exon-exon junction of each control housekeeping gene,

wherein each forward primer of the plurality of forward and reverse primer pairs specific to the plurality of control housekeeping genes is complementary to a sequence about 100 base pairs downstream of the sequence spanning the exon-exon junction of each control housekeeping gene,

wherein each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of control housekeeping genes comprises a barcode sequence on its 5' end, wherein the barcode sequence of each reverse primer corresponding to each control housekeeping gene is different;

(iii) each forward and each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of control housekeeping genes is complementary to consecutive sequences spanning an exon-exon junction of each control housekeeping gene,

wherein each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of control housekeeping genes comprises a barcode sequence on its 5' end, wherein the barcode sequence of each reverse primer corresponding to each control housekeeping gene is different, and/or

(III) a plurality of primer sets specific to a plurality of **target genes related to protein expression**,

wherein each primer set comprises a plurality of forward and reverse primer pairs specific to each target gene related to protein expression, wherein:

(i) each forward primer of the of the plurality of forward and reverse primer pairs specific to each target gene related to protein expression is complementary to a sequence spanning an exon-exon junction of each target gene related to protein expression,

wherein each reverse primer of the of the plurality of forward and reverse primer pairs specific to each target gene related to protein expression is complementary to a sequence about 100 base pairs downstream of the sequence spanning the exon-exon junction of each target gene related to protein expression,

5 wherein each reverse primer of the plurality of forward and reverse primer pairs specific to each target gene related to protein expression comprises a barcode sequence on its 5' end, wherein the barcode sequence of each reverse primer corresponding to each target gene related to protein expression is different,

(ii) each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of target genes related to protein expression is complementary to a sequence spanning an exon-exon junction of each target gene related to protein expression,

10 wherein each forward primer of the plurality of forward and reverse primer pairs specific to the plurality of target genes related to protein expression is complementary to a sequence about 100 base pairs downstream of the sequence spanning the exon-exon junction of each target gene related to protein expression,

15 wherein each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of target genes related to protein expression comprises a barcode sequence on its 5' end, wherein the barcode sequence of each reverse primer corresponding to each target gene related to protein expression is different;

20 (iii) each forward and each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of target genes related to protein expression is complementary to consecutive sequences spanning an exon-exon junction of each target gene related to protein expression,

25 wherein each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of target genes related to protein expression comprises a barcode sequence on its 5' end, wherein the barcode sequence of each reverse primer corresponding to each target gene related to protein expression is different,

thereby generating a plurality of amplicons;

- 30 (c) purifying the plurality of amplicons from step (b);
(d) amplifying the purified product from step (c) by using universal indexed adapter primers to generate a sequencing library;
(e) purifying the sequencing library obtained from step (d);

- (f) subjecting the purified sequencing library from step (e) to multiplex sequencing on a next-generation sequencing platform to obtain a plurality of sequencing reads;
- (g) deriving a consensus read of each sequence from the plurality of sequencing reads obtained from step (f);
- 5 (h) performing a sequence alignment of the consensus read obtained from step (g) to a reference genome,
- (I) if the sequence alignment results in a partial alignment to the reference genome of an exon from a first gene and a partial alignment to the reference genome of an exon from a second gene, then:
- 10 (i) determining the sequence alignment as a split read,
- (ii) counting/enumerating the number of split reads from step (h)(I)(i) that supports a fusion junction, and
- (iii) if the number of split reads from step (h)(I)(ii) is two or more, then determining the first gene and the second gene as fusion partners,
- 15 (II) if the sequence alignment results in an alignment to the reference genome of the control housekeeping gene, then:
- (i) determining the sequence alignment as a consensus read of the control housekeeping gene,
- (ii) counting/enumerating consensus read pairs of the control housekeeping gene from step (h)(II)(i), and
- 20 (iii) determining the level of gene expression of the control housekeeping gene,
- (III) if the sequence alignment results in an alignment to the reference genome of the target gene related to protein expression,
- 25 (i) determining the sequence alignment as a consensus read of the target gene related to protein expression,
- (ii) counting/enumerating consensus read pairs of the target gene related to protein expression from step (h)(III)(i), and
- (iii) determining the level of gene expression of the target gene related to protein expression;
- 30 (i) determining presence or absence of the genomic alteration and/or determining presence or absence of the gene expression and/or quantifying the level of the gene expression based on the sequence alignment from step (h).

[0008] In another aspect, the present disclosure refers to a kit for detecting genomic alteration and/or detecting gene expression and/or quantifying a level of gene expression using RNA in a biological sample according to the method disclosed herein, wherein the kit comprises:

- a plurality of forward and reverse primer pairs specific to a plurality of **target genes that are capable of undergoing genomic alteration** as defined in the method disclosed herein,
- a plurality of forward and reverse primer pairs specific to a plurality of **control housekeeping genes** as defined in the method disclosed herein, and
- a plurality of primer sets specific to a plurality of **target genes related to protein expression** as defined in the method disclosed herein.

BRIEF DESCRIPTION OF THE DRAWINGS

[0009] The invention will be better understood with reference to the detailed description when considered in conjunction with the non-limiting examples and the accompanying drawings, in which:

[0010] **Fig. 1** is a general overview of the cfRNA-based detection method of a gene fusion event resulting from the intronic DNA rearrangement between two genes as described herein. Primers (represented by arrows) are designed to flank exonic junctions of genes known to undergo fusions. Primers (→) are designed such that the if a fusion product is present, the resulting amplicon would be of ~100 base pairs in length to be compatible with the observed cfRNA fragment size in plasma samples.

[0011] **Fig. 2**, comprising **Figs. 2A and 2B**, illustrates examples of primer design of the disclosed method, wherein **Fig. 2A** illustrates examples of primer design for capturing control housekeeping genes (left panel) and expression genes (right panel) in cfRNA. At least one primer of a primer pair spans an exon-exon junction, to prevent unintended amplification of cfDNA and the resulting amplicons are approximately 100 base pairs in length. It should be noted that the primer pairs for the housekeeping genes differ from that of the expression genes, and **Fig. 2B** illustrates an example of a forward primer and a reverse primer designed to bind to two different exons, intervened by an intron >5000 base pairs in length.

[0012] **Fig. 3**, comprising **Figs. 3A and 3B**, shows the size and concentration analysis of cfRNA from plasma total nucleic acid extracts from cancer patients and healthy individuals, wherein **Fig. 3A** shows the size and concentration analysis of cfRNA from plasma total nucleic acid extracts from a cancer patient (sample A), **Fig. 3B** shows the size and concentration analysis of cfRNA from plasma total nucleic acid extracts from another cancer patient (sample

B), **Fig. 3C** shows the size and concentration analysis of cfRNA from plasma total nucleic acid extracts from a healthy individual (sample C), and **Fig. 3D** shows the size and concentration analysis of cfRNA from plasma total nucleic acid extracts from another healthy individual (sample D). Bioanalyzer RNA 6000 Pico kit or the High Sensitivity RNA Screentape on the 4200 Tapestation were used to quantify and profile the samples. The total concentration of cfRNA (representative of abundance) is generally higher in the representative plasma extracted from cancer patients, relative to those extracted from healthy individuals.

[0013] **Fig. 4** shows the comparison of yield for cfDNA and cfRNA in total nucleic acid extracts from plasma extracted from cancer patients and healthy individuals.

[0014] **Fig. 5**, comprising **Figs. 5A, 5B and 5C**, shows an example of the fragmentation of extracted H2228 cell line RNA by physical shearing of large size nucleotides (>1500 nucleotides) into smaller size to mimic cfRNA fragment size. Bioanalyzer RNA 6000 Pico kit or the High Sensitivity RNA Screentape on the 4200 Tapestation were used to quantify and profile the samples, wherein **Fig. 5A** shows the fragmentation profile of extracted H2228 cell line RNA, **Fig. 5B** shows the resulting fragmentation profile of fragmented H2228 cell line RNA, and **Fig. 5C** shows the fragmentation profile of plasma cfRNA. The resulting fragmentation profile of H2228 cell line RNA is similar to that of plasma cfRNA, with a dominant RNA peak at 119 nucleotides (represented by arrows).

[0015] **Fig. 6**, comprising **Figs. 6A and 6B**, illustrates the detection of EML4-ALK fusion in 1 ng of fragmented H2228 RNA showing the alignment of split reads capturing the fusion break points of exon 6b of EML4 and exon 20 of ALK, wherein **Fig. 6A** is a visualization of the split read on Integrated Genome Viewer (IGV), and **Fig. 6B** is a diagrammatic representation showing exonic fusion (from Arriba tool for detection of gene fusions).

[0016] **Fig. 7** is a diagrammatic representation from Arriba tool showing the detection of various exonic fusion in cancer cell lines: NCI-H660 cell line (CRL-5813, ATCC), VCaP cell line (CRL-2876, ATCC), MV-4-11 cell line (CRL-9591, ATCC) and Kasumi-1 cell line (CRL-2724, ATCC), using the multiplex amplicon sequencing method as described herein for fragmented RNA samples.

[0017] **Fig. 8**, comprising **Figs. 8A, 8B and 8C**, shows detection of TMPRSS2-ERG gene fusion in nucleic acid extracts from a metastatic prostate patient using the cfRNA-based method described herein, compared to a cfDNA-based method, wherein **Fig. 8A** is an IGV graphic view showing 17 split reads, which supported the presence of intronic breakpoints, detected with the cfDNA-based detection method, **Fig. 8B** is an IGV graphic report showing 4123 split reads, which supported the presence of corresponding exonic breakpoints, detected with the

cfRNA-based method described herein, and **Fig. 8C** is a diagrammatic representation from Arriba tool showing TMPRSS2-ERG gene fusion.

[0018] **Fig. 9**, comprising **Figs. 9A, 9B and 9C**, shows detection of CCDC6-RET gene fusion in a nucleic acid extract from a metastatic lung cancer patient using the cfRNA-based method described herein, compared to a cfDNA-based method, wherein **Fig. 9A** is an IGV graphic report showing 12 split reads, which supported the presence of intronic breakpoints, detected with the cfDNA-based detection method, **Fig. 9B** is an IGV graphic report showing 1474 split reads, which supported the presence of corresponding exonic breakpoints, detected with the cfRNA-based method described herein, and **Fig. 9C** is a diagrammatic representation from Arriba tool showing CCDC6-RET gene fusion.

[0019] **Fig. 10**, comprising **Figs. 10A and 10B**, shows the detection of BCR-ABL1 gene fusion in an RNA sample extracted from the peripheral blood cell fraction of an acute lymphoblastic leukaemia clinical sample using the cfRNA-based method described herein, wherein **Fig. 10A** is an IGV graphic report showing BCR-ABL1 gene fusion, and **Fig. 10B** is a diagrammatic representation from Arriba tool showing BCR-ABL1 gene fusion.

[0020] **Fig. 11** shows the result from quantitating the number of EML4-ALK fusion transcript copies per nanogram RNA from H2228 cell line, for determining the sensitivity limits of detection of the cfRNA-based method described herein.

[0021] **Fig. 12** shows the detection and quantification of expression of control genes and other target genes in cfRNA from both cancer and healthy samples, using the method described herein. The table (top panel) describes the amount of cfRNA input for each sample tested, including repeats of two samples with differing input cfRNA amounts. The expression heatmap (bottom panel) demonstrates the distribution of the expression read counts, as derived from the method described herein, for each sample. Fusion detection in the same sample is feasible and shown for C_20-347, and C_20-146, which were simultaneously positive for CCDC6-RET and TMPRSS2-ERG fusion, respectively, as depicted in **Fig. 8 and Fig. 9**.

[0022] **Fig. 13**, comprising **Figs. 13A, 13B and 13C**, shows the identification of actionable driver fusions in untreated lung cancer cases using cfRNA using the method described herein, wherein **Fig. 13A** shows detection of LMNA-NTRK1 fusion, **Fig. 13B** shows detection of CD74-NRG1 fusion, and **Fig. 13C** shows detection of ETV6-NTRK3 fusion in cfRNA in three lung cancer cases, respectively, that were negative for the presence of other driver gene mutations in cfDNA.

[0023] **Fig. 14**, comprising **Figs. 14A and 14B**, shows fusion detection in 45 lung cancer cases by cfDNA and cfRNA using the method described herein and that additional fusions were identified when cfRNA fraction was used, compared to cfDNA. Clinical samples processed simultaneously using cfRNA and cfDNA were compared for fusion detection, wherein **Fig. 14A** shows concordance of fusion detection based on cfDNA and cfRNA, showing cfRNA identified additional fusions in 5 cases, and missed 1 fusion detectable by cfDNA. There were 12 cases with concordant fusion detection by both methods, and **Fig. 14B** lists the range of fusions detected by both cfDNA and cfRNA methods, or by one of the two methods and the detection of multiple co-occurring fusions detected by cfRNA. (* = fusion detected by both cfDNA and cfRNA).

[0024] **Fig. 15** illustrates a typical library profile for a cfRNA sample converted to a sequencing library as seen on the High Sensitivity DNA Screentape. The multiple peaks >200 base pairs correspond to the multiple products encompassing potential fusion products, control gene products and other gene expression products for which the multiplicity of forward and reverse primers are included. Qualified libraries will have prominent peaks of size >200 base pairs.

[0025] **Fig. 16** is an IGV graphic report showing the detection of an 18-bp deletion in RNA extracted from FFPE lung tumor tissue using cfRNA-based method described herein. The expression of EGFR c.2240_2257del p.L747_P753delinsS mutant transcript (comprising the deletion) was supported by 4266 reads.

[0026] **Fig. 17** is an IGV graphic report showing the detection of a single nucleotide variation in cfRNA extracted from the plasma of a metastatic lung cancer patient using cfRNA-based method described herein. Expression of EGFR c.2573T>G p.L858R mutant transcript (comprising the single nucleotide variation) was supported by 112 reads.

[0027] **Fig. 18**, comprising **Figs. 18A and 18B**, shows the detection of expressed transcripts containing single nucleotide variation, insertion (e.g. duplication) or deletion mutations using cfRNA-based method described herein, wherein **Fig. 18A** shows single nucleotide variation, insertion or deletion mutations detected in tissue RNA extracted from FFPE tumor samples, and **Fig. 18B** shows single nucleotide variation detected in cfRNA extracted from plasma.

DETAILED DESCRIPTION

[0028] The disclosed method allows for detection of genomic alteration and gene expression as well as quantifying the level of gene expression of RNA (such as cfRNA) in a biological

sample, for the purpose of non-invasive cancer detection, prognostication, and prediction of treatment response. The present disclosure describes a method based on highly multiplexed amplicon-based NGS, that involves the tagging of individual cfRNA molecules using barcode sequences, and the optimized design of amplicons to be compatible with the fragmented nature of cfRNA. The method described herein can be applied to circulating nucleic acid extracts containing both cfDNA and cfRNA, and can detect and quantitate fusion RNA transcript and gene expression simultaneously, in nucleic acid extracts samples. The applicability of cfRNA is extended in the present disclosure with a novel amplicon-based NGS assay combining fusion detection and gene expression monitoring. With hybridization-based library preparation methods sequence-specific biases due to enzymatic ligation during library construction step lead to biased representation of transcripts, particularly during analysis of small amount of input RNA. Targeted NGS assays such as hybridization capture or amplicon sequencing can allow for sensitive quantification of cfRNA. A targeted NGS-based method has a higher conversion efficiency as compared to whole transcriptome analysis, which has disadvantages such as cost and manpower.

[0029] In a first aspect, the present disclosure refers to a method of detecting genomic alteration and/or detecting gene expression and/or quantifying a level of gene expression using RNA in a biological sample, comprising:

(a) extracting RNA from the biological sample and converting the RNA to complementary DNA (cDNA);

(b) performing a plurality of multiplexed PCR reaction on the converted cDNA using

(I) a plurality of forward and reverse primer pairs specific to a plurality of **target genes that are capable of undergoing genomic alteration**,

wherein each forward primer of the plurality of forward and reverse primer pairs specific to the plurality of target genes that are capable of undergoing genomic alteration is complementary to a sequence located about 50 base pairs upstream of an exonic junction of each target gene that is capable of undergoing genomic alteration,

wherein each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of target genes that are capable of undergoing genomic alteration is complementary to a sequence located about 50 base pairs downstream of an exonic junction of each target gene that is capable of undergoing genomic alteration,

wherein each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of target genes that are capable of undergoing genomic alteration comprises a barcode sequence on its 5' end, wherein the barcode sequence of each reverse primer corresponding to each target gene that is capable of undergoing genomic alteration is different, and/or

(II) a plurality of forward and reverse primer pairs specific to a plurality of **control housekeeping genes**, wherein:

(i) each forward primer of the plurality of forward and reverse primer pairs specific to the plurality of control housekeeping genes is complementary to a sequence spanning an exon-exon junction of each control housekeeping gene,

wherein each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of control housekeeping genes is complementary to a sequence about 100 base pairs downstream of the sequence spanning the exon-exon junction of each control housekeeping gene,

wherein each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of control housekeeping genes comprises a barcode sequence on its 5' end, wherein the barcode sequence of each reverse primer corresponding to each control housekeeping gene is different;

(ii) each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of control housekeeping genes is complementary to a sequence spanning an exon-exon junction of each control housekeeping gene,

wherein each forward primer of the plurality of forward and reverse primer pairs specific to the plurality of control housekeeping genes is complementary to a sequence about 100 base pairs downstream of the sequence spanning the exon-exon junction of each control housekeeping gene,

wherein each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of control housekeeping genes comprises a barcode sequence on its 5' end, wherein the barcode

sequence of each reverse primer corresponding to each control housekeeping gene is different;

- (iii) each forward and each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of control housekeeping genes is complementary to consecutive sequences spanning an exon-exon junction of each control housekeeping gene,

wherein each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of control housekeeping genes comprises a barcode sequence on its 5' end, wherein the barcode sequence of each reverse primer corresponding to each control housekeeping gene is different, and/or

- (III) a plurality of primer sets specific to a plurality of **target genes related to protein expression**,

wherein each primer set comprises a plurality of forward and reverse primer pairs specific to each target gene related to protein expression, wherein:

- (i) each forward primer of the of the plurality of forward and reverse primer pairs specific to each target gene related to protein expression is complementary to a sequence spanning an exon-exon junction of each target gene related to protein expression

wherein each reverse primer of the of the plurality of forward and reverse primer pairs specific to each target gene related to protein expression is complementary to a sequence about 100 base pairs downstream of the sequence spanning the exon-exon junction of each target gene related to protein expression,

wherein each reverse primer of the plurality of forward and reverse primer pairs specific to each target gene related to protein expression comprises a barcode sequence on its 5' end, wherein the barcode sequence of each reverse primer corresponding to each target gene related to protein expression is different,

- (ii) each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of target genes related to protein expression is complementary to a sequence spanning an exon-exon junction of each target gene related to protein expression,

wherein each forward primer of the plurality of forward and reverse primer pairs specific to the plurality of target genes related to protein expression is complementary to a sequence about 100 base pairs downstream of the sequence spanning the exon-exon junction of each target gene related to protein expression,

wherein each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of target genes related to protein expression comprises a barcode sequence on its 5' end, wherein the barcode sequence of each reverse primer corresponding to each target gene related to protein expression is different;

(iii) each forward and each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of target genes related to protein expression is complementary to consecutive sequences spanning an exon-exon junction of each target gene related to protein expression,

wherein each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of target genes related to protein expression comprises a barcode sequence on its 5' end, wherein the barcode sequence of each reverse primer corresponding to each target gene related to protein expression is different,

thereby generating a plurality of amplicons;

- (c) purifying the plurality of amplicons from step (b);
- (d) amplifying the purified product from step (c) by using universal indexed adapter primers to generate a sequencing library;
- (e) purifying the sequencing library obtained from step (d);
- (f) subjecting the purified sequencing library from step (e) to multiplex sequencing on a next-generation sequencing platform to obtain a plurality of sequencing reads;
- (g) deriving a consensus read of each sequence from the plurality of sequencing reads obtained from step (f);
- (h) performing a sequence alignment of the consensus read obtained from step (g) to a reference genome,

(I) if the sequence alignment results in a partial alignment to the reference genome of an exon from a first gene and a partial alignment to the reference genome of an exon from a second gene, then:

- (i) determining the sequence alignment as a split read,

- (ii) counting/enumerating the number of split reads from step (h)(I)(i) that supports a fusion junction, and
 - (iii) if the number of split reads from step (h)(I)(ii) is two or more, then determining the first gene and the second gene as fusion partners,
 - 5 (II) if the sequence alignment results in an alignment to the reference genome of the control housekeeping gene, then:
 - (i) determining the sequence alignment as a consensus read of the control housekeeping gene,
 - (ii) counting/enumerating consensus read pairs of the control housekeeping gene from step (h)(II)(i), and
 - 10 (iii) determining the level of gene expression of the control housekeeping gene,
 - (III) if the sequence alignment results in an alignment to the reference genome of the target gene related to protein expression,
 - 15 (i) determining the sequence alignment as a consensus read of the target gene related to protein expression,
 - (ii) counting/enumerating consensus read pairs of the target gene related to protein expression from step (h)(III)(i), and
 - (iii) determining the level of gene expression of the target gene related to protein expression,
 - 20 (i) determining presence or absence of the genomic alteration and/or determining presence or absence of the gene expression and/or quantifying the level of the gene expression based on the sequence alignment from step (h).
- [0030] In one example, the disclosed method is used to detect genomic alteration of RNA in a biological sample. For example, the method may be used to detect known and unknown fusions and their quantification, relative to the quantity of control housekeeping genes expression in a given sample. In another example, the disclosed method is used to detect gene expression of RNA in a biological sample. In yet another example, the disclosed method is used to quantify the level of gene expression of RNA in a biological sample. In a further example, the disclosed method is used to simultaneously detect genomic alteration of RNA and detect gene expression of RNA in a biological sample. In a further example, the disclosed method is used to simultaneously detect genomic alteration of RNA and quantify gene expression of RNA in a biological sample. In a further example, the disclosed method is used to simultaneously detect

genomic alteration of RNA, detect gene expression of RNA, and quantify gene expression of RNA in a biological sample.

[0031] In one example, the disclosed method is used to detect genomic alteration of cfRNA in a biological sample. For example, the method may be used to detect known and unknown fusions and their quantification, relative to the quantity of control housekeeping genes expression in a given sample. In another example, the disclosed method is used to detect gene expression of cfRNA in a biological sample. In yet another example, the disclosed method is used to quantify the level of gene expression of cfRNA in a biological sample. In a further example, the disclosed method is used to simultaneously detect genomic alteration of cfRNA and detect gene expression of cfRNA in a biological sample. In a further example, the disclosed method is used to simultaneously detect genomic alteration of cfRNA and quantify gene expression of cfRNA in a biological sample. In a further example, the disclosed method is used to simultaneously detect genomic alteration of cfRNA, detect gene expression of cfRNA, and quantify gene expression of cfRNA in a biological sample.

[0032] In one example, the design of the primers to capture fusion transcripts has two main features – 1) the presence of a random barcode sequence in the downstream primers (downstream relative to the target gene (e.g. fusion) transcript) to individually tag each copy of the RNA transcript if present, and 2) the location of each primer approximately 50 base pairs from each exonic junction in the panel, such that the expected total amplicon length would be close to 90-110 base pairs. This was done in order to meet the sample cfRNA size distribution observed which peaked at 110-120 nucleotides.

[0033] In one example, the plurality of forward and reverse primer pairs specific to a plurality of **target genes that are capable of undergoing genomic alteration** as disclosed in step (b)(I) is designed as shown in **Fig. 1**:

wherein each forward primer of the plurality of forward and reverse primer pairs specific to the plurality of target genes that are capable of undergoing genomic alteration is complementary to a sequence located about 50 base pairs upstream of an exonic junction of each target gene that is capable of undergoing genomic alteration,

wherein each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of target genes that are capable of undergoing genomic alteration is complementary to a sequence located about 50 base pairs downstream of an exonic junction of each target gene that is capable of undergoing genomic alteration,

wherein each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of target genes that are capable of undergoing genomic

alteration comprises a barcode sequence on its 5' end, wherein the barcode sequence of each reverse primer corresponding to each target gene that is capable of undergoing genomic alteration is different.

In one example, the plurality of forward and reverse primer pairs specific to a plurality of
5 **control housekeeping genes** as disclosed in step (b)(II) is designed, wherein:

(i) each forward primer of the plurality of forward and reverse primer pairs specific to the plurality of control housekeeping genes is complementary to a sequence spanning an exon-exon junction of each control housekeeping gene,

wherein each reverse primer of the plurality of forward and reverse primer pairs
10 specific to the plurality of control housekeeping genes is complementary to a sequence about 100 base pairs downstream of the sequence spanning the exon-exon junction of each control housekeeping gene,

wherein each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of control housekeeping genes comprises a barcode
15 sequence on its 5' end, wherein the barcode sequence of each reverse primer corresponding to each control housekeeping gene is different, as shown in **Fig. 2A (left)**;

(ii) each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of control housekeeping genes is complementary to a sequence
20 spanning an exon-exon junction of each control housekeeping gene,

wherein each forward primer of the plurality of forward and reverse primer pairs specific to the plurality of control housekeeping genes is complementary to a sequence about 100 base pairs downstream of the sequence spanning the exon-exon junction of each control housekeeping gene,

wherein each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of control housekeeping genes comprises a barcode
25 sequence on its 5' end, wherein the barcode sequence of each reverse primer corresponding to each control housekeeping gene is different;

(iii) each forward and each reverse primer of the plurality of forward and reverse primer
30 pairs specific to the plurality of control housekeeping genes is complementary to consecutive sequences spanning an exon-exon junction of each control housekeeping gene,

wherein each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of control housekeeping genes comprises a barcode

sequence on its 5' end, wherein the barcode sequence of each reverse primer corresponding to each control housekeeping gene is different.

In one example, the plurality primer sets specific to a plurality of **target genes related to protein expression** as disclosed in step (b)(III) is designed,

5 wherein each primer set comprises a plurality of forward and reverse primer pairs specific to each target gene related to protein expression, wherein:

(i) each forward primer of the of the plurality of forward and reverse primer pairs specific to each target gene related to protein expression is complementary to a sequence spanning an exon-exon junction of each target gene related to protein expression,

10 wherein each reverse primer of the of the plurality of forward and reverse primer pairs specific to each target gene related to protein expression is complementary to a sequence about 100 base pairs downstream of the sequence spanning the exon-exon junction of each target gene related to protein expression,

15 wherein each reverse primer of the plurality of forward and reverse primer pairs specific to each target gene related to protein expression comprises a barcode sequence on its 5' end, wherein the barcode sequence of each reverse primer corresponding to each target gene related to protein expression is different, as shown in **Fig. 2A (right)**;

20 (ii) each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of target genes related to protein expression is complementary to a sequence spanning an exon-exon junction of each target gene related to protein expression,

25 wherein each forward primer of the plurality of forward and reverse primer pairs specific to the plurality of target genes related to protein expression is complementary to a sequence about 100 base pairs downstream of the sequence spanning the exon-exon junction of each target gene related to protein expression,

30 wherein each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of target genes related to protein expression comprises a barcode sequence on its 5' end, wherein the barcode sequence of each reverse primer corresponding to each target gene related to protein expression is different;

(iii) each forward and each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of target genes related to protein expression is

complementary to consecutive sequences spanning an exon-exon junction of each target gene related to protein expression,

wherein each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of target genes related to protein expression comprises a barcode sequence on its 5' end, wherein the barcode sequence of each reverse primer corresponding to each target gene related to protein expression is different.

[0034] In one example, the forward primer of the plurality of forward and reverse primer pairs specific to a plurality of control housekeeping genes as disclosed in step (b)(II) is complementary to a sequence in a first exon and the reverse primer of the plurality of forward and reverse primer pairs specific to a plurality of control housekeeping genes as disclosed in step (b)(II) is complementary to a sequence in a second exon as shown in **Fig. 2B**, wherein the first exon and the second exon is intervened by an intron of >5000 base pairs in length, thereby avoiding unintended amplification of any genomic DNA during the plurality of multiplexed PCR reactions.

[0035] In one example, at least one of the primers of each forward and reverse primer pair of the plurality of forward and reverse primer pairs specific to a plurality of control housekeeping genes as disclosed in step (b)(II) span an exon-exon junction. In one example, at least one of the primers of each forward and reverse primer pair of the plurality of forward and reverse primer pairs specific to a plurality of target genes related to protein expression as disclosed in step (b)(III) span an exon-exon junction. In one example, at least one of the primers of each forward and reverse primer pair of the plurality of forward and reverse primer pairs specific to a plurality of control housekeeping genes as disclosed in step (b)(II), and/or at least one of the primers of each forward and reverse primer pair of the plurality of forward and reverse primer pairs specific to a plurality of target genes related to protein expression as disclosed in step (b)(III) span an exon-exon junction. In one example, a forward primer or a reverse primer of the plurality of forward and reverse primer pairs specific to a plurality of control housekeeping genes as disclosed in step (b)(II), and/or a forward primer or a reverse primer of the plurality of forward and reverse primer pairs specific to a plurality of target genes related to protein expression as disclosed in step (b)(III) span an exon-exon junction. In another example, both the forward primer and the reverse primer of the plurality of forward and reverse primer pairs specific to a plurality of control housekeeping genes as disclosed in step (b)(II), and/or both the forward primer and the reverse primer of the plurality of forward and reverse primer pairs specific to a plurality of target genes related to protein expression

as disclosed in step (b)(III) span an exon-exon junction, wherein the exon length is about 100 base pairs.

[0036] In one example, each reverse primer of the plurality of forward and reverse primer pairs specific to a plurality of **target genes that are capable of undergoing genetic alteration** as disclosed in step (b)(I), each reverse primer of the plurality of forward and reverse primer pairs specific to a plurality of **control housekeeping genes** as disclosed in step (b)(II), and each reverse primer of the plurality of forward and reverse primer pairs specific to a plurality of **target genes related to protein expression** as disclosed in step (b)(III) comprise a barcode sequence on its 5' end, wherein each barcode sequence is different. As used herein, the term "barcode sequence" refers to an encoded molecule or barcode that includes variable amount of information within the nucleic acid sequence. For example, the barcode sequence is a tag that can be read out using any of a variety of sequence identification techniques, for example, nucleic acid sequencing, probe hybridization-based assay, and the like. In some examples, the barcode sequence is used in the method as described herein to tag different converted cDNA sequences of target regions of a sample, such that when the barcode sequence tags to the converted DNA sequences of target regions, each different converted cDNA sequence of target region would then have a unique barcode sequence that is attached to it and read out with the converted cDNA sequence of target region from the sample.

[0037] The barcode sequence allows the pooled analysis of multiple unique target sequences, where the resulting sequence information from the pool can be later attributed back to each starting target sequence. That is, after the process of amplification, the barcode sequence is used to group amplicons to form a family of amplicons having the same barcode sequence. In some examples, the barcode sequence is an overhang that does not complement any sequence within the target region. As each reverse primer carries on its 5' end a randomly assigned barcode sequence as disclosed herein, the barcode sequence allows individual cDNA molecules to be tagged uniquely in the step of sequencing library formation.

[0038] In one example, the barcode sequence is an oligonucleotide comprising 10 to 16 random nucleotides, or 10 to 15 random nucleotides, or 10 to 13 random nucleotides, or 10 random nucleotides, or 11 random nucleotides, or 12 random nucleotides, or 13 random nucleotides, or 14 random nucleotides, or 15 random nucleotides, or 16 random nucleotides. In one example, the barcode sequence is an oligonucleotide comprising 10 to 16 random nucleotides. In one example, the barcode sequence is an oligonucleotide comprising 10 random nucleotides. In one specific example, the barcode sequence is an oligonucleotide comprising 10 random nucleotides which can be represented as NNNNNNNNNN (SEQ ID NO: 615).

[0039] In one example, the typical length of each forward primer of the plurality of forward and reverse primers pairs as disclosed in step (b), excluding the barcode sequence and partial adapter sequence, is about 20 base pairs. In one example, the typical length of each reverse primer of the plurality of forward and reverse primers pairs as disclosed in step (b), excluding the barcode sequence and partial adapter sequence, is about 20 base pairs. In one example, the typical length of each forward primer of the plurality of forward and reverse primer pairs as disclosed in step (b), including the barcode sequence and partial adapter sequence, is about 45 base pairs, wherein the length of the barcode sequence is about 10 base pairs, wherein the length of the partial adapter sequence is about 20 base pairs. In one example, the typical length of each reverse primer of the plurality of forward and reverse primer pairs as disclosed in step (b), including the barcode sequence and partial adapter sequence, is about 45 base pairs, wherein the length of the barcode sequence is about 10 base pairs, wherein the length of the partial adapter sequence is about 20 base pairs.

[0040] In one example, the biological sample contains RNA. In one example, the RNA is cfRNA. In one example, the cfRNA is present freely in the biological sample and can be converted to cDNA directly as disclosed in step (a) of the disclosed method.

[0041] In one example, the cfRNA is extracted from the biological sample prior to step (a) of the disclosed method. In a further example, the RNA may be those that are originally encapsulated within cells and needs to be extracted prior to step (a) of the disclosed method. In one example, the cell may be any type of cell in the body. In one example, the cell is from bone, epithelial, cartilage, adipose tissue, nerves, muscle, connective tissue, esophagus, stomach, liver, gallbladder, pancreas, adrenal glands, bladder, gallbladder, large intestine, small intestine, kidneys, liver, pancreas, colon, stomach, thymus, spleen, brain, spinal cord, heart, lungs, eyes, corneal, skin, or islet tissue or organs. In one example, the cell may be a cancer cell, a stem cell, an endothelial cell, or a fat cell. In one example, the cell is a blood cell. The blood cell may be a white blood cell, or a platelet. In one example, the cell is selected from cancer cells known to harbour genomic alterations. In one example, the cell is selected from cancer cell lines known to harbour fusion genes. In one example, the cancer cell lines harbouring fusion genes may include, but are not limited to, CRL-9591, H-2228, CRL-2724, VCaP, CRL-5813, etc. Various methods for RNA extraction are known in the art and may be used for the purpose of the disclosed method. Various methods for RNA extraction are known in the art and may be used for the purpose of the disclosed method. In one example, the cfRNA is extracted from the biological sample before step (a) using a kit such as, but not limited to Zymo Quick-cfRNA Serum & Plasma Kit (Zymo Research), NextPrep™ Magnazol™ cfRNA

Isolation Kit (PerkinElmer), Isopure Plasma cfDNA/RNA Isolation Kit (Aline Biosciences), QIAmp Circulating Nucleic Acid Kit (Qiagen), QIAamp ccfDNA/RNA Kit (Qiagen), MagMAX™ Cell-Free Total Nucleic Acid Isolation Kit (Applied Biosystems), etc.

[0042] In one example, the RNA extracted from cells are subjected to ultrasonification to thereby resemble the size of cfRNA more closely. In another example, the ultrasonification is achieved using Covaris, Qsonica, Diagenode Bioruptor, etc. In another example, the RNA extracted from the cells are subjected to heat and divalent cation-based fragmentation. In yet another example, the fragmentation is achieved using NEBNext® Magnesium RNA Fragmentation Module.

[0043] In one example, the biological sample contains both cfRNA and cfDNA. As used herein, cfDNA refers to non-encapsulated DNA which is present freely in a liquid sample disclosed herein and not contained within cells. The presence of long intervening introns which have undergone rearrangements prevents rearranged cfDNA from forming sequenceable products.

[0044] In the disclosed method, cfRNA present freely in the biological sample or those extracted from the biological sample, is first converted to cDNA as disclosed in step (a) of the method of the first aspect. In one example, cfRNA is converted to cDNA by reverse transcription. The term “reverse transcription” and its grammatical variants as used herein refers to the enzyme-mediated synthesis of a DNA molecule from an RNA template. The resulting DNA, known as complementary DNA (cDNA), can be used as a template for PCR amplification. Methods of reverse transcription, which typically involve the use of non-target specific primers (random primers), are well known in the art. In one example, cfRNA is converted to cDNA using a reverse transcription kit, wherein the reverse transcription kit comprises a reverse transcriptase enzyme and a plurality of random primers. In one example, the random primers are 6-mer primers, 7-mer primers, 8-mer primers, 9-mer primers or combinations thereof. In one example, the random primers are 6-mer (hexamer/hexanucleotide) primers. In one example, the reverse transcription kit is selected from, but is not limited to, High-Capacity cDNA Reverse Transcription Kit (Thermo Fisher Scientific), SuperScript IV One-Step RT-PCR System (Invitrogen), etc.

[0045] In one example, the biological sample containing the RNA is a liquid sample, a tissue sample, or a cell sample. In yet another example, the tissue sample is a frozen tissue sample or a fixed tissue sample. In another example, the fixed tissue sample is a Formalin-Fixed Paraffin-Embedded (FFPE) tissue sample. In another example, the liquid sample is a bodily fluid. In one example, the bodily fluid is selected from the group consisting of blood, bone marrow,

cerebral spinal fluid, peritoneal fluid, pleural fluid, lymph fluid, ascites, serous fluid, sputum, lacrimal fluid, stool, urine, saliva, ductal fluid from breast, gastric juice, and pancreatic juice. In one example, the bodily fluid is blood. In one example, the blood is plasma.

[0046] In another example, the biological sample is obtained from a subject having and/or
5 suspected of having a disease. In another example, the disease is cancer. In yet another example, the cancer is selected from the group consisting of leukemia, lung cancer, colorectal cancer, breast cancer, pancreatic cancer, prostate cancer, nasopharyngeal cancer, liver cancer, cholangiocarcinoma, esophageal cancer, urothelial cancer, and gastrointestinal cancer. In one example, the cancer is an early-stage cancer. In another example, the cancer is a late-stage or
10 metastatic cancer. In one example, the cancer is selected from the group consisting of metastatic prostate cancer, metastatic lung cancer, metastatic breast cancer and leukemia.

[0047] In one example, the genomic alteration(s) detected using the disclosed method comprise(s) structural rearrangement(s). In one example, the term “rearrangement” refers to – rearrangement in the order of sections of the DNA. In one example, the structural
15 rearrangement is a fusion, such as a gene fusion. In one example, the term “fusion” refers to structural variations produced through structural rearrangements, such as interchromosomal or intrachromosomal rearrangements. In one example, the structural rearrangement may include, but are not limited to, deletion, insertion (such as duplication), inversion, transversion, translocation, alternative splicing, and the like. In one example, the structural rearrangement
20 results in formation of a fusion gene, such as one that is detectable using the methods disclosed herein. In one example, the “deletion” is a sequence change where at least one nucleotide is removed. In one example, the “deletion” is a sequence change where more than 10 nucleotides are removed. In one example, the “deletion” is a sequence change where more than 20 nucleotides are removed. In one example, the “deletion” is a sequence change where more than
25 30 nucleotides are removed. In one example, the “deletion” is a sequence change where more than 40 nucleotides are removed. In one example, the “deletion” is a sequence change where more than 50 nucleotides are removed. In one example, the “deletion” may be a “small deletion” where less than 50 nucleotides are removed. In one example, the “insertion” is a sequence change where at least one nucleotide is inserted between two nucleotides. In one
30 example, the “insertion” is a sequence change where more than 10 nucleotides are inserted between two nucleotides. In one example, the “insertion” is a sequence change where more than 20 nucleotides are inserted between two nucleotides. In one example, the “insertion” is a sequence change where more than 30 nucleotides are inserted between two nucleotides. In one example, the “insertion” is a sequence change where more than 40 nucleotides are inserted

between two nucleotides. In one example, the “insertion” is a sequence change where more than 50 nucleotides are inserted between two nucleotides. In one example, the “insertion” may be a “small insertion” where less than 50 nucleotides are inserted between two nucleotides. In one example, the “insertion” is a “duplication”. In one example, the “duplication” is a sequence change where a copy of one or more nucleotides are inserted directly 3’-flanking of the original copy. In one example, the term “inversion” refers to a sequence change where more than one nucleotide replacing the original sequence are the reverse complement of the original sequence. In one example, the term “translocation” refers to rearrangement of parts between non-homologous chromosomes, which can result in “fusion”. In one example, “altered splicing” refers to aberrant splicing of a single gene transcript that may cause one or more exons in sequence to be spliced out of the RNA, bringing usually more distant exons of the same gene in juxtaposition. Altered splicing involves the same gene, compared to fusion which is a definition reserved for two genes. In one example, altered splicing included MET exon 14 skipping where exon 14 of MET gene is spliced out bringing exon 13 and exon 15 in proximity, which is detectable using the method described herein (**Fig. 14**). In one example, the genomic alteration(s) detected using the disclosed method comprise(s) single nucleotide variations. In one example, “single nucleotide variations” refer to variation in a single nucleotide that occurs at a specific position in the genome, differing from the nucleotide defining the position in the reference genome.

[0048] In one example, “housekeeping genes” refer to highly conserved genes which are essential for maintaining cellular function. In one example, the control housekeeping gene comprises Glucose-6-phosphate isomerase (GPI), FERM domain containing 8 (FRMD8), Small nuclear ribonucleoprotein D3 (SNRPD3), Proteasome subunit, beta type, 2 (PSMB2), TATA box binding protein (TBP), REL proto-oncogene, NF-kB subunit (REL), synaptosome associated protein 29 (SNAP29), Tubulin gamma complex associated protein 2 (TUBGCP2), Receptor accessory protein 5 (REEP5), Solute carrier family 4 member 1 adaptor protein (SLC4A1AP), Integrin subunit beta 7 (ITGB7), Protein-O-mannose kinase (POMK), ER membrane protein complex subunit 7 (EMC7), Nuclear autoantigenic sperm protein (NASP), Checkpoint with forkhead and ring finger domains (CHFR), Ribosomal RNA processing 1 (RRP1), Cytosolic iron-sulfur assembly component 1 (CIAO1), Pumilio RNA binding family member 1 (PUM1), Retention in endoplasmic reticulum sorting receptor 1 (RER1), Serine and arginine rich splicing factor 4 (SRSF4) (see **Fig. 12B**). The expression of housekeeping genes is assumed to be relatively constant across samples. For example, for samples containing the same amount of RNA, the number and expression of housekeeping

genes will be similar. For example, for samples containing a smaller amount of RNA, the number and expression of housekeeping genes will be fewer than samples containing a larger amount of RNA, or *vice versa*. Therefore, the enumeration of RNA molecules of housekeeping genes on average can be used for the normalisation of RNA molecules of gene alteration targets and target genes related to protein expression.

[0049] In one example, the amount of cfRNA used in the method disclosed herein is at least 6 ng. In another example, the amount of cfRNA used in the method disclosed herein is about 6 ng to about 100 ng, or about 10ng, or about 20 ng, or about 30 ng, or about 40 ng, or about 50 ng, or about 60 ng, or about 70 ng, or about 80 ng, or about 90 ng, or about 100 ng. In one example, the amount of cfRNA used in the method disclosed herein is 20 ng to 50 ng.

[0050] A multiplexed PCR reaction is then performed on the converted cDNA as disclosed in step (b) of the first aspect, using a plurality of forward and reverse primers pairs specific to a plurality of **target genes that are capable of undergoing genomic alteration** as disclosed in (b)(I), and/or a plurality of forward and reverse primer pairs specific to a plurality of **control housekeeping genes** as disclosed in (b)(II), and/or a plurality of forward and reverse primer pairs specific to a plurality of **target genes related to protein expression** as disclosed in (b)(III), wherein the plurality of forward and reverse primer pairs specific to a plurality of **target genes that are capable of undergoing genomic alteration** differ from that of a plurality of **control housekeeping genes**, and differ from that of a plurality of **target genes related to protein expression**.

[0051] In one example, the plurality of multiplexed PCR reaction on the converted cDNA in step (b) is performed using a plurality of forward and reverse primer pairs specific to a plurality of **target genes that are capable of undergoing genomic alteration** as disclosed in step (b)(I), a plurality of forward and reverse primer pairs specific to a plurality of **control housekeeping genes** as disclosed in step (b)(II), and a plurality of primer sets specific to a plurality of **target genes related to protein expression** as disclosed in step (b)(III). In one example, the plurality of multiplexed PCR reactions on the converted cDNA in step (b) is performed using a plurality of forward and reverse primer pairs specific to a plurality of **target genes that are capable of undergoing genomic alteration** disclosed in step (b)(I) and a plurality of forward and reverse primer pairs specific to a plurality of **control housekeeping genes** as disclosed in step (b)(II). In another example, the plurality of multiplexed PCR reactions on the converted cDNA in step (b) is performed using a plurality of forward and reverse primers specific to a plurality of **control housekeeping genes** as disclosed in step (b)(II) and a plurality of primer sets specific to a plurality of **target genes related to protein expression** as disclosed in step (b)(III). In one

example, the plurality of multiplexed PCR reactions on the converted cDNA in step (b) is performed using a plurality of forward and reverse primer pairs specific to a plurality of **target genes that are capable of undergoing genomic alteration** as disclosed in step (b)(I) and a plurality of forward and reverse primer pairs specific to a plurality of **target genes related to**
 5 **protein expression** as disclosed in step (b)(III).

[0052] In one example, the multiplexed PCR reaction is performed on the converted cDNA using Platinum SuperFi II DNA Polymerase (Invitrogen), KAPA HiFi DNA Polymerase (Roche), Platinum Taq DNA Polymerase or Platinum SuperFi DNA Polymerase (Invitrogen) and Q5 High-Fidelity DNA Polymerase (NEB), etc.

10 [0053] In one example, the plurality of multiplexed PCR reactions performed on the converted cDNA comprises 3 to 15 PCR cycles. In one example, the PCR amplification comprises 3 PCR cycles. In one example, the PCR amplification comprises 4 PCR cycles. In one example, the PCR amplification comprises 5 PCR cycles. In one example, the PCR amplification comprises 6 PCR cycles. In one example, the PCR amplification comprises 7 PCR cycles. In one example,
 15 the PCR amplification comprises 8 PCR cycles. In one example, the PCR amplification comprises 9 PCR cycles. In one example, the PCR amplification comprises 10 PCR cycles. In one example, the PCR amplification comprises 11 PCR cycles. In one example, the PCR amplification comprises 12 PCR cycles. In one example, the PCR amplification comprises 13 PCR cycles.

20 [0054] In one example, the number of the forward and reverse primer pairs specific to the plurality of **target genes that are capable of undergoing genomic alteration** as disclosed in step (b)(I) is at least 100. In another example, the number of the plurality of forward and reverse primer pairs specific to the plurality of **target genes that are capable of undergoing genomic alteration** as disclosed in step (b)(I) is from 100 to 2000. In one example, the number of the
 25 forward and reverse primer pairs specific to the plurality of **target genes that are capable of undergoing genomic alteration** as disclosed in step (b)(I) is from 200 to 1900, or from 300 to 1800, or from 400 to 1700, or from 500 to 1600, or from 600 to 1500, or from 700 to 1400, or from 800 to 1300, or from 900 to 1200, or from 1000 to 1100. In one example, the number of the plurality of forward and reverse primer pairs specific to the plurality of **target genes that**
 30 **are capable of undergoing genomic alteration** as disclosed in step (b)(I) is about 100, about 200, about 300, or about 400, or about 500, or about 600, or about 700, or about 800, or about 900, or about 1000, or about 1100, or about 1200, or about 1300, or about 1400, or about 1500, or about 1600, or about 1700, or about 1800, or about 1900, or about 2000. In one example, there is no upper limit on the number of the plurality of forward and reverse primer pairs

specific to the plurality of **target genes that are capable of undergoing genomic alteration** as disclosed in step (b)(I).

[0055] In one example, the number of the plurality of forward and reverse primer pairs specific to the plurality of **control housekeeping genes** as disclosed in step (b)(II) is at least 20. In one example, the number of the plurality of forward and reverse primer pairs specific to the plurality of **control housekeeping genes** as disclosed in step (b)(II) is from 20 to 300. In one example, the number of the plurality of forward and reverse primer pairs specific to the plurality of **control housekeeping genes** as disclosed in step (b)(II) is from 30 to 290, or from 40 to 280, or from 50 to 260, or from 60 to 250, or from 70 to 240, or from 80 to 230, or from 90 to 220, or from 100 to 210, or from 110 to 200, or from 120 to 190, or from 130 to 180, or from 140 to 170. In one example, the number of the plurality of forward and reverse primer pairs specific to the plurality of **control housekeeping genes** as disclosed in step (b)(II) is about 20, or about 30, or about 40, or about 50, or about 60, or about 70, or about 80, or about 90, or about 100, or about 110, or about 120, or about 130, or about 140, or about 150, or about 160, or about 170, or about 180, or about 190, or about 200, or about 210, or about 220, or about 230, or about 240, or about 250, or about 260, or about 270, or about 280, or about 290, or about 300. In one example, there is no upper limit on the number of the plurality of forward and reverse primer pairs specific to the plurality of **control housekeeping genes** as disclosed in step (b)(II).

[0056] In one example, the number of the plurality of forward and reverse primer pairs specific to the plurality of **target genes related to protein expression** as disclosed in step (b)(III) is at least 10. In one example, the number of the plurality of forward and reverse primer pairs specific to the plurality of **target genes related to protein expression** as disclosed in step (b)(III) is from 10 to 1700. In one example, the number of the plurality of forward and reverse primer pairs specific to the plurality of **target genes related to protein expression** as disclosed in step (b)(III) is from 10 to 1700, or from 100 to 1600, or from 200 to 1500, or from 300 to 1400, or from 400 to 1300, or from 500 to 1200, or from 600 to 1100, or from 700 to 1000. In one example, the number of the plurality of forward and reverse primer pairs specific to the plurality of **target genes related to protein expression** as disclosed in step (b)(III) is about 10, or about 100, or about 200, or about 300, or about 400, or about 500, or about 600, or about 700, or about 800, or about 900, or about 1000, or about 1100, or about 1200, or about 1300, or about 1400, or about 1500, or about 1600, or about 1700. In one example, there is no upper limit on the number of the plurality of forward and reverse primer pairs specific to the plurality of **target genes related to protein expression** as disclosed in step (b)(III).

[0057] In another example, the maximum total number of plurality of forward and reverse primer pairs in the multiplexed PCR reaction is about 4000, wherein the number of plurality of forward and reverse primer pairs specific to a plurality of **target genes that are capable of undergoing genomic alteration** as disclosed in step (b)(I) is about 2000, wherein the number of plurality of forward and reverse primer pairs specific to a plurality of **control housekeeping genes** as disclosed in step (b)(II) is about 300, and wherein the number of plurality of forward and reverse primer pairs specific to a plurality of **target genes related to protein expression** as disclosed in step (b)(III) is about 1700.

[0058] In one example, the plurality of **target genes that are capable of undergoing genomic alterations** comprise an exon from a gene known to undergo fusion fused to an exon from a partner gene of the gene known to undergo fusion. In one example, the gene known to undergo fusion is selected from the group consisting ALK receptor tyrosine kinase, RET proto-oncogene, ROS proto-oncogene 1, fibroblast growth factor receptor 1 (FGFR1), fibroblast growth factor receptor 2 (FGFR2), fibroblast growth factor receptor 3 (FGFR3), neurotrophic receptor tyrosine kinase 1 (NTRK1), neurotrophic receptor tyrosine kinase 2 (NTRK2), neurotrophic receptor tyrosine kinase 3 (NTRK3), neuregulin 1 (NRG1), B-Raf proto-oncogene, serine/threonine kinase (BRAF), transmembrane serine protease 2 (TMPRSS2), MET proto-oncogene, receptor tyrosine kinase (MET), epidermal growth factor receptor (EGFR), estrogen receptor 1 (ESR1), platelet derived growth factor receptor alpha (PDGFRA), androgen receptor (AR), BCR activator of RhoGEF and GTPase (BCR), core-binding factor subunit beta (CBFB), lysine methyltransferase 2A (KMT2A), nucleophosmin 1 (NPM1), PML nuclear body scaffold (PML), and RUNX family transcription factor 1 (RUNX1).. In one example, the partner gene of the gene known to undergo fusion is selected from the group consisting of EMAP like 4 (EML4), kinesin family member 5B (KIF5B), coiled-coil domain containing 6 (CCDC6), CD74 molecule (CD74), transforming acidic coiled-coil containing protein 3 (TACC3), ezrin (EZR), ETS transcription factor ERG (ERG), ArfGAP with GTPase domain, ankyrin repeat and PH domain 3 (AGAP3), A-kinase anchoring protein 9 (AKAP9), KIAA1549, tropomyosin 3 (TMP3), translocated promoter region, nuclear basket protein (TPR), trafficking from ER to golgi regulator (TFG), lamin A/C (LMNA), BicC family RNA binding protein 1 (BICC1), RAD51 recombinase (RAD51), CD47 molecule (CD47), Yes1 associated transcriptional regulator (YAP1), ETS variant transcription factor 1 (ETV1), ETS variant transcription factor 4 (ETV4), ETS variant transcription factor 5 (ETV5), ETS variant transcription factor 6 (ETV6), factor interacting with PAPOLA and CPSF1 (FIP1L1), centriolin (CNTRL), ABL proto-oncogene 1, non-receptor tyrosine kinase (ABL1),

AF4/FMR2 family member 1 (AFF1), MDS1 and EVI1 complex locus (MECOM), MLLT3 super elongation complex subunit (MLLT3), myosin heavy chain 11 (MYH11), PBX homeobox 1 (PBX1), retinoic acid receptor alpha (RARA), and RUNX1 partner transcriptional co-repressor 1 (RUNX1T1).

5 [0059] The method of the present disclosure is optimized for generating amplicons having certain sizes. The chosen length of 90-110 base pairs was considered optimal because products of shorter amplicons (<80 base pairs) would be less effectively retained through the multi-step library preparation method for amplicon sequencing. In one example, the length of the plurality of amplicons derived from cDNA in step (b) is 90 to 110 base pairs. In one example, the length
10 of the plurality of amplicons derived from cDNA in step (b) is about 90 base pairs, or about 100 base pairs, or about 110 base pairs.

[0060] The plurality of amplicons derived from the cDNA in step (b) are then purified, as disclosed in step (c) of the first aspect.

[0061] The method of the present disclosure is designed to involve size-based separation
15 (magnetic bead based) of smaller primer dimer artefacts to be removed and desired products to be retained, and excess primers to be enzymatically digested (e.g. using endonucleases and exonucleases). In one example, the purification of DNA is performed using an agent such as paramagnetic beads. In one example, the paramagnetic beads are selected from the group consisting of AMPure XP beads, SPRI beads, and Dynabeads. In one example, the
20 paramagnetic beads are AMPure XP beads.

[0062] Next, the purified plurality of amplicons is amplified using universal indexed adapter primers to generate a plurality of sequencing library, as disclosed in step (d) of the first aspect.

[0063] In one example, the amplification is performed by using KAPA Hifi HotStart ReadyMix, Phusion U Hot Start DNA Polymerase (Thermo Scientific), ZymoTaq DNA
25 Polymerase (Zymo Research) and Q5U Hot Start High-Fidelity DNA Polymerase (NEB), etc.

[0064] In one example, each universal indexed adapter primer as disclosed in step (d) comprises an adapter sequence. In one example, the term “adapter sequence” refers to any nucleotide sequence which can be added to an oligonucleotide of interest to prepare said oligonucleotide of interest for various purposes. The adapter sequences are complementary to
30 the plurality of oligonucleotides present on the surface of the flow cells of the sequencing tools thereby allowing the DNA fragment to attach to the sequencing tool. In some examples, an adapter sequence allows for the sequencing of the oligonucleotide of interest. Sequencing platform specific adapter sequences are known in the art, and include, for example, the Illumina P5/P7 adapter sequences.

[0065] In one example, the universal indexed adapter primers as disclosed in step (d) of the method of the first aspect comprise:

a forward primer comprising the sequence of

AATGATACGGCGACCACCGAGATCTACACCTAGCGCTACACTCTTTCCCTACACG

5 ACGCTCTTCCGATC*T (SEQ ID NO: 616); and

a reverse primer comprising the sequence of

CAAGCAGAAGACGGCATACGAGATAAACCGCGGGTGACTGGAGTTCAGACGTGTGCTCTTCCGATC*T, (SEQ ID NO: 617),

wherein “*” represents a phosphorothioate bond, and wherein the underlined sequences are the barcode sequences. The plurality of sequencing library formed is then purified, as disclosed in step (e) of the first aspect.

[0066] In one example, the purification of the plurality of sequencing library is performed using an agent such as paramagnetic beads. In one example, the paramagnetic beads are selected from the group consisting of AMPure XP beads, SPRI beads, and Dynabeads. In one example, the paramagnetic beads are AMPure XP beads.

[0067] The purified plurality of sequencing library is then subjected to multiplex sequencing on a next-generation sequencing platform, as disclosed in step (f) of the first aspect, to obtain a plurality of sequencing reads.

[0068] In one example, the plurality of sequencing library is sequenced on NextSeq 550, NovaSeq 6000, or BGI MGISEQ-2000, DNBSEQ-G400, DNBSEQ-T7.

[0069] In one example, the plurality of sequencing library is qualified using the Agilent High Sensitivity DNA Screentape and quantified using KAPA Library Quantification Kit. In one example, the plurality of sequencing library is qualified by determining the size profile of the sequencing library which if successful will have a typical size profile of multiple prominent peaks greater than 200 base pairs (for example, as shown in **Fig. 15**).

[0070] Subsequently, a plurality of consensus reads is derived from each sequence of the plurality of sequencing reads obtained from step (f), as disclosed in step (g) of the first aspect.

[0071] In one example, step (g) of the first aspect further comprises:

- (g)(I) detecting the presence of the barcode sequence from each sequencing read,
- 30 (g)(II) performing cluster reassignment for the plurality of sequencing reads having the same barcode sequence to generate a plurality of barcode clusters, wherein each barcode cluster contains reads from the same amplicon and with the same barcode sequence, and

(g)(III) performing consensus calling for each barcode cluster to obtain the consensus read of each sequence.

[0072] The derived consensus sequence is aligned to the reference genome, as disclosed in step (h) of the first aspect. In one example, the term “reference genome” refers to DNA sequences known in the art that may be obtainable from public databases. In one example, the term “consensus read” refers to a nucleotide sequence obtained from consensus calling. In one example, consensus calling is performed by identifying the nucleotide at each position for each sequencing result within the subgroup, comparing the identity for the nucleotide at each position across the plurality of sequencing results, and determining a majority nucleotide at each position. If the majority nucleotide count is above a threshold set for determining majority for a specific position, the assignment for said position is the majority nucleotide. If the majority nucleotide count is below this threshold, no assignment is made for said position. The threshold is variable for every position and is a function of the total number of sequencing results corresponding to a specific position.

[0073] In one example, step (h) of the disclosed method further comprises, if the sequence alignment results in a partial alignment to the reference genome of an exon from a first gene and a partial alignment to the reference genome of an exon from a second gene as disclosed in step (h)(I), then the result is used to (i) determine the sequence alignment as a split read, (ii) count/enumerate the number of split reads from step (h)(I)(i) that supports a fusion junction, and (iii) if the number of split reads from step (h)(I)(ii) is two or more, determine the first gene and the second gene as fusion partners. In one example, step (h) of the disclosed method further comprises, if the sequence alignment results in an alignment to the reference genome of the control housekeeping gene as disclosed in step (h)(II), then the result is used to (i) determine the sequence alignment as a consensus read of the control housekeeping gene and (ii) count/enumerate consensus read pairs of the control housekeeping gene from step (h)(II)(i) to determine the level of gene expression of the control housekeeping gene. In one example, step (h) of the disclosed method further comprises, if the sequence alignment results in an alignment to the reference genome of the target gene related to protein expression as disclosed in step (h)(III), then the result is used to (i) determine the sequence alignment as a consensus read of the target gene related to protein expression and (ii) count/enumerate consensus read pairs of the target gene related to protein expression from step (h)(III)(i) to determine the level of gene expression of the target gene related to protein expression.

[0074] In one example, “consensus read pairs” refers to the consensus sequence called after collapsing all sequencing reads containing the same barcode sequence and primer pair. Each

consensus read pair, for example, is presumed to belong to an original RNA molecule converted to cDNA. In one example, counting/enumerating as disclosed in step (h) is achieved based on the barcode sequence-based consensus counting, wherein each RNA molecule comprising the same barcode sequence and primer pair combination represents a unique RNA molecule. In one example, all reverse primers of the plurality of forward and reverse primer pairs as disclosed in step (b) of the first aspect comprise a barcode sequence. Therefore, all RNA molecules captured by a given barcode sequence and primer pair combination can be detected and counted/enumerated.

[0075] In one example, the alignment of the derived plurality of consensus sequence to the reference genome is performed using a sequence alignment tool. In one example, the alignment tool is STAR, HISAT2, bwa, CLC, RSEM, kallisto, salmon, etc.

[0076] The results of sequence alignment from step (h) is used to determine presence or absence of the genomic alteration and/or determine presence or absence of gene expression and/or quantify the level of gene expression as disclosed in step (i) of the first aspect.

[0077] In one example, the disclosed method further comprises visualisation and fusion calling of the sequence alignment from step (h)(I). In one example, the visualisation is performed using Integrated Genome Viewer, or Savant Genome Browser, etc. In one example, the fusion calling is performed using Arriba and Fusion Catcher, etc.

[0078] In one example, the step of determining presence or absence of the genomic alteration and/or determining presence or absence of the gene expression and/or quantifying the level of the gene expression, further comprises performing variant calling of the sequence alignment from step (h). In one example, the step of determining presence or absence of the genomic alteration and/or determining presence or absence of the gene expression and/or quantifying the level of the gene expression, further comprises performing variant calling of the sequence alignment from step (h)(II). In one example, the step of determining presence or absence of the genomic alteration and/or determining presence or absence of the gene expression and/or quantifying the level of the gene expression, further comprises performing variant calling of the sequence alignment from step (h)(III). In one example, the step of variant calling comprises: (i) identifying differences between a consensus read and a reference genome based on the sequence alignment from step (h); and ii) determining the read count of sequence alignments comprising genomic alteration. In one example, the step of variant calling comprises: (i) identifying differences between a consensus read and a reference genome based on the sequence alignment from step (h)(II); and ii) determining the read count of sequence alignments comprising genomic alteration. In one example, the step of variant calling

comprises: (i) identifying differences between a consensus read and a reference genome based on the sequence alignment from step (h)(III); and ii) determining the read count of sequence alignments comprising genomic alteration. In one example, the genomic alteration is selected from the group comprising of insertions (e.g., duplications), deletions, and single nucleotide variants. In one example, the variant calling is performed using Mutect2 and a custom variant caller.

[0079] In one example, wherein the disclosed method of the first aspect is used to simultaneously detect gene expression, structural rearrangements and quantify gene expression in cfRNA from a biological sample, the expression level of genes that are known to be overexpressed in cancer cells is quantified. In one example, wherein the disclosed method of the first aspect is used to simultaneously detect genomic alteration in cfRNA and quantify gene expression in cfRNA from a biological sample, the expression level of target genes that have undergone genomic alterations is quantified. In one example, wherein the disclosed method of the first aspect is used to simultaneously detect gene expression and quantify gene expression of cfRNA, the expression level of target genes related to protein expression is quantified.

[0080] In one example, the statistical modelling technique used to visualise the level of expression of genes related to protein expression is heatmap visualisation, principal component analysis, hierarchical clustering, etc.

[0081] In a second aspect, the present disclosure refers to a kit for detecting genomic alteration and/or detecting gene expression and/or quantifying the level of gene expression using RNA in a biological sample according to the method of the first aspect, wherein the kit comprises:

(a) a plurality of forward and reverse primer pairs specific to a plurality of **target genes that are capable of undergoing genomic alteration as defined in step (b)(I) of the method of the first aspect;**

(b) a plurality of forward and reverse primer pairs specific to a plurality of **control housekeeping genes as defined in step (b)(II) of the method of the first aspect;** and

(c) a plurality of primer sets specific to a plurality of **genes related to protein expression as defined in step (b)(III) of the method of the first aspect.**

[0082] In one example, a person skilled in the art would be able to design the plurality of primer pairs and primer sets in (a), (b) and (c) of the kit of the second aspect based on the disclosure herein, for example, as described in steps (b)(I), (b)(II) and (b)(III) of the method of the first aspect. In one example, the plurality of primer sets specific to a plurality of genes related to protein expression as defined in step (b)(III) of the method of the first aspect provided in the kit as described herein may be used for determining presence or absence of the genomic

alteration. In one example, the plurality of primer sets specific to a plurality of genes related to protein expression as defined in step (b)(III) of the method of the first aspect provided in the kit as described herein may be used for determining presence or absence of the genomic alteration such as deletions, insertions (e.g., duplications) and single nucleotide variations. In one example, the plurality of primer sets specific to a plurality of genes related to protein expression as defined in step (b)(III) of the method of the first aspect provided in the kit as described herein may be used for determining presence or absence of the genomic alteration by further performing the step of variant calling as described herein. In one example, the genomic alteration may be single nucleotide variation, insertion (e.g., duplications) or deletion.

10 In one example, the kit for detecting genomic alteration and/or detecting gene expression and/or quantifying the level of gene expression of cfRNA in a biological sample according to the method of the first aspect further comprises a buffer for performing a plurality of multiplexed PCR reactions, a reverse transcriptase, a DNA polymerase, and a plurality of deoxynucleotide triphosphates (dNTPs). In some examples, the reagents provided in the kit as described herein may be provided in separate containers comprising the components independently distributed in one or more containers. As the method as described herein relates to sequencing (such as high-throughput sequencing), further components required in sequencing process could be easily determined by the person skilled in the art.

[0083] As used in this application, the singular form “a,” “an,” and “the” include plural references unless the context clearly dictates otherwise. For example, the term “a primer” includes a plurality of primers, including mixtures and combinations thereof.

[0084] As used herein, the terms “increase” and “decrease” refer to the relative alteration of a chosen trait or characteristic in a subset of a population in comparison to the same trait or characteristic as present in the whole population. An increase thus indicates a change on a positive scale, whereas a decrease indicates a change on a negative scale. The term “change”, as used herein, also refers to the difference between a chosen trait or characteristic of an isolated population subset in comparison to the same trait or characteristic in the population as a whole. However, this term is without valuation of the difference seen.

[0085] As used herein, the term “about” in the context of concentration of a substance, size of a substance, length of time, or other stated values means +/- 5% of the stated value, or +/- 4% of the stated value, or +/- 3% of the stated value, or +/- 2% of the stated value, or +/- 1% of the stated value, or +/- 0.5% of the stated value.

[0086] Throughout this disclosure, certain embodiments may be disclosed in a range format. It should be understood that the description in range format is merely for convenience and

brevity and should not be construed as an inflexible limitation on the scope of the disclosed ranges. Accordingly, the description of a range should be considered to have specifically disclosed all the possible sub-ranges as well as individual numerical values within that range. For example, description of a range such as from 1 to 6 should be considered to have

5 specifically disclosed sub-ranges such as from 1 to 3, from 1 to 4, from 1 to 5, from 2 to 4, from 2 to 6, from 3 to 6 etc., as well as individual numbers within that range, for example, 1, 2, 3, 4, 5, and 6. This applies regardless of the breadth of the range.

[0087] The present disclosure illustratively described herein may suitably be practiced in the absence of any element or elements, limitation or limitations, not specifically disclosed herein.

10 Thus, for example, the terms "comprising", "including", "containing", etc. shall be read expansively and without limitation. Additionally, the terms and expressions employed herein have been used as terms of description and not of limitation, and there is no intention in the use of such terms and expressions of excluding any equivalents of the features shown and described or portions thereof, but it is recognized that various modifications are possible within the scope

15 of the disclosure claimed. Thus, it should be understood that although the present disclosure has been specifically disclosed by preferred embodiments and optional features, modification and variation of the present disclosure embodied therein herein disclosed may be resorted to by those skilled in the art, and that such modifications and variations are considered to be within the scope of this present disclosure.

20 [0088] The invention has been described broadly and generically herein. Each of the narrower species and subgeneric groupings falling within the generic disclosure also form part of the present disclosure. This includes the generic description of the present disclosure with a proviso or negative limitation removing any subject matter from the genus, regardless of whether or not the excised material is specifically recited herein.

25 [0089] Other embodiments are within the following claims and non-limiting examples.

EXAMPLES

[0090] **Methods**

[0091] ***Sample collection and Processing***

[0092] Blood collected into Streck Cell-free DNA BCT® was shipped at ambient temperature

30 before plasma separation. Briefly, blood plasma was prepared using a 2-step centrifugation process: first centrifugation was done at 1500 x g for 10 min at 4 °C to separate plasma. The plasma layer was transferred to a separate tube and centrifuged at 15,000 × g for 10 min at 4 °C to further remove cellular contaminants, and immediately processed for nucleic acid

extraction or stored at -80°C until used for extraction. If frozen, the plasma was fully thawed at room temperature before extraction.

[0093] Plasma cell-free total nucleic acids were extracted using the QIAamp Circulating Nucleic Acids kit (Qiagen). The nucleic acid extract contains co-eluted cfDNA and cfRNA fractions. The cfDNA was quantified using the Qubit Fluorometer (Thermo Fisher Scientific), and sized using the Genomic DNA ScreenTape on the 4200 TapeStation (Agilent). cfRNA was quantified and profiled using the Bioanalyzer RNA 6000 Pico kit or the High Sensitivity RNA Screentape on the 4200 TapeStation.

[0094] *Design of primers for fusions and expression in a sequencing library*

[0095] A highly multiplex amplicon-based NGS assay was designed to capture potential fusions in cfRNA sample. Depending on the expected orientation of a partner exon in a fusion gene, a primer upstream of the exonic fusion junction ("forward" primer) or downstream of the fusion junction ("reverse" primer) was designed for the target gene's exon. Broadly, multiple exon-flanking primers were designed for target genes that are known to participate in fusion events in cancer. For all downstream primers, a random 10-base pairs barcode sequence was incorporated upstream of the gene-specific sequence for consensus calling and unique molecule enumeration. A pool of >300 "forward" primers and >300 "reverse" primers was prepared. A multiplicity of "upstream" and "downstream" primers were included in the multiplex PCR to optimally capture potential fusions known to occur between genes. The design of primers included exons of well-characterized genes known to undergo fusions and the addition of barcode sequence primers allowed for accurate enumeration of copies of RNA transcript as per method of enumeration (**Fig. 1**).

[0096] For the capture of transcripts corresponding to control genes and other genes for which expression was to be quantified, primers were designed such that at least one primer of a pair landed on an exon-exon junction, or the primer pairs were within two exons intervened by an intron >5000 base pairs in length. These primers were also included in the final primer pools. The specificity of cfRNA amplification was verified by performing the whole cfRNA sequencing workflow, but with leaving out the reverse transcriptase enzyme during the complementary DNA preparation. Any sequencing for intended regions, particularly control and expression genes, when no reverse transcription was performed, could then be attributed to the primers amplifying cfDNA. Any such primers were redesigned to improve specificity for RNA by reducing the 3' exon span of the exon-exon spanning primer. The design of primers for target genes related to expression were similar to the control gene targets, and at least one primer of primer pair spanned an exon-exon junction, and two or more primer pairs were

designed per target gene covering both 5' and 3' end exons, to more reliably capture expression of target genes for expression, by allowing one or more amplicons to represent a given target gene. A highly multiplexed primer pool was employed with a plurality of upstream and downstream primers, some of which are expected to generate sequenceable targets in most samples depending on expression variability, and some primers which are expected to generate a product only when a sample is positive for structural rearrangement, generating a fusion gene that is productively expressed. The primers additionally carried the appropriate extensions necessary for generating sequenceable libraries with sequencing adapters for Illumina sequencing (**Figs. 2A and 2B**).

10 [0097] ***Preparation of cfRNA Sequencing library***

[0098] Between 20-50 ng of cfRNA was converted to complementary DNA (cDNA) using the High-Capacity cDNA Reverse Transcription Kit (Thermo Fisher Scientific) in a total volume of 20 ul using random primers. The converted cDNA was used as a template in a highly multiplexed PCR reaction for target capture using the Platinum™ SuperFi II DNA Polymerase (Thermo Fisher Scientific). Briefly, cDNA was combined with primers and DNA Polymerase in a single reaction, and subjected to 3 to 15 cycles of PCR with the following conditions: 98°C, 1 min; 60°C, 1 min; 72°C, 1 min, followed by a final extension for 5 min at 72°C. The amplification product was subjected to one round of enzymatic digestion (using exonucleases, ExoI and ExoT) and two rounds of clean-up using 1.8x volume of AMPure XP beads and eluted in Buffer EB or nuclease-free water. The purified PCR products were then amplified with universal indexed adapter primers, compatible for sequencing with Illumina platform, with primers using KAPA HiFi HotStart ReadyMix. The final amplified library was purified with two rounds of 0.8x volume AMPure XP beads to remove excess adapters and size-select the final sequencing library. Library was quantified using the High Sensitivity DNA Screentape and quantified using KAPA Library Quantification Kit. Each library was sequenced on a Nextseq 550 to a depth of 3 million paired-end reads per sample.

[0099] ***Data Analysis***

[00100] FASTQ files were processed using a custom pipeline. First, sequenced amplicons were identified and labelled in the FASTQ files based on the presence of any potential primer sequences in the right directionality, upstream or downstream (from a predetermined list of primer sequences based on panel design) in Read 1 and paired Read 2. Barcode sequence sequences in read 1 were identified upstream of primer in Read 1 and trimmed using cutadapt. The extracted molecular tag sequences were used to derive the consensus read sequence for all duplicate reads of a sequence identifiable by a given primer

pair and unique barcode sequence. The consensus reads were then written to a new FASTQ file and aligned to human genome reference hg19 using STAR aligner. Fusion reads in which non-contiguous regions of the genome are captured within a read were identified as split reads and fusion partners were identified based on the sequence alignment. The presence of split read sequences mapping to two reciprocal partner genes were additionally confirmed to have been captured by primers specific to identified genes. Number of split reads (read pairs) supporting a fusion junction were enumerated. Visualization and fusion calling were also performed using Arriba and FusionCatcher. At least 2 supporting split reads were required for calling fusion and exon skipping variants (transcript variants). With molecular barcoding, the sequencing is error-free and increases confidence fusion calls due to the high quality of sequencing data.

[00101] Expression-level analysis was done by enumerating consensus read pairs which supported a given amplicon predefined by primer pairs for expression. Read pair counts were enumerated and tabulated for downstream analysis as control genes or target genes. Variant calling was performed on consensus BAM files using Mutect2 and a custom variant caller to identify single nucleotide variations, insertion and deletion mutations relative to the reference sequence. Expression of mutant transcripts comprising single nucleotide variations, insertion and deletion was quantified based on the number of reads containing the particular single nucleotide variation, insertion or deletion mutation and mapping to the intended target region. Expression of wild-type transcripts was quantified based on the number of reads matching the reference sequence and mapping to the intended target region. The relative expression of each mutation was also determined based on the proportion of mutant read counts relative to total read counts.

[00102] **Results**

[00103] The present disclosure describes a method for the simultaneous detection and quantification of clinically relevant genomic and gene expression alteration using cfRNA, with high sensitivity, specificity, and minimal invasive procedures.

[00104] ***Validation of cfRNA-based detection assay: relative abundance of cell-free nucleic acid in plasma***

[00105] Total cfRNA concentration from plasma of healthy individuals and cancer patients were characterized for presence of cfRNA and analyzed for fragment size distribution using Bioanalyzer RNA 6000 Pico assay. cfRNA was present in all cancer samples and showed a predominant peak at 110 to 120 nucleotides in size and a second population of RNA in the 200 to 300 nucleotides range (**Fig. 3A and 3B**). In terms of relative abundance, the shorter fragments (110 to 120 nucleotides) were about 5 to 10 times more abundant than the larger size

RNA fragments (200 to 300 nucleotides). cfRNA from healthy individuals also showed the same pattern of size distribution, but at significantly lower cfRNA concentrations (**Fig. 3C and 3D**).

[00106] Total nucleic acid extracts comprising cfDNA and cfRNA from plasma of healthy and cancer individuals were analyzed. Relative to each extract's cfDNA concentration, cfRNA concentrations were generally lower, and differed most significantly when the concentration of cfDNA exceeded 10 ng/ml plasma (**Fig. 4**).

[00107] *Technical validation of cfRNA-based multiplex amplicon sequencing detection with RNA extracted from cancerous cell lines*

[00108] The method described herein showed the ability to detect fusions using RNA extracted from cultured cancer cell lines known to harbor fusion genes, such as CRL-9591 (KMT2A-AFF1), H2228 (EML4-ALK), CRL-2724 (RUNX1-RUNX1T1), VCaP (TMPRSS2-ERG) and CRL-5813 (TMPRSS2-ERG). As RNA from cultured cells is relatively intact compared for plasma cfDNA, the cell line RNA was subjected to ultrasonication (using Covaris) in order to more closely resemble the size of cfRNA. The resulting fraction as used to mimic cfRNA to demonstrate the performance of the multiplex amplicon sequencing for the detection of a variety of known fusions (**Fig. 5A, 5B and 5C**). This was used to provide adequate material to mimic cfRNA to demonstrate the performance of the multiplex amplicon sequencing for the detection of a variety of known fusions. RNA-based detection of fusions in all five cancer cell lines was successful (**Fig. 7**). The plurality of sequencing library obtained can be qualified using the Agilent High Sensitivity DNA Screentape as shown in **Fig. 15**, which illustrates a typical library profile for a cfRNA sample converted to a sequencing library as seen on the High Sensitivity DNA Screentape. The multiple peaks >200 base pairs correspond to the multiple products encompassing potential fusion products, control gene products and other gene expression products for which the multiplicity of forward and reverse primers are included. Qualified libraries will have prominent peaks of size >200 base pairs.

[00109] Sequence alignments to the reference genome showed the capture of sequencing reads with partial alignment to the target exon, and partial alignment to another part of the genome sequence of which corresponds to the partner gene exon, known as a split read, and confirmed the detection of the EML4-ALK fusion transcript in H2228 cell line with as little as 1 ng of fragmented RNA with 8364 reads supporting the split configuration. (**Figs. 6A and 6B**). The alignment of split reads showed that fusions in cancer cell lines: NCI-H660 cell line (CRL-5813, ATCC), VCaP cell line (CRL-2876, ATCC), Human MV-4-11 cell line (CRL-

9591, ATCC) and Kasumi-1 (CRL-2724, ATCC), were accurately detected as visualized by Arriba tool for detection of fusions in RNA sequencing data using the multiplex amplicon sequencing method in fragmented RNA. (**Fig. 7**).

[00110] ***Data comparison between cfDNA and cfRNA-based detection assay***

5 [00111] Nucleic acid extracts from plasma of two cancer patients previously characterized to be positive for fusions using a DNA-based method (Liquid Hallmark) were tested. In the first case of a metastatic prostate cancer, TMPRSS2-ERG fusion was detected in cfDNA (using 70 ng of cfDNA) supported by 17 split reads mapping to intronic position chr21:42867069 within TMPRSS2 (intron 2 of TMPRSS2-NM_005656.4) and intronic
10 position chr21:39818058 within ERG (intron 3 of ERG- NM_001291391.1) (**Fig. 8A**). Using the same circulating nucleic acid extract, fusion in the cfRNA (equivalent of just 24 ng cfRNA) was detected with 4123 supporting split reads, fusing exon 2 of TMPRSS2 (chr21: 21:42870045) with exon 4 of NM_001291391.1 (or exon 2 of ERG NM_182918.4) (chr 21:39817544) (**Fig. 8B**).

15 [00112] In a second case of metastatic lung cancer, a CCDC6-RET fusion detected using cfDNA (breakpoints CCDC6 Intron 1 (chr10:61623181) and RET intron 11 (chr10:43611035) and cfRNA CCDC6 exon 1 (10:61665879) and RET exon 12 (10:43612031). cfDNA was detected with 12 supporting reads, while fusion in cfRNA was supported by 13 split reads (**Figs. 9A and 9B**).

20 [00113] In a third clinical sample from a hematological malignancy (acute lymphoblastic leukemia) with BCR-ABL1 rearrangement confirmed in DNA from the peripheral blood cells, RNA was extracted from another fraction of archived buffy coat and tested with the multiplex amplicon sequencing method described here. The fusion between exon 14 of BCR, and exon 2 of ABL1 was readily detectable in the RNA fraction with an
25 abundant 159,106 supporting reads. The large number of supporting reads is indicative of the enrichment of transcripts with BCR-ABL1 fusion, due to increased expression and secondary enrichment of cancer cells positive for the fusion in sample tested (buffy coat RNA) (**Fig. 10A and Fig. 10B**).

[00114] Additional fusion events are shown in **Fig. 13** illustrating the identification of
30 actionable driver fusions in untreated lung cancer cases using cfRNA using the method described herein. **Fig. 13A**, **Fig. 13B** and **Fig 13C** show the detection of various gene fusion events namely LMNA-NTRK1 fusion, CD74-NRG1 and ETV6-NTRK3 fusion in cfRNA samples in three lung cancer cases, respectively. These mutations were otherwise undetectable using a DNA-based assay and appear as negative for the presence of other

driver gene mutations in cfDNA. In addition, when the method described used cfDNA and cfRNA for fusion detection in 45 lung cancer cases, additional fusions were identified when cfRNA fraction was used compared to cfDNA (**Fig. 14**). When testing for fusion was performed with DNA and RNA as sample input orthogonally, there were 12 cases with concordant fusion detection based on cfDNA and cfRNA as sample input. There were additional fusions detected in 5 cases and 1 fusion that was missed and not detected when cfRNA was used as sample input instead of cfDNA. The list of range of fusions detected by both cfDNA and cfRNA methods or by one of the two methods shown in **Fig. 14B**.

[00115] ***Limit of detection***

[00116] The limit of detection is defined as the lowest RNA concentration at which fusion events can be readily detectable. Initial determination of limit of detection of RNA-based fusion was done by quantitating the number of EML4-ALK fusion transcripts present in 1 ng of H2228 cell line RNA, from which EML4-ALK fusion was readily detectable using the method described herein (**Fig. 6 and Fig. 7**). The number of EML4-ALK fusion RNA transcripts was determined to be ~13.7 copies per 5 ng of RNA using a qRT-PCR assay designed specifically for the EML4-ALK transcript present in H2228 cells (**Fig. 11**). Therefore, the method described herein is shown to be able to detect down to 2.72 copies of EML4-ALK fusion (in 1 ng of H2228 RNA), suggesting a very sensitive detection for RNA-based fusions.

[00117] ***Simultaneous detection and quantification of expression cfRNA-based fusion events***

[00118] Besides the detection of fusions in cfRNA, simultaneous detection of targets genes intended for non-invasive expression monitoring was also done for cfRNA from cancer and healthy samples. In the same multiplex reaction, primers for 22 control genes and 13 amplicons for 6 genes related to immunotherapy response (CD274, PDCD1, CTLA4, LAG3, HAVCR2 and CD47) were included, and the combined target capture was performed. Based on read counts mapping to the intended target regions, the determination of expression level of each target was done. The range of expression levels were visualized in an expression heatmap (**Fig. 12**).

[00119] As healthy samples typically had very low yields of both cfRNA and cfDNA, the expression was low for control genes and immunotherapy response genes across along healthy samples as expected. However, among cancer samples, a range of expression patterns was observed, with some samples showing limited expression of nearly all targets, despite an equivalent amount of cfRNA material used in the method. The reliability of detection and the

quantitative ability of the method was demonstrated by the performing repeats of the same sample with differing amounts of cfRNA, which showed an increase in the expression read count, but similarity in pattern among the sample repeats (**Fig. 12**). The repeats are represented by C_20.126.1 and C_20.126.2 (sample 20.126 repeated) and C_20.1069.1 and C_20.1069.2 (sample 20.1069 repeated). In the heatmap, the two repeats are closest to each other indicating greater similarity between two repeats of same sample, compared to other samples.

[00120] ***Detection of expressed transcripts comprising deletion mutation in RNA sample***

[00121] The method described herein showed the ability to detect an 18-nucleotide deletion in an RNA sample extracted from FFPE lung tumor tissue. The expression of EGFR c.2240_2257del p.L747_P753delinsS mutant transcript (comprising the deletion) was detected with 4266 supporting reads (**Fig. 16**).

[00122] ***Detection of expressed transcripts comprising single nucleotide variations in RNA sample***

[00123] The method described herein showed the ability to detect single nucleotide variation in a cfRNA sample extracted from the plasma of a metastatic lung cancer patient. The expression of EGFR c.2573T>G p.L858R mutant transcript (comprising the single nucleotide variation) was supported by 112 reads. (**Fig. 17**)

[00124] ***Detection of expressed transcripts comprising single nucleotide variations, insertion and deletion mutation in RNA sample***

[00125] The method described herein showed the ability to detect single nucleotide variations, insertion and deletion mutations in tissue RNA extracted from FFPE tumor samples (**Fig. 18A**) and cfRNA extracted from plasma (**Fig. 18B**). Simultaneous detection of target genes intended for detection of expressed transcripts containing single nucleotide variations, insertion (e.g., duplications) and deletion mutations was performed for tumor tissue RNA from 4 cancer samples and plasma cfRNA from 3 cancer samples. In the same multiplex PCR reaction, primers for desired targets were included, and combined target capture was performed. The variant allele frequency (VAF) was determined based on the proportion of mutant read counts relative to total read counts detected from the method described herein. The validity of the RNA-based method described herein is shown by the VAF percentage depicted in **Figs. 18 A and 18B**.

[00126] **Discussion**

[00127] In the present disclosure, a method to simultaneously detect genomic alterations such as structural rearrangements, and gene expression using circulating cell-free RNA (cfRNA) is described. Such detection and quantification non-invasively is envisioned to allow
5 the detection of cancer, determination of prognosis and prediction of therapy response. The method is based on highly multiplexed amplicon-based NGS, and involves the tagging of individual cfRNA molecules using barcode sequences, and the optimized design of amplicons to be compatible with the fragmented nature of cfRNA. The inventors have shown that the method can be applied to circulating nucleic acid extracts containing both cfDNA and cfRNA,
10 and can detect and quantitate fusion RNA transcript and gene expression simultaneously, in such samples.

[00128] In order to detect structural rearrangements, such as gene fusions- that result in the juxtaposition of exons from different genes, resulting in a fusion transcript- from cfRNA analyte, a targeted multiplex amplicon panel for detection of fusions by next-generation
15 sequencing (NGS) was designed. The juxtaposition of gene exons was exploited to amplify fusion transcripts, by a pair of primers flanking the exonic junctions involved in the fusion. Primers specific for exons of fusion and partner genes known to undergo fusions were designed just flanking the exonic junction sites. Such juxtaposition of exons from different genes can only happen when processed mRNA is generated (by splicing) bringing fused exons together,
20 and so equivalent DNA sequences are unlikely to contribute to productive amplification with the same primers, because of the intervening fused relatively long introns separating the exons in the DNA.

[00129] The design of the primers to capture fusion transcripts had two main features –
1) the presence of a random barcode sequence in the downstream primers (downstream relative
25 to the fusion transcript) to individually tag each copy of the RNA fusion transcript if present, and 2) the location of each primer approximately 50 base pairs from each exonic junction in the panel, such that the expected total amplicon length would be close to 90-110 base pairs. This was done in order to meet the sample cfRNA size distribution observed which peaked at 110-120 nucleotides. The chosen length of 90-110 base pairs was considered optimal because
30 products of shorter amplicons (<80 base pairs) would be less effectively retained through the multi-step library preparation method for amplicon sequencing, involving size-based separation (magnetic bead based) of smaller primer dimer artefacts to be removed and desired products to be retained. A multiplicity of “upstream” and “downstream” primers were included in the multiplex PCR to optimally capture potential fusions known to occur between genes.

The design of primers includes exons of well-characterized genes known to undergo fusions, such as ALK, RET, ROS1, FGFR2, FGFR3 and exons of their partner genes, such as EML4, KIF5B, CCDC6, CD74, TACC3, among others. Potential fusions between any upstream and any downstream exon (not limited to gene pairs for which design was intended) can theoretically be detected if present in a sample, if the capture reaction simultaneously includes the multiplicity of primers. Broadly, primers to capture all exonic junctions known to undergo fusions (and intervening exons which may not have been previously reported to be involved in fusions) in target and partner genes were designed. The barcode sequence primers allow for accurate enumeration of copies of RNA transcript as per method of enumeration.

10 [00130] The first step in the process of preparing a cfRNA NGS library based on this method is the conversion of cfRNA (naturally fragmented) into complementary DNA (cDNA) using reverse transcriptase enzyme with random primers. The result of the reverse transcription reaction is a total complement of the cfRNA molecules present in the sample. In addition to the exon flanking primers for fusion detection, and in order to provide a quantitative measure of amount of cfRNA included in a reaction, primers were also included for several (>20) control housekeeping genes in the multiplex reaction. The purpose of capturing transcripts of genes expressed at some baseline line across all sample types, was to estimate an average abundance of cellular material going into the multiplex PCR reaction, and to serve as a control for the whole process of preparation of cfRNA sequencing libraries, including the sample extraction, reverse transcription, and PCR steps. The design of primers intended for control target genes differed from that of fusion targets, in that at least one primer of a control gene primer pair was designed to span an exon-exon junction, in order to prevent unintended amplification of DNA of the control target gene, and the resulting amplicon was ~100 base pairs in length (**Fig. 1**). The design of primers for target genes related to expression were similar to the control gene targets, and at least one primer of primer pair spanned an exon-exon junction, and two or more primer pairs were designed per target gene covering both 5' and 3' end exons, to more reliably capture expression of target genes for expression, by allowing one or more amplicons to represent a given target gene. A highly multiplexed primer pool was employed with a plurality of upstream and downstream primers, some of which are expected to generate sequenceable targets in most samples depending on expression variability, and some primers which are expected to generate a product only when a sample is positive for structural rearrangement, generating a fusion gene that is productively expressed. The primers additionally carried the appropriate extensions necessary for generating sequenceable libraries with sequencing adapters for Illumina sequencing.

[00131] In this disclosure, the use of cfRNA analyte for the enhanced detection of structural rearrangements and gene expression simultaneously, was demonstrated. This was achieved by the design of multiplex amplicon NGS assay encompassing the exons of genes involved in fusion and the design of amplicons for the expression of target genes, with use of barcode sequences and optimal size selection of amplicons for cfRNA applications. Overall presence of abundance was quantified by read density of accumulated read numbers. In this disclosure, issues related to whole-transcriptome sequencing including cost and manpower were partially overcome with the application of targeted sequencing for plasma cfRNA.

[00132] In the present disclosure, clinically relevant altered splicing events such as MET proto-oncogene, receptor tyrosine kinase (MET) exon 14 skipping, Androgen receptor (AR) transcript variants are approached as intra-gene fusions events and are designed to be captured if present using a combination of primers which would capture the aberrant splicing as the juxtaposition of exons of the same gene not normally observed, but that which can occur in cancers. The ability to quantify expression of relevant genes, non-invasively, for the prediction of response to various treatments is valuable, as it allows the longitudinal monitoring of response, and informs clinical decision. However, this has not been routinely implemented in clinical practice, and is largely limited to the detection of DNA level alterations such as mutations and genomic copy number changes. Using sequencing technology such as NGS, mutations are identified by comparing sequencing reads to reference sequences (genomes). Genomic copy number changes are quantified by counting the number of reads corresponding to a gene, and quantifying the deviation from normal copy number count expected from cells or samples having two copies of DNA per gene. In one example, DNA level alterations include single nucleotide variants leading to missense mutations, frameshift mutations, insertion-deletions, splice site mutations. The non-invasive monitoring of expression changes by accessing the cfRNA analyte can exploit the overexpression of tumor-specific transcripts, lead to amplification of tumor-derived RNA signals in blood, thereby increasing sensitivity of detection. For the non-invasive characterization of structural rearrangements, for example, gene fusions in plasma, typically targeted cfDNA-based next-generation sequencing (NGS)-based methods are utilized.

[00133] To overcome the issues of stability, appropriate RNA isolation procedures, removing DNA contamination and the use of endogenous housekeeping control genes, has been applied in this disclosure. Combined together, cfRNA can be used to provide precise information related to cancer diagnosis, prognosis and prediction of therapy response.

The novel features of the present disclosure and the reason why they are technologically significant are as follows:

1. The specific design of primers to allow amplification of consistently short amplicons to be able to amplify targets from cfRNA which is usually about 100 nucleotides in length when isolated from plasma.
2. The inclusion of barcode sequences in the primer design for accurate enumeration of specific targets, whether containing fusions or not.
3. The combination of designs for the simultaneous capture of fusions (if any) and target gene expression.
4. The ability to detect novel fusions with any potential primer combination included in the multiplex panel.
5. The design of data analysis workflows which allow the parallel analysis of RNA-based fusions and expression.

[00134] The method of the present disclosure has the following advantages:

1. The method of the present disclosure uses cfRNA (which lacks introns) for sample input, thereby allowing the identification of genetic fusions involving long introns which are typically excluded from conventional DNA-based assays.
2. The method of the present disclosure allows for identification of both fully characterised genomic alteration targets and novel genomic alteration targets (i.e. genomic alteration targets which are not previously characterised). Novel genomic alteration targets can be detected with any potential primer combination included in the multiplexed panel. The design of data analysis workflows which allow the parallel analysis of RNA-based fusions and expression.
3. The method of the present disclosure allows for simultaneous detection of structural rearrangement and determination of expression level of cfRNA. For cancer-related genes that are expressed, ctRNA provides the same mutational information as ctDNA; additionally, it can provide quantitative information about the expression levels of target genes of interest, and can potentially increase the sensitivity of detection of variants with low allelic frequencies due to the overexpression of tumor-specific transcripts. The ability to quantitate the expression of these targets non-invasively can be very useful for the monitoring treatment response and making treatment decisions.
4. The method of the present disclosure may be used on a blood-based test (for example, to detect fusion targets in cfRNA in the blood) that is fast and non-invasive (only one

draw of blood is needed). In addition, the method is scalable for the detection of multiple cancers in a single test and is suitable for cancer screening in an asymptomatic population.

- 5 5. The method of the present disclosure is highly sensitive compared to conventional methods of genomic structural alteration detection. Smaller starting material (cfRNA) is required for equivalent or better detection capabilities. For example, only 24 ng of cfRNA is required for detection of TMPRSS2-ERG fusion in a metastatic prostate cancer sample, as compared to using 70 ng of cfDNA to generate similar sequencing reads.
- 10 6. The technological significance lies in the generalizable use of primers for target capture, which allows working with smaller, limiting amounts of input of nucleic acid sample. In addition, the unique combination of targets is selected for the sensitivity and specific detection of multiple cancers.
- 15 7. The method of the present disclosure is scalable and allows the capture of multiple genomic regions for the identification of several cancer types in a single assay. The target gene coverage can be expanded by the addition of forward and reverse primer pairs.
- 20 8. The method of the present disclosure may be used in the following applications:
 - Detection, identification and quantification of well-characterised genomic alterations (such as gene fusions) that are clinically relevant, for example those associated with to cancers.
 - Identification of novel genomic alterations specific to cancers.
 - Cancer screening in healthy individuals and individuals at high risk for the tested cancers.
 - 25 • Disease monitoring in cancer patients, including monitoring response to treatment, such as immunotherapy.
- 30 9. Shorter fragments are more challenging as starting material for sequencing-based assays, due to restrictions on primer design and the sequence information that can be optimally captured. The method of the present disclosure uses cfRNA, which is shorter (about 100 nucleotides) in length compared to cfDNA (about 160 base pairs) in length. The primers described herein have been optimally designed to capture fragmented cfRNA of about 100 nucleotides in length to maximize sensitivity of detection of fusions and expression changes.

10. The method of the present disclosure uses RNA and not DNA as the sample input for detection of genomic alteration events. This allows for detection of genomic alteration events that would have been excluded in a typical DNA-based detection assay. Examples of such genomic alterations include:

- Copy number gains in DNA leading to overexpression of RNA;
- Structural rearrangements involving very long introns of two or more genes; and
- Changes in gene expression patterns corresponding to drug response or resistance.

SEQUENCE LISTING

Table of forward primers specific to genes that are capable of undergoing genomic alteration.

SEQ ID NO	Primer name	Sequence
1	AGAP3_exon9F	AGAAGAAGGCTGCCGAGTG
2	AGK_exon2F	GCTCTGCCTGCTGACCTG
3	AGTRAP_exon5F	CAGAGCACAGCATTAAGTTTGG
4	AKAP9_exon21F	AGGCATCTGTAAAGTCATGTGTC
5	AKAP9_exon8F	GAGCAACTCAACCAAGTGAAAATG
6	ARMC10_exon4F	GCACTAAATAACCTGAGTGTGAATG
7	ATIC_exon7F	GTACACACTGCAGCCCAAG
8	BCR_E1_F	GCAGATCTGGCCCAACGAT
9	BCR_E13_F	CTGACCAACTCGTGTGTGAAACTCC
10	BCR_E14_F	CGGGGCTCTATGGGTTTCTG
11	BCR_E18_F	GTCTTCGGAGTCAAGATTGCTG
12	BCR_E19_F	ATCTACCGCGTGTCCGGT
13	BCR_E2_F_rdsgn	ACATTGATGACTCGCCCTCC
14	BCR_E6_F	AAGATGCCAAGGATCCAACGAC
15	BCR_E8_F	CAATGAGGAGATCACACCCCG
16	BRAF_exon1F	TCTTCGGCTGCGGACCCT
17	BRAF_exon2F	GAACATATAGAGGCCCTATTGGAC
18	BRAF_exon3F	GAGCAACCCCAAGTCACCA
19	BRAF_exon4F	TGAGAGGTCTAATCCCAGAGT
20	BTF3L4_exon3F	CAGAGTTCTCTAAAAAACTGGCTG
21	CARS_exon17F	GAGAAGGAGTGCGGAAGATTG
22	CBFB_E4_F	TGGTATGGGCTGTCTGGAGT
23	CBFB_E5_F	CGGAGAAGGACACGCGAAT
24	CCDC6_exon1FS	CAAGGCACTGCAGGAGGAGAAC
25	CCDC6_exon2F	GAATTCCTCACTAATGAGCTCTCCAG

26	CCDC6_exon8F	CTTCACGTGCAGCACATGG
27	CCDC91_exon11F	GGCAGTGAAAAGAACAAGAGATG
28	CD74_exon6FS	TCCTTGAGCAAAAGCCCCTG
29	CDC27_exon16F	TTGATCCCAAGAACCCTCTATG
30	CLTC_exon30F	CATGCCCTATTTTCATCCAGGT
31	CLTC_exon31F	GAAGAAGAACAAGCTACAGAGACAC
32	CUX1_exon10F_new	GCCAATCACTCCCTCCAG
33	DCTN1_exon16F	ATGACTGCGTTCTGGTGCTG
34	DCTN1_exon26F	CATTGCTACTCTGGTCTCTGG
35	EML4_exon13F	CTACTGTAGAGCCACACCTG
36	EML4_exon14F	ATTAAGTGGAGGAGGGAAAGACAG
37	EML4_exon15F	CGAGGAACATTTAATGATGGCTTC
38	EML4_exon16F	TGCTCTTGACATGTGCTCAGGACAG
39	EML4_exon17F	CTGTGCAGATTTTCATCCAAGTGGC
40	EML4_exon18F	TCTATCCACACAGACGGGAATGAAC
41	EML4_exon20FS	ATAATGTCTAACTCGGGAGACTATG
42	EML4_exon2F	GCAATCTCTGAAGATCATGTGG
43	EML4_exon5F	GCAGACAAGCATAAAGATGTCATCATC
44	EPS15_exon22F	AATCATTTGGAGGTGGATTTGCTG
45	ERC1_exon12F	GAGGTGGAAAATGAGAAGAATGAC
46	EWSR1_exon7F	CTACAGCCAAGCTCCAAGTC
47	EWSR1_exon9_10F	GCTTCAATAAGCCTGGTGGA
48	EZR_exon10F	GGCTGCAGGACTATGAGG
49	FAM131B_exon2F	CATGGACAGCACCAGCTCA
50	FCHSD1_exon13F	GATGAGGTGGAGCAGGAG
51	FGFR1OP_exon12FS	GTGGAAATAGATGACATCAATACCAGTG
52	GHR_exon1F	CGAACCCGCGCTCTCTGA
53	GOLGA5_exon7F	GGCCAGATACATCAGCTCAG
54	GOPC_exon4F	TGTTCTCCAGGCTGAAGTATATG
55	GOPC_exon8F	CAAGTGGGGAAATCAAAGTATTACAAG
56	GTF2I_exon4F	CAGTTGAGGACTATTTCTGCTTTTG
57	HIP1_exon21F	ACCACCTGCCTCAGAGCC
58	HIP1_exon28F	CTCAACCATTTCCGGCAAATCAC
59	HIP1_exon30F	CTTGCTGGTGTGTGCTGAGG
60	IRF2BP2_E1a_F	GAGCAAGTTTAAGAAGGAGCCG
61	IRF2BP2_E1b_F	GCAGGTTGTTGGGTTTCGAG
62	IRF2BP2_E2_F	GGAGAGGTCTATTGTCCCAGTG
63	KIAA1468_exon10FS	CTGCCTGCCACACATTGTTC
64	KIAA1549_exon12F	ACGAGGAGATAAGACGCC
65	KIAA1549_exon13F	CTTATCGCCATGCAGCCGA
66	KIAA1549_exon14F	ACAAGATCCTGGACCCAC
67	KIAA1549_exon15F	AGCGATGGCACCTACAGGA
68	KIAA1549_exon16F	AAGAGAGGCGAGCCACCC
69	KIAA1549_exon18F	GGAGGAGATGCCGTCGGT

70	KIAA1549_exon19F	AAGCAGAGGCAGCCAGTAT
71	KIF5B_exon15F	CTTGCAGAAATAGGAATTGCTGTGG
72	KIF5B_exon16FS	TGAAAAGGAGTTAGCAGCATGTCAG
73	KIF5B_exon17F	ATGCCCTCAGTGAAGAACTAGTCC
74	KIF5B_exon22F	GAACCTCAGACTTTACACAACCTGCG
75	KIF5B_exon23FS	ATCTTGAACAGCTCACTAAAGTGC
76	KIF5B_exon24F_new	GAAGCAGTCAGGTCAAAGAATATGG
77	KLC1_exon9F	TGCACATGAAAGGGAGTTTGGTTC
78	KMT2A_E10_F	CCA GGG TGG TTT GCT TTC TCT
79	KMT2A_E11_F	TCT GTC ACG TTT GTG GAA GG
80	KMT2A_E5_F	AAG CCC AAG TTT GGT GGT CG
81	KMT2A_E6_F	CAATGGATGCCTTCCAAAGCC
82	KMT2A_E7_F	AAACCACTCCTAGTGAGCCC
83	KMT2A_E8_F	GCT CCC CGC CCA AGT ATC
84	KMT2A_E9_F	GCAGATGGAGTCCACAGGAT
85	KTN1_exon29F	CATGCTAAAAGAGAGGGAGAGTG
86	LRIG3_exon16F	TTCTTACCACAACATGACAGTAGTG
87	MKRN1_exon4F	ATCCAATGGATGCTGCCCAGA
88	MSN_exon11F	GACAGAAGAAGGAGAGTGAGG
89	NCOA4_exon6FS	TTGAAGCTGACACAATTACTCTGC
90	NCOA4_exon7F	CCTGGAGAAGAGAGGCTGTATC
91	NCOA4_exon8F_new	AGGACTGGCTTACCCAAAAGCAG
92	NPM1_E4_F	AAGTGTGGTTCAGGGCCAGT
93	NPM1_E5_F	TATCTGGAAAGCGGTCTGCC
94	NUP214_exon21F	AAGACCCCACCAAGTGAGAT
95	PAPSS1_exon5F	CTCCTGTGATGTAAATGACTGTG
96	PML_E3_F	AGTTCAAGGTGCGCCTGC
97	PML_E6a_F	CTTCCTGCCCAACAGCAAC
98	PML_E6b_F	TGCCCCAGGAAGGTCATCAAG
99	PPFIBP1_exon12F	ATGCAAGACACGGTGGTACTG
100	PPFIBP1_exon8F	GGACAGTGAGAGACTTCAGTATG
101	PPFIBP1_exon9F	TGGTTTGCAAGATGAAAGGAGAAG
102	PRKAR1A_7F	CATCGACCGAGACAGCTATAGAAG
103	PWWP2A_exon1F	TTGTTCGTGTCGTTCCGCTT
104	RANBP2_exon18F	GGGTCACAGACATTTTCATGGG
105	RUNX1_E5_F	ATGACCTCAGGTTTGTGCGGC
106	RUNX1_E6_F	ACCTACCACAGAGCCATCAAA
107	RUNX1_E7_F	ACTGCCTTTAACCCTCAGCC
108	SDC4_exon2F	ATCTGATGACTTTGAGCTGTCTGGC
109	SDC4_exon4F	GCAGCAACATCTTTGAGAGAACGG
110	SLC34A2_exon13delF	TGTCAAGGCTCCTGAGACCTTTGAT
111	SLC34A2_exon4F	TCGTGTGCTCCCTGGATATTCTTAG
112	SND1_exon10F	GATTACCTGTCCAGCATCC
113	SND1_exon11F	CCTTACATGTTTGAGGCC

114	SND1_exon14F	AGGATTGCATAGCAAGAAGGAAG
115	SND1_exon16F	CTTGGTGCAGGAAGGAGAG
116	SND1_exon9F	GCTCCACAGCTAATTTGGAC
117	SQSTM1_exon5F	CGAGTGTGAATTTCTGAAGAAC
118	STAT5B_E15_F	GTGACTCAGAAATTGGCGGC
119	STRN_exon3F	TGAATCAGGGAGATATGAAGCCTCC
120	TAX1BP1_exon8F	GCAGTTATGTTTGGCTGAAAAGG
121	TFG_exon3F	GTGCAGTAGGATACTGAAACTGAC
122	TFG_exon4F	GAGAACCAGGACCTTCCAC
123	TFG_exon5F_new	ATGTTATGTCAGCGTTTGGCTTAAC
124	TFG_exon6F_new	CAGCAGCCACCATATACAGG
125	TMPRSS2_exon1F	TAAGCAGGAGGCGGAGGC
126	TMPRSS2_exon2F	CAGATACCTATCATTACTCGATGCTG
127	TMPRSS2_exon3F	TCCTGACGCAGGCTTCCA
128	TMPRSS2_exon5F	CTCTAACTGGTGTGATGGCGT
129	TPM3_exon2F	GCAAAAGCTGGAAGAAGCTGA
130	TPM3_exon8F	AGTTTGCTGAGAGATCGGTAGC
131	TPM4_exon8F	GGAAAAGACAATTGATGACCTGGA
132	TPR_exon15F	CAAACAACAGGAGTTGCCATTCC
133	TRIM24_exon3F	GTTCACAAAAGACCACACTGTC
134	TRIM24_exon8F	TCACTGTGATCCTAGTTTCTGG
135	TRIM24_exon9F	TCCCAACACAGATCAGCCT
136	VCL_exon16F	CGATGGTGATGGATGCAAAAG
137	ZBTB16_E2_F	TGGGGTCGAGCTTCCTGATA
138	ZBTB16_E3_F	GTTCTGGATAGTTTGCGGC
139	ZBTB16_E4_F	TTTCGAAGGAGGATGCCCTG
140	ZC3HAV1_exon3F	CATCTGCAACAGCAAGCACA
141	AGBL4_exon5_6F	CCAGAGTGTGCAACAACGAAAG
142	AR_exon2_F_rdsgn	TGTGGAAGCTGCAAGGTCTTC
143	AR_exon3_F	CCATCTTGTCGTCTTCGGAAATG
144	BAG4_exon1_F	ATGGCTACTATCCCTCGGGAG
145	BAG4_exon2_F	GCTCCTTACCCAAGTACATATCCTG
146	BAIAP2L1_exon9_F	CGATATGTTTAATAACCCAGCCAC
147	BCAN_exon13F	AAGAGAACGGTCGTTGGGAG
148	BCR_exon4_F	TTCTATGATGGGCTCTTCCCCC
149	BCR_exon7_F	CTCTGCTCTACAAGCCTGTGGAC
150	BCR_exon9_F	CGTCTTCCTGTTCACCGACCT
151	BTBD1_exon4F	AAGCACTTTCCTTAATCCGGTTC
152	HMG2P46_exon1_F	AGGTGAATCTTTGGTTGGTGA
153	HMG2P46_exon2_F	CATGCTTGTCAAAAATCAGAGGC
154	C8orf34_exon2_F_rdsgn	GGAACCGTGGACAACTTCAAA
155	CAPZA2_exon4_F	GCATTTGCACAGTATAACTTGGAC
156	CD47_exon7F	GGTTTGAGTATCTTAGCTCTAGCAC
157	CD74_exon8F	GAAAGAGTCACTGGAAGTGGAGG

158	CNTRL_exon15_F	GCCAACCAGCTCAAGGAAGAGTT
159	CNTRL_exon38_F	GCAGGAAGAGGAGAGGTGG
160	CNTRL_exon39_F	CCAGCCTGAAGGAAGCACTTAA
161	CNTRL_exon40_F	AGCTCAACCAGATGCAGTATGAG
162	CTNNB1_exon1_F	AGGTCGAGGACGGTCGG
163	CUX1_exon1_F_rdsgn	TCTCAAGATGGCGGCCAATGTG
164	CUX1_exon7_F	ATGTCCACCACCTCAAAGCTGG
165	CUX1_exon8_9_F	CAAAGGCCGACGAGATTGAAATGAT
166	CUX1_exon8_F	AACTCGAACAGAATTATTTGACCTGAA
167	DDX5_exon2_F	GAGAGAATTTTTATCAAGAGCACCCCTG
168	EGFR_exon15F	TGCCATCCAAACTGCACCTAC
169	EGFR_exon24F_rdsgn	TCTCCAAAATGGCCCGAGAC
170	ERBB2_exon14_15_F	TTTGGACCGGAGGCTGACC
171	ERBB2_exon15F	GAAGTTTCCAGATGAGGAGGG
172	ERBB2_exon7-8_F	CCTGTCCCTACAACCTTTCTAC
173	ESR1_exon3_F	AACGAGCCCAGCGGCTAC
174	ESR1_exon4_F	CCATTATGGAGTCTGGTCCTGT
175	ESR1_exon5_F	GGCTCCGTAAATGCTACGAAGTG
176	ESR1_exon6_F_rdsgn	CAGACAGGGAGCTGGTTCACA
177	ESR1_exon7_F	ACCCAGGGAAGCTACTGTTTG
178	ETV6_exon4F	TGGAACTCTATACACACACAGCC
179	ETV6_exon5F	CACATCATGGTCTCTGTCTCCC
180	ETV6_exon7F	ACATTATCAGGAAGGAGCCAGG
181	FGFR1_exon1_F	ACAAGCCACGGCGGACTCT
182	FGFR1_exon17_F	TTCAAGCAGCTGGTGGAAGAC
183	FGFR1_exon18_F	CACAAATTTCCCCAAAGACTGCG
184	FGFR1_exon2_F	GTCACAGCCACACTCTGCAC
185	FGFR1OP_exon5_F	TTAGAAGTGATCAGGCGCTGTC
186	FGFR1OP_exon6_F	TCCACCAAAGTCACCAGAGG
187	FGFR1OP_exon7_F	GACAAGGTAAGAAGAAGACAAGCG
188	FGFR2_exon17F	AGAAGACTTGGATCGAATTCTCAC
189	FGFR2_exon18F	CGAACCATGCCTTCCTCAGTATCC
190	FGFR3_exon17F	TCCTTACCGTGACGTCCA
191	FIP1L1_exon10_F	GTTGGGAAGTGGCAGGATCGA
192	FIP1L1_exon11_F	CTATAACTATCAGCCGAGTAGAAGGCAG
193	FIP1L1_exon12_F	TCTTCCACCTCCTCCGACTGT
194	FIP1L1_exon14_F	TTTTCTCCTCCACCAGGCG
195	FIP1L1_exon15_F	GGACATTCTCTGGTTATGATAGTCG
196	FIP1L1_exon16_F	GACAGAGAAAGAGAACGCACCA
197	FIP1L1_exon17_F	ACGACACAGGGAGAAAGAGG
198	FIP1L1_exon18_F	AGAAGGAAAAGAAGCGGGCA
199	FIP1L1_exon9_F	TGCCCTTCCATCTACAAAAGC
200	HERV-K_F	GATTCCCGAGTACGTCTACAGTGA
201	HERVK17_F	TTTCCACACTCTCATTCCGGA

202	HNRNPA2B1_exon1_F	TGCGGGAAATCGGGCTGAA
203	LMNA_exon2F	GAGCTGCATGATCTGCGG
204	MDK_exon4F	CCCAAGACCAAAGCAAAGG
205	MET_exon13F	ACAGCACTGTTATTACTACTTGGG
206	MPRIIP_exon21F	AGAAGTCCCCCTGACAGTGCC
207	NACC2_exon4F_rdsgn	GACTGGCATCCGCTCGTC
208	NFASC_exon21F	TCCAGGCTGAAAATGACTTCGGG
209	PAX8_exon10F	ACATCCCCACCAGCGGAC
210	PAX8_exon7F	GAGTGCCCATTTGAGCGG
211	PAX8_exon8F	CTTCCAACACGCCACTGG
212	PAX8_exon9F	CGCCCTTCAATGCCTTTCC
213	PCM1_E23_F	CTCCCAAGTCAAAAAGTAAGAAGAGGA
214	PCM1_E24_F	CAGCAGAAAGAATCATGAGCAACTGGAAA
215	PCM1_E25_F	CCATGAGCTGCAGCTACTAAAC
216	PCM1_E26_F	ATAGCATCAAACTCAGAACTTACTCCT
217	PCM1_E36_F	ACTCCTGAAAGCTCTCTGGCTG
218	PCM1_E37_F	GATTTTGTAAGAGTTGAAGATTTACCACTG
219	QKI_exon6F	ACACATTGGCACCAGCTACATC
220	RBPMS_exon5F	CAAACCTCGTAGGGACTCCAAAC
221	SLC45A3_exon1_F	AACCAGCCTGCACGCGCT
222	SEC61G_5'UTRF_rdsgn	CTCCAGTGCTACGTGTCCCTG
223	TCF3_exon16_F	ACCCTCCCTGACCTGTCTC
224	TPM4_exon6F	CTGAAAAGGAGGACAAATATGAAGAAG
225	TPR_exon21F	TTGAAACAGCACCTCAGTAATATGG
226	TPR_exon6F	GGGAATGAGATTCTAGAGCTTAAATG
227	UBTF_exon2_F	AACGGAGAAGCCGACTGC
228	ZMYM2_exon16_F	TGTATATCCCAGTTCCTATGCACATG
229	ZMYM2_exon17_F	AGTTGCTTACAATGACGGATATGATGAGT
230	AFAP1_exon13_F	AGCAGTCAGCCAAAGAAAGC
231	AFAP1_exon14_F	GCTCCGCAAGGAAAGAAAAGA
232	AKAP9_exon18F	CAGCTACAAGAAGAGATTAAGAGACTT
233	AKAP9_exon7F	GACTTCACAATGCAAATTAGTTTCTTG
234	AKAP9_exon9F	GAACCTCAGAAAATACACCAGTTAGAACT
235	AXL_exon20_F	TGCTGATAGGGGGCTCCCCA
236	BCR_exon10_F	TGGACGCTTTGAAGATCAAGATC
237	BCR_exon11_F	CTGCTTATGTCTCCCAGCATG
238	BCR_exon12_F	CGTGCAGAGTGGAGGGAGAA
239	BCR_exon15_F	GCAAAGACGCGCGTCTACAG
240	BCR_exon16_F	AAGGAGGACGGCGAGAGC
241	BCR_exon17_F	AGGACAGAGACTGGCAGCG
242	BCR_exon20_F	TTCACTGACGAGTTCTACCCCAA
243	BCR_exon3_F	CTAGCGAGGAGACTTACCTGAG
244	BCR_exon5_F	AAATGGCTGAGAAGTGCTGTC
245	HMG2N2P46_exon2_F_rdsgn	GGAACCACACTTCGAGAATCAC

246	CCDC6_exon3F	GGAGAATGACACCATTCTTAAGCA
247	CCDC6_exon4F	AGCACTAGTTAATCGCCTCTGGAAA
248	CCDC6_exon5F	TGAAGTGGAACGGCTGAAGA
249	CCDC6_exon6F	TCTCCGAGAGTGAGTCCAGC
250	CCDC6_exon7F	CAGCCCAGATCCCTTACACAC
251	DCBLD1_exon2_F	CCAGACCTGTGCTTCTGACTAT
252	EGFR_exon1_F	AGTCGGGCTCTGGAGGAAAAG
253	EML4_exon10F	GGACTTGGCACTTTTGAGCG
254	EML4_exon11F	GCTTACTGTATGGGACTGGCA
255	EML4_exon12F	TTTCTTCTGGACCTGGAGCG
256	EML4_exon19F	CTATGTAGTCTCTGAAAATGGAAGAAAAT A
257	EML4_exon1F	CGCTTTCCCCGCAAGATGGA
258	EML4_exon21F	AAGGACATTGATTGGACGACA
259	EML4_exon22F	CGATGACTTTTGTAAGTCCATCTGTTTCA
260	EML4_exon23F	AAAGCCACCCTTCTGGAGGA
261	EML4_exon3F	GTGCTGTCTCAATTGCAGGAAAAG
262	EML4_exon4F	CACAGACAACTCCAGAAAGCA
263	EML4_exon7F	ATCAGAACGGAACTGCCTCC
264	EML4_exon8F	GAACTCAGCGACACTACCTGG
265	EML4_exon9F	GCAACTGGACAGATAGCTGG
266	ESR1_exon8_F	GGAGAGGAGTTTGTGTGCCT
267	ESR1_exon9_F	GCCCAGCTCCTCCTCATC
268	ETV6_exon6F	GATCCCAACGGACTGGCTC
269	EWSR1_exon8_F	TGGAGGCATGAGCAGAGGT
270	EZR_exon11F	CTAAGGAGGAGCTGGAGAGACA
271	EZR_exon12F	GAGGCGCAAGGAGGATGAA
272	FIP1L1_exon8_F	ATGGAAGTTACACCAGGTGCAGAG
273	FRMD4A_exon2_F	ATGGGAATCGGAGACCATG
274	GOPC_exon5F	TGCAGAGGACGTAATGACTTGA
275	GOPC_exon6F	GGTCCAATTAGAAAAGTTCTCCTCCTTA
276	GOPC_exon7F	ACCTAAGGGACACAAAGCATA
277	GOPC_exon8F_rdsgn	AGGTGGTAACCCTGGTGCTAGTT
278	HIP1_exon29F	AAGCATGACGCTGACACAGA
279	IRF2BP2_exon1_F	CAGGCAGGTTGTTGGGTTTCG
280	KIAA1549_exon17F	GATACGAAGACTATGGAATGACTCC
281	KIF5B_exon18F	CATGAAATGGAAAAGGAGCACTTA
282	KIF5B_exon19F	AGTTTGAGAGATGAAGTAGAAGCA
283	KIF5B_exon20F	AGTTGAAAGCCACAGATCAGGA
284	KIF5B_exon21F	ACGAGAACAAGCAAGACAAGAC
285	MECOM_exon2_F	AGAGCAGAGGTCAAACCTGA
286	MET_exon11F	AGCCAGTGATGATCTCAATGGG
287	MYB_exon10_F	AGTTCAACTCCCAAGCGTTCCC
288	MYB_exon11_F	AACACCATTTCATAGAGACCAGACTG
289	MYB_exon12_F	TCAAACATGCACTTGCAGCTCAAG

290	MYB_exon13_F	TGCTGAGTTTCAAGAAAATGGACC
291	MYB_exon14_F	AGGGGACAGTCTGAATACCCA
292	MYB_exon15_F	TGTTCTCAAAGCATTTACAGTACCT
293	MYB_exon8_F	TTAGAATTGCTCCTAATGTCAACCGAGA
294	MYB_exon9_F	ACCTCTTAGAATTTGCAGAAACACT
295	NCOA4_exon9F	CAACCTCAGCCAGTTATCTTCTGG
296	PAN3_exon1_F	AGGCAGTAGCGGGGGAC
297	PAX3_exon7_F	TCCAACCCCATGAACCCAC
298	PPFIBP1_exon11F	CAAAAAATGAAAAAAGCTGTGGAGTCC
299	PPFIBP1_exon9F_rdsgn	GGTCAGATGCAGTATGAAAAGCA
300	SCAF11_exon1_F	CCTGGGAACCTACTGTGGGG
301	SDC4_exon3F	TGGAAGACTCCATGATCGGC
302	SND1_exon12F	GCCTTTTCAGAGCGTACCTG
303	SND1_exon13F	GACCAGAGATCATCACACTACG
304	SND1_exon15F	GTCTCAAACCTCTATTTGCCAAAGGAACTT
305	SPTBN1_exon6_F	AGGGACGGCATGGCCTTC
306	SS18_exon10_F	AACACAGCCTGGACCACCA
307	SS18_exon9_F	AGTACCCAGGGCAGCAAGG
308	TMPRSS2_exon4F	ACCTTCCTCGTGGGAGCT
309	TPM3_exon2F_rdsgn	TTTGAAGGATGCCCAGGAGAAG
310	TPM3_exon4F	CTAAGCACATTGCAGAAGAGGC
311	TPM3_exon5F	AGGAGACTTGGAACGCACAG
312	TPM3_exon6F	GTCACCAACAACCTCAAGTCTC
313	TPM3_exon7F_rdsgn	GAAGATAAATATGAGGAAGAAATCAAGAT T
314	TRIM24_exon12_F	CAGAACGGTCCAGTCACCAA
315	FGFR3_exon17F_new	GAGGCCCACCTTCAAGCAG

Table of reverse primers specific to genes that are capable of undergoing genomic alteration.

SEQ ID NO	Primer name	Sequence
316	ABL1_E2a_R	CTTCACTCAGACCCTGAGGC
317	ABL1_E2b_R	CAACGAAAAGGTTGGGGTCA
318	ABL1_E3_R	GCTTCACACCATTCCCCATTG
319	ABL1_E4_R	GATGATGAACCAACTCGGCCA
320	AFF1_E11_R	ATGTGCTGATGCCACTGGTT
321	AFF1_E6_R	GGCTCAGCTGTACTAGGCG
322	AFF1_E9-10_R_rdsgn	GCTTCTCTGGGGTTTGTTCCT
323	ALK_exon19R	AGAGAGGATCAGCGAGAGTG
324	ALK_exon20RL	TCAGCTTGTACTCAGGGCTCT
325	ALK_exon21R	CTTGGGTCGTTGGGCATTC
326	BRAF_exon10R	CGTTAGTTAGTGAGCCAGGT
327	BRAF_exon11R	CAGGAATCTCCAATCATCACTC

328	BRAF_exon8R	AGGGCTGTGGAATTGGAATG
329	BRAF_exon9R	CTCCATCACCACGAAATCCTTG
330	ERG_exon10R	AAGGCGGCTACTTGTTGGTC
331	ERG_exon11R	GAGGAACTGCCAAAGCTGGAT
332	ERG_exon2R	CTTTCCTCGGGTCTCCAAAG
333	ERG_exon3R	CCCATCTACCAGCTGTTTCAG
334	ERG_exon4RS	TGGTCCTCACTCACAAGTATAAGG
335	ERG_exon5RS	CCACCATCTTCCCGCCTTTG
336	ERG_exon8R	CAGGAGATCAGCCTGGAC
337	ERG_exon9R	GGTCTTCAGTTTTGGGCACTG
338	MECOM_E2_R	TGCATCTGGCATTCTCTCCAAAG
339	MLLT3_E10_R	ATGTCATTAACCTTCTGTGAAGCTCTAC
340	MLLT3_E5_R	TGCTGCTGCTGCTGGTATGAATAC
341	MLLT3_E7_R	TCTGATTCTCCTCATTTGTCATCA
342	MLLT3_E9_R	TTTGCTTATCTGATTTGCTTTGCTTTATTG
343	MYH11_E28_R	TCTCTTTCTCCAGCGTCTGC
344	MYH11_E29_R	CTCGGCCTCGTTAAGCATCC
345	MYH11_E30_R	GCAGCTTCGTAGACACGTTG
346	MYH11_E32_R	CCTCTCATCCGCGTATTTGGA
347	MYH11_E33_R	TCCATCTGGGTCTCCAGGG
348	MYH11_E34_R	GTTGCTTTTCGCTCGTCTTCC
349	RARA_E3_R	GGCTGGGCACTATCTCTTCA
350	RET_exon11RS	ATGAAGGAGAAGAGGACAGCGG
351	RET_exon12RL	TCCTAGAGTTTTTCCAAGAACCAAG
352	RET_exon7R	GGACGTTGATGCCACTGAAT
353	RET_exon8R	CTCTTGCTGACTGCACAGGACAG
354	ROS1_exon34R	GGTCAGTGGGATTGTAACAACCAGA
355	ROS1_exon35R	CTTCGTTTATAAGCACTGTCACC
356	ROS1_exon36R	GAGGGAAGGCAGGAAGATT
357	RUNX1T1_E2_R	GCGTCTTCACATCCACAGGT
358	ACTR1A_exon3_R	AAGATGTCGCCTTCAAGGGCT
359	AHCYL1_exon2R	GTTTTGGTGGGGAATTTGGTGAAC
360	AKAP12_exon4_R	AGTAGCCATCTCTTTATCGGAGTCTC
361	AR_CE3_R	GGTCTGGTCATTTTGAGATGCTTG
362	AR_CE5_R	ACAACGTGATCCCAAAGATGTG
363	AR_CE4_R	ATGCAGTATGGCTTGGGGTT
364	BAIAP2L1_exon2_3R	AGGATCATAGCGTTTACAGCTTTCT
365	BICC1_exon2_3R	TGCGTATTTGTTTCCTCCATGATC
366	CCDC170_exon2_R	AGTGGTTTAACTGCTCCCGC
367	CLDN7_exon2_R	GCACCAGGGAGACCACCATT
368	EGFR_exon14R	GGCACTGTATGCACTCAGAGTT
369	EGFR_exon15R	CAGGTCTTGACGCAGTGG
370	EGFR_exon18R	GCTTGGTTGGGAGCTTCTC
371	ERBB2_exon17R	AAGACCACGACCAGCAGAAT

372	ERBB2_ex8-int8_R	TGAGTGGGTACCTCACACC
373	ETV1_exon4_R	CTTCTGCAAGCCATGTTTCT
374	ETV1_exon6_R	TGAACATGGACTGTGGGGTTCTTTC
375	ETV1_exon7_R	GGAGGGCCTCATTCCTTCT
376	ETV1_exon2_R	TGGTGACCATGTAAGGCACT
377	ETV1_exon2b_R	AGCATTTAGCTGGAGATTTCCTCA
378	ETV1_exon5_R	AACTTTCAGCCTGATAGTCTGGTACAA
379	ETV1_exon3_R	TTTTCTTTTCCTGACATTTGTTGGTTTC
380	ETV1_exon11_R	AGTTCATAGCTGGCCTGTTTTTCTG
381	ETV1_exon12_R	AGAAAAGGGCTTCTGGATCACACA
382	ETV4_exon1_2_R	CGAGACCTGCTCCCAGGA
383	ETV4_exon3_R	CATGAGCTTCCCCAGCGG
384	ETV4_exon5_R	CTGAATGGAAATCAGGAACAACTGC
385	ETV4_exon4_R	AGCGAGCCACGTCTCCTG
386	ETV5_exon2_R	CAGCATTGAGTAATTTCTGGGGG
387	ETV5_exon8_R	AACTCCTGGCTGAGGAGGGAA
388	FGFR1_exon8_R	AGCACCTCCATCTCTTTGTCTG
389	FGFR1_exon6_R	TGTCAGAGGGCACCACAGAG
390	FGFR1_exon9_R	AGATGATGATCTCCAGGTACAGG
391	FGFR1_exon10_R	GAACCAGAAGAACCCCAGAGTTC
392	JAK2_exon9_R	CCATCAATTAATGACACGAAAGACAAAGC
393	JAK2_exon10_R	ACAGTCCAGTCTGATTACCTGC
394	JAK2_exon11_R	TGTCCCACTGAGGTTGTACT
395	JAK2_exon12_R	TGGTGAGGTTGGTACATCAGAA
396	JAK2_exon13_R	TGACCGTAGTCTCCTACTTCTCT
397	JAK2_exon14_R	TTAAAACCAAATGCTTGTGAGAAAGCT
398	JAK2_exon15_R	TCTTTTTTCAGATATGTATCTAGTGATCCAA
399	JAK2_exon16_R	TCTCTGATAAGCAGAATATTTTTGGCACA
400	JAK2_exon17_R	GCATTACAGGTGGTACCCATGGTATT
401	JAK2_exon18_R	CCACTTTGGTGCAGGAAGCTGA
402	JAK2_exon19_R	AGGGCACCTATCCTCATATTTGGTAAC
403	MET_exon11R	GGAGTGGTACAACAGATTATCTCTG
404	MET_exon15R	CTGCACTTGTCGGCATGAAC
405	NOP2_exon16_R	GGATGACCTGAGGCAAGTCTA
406	NRG1_exon6R	CTCCGCACATTTTACAAGATGG
407	NRG1_exon5_6R	GTGGTGGATGTAGATGTAGATGAAGAAG
408	NRG1_exon2R	AACCTGCAGCCGATTCTCTG
409	NRG1_exon3R	GAATCAGCCAGTGATGCTTTGT
410	NRG1_exon4R	TGCTCCTTCAGTTGAGGCTG
411	NTRK1-exon10R_rdsgn	GTGTTTCGTCCTTCTTCTCCACC
412	NTRK1-exon12R	TGTCATGAAATGCAGGGACATGG
413	NTRK1-exon9R	ACCAGTGGTGCATCTCCA
414	NTRK1-exon11R	AGCGTAGAAAGGAAGAGGCAGG
415	NTRK2_exon13_R_rdsgn	TGGTGTCCCCGATGTCATTC

416	NTRK2_exon16_R_rdsgn	GCTGGCAGAGTCATCATCATTG
417	NTRK3_exon14R	GTCTCCTCACCCTGATGAC
418	NTRK3_exon15R	CTTCAGCACGATGTCTCTCTCTTA
419	NTRK3_exon11_12R	GGACTCACTTCGTCAAACAAGAT
420	PAWR_exon3_R	TAATTGCATCTTCTCGTTTCCGCT
421	PBX1_exon3_R	CTGGGGGTCTGTGGGTTC
422	PCDH11X_exon11_R	CTTGAGTGCAGTTGTCAGAGGC
423	PDGFRA_exon12_R	TGATTCAATGACCCTCCAGCG
424	PDGFRB_exon11_R	GGATGATAAGGGAGATGATGGTGAG
425	PDGFRB_exon9_R	TGTCTGTTCCCCACTGTCAGG
426	PDGFRB_exon10_R	AGCTGGCTCTCCTCTTCGGA
427	PDGFRB_exon12_R	AGCTCACAGACTCAATCACCTTC
428	PLAG1_exon4_R	GCTTTAGGTGGCTTCTCAAGTTTC
429	PLAG1_exon3_R	GACTCTTCGTGGAAGAGAGTGG
430	PLAG1_exon2_R	GAATGAAGCATTCTGGGTGCC
431	POLH_exon2_R	GTCCATGTCCACGAGAGCAA
432	PPARG_exon2R	AATGGCATCTCTGTGTCAACC
433	PPARGC1A_exon2R	AGGAAGATCTGGGCAAAGAGG
434	PPLN1_exon3R	GTCTAGCAGTGGTGGTTTCTTTG
435	RAD51_exon4R_rdsg	GGTGGAAATTCAGTTGCAGTGG
436	RUNX1_Exon4_R	CCTCGCTCATCTTGCCTGG
437	SHC1_exon2R_rdsgn	CCCTTCCACCCGAGTCCT
438	KIAA1598_exon7_R	TTCTCTGTTCAAGAACTTCTGAATTTAA
439	KIAA1598_exon8_R	GCTCCAGGTTTACTTGCATCTC
440	KIAA1598_exon9_R	AGCAGAAGGTGGCTTTGTCT
441	TACC1_exon7_R	TCCACAGGACACCGACACA
442	TACC3_exon11R	TTCTTCCCGTGGAGCTCCTC
443	TACC3_exon10R	GAGCAGGTCCACTATAGGTC
444	TACC3_exon8R	TCTACCAGGACTGTCCCTCA
445	TACC3_exon14R_rdsgn2	GGGTGATCCTTGCCAGGTAAT
446	TACC3_exon6R	TCACTGCCTGGACAGCTTGTG
447	YAP1_exon4_R	TCCTGAGTCATGGCTTGTTC
448	ABL1_E5_R_rdsgn	GCACCAGGTTAGGGTGTGTTGA
449	AFF1_E4_R_redsgn	CCAGGCGATGAGTGTGAGAC
450	AFF1_E8_R	CGTTCCTTGCTGAGAATTTGAGT
451	CCAR2_exon4_R	AGTCATGCAAGCTGGTAACAA
452	CCDC170_exon10_R	GGATTTGTTTAGATCTTCAATGGCTTTA
453	CCDC170_exon6_R	TAACCTCCCTTTCAAGAGCTTCTTTG
454	CCDC170_exon7_R	CAACTGTTCAACAAGCTCAGAT
455	CCDC170_exon8_R	GCCATCTGGTCCAACCTTCAATTTCT
456	CCDC170_exon9_R	GGAGGCTCATGTGTAATTTGCTCT
457	CIT_exon23_R	AGCTGTTACGAAGAGCATCAA
458	EGFR_exon17a_R	GTGGCGATGGACGGGATCT
459	EGFR_exon17b_R	GCATGAAGAGGCCGATCCC

460	EGFR_exon8_R	TCCTCCATCTCATAGCTGTCTG
461	ERBB4_exon18_R	GAGCTTGATTGGGTGCTGTG
462	ERG_exon6R	TTCTTTCACCGCCCACTCCAG
463	ERG_exon7R	CCGTGGAGAGTTTTGTAAAGGCTT
464	ETV1_exon10R	ATCCTCGCCGTTGGTATGTGG
465	ETV1_exon8R	TCGTGCGCAAAGGAGGAAAG
466	ETV1_exon9R	GGACAACACAGGTGTCATCAT
467	FGFR1_exon7R	GCCACTGTTTTGTTGGCGG
468	FLI1_exon3_R	AGCTTGCTGCATTTGCTAAC
469	FLI1_exon4_R	TTATGGCCCACTCCAGCCATT
470	FLI1_exon5_R	ATCGTGAGGATTGGTCCGGTG
471	FLI1_exon6_R	GTTATTGCCCAAGCTCCTCT
472	FLI1_exon7_R	TATTCTTACTGATCGTTTGTGCC
473	FLI1_exon8_R	GTTGGCTAGGCGACTGCT
474	FLI1_exon9_R	GAGAGCAGCTCCAGGAGGAATTG
475	FOXO1_exon2_R	TCTGCACACGAATGAACCTGC
476	GPHN_exon11_R	GACATGCGATGTCTTCTAGCCAC
477	MBIP_exon4_R	CATTGATTTTCAGCTTGCTTTCTTTC
478	MLLT3_E6_R_redsgn	TGGTCTGGGATGGTGTGAAG
479	MX1_exon12_R	CCACGATACTGATTTTCAAATTTCTGG
480	MX1_exon9_R	AAGTTTTTCTGCCAGGCAGGG
481	MYH11_E31_R_rdsgn	CTCTTCCAGAGCTTCCACGG
482	NCOA3_exon14_R	CTGCTCGGTTATATGGAGGACGAA
483	NCOA3_exon15_R	TAAGCCCCAGTCTCCTGAGGAA
484	NFIB_exon10_R	GGGCTTAGTCCCACATATCG
485	NFIB_exon11_R	GGGGTATAAATGCCTGCCGT
486	NFIB_exon12_R	AGATGGGTGTCTTATTTGACACTTGG
487	NFIB_exon9_R	GCCAGGCACTTTCCCTACTA
488	NTRK2_exon14R	AAAGGCAAAATCCCACCACAGA
489	NTRK2_exon15R	CAACACCTTGTCTTGATTTTACTTTCCC
490	NTRK2_exon17_R	TTCGCCTAGCTCCCTTTTCA
491	NTRK3_exon13R	AGAACCACCAACAGGACACAG
492	PDGFRA_exon2_R	ACAGCCTAAGACCAGGAACGC
493	PPARG_exon7_R	AGGTTGTCTTGAATGTCTTCAATGGGC
494	RET_exon10_R	GTGCCATAGCCAGCTTTAATCC
495	RET_exon9_R	CAGGTCTTGGTGCTGGGAG
496	ROS1_exon32_R_rdsgn	GAATTTTTACTCCCTTCTAGTAATTTGG
497	ROS1_exon33R	TTCCATGTGCAAACACTACTGC
498	SEPT14_exon10_R	GCTTCCTTATCTCCTCCTGTTGA
499	SSX1_exon4_R	CTGGAAGTCTGTGGCCTGTTT
500	SSX1_exon5_R	GGGATGATTCTGTGGAGCCT
501	SSX1_exon6_R	TGCTTCTGACACTCCCTTCG
502	SSX2_exon3_R	TCATCTTTTCCCACTCTCCTTAGAGA
503	SSX2_exon4_R	AGTCTTCGGCCCGTTTATTACA

504	SSX2_exon5_R	CTTCGGGGAGATTCCCTGGAG
505	SSX2_exon6_R	CTGGCACTTCCTCCGAATCA
506	TACC3_exon13a_R	CCTTCTGCTTCTGAACTTCCT
507	TACC3_exon13b_R	TTGGTCTTTTTCTTTTAGAACTTTCTGGAT
508	TACC3_exon5_R	TGCCAACTGCACCACAGG
509	TACC3_exon7R	AGGAAGTTCCAAACTGCTCCAGGTA
510	TACC3_exon9R	AAGAAATCGAACTCCACAAGC
511	TACC3_midExon4_R	TCTCCGCTTTGCATTCTTCCT
512	YAP1_exon5_R	GTGGCTGTTTCACTGGAGCA

Table of forward primers specific to control housekeeping genes.

SEQ ID NO	Primer name	Sequence
513	TBP_F_rdsgn	CTTTGCAGTGACCCAGCATCACT
514	ITGB7-F	GCACGCACCTATGTGGAAAC
515	PSMB2_F_rdsgn	TCTTCGGAGTCGGACCCCATAT
516	EMC7_A_F	TCGGTTTCCTTAAGACAGATGG
517	EMC7_B_F_rdsgn	ACTTTCTAATGAACCCAATGGTTAT
518	GPI_F	GCATCACAAGATCCTCCTGG
519	REEP5_F	CCAGCCTACATCTCAATTAAAGC
520	RRP1_F_rdsgn	CGGGCCGCAGGTGGTTTT
521	CHFR_F	AGTATTGTGATTACAGGGTCTGG
522	REL_F	GAATCAATCCATTCAATGTCCCTG
523	NASP_F_rdsgn	CAGGAAGCAGCTAGTCTTTTAGGTAAG
524	POMK_F_rdsgn_2	GAAGGAGCTGTAAAGAGAGTCTT
525	FRMD8_F	GATAGCAGAGAGAAGCATGTCC
526	SLC4A1AP_A_F_rdsgn	TCGGCAGGAAGCAGTATCT
527	SNAP29_F	GATCGACAGCAACCTAGATGAG
528	TUBGCP2_A_F_rdsgn	CATCCACGACCCATACAGTGAGTTTATG
529	SNRPD3_F_rdsgn	GACAACATGAACTGCCAGATGTCCAA
530	PUM1_F	TCAGACCAGCAGGTAATTAATGAGA
531	CIAO1_F	CGTTTGGGTCTGGGAAGTTGATGA
532	SRSF4_F	TGGAACTGAAGTCAATGGGAG
533	VCP_E1-2_F_new	GCTTCTGGAGCCGATTCAAA
534	RER1_E3_F_new	GGCTAGACAAGTCCACACCC
535	CHMP2A_E2-3_F	AAGCAAGGCCAGATGGATGC
536	RAB7A_E1_F	GTTTAGTCTCCTCCTCGGCG
537	ACTB_F	GAGACCGCGTCCGCC

5

10

Table of reverse primers specific to control housekeeping genes.

SEQ ID NO	Primer name	Sequence
538	TBP-R	GCATCTCCAGCACACTCTTC
539	ITGB7-R	AGCCAAACAGGAAACAGACCAG
540	PSMB2-R	AGGTAGTCCATGTAATACAGCG
541	EMC7_A-R	ACGGGATCAAATCTGTAAGCTG
542	EMC7_B-R	GTCTCATGTCAGGATCACTTGT
543	GPI-R	AAGGTCCTCTGGACTCTTGC
544	REEP5-R	CCATGACAGGAAGATATCAGAGAAG
545	RRP1_R	CTGGAGGAGTGGCTTGTG
546	CHFR-R	CTTTCTGTCTGGGAGAGCTG
547	REL-R	CATGTTTCATCAGGGAGAAAACTTG
548	NASP-R	CTTGCCAACTCCAGAAGTGA
549	POMK-R	CCTTGGAGAGATTTCAGCATCT
550	FRMD8-R	GAACTCCAGCCACAAGATG
551	SLC4A1AP_A-R	TCAGACGCTTCTTCTCAATCAG
552	SNAP29-R	TGTCATCTTGCTCCTCAATTTCTG
553	TUBGCP2_A-R	ACTTGTCGTTGTAATCCTCCTG
554	SNRPD3-R	GATTTTGCTGCCACGGATG
555	PUM1_R	ACCACGTGATTGCCATTCTG
556	CIAO1_R	GCCAAACCACATGCTTGACA
557	SRSF4_R	CTTCGAGAGCGAGACCTTGAAT
558	VCP_E2_R_new	ACAATTAACCGATTGGGACGG
559	RER1_E3-4_R_new	GTCACAATGTACCAACCTGC
560	CHMP2A_E3_R	TTGAGGGACACAGCCTGGAT
561	RAB7A_E1-2_R	AGGTCATCCTTCAAACGCGG
562	ACTB_R	ATCATCCATGGTGAGCTGGC

5 Table of forward primers specific to target genes related to protein expression

SEQ ID NO	Primer name	Sequence
563	CD274_3UTR_F	CAAAGAAGCAAAGTGATACACATTTG
564	CD274_EXON4_F	GTGAAAGTCAATGCCCCATACAAC
565	CD274_EXON3_F	GGCATTTGCTGAACGCATTTACTG
566	PDCD1-EXON3_4_5_F	CCGCACGAGGGACAATAGGA
567	CTLA4-EXON1_2_F	CTTCTCTTCATCCCTGTCTTCTG
568	CTLA4-EXON4_F	GCTGTTTCTTTGAGCAAAATGCTAAAGA
569	CD47-EXON3_F	GAAGGTGAAACGATCATCGAGC
570	CD47-EXON9_10_11_F	GAGGAACCCCTTAATGCATTCAAAG
571	LAG3-EXON8_F	CTTTGGAGAAGACAGTGCGAC
572	LAG3_EXON2-F_rdsgn	TTTGGGTGGCTCCAGTGAAG

573	HAVCR2_EXON3-F_rdsgn	AGTTGGTCATCAAACCAGCCAAG
574	HAVCR2_EXON7-F_rdsgn	CAAAGAGAAGATACAGAATTTAAGCCTCAT
575	PDCD1_EXON2-F_rdsgn	GCCAGGATGGTTCTTAGACTCC
576	CD14_exon1_1F	GAAGACTTATCGACCATGGAGC
577	CD27_exon1-2_1F	CAGATGTGTGAGCCAGGAAC
578	CD39_exon1_1F	CTTGAGAAAGGATTGCTGGTCA
579	CD40_exon6-7_1F	GACTGATGTTGTCTGTGGTCC
580	CD70_exon1_2F	CTTGGTGATCTGCCTCGTG
581	CD80_exon1_1F	TCTCAGAAGTGGAGTCTTACCC
582	FOXP3_exon6-7_1F	AGAGGACTTCCTCAAGCACTG
583	ICOS_exon3_1F	TGTGCAGCCTTTGTTGTAGT
584	ICOS_exon1-2_2F	GCATTAAAGTTTTAACAGGAGA
585	TIGIT_exon3_1F	AGATTCCATTGCTTGGAGCC
586	TNFRSF18_exon2-3_1F	TACAGTCCCAGGGGAAATTCAG
587	VISTA_exon4-5_1F	TGCGGATGGACAGCAACATT
588	VISTA_exon2-3_2F	CCTCCCAGGATAGTGAAAACATC

Table of reverse primers specific to target genes related to protein expression

SEQ ID NO	Primer name	Sequence
589	CD274_3UTR_R	GAACCCCTAAACCACAGGTTGAG
590	CD274_EXON4_R	CTCAGCCTGACATGTCAGTTCATG
591	CD274_EXON3_R	CTACTGGGAATTTGCATTCAATTGTC
592	PDCD1_EXON2_R	AGAAGCTGCAGGTGAAGGTG
593	PDCD1-EXON3_4_5_R	GGAAATCCAGCTCCCCATAGTC
594	CTLA4-EXON1_2_R	GCATACTCACACAAAGCTGG
595	CTLA4-EXON4_R	TTGCTTTTCACATTCTGGCTCTG
596	CD47-EXON3_R	CTGTCCCCAGAACAGGAGTATAG
597	CD47-EXON9_10_11_R	CCTTTCACGTCTTACTACTCTCCA
598	LAG3-EXON2_R	CAGAAGGCTGAGATCCTGGAG
599	LAG3-EXON8_R	GGTTCTTGCTCCAGCTCCTC
600	HAVCR2-EXON3_R	CCATGTCCCCTGGTGGTAAG
601	HAVCR2-EXON7_R	GGTATAGATGTTTTCTTCTGAGCGA
602	CD14_exon2_1R	CGCAGCGGAAATCTTCATCG
603	CD27_exon2_1R	GTCAGGAGAGAAGGAGACCC
604	CD39_exon1-2_1R	TCACGTTAGACTCTTGAAACCC
605	CD40_exon7_1R	GATAAAGACCAGCACCAAGAGG
606	CD70_exon1-2_2R	AGCTACGTCCCACCCAAG
607	CD80_exon1-2_1R	GTGGATTTAGTTTCACAGCTTGC
608	FOXP3_exon7_1R	TCTCTCTCTGGAGGAGACATTG
609	ICOS_exon3-4_1R	GCACACTGGATGAATACTTCTTT
610	ICOS_exon2_2R	TTTGTACACCTCCGTTGTGA
611	TIGIT_exon3-4_1R	GGATTCTGAGGGCTTTCTTCTT

612	TNFRSF18_exon3-4_1R	CAGCAGTCTGTCCAAGGTTTG
613	VISTA_exon5_1R	AGGACAGGGGGTGCCTGA
614	VISTA_exon3_2R	CTGCCTTTGCTTGTAGACCA

Table of other sequences

SEQ ID NO	Sequence name	Sequence
615	Barcode sequence	NNNNNNNNNN
616	Universal indexed forward primer	AATGATACGGCGACCACCGAG ATCTACACCTAGCGCTACACT CTTCCCTACACGACGCTCTTCCGATC*T
617	Universal indexed reverse primer	CAAGCAGAAGACGGCATACGA GATAACCGCGGGTGACTGGAG TTCAGACGTGTGCTCTTCCGATC*T

5

CLAIMS

1. A method of detecting genomic alteration and/or detecting gene expression and/or quantifying a level of gene expression using RNA in a biological sample, comprising:

(a) extracting RNA from the biological sample and converting the RNA to complementary DNA (cDNA);

(b) performing a plurality of multiplexed PCR reactions on the converted cDNA using

(I) a plurality of forward and reverse primer pairs specific to a plurality of **target genes that are capable of undergoing genomic alteration**,

wherein each forward primer of the plurality of forward and reverse primer pairs specific to the plurality of target genes that are capable of undergoing genomic alteration is complementary to a sequence located about 50 base pairs upstream of an exonic junction of each target gene that is capable of undergoing genomic alteration,

wherein each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of target genes that are capable of undergoing genomic alteration is complementary to a sequence located about 50 base pairs downstream of an exonic junction of each target gene that is capable of undergoing genomic alteration,

wherein each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of target genes that are capable of undergoing genomic alteration comprises a barcode sequence on its 5' end, wherein the barcode sequence of each reverse primer corresponding to each target gene that is capable of undergoing genomic alteration is different, and/or

(II) a plurality of forward and reverse primer pairs specific to a plurality of **control housekeeping genes**, wherein:

(i) each forward primer of the plurality of forward and reverse primer pairs specific to the plurality of control housekeeping genes is complementary to a sequence spanning an exon-exon junction of each control housekeeping gene,

wherein each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of control housekeeping genes is complementary to a sequence about 100 base pairs downstream of the sequence spanning the exon-exon junction of each control housekeeping gene,

5

wherein each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of control housekeeping genes comprises a barcode sequence on its 5' end, wherein the barcode sequence of each reverse primer corresponding to each control housekeeping gene is different;

10

- (ii) each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of control housekeeping genes is complementary to a sequence spanning an exon-exon junction of each control housekeeping gene,

15

wherein each forward primer of the plurality of forward and reverse primer pairs specific to the plurality of control housekeeping genes is complementary to a sequence about 100 base pairs downstream of the sequence spanning the exon-exon junction of each control housekeeping gene,

20

wherein each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of control housekeeping genes comprises a barcode sequence on its 5' end, wherein the barcode sequence of each reverse primer corresponding to each control housekeeping gene is different;

25

- (iii) each forward and each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of control housekeeping genes is complementary to consecutive sequences spanning an exon-exon junction of each control housekeeping gene,

30

wherein each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of control housekeeping genes comprises a barcode sequence on its 5' end, wherein the barcode sequence of each reverse primer corresponding to each control housekeeping gene is different, and/or

- (III) a plurality of primer sets specific to a plurality of **target genes related to protein expression**,

wherein each primer set comprises a plurality of forward and reverse primer pairs specific to each target gene related to protein expression, wherein:

- (i) each forward primer of the of the plurality of forward and reverse primer pairs specific to each target gene related to protein expression is complementary to a sequence spanning an exon-exon junction of each target gene related to protein expression,

wherein each reverse primer of the of the plurality of forward and reverse primer pairs specific to each target gene related to protein expression is complementary to a sequence about 100 base pairs downstream of the sequence spanning the exon-exon junction of each target gene related to protein expression,

wherein each reverse primer of the plurality of forward and reverse primer pairs specific to each target gene related to protein expression comprises a barcode sequence on its 5' end, wherein the barcode sequence of each reverse primer corresponding to each target gene related to protein expression is different,

- (ii) each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of target genes related to protein expression is complementary to a sequence spanning an exon-exon junction of each target gene related to protein expression,

wherein each forward primer of the plurality of forward and reverse primer pairs specific to the plurality of target genes related to protein expression is complementary to a sequence about 100 base pairs

downstream of the sequence spanning the exon-exon junction of each target gene related to protein expression,

5 wherein each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of target genes related to protein expression comprises a barcode sequence on its 5' end, wherein the barcode sequence of each reverse primer corresponding to each target gene related to protein expression is different;

10 (iii) each forward and each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of target genes related to protein expression is complementary to consecutive sequences spanning an exon-exon junction of each target gene related to protein expression,

15 wherein each reverse primer of the plurality of forward and reverse primer pairs specific to the plurality of target genes related to protein expression comprises a barcode sequence on its 5' end, wherein the barcode sequence of each reverse primer corresponding to each target gene related to protein expression is different,

thereby generating a plurality of amplicons;

- 20 (c) purifying the plurality of amplicons from step (b);
 (d) amplifying the purified product from step (c) by using universal indexed adapter primers to generate a sequencing library;
 (e) purifying the sequencing library obtained from step (d);
 (f) subjecting the purified sequencing library from step (e) to multiplex sequencing on a
 25 next-generation sequencing platform to obtain a plurality of sequencing reads;
 (g) deriving a consensus read of each sequence from the plurality of sequencing reads obtained from step (f);
 (h) performing a sequence alignment of the consensus read obtained from step (g) to a reference genome,
 30 (I) if the sequence alignment results in a partial alignment to the reference genome of an exon from a first gene and a partial alignment to the reference genome of an exon from a second gene, then:
 (i) determining the sequence alignment as a split read,

- (ii) counting/enumerating the number of split reads from step (h)(I)(i) that supports a fusion junction, and
 - (iii) if the number of split reads from step (h)(I)(ii) is two or more, then determining the first gene and the second gene as fusion partners,
- 5 (II) if the sequence alignment results in an alignment to the reference genome of the control housekeeping gene, then:
 - (i) determining the sequence alignment as a consensus read of the control housekeeping gene,
 - 10 (ii) counting/enumerating consensus read pairs of the control housekeeping gene from step (h)(II)(i), and
 - (iii) determining the level of gene expression of the control housekeeping gene,
- (III) if the sequence alignment results in an alignment to the reference genome of the target gene related to protein expression,
 - 15 (i) determining the sequence alignment as a consensus read of the target gene related to protein expression,
 - (ii) counting/enumerating consensus read pairs of the target gene related to protein expression from step (h)(III)(i), and
 - 20 (iii) determining the level of gene expression of the target gene related to protein expression;
- (i) determining presence or absence of the genomic alteration and/or determining presence or absence of the gene expression and/or quantifying the level of the gene expression based on the sequence alignment from step (h).
- 25 2. The method of claim 1, wherein the RNA is selected from the group consisting of cell-free RNA (cfRNA) and RNA encapsulated within tissues and/or cells.
- 3. The method of claim 1 or 2, wherein the biological sample is selected from the group consisting of a liquid sample, a tissue sample, and a cell sample
- 30 4. The method of claim 3, wherein the liquid sample is a bodily fluid, wherein optionally the bodily fluid is selected from the group consisting of blood, bone marrow, cerebral spinal fluid, peritoneal fluid, pleural fluid, lymph fluid, ascites, serous fluid, sputum, lacrimal fluid, stool, urine, saliva, ductal fluid from breast, gastric juice and pancreatic

juice, wherein optionally the bodily fluid is blood, and wherein optionally the blood is plasma.

5. The method of claim 3, wherein the tissue sample is a frozen tissue sample or a fixed tissue sample, and wherein optionally the fixed tissue sample is a Formalin-Fixed Paraffin-Embedded (FFPE) tissue sample.
6. The method of any one of claims 1 to 5, wherein the biological sample is to be obtained from a subject having and/or suspected of having cancer.
7. The method of claim 6, wherein the cancer is selected from the group consisting of leukemia, lung cancer, colorectal cancer, breast cancer, pancreatic cancer, prostate cancer, nasopharyngeal cancer, liver cancer, cholangiocarcinoma, esophageal cancer, urothelial cancer, and gastrointestinal cancer.
8. The method of claim 6 or 7, wherein the cancer is selected from the group consisting of metastatic prostate cancer, metastatic lung cancer, metastatic breast cancer, and leukemia.
9. The method of any one of claims 1 to 8, wherein the amount of RNA used in step (a) is from 6 ng to 100 ng.
10. The method of any one of claims 1 to 9, wherein step (a) is performed using a reverse transcription kit, wherein the reverse transcription kit comprises a buffer for performing reverse transcription, a reverse transcriptase enzyme and a plurality of random primers.
11. The method of any one of claims 1 to 10, wherein the plurality of multiplexed PCR reactions performed on the converted cDNA comprises 3 to 15 PCR cycles.
12. The method of any one of claims 1 to 11, wherein the barcode sequence is an oligonucleotide comprising 10 to 16 random nucleotides.
13. The method of any one of claims 1 to 12, wherein the barcode sequence is an oligonucleotide comprising 10 random nucleotides.

14. The method of any one of claims 1 to 13, wherein the target gene that is capable of undergoing genomic alteration comprises an exon from a gene known to undergo fusion fused to an exon from a partner gene of the gene known to undergo fusion.
- 5 15. The method of claim 14, wherein the gene known to undergo fusion is selected from the group consisting of ALK receptor tyrosine kinase, RET proto-oncogene, ROS proto-oncogene 1, fibroblast growth factor receptor 1 (FGFR1), fibroblast growth factor receptor 2 (FGFR2), fibroblast growth factor receptor 3 (FGFR3), neurotrophic receptor tyrosine kinase 1 (NTRK1), neurotrophic receptor tyrosine kinase 2 (NTRK2),
10 neurotrophic receptor tyrosine kinase 3 (NTRK3), neuregulin 1 (NRG1), B-Raf proto-oncogene, serine/threonine kinase (BRAF), transmembrane serine protease 2 (TMPRSS2), MET proto-oncogene, receptor tyrosine kinase (MET), epidermal growth factor receptor (EGFR), estrogen receptor 1 (ESR1), platelet derived growth factor receptor alpha (PDGFRA), androgen receptor (AR), BCR activator of RhoGEF and
15 GTPase (BCR), core-binding factor subunit beta (CBFB), lysine methyltransferase 2A (KMT2A), nucleophosmin 1 (NPM1), PML nuclear body scaffold (PML), and RUNX family transcription factor 1 (RUNX1).
16. The method of claim 14, wherein the partner gene of the gene known to undergo fusion
20 is selected from the group consisting of EMAP like 4 (EML4), kinesin family member 5B (KIF5B), coiled-coil domain containing 6 (CCDC6), CD74 molecule (CD74), transforming acidic coiled-coil containing protein 3 (TACC3), ezrin EZR), ETS transcription factor ERG (ERG), ArfGAP with GTPase domain, ankyrin repeat and PH domain 3 (AGAP3), A-kinase anchoring protein 9 (AKAP9), KIAA1549, tropomyosin
25 3 (TPM3), translocated promoter region, nuclear basket protein (TPR), trafficking from ER to golgi regulator (TFG), lamin A/C (LMNA), BicC family RNA binding protein 1 (BICC1), RAD51 recombinase (RAD51), CD47 molecule (CD47), Yes1 associated transcriptional regulator (YAP1), ETS variant transcription factor 1 (ETV1), ETS variant transcription factor 4 (ETV4), ETS variant transcription factor 5 (ETV5), ETS
30 variant transcription factor 6 (ETV6), factor interacting with PAPOLA and CPSF1 (FIP1L1), centriolin (CNTRL), ABL proto-oncogene 1, non-receptor tyrosine kinase (ABL1), AF4/FMR2 family member 1 (AFF1), MDS1 and EVI1 complex locus (MECOM), MLLT3 super elongation complex subunit (MLLT3), myosin heavy chain

11 (MYH11), PBX homeobox 1 (PBX1), retinoic acid receptor alpha (RARA), RUNX1 partner transcriptional co-repressor 1 (RUNX1T1).

- 5 17. The method of any one of claims 1 to 16, wherein the number of the plurality of forward and reverse primer pairs specific to the plurality of **target genes that are capable of undergoing genomic alteration**, the plurality of forward and reverse primer pairs specific to the plurality of **control housekeeping genes**, and the plurality of primer sets specific to a plurality of **target genes related to protein expression** is at least 300.
- 10 18. The method of any one of claims 1 to 17, wherein the length of the plurality of amplicons generated in step (b) is 90 to 110 base pairs.
- 15 19. The method of any one of claims 1 to 18, wherein the purification in step (c) and/or (e) is performed using a plurality of paramagnetic beads; and wherein optionally the paramagnetic beads are selected from the group consisting of AMPure XP beads, SPRI beads, and dynabeads.
- 20 20. The method of any one of claims 1 to 19, wherein step (g) further comprises:
 - (g)(I) detecting the presence of the barcode sequence from each sequencing read,
 - 20 (g)(II) performing cluster reassignment for the plurality of sequencing reads having the same barcode sequence to generate a plurality of barcode clusters, wherein each barcode cluster contains reads from the same amplicon and with the same barcode sequence, and
 - (g)(III) performing consensus calling for each barcode cluster to obtain the consensus
 - 25 read of each sequence.
21. The method of claim 1-13 and 17-20, wherein the step of determining presence or absence of the genomic alteration and/or determining presence or absence of the gene expression and/or quantifying the level of the gene expression, further comprises
 - 30 performing variant calling of the sequence alignment from step (h).
22. The method of claim 21, wherein the step of variant calling comprises:
 - (i) identifying differences between a consensus read and a reference genome based on the sequence alignment from step (h); and

(ii) determining the read count of sequence alignments comprising genomic alteration.

23. The method of claim 21 or 22, wherein the genomic alteration is selected from the group comprising of insertions, deletions, and single nucleotide variants, wherein optionally the insertion is a duplication.
24. A kit for detecting genomic alteration and/or detecting gene expression and/or quantifying a level of gene expression using RNA in a biological sample according to the method of claim 1-23, wherein the kit comprises:
- a plurality of forward and reverse primer pairs specific to a plurality of **target genes that are capable of undergoing genomic alteration** as defined in claim 1,
 - a plurality of forward and reverse primer pairs specific to a plurality of **control housekeeping genes** as defined in claim 1, and
 - a plurality of primer sets specific to a plurality of **target genes related to protein expression** as defined in claim 1.
25. The kit of claim 24, wherein the kit further comprises:
- a buffer for performing a plurality of multiplexed PCR reactions,
 - a reverse transcriptase,
 - a buffer for performing reverse transcription,
 - random primers,
 - a DNA polymerase, and
 - a plurality of deoxynucleoside triphosphates (dNTPs).