

DIAGNOSTIC METHODS FOR DETERMINING PROGNOSIS OF NON-SMALL CELL LUNG CANCER

CROSS REFERENCE TO A RELATED APPLICATION

5 This application claims priority from U.S. provisional patent application no. 61/254,968 filed on October 26, 2009, the disclosure of which is herein incorporated by reference in its entirety.

FIELD OF THE INVENTION

10 The present disclosure relates to *in vitro* diagnostic assays of tissue samples from lung cancer patients for determining patient prognosis, and in particular relates to an *in vitro* assay for determining prognosis of early stage patients, such as those diagnosed with Stage I or Stage II non-small cell lung cancer.

15 BACKGROUND OF THE INVENTION

 Lung cancer accounted for almost one third of cancer deaths in the United States in 2005, and is broadly classified into two types: non-small cell lung cancer and small cell lung cancer. Non-small cell lung cancer (NSCLC) comprises 80-85% of lung cancer cases in the United States. NSCLC comprises three major types: (i) Squamous cell carcinoma, which begins in
20 squamous cells, that are thin, flat cells that look like fish scales. Squamous cell carcinoma is also called epidermoid carcinoma; (ii) Large cell carcinoma, which begins in several types of large lung cells; (iii) Adenocarcinoma, which begins in the cells that line the alveoli of the lung and make substances such as mucus. Other less common types of NSCLC include pleomorphic carcinoma, carcinoid tumor and unclassified carcinoma.

25 Diagnosis of NSCLC is done by a pathologist's examination of suspected tissue, such as a biopsy sample. After NSCLC diagnosis, the patient's disease is assigned a prognosis (the chance of recovery) using the patient's overall health and age, the severity of symptoms such as coughing and difficulty in breathing, the particular type of NSCLC, and the staging of the cancer. Staging takes into account the size of the tumor and whether the tumor is present in the lung only
30 or has spread to other places in the body. The particular treatment options for a NSCLC patient

are then selected based upon these considerations, and the cancer staging is an important component for treatment selection. Patients with early stage NSCLC can potentially be cured by surgical resection to remove the tumor, but the current diagnostic modalities are not able to predict which patients will recur after surgery. Cancer is a frequently fatal disease with a low cure rate, for which the majority of treatments are directed at improving the quality and duration of life. Because cancer cells are human cells, frequently distinguished only by the accumulation of a relatively small number of genetic aberrations or protein mutations, drug therapies that are useful in killing cancer cells are commonly also detrimental to many normal human cells and cause typically significant toxicities in patients who are treated. Furthermore, because cancers frequently recur locally or metastasize to tissues and organs remote from their tissue of origin, it is critical to know which patients with early stage cancers need drug treatment after surgical removal of their primary tumor. This is an especially critical issue in patients with early stage NSCLC, whose tumors were detected early and removed surgically, specifically patients with Stage I and IIa disease. Under-treating these patients with anti-cancer drugs results in an unacceptably high rate of patients developing recurrent or metastatic disease, ultimately leading to increased morbidity and death. Over-treating this population results in an unacceptably high number of patients who, not needing drug therapy, experience the toxic side effects from the drugs given to them.

The National Comprehensive Cancer Network internet web site describes NSCLC staging as follows. “The system most often used in United States clinical practice to describe the growth and spread of non-small cell lung cancer (NSCLC) is the TNM staging system, also known as the American Joint Committee on Cancer (AJCC) system. In TNM staging, information about the tumor (T), any spread into nearby lymph nodes (N), and any distant organ metastases (M) is combined and a stage is assigned to specific TNM groupings. The grouped stages are described using the number 0 and Roman numerals from I to IV.

“T categories are based on the lung cancer’s size, its spread and location within the lungs, and its spread to nearby tissues. In the Tis category, the cancer is found only in the layer of cells lining the air passages. It has not spread into other lung tissues. This category is also known as *carcinoma in situ*.

“In the T1 category, the cancer is no larger than 3 centimeters (slightly less than 1 to 1 ¼ inches), has not spread to the visceral pleura (membranes that surround the lungs), and does not affect the main branches of the bronchi.

5 “In the T2 category, the cancer has one or more of the following features: (i) it is larger than 3 cm; (ii) it involves a main bronchus of a lung but is not closer than 2 cm (about 3 ¼ to 4 inches) to the point where the trachea (windpipe) branches into the left and right main bronchi; or (iii) has spread to the visceral pleura. The cancer may partially block the airways, but this has not caused the entire lung to collapse or develop pneumonia.

10 “In the T3 category, the cancer has one or more of the following features: (i) it has spread to the chest wall, the diaphragm (the breathing muscle that separates the chest from the abdomen), the mediastinal pleura (the membranes surrounding the space between the 2 lungs), or parietal pericardium (the membranes of the sac surrounding the heart); (ii) it involves a main bronchus of a lung, and it is closer than 2 cm to the point where the trachea (or windpipe) branches into the left and right main bronchi, but does not involve this area; or (iii) It has grown
15 into the airways enough to cause one lung to entirely collapse or to cause pneumonia of the entire lung.

“In the T4 category, the cancer has one or more of the following features: (i) It has spread to the mediastinum (the space behind the chest bone and in front of the heart), the heart, the trachea (windpipe), the esophagus (the tube connecting the throat to the stomach), the
20 backbone, or the point where the trachea branches into the left and right main bronchi; (ii) Two or more separate tumor nodules are present in the same lobe; or (iii) a malignant pleural effusion is present, which is the existence of fluid containing cancer cells in the space surrounding the lung.

“The N category depends on which, if any, of the lymph nodes near the lungs are affected
25 by the cancer. In the N0 category, the cancer has not spread to any lymph node. In the N1 category, the cancer has spread to lymph nodes within the lung or into the hilar lymph nodes (those located around the area where the bronchus enters the lung). In N1 category the affected lymph nodes are only on the same side as the cancerous lung. In the N2 category, the cancer has spread to subcarinal lymph nodes (those which are around the point where the trachea branches
30 into the left and right bronchi) or to lymph nodes in the mediastinum (the space behind the chest

bone and in front of the heart). In the N2 category, the affected lymph nodes are on the same side of the cancerous lung. In the N3 category, the cancer has spread to lymph nodes near the collarbone on either side, and/or to the hilar or mediastinal lymph nodes on the side opposite the cancerous lung.

5 “The M category depends on whether the cancer has metastasized and spread to any distant tissues and organs. In the M0 category, there is no distant cancer spread. In the M1 category, the cancer has spread to 1 or more distant sites. Sites which are considered distant include other lobes of the lungs, lymph nodes further than those used to determine the N category of the cancer, and other organs or tissues such as the liver, bones, or brain.

10 Once the T, N, and M categories have been assigned for the particular NSCLC, this information is combined (stage grouping) to assign an overall stage of 0, I, II, III, or IV (see Table 1). Various combinations of the T and N categories are combined into stages. The stages identify tumor types that have a similar prognosis and are treated in a similar way. As noted in Table 1, a tumor with distant spread (i.e., an M1 category cancer) is considered Stage IV,
15 regardless of tumor size or involvement of lymph nodes.” The following Table from the NCCN internet web site shows the combined category and stage classification for NSCLC.

TABLE 1

Overall Stage	T Category	N Category	M Category
Stage 0	Tis	N0	M0
Stage IA	T1	N0	M0
Stage IB	T2	N0	M0
Stage IIA	T1	N1	M0
Stage IIB	T2	N1	M0
	T3	N0	M0
Stage IIIA	T1	N2	M0
	T2	N2	M0
	T3	N1	M0
	T3	N2	M0
Stage IIIB	Any T	N3	M0
	T4	Any N	M0
Stage IV	Any T	Any N	M1

NSCLC patients with lower stage numbers generally have a more favorable prognosis and outlook for survival, and these patients are generally treated by surgical resection of the tumor. However, even for early stage patients, such as those with Stage 1B, Stage IIA or IIB
5 NSCLC, a significant percentage of these patients will recur after surgical resection with more aggressive disease and die. The current clinical diagnostic methods are incapable of identifying early stage NSCLC prognosis with sufficient accuracy to direct more aggressive therapy against those patients more likely to recur. Better *in vitro* diagnostic methods are needed to identify
10 higher risk, early stage NSCLC patients who should receive neoadjuvant or adjuvant chemotherapy or generally have treatment opinions re-evaluated.

Molecular *in vitro* diagnostic assays based on fluorescence *in situ* hybridization (FISH) using fluorescently labeled DNA hybridization probes to identify chromosomal abnormalities have been disclosed for use in the selection of chemotherapy for NSCLC patients,
(PCT/US2005/018879, "Methods for prediction of clinical outcome to epidermal growth factor
15 inhibitors by cancer patients", M. Garcia et al.). FISH assays have been described as an initial diagnostic assay for NSCLC in U.S. Patent Application 20060063194, "Methods and probes for the detection of cancer", L. Morrison et al., published March 23, 2006 (hereafter referred to as "Morrison '194"), the disclosure of which is incorporated herein by reference in its entirety. The Morrison '194 application describes multiple FISH probe sets useful for screening and diagnosis
20 of NSCLC, and one probe set described in Morrison '194 is commercially available as the LAVysion™ probe set from Abbott Molecular, Inc. (Des Plaines, Illinois, U.S.A.) under ASR (Analyte Specific Reagent) labeling for use by clinical laboratories to produce clinical diagnostic assays. Under the U.S. Food and Drug Administration ASR labeling requirements, the ASR labeling must not include any claims as to the medical utility of the ASR. The LAVysion ASR
25 probe set comprises four FISH probes: a chromosome 5p15 locus specific probe labeled with the SpectrumGreen green fluorophore, a chromosome 8q24 locus specific probe labeled with the SpectrumGold yellow fluorophore, a chromosome 6 enumeration probe labeled with the SpectrumAqua blue fluorophore, and a chromosome 7p12 locus specific probe labeled with the SpectrumRed red fluorophore. Research performed using the LAVysion probe set has been

described and is reviewed for example in K. Halling et al., “Fluorescence *in situ* hybridization in diagnostic cytology”, Hum. Path. (2007) 38: 1137-1144.

Overexpression of cyclin E has previously been associated with poor outcome in lung cancer (reviewed in Singhal et al., Clin. Cancer Res., 2005, 11, pp.3974—3986). However, no
5 copy number alterations at the cyclin E locus have been established as predictive markers. Moreover, no previous reports on FISH assays for NSCLC have disclosed the use of FISH probes to more accurately identify prognosis for early stage NSCLC, in particular, those classified as Stage IB or Stage II.

10 SUMMARY OF THE INVENTION

In one aspect, the present disclosure provides a method of predicting disease outcome in a patient being treated for lung cancer, the method comprising the steps of: a) providing a test sample from a patient; b) determining a copy number of a cancer outcome marker in the test sample; c) comparing the copy number of the cancer outcome marker in the test sample against a
15 baseline copy number of two, thereby determining the presence or absence of a copy number change for the cancer outcome marker in the test sample; and d) based on the presence or absence of a copy number change for the cancer outcome marker in the test sample, identifying the patient as having an increased risk of a poor disease outcome when compared to a baseline measure of disease outcome in patients having no copy number change in the cancer outcome
20 marker, wherein the presence of a copy number change in the cancer outcome marker is predictive of poor disease outcome. In one embodiment, the poor disease outcome is for example at least one of a decreased overall survival time, when compared to an overall survival time of patients having no copy number change for the cancer outcome marker, and a shorter time to recurrence, when compared to the time to recurrence for patients having no copy number
25 change for the cancer outcome marker.

In another aspect, the present disclosure provides a method of predicting treatment outcome in a patient being treated for lung cancer, the method comprising: a) providing a test sample from a patient; b) determining the presence or absence of a copy number change for a cancer outcome marker in the test sample, wherein the cancer outcome marker is a region of
30 chromosomal DNA, the change in copy number of which is associated with a poor disease

outcome; and c) based on the presence or absence of a copy number change for the cancer outcome marker, determining whether the patient has a higher risk of a decreased overall survival time or a shorter time to recurrence, when compared to an overall survival time of patients having no copy number gain for the cancer outcome marker.

5 In any of the methods, the cancer outcome marker is for example a region of chromosomal DNA, the amplification of which produces a copy number gain of the cancer outcome marker, wherein the copy number gain is associated with a poor disease outcome. Such cancer outcome markers include any selected from the group consisting of Chr 19, 34.7 Mb—35.6 Mb; Chr 19, 38.9—40.7 Mb; Chr 17, 69.2—71.3 Mb; Chr 6, 70.8—71.1 Mb; Chr 12, 93.7 kb—1.9Mb; Chr 11, 64.3—64.8 Mb; Chr 19, 57.0—62.2 Mb; Chr 6, 39.1—39.9 Mb; Chr 11, 64.8—65.7 Mb; Chr 11, 61.4—64.3 Mb; Chr 17, 51.5—53.2 Mb; Chr 17, 43.5—44.9 Mb; Chr 2, 147.6—151.1 Mb; Chr 6, 123.7—135.6 Mb; Chr 8, 6.9—8.8 Mb; Chr 2, 159.9—161.4 Mb; Chr 2, 200.9—204.2 Mb; Chr 6, 36.3—36.7 Mb; Chr 2, 205.9—208.1 Mb; and Chr 1, 109.5—111.1 Mb. In a method wherein the cancer outcome marker is Chr 19, 34.7 Mb—35.6 Mb, the marker includes nucleotide sequences encoding C19orf12; C19orf12; cyclin E1; PLEKHF1; POP4; and ZNF536. In a method wherein the cancer outcome marker is Chr 19, 38.9—40.7 Mb, the marker includes nucleotide sequences encoding ATP4A ATPase; CHST8, DMKN FAR1,2,3; FXYD1,3,5,7; GAPDHS; GPI; GPR42; GRAMD1A; HAMP; HPN; KCTD15 KIAA0355; KRTDAP; LGI4; LSM14A; LSR; MAG; PDCD2L; SAE2 SUMO1; SBSN; SCN1B; TMEM147,162; USF2; WTIP; and ZNF181,30,302,599,792. In a method wherein the cancer outcome marker is Chr 17, 69.2—71.3 Mb, the marker includes nucleotide sequences encoding ARMC7 (armadillo repeat containing 7); ATP5H ATP synthase (H⁺ transporting, mitochondrial F0 complex, subunit d); CASKIN2 (CASK interacting protein 2); CD300A (CD300a molecule); CD300C (CD300c molecule); CD300E (CD300e molecule); CD300LB (CD300 molecule-like family member b); CD300LF (CD300 molecule-like family member f); CDR2L (cerebellar degeneration-related protein 2-like); DNAI2 (dynein, axonemal, intermediate chain 2); (FADS6 fatty acid desaturase domain family, member 6); FDXR (ferredoxin reductase); GALK1 (galactokinase 1); GGA3 (golgi associated, gamma adaptin ear containing, ARF binding protein); GPR142 (G protein-coupled receptor 142); GPRC5C (G protein-coupled receptor, family C, group 5, member C); GRB2 (growth factor receptor-bound protein 2); GRIN2C

(glutamate receptor, ionotropic, N-methyl D-aspartate 2C); H3F3B (H3 histone, family 3B (H3.3B)); HN1 (hematological and neurological expressed 1 ICT1 immature colon carcinoma transcript 1); ITGB4 (integrin, beta 4); KCTD2 (potassium channel tetramerisation domain containing 2); KIAA0195; KIF19 (kinesin family member 19); LLGL2 (lethal giant larvae homolog 2 (Drosophila)); LOC388419 (galectin-3-binding protein-like); MIF4GD (MIF4G domain containing); MRPS7 (mitochondrial ribosomal protein S7); NAT9 (N-acetyltransferase 9); NT5C (5', 3'-nucleotidase, cytosolic); NUP85 (nucleoporin 85kDa); OTOP2 (otopetrin 2); OTOP3 (otopetrin 3); RAB37 (RAB37, member RAS oncogene family); RECQL5 (RecQ protein-like 5); RPL38 ribosomal protein L38; SAP30BP (SAP30 binding protein); SLC16A5 (solute carrier family 16, member 5 (monocarboxylic acid transporter 6)); SLC25A19 (solute carrier family 25 (mitochondrial thiamine pyrophosphate carrier), member 19); SLC9A3R1 (solute carrier family 9 (sodium/hydrogen exchanger), member 3 regulator 1); SUMO2 (SMT3 suppressor of mif two 3 homolog 2 (S. cerevisiae)); TMEM104 (transmembrane protein 104); TTYH2 (tweety homolog 2 (Drosophila)); UNK (unkempt homolog (Drosophila)); and USH1G (Usher syndrome 1G (autosomal recessive) [Marker 3]. In a method wherein the cancer outcome marker is Chr 6, 70.8—71.1 Mb, the marker includes nucleotide sequences encoding COL19A1 (collagen, type XIX, alpha 1), and COL9A1 (collagen, type IX, alpha 1). [Marker 4]. In a method wherein the cancer outcome marker is Chr 12, 93.7 kb—1.9Mb, the marker includes nucleotide sequences encoding ADIPOR2 (adiponectin receptor 2); B4GALNT3 (beta-1,4-N-acetyl-galactosaminyl transferase 3); CACNA2D4 (calcium channel, voltage-dependent, alpha 2/delta subunit 4); CCDC77 (coiled-coil domain containing 77); ERC1 (ELKS/RAB6-interacting/CAST family member 1); FBXL14 (F-box and leucine-rich repeat protein 14); HSN2 (hereditary sensory neuropathy, type II); IQSEC3 (IQ motif and Sec7 domain 3); JARID1A (jumonji, AT rich interactive domain 1A); LRTM2 (leucine-rich repeats and transmembrane domains 2); NINJ2 (ninjurin 2); RAD52 (RAD52 homolog (S. cerevisiae)); SLC6A12 (solute carrier family 6 (neurotransmitter transporter, betaine/GABA), member 12); SLC6A13 (solute carrier family 6 (neurotransmitter transporter, GABA), member 13); WNK1 (WNK lysine deficient protein kinase 1); and WNT5B (wingless-type MMTV integration site family, member 5B). [Marker 5]. In a method wherein the cancer outcome marker is Chr 11, 64.3—64.8 Mb, the marker includes nucleotide sequences encoding ARL2 (ADP-ribosylation factor-like 2); ATG2A

ATG2 (autophagy related 2 homolog A (*S. cerevisiae*)); BATF2 (basic leucine zipper transcription factor, ATF-like 2; CAPN1 calpain 1, (μ /I) large subunit); CDC42BPG (CDC42 binding protein kinase gamma (DMPK-like)); CDCA5 (cell division cycle associated 5); EHD1 (EH-domain containing 1); FAU (Finkel-Biskis-Reilly murine sarcoma virus (FBR-MuSV) ubiquitously expressed); GPHA2 (glycoprotein hormone alpha 2); MAP4K2 mitogen-activated protein kinase kinase kinase 2; MEN1 multiple endocrine neoplasia I; MRPL49 mitochondrial ribosomal protein L49; NAALADL1 N-acetylated alpha-linked acidic dipeptidase-like 1; POLA2 polymerase (DNA directed), alpha 2 (70kD subunit); PPP2R5B protein phosphatase 2, regulatory subunit B', beta isoform; SAC3D1 SAC3 domain containing 1
 10 SLC22A20 solute carrier family 22, member 20; SNX15 sorting nexin 15; SPDYC speedy homolog C (*Drosophila*); SYVN1 synovial apoptosis inhibitor 1, synoviolin; TM7SF2 transmembrane 7 superfamily member 2; ZFPL1 zinc finger protein-like 1; ZNHIT2 zinc finger, HIT type 2; hsa-mir-192; and hsa-mir-194-2. [Marker 6]. In a method wherein the cancer outcome marker is Chr 19, 57.0—62.2 Mb, the marker includes nucleotide sequences encoding
 15 BIRC8 (baculoviral IAP repeat-containing 8); BRSK1 (BR serine/threonine kinase 1); CACNG6,7,8 calcium channel, voltage-dependent, gamma subunit 6,7,8; CCDC106 coiled-coil domain containing 106; CDC42EP5 CDC42 effector protein (Rho GTPase binding) 5 ; CNOT3 CCR4-NOT transcription complex, subunit 3; COX6B2 cytochrome c oxidase subunit VIb polypeptide 2 (testis); DPRX divergent-paired related homeobox ; EPN1 epsin 1; EPS8L1
 20 EPS8-like 1; FCAR Fc fragment of IgA, receptor for; FIZ1 FLT3-interacting zinc finger 1; GALP galanin-like peptide; GP6 glycoprotein VI (platelet); HSPBP1 hsp70-interacting protein; IL11 interleukin 11; ISOC2 isochorismatase domain containing 2; KIR2DL1, KIR2DL4, KIR2DS4 KIR3DL1, KIR3DL3, KIR3DX1 killer cell immunoglobulin-like receptor; LAIR1,2 leukocyte-associated immunoglobulin-like receptor 1,2; LENG1,4,8,9 leukocyte receptor cluster
 25 (LRC) member 1,4,8,9; LILRA2,3,4 leukocyte immunoglobulin-like receptor, subfamily A (with TM domain), member 2,3,4; LILRB1,2,3,4,5 leukocyte immunoglobulin-like receptor, subfamily B (with TM and ITIM domains), member 1,2,3,4,5; MYADM myeloid-associated differentiation marker; NAT14 N-acetyltransferase 14; NCR1 natural cytotoxicity triggering receptor 1; NDUFA3 NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 3, 9kDa;
 30 NLRP2,4,5,7,8,9,11,12,13 NLR family, pyrin domain containing 2,4,5,7,8,9,11,12,13; OSCAR

osteoclast associated, immunoglobulin-like receptor; PEG3 paternally expressed 3; PPP1R12C protein phosphatase 1, regulatory (inhibitor) subunit 12C; PPP2R1A protein phosphatase 2 (formerly 2A), regulatory subunit A, alpha isoform; PRKCG protein kinase C, gamma; PRPF31 PRP31 pre-mRNA processing factor 31 homolog (*S. cerevisiae*); PTPRH protein tyrosine
5 phosphatase, receptor type, H; RDH13 retinol dehydrogenase 13 (all-trans/9-cis); RPL28 ribosomal protein L28; RPS9 ribosomal protein S9; SAPS1 SAPS domain family, member 1; SUV420H2 suppressor of variegation 4-20 homolog 2 (*Drosophila*); SYT5 synaptotagmin V; TFPT TCF3 (E2A) fusion partner (in childhood Leukemia); TMC4 transmembrane channel-like
10 4; TMEM190 transmembrane protein 190; TMEM86B transmembrane protein 86B; TNNT3 troponin I type 3 (cardiac); TNNT1 troponin T type 1 (skeletal, slow); TSEN34 tRNA splicing endonuclease 34 homolog (*S. cerevisiae*); TTYH1 tweety homolog 1 (*Drosophila*); U2AF2 U2 small nuclear RNA auxiliary factor 2; UBE2S ubiquitin-conjugating enzyme E2S; VN1R2 vomeronasal 1 receptor 2; VN1R4 vomeronasal 1 receptor 4; VSTM1 V-set and transmembrane domain containing 1; ZNF28,160,320,321,331,347,350,415,432,444,468,470 zinc finger protein
15 28,160,320,321,331,347,350,415,432,444,468,470; and miRNA's including hsa-mir-643, hsa-mir-512-1, hsa-mir-512-2, hsa-mir-498, hsa-mir-520e, hsa-mir-515-1, hsa-mir-519e, hsa-mir-520f, hsa-mir-515-2, hsa-mir-519c, hsa-mir-520a, hsa-mir-526b, hsa-mir-519b, hsa-mir-525, hsa-mir-523, hsa-mir-518f, hsa-mir-520b, hsa-mir-518b, hsa-mir-526a-1, hsa-mir-520c, hsa-mir-518c, hsa-mir-524, hsa-mir-517a, hsa-mir-519d, hsa-mir-521-2, hsa-mir-520d, hsa-mir-517b,
20 hsa-mir-520g, hsa-mir-516-3, hsa-mir-526a-2, hsa-mir-518e, hsa-mir-518a-1, hsa-mir-518d, hsa-mir-516-4, hsa-mir-518a-2, hsa-mir-517c, hsa-mir-520h, hsa-mir-521-1, hsa-mir-522, hsa-mir-519a-1, hsa-mir-527, hsa-mir-516-1, hsa-mir-516-2, hsa-mir-519a-2, hsa-mir-371, hsa-mir-372, hsa-mir-373, hsa-mir-516a-1, hsa-mir-516a-2, hsa-mir-516b-1, hsa-mir-516b-2, hsa-mir-517a-1, hsa-mir-517a-2, hsa-mir-520c-1, and hsa-mir-520c-2 [Marker 7]. In a method wherein the cancer
25 outcome marker is Chr 6, 39.1—39.9 Mb, the marker includes nucleotide sequences encoding C6orf64 (chromosome 6 open reading frame 64); DNAH8 dynein, axonemal, heavy chain 8; GLP1R glucagon-like peptide 1 receptor; KCNK16 potassium channel, subfamily K, member 16; KCNK17 potassium channel, subfamily K, member 17; KCNK5 potassium channel, subfamily K, member 5; and KIF6 kinesin family member 6. [Marker 8]. In a method wherein
30 the cancer outcome marker is Chr 11, 64.8—65.7 Mb, the marker includes nucleotide sequences

encoding BANF1 (barrier to autointegration factor 1); CATSPER1 cation channel, sperm
 associated 1 CCDC85B coiled-coil domain containing 85B; CDC42EP2CDC42 effector protein
 (Rho GTPase binding) 2; CFL1 cofilin 1 (non-muscle); CST6 cystatin E/M; CTSW cathepsin W;
 DPF2 D4, zinc and double PHD fingers family 2; DRAP1 DR1-associated protein 1 (negative
 5 cofactor 2 alpha); EFEMP2 EGF-containing fibulin-like extracellular matrix protein 2;
 EHBP1L1 EH domain binding protein 1-like 1; FAM89B family with sequence similarity 89,
 member B; FIBP fibroblast growth factor (acidic) intracellular binding protein; FOSL1 FOS-like
 antigen 1; FRMD8 FERM domain containing 8; GAL3ST3 galactose-3-O-sulfotransferase 3;
 HTATIP HIV-1 Tat interacting protein, 60kDa. KCNK7 potassium channel, subfamily K,
 10 member 7; LTBP3 latent transforming growth factor beta binding protein 3; MAP3K11 mitogen-
 activated protein kinase kinase kinase 11; MGC11102 hypothetical protein MGC11102; MUS81
 MUS81 endonuclease homolog (*S. cerevisiae*); OVOL1 ovo-like 1 (*Drosophila*);
 PACS1 phosphofurin acidic cluster sorting protein 1; PCNXL3 pecanex-like 3 (*Drosophila*);
 POLA2 polymerase (DNA directed) alpha 2 (70kD subunit); RELA v-rel reticuloendotheliosis
 15 viral oncogene homolog A, nuclear factor of kappa light polypeptide gene enhancer in B-cells 3,
 p65 (avian); RNASEH2C ribonuclease H2, subunit C; SART1 squamous cell carcinoma antigen
 recognized by T cells; SCYL1 SCY1-like 1 (*S. cerevisiae*); SF3B2 splicing factor 3b, subunit 2,
 145kDa; SIPA1 signal-induced proliferation-associated gene 1; SLC25A45 solute carrier family
 25, member 45; SSSCA1 Sjogren syndrome/scleroderma autoantigen 1; TIGD3 tigger
 20 transposable element derived 3; and TSGA10IP testis specific, 10 interacting protein [Marker 9].
 In a method wherein the cancer outcome marker is Chr 11, 61.4—64.3 Mb, the marker includes
 nucleotide sequences encoding AHNAK (AHNAK nucleoprotein); ASRGL1 asparaginase like 1;
 B3GAT3 beta-1,3-glucuronyltransferase 3 (glucuronosyltransferase I); BAD BCL2-antagonist of
 cell death; BEST1 bestrophin 1; BSCL2 Bernardinelli-Seip congenital lipodystrophy 2 (seipin);
 25 CCDC88B coiled-coil domain containing 88B; CHRM1 cholinergic receptor, muscarinic 1;
 COX8A cytochrome c oxidase subunit 8A (ubiquitous); DKFZP564J0863 DKFZP564J0863
 protein; DKFZP566E164 DKFZP566E164 protein; DNAJC4 DnaJ (Hsp40) homolog, subfamily
 C, member 4; EEF1G eukaryotic translation elongation factor 1 gamma; EML3 echinoderm
 microtubule associated protein like 3; ESRRA estrogen-related receptor alpha; FADS2,3 fatty
 30 acid desaturase 2,3; FKBP2 FK506 binding protein 2, 13kDa; FLRT1 fibronectin leucine rich

transmembrane protein1; FTH1 ferritin, heavy polypeptide 1; GANAB glucosidase, alpha; neutral AB; GNG3 guanine nucleotide binding protein (G protein), gamma 3; GPR137 G protein-coupled receptor 137; HRASLS2,3,5 HRAS-like suppressor 2,3,5; INCENP inner centromere protein antigens 135/155kDa; INTS5 integrator complex subunit 5; KCNK4

5 potassium channel, subfamily K, member 4; LGALS12 lectin, galactoside-binding, soluble, 12 (galectin 12); MACROD1 MACRO domain containing 1; MARK2 MAP/microtubule affinity-regulating kinase 2; MGC3196 hypothetical protein MGC3196; MTA2 metastasis associated 1 family, member 2; NAT11 N-acetyltransferase 11; NRXN2 neurexin 2; NUDT22 nudix (nucleoside diphosphate linked moiety X)-type motif 22; NXF1 nuclear RNA export factor 1;

10 OTUB1 OTU domain, ubiquitin aldehyde binding 1; PLCB3 phospholipase C, beta 3 (phosphatidylinositol-specific); POLR2G polymerase (RNA) II (DNA directed) polypeptide G; PPP1R14B protein phosphatase 1, regulatory (inhibitor) subunit 14B; PRDX5 peroxiredoxin 5; PYGM phosphorylase, glycogen; muscle (McArdle syndrome, glycogen storage disease type V); RAB3IL1 RAB3A interacting protein (rabin3)-like 1; RARRES3 retinoic acid receptor

15 responder (tazarotene induced) 3; RASGRP2 RAS guanyl releasing protein 2 (calcium and DAG-regulated); RCOR2 REST corepressor 2; ROM1 retinal outer segment membrane protein 1; RPS6KA4 ribosomal protein S6 kinase, 90kDa, polypeptide 4; RTN3 reticulon 3; SCGB1A1, 1D1, 1D2, 1D4, 2A1, 2A1 secretoglobin, family; SF1 splicing factor 1; SLC22A10, 11, 12, 6, 8, 9 solute carrier family 22 (organic anion/cation transporter) SLC3A2 solute carrier family 3

20 (activators of dibasic and neutral amino acid transport), member 2; STIP1 stress-induced-phosphoprotein 1 (Hsp70/Hsp90-organizing protein); STX5 syntaxin 5; TAF6L TAF6-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa; TRPT1 tRNA phosphotransferase 1; TTC9C tetratricopeptide repeat domain 9C; TUT1 terminal uridylyl transferase 1; U6 snRNA-specific RP2 UNC-112 related protein 2; UST6 putative UST1-like

25 organic anion transporter; VEGFB vascular endothelial growth factor B; WDR74 WD repeat domain 74; and ZBTB3 zinc finger and BTB domain containing 3. [Marker 10]. In a method wherein the cancer outcome marker is Chr 17, 51.5—53.2 Mb, the marker includes nucleotide sequences encoding AKAP1(A kinase (PRKA) anchor protein 1); ANKFN1 (ankyrin-repeat and fibronectin type III domain containing 1); C17orf67 chromosome 17 open reading frame 67;

30 COIL coilin; DGKE diacylglycerol kinase, epsilon 64kDa; MSI2 musashi homolog 2

(*Drosophila*); NOG noggin; SCPEP1 serine carboxypeptidase 1; and TRIM25 tripartite motif-containing 25. [Marker 11]. In a method wherein the cancer outcome marker is Chr 17, 43.5—44.9 Mb, the marker includes nucleotide sequences encoding hsa-mir-10a; hsa-mir-196a-1; ABI3 (ABI gene family, member 3); ATP5G1 (ATP synthase, H⁺ transporting, mitochondrial F0 complex, subunit C1 (subunit 9)); B4GALNT2 beta-1,4-N-acetyl-galactosaminyl transferase 2; 5 CALCOCO2 calcium binding and coiled-coil domain 2; CBX1 chromobox homolog 1 (HP1 beta homolog *Drosophila*); GIP gastric inhibitory polypeptide; GNGT2 guanine nucleotide binding protein (G protein), gamma transducing activity polypeptide 2; HOXB1,2,3,4,5,6,7,8,9,13 homeobox B1,2,3,4,5,6,7,8,9,13; IGF2BP1 insulin-like growth factor 10 2 mRNA binding protein 1; NFE2L1 nuclear factor (erythroid-derived 2)-like 1; NGFR nerve growth factor receptor (TNFR superfamily, member 16); PHB prohibitin PHOSPHO1 phosphatase, orphan 1; PRAC small nuclear protein PRAC; SKAP1 src kinase associated phosphoprotein 1; SNF8 SNF8, ESCRT-II complex subunit, homolog (*S. cerevisiae*); SNX11 sorting nexin 11; TTL6 tubulin tyrosine ligase-like family, member 6; UBE2Z (ubiquitin-conjugating enzyme E2Z); and ZNF652 (zinc finger protein 652). [Marker 12]. In a method 15 wherein the cancer outcome marker is Chr 2, 147.6—151.1 Mb, the marker includes nucleotide sequences encoding ACVR2A activin A receptor, type IIA; C2orf25 chromosome 2 open reading frame 25; EPC2 enhancer of polycomb homolog 2 (*Drosophila*); KIF5C kinesin family member 5C; LOC130576 hypothetical protein LOC130576; LYPD6 LY6/PLAUR domain 20 containing 6; MBD5 methyl-CpG binding domain protein 5; ORC4L origin recognition complex, subunit 4-like (yeast); and RND3 Rho family GTPase 3. [Marker 13]. In a method wherein the cancer outcome marker is Chr 6, 123.7—135.6 Mb, the marker includes nucleotide sequences encoding hsa-mir-588; AKAP7 (A kinase (PRKA) anchor protein 7); ALDH8A1 aldehyde dehydrogenase 8 family, member A1; ARG1 arginase, liver; ARHGAP18 Rho GTPase 25 activating protein 18; CTGF connective tissue growth factor; ECHDC1 enoyl Coenzyme A hydratase domain containing 1; ENPP1,3 ectonucleotide pyrophosphatase/phosphodiesterase 1,3; EPB41L2 erythrocyte membrane protein band 4.1-like 2; EYA4 eyes absent homolog 4 (*Drosophila*); HDDC2 HD domain containing 2; HEY2 hairy/enhancer-of-split related with YRPW motif 2; HINT3 histidine triad nucleotide binding protein 3; KIAA1913 KIAA1913; 30 LAMA2 laminin, alpha 2 (merosin, congenital muscular dystrophy); MED23 mediator complex

subunit 23; MOXD1 monooxygenase, DBH-like 1; MYB v-myb myeloblastosis viral oncogene homolog (avian); NCOA7 nuclear receptor coactivator 7; NKAIN2 Na⁺/K⁺ transporting ATPase interacting 2; OR2A4 olfactory receptor, family 2, subfamily A, member 4; PTPRK protein tyrosine phosphatase, receptor type, K; RNF146 ring finger protein 146; RNF217 ring finger protein 217; RPS12 ribosomal protein S12; SAMD3 sterile alpha motif domain containing 3; SGK serum/glucocorticoid regulated kinase; SLC2A12 solute carrier family 2 (facilitated glucose transporter), member 12; STX7 syntaxin 7; TAAR1,2,5,6,8,9 trace amine associated receptor 1,2,5,6,8,9; TBPL1 TBP-like 1; TCF21 transcription factor 21; TPD52L1 tumor protein D52-like 1; TRDN triadin; TRMT11 tRNA methyltransferase 11 homolog (*S. cerevisiae*)); and VNN1,2,3 (vanin 1,2,3). [Marker 14]. In a method wherein the cancer outcome marker is Chr 8, 6.9—8.8 Mb, the marker includes nucleotide sequences encoding CLDN23 claudin 23; DEFA5 defensin, alpha 5, Paneth cell-specific; DEFB103B defensin, beta 103B; DEFB104A defensin, beta 104A; DEFB104B defensin, beta 104B; DEFB105B defensin, beta 105B; DEFB106A defensin, beta 106A; DEFB106B defensin, beta 106B; DEFB107A defensin, beta 107A; DEFB107B defensin, beta 107B; DEFB4 defensin, beta 4; MFHAS1 malignant fibrous histiocytoma amplified sequence 1; PRAGMIN homolog of rat pragra of Rnd2; SPAG11A sperm associated antigen 11A; and SPAG11B sperm associated antigen 11B. [Marker 15]. In a method wherein the cancer outcome marker is Chr 2, 159.9—161.4 Mb, the marker includes nucleotide sequences encoding BAZ2B bromodomain adjacent to zinc finger domain, 2B; CD302 CD302 molecule; ITGB6 integrin, beta 6; LY75 lymphocyte antigen 75; MARCH7 (membrane-associated ring finger (C3HC4) 7); PLA2R1 (phospholipase A2 receptor 1, 180kDa); and RBMS1 (RNA binding motif, single stranded interacting protein 1). [Marker 16]. In a method wherein the cancer outcome marker is Chr 2, 200.9—204.2 Mb, the marker includes nucleotide sequences encoding ABI2 abl interactor 2; ALS2 amyotrophic lateral sclerosis 2 (juvenile); ALS2CR2, 4, 7, 8, 11, 12, 13 amyotrophic lateral sclerosis 2 (juvenile) chromosome region, candidate 2, 4, 7, 8, 11, 12, 13; AOX1 aldehyde oxidase 1; BMPR2 bone morphogenetic protein receptor, type II (serine/threonine kinase); BZW1 basic leucine zipper and W2 domains 1; CASP10 caspase 10, apoptosis-related cysteine peptidase; CASP8 caspase 8, apoptosis-related cysteine peptidase; CFLAR CASP8 and FADD-like apoptosis regulator; CLK1 CDC-like kinase 1; CYP20A1 cytochrome P450, family 20, subfamily A, polypeptide 1; FAM126B family with

sequence similarity 126, member B; FZD7 frizzled homolog 7 (*Drosophila*) ICA1L islet cell autoantigen 1,69kDa-like; KCTD18 potassium channel tetramerisation domain containing 18; LOC26010 viral DNA polymerase-transactivated protein 6; MPP4 membrane protein, palmitoylated 4 (MAGUK p55 subfamily member 4). NBEAL1 neurobeachin-like 1; NDUFB3

5 NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 3, 12kDa; NIF3L1 NIF3 NGG1 interacting factor 3-like 1 (*S. pombe*); NOP5/NOP58 nucleolar protein NOP5/NOP58; ORC2L origin recognition complex, subunit 2-like (yeast); PPIL3 peptidylprolyl isomerase (cyclophilin)-like 3; RAPH1 Ras association (RalGDS/AF-6) and pleckstrin homology domains 1; SGOL2 shugoshin-like 2 (*S. pombe*); SUMO1 SMT3 suppressor of mif two 3 homolog 1 (*S. cerevisiae*);

10 TRAK2 trafficking protein, kinesin binding 2; and WDR12 (WD repeat domain 12). [Marker 17]. In a method wherein the cancer outcome marker is Chr 6, 36.3—36.7 Mb, the marker includes nucleotide sequences encoding BRPF3 (bromodomain and PHD finger containing, 3); DKFZp779B1540 hypothetical protein DKFZp779B1540; ETV7 ets variant gene 7 (TEL2 oncogene); KCTD20 potassium channel tetramerisation domain containing 20; PNPLA1 patatin-like phospholipase domain containing 1; PXT1 peroxisomal, testis specific 1; SFRS3 splicing

15 factor, arginine/serine-rich 3; and STK38 (serine/threonine kinase 38). [Marker 18]. In a method wherein the cancer outcome marker is Chr 2, 205.9—208.1 Mb, the marker includes nucleotide sequences encoding ADAM23 (ADAM metallopeptidase domain 23); CPO carboxypeptidase O; DYT1 dystrotelin; EEF1B2 eukaryotic translation elongation factor 1 beta

20 2; FASTKD2 FAST kinase domains 2; FLJ20309 hypothetical protein FLJ20309; GPR1 G protein-coupled receptor 1; KLF7 Kruppel-like factor 7 (ubiquitous); MDH1B malate dehydrogenase 1B, NAD (soluble); NDUFS1 NADH dehydrogenase (ubiquinone) Fe-S protein 1, 75kDa (NADH-coenzyme Q reductase); NRP2 neuropilin 2; PARD3B par-3 partitioning defective 3 homolog B (*C. elegans*); ZDBF2 (zinc finger, DBF-type containing 2); and

25 hCG_1657980 hCG1657980. [Marker 19]. In a method wherein the cancer outcome marker is Chr 1, 109.5—111.1 Mb, the marker includes nucleotide sequences encoding hsa-mir-197; AHCYL1 S-adenosylhomocysteine hydrolase-like 1; ALX3 aristaless-like homeobox 3; AMIGO1 adhesion molecule with Ig-like domain 1; AMPD2 adenosine monophosphate deaminase 2 (isoform L); ATXN7L2 ataxin 7-like 2; CELSR2 cadherin, EGF LAG seven-pass

30 G-type receptor 2 (flamingo homolog, *Drosophila*); CSF1 colony stimulating factor 1

(macrophage); CYB561D1 cytochrome b-561 domain containing 1; EPS8L3 EPS8-like 3; FAM40A family with sequence similarity 40, member A; GNAI3 guanine nucleotide binding protein (G protein), alpha inhibiting activity polypeptide 3; GNAT2 guanine nucleotide binding protein (G protein), alpha transducing activity polypeptide 2; GPR61 G protein-coupled receptor
 5 61; GSTM1,M2,M3,M4,M5 glutathione S-transferase M1, M2 (muscle), M3 (brain), M4, M5; HBXIP hepatitis B virus x interacting protein; KCNA2,3,4,10 potassium voltage-gated channel, shaker-related subfamily, member 2,3,4,10; KIAA1324 KIAA1324; MYBPHL myosin binding protein H-like; PROK1 prokineticin 1; PSMA5 proteasome (prosome, macropain) subunit, alpha type, 5; PSRC1 proline/serine-rich coiled-coil 1; RBM15 RNA binding motif protein 15; SARS
 10 seryl-tRNA synthetase; SLC16A4 solute carrier family 16, member 4 (monocarboxylic acid transporter 5) ; SLC6A17 solute carrier family 6, member 17; SORT1 sortilin 1; SYPL2 synaptophysin-like 2;and UBL4B (ubiquitin-like 4B). [Marker 20].

Alternatively, in any of the methods the cancer outcome marker is for example a region of chromosomal DNA, the deletion of which produces a copy number loss of the cancer outcome
 15 marker, wherein the copy number loss is associated with a poor disease outcome. Such cancer outcome markers can be selected from the group consisting of Chr 5, 62.9 – 67.8 Mb; Chr 5, 53.3 – 53.8 Mb; Chr 4, 105.8 – 107.2 Mb; Chr 16, 45.8 – 46.3 Mb; Chr 5, 50.7 – 52.0 Mb; Chr 5, 94.2 – 96.1 Mb; Chr 9, 36.1 – 37.0 Mb; Chr 5, 94.2 – 96.1 Mb; Chr14, 51.1 – 52.8 Mb; Chr 14, 61.5 – 68.6 Mb; Chr 9, 28.1 Mb; Chr 4, 43.7 – 44.2 Mb; Chr 5, 60.8 – 62.9 Mb; Chr 3, 120.0 –
 20 121.1 Mb; Chr 4, 46.2 – 48.0 Mb; Chr 14, 38.9 – 40.0 Mb; Chr 4, 44.2 – 44.6 Mb; Chr 2, 213.7 – 214.3 Mb; Chr14, 43.9 – 46.6 Mb; Chr 14, 27.6 – 28.6 Mb; Chr 3, 98.0 – 98.3 Mb; Chr14, 55.2 – 60.0 Mb; Chr14, 48.7 – 51.1 Mb; Chr 4, 81.4 – 83.2 Mb; Chr 10, 51.9 – 54.2 Mb; Chr 5, 55.2 – 58.6 Mb; and Chr 5, 67.8 – 68.5 Mb. In a method wherein the cancer outcome marker is Chr 5, 62.9 – 67.8 Mb, the marker includes nucleotide sequences encoding ADAMTS6 ADAM
 25 metallopeptidase with thrombospondin type 1 motif, 6; CD180 CD180 molecule; CENPK centromere protein K; ERBB2IP erbb2 interacting protein; FLJ13611 hypothetical protein FLJ13611; HTR1A 5-hydroxytryptamine (serotonin) receptor 1A; MAST4 microtubule associated serine/threonine kinase family member 4; NLN neurolysin (metallopeptidase M3 family); P18SRP P18SRP protein; PIK3R1 phosphoinositide-3-kinase, regulatory subunit 1 (p85
 30 alpha); PPWD1 peptidylprolyl isomerase domain and WD repeat containing 1; RGS7BP

regulator of G-protein signaling 7 binding protein; RNF180 ring finger protein 180; SDCCAG10 serologically defined colon cancer antigen 10; SFRS12 splicing factor, arginine/serine-rich 12; SGTB small glutamine-rich tetratricopeptide repeat (TPR)-containing, beta0; and TRIM23 tripartite motif-containing 23. [Deletion Marker 1]. In a method wherein the cancer outcome marker is Chr 5, 53.3 – 53.8 Mb, the marker includes nucleotide sequences encoding ARL15 (ADP-ribosylation factor-like 15); HSPB3 (heat shock 27kDa protein 3) and hsa-miR-581. [Deletion Marker 2]. In a method wherein the cancer outcome marker is Chr 4, 105.8 – 107.2 Mb, the marker includes nucleotide sequences encoding FLJ20184 (hypothetical protein FLJ20184); GSTCD (glutathione S-transferase, C-terminal domain containing); INTS12 integrator complex subunit 12; KIAA1546 KIAA1546; MGC16169 hypothetical protein MGC16169; NPNT (nephronectin); and PPA2 pyrophosphatase (inorganic) 2. [Deletion Marker 3]. In a method wherein the cancer outcome marker is Chr 16, 45.8 – 46.3 Mb, the marker includes nucleotide sequences encoding ITFG1 (integrin alpha FG-GAP repeat containing 1) and PHKB (phosphorylase kinase, beta). [Deletion Marker 4]. In a method wherein the cancer outcome marker is Chr 5, 50.7 – 52.0 Mb, the marker includes a nucleotide sequence encoding ISL1 (ISL LIM homeobox). [Deletion Marker 5]. In a method wherein the cancer outcome marker is Chr 5, 94.2 – 96.1 Mb, the marker includes nucleotide sequences encoding ARSK (arylsulfatase family, member K); CAST (calpastatin); ELL2 (elongation factor, RNA polymerase II, 2); FAM81B family with sequence similarity 81, member B; GLRX glutaredoxin (thioltransferase); GPR150 G protein-coupled receptor 150; KIAA0372 KIAA0372; MCTP1 multiple C2 domains, transmembrane 1; PCSK1 proprotein convertase subtilisin/kexin type 1; RFESD (Rieske (Fe-S) domain containing); RHOBTB3 Rho-related BTB domain containing 3; SPATA9 (spermatogenesis associated 9); and hsa-miR-583. [Deletion Marker 6]. In a method wherein the cancer outcome marker is Chr 9, 36.1 – 37.0 Mb, the marker includes nucleotide sequences encoding C9orf19 chromosome 9 open reading frame 19; CCIN calicin; CLTA clathrin, light chain (Lca); GNE glucosamine (UDP-N-acetyl)-2-epimerase/N-acetylmannosamine kinase; MELK maternal embryonic leucine zipper kinase; PAX5 paired box 5; RECK reversion-inducing-cysteine-rich protein with kazal motifs; and RNF38 ring finger protein 38. [Deletion Marker 7]. In a method wherein the cancer outcome marker is Chr 5, 94.2 – 96.1 Mb, the marker includes nucleotide sequences encoding ARSK arylsulfatase family,

member K; CAST calpastatin; ELL2 elongation factor, RNA polymerase II, 2; FAM81B family with sequence similarity 81, member B; GLRX glutaredoxin (thioltransferase); GPR150 G protein-coupled receptor 150; KIAA0372 KIAA0372; MCTP1 multiple C2 domains, transmembrane 1; PCSK1 proprotein convertase subtilisin/kexin type 1; RFESD Rieske (Fe-S) domain containing; RHOBTB3 Rho-related BTB domain containing 3; SPATA9 spermatogenesis associated 9. [Deletion Marker 8]. In a method wherein the cancer outcome marker is Chr14, 51.1 – 52.8 Mb, the marker includes nucleotide sequences encoding; C14orf166 chromosome 14 open reading frame 166; DDHD1 DDHD domain containing 1; ERO1L ERO1-like (*S. cerevisiae*); FRMD6 FERM domain containing 6; GNG2 guanine nucleotide binding protein (G protein), gamma 2; GNPAT1 glucosamine-phosphate N-acetyltransferase 1; GPR137C G protein-coupled receptor 137C; NID2 nidogen 2 (osteonidogen); PLEKHC1 pleckstrin homology domain containing, family C (with FERM domain) member 1; PSMC6 proteasome (prosome, macropain) 26S subunit, ATPase, 6; PTGDR prostaglandin D2 receptor (DP); PTGER2 prostaglandin E receptor 2 (subtype EP2), 53kDa; STYX serine/threonine/tyrosine interacting protein; TXNDC16 thioredoxin domain containing 16. [Deletion Marker 9]. In a method wherein the cancer outcome marker is Chr 14, 61.5 – 68.6 Mb, the marker includes nucleotide sequences encoding ACTN1 actinin, alpha 1; AKAP5 A kinase (PRKA) anchor protein 5; ARG2 arginase, type II; ATP6V1D ATPase, H⁺ transporting, lysosomal 34kDa, V1 subunit D; C14orf50 chromosome 14 open reading frame 50; C14orf54 chromosome 14 open reading frame 54; C14orf83 chromosome 14 open reading frame 83; CHURC1 churchill domain containing 1; EIF2S1 eukaryotic translation initiation factor 2, subunit 1 alpha, 35kDa; ESR2 estrogen receptor 2 (ER beta); FLJ39779 FLJ39779 protein; FNTB farnesyltransferase, CAAX box, beta; FUT8 fucosyltransferase 8 (alpha (1,6) fucosyltransferase); GPHB5 glycoprotein hormone beta 5; GPHN gephyrin; GPX2 glutathione peroxidase 2 (gastrointestinal); HSPA2 heat shock 70kDa protein 2; KCNH5 potassium voltage-gated channel, subfamily H (eag-related), member 5; MAX MYC associated factor X; MPP5 membrane protein, palmitoylated 5 (MAGUK p55 subfamily member 5); MTHFD1 methylenetetrahydrofolate dehydrogenase (NADP⁺ dependent) 1, methenyltetrahydrofolate cyclohydrolase, formyltetrahydrofolate synthetase; PIGH phosphatidylinositol glycan anchor biosynthesis, class H; PLEK2 pleckstrin 2; PLEKHG3 pleckstrin homology domain containing,

family G (with RhoGef domain) member 3; PLEKHH1 pleckstrin homology domain containing, family H (with MyTH4 domain) member 1; PPP2R5E protein phosphatase 2, regulatory subunit B', epsilon isoform; RAB15 RAB15, member RAS oncogene family; RAD51L1 RAD51-like 1 (*S. cerevisiae*); RDH11 retinol dehydrogenase 11 (all-trans/9-cis/11-cis); RDH12 retinol dehydrogenase 12 (all-trans/9-cis/11-cis); RHOJ ras homolog gene family, member J; SGPP1 sphingosine-1-phosphate phosphatase 1; SPTB spectrin, beta, erythrocytic (includes spherocytosis, clinical type I); SYNE2 spectrin repeat containing, nuclear envelope 2; SYT16 synaptotagmin XVI; VTI1B vesicle transport through interaction with t-SNAREs homolog 1B (yeast); WDR22 WD repeat domain 22; WDR89 WD repeat domain 89; ZBTB1 zinc finger and BTB domain containing 1; ZBTB25 zinc finger and BTB domain containing 25; ZFP36L1 zinc finger protein 36, C3H type-like 1; ZFYVE26 zinc finger, FYVE domain containing 26 and hsa-miR-625. [Deletion Marker 10]. In a method wherein the cancer outcome marker is Chr 9, 28.1 Mb, the marker includes a nucleotide sequence encoding LINGO2 (leucine rich repeat and Ig domain containing 2). [Deletion Marker 11]. In a method wherein the cancer outcome marker is Chr 4, 43.7 – 44.2 Mb, the marker includes a nucleotide sequence encoding KCTD8 (potassium channel tetramerisation domain containing 8). [Deletion Marker 12]. In a method wherein the cancer outcome marker is Chr 5, 60.8 – 62.9 Mb, the marker includes nucleotide sequences encoding DIMT1L DIM1 dimethyladenosine transferase 1-like (*S. cerevisiae*); FLJ37543 hypothetical protein FLJ37543; IPO11 importin 11; ISCA1L iron-sulfur cluster assembly 1 homolog (*S. cerevisiae*)-like; and KIF2A kinesin heavy chain member 2A. [Deletion Marker 13]. In a method wherein the cancer outcome marker is Chr 3, 120.0 – 121.1 Mb, the marker includes nucleotide sequences encoding ADPRH ADP-ribosylarginine hydrolase; B4GALT4 UDP-Gal:betaGlcNAc beta 1,4- galactosyltransferase, polypeptide 4; C3orf1 chromosome 3 open reading frame 1; C3orf15 chromosome 3 open reading frame 15; C3orf30 chromosome 3 open reading frame 30; CD80 CD80 molecule; CDGAP Cdc42 GTPase-activating protein; COX17 COX17 cytochrome c oxidase assembly homolog (*S. cerevisiae*); GSK3B glycogen synthase kinase 3 beta; IGSF11 immunoglobulin superfamily, member 11; KTELC1 KTEL (Lys-Tyr-Glu-Leu) containing 1; NR1I2 nuclear receptor subfamily 1, group I, member 2; PLA1A phospholipase A1 member A; POPDC2 popeye domain containing 2; TMEM39A transmembrane protein 39A; and UPK1B uroplakin 1B. [Deletion Marker 14]. In a method

wherein the cancer outcome marker is Chr 4, 46.2 – 48.0 Mb, the marker includes nucleotide sequences encoding ATP10D ATPase, class V, type 10D; CNGA1 cyclic nucleotide gated channel alpha 1; COMMD8 COMM domain containing 8; CORIN corin, serine peptidase; COX7B2 cytochrome c oxidase subunit VIIb2; GABRA4 gamma-aminobutyric acid (GABA) A receptor, alpha 4; GABRB1 gamma-aminobutyric acid (GABA) A receptor, beta 1; NFXL1 nuclear transcription factor, X-box binding-like 1; NPAL1 NIPA-like domain containing 1; TEC tec protein tyrosine kinase; and TXK TXK tyrosine kinase. [Deletion Marker 15]. In a method wherein the cancer outcome marker is Chr 14, 38.9 – 40.0 Mb, the marker includes a nucleotide sequence encoding FBXO33 (F-box protein 33). [Deletion Marker 16]. In a method wherein the cancer outcome marker is Chr 4, 44.2 – 44.6 Mb, the marker includes nucleotide sequences encoding GNPDA2 (glucosamine-6-phosphate deaminase 2); GUF1 (GUF1 GTPase homolog (*S. cerevisiae*)); and YIPF7 (Yip1 domain family, member 7). [Deletion Marker 17]. In a method wherein the cancer outcome marker is Chr 2, 213.7 – 214.3 Mb, the marker includes nucleotide sequences encoding IKZF2 IKAROS family zinc finger 2 (Helios); and SPAG16 sperm associated antigen 16. [Deletion Marker 18]. In a method wherein the cancer outcome marker is Chr14, 43.9 – 46.6 Mb, the marker includes nucleotide sequences encoding C14orf106 chromosome 14 open reading frame 106; C14orf155 chromosome 14 open reading frame 155; C14orf28 chromosome 14 open reading frame 28; FANCM Fanconi anemia, complementation group M; FKBP3 FK506 binding protein 3, 25kDa; KIAA0423 KIAA0423; KLHL28 kelch-like 28 (*Drosophila*); MDGA2 MAM domain containing glycosylphosphatidylinositol anchor 2; PRPF39 PRP39 pre-mRNA processing factor 39 homolog (*S. cerevisiae*); and RPL10L ribosomal protein L10-like. [Deletion Marker 19]. In a method wherein the cancer outcome marker is Chr 14, 27.6 – 28.6 Mb, the marker includes a nucleotide sequence encoding FOXG1 (forkhead box G1). [Deletion Marker 20]. In a method wherein the cancer outcome marker is Chr 3, 98.0 – 98.3 Mb, the marker includes nucleotide sequences encoding EPHA6 (EPH receptor A6). [Deletion Marker 21]. In a method wherein the cancer outcome marker is Chr14, 55.2 – 60.0 Mb, the marker includes nucleotide sequences encoding ACTR10 actin-related protein 10 homolog (*S. cerevisiae*); ARID4A AT rich interactive domain 4A (RBP1-like); C14orf100 chromosome 14 open reading frame 100; C14orf101 chromosome 14 open reading frame 101; C14orf105 chromosome 14 open reading frame 105; C14orf108 chromosome 14

open reading frame 108; C14orf135 chromosome 14 open reading frame 135; C14orf149 chromosome 14 open reading frame 149; C14orf37 chromosome 14 open reading frame 37; C14orf39 chromosome 14 open reading frame 39; DAAM1 dishevelled associated activator of morphogenesis 1; DACT1 dapper, antagonist of beta-catenin, homolog 1 (*Xenopus laevis*);

5 DHRS7 dehydrogenase/reductase (SDR family) member 7; EXOC5 exocyst complex component 5; GPR135 G protein-coupled receptor 135; KIAA0586 KIAA0586; NAT12 N-acetyltransferase 12; OTX2 orthodenticle homeobox 2; PELI2 pellino homolog 2 (*Drosophila*); PPM1A protein phosphatase 1A (formerly 2C), magnesium-dependent, alpha isoform; PSMA3 proteasome (prosome, macropain) subunit, alpha type, 3; RTN1 reticulon 1; SLC35F4 solute carrier family

10 35, member F4; TIMM9 translocase of inner mitochondrial membrane 9 homolog (yeast); and UNQ9438 TIMM. [Deletion Marker 22]. In a method wherein the cancer outcome marker is Chr14, 48.7 – 51.1 Mb, the marker includes nucleotide sequences encoding ABHD12B abhydrolase domain containing 12B; ARF6 ADP-ribosylation factor 6; ATP5S ATP synthase, H⁺ transporting, mitochondrial F0 complex, subunit s (factor B); C14orf104 chromosome 14

15 open reading frame 104; C14orf138 chromosome 14 open reading frame 138; CDKL1 cyclin-dependent kinase-like 1 (CDC2-related kinase); FRMD6 FERM domain containing 6; KLHDC1 kelch domain containing 1; KLHDC2 kelch domain containing 2; L2HGDH L-2-hydroxyglutarate dehydrogenase; LOC196913 hypothetical protein LOC196913; LOC283551 hypothetical protein LOC283551; MAP4K5 mitogen-activated protein kinase kinase kinase

20 kinase 5; MGAT2 mannosyl (alpha-1,6-)-glycoprotein beta-1,2-N-acetylglucosaminyltransferase; NIN ninein (GSK3B interacting protein); POLE2 polymerase (DNA directed), epsilon 2 (p59 subunit); PPIL5 peptidylprolyl isomerase (cyclophilin)-like 5 PYGL phosphorylase, glycogen; liver (Hers disease, glycogen storage disease type VI); RPL36AL ribosomal protein L36a-like; and RPS29 ribosomal protein S29. [Deletion Marker

25 23]. In a method wherein the cancer outcome marker is Chr 4, 81.4 – 83.2 Mb, the marker includes nucleotide sequences encoding BMP3 bone morphogenetic protein 3 (osteogenic); C4orf22 chromosome 4 open reading frame 22; FGF5 fibroblast growth factor 5; PRKG2 protein kinase, cGMP-dependent, type II; and RASGEF1B RasGEF domain family, member 1B. [Deletion Marker 24]. In a method wherein the cancer outcome marker is Chr 10, 51.9 – 54.2

30 Mb, the marker includes nucleotide sequences encoding ACF apobec-1 complementation factor;

ASAH2B N-acylsphingosine amidohydrolase (non-lysosomal ceramidase) 2B; CSTF2T
 cleavage stimulation factor, 3' pre-RNA, subunit 2, 64kDa, tau variant; DKK1 dickkopf homolog
 1 (*Xenopus laevis*); MBL2 mannose-binding lectin (protein C) 2, soluble (opsonic defect);
 PRKG1 protein kinase, cGMP-dependent, type I; SGMS1 sphingomyelin synthase 1; and hsa-
 5 miR-605. [Deletion Marker 25]. In a method wherein the cancer outcome marker is Chr 5, 55.2
 – 58.6 Mb, the marker includes nucleotide sequences encoding ANKRD55 ankyrin repeat
 domain 55; C5orf29 chromosome 5 open reading frame 29; C5orf35 chromosome 5 open
 reading frame 35; DKFZp686D0972 similar to RIKEN cDNA 4732495G21 gene; GPBP1 GC-
 rich promoter binding protein 1; IL31RA interleukin 31 receptor A; IL6ST interleukin 6 signal
 10 transducer (gp130, oncostatin M receptor); MAP3K1 mitogen-activated protein kinase kinase
 kinase 1; MIER3 mesoderm induction early response 1, family member 3; PDE4D
 phosphodiesterase 4D, cAMP-specific (phosphodiesterase E3 dunce homolog, *Drosophila*);
 PLK2 polo-like kinase 2 (*Drosophila*); and RAB3C RAB3C, member RAS oncogene family.
 [Deletion Marker 26]. In a method wherein the cancer outcome marker is Chr 5, 67.8 – 68.5 Mb,
 15 the marker includes nucleotide sequences encoding CCNB1 (cyclin B1) and SLC30A5 (solute
 carrier family 30 (zinc transporter), member 5). [Deletion Marker 27].

In any of the methods, the test sample can be a tissue sample that may contain tumor
 cells, such as for example a blood sample, a tumor tissue or a suspected tumor tissue, a thin layer
 cytological sample, a fine needle aspirate sample, a lung wash sample, a pleural effusion sample,
 20 a fresh frozen tissue sample, a paraffin embedded tissue sample or an extract or processed
 sample produced from any of the preceding. In exemplary embodiments, the tissue sample is a
 lung tissue sample or a peripheral blood sample comprising circulating tumor cells. In any of the
 methods, the determining step (b) can be performed by *in situ* hybridization. The *in situ*
 hybridization can be performed with a nucleic acid probe that is fluorescently labeled, with at
 25 least two nucleic acid probes, or a peptide nucleic acid probe. The determining step (b) may be
 performed by polymerase chain reaction, a nucleic acid sequencing assay, or nucleic acid
 microarray assay. In an exemplary embodiment, the lung cancer is non-small-cell lung cancer,
 which can be for example squamous cell carcinoma, large cell carcinoma or adenocarcinoma. In
 any of the methods, the patient can receiving treatment with chemotherapy, radiation, surgery or
 30 any combination thereof.

In another aspect, the present disclosure provides a method of selecting a treatment for a patient suffering from lung cancer, the method comprising the steps of: a) providing a test sample from the patient wherein treatment with a chemotherapy agent is at least one treatment option for the patient; b) determining a copy number of the cancer outcome marker in the test sample; c) comparing the copy number of the cancer outcome marker in the test sample against a baseline copy number of two thereby determining the presence or absence of a copy number change for the cancer outcome marker in the test sample; and d) determining a chemotherapy treatment regimen based on the comparison in step c). The step of determining a treatment regimen based on the comparison in step c) includes for example selecting a chemotherapy agent and determining a frequency of chemotherapy treatment when a copy number change is present for the cancer outcome marker.

In another aspect, the present disclosure provides a method of classifying a patient as having a lung cancer that is resistant to treatment comprising the steps of: a) providing a test sample from a patient; b) determining a copy number for the cancer outcome marker; c) comparing the copy number for the cancer outcome marker in the test sample against a baseline copy number of two for the cancer outcome marker to determine the presence or absence of a copy number change in the cancer outcome marker in the patient; and d) classifying the patient as having a lung cancer that is resistant to treatment based on the presence of a copy number change in the cancer outcome marker.

In another aspect, the present disclosure provides a kit comprising: a) reagents for determining the presence or absence of a copy number change for the cancer outcome marker; and b) instructions for performing the test. The reagents to determine the presence or absence of a copy number change for the cancer outcome marker can include for example detectably-labeled polynucleotides that hybridize to at least a portion of the cancer outcome marker. The cancer outcome marker can be a region of chromosomal DNA, the amplification of which produces a copy number gain of the cancer outcome marker, wherein the copy number gain is associated with a poor disease outcome. Such cancer outcome markers can be selected from the group consisting of Chr 19, 34.7 Mb—35.6 Mb; Chr 19, 38.9—40.7 Mb; Chr 17, 69.2—71.3 Mb; Chr 6, 70.8—71.1 Mb; Chr 12, 93.7 kb—1.9Mb; Chr 11, 64.3—64.8 Mb; Chr 19, 57.0—62.2 Mb; Chr 6, 39.1—39.9 Mb; Chr 11, 64.8—65.7 Mb; Chr 11, 61.4—64.3 Mb; Chr 17,

51.5—53.2 Mb; Chr 17, 43.5—44.9 Mb; Chr 2, 147.6—151.1 Mb; Chr 6, 123.7—135.6 Mb; Chr 8, 6.9—8.8 Mb; Chr 2, 159.9—161.4 Mb; Chr 2, 200.9—204.2 Mb; Chr 6, 36.3—36.7 Mb; Chr 2, 205.9—208.1 Mb; and Chr 1, 109.5—111.1 Mb. The cancer outcome marker can be a region of chromosomal DNA, the deletion of which produces a copy number loss of the cancer outcome marker, wherein the copy number loss is associated with a poor disease outcome. Such cancer outcome markers can be selected from the group consisting of Chr 5, 62.9 – 67.8 Mb; Chr 5, 53.3 – 53.8 Mb; Chr 4, 105.8 – 107.2 Mb; Chr 16, 45.8 – 46.3 Mb; Chr 5, 50.7 – 52.0 Mb; Chr 5, 94.2 – 96.1 Mb; Chr 9, 36.1 – 37.0 Mb; Chr 5, 94.2 – 96.1 Mb; Chr14, 51.1 – 52.8 Mb; Chr 14, 61.5 – 68.6 Mb; Chr 9, 28.1 Mb; Chr 4, 43.7 – 44.2 Mb; Chr 5, 60.8 – 62.9 Mb; Chr 3, 120.0 – 121.1 Mb; Chr 4, 46.2 – 48.0 Mb; Chr 14, 38.9 – 40.0 Mb; Chr 4, 44.2 – 44.6 Mb; Chr 2, 213.7 – 214.3 Mb; Chr14, 43.9 – 46.6 Mb; Chr 14, 27.6 – 28.6 Mb; Chr 3, 98.0 – 98.3 Mb; Chr14, 55.2 – 60.0 Mb; Chr14, 48.7 – 51.1 Mb; Chr 4, 81.4 – 83.2 Mb; Chr 10, 51.9 – 54.2 Mb; Chr 5, 55.2 – 58.6 Mb; and Chr 5, 67.8 – 68.5 Mb.

BRIEF DESCRIPTION OF THE DRAWINGS

The patent or application file contains at least one drawing executed in color. Copies of this patent or patent application publication with color drawing(s) will be provided by the Office upon request and payment of the necessary fee.

Figure 1 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr19, 34.7 Mb—35.6 Mb. [marker 1]

Figure 2 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 19; 38.9—40.7 Mb. [marker 2]

Figure 3 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 17; 69.2—71.3 Mb. [marker 3]

Figure 4 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 71 patient cohort with NSCLC stage Ib-IIb, classified by presence or absence of a copy number gain in Chr 6, 70.8—71.1 Mb. [marker 4]

Figure 5 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 102 patient cohort with NSCLC stage Ia-IIb, classified by presence or absence of a copy number gain in Chr 6, 70.8—71.1 Mb. [marker 4]

Figure 6 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 12, 93.7 kb—1.9Mb. [marker 5]

Figure 7 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 11, 64.3—64.8 Mb. [marker 6]

Figure 8 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 11, 64.3—64.8 Mb. [marker 6]

Figure 9 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 19, 57.0—62.2 Mb. [marker 7]

Figure 10 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 6, 39.1—39.9 Mb. [marker 8]

Figure 11 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 11, 64.8—65.7 Mb. [marker 9]

Figure 12 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 11, 64.8—65.7 Mb. [marker 9]

Figure 13 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 11, 61.4—64.3 Mb. [marker 10]

Figure 14 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 17, 51.5—53.2 Mb. [marker 11]

Figure 15 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 17, 43.5—44.9 Mb. [marker 12]

Figure 16 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 17, 43.5—44.9 Mb. [marker 12]

Figure 17 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 2, 147.6—151.1 Mb. [marker 13]

Figure 18 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 2, 147.6—151.1 Mb. [marker 13]

Figure 19 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIb, classified by presence or absence of a copy number gain in Chr 6, 123.7—135.6 Mb. [marker 14]

Figure 20 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 8, 6.9—8.8 Mb. [marker 15]

Figure 21 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 2, 159.9—161.4 Mb. [marker 16]

Figure 22 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 2, 159.9—161.4 Mb. [marker 16]

Figure 23 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 2, 200.9—204.2 Mb. [marker 17]

Figure 24 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 6, 36.3—36.7 Mb. [marker 18]

Figure 25 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 2, 205.9—208.1 Mb. [marker 19]

Figure 26 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 66 patient cohort with NSCLC stage Ib-IIb, classified by presence or absence of a copy number gain in Chr 2, 205.9—208.1 Mb. [marker 19]

Figure 27 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 2, 205.9—208.1 Mb. [marker 19]

Figure 28 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 71 patient cohort with NSCLC stage Ib-IIb, classified by presence or absence of a copy number gain in Chr 1, 109.5—111.1 Mb. [marker 20]

Figure 29 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 102 patient cohort with NSCLC stage Ia-IIb, classified by presence or absence of a copy number loss (deletion) of Chr 5, 62.9 – 67.8 Mb. [deletion marker 1]

Figure 30 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 102 patient cohort with NSCLC stage Ia-IIb, classified by presence or absence of a copy number loss (deletion) of Chr 5, 53.3 – 53.8 Mb. [deletion marker 2]

Figure 31 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number loss (deletion) of Chr 4, 105.8 – 107.2 Mb. [deletion marker 3]

Figure 32 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number loss (deletion) of Chr 16, 45.8 – 46.3 Mb. [deletion marker 4]

Figure 33 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number loss (deletion) of Chr 5, 50.7 – 52.0 Mb. [deletion marker 5]

Figure 34 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number loss (deletion) of Chr 5, 94.2 – 96.1 Mb. [deletion marker 6]

Figure 35 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 102 patient cohort with NSCLC stage Ia-IIb, classified by presence or absence of a copy number loss (deletion) of Chr 5, 94.2 – 96.1 Mb. [deletion marker 6]

Figure 36 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number loss (deletion) of Chr 9, 36.1 – 37.0 Mb. [deletion marker 7]

Figure 37 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 102 patient cohort with NSCLC stage Ia-IIb, classified by presence or absence of a copy number loss (deletion) of Chr 5, 94.2 – 96.1 Mb. [deletion marker 8]

Figure 38 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 97 patient cohort with NSCLC stage Ia-IIb, classified by presence or absence of a copy number loss (deletion) of Chr14, 51.1 – 52.8 Mb. [deletion marker 9]

Figure 39 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 97 patient cohort with NSCLC stage Ia-IIb, classified by presence or absence of a copy number loss (deletion) of Chr 14, 61.5 – 68.6 Mb. [deletion marker 10]

Figure 40 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 97 patient cohort with NSCLC stage Ia-IIb, classified by presence or absence of a copy number loss (deletion) of Chr 9, 28.1 Mb. [deletion marker 11]

Figure 41 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number loss (deletion) of Chr 4, 43.7 – 44.2 Mb. [deletion marker 12]

Figure 42 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number loss (deletion) of Chr 5, 60.8 – 62.9 Mb. [deletion marker 13]

Figure 43 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 102 patient cohort with NSCLC stage Ia-IIb, classified by presence or absence of a copy number loss (deletion) of Chr 5, 60.8 – 62.9 Mb. [deletion marker 13]

Figure 44 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 97 patient cohort with NSCLC stage Ia-IIb, classified by presence or absence of a copy number loss (deletion) of Chr 5, 60.8 – 62.9 Mb. [deletion marker 13]

Figure 45 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number loss (deletion) of Chr 3, 120.0 – 121.1 Mb. [deletion marker 14]

Figure 46 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number loss (deletion) of Chr 4, 46.2 – 48.0 Mb. [deletion marker 15]

Figure 47 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 97 patient cohort with NSCLC stage Ia-IIb, classified by presence or absence of a copy number loss (deletion) of Chr 14, 38.9 – 40.0 Mb. [deletion marker 16]

Figure 48 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number loss (deletion) of Chr 4, 44.2 – 44.6 Mb. [deletion marker 17]

Figure 49 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number loss (deletion) of Chr 2, 213.7 – 214.3 Mb. [deletion marker 18]

Figure 50 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 97 patient cohort with NSCLC stage Ia-IIb, classified by presence or absence of a copy number loss (deletion) of Chr14, 43.9 – 46.6 Mb. [deletion marker 19]

Figure 51 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number loss (deletion) of Chr 14, 27.6 – 28.6 Mb. [deletion marker 20]

Figure 52 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number loss (deletion) of Chr 14, 27.6 – 28.6 Mb. [deletion marker 20]

Figure 53 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number loss (deletion) of Chr 3, 98.0 – 98.3 Mb. [deletion marker 21]

Figure 54 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number loss (deletion) of Chr 3, 98.0 – 98.3 Mb. [deletion marker 21]

Figure 55 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 97 patient cohort with NSCLC stage Ia-IIb, classified by presence or absence of a copy number loss (deletion) of Chr14, 55.2 – 60.0 Mb. [deletion marker 22]

5 **Figure 56** is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 97 patient cohort with NSCLC stage Ia-IIb, classified by presence or absence of a copy number loss (deletion) of Chr14, 48.7 – 51.1 Mb. [deletion marker 23]

Figure 57 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 97 patient cohort with NSCLC stage Ia-IIb, classified by presence or absence of a copy number loss (deletion) of Chr 4, 81.4 – 83.2 Mb. [deletion marker 24]

10 **Figure 58** is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number loss (deletion) of Chr 10, 51.9 – 54.2 Mb. [deletion marker 25]

15 **Figure 59** is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number loss (deletion) of Chr 5, 55.2 – 58.6 Mb. [deletion marker 26]

Figure 60 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 102 patient cohort with NSCLC stage Ia-IIb, classified by presence or absence of a copy number loss (deletion) of Chr 5, 67.8 – 68.5 Mb. [deletion marker 27].

20 **Figure 61** is a plot indicating average copy number, comparing the average copy number pattern obtained using a training (100K array) data set and a validating (SNP 6.0 array) data set.

Figure 62 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in ARSK.

25 **Figure 63** is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in ARSK.

Figure 64 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in C5orf27 .

Figure 65 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in C5orf27.

Figure 66 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in CAST.

Figure 67 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in CAST.

Figure 68 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in ELL2.

Figure 69 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in FAM81B.

Figure 70 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in FAM81B.

Figure 71 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in GLRX.

Figure 72 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in GLRX.

Figure 73 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in GPR150.

Figure 74 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in GPR150.

Figure 75 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in MCTP1.

Figure 76 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in MCTP1.

Figure 77 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in PCSK1.

Figure 78 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in PCSK1.

Figure 79 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in RFESD.

Figure 80 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in RFESD.

Figure 81 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in RHOBTB3.

Figure 82 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in RHOBTB3.

Figure 83 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in SPATA9.

Figure 84 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in SPATA9.

Figure 85 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in TTC37.

Figure 86 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in TTC37.

Figure 87 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in CCNB1.

Figure 88 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in SLC30A5.

Figure 89 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in MYO15B.

Figure 90 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in SLC16A5.

Figure 91 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in DKFZp761E198.

Figure 92 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in DKFZp761E198.

Figure 93 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in EHBP1L1.

Figure 94 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in EHBP1L1.

Figure 95 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in FAM89B.

Figure 96 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in FAM89B.

Figure 97 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in KAT5.

Figure 98 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in KAT5.

Figure 99 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in KCNK7.

Figure 100 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in KCNK7.

Figure 101 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in LTBP3.

Figure 102 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in LTBP3.

Figure 103 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in MALAT1.

Figure 104 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in MALAT1.

Figure 105 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in MAP3K11.

Figure 106 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in MAP3K11.

Figure 107 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in PACS1.

Figure 108 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in PACS1.

Figure 109 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in PCNXL3.

Figure 110 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in PCNXL3.

Figure 111 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in RELA.

Figure 112 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in RELA.

Figure 113 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in RNASEH2C.

Figure 114 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in RNASEH2C.

Figure 115 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in SCYL1.

Figure 116 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in SCYL1.

Figure 117 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in SIPA1.

Figure 118 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in SIPA1.

Figure 119 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in SSSCA1.

Figure 120 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in SSSCA1.

Figure 121 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in BAD.

Figure 122 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in C11orf20.

Figure 123 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in BAD.

Figure 124 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in C11orf20.

Figure 125 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in DNAJC4.

Figure 126 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in DNAJC4.

Figure 127 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in ESRRA.

Figure 128 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in ESRRA.

Figure 129 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in FADS2.

Figure 130 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in FADS3.

Figure 131 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in FKBP2.

Figure 132 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in FKBP2.

Figure 133 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in FLRT1.

Figure 134 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in GPR137.

Figure 135 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in HSPC152.

Figure 136 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in HSPC152.

Figure 137 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in KCNK4.

Figure 138 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in KCNK4.

Figure 139 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in NUDT22.

Figure 140 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in NUDT22.

Figure 141 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in PLCB3.

Figure 142 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in PLCB3.

Figure 143 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in PPP1R14B.

Figure 144 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in PPP1R14B.

Figure 145 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in PRDX5.

Figure 146 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in PRDX5.

Figure 147 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in RAB3IL1.

Figure 148 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in TRPT1.

Figure 149 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in TRPT1.

Figure 150 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in VEGFB.

Figure 151 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in VEGFB.

Figure 152 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in AKAP1.

Figure 153 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in ANKFN1.

Figure 154 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in C17orf67.

Figure 155 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in COIL.

Figure 156 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in DGKE.

Figure 157 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in MSI2.

Figure 158 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in MTVR2.

Figure 159 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in NOG.

Figure 160 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in RNF126P1.

Figure 161 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in SCPEP1.

Figure 162 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in TRIM25.

DETAILED DESCRIPTION OF THE INVENTION

Previously described expression-based markers of poor outcome in cancer cannot be measured with FISH, a well-established clinical diagnostic tool. Until now, no gene amplifications/deletions have been identified that can predict disease outcome. The inventors have discovered copy number changes in certain chromosomal sequences in certain cancer

patients. Moreover, the inventors have determined that the copy number changes are statistically significantly associated with shorter overall survival or reduced time to recurrence in stage I—II NSCLC.

Accordingly, the present disclosure provides methods of determining prognosis of early stage non-small cell lung cancer (NSCLC) in a human by assessment of the copy number of chromosomal DNA at any one of forty-seven markers. For each of these markers, a change in copy number is associated with a poorer prognosis in cancer. Copy number change was either a copy number gain of one or more, or a copy number loss. Poorer prognosis was assessed in patients having either a copy number gain or a copy number loss when compared to the normal baseline complement of two copies. Poorer prognosis was determined using measures of Overall Survival and Time to Recurrence. The present disclosure is particularly beneficial for providing improved prognostic information for early stage NSCLC patients and enables improved therapy selection for those early-stage NSCLC patients at higher risk of cancer recurrence.

The present disclosure includes methods for determining prognosis of NSCLC patients classified as early-stage cancers, in particular those classified as Stage IA, Stage IB, Stage IIA or Stage IIB (Stage IIA and IIB are collectively referred to as Stage II) using the widely used TNM staging system. Alternate NSCLC staging systems based upon other diagnostic classifications can be used to identify the patients whose tissue sample may be assayed by the methods disclosed herein. As used herein, an early stage NSCLC refers to a NSCLC tumor that has not spread to more than one lymph node, nor metastasized to any other organ. Early-stage NSCLC patients are almost always treated by surgical resection seeking complete tumor removal, yet a significant risk of recurrence exists for these early stage patients even where the tumor is believed to be completely resected. Current diagnostic modalities do not permit accurate prediction of which of these early stage cancers are high risk for recurrence and thus should be treated post-resection with adjuvant chemotherapy or before the resection using neoadjuvant chemotherapy. The present disclosure provides prognostic identification of those early stage patients at higher risk by determining the gene copy number in the patient sample.

Thus in one aspect, the methods encompass a method of predicting disease outcome in a patient being treated for lung cancer. A test sample, which is a biological sample from the patient, is provided, and a copy number for a selected cancer outcome marker in the test sample

is determined. The copy number from the test sample is compared against a baseline copy number of two, thereby determining the presence or absence of a copy number change for the cancer outcome marker. Based on the presence or absence of a copy number change for the cancer outcome marker in the test sample, the patient is identified as having an increased risk of a poor disease outcome when compared to a baseline measure of disease outcome in patients having no copy number change for the cancer outcome marker. The presence of a copy number change for the cancer outcome marker, i.e., a copy number of greater than 2 due to amplification, or less than 2 due to deletion, is predictive of poor disease outcome. The poor disease outcome is at least one of a decreased overall survival time when compared to an overall survival time of patients having no copy number change in the cancer outcome marker, and a shorter time to recurrence when compared to the time to recurrence for patients having no copy number change in the cancer outcome marker. The methods also encompass a method of predicting disease outcome in a patient being treated for lung cancer, in which based on the presence or absence of a copy number change in the cancer outcome marker, a determination is made as to whether the patient has a higher risk of a decreased overall survival time or a shorter time to recurrence when compared to an overall survival time of patients having no copy number change in the cancer outcome marker.

In any of the methods, the cancer outcome marker can be a region of chromosomal DNA, the amplification of which produces a copy number gain of the cancer outcome marker, wherein the copy number gain is associated with a poor disease outcome. Such cancer outcome markers include Chr 19, 34.7 Mb—35.6 Mb; Chr 19, 38.9—40.7 Mb; Chr 17, 69.2—71.3 Mb; Chr 6, 70.8—71.1 Mb; Chr 12, 93.7 kb—1.9Mb; Chr 11, 64.3—64.8 Mb; Chr 19, 57.0—62.2 Mb; Chr 6, 39.1—39.9 Mb; Chr 11, 64.8—65.7 Mb; Chr 11, 61.4—64.3 Mb; Chr 17, 51.5—53.2 Mb; Chr 17, 43.5—44.9 Mb; Chr 2, 147.6—151.1 Mb; Chr 6, 123.7—135.6 Mb; Chr 8, 6.9—8.8 Mb; Chr 2, 159.9—161.4 Mb; Chr 2, 200.9—204.2 Mb; Chr 6, 36.3—36.7 Mb; Chr 2, 205.9—208.1 Mb; and Chr 1, 109.5—111.1 Mb. Alternatively, the cancer outcome marker can be a region of chromosomal DNA, the deletion of which produces a copy number loss of the cancer outcome marker, wherein the copy number loss is associated with a poor disease outcome. Such cancer outcome markers include Chr 5, 62.9 – 67.8 Mb; Chr 5, 53.3 – 53.8 Mb; Chr 4, 105.8 – 107.2 Mb; Chr 16, 45.8 – 46.3 Mb; Chr 5, 50.7 – 52.0 Mb; Chr 5, 94.2 – 96.1 Mb; Chr 9, 36.1 – 37.0

Mb; Chr 5, 94.2 – 96.1 Mb; Chr14, 51.1 – 52.8 Mb; Chr 14, 61.5 – 68.6 Mb; Chr 9, 28.1 Mb; Chr 4, 43.7 – 44.2 Mb; Chr 5, 60.8 – 62.9 Mb; Chr 3, 120.0 – 121.1 Mb; Chr 4, 46.2 – 48.0 Mb; Chr 14, 38.9 – 40.0 Mb; Chr 4, 44.2 – 44.6 Mb; Chr 2, 213.7 – 214.3 Mb; Chr14, 43.9 – 46.6 Mb; Chr 14, 27.6 – 28.6 Mb; Chr 3, 98.0 – 98.3 Mb; Chr14, 55.2 – 60.0 Mb; Chr14, 48.7 – 51.1 Mb; Chr 4, 81.4 – 83.2 Mb; Chr 10, 51.9 – 54.2 Mb; Chr 5, 55.2 – 58.6 Mb; and Chr 5, 67.8 – 68.5 Mb.

The methods can also be applied to the problem of selecting a treatment for a patient suffering from lung cancer. For example, the method can include determining a chemotherapy treatment regimen based on the presence or absence of a copy number change of a cancer outcome marker in the sample from the patient. The step of determining a treatment regimen based on the comparison in step c) includes for example selecting a chemotherapy agent and determining a frequency of chemotherapy treatment when a copy number change is present for the cancer outcome marker. For example, an unfavorable copy number change of a cancer outcome marker in an individual patient, which may be a copy number gain or loss depending on the marker as described herein, indicates a lung cancer that is resistant to treatment. In such cases a treatment provider may wish to consider a more aggressive than usual chemotherapy treatment regimen, including use of a stronger chemotherapy agent or more frequent chemotherapy treatment, or both.

Accordingly, the methods can also be used to classify a patient as having a lung cancer that is resistant to treatment. For example, given a determination that a copy change is present in the sample from the patient, the patient is classified as having a lung cancer that is resistant to further treatment. The patient may be currently undergoing treatment with chemotherapy, radiation, surgery or any combination thereof, or may be currently under consideration for any one of chemotherapy, radiation, surgery treatment or any combination thereof.

The determining step (b) can be performed, for example, using *in situ* hybridization and, more preferably, fluorescent *in situ* hybridization (FISH) with fluorescently labeled nucleic acid probes or fluorescently labeled probes comprising nucleic acid analogs. Preferably at least two nucleic acid probes are used. A peptide nucleic acid probe can be used. The determining step (b) can also be performed by polymerase chain reaction, a nucleic acid sequencing assay, or a nucleic acid microarray assay as known in the art.

The testing of early stage NSCLC is preferably done on an appropriate biological sample obtained from the patient, preferably by *in situ* hybridization. In general, *in situ* hybridization includes the steps of fixing a biological sample, hybridizing one or more chromosomal probes to target DNA contained within the fixed sample, washing to remove non-specifically bound probe, and detecting the hybridized probe. The *in situ* hybridization can also be carried out with the specimen cells from the biological sample in liquid suspension, followed by detection by flow cytometry. The method preferably uses a FISH assay with a two probe set comprising a probe specific to the marker region to evaluate chromosomal copy number abnormalities in a biological sample from a patient. Preferred FISH probes for use according to the present disclosure comprise a pair of probes specific to the cancer outcome marker nucleotide sequence, which may include any portion of the nucleotide sequence.

The identification of NSCLC prognosis can also be used with other prognostic *in vitro* diagnostic assays, such as evaluating the expression in the patient sample of suitable proteins known to be encoded in the marker region. Patients whose samples are found with expression of such proteins in conjunction with an abnormal chromosomal copy number pattern, that is associated with an unfavorable outcome (poor prognosis), may be eligible for more aggressive post-surgery treatment, such as chemotherapy.

Typically for a lung cancer patient the biological sample is a tissue sample such as a peripheral blood sample that contains circulating tumor cells, or a lung tumor tissue biopsy or resection. Other suitable tissue samples include for example a thin layer cytological sample, a fine needle aspirate sample, a lung wash sample, a pleural effusion sample, a fresh frozen tissue sample, a paraffin embedded tissue sample or an extract or processed sample produced from any of a peripheral blood sample. Preferably, the sample has been classified as an early stage cancer, for example, such as any of Stage IA, Stage IB, Stage IIA or Stage IIB, according to generally accepted staging practice, for example using pathological stages.

Probes constructed according to the polynucleotide sequences of the cancer outcome markers described herein can be used in various assay methods to provide various types of analysis. For example, such probes can be used in fluorescent *in situ* hybridization (FISH) technology to perform chromosomal analysis, including copy number profiling, and used to identify cancer-specific copy number changes in the cancer outcome markers. Probes also can

be labeled with radioisotopes, directly- or indirectly- detectable haptens, or fluorescent molecules, and utilized for *in situ* hybridization studies to evaluate copy number of the cancer outcome markers in tissue specimens or cells.

Probes bind selectively to a target polynucleotide sequence, which is at least a portion of the sequence of a cancer outcome marker as described herein, i.e., a chromosomal region that is amplified in certain individuals. The nucleotide sequences of the cancer outcome markers provided herein, or any portion thereof, may be used to produce probes which can be used in various assays for copy number profiling in test samples. The probes may be designed from conserved nucleotide regions of the cancer outcome marker of interest, or from non-conserved nucleotide regions of the cancer outcome marker of interest, or any portion thereof including genes contained therein and portions thereof. The design of such probes for optimization in assays is within the skill of the routineer. Generally, nucleic acid probes are developed from non-conserved or unique regions when maximum specificity is desired, and nucleic acid probes are developed from conserved regions when assaying for nucleotide regions that are closely related to, for example, different members of a multi-gene family or in related species like mouse and man.

The polymerase chain reaction (PCR) is a technique for amplifying a desired nucleic acid sequence (target) contained in a nucleic acid or mixture thereof. In PCR, a pair of primers is employed in excess to hybridize to the complementary strands of the target nucleic acid. The primers are each extended by a polymerase using the target nucleic acid as a template. The extension products become target sequences themselves, following dissociation from the original target strand. New primers then are hybridized and extended by a polymerase, and the cycle is repeated to geometrically increase the number of target sequence molecules. PCR is disclosed in U.S. Pat. Nos. 4,683,195 and 4,683,202, which are incorporated herein by reference.

The Ligase Chain Reaction (LCR) is an alternate method for nucleic acid amplification. In LCR, probe pairs are used which include two primary (first and second) and two secondary (third and fourth) probes, all of which are employed in molar excess to target. The first probe hybridizes to a first segment of the target strand, and the second probe hybridizes to a second segment of the target strand, the first and second segments being contiguous so that the primary probes abut one another in 5' phosphate-3' hydroxyl relationship, and so that a ligase can

covalently fuse or ligate the two probes into a fused product. In addition, a third (secondary) probe can hybridize to a portion of the first probe and a fourth (secondary) probe can hybridize to a portion of the second probe in a similar abutting fashion. Of course, if the target is initially double stranded, the secondary probes also will hybridize to the target complement in the first instance. Once the ligated strand of primary probes is separated from the target strand, it will hybridize with the third and fourth probes that can be ligated to form a complementary, secondary ligated product. It is important to realize that the ligated products are functionally equivalent to either the target or its complement. By repeated cycles of hybridization and ligation, amplification of the target sequence is achieved. This technique is described more completely in EP-A-320 308 to K. Backman published Jun. 16, 1989 and EP-A-439 182 to K. Backman et al., published Jul. 31, 1991, both of which are incorporated herein by reference.

For amplification of mRNAs, it is within the scope of the present disclosure to reverse transcribe mRNA into cDNA followed by polymerase chain reaction (RT-PCR); or, to use a single enzyme for both steps as described in U.S. Pat. No. 5,322,770, which is incorporated herein by reference; or reverse transcribe mRNA into cDNA followed by asymmetric gap ligase chain reaction (RT-AGLCR) as described by R. L. Marshall et al., PCR Methods and Applications 4:80-84 (1994), which also is incorporated herein by reference.

Chromosomal Probes. Suitable probes for *in situ* hybridization techniques fall into three broad groups: chromosome enumeration probes, which hybridize to a chromosomal region and indicate the presence or absence of a chromosome; chromosome arm probes, which hybridize to a chromosomal region and indicate the presence or absence of an arm of a chromosome; and locus specific probes, which hybridize to a specific locus on a chromosome and detect the presence or absence of a specific locus. Chromosomal probes and combinations thereof are chosen for sensitivity and/or specificity when used in the methods. Probe sets can include any number of probes, e.g., at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, or 20 probes. The selection of individual probes and probe sets can be performed according to routine in the art, for example as described in US 20060063194, the entire disclosure of which is herein incorporated by reference. Such selection methods make use of discriminate and/or combinatorial analysis to select probes and probes sets that are useful for copy number profiling of the cancer outcome markers.

Suitable probes for use in *in situ* hybridization methods according to the present disclosure for the detection of abnormal copy number pattern (aneusomy or polysomy) are a combination of a chromosome enumeration probe and a chromosome locus-specific probe hybridizable to a portion of the marker sequence, with each probe labeled to be distinguishable from the other. As is well known in the art, a chromosome enumeration probe can hybridize to a repetitive sequence, located either near or removed from a centromere, or can hybridize to a unique sequence located at any position on a chromosome. For example, a chromosome enumeration probe can hybridize with repetitive DNA associated with the centromere of a chromosome. Centromeres of primate chromosomes contain a complex family of long tandem repeats of DNA comprised of a monomer repeat length of about 171 base pairs, that are referred to as alpha-satellite DNA. A non-limiting example of a specific chromosome enumeration probe is the SpectrumGreen™ CEP® 10 probe (Abbott Molecular, Inc., Des Plaines, Illinois). For example, the chromosome 19 enumeration probe is used with a locus specific probe for detecting copy number abnormalities at Chr19, 34.7 Mb—35.6 Mb, for example to determine the status of deletion and/ or polysomy of loci contained therein. A locus specific probe hybridizes to a specific, non-repetitive locus on a chromosome. A suitable probe includes for example at least a portion of the any gene for which the marker sequence includes the nucleotide sequence encoding the gene. Locus specific probes are available commercially from Abbott Molecular Inc. in a probe set, for example mixed with the Vysis CEP® 10 SpectrumGreen probe.

Probes that hybridize with centromeric DNA are available commercially from Abbott Molecular Inc. (Des Plaines, IL) and Molecular Probes, Inc. (Eugene, OR). Alternatively, probes can be made non-commercially using well known techniques. Sources of DNA for use in constructing DNA probes include genomic DNA, cloned DNA sequences such as bacterial artificial chromosomes (BAC), somatic cell hybrids that contain one or a part of a human chromosome along with the normal chromosome complement of the host, and chromosomes purified by flow cytometry or microdissection. The region of interest can be isolated through cloning or by site-specific amplification via the polymerase chain reaction (PCR). See, for example, Nath, et al., *Biotechnic Histochem*, 1998, 73 (1): 6-22; Wheelless, et al., *Cytometry*, 1994, 17:319-327; and U.S. Pat. No. 5,491,224. The starting human DNA used to manufacture useful locus specific probes can be obtained by obtaining a nucleic acid sequence for the locus

from the Human Genome database, such as that maintained by the University of California Santa Cruz, and then using that sequence to screen in silico a BAC human DNA library, such as that maintained by the Roswell Park Cancer Center or Invitrogen, to identify useful BAC clones. Synthesized oligomeric DNA probes or probes made from nucleic acid analogs, such as peptide
5 nucleic acid (PNA) probes, can also be used.

The size of the chromosomal region detected by the probes according to the present disclosure can vary in size, for example, from a short couple hundred base pair probe sequence to a large segment of 900,000 bases. Locus-specific probes that are directly labeled are preferably at least 100,000 bases in complexity, and use unlabeled blocking nucleic acid, as disclosed in
10 U.S. Pat. No. 5,756,696, herein incorporated by reference, to avoid non-specific binding of the probe. It is also possible to use unlabeled, synthesized oligomeric nucleic acid or unlabeled nucleic acid analogs, such as a peptide nucleic acid, as the blocking nucleic acid.

The chromosomal probes can contain any detection moiety that facilitates the detection of the probe when hybridized to a chromosome. Effective detection moieties include both direct and
15 indirect labels as described herein. Examples of detectable labels include fluorophores (i.e., organic molecules that fluoresce after absorbing light), radioactive isotopes (e.g., ^{32}P , and ^3H) and chromophores (e.g., enzymatic markers that produce a visually detectable marker). Fluorophores are preferred and can be directly labeled following covalent attachment to a nucleotide by incorporating the labeled nucleotide into the probe with standard techniques such
20 as nick translation, random priming, and PCR labeling. Alternatively, deoxycytidine nucleotides within the probe can be transaminated with a linker. The fluorophore can then be covalently attached to the transaminated deoxycytidine nucleotides. See, e.g., U.S. Pat. No. 5,491,224 to Bittner, et al., which is incorporated herein by reference. Useful probe labeling techniques are described in *Molecular Cytogenetics: Protocols and Applications*, Y.-S. Fan, Ed., Chap. 2,
25 "Labeling Fluorescence *In Situ* Hybridization Probes for Genomic Targets", L. Morrison et. al., p. 21-40, Humana Press, © 2002, incorporated herein by reference.

Examples of fluorophores that can be used in the methods described herein are: 7-amino-4-methylcoumarin-3-acetic acid (AMCA), Texas Red™ (Molecular Probes, Inc., Eugene, OR); 5-(and-6)-carboxy-X-rhodamine, lissamine rhodamine B, 5-(and-6)-carboxyfluorescein;
30 fluorescein-5-isothiocyanate (FITC); 7-diethylaminocoumarin-3-carboxylic acid,

tetramethylrhodamine-5-(and-6)-isothiocyanate; 5-(and-6)-carboxytetramethylrhodamine; 7-hydroxycoumarin-3-carboxylic acid; 6-[fluorescein 5-(and-6)-carboxamido]hexanoic acid; N-(4,4-difluoro-5,7-dimethyl-4-bora-3a, 4a diaza-3-indacenepropionic acid; eosin-5-isothiocyanate; erythrosine-5-isothiocyanate; 5-(and-6)-carboxyrhodamine 6G; and Cascade™ blue acetylazide
5 (Molecular Probes, Inc., Eugene, OR). In the preferred probe set, fluorophores of different colors are used such that each chromosomal probe in the set can be distinctly visualized.

After hybridization, the probes can be viewed with a fluorescence microscope and an appropriate filter for each fluorophore, or by using dual or triple band-pass filter sets to observe multiple fluorophores. See, e.g., U.S. Pat. No. 5,776,688 to Bittner, et al., which is incorporated
10 herein by reference. Any suitable microscopic imaging method can be used to visualize the hybridized probes, including automated digital imaging systems, such as those available from MetaSystems or Applied Imaging. Alternatively, techniques such as flow cytometry can be used to examine the hybridization pattern of the chromosomal probes.

Probes can also be labeled indirectly, e.g., with biotin or digoxigenin by means well
15 known in the art. However, secondary detection molecules or further processing are then required to visualize the labeled probes. For example, a probe labeled with biotin can be detected by avidin (e.g. streptavidin) conjugated to a detectable marker, e.g., a fluorophore. Additionally, avidin can be conjugated to an enzymatic marker such as alkaline phosphatase or horseradish peroxidase. Such enzymatic markers can be detected in standard colorimetric reactions using a
20 substrate for the enzyme. Substrates for alkaline phosphatase include 5-bromo-4-chloro-3-indolylphosphate and nitro blue tetrazolium. Diaminobenzidine can be used as a substrate for horseradish peroxidase.

The probes and probe sets useful with the methods can be packaged with other reagents into kits to be used in carrying out the methods as described herein.

25 ***Preferred Probe Set.*** An exemplary probe composition comprises a mixture of directly labeled DNA FISH probes. For example, such a probe set would include a Vysis SpectrumOrange probe and a Vysis SpectrumGreen probe. Suitable probe sets are available commercially premixed in a suitable hybridization buffer.

Preparation of Samples. A biological sample is a sample that contains cells or cellular
30 material, including cell-containing extracts from a patient sample. For example, lung samples

are typically cells or cellular material derived from pulmonary structures, including but not limited to lung parenchyma, bronchioles, bronchial, bronchi, and trachea. Non-limiting examples of biological samples useful for the detection of lung cancer include bronchial specimens, resected lung tissue, lung biopsies, and sputum samples. Examples of bronchial specimens
5 include bronchial secretions, washings, lavage, aspirations, and brushings. Lung biopsies can be obtained by methods including surgery, bronchoscopy, fine needle aspiration (FNA), and transthoracic needle biopsy. In one example, touch preparations can be made from lung biopsies. The inventive assays can also be performed on a circulating tumor cell sample derived from a blood sample from an early stage NSCLC patient. A circulating tumor cell sample can be
10 prepared using the immunomagnetic separation technology available from Immunicon.

Tissues can be fixed with a fixative such as formaldehyde and then embedded in paraffin. Sections are then cut using a microtome and are applied to a microscope slide. Cytology specimens can be prepared from cellular suspensions derived from FNA, bronchial washings, bronchial lavage, or sputum, or disseminated tissue cells. Cytology specimens can be prepared
15 by fixation of cells in ethanol or methanol:acetic acid combined with cytocentrifugation, thin layer deposition methods (e.g. ThinPrep, Cytoc Corp.), smears, or pipetting onto microscope slides. In addition, biological samples can include effusions, e.g., pleural effusions, pericardial effusions, or peritoneal effusions.

Hybridization Methods. Any suitable *in situ* hybridization method can be used. Prior to
20 *in situ* hybridization, chromosomal probes and chromosomal DNA contained within the cell each are denatured. If the chromosomal probes are prepared as a single-stranded nucleic acid, then denaturation of the probe is not required. Denaturation typically is performed by incubating in the presence of high pH, heat (e.g., temperatures from about 70°C to about 95°C), organic solvents such as formamide and tetraalkylammonium halides, or combinations thereof. For
25 example, chromosomal DNA can be denatured by a combination of temperatures above 70°C (e.g., about 73°C) and a denaturation buffer containing 70% formamide and 2X SSC (0.3M sodium chloride and 0.03 M sodium citrate). Denaturation conditions typically are established such that cell morphology is preserved. For example, chromosomal probes can be denatured by heat, e.g., by heating the probes to about 73°C for about five minutes.

After removal of denaturing chemicals or conditions, probes are annealed to the chromosomal DNA under hybridizing conditions. "Hybridizing conditions" are conditions that facilitate annealing between a probe and target chromosomal DNA. Hybridization conditions vary, depending on the concentrations, base compositions, complexities, and lengths of the probes, as well as salt concentrations, temperatures, and length of incubation. For example, *in situ* hybridizations are typically performed in hybridization buffer containing 1-2.X.SSC, 50-55% formamide, a hybridization acceleratant (e.g. 10% dextran sulfate), and unlabeled blocking DNA to suppress non-specific hybridization. In general, hybridization conditions, as described above, include temperatures of about 25°C to about 55°C, and incubation lengths of about 0.5 hours to about 96 hours. More particularly, hybridization can be performed at about 32°C to about 45°C for about 2 to about 16 hours.

Non-specific binding of chromosomal probes to DNA outside of the target region can be removed by a series of washes with a salt solution. Temperature and concentration of salt in each wash depend on the desired stringency. For example, for high stringency conditions, washes can be carried out at about 65°C to about 80°C, using 0.2.X to about 2.X.SSC, and about 0.1% to about 1% of a non-ionic detergent such as Nonidet P-40 (NP40). Stringency can be lowered by decreasing the temperature of the washes or by increasing the concentration of salt in the washes.

The hybridization of the probes to the tissue sample can be performed manually, or with the assistance of instruments, such as the ThermoBrite hybridization oven, the VP 2000 Processor, or the XMatrix™ processing instrument (all available commercially from Abbott Molecular, Inc.).

Pre-Selection of Cells. Cell samples can be evaluated preliminarily by a variety of methods and using a variety of criteria. The probes and methods described herein are not limited to usage with a particular screening methodology. One example is the "scanning method" wherein the observer scans hundreds to thousands of cells for cytologic abnormalities, e.g., as viewed with a DAPI filter. The number of cells assessed will depend on the cellularity of the specimen, which varies from patient to patient. Cytologic abnormalities commonly but not invariably associated with dysplastic and neoplastic cells include nuclear enlargement, nuclear irregularity, and abnormal DAPI staining (frequently mottled and lighter in color). In the scanning step, the observer preferably focuses the evaluation of the cells for chromosomal

abnormalities (as demonstrated by FISH) to those cells that also exhibit cytological abnormalities. In addition, a proportion of the cells that do not have obvious cytologic abnormalities can be evaluated since chromosomal abnormalities also occur in the absence of cytologic abnormalities. This scanning method is described in further detail in U.S. Pat. No. 6,174,681 to Halling, et al., which is incorporated herein by reference. Lung cancer cells can be selected for evaluation using the method described in US Patent Pub. 2003/0087248 A1 by Morrison, et al., which is incorporated herein by reference.

Regions of the specimen may also be selected for evaluation using conventional stains, such as stains containing hematoxylin and eosin. For example, a pathologist can stain a section of a paraffin-embedded specimen with a hematoxylin/eosin stain, identify a region as probably cancerous by tissue morphology and staining pattern, and outline that region with a felt tip ink pen or glass scribe. The marked region is then transferred to the corresponding location on a serial section of the paraffin-embedded specimen with a glass scribe, and FISH is performed on that slide. Cells within the scribed region are then evaluated for FISH signals.

Detection of Classification Patterns of Chromosomal Abnormality. Abnormal cells are characterized by the presence of one or more patterns of chromosomal copy number abnormalities. The presence of a copy number abnormality pattern in a cell in the patient sample is assessed by examining the hybridization pattern of the chromosomal probe (e.g., the number of signals for each probe) in the cell, and recording the number of signals. Aneusomy is typically intended to mean abnormal copy number, either of the whole chromosome or a locus on a chromosome. Abnormal copy number includes both monosomy (one copy) and nullsomy (zero copies) of the autosomes, also referred to as a deletion, and greater than 2 copies, which for a particular chromosomal locus is sometimes referred to as gene amplification (alternatively, amplification is reserved for the situation in which the gene copy number exceeds the copy number of the chromosome in which it is contained). However, sectioning of paraffin-embedded specimens (typically 4-6 μm) may result in truncation of cell nuclei such that the number of FISH signals per cell for some cells will be somewhat lower than the actual number of copies in an intact nucleus. In the methods as described herein, the absolute number of particular FISH probe hybridization signals for each probe is determined and then used in various ratio comparisons.

Test samples can comprise any number of cells that is sufficient for a clinical diagnosis, and in a preferred paraffin-embedded tissue sample, the hybridization pattern is typically assessed in about 20 to about 200 cells. It is preferred to assess the hybridization pattern in about 40 to about 120 cells per sample.

5 The present disclosure thus describes new findings (DNA copy number changes of a cancer outcome marker) that may solve recognized treatment dilemmas by providing methods of determining which patients with early stage disease are at highest risk of disease recurrence or metastasis and who should be definitively treated with drug (or alternatives like radiation) therapies to maximize their chances of long term survival. In turn, the present disclosure
10 describes findings enabling of a specific DNA test that detects a chromosomal copy number change in any one of a number of cancer outcome markers. Consequently when a test for a copy number change is negative, or normal copy number is present, this identifies patients who have low or no risk of disease recurrence or metastasis who do not need follow-up therapy after resection of their initial tumors. These testing strategies can significantly impact both the
15 morbidity and mortality in patients with early stage NSCLC. The methods used herein also suggest application to other cancers to similarly detect DNA copy number gains of the cancer outcome markers that significantly associate with time to disease progression and/or overall survival. As such, the methods described herein have the potential to solve the question of which early-stage NSCLC patients should receive drug therapy after surgery and can broadly impact
20 cancer treatment decisions and patient outcomes.

Kits. In another aspect, the present disclosure provides a kit comprising reagents for determining the presence or absence of a copy number change for the cancer outcome marker. The kit includes instructions for using the reagents to perform the test. The reagents to determine the presence or absence of a copy number change for the cancer outcome marker can
25 include for example detectably-labeled polynucleotides that hybridize to at least a portion of the cancer outcome marker. The polynucleotides (probes) can be selected for sequences that hybridize, for example, to any portion of the cancer outcome marker. The cancer outcome marker can be a region of chromosomal DNA, the amplification of which produces a copy number gain of the cancer outcome marker, wherein the copy number gain is associated with a
30 poor disease outcome. Such cancer outcome markers can be selected from the group consisting

of Chr 19, 34.7 Mb—35.6 Mb; Chr 19, 38.9—40.7 Mb; Chr 17, 69.2—71.3 Mb; Chr 6, 70.8—71.1 Mb; Chr 12, 93.7 kb—1.9Mb; Chr 11, 64.3—64.8 Mb; Chr 19, 57.0—62.2 Mb; Chr 6, 39.1—39.9 Mb; Chr 11, 64.8—65.7 Mb; Chr 11, 61.4—64.3 Mb; Chr 17, 51.5—53.2 Mb; Chr 17, 43.5—44.9 Mb; Chr 2, 147.6—151.1 Mb; Chr 6, 123.7—135.6 Mb; Chr 8, 6.9—8.8 Mb; Chr 2, 159.9—161.4 Mb; Chr 2, 200.9—204.2 Mb; Chr 6, 36.3—36.7 Mb; Chr 2, 205.9—208.1 Mb; and Chr 1, 109.5—111.1 Mb. The cancer outcome marker can instead be a region of chromosomal DNA, the deletion of which produces a copy number loss of the cancer outcome marker, wherein the copy number loss is associated with a poor disease outcome. Such cancer outcome markers can be selected from the group consisting of Chr 5, 62.9 – 67.8 Mb; Chr 5, 53.3 – 53.8 Mb; Chr 4, 105.8 – 107.2 Mb; Chr 16, 45.8 – 46.3 Mb; Chr 5, 50.7 – 52.0 Mb; Chr 5, 94.2 – 96.1 Mb; Chr 9, 36.1 – 37.0 Mb; Chr 5, 94.2 – 96.1 Mb; Chr14, 51.1 – 52.8 Mb; Chr 14, 61.5 – 68.6 Mb; Chr 9, 28.1 Mb; Chr 4, 43.7 – 44.2 Mb; Chr 5, 60.8 – 62.9 Mb; Chr 3, 120.0 – 121.1 Mb; Chr 4, 46.2 – 48.0 Mb; Chr 14, 38.9 – 40.0 Mb; Chr 4, 44.2 – 44.6 Mb; Chr 2, 213.7 – 214.3 Mb; Chr14, 43.9 – 46.6 Mb; Chr 14, 27.6 – 28.6 Mb; Chr 3, 98.0 – 98.3 Mb; Chr14, 55.2 – 60.0 Mb; Chr14, 48.7 – 51.1 Mb; Chr 4, 81.4 – 83.2 Mb; Chr 10, 51.9 – 54.2 Mb; Chr 5, 55.2 – 58.6 Mb; and Chr 5, 67.8 – 68.5 Mb. Accordingly, the probes can be selected for sequences that hybridize, for example, to any portion of any of the listed markers, including any portion of any of the genes encoded therein as set forth elsewhere herein.

Details of the present disclosure are further described in the following example, which is not intended to limit the scope of the invention as claimed. One of skill in the art will recognize that variations and modifications of the invention may be apparent upon reviewing the instant specification. It is therefore an object to provide for such modifications and variations of the embodiments described herein, without departing from the scope or the spirit of the invention.

EXAMPLES

Example 1: Analysis of NSCLC patient samples

Experimental Methods: *Specimens.* A total of 178 NSCLC clinically annotated samples were profiled for copy number alterations using high-density SNP genotyping microarrays (100K array set by Affymetrix). All samples were carefully dissected to maximize tumor/normal tissue ratio and verify histopathological type and stage. Only samples from patients with stage I

and II samples were analyzed. All of these were from patients treated with surgical resection without any follow-up or neoadjuvant chemotherapy. Clinical information collected for each patient included race, age, date of birth, sex, clinical stage, pathological stage, location, surgical procedure (SP) date, histology, differentiation, diagnosis date, node positivity, smoking status, chemotherapy status, radiation status, recurrence status, recurrence date, recurrence location, time to recurrence, date of last follow up, status at the last follow up, alive/dead, overall survival and cause of death. Time to Recurrence (TTR) and Overall Survival (OS) were chosen as the parameters of outcome. Other clinical parameters (node status, stage, etc) were considered as confounding variables. Times to recurrence of lung cancer and the overall survival times were obtained from the patient charts.

Tables 2 and 3 provide the figures for Overall Survival and Total Time to Recurrence, respectively, for the patient cohort studied.

TABLE 2:

OS

Stage	deaths	alive (censored)	total
1a	6	25	31
1b	6	40	46
2a	0	1	1
2b	7	17	24
total	19	83	102

TABLE 3:

TTR

Stage	recurred	recurrence free (censored)	total
1a	10	21	31
1b	9	34	43
2a	1	0	1
2b	9	13	22
total	29	68	97

5 **Copy number profiling.** Approximately 30 mg tissue from each tumor were used to
extract high molecular weight, genomic DNA using the Qiagen DNAeasy kit (Qiagen, Valencia,
CA) following the instructions by the manufacturer. The quality of DNA was checked by
agarose gel electrophoresis. Two hundred and fifty nanograms of DNA were processed for
hybridization to each of the two microarrays comprising the Genechip Human Mapping 100K set
10 (Matsuzaki H, Dong S, Loi H, et al. Genotyping over 100,000 SNPs on a pair of oligonucleotide
arrays. Nat Methods 2004; 1:109-11) -arrays (Affymetrix, Inc., Santa Clara, CA), which covers
116,204 single-nucleotide polymorphism (SNP) loci in the human genome with a mean
intermarker distance of 23.6 kb. The microarrays were processed according to recommendations
of the manufacturer (www.affymetrix.com). Copy number was calculated by comparing the chip
15 signal to the average of 48 normal female samples. Samples with normal tissue contamination
were removed by QC.

Statistical Methods. Univariate analysis was used to test the following parameters as
potential confounding factors: Pathological stage, Clinical stage, Smoking status, Age, Sex,
Node status, Histology (adenocarcinoma vs squamous cell carcinoma). No significant effects
20 were detected. In survival analysis, interaction of clinical stage and marker regions was tested.
No copy number abnormalities had significant interaction with stage (FDR p value<0.05).

Results: Only patients with stage I-II disease were analyzed. **Figures 1-28** are each a Kaplan-Meier plot showing the difference in OS or TTR between patients with and without amplification of a selected marker as indicated (i.e., a copy number gain of at least one).

Figures 29-60 are each a Kaplan-Meier plot showing the difference in OS or TTR between

patients with and without deletion (i.e., a copy number loss of at least one) of a selected marker as indicated. In all Figures 1-60, the x-axis represents time in days, and the y-axis the probability of patient survival (for OS), or of patient being free of disease recurrence (for TTR).

Whenever a relevant event occurred (either death for OS, or disease recurrence for TTR), the curve drops. In all Figures 1-60, the upper line always shows data from patients with a normal

baseline complement of two. In **Figures 1-28**, data for patients with a copy number **gain** for the marker are shown in **red**, and data for patients with the **normal** baseline complement of two are shown in **green**. In **Figures 29-60**, data for patients with the normal baseline complement of

two are always shown in the upper line, in some figures shown in **green** (Figures 29, 30, 35, 37-40, 43, 44, 47, 50, 55-57 and 60), and in other figures shown in **red** (Figures 31-34, 36, 41, 42, 45, 46, 48, 49, 51-54, 58 and 59), while data for patients with a copy number **loss** for the marker are shown in certain figures in **blue** (Figures 29, 30, 35, 37-40, 43, 44, 47, 50, 55-57 and 60), and in other figures in **green** (Figures 31-34, 36, 41, 42, 45, 46, 48, 49, 51-54, 58 and 59). More specifically, the figures show results as follows:

Figure 1 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr19, 34.7 Mb—35.6 Mb. [marker 1]. FDR adjusted p-value = 0.0299. 17 samples: 3 copies, 3 samples: 4 copies, 7 samples: 5 or more copies.

Figure 2 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 19; 38.9—40.7 Mb. [marker 2]. FDR p-value = 0.0085. 17 samples: 3 copies; 3 samples: 4 copies; 4 samples: ≥ 5 copies.

Figure 3 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 17; 69.2—71.3 Mb. [marker 3]. FDR p-value= 0.0304. 16 samples: 3 copies; 5 samples: 4 copies.

Figure 4 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 71 patient cohort with NSCLC stage Ib-IIb, classified by presence or absence of a copy number gain in Chr 6, 70.8—71.1 Mb. [marker 4]. FDR p-value = 0.0116. 15 samples: 3 copies, 1 sample: 4 copies, 1 sample: ≥ 5 copies.

5 **Figure 5** is a Kaplan-Meier plot showing the overall survival (OS) in days for a 102 patient cohort with NSCLC stage Ia-IIb, classified by presence or absence of a copy number gain in Chr 6, 70.8—71.1 Mb. [marker 4]. FDR p-value 0.0110. 15 samples: 3 copies, 1 sample: 4 copies, 1 sample: ≥ 5 copies.

10 **Figure 6** is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 12, 93.7 kb—1.9Mb. [marker 5]. FDR p-value = 0.0493. 5 samples: 3 copies, 5 samples: 4 copies, 1 sample: ≥ 5 copies.

15 **Figure 7** is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 11, 64.3—64.8 Mb. [marker 6]. FDR p-value = 0.0413. 9 samples: 3 copies, 2 samples: ≥ 5 copies.

20 **Figure 8** is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 11, 64.3—64.8 Mb. [marker 6]. FDR p-value = 0.0040. 9 samples: 3 copies, 2 sample: ≥ 5 copies.

Figure 9 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 19, 57.0—62.2 Mb. [marker 7]. FDR p-value = 0.0091. 10 samples: 3 copies, 3 samples: 4 copies.

25 **Figure 10** is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 6, 39.1—39.9 Mb. [marker 8]. FDR p-value = 0.0356. 13 samples: 3 copies, 1 sample: 4 copies.

30 **Figure 11** is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain

in Chr 11, 64.8—65.7 Mb. [marker 9]. FDR p-value = 0.0484. 5 samples: 3 copies, 2 samples: ≥ 5 copies.

Figure 12 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 11, 64.8—65.7 Mb. [marker 9]. FDR p-value = 0.0004. 5 samples: 3 copies, 2 samples: ≥ 5 copies.

Figure 13 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 11, 61.4—64.3 Mb. [marker 10]. FDR p-value = 0.0004; 8 samples; 3 copies, 1 sample: 4 copy.

Figure 14 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 17, 51.5—53.2 Mb. [marker 11]. FDR p-value = 0.0054. 8 samples: 3 copies, 1 sample: 4 copies.

Figure 15 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 17, 43.5—44.9 Mb. [marker 12]. FDR p-value = 0.0269. 4 samples: 3 copies, 2 samples: 4 copies.

Figure 16 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 17, 43.5—44.9 Mb. [marker 12]. FDR p-value = 0.0040. 5 samples: 3 copies, 2 samples: 4 copies.

Figure 17 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 2, 147.6—151.1 Mb. [marker 13]. FDR p-value = 0.0210. 7 samples: 3 copies.

Figure 18 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 2, 147.6—151.1 Mb. [marker 13]. FDR p-value = 0.0233. 7 samples: 3 copies.

Figure 19 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIb, classified by presence or absence of a copy number gain in Chr 6, 123.7—135.6 Mb. [marker 14]. FDR p-value = 0.0377. 7 samples: 3 copies.

Figure 20 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 8, 6.9—8.8 Mb. [marker 15]. FDR p-value = 0.0166. 7 samples: 3 copy.

Figure 21 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 2, 159.9—161.4 Mb. [marker 16]. FDR p-value = 0.0013. 4 samples: 3 copies, 1 sample: 4 copies.

Figure 22 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 2, 159.9—161.4 Mb. [marker 16]. FDR p-value = 0.0001. 4 samples: 3 copy, 1 sample: 4 copies.

Figure 23 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 2, 200.9—204.2 Mb. [marker 17]. FDR p-value = 0.0398. 6 samples: 3 copies.

Figure 24 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 6, 36.3—36.7 Mb. [marker 18]. FDR p-value 0.0347. 6 samples: 3 copies.

Figure 25 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 2, 205.9—208.1 Mb. [marker 19]. FDR p-value = 0.04. 5 samples: 3 copies.

Figure 26 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 66 patient cohort with NSCLC stage Ib-IIb, classified by presence or absence of a copy number gain in Chr 2, 205.9—208.1 Mb. [marker 19]. FDR p-value = 0.0351. 6 samples: 3 copies.

Figure 27 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number gain in Chr 2, 205.9—208.1 Mb. [marker 19]. FDR p-value = 0.0075. 5 samples: 3 copies.

Figure 28 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 71 patient cohort with NSCLC stage Ib-IIb, classified by presence or absence of a copy number gain in Chr 1, 109.5—111.1 Mb. [marker 20]. FDR p-value = 0.0224. 5 samples: 3 copies.

Figure 29 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 102 patient cohort with NSCLC stage Ia-IIb, classified by presence or absence of a copy number loss (deletion) of Chr 5, 62.9 – 67.8 Mb. [deletion marker 1]. FDR p-value = 0.0282. Deleted in 10 samples.

Figure 30 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 102 patient cohort with NSCLC stage Ia-IIb, classified by presence or absence of a copy number loss (deletion) of Chr 5, 53.3 – 53.8 Mb. [deletion marker 2]. FDR p-value = 0.0409. Deleted in 12 samples.

Figure 31 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number loss (deletion) of Chr 4, 105.8 – 107.2 Mb. [deletion marker 3]. FDR p-value = 0.0469. Deleted in 21 samples.

Figure 32 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number loss (deletion) of Chr 16, 45.8 – 46.3 Mb. [deletion marker 4]. FDR p-value = 0.0039. Deleted in 11 samples.

Figure 33 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number loss (deletion) of Chr 5, 50.7 – 52.0 Mb. [deletion marker 5]. FDR p-value = 0.0000. Deleted in 10 samples.

Figure 34 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number loss (deletion) of Chr 5, 94.2 – 96.1 Mb. [deletion marker 6]. FDR p-value = 0.0202. Deleted in 7 samples.

Figure 35 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 102 patient cohort with NSCLC stage Ia-IIb, classified by presence or absence of a copy number loss

(deletion) of Chr 5, 94.2 – 96.1 Mb. [deletion marker 6]. FDR p-value = 0.0282. Deleted in 10 samples.

Figure 36 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number loss (deletion) of Chr 9, 36.1 – 37.0 Mb. [deletion marker 7]. FDR p-value = 0.0468. Deleted in 24 samples.

Figure 37 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 102 patient cohort with NSCLC stage Ia-IIb, classified by presence or absence of a copy number loss (deletion) of Chr 5, 94.2 – 96.1 Mb. [deletion marker 8]. FDR p-value = 0.0282. Deleted in 10 samples.

Figure 38 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 97 patient cohort with NSCLC stage Ia-IIb, classified by presence or absence of a copy number loss (deletion) of Chr14, 51.1 – 52.8 Mb. [deletion marker 9]. FDR p-value = 0.0008. Deleted in 9 samples.

Figure 39 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 97 patient cohort with NSCLC stage Ia-IIb, classified by presence or absence of a copy number loss (deletion) of Chr 14, 61.5 – 68.6 Mb. [deletion marker 10]. FDR p-value = 0.0034. Deleted in 9 samples.

Figure 40 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 97 patient cohort with NSCLC stage Ia-IIb, classified by presence or absence of a copy number loss (deletion) of Chr 9, 28.1 Mb. [deletion marker 11]. FDR p-value = 0.0270. Deleted in 9 samples.

Figure 41 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number loss (deletion) of Chr 4, 43.7 – 44.2 Mb. [deletion marker 12]. FDR p-value = 0.0053. Deleted in 8 samples.

Figure 42 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number loss (deletion) of Chr 5, 60.8 – 62.9 Mb. [deletion marker 13]. FDR p-value = 0.0121. Deleted in 7 samples.

Figure 43 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 102 patient cohort with NSCLC stage Ia-IIb, classified by presence or absence of a copy number loss (deletion) of Chr 5, 60.8 – 62.9 Mb. [deletion marker 13]. FDR p-value = 0.0425. Deleted in 8 samples.

5 **Figure 44** is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 97 patient cohort with NSCLC stage Ia-IIb, classified by presence or absence of a copy number loss (deletion) of Chr 5, 60.8 – 62.9 Mb. [deletion marker 13]. FDR p-value = 0.0320. Deleted in 8 samples.

10 **Figure 45** is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number loss (deletion) of Chr 3, 120.0 – 121.1 Mb. [deletion marker 14]. FDR p-value = 0.0228. Deleted in 8 samples.

15 **Figure 46** is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number loss (deletion) of Chr 4, 46.2 – 48.0 Mb. [deletion marker 15]. FDR p-value = 0.0341. Deleted in 8 samples.

20 **Figure 47** is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 97 patient cohort with NSCLC stage Ia-IIb, classified by presence or absence of a copy number loss (deletion) of Chr 14, 38.9 – 40.0 Mb. [deletion marker 16]. FDR p-value = 0.0451. Deleted in 8 samples.

Figure 48 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number loss (deletion) of Chr 4, 44.2 – 44.6 Mb. [deletion marker 17]. FDR p-value = 0.0053. Deleted in 7 samples.

25 **Figure 49** is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number loss (deletion) of Chr 2, 213.7 – 214.3 Mb. [deletion marker 18]. FDR p-value = 0.0286. Deleted in 7 samples.

30 **Figure 50** is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 97 patient cohort with NSCLC stage Ia-IIb, classified by presence or absence of a copy number loss

(deletion) of Chr14, 43.9 – 46.6 Mb. [deletion marker 19]. FDR p-value = 0.0009. Deleted in 6 samples.

Figure 51 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number loss (deletion) of Chr 14, 27.6 – 28.6 Mb. [deletion marker 20]. FDR p-value = 0.0021. Deleted in 6 samples.

Figure 52 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number loss (deletion) of Chr 14, 27.6 – 28.6 Mb. [deletion marker 20]. FDR p-value = 0.0101. Deleted in 5 samples.

Figure 53 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number loss (deletion) of Chr 3, 98.0 – 98.3 Mb. [deletion marker 21]. FDR p-value = 0.0316. Deleted in 6 samples.

Figure 54 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 74 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number loss (deletion) of Chr 3, 98.0 – 98.3 Mb. [deletion marker 21]. FDR p-value = 0.0416. Deleted in 5 samples.

Figure 55 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 97 patient cohort with NSCLC stage Ia-IIb, classified by presence or absence of a copy number loss (deletion) of Chr14, 55.2 – 60.0 Mb. [deletion marker 22]. FDR p-value = 0.0345. Deleted in 6 samples.

Figure 56 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 97 patient cohort with NSCLC stage Ia-IIb, classified by presence or absence of a copy number loss (deletion) of Chr14, 48.7 – 51.1 Mb. [deletion marker 23]. FDR p-value = 0.0006. Deleted in 5 samples.

Figure 57 is a Kaplan-Meier plot showing the time to recurrence (TTR) in days for a 97 patient cohort with NSCLC stage Ia-IIb, classified by presence or absence of a copy number loss (deletion) of Chr 4, 81.4 – 83.2 Mb. [deletion marker 24]. FDR p-value = 0.0047. Deleted in 5 samples.

Figure 58 is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number loss (deletion) of Chr 10, 51.9 – 54.2 Mb. [deletion marker 25]. FDR p-value = 0.0067. Deleted in 5 samples.

5 **Figure 59** is a Kaplan-Meier plot showing the overall survival (OS) in days for a 78 patient cohort with NSCLC stage Ia-IIa, classified by presence or absence of a copy number loss (deletion) of Chr 5, 55.2 – 58.6 Mb. [deletion marker 26]. FDR p-value = 0.0130 Deleted in 5 samples.

10 **Figure 60** is a Kaplan-Meier plot showing the overall survival (OS) in days for a 102 patient cohort with NSCLC stage Ia-IIb, classified by presence or absence of a copy number loss (deletion) of Chr 5, 67.8 – 68.5 Mb. [deletion marker 27]. FDR p-value = 0.0475 Deleted in 5 samples.

15 As can be seen from the Kaplan-Meier plots in **Figures 1-60**, copy number alterations in the specified markers are associated with short OS and/or shorter TTR in NSCLC stage I—II patients. Table 4 lists overall survival data for several markers. (Markers for which data is shown in red are shared between different clinical stages).

TABLE 4: Overall Survival for markers on Chr1, Chr2, Chr6, Chr8, Chr11, Chr12, Chr17, and Chr19

Stages	chrom	start.pos	length.seg	FDR	n.amp	mean.amp	n.SNP
1a-2a	2	147604021	3513659	0.0233	7	2.8516132	166
	2	159911944	1511940	0.0001	5	3.2498274	67
	2	200924525	3320890	0.0398	6	3.006085	79
	2	205893481	2160144	0.0075	5	2.9990652	101
	3	88399682	386599	0.0140	5	3.5534647	12
	6	36255222	423122	0.0347	6	2.9201916	8
	6	39088059	762306	0.0356	15	3.1071308	30
	6	123724457	11850520	0.0377	7	2.9452862	667
	8	4115551	55428	0.0126	7	2.8073117	19
	8	6895465	1889190	0.0166	7	3.0262839	36
	11	61374252	2935902	0.0004	9	3.2120357	46
	11	64310154	493823	0.0040	12	3.5343537	6
	11	64803977	880941	0.0004	7	3.6506583	9
	12	93683	1774306	0.0493	11	3.604318	50
	17	43477124	1455714	0.0219	7	3.1622542	24
	17	51532820	1678229	0.0054	10	3.1730034	54
	17	69173224	2131396	0.0304	23	3.1612824	32
	19	32693527	387442	0.0183	18	4.0913848	8
	19	33195577	113123	0.0459	22	3.841479	6
	19	34722418	921516	0.0299	27	4.1530261	20
	19	38853838	1895624	0.0085	24	3.895232	34
	19	57033283	5156456	0.0091	14	3.1469281	83
1b-2b	1	109538586	1580066	0.0224	5	2.9805551	58
	6	70761833	382704	0.0116	17	3.2107404	28
1a-2b	6	70761833	382704	0.0110	24	3.0754468	28

Table 5 lists the genes and miRNA's that are encoded by nucleotide sequences within each cancer outcome marker sequence. In any of the methods, the cancer outcome marker can be selected from among those listed in Table 5. Those markers designated "M1" through "M20" are each a region of chromosomal DNA, the amplification of which produces a copy number gain in the cancer outcome marker, wherein the copy number gain is associated with a poor disease outcome. Those markers designated "DM1" through "DM27" are each a region of chromosomal DNA, the deletion of which produces a copy number loss in the cancer outcome marker, wherein the copy number loss is associated with a poor disease outcome.

TABLE 5: Cancer outcome markers and corresponding genes and miRNA's

ID No.	Cancer outcome marker	Genes and miRNA's
M1	Chr 19, 34.7 Mb—35.6 Mb;	C19orf12; C19orf12; cyclin E1; PLEKHF1; POP4; and ZNF536
M2	Chr 19, 38.9—40.7 Mb;	ATP4A ATPase; CHST8, DMKN FAR1,2,3; FXYD1,3,5,7; GAPDHS; GPI; GPR42; GRAMD1A; HAMP; HPN; KCTD15 KIAA0355; KRTDAP; LGI4; LSM14A; LSR; MAG; PDCD2L; SAE2 SUMO1; SBSN; SCN1B; TMEM147,162; USF2; WTIP; and ZNF181,30,302,599,792
M3	Chr 17, 69.2—71.3 Mb;	ARMC7 (armadillo repeat containing 7); ATP5H ATP synthase (H ⁺ transporting, mitochondrial F0 complex, subunit d); CASKIN2 (CASK interacting protein 2); CD300A (CD300a molecule); CD300C (CD300c molecule); CD300E (CD300e molecule); CD300LB (CD300 molecule-like family member b); CD300LF (CD300 molecule-like family member f); CDR2L (cerebellar degeneration-related protein 2-like); DNAI2 (dynein, axonemal, intermediate chain 2); (FADS6 fatty acid desaturase domain family, member 6); FDXR (ferredoxin reductase); GALK1 (galactokinase 1); GGA3 (golgi associated, gamma adaptin ear containing, ARF binding protein); GPR142 (G protein-coupled receptor 142); GPRC5C (G protein-coupled receptor, family C, group 5, member C); GRB2 (growth factor receptor-bound protein 2); GRIN2C (glutamate receptor, ionotropic, N-methyl D-aspartate 2C); H3F3B (H3 histone, family 3B (H3.3B)); HN1 (hematological and neurological expressed 1 ICT1 immature colon carcinoma transcript 1); ITGB4 (integrin, beta 4); KCTD2 (potassium channel tetramerisation domain containing 2); KIAA0195; KIF19 (kinesin family member 19); LLGL2 (lethal giant larvae homolog 2 (Drosophila)); LOC388419 (galectin-3-binding protein-like); MIF4GD (MIF4G domain containing); MRPS7 (mitochondrial ribosomal protein S7); NAT9 (N-acetyltransferase 9); NT5C (5', 3'-nucleotidase, cytosolic); NUP85 (nucleoporin 85kDa); OTOP2 (otopetrin 2); OTOP3 (otopetrin 3); RAB37 (RAB37, member RAS oncogene family); RECQL5 (RecQ protein-like 5); RPL38 ribosomal protein L38; SAP30BP (SAP30 binding protein); SLC16A5 (solute carrier family 16, member 5 (monocarboxylic acid transporter 6)); SLC25A19 (solute carrier family 25 (mitochondrial thiamine pyrophosphate carrier), member 19); SLC9A3R1 (solute carrier family 9 (sodium/hydrogen exchanger), member 3 regulator 1); SUMO2 (SMT3 suppressor of mif two 3 homolog 2 (S.

		cerevisiae)); TMEM104 (transmembrane protein 104); TTYH2 (tweety homolog 2 (Drosophila)); UNK (unkempt homolog (Drosophila)); and USH1G (Usher syndrome 1G (autosomal recessive))
M4	Chr 6, 70.8—71.1 Mb;	COL19A1 (collagen, type XIX, alpha 1), and COL9A1 (collagen, type IX, alpha 1)
M5	Chr 12, 93.7 kb—1.9Mb;	ADIPOR2 (adiponectin receptor 2); B4GALNT3 (beta-1,4-N-acetyl-galactosaminyl transferase 3); CACNA2D4 (calcium channel, voltage-dependent, alpha 2/delta subunit 4); CCDC77 (coiled-coil domain containing 77); ERC1 (ELKS/RAB6-interacting/CAST family member 1); FBXL14 (F-box and leucine-rich repeat protein 14); HSN2 (hereditary sensory neuropathy, type II); IQSEC3 (IQ motif and Sec7 domain 3); JARID1A (jumonji, AT rich interactive domain 1A); LRTM2 (leucine-rich repeats and transmembrane domains 2); NINJ2 (ninjurin 2); RAD52 (RAD52 homolog (S. cerevisiae)); SLC6A12 (solute carrier family 6 (neurotransmitter transporter, betaine/GABA), member 12); SLC6A13 (solute carrier family 6 (neurotransmitter transporter, GABA), member 13); WNK1 (WNK lysine deficient protein kinase 1); and WNT5B (wingless-type MMTV integration site family, member 5B)
M6	Chr 11, 64.3—64.8 Mb;	ARL2 (ADP-ribosylation factor-like 2); ATG2A ATG2 (autophagy related 2 homolog A (S. cerevisiae)); BATF2 (basic leucine zipper transcription factor, ATF-like 2); CAPN1 calpain 1, (mu/I) large subunit); CDC42BPG (CDC42 binding protein kinase gamma (DMPK-like)); CDCA5 (cell division cycle associated 5); EHD1 (EH-domain containing 1); FAU (Finkel-Biskis-Reilly murine sarcoma virus (FBR-MuSV) ubiquitously expressed); GPHA2 (glycoprotein hormone alpha 2); MAP4K2 mitogen-activated protein kinase kinase kinase kinase 2 MEN1 multiple endocrine neoplasia I MRPL49 mitochondrial ribosomal protein L49 NAALADL1 N-acetylated alpha-linked acidic dipeptidase-like 1 POLA2 polymerase (DNA directed), alpha 2 (70kD subunit) PPP2R5B protein phosphatase 2, regulatory subunit B', beta isoform SAC3D1 SAC3 domain containing 1 SLC22A20 solute carrier family 22, member 20 SNX15 sorting nexin 15 SPDYC speedy homolog C (Drosophila) SYVN1 synovial apoptosis inhibitor 1, synoviolin TM7SF2 transmembrane 7 superfamily member 2 ZFPL1 zinc finger protein-like 1 ZNHIT2 zinc finger, HIT type 2;

		hsa-mir-192; and hsa-mir-194-2
M7	Chr 19, 57.0—62.2 Mb;	<p> BIRC8 (baculoviral IAP repeat-containing 8); BRSK1 (BR serine/threonine kinase 1); CACNG6,7,8 calcium channel, voltage-dependent, gamma subunit 6,7,8 CCDC106 coiled-coil domain containing 106 CDC42EP5 CDC42 effector protein (Rho GTPase binding) 5 CNOT3 CCR4-NOT transcription complex, subunit 3 COX6B2 cytochrome c oxidase subunit VIb polypeptide 2 (testis) DPRX divergent-paired related homeobox EPN1 epsin 1 EPS8L1 EPS8-like 1 FCAR Fc fragment of IgA, receptor for FIZ1 FLT3-interacting zinc finger 1 GALP galanin-like peptide GP6 glycoprotein VI (platelet) HSPBP1 hsp70-interacting protein IL11 interleukin 11 ISOC2 isochorismatase domain containing 2 KIR2DL1, KIR2DL4, KIR2DS4 KIR3DL1, KIR3DL3, KIR3DX1 killer cell immunoglobulin-like receptor LAIR1,2 leukocyte-associated immunoglobulin-like receptor 1,2 LENG1,4,8,9 leukocyte receptor cluster (LRC) member 1,4,8,9 LILRA2,3,4 leukocyte immunoglobulin-like receptor, subfamily A (with TM domain), member 2,3,4 LILRB1,2,3,4,5 leukocyte immunoglobulin-like receptor, subfamily B (with TM and ITIM domains), member 1,2,3,4,5 MYADM myeloid-associated differentiation marker NAT14 N-acetyltransferase 14 NCR1 natural cytotoxicity triggering receptor 1 NDUFA3 NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 3, 9kDa NLRP2,4,5,7,8,9,11,12,13 NLR family, pyrin domain containing 2,4,5,7,8,9,11,12,13. OSCAR osteoclast associated, immunoglobulin-like receptor PEG3 paternally expressed 3 PPP1R12C protein phosphatase 1, regulatory (inhibitor) subunit 12C PPP2R1A protein phosphatase 2 (formerly 2A), regulatory subunit A, alpha isoform PRKCG protein kinase C, gamma PRPF31 PRP31 pre-mRNA processing factor 31 homolog (S. </p>

		<p>cerevisiae) PTPRH protein tyrosine phosphatase, receptor type, H RDH13 retinol dehydrogenase 13 (all-trans/9-cis) RPL28 ribosomal protein L28 RPS9 ribosomal protein S9 SAPS1 SAPS domain family, member 1 SUV420H2 suppressor of variegation 4-20 homolog 2 (Drosophila) SYT5 synaptotagmin V TFPT TCF3 (E2A) fusion partner (in childhood Leukemia) TMC4 transmembrane channel-like 4 TMEM190 transmembrane protein 190 TMEM86B transmembrane protein 86B TNNT3 troponin I type 3 (cardiac) TNNT1 troponin T type 1 (skeletal, slow) TSEN34 tRNA splicing endonuclease 34 homolog (S. cerevisiae) TTYH1 tweety homolog 1 (Drosophila) U2AF2 U2 small nuclear RNA auxiliary factor 2 UBE2S ubiquitin-conjugating enzyme E2S VN1R2 vomeronasal 1 receptor 2 VN1R4 vomeronasal 1 receptor 4 VSTM1 V-set and transmembrane domain containing 1 ZNF28,160,320,321,331,347,350,415,432,444,468,470 zinc finger protein 28,160,320,321,331,347,350,415,432,444,468,470; and miRNA's including hsa-mir-643, hsa-mir-512-1, hsa-mir-512-2, hsa-mir-498, hsa-mir-520e, hsa-mir-515-1, hsa-mir-519e, hsa-mir-520f, hsa-mir-515-2, hsa-mir-519c, hsa-mir-520a, hsa-mir-526b, hsa-mir-519b, hsa-mir-525, hsa-mir-523, hsa-mir-518f, hsa-mir-520b, hsa-mir-518b, hsa-mir-526a-1, hsa-mir-520c, hsa-mir-518c, hsa-mir-524, hsa-mir-517a, hsa-mir-519d, hsa-mir-521-2, hsa-mir-520d, hsa-mir-517b, hsa-mir-520g, hsa-mir-516-3, hsa-mir-526a-2, hsa-mir-518e, hsa-mir-518a-1, hsa-mir-518d, hsa-mir-516-4, hsa-mir-518a-2, hsa-mir-517c, hsa-mir-520h, hsa-mir-521-1, hsa-mir-522, hsa-mir-519a-1, hsa-mir-527, hsa-mir-516-1, hsa-mir-516-2, hsa-mir-519a-2, hsa-mir-371, hsa-mir-372, hsa-mir-373, hsa-mir-516a-1, hsa-mir-516a-2, hsa-mir-516b-1, hsa-mir-516b-2, hsa-mir-517a-1, hsa-mir-517a-2, hsa-mir-520c-1, hsa-mir-520c-2</p>
M8	Chr 6, 39.1—39.9 Mb;	<p>C6orf64 (chromosome 6 open reading frame 64); DNAH8 dynein, axonemal, heavy chain 8 GLP1R glucagon-like peptide 1 receptor KCNK16 potassium channel, subfamily K, member 16</p>

		KCNK17 potassium channel, subfamily K, member 17 KCNK5 potassium channel, subfamily K, member 5 KIF6 kinesin family member 6.
M9	Chr 11, 64.8—65.7 Mb;	BANF1 (barrier to autointegration factor 1); CATSPER1 cation channel, sperm associated 1 CCDC85B coiled-coil domain containing 85B CDC42EP2CDC42 effector protein (Rho GTPase binding) 2 CFL1 cofilin 1 (non-muscle) CST6 cystatin E/M CTSW cathepsin W DPF2 D4, zinc and double PHD fingers family 2 DRAP1 DR1-associated protein 1 (negative cofactor 2 alpha) EFEMP2 EGF-containing fibulin-like extracellular matrix protein 2 EHBP1L1 EH domain binding protein 1-like 1 FAM89B family with sequence similarity 89, member B FIBP fibroblast growth factor (acidic) intracellular binding protein FOSL1 FOS-like antigen 1 FRMD8 FERM domain containing 8 GAL3ST3 galactose-3-O-sulfotransferase 3 HTATIP HIV-1 Tat interacting protein, 60kDa. KCNK7 potassium channel, subfamily K, member 7 LTBP3 latent transforming growth factor beta binding protein 3 MAP3K11 mitogen-activated protein kinase kinase kinase 11 MGC11102 hypothetical protein MGC11102 MUS81 MUS81 endonuclease homolog (S. cerevisiae) OVOL1 ovo-like 1(Drosophila) PACS1 phosphofurin acidic cluster sorting protein 1 PCNXL3 pecanex-like 3 (Drosophila) POLA2 polymerase (DNA directed), alpha 2 (70kD subunit) RELA v-rel reticuloendotheliosis viral oncogene homolog A, nuclear factor of kappa light polypeptide gene enhancer in B-cells 3, p65 (avian) RNASEH2C ribonuclease H2, subunit C SART1 squamous cell carcinoma antigen recognized by T cells SCYL1 SCY1-like 1 (S. cerevisiae) SF3B2 splicing factor 3b, subunit 2, 145kDa SIPA1 signal-induced proliferation-associated gene 1 SLC25A45 solute carrier family 25, member 45 SSSCA1 Sjogren syndrome/scleroderma autoantigen 1 TIGD3 tigger transposable element derived 3 TSGA10IP testis specific, 10 interacting protein
M10	Chr 11, 61.4—64.3 Mb;	AHNAK (AHNAK nucleoprotein); ASRGL1 asparaginase like 1

	<p> B3GAT3 beta-1,3-glucuronyltransferase 3 (glucuronosyltransferase I) BAD BCL2-antagonist of cell death BEST1 bestrophin 1 BSCL2 Bernardinelli-Seip congenital lipodystrophy 2 (seipin) CCDC88B coiled-coil domain containing 88B CHRM1 cholinergic receptor, muscarinic 1 COX8A cytochrome c oxidase subunit 8A (ubiquitous) DKFZP564J0863 DKFZP564J0863 protein DKFZP566E164 DKFZP566E164 protein DNAJC4 DnaJ (Hsp40) homolog, subfamily C, member 4 EEF1G eukaryotic translation elongation factor 1 gamma EML3 echinoderm microtubule associated protein like 3 ESRRA estrogen-related receptor alpha FADS2,3 fatty acid desaturase 2,3 FKBP2 FK506 binding protein 2, 13kDa FLRT1 fibronectin leucine rich transmembrane protein 1 FTH1 ferritin, heavy polypeptide 1 GANAB glucosidase, alpha; neutral AB GNG3 guanine nucleotide binding protein (G protein), gamma 3 GPR137 G protein-coupled receptor 137 HRASLS2,3,5 HRAS-like suppressor 2,3,5 INCENP inner centromere protein antigens 135/155kDa INTS5 integrator complex subunit 5 KCNK4 potassium channel, subfamily K, member 4 LGALS12 lectin, galactoside-binding, soluble, 12 (galectin 12) MACROD1 MACRO domain containing 1 MARK2 MAP/microtubule affinity-regulating kinase 2 MGC3196 hypothetical protein MGC3196 MTA2 metastasis associated 1 family, member 2 NAT11 N-acetyltransferase 11 NRXN2 neurexin 2 NUDT22 nudix (nucleoside diphosphate linked moiety X)-type motif 22 NXF1 nuclear RNA export factor 1 OTUB1 OTU domain, ubiquitin aldehyde binding 1 PLCB3 phospholipase C, beta 3 (phosphatidylinositol-specific) POLR2G polymerase (RNA) II (DNA directed) polypeptide G PPP1R14B protein phosphatase 1, regulatory (inhibitor) subunit 14B PRDX5 peroxiredoxin 5 PYGM phosphorylase, glycogen; muscle (McArdle syndrome, glycogen storage disease type V) RAB3IL1 RAB3A interacting protein (rabin3)-like 1 RARRES3 retinoic acid receptor responder (tazarotene induced) </p>
--	--

		<p>3</p> <p>RASGRP2 RAS guanyl releasing protein 2 (calcium and DAG-regulated)</p> <p>RCOR2 REST corepressor 2</p> <p>ROM1 retinal outer segment membrane protein 1</p> <p>RPS6KA4 ribosomal protein S6 kinase, 90kDa, polypeptide 4</p> <p>RTN3 reticulon 3</p> <p>SCGB1A1, 1D1, 1D2, 1D4, 2A1, 2A1 secretoglobin, family</p> <p>SF1 splicing factor 1</p> <p>SLC22A10, 11, 12, 6, 8, 9 solute carrier family 22 (organic anion/cation transporter) SLC3A2 solute carrier family 3 (activators of dibasic and neutral amino acid transport), member 2</p> <p>STIP1 stress-induced-phosphoprotein 1 (Hsp70/Hsp90-organizing protein)</p> <p>STX5 syntaxin 5</p> <p>TAF6L TAF6-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa</p> <p>TRPT1 tRNA phosphotransferase 1</p> <p>TTC9C tetratricopeptide repeat domain 9C</p> <p>TUT1 terminal uridylyl transferase 1, U6 snRNA-specific</p> <p>URP2 UNC-112 related protein 2</p> <p>UST6 putative UST1-like organic anion transporter</p> <p>VEGFB vascular endothelial growth factor B</p> <p>WDR74 WD repeat domain 74; and</p> <p>ZBTB3 zinc finger and BTB domain containing 3</p>
M11	Chr 17, 51.5—53.2 Mb;	<p>AKAP1 (A kinase (PRKA) anchor protein 1);</p> <p>ANKFN1 (ankyrin-repeat and fibronectin type III domain containing 1);</p> <p>C17orf67 chromosome 17 open reading frame 67</p> <p>COIL coilin</p> <p>DGKE diacylglycerol kinase, epsilon 64kDa</p> <p>MSI2 musashi homolog 2 (Drosophila)</p> <p>NOG noggin</p> <p>SCPEP1 serine carboxypeptidase 1; and</p> <p>TRIM25 tripartite motif-containing 25</p>
M12	Chr 17, 43.5—44.9 Mb;	<p>hsa-mir-10a; hsa-mir-196a-1; ABI3 (ABI gene family, member 3); ATP5G1 (ATP synthase, H⁺ transporting, mitochondrial F0 complex, subunit C1 (subunit 9));</p> <p>B4GALNT2 beta-1,4-N-acetyl-galactosaminyl transferase 2</p> <p>CALCOCO2 calcium binding and coiled-coil domain 2</p> <p>CBX1 chromobox homolog 1 (HP1 beta homolog Drosophila)</p> <p>GIP gastric inhibitory polypeptide</p> <p>GNGT2 guanine nucleotide binding protein (G protein), gamma transducing activity polypeptide 2</p>

		<p>HOXB1,2,3,4,5,6,7,8,9,13 homeobox B1,2,3,4,5,6,7,8,9,13</p> <p>IGF2BP1 insulin-like growth factor 2 mRNA binding protein 1</p> <p>NFE2L1 nuclear factor (erythroid-derived 2)-like 1</p> <p>NGFR nerve growth factor receptor (TNFR superfamily, member 16)</p> <p>PHB prohibitin</p> <p>PHOSPHO1 phosphatase, orphan 1</p> <p>PRAC small nuclear protein PRAC</p> <p>SKAP1 src kinase associated phosphoprotein 1</p> <p>SNF8 SNF8, ESCRT-II complex subunit, homolog (S. cerevisiae)</p> <p>SNX11 sorting nexin 11</p> <p>TTL6 tubulin tyrosine ligase-like family, member 6</p> <p>UBE2Z (ubiquitin-conjugating enzyme E2Z); and</p> <p>ZNF652 (zinc finger protein 652).</p>
M13	Chr 2, 147.6—151.1 Mb;	<p>ACVR2A activin A receptor, type IIA; C2orf25 chromosome 2 open reading frame 25</p> <p>EPC2 enhancer of polycomb homolog 2 (Drosophila)</p> <p>KIF5C kinesin family member 5C</p> <p>LOC130576 hypothetical protein LOC130576</p> <p>LYPD6 LY6/PLAUR domain containing 6</p> <p>MBD5 methyl-CpG binding domain protein 5</p> <p>ORC4L origin recognition complex, subunit 4-like (yeast)</p> <p>RND3 Rho family GTPase 3</p>
M14	Chr 6, 123.7—135.6 Mb;	<p>hsa-mir-588;</p> <p>AKAP7 (A kinase (PRKA) anchor protein 7);</p> <p>ALDH8A1 aldehyde dehydrogenase 8 family, member A1</p> <p>ARG1 arginase, liver</p> <p>ARHGAP18 Rho GTPase activating protein 18</p> <p>CTGF connective tissue growth factor</p> <p>ECHDC1 enoyl Coenzyme A hydratase domain containing 1</p> <p>ENPP1,3 ectonucleotide pyrophosphatase/phosphodiesterase 1,3</p> <p>EPB41L2 erythrocyte membrane protein band 4.1-like 2</p> <p>EYA4 eyes absent homolog 4 (Drosophila)</p> <p>HDDC2 HD domain containing 2</p> <p>HEY2 hairy/enhancer-of-split related with YRPW motif 2</p> <p>HINT3 histidine triad nucleotide binding protein 3</p> <p>KIAA1913 KIAA1913</p> <p>LAMA2 laminin, alpha 2 (merosin, congenital muscular dystrophy)</p> <p>MED23 mediator complex subunit 23</p> <p>MOXD1 monooxygenase, DBH-like 1</p> <p>MYB v-myb myeloblastosis viral oncogene homolog (avian)</p> <p>NCOA7 nuclear receptor coactivator 7</p> <p>NKAIN2 Na⁺/K⁺ transporting ATPase interacting 2</p>

		<p>OR2A4 olfactory receptor, family 2, subfamily A, member 4</p> <p>PTPRK protein tyrosine phosphatase, receptor type, K. RNF146</p> <p>ring finger protein 146</p> <p>RNF217 ring finger protein 217</p> <p>RPS12 ribosomal protein S12</p> <p>SAMD3 sterile alpha motif domain containing 3</p> <p>SGK serum/glucocorticoid regulated kinase</p> <p>SLC2A12 solute carrier family 2 (facilitated glucose transporter), member 12</p> <p>STX7 syntaxin 7</p> <p>TAAR1,2,5,6,8,9 trace amine associated receptor 1,2,5,6,8,9</p> <p>TBPL1 TBP-like 1</p> <p>TCF21 transcription factor 21</p> <p>TPD52L1 tumor protein D52-like 1</p> <p>TRDN triadin</p> <p>TRMT11 tRNA methyltransferase 11 homolog (<i>S. cerevisiae</i>)); and</p> <p>VNN1,2,3 (vanin 1,2,3).</p>
M15	Chr 8, 6.9—8.8 Mb;	<p>CLDN23 claudin 23;</p> <p>DEFA5 defensin, alpha 5, Paneth cell-specific;</p> <p>DEFB103B defensin, beta 103B</p> <p>DEFB104A defensin, beta 104A</p> <p>DEFB104B defensin, beta 104B</p> <p>DEFB105B defensin, beta 105B</p> <p>DEFB106A defensin, beta 106A</p> <p>DEFB106B defensin, beta 106B</p> <p>DEFB107A defensin, beta 107A</p> <p>DEFB107B defensin, beta 107B</p> <p>DEFB4 defensin, beta 4</p> <p>MFHAS1 malignant fibrous histiocytoma amplified sequence 1</p> <p>PRAGMIN homolog of rat pragma of Rnd2</p> <p>SPAG11A sperm associated antigen 11A; and</p> <p>SPAG11B sperm associated antigen 11B</p>
M16	Chr 2, 159.9—161.4 Mb;	<p>BAZ2B bromodomain adjacent to zinc finger domain, 2B;</p> <p>CD302 CD302 molecule</p> <p>ITGB6 integrin, beta 6</p> <p>LY75 lymphocyte antigen 75</p> <p>MARCH7 (membrane-associated ring finger (C3HC4) 7);</p> <p>PLA2R1 (phospholipase A2 receptor 1, 180kDa); and</p> <p>RBMS1 (RNA binding motif, single stranded interacting protein 1).</p>
M17	Chr 2, 200.9—204.2 Mb;	<p>ABI2 abl interactor 2;</p> <p>ALS2 amyotrophic lateral sclerosis 2 (juvenile)</p> <p>ALS2CR2, 4, 7, 8, 11, 12, 13 amyotrophic lateral sclerosis 2 (juvenile) chromosome region, candidate 2, 4, 7, 8, 11, 12, 13</p>

		<p> AOX1 aldehyde oxidase 1 BMPR2 bone morphogenetic protein receptor, type II (serine/threonine kinase) BZW1 basic leucine zipper and W2 domains 1 CASP10 caspase 10, apoptosis-related cysteine peptidase CASP8 caspase 8, apoptosis-related cysteine peptidase CFLAR CASP8 and FADD-like apoptosis regulator CLK1 CDC-like kinase 1 CYP20A1 cytochrome P450, family 20, subfamily A, polypeptide 1 FAM126B family with sequence similarity 126, member B FZD7 frizzled homolog 7 (Drosophila) ICA1L islet cell autoantigen 1,69kDa-like KCTD18 potassium channel tetramerisation domain containing 18 LOC26010 viral DNA polymerase-transactivated protein 6 MPP4 membrane protein, palmitoylated 4 (MAGUK p55 subfamily member 4). NBEAL1 neurobeachin-like 1 NDUFB3 NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 3, 12kDa NIF3L1 NIF3 NGG1 interacting factor 3-like 1 (S. pombe) NOP5/NOP58 nucleolar protein NOP5/NOP58 ORC2L origin recognition complex, subunit 2-like (yeast) PPIL3 peptidylprolyl isomerase (cyclophilin)-like 3 RAPH1 Ras association (RalGDS/AF-6) and pleckstrin homology domains 1 SGOL2 shugoshin-like 2 (S. pombe) SUMO1 SMT3 suppressor of mif two 3 homolog 1 (S. cerevisiae) TRAK2 trafficking protein, kinesin binding 2; and WDR12 (WD repeat domain 12) </p>
M18	Chr 6, 36.3—36.7 Mb;	<p> BRPF3 (bromodomain and PHD finger containing, 3) DKFZp779B1540 hypothetical protein DKFZp779B1540 ETV7 ets variant gene 7 (TEL2 oncogene) KCTD20 potassium channel tetramerisation domain containing 20 PNPLA1 patatin-like phospholipase domain containing 1 PXT1 peroxisomal, testis specific 1 SFRS3 splicing factor, arginine/serine-rich 3; and STK38 (serine/threonine kinase 38) </p>
M19	Chr 2, 205.9—208.1 Mb; and	<p> ADAM23 (ADAM metallopeptidase domain 23); CPO carboxypeptidase O; DYT1 dystrotelin EEF1B2 eukaryotic translation elongation factor 1 beta 2 FASTKD2 FAST kinase domains 2 </p>

		<p>FLJ20309 hypothetical protein FLJ20309</p> <p>GPR1 G protein-coupled receptor 1</p> <p>KLF7 Kruppel-like factor 7 (ubiquitous)</p> <p>MDH1B malate dehydrogenase 1B, NAD (soluble)</p> <p>NDUFS1 NADH dehydrogenase (ubiquinone) Fe-S protein 1, 75kDa (NADH-coenzyme Q reductase)</p> <p>NRP2 neuropilin 2</p> <p>PARD3B par-3 partitioning defective 3 homolog B (C. elegans)</p> <p>ZDBF2 (zinc finger, DBF-type containing 2); and</p> <p>HCG_1657980 hCG1657980</p>
M20	Chr 1, 109.5—111.1 Mb.	<p>hsa-mir-197;</p> <p>AHCYL1 S-adenosylhomocysteine hydrolase-like 1);</p> <p>ALX3 aristaless-like homeobox 3</p> <p>AMIGO1 adhesion molecule with Ig-like domain 1</p> <p>AMPD2 adenosine monophosphate deaminase 2 (isoform L)</p> <p>ATXN7L2 ataxin 7-like 2</p> <p>CELSR2 cadherin, EGF LAG seven-pass G-type receptor 2 (flamingo homolog, Drosophila)</p> <p>CSF1 colony stimulating factor 1 (macrophage)</p> <p>CYB561D1 cytochrome b-561 domain containing 1</p> <p>EPS8L3 EPS8-like 3</p> <p>FAM40A family with sequence similarity 40, member A</p> <p>GNAI3 guanine nucleotide binding protein (G protein), alpha inhibiting activity polypeptide 3</p> <p>GNAT2 guanine nucleotide binding protein (G protein), alpha transducing activity polypeptide 2</p> <p>GPR61 G protein-coupled receptor 61</p> <p>GSTM1,M2,M3,M4,M5 glutathione S-transferase M1, M2 (muscle), M3 (brain), M4, M5</p> <p>HBXIP hepatitis B virus x interacting protein</p> <p>KCNA2,3,4,10 potassium voltage-gated channel, shaker-related subfamily, member 2,3,4,10; KIAA1324 KIAA1324</p> <p>MYBPHL myosin binding protein H-like</p> <p>PROK1 prokineticin 1</p> <p>PSMA5 proteasome (prosome, macropain) subunit, alpha type, 5</p> <p>PSRC1 proline/serine-rich coiled-coil 1</p> <p>RBM15 RNA binding motif protein 15</p> <p>SARS seryl-tRNA synthetase</p> <p>SLC16A4 solute carrier family 16, member 4 (monocarboxylic acid transporter 5)</p> <p>SLC6A17 solute carrier family 6, member 17</p> <p>SORT1 sortilin 1</p> <p>SYPL2 synaptophysin-like 2</p> <p>UBL4B (ubiquitin-like 4B)</p>
DM1	Chr 5, 62.9 – 67.8 Mb	ADAMTS6 ADAM metalloproteinase with thrombospondin type

		<p>1 motif, 6 CD180 CD180 molecule CENPK centromere protein K ERBB2IP erbb2 interacting protein FLJ13611 hypothetical protein FLJ13611 HTR1A 5-hydroxytryptamine (serotonin) receptor 1A MAST4 microtubule associated serine/threonine kinase family member 4 NLN neurolysin (metallopeptidase M3 family) P18SRP P18SRP protein PIK3R1 phosphoinositide-3-kinase, regulatory subunit 1 (p85 alpha) PPWD1 peptidylprolyl isomerase domain and WD repeat containing 1 RGS7BP regulator of G-protein signaling 7 binding protein RNF180 ring finger protein 180 SDCCAG10 serologically defined colon cancer antigen 10 SFRS12 splicing factor, arginine/serine-rich 12 SGTB small glutamine-rich tetratricopeptide repeat (TPR)-containing, beta0; and TRIM23 tripartite motif-containing 23.</p>
DM2	Chr 5, 53.3 – 53.8 Mb	<p>ARL15 (ADP-ribosylation factor-like 15); HSPB3 (heat shock 27kDa protein 3) and hsa-miR-581.</p>
DM3	Chr 4, 105.8 – 107.2 Mb	<p>FLJ20184 (hypothetical protein FLJ20184); GSTCD (glutathione S-transferase, C-terminal domain containing); INTS12 integrator complex subunit 12 KIAA1546 KIAA1546 MGC16169 hypothetical protein MGC16169 NPNT (nephronectin); and PPA2 pyrophosphatase (inorganic) 2.</p>
DM4	Chr 16, 45.8 – 46.3 Mb	<p>ITFG1 (integrin alpha FG-GAP repeat containing 1) and PHKB (phosphorylase kinase, beta).</p>
DM5	Chr 5, 50.7 – 52.0 Mb	<p>ISL1 (ISL LIM homeobox).</p>
DM6	Chr 5, 94.2 – 96.1 Mb	<p>ARSK (arylsulfatase family, member K); CAST (calpastatin); ELL2 (elongation factor, RNA polymerase II, 2); FAM81B family with sequence similarity 81, member B GLRX glutaredoxin (thioltransferase) GPR150 G protein-coupled receptor 150 KIAA0372 KIAA0372 MCTP1 multiple C2 domains, transmembrane 1 PCSK1 proprotein convertase subtilisin/kexin type 1 RFESD (Rieske (Fe-S) domain containing) RHOBTB3 Rho-related BTB domain containing 3</p>

		SPATA9 (spermatogenesis associated 9); and hsa-miR-583.
DM7	Chr 9, 36.1 – 37.0 Mb	C9orf19 chromosome 9 open reading frame 19 CCIN calicin CLTA clathrin, light chain (Lca) GNE glucosamine (UDP-N-acetyl)-2-epimerase/N- acetylmannosamine kinase MELK maternal embryonic leucine zipper kinase PAX5 paired box 5 RECK reversion-inducing-cysteine-rich protein with kazal motifs RNF38 ring finger protein 38.
DM8	Chr 5, 94.2 – 96.1 Mb	ARSK arylsulfatase family, member K CAST calpastatin ELL2 elongation factor, RNA polymerase II, 2 FAM81B family with sequence similarity 81, member B GLRX glutaredoxin (thioltransferase) GPR150 G protein-coupled receptor 150 KIAA0372 KIAA0372 MCTP1 multiple C2 domains, transmembrane 1 PCSK1 proprotein convertase subtilisin/kexin type 1 RFESD Rieske (Fe-S) domain containing RHOBTB3 Rho-related BTB domain containing 3 SPATA9 spermatogenesis associated
DM9	Chr14, 51.1 – 52.8 Mb	C14orf166 chromosome 14 open reading frame 166; DDHD1 DDHD domain containing 1 ERO1L ERO1-like (S. cerevisiae) FRMD6 FERM domain containing 6 GNG2 guanine nucleotide binding protein (G protein), gamma 2 GNPNAT1 glucosamine-phosphate N-acetyltransferase 1 GPR137C G protein-coupled receptor 137C NID2 nidogen 2 (osteonidogen) PLEKHC1 pleckstrin homology domain containing, family C (with FERM domain) member 1 PSMC6 proteasome (prosome, macropain) 26S subunit, ATPase, 6 PTGDR prostaglandin D2 receptor (DP) PTGER2 prostaglandin E receptor 2 (subtype EP2), 53kDa STYX serine/threonine/tyrosine interacting protein TXNDC16 thioredoxin domain containing 16.
DM10	Chr 14, 61.5 – 68.6 Mb	ACTN1 actinin, alpha 1 AKAP5 A kinase (PRKA) anchor protein 5 ARG2 arginase, type II ATP6V1D ATPase, H ⁺ transporting, lysosomal 34kDa, V1 subunit D

	<p> C14orf50 chromosome 14 open reading frame 50 C14orf54 chromosome 14 open reading frame 54 C14orf83 chromosome 14 open reading frame 83 CHURC1 churchill domain containing 1 EIF2S1 eukaryotic translation initiation factor 2, subunit 1 alpha, 35kDa ESR2 estrogen receptor 2 (ER beta) FLJ39779 FLJ39779 protein FNTB farnesyltransferase, CAAX box, beta FUT8 fucosyltransferase 8 (alpha (1,6) fucosyltransferase) GPHB5 glycoprotein hormone beta 5 GPHN gephyrin GPX2 glutathione peroxidase 2 (gastrointestinal) HSPA2 heat shock 70kDa protein 2 KCNH5 potassium voltage-gated channel, subfamily H (eag-related), member 5 MAX MYC associated factor X MPP5 membrane protein, palmitoylated 5 (MAGUK p55 subfamily member 5) MTHFD1 methylenetetrahydrofolate dehydrogenase (NADP+ dependent) 1, methenyltetrahydrofolate cyclohydrolase, formyltetrahydrofolate synthetase PIGH phosphatidylinositol glycan anchor biosynthesis, class H PLEK2 pleckstrin 2 PLEKHG3 pleckstrin homology domain containing, family G (with RhoGef domain) member 3 PLEKHH1 pleckstrin homology domain containing, family H (with MyTH4 domain) member 1 PPP2R5E protein phosphatase 2, regulatory subunit B', epsilon isoform. RAB15 RAB15, member RAS oncogene family RAD51L1 RAD51-like 1 (S. cerevisiae) RDH11 retinol dehydrogenase 11 (all-trans/9-cis/11-cis) RDH12 retinol dehydrogenase 12 (all-trans/9-cis/11-cis) RHOJ ras homolog gene family, member J SGPP1 sphingosine-1-phosphate phosphatase 1 SPTB spectrin, beta, erythrocytic (includes spherocytosis, clinical type I) SYNE2 spectrin repeat containing, nuclear envelope 2 SYT16 synaptotagmin XVI VTI1B vesicle transport through interaction with t-SNAREs homolog 1B (yeast) WDR22 WD repeat domain 22 WDR89 WD repeat domain 89 ZBTB1 zinc finger and BTB domain containing 1 </p>
--	--

		ZBTB25 zinc finger and BTB domain containing 25 ZFP36L1 zinc finger protein 36, C3H type-like 1 ZFYVE26 zinc finger, FYVE domain containing 26 and hsa-miR-625.
DM11	Chr 9, 28.1 Mb	LINGO2 (leucine rich repeat and Ig domain containing 2).
DM12	Chr 4, 43.7 – 44.2 Mb	KCTD8 (potassium channel tetramerisation domain containing 8).
DM13	Chr 5, 60.8 – 62.9 Mb	DIMT1L DIM1 dimethyladenosine transferase 1-like (S. cerevisiae) FLJ37543 hypothetical protein FLJ37543 IPO11 importin 11 ISCA1L iron-sulfur cluster assembly 1 homolog (S. cerevisiae)-like KIF2A kinesin heavy chain member 2A .
DM14	Chr 3, 120.0 – 121.1 Mb	ADPRH ADP-ribosylarginine hydrolase; B4GALT4 UDP-Gal:betaGlcNAc beta 1,4-galactosyltransferase, polypeptide 4 C3orf1 chromosome 3 open reading frame 1 C3orf15 chromosome 3 open reading frame 15 C3orf30 chromosome 3 open reading frame 30 CD80 CD80 molecule CDGAP Cdc42 GTPase-activating protein COX17 COX17 cytochrome c oxidase assembly homolog (S. cerevisiae) GSK3B glycogen synthase kinase 3 beta IGSF11 immunoglobulin superfamily, member 11 KTELC1 KTEL (Lys-Tyr-Glu-Leu) containing 1 NR1I2 nuclear receptor subfamily 1, group I, member 2 PLA1A phospholipase A1 member A POPDC2 popeye domain containing 2 TMEM39A transmembrane protein 39A; and UPK1B uroplakin 1B.
DM15	Chr 4, 46.2 – 48.0 Mb	CLDN23 claudin 23; DEFA5 defensin, alpha 5, Paneth cell-specific; DEFB103B defensin, beta 103B DEFB104A defensin, beta 104A DEFB104B defensin, beta 104B DEFB105B defensin, beta 105B DEFB106A defensin, beta 106A DEFB106B defensin, beta 106B DEFB107A defensin, beta 107A DEFB107B defensin, beta 107B DEFB4 defensin, beta 4 MFHAS1 malignant fibrous histiocytoma amplified sequence 1 PRAGMIN homolog of rat pragma of Rnd2

		SPAG11A sperm associated antigen 11A; and SPAG11B sperm associated antigen 11B.
DM16	Chr 14, 38.9 – 40.0 Mb	FBXO33 (F-box protein 33).
DM17	Chr 4, 44.2 – 44.6 Mb	GNPDA2 (glucosamine-6-phosphate deaminase 2); GUF1 (GUF1 GTPase homolog (S. cerevisiae)); and YIPF7 (Yip1 domain family, member 7).
DM18	Chr 2, 213.7 – 214.3 Mb	IKZF2 IKAROS family zinc finger 2 (Helios) SPAG16 sperm associated antigen 16.
DM19	Chr14, 43.9 – 46.6 Mb	C14orf106 chromosome 14 open reading frame 106 C14orf155 chromosome 14 open reading frame 155 C14orf28 chromosome 14 open reading frame 28 FANCM Fanconi anemia, complementation group M FKBP3 FK506 binding protein 3, 25kDa KIAA0423 KIAA0423 KLHL28 kelch-like 28 (Drosophila) MDGA2 MAM domain containing glycosylphosphatidylinositol anchor 2 PRPF39 PRP39 pre-mRNA processing factor 39 homolog (S. cerevisiae) RPL10L ribosomal protein L10-like.
DM20	Chr 14, 27.6 – 28.6 Mb	FOXP1 (forkhead box G1).
DM21	Chr 3, 98.0 – 98.3 Mb	EPHA6 (EPH receptor A6).
DM22	Chr14, 55.2 – 60.0 Mb	ACTR10 actin-related protein 10 homolog (S. cerevisiae) ARID4A AT rich interactive domain 4A (RBP1-like) C14orf100 chromosome 14 open reading frame 100 C14orf101 chromosome 14 open reading frame 101 C14orf105 chromosome 14 open reading frame 105 C14orf108 chromosome 14 open reading frame 108 C14orf135 chromosome 14 open reading frame 135 C14orf149 chromosome 14 open reading frame 149 C14orf37 chromosome 14 open reading frame 37 C14orf39 chromosome 14 open reading frame 39 DAAM1 dishevelled associated activator of morphogenesis 1 DACT1 dapper, antagonist of beta-catenin, homolog 1 (Xenopus laevis) DHRS7 dehydrogenase/reductase (SDR family) member 7 EXOC5 exocyst complex component 5 GPR135 G protein-coupled receptor 135 KIAA0586 KIAA0586 NAT12 N-acetyltransferase 12 OTX2 orthodenticle homeobox 2 PELI2 pellino homolog 2 (Drosophila) PPM1A protein phosphatase 1A (formerly 2C), magnesium- dependent, alpha isoform PSMA3 proteasome (prosome, macropain) subunit, alpha type, 3

		RTN1 reticulon 1 SLC35F4 solute carrier family 35, member F4 TIMM9 translocase of inner mitochondrial membrane 9 homolog (yeast) UNQ9438 TIMM.
DM23	Chr14, 48.7 – 51.1 Mb	ABHD12B abhydrolase domain containing 12B ARF6 ADP-ribosylation factor 6 ATP5S ATP synthase, H ⁺ transporting, mitochondrial F0 complex, subunit s (factor B) C14orf104 chromosome 14 open reading frame 104 C14orf138 chromosome 14 open reading frame 138 CDKL1 cyclin-dependent kinase-like 1 (CDC2-related kinase) FRMD6 FERM domain containing 6 KLHDC1 kelch domain containing 1 KLHDC2 kelch domain containing 2 L2HGDH L-2-hydroxyglutarate dehydrogenase LOC196913 hypothetical protein LOC196913 LOC283551 hypothetical protein LOC283551 MAP4K5 mitogen-activated protein kinase kinase kinase 5 MGAT2 mannosyl (alpha-1,6-)-glycoprotein beta-1,2-N-acetylglucosaminyltransferase NIN ninein (GSK3B interacting protein) POLE2 polymerase (DNA directed), epsilon 2 (p59 subunit) PPIL5 peptidylprolyl isomerase (cyclophilin)-like 5 PYGL phosphorylase, glycogen; liver (Hers disease, glycogen storage disease type VI) RPL36AL ribosomal protein L36a-like RPS29 ribosomal protein S29.
DM24	Chr 4, 81.4 – 83.2 Mb	BMP3 bone morphogenetic protein 3 (osteogenic) C4orf22 chromosome 4 open reading frame 22 FGF5 fibroblast growth factor 5 PRKG2 protein kinase, cGMP-dependent, type II RASGEF1B RasGEF domain family, member 1B.
DM25	Chr 10, 51.9 – 54.2 Mb	ACF apobec-1 complementation factor ASAH2B N-acylsphingosine amidohydrolase (non-lysosomal ceramidase) 2B CSTF2T cleavage stimulation factor, 3' pre-RNA, subunit 2, 64kDa, tau variant DKK1 dickkopf homolog 1 (Xenopus laevis) MBL2 mannose-binding lectin (protein C) 2, soluble (opsonic defect) PRKG1 protein kinase, cGMP-dependent, type I SGMS1 sphingomyelin synthase 1 hsa-miR-605.

DM26	Chr 5, 55.2 – 58.6 Mb	ANKRD55 ankyrin repeat domain 55 C5orf29 chromosome 5 open reading frame 29 C5orf35 chromosome 5 open reading frame 35 DKFZp686D0972 similar to RIKEN cDNA 4732495G21 gene GPBP1 GC-rich promoter binding protein 1 IL31RA interleukin 31 receptor A IL6ST interleukin 6 signal transducer (gp130, oncostatin M receptor) MAP3K1 mitogen-activated protein kinase kinase kinase 1 MIER3 mesoderm induction early response 1, family member 3 PDE4D phosphodiesterase 4D, cAMP-specific (phosphodiesterase E3 dunce homolog, Drosophila) PLK2 polo-like kinase 2 (Drosophila) RAB3C RAB3C, member RAS oncogene family.
DM27	Chr 5, 67.8 – 68.5 Mb.	CCNB1 (cyclin B1) and SLC30A5 (solute carrier family 30 (zinc transporter), member 5).

Table 6 lists the coordinates for each cancer outcome marker, using the same reference numerals listed in Table 5. All coordinates are based on human genome assembly hg18 (NCBI Build 36).

5 **TABLE 6:**

MarkerID	chrom	start.pos	end.pos
M1	chr19	34722418	35643933
M2	chr19	38853838	40749461
M3	chr17	69173224	71304619
M4	chr6	70761833	71144537
M5	chr12	93683	1867988
M6	chr11	64310154	64803976
M7	chr19	57033283	62189738
M8	chr6	39088059	39850364
M9	chr11	64803977	65684917
M10	chr11	61374252	64310153
M11	chr17	51532820	53211048
M12	chr17	43477124	44932837
M13	chr2	147604021	151117679
M14	chr6	123724457	135574976
M15	chr8	6895465	8784654
M16	chr2	159911944	161423883
M17	chr2	200924525	204245414
M18	chr6	36255222	36678343
M19	chr2	205893481	208053624
M20	chr1	109538586	111118652
DM1	chr9	36056899	36988415

DM2	chr4	105818261	107238628
DM3	chr5	53264432	53790965
DM4	chr16	45791880	46313827
DM5	chr5	50706878	52008065
DM6	chr5	94204208	96112445
DM7	chr5	62942847	67798156
DM9	chr14	51108156	52752331
DM10	chr14	61456273	68632720
DM11	chr9	28057491	28114180
DM12	chr4	43689020	44161565
DM13	chr5	60797829	62942846
DM14	chr3	119993321	121112610
DM15	chr4	46246303	47955581
DM16	chr14	38939630	40021400
DM17	chr4	44161566	44606114
DM18	chr2	213677020	214308243
DM19	chr14	43899026	46591909
DM20	chr14	27646449	28630571
DM21	chr3	97988751	98257089
DM22	chr14	55249852	60045332
DM23	chr14	48734855	51108156
DM24	chr4	81371219	83187388
DM25	chr10	51929419	54199330
DM26	chr5	55221121	58648144
DM27	chr5	67798156	68516077

Unlike previously identified predictors (expression signatures), the biomarkers described herein represent DNA gains and losses (stable events measurable by FISH). FISH probes can be used to enable validation/use of the markers, and the markers are strong candidates for use as stratification biomarkers in clinical trials. They can be used for example to define molecular subgroups of disease with distinct outcomes. As such they are likely to correlate with drug response.

These data indicate that use of genomic copy number assessment of the genetic markers measured by FISH, and with use of an appropriate classifier, is of prognostic importance in early stage NSCLC. The classifier was able to produce statistically significant classification of patients who had been treated with surgery without neoadjuvant or follow-up chemotherapy into favorable and unfavorable recurrence categories. No present clinical *in vitro* diagnostic assay provides this capability. Thus, FISH assays to the listed markers performed on early stage

NSCLC biopsy specimens or resected tumors appear valuable in decisions related to adjuvant therapy.

Example 2: Validation of prognostic markers using a Korean sample set

To validate forty-six (46) of the biomarkers that correlated with the clinical outcome of low stage NSCLC patients, an additional set of low stage NSCLC tumor tissues was collected from the Samsung Cancer Center in Korea, together with associated clinical outcome information.

All samples were carefully dissected to maximize tumor/normal tissue ratio and verify histopathological type and stage. Only samples from patients with stage I and II samples were analyzed. All of these were from patients treated with surgical resection without any follow-up or neoadjuvant chemotherapy. Clinical information collected for each patient included age, sex, clinical stage, pathological stage, location, histology, differentiation, smoking status, chemotherapy status, radiation status, recurrence status, recurrence date, recurrence location, brain metastasis status, time to recurrence, date of last follow up, status at the last follow up, alive/dead, overall survival and cause of death. Time to Recurrence (TTR) and Overall Survival (OS) were chosen as the parameters of outcome. Other clinical parameters (node status, stage, etc) were considered as confounding variables. Times to recurrence of lung cancer and the overall survival times were obtained from the patient charts. Tables 7 and 8 provide the figures for Overall Survival and Total Time to Recurrence, respectively, for the patient cohort studied.

TABLE 7:

OS

Stage	deaths	Alive (censored)	total
1a	0	10	10
1b	22	33	55
2a	0	0	0
2b	6	2	8
Total	28	45	73

TABLE 8:**TTR**

Stage	recurred	Recurrence free (censored)	total
1a	0	10	10
1b	24	31	55
2a	0	0	0
2b	6	2	8
total	30	43	73

5 The samples were processed, DNA extracted, amplified and hybridized to Affymetrix SNP 6.0 arrays (Affymetrix, Inc., Santa Clara, CA) which contains more than 906,600 single nucleotide polymorphisms (SNPs) and more than 946,000 probes for the detection of copy number variation with a median intermarker distance over all 1.8 million SNP and copy number markers combined of less than 700 bases. The microarrays were processed according to recommendations of the manufacturer (Affymetrix). Copy number of these tumors was calculated by comparing to a HapMap set of 270 normal controls. The copy number was segmented using Partek software 6.09.0310.

15 The average copy number of the validation set showed a pattern similar to the previous training data set, but with a much higher density, as shown in Figure 61. Figure 61 compares the average copy number pattern between training and validating data set. The log transformed copy numbers of each marker were averaged across all samples in the training (above) and testing (below) sets, where 0 represents the normal two copies, and red and blue represent on average gain or loss of copy numbers, respectively. Each dot represents one marker on the array, and the x-axis represents the genomic locations ranked by chromosome 1 to 22.

20 The validation data presented in this example is based on eighteen (18) times greater the coverage of SNPs and CNV markers as compared to the 100K microarray data generated and used to identify the diagnostic markers, and therefore more small scale copy number changing events could be identified. Therefore, instead of calculating the copy number of each biomarker,

the copy number of each gene within these biomarkers was calculated, and then correlated to the Overall Survival or Time to Recurrence of the patients.

Sixty-one (61) genes in total, in six (6) different markers were validated with the criterion of p-value by logrank test below 0.05. The genes are listed in **Table 9** with significant p-values high-lighted by gray shading. Figures 62-162 are Kaplan-Meier plots showing either the overall survival (OS) or the time to recurrence (TTR) in days for the 73 patient cohort, classified by presence or absence of a copy number gain in the particular gene as indicated on each plot. As with the Kaplan-Meier plots of Figures 1-60, the x-axis represents time in days, and the y-axis the probability of patient survival (for OS), or of patient being being free of disease recurrence (for TTR). Whenever a relevant event occurred (either death for OS, or disease recurrence for TTR), the curve drops. When a patient was lost to follow up without a relevant event occurring, a mark in the horizontal lines was made, indicating the last event-free time. P-values were obtained by comparing the decrease in patient population with and without the biomarkers.

The number of markers ultimately validated as described is relatively small, which may be attributable in part to the fact that the two patient populations are from different ethnic groups. Previous large samplings were collected in Chicago, Illinois, USA and included a mix of Asian, Caucasian, African and Hispanic patients, while the validation samples were collected in Korea from a homogeneously Asian population. Additionally, the two samples sets were processed at different locations, Abbott Park for the first sample set and the Samsung Cancer Center for the second sample set. Although the sample processing in both locations followed the same suggested protocol, potential systems bias cannot be totally ruled out. Also, the two sample sets were assayed using different versions of Affymetrix SNPs arrays, between which density differed by a factor of eighteen (18). The additional probes included in the new array may reveal more detailed copy number variation events that were not observable using the older version of SNPs arrays.

TABLE 9: Validated genes within the identified biomarkers that correlate to clinical outcome of NSCLC patients

GeneSymbol	Chromosome	Type	MarkerID	Event	Normal	PvalTTR	PvalOS
C11orf20	chr11	Amp	Marker10	6	67	0.0024	0.0477
ESRRA	chr11	Amp	Marker10	6	67	0.0024	0.0477
HSPC152	chr11	Amp	Marker10	6	67	0.0024	0.0477
KCNK4	chr11	Amp	Marker10	6	67	0.0024	0.0477
PRDX5	chr11	Amp	Marker10	6	67	0.0024	0.0477
GPR137	chr11	Amp	Marker10	5	68	0.0132	0.1138
BAD	chr11	Amp	Marker10	4	69	0.0232	0.0278
DNAJC4	chr11	Amp	Marker10	4	69	0.0232	0.0278
FKBP2	chr11	Amp	Marker10	4	69	0.0232	0.0278
NUDT22	chr11	Amp	Marker10	4	69	0.0232	0.0278
PLCB3	chr11	Amp	Marker10	4	69	0.0232	0.0278
PPP1R14B	chr11	Amp	Marker10	4	69	0.0232	0.0278
TRPT1	chr11	Amp	Marker10	4	69	0.0232	0.0278
VEGFB	chr11	Amp	Marker10	4	69	0.0232	0.0278
FLRT1	chr11	Amp	Marker10	4	69	0.1288	0.0069
FADS2	chr11	Amp	Marker10	3	70	0.4208	0.0482
FADS3	chr11	Amp	Marker10	3	70	0.4208	0.0482
RAB3IL1	chr11	Amp	Marker10	3	70	0.4208	0.0482
AKAP1	chr17	Amp	Marker11	3	70	0.0370	0.3782
ANKFN1	chr17	Amp	Marker11	3	70	0.0370	0.3782
C17orf67	chr17	Amp	Marker11	3	70	0.0370	0.3782
COIL	chr17	Amp	Marker11	3	70	0.0370	0.3782
DGKE	chr17	Amp	Marker11	3	70	0.0370	0.3782
MSI2	chr17	Amp	Marker11	3	70	0.0370	0.3782
MTVR2	chr17	Amp	Marker11	3	70	0.0370	0.3782
NOG	chr17	Amp	Marker11	3	70	0.0370	0.3782
RNF126P1	chr17	Amp	Marker11	3	70	0.0370	0.3782
SCPEP1	chr17	Amp	Marker11	3	70	0.0370	0.3782
TRIM25	chr17	Amp	Marker11	3	70	0.0370	0.3782
MYO15B	chr17	Amp	Marker3	4	69	0.0312	0.1535
SLC16A5	chr17	Amp	Marker3	4	69	0.0342	0.1126
PACS1	chr11	Amp	Marker9	2	71	0.0015	0.0004
DKFZp761E198	chr11	Amp	Marker9	4	69	0.0189	0.0136
KAT5	chr11	Amp	Marker9	4	69	0.0189	0.0136
LTBP3	chr11	Amp	Marker9	4	69	0.0189	0.0136
MALAT1	chr11	Amp	Marker9	4	69	0.0189	0.0136
RELA	chr11	Amp	Marker9	4	69	0.0189	0.0136
RNASEH2C	chr11	Amp	Marker9	4	69	0.0189	0.0136
SCYL1	chr11	Amp	Marker9	4	69	0.0189	0.0136
EHBP1L1	chr11	Amp	Marker9	5	68	0.0224	0.0004
FAM89B	chr11	Amp	Marker9	5	68	0.0224	0.0004
KCNK7	chr11	Amp	Marker9	5	68	0.0224	0.0004

MAP3K11	chr11	Amp	Marker9	5	68	0.0224	0.0004
PCNXL3	chr11	Amp	Marker9	5	68	0.0224	0.0004
SIPA1	chr11	Amp	Marker9	5	68	0.0224	0.0004
SSSCA1	chr11	Amp	Marker9	5	68	0.0224	0.0004
CCNB1	chr5	Del	DelMarker27	19	54	0.3161	0.0495
SLC30A5	chr5	Del	DelMarker27	19	54	0.3161	0.0495
ARSK	chr5	Del	DelMarker6	9	64	0.0033	0.0135
FAM81B	chr5	Del	DelMarker6	9	64	0.0033	0.0135
GPR150	chr5	Del	DelMarker6	9	64	0.0033	0.0135
MCTP1	chr5	Del	DelMarker6	10	63	0.0114	0.0051
C5orf27	chr5	Del	DelMarker6	10	63	0.0122	0.0406
CAST	chr5	Del	DelMarker6	10	63	0.0122	0.0406
GLRX	chr5	Del	DelMarker6	10	63	0.0122	0.0406
PCSK1	chr5	Del	DelMarker6	10	63	0.0122	0.0406
RFESD	chr5	Del	DelMarker6	10	63	0.0122	0.0406
RHOBTB3	chr5	Del	DelMarker6	10	63	0.0122	0.0406
SPATA9	chr5	Del	DelMarker6	10	63	0.0122	0.0406
TTC37	chr5	Del	DelMarker6	10	63	0.0193	0.0491
ELL2	chr5	Del	DelMarker6	11	62	0.0467	0.1066

It is to be understood that the foregoing description is intended to illustrate and not limit the scope of the invention. Other aspects, advantages, and modifications of the invention are within the intended scope of the claims set forth below.

5

WHAT IS CLAIMED IS:

1. A method of predicting disease outcome in a patient being treated for lung cancer, the method comprising the steps of:

- a) providing a test sample from a patient;
- b) determining a copy number of a cancer outcome marker in the test sample;
- c) comparing the copy number of the cancer outcome marker in the test sample against a baseline copy number of two, thereby determining the presence or absence of a copy number change for the cancer outcome marker in the test sample; and
- d) based on the presence or absence of a copy number change for the cancer outcome marker in the test sample, identifying the patient as having an increased risk of a poor disease outcome when compared to a baseline measure of disease outcome in patients having no copy number change in the cancer outcome marker, wherein the presence of a copy number change in the cancer outcome marker is predictive of poor disease outcome.

2. The method of claim 1 wherein the poor disease outcome is at least one of a decreased overall survival time when compared to an overall survival time of patients having no copy number change for the cancer outcome marker, and a shorter time to recurrence when compared to an overall survival time of patients having no copy number change for the cancer outcome marker.

3. A method of predicting treatment outcome in a patient being treated for lung cancer, the method comprising the steps of:

- a) providing a test sample from a patient;
- b) determining the presence or absence of a copy number change for a cancer outcome marker in the test sample, wherein the cancer outcome marker is a region of chromosomal DNA, the change in copy number of which is associated with a poor disease outcome; and
- c) based on the presence or absence of a copy number change for the cancer outcome marker, determining whether the patient has a higher risk of a decreased overall survival time or a shorter time to recurrence when compared to an overall survival time of patients having no copy number gain for the cancer outcome marker.

4. The method of any of claims 1-3 wherein the cancer outcome marker is a region of chromosomal DNA, the amplification of which produces a copy number gain of the cancer outcome marker, wherein the copy number gain is associated with a poor disease outcome.

5. The method of claim 4, wherein the cancer outcome marker is selected from the group consisting of Chr 19, 34.7 Mb—35.6 Mb; Chr 19, 38.9—40.7 Mb; Chr 17, 69.2—71.3 Mb; Chr 6, 70.8—71.1 Mb; Chr 12, 93.7 kb—1.9Mb; Chr 11, 64.3—64.8 Mb; Chr 19, 57.0—62.2 Mb; Chr 6, 39.1—39.9 Mb; Chr 11, 64.8—65.7 Mb; Chr 11, 61.4—64.3 Mb; Chr 17, 51.5—53.2 Mb; Chr 17, 43.5—44.9 Mb; Chr 2, 147.6—151.1 Mb; Chr 6, 123.7—135.6 Mb; Chr 8, 6.9—8.8 Mb; Chr 2, 159.9—161.4 Mb; Chr 2, 200.9—204.2 Mb; Chr 6, 36.3—36.7 Mb; Chr 2, 205.9—208.1 Mb; and Chr 1, 109.5—111.1 Mb.

6. The method of claim 4, wherein the cancer outcome marker is Chr 19, 34.7 Mb—35.6 Mb and includes nucleotide sequences encoding C19orf12; C19orf12; cyclin E1; PLEKHF1; POP4; and ZNF536. [Marker 1]

7. The method of claim 4, wherein the cancer outcome marker is Chr 19, 38.9—40.7 Mb and includes nucleotide sequences encoding ATP4A ATPase; CHST8, DMKN FAR1,2,3; FXYD1,3,5,7; GAPDHS; GPI; GPR42; GRAMD1A; HAMP; HPN; KCTD15 KIAA0355; KRTDAP; LGI4; LSM14A; LSR; MAG; PDCD2L; SAE2 SUMO1; SBSN; SCN1B; TMEM147,162; USF2; WTIP; and ZNF181,30,302,599,792. [Marker 2]

8. The method of claim 4, wherein the cancer outcome marker is Chr 17, 69.2—71.3 Mb and includes nucleotide sequences encoding ARMC7 (armadillo repeat containing 7); ATP5H ATP synthase (H⁺ transporting, mitochondrial F0 complex, subunit d); CASKIN2 (CASK interacting protein 2); CD300A (CD300a molecule); CD300C (CD300c molecule); CD300E (CD300e molecule); CD300LB (CD300 molecule-like family member b); CD300LF (CD300 molecule-like family member f); CDR2L (cerebellar degeneration-related protein 2-like); DNAI2 (dynein, axonemal, intermediate chain 2); (FADS6 fatty acid desaturase domain family, member 6); FDXR (ferredoxin reductase); GALK1 (galactokinase 1); GGA3 (golgi associated, gamma adaptin ear containing, ARF binding protein); GPR142 (G protein-coupled receptor 142); GPRC5C (G protein-coupled receptor, family C, group 5, member C); GRB2 (growth factor receptor-bound protein 2); GRIN2C (glutamate receptor, ionotropic, N-methyl D-aspartate 2C); H3F3B (H3 histone, family 3B (H3.3B)); HN1 (hematological and neurological

expressed 1 ICT1 (immature colon carcinoma transcript 1); ITGB4 (integrin, beta 4); KCTD2 (potassium channel tetramerisation domain containing 2); KIAA0195; KIF19 (kinesin family member 19); LLGL2 (lethal giant larvae homolog 2 (*Drosophila*)); LOC388419 (galectin-3-binding protein-like); MIF4GD (MIF4G domain containing); MRPS7 (mitochondrial ribosomal protein S7); NAT9 (N-acetyltransferase 9); NT5C (5', 3'-nucleotidase, cytosolic); NUP85 (nucleoporin 85kDa); OTOP2 (otopetrin 2); OTOP3 (otopetrin 3); RAB37 (RAB37, member RAS oncogene family); RECQL5 (RecQ protein-like 5); RPL38 (ribosomal protein L38); SAP30BP (SAP30 binding protein); SLC16A5 (solute carrier family 16, member 5 (monocarboxylic acid transporter 6)); SLC25A19 (solute carrier family 25 (mitochondrial thiamine pyrophosphate carrier), member 19); SLC9A3R1 (solute carrier family 9 (sodium/hydrogen exchanger), member 3 regulator 1); SUMO2 (SMT3 suppressor of mif two homolog 2 (*S. cerevisiae*)); TMEM104 (transmembrane protein 104); TTYH2 (tweety homolog 2 (*Drosophila*)); UNK (unkempt homolog (*Drosophila*)); and USH1G (Usher syndrome 1G (autosomal recessive)) [Marker 3].

9. The method of claim 4, wherein the cancer outcome marker is Chr 6, 70.8—71.1 Mb and includes nucleotide sequences encoding COL19A1 (collagen, type XIX, alpha 1), and COL9A1 (collagen, type IX, alpha 1). [Marker 4].

10. The method of claim 4, wherein the cancer outcome marker is Chr 12, 93.7 kb—1.9Mb and includes nucleotide sequences encoding ADIPOR2 (adiponectin receptor 2); B4GALNT3 (beta-1,4-N-acetyl-galactosaminyl transferase 3); CACNA2D4 (calcium channel, voltage-dependent, alpha 2/delta subunit 4); CCDC77 (coiled-coil domain containing 77); ERC1 (ELKS/RAB6-interacting/CAST family member 1); FBXL14 (F-box and leucine-rich repeat protein 14); HSN2 (hereditary sensory neuropathy, type II); IQSEC3 (IQ motif and Sec7 domain 3); JARID1A (jumonji, AT rich interactive domain 1A); LRTM2 (leucine-rich repeats and transmembrane domains 2); NINJ2 (ninjurin 2); RAD52 (RAD52 homolog (*S. cerevisiae*)); SLC6A12 (solute carrier family 6 (neurotransmitter transporter, betaine/GABA), member 12); SLC6A13 (solute carrier family 6 (neurotransmitter transporter, GABA), member 13); WNK1 (WNK lysine deficient protein kinase 1); and WNT5B (wingless-type MMTV integration site family, member 5B). [Marker 5].

11. The method of claim 4, wherein the cancer outcome marker is Chr 11, 64.3—64.8 Mb and includes nucleotide sequences encoding ARL2 (ADP-ribosylation factor-like 2); ATG2A ATG2 (autophagy related 2 homolog A (*S. cerevisiae*)); BATF2 (basic leucine zipper transcription factor, ATF-like 2); CAPN1 calpain 1, (mu/I) large subunit); CDC42BPG (CDC42 binding protein kinase gamma (DMPK-like)); CDCA5 (cell division cycle associated 5); EHD1 (EH-domain containing 1); FAU (Finkel-Biskis-Reilly murine sarcoma virus (FBR-MuSV) ubiquitously expressed); GPHA2 (glycoprotein hormone alpha 2); MAP4K2 (mitogen-activated protein kinase kinase kinase kinase 2); MEN1 (multiple endocrine neoplasia I); MRPL49 (mitochondrial ribosomal protein L49); NAALADL1 (N-acetylated alpha-linked acidic dipeptidase-like 1); POLA2 (polymerase (DNA directed), alpha 2 (70kD subunit)); PPP2R5B (protein phosphatase 2, regulatory subunit B', beta isoform); SAC3D1 SAC3 (domain containing 1); SLC22A20 (solute carrier family 22, member 20); SNX15 (sorting nexin 15); SPDYC (speedy homolog C (*Drosophila*)); SYVN1 (synovial apoptosis inhibitor 1, synoviolin); TM7SF2 (transmembrane 7 superfamily member 2); ZFPL1 (zinc finger protein-like 1); ZNHIT2 (zinc finger, HIT type 2); (hsa-mir-192); and (hsa-mir-194-2). [Marker 6].

12. The method of claim 4, wherein the cancer outcome marker is Chr 19, 57.0—62.2 Mb and includes nucleotide sequences encoding BIRC8 (baculoviral IAP repeat-containing 8); BRSK1 (BR serine/threonine kinase 1); CACNG6,7,8 (calcium channel, voltage-dependent, gamma subunit 6,7,8); CCDC106 (coiled-coil domain containing 106); CDC42EP5 CDC42 (effector protein (Rho GTPase binding) 5); CNOT3 CCR4-NOT (transcription complex, subunit 3); COX6B2 (cytochrome c oxidase subunit VIb polypeptide 2 (testis)); DPRX (divergent-paired related homeobox); EPN1 (epsin 1); EPS8L1 (EPS8-like 1); FCAR (Fc fragment of IgA, receptor for); FIZ1 (FLT3-interacting zinc finger 1); GALP (galanin-like peptide GP6 glycoprotein VI (platelet)); HSPBP1 (hsp70-interacting protein); IL11 (interleukin 11); ISOC2 (isochorismatase domain containing 2); KIR2DL1, KIR2DL4, KIR2DS4 KIR3DL1, KIR3DL3, KIR3DX1 (killer cell immunoglobulin-like receptor); LAIR1,2 (leukocyte-associated immunoglobulin-like receptor 1,2); LENG1,4,8,9 (leukocyte receptor cluster (LRC) member 1,4,8,9); LILRA2,3,4 (leukocyte immunoglobulin-like receptor, subfamily A (with TM domain), member 2,3,4); LILRB1,2,3,4,5 (leukocyte immunoglobulin-like receptor, subfamily B (with TM and ITIM domains), member 1,2,3,4,5); MYADM (myeloid-associated

differentiation marker); NAT14 (N-acetyltransferase 14); NCR1 (natural cytotoxicity triggering receptor 1); NDUFA3 (NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 3, 9kDa); NLRP2,4,5,7,8,9,11,12,13 (NLR family, pyrin domain containing 2,4,5,7,8,9,11,12,13.); OSCAR (osteoclast associated, immunoglobulin-like receptor); PEG3 (paternally expressed 3); PPP1R12C (protein phosphatase 1, regulatory (inhibitor) subunit 12C); PPP2R1A (protein phosphatase 2 (formerly 2A), regulatory subunit A, alpha isoform); PRKCG (protein kinase C, gamma); PRPF31 (PRP31 pre-mRNA processing factor 31 homolog (S. cerevisiae)); PTPRH (protein tyrosine phosphatase, receptor type, H); RDH13 (retinol dehydrogenase 13 (all-trans/9-cis)); RPL28 (ribosomal protein L28); RPS9 (ribosomal protein S9); SAPS1 (SAPS domain family, member 1); SUV420H2 (suppressor of variegation 4-20 homolog 2 (Drosophila)); SYT5 (synaptotagmin V); TFPT (TCF3 (E2A) fusion partner (in childhood Leukemia)); TMC4 (transmembrane channel-like 4); TMEM190 (transmembrane protein 190 TMEM86B transmembrane protein 86B); TNNT1 (troponin T type 1 (skeletal, slow)); TSEN34 (tRNA splicing endonuclease 34 homolog (S. cerevisiae)); TTYH1 (tweety homolog 1 (Drosophila)); U2AF2 (U2 small nuclear RNA auxiliary factor 2); UBE2S (ubiquitin-conjugating enzyme E2S); VN1R2 (vomeronasal 1 receptor 2); VN1R4 (vomeronasal 1 receptor 4); VSTM1 (V-set and transmembrane domain containing 1); ZNF28,160,320,321,331,347,350,415,432,444,468,470 (zinc finger protein 28,160,320,321,331,347,350,415,432,444,468,470); and miRNA's including hsa-mir-643, hsa-mir-512-1, hsa-mir-512-2, hsa-mir-498, hsa-mir-520e, hsa-mir-515-1, hsa-mir-519e, hsa-mir-520f, hsa-mir-515-2, hsa-mir-519c, hsa-mir-520a, hsa-mir-526b, hsa-mir-519b, hsa-mir-525, hsa-mir-523, hsa-mir-518f, hsa-mir-520b, hsa-mir-518b, hsa-mir-526a-1, hsa-mir-520c, hsa-mir-518c, hsa-mir-524, hsa-mir-517a, hsa-mir-519d, hsa-mir-521-2, hsa-mir-520d, hsa-mir-517b, hsa-mir-520g, hsa-mir-516-3, hsa-mir-526a-2, hsa-mir-518e, hsa-mir-518a-1, hsa-mir-518d, hsa-mir-516-4, hsa-mir-518a-2, hsa-mir-517c, hsa-mir-520h, hsa-mir-521-1, hsa-mir-522, hsa-mir-519a-1, hsa-mir-527, hsa-mir-516-1, hsa-mir-516-2, hsa-mir-519a-2, hsa-mir-371, hsa-mir-372, hsa-mir-373, hsa-mir-516a-1, hsa-mir-516a-2, hsa-mir-516b-1, hsa-mir-516b-2, hsa-mir-517a-1, hsa-mir-517a-2, hsa-mir-520c-1, hsa-mir-520c-2 [Marker 7].

13. The method of claim 4, wherein the cancer outcome marker is Chr 6, 39.1—39.9 Mb and includes nucleotide sequences encoding C6orf64 (chromosome 6 open reading frame 64); DNAH8 (dynein, axonemal, heavy chain 8); GLP1R (glucagon-like peptide 1 receptor); KCNK16 (potassium channel, subfamily K, member 16); KCNK17 (potassium channel, subfamily K, member 17); KCNK5 (potassium channel, subfamily K, member 5); KIF6 (kinesin family member 6). [Marker 8].

14. The method of claim 4, wherein the cancer outcome marker is Chr 11, 64.8—65.7 Mb and includes nucleotide sequences encoding BANF1 (barrier to autointegration factor 1); CATSPER1 (cation channel, sperm associated 1); CCDC85B (coiled-coil domain containing 85B); CDC42EP2CDC42 (effector protein (Rho GTPase binding) 2); CFL1 (cofilin 1 (non-muscle)); CST6 (cystatin E/M); CTSW (cathepsin W); DPF2 D4, (zinc and double PHD fingers family 2); DRAP1 (DR1-associated protein 1 (negative cofactor 2 alpha)); EFEMP2 (EGF-containing fibulin-like extracellular matrix protein 2); EHBP1L1 (EH domain binding protein 1-like 1); FAM89B (family with sequence similarity 89, member B); FIBP (fibroblast growth factor (acidic) intracellular binding protein); FOSL1 (FOS-like antigen 1); FRMD8 (FERM domain containing 8); GAL3ST3 (galactose-3-O-sulfotransferase 3); HTATIP (HIV-1 Tat interacting protein, 60kDa); KCNK7 (potassium channel, subfamily K, member 7); LTBP3 (latent transforming growth factor beta binding protein 3); MAP3K11 (mitogen-activated protein kinase kinase kinase 11); MGC11102 (hypothetical protein MGC11102); MUS81 MUS81 (endonuclease homolog (*S. cerevisiae*)); OVOL1 (ovo-like 1(*Drosophila*)); PACS1(phosphofurin acidic cluster sorting protein 1); PCNXL3 (pecanex-like 3 (*Drosophila*)); POLA2 (polymerase (DNA directed), alpha 2 (70kD subunit)); RELA (v-rel reticuloendotheliosis viral oncogene homolog A, nuclear factor of kappa light polypeptide gene enhancer in B-cells 3, p65 (avian)); RNASEH2C (ribonuclease H2, subunit C); SART1 (squamous cell carcinoma antigen recognized by T cells); SCYL1 (SCY1-like 1 (*S. cerevisiae*)); SF3B2 (splicing factor 3b, subunit 2, 145kDa); SIPA1 (signal-induced proliferation-associated gene 1); SLC25A45 (solute carrier family 25, member 45); SSSCA1 (Sjogren syndrome/scleroderma autoantigen 1); TIGD3 (tigger transposable element derived 3); TSGA10IP (testis specific, 10 interacting protein). [Marker 9].

15. The method of claim 4, wherein the cancer outcome marker is Chr 11, 61.4—64.3 Mb and includes nucleotide sequences encoding AHNAK (AHNAK nucleoprotein); ASRGL1 (asparaginase like 1); B3GAT3 (beta-1,3-glucuronyltransferase 3 (glucuronosyltransferase I)); BAD (BCL2-antagonist of cell death); BEST1 (bestrophin 1); BSCL2 (Bernardinelli-Seip congenital lipodystrophy 2 (seipin)); CCDC88B (coiled-coil domain containing 88B); CHRM1 (cholinergic receptor, muscarinic 1); COX8A (cytochrome c oxidase subunit 8A (ubiquitous)); DKFZP564J0863 (DKFZP564J0863 protein); DKFZP566E164 (DKFZP566E164 protein); DNAJC4 (DnaJ (Hsp40) homolog, subfamily C, member 4); EEF1G (eukaryotic translation elongation factor 1 gamma); EML3 (echinoderm microtubule associated protein like 3); ESRRA (estrogen-related receptor alpha); FADS2,3 (fatty acid desaturase 2,3); FKBP2 (FK506 binding protein 2, 13kDa); FLRT1 (fibronectin leucine rich transmembrane protein 1); FTH1 (ferritin, heavy polypeptide 1); GANAB (glucosidase, alpha; neutral AB); GNG3 (guanine nucleotide binding protein (G protein), gamma 3); GPR137 (G protein-coupled receptor 137); HRASLS2,3,5 (HRAS-like suppressor 2,3,5); INCENP (inner centromere protein antigens 135/155kDa); INTS5 (integrator complex subunit 5); KCNK4 (potassium channel, subfamily K, member 4); LGALS12 (lectin, galactoside-binding, soluble, 12 (galectin 12)); MACROD1 (MACRO domain containing 1); MARK2 (MAP/microtubule affinity-regulating kinase 2); MGC3196 (hypothetical protein MGC3196); MTA2 (metastasis associated 1 family, member 2); NAT11 (N-acetyltransferase 11); NRXN2 (neurexin 2); NUDT22 (nudix (nucleoside diphosphate linked moiety X)-type motif 22); NXF1 (nuclear RNA export factor 1); OTUB1 (OTU domain, ubiquitin aldehyde binding 1); PLCB3 (phospholipase C, beta 3 (phosphatidylinositol-specific)); POLR2G (polymerase (RNA) II (DNA directed) polypeptide G); PPP1R14B (protein phosphatase 1, regulatory (inhibitor) subunit 14B); PRDX5 (peroxiredoxin 5); PYGM (phosphorylase, glycogen; muscle (McArdle syndrome, glycogen storage disease type V)); RAB3IL1 (RAB3A interacting protein (rabin3)-like 1); RARRES3 (retinoic acid receptor responder (tazarotene induced) 3); RASGRP2 (RAS guanyl releasing protein 2 (calcium and DAG-regulated)); RCOR2 REST (corepressor 2); ROM1 (retinal outer segment membrane protein 1); RPS6KA4 (ribosomal protein S6 kinase, 90kDa, polypeptide 4 RTN3 reticulon 3); SCGB1A1, 1D1, 1D2, 1D4, 2A1, 2A1 (secretoglobin, family

SF1 splicing factor 1); SLC22A10, 11, 12, 6, 8, 9 (solute carrier family 22 (organic anion/cation transporter) SLC3A2 solute carrier family 3 (activators of dibasic and neutral amino acid transport), member 2); STIP1 (stress-induced-phosphoprotein 1 (Hsp70/Hsp90-organizing protein)); STX5 (syntaxin 5); TAF6L (TAF6-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa); TRPT1 (tRNA phosphotransferase 1); TTC9C (tetratricopeptide repeat domain 9C); TUT1 (terminal uridylyl transferase 1, U6 snRNA-specific URP2 UNC-112 related protein 2); UST6 (putative UST1-like organic anion transporter VEGFB vascular endothelial growth factor B); WDR74 (WD repeat domain 74); and ZBTB3 (zinc finger and BTB domain containing 3). [Marker 10].

16. The method of claim 4, wherein the cancer outcome marker is Chr 17, 51.5—53.2 Mb and includes nucleotide sequences encoding AKAP1 (A kinase (PRKA) anchor protein 1); ANKFN1 (ankyrin-repeat and fibronectin type III domain containing 1); C17orf67 (chromosome 17 open reading frame 67); COIL (coilin); DGKE (diacylglycerol kinase, epsilon 64kDa MSI2 musashi homolog 2 (Drosophila)); NOG (noggin); SCPEP1 (serine carboxypeptidase 1); and TRIM25 (tripartite motif-containing 25). [Marker 11].

17. The method of claim 4, wherein the cancer outcome marker is Chr 17, 43.5—44.9 Mb and includes nucleotide sequences encoding hsa-mir-10a; hsa-mir-196a-1; ABI3 (ABI gene family, member 3); ATP5G1 (ATP synthase, H⁺ transporting, mitochondrial F0 complex, subunit C1 (subunit 9)); B4GALNT2 (beta-1,4-N-acetyl-galactosaminyl transferase 2); CALCOCO2 (calcium binding and coiled-coil domain 2); CBX1 (chromobox homolog 1 (HP1 beta homolog Drosophila)); GIP (gastric inhibitory polypeptide); GNGT2 (guanine nucleotide binding protein (G protein), gamma transducing activity polypeptide 2); HOXB1,2,3,4,5,6,7,8,9,13 (homeobox B1,2,3,4,5,6,7,8,9,13); IGF2BP1 (insulin-like growth factor 2 mRNA binding protein 1); NFE2L1 (nuclear factor (erythroid-derived 2)-like 1); NGFR (nerve growth factor receptor (TNFR superfamily, member 16)); PHB (prohibitin PHOSPHO1 phosphatase, orphan 1); PRAC (small nuclear protein PRAC); SKAP1 (src kinase associated phosphoprotein 1); SNF8 (SNF8, ESCRT-II complex subunit, homolog (S. cerevisiae)); SNX11 (sorting nexin 11); TTL6 (tubulin tyrosine ligase-like family, member 6); UBE2Z (ubiquitin-conjugating enzyme E2Z); and ZNF652 (zinc finger protein 652). [Marker 12].

18. The method of claim 4, wherein the cancer outcome marker is Chr 2, 147.6—151.1 Mb and includes nucleotide sequences encoding ACVR2A (activin A receptor, type IIA); C2orf25 (chromosome 2 open reading frame 25); EPC2 (enhancer of polycomb homolog 2 (*Drosophila*)); KIF5C (kinesin family member 5C); LOC130576 (hypothetical protein LOC130576); LYPD6 (LY6/PLAUR domain containing 6); MBD5 (methyl-CpG binding domain protein 5); ORC4L (origin recognition complex, subunit 4-like (yeast)); RND3 (Rho family GTPase 3). [Marker 13].

19. The method of claim 4, wherein the cancer outcome marker is Chr 6, 123.7—135.6 Mb and includes nucleotide sequences encoding hsa-mir-588; AKAP7 (A kinase (PRKA) anchor protein 7); ALDH8A1 (aldehyde dehydrogenase 8 family, member A1); ARG1 (arginase, liver); ARHGAP18 (Rho GTPase activating protein 18); CTGF (connective tissue growth factor ECHDC1 enoyl Coenzyme A hydratase domain containing 1); ENPP1,3 (ectonucleotide pyrophosphatase/phosphodiesterase 1,3); EPB41L2 (erythrocyte membrane protein band 4.1-like 2); EYA4 (eyes absent homolog 4 (*Drosophila*)); HDHC2 (HD domain containing 2); HEY2 (hairy/enhancer-of-split related with YRPW motif 2); HINT3 (histidine triad nucleotide binding protein 3); KIAA1913 KIAA1913 LAMA2 (laminin, alpha 2 (merosin, congenital muscular dystrophy)); MED23 (mediator complex subunit 23); MOXD1 (monooxygenase, DBH-like 1 MYB v-myb myeloblastosis viral oncogene homolog (avian)); NCOA7 (nuclear receptor coactivator 7); NKAIN2 (Na⁺/K⁺ transporting ATPase interacting 2); OR2A4 (olfactory receptor, family 2, subfamily A, member 4); PTPRK (protein tyrosine phosphatase, receptor type, K. RNF146 ring finger protein 146); RNF217 (ring finger protein 217); RPS12 (ribosomal protein S12); SAMD3 (sterile alpha motif domain containing 3); SGK (serum/glucocorticoid regulated kinase); SLC2A12 (solute carrier family 2 (facilitated glucose transporter), member 12 STX7 syntaxin 7); TAAR1,2,5,6,8,9 (trace amine associated receptor 1,2,5,6,8,9); TBPL1 (TBP-like 1); TCF21 (transcription factor 21); TPD52L1 (tumor protein D52-like 1); TRDN (triadin TRMT11 tRNA methyltransferase 11 homolog (*S. cerevisiae*)); and VNN1,2,3 (vanin 1,2,3). [Marker 14].

20. The method of claim 4, wherein the cancer outcome marker is Chr 8, 6.9—8.8 Mb and includes nucleotide sequences encoding CLDN23 (claudin 23); DEFA5 (defensin, alpha 5, Paneth cell-specific); DEFB103B (defensin, beta 103B); DEFB104A (defensin, beta 104A);

DEFB104B (defensin, beta 104B); DEFB105B (defensin, beta 105B); DEFB106A (defensin, beta 106A); DEFB106B (defensin, beta 106B); DEFB107A (defensin, beta 107A); DEFB107B (defensin, beta 107B); DEFB4 (defensin, beta 4); MFHAS1 (malignant fibrous histiocytoma amplified sequence 1); PRAGMIN (homolog of rat pragma of Rnd2); SPAG11A (sperm associated antigen 11A); and SPAG11B (sperm associated antigen 11B). [Marker 15].

21. The method of claim 4, wherein the cancer outcome marker is Chr 2, 159.9—161.4 Mb and includes nucleotide sequences encoding BAZ2B (bromodomain adjacent to zinc finger domain, 2B); CD302 (CD302 molecule); ITGB6 (integrin, beta 6); LY75 (lymphocyte antigen 75); MARCH7 (membrane-associated ring finger (C3HC4) 7); PLA2R1 (phospholipase A2 receptor 1, 180kDa); and RBMS1 (RNA binding motif, single stranded interacting protein 1). [Marker 16].

22. The method of claim 4, wherein the cancer outcome marker is Chr 2, 200.9—204.2 Mb and includes nucleotide sequences encoding ABI2 (abl interactor 2); ALS2 (amyotrophic lateral sclerosis 2 (juvenile)); ALS2CR2, 4, 7, 8, 11, 12, 13 (amyotrophic lateral sclerosis 2 (juvenile) chromosome region, candidate 2, 4, 7, 8, 11, 12, 13); AOX1 (aldehyde oxidase 1); BMPR2 (bone morphogenetic protein receptor, type II (serine/threonine kinase)); BZW1 (basic leucine zipper and W2 domains 1); CASP10 (caspase 10, apoptosis-related cysteine peptidase); CASP8 (caspase 8, apoptosis-related cysteine peptidase); CFLAR (CASP8 and FADD-like apoptosis regulator); CLK1 (CDC-like kinase 1); CYP20A1 (cytochrome P450, family 20, subfamily A, polypeptide 1); FAM126B (family with sequence similarity 126, member B); FZD7 (frizzled homolog 7 (Drosophila) ICA1L islet cell autoantigen 1,69kDa-like); KCTD18 (potassium channel tetramerisation domain containing 18); LOC26010 (viral DNA polymerase-transactivated protein 6); MPP4 (membrane protein, palmitoylated 4 (MAGUK p55 subfamily member 4); NBEAL1 (neurobeachin-like 1); NDUFB3 (NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 3, 12kDa); NIF3L1 (NIF3 NGG1 interacting factor 3-like 1 (S. pombe)); NOP5/NOP58 (nucleolar protein NOP5/NOP58); ORC2L (origin recognition complex, subunit 2-like (yeast)); PPIL3 (peptidylprolyl isomerase (cyclophilin)-like 3); RAPH1 (Ras association (RalGDS/AF-6) and pleckstrin homology domains 1); SGOL2 (shugoshin-like 2 (S. pombe)); SUMO1 (SMT3 suppressor of mif two 3 homolog 1 (S. cerevisiae)); TRAK2 (trafficking protein, kinesin binding 2); and WDR12 (WD repeat domain 12). [Marker 17].

23. The method of claim 4, wherein the cancer outcome marker is Chr 6, 36.3—36.7 Mb and includes nucleotide sequences encoding BRPF3 (bromodomain and PHD finger containing, 3); DKFZp779B1540 (hypothetical protein DKFZp779B1540); ETV7 (ets variant gene 7 (TEL2 oncogene)); KCTD20 (potassium channel tetramerisation domain containing 20 PNPLA1 patatin-like phospholipase domain containing 1); PXT1 (peroxisomal, testis specific 1 SFRS3 splicing factor, arginine/serine-rich 3); and STK38 (serine/threonine kinase 38). [Marker 18].

24. The method of claim 4, wherein the cancer outcome marker is Chr 2, 205.9—208.1 Mb and includes nucleotide sequences encoding ADAM23 (ADAM metallopeptidase domain 23); CPO (carboxypeptidase O); DYT1 (dystrotelin); EEF1B2 (eukaryotic translation elongation factor 1 beta 2); FASTKD2 (FAST kinase domains 2); FLJ20309 (hypothetical protein FLJ20309); GPR1 (G protein-coupled receptor 1); KLF7 (Kruppel-like factor 7 (ubiquitous)); MDH1B (malate dehydrogenase 1B, NAD (soluble)); NDUFS1 (NADH dehydrogenase (ubiquinone) Fe-S protein 1, 75kDa (NADH-coenzyme Q reductase); NRP2 (neuropilin 2); PARD3B (par-3 partitioning defective 3 homolog B (C. elegans)); ZDBF2 (zinc finger, DBF-type containing 2); and hCG_1657980 (hCG1657980). [Marker 19].

25. The method of claim 4, wherein the cancer outcome marker is Chr 1, 109.5—111.1 Mb and includes nucleotide sequences encoding hsa-mir-197; AHCYL1 (S-adenosylhomocysteine hydrolase-like 1); ALX3 (aristaless-like homeobox 3); AMIGO1 (adhesion molecule with Ig-like domain 1); AMPD2 (adenosine monophosphate deaminase 2 (isoform L)); ATXN7L2 (ataxin 7-like 2); CELSR2 (cadherin, EGF LAG seven-pass G-type receptor 2 (flamingo homolog, Drosophila)); CSF1 (colony stimulating factor 1 (macrophage)); CYB561D1 (cytochrome b-561 domain containing 1); EPS8L3 (EPS8-like 3); FAM40A (family with sequence similarity 40, member A); GNAI3 (guanine nucleotide binding protein (G protein), alpha inhibiting activity polypeptide 3); GNAT2 (guanine nucleotide binding protein (G protein), alpha transducing activity polypeptide 2); GPR61 (G protein-coupled receptor 61); GSTM1,M2,M3,M4,M5 (glutathione S-transferase M1, M2 (muscle), M3 (brain), M4, M5); HBXIP (hepatitis B virus x interacting protein); KCNA2,3,4,10 (potassium voltage-gated channel, shaker-related subfamily, member 2,3,4,10); KIAA1324 (KIAA1324); MYBPHL (myosin binding protein H-like); PROK1 (prokineticin 1); PSMA5 (proteasome (prosome,

macropain) subunit, alpha type, 5); PSRC1 (proline/serine-rich coiled-coil 1); RBM15 (RNA binding motif protein 15); SARS (seryl-tRNA synthetase); SLC16A4 (solute carrier family 16, member 4 (monocarboxylic acid transporter 5)); SLC6A17 (solute carrier family 6, member 17); SORT1 (sortilin 1); SYPL2 (synaptophysin-like 2); UBL4B (ubiquitin-like 4B). [Marker 20].

26. The method of any of claims 1-3 wherein the cancer outcome marker is a region of chromosomal DNA, the deletion of which produces a copy number loss of the cancer outcome marker, wherein the copy number loss is associated with a poor disease outcome.

27. The method of claim 26, wherein the cancer outcome marker is selected from the group consisting of Chr 5, 62.9 – 67.8 Mb; Chr 5, 53.3 – 53.8 Mb; Chr 4, 105.8 – 107.2 Mb; Chr 16, 45.8 – 46.3 Mb; Chr 5, 50.7 – 52.0 Mb; Chr 5, 94.2 – 96.1 Mb; Chr 9, 36.1 – 37.0 Mb; Chr 5, 94.2 – 96.1 Mb; Chr14, 51.1 – 52.8 Mb; Chr 14, 61.5 – 68.6 Mb; Chr 9, 28.1 Mb; Chr 4, 43.7 – 44.2 Mb; Chr 5, 60.8 – 62.9 Mb; Chr 3, 120.0 – 121.1 Mb; Chr 4, 46.2 – 48.0 Mb; Chr 14, 38.9 – 40.0 Mb; Chr 4, 44.2 – 44.6 Mb; Chr 2, 213.7 – 214.3 Mb; Chr14, 43.9 – 46.6 Mb; Chr 14, 27.6 – 28.6 Mb; Chr 3, 98.0 – 98.3 Mb; Chr14, 55.2 – 60.0 Mb; Chr14, 48.7 – 51.1 Mb; Chr 4, 81.4 – 83.2 Mb; Chr 10, 51.9 – 54.2 Mb; Chr 5, 55.2 – 58.6 Mb; and Chr 5, 67.8 – 68.5 Mb.

28. The method of claim 26, wherein the cancer outcome marker is Chr 5, 62.9 – 67.8 Mb and includes nucleotide sequences encoding ADAMTS6 (ADAM metalloproteinase with thrombospondin type 1 motif, 6); CD180 (CD180 molecule); CENPK (centromere protein K); ERBB2IP (erbb2 interacting protein); FLJ13611 (hypothetical protein FLJ13611); HTR1A (5-hydroxytryptamine (serotonin) receptor 1A); MAST4 (microtubule associated serine/threonine kinase family member 4); NLN (neurolysin (metalloproteinase M3 family)); P18SRP (P18SRP protein); PIK3R1 (phosphoinositide-3-kinase, regulatory subunit 1 (p85 alpha)); PPWD1 (peptidylprolyl isomerase domain and WD repeat containing 1); RGS7BP (regulator of G-protein signaling 7 binding protein); RNF180 (ring finger protein 180); SDCCAG10 (serologically defined colon cancer antigen 10); SFRS12 (splicing factor, arginine/serine-rich 12); SGTB (small glutamine-rich tetratricopeptide repeat (TPR)-containing, beta0); and TRIM23 (tripartite motif-containing 23). [Deletion Marker 1].

29. The method of claim 26, wherein the cancer outcome marker is Chr 5, 53.3 – 53.8 Mb and includes nucleotide sequences encoding ARL15 (ADP-ribosylation factor-like 15); HSPB3 (heat shock 27kDa protein 3) and hsa-miR-581. [Deletion Marker 2].

30. The method of claim 26, wherein the cancer outcome marker is Chr 4, 105.8 – 107.2 Mb and includes nucleotide sequences encoding FLJ20184 (hypothetical protein FLJ20184); GSTCD (glutathione S-transferase, C-terminal domain containing); INTS12 (integrator complex subunit 12); KIAA1546 (KIAA1546); MGC16169 (hypothetical protein MGC16169); NPNT (nephronectin); and PPA2 (pyrophosphatase (inorganic) 2). [Deletion Marker 3].

31. The method of claim 26, wherein the cancer outcome marker is Chr 16, 45.8 – 46.3 Mb and includes nucleotide sequences encoding ITFG1 (integrin alpha FG-GAP repeat containing 1) and PHKB (phosphorylase kinase, beta). [Deletion Marker 4].

32. The method of claim 26, wherein the cancer outcome marker is Chr 5, 50.7 – 52.0 Mb and includes a nucleotide sequence encoding ISL1 (ISL LIM homeobox). [Deletion Marker 5].

33. The method of claim 26, wherein the cancer outcome marker is Chr 5, 94.2 – 96.1 Mb and includes nucleotide sequences encoding ARSK (arylsulfatase family, member K); CAST (calpastatin); ELL2 (elongation factor, RNA polymerase II, 2); FAM81B (family with sequence similarity 81, member B); GLRX (glutaredoxin (thioltransferase)); GPR150 (G protein-coupled receptor 150); KIAA0372 (KIAA0372); MCTP1 (multiple C2 domains, transmembrane 1); PCSK1 (proprotein convertase subtilisin/kexin type 1); RFESD (Rieske (Fe-S) domain containing); RHOBTB3 (Rho-related BTB domain containing 3); SPATA9 (spermatogenesis associated 9); and hsa-miR-583. [Deletion Marker 6].

34. The method of claim 26, wherein the cancer outcome marker is Chr 9, 36.1 – 37.0 Mb and includes nucleotide sequences encoding C9orf19 (chromosome 9 open reading frame 19); CCIN (calicin); CLTA (clathrin, light chain (Lca)); GNE (glucosamine (UDP-N-acetyl)-2-epimerase/N-acetylmannosamine kinase); MELK (maternal embryonic leucine zipper kinase); PAX5 (paired box 5); RECK (reversion-inducing-cysteine-rich protein with kazal motifs); RNF38 (ring finger protein 38). [Deletion Marker 7].

35. The method of claim 26, wherein the cancer outcome marker is Chr 5, 94.2 – 96.1 Mb and includes nucleotide sequences encoding ARSK (arylsulfatase family, member K); CAST (calpastatin); ELL2 (elongation factor, RNA polymerase II, 2); FAM81B (family with sequence similarity 81, member B); GLRX (glutaredoxin (thioltransferase)); GPR150 (G

protein-coupled receptor 150); KIAA0372 (KIAA0372); MCTP1 (multiple C2 domains, transmembrane 1); PCSK1 (proprotein convertase subtilisin/kexin type 1); RFESD (Rieske (Fe-S) domain containing); RHOBTB3 (Rho-related BTB domain containing 3); SPATA9 (spermatogenesis associated 9). [Deletion Marker 8].

36. The method of claim 26, wherein the cancer outcome marker is Chr14, 51.1 – 52.8 Mb and includes nucleotide sequences encoding C14orf166 (chromosome 14 open reading frame 166); DDHD1 (DDHD domain containing 1); ERO1L (ERO1-like (*S. cerevisiae*)); FRMD6 (FERM domain containing 6); GNG2 (guanine nucleotide binding protein (G protein), gamma 2); GNPAT1 (glucosamine-phosphate N-acetyltransferase 1); GPR137C (G protein-coupled receptor 137C); NID2 (nidogen 2 (osteonidogen)); PLEKHC1 (pleckstrin homology domain containing, family C (with FERM domain) member 1); PSMC6 (proteasome (prosome, macropain) 26S subunit, ATPase, 6); PTGDR (prostaglandin D2 receptor (DP)); PTGER2 (prostaglandin E receptor 2 (subtype EP2), 53kDa); STYX (serine/threonine/tyrosine interacting protein); TXNDC16 (thioredoxin domain containing 16). [Deletion Marker 9].

37. The method of claim 26, wherein the cancer outcome marker is Chr 14, 61.5 – 68.6 Mb and includes nucleotide sequences encoding ACTN1 (actinin, alpha 1); AKAP5 (A kinase (PRKA) anchor protein 5); ARG2 (arginase, type II); ATP6V1D (ATPase, H⁺ transporting, lysosomal 34kDa, V1 subunit D); C14orf50 (chromosome 14 open reading frame 50); C14orf54 (chromosome 14 open reading frame 54); C14orf83 (chromosome 14 open reading frame 83); CHURC1 (churchill domain containing 1); EIF2S1 (eukaryotic translation initiation factor 2, subunit 1 alpha, 35kDa); ESR2 (estrogen receptor 2 (ER beta)); FLJ39779 (FLJ39779 protein); FNTB (farnesyltransferase, CAAX box, beta); FUT8 (fucosyltransferase 8 (alpha (1,6) fucosyltransferase)); GPHB5 (glycoprotein hormone beta 5); GPHN (gephyrin); GPX2 (glutathione peroxidase 2 (gastrointestinal)); HSPA2 (heat shock 70kDa protein 2); KCNH5 (potassium voltage-gated channel, subfamily H (eag-related), member 5); MAX MYC (associated factor X); MPP5 (membrane protein, palmitoylated 5 (MAGUK p55 subfamily member 5)); MTHFD1 (methylenetetrahydrofolate dehydrogenase (NADP⁺ dependent) 1, methenyltetrahydrofolate cyclohydrolase, formyltetrahydrofolate synthetase); PIGH (phosphatidylinositol glycan anchor biosynthesis, class H); PLEK2 (pleckstrin 2); PLEKHG3 (pleckstrin homology domain containing, family G (with RhoGef domain) member 3);

PLEKHH1 (pleckstrin homology domain containing, family H (with MyTH4 domain) member 1); PPP2R5E (protein phosphatase 2, regulatory subunit B', epsilon isoform); RAB15 (RAB15, member RAS oncogene family); RAD51L1 (RAD51-like 1 (*S. cerevisiae*)); RDH11 (retinol dehydrogenase 11 (all-trans/9-cis/11-cis)); RDH12 (retinol dehydrogenase 12 (all-trans/9-cis/11-cis)); RHOJ (ras homolog gene family, member J); SGPP1 (sphingosine-1-phosphate phosphatase 1); SPTB (spectrin, beta, erythrocytic (includes spherocytosis, clinical type I)); SYNE2 (spectrin repeat containing, nuclear envelope 2); SYT16 (synaptotagmin XVI); VTI1B (vesicle transport through interaction with t-SNAREs homolog 1B (yeast)); WDR22 (WD repeat domain 22); WDR89 (WD repeat domain 89); ZBTB1 (zinc finger and BTB domain containing 1); ZBTB25 (zinc finger and BTB domain containing 25); ZFP36L1 (zinc finger protein 36, C3H type-like 1); ZFYVE26 (zinc finger, FYVE domain containing 26); and hsa-miR-625. [Deletion Marker 10]

38. The method of claim 26, wherein the cancer outcome marker is Chr 9, 28.1 Mb and includes a nucleotide sequence encoding LINGO2 (leucine rich repeat and Ig domain containing 2). [Deletion Marker 11]

39. The method of claim 26, wherein the cancer outcome marker is Chr 4, 43.7 – 44.2 Mb and includes a nucleotide sequence encoding KCTD8 (potassium channel tetramerisation domain containing 8). [Deletion Marker 12]

40. The method of claim 26, wherein the cancer outcome marker is Chr 5, 60.8 – 62.9 Mb and includes nucleotide sequences encoding DIMT1L DIM1 (dimethyladenosine transferase 1-like (*S. cerevisiae*)); FLJ37543 (hypothetical protein FLJ37543); IPO11 (importin 11); ISCA1L (iron-sulfur cluster assembly 1 homolog (*S. cerevisiae*)-like); KIF2A (kinesin heavy chain member 2A). [Deletion Marker 13]

41. The method of claim 26, wherein the cancer outcome marker is Chr 3, 120.0 – 121.1 Mb and includes nucleotide sequences encoding ADPRH (ADP-ribosylarginine hydrolase); B4GALT4 (UDP-Gal:betaGlcNAc beta 1,4- galactosyltransferase, polypeptide 4); C3orf1 (chromosome 3 open reading frame 1); C3orf15 (chromosome 3 open reading frame 15); C3orf30 (chromosome 3 open reading frame 30); CD80 (CD80 molecule); CDGAP (Cdc42 GTPase-activating protein); COX17 (COX17 cytochrome c oxidase assembly homolog (*S. cerevisiae*)); GSK3B (glycogen synthase kinase 3 beta); IGSF11 (immunoglobulin superfamily,

member 110); KTELC1 (KTEL (Lys-Tyr-Glu-Leu) containing 1); NR1I2 (nuclear receptor subfamily 1, group I, member 2); PLA1A (phospholipase A1 member A); POPDC2 (popeye domain containing 2); TMEM39A (transmembrane protein 39A); and UPK1B (uroplakin 1B). [Deletion Marker 14]

42. The method of claim 26, wherein the cancer outcome marker is Chr 4, 46.2 – 48.0 Mb and includes nucleotide sequences encoding ATP10D (ATPase, class V, type 10D); CNGA1 (cyclic nucleotide gated channel alpha 1); COMMD8 (COMM domain containing 8); CORIN (corin, serine peptidase); COX7B2 (cytochrome c oxidase subunit VIIb2); GABRA4 (gamma-aminobutyric acid (GABA) A receptor, alpha 4); GABRB1 (gamma-aminobutyric acid (GABA) A receptor, beta 1); NFXL1 (nuclear transcription factor, X-box binding-like 1); NPAL1 (NIPA-like domain containing 1); TEC (tec protein tyrosine kinase); and TXK (TXK tyrosine kinase). [Deletion Marker 15]

43. The method of claim 26, wherein the cancer outcome marker is Chr 14, 38.9 – 40.0 Mb and includes a nucleotide sequence encoding FBXO33 (F-box protein 33). [Deletion Marker 16]

44. The method of claim 26, wherein the cancer outcome marker is Chr 4, 44.2 – 44.6 Mb and includes nucleotide sequences encoding GNPDA2 (glucosamine-6-phosphate deaminase 2); GUF1 (GUF1 GTPase homolog (*S. cerevisiae*)); and YIPF7 (Yip1 domain family, member 7). [Deletion Marker 17]

45. The method of claim 26, wherein the cancer outcome marker is Chr 2, 213.7 – 214.3 Mb and includes nucleotide sequences encoding IKZF2 (IKAROS family zinc finger 2 (*Helios*)); SPAG16 (sperm associated antigen 16). [Deletion Marker 18]

46. The method of claim 26, wherein the cancer outcome marker is Chr 14, 43.9 – 46.6 Mb and includes nucleotide sequences encoding C14orf106 (chromosome 14 open reading frame 106); C14orf155 (chromosome 14 open reading frame 155); C14orf28 (chromosome 14 open reading frame 28); FANCM (Fanconi anemia, complementation group M); FKBP3 (FK506 binding protein 3, 25kDa); KIAA0423 (KIAA0423); KLHL28 (kelch-like 28 (*Drosophila*)); MDGA2 (MAM domain containing glycosylphosphatidylinositol anchor 2); PRPF39 (PRP39 pre-mRNA processing factor 39 homolog (*S. cerevisiae*)); RPL10L (ribosomal protein L10-like). [Deletion Marker 19]

47. The method of claim 26, wherein the cancer outcome marker is Chr 14, 27.6 – 28.6 Mb and includes a nucleotide sequence encoding FOXG1 (forkhead box G1). [Deletion Marker 20]

48. The method of claim 26, wherein the cancer outcome marker is Chr 3, 98.0 – 98.3 Mb and includes nucleotide sequences encoding EPHA6 (EPH receptor A6). [Deletion Marker 21]

49. The method of claim 26, wherein the cancer outcome marker is Chr14, 55.2 – 60.0 Mb and includes nucleotide sequences encoding ACTR10 (actin-related protein 10 homolog (*S. cerevisiae*)); ARID4A (AT rich interactive domain 4A (RBP1-like)); C14orf100 (chromosome 14 open reading frame 100); C14orf101 (chromosome 14 open reading frame 101); C14orf105 (chromosome 14 open reading frame 105); C14orf108 (chromosome 14 open reading frame 108); C14orf135 (chromosome 14 open reading frame 135); C14orf149 (chromosome 14 open reading frame 149); C14orf37 (chromosome 14 open reading frame 37); C14orf39 (chromosome 14 open reading frame 39); DAAM1 (dishevelled associated activator of morphogenesis 1); DACT1 (dapper, antagonist of beta-catenin, homolog 1 (*Xenopus laevis*)); DHRS7 (dehydrogenase/reductase (SDR family) member 7); EXOC5 (exocyst complex component 5); GPR135 (G protein-coupled receptor 135); KIAA0586 (KIAA0586); NAT12 (N-acetyltransferase 12); OTX2 (orthodenticle homeobox 2); PELI2 (pellino homolog 2 (*Drosophila*)); PPM1A (protein phosphatase 1A (formerly 2C), magnesium-dependent, alpha isoform); PSMA3 (proteasome (prosome, macropain) subunit, alpha type, 3); RTN1 (reticulon 1); SLC35F4 (solute carrier family 35, member F4); TIMM9 (translocase of inner mitochondrial membrane 9 homolog (yeast)); UNQ9438 (TIMM). [Deletion Marker 22]

50. The method of claim 26, wherein the cancer outcome marker is Chr14, 48.7 – 51.1 Mb and includes nucleotide sequences encoding ABHD12B (abhydrolase domain containing 12B); ARF6 (ADP-ribosylation factor 6); ATP5S (ATP synthase, H⁺ transporting, mitochondrial F0 complex, subunit s (factor B)); C14orf104 (chromosome 14 open reading frame 104); C14orf138 (chromosome 14 open reading frame 138); CDKL1 (cyclin-dependent kinase-like 1 (CDC2-related kinase)); FRMD6 (FERM domain containing 6); KLHDC1 (kelch domain containing 1); KLHDC2 (kelch domain containing 2); L2HGDH (L-2-hydroxyglutarate dehydrogenase); LOC196913 (hypothetical protein LOC196913); LOC283551 (hypothetical

protein LOC283551); MAP4K5 (mitogen-activated protein kinase kinase kinase 5); MGAT2 (mannosyl (alpha-1,6-)-glycoprotein beta-1,2-N-acetylglucosaminyltransferase); NIN (ninein (GSK3B interacting protein)); POLE2 (polymerase (DNA directed), epsilon 2 (p59 subunit)); PPIL5 (peptidylprolyl isomerase (cyclophilin)-like 5); PYGL (phosphorylase, glycogen; liver (Hers disease, glycogen storage disease type VI)); RPL36AL (ribosomal protein L36a-like); RPS29 (ribosomal protein S29). [Deletion Marker 23]

51. The method of claim 26, wherein the cancer outcome marker is Chr 4, 81.4 – 83.2 Mb and includes nucleotide sequences encoding BMP3 (bone morphogenetic protein 3 (osteogenic)); C4orf22 (chromosome 4 open reading frame 22); FGF5 (fibroblast growth factor 5); PRKG2 (protein kinase, cGMP-dependent, type II); RASGEF1B (RasGEF domain family, member 1B). [Deletion Marker 24]

52. The method of claim 26, wherein the cancer outcome marker is Chr 10, 51.9 – 54.2 Mb and includes nucleotide sequences encoding ACF (apobec-1 complementation factor); ASAH2B (N-acylsphingosine amidohydrolase (non-lysosomal ceramidase) 2B); CSTF2T (cleavage stimulation factor, 3' pre-RNA, subunit 2, 64kDa, tau variant); DKK1 (dickkopf homolog 1 (*Xenopus laevis*)); MBL2 (mannose-binding lectin (protein C) 2, soluble (opsonic defect)); PRKG1 (protein kinase, cGMP-dependent, type I); SGMS1 (sphingomyelin synthase 1); hsa-miR-605. [Deletion Marker 25]

53. The method of claim 26, wherein the cancer outcome marker is Chr 5, 55.2 – 58.6 Mb and includes nucleotide sequences encoding ANKRD55 (ankyrin repeat domain 55); C5orf29(chromosome 5 open reading frame 29); C5orf35 (chromosome 5 open reading frame 35); DKFZp686D0972 (similar to RIKEN cDNA 4732495G21 gene); GPBP1 (GC-rich promoter binding protein 10); IL31RA (interleukin 31 receptor A); IL6ST (interleukin 6 signal transducer (gp130, oncostatin M receptor)); MAP3K1 (mitogen-activated protein kinase kinase kinase 1); MIER3 (mesoderm induction early response 1, family member 3); PDE4D (phosphodiesterase 4D, cAMP-specific (phosphodiesterase E3 dunce homolog, *Drosophila*)); PLK2 (polo-like kinase 2 (*Drosophila*)); RAB3C RAB3C (member RAS oncogene family). [Deletion Marker 26]

54. The method of claim 26, wherein the cancer outcome marker is Chr 5, 67.8 – 68.5 Mb and includes nucleotide sequences encoding CCNB1 (cyclin B1) and SLC30A5 (solute carrier family 30 (zinc transporter), member 5). [Deletion Marker 27]

55. The method of any of claims 1-4, and 26 wherein the test sample comprises a tissue sample.

56. The method of claim 55, wherein the tissue sample comprises a blood sample, a tumor tissue or a suspected tumor tissue, a thin layer cytological sample, a fine needle aspirate sample, a lung wash sample, a pleural effusion sample, a fresh frozen tissue sample, a paraffin embedded tissue sample or an extract or processed sample produced from any thereof.

57. The method of claim 55, wherein the tissue sample comprises a lung tissue sample or a peripheral blood sample comprising circulating tumor cells.

58. The method of any of claims 1-4, and 26 wherein the determining step (b) is performed by *in situ* hybridization.

59. The method of claim 58, wherein the *in situ* hybridization is performed with a nucleic acid probe that is fluorescently labeled.

60. The method of claim 58, wherein the *in situ* hybridization is performed with at least two nucleic acid probes.

61. The method of claim 58, wherein the *in situ* hybridization is performed with a peptide nucleic acid probe.

62. The method of any of claims 1-4, and 26 wherein the determining step (b) is performed by polymerase chain reaction.

63. The method of any of claims 1-4, and 26 wherein the determining step (b) is performed by a nucleic acid sequencing assay.

64. The method of any of claims 1-4, and 26 wherein the determining step (b) is performed by a nucleic acid microarray assay.

65. The method of any of claims 1-4, and 26 wherein the lung cancer is non-small-cell lung cancer.

66. The method of any of claims 1-4, and 26 wherein the cancer is selected from the group consisting of squamous cell carcinoma, large cell carcinoma and adenocarcinoma.

67. The method of any of claims 1-4, and 26 wherein the patient is being treated with chemotherapy, radiation, surgery or any combination thereof.

68. A method of selecting a treatment for a patient suffering from lung cancer, the method comprising the steps of:

- a) providing a test sample from the patient wherein treatment with a chemotherapy agent is at least one treatment option for the patient;
- b) determining a copy number of the cancer outcome marker in the test sample;
- c) comparing the copy number of the cancer outcome marker in the test sample against a baseline copy number of two thereby determining the presence or absence of a copy number change for the cancer outcome marker in the test sample; and
- d) determining a chemotherapy treatment regimen based on the comparison in step c).

69. The method of claim 68 wherein determining a treatment regimen based on the comparison in step c) comprises selecting a chemotherapy agent and determining a frequency of chemotherapy treatment when a copy number change is present for the cancer outcome marker.

70. The method of claim 68, wherein the cancer outcome marker is a region of chromosomal DNA, the amplification of which produces a copy number gain of the cancer outcome marker, wherein the copy number gain is associated with a poor disease outcome.

71. The method of claim 70, wherein the cancer outcome marker is selected from the group consisting of Chr 19, 34.7 Mb—35.6 Mb; Chr 19, 38.9—40.7 Mb; Chr 17, 69.2—71.3 Mb; Chr 6, 70.8—71.1 Mb; Chr 12, 93.7 kb—1.9Mb; Chr 11, 64.3—64.8 Mb; Chr 19, 57.0—62.2 Mb; Chr 6, 39.1—39.9 Mb; Chr 11, 64.8—65.7 Mb; Chr 11, 61.4—64.3 Mb; Chr 17, 51.5—53.2 Mb; Chr 17, 43.5—44.9 Mb; Chr 2, 147.6—151.1 Mb; Chr 6, 123.7—135.6 Mb; Chr 8, 6.9—8.8 Mb; Chr 2, 159.9—161.4 Mb; Chr 2, 200.9—204.2 Mb; Chr 6, 36.3—36.7 Mb; Chr 2, 205.9—208.1 Mb; and Chr 1, 109.5—111.1 Mb.

72. The method of claim 68, wherein the cancer outcome marker is a region of chromosomal DNA, the deletion of which produces a copy number loss of the cancer outcome marker, wherein the copy number loss is associated with a poor disease outcome.

73. The method of claim 72, wherein the cancer outcome marker is selected from the group consisting of Chr 5, 62.9 – 67.8 Mb; Chr 5, 53.3 – 53.8 Mb; Chr 4, 105.8 – 107.2 Mb; Chr 16, 45.8 – 46.3 Mb; Chr 5, 50.7 – 52.0 Mb; Chr 5, 94.2 – 96.1 Mb; Chr 9, 36.1 – 37.0 Mb; Chr 5, 94.2 – 96.1 Mb; Chr 14, 51.1 – 52.8 Mb; Chr 14, 61.5 – 68.6 Mb; Chr 9, 28.1 Mb; Chr 4, 43.7 – 44.2 Mb; Chr 5, 60.8 – 62.9 Mb; Chr 3, 120.0 – 121.1 Mb; Chr 4, 46.2 – 48.0 Mb; Chr 14, 38.9 – 40.0 Mb; Chr 4, 44.2 – 44.6 Mb; Chr 2, 213.7 – 214.3 Mb; Chr 14, 43.9 – 46.6 Mb; Chr 14, 27.6

– 28.6 Mb; Chr 3, 98.0 – 98.3 Mb; Chr14, 55.2 – 60.0 Mb; Chr14, 48.7 – 51.1 Mb; Chr 4, 81.4 – 83.2 Mb; Chr 10, 51.9 – 54.2 Mb; Chr 5, 55.2 – 58.6 Mb; and Chr 5, 67.8 – 68.5 Mb.

74. The method of claim 68, wherein the test sample comprises a tissue sample.

75. The method of claim 74, wherein the tissue sample comprises a blood sample, a tumor tissue or a suspected tumor tissue, a thin layer cytological sample, a fine needle aspirate sample, a lung wash sample, a pleural effusion sample, a fresh frozen tissue sample, a paraffin embedded tissue sample or an extract or processed sample produced from any thereof.

76. The method of claim 74, wherein the tissue sample comprises a lung tissue sample or a peripheral blood sample comprising circulating tumor cells.

77. The method of claim 68, wherein the determining step (b) is performed by *in situ* hybridization.

78. The method of claim 77, wherein the *in situ* hybridization is performed with a nucleic acid probe that is fluorescently labeled.

79. The method of claim 77, wherein the *in situ* hybridization is performed with at least two nucleic acid probes.

80. The method of claim 77, wherein the *in situ* hybridization is performed with a peptide nucleic acid probe.

81. The method of claim 68, wherein the determining step (b) is performed by polymerase chain reaction.

82. The method of any of claim 68, wherein the determining step (b) is performed by a nucleic acid sequencing assay.

83. The method of claim 68, wherein the determining step (b) is performed by a nucleic acid microarray assay.

84. The method of claim 68, wherein the lung cancer is non-small-cell lung cancer.

85. The method of claim 68, wherein the cancer is selected from the group consisting of squamous cell carcinoma, large cell carcinoma and adenocarcinoma.

86. The method of claim 68, wherein the patient is also being treated with radiation or surgery or a combination thereof.

87. A method of classifying a patient as having a lung cancer that is resistant to treatment comprising the steps of:

- a) providing a test sample from a patient;
- b) determining a copy number for the cancer outcome marker;
- c) comparing the copy number for the cancer outcome marker in the test sample against a baseline copy number of two for the cancer outcome marker to determine the presence or absence of a copy number change in the cancer outcome marker in the patient; and
- d) classifying the patient as having a lung cancer that is resistant to treatment based on the presence of a copy number change in the cancer outcome marker.

88. The method of claim 87, wherein the cancer outcome marker is a region of chromosomal DNA, the amplification of which produces a copy number gain of the cancer outcome marker, wherein the copy number gain is associated with a poor disease outcome.

89. The method of claim 88, wherein the cancer outcome marker is selected from the group consisting of Chr 19, 34.7 Mb—35.6 Mb; Chr 19, 38.9—40.7 Mb; Chr 17, 69.2—71.3 Mb; Chr 6, 70.8—71.1 Mb; Chr 12, 93.7 kb—1.9Mb; Chr 11, 64.3—64.8 Mb; Chr 19, 57.0—62.2 Mb; Chr 6, 39.1—39.9 Mb; Chr 11, 64.8—65.7 Mb; Chr 11, 61.4—64.3 Mb; Chr 17, 51.5—53.2 Mb; Chr 17, 43.5—44.9 Mb; Chr 2, 147.6—151.1 Mb; Chr 6, 123.7—135.6 Mb; Chr 8, 6.9—8.8 Mb; Chr 2, 159.9—161.4 Mb; Chr 2, 200.9—204.2 Mb; Chr 6, 36.3—36.7 Mb; Chr 2, 205.9—208.1 Mb; and Chr 1, 109.5—111.1 Mb.

90. The method of claim 87, wherein the cancer outcome marker is a region of chromosomal DNA, the deletion of which produces a copy number loss of the cancer outcome marker, wherein the copy number loss is associated with a poor disease outcome.

91. The method of claim 90, wherein the cancer outcome marker is selected from the group consisting of Chr 5, 62.9 – 67.8 Mb; Chr 5, 53.3 – 53.8 Mb; Chr 4, 105.8 – 107.2 Mb; Chr 16, 45.8 – 46.3 Mb; Chr 5, 50.7 – 52.0 Mb; Chr 5, 94.2 – 96.1 Mb; Chr 9, 36.1 – 37.0 Mb; Chr 5, 94.2 – 96.1 Mb; Chr14, 51.1 – 52.8 Mb; Chr 14, 61.5 – 68.6 Mb; Chr 9, 28.1 Mb; Chr 4, 43.7 – 44.2 Mb; Chr 5, 60.8 – 62.9 Mb; Chr 3, 120.0 – 121.1 Mb; Chr 4, 46.2 – 48.0 Mb; Chr 14, 38.9 – 40.0 Mb; Chr 4, 44.2 – 44.6 Mb; Chr 2, 213.7 – 214.3 Mb; Chr14, 43.9 – 46.6 Mb; Chr 14, 27.6 – 28.6 Mb; Chr 3, 98.0 – 98.3 Mb; Chr14, 55.2 – 60.0 Mb; Chr14, 48.7 – 51.1 Mb; Chr 4, 81.4 – 83.2 Mb; Chr 10, 51.9 – 54.2 Mb; Chr 5, 55.2 – 58.6 Mb; and Chr 5, 67.8 – 68.5 Mb.

92. The method of claim 87, wherein the test sample comprises a tissue sample.

93. The method of claim 92, wherein the tissue sample comprises a blood sample, a tumor tissue or a suspected tumor tissue, a thin layer cytological sample, a fine needle aspirate sample, a lung wash sample, a pleural effusion sample, a fresh frozen tissue sample, a paraffin embedded tissue sample or an extract or processed sample produced from any of a peripheral blood sample.

94. The method of claim 92, wherein the tissue sample comprises a lung tissue sample or a peripheral blood sample comprising circulating tumor cells.

95. The method of claim 86, wherein the determining step (b) is performed by *in situ* hybridization.

96. The method of claim 94, wherein the *in situ* hybridization is performed with a nucleic acid probe that is fluorescently labeled.

97. The method of claim 94, wherein the *in situ* hybridization is performed with at least two nucleic acid probes.

98. The method of claim 94, wherein the *in situ* hybridization is performed with a peptide nucleic acid probe.

99. The method of claim 87, wherein the determining step (b) is performed by polymerase chain reaction.

100. The method of claim 87, wherein the determining step (b) is performed by a nucleic acid sequencing assay.

101. The method of claim 87, wherein the determining step (b) is performed by a nucleic acid microarray assay.

102. The method of claim 87, wherein the lung cancer is non-small-cell lung cancer.

103. The method of claim 87, wherein the cancer is selected from the group consisting of squamous cell carcinoma, large cell carcinoma and adenocarcinoma.

104. The method of claim 87, wherein the patient is being treated with chemotherapy, radiation, surgery or any combination thereof.

105. A kit comprising:

- a) reagents for determining the presence or absence of a copy number change for the cancer outcome marker; and
- b) instructions for performing the test.

106. The kit of claim 104, wherein the reagents to determine the presence or absence of a copy number change for the cancer outcome marker comprise detectably-labeled polynucleotides that hybridize to at least a portion of the cancer outcome marker.

107. The kit of claim 105, wherein the cancer outcome marker is a region of chromosomal DNA, the amplification of which produces a copy number gain of the cancer outcome marker, wherein the copy number gain is associated with a poor disease outcome.

108. The kit of claim 106, wherein the cancer outcome marker is selected from the group consisting of Chr 19, 34.7 Mb—35.6 Mb; Chr 19, 38.9—40.7 Mb; Chr 17, 69.2—71.3 Mb; Chr 6, 70.8—71.1 Mb; Chr 12, 93.7 kb—1.9Mb; Chr 11, 64.3—64.8 Mb; Chr 19, 57.0—62.2 Mb; Chr 6, 39.1—39.9 Mb; Chr 11, 64.8—65.7 Mb; Chr 11, 61.4—64.3 Mb; Chr 17, 51.5—53.2 Mb; Chr 17, 43.5—44.9 Mb; Chr 2, 147.6—151.1 Mb; Chr 6, 123.7—135.6 Mb; Chr 8, 6.9—8.8 Mb; Chr 2, 159.9—161.4 Mb; Chr 2, 200.9—204.2 Mb; Chr 6, 36.3—36.7 Mb; Chr 2, 205.9—208.1 Mb; and Chr 1, 109.5—111.1 Mb.

109. The kit of claim 105, wherein the cancer outcome marker is a region of chromosomal DNA, the deletion of which produces a copy number loss of the cancer outcome marker, wherein the copy number loss is associated with a poor disease outcome.

110. The kit of claim 108, wherein the cancer outcome marker is selected from the group consisting of Chr 5, 62.9 – 67.8 Mb; Chr 5, 53.3 – 53.8 Mb; Chr 4, 105.8 – 107.2 Mb; Chr 16, 45.8 – 46.3 Mb; Chr 5, 50.7 – 52.0 Mb; Chr 5, 94.2 – 96.1 Mb; Chr 9, 36.1 – 37.0 Mb; Chr 5, 94.2 – 96.1 Mb; Chr14, 51.1 – 52.8 Mb; Chr 14, 61.5 – 68.6 Mb; Chr 9, 28.1 Mb; Chr 4, 43.7 – 44.2 Mb; Chr 5, 60.8 – 62.9 Mb; Chr 3, 120.0 – 121.1 Mb; Chr 4, 46.2 – 48.0 Mb; Chr 14, 38.9 – 40.0 Mb; Chr 4, 44.2 – 44.6 Mb; Chr 2, 213.7 – 214.3 Mb; Chr14, 43.9 – 46.6 Mb; Chr 14, 27.6 – 28.6 Mb; Chr 3, 98.0 – 98.3 Mb; Chr14, 55.2 – 60.0 Mb; Chr14, 48.7 – 51.1 Mb; Chr 4, 81.4 – 83.2 Mb; Chr 10, 51.9 – 54.2 Mb; Chr 5, 55.2 – 58.6 Mb; and Chr 5, 67.8 – 68.5 Mb.