

DETECTION OF COLORECTAL CANCER

DESCRIPTION

BACKGROUND

Cancer screening is a critical component of cancer prevention, diagnosis, and treatment.

5 Colorectal cancer (CRC) has been identified, according to some reports, as the third most common type of cancer and the second most frequent cause of cancer mortality in the world. According to some reports, there are over 1.8 million new cases of colorectal cancer per year and about 881,000 deaths from colorectal cancer, accounting for about 1 in 10 cancer deaths. Regular colorectal cancer screening is recommended, particular for individuals over

10 age 50. Moreover, incidence of colorectal cancer in individuals below 50 has increased over time. Statistics suggest that current colorectal cancer screening techniques are insufficient.

SUMMARY

Despite improvements over time, only about 40-44% of colorectal cancers are currently

15 detected by screening in an early, localized stage. This is at least in part due to insufficient sensitivity and/or specificity of current screening techniques. Currently recommended techniques include colonoscopy and/or fecal blood testing for those over age 50.

The present disclosure provides, among other things, methods for colorectal cancer screening and compositions related thereto. In various embodiments, the present disclosure

20 provides methods for colorectal cancer screening that include analysis of methylation status of one or more methylation biomarkers, and compositions related thereto. In various embodiments, the present disclosure provides methods for colorectal cancer screening that include screening methylation status of one or more methylation biomarkers in cfDNA, e.g., in ctDNA. In various embodiments, the present disclosure provides methods for colorectal

25 cancer screening that include screening methylation status of one or more methylation biomarkers in cfDNA, e.g., in ctDNA, using MSRE-qPCR. Various compositions and methods provided herein provide sensitivity and specificity sufficient for clinical application in colorectal cancer screening. Various compositions and methods provided herein are useful in colorectal cancer screening by analysis of an accessible tissue sample of a subject, e.g., a

30 tissue sample that is blood or a blood component (e.g., cfDNA, e.g., ctDNA), or stool.

In certain aspects, the present disclosure provides, among other things, a method of screening for colorectal cancer, the method including determining the methylation status of one or more colorectal cancer methylation loci selected from: (a) one or more genes according to Table 1 (which includes, e.g., ALK, LONRF2, ADAMTS2, FGF14, DMRT1,

35 ST6GALNAC5, MCIDAS, PDGFD, GSG1L, ZNF492, ZNF568, ZNF542, ZNF471, ZNF132,

JAM2, and CNRIP1); and/or (b) one or more differentially methylated regions (DMRs) according to Table 7 (which includes, e.g., ALK '434, CNRIP1 '232, CNRIP1 '272, LONRF2 '281, LONRF2 '387, ADAMTS2 '254, ADAMTS2 '284, ADAMTS2 '328, FGF14 '577, DMRT1 '934, ST6GALNAC5 '456, MCIDAS '855, MCIDAS '003, PDGFD '388, PDGFD '921, GSG1L '861, ZNF492 '499, ZNF492 '069, ZNF568 '252, ZNF568 '405, ZNF542 '525, ZNF542 '502, ZNF471 '527, ZNF471 '558, ZNF471 '662, ZNF132 '268, , ZNF132 '415, and JAM2 '320), in DNA of a human subject, and diagnosing colorectal cancer in the subject. For instance, in some embodiments, methylation status can be determined using one or more members selected from the group consisting of methylation specific restriction enzyme quantitative polymerase chain reaction (MSRE-qPCR), methylation-specific PCR, methylation specific nuclease-assisted minor-allele enrichment PCR, and next-generation sequencing. In various embodiments, a diagnosis-confirmatory assay is performed on the human subject, e.g., a diagnosis-confirmatory assay provided in the present disclosure, subsequent to screening or diagnosis. In some instances, a diagnosis-confirmatory assay is a colonoscopy. In some instances, a diagnosis-confirmatory assay is performed within two weeks of screening or diagnosis, within one month of screening or diagnosis, within two months of screening or diagnosis, within three months of screening or diagnosis, or within one year of screening or diagnosis.

In certain aspects, the present disclosure provides, among other things, a method of methylation specific restriction enzyme quantitative polymerase chain reaction (MSRE-qPCR) for colorectal cancer screening, the method including: (a) contacting DNA of a human subject with one or more methylation specific restriction enzymes; and (b) performing qPCR of enzyme-digested DNA, or amplicons thereof, to determine the methylation status of one or more colorectal cancer methylation loci selected from: (i) one or more genes according to Table 1; and/or (ii) one or more differentially methylated regions (DMRs) according to Table 7. In various embodiments, e.g., as specifically referred to above, the DNA of the human subject is contacted with the methylation specific restriction enzymes, e.g., to digest unmethylated non-tumor derived DNA. In various embodiments, a digested DNA and/or DNA remaining after digestion, is preamplified (e.g., amplified in an amplification step that is prior to qPCR) for all or a portion of one or more of the selected colorectal cancer methylation loci. In certain embodiments, preamplification includes amplification of one or more samples of digested DNA, or DNA remaining after digestion, in an amplification reaction including one or more or all oligonucleotide primer pairs of Table 13, e.g., where the oligonucleotide primers are provided in a single pool. In subsequent qPCR, those of skill in the art will appreciate that methylation status will be separately, individually determined for

each individual methylation loci (e.g., each DMR) . In various embodiments, e.g., as specifically referred to above, the DNA is an aliquot of a sample, such that a separate aliquot of the sample that has not been contacted with the methylation specific restriction enzymes can provide a reference pool of DNA for qPCR, which reference pool of DNA can be

5 subjected to the same or comparable qPCR conditions as the digested DNA (e.g., preamplification using the same oligonucleotide primer pairs and quantification with qPCR).

In certain aspects, the present disclosure provides, among other things, a method of treating colorectal cancer (e.g., a colorectal cancer that is an early stage colorectal cancer), the method including: (a) diagnosing an colorectal cancer (e.g., a colorectal cancer that is an

10 early stage colorectal cancer) based on methylation status of one or more colorectal cancer methylation loci selected from: (i) one or more genes according to Table 1; and/or (ii) one or more differentially methylated regions (DMRs) according to Table 7, in DNA of a human subject; and (b) treating the colorectal cancer (e.g., the colorectal cancer that is an early stage colorectal cancer). In various embodiments, e.g., as specifically referred to above, a

15 diagnosis-confirmatory assay is performed on the human subject, e.g., a diagnosis-confirmatory assay provided in the present disclosure, subsequent to diagnosis based on methylation status. In some embodiments, e.g., as specifically referred to above, a diagnosis-confirmatory assay is a colonoscopy. In some embodiments, e.g., as specifically referred to above, a diagnosis-confirmatory assay is performed within two weeks of diagnosis

20 based on methylation status, within one month of diagnosis based on methylation status, within two months of diagnosis based on methylation status, within three months of diagnosis based on methylation status, or within one year of diagnosis based on methylation status.

In certain aspects, the present disclosure provides, among other things, a method of treating early stage colorectal cancer, the method including: (a) diagnosing an early stage colorectal

25 cancer based on methylation status of one or more colorectal cancer methylation loci selected from: (i) one or more genes according to Table 1; and/or (ii) one or more differentially methylated regions (DMRs) according to Table 7, in DNA of a human subject; and (b) treating the early stage colorectal cancer. In various embodiments, e.g., as specifically referred to above, a diagnosis-confirmatory assay is performed on the human

30 subject, e.g., a diagnosis-confirmatory assay provided in the present disclosure, subsequent to diagnosis based on methylation status. In some embodiments, e.g., as specifically referred to above, a diagnosis-confirmatory assay is a colonoscopy. In some embodiments, e.g., as specifically referred to above, a diagnosis-confirmatory assay is performed within two weeks of diagnosis based on methylation status, within one month of diagnosis based on

35 methylation status, within two months of diagnosis based on methylation status, within three

months of diagnosis based on methylation status, or within one year of diagnosis based on methylation status.

In certain aspects, the present disclosure provides a method of analyzing or using methylation sensitive restriction enzyme-digested human DNA, e.g., methylation sensitive restriction enzyme-digested human cfDNA, in an in vitro method of screening for colorectal cancer, the method including determining the methylation status of one or more colorectal cancer methylation loci selected from: (a) one or more genes according to Table 1 (which includes, e.g., ALK, LONRF2, ADAMTS2, FGF14, DMRT1, ST6GALNAC5, MCIDAS, PDGFD, GSG1L, ZNF492, ZNF568, ZNF542, ZNF471, ZNF132, JAM2, and CNRIP1);
5 and/or (b) one or more differentially methylated regions (DMRs) according to Table 7 (which includes, e.g., ALK '434, CNRIP1 '232, CNRIP1 '272, LONRF2 '281, LONRF2 '387, ADAMTS2 '254, ADAMTS2 '284, ADAMTS2 '328, FGF14 '577, DMRT1 '934, ST6GALNAC5 '456, MCIDAS '855, MCIDAS '003, PDGFD '388, PDGFD '921, GSG1L '861, ZNF492 '499, ZNF492 '069, ZNF568 '252, ZNF568 '405, ZNF542 '525, ZNF542 '502, ZNF471 '527,
15 ZNF471 '558, ZNF471 '662, ZNF132 '268, , ZNF132 '415, and JAM2 '320), in the DNA, and diagnosing colorectal cancer in the subject. For instance, in some embodiments, methylation status can be determined using one or more members selected from the group consisting of methylation specific restriction enzyme quantitative polymerase chain reaction (MSRE-qPCR), methylation-specific PCR, methylation specific nuclease-assisted minor-allele
20 enrichment PCR, and next-generation sequencing. In various embodiments, a diagnosis-confirmatory assay is performed on the human subject, e.g., a diagnosis-confirmatory assay provided in the present disclosure, subsequent to screening or diagnosis. In some instances, a diagnosis-confirmatory assay is a colonoscopy. In some instances, a diagnosis-confirmatory assay is performed within two weeks of screening or diagnosis, within one
25 month of screening or diagnosis, within two months of screening or diagnosis, within three months of screening or diagnosis, or within one year of screening or diagnosis.

In various embodiments, e.g., as specifically referred to above, an aspect of the present disclosure can include determining whether the one or more methylation loci are hypermethylated as compared to a reference, where hypermethylation is indicative of
30 colorectal cancer.

In various embodiments, e.g., as specifically referred to above, an aspect of the present disclosure can include determining the methylation status of at least one DMR of each of ZNF471 and FGF14. In various embodiments, e.g., as specifically referred to above, an aspect of the present disclosure can include determining the methylation status of at least
35 one DMR of each of ZNF471, FGF14, and PDGFD. In various embodiments, e.g., as

specifically referred to above, an aspect of the present disclosure can include determining the methylation status of at least one DMR of each of ZNF471, FGF14, PDGFD, and ADAMTS2. In various embodiments, e.g., as specifically referred to above, an aspect of the present disclosure can include determining the methylation status of at least one DMR of each of ZNF471, FGF14, PDGFD, ADAMTS2, ZNF492, ST6GALNAC5. In various
5 embodiments, e.g., as specifically referred to above, an aspect of the present disclosure can include determining the methylation status of at least one DMR of each of ZNF471, FGF14, PDGFD, ADAMTS2, ZNF492, ST6GALNAC5, ZNF542, LONRF2, ZNF132, and CNRIP1, ALK. In various embodiments, e.g., as specifically referred to above, an aspect of the present
10 disclosure can include determining the methylation status of at least one DMR of each of ALK, LONRF2, ADAMTS2, FGF14, DMRT1, ST6GALNAC5, MCIDAS, PDGFD, GSG1L, ZNF568, ZNF542, ZNF471, ZNF132, JAM2, ZNF492, and CNRIP1.

In various embodiments, e.g., as specifically referred to above, an aspect of the present disclosure can expressly and/or particularly exclude use of one or more colorectal cancer
15 methylation loci selected from: (i) one or more genes according to Table 1; and/or (ii) one or more differentially methylated regions (DMRs) according to Table 7. In some embodiments, e.g., as specifically referred to above, an aspect of the present disclosure can expressly and/or particularly exclude use of one or more colorectal cancer methylation loci selected
20 from FGF14, ZNF471, PDGFD, and ALK and combinations thereof. In some embodiments, e.g., as specifically referred to above, an aspect of the present disclosure can expressly and/or particularly exclude use of FGF14 as a methylation locus for screening colorectal cancer, which exclusion can be exclusion of FGF14 alone or exclusion of a plurality of methylation loci including FGF14. In some embodiments, e.g., as specifically referred to
25 above, an aspect of the present disclosure can expressly and/or particularly exclude use of ZNF471 as a methylation locus for screening colorectal cancer, which exclusion can be exclusion of ZNF471 alone or exclusion of a plurality of methylation loci including ZNF471. In some embodiments, e.g., as specifically referred to above, an aspect of the present
30 disclosure can expressly and/or particularly exclude use of PDGFD as a methylation locus for screening colorectal cancer, which exclusion can be exclusion of PDGFD alone or exclusion of a plurality of methylation loci including PDGFD. In some embodiments, e.g., as specifically referred to above, an aspect of the present disclosure can expressly and/or particularly exclude use of ALK as a methylation locus for screening colorectal cancer, which exclusion can be exclusion of ALK alone or exclusion of a plurality of methylation loci including ALK.

35 In various embodiments, e.g., as specifically referred to above, an aspect of the present

disclosure can include determining the methylation status of DMRs ZNF471 '558 and FGF14 '577. In various embodiments, e.g., as specifically referred to above, an aspect of the present disclosure can include determining the methylation status of DMRs ZNF471 '558, FGF14 '577, and PDGFD '388. In various embodiments, e.g., as specifically referred to above, an aspect of the present disclosure can include determining the methylation status of DMRs ZNF471 '558, FGF14 '577, PDGFD '388, ZNF471 '527, and ADAMTS2 '284. In various embodiments, e.g., as specifically referred to above, an aspect of the present disclosure can include determining the methylation status of DMRs ZNF471 '558, FGF14 '577, PDGFD '388, ZNF471 '527, ADAMTS2 '284, ADAMTS2 '254, ZNF492 '069, and ST6GALNAC5 '456. In various embodiments, e.g., as specifically referred to above, an aspect of the present disclosure can include determining the methylation status of DMRs ZNF471 '558, FGF14 '577, PDGFD '388, ZNF471 '527, ADAMTS2 '284, ADAMTS2 '254, ZNF492 '069, ST6GALNAC5 '456, ZNF542 '502, LONRF2 '281, ZNF132 '415, PDGFD '921, ZNF132 '268, CNRIP1 '272, and ALK '434. In various embodiments, e.g., as specifically referred to above, an aspect of the present disclosure can include determining the methylation status of DMRs ALK '434, CNRIP1 '232, CNRIP1 '272, LONRF2 '281, LONRF2 '387, ADAMTS2 '254, ADAMTS2 '284, ADAMTS2 '328, FGF14 '577, DMRT1 '934, ST6GALNAC5 '456, MCIDAS '855, MCIDAS '003, PDGFD '388, PDGFD '921, GSG1L '861, ZNF492 '499, ZNF492 '069, ZNF568 '252, ZNF568 '405, ZNF542 '525, ZNF542 '502, ZNF471 '527, ZNF471 '558, ZNF471 '662, ZNF132 '268, , ZNF132 '415, and JAM2 '320. In various embodiments, e.g., as specifically referred to above, an aspect of the present disclosure can expressly and/or particularly exclude use of one or more colorectal cancer DMRs selected from Table 7. In some embodiments, e.g., as specifically referred to above, an aspect of the present disclosure can expressly and/or particularly exclude use of one or more colorectal cancer DMRs selected from FGF14 '577, ZNF471 '527, ZNF471 '558, ZNF471 '662, PDGFD '388, PDGFD '921, and ALK '434 and combinations thereof. In some embodiments, e.g., as specifically referred to above, an aspect of the present disclosure can expressly and/or particularly exclude use of FGF14 '577 as a DMR for screening colorectal cancer, which exclusion can be exclusion of FGF14 '577 alone or exclusion of a plurality of DMRs including FGF14 '577. In some embodiments, e.g., as specifically referred to above, an aspect of the present disclosure can expressly and/or particularly exclude use of ZNF471 '527, ZNF471 '558, and/or ZNF471 '662 as DMRs for screening colorectal cancer, which exclusion can be exclusion of ZNF471 '527, ZNF471 '558, and/or ZNF471 '662 alone or exclusion of a plurality of DMRs including ZNF471 '527, ZNF471 '558, and/or ZNF471 '662. In some embodiments, e.g., as specifically referred to above, an aspect of the present

disclosure can expressly and/or particularly exclude use of PDGFD '388 and/or PDGFD '921 as DMRs for screening colorectal cancer, which exclusion can be exclusion of PDGFD '388 and/or PDGFD '921 alone or exclusion of a plurality of DMRs including PDGFD '388 and/or PDGFD '921. In some embodiments, e.g., as specifically referred to above, an aspect of the present disclosure can expressly and/or particularly exclude use of ALK '434 as a DMR for screening colorectal cancer, which exclusion can be exclusion of ALK '434 alone or exclusion of a plurality of DMRs including ALK '434.

In various embodiments, e.g., as specifically referred to above, an aspect of the present disclosure can include amplification of one or more of the DMRs by an oligonucleotide primer pair or set of oligonucleotide primer pairs provided in Table 13. In various embodiments, e.g., as specifically referred to above, of one or more aspects of the present disclosure, subject DNA is isolated from blood, plasma, urine, saliva, or stool of the human subject. In various embodiments, e.g., as specifically referred to above, of one or more aspects of the present disclosure, DNA is cell-free DNA of the human subject.

In various embodiments, e.g., as specifically referred to above, of one or more aspects of the present disclosure, a screened subject was asymptomatic for colorectal cancer at the time of screening. In various embodiments, e.g., as specifically referred to above, of one or more aspects of the present disclosure, a screened subject had been previously screened for colorectal cancer. In various embodiments, e.g., as specifically referred to above, of one or more aspects of the present disclosure, a screened subject had been screened for colorectal cancer with in the last 10 years, within the last 5 years, within the last 4 years, within the last 3 years, within the last 2 years, or within the last year. In some embodiments, e.g., as specifically referred to above, a previous screen for colorectal cancer in the subject had diagnosed the subject as not having colorectal cancer, e.g., where the previous screen for colorectal cancer that had diagnosed the subject as not having colorectal cancer was within one year and/or where the previous screen for colorectal cancer that had diagnosed the subject as not having colorectal cancer was a colonoscopy.

In various embodiments, e.g., as specifically referred to above, of one or more aspects of the present disclosure, a method of screening achieves or includes diagnosis of early stage colorectal cancer, optionally wherein the colorectal cancer is a stage 0, stage I, stage IIA, stage IIB, or stage IIC colorectal cancer. In various embodiments, e.g., as specifically referred to above, of one or more aspects of the present disclosure, a method of screening achieves or includes diagnosis of early stage colorectal cancer, e.g., where the cancer has not metastasized. In various embodiments, e.g., as specifically referred to above, of one or more aspects of the present disclosure, a method of screening achieves or includes

diagnosis of non-early stage colorectal cancer, optionally wherein the colorectal cancer is a stage IIIA, stage IIIB, stage IIIC, stage IVA, stage IVB, or a stage IVC colorectal cancer. In various embodiments, e.g., as specifically referred to above, of one or more aspects of the present disclosure, a method of screening achieves or includes diagnosis of non-early stage
5 colorectal cancer, e.g., where the cancer has metastasized.

In various embodiments, e.g., as specifically referred to above, of one or more aspects of the present disclosure, a method of screening achieves or provides a sensitivity for colorectal cancer of at least 0.6, e.g., at least 0.7. In some embodiments, e.g., as specifically referred to above, a method of screening provided herein achieves or provides a sensitivity for
10 colorectal cancer equal to or greater than 0.5, 0.55, 0.6, 0.65, 0.7, 0.75, 0.8, 0.85, 0.9, or 0.95.

In various embodiments, e.g., as specifically referred to above, of one or more aspects of the present disclosure, a method of screening achieves or provides a specificity for colorectal cancer of at least 0.7, e.g., at least 0.8. In some embodiments, e.g., as specifically referred to above, a method of screening provided herein achieves or provides a specificity for
15 colorectal cancer equal to or greater than 0.5, 0.55, 0.6, 0.65, 0.7, 0.75, 0.8, 0.85, 0.9, or 0.95.

In various embodiments, e.g., as specifically referred to above, one or more aspects of the present invention can include determining whether a subject includes an oncogenic mutation.
20 For instance, in various embodiments, e.g., as specifically referred to above, diagnosis can include determining the presence or absence of an oncogenic mutation in the subject, e.g., by analysis of cfDNA, cell sample DNA, or tissue sample DNA. In some embodiments, e.g., as specifically referred to above, the oncogenic mutation is in Kras, NRAS, PIK3CA, PTEN, TP53, BRAF, and APC. In some embodiments, e.g., as specifically referred to above, a
25 subject may include oncogenic mutations in two or more of Kras, NRAS, PIK3CA, PTEN, TP53, BRAF, and APC.

In some aspects, the present disclosure provides, among other things, a kit for use in colorectal cancer screening, the kit including: (a) at least one oligonucleotide primer pair of Table 13, and optionally further including: (b) at least one methylation specific restriction
30 enzyme and/or (c) a bisulfite reagent. In some aspects, the present disclosure provides, among other things, a diagnostic qPCR reaction for screening of colorectal cancer, the diagnostic qPCR reaction including: (a) human DNA; (b) a polymerase; and (c) at least one oligonucleotide primer pair of Table 13, optionally wherein the human DNA is bisulfite-treated human DNA or methylation specific restriction enzyme-digested human DNA. In various
35 embodiments, e.g., as specifically referred to above, the oligonucleotide primer pairs include

oligonucleotide primer pairs for amplification of DMRs ZNF471 '558 and FGF14 '577. In various embodiments, e.g., as specifically referred to above, the oligonucleotide primer pairs include oligonucleotide primer pairs for amplification of DMRs ZNF471 '558, FGF14 '577, and PDGFD '388. In various embodiments, e.g., as specifically referred to above, the oligonucleotide primer pairs include oligonucleotide primer pairs for amplification of DMRs ZNF471 '558, FGF14 '577, PDGFD '388, ZNF471 '527, and ADAMTS2 '284. In various embodiments, e.g., as specifically referred to above, the oligonucleotide primer pairs include oligonucleotide primer pairs for amplification of DMRs ZNF471 '558, FGF14 '577, PDGFD '388, ZNF471 '527, ADAMTS2 '284, ADAMTS2 '254, ZNF492 '069, and ST6GALNAC5 '456. In various embodiments, e.g., as specifically referred to above, the oligonucleotide primer pairs include oligonucleotide primer pairs for amplification of DMRs ZNF471 '558, FGF14 '577, PDGFD '388, ZNF471 '527, ADAMTS2 '284, ADAMTS2 '254, ZNF492 '069, ST6GALNAC5 '456, ZNF542 '502, LONRF2 '281, ZNF132 '415, PDGFD '921, ZNF132 '268, CNRIP1 '272, and ALK '434. In various embodiments, e.g., as specifically referred to above, the oligonucleotide primer pairs include oligonucleotide primer pairs for amplification of DMRs ALK '434, CNRIP1 '232, CNRIP1 '272, LONRF2 '281, LONRF2 '387, ADAMTS2 '254, ADAMTS2 '284, ADAMTS2 '328, FGF14 '577, DMRT1 '934, ST6GALNAC5 '456, MCIDAS '855, MCIDAS '003, PDGFD '388, PDGFD '921, GSG1L '861, ZNF492 '499, ZNF492 '069, ZNF568 '252, ZNF568 '405, ZNF542 '525, ZNF542 '502, ZNF471 '527, ZNF471 '558, ZNF471 '662, ZNF132 '268, , ZNF132 '415, and JAM2 '320.

In various embodiments, e.g., as specifically referred to above, an aspect of the present disclosure can expressly and/or particularly exclude use of oligonucleotide primer pairs for amplification of one or more colorectal cancer DMRs selected from Table 7. In some embodiments, e.g., as specifically referred to above, an aspect of the present disclosure can expressly and/or particularly exclude use of oligonucleotide primer pairs for amplification of one or more colorectal cancer DMRs selected from FGF14 '577, ZNF471 '527, ZNF471 '558, ZNF471 '662, PDGFD '388, PDGFD '921, and ALK '434 and combinations thereof. In some embodiments, e.g., as specifically referred to above, an aspect of the present disclosure can expressly and/or particularly exclude use of oligonucleotide primer pairs for amplification of FGF14 '577 as a DMR for screening colorectal cancer, which exclusion can be exclusion of FGF14 '577 alone or exclusion of a plurality of DMRs including FGF14 '577. In some embodiments, e.g., as specifically referred to above, an aspect of the present disclosure can expressly and/or particularly exclude use of oligonucleotide primer pairs for amplification of ZNF471 '527, ZNF471 '558, and/or ZNF471 '662 as DMRs for screening colorectal cancer, which exclusion can be exclusion of

ZNF471 '527, ZNF471 '558, and/or ZNF471 '662 alone or exclusion of a plurality of DMRs including ZNF471 '527, ZNF471 '558, and/or ZNF471 '662. In some embodiments, e.g., as specifically referred to above, an aspect of the present disclosure can expressly and/or particularly exclude use of oligonucleotide primer pairs for amplification of PDGFD '388 and/or PDGFD '921 as DMRs for screening colorectal cancer, which exclusion can be exclusion of PDGFD '388 and/or PDGFD '921 alone or exclusion of a plurality of DMRs including PDGFD '388 and/or PDGFD '921. In some embodiments, e.g., as specifically referred to above, an aspect of the present disclosure can expressly and/or particularly exclude use of oligonucleotide primer pairs for amplification of ALK '434 as a DMR for screening colorectal cancer, which exclusion can be exclusion of ALK '434 alone or exclusion of a plurality of DMRs including ALK '434.

In various embodiments, e.g., as specifically referred to above, of various aspects of the present disclosure a method, kit, or other composition of the present disclosure can further include or be useful in using a determined methylation status of the one or more methylation loci to identify at least one of (i) to (iv) as follows: (i) a presence of colorectal cancer in the human subject; (ii) a predisposition for colorectal cancer in the human subject; (iii) an increased risk of colorectal cancer in the human subject, and (iv) a stage of colorectal cancer in the human subject.

In various aspects and embodiments, including without limitation all aspects and embodiments provided above or otherwise provided herein, methods of colorectal cancer diagnosis provided herein are in vitro diagnostic methods, and compositions provided herein for use colorectal cancer diagnosis are compositions for in vitro use.

In various aspects, methods and compositions of the present invention can be used in combination with biomarkers known in the art, e.g., as disclosed in U.S. Patent No. 10,006,925, which is herein incorporated by reference in its entirety.

DEFINITIONS

A or An: The articles "a" and "an" are used herein to refer to one or to more than one (*i.e.*, to at least one) of the grammatical object of the article. By way of example, "an element" refers to one element or more than one element.

About: The term "about", when used herein in reference to a value, refers to a value that is similar, in context, to the referenced value. In general, those skilled in the art, familiar with the context, will appreciate the relevant degree of variance encompassed by "about" in that context. For example, in some embodiments, the term "about" can encompass a range of values that within 25%, 20%, 19%, 18%, 17%, 16%, 15%, 14%, 13%, 12%, 11%, 10%, 9%,

8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, or with a fraction of a percent, of the referred value.

Administration: As used herein, the term “administration” typically refers to the administration of a composition to a subject or system, for example to achieve delivery of an agent that is, is included in, or is otherwise delivered by, the composition.

- 5 *Agent:* As used herein, the term “agent” refers to an entity (e.g., for example, a small molecule, peptide, polypeptide, nucleic acid, lipid, polysaccharide, complex, combination, mixture, system, or phenomenon such as heat, electric current, electric field, magnetic force, magnetic field, etc.).

- Amelioration:* As used herein, the term “amelioration” refers to the prevention, reduction, 10 palliation, or improvement of a state of a subject. Amelioration includes, but does not require, complete recovery or complete prevention of a disease, disorder or condition.

- Amplicon or amplicon molecule:* As used herein, the term “amplicon” or “amplicon molecule” refers to a nucleic acid molecule generated by transcription from a template nucleic acid molecule, or a nucleic acid molecule having a sequence complementary thereto, or a double- 15 stranded nucleic acid including any such nucleic acid molecule. Transcription can be initiated from a primer.

- Amplification:* As used herein, the term “amplification” refers to the use of a template nucleic acid molecule in combination with various reagents to generate further nucleic acid molecules from the template nucleic acid molecule, which further nucleic acid molecules may 20 be identical to or similar to (e.g., at least 70% identical, e.g., at least 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to) a segment of the template nucleic acid molecule and/or a sequence complementary thereto.

- Amplification reaction mixture:* As used herein, the terms “amplification reaction mixture” or “amplification reaction” refer to a template nucleic acid molecule together with reagents 25 sufficient for amplification of the template nucleic acid molecule.

- Biological Sample:* As used herein, the term “biological sample” typically refers to a sample obtained or derived from a biological source (e.g., a tissue or organism or cell culture) of interest, as described herein. In some embodiments, a biological source is or includes an organism, such as an animal or human. In some embodiments, a biological sample is or 30 include biological tissue or fluid. In some embodiments, a biological sample can be or include cells, tissue, or bodily fluid. In some embodiments, a biological sample can be or include blood, blood cells, cell-free DNA, free floating nucleic acids, ascites, biopsy samples, surgical specimens, cell-containing body fluids, sputum, saliva, feces, urine, cerebrospinal fluid, peritoneal fluid, pleural fluid, lymph, gynecological fluids, secretions, excretions, skin 35 swabs, vaginal swabs, oral swabs, nasal swabs, washings or lavages such as a ductal

lavages or bronchoalveolar lavages, aspirates, scrapings, bone marrow. In some embodiments, a biological sample is or includes cells obtained from a single subject or from a plurality of subjects. A sample can be a "primary sample" obtained directly from a biological source, or can be a "processed sample." A biological sample can also be referred to as a "sample."

Biomarker: As used herein, the term "biomarker," consistent with its use in the art, refers to a to an entity whose presence, level, or form, correlates with a particular biological event or state of interest, so that it is considered to be a "marker" of that event or state. Those of skill in the art will appreciate, for instance, in the context of a DNA biomarker, that a biomarker can be or include a locus (such as one or more methylation loci) and/or the status of a locus (e.g., the status of one or more methylation loci). To give but a few examples of biomarkers, in some embodiments, a biomarker can be or include a marker for a particular disease, disorder or condition, or can be a marker for qualitative or quantitative probability that a particular disease, disorder or condition can develop, occur, or reoccur, e.g., in a subject. In some embodiments, a biomarker can be or include a marker for a particular therapeutic outcome, or qualitative or quantitative probability thereof. Thus, in various embodiments, a biomarker can be predictive, prognostic, and/or diagnostic, of the relevant biological event or state of interest. A biomarker can be an entity of any chemical class. For example, in some embodiments, a biomarker can be or include a nucleic acid, a polypeptide, a lipid, a carbohydrate, a small molecule, an inorganic agent (e.g., a metal or ion), or a combination thereof. In some embodiments, a biomarker is a cell surface marker. In some embodiments, a biomarker is intracellular. In some embodiments, a biomarker is found outside of cells (e.g., is secreted or is otherwise generated or present outside of cells, e.g., in a body fluid such as blood, urine, tears, saliva, cerebrospinal fluid, and the like). In some embodiments a biomarker is methylation status of a methylation locus. In some instances, a biomarker may be referred to as a "marker."

To give but one example of a biomarker, in some embodiments, the term refers to expression of a product encoded by a gene, expression of which is characteristic of a particular tumor, tumor subclass, stage of tumor, etc. Alternatively or additionally, in some embodiments, presence or level of a particular marker can correlate with activity (or activity level) of a particular signaling pathway, for example, of a signaling pathway the activity of which is characteristic of a particular class of tumors.

Those of skill in the art will appreciate that a biomarker may be individually determinative of a particular biological event or state of interest, or may represent or contribute to a determination of the statistical probability of a particular biological event or state of interest.

Those of skill in the art will appreciate that markers may differ in their specificity and/or sensitivity as related to a particular biological event or state of interest.

Blood component: As used herein, the term “blood component” refers to any component of whole blood, including red blood cells, white blood cells, plasma, platelets, endothelial cells, mesothelial cells, epithelial cells, and cell-free DNA. Blood components also include the components of plasma, including proteins, metabolites, lipids, nucleic acids, and carbohydrates, and any other cells that can be present in blood, e.g., due to pregnancy, organ transplant, infection, injury, or disease.

Cancer: As used herein, the terms “cancer,” “malignancy,” “neoplasm,” “tumor,” and “carcinoma,” are used interchangeably to refer to a disease, disorder, or condition in which cells exhibit or exhibited relatively abnormal, uncontrolled, and/or autonomous growth, so that they display or displayed an abnormally elevated proliferation rate and/or aberrant growth phenotype. In some embodiments, a cancer can include one or more tumors. In some embodiments, a cancer can be or include cells that are precancerous (e.g., benign), malignant, pre-metastatic, metastatic, and/or non-metastatic. In some embodiments, a cancer can be or include a solid tumor. In some embodiments, a cancer can be or include a hematologic tumor. In general, examples of different types of cancers known in the art include, for example, colorectal cancer, hematopoietic cancers including leukemias, lymphomas (Hodgkin’s and non-Hodgkin’s), myelomas and myeloproliferative disorders; sarcomas, melanomas, adenomas, carcinomas of solid tissue, squamous cell carcinomas of the mouth, throat, larynx, and lung, liver cancer, genitourinary cancers such as prostate, cervical, bladder, uterine, and endometrial cancer and renal cell carcinomas, bone cancer, pancreatic cancer, skin cancer, cutaneous or intraocular melanoma, cancer of the endocrine system, cancer of the thyroid gland, cancer of the parathyroid gland, head and neck cancers, breast cancer, gastro-intestinal cancers and nervous system cancers, benign lesions such as papillomas, and the like.

Chemotherapeutic agent: As used herein, the term “chemotherapeutic agent,” consistent with its use in the art, refers to one or more agents known, or having characteristics known to, treat or contribute to the treatment of cancer. In particular, chemotherapeutic agents include pro-apoptotic, cytostatic, and/or cytotoxic agents. In some embodiments, a chemotherapeutic agent can be or include alkylating agents, anthracyclines, cytoskeletal disruptors (e.g. microtubule targeting moieties such as taxanes, maytansine, and analogs thereof, of), epothilones, histone deacetylase inhibitors HDACs), topoisomerase inhibitors (e.g., inhibitors of topoisomerase I and/or topoisomerase II), kinase inhibitors, nucleotide analogs or nucleotide precursor analogs, peptide antibiotics, platinum-based agents,

retinoids, vinca alkaloids, and/or analogs that share a relevant anti-proliferative activity. In some particular embodiments, a chemotherapeutic agent can be or include of Actinomycin, All-trans retinoic acid, an Auristatin, Azacitidine, Azathioprine, Bleomycin, Bortezomib, Carboplatin, Capecitabine, Cisplatin, Chlorambucil, Cyclophosphamide, Curcumin, Cytarabine, Daunorubicin, Docetaxel, Doxifluridine, Doxorubicin, Epirubicin, Epothilone, Etoposide, Fluorouracil, Gemcitabine, Hydroxyurea, Idarubicin, Imatinib, Irinotecan, Maytansine and/or analogs thereof (e.g. DM1) Mechlorethamine, Mercaptopurine, Methotrexate, Mitoxantrone, a Maytansinoid, Oxaliplatin, Paclitaxel, Pemetrexed, Teniposide, Tioguanine, Topotecan, Valrubicin, Vinblastine, Vincristine, Vindesine, Vinorelbine, or a combination thereof. In some embodiments, a chemotherapeutic agent can be utilized in the context of an antibody-drug conjugate. In some embodiments, a chemotherapeutic agent is one found in an antibody-drug conjugate selected from the group consisting of: hLL1-doxorubicin, hRS7-SN-38, hMN-14-SN-38, hLL2-SN-38, hA20-SN-38, hPAM4-SN-38, hLL1-SN-38, hRS7-Pro-2-P-Dox, hMN-14-Pro-2-P-Dox, hLL2-Pro-2-P-Dox, hA20-Pro-2-P-Dox, hPAM4-Pro-2-P-Dox, hLL1-Pro-2-P-Dox, P4/D10-doxorubicin, gemtuzumab ozogamicin, brentuximab vedotin, trastuzumab emtansine, inotuzumab ozogamicin, glembatumomab vedotin, SAR3419, SAR566658, BIIB015, BT062, SGN-75, SGN-CD19A, AMG-172, AMG-595, BAY-94-9343, ASG-5ME, ASG-22ME, ASG-16M8F, MDX-1203, MLN-0264, anti-PSMA ADC, RG-7450, RG-7458, RG-7593, RG-7596, RG-7598, RG-7599, RG-7600, RG-7636, ABT-414, IMGN-853, IMGN-529, vorsetuzumab mafodotin, and lorvotuzumab mertansine. In some embodiments, a chemotherapeutic agent can be or comprise of farnesyl-thiosalicylic acid (FTS), 4-(4-Chloro-2-methylphenoxy)-N-hydroxybutanamide (CMH), estradiol (E2), tetramethoxystilbene (TMS), δ -tocatrienol, salinomycin, or curcumin.

Combination therapy: As used herein, the term “combination therapy” refers to administration to a subject of two or more agents or regimens such that the two or more agents or regimens together treat a disease, condition, or disorder of the subject. In some embodiments, the two or more therapeutic agents or regimens can be administered simultaneously, sequentially, or in overlapping dosing regimens. Those of skill in the art will appreciate that combination therapy includes but does not require that the two agents or regimens be administered together in a single composition, nor at the same time.

Comparable: As used herein, the term “comparable” refers to members within sets of two or more conditions, circumstances, agents, entities, populations, etc., that may not be identical to one another but that are sufficiently similar to permit comparison there between, such that one of skill in the art will appreciate that conclusions can reasonably be drawn based on differences or similarities observed. In some embodiments, comparable sets of conditions,

circumstances, agents, entities, populations, etc. are typically characterized by a plurality of substantially identical features and zero, one, or a plurality of differing features. Those of ordinary skill in the art will understand, in context, what degree of identity is required to render members of a set comparable. For example, those of ordinary skill in the art will appreciate that members of sets of conditions, circumstances, agents, entities, populations, etc., are comparable to one another when characterized by a sufficient number and type of substantially identical features to warrant a reasonable conclusion that differences observed can be attributed in whole or part to non-identical features thereof.

Detectable moiety: The term “detectable moiety” as used herein refers to any element, molecule, functional group, compound, fragment, or other moiety that is detectable. In some embodiments, a detectable moiety is provided or utilized alone. In some embodiments, a detectable moiety is provided and/or utilized in association with (e.g., joined to) another agent. Examples of detectable moieties include, but are not limited to, various ligands, radionuclides (e.g., ^3H , ^{14}C , ^{18}F , ^{19}F , ^{32}P , ^{35}S , ^{135}I , ^{125}I , ^{123}I , ^{64}Cu , ^{187}Re , ^{111}In , ^{90}Y , $^{99\text{m}}\text{Tc}$, ^{177}Lu , ^{89}Zr etc.), fluorescent dyes, chemiluminescent agents, bioluminescent agents, spectrally resolvable inorganic fluorescent semiconductor nanocrystals (i.e., quantum dots), metal nanoparticles, nanoclusters, paramagnetic metal ions, enzymes, colorimetric labels, biotin, dioxigenin, haptens, and proteins for which antisera or monoclonal antibodies are available.

Diagnosis: As used herein, the term “Diagnosis” refers to determining whether, and/or the qualitative or quantitative probability that, a subject has or will develop a disease, disorder, condition, or state. For example, in diagnosis of cancer, diagnosis can include a determination regarding the risk, type, stage, malignancy, or other classification of a cancer. In some instances, a diagnosis can be or include a determination relating to prognosis and/or likely response to one or more general or particular therapeutic agents or regimens.

Diagnostic information: As used herein, the term “diagnostic information” refers to information useful in providing a diagnosis. Diagnostic information can include, without limitation, biomarker status information.

Differentially methylated: As used herein, the term “differentially methylated” describes a methylation site for which the methylation status differs between a first condition and a second condition. A methylation site that is differentially methylated can be referred to as a differentially methylated site. In some instances, a DMR is defined by the amplicon produced by amplification using oligonucleotide primers, e.g., a pair of oligonucleotide primers selected for amplification of the DMR or for amplification of a DNA region of interest present in the amplicon. In some instances, a DMR is defined as a DNA region amplified by a pair of oligonucleotide primers, including the region having the sequence of, or a sequence

complementary to, the oligonucleotide primers. In some instances, a DMR is defined as a DNA region amplified by a pair of oligonucleotide primers, excluding the region having the sequence of, or a sequence complementary to, the oligonucleotide primers. As used herein, a specifically provided DMR can be unambiguously identified by the name of an associated gene followed by three digits of a starting position, such that, for example, a DMR starting at position 29921434 of ALK can be identified as ALK '434.

Differentially methylated region: As used herein, the term "differentially methylated region" (DMR) refers to a DNA region that includes one or more differentially methylated sites. A DMR that includes a greater number or frequency of methylated sites under a selected condition of interest, such as a cancerous state, can be referred to as a hypermethylation DMR. A DMR that includes a smaller number or frequency of methylated sites under a selected condition of interest, such as a cancerous state, can be referred to as a hypomethylation DMR. A DMR that is a methylation biomarker for colorectal cancer can be referred to as a colorectal cancer DMR. In some instances, a DMR can be a single nucleotide, which single nucleotide is a methylation site.

DNA region: As used herein, "DNA region" refers to any contiguous portion of a larger DNA molecule. Those of skill in the art will be familiar with techniques for determining whether a first DNA region and a second DNA region correspond, based, e.g., on sequence similarity (e.g., sequence identity or homology) of the first and second DNA regions and/or context (e.g., the sequence identity or homology of nucleic acids upstream and/or downstream of the first and second DNA regions).

Except as otherwise specified herein, sequences found in or relating to humans (e.g., that hybridize to human DNA) are found in, based on, and/or derived from the example representative human genome sequence commonly referred to, and known to those of skill in the art, as Homo sapiens (human) genome assembly GRCh38, hg38, and/or Genome Reference Consortium Human Build 38. Those of skill in the art will further appreciate that DNA regions of hg38 can be referred to by a known system including identification of particular nucleotide positions or ranges thereof in accordance with assigned numbering.

Dosing regimen: As used herein, the term "dosing regimen" can refer to a set of one or more same or different unit doses administered to a subject, typically including a plurality of unit doses administration of each of which is separated from administration of the others by a period of time. In various embodiments, one or more or all unit doses of a dosing regimen may be the same or can vary (e.g., increase over time, decrease over time, or be adjusted in accordance with the subject and/or with a medical practitioner's determination). In various embodiments, one or more or all of the periods of time between each dose may be the same

or can vary (e.g., increase over time, decrease over time, or be adjusted in accordance with the subject and/or with a medical practitioner's determination). In some embodiments, a given therapeutic agent has a recommended dosing regimen, which can involve one or more doses. Typically, at least one recommended dosing regimen of a marketed drug is known to those of skill in the art. In some embodiments, a dosing regimen is correlated with a desired or beneficial outcome when administered across a relevant population (i.e., is a therapeutic dosing regimen).

Downstream: As used herein, the term "downstream" means that a first DNA region is closer, relative to a second DNA region, to the C-terminus of a nucleic acid that includes the first DNA region and the second DNA region.

Gene: As used herein, the term "gene" refers to a single DNA region, e.g., in a chromosome, that includes a coding sequence that encodes a product (e.g., an RNA product and/or a polypeptide product), together with all, some, or none of the DNA sequences that contribute to regulation of the expression of coding sequence. In some embodiments, a gene includes one or more non-coding sequences. In some particular embodiments, a gene includes exonic and intronic sequences. In some embodiments, a gene includes one or more regulatory elements that, for example, can control or impact one or more aspects of gene expression (e.g., cell-type-specific expression, inducible expression, etc.). In some embodiments a gene includes a promoter. In some embodiments, a gene includes one or both of a (i) DNA nucleotides extending a predetermined number of nucleotides upstream of the coding sequence and (ii) DNA nucleotides extending a predetermined number of nucleotides downstream of the coding sequence. In various embodiments, the predetermined number of nucleotides can be 500 bp, 1 kb, 2 kb, 3 kb, 4 kb, 5 kb, 10 kb, 20 kb, 30 kb, 40 kb, 50 kb, 75 kb, or 100 kb.

Homology: As used herein, the term "homology" refers to the overall relatedness between polymeric molecules, e.g., between nucleic acid molecules (e.g., DNA molecules and/or RNA molecules) and/or between polypeptide molecules. Those of skill in the art will appreciate that homology can be defined, e.g., by a percent identity or by a percent homology (sequence similarity). In some embodiments, polymeric molecules are considered to be "homologous" to one another if their sequences are at least 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 99% identical. In some embodiments, polymeric molecules are considered to be "homologous" to one another if their sequences are at least 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 99% similar.

Hybridize: As used herein, "hybridize" refers to the association of a first nucleic acid with a

second nucleic acid to form a double-stranded structure, which association occurs through complementary pairing of nucleotides. Those of skill in the art will recognize that complementary sequences, among others, can hybridize. In various embodiments, hybridization can occur, for example, between nucleotide sequences having at least 70% complementarity, e.g., at least 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% complementarity. Those of skill in the art will further appreciate that whether hybridization of a first nucleic acid and a second nucleic acid does or does not occur can depend upon various reaction conditions. Conditions under which hybridization can occur are known in the art.

10 *Hypomethylation*: As used herein, the term “hypomethylation” refers to the state of a methylation locus having at least one fewer methylated nucleotides in a state of interest as compared to a reference state (e.g., at least one fewer methylated nucleotides in colorectal cancer than in healthy control).

Hypermethylation: As used herein, the term “hypermethylation” refers to the state of a methylation locus having at least one more methylated nucleotide in a state of interest as compared to a reference state (e.g., at least one more methylated nucleotide in colorectal cancer than in healthy control).

15 *Identity*: As used herein, the term “identity” refers to the overall relatedness between polymeric molecules, e.g., between nucleic acid molecules (e.g., DNA molecules and/or RNA molecules) and/or between polypeptide molecules. Methods for the calculation of a percent identity as between two provided sequences are known in the art. Calculation of the percent identity of two nucleic acid or polypeptide sequences, for example, can be performed by aligning the two sequences (or the complement of one or both sequences) for optimal comparison purposes (e.g., gaps can be introduced in one or both of a first and a second sequences for optimal alignment and non-identical sequences can be disregarded for comparison purposes). The nucleotides or amino acids at corresponding positions are then compared. When a position in the first sequence is occupied by the same residue (e.g., nucleotide or amino acid) as the corresponding position in the second sequence, then the molecules are identical at that position. The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, optionally taking into account the number of gaps, and the length of each gap, which may need to be introduced for optimal alignment of the two sequences. The comparison of sequences and determination of percent identity between two sequences can be accomplished using a computational algorithm, such as BLAST (basic local alignment search tool).

20 25 30 35 *“Improved,” “increased,” or “reduced”*: As used herein, these terms, or grammatically

- comparable comparative terms, indicate values that are relative to a comparable reference measurement. For example, in some embodiments, an assessed value achieved with an agent of interest may be “improved” relative to that obtained with a comparable reference agent or with no agent. Alternatively or additionally, in some embodiments, an assessed
- 5 value in a subject or system of interest may be “improved” relative to that obtained in the same subject or system under different conditions or at a different point in time (e.g., prior to or after an event such as administration of an agent of interest), or in a different, comparable subject (e.g., in a comparable subject or system that differs from the subject or system of interest in presence of one or more indicators of a particular disease, disorder or condition of
- 10 interest, or in prior exposure to a condition or agent, etc.). In some embodiments, comparative terms refer to statistically relevant differences (e.g., differences of a prevalence and/or magnitude sufficient to achieve statistical relevance). Those of skill in the art will be aware, or will readily be able to determine, in a given context, a degree and/or prevalence of difference that is required or sufficient to achieve such statistical significance.
- 15 *Methylation*: As used herein, the term “methylation” includes methylation at any of (i) C5 position of cytosine; (ii) N4 position of cytosine; and (iii) the N6 position of adenine. Methylation also includes (iv) other types of nucleotide methylation. A nucleotide that is methylated can be referred to as a “methylated nucleotide” or “methylated nucleotide base.” In certain embodiments, methylation specifically refers to methylation of cytosine residues.
- 20 In some instances, methylation specifically refers to methylation of cytosine residues present in CpG sites.
- Methylation assay*: As used herein, the term “methylation assay” refers to any technique that can be used to determine the methylation status of a methylation locus.
- Methylation biomarker*: As used herein, the term “methylation biomarker” refers to a
- 25 biomarker that is or includes at least one methylation locus and/or the methylation status of at least one methylation locus, e.g., a hypermethylation locus. In particular, a methylation biomarker is a biomarker characterized by a change between a first state and a second state (e.g., between a cancerous state and a non-cancerous state) in methylation status of one or more nucleic acid loci.
- 30 *Methylation locus*: As used herein, the term “methylation locus” refers to a DNA region that includes at least one differentially methylated region. A methylation locus that includes a greater number or frequency of methylated sites under a selected condition of interest, such as a cancerous state, can be referred to as a hypermethylation locus. A methylation locus that includes a smaller number or frequency of methylated sites under a selected condition of
- 35 interest, such as a cancerous state, can be referred to as a hypomethylation locus.

Methylation site: As used herein, a methylation site refers to a nucleotide or nucleotide position that is methylated in at least one condition. In its methylated state, a methylation site can be referred to as a methylated site.

Methylation status: As used herein, "methylation status," "methylation state," or "methylation profile" refer to the number, frequency, or pattern of methylation at methylation sites within a methylation locus. Accordingly, a change in methylation status between a first state and a second state can be or include an increase in the number, frequency, or pattern of methylated sites, or can be or include a decrease in the number, frequency, or pattern of methylated sites. In various instances, a change in methylation status is a change in methylation value.

Methylation value: As used herein, the term "methylation value" refers to a numerical representation of a methylation status, e.g., in the form of number that represents the frequency or ratio of methylation of a methylation locus. In some instances, a methylation value can be generated by a method that includes quantifying the amount of intact nucleic acid present in a sample following restriction digestion of the sample with a methylation dependent restriction enzyme. In some instances, a methylation value can be generated by a method that includes comparing amplification profiles after bisulfite reaction of a sample. In some instances, a methylation value can be generated by comparing sequences of bisulfite-treated and untreated nucleic acids. In some instances a methylation value is, includes, or is based on a quantitative PCR result.

Nucleic acid: As used herein, in its broadest sense, the term "nucleic acid" refers to any compound and/or substance that is or can be incorporated into an oligonucleotide chain. In some embodiments, a nucleic acid is a compound and/or substance that is or can be incorporated into an oligonucleotide chain via a phosphodiester linkage. As will be clear from context, in some embodiments, the term nucleic acid refers to an individual nucleic acid residue (e.g., a nucleotide and/or nucleoside), and in some embodiments refers to an polynucleotide chain comprising a plurality of individual nucleic acid residues. A nucleic acid can be or include DNA, RNA, or a combinations thereof. A nucleic acid can include natural nucleic acid residues, nucleic acid analogs, and/or synthetic residues. In some embodiments, a nucleic acid includes natural nucleotides (e.g., adenosine, thymidine, guanosine, cytidine, uridine, deoxyadenosine, deoxythymidine, deoxy guanosine, and deoxycytidine). In some embodiments, a nucleic acid is or includes of one or more nucleotide analogs (e.g., 2-aminoadenosine, 2-thiothymidine, inosine, pyrrolo-pyrimidine, 3 - methyl adenosine, 5-methylcytidine, C-5 propynyl-cytidine, C-5 propynyl-uridine, 2-aminoadenosine, C5-bromouridine, C5-fluorouridine, C5-iodouridine, C5-propynyl-uridine, C5

-propynyl-cytidine, C5-methylcytidine, 2-aminoadenosine, 7-deazaadenosine, 7-deazaguanosine, 8-oxoadenosine, 8-oxoguanosine, O(6)-methylguanine, 2-thiocytidine, methylated bases, intercalated bases, and combinations thereof).

In some embodiments, a nucleic acid has a nucleotide sequence that encodes a functional gene product such as an RNA or protein. In some embodiments, a nucleic acid includes one or more introns. In some embodiments, a nucleic acid includes one or more genes. In some embodiments, nucleic acids are prepared by one or more of isolation from a natural source, enzymatic synthesis by polymerization based on a complementary template (*in vivo* or *in vitro*), reproduction in a recombinant cell or system, and chemical synthesis.

In some embodiments, a nucleic acid analog differs from a nucleic acid in that it does not utilize a phosphodiester backbone. For example, in some embodiments, a nucleic acid can include one or more peptide nucleic acids, which are known in the art and have peptide bonds instead of phosphodiester bonds in the backbone. Alternatively or additionally, in some embodiments, a nucleic acid has one or more phosphorothioate and/or 5'-N-phosphoramidite linkages rather than phosphodiester bonds. In some embodiments, a nucleic acid comprises one or more modified sugars (e.g., 2'-fluororibose, ribose, 2'-deoxyribose, arabinose, and hexose) as compared with those in natural nucleic acids.

In some embodiments, a nucleic acid is or includes at least 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 225, 250, 275, 300, 325, 350, 375, 400, 425, 450, 475, 500, 600, 700, 800, 900, 1000, 1500, 2000, 2500, 3000, 3500, 4000, 4500, 5000 or more residues. In some embodiments, a nucleic acid is partly or wholly single stranded, or partly or wholly double stranded.

Nucleic acid detection assay: As used herein, the term "nucleic acid detection assay" refers to any method of determining the nucleotide composition of a nucleic acid of interest. Nucleic acid detection assays include but are not limited to, DNA sequencing methods, polymerase chain reaction-based methods, probe hybridization methods, ligase chain reaction, etc.

Nucleotide: As used herein, the term "nucleotide" refers to a structural component, or building block, of polynucleotides, e.g., of DNA and/or RNA polymers. A nucleotide includes of a base (e.g., adenine, thymine, uracil, guanine, or cytosine) and a molecule of sugar and at least one phosphate group. As used herein, a nucleotide can be a methylated nucleotide or an un-methylated nucleotide. Those of skill in the art will appreciate that nucleic acid terminology, such as, as examples, "locus" or "nucleotide" can refer to both a locus or nucleotide of a single nucleic acid molecule and/or to the cumulative population of loci or

nucleotides within a plurality of nucleic acids (e.g., a plurality of nucleic acids in a sample and/or representative of a subject) that are representative of the locus or nucleotide (e.g., having the same identical nucleic acid sequence and/or nucleic acid sequence context, or having a substantially identical nucleic acid sequence and/or nucleic acid context).

5 *Oligonucleotide primer*: As used herein, the term oligonucleotide primer, or primer, refers to a nucleic acid molecule used, capable of being used, or for use in, generating amplicons from a template nucleic acid molecule. Under transcription-permissive conditions (e.g., in the presence of nucleotides and a DNA polymerase, and at a suitable temperature and pH), an oligonucleotide primer can provide a point of initiation of transcription from a template to
10 which the oligonucleotide primer hybridizes. Typically, an oligonucleotide primer is a single-stranded nucleic acid between 5 and 200 nucleotides in length. Those of skill in the art will appreciate that optimal primer length for generating amplicons from a template nucleic acid molecule can vary with conditions including temperature parameters, primer composition, and transcription or amplification method. A pair of oligonucleotide primers, as used herein,
15 refers to a set of two oligonucleotide primers that are respectively complementary to a first strand and a second strand of a template double-stranded nucleic acid molecule. First and second members of a pair of oligonucleotide primers may be referred to as a “forward” oligonucleotide primer and a “reverse” oligonucleotide primer, respectively, with respect to a template nucleic acid strand, in that the forward oligonucleotide primer is capable of
20 hybridizing with a nucleic acid strand complementary to the template nucleic acid strand, the reverse oligonucleotide primer is capable of hybridizing with the template nucleic acid strand, and the position of the forward oligonucleotide primer with respect to the template nucleic acid strand is 5' of the position of the reverse oligonucleotide primer sequence with respect to the template nucleic acid strand. It will be understood by those of skill in the art that the
25 identification of a first and second oligonucleotide primer as forward and reverse oligonucleotide primers, respectively, is arbitrary inasmuch as these identifiers depend upon whether a given nucleic acid strand or its complement is utilized as a template nucleic acid molecule.

Pharmaceutical composition: As used herein, the term “pharmaceutical composition” refers
30 to a composition in which an active agent is formulated together with one or more pharmaceutically acceptable carriers. In some embodiments, the active agent is present in a unit dose amount appropriate for administration to a subject, e.g., in a therapeutic regimen that shows a statistically significant probability of achieving a predetermined therapeutic effect when administered to a relevant population. In some embodiments, a pharmaceutical
35 composition can be formulated for administration in a particular form (e.g., in a solid form or a

liquid form), and/or can be specifically adapted for, for example: oral administration (for example, as a drench (aqueous or non-aqueous solutions or suspensions), tablet, capsule, bolus, powder, granule, paste, etc., which can be formulated specifically for example for buccal, sublingual, or systemic absorption); parenteral administration (for example, by subcutaneous, intramuscular, intravenous or epidural injection as, for example, a sterile solution or suspension, or sustained-release formulation, etc.); topical application (for example, as a cream, ointment, patch or spray applied for example to skin, lungs, or oral cavity); intravaginal or intrarectal administration (for example, as a pessary, suppository, cream, or foam); ocular administration; nasal or pulmonary administration, etc.

10 *Pharmaceutically acceptable*: As used herein, the term “pharmaceutically acceptable,” as applied to one or more, or all, component(s) for formulation of a composition as disclosed herein, means that each component must be compatible with the other ingredients of the composition and not deleterious to the recipient thereof.

Pharmaceutically acceptable carrier: As used herein, the term “pharmaceutically acceptable carrier” refers to a pharmaceutically-acceptable material, composition, or vehicle, such as a liquid or solid filler, diluent, excipient, or solvent encapsulating material, that facilitates formulation and/or modifies bioavailability of an agent, e.g., a pharmaceutical agent. Some examples of materials which can serve as pharmaceutically-acceptable carriers include: sugars, such as lactose, glucose and sucrose; starches, such as corn starch and potato starch; cellulose, and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; powdered tragacanth; malt; gelatin; talc; excipients, such as cocoa butter and suppository waxes; oils, such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; glycols, such as propylene glycol; polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol; esters, such as ethyl oleate and ethyl laurate; agar; buffering agents, such as magnesium hydroxide and aluminum hydroxide; alginic acid; pyrogen-free water; isotonic saline; Ringer’s solution; ethyl alcohol; pH buffered solutions; polyesters, polycarbonates and/or polyanhydrides; and other non-toxic compatible substances employed in pharmaceutical formulations.

30 *Prevent or prevention*: The terms “prevent” and “prevention,” as used herein in connection with the occurrence of a disease, disorder, or condition, refers to reducing the risk of developing the disease, disorder, or condition; delaying onset of the disease, disorder, or condition; delaying onset of one or more characteristics or symptoms of the disease, disorder, or condition; and/or to reducing the frequency and/or severity of one or more characteristics or symptoms of the disease, disorder, or condition. Prevention can refer to prevention in a particular subject or to a statistical impact on a population of subjects.

Prevention can be considered complete when onset of a disease, disorder, or condition has been delayed for a predefined period of time.

Probe: As used herein, the term “probe” refers to a single- or double-stranded nucleic acid molecule that is capable of hybridizing with a complementary target and includes a detectable moiety. In certain embodiments, a probe is a restriction digest product or is a synthetically produced nucleic acid, e.g., a nucleic acid produced by recombination or amplification. In some instances a probe is a capture probe useful in detection, identification, and/or isolation of a target sequence, such as a gene sequence. In various instances, a detectable moiety of probe can be, e.g., an enzyme (e.g., ELISA, as well as enzyme-based histochemical assays), fluorescent moiety, radioactive moiety, or moiety associated with a luminescence signal.

Prognosis: As used herein, the term “prognosis” refers to determining the qualitative or quantitative probability of at least one possible future outcome or event. As used herein, a prognosis can be a determination of the likely course of a disease, disorder, or condition such as cancer in a subject, a determination regarding the life expectancy of a subject, or a determination regarding response to therapy, e.g., to a particular therapy.

Prognostic information: As used herein, the term “prognostic information” refers to information useful in providing a prognosis. Prognostic information can include, without limitation, biomarker status information.

Promoter: As used herein, a “promoter” can refer to a DNA regulatory region that directly or indirectly (e.g., through promoter-bound proteins or substances) associates with an RNA polymerase and participates in initiation of transcription of a coding sequence.

Reference: As used herein describes a standard or control relative to which a comparison is performed. For example, in some embodiments, an agent, subject, animal, individual, population, sample, sequence, or value of interest is compared with a reference or control agent, subject, animal, individual, population, sample, sequence, or value. In some embodiments, a reference or characteristic thereof is tested and/or determined substantially simultaneously with the testing or determination of the characteristic in a sample of interest. In some embodiments, a reference is a historical reference, optionally embodied in a tangible medium. Typically, as would be understood by those of skill in the art, a reference is determined or characterized under comparable conditions or circumstances to those under assessment, e.g., with regard to a sample. Those skilled in the art will appreciate when sufficient similarities are present to justify reliance on and/or comparison to a particular possible reference or control.

Risk: As used herein with respect to a disease, disorder, or condition, the term “risk” refers to

- the qualitative of quantitative probability (whether expressed as a percentage or otherwise) that a particular individual will develop the disease, disorder, or condition. In some embodiments, risk is expressed as a percentage. In some embodiments, a risk is a qualitative of quantitative probability that is equal to or greater than 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 60, 70, 80, 90, or 100%. In some embodiments risk is expressed as a qualitative of quantitative level of risk relative to a reference risk or level or the risk of the same outcome attributed to a reference. In some embodiments, relative risk is increased or decreased in comparison to the reference sample by a factor of 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more.
- 10 *Sample:* As used herein, the term "sample" typically refers to an aliquot of material obtained or derived from a source of interest. In some embodiments, a source of interest is a biological or environmental source. In some embodiments, a sample is a "primary sample" obtained directly from a source of interest. In some embodiments, as will be clear from context, the term "sample" refers to a preparation that is obtained by processing of a primary
- 15 sample (e.g., by removing one or more components of and/or by adding one or more agents to a primary sample). Such a "processed sample" can include, for example cells, nucleic acids, or proteins extracted from a sample or obtained by subjecting a primary sample to techniques such as amplification or reverse transcription of nucleic acids, isolation and/or purification of certain components, *etc.*
- 20 In certain instances, a processed sample can be a DNA sample that has been amplified (e.g., pre-amplified). Thus, in various instances, an identified sample can refer to a primary form of the sample or to a processed form of the sample. In some instances, a sample that is enzyme-digested DNA can refer to primary enzyme-digested DNA (the immediate product of enzyme digestion) or a further processed sample such as enzyme-digested DNA that has
- 25 been subject to an amplification step (e.g., an intermediate amplification step, e.g., pre-amplification) and/or to a filtering step, purification step, or step that modifies the sample to facilitate a further step, e.g., in a process of determining methylation status (e.g., methylation status of a primary sample of DNA and/or of DNA as it existed in its original source context).
- Screening:* As used herein, the term "screening" refers to any method, technique, process, or undertaking intended to generate diagnostic information and/or prognostic information.
- 30 Accordingly, those of skill in the art will appreciate that the term screening encompasses method, technique, process, or undertaking that determines whether an individual has, is likely to have or develop, or is at risk of having or developing a disease, disorder, or condition, e.g., colorectal cancer.
- 35 *Specificity:* As used herein, the "specificity" of a biomarker refers to the percentage of

samples that are characterized by absence of the event or state of interest for which measurement of the biomarker accurately indicates absence of the event or state of interest (true negative rate). In various embodiments, characterization of the negative samples is independent of the biomarker, and can be achieved by any relevant measure, e.g., any relevant measure known to those of skill in the art. Thus, specificity reflects the probability that the biomarker would detect the absence of the event or state of interest when measured in a sample not characterized that event or state of interest. In particular embodiments in which the event or state of interest is colorectal cancer, specificity refers to the probability that a biomarker would detect the absence of colorectal cancer in a subject lacking colorectal cancer. Lack of colorectal cancer can be determined, e.g., by histology.

Sensitivity: As used herein, the "sensitivity" of a biomarker refers to the percentage of samples that are characterized by the presence of the event or state of interest for which measurement of the biomarker accurately indicates presence of the event or state of interest (true positive rate). In various embodiments, characterization of the positive samples is independent of the biomarker, and can be achieved by any relevant measure, e.g., any relevant measure known to those of skill in the art. Thus, sensitivity reflects the probability that a biomarker would detect the presence of the event or state of interest when measured in a sample characterized by presence of that event or state of interest. In particular embodiments in which the event or state of interest is colorectal cancer, sensitivity refers to the probability that a biomarker would detect the presence of colorectal cancer in a subject that has colorectal cancer. Presence of colorectal cancer can be determined, e.g., by histology.

Solid Tumor: As used herein, the term "solid tumor" refers to an abnormal mass of tissue including cancer cells. In various embodiments, a solid tumor is or includes an abnormal mass of tissue that does not contain cysts or liquid areas. In some embodiments, a solid tumor can be benign; in some embodiments, a solid tumor can be malignant. Examples of solid tumors include carcinomas, lymphomas, and sarcomas. In some embodiments, solid tumors can be or include adrenal, bile duct, bladder, bone, brain, breast, cervix, colon, endometrium, esophagus, eye, gall bladder, gastrointestinal tract, kidney, larynx, liver, lung, nasal cavity, nasopharynx, oral cavity, ovary, penis, pituitary, prostate, retina, salivary gland, skin, small intestine, stomach, testis, thymus, thyroid, uterine, vaginal, and/or vulval tumors.

Stage of cancer: As used herein, the term "stage of cancer" refers to a qualitative or quantitative assessment of the level of advancement of a cancer. In some embodiments, criteria used to determine the stage of a cancer can include, but are not limited to, one or more of where the cancer is located in a body, tumor size, whether the cancer has spread to

lymph nodes, whether the cancer has spread to one or more different parts of the body, etc. In some embodiments, cancer can be staged using the so-called TNM System, according to which T refers to the size and extent of the main tumor, usually called the primary tumor; N refers to the number of nearby lymph nodes that have cancer; and M refers to whether the cancer has metastasized. In some embodiments, a cancer can be referred to as Stage 0 (abnormal cells are present but have not spread to nearby tissue, also called carcinoma *in situ*, or CIS; CIS is not cancer, but it can become cancer), Stage I-III (cancer is present; the higher the number, the larger the tumor and the more it has spread into nearby tissues), or Stage IV (the cancer has spread to distant parts of the body). In some embodiments, a cancer can be assigned to a stage selected from the group consisting of: *in situ* (abnormal cells are present but have not spread to nearby tissue); localized (cancer is limited to the place where it started, with no sign that it has spread); regional (cancer has spread to nearby lymph nodes, tissues, or organs); distant (cancer has spread to distant parts of the body); and unknown (there is not enough information to identify cancer stage).

Susceptible to: An individual who is "susceptible to" a disease, disorder, or condition is at risk for developing the disease, disorder, or condition. In some embodiments, an individual who is susceptible to a disease, disorder, or condition does not display any symptoms of the disease, disorder, or condition. In some embodiments, an individual who is susceptible to a disease, disorder, or condition has not been diagnosed with the disease, disorder, and/or condition. In some embodiments, an individual who is susceptible to a disease, disorder, or condition is an individual who has been exposed to conditions associated with, or presents a biomarker status (e.g., a methylation status) associated with, development of the disease, disorder, or condition. In some embodiments, a risk of developing a disease, disorder, and/or condition is a population-based risk (e.g., family members of individuals suffering from the disease, disorder, or condition).

Subject: As used herein, the term "subject" refers to an organism, typically a mammal (e.g., a human). In some embodiments, a subject is suffering from a disease, disorder or condition. In some embodiments, a subject is susceptible to a disease, disorder, or condition. In some embodiments, a subject displays one or more symptoms or characteristics of a disease, disorder or condition. In some embodiments, a subject is not suffering from a disease, disorder or condition. In some embodiments, a subject does not display any symptom or characteristic of a disease, disorder, or condition. In some embodiments, a subject is someone with one or more features characteristic of susceptibility to or risk of a disease, disorder, or condition. In some embodiments, a subject is a patient. In some embodiments, a subject is an individual to whom diagnosis has been performed

and/or to whom therapy has been administered. In some instances, a human subject can be interchangeably referred to as an "individual."

Therapeutic agent: As used herein, the term "therapeutic agent" refers to any agent that elicits a desired pharmacological effect when administered to a subject. In some embodiments, an agent is considered to be a therapeutic agent if it demonstrates a statistically significant effect across an appropriate population. In some embodiments, the appropriate population can be a population of model organisms or a human population. In some embodiments, an appropriate population can be defined by various criteria, such as a certain age group, gender, genetic background, preexisting clinical conditions, etc. In some embodiments, a therapeutic agent is a substance that can be used for treatment of a disease, disorder, or condition. In some embodiments, a therapeutic agent is an agent that has been or is required to be approved by a government agency before it can be marketed for administration to humans. In some embodiments, a therapeutic agent is an agent for which a medical prescription is required for administration to humans.

Therapeutically effective amount: As used herein, the term "therapeutically effective amount" refers to an amount that produces a desired effect for which it is administered. In some embodiments, the term refers to an amount that is sufficient, when administered to a population suffering from or susceptible to a disease, disorder, or condition, in accordance with a therapeutic dosing regimen, to treat the disease, disorder, or condition. Those of ordinary skill in the art will appreciate that the term therapeutically effective amount does not in fact require successful treatment be achieved in a particular individual. Rather, a therapeutically effective amount can be an amount that provides a particular desired pharmacological response in a significant number of subjects when administered to individuals in need of such treatment. In some embodiments, reference to a therapeutically effective amount can be a reference to an amount as measured in one or more specific tissues (e.g., a tissue affected by the disease, disorder or condition) or fluids (e.g., blood, saliva, serum, sweat, tears, urine, etc.). Those of ordinary skill in the art will appreciate that, in some embodiments, a therapeutically effective amount of a particular agent can be formulated and/or administered in a single dose. In some embodiments, a therapeutically effective agent can be formulated and/or administered in a plurality of doses, for example, as part of a multi-dose dosing regimen.

Treatment: As used herein, the term "treatment" (also "treat" or "treating") refers to administration of a therapy that partially or completely alleviates, ameliorates, relieves, inhibits, delays onset of, reduces severity of, and/or reduces incidence of one or more symptoms, features, and/or causes of a particular disease, disorder, or condition, or is

administered for the purpose of achieving any such result. In some embodiments, such treatment can be of a subject who does not exhibit signs of the relevant disease, disorder, or condition and/or of a subject who exhibits only early signs of the disease, disorder, or condition. Alternatively or additionally, such treatment can be of a subject who exhibits one or more established signs of the relevant disease, disorder and/or condition. In some embodiments, treatment can be of a subject who has been diagnosed as suffering from the relevant disease, disorder, and/or condition. In some embodiments, treatment can be of a subject known to have one or more susceptibility factors that are statistically correlated with increased risk of development of the relevant disease, disorder, or condition. In various examples, treatment is of a cancer.

Upstream: As used herein, the term “upstream” means a first DNA region is closer, relative to a second DNA region, to the N-terminus of a nucleic acid that includes the first DNA region and the second DNA region.

Unit dose: As used herein, the term “unit dose” refers to an amount administered as a single dose and/or in a physically discrete unit of a pharmaceutical composition. In many embodiments, a unit dose contains a predetermined quantity of an active agent. In some embodiments, a unit dose contains an entire single dose of the agent. In some embodiments, more than one unit dose is administered to achieve a total single dose. In some embodiments, administration of multiple unit doses is required, or expected to be required, in order to achieve an intended effect. A unit dose can be, for example, a volume of liquid (e.g., an acceptable carrier) containing a predetermined quantity of one or more therapeutic moieties, a predetermined amount of one or more therapeutic moieties in solid form, a sustained release formulation or drug delivery device containing a predetermined amount of one or more therapeutic moieties, etc. It will be appreciated that a unit dose can be present in a formulation that includes any of a variety of components in addition to the therapeutic agent(s). For example, acceptable carriers (e.g., pharmaceutically acceptable carriers), diluents, stabilizers, buffers, preservatives, etc., can be included. It will be appreciated by those skilled in the art, in many embodiments, a total appropriate daily dosage of a particular therapeutic agent can comprise a portion, or a plurality, of unit doses, and can be decided, for example, by a medical practitioner within the scope of sound medical judgment. In some embodiments, the specific effective dose level for any particular subject or organism can depend upon a variety of factors including the disorder being treated and the severity of the disorder; activity of specific active compound employed; specific composition employed; age, body weight, general health, sex and diet of the subject; time of administration, and rate of excretion of the specific active compound employed; duration of

the treatment; drugs and/or additional therapies used in combination or coincidental with specific compound(s) employed, and like factors well known in the medical arts

Unmethylated: As used herein, the terms “unmethylated” and “non-methylated” are used interchangeable and mean that an identified DNA region includes no methylated nucleotides.

5 **Variant**: As used herein, the term “variant” refers to an entity that shows significant structural identity with a reference entity but differs structurally from the reference entity in the presence, absence, or level of one or more chemical moieties as compared with the reference entity. In some embodiments, a variant also differs functionally from its reference entity. In general, whether a particular entity is properly considered to be a “variant” of a
10 reference entity is based on its degree of structural identity with the reference entity. A variant can be a molecule comparable, but not identical to, a reference. For example, a variant nucleic acid can differ from a reference nucleic acid at one or more differences in nucleotide sequence. In some embodiments, a variant nucleic acid shows an overall sequence identity with a reference nucleic acid that is at least 85%, 86%, 87%, 88%, 89%,
15 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, or 99%. In many embodiments, a nucleic acid of interest is considered to be a “variant” of a reference nucleic acid if the nucleic acid of interest has a sequence that is identical to that of the reference but for a small number of sequence alterations at particular positions. In some embodiments, a variant has 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 substituted residues as compared with a reference. In some embodiments,
20 a variant has not more than 5, 4, 3, 2, or 1 residue additions, substitutions, or deletions as compared with the reference. In various embodiments, the number of additions, substitutions, or deletions is fewer than about 25, about 20, about 19, about 18, about 17, about 16, about 15, about 14, about 13, about 10, about 9, about 8, about 7, about 6, and commonly are fewer than about 5, about 4, about 3, or about 2 residues.

25

BRIEF DESCRIPTION OF THE DRAWING

Figure 1 is a schematic showing an example MSRE-qPCR approach.

Figure 2 is a table showing characteristics of a first subject group of 70 human subjects.

Figure 2 provides the percent female, percent male, age range, and BMI of subjects. Figure
30 2 further distinguishes the types of colorectal cancer identified in the first subject group as localized or advanced based on histological evaluation, and as a proximal or distal based on colonoscopy evaluation of colon.

Figure 3 is a table showing characteristics of a second subject group of 63 human subjects.

Figure 3 provides the percent female, percent male, age range, and BMI of subjects. Figure
35 3 further distinguishes the types of colorectal cancer identified in the second subject group as

localized or advanced based on histological evaluation, and as a proximal or distal based on colonoscopy evaluation of colon.

Figure 4 includes panels A and B. Panel A of Figure 4 is a graph showing performance of colorectal cancer screening using a representative proof-of-principle panel of DMRs on the second subject group. ROC curve and AUC for all subjects of the second subject group are shown. Panel B of Figure 4 is a chart showing accuracy values, including, from left to right, overall sensitivity of colorectal screening for colorectal cancer, sensitivity of colorectal screening for localized colorectal cancer, sensitivity of colorectal screening for advanced colorectal cancer, sensitivity of colorectal screening for proximal colorectal cancer, sensitivity of colorectal screening for distal colorectal cancer, and specificity of colorectal screening for control subjects (healthy subjects and subjects with non-advanced adenoma).

Figure 5 is a graph representing Ct values from MSRE-qPCR of ALK '434 for subjects with colorectal cancer and control subjects (healthy subjects and subjects with non-advanced adenoma). Data represent the second subject group (63 subjects) used for testing. For display purposes, Ct values are subtracted from 45 (45– dCt). Higher 45 – dCt values correspond to higher methylation status, demonstrating hypermethylation in subjects with colorectal cancer.

Figure 6 is a graph representing Ct values from MSRE-qPCR of FGF14 '577 DMR for subjects with colorectal cancer and control subjects (healthy subjects and subjects with non-advanced adenoma). Data represent the second subject group (63 subjects) used for testing. For display purposes, Ct values are subtracted from 45 (45– dCt). Higher 45 – dCt values correspond to higher methylation status, demonstrating hypermethylation in subjects with colorectal cancer.

Figure 7 is a graph representing Ct values from MSRE-qPCR of PDGFD '388 for subjects with colorectal cancer and control subjects (healthy subjects and subjects with non-advanced adenoma). Data represent the second subject group (63 subjects) used for testing. For display purposes, Ct values are subtracted from 45 (45– dCt). Higher 45 – dCt values correspond to higher methylation status, demonstrating hypermethylation in subjects with colorectal cancer.

Figure 8 is a graph representing Ct values from MSRE-qPCR of JAM2 '320 for subjects with colorectal cancer and control subjects (healthy subjects and subjects with non-advanced adenoma). Data represent the second subject group (63 subjects) used for testing. For display purposes, Ct values are subtracted from 45 (45– dCt). Higher 45 – dCt values correspond to higher methylation status, demonstrating hypermethylation in subjects with colorectal cancer.

Figure 9 is a graph representing Ct values from MSRE-qPCR of LONRF2 '281 for subjects with colorectal cancer and control subjects (healthy subjects and subjects with non-advanced adenoma). Data represent the second subject group (63 subjects) used for testing. For display purposes, Ct values are subtracted from 45 (45– dCt). Higher 45 – dCt values correspond to higher methylation status, demonstrating hypermethylation in subjects with colorectal cancer.

Figure 10 is a table showing characteristics of a third subject group of 82 human subjects. Figure 10 provides the percent female, percent male, age range, and BMI of subjects diagnosed by screening using 28 colorectal cancer DMRs of the present disclosure. Figure 10 further distinguishes the types of colorectal cancer identified in the third subject group as localized or advanced based on histological evaluation, and as a proximal or distal based on colonoscopy evaluation of colon.

Figure 11 is a graph showing performance of colorectal cancer screening using a 28 DMR panel in the third subject group. ROC curve and AUC for all subjects of the third subject group are shown. ROC-curve analysis showed that a 28 DMR panel achieved general colorectal cancer sensitivity of 79%, with 75% sensitivity for localized (early) cancer and 84% sensitivity for advanced cancer, on a very stable specificity of 87% at AUC 82% (see also Table 15).

Figure 12 is a graph representing Ct values from MSRE-qPCR of ZNF471 '527 for subjects with colorectal cancer and control subjects (healthy subjects and subjects with non-advanced adenoma). Data represent the third subject group (82 subjects). For display purposes, Ct values are subtracted from 45 (45– dCt). Higher 45 – dCt values correspond to higher methylation status, demonstrating hypermethylation in subjects with colorectal cancer.

Figure 13 is a graph representing Ct values from MSRE-qPCR of FGF14 '577 for subjects with colorectal cancer and control subjects (healthy subjects and subjects with non-advanced adenoma). Data represent the third subject group (82 subjects). For display purposes, Ct values are subtracted from 45 (45– dCt). Higher 45 – dCt values correspond to higher methylation status, demonstrating hypermethylation in subjects with colorectal cancer.

Figure 14 is a graph representing Ct values from MSRE-qPCR of PDGFD '388 for subjects with colorectal cancer and control subjects (healthy subjects and subjects with non-advanced adenoma). Data represent the third subject group (82 subjects). For display purposes, Ct values are subtracted from 45 (45– dCt). Higher 45 – dCt values correspond to higher methylation status, demonstrating hypermethylation in subjects with colorectal cancer.

Figure 15 is a graph representing Ct values from MSRE-qPCR of ADAMTS2 '254 for subjects with colorectal cancer and control subjects (healthy subjects and subjects with non-

- advanced adenoma). Data represent the third subject group (82 subjects). For display purposes, Ct values are subtracted from 45 (45– dCt). Higher 45 – dCt values correspond to higher methylation status, demonstrating hypermethylation in subjects with colorectal cancer. Figure 16 is a graph representing Ct values from MSRE-qPCR of ZNF471 '558(which DMR
- 5 is alternatively referred to herein as ZNF471_2) for subjects with colorectal cancer and control subjects (healthy subjects and subjects with non-advanced adenoma). Data represent the third subject group (82 subjects). For display purposes, Ct values are subtracted from 45 (45– dCt). Higher 45 – dCt values correspond to higher methylation status, demonstrating hypermethylation in subjects with colorectal cancer.
- 10 Figure 17 is a graph representing Ct values from MSRE-qPCR of ST6GALNAC5 '456 for subjects with colorectal cancer and control subjects (healthy subjects and subjects with non-advanced adenoma). Data represent the third subject group (82 subjects). For display purposes, Ct values are subtracted from 45 (45– dCt). Higher 45 – dCt values correspond to higher methylation status, demonstrating hypermethylation in subjects with colorectal cancer.
- 15 Figure 18 is a graph representing Ct values from MSRE-qPCR of ZNF542 '525 for subjects with colorectal cancer and control subjects (healthy subjects and subjects with non-advanced adenoma). Data represent the third subject group (82 subjects). For display purposes, Ct values are subtracted from 45 (45– dCt). Higher 45 – dCt values correspond to higher methylation status, demonstrating hypermethylation in subjects with colorectal cancer.
- 20 Figure 19 is a graph representing Ct values from MSRE-qPCR of LONRF2 '281 for subjects with colorectal cancer and control subjects (healthy subjects and subjects with non-advanced adenoma). Data represent the third subject group (82 subjects). For display purposes, Ct values are subtracted from 45 (45– dCt). Higher 45 – dCt values correspond to higher methylation status, demonstrating hypermethylation in subjects with colorectal cancer.
- 25 Figure 20 is a schematic showing example methylation changes in methylation status between normal and cancer cells, and further indicates how changes in methylation status can impact gene expression differences between normal and cancer cells.

DETAILED DESCRIPTION

30 Screening for Colorectal Cancer

There is a need for improved methods of screening for colorectal cancer, including screening for diagnosis of early-stage colorectal cancer. Despite recommendations for screening of individuals, e.g., over age 50, colorectal cancer screening programs are often ineffective or unsatisfactory. Improved colorectal cancer screening improves diagnosis and reduces

35 colorectal cancer mortality.

DNA methylation (e.g., hypermethylation or hypomethylation) can activate or inactivate genes, including genes that impact cancer development (see, e.g., Fig. 20). Thus, for example, hypermethylation can inactivate one or more genes that typically act to suppress cancer, causing or contributing to development of cancer in a sample or subject.

- 5 The present disclosure includes the discovery that determination of the methylation status of one or more methylation loci provided herein, and/or the methylation status of one or more DMRs provided herein, and/or the methylation status of one or more methylation sites provided herein, provides screening for colorectal cancer, e.g., with a high degree of sensitivity and/or specificity. The present disclosure provides compositions and methods
- 10 including or relating to colorectal cancer methylation biomarkers that, individually or in various panels comprising two or more colorectal cancer methylation biomarkers, provide for screening of colorectal cancer, e.g., with a high degree of specificity and/or sensitivity.

In various embodiments, a colorectal cancer methylation biomarker of the present disclosure is selected from a methylation locus that is or includes ALK, LONRF2, ADAMTS2, FGF14, DMRT1, ST6GALNAC5, MCIDAS, PDGFD, GSG1L, ZNF492, ZNF568, ZNF542, ZNF471, ZNF132, JAM2, and CNRIP1 (see, e.g., Table 1). In various embodiments, a colorectal cancer DMR is selected from ALK '434, CNRIP1 '232, CNRIP1 '272, LONRF2 '281, LONRF2 '387, ADAMTS2 '254, ADAMTS2 '284, ADAMTS2 '328, FGF14 '577, DMRT1 '934, ST6GALNAC5 '456, MCIDAS '855, MCIDAS '003, PDGFD '388, PDGFD '921, GSG1L '861, ZNF492 '499, ZNF492 '069, ZNF568 '252, ZNF568 '405, ZNF542 '525, ZNF542 '502, ZNF471 '527, ZNF471 '558, ZNF471 '662, ZNF132 '268, , ZNF132 '415, and JAM2 '320 (see, e.g., Table 7)

For the avoidance of any doubt, any methylation biomarker provided herein can be, or be included in, among other things, a colorectal cancer methylation biomarker.

- 25 In some embodiments, a colorectal cancer methylation biomarker can be or include a single methylation locus. In some embodiments, a colorectal cancer methylation biomarker can be or include two or more methylation loci. In some embodiments, a colorectal cancer methylation biomarker can be or include a single differentially methylated region (DMR). In some embodiments, a methylation locus can be or include two or more DMRs. In some
- 30 embodiments, a methylation biomarker can be or include a single methylation site. In other embodiments, a methylation biomarker can be or include two or more methylation sites. In some embodiments, a methylation locus can include two or more DMRs and further include DNA regions adjacent to one or more of the included DMRs.

In some instances, a methylation locus is or includes a gene, such as a gene provided in Table 1. In some instances a methylation locus is or includes a portion of a gene, e.g., a

35

portion of a gene provided in Table 1. In some instances, a methylation locus includes but is not limited to identified nucleic acid boundaries of a gene.

In some instances, a methylation locus is or includes a coding region of a gene, such as a coding region of a gene provided in Table 1. In some instances a methylation locus is or includes a portion of the coding region of gene, e.g., a portion of the coding region a gene provided in Table 1. In some instances, a methylation locus includes but is not limited to identified nucleic acid boundaries of a coding region of gene.

In some instances, a methylation locus is or includes a promoter and/or other regulatory region of a gene, such as a promoter and/or other regulatory region of a gene provided in Table 1. In some instances a methylation locus is or includes a portion of the promoter and/or regulatory region of gene, e.g., a portion of promoter and/or regulatory region a gene provided in Table 1. In some instances, a methylation locus includes but is not limited to identified nucleic acid boundaries of a promoter and/or other regulatory region of gene. In some embodiments a methylation locus is or includes a high CpG density promoter, or a portion thereof.

In some embodiments, a methylation locus is or includes non-coding sequence. In some embodiments, a methylation locus is or includes one or more exons, and/or one or more introns.

In some embodiments, a methylation locus includes a DNA region extending a predetermined number of nucleotides upstream of a coding sequence, and/or a DNA region extending a predetermined number of nucleotides downstream of a coding sequence. In various instances, a predetermined number of nucleotides upstream and/or downstream and be or include, e.g., 500 bp, 1 kb, 2 kb, 3 kb, 4 kb, 5 kb, 10 kb, 20 kb, 30 kb, 40 kb, 50 kb, 75 kb, or 100 kb. Those of skill in the art will appreciate that methylation biomarkers capable of impacting expression of a coding sequence may typically be within any of these distances of the coding sequence, upstream and/or downstream.

Those of skill in the art will appreciate that a methylation locus identified as a methylation biomarker need not necessarily be assayed in a single experiment, reaction, or amplicon. A single methylation locus identified as a colorectal cancer methylation biomarker can be assayed, e.g., in a method including separate amplification (or providing oligonucleotide primers and conditions sufficient for amplification of) of one or more distinct or overlapping DNA regions within a methylation locus, e.g., one or more distinct or overlapping DMRs. Those of skill in the art will further appreciate that a methylation locus identified as a methylation biomarker need not be analyzed for methylation status of each nucleotide, nor each CpG, present within the methylation locus. Rather, a methylation locus that is a

methylation biomarker may be analyzed, e.g., by analysis of a single DNA region within the methylation locus, e.g., by analysis of a single DMR within the methylation locus.

DMRs of the present disclosure can be a methylation locus or include a portion of a methylation locus. In some instances, a DMR is a DNA region with a methylation locus that is, e.g., 1 to 5,000 bp in length. In various embodiments, a DMR is a DNA region with a methylation locus that is equal to or less than 5000 bp, 4,000 bp, 3,000 bp, 2,000 bp, 1,000 bp, 950 bp, 900 bp, 850 bp, 800 bp, 750 bp, 700 bp, 650 bp, 600 bp, 550 bp, 500 bp, 450 bp, 400 bp, 350 bp, 300 bp, 250 bp, 200 bp, 150 bp, 100 bp, 50 bp, 40 bp, 30 bp, 20 bp, or 10 bp in length. In some embodiments, a DMR is 1, 2, 3, 4, 5, 6, 7, 8 or 9 bp in length.

10 Methylation biomarkers, including without limitation methylation loci and DMRs provided herein, can include at least one methylation site that is a colorectal cancer methylation biomarker.

For clarity, those of skill in the art will appreciate that term methylation biomarker is used broadly, such that a methylation locus can be a methylation biomarker that includes one or more DMRs, each of which DMRs is also itself a methylation biomarker, and each of which DMRs can include one or more methylation sites, each of which methylation sites is also itself a methylation biomarker. Moreover, a methylation biomarker can include two or more methylation loci. Accordingly, status as a methylation biomarker does not turn on the contiguousness of nucleic acids included in a biomarker, but rather on the existence of a change in methylation status for included DNA region(s) between a first state and a second state, such as between colorectal cancer and controls.

15

20

As provided herein, a methylation locus can be any of one or more methylation loci each of which methylation loci is or includes a gene identified in Table 1. In some particular embodiments, a colorectal cancer methylation biomarker includes a single methylation locus that is or includes a gene identified in Table 1. For example, in various embodiments, a colorectal cancer methylation biomarker can include a single methylation locus that is or includes a gene selected from LONRF2, ADAMTS2, FGF14, ST6GALNAC5, PDGFD, ZNF492, ZNF542, ZNF471, JAM2, GSG1L, DMRT1, and MCIDAS .

25

In some particular embodiments, a colorectal cancer methylation biomarker includes two or more methylation loci, each of which is or includes a gene identified in Table 1. In some embodiments, a colorectal cancer methylation biomarker includes 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, or 16, methylation loci, each of which is or includes a gene identified in Table 1.

30

In some particular embodiments, a colorectal cancer methylation biomarker includes two or more methylation loci, each of which two or more methylation loci is or includes a gene

35

identified in any one of tables Table 1 to 6, including without limitation and combinations of two or more methylation loci that respectively are or include genes identified in a one of Tables 2 to 6. In some particular embodiments, a colorectal cancer methylation biomarker includes two methylation loci, which two methylation loci include methylation loci that are or include the genes identified in Table 2. In some particular embodiments, a colorectal cancer methylation biomarker includes three methylation loci, which three methylation loci include methylation loci that are or include the genes identified in Table 3. In some particular embodiments, a colorectal cancer methylation biomarker includes four methylation loci, which four methylation loci include methylation loci that are or include the genes identified in Table 4. In some particular embodiments, a colorectal cancer methylation biomarker includes six methylation loci, which six methylation loci include methylation loci that are or include the genes identified in Table 5. In some particular embodiments, a colorectal cancer methylation biomarker includes eleven methylation loci, which eleven methylation loci include methylation loci that are or include the genes identified in Table 6. In various particular embodiments, a colorectal cancer methylation biomarker or colorectal cancer methylation biomarker panel includes one or more methylation loci of the present disclosure, but does not include a methylation locus that is or includes all or a portion of one or more of FGF14, ZNF471, PDGFD, and ALK.

Table 1. Methylation loci identified by gene name

Gene	Example DNA Region of Homo sapiens (human) genome assembly GRCh38 (hg38)	SEQ ID NO
ZNF132	ZNF132 chr19, bp 58439728 to 58440994	SEQ ID NO: 1
DMRT1	DMRT1 chr9, bp 841340 to 968090	SEQ ID NO: 2
ALK	ALK chr2, bp 29193215 to 29922286	SEQ ID NO: 3
JAM2	JAM2 chr21, bp 25637848 to 25714704	SEQ ID NO: 4
FGF14	FGF14 chr13, bp 101919879 to 102403137	SEQ ID NO: 5
MCIDAS	MCIDAS chr5, bp 55220951 to 55221051	SEQ ID NO: 6
ST6GALNAC5	ST6GALNAC5 chr1, bp 76866255 to 77063388	SEQ ID NO: 7
LONRF2	LONRF2 chr2, bp 100285667 to 100323015	SEQ ID NO: 8
PDGFD	PDGFD chr11, bp 104163499 to 104164026	SEQ ID NO: 9
GSG1L	GSG1L chr16, bp 27920615 to 28064275	SEQ ID NO: 10
ZNF492	ZNF492 chr19, bp 22633051 to 22666433	SEQ ID NO: 11
ZNF568	ZNF568 chr19, bp 36916312 to 36943940	SEQ ID NO: 12
ADAMTS2	ADAMTS2 chr5, bp 179118114 to 179344392	SEQ ID NO: 13
ZNF542	ZNF542 chr19, bp 56367838 to 56370986	SEQ ID NO: 14
ZNF471	ZNF471 chr19, bp 56507245 to 56508589	SEQ ID NO: 15
CNRIP1	CNRIP1 chr2, bp 68293114 to 68320928	SEQ ID NO: 16

Table 2. Combination of 2 methylation loci

ZNF471
FGF14

5

Table 3. Combination of 3 methylation loci

ZNF471
FGF14
PDGFD

Table 4. Combination of 4 methylation loci

ZNF471
FGF14
PDGFD
ADAMTS2

Table 5. Combination of 6 methylation loci

ZNF471
FGF14
PDGFD
ADAMTS2
ZNF492
ST6GALNAC5

5 Table 6. Combination of 11 methylation loci

ZNF471
FGF14
PDGFD
ADAMTS2
ZNF492
ST6GALNAC5
ZNF542
LONRF2
ZNF132
CNRIP1
ALK

As provided herein, a DMR can be any of one or more DMRs each of which DMRs is present in a methylation locus that is or includes a gene identified in Table 1. In some particular embodiments, a colorectal cancer methylation biomarker is or includes a single DMR that is, includes all or a portion of, or is present in a gene identified in Table 1. For example, in various embodiments, a colorectal cancer methylation biomarker can include a single DMR that is, includes all or a portion of, or is present in a gene selected from ALK, LONRF2, ADAMTS2, FGF14, DMRT1, ST6GALNAC5, MCIDAS, PDGFD, GSG1L, ZNF568, ZNF542, ZNF471, ZNF132, JAM2, ZNF492, and CNRIP1

In some particular embodiments, a colorectal cancer methylation biomarker includes two or more DMRs, each of which is, includes all or a portion of, or is present in a gene identified in Table 1. In some embodiments, a colorectal cancer methylation biomarker includes 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, or 16, DMRs, each of which is, includes all or a portion of, or is present in a gene identified in Table 1.

In some particular embodiments, a colorectal cancer methylation biomarker includes two or more DMRs, each of which two or more DMRs is, includes all or a portion of, or is present in a gene identified in any one of Tables 1-6. In some particular embodiments, a colorectal cancer methylation biomarker includes two DMRs, which two DMRs include DMRs that are, include all or a portion of, or are present in the genes identified in Table 2. In some particular embodiments, a colorectal cancer methylation biomarker includes three DMRs, which three DMRs include DMRs that are, include all or a portion of, or are present in the genes identified in Table 3. In some particular embodiments, a colorectal cancer methylation biomarker includes four DMRs, which four DMRs include DMRs that are, include all or a portion of, or are present in the genes identified in Table 4. In some particular embodiments, a colorectal cancer methylation biomarker includes six DMRs, which six DMRs include DMRs that are, include all or a portion of, or are present in the genes identified in Table 5. In some particular embodiments, a colorectal cancer methylation biomarker includes eleven DMRs, which eleven DMRs include DMRs that are, include all or a portion of, or are present in the genes identified in Table 6. In various particular embodiments, a colorectal cancer methylation biomarker or colorectal cancer methylation biomarker panel includes one or more DMRs, but the one or more DMRs do not include a DMR that is, includes all or a portion of, or is present in one or more of FGF14, ZNF471, PDGFD, and ALK.

As provided herein, a DMR can be any of one or more DMRs, each of which DMRs is, includes all of, or includes a portion of a DMR identified in Table 7, including without limitation DMRs specifically as identified in Table 7. In some particular embodiments, a colorectal cancer methylation biomarker is or includes a single DMR that is, includes all of, or includes a portion of a DMR identified in Table 7, including without limitation a DMR specifically as identified in Table 7, e.g., a DMR of Table Q selected from the group of DMRs including, without limitation, ALK '434, CNRIP1 '232, CNRIP1 '272, LONRF2 '281, LONRF2 '387, ADAMTS2 '254, ADAMTS2 '284, ADAMTS2 '328, FGF14 '577, DMRT1 '934, ST6GALNAC5 '456, MCIDAS '855, MCIDAS '003, PDGFD '388, PDGFD '921, GSG1L '861, ZNF492 '499, ZNF492 '069, ZNF568 '252, ZNF568 '405, ZNF542 '525, ZNF542 '502, ZNF471 '527, ZNF471 '558, ZNF471 '662, ZNF132 '268, , ZNF132 '415, and JAM2 '320. For example, in various embodiments, a colorectal cancer methylation biomarker can include

a single DMR that is, includes all of, or includes a portion of a DMR selected from LONRF2 '281, LONRF2 '387, ADAMTS2 '254, ADAMTS2 '284, ADAMTS2 '328, FGF14 '577, ST6GALNAC5 '456, PDGFD '388, PDGFD '921, ZNF492 '499, ZNF492 '069, ZNF542 '525, ZNF542 '502, ZNF471 '527, ZNF471 '558, and ZNF471 '662.

- 5 In some particular embodiments, a colorectal cancer methylation biomarker includes two or more DMRs, each of which DMRs is, includes all of, or includes a portion of a DMR identified in Table 7, including without limitation DMRs specifically as identified in Table 7. In some embodiments, a colorectal cancer methylation biomarker includes 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, or 28 DMRs, each of which
- 10 DMRs is, includes all of, or includes a portion of a DMR identified in Table 7, including without limitation DMRs specifically as identified in Table 7.

- In some particular embodiments, a colorectal cancer methylation biomarker includes two or more DMRs, each of which two or more DMRs is, includes all of, or includes a portion of a DMR identified in any one of Tables 7 to 12, including without limitation DMRs and
- 15 combinations thereof specifically as identified in Tables 8 to 12. In some particular embodiments, a colorectal cancer methylation biomarker includes two DMRs, which two DMRs are the DMRs identified in Table 8. In some particular embodiments, a colorectal cancer methylation biomarker includes three DMRs, which three DMRs are the DMRs identified in Table 9. In some particular embodiments, a colorectal cancer methylation
- 20 biomarker includes five DMRs, which five DMRs are the DMRs identified in Table 10. In some particular embodiments, a colorectal cancer methylation biomarker includes eight DMRs, which eight DMRs are the DMRs identified in Table 11. In some particular embodiments, a colorectal cancer methylation biomarker includes fifteen DMRs, which fifteen DMRs are the DMRs identified in Table 12. In various particular embodiments, a colorectal
- 25 cancer methylation biomarker or colorectal cancer methylation biomarker panel includes one or more DMRs of Table 7, but the one or more DMRs do not include one or more, or all, DMRs of FGF14, ZNF471, PDGFD, and ALK, e.g., do not include one or more, or all, DMRs of FGF14, ZNF471, PDGFD, and ALK as provided in Table 7.

Table 7: Colorectal cancer DMRs

gene name	chr	start	end	width	SEQ ID NO
ALK	2	29921434	29921541	108	SEQ ID NO:17
CNRIP1	2	68319232	68319342	111	SEQ ID NO:18
LONRF2	2	100321281	100321395	115	SEQ ID NO:19
LONRF2	2	100322387	100322463	77	SEQ ID NO:20
ADAMTS2	5	179344254	179344348	95	SEQ ID NO:21
ADAMTS2	5	179344284	179344383	100	SEQ ID NO:22
FGF14	13	102394577	102394651	75	SEQ ID NO:23
DMRT1	9	841934	842046	113	SEQ ID NO:24
ST6GALNAC5	1	76868456	76868525	70	SEQ ID NO:25
MCIDAS	5	55220855	55220971	117	SEQ ID NO:26
MCIDAS	5	55221003	55221122	120	SEQ ID NO:27
PDGFD	11	104163388	104163503	116	SEQ ID NO:28
PDGFD	11	104163921	104164058	138	SEQ ID NO:29
GSG1L	16	28063861	28063964	104	SEQ ID NO:30
ZNF492	19	22634499	22634596	98	SEQ ID NO:31
ZNF568	19	36916252	36916371	120	SEQ ID NO:32
ZNF568	19	36916405	36916476	72	SEQ ID NO:33
ZNF542	19	56368525	56368610	86	SEQ ID NO:34
ZNF542	19	56368502	56368591	90	SEQ ID NO:35
ZNF471	19	56507527	56507675	149	SEQ ID NO:36
ZNF471	19	56507558	56507675	118	SEQ ID NO:37
ZNF471	19	56507662	56507750	89	SEQ ID NO:38
ZNF132	19	58440268	58440435	168	SEQ ID NO:39
ZNF132	19	58440415	58440523	109	SEQ ID NO:40
JAM2	21	25640320	25640399	80	SEQ ID NO:41
ZNF492	19	22634069	22634174	106	SEQ ID NO:42
CNRIP1	2	68319272	68319342	71	SEQ ID NO:43
ADAMTS2	5	179344328	179344412	85	SEQ ID NO:44

Table 8. Combination of 2 DMRs

gene name	chr	start site	end site
ZNF471	19	56507558	56507675
FGF14	13	102394577	102394651

Table 9 Combination of 3 DMRs

gene name	chr	start site	end site
ZNF471	19	56507558	56507675
FGF14	13	102394577	102394651
PDGFD	11	104163388	104163503

Table 10. Combination of 5 DMRs

gene name	chr	start site	end site
ZNF471	19	56507558	56507675
FGF14	13	102394577	102394651
PDGFD	11	104163388	104163503
ZNF471	19	56507527	56507675
ADAMTS2	5	179344284	179344383

5

Table 11. Combination of 8 DMRs

gene name	chr	start site	end site
ZNF471	19	56507558	56507675
FGF14	13	102394577	102394651
PDGFD	11	104163388	104163503
ZNF471	19	56507527	56507675
ADAMTS2	5	179344284	179344383
ADAMTS2	5	179344254	179344348
ZNF492	19	22634069	22634174
ST6GALNAC5	1	76868456	76868525

10 Table 12. Combination of 15 DMRs

gene name	chr	start site	end site
ZNF471	19	56507558	56507675
FGF14	13	102394577	102394651
PDGFD	11	104163388	104163503
ZNF471	19	56507527	56507675
ADAMTS2	5	179344284	179344383
ADAMTS2	5	179344254	179344348
ZNF492	19	22634069	22634174
ST6GALNAC5	1	76868456	76868525
ZNF542	19	56368502	56368591
LONRF2	2	100321281	100321395
ZNF132	19	58440415	58440523
PDGFD	11	104163921	104164058
ZNF132	19	58440268	58440435
CNRIP1	2	68319272	68319342
ALK	2	29921434	29921541

In various embodiments, a methylation biomarker can be or include one or more individual nucleotides (e.g., a single individual cysteine residue in the context of CpG) or a plurality of individual cysteine residues (e.g., of a plurality of CpGs) present within one or more methylation loci (e.g., one or more DMRs) provided herein. Thus, in certain embodiments a methylation biomarker is or includes methylation status of a plurality of individual methylation sites.

In various embodiments, a methylation biomarker is, includes, or is characterized by change in methylation status that is a change in the methylation of one or more methylation sites within one or more methylation loci (e.g., one or more DMRs). In various embodiments, a methylation biomarker is or includes a change in methylation status that is a change in the number of methylated sites within a one or more methylation loci (e.g., one or more DMRs). In various embodiments, a methylation biomarker is or includes a change in methylation status that is a change in the frequency of methylation sites within one or more methylation loci (e.g., one or more DMRs). In various embodiments, a methylation biomarker is or includes a change in methylation status that is a change in the pattern of methylation sites within one or more methylation loci (e.g., one or more DMRs).

In various embodiments, methylation status of one or more methylation loci (e.g., one or more DMRs) is expressed as a fraction or percentage of the one or more methylation loci (e.g., the one or more DMRs) present in a sample that are methylated, e.g., as a fraction of the number of individual DNA strands of DNA in a sample that are methylated at a one or more particular methylation loci (e.g., one or more particular DMRs). Those of skill in the art will appreciate that, in some instances, the fraction or percentage of methylation can be calculated from the ratio of methylated DMRs to unmethylated DMRs for one or more analyzed DMRs, e.g., within a sample.

In various embodiments, methylation status of one or more methylation loci (e.g., one or more DMRs) is compared to a reference methylation status value and/or to methylation status of the one or more methylation loci (e.g., one or more DMRs) in a reference sample. In certain instances, a reference is a non-contemporaneous sample from the same source, e.g., a prior sample from the same source, e.g., from the same subject. In certain instances, a reference for the methylation status of one or more methylation loci (e.g., one or more DMRs) is the methylation status of the one or more methylation loci (e.g., one or more DMRs) in a sample (e.g., a sample from a subject), or a plurality of samples, known to represent a particular state (e.g., a cancer state or a non-cancer state). Thus, a reference can be or include one or more predetermined thresholds, which thresholds can be

quantitative (e.g., a methylation value) or qualitative. In certain instances, a reference for methylation status of a DMR is the methylation status of a nucleotide or plurality of nucleotides (e.g., a plurality of contiguous oligonucleotides) present in the same sample that does not include nucleotides of the DMR. Those of skill in the art will appreciate that a
5 reference measurement is typically produced by measurement using a methodology identical to, similar to, or comparable to that by which the non-reference measurement was taken.

Without wishing to be bound by any particular scientific theory, Figure 20 provides a schematic of one possible mechanism by which hypermethylation or hypomethylation of a regulatory sequence of gene can impact expression. As shown in Figure 20,
10 hypomethylation can result in increased expression and/or hypermethylation can result in suppression of expression. In various instances, increased methylation of expression-regulatory regions, such as promoter regions and enhancer regions, as compared to a reference can reduce or silence expression of an operably linked gene, e.g., of an operably linked gene that typically acts to suppress cancer. In various embodiments, decreased
15 methylation of expression-regulatory regions, such as promoter regions and enhancer regions, as compared to a reference can increase expression of an operably linked gene, e.g., of an operably linked gene having an activity that contributes to oncogenesis. Without wishing to be bound by any particular scientific theory, DNA methylation may provide a more chemically and biologically stable indicator of cancer status than RNA expression or protein
20 expression *per se*.

Methylation is typically thought to be highly tissue-specific, providing a dimension of information not necessarily present in DNA sequence analysis.

Methylation events that substantially contribute to oncogenesis can occur, e.g., in expression-regulatory regions of DNA (e.g., at a promoter region, enhancer region,
25 transcription factor binding site, CTCF-binding site, CpG island, or other sequence) operably linked with cancer-associated genes such as genes that typically act to suppress cancer. Accordingly, inactivation of genes that typically act to suppress cancer results in or contribute to oncogenesis. Moreover, hyper methylation is typically found at CpG islands.

30 Cancers

Methods and compositions of the present disclosure are useful for screening for cancer, particularly colorectal cancer. Colorectal cancers include, without limitation, colon cancer, rectal cancer, and combinations thereof. Colorectal cancers include metastatic colorectal cancers and non-metastatic colorectal cancers. Colorectal cancers include cancer located in
35 the proximal part of the colon cancer and cancer located the distal part of the colon.

Colorectal cancers include colorectal cancers at any of the various possible stages known in the art, including, e.g., Stage I, Stage II, Stage III, and Stage IV colorectal cancers (e.g., stages 0, I, IIA, IIB, IIC, IIIA, IIIB, IIIC, IVA, IVB, and IVC). Colorectal cancers include all stages of the Tumor/Node/Metastasis (TNM) staging system. With respect to colorectal cancer, T can refer to whether the tumor grown into the wall of the colon or rectum, and if so by how many layers; N can refer to whether the tumor has spread to lymph nodes, and if so how many lymph nodes and where they are located; and M can refer to whether the cancer has spread to other parts of the body, and if so which parts and to what extent. Particular stages of T, N, and M are known in the art. T stages can include TX, T0, Tis, T1, T2, T3, T4a, and T4b; N stages can include NX, N0, N1a, N1b, N1c, N2a, and N2b; M stages can include M0, M1a, and M1b. Moreover, grades of colorectal cancer can include GX, G1, G2, G3, and G4. Various means of staging cancer, and colorectal cancer in particular, are well known in the art summarized, e.g., on the world wide web at cancer.net/cancer-types/colorectal-cancer/stages.

In certain instances, the present disclosure includes screening of early stage colorectal cancer. Early stage colorectal cancers can include, e.g., colorectal cancers localized within a subject, e.g., in that they have not yet spread to lymph nodes of the subject, e.g., lymph nodes near to the cancer (stage N0), and have not spread to distant sites (stage M0). Early stage cancers include colorectal cancers corresponding to, e.g., Stages 0 to II C.

Thus, colorectal cancer s of the present disclosure include, among other things, pre-malignant colorectal cancer and malignant colorectal cancer. Methods and compositions of the present disclosure are useful for screening of colorectal cancer in all of its forms and stages, including without limitation those named herein or otherwise known in the art, as well as all subsets thereof. Accordingly, the person of skill in art will appreciate that all references to colorectal cancer provided here include, without limitation, colorectal cancer in all of its forms and stages, including without limitation those named herein or otherwise known in the art, as well as all subsets thereof.

Subjects and Samples

A sample analyzed using methods and compositions provided herein can be any biological sample and/or any sample including nucleic acid. In various particular embodiments, a sample analyzed using methods and compositions provided herein can be a sample from a mammal. In various particular embodiments, a sample analyzed using methods and compositions provided herein can be a sample from a human subject. In various particular embodiments, a sample analyzed using methods and compositions provided herein can be a

sample from a mouse, rat, pig, horse, chicken, or cow.

In various instances, a human subject is a subject diagnosed or seeking diagnosis as having, diagnosed as or seeking diagnosis as at risk of having, and/or diagnosed as or seeking diagnosis as at immediate risk of having, a cancer such as a colorectal cancer. In various
5 instances, a human subject is a subject identified as a subject in need of colorectal cancer screening. In certain instances, a human subject is a subject identified as in need of colorectal cancer screening by a medical practitioner. In various instances, a human subject is identified as in need of colorectal cancer screening due to age, e.g., due to an age equal to or greater than 50 years, e.g., an age equal to or greater than 50, 55, 60, 65, 70, 75, 80, 85,
10 or 90 years. In various instances, a human subject is a subject not diagnosed as having, not at risk of having, not at immediate risk of having, not diagnosed as having, and/or not seeking diagnosis for a cancer such as a colorectal cancer, or any combination thereof.

A sample from a subject, e.g., a human or other mammalian subject, can be a sample of, e.g., blood, blood component, cfDNA, ctDNA, stool, or colorectal tissue. In some particular
15 embodiments, a sample is an excretion or bodily fluid of a subject (e.g., stool, blood, lymph, or urine of a subject) or a colorectal cancer tissue sample. A sample from a subject can be a cell or tissue sample, e.g., a cell or tissue sample that is of a cancer or includes cancer cells, e.g., of a tumor or of a metastatic tissue. In various embodiments, a sample from a subject, e.g., a human or other mammalian subject, can be obtained by biopsy (e.g., fine needle
20 aspiration or tissue biopsy) or surgery.

In various particular embodiments, a sample is a sample of cell-free DNA (cfDNA). cfDNA is typically found in human biofluids (e.g., plasma, serum, or urine) in short, double-stranded fragments. The concentration of cfDNA is typically low, but can significantly increase under particular conditions, including without limitation pregnancy, autoimmune disorder,
25 myocardial infarction, and cancer. Circulating tumor DNA (ctDNA) is the component of circulating DNA specifically derived from cancer cells. ctDNA can be present in human biofluids bound to leukocytes and erythrocytes or not bound to leukocytes and erythrocytes. Various tests for detection of tumor-derived cfDNA are based on detection of genetic or epigenetic modifications that are characteristic of cancer (e.g., of a relevant cancer). Genetic
30 or epigenetic modifications characteristic of cancer can include, without limitation, oncogenic or cancer-associated mutations in tumor-suppressor genes, activated oncogenes, hypermethylation, and/or chromosomal disorders. Detection of genetic or epigenetic modifications characteristic of cancer can confirm that detected cfDNA is ctDNA.

cfDNA and ctDNA provide a real-time or nearly real time metric of the methylation status of a
35 source tissue. cfDNA and ctDNA demonstrate a half-life in blood of about 2 hours, such that

a sample taken at a given time provides a relatively timely reflection of the status of a source tissue.

Various methods of isolating nucleic acids from a sample (e.g., of isolating cfDNA from blood or plasma) are known in the art. Nucleic acids can be isolated, e.g., without limitation, standard DNA purification techniques, by direct gene capture (e.g., by clarification of a sample to remove assay-inhibiting agents and capturing a target nucleic acid, if present, from the clarified sample with a capture agent to produce a capture complex, and isolating the capture complex to recover the target nucleic acid).

10 Methods of measuring methylation status

Methylation status can be measured by a variety of methods known in the art and/or by methods provided herein. Those of skill in the art will appreciate that a method for measuring methylation status can generally be applied to samples from any source and of any kind, and will further be aware of processing steps available to modify a sample into a form suitable for measurement by a given methodology. Methods of measuring methylation status include, without limitation, methods including methylation-status-specific polymerase chain reaction (PCR), methods including nucleic acid sequencing, methods including mass spectrometry, methods including methylation-specific nucleases, methods including mass-based separation, methods including target-specific capture, and methods including methylation-specific oligonucleotide primers. Certain particular assays for methylation utilize a bisulfite reagent (e.g., hydrogen sulfite ions).

Bisulfite reagents can include, among other things, bisulfite, disulfite, hydrogen sulfite, or combinations thereof, which reagents can be useful in distinguishing methylated and unmethylated nucleic acids. Bisulfite interacts differently with cytosine and 5-methylcytosine. In typical bisulfite-based methods, contacting of DNA with bisulfite deaminates unmethylated cytosine to uracil, while methylated cytosine remains unaffected; methylated cytosines, but not unmethylated cytosines, are selectively retained. Thus, in a bisulfite processed sample, uracil residues stand in place of, and thus provide an identifying signal for, unmethylated cytosine residues, while remaining (methylated) cytosine residues thus provide an identifying signal for methylated cytosine residues. Bisulfite processed samples can be analyzed, e.g., by PCR.

Various methylation assay procedures can be used in conjunction with bisulfite treatment to determine methylation status of a target sequence such as a DMR. Such assays can include, among others, Methylation-Specific Restriction Enzyme qPCR, sequencing of bisulfite-treated nucleic acid, PCR (e.g., with sequence-specific amplification), Methylation

Specific Nuclease-assisted Minor-allele Enrichment PCR, and Methylation-Sensitive High Resolution Melting. In some embodiments, DMRs are amplified from a bisulfite-treated DNA sample and a DNA sequencing library is prepared for sequencing according to, e.g., an Illumina protocol or transpose-based Nextera XT protocol. In certain embodiments, high-throughput and/or next-generation sequencing techniques are used to achieve base-pair level resolution of DNA sequence, permitting analysis of methylation status.

In various embodiments, methylation status is detected by a method including PCR amplification with methylation-specific oligonucleotide primers (MSP methods), e.g., as applied to bisulfite-treated sample (see, e.g., Herman 1992 Proc. Natl. Acad. Sci. USA 93: 9821-9826, which is herein incorporated by reference with respect to methods of determining methylation status). Use of methylation-status-specific oligonucleotide primers for amplification of bisulfite-treated DNA allows differentiation between methylated and unmethylated nucleic acids. Oligonucleotide primer pairs for use in MSP methods include at least one oligonucleotide primer capable of hybridizing with sequence that includes a methylation site, e.g., a CpG. An oligonucleotide primer that includes a T residue at a position complementary to a cytosine residue will selectively hybridize to templates in which the cytosine was unmethylated prior to bisulfite treatment, while an oligonucleotide primer that includes a G residue at a position complementary to a cytosine residue will selectively hybridize to templates in which the cytosine was methylated prior to bisulfite treatment. MSP results can be obtained with or without sequencing amplicons, e.g., using gel electrophoresis. MSP (methylation-specific PCR) allows for highly sensitive detection (detection level of 0.1% of the alleles, with full specificity) of locus-specific DNA methylation, using PCR amplification of bisulfite-converted DNA.

Another method that can be used to determine methylation status after bisulfite treatment of a sample is Methylation-Sensitive High Resolution Melting (MS-HRM) PCR (see, e.g., Hussmann 2018 Methods Mol Biol. 1708:551-571, which is herein incorporated by reference with respect to methods of determining methylation status). MS-HRM is an in-tube, PCR-based method to detect methylation levels at specific loci of interest based on hybridization melting. Bisulfite treatment of the DNA prior to performing MS-HRM ensures a different base composition between methylated and unmethylated DNA, which is used to separate the resulting amplicons by high resolution melting. A unique primer design facilitates a high sensitivity of the assays enabling detection of down to 0.1–1% methylated alleles in an unmethylated background. Oligonucleotide primers for MS-HRM assays are designed to be complementary to the methylated allele, and a specific annealing temperature enables these primers to anneal both to the methylated and the unmethylated alleles thereby increasing the

sensitivity of the assays.

Another method that can be used to determine methylation status after bisulfite treatment of a sample is Quantitative Multiplex Methylation-Specific PCR (QM-MSP). QM-MSP uses methylation specific primers for sensitive quantification of DNA methylation (see, e.g.,
5 Fackler 2018 *Methods Mol Biol.* 1708:473-496, which is herein incorporated by reference with respect to methods of determining methylation status). QM-MSP is a two-step PCR approach, where in the first step, one pair of gene-specific primers (forward and reverse) amplifies the methylated and unmethylated copies of the same gene simultaneously and in multiplex, in one PCR reaction. This methylation-independent amplification step produces
10 amplicons of up to 10^9 copies per μL after 36 cycles of PCR. In the second step, the amplicons of the first reaction are quantified with a standard curve using real-time PCR and two independent fluorophores to detect methylated/unmethylated DNA of each gene in the same well (e.g., 6FAM and VIC). One methylated copy is detectable in 100,000 reference gene copies.

15 Another method that can be used to determine methylation status after bisulfite treatment of a sample is Methylation Specific Nuclease-assisted Minor-allele Enrichment (MS-NaME) (see, e.g., Liu 2017 *Nucleic Acids Res.* 45(6):e39, which is herein incorporated by reference with respect to methods of determining methylation status). Ms-NaME is based on selective hybridization of probes to target sequences in the presence of DNA nuclease specific to
20 double-stranded (ds) DNA (DSN), such that hybridization results in regions of double-stranded DNA that are subsequently digested by the DSN. Thus, oligonucleotide probes targeting unmethylated sequences generate local double stranded regions resulting to digestion of unmethylated targets; oligonucleotide probes capable of hybridizing to methylated sequences generate local double-stranded regions that result in digestion of
25 methylated targets, leaving methylated targets intact. Moreover, oligonucleotide probes can direct DSN activity to multiple targets in bisulfite-treated DNA, simultaneously. Subsequent amplification can enrich non-digested sequences. Ms-NaME can be used, either independently or in combination with other techniques provided herein.

Another method that can be used to determine methylation status after bisulfite treatment of
30 a sample is Methylation-sensitive Single Nucleotide Primer Extension (Ms-SNuPE™) (see, e.g., Gonzalgo 2007 *Nat Protoc.* 2(8):1931-6, which is herein incorporated by reference with respect to methods of determining methylation status). In Ms-SNuPE, strand-specific PCR is performed to generate a DNA template for quantitative methylation analysis using Ms-SNuPE. SNuPE is then performed with oligonucleotide(s) designed to hybridize immediately
35 upstream of the CpG site(s) being interrogated. Reaction products can be electrophoresed

on polyacrylamide gels for visualization and quantitation by phosphor-image analysis. Amplicons can also carry a directly or indirectly detectable labels such as a fluorescent label, radionuclide, or a detachable molecule fragment or other entity having a mass that can be distinguished by mass spectrometry. Detection may be carried out and/or visualized by
5 means of, e.g., matrix assisted laser desorption/ionization mass spectrometry (MALDI) or using electron spray mass spectrometry (ESI).

Certain methods that can be used to determine methylation status after bisulfite treatment of a sample utilize a first oligonucleotide primer, a second oligonucleotide primer, and an oligonucleotide probe in an amplification-based method. For instance, the oligonucleotide
10 primers and probe can be used in a method of real-time polymerase chain reaction (PCR) or droplet digital PCR (ddPCR). In various instances, the first oligonucleotide primer, the second oligonucleotide primer, and/or the oligonucleotide probe selectively hybridize methylated DNA and/or unmethylated DNA, such that amplification or probe signal indicate methylation status of a sample.

15 Other bisulfite-based methods for detecting methylation status (e.g., the presence of level of 5-methylcytosine) are disclosed, e.g., in Frommer (1992 Proc Natl Acad Sci U S A. 1;89(5):1827-31, which is herein incorporated by reference with respect to methods of determining methylation status).

Certain methods that can be used to determine methylation status do not include bisulfite
20 treatment of a sample. For instance, changes in methylation status can be detected by a PCR-based process in which DNA is digested with one or more methylation-sensitive restriction enzymes (MSREs) prior to PCR amplification (e.g., by MSRE-qPCR). Typically, MSREs have recognition sites that include at least one CpG motif, such that activity of the MSRE is blocked from cleaving a possible recognition site if the site includes 5-
25 methylcytosine. (see, e.g., Beikircher 2018 Methods Mol Biol. 1708:407-424, which is herein incorporated by reference with respect to methods of determining methylation status). Thus, MSREs selectively digest nucleic acids based upon methylation status of the recognition site of the MSRE; they can digest DNA at MSRE recognition sites that are unmethylated, but not digest DNA in MSRE recognition sites that are methylated. In certain embodiments, an
30 aliquot of sample can be digested with MSREs, generating a processed sample in which unmethylated DNA has been cleaved by the MSREs, such that, the proportion of uncleaved and/or amplifiable DNA with at least one methylated site within MSRE recognition sites (e.g., at least one methylated site within each MSRE recognition site of the DNA molecule) is increased relative to uncleaved and/or amplifiable DNA that did not include at least one
35 methylated site within MSRE recognition sites (e.g., did not include at least one methylated

- site within each MSRE recognition site of the DNA molecule). Uncleaved sequences of a restriction-enzyme-digested sample can then be preamplified, e.g. in PCR, and quantified e.g. by qPCR, real-time PCR, or digital PCR. Oligonucleotide primers for MSRE-qPCR amplify regions that include one or more MSRE cleavage sites, and/or a plurality of MSRE
- 5 cleavage sites. Amplicons including a plurality of MSRE cleavage sites are typically more likely to yield robust results. The number of cleavage sites within a DMR amplicon, and in some instances the resulting robustness of methylation status determination for the DMR, can be increased by design of DMRs that include a plurality of MSRE recognition sites (as opposed to a single recognition site) in a DMR amplicon. In various instances, a plurality of
- 10 MSREs can be applied to the same sample, including, e.g., two or more of Acil, Hin6I, HpyCH4IV, and HpaII (e.g., including Acil, Hin6I, and HpyCH4IV). A plurality of MSREs (e.g., the combination of Acil, Hin6I, HpyCH4IV, and HpaII, or the combination of Acil, Hin6I, and HpyCH4IV) can provide improved frequency of MSRE recognition sites within DMR amplicons.
- 15 MSRE-qPCR can also include a pre-amplification step following sample digestion by MSREs but before qPCR in order to improve the amount of available sample, given the low prevalence of cfDNA in blood.
- In certain MSRE-qPCR embodiments, the amount of total DNA is measured in an aliquot of sample in native (e.g., undigested) form using, e.g., real-time PCR or digital PCR.
- 20 Various amplification technologies can be used alone or in conjunction with other techniques described herein for detection of methylation status. Those of skill in the art, having reviewed the present specification, will understand how to combine various amplification technologies known in the art and/or described herein together with various other technologies for methylation status determination known in the art and/or provided herein.
- 25 Amplification technologies include, without limitation, PCR, e.g., quantitative PCR (qPCR), real-time PCR, and/or digital PCR. Those of skill in the art will appreciate that polymerase amplification can multiplex amplification of multiple targets in a single reaction. PCR amplicons are typically 100 to 2000 base pairs in length. In various instances, an amplification technology is sufficient to determine methylations status.
- 30 Digital PCR (dPCR) based methods involve dividing and distributing a sample across wells of a plate with 96-, 384-, or more wells, or in individual emulsion droplets (ddPCR) e.g., using a microfluidic device, such that some wells include one or more copies of template and others include no copies of template. Thus, the average number of template molecules per well is less than one prior to amplification. The number of wells in which amplification of template
- 35 occurs provides a measure of template concentration. If the sample has been contacted with

MSRE, the number of wells in which amplification of template occurs provides a measure of the concentration of methylated template.

In various embodiments a fluorescence-based real-time PCR assay, such as MethyLight™, can be used to measure methylation status (see, e.g., Campan 2018 Methods Mol Biol. 1708:497-513, which is herein incorporated by reference with respect to methods of determining methylation status) MethyLight is a quantitative, fluorescence-based, real-time PCR method to sensitively detect and quantify DNA methylation of candidate regions of the genome. MethyLight is uniquely suited for detecting low-frequency methylated DNA regions against a high background of unmethylated DNA, as it combines methylation-specific priming with methylation-specific fluorescent probing. Additionally, MethyLight can be combined with Digital PCR, for the highly sensitive detection of individual methylated molecules, with use in disease detection and screening.

Real-time PCR-based methods for use in determining methylation status typically include a step of generating a standard curve for unmethylated DNA based on analysis of external standards. A standard curve can be constructed from at least two points and can permit comparison of a real-time Ct value for digested DNA and/or a real-time Ct value for undigested DNA to known quantitative standards. In particular instances, sample Ct values can be determined for MSRE-digested and/or undigested samples or sample aliquots, and the genomic equivalents of DNA can be calculated from the standard curve. Ct values of MSRE-digested and undigested DNA can be evaluated to identify amplicons digested (e.g., efficiently digested; e.g., yielding a Ct value of 45). Amplicons not amplified under either digested or undigested conditions can also be identified. Corrected Ct values for amplicons of interest can then be directly compared across conditions to establish relative differences in methylation status between conditions. Alternatively or additionally, delta-difference between the Ct values of digested and undigested DNA can be used to establish relative differences in methylation status between conditions.

Methods of measuring methylation status can include, without limitation, massively parallel sequencing (e.g., next-generation sequencing) to determine methylation state, e.g., sequencing by- synthesis, real-time (e.g., single-molecule) sequencing, bead emulsion sequencing, nanopore sequencing, or other sequencing techniques known in the art. In some embodiments, a method of measuring methylation status can include whole-genome sequencing, e.g., with base-pair resolution.

In certain particular embodiments, MSRE-qPCR, among other techniques, can be used to determine the methylation status of a colorectal cancer methylation biomarker that is or includes a single methylation locus. In certain particular embodiments, MSRE-qPCR, among

other techniques, can be used to determine the methylation status of a colorectal cancer methylation biomarker that is or includes two or more methylation loci. In certain particular embodiments, MSRE-qPCR, among other techniques, can be used to determine the methylation status of a colorectal cancer methylation biomarker that is or includes a single differentially methylated region (DMR). In certain particular embodiments, MSRE-qPCR, among other techniques, can be used to determine the methylation status of a colorectal cancer methylation biomarker that is or includes two or more DMRs. In certain particular embodiments, MSRE-qPCR, among other techniques, can be used to determine the methylation status of a colorectal cancer methylation biomarker that is or includes a single methylation site. In certain particular embodiments, MSRE-qPCR, among other techniques, can be used to determine the methylation status of a colorectal cancer methylation biomarker that is or includes two or more methylation sites. In various embodiments, a colorectal cancer methylation biomarker can be any colorectal cancer methylation biomarker provided herein. The present disclosure includes, among other things, oligonucleotide primer pairs for amplification of DMRs, e.g., for amplification of DMRs identified in Table 7.

In certain particular embodiments, a cfDNA sample is derived from subject plasma and contacted with MSREs that are or include one or more of Acil, Hin6I, HpyCH4IV, and HpaII (e.g., Acil, Hin6I, and HpyCH4IV). The digested sample can be preamplified with oligonucleotide primer pairs of one or more DMRs, e.g., with one or more oligonucleotide primer pairs provided in Table 13. Digested DNA, e.g., preamplified digested DNA, can be quantified with qPCR with oligonucleotide primer pairs of one or more DMRs, e.g., with one or more oligonucleotide primer pairs provided in Table 13. qPCR ct values can then be determined and used to determine methylation status of each DMR amplicon.

It will be appreciated by those of skill in the art that oligonucleotide primer pairs provided in Table 13 can be used in accordance with any combination of colorectal cancer methylation biomarkers identified herein. The skilled artisan will be aware that the oligonucleotide primer pairs of Table 2 may be individual included or not included in a given analysis in order to analyze a particular desired combination of DMRs.

The person of skill in the art will further appreciate that while other oligonucleotide primer pairs may be used, selection and pairing of oligonucleotide primers to produce useful DMR amplicons is non-trivial and represents a substantial contribution.

Those of skill in the art will further appreciate that methods, reagents, and protocols for qPCR are well-known in the art. Unlike traditional PCR, qPCR is able to detect the production of amplicons over time in amplification (e.g., at the end of each amplification cycle), often by use of an amplification-responsive fluorescence system, e.g., in combination

with a thermocycler with fluorescence-detection capability. Two common types of fluorescent reporters used in qPCR include (i) double-stranded DNA binding dyes that fluoresce substantially more brightly when bound than when unbound; and (ii) labeled oligonucleotides (e.g., labeled oligonucleotide primers or labeled oligonucleotide probes).

- 5 Those of skill in the art will appreciate that in embodiments in which a plurality of methylation loci (e.g., a plurality of DMRs) are analyzed for methylation status in a method of screening for colorectal cancer provided herein, methylation status of each methylation locus can be measured or represented in any of a variety of forms, and the methylation statuses of a plurality of methylation loci (preferably each measured and/or represented in a same, similar, or comparable manner) be together or cumulatively analyzed or represented in any of a variety of forms. In various embodiments, methylation status of each methylation locus can be measured as a ct value. In various embodiments, methylation status of each methylation locus can be represented as the difference in ct value between a measured sample and a reference. In various embodiments, methylation status of each methylation locus can be represented as a qualitative comparison to a reference, e.g., by identification of each methylation locus as hypermethylated or not hypermethylated.

- In some embodiments in which a single methylation locus is analyzed, hypermethylation of the single methylation locus constitutes a diagnosis that a subject is suffering from or possibly suffering from colorectal cancer, while absence of hypermethylation of the single methylation locus constitutes a diagnosis that the subject is likely not suffering from colorectal cancer. In some embodiments, hypermethylation of a single methylation locus (e.g., a single DMR) of a plurality of analyzed methylation loci constitutes a diagnosis that a subject is suffering from or possibly suffering from colorectal cancer, while the absence of hypermethylation at any methylation locus of a plurality of analyzed methylation loci constitutes a diagnosis that a subject is likely not suffering from colorectal cancer. In some embodiments, hypermethylation of a determined percentage (e.g., a predetermined percentage) of methylation loci (e.g., at least 10% (e.g., at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, or 100%)) of a plurality of analyzed methylation loci constitutes a diagnosis that a subject is suffering from or possibly suffering from colorectal cancer, while the absence of hypermethylation of a determined percentage (e.g., a predetermined percentage) of methylation loci (e.g., at least 10% (e.g., at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, or 100%)) of a plurality of analyzed methylation loci constitutes a diagnosis that a subject is not likely suffering from colorectal cancer. In some embodiments, hypermethylation of a determined

number (e.g., a predetermined number) of methylation loci (e.g., at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, or 28 DMRs) of a plurality of analyzed methylation loci (e.g., 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, or 28 DMRs) constitutes a diagnosis that a subject is
5 suffering from or possibly suffering from colorectal cancer, while the absence of hypermethylation of a determined number (e.g., a predetermined number) of methylation loci (e.g., at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, or 28 DMRs) of a plurality of analyzed methylation loci (e.g., 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, or 28 DMRs) constitutes
10 a diagnosis that a subject is not likely suffering from colorectal cancer.

In some embodiments, methylation status of a plurality of methylation loci (e.g., a plurality of DMRs) is measured qualitatively or quantitatively and the measurement for each of the plurality of methylation loci are combined to provide a diagnosis. In some embodiments, the qualitative or quantitatively measured methylation status of each of a plurality of methylation
15 loci is individually weighted, and weighted values are combined to provide a single value that can be comparative to a reference in order to provide a diagnosis. To provide but one example of such an approach, support vector machine (SVM) algorithm can be used to analyze the methylation statuses of a plurality of methylation loci of the present disclosure to produce a diagnosis. At least one objective of the support vector machine algorithm is to
20 identify a hyperplane in an N-dimensional space (N—the number of features) that distinctly classifies the data points with the objective to find a plane that has the maximum margin, i.e. the maximum distance between data points of both classes. As discussed in the present Examples, an SVM model is built on marker values (e.g., Ct values) derived from a training sample set (e.g., the first subject group and/or the second subject group) that are
25 transformed to support vector values upon which a prediction is made. . In application of the SVM model to new samples, samples will be mapped onto vectorial space the model and categorized as having a probability of belonging to the first condition or the second condition, e.g., based on each new sample's location relative to the gap between the two conditions. Those of skill in the art will appreciate that, once relevant compositions and methods have
30 been identified, vector values can be used in conjunction with an SVM algorithm defined by predict () function of R-package(see Hypertext Transfer Protocol Secure (HTTPS)://cran.r-project.org/web/packages/e1071/index.html, the SVM of which is hereby incorporated by reference) to easily generate a prediction on a new sample. Accordingly, with compositions and methods for colorectal cancer diagnosis disclosed herein in hand (and only then),
35 generation of a predictive model utilizing algorithm input information in combination to

predict () function of R-package (see Hypertext Transfer Protocol Secure (HTTPS) ://cran.r-project.org/web/packages/e1071/index.html, the SVM of which is hereby incorporated by reference) to provide colorectal cancer diagnosis would be straightforward. By way of example, one non-limiting example of SVM vectors for use in diagnosis of colorectal cancer by analysis of methylation status of a plurality of DMRs provided herein is provided in Table 17. Those of skill in the art will appreciate that, with the present disclosure in hand, generation of SVM vectors can be accomplished according to methods provided herein and otherwise known in the art.

Table 13: Colorectal cancer DMR oligonucleotide primer pairs, e.g., for MSRE-qPCR

gene name	Fp_Seq	Rp_Seq	Fp SEQ ID	RP SEQ ID
ALK	CCTCCTCACCACATCATCAGCGCCC	GGTACCTCCCGCCGCTCTGTTC	45	46
CNRIP1	GCGTGCTGGGTTAATCTTCACCTCAA	ACGGCCCGGTCTTTTACAAGGTGG	47	48
LONRF2	AGGAAGCAAAGTGACCCCTAAGCCT	GGTCCGCTCCCTACACCT	49	50
LONRF2	CTCTCAGTCCCGCCGGCTTAGGTA	GCAAGAGACGCGGACCTGGAGC	51	52
ADAMTS2	CCACTGCGAAGGGAAGGGGCA	CCCTGTAAACGCCCTTCCCGTT	53	54
ADAMTS2	GCGACCCAGAAAGCCAGCCT	AACGGCTGGGGAGTGGCGGA	55	56
FGF14	CAACGAAACTTCCGCGCTAC	CTCGCCGGGGCTTCGCTAC	57	58
DMRT1	CAAAGCGTCTGGGGCGCTAGT	ACTTCTTGCTCCCGGCACCCAGGTC	59	60
ST6GALNAC5	CGCTCAGCCGCTCTCTCTTCTCT	AGCGCTAAACACACTGCCAGACCA	61	62
MCIDAS	GGGTTCGGAGCGTGCAAAAGGTGA	GAACAGTTCAGTGCATCCCCGCC	63	64
MCIDAS	GCGCCCACTTACATCCAGCACC	ACGTGACATTGACCCAGAAACAGGAGGA	65	66
PDGFD	AACGTCTATCACCAGGGAAGCT	TCCCGGAGTTGGCGAAAGTTGCAA	67	68
PDGFD	GGTGCAATTGGGATCAGCGACTAGAGAC	GATTAGCACAGCGACCCGGGCCAG	69	70
GSG1L	CCGAAAGAAATCCGAGCCAGGGTGA	GGTTTGTGTGCCCCACGTCC	71	72
ZNF492	CGAGAGAGGGGAAGGGGTGGTTG	CGAACTTGGGGCGCAGATTGTGG	73	74
ZNF568	GCCCAAGCCTACCCCTACACAG	CGAACCATCCCTCCGCGCCA	75	76
ZNF568	GGTCGCTTACCCAGCATCTCAG	CAGCGTACCTGCCGGAACACC	77	78
ZNF542	CCAGAGGCCAGGGATCCGTTTACAG	ACGCGAGCATCTTGTAAAGGCACCC	79	80
ZNF542	GGGAGGAGTGGGCGGTGAATGG	GCACCCGCCACCTCCAACTCAG	81	82
ZNF471	CCCCACGCTACTCACACCGAAG	GCGGGTAAGAGCAGGAGTGTG	83	84
ZNF471	GTCGCGGTTTCCCTCCAG	GCGGGTAAGAGCAGGAGTGTG	85	86
ZNF471	CTGCTCTTACCCGCCGAACCTG	GAGGGACCTTAGAGCAGAGCGGGC	87	88
ZNF132	CTACTGCTAGGTGCTGCAAGG	TGATTGGCCAGCGTCTTACACTCCG	89	90
ZNF132	GTGTAAGACGCTGGCCAATCACA	ACAACGCGGTCCCTTCAGAAGCAG	91	92
JAM2	CCGCGTGGTCTGGGCTCTGTAG	GAATCCCTCCACCTCCGCCCCAC	93	94
ZNF492	CAACGTTAAAGGCAACACCTTCTGC	GGCCGAATGAGGACAGAGTGACAG	95	96
CNRIP1	GCCGGTGAGCAGCTTGATGGT	ACGGCCCGGTCTTTACAAGG	97	98
ADAMTS2	CGGGAAGGGGCGTTAACAGGGC	TCTTGGCAGGAAGGTCTCCGGAG	99	100

10

Applications

Methods and compositions of the present disclosure can be used in any of a variety of

applications. For example, methods and compositions of the present disclosure can be used to screen, or aid in screening for, colorectal cancer. In various instances, screening using methods and compositions of the present disclosure can detect any stage of colorectal cancer, including without limitation early-stage colorectal cancer. In some embodiments, colorectal cancer screening using methods and compositions of the present disclosure is applied to individuals 50 years of age or older, e.g., 50, 55, 60, 65, 70, 75, 80, 85, or 90 years or older. In some embodiments, colorectal cancer screening using methods and compositions of the present disclosure is applied to individuals 20 years of age or older, e.g., 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, or 90 years or older. In some embodiments, colorectal cancer screening using methods and compositions of the present disclosure is applied to individuals 20 to 50 years of age, e.g., 20 to 30 years of age, 20 to 40 years of age, 20 to 50 years of age, 30 to 40 years of age, 30 to 50 years of age, or 40 to 50 years of age. In various embodiments, colorectal cancer screening using methods and compositions of the present disclosure is applied to individuals experiencing abdominal pain or discomfort, e.g., experiencing undiagnosed or incompletely diagnosed abdominal pain or discomfort. In various embodiments, colorectal cancer screening using methods and compositions of the present disclosure is applied to individuals experiencing no symptoms likely to be associated with colorectal cancer. Thus, in certain embodiments, colorectal cancer screening using methods and compositions of the present disclosure is fully or partially preventative or prophylactic, at least with respect to later or non-early stages of colorectal cancer.

In various embodiments, colorectal cancer screening using methods and compositions of the present disclosure can be applied to an asymptomatic human subject. As used herein, a subject can be referred to as "asymptomatic" if the subject does not report, and/or demonstrate by non-invasively observable indicia (e.g., without one, several, or all of device-based probing, tissue sample analysis, bodily fluid analysis, surgery, or colorectal cancer screening), sufficient characteristics of colorectal cancer to support a medically reasonable suspicion that the subject is likely suffering from colorectal cancer, and/or from cancer. Detection of early stage colorectal cancer is particularly likely in asymptomatic individuals screened in accordance with methods and compositions of the present disclosure.

In various embodiments, colorectal cancer screening using methods and compositions of the present disclosure can be applied to a symptomatic human subject. As used herein, a subject can be referred to as "symptomatic" if the subject report, and/or demonstrates by non-invasively observable indicia (e.g., without one, several, or all of device-based probing, tissue sample analysis, bodily fluid analysis, surgery, or colorectal cancer screening),

sufficient characteristics of colorectal cancer to support a medically reasonable suspicion that the subject is likely suffering from colorectal cancer, and/or from cancer. Symptoms of colorectal cancer can include, without limitation, change in bowel habits (diarrhea, constipation, or narrowing of the stool) that are persistent (e.g., lasting more than 3 days),
5 feeling of a need to have a bowel movement which feeling is not relieved upon bowel movement, rectal bleeding (e.g., with bright red blood), blood in stool (which can cause stool to appear dark), abdominal cramping, abdominal pain, weakness, fatigue, unintended weight loss, anemia, and combinations thereof. Those of skill in the art will appreciate that individual symptoms that would not alone indicate or raise a suspicion of colorectal cancer
10 may do so when presented in combination, e.g., a combination of abdominal cramping and blood in stool, to provide but one non-limiting example.

Those of skill in the art will appreciate that regular, preventative, and/or prophylactic screening for colorectal cancer improves diagnosis of colorectal cancer, including and/or particularly early stage cancer. As noted above, early stage cancers include, according to at
15 least one system of cancer staging, Stages 0 to II C of colorectal cancer. Thus, the present disclosure provides, among other things, methods and compositions particularly useful for the diagnosis and treatment of early stage colorectal cancer. Generally, and particularly in embodiments in which colorectal cancer screening in accordance with the present disclosure is carried out annually, and/or in which a subject is asymptomatic at time of screening,
20 methods and compositions of the present invention are especially likely to detect early stage colorectal cancer.

In various embodiments colorectal cancer screening in accordance with the present disclosure is performed once for a given subject or multiple times for a given subject. In various embodiments, colorectal cancer screening in accordance with the present disclosure
25 is performed on a regular basis, e.g., every six months, annually, every two years, every three years, every four years, every five years, or every ten years.

In various embodiments, screening for colorectal cancer using methods and compositions disclosed herein will be provide a diagnosis of colorectal cancer. In other instances, screening for colorectal cancer using methods and compositions disclosed herein will be
30 indicative of colorectal cancer diagnosis but not definitive for colorectal cancer diagnosis. In various instances in which methods and compositions of the present disclosure are used to screen for colorectal cancer, screening using methods and compositions of the present disclosure can be followed by a further diagnosis-confirmatory assay, which further assay can confirm, support, undermine, or reject a diagnosis resulting from prior screening, e.g.,
35 screening in accordance with the present disclosure. As used herein, a diagnosis-

confirmatory assay can be a colorectal cancer assay that provides a diagnosis recognized as definitive by medical practitioners, e.g., a colonoscopy-based diagnosis, or a colorectal cancer assay that substantially increases or decreases the likelihood that a prior diagnosis was correct, e.g., a diagnosis resulting from screening in accordance with the present disclosure. Diagnosis-confirmatory assays could include existing screening technologies, which are generally in need of improvement with respect to one or more of sensitivity, specificity, and non-invasiveness, particularly in the detection of early stage colorectal cancers.

In some instances, a diagnosis-confirmatory assay is a test that is or includes a visual or structural inspection of subject tissues, e.g., by colonoscopy. In some embodiments, colonoscopy includes or is followed by histological analysis. Visual and/or structural assays for colorectal cancer can include inspection of the structure of the colon and/or rectum for any abnormal tissues and/or structures. Visual and/or structural inspection can be conducted, for example, by use of a scope via the rectum or by CT-scan. In some instances, a diagnosis-confirmatory assay is a colonoscopy, e.g., including or followed by histological analysis. According to some reports, colonoscopy is currently the predominant and/or most relied upon diagnosis-confirmatory assay.

Another visual and/or structural diagnosis confirmatory assay based on computer tomography (CT) is CT colonography, sometimes referred to as virtual colonoscopy. A CT scan utilizes numerous x-ray images of the colon and/or rectum to produce dimensional representations of the colon. Although useful as a diagnosis-confirmatory assay, some reports suggest that CT colonography is not sufficient for replacement of colonoscopy, at least in part because a medical practitioner has not physically accessed the subject's colon to obtain tissue for histological analysis.

Another diagnosis-confirmatory assay can be a sigmoidoscopy. In sigmoidoscopy, a sigmoidoscope is used via the rectum to image portions of the colon and/or rectum. According to some reports, sigmoidoscopy is not widely used.

In some instances, a diagnosis-confirmatory assay is a stool-based assay. Typically, stool-based assays, when used in place of visual or structural inspection, are recommended to be utilized at a greater frequency than would be required if using visual or structural inspection. In some instances, a diagnosis-confirmatory assay is a guaiac-based fecal occult blood test or a fecal immunochemical test (gFOBTs/FITs) (see, e.g., Navarro 2017 World J Gastroenterol. 23(20):3632-3642, which is herein incorporated by reference with respect to colorectal cancer assays). FOBTs and FITs are sometimes used for diagnosis of colorectal cancer (see, e.g., Nakamura 2010 J Diabetes Investig. Oct 19;1(5):208-11, which is herein

incorporated by reference with respect to colorectal cancer assays). FIT is based on detection of occult blood in stool, the presence of which is often indicative of colorectal cancer but is often not in sufficient volume to permit identification by the unaided eye. For example, in a typical FIT, the test utilizes hemoglobin-specific reagent to test for occult blood

5 in a stool sample. In various instances, FIT kits are suitable for use by individuals in their own homes. When used in the absence of other diagnosis-confirmatory assays, FIT may be recommended for use on an annual basis. FIT is generally not relied upon to provide sufficient diagnostic information for conclusive diagnosis of colorectal cancer.

Diagnosis-confirmatory assays also include gFOBT, which is designed to detect occult blood

10 in stool by chemical reaction. Like FIT, when used in the absence of other diagnosis-confirmatory assays, gFOBT may be recommended for use on an annual basis. gFOBT is generally not relied upon to provide sufficient diagnostic information for conclusive diagnosis of colorectal cancer.

Diagnosis-confirmatory assays can also include stool DNA testing. Stool DNA testing for

15 colorectal cancer can be designed to identify DNA sequences characteristic of cancer in stool samples. When used in the absence of other diagnosis-confirmatory assays, stool DNA testing may be recommended for use every three years. Stool DNA testing is generally not relied upon to provide sufficient diagnostic information for conclusive diagnosis of colorectal cancer.

20 One particular screening technology is a stool-based screening test (Cologuard® (Exact Sciences Corporation, Madison, WI, United States), which combines an FIT assay with analysis of DNA for abnormal modifications, such as mutation and methylation. The Cologuard® test demonstrates improved sensitivity as compared to FIT assay alone, but can be clinically impracticable or ineffective due to low compliance rates, which low compliance

25 rates are at least in part due to subject dislike of using stool-based assays (see, e.g., doi: 10.1056/NEJMc1405215 (e.g., 2014 N Engl J Med. 371(2):184-188)). The Cologuard® test appears to leave almost half of the eligible population out of the screening programs (see, e.g., van der Vlugt 2017 Br J Cancer. 116(1):44-49). Use of screening as provided herein, e.g., by a blood-based analysis, would increase the number of individuals electing to screen

30 for colorectal cancer (see, e.g., Adler 2014 BMC Gastroenterol. 14:183; Liles 2017 Cancer Treatment and Research Communications 10: 27-31). To present knowledge, only one existing screening technology for colorectal cancer, Epirocolon, is FDA-approved and CE-IVD marked and is blood-based. Epirocolon is based on hypermethylation of SEPT9 gene. The Epirocolon test suffers from low accuracy for colorectal cancer detection with sensitivity

35 of 68% and advanced adenoma sensitivity of only 22% (see, e.g., Potter 2014 Clin Chem.

60(9):1183-91). There is need in the art for, among other things, a non-invasive colorectal cancer screen that will likely achieve high subject adherence with high and/or improved specificity and/or sensitivity.

In various embodiments, screening in accordance with methods and compositions of the present disclosure reduces colorectal cancer mortality, e.g., by early colorectal cancer diagnosis. Data supports that colorectal cancer screening reduces colorectal cancer mortality, which effect persisted for over 30 years (see, e.g., Shaukat 2013 N Engl J Med. 369(12):1106-14). Moreover, colorectal cancer is particularly difficult to treat at least in part because colorectal cancer, absent timely screening, may not be detected until cancer is past early stages. For at least this reason, treatment of colorectal cancer is often unsuccessful. To maximize population-wide improvement of colorectal cancer outcomes, utilization of screening in accordance with the present disclosure can be paired with, e.g., recruitment of eligible subjects to ensure widespread screening.

In various embodiments, screening of colorectal cancer including one or more methods and/or composition s disclosed herein is followed by treatment of colorectal cancer, e.g., treatment of early stage colorectal cancer. In various embodiments, treatment of colorectal cancer, e.g., early stage colorectal cancer, includes administration of a therapeutic regimen including one or more of surgery, radiation therapy, and chemotherapy. In various embodiments, treatment of colorectal cancer, e.g., early stage colorectal cancer, includes administration of a therapeutic regimen including one or more of treatments provided herein for treatment of stage 0 colorectal cancer, stage I colorectal cancer, and/or stage II colorectal cancer.

In various embodiments, treatment of colorectal cancer includes treatment of early stage colorectal cancer, e.g., stage 0 colorectal cancer or stage I colorectal cancer, by one or more of surgical removal of cancerous tissue e.g., by local excision (e.g., by colonoscopy), partial colectomy, or complete colectomy.

In various embodiments, treatment of colorectal cancer includes treatment of early stage colorectal cancer, e.g., stage II colorectal cancer, by one or more of surgical removal of cancerous tissue (e.g., by local excision (e.g., by colonoscopy), partial colectomy, or complete colectomy), surgery to remove lymph nodes near to identified colorectal cancer tissue, and chemotherapy(e.g., administration of one or more of 5-FU and leucovorin, oxaliplatin, or capecitabine).

In various embodiments, treatment of colorectal cancer includes treatment of stage III colorectal cancer, by one or more of surgical removal of cancerous tissue (e.g., by local excision (e.g., by colonoscopy-based excision), partial colectomy, or complete colectomy),

surgical removal of lymph nodes near to identified colorectal cancer tissue, chemotherapy(e.g., administration of one or more of 5-FU, leucovorin, oxaliplatin, capecitabine, e.g., in a combination of (i) 5-FU and leucovorin, (ii) 5-FU, leucovorin, and oxaliplatin (e.g., FOLFOX), or (iii) capecitabine and oxaliplatin (e.g., CAPEOX)), and
5 radiation therapy.

In various embodiments, treatment of colorectal cancer includes treatment of stage IV colorectal cancer, by one or more of surgical removal of cancerous tissue (e.g., by local excision (e.g., by colonoscope), partial colectomy, or complete colectomy), surgical removal of lymph nodes near to identified colorectal cancer tissue, surgical removal of metastases,
10 chemotherapy (e.g., administration of one or more of 5-FU, leucovorin, oxaliplatin, capecitabine, irinotecan, VEGF- targeted therapeutic agent (e.g., bevacizumab, ziv-aflibercept, or ramucirumab), EGFR-targeted therapeutic agent (e.g., cetuximab or panitumumab), Regorafenib, trifluridine, and tipiracil, e.g., in a combination of or including (i) 5-FU and leucovorin, (ii) 5-FU, leucovorin, and oxaliplatin (e.g., FOLFOX), (iii) capecitabine
15 and oxaliplatin (e.g., CAPEOX), (iv) leucovorin, 5-FU, oxaliplatin, and irinotecan (FOLFOXIRI), and (v) trifluridine and tipiracil (Lonsurf)), radiation therapy, hepatic artery infusion (e.g., if cancer has metastasized to liver), ablation of tumors, embolization of tumors, colon stent, colectomy, colostomy (e.g., diverting colostomy), and immunotherapy (e.g., pembrolizumab).

Those of skill in the art that treatments of colorectal cancer provided herein can be utilized, e.g., as determined by a medical practitioner, alone or in any combination, in any order, regimen, and/or therapeutic program. Those of skill in the art will further appreciate that advanced treatment options may be appropriate for earlier stage cancers in subjects previously having suffered a cancer or colorectal cancer, e.g., subjects diagnosed as having
25 a recurrent colorectal cancer.

In some embodiments, methods and compositions for colorectal cancer screening provided herein can inform treatment and/or payment (e.g., reimbursement for or reduction of cost of medical care, such as screening or treatment) decisions and/or actions, e.g., by individuals, healthcare facilities, healthcare practitioners, health insurance providers, governmental
30 bodies, or other parties interested in healthcare cost.

In some embodiments, methods and compositions for colorectal cancer screening provided herein can inform decision making relating to whether health insurance providers reimburse a healthcare cost payer or recipient (or not), e.g., for (1) screening itself (e.g., reimbursement for screening otherwise unavailable, available only for periodic/regular screening, or available
35 only for temporally- and/or incidentally- motivated screening); and/or for (2) treatment,

including initiating, maintaining, and/or altering therapy, e.g., based on screening results. For example, in some embodiments, methods and compositions for colorectal cancer screening provided herein are used as the basis for, to contribute to, or support a determination as to whether a reimbursement or cost reduction will be provided to a healthcare cost payer or recipient. In some instances, a party seeking reimbursement or cost reduction can provide results of a screen conducted in accordance with the present specification together with a request for such reimbursement or cost reduction of a healthcare cost. In some instances, a party making a determination as to whether or not to provide a reimbursement or cost reduction of a healthcare cost will reach a determination based in whole or in part upon receipt and/or review of results of a screen conducted in accordance with the present specification.

For the avoidance of any doubt, those of skill in the art will appreciate from the present disclosure that methods and compositions for colorectal cancer diagnosis of the present specification are at least for in vitro use. Accordingly, all aspects and embodiments of the present disclosure can be performed and/or used at least in vitro.

Kits

The present disclosure includes, among other things, kits including one or more compositions for use in colorectal cancer screening as provided herein, optionally in combination with instructions for use thereof in colorectal cancer screening. In various embodiments, a kit for screening of colorectal cancer can include one or more of: one or more oligonucleotide primers (e.g., one or more oligonucleotide primer pairs, e.g., as found in Table 13), one or more MSREs, one or more reagents for qPCR (e.g., reagents sufficient for a complete qPCR reaction mixture, including without limitation dNTP and polymerase), and instructions for use of one or more components of the kit for colorectal cancer screening. In various embodiments, a kit for screening of colorectal cancer can include one or more of: one or more oligonucleotide primers (e.g., one or more oligonucleotide primer pairs, e.g., as found in Table 13), one or more bisulfite reagents, one or more reagents for qPCR (e.g., reagents sufficient for a complete qPCR reaction mixture, including without limitation dNTP and polymerase), and instructions for use of one or more components of the kit for colorectal cancer screening.

In certain embodiments, a kit of the present disclosure includes at least one oligonucleotide primer pair for amplification of a methylation locus and/or DMR as disclosed herein.

In some instances, a kit of the present disclosure includes one or more oligonucleotide primer pairs for amplification of one or more methylation loci of the present disclosure. In

some instances, kit of the present disclosure includes one or more oligonucleotide primer pairs for amplification of one or more methylation loci that are or include all or a portion of one or more genes provided in Table 1. In some particular instances, a kit of the present disclosure includes oligonucleotide primer pairs for a plurality of methylation loci that each
5 are or include all or a portion of a gene identified in Table 1, the plurality of methylation loci including, e.g., 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, or 16 methylation loci, e.g., as provided in any of Tables 1 to 6.

In some instances, a kit of the present disclosure includes one or more oligonucleotide primer pairs for amplification of one or more DMRs of the present disclosure. In some
10 instances, kit of the present disclosure includes one or more oligonucleotide primer pairs for amplification of one or more DMRs that are, include all or a portion of, or are within a gene identified in Table 1. In some particular embodiments, a kit of the present disclosure includes oligonucleotide primer pairs for a plurality of DMRs each of which is, includes all or a portion of, or is within a gene identified in Table 1, e.g., 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13,
15 14, 15, or 16 DMRs, e.g., in accordance with any one of Tables 1 to 6..

In some instances, kit of the present disclosure includes one or more oligonucleotide primer pairs for amplification of one or more DMRs of Table 7. In some particular instances, a kit of the present disclosure includes oligonucleotide primer pairs for a plurality of DMRs of Table 7, the plurality of DMRs including, e.g., 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17,
20 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, or 28 DMRs of Table 7, e.g., as provided in any of Tables 8 to 12.

In various embodiments, a kit of the present disclosure includes one or more oligonucleotide primer pairs provided in Table 13. Those of skill in the art will appreciate that oligonucleotide primer pairs provided in Table 13 can be provided in any combination of one or more
25 oligonucleotide primer pairs, e.g., in a combination as provided in any one of Tables 1-12.

In various particular embodiments, kit of the present disclosure does not include oligonucleotide primer pairs that amplify all or a portion of one or more of FGF14, ZNF471, PDGFD, and ALK.

A kit of the present disclosure can further include one or more MSREs individually or in a single solution. In various embodiments, one or more MSREs are selected from the set of
30 MSREs including Acil, Hin6I, HpyCH4IV, and HpaII (e.g., such that the kit includes Acil, Hin6I, and HpyCH4IV, either individually or in a single solution). In certain embodiments, a kit of the present disclosure includes one or more reagents for qPCR (e.g., reagents sufficient for a complete qPCR reaction mixture, including without limitation dNTP and
35 polymerase).

EXAMPLES

The present Examples confirm that the present disclosure provides methods and compositions for, among other things, screening for and treatment of colorectal cancer. The present Examples further demonstrate that compositions and methods provided herein provide a remarkably high degree of sensitivity and specificity in screening and/or treatment of colorectal cancer. Also provided are clinical studies comparing methylation of biomarkers in samples from subjects diagnosed as having colorectal cancer and methylation of biomarkers in samples from control subjects, further demonstrating screening for colorectal cancer including methods and/or compositions of the present disclosure. Except as specifically stated otherwise, samples of the present Examples are humans or of human origin. With the exception of Example 1, all experiments were performed using plasma samples.

Example 1. Identification of methylation biomarkers associated with colorectal cancer

The present Example includes identification of CpG loci that are hypermethylated in one or more of colon cancer and rectal cancer as compared to healthy controls. In particular, experiments of the present Example examined CpG methylation in samples from (i) colon cancers of 341 subjects previously diagnosed as suffering from colon cancer, which subjects had not been previously treated by chemotherapy or radiotherapy; (ii) rectal cancers of 118 subjects previously diagnosed as suffering from rectal cancer, which subjects had not been previously treated by chemotherapy or radiotherapy; (iii) colons of 40 healthy control subjects not diagnosed as suffering from colorectal cancer; and (iv) leukocytes of 10 healthy control subjects not diagnosed as suffering from a colorectal cancer. Tissue samples were of fresh frozen tissue.

Samples were analyzed for DNA methylation by a global methylomics analysis platform (Infinium HumanMethylation450 (HM450) beadarray). The Infinium HumanMethylation450 array assesses methylation status of >450 000 CpGs located throughout the genome. DNA methylation profiles were obtained from all the tissue samples.

CpG methylation sites for which methylation status did not substantially differ between a colorectal cancer and healthy controls were identified and removed from consideration (mean b-value < 0.25 and b-value > 0.3 in no more than five samples across the entire set). This filtering produced a list of CpG methylation sites for which methylation status substantially differed between a colorectal cancer and healthy controls. The resulting set of CpG methylation sites was then further filtered by excluding CpG methylation sites with a mean b-value difference equal to or less than 0.1, yielding 253 CpG methylation sites. Each

of the 253 CpG methylation sites was associated with a hypermethylated status in colorectal cancer status as compared to controls.

Thus, the present Example generated a set of 253 individual CpG methylation sites that are methylation biomarkers for colorectal cancer. The 253 methylation biomarkers represent a plurality of DMRs together found within 36 genes, i.e., within 36 methylation loci. Each of the 36 DMRs is hypermethylated in colorectal cancer as compared to healthy controls.

Example 2: Development of cell-free DNA assay for methylation biomarkers by MSRE-qPCR

The present Example develops an assay for determining the methylation status of colorectal cancer methylation biomarkers based on circulating cell free DNA (cfDNA). cfDNA is incomplete and fragmented, and the mechanism by which the cfDNA is transmitted from cancer cells to blood (as a portion called circulating tumor DNA) is unknown. At least because the 253 methylation biomarkers of Example 1 were identified from tissue samples, it was not known prior to the experiments of the present Example whether identified colorectal cancer methylation biomarkers could be sufficiently analyzed from cfDNA to successfully capture the ctDNA portion that allows for identifying subjects or samples for colorectal cancer.

As a critical step toward determining whether colorectal cancer methylation biomarkers identified in Example 1 could be sufficiently analyzed from cfDNA to successfully capture the ctDNA portion that allows for identification of subjects or samples for colorectal cancer, a sensitive assay was developed for screening of these biomarkers. In particular, a Methylation-Sensitive Restriction Enzyme (MSRE)-qPCR methodology was developed. The MSRE-qPCR methodology was developed to measure methylation of DMRs covering identified CpG sites in blood samples, in particular in cell-free DNA (cfDNA) of tumors present in blood.

Development of the MSRE-qPCR methodology was significant at least in part because analyzing CpG methylation biomarkers derived from tumor tissue by analysis of cfDNA is challenging due to the low concentration of tumor-derived DNA circulating in blood (0.1 - 1%) as compared to the non-tumor DNA background of the sample. Thus, while it is generally preferred to develop biomarker analyses that rely on readily obtainable samples such as blood, urine, or stool, use of blood for analysis of tumor derived methylation biomarkers is challenging. Thus, even after identification of methylation biomarkers characteristic of colorectal cancer in tissue, as discussed above, it cannot be predicted whether the fragmented and poorly understood nature of ctDNA will permit successful screening using

methylation biomarkers identified in tissue.

MSRE-qPCR requires design of oligonucleotide primers (MSRE-qPCR oligonucleotide primer pairs) that amplify loci that each include at least one colorectal cancer MSRE cleavage site (i.e., an MSRE cleavage site that covers at least one colorectal cancer methylation biomarker site, such that cleavage of the MSRE cleavage site is permitted in nucleic acid molecules where all of the at least one colorectal cancer methylation biomarker sites are unmethylated and blocked in nucleic acid molecules where at least one of the at least one colorectal cancer methylation biomarker sites is methylated). MSRE-qPCR assays can utilize multiple restriction enzymes to enhance the range of colorectal cancer methylation biomarker sites that can be assayed by a single MSRE-qPCR reaction, as a single MSRE is unlikely to cleavage sites that together include all methylation biomarker sites of interest. MSRE-qPCR assays of the present Examples utilize the MSREs Acil, Hin6I, and HpyCH4IV, which together were found to provide sufficient coverage.

An exemplary schematic work flow for MSRE-qPCR is provided in Figure 1. As performed in the present Examples, circulating cell-free tumor DNA was extracted from subject blood (typically a plasma sample of approximately 4 mL) by QIAamp MinElute ccfDNA Kit in accordance with manufacturer protocol (QIAamp MinElute ccfDNA Handbook 08/2018, Qiagen). As shown in Figure 1, isolated cfDNA was divided into two aliquots, a first of which aliquots is utilized in a qPCR quality control analysis, and a second of which aliquots is used in MSRE-qPCR.

For MSRE-qPCR, 2/3 of eluted cfDNA by volume was digested with MSREs. Because non-methylated DNA is selectively cleaved, contacting the cfDNA with the MSREs enriches the sample for methylation-derived signal; methylated DNA remains intact and quantifiable. The remaining 1/3 of eluted cfDNA by volume was used for qPCR using the MSRE-qPCR oligonucleotide primers to confirm that amplicons were successfully amplified from cfDNA, which amplification confirms that template is present, hence providing technical quality control.

As applied herein, MSRE-qPCR oligonucleotide primer pairs were successfully developed for amplification of DMRs together including 180 of the 253 CpG methylation biomarker sites identified in Example 1. DMRs typically included 1 to 15 MSRE cleavage sites, which MSRE cleavage sites together covered each of the 180 methylation biomarker sites. As applied herein, methylation status of six genes (JUB, H19, TBP, TCEB2, SNRPN, IRF4) provided a methylation control, which control permitted monitoring of assay robustness and reproducibility.

Example 3: MSRE-qPCR of cfDNA Successfully Distinguishes Subjects by Colorectal Cancer Status

To probe clinical diagnostic and prognostic power of identified methylation biomarkers, the DMRs amplified by the MSRE-qPCR oligonucleotide primer pairs covering the 180 methylation biomarker sites, and appropriate controls, were assayed in cfDNA extracted from plasma of human subjects. Subjects were undiagnosed individuals seeking, or in the process of obtaining, a diagnosis regarding possible colorectal cancer, such that methylation biomarker analysis could be performed prior to traditional diagnostic testing for colorectal cancer and then compared to a subsequent traditional diagnosis. In particular, cfDNA was sampled from undiagnosed individuals seeking, or in the process of obtaining, a diagnosis regarding possible colorectal cancer at screening centers and oncology clinics in Spain and the United States between 2017 and 2018. A first subject group included 70 such individuals (see description of the first subject group in Figure 2), and a second subject group included 63 such individuals (see description of second subject group in Figure 3). Initial results based on MSRE-qPCR analysis of a small panel of tested DMRs of genes shown in Table 14 in the second subject group provided proof-of-principle for colorectal cancer diagnosis: results demonstrated overall diagnostic sensitivity of 80%, for colorectal cancer, with diagnostic sensitivity of up to 75% for early localized colorectal cancer, and 90% specificity (Figure 4). The representative proof-of-principle panel of DMRs performed similarly, and/or statistically equally, well with respect to proximal cancers and distal cancers (Figure 4). The proof-of-principle panel of DMRs also performed similarly, and/or statistically equally, well with respect to localized cancer and advanced cancer (Figure 4). Moreover, MSRE-qPCR analysis of methylation of MSRE-qPCR control genes and of undigested DNA controls showed high technical reliability of the developed MSRE-qPCR assays in measuring colorectal cancer biomarker methylation status in plasma cfDNA.

Figures 5-9 show the association of methylation status of colorectal cancer DMRs with colorectal cancer. Results are displayed as the MSRE-qPCR Ct value subtracted from 45 (i.e., $45 - \text{Ct value}$) for display purposes. Results demonstrate the surprisingly high predictive power of individual colorectal cancer methylation biomarkers, or of as few as three individual colorectal cancer methylation biomarkers, of the present disclosure for colorectal cancer, e.g., for use in determining screening for colorectal cancer in a subject.

Table 14: Proof-of-principle DMR panel

gene	chr	start	end	width	uid
PDGFD	chr11	104163499	104164026	528	PDGFD_chr11_104163499_104164026
FGF14	chr13	101919879	102403137	483259	FGF14_chr13_101919879_102403137
ALK	chr19	58439728	58440994	1267	ALK_chr19_58439728_58440994
LONRF2	chr2	29193215	29922286	729072	LONRF2_chr2_29193215_29922286
JAM2	chr2	68293114	68320928	27815	JAM2_chr2_68293114_68320928

Example 4. Further validation of methylation biomarkers by MSRE-qPCR

To verify the predictive power of methylation biomarker DMRs for colorectal cancer, data derived from MSRE-qPCR analysis of samples from the 133 subjects of the first and second subject groups identified in Example 3 (see Figures 2 and 3) were further analyzed. Monte-Carlo cross-validation was used over 50 runs and random forest algorithm was used for feature ranking and markers with VIP >2 were used for building a support-vector machine (SVM) algorithm-based classification model. This analysis identified several subsets of markers (2,3,5,8,15,28 as described in Tables 7-12) that in SVM-model gave a good prediction.

All models (2, 3, 5, 8, 15 and 28 colorectal cancer DMR panels) were applied to cfDNA extracted from plasma of a third subject group. The third subject group included 82 subjects who had either previously received a confirmed diagnosis of colorectal cancer or were control subjects known to not have colorectal cancer deemed (the control group including subjects having hyperplastic polyps and/or non-advanced adenoma, but not colorectal cancer), based on colonoscopy screening. . The 82 subjects were subjects attending colorectal cancer screening and oncology units in Spain and the United States. Further description of the third subject group is shown in Figure 10.

Oligonucleotide primer pairs (Table 13) for amplification of the 28 DRMs in MSRE-qPCR cover at least one MSRE cleavage site, typically 3 to 15 MSRE cleavage sites. MSRE-qPCR was carried out according to the methodology described in Example 2.

Notwithstanding the sufficiency and utility of all tested panels for screening for colorectal cancer, those of skill in the art will appreciate that the 28-DMR panel provided increased sensitivity and comparable specificity as compared to all other DMR panels indicated in Table 15. For the avoidance of any doubt, all of the panels tested, e.g., as described in Tables 7-14, are individually alone sufficient (e.g., in both sensitivity and specificity), and useful, for clinical screening of colorectal cancer. Analysis of the third subject group using the 28 colorectal cancer DMR panel showed general sensitivity for diagnosis of colorectal cancer of 79%, with 75% sensitivity for localized (early) cancer and 84% sensitivity for advanced cancer. Data also revealed specificity of 87% at AUC 82% (Figure 11). A ROC curve analysis of the second validation group data based on a 28-marker panel identified by

the SVM model is provided in Figure 11.

- Thus, evaluation of the performance of the 28 colorectal cancer DMR panel and subsets thereof reveal that both the full panel and each of the various subsets of 2, 3, 5, 8, and 15 of the 28 colorectal cancer DMRs are individually sufficient for clinical screening of colorectal cancer (See Tables 7-14). For instance, to highlight just one example, the 3-DMR subset (Table 9) achieved good separation of colorectal cancer subjects from control subjects, demonstrating sufficient performance for clinical screening of colorectal cancer, at least in part as demonstrated by the determined sensitivity of 60% and specificity of 87% (Table 15). SVM- model characteristics are described in Table 16. Input support SVM Vectors and their
- coefficient (weight) values are given in Table 17 (due to the size of Table 17, Table 17 is presented in several portions, with the coefficients and gene names repeated in each portion for reference). For prediction purposes the provided information was used in combination with predict() function in R-package (see Hypertext Transfer Protocol Secure (HTTPS) [://cran.r-project.org/web/packages/e1071/index.html](https://cran.r-project.org/web/packages/e1071/index.html)).
- Table 15. Accuracy metrics for application of 28 colorectal cancer DMR panel and subsets thereof to third subject group

	2	3	5	8	15	28
AUC	0.77	0.82	0.83	0.83	0.82	0.82
AUC_CI_LOW	0.67	0.73	0.73	0.74	0.73	0.72
AUC_CI_HIGH	0.87	0.91	0.92	0.92	0.92	0.92
Sensitivity	0.63	0.60	0.65	0.53	0.72	0.79
Specificity	0.79	0.87	0.82	0.87	0.79	0.87

Table 16 SVM-model input characteristics

vars	value
type	0
kernel	2
cost	1
degree	3
gamma	0.035714
coef0	0
nu	0.5
epsilon	0.1
nclasses	2
rho	-0.73551
probA	-2.14936
probB	-0.10114
sigma	0

Table 17: SVM Vectors

coefs	GSG1L	ZNF492_2	ZNF568_2	ZNF568_1	ZNF542_2
	GSG1L '861	ZNF492 '499	ZNF568 '252	ZNF568 '405	ZNF542 '525
1	0.166333806	0.423405118	-0.939531248	0.275236404	-0.052862411
1	0.183718034	-2.19255911	-0.092363545	-0.327541264	0.309450278
0.160992568	-0.094429617	-2.19255911	-0.939531248	-0.413652359	0.137466406
1	0.475386751	0.318930431	-0.147777884	0.019978514	0.164983826
1	-0.152377044	0.473593938	1.146062978	0.429006217	0.574305439
1	0.315065536	0.442866089	0.835406834	0.041506288	0.55940017
0.539110203	-0.428593114	0.344536972	0.864793226	-0.0507556	0.174156299
1	0.143154835	-2.19255911	0.848840613	-0.373672208	0.025103611
0.228841426	-0.16396653	0.392677269	-0.939531248	-0.0507556	0.161544148
1	0.691723813	0.288202582	-0.939531248	0.232180856	0.641952428
1	0.152812739	0.548365038	0.897538063	0.321367348	0.6259006
0.086502196	0.152812739	0.413162502	-0.939531248	-0.318315075	0.260148235
1	0.058165275	0.455157229	-0.939531248	-0.035378619	0.421813073
0.022981588	-0.055797999	-2.19255911	-0.939531248	-0.469009492	0.310596837
0.199753156	0.029191561	0.425453641	-0.939531248	-0.192223828	0.392002536
0.640672506	0.401986676	0.512515881	0.731295044	-0.352144434	-2.050168434
0.488436741	0.11224954	-2.19255911	-0.939531248	-0.327541264	-2.050168434
0.290813367	0.038849466	0.455157229	0.842963334	0.038430892	0.370217912
0.93438076	-0.165898111	0.292299628	0.996612184	-0.293711905	0.182182213
1	0.34597083	0.531976852	-0.939531248	0.622756182	0.534175869
0.177133374	0.291886565	0.201140342	0.931122511	0.312141159	0.439011461
0.363673626	-0.252819252	0.199091819	-0.939531248	-0.232203979	0.408054364
0.372869951	-0.022961123	-2.19255911	-0.939531248	-0.438255529	-2.050168434
0.180416799	-0.0152348	0.246207854	-0.939531248	-0.247580961	0.130587052
1	1.837151292	1.131169911	-0.939531248	1.084065621	1.180835225
0.236974727	-4.54286045	0.032137172	-0.240135117	-0.619703909	-2.050168434
0.433173404	-4.54286045	-2.19255911	-0.939531248	-0.40442617	0.427545869
0.037394649	0.15474432	-2.19255911	-0.939531248	-0.204525413	0.471115116
1	0.245528623	0.455157229	-0.939531248	-0.207600809	0.380536944
0.263636311	-4.54286045	-2.19255911	-0.939531248	-1.72992196	-2.050168434
1	0.682065908	0.557583393	1.034394689	0.828807731	0.838014042
1	-0.042276933	0.55553487	0.775794438	0.158371346	0.349579848
1	0.025328399	0.467448369	0.848001002	-0.524366624	0.018224256
0.26169536	-4.54286045	-2.19255911	-0.939531248	-7.963750185	-2.050168434

Table 17 cont'd: SVM Vectors

coefs	GSG1L	ZNF492_2	ZNF568_2	ZNF568_1	ZNF542_2
	GSG1L '861	ZNF492 '499	ZNF568 '252	ZNF568 '405	ZNF542 '525
0.1799778	0.295749727	0.267717349	-0.939531248	0.226030064	0.467675439
0.898868481	-0.18521392	0.324051739	-0.939531248	-0.023077034	0.29339845
0.042908076	0.019533657	0.191921987	-0.939531248	-0.622779305	0.275053503
0.140898434	-4.54286045	0.204213127	-0.939531248	-0.567422172	0.123707697
0.140415448	0.293818146	-2.19255911	-0.939531248	-0.699664211	0.061793503
1	0.166333806	0.135587597	-0.939531248	-0.044604807	0.459649525
0.108835219	-0.378372011	0.333270094	-0.939531248	-0.604326927	-2.050168434
1	0.608665834	0.370143513	0.901736119	0.641208559	0.750875547
-1	0.322791859	-2.19255911	1.066299914	0.186049912	0.550227697
-1	0.025328399	-2.19255911	-0.497895756	0.026129307	0.433278665
-1	0.311202374	0.387555961	-0.939531248	-0.182997639	0.5192706
-1	-0.059661161	0.322003216	1.020121298	0.422855425	0.403468127
-1	0.403918257	0.612893521	0.959669292	0.468986368	0.589210708
-1	0.172128548	0.145830214	0.701908652	0.182974516	0.491753181
-1	-0.16396653	0.411113978	-0.939531248	-0.336767452	0.473408235
-1	0.336312926	0.694834453	-0.939531248	0.819581542	0.740556515
-0.27309104	1.360050807	0.923244798	1.68509337	2.378807448	1.22669759
-1	2.101777876	1.346264856	1.939495565	3.255295383	1.550027268
-1	0.712971203	0.404968409	0.776634049	0.475137161	-2.050168434
-0.675668601	0.382670867	0.677422005	1.048668079	0.579700634	0.674056084
-1	0.245528623	0.403944147	-0.939531248	-0.545894398	-2.050168434
-0.228886868	1.516508861	0.993918852	1.713640151	2.252716201	1.264534042
-1	0.523676274	0.505346049	0.922726399	0.272161008	0.602969418
-0.253709663	0.53719734	0.671276435	-0.939531248	0.186049912	0.764634257
-1	-0.007508476	-2.19255911	-0.939531248	-0.284485716	0.336967697
-1	0.465728846	0.297420936	-0.939531248	0.075335647	0.648831783
-1	0.206897005	0.52480702	0.981499183	0.066109458	0.583477912
-1	0.626050062	0.583189934	-0.939531248	0.804204561	0.654564579
-1	0.094865312	0.356828111	0.968065404	0.112240402	0.397735331
-1	0.04850737	-2.19255911	-0.939531248	-1.041033197	0.29339845
-1	-0.241229766	-2.19255911	0.913490675	-0.14916828	0.136319847
-1	0.419370905	-2.19255911	-0.939531248	0.099938817	0.659150816
-1	0.228144395	0.438769043	0.834567222	0.413629236	0.549081138
-1	0.45220778	0.585238457	0.886623117	-0.103037336	0.838014042
-1	0.237802299	0.444914613	-0.939531248	0.078411043	0.295691568

Table 17 cont'd: SVM Vectors

coefs	ZNF542_1	ZNF471_2	ZNF471_1	ZNF471_3	ZNF132_2
	ZNF542 '502	ZNF471 '527	ZNF471 '558	ZNF471 '662	ZNF132 '268
1	0.210023302	0.448781467	0.381358009	0.276869984	-0.079808848
1	0.096511033	-1.701659498	0.111027436	0.245615084	0.220719423
0.160992568	0.011755205	0.491015986	0.192253523	0.011882788	0.004011962
1	0.075322076	0.362552655	0.179561947	0.342097602	0.355650484
1	0.438561338	0.531490735	0.353436542	0.351609963	0.275918493
1	0.322022075	0.502454502	0.314092655	0.193976554	0.091921593
0.539110203	0.007214715	0.395108431	0.091990071	0.143696932	-0.206562268
1	0.010241709	0.604521258	0.298862764	0.238820541	0.200275323
0.228841426	0.010241709	-1.701659498	-0.009542539	-3.291624259	0.120543333
1	0.477912258	0.220891037	0.146563849	0.172234015	0.333161974
1	0.228185265	0.381030258	0.097066702	0.160003836	0.380183404
0.086502196	0.116186493	-1.701659498	-0.38775151	-0.122649173	-0.237228418
1	0.194888333	0.592202857	0.357244014	0.299971432	0.053077803
0.022981588	-0.13808099	-1.701659498	-0.060308843	-3.291624259	-0.186118168
0.199753156	0.15705091	0.495415416	0.260788035	0.227949271	-0.034831828
0.640672506	0.120726984	-1.701659498	0.215098361	0.210283458	0.183920043
0.488436741	0.05867361	-1.701659498	0.02853219	0.007806062	-0.341493329
0.290813367	0.063214101	0.449661352	0.10848912	0.248332902	0.433338064
0.93438076	0.043538641	0.359912998	0.034877978	-0.051985921	0.051033393
1	0.211536799	-1.701659498	-0.187224606	-0.012577568	0.482403904
0.177133374	0.264509192	0.375750943	0.046300397	-3.291624259	-7.69E-05
0.363673626	0.073808579	0.258726128	0.171947002	-0.049268103	-0.204517858
0.372869951	-3.407234345	0.527971191	0.279825399	0.138261297	-0.406914449
0.180416799	-3.407234345	-1.701659498	-0.136458301	-3.291624259	-0.406914449
1	1.345145995	1.182606248	1.282459921	1.285180239	1.347189337
0.236974727	-3.407234345	-1.701659498	-2.980640533	-0.383559643	-0.085942078
0.433173404	0.001160727	0.382790029	0.060261131	-0.069651734	-0.110474998
0.037394649	-0.468023319	0.373111285	0.013302299	-0.008500842	-0.257672518
1	0.455209804	0.5957224	0.409279477	0.446733572	0.001967552
0.263636311	-3.407234345	-1.701659498	-2.980640533	0.049932232	-5.066124854
1	0.721585263	0.52181199	0.325515074	0.344815419	0.791109815
1	0.060187107	0.342315281	0.10848912	-0.046550286	0.071477493
1	-0.038190193	0.419745234	0.317900128	0.173592923	-0.413047679
0.26169536	-3.407234345	0.529730963	0.213829203	-0.096829908	-0.366026249

Table 17 cont'd: SVM Vectors

coefs	ZNF542_1	ZNF471_2	ZNF471_1	ZNF471_3	ZNF132_2
	ZNF542 '502	ZNF471 '527	ZNF471 '558	ZNF471 '662	ZNF132 '268
0.1799778	0.155537413	-1.701659498	-0.034925691	0.191258736	-5.066124854
0.898868481	0.278130664	0.655554636	0.5463485	0.405966311	-0.192251398
0.042908076	0.079862567	0.491895872	0.225251622	0.179028558	0.020367243
0.140898434	0.051106125	-1.701659498	-0.243067541	-3.291624259	-5.066124854
0.140415448	-0.074514119	-1.701659498	-2.980640533	0.154568201	-0.304693949
1	0.128294469	0.220011151	-0.135189143	0.202130006	0.106232463
0.108835219	-3.407234345	-1.701659498	-2.980640533	-3.291624259	0.026500473
1	0.526344159	-1.701659498	-0.140265774	0.233384906	0.603024094
-1	0.368940479	0.572845368	0.377550536	0.036343145	0.177786813
-1	0.31445459	0.503334388	0.298862764	0.181746375	0.132809793
-1	0.240293241	0.61683966	0.287440345	0.473911746	0.378138994
-1	0.134348456	0.407426833	0.248096459	0.279587802	-0.165674068
-1	0.424939866	0.570205711	0.39151127	0.305407067	0.523292104
-1	0.347751522	0.558767195	0.404202846	0.324431789	0.210497373
-1	0.325049069	0.280723273	0.102143332	0.295894706	-0.286294258
-1	0.577803055	0.540289593	0.434662629	0.505166646	0.881063855
-0.27309104	1.379956424	0.847369747	0.820486547	0.710361859	1.046661066
-1	1.806762557	1.486166858	1.711435198	1.720031023	2.25286297
-1	0.343211032	0.614200002	0.400395374	0.426349941	0.654134345
-0.675668601	0.455209804	0.547328679	0.424509368	0.408684128	0.776798945
-1	-0.012460745	0.328237108	-0.031118218	-0.007141934	0.077610723
-0.228886868	1.384496915	1.151810244	1.269768345	1.328665317	1.330834057
-1	0.391642933	0.754981735	0.631382061	-3.291624259	0.118498923
-0.253709663	0.741260723	1.184366019	1.220271198	1.031064312	0.367916944
-1	-0.147161972	-1.701659498	-0.187224606	-0.133520443	0.065344263
-1	0.536938638	0.610680459	0.408010319	0.446733572	0.425160424
-1	0.188834345	-1.701659498	0.259518877	0.237461632	0.480359494
-1	0.421912872	0.484856786	0.344552438	0.318996154	0.692978135
-1	0.190347842	0.604521258	0.31663097	0.270075441	-0.024609778
-1	0.060187107	0.410946376	0.26332635	0.282305619	0.108276873
-1	0.043538641	0.546448794	0.279825399	0.100211854	-0.063453568
-1	0.443101829	0.402147518	0.429585999	0.274152167	0.055122213
-1	0.446128822	0.451421124	0.211290888	0.308124884	0.220719423
-1	0.774557655	0.410946376	0.221444149	0.145055841	0.717511055
-1	0.110132505	0.549968337	0.243019828	-0.111777904	0.273874083

Table 17 cont'd: SVM Vectors

coefs	ZNF132_1	JAM2	MCIDAS_1	MCIDAS_2	PDGFD_1
	ZNF132 '415	JAM2 '320	MCIDAS '855	MCIDAS '003	PDGFD '388
1	0.238896666	0.406419198	0.492859834	0.416450063	0.046762915
1	-0.397730773	-2.423671354	0.320276073	0.121123394	-0.124572283
0.160992568	-0.170552517	-2.423671354	-2.052243039	0.091026409	-0.261640442
1	0.347202579	0.464995238	0.485753444	0.245273459	0.426555938
1	0.120024323	0.545266109	0.761887461	0.544362251	-0.027482338
1	0.006435195	0.297945048	0.524330991	0.190722673	0.235231633
0.539110203	-0.693590827	0.228521593	0.262409754	0.038356684	-0.53577676
1	-0.550944015	0.287097633	0.044142056	-0.183608583	0.20382018
0.228841426	-0.352823443	0.214419954	0.199467441	0.028951376	0.121008168
1	0.299653641	0.447639374	0.58219731	0.501097835	0.423700352
1	0.26795435	0.317470395	0.572045325	0.252797705	0.266643087
0.086502196	-0.487545432	0.333741518	0.359868819	0.213295412	-0.16169491
1	0.088325031	0.395571783	0.40961355	0.333683353	0.126719341
0.022981588	-1.155872162	-2.423671354	-2.052243039	0.019546069	-0.669989332
0.199753156	-0.231309492	-2.423671354	0.225862604	0.043999869	-0.17882843
0.640672506	0.407959554	0.378215919	0.320276073	0.141815072	0.152419621
0.488436741	-0.410938811	-0.105578787	0.311139286	-4.590935879	-0.375863908
0.290813367	-0.049038566	0.15367443	-2.052243039	0.168149934	0.358021859
0.93438076	-0.614342598	0.187301416	-0.001541881	-0.078269134	-0.624299946
1	0.415884377	0.216589437	0.588288502	0.241511336	0.469389738
0.177133374	0.135873968	0.230691076	0.443115103	0.15122038	0.132430514
0.363673626	-0.025264097	0.130894859	0.318245676	0.072215793	-0.247362509
0.372869951	-0.281500037	0.267572287	0.200482639	-0.042528964	-0.190250776
0.180416799	-6.542110115	0.37930066	0.199467441	-0.245683615	-0.435831227
1	1.855560534	1.410889819	1.44105532	1.953277379	1.120463492
0.236974727	-0.71208208	-2.423671354	-2.052243039	-4.590935879	-0.344452455
0.433173404	0.104174677	-0.128358358	0.297941704	-0.078269134	0.292343366
0.037394649	-0.226026277	0.150420206	-2.052243039	0.049643054	-0.104583177
1	-0.424146849	0.275165477	-2.052243039	0.072215793	0.063896435
0.263636311	-6.542110115	-0.08605344	-2.052243039	-0.591798947	-7.103626036
1	0.69061157	0.407503939	0.605546878	0.314872738	1.134741426
1	-0.038472135	0.309877205	0.161905093	-0.241921492	0.295198953
1	-0.014697667	0.199233573	0.256318562	-0.24004043	-0.407275361
0.26169536	-0.878503361	0.122216927	0.038050865	-4.590935879	-0.415842121

Table 17 cont'd: SVM Vectors

coefs	ZNF132_1	JAM2	MCIDAS_1	MCIDAS_2	PDGFD_1
	ZNF132 '415	JAM2 '320	MCIDAS '855	MCIDAS '003	PDGFD '388
0.1799778	-0.125645187	0.492113775	0.581182112	0.352493969	-7.103626036
0.898868481	0.101533069	-2.423671354	0.41671994	0.02142713	-0.470098267
0.042908076	-0.281500037	-2.423671354	0.402507159	-0.345379879	-0.235940162
0.140898434	-0.532452762	0.177538743	-2.052243039	-0.040647902	-0.707111958
0.140415448	-0.136211617	-2.423671354	0.160889894	0.143696133	-0.147416977
1	0.006435195	0.180792967	0.542604565	0.326159107	-0.304474242
0.108835219	-0.292066467	-2.423671354	0.051248446	-0.388644296	-0.632866706
1	0.574380834	0.446554633	0.667473992	0.598913037	0.812060135
-1	0.196630944	0.384724368	0.345656038	0.218938597	0.363733032
-1	0.180781298	0.228521593	0.117236355	-0.119652489	0.546490577
-1	0.471358137	0.249131681	0.483723047	-0.149749474	0.483667671
-1	-0.022622489	0.469334204	0.406567954	0.145577195	-0.007493231
-1	0.574380834	0.309877205	0.617729261	0.132409764	0.466534151
-1	0.030209663	0.205742022	0.451236692	-0.003026671	-0.010348818
-1	0.02228484	0.213335212	0.123327546	0.237749213	0.078174368
-1	0.970621979	0.511639122	0.606562077	0.311110614	0.689269909
-0.27309104	0.981188409	1.000857535	1.309079503	1.5507302	1.465989476
-1	2.840880179	1.538889315	1.185225275	1.65795071	2.031395631
-1	0.574380834	0.560452489	0.650215616	0.561291805	0.777793095
-0.675668601	1.139684867	0.64072336	0.720264319	0.572578175	0.892016561
-1	-0.178477339	0.079912009	0.120281951	0.008259699	0.123863754
-0.228886868	1.945375193	1.1646535	1.014671911	1.069178434	1.683014061
-1	0.064550562	0.462825755	0.547680558	0.290418937	0.395144485
-0.253709663	0.79099266	0.783909237	0.578136516	0.538719066	0.709259016
-1	-0.276216822	-2.423671354	0.205558632	-0.010550917	-0.298763069
-1	0.36833544	0.443300408	0.530422182	0.094788532	0.400855659
-1	0.36833544	0.212250471	0.372051202	0.143696133	0.226664874
-1	0.761934976	0.514893347	0.615698864	0.350612908	0.717825776
-1	0.283803995	0.427029286	0.48169265	0.243392397	0.149564034
-1	-0.659249928	0.18296245	0.281698527	-0.371714741	0.118152581
-1	-0.241875922	0.190555641	0.327382464	-0.001145609	-0.116005524
-1	0.149082006	0.290351858	0.572045325	0.388234139	0.452256218
-1	0.347202579	0.420520837	0.494890231	0.164387811	0.466534151
-1	1.055153423	0.579977836	0.862392122	0.811472996	0.186686661
-1	-0.123003579	-2.423671354	0.184239462	0.29418106	0.443689458

Table 17 cont'd: SVM Vectors

coefs	PDGFD_2	ST6GALNAC5	ZNF492_1	CNRIP1_1	LONRF2_1
	PDGFD '921	ST6GALNAC5 '456	ZNF492 '069	CNRIP1 '272	LONRF2 '281
1	1.302789852	0.466406944	-0.240707793	-0.064165795	0.182108844
1	-0.682617737	0.569041467	-0.17761095	-0.307307386	-0.063816882
0.160992568	-0.682617737	0.705281983	-0.150569446	0.093547669	-0.189987472
1	-0.682617737	0.569041467	0.176933218	0.189928119	0.325387311
1	1.475318586	0.828806718	-0.132541776	0.498783653	0.197078236
1	-0.682617737	-1.277471664	0.08679487	0.38268902	0.201355206
0.539110203	-0.682617737	-1.277471664	-0.210661677	0.019071866	0.100846431
1	-0.682617737	0.367405503	0.056748754	-0.149593922	0.090154008
0.228841426	-0.682617737	0.246605578	0.194960887	0.161452077	-0.098032635
1	1.096640133	0.76522781	0.480398989	0.178975795	0.534958799
1	-0.682617737	0.792475913	0.065762589	0.119833246	0.338218218
0.086502196	1.281555546	-1.277471664	-0.327841529	-0.079499049	-0.207095348
1	-0.682617737	0.638978265	0.149891714	0.369546231	0.333941249
0.022981588	-0.682617737	-1.277471664	-0.360892257	-3.744146627	-0.585607118
0.199753156	-0.682617737	-1.277471664	-0.162587892	-0.042261148	-5.029378066
0.640672506	-0.682617737	0.586298599	0.23702545	-0.07073719	-0.22634171
0.488436741	1.209005001	-1.277471664	0.017688804	-0.348926217	-0.160048688
0.290813367	-0.682617737	0.638069995	0.110831763	-0.007213711	0.165000968
0.93438076	1.368262294	0.469131755	-0.57421968	-0.379592724	-0.014631737
1	-0.682617737	-1.277471664	0.423311368	0.316975076	0.498604562
0.177133374	-0.682617737	0.554509145	0.243034673	0.222785091	0.154308545
0.363673626	-0.682617737	0.566316656	-0.369906092	-0.186831823	-0.068093851
0.372869951	-0.682617737	-1.277471664	-0.231693959	-0.06635626	-0.435913198
0.180416799	-0.682617737	-1.277471664	-1.154109717	-2.41672497	-0.136525357
1	-0.682617737	1.508192759	1.468916202	2.071537366	0.917747538
0.236974727	-0.682617737	-1.277471664	-0.330846141	-4.848140877	-5.029378066
0.433173404	-0.682617737	0.542701634	0.056748754	-0.243783907	0.263371258
0.037394649	-0.682617737	-1.277471664	0.149891714	-0.362069006	-0.068093851
1	-0.682617737	0.620812863	0.336177632	0.045357444	-0.311881093
0.263636311	-0.682617737	-1.277471664	-7.154319065	-0.734448018	-5.029378066
1	1.438158551	-1.277471664	1.072307472	0.660878047	0.791576948
1	1.220506917	0.624445943	0.131864044	-0.048832542	-0.016770221
1	-0.682617737	0.609913622	-0.204652454	-0.160546246	-0.269111401
0.26169536	-0.682617737	-1.277471664	-7.154319065	0.207451837	-5.029378066

Table 17 cont'd: SVM Vectors

coefs	PDGFD_2	ST6GALNAC5	ZNF492_1	CNRIP1_1	LONRF2_1
	PDGFD '921	ST6GALNAC5 '456	ZNF492 '069	CNRIP1 '272	LONRF2 '281
0.1799778	-0.682617737	-1.277471664	0.534481997	0.062881162	0.607667275
0.898868481	-0.682617737	0.698015822	-0.021371147	0.268784851	-0.040293551
0.042908076	-0.682617737	-1.277471664	-0.339859976	-0.567972695	-0.326850485
0.140898434	-0.682617737	-1.277471664	-0.571215068	-0.537306188	-0.442328652
0.140415448	-0.682617737	-1.277471664	-0.144560222	-3.146149742	0.111538853
1	-0.682617737	-1.277471664	-0.021371147	0.178975795	0.505020015
0.108835219	-0.682617737	-1.277471664	-0.318827695	-4.848140877	-0.463713497
1	-0.682617737	-1.277471664	0.429320592	0.560116667	0.757361195
-1	1.29571175	-1.277471664	0.092804093	-0.136451133	0.107261884
-1	-0.682617737	0.76613608	0.393265253	-0.094832302	0.220601567
-1	-0.682617737	-1.277471664	0.324159186	0.260022992	0.415203663
-1	1.442582365	0.811549586	-0.108504883	0.424307851	0.248401866
-1	-0.682617737	0.770677431	0.579551171	0.380498555	0.637606059
-1	1.147071609	-1.277471664	-0.084467991	-0.241593443	0.094430977
-1	-0.682617737	0.633528644	0.342186855	0.216213697	0.081600069
-1	-0.682617737	0.821540557	0.792878594	0.619259216	0.872839362
-0.27309104	2.042451502	1.427356719	1.264602614	1.732015325	1.51224625
-1	-0.682617737	1.5981115	2.0127509	2.529344505	2.128129808
-1	1.308983191	0.435525761	0.480398989	0.316975076	0.733837865
-0.675668601	-0.682617737	0.837889419	0.726777139	0.551354808	0.63332909
-1	-0.682617737	-1.277471664	0.083790259	-0.06635626	-0.040293551
-0.228886868	1.923893295	1.317456036	1.381782466	1.839348099	1.349721423
-1	-0.682617737	0.856963091	0.315145351	0.343260654	0.259094289
-0.253709663	1.235547883	1.01500209	0.964141455	1.026685666	0.729560896
-1	1.354106091	-1.277471664	-0.048412651	-0.099213232	-0.076647789
-1	-0.682617737	0.617179782	0.393265253	0.303832288	0.505020015
-1	1.269168867	0.629895564	0.342186855	0.481259935	0.421619117
-1	1.248819324	0.831531528	0.315145351	0.308213217	0.673960297
-1	1.336410836	0.788842833	-0.168597115	0.371736696	0.188524298
-1	1.133800168	0.567224926	-0.126532553	-0.134260668	-0.001800829
-1	-0.682617737	-1.277471664	-0.105500272	-0.090451373	0.216324598
-1	-0.682617737	0.717997765	0.390260641	0.411165062	0.513573954
-1	-0.682617737	0.735254897	0.231016227	0.384879485	0.541374253
-1	-0.682617737	1.01681863	0.231016227	1.147161228	0.703899081
-1	1.284209834	-1.277471664	0.158905548	-0.103594161	-0.574914695

Table 17 cont'd: SVM Vectors

coefs	LONRF2_2	ADAMTS2_2	ADAMTS2_1	ADAMTS2_3	ALK
	LONRF2 '387	ADAMTS2 '254	ADAMTS2 '284	ADAMTS2 '328	ALK '434
1	0.816278101	-1.226540092	0.479237716	0.308464228	1.224961864
1	0.707526931	0.608032885	0.377972809	0.108720604	-0.67135809
0.160992568	-1.17952336	-1.226540092	-1.402761026	-3.569757964	-0.67135809
1	0.644016248	0.715511247	0.413319994	0.342021157	-0.67135809
1	1.028560383	0.889956898	0.805960341	0.757487893	-0.67135809
1	0.704916903	0.774210969	0.512674242	0.359598595	-0.67135809
0.539110203	-1.17952336	0.74444773	0.34931293	-0.159734825	-0.67135809
1	-1.17952336	-1.226540092	-1.344485938	-0.001537876	1.290071562
0.228841426	-1.17952336	0.651850986	0.265243951	0.155061125	-0.67135809
1	0.849338456	0.647717203	0.611073161	0.319649871	1.311232214
1	0.788437801	0.675826929	0.455354484	0.27330935	-0.67135809
0.086502196	0.548315219	-1.226540092	0.357910894	0.155061125	1.30146576
1	0.564845397	0.690708548	-1.172526662	0.340423208	1.271352524
0.022981588	-1.17952336	-1.226540092	-1.818329276	-0.437777949	-0.67135809
0.199753156	0.650976323	0.717164761	0.579547294	0.310062177	-0.67135809
0.640672506	-1.17952336	0.763463132	0.44580119	0.290886789	-0.67135809
0.488436741	-1.17952336	0.617127208	0.287216525	-0.153343029	-0.67135809
0.290813367	0.570935463	0.735353407	0.491656997	0.316453973	-0.67135809
0.93438076	-1.17952336	0.666732606	0.444845861	0.169442666	-0.67135809
1	-0.840219712	0.793226371	0.47350574	0.389959626	-0.67135809
0.177133374	-1.17952336	-1.226540092	0.355044906	0.129493941	-0.67135809
0.363673626	0.538745116	-1.226540092	0.304412453	-0.142157387	-0.67135809
0.372869951	-1.17952336	-1.226540092	0.354089577	0.111916502	-0.67135809
0.180416799	0.499594695	-1.226540092	-1.818329276	-3.569757964	-0.67135809
1	0.880658793	1.437269791	1.395398525	1.705071643	1.21519541
0.236974727	-1.17952336	-1.226540092	-1.818329276	-3.569757964	-0.67135809
0.433173404	-1.17952336	-1.226540092	0.347402271	0.057586237	-0.67135809
0.037394649	-1.17952336	-1.226540092	-1.818329276	-3.569757964	-0.67135809
1	-1.17952336	0.671693146	0.532736158	-0.081435325	-0.67135809
0.263636311	-1.17952336	-1.226540092	-1.818329276	-3.569757964	-0.67135809
1	0.846728428	0.660118553	0.491656997	0.417124759	1.282746721
1	0.650106314	-1.226540092	0.401856042	-0.322725622	-0.67135809
1	0.640536211	-1.226540092	0.567128013	0.254133962	-0.67135809
0.26169536	0.519604911	-1.226540092	0.536557475	0.284494993	-0.67135809

Table 17 cont'd: SVM Vectors

coefs	LONRF2_2	ADAMTS2_2	ADAMTS2_1	ADAMTS2_3	ALK
	LONRF2 '387	ADAMTS2 '254	ADAMTS2 '284	ADAMTS2 '328	ALK '434
0.1799778	0.525694976	0.613820181	0.533691487	0.266917554	-0.67135809
0.898868481	-1.17952336	0.719645031	0.407588018	-0.172518417	1.353553518
0.042908076	0.605735837	0.541065597	0.508852925	0.28928884	-0.67135809
0.140898434	-1.17952336	-1.226540092	-1.818329276	-3.569757964	-0.67135809
0.140415448	-1.17952336	-1.226540092	-1.818329276	-0.332313316	-0.67135809
1	0.771037614	-1.226540092	0.249958682	-0.099012764	-0.67135809
0.108835219	0.456094228	0.611339911	-1.818329276	-0.242828173	-0.67135809
1	0.865868634	-1.226540092	0.373196163	0.410732963	-0.67135809
-1	-1.17952336	0.717991517	0.575725977	0.27330935	1.293327047
-1	-0.466985698	0.596458292	0.295814489	0.17903036	-0.67135809
-1	0.741457296	0.675000172	0.399945384	0.287690891	-0.67135809
-1	0.838898344	0.763463132	0.587189928	0.35160885	-0.67135809
-1	0.764077539	0.795706641	0.609162503	0.4315063	-0.67135809
-1	0.920679223	-1.226540092	0.530825499	0.139081635	1.257516713
-1	-1.17952336	0.608859641	0.236584071	0.044802645	-0.67135809
-1	0.706656922	0.8155488	0.656928968	0.760683791	1.349484162
-0.27309104	1.371344069	1.329791428	1.262607751	1.433420315	1.942796285
-1	1.875079486	1.591046526	1.560670496	2.013475797	2.200793463
-1	0.539615126	0.697322601	0.524138194	0.247742166	-0.67135809
-0.675668601	0.845858419	0.760982862	0.572859989	0.501816055	1.410524504
-1	-1.17952336	0.765943402	0.286261196	-0.031898906	1.241239289
-0.228886868	1.433114734	1.358727911	1.26451841	1.481358784	1.862223034
-1	0.513514845	0.884996358	0.706606092	0.567331964	1.322626411
-0.253709663	1.074670879	0.993301477	0.790675072	0.69996173	1.619689409
-1	-1.17952336	0.583230186	0.338804308	0.166246768	-0.67135809
-1	0.977229831	0.713030978	0.603430527	0.370784238	-0.67135809
-1	0.684036679	0.742794216	0.471595082	0.263721656	-0.67135809
-1	0.950259541	0.777517995	0.650241663	0.568929913	1.333206737
-1	0.843248391	0.738660433	0.582413282	0.319649871	-0.67135809
-1	-1.17952336	-1.226540092	0.244226706	-0.234838428	-0.67135809
-1	0.717097034	-1.226540092	0.458220472	0.107122655	-0.67135809
-1	0.689256735	-1.226540092	0.342625625	0.242948319	-0.67135809
-1	0.864128615	0.785785562	0.597698551	0.452279637	-0.67135809
-1	1.023340327	-1.226540092	0.275752573	0.006451869	1.64085006
-1	-1.17952336	0.702283141	0.516495559	0.334031412	-0.67135809

Table 17 cont'd: SVM Vectors

coefs	FGF14	DMRT1	CNRIP1_2
	FGF14 '577	DMRT1 '934	CNRIP1 '232
1	0.955300844	1.131612847	0.296394201
1	-0.93950462	-0.701307401	0.101966839
0.160992568	0.846924723	-0.701307401	-2.336804129
1	-0.93950462	-0.701307401	0.230086604
1	1.296419127	1.516680127	0.443619546
1	0.974844079	-0.701307401	0.407656103
0.539110203	-0.93950462	1.17300758	0.203114022
1	-0.93950462	-0.701307401	0.261554616
0.228841426	-0.93950462	1.199962289	0.268297762
1	-0.93950462	-0.701307401	0.368321087
1	0.734106957	-0.701307401	0.228962746
0.086502196	0.761645152	-0.701307401	0.236829749
1	-0.93950462	-0.701307401	0.536899725
0.022981588	-0.93950462	-0.701307401	-2.336804129
0.199753156	0.771416769	-0.701307401	0.216600313
0.640672506	0.814944884	-0.701307401	0.384055093
0.488436741	-0.93950462	-0.701307401	0.073870399
0.290813367	-0.93950462	-0.701307401	0.268297762
0.93438076	-0.93950462	-0.701307401	0.027792238
1	0.960630817	-0.701307401	0.3997891
0.177133374	-0.93950462	-0.701307401	0.092975978
0.363673626	0.769640112	-0.701307401	0.317747496
0.372869951	-0.93950462	-0.701307401	0.225591173
0.180416799	-0.93950462	1.108508811	-2.336804129
1	1.728146955	1.533045486	1.467453809
0.236974727	-0.93950462	-0.701307401	-2.336804129
0.433173404	-0.93950462	-0.701307401	0.284031768
0.037394649	-0.93950462	-0.701307401	-2.336804129
1	0.834488119	-0.701307401	0.25031604
0.263636311	-0.93950462	-0.701307401	-2.336804129
1	0.798066635	-0.701307401	0.540271298
1	-0.93950462	-0.701307401	-0.045258506
1	0.853143025	1.17300758	-2.336804129
0.26169536	-0.93950462	-0.701307401	0.450362691

Table 17 cont'd: SVM Vectors

coefs	FGF14	DMRT1	CNRIP1_2
	FGF14 '577	DMRT1 '934	CNRIP1 '232
0.1799778	-0.93950462	1.108508811	0.412151533
0.898868481	-0.93950462	-0.701307401	0.412151533
0.042908076	-0.93950462	-0.701307401	0.205361737
0.140898434	-0.93950462	-0.701307401	-2.336804129
0.140415448	0.748320219	-0.701307401	-2.336804129
1	0.726111997	-0.701307401	0.384055093
0.108835219	-0.93950462	-0.701307401	-2.336804129
1	-0.93950462	1.12487417	0.598711893
-1	0.830046474	-0.701307401	0.314375923
-1	-0.93950462	-0.701307401	0.078365829
-1	-0.93950462	-0.701307401	0.552633732
-1	0.867356287	1.072890087	0.532404295
-1	1.115200039	-0.701307401	0.452610406
-1	0.632837467	1.202850294	0.307632777
-1	0.974844079	-0.701307401	0.05476482
-1	0.946417555	-0.701307401	0.585225602
-0.27309104	1.777893371	2.092355709	1.213461994
-1	1.891599465	2.29451603	1.66750046
-1	0.822939843	-0.701307401	-2.336804129
-0.675668601	0.790071676	-0.701307401	0.593092605
-1	-0.93950462	-0.701307401	0.252563755
-0.228886868	1.602004256	1.983574202	1.266283301
-1	1.124971657	1.209588971	0.542519013
-0.253709663	1.057458663	-0.701307401	0.829102699
-1	1.010377234	-0.701307401	0.017677519
-1	0.846924723	1.254834377	0.354834796
-1	1.049463703	1.121023497	0.557129162
-1	-0.93950462	-0.701307401	0.488573849
-1	0.948194213	0.985287281	0.479582988
-1	-0.93950462	-0.701307401	-2.336804129
-1	-0.93950462	-0.701307401	0.325614499
-1	0.225094519	-0.701307401	0.389674381
-1	0.870909602	1.213439644	0.285155626
-1	-0.93950462	1.568664209	0.829102699
-1	-0.93950462	-0.701307401	0.224467316

Example 5. Various Individual Methylation Biomarkers are Each Highly Informative

Evaluation of the performance of individual colorectal cancer DMRs from among the 28 colorectal cancer DMR panel reveal that various individual colorectal cancer DMRs are sufficient for screening of colorectal cancer (See Figures 12-19). For selected colorectal cancer DMRs, Figures 12-19 show methylation status of the indicated DMR in colorectal cancer samples and control samples. Results are displayed as the MSRE-qPCR Ct value subtracted from 45(i.e., $45 - \text{Ct value}$) for display purposes. Data provided in this Example, as well as data provided by the present Examples cumulatively (including, e.g., Figures 5-9), demonstrate that for each individual colorectal cancer DMR the methylation status signal is sufficiently stable across subject groups to permit clinical screening. Results presented in Figures 12-19 therefore confirm that methylation markers of colorectal cancer provided herein can provide a robust signal for screening of colorectal cancer. Moreover, those of skill in the art will appreciate that the present disclosure provides methylation biomarkers that are individually independently useful in screening for colorectal cancer, and specifically that methylation biomarkers provided herein are useful both individually or in combination.

SEQUENCES

- SEQ ID NO: 1 (ZNF132) (see Table 1)
ZNF132 chr19, bp 58439728 to 58440994 of hg38
5
- SEQ ID NO: 2 (DMRT1) (see Table 1)
DMRT1 chr9, bp 841340 to 968090 of hg38
- SEQ ID NO: 3 (ALK) (see Table 1)
10 ALK chr2, bp 29193215 to 29922286 of hg38
- SEQ ID NO: 4 (JAM2) (see Table 1)
JAM2 chr21, bp 25637848 to 25714704 of hg38
- 15 SEQ ID NO: 5 (FGF14) (see Table 1)
FGF14 chr13, bp 101919879 to 102403137 of hg38
- SEQ ID NO: 6 (MCIDAS) (see Table 1)
MCIDAS chr5, bp 55220951 to 55221051 of hg38
20
- SEQ ID NO: 7 (ST6GALNAC5) (see Table 1)
ST6GALNAC5 chr1, bp 76866255 to 77063388 of hg38
- SEQ ID NO: 8 (LONRF2) (see Table 1)
25 LONRF2 chr2, bp 100285667 to 100323015 of hg38
- SEQ ID NO: 9 (PDGFD) (see Table 1)
PDGFD chr11, bp 104163499 to 104164026 of hg38
- 30 SEQ ID NO: 10 (GSG1L) (see Table 1)
GSG1L chr16, bp 27920615 to 28064275 of hg38
- SEQ ID NO: 11 (ZNF492) (see Table 1)
ZNF492 chr19, bp 22633051 to 22666433 of hg38
35

SEQ ID NO: 12 (ZNF568) (see Table 1)
ZNF568 chr19, bp 36916312 to 36943940 of hg38

SEQ ID NO: 13 (ADAMTS2) (see Table 1)
5 ADAMTS2 chr5, bp 179118114 to 179344392 of hg38

SEQ ID NO: 14 (ZNF542) (see Table 1)
ZNF542 chr19, bp 56367838 to 56370986 of hg38

10 SEQ ID NO: 15 (ZNF471) (see Table 1)
ZNF471 chr19, bp 56507245 to 56508589 of hg38

SEQ ID NO: 16 (CNRIP1) (see Table 1)
CNRIP1 chr2, bp 68293114 to 68320928 of hg38

15
SEQ ID NO: 17 (see Table 7)
CCTCCTCACCCATCATCAGCGCCCGCGGCTTTGGGTGGCCGACCAGAGGGCGGCCGG
AAAGCACCTCGGTGCCCCGCGACCCTCCGAACAGAGGCGGCGGGAGGTACC

20 SEQ ID NO: 18 (see Table 7)
GCGTGCTGGGTTTAATCTTCACCTCAACCTTGTAAGGAGGAGCCGGTGAGCAGCTTGAT
GGTGCGGTTCTGGCCGAAGCGCTGCCCGTCCACCTTGTAAGACCGGGCCGT

SEQ ID NO: 19 (see Table 7)
25 AGGAAGCAAAGTGACCCCTAAGCCTAGACAAAGCTCTCGAAAGCCCAAAGCCTCGGGC
CCACCGGCCAGCTCCCCACCCCGCTGCTGGGCGGACAGGTGTAGGGGAGGCGGAC
C

SEQ ID NO: 20 (see Table 7)
CTCTCAGTCCCGCCGGCTTAGGTAACCCAGGTCGCTGCGGTAACGCAGTGACCGCGCT
30 CCAGGTCCGCGTCTCTTGC

SEQ ID NO: 21 (see Table 7)
CCACTGCGAAGGGAAGGGGCATTCCGCCAGGCGACCCCAGAAGCCAGCCTGCACCTC
CCCGGCTTTCTGCAACCGGGAAGGGGCGTTAACAGGG

35

SEQ ID NO: 22 (see Table 7)

GCGACCCCAGAAGCCAGCCTGCACCTCCCCGGCTTTCCTGCAACCGGGAAGGGGCGT
TAACAGGGCCACCACTCCGGGGCTCCGCCACTCCCCAGCCGTT

5 SEQ ID NO: 23 (see Table 7)

CAACGGAAACTTCCCGCGCTACGGCGGCTCCAACGGGCGCTTCCGCCGCATTGCGTA
GCGAAGCCCCCGGCGAG

SEQ ID NO: 24 (see Table 7)

10 CAAAGCGTCTGGGGCGCTAGTGGGGGCGGCCAGCGGCTCGAGCGCCGGGGGCGAGCA
GCAGAGGAGGCGGCTCCGGCTCCGGGGCGTCGGACCTGGGTGCCGGGAGCAAGAAG
T

SEQ ID NO: 25 (see Table 7)

15 CGCTCAGCCGCTCTCCTCTTCTCTCTCCCGCCCGCCCGCAGCGCCATGGTCTGGCAGT
GTGTTTAGCGCT

SEQ ID NO: 26 (see Table 7)

GGGTTCGGAGCGTGCAAAGGTGACCTAGGCGCGCTACGCACCACGCACTCAGCGGT
20 ACTCTCCTCTCCCGGGCCCCCACGGGTCCCGATGCTGGGCGGGGATGCACTGAACTG
TTC

SEQ ID NO: 27 (see Table 7)

GCGCCCCACTTACATCCAGCACCGAGGCCAGGTGCCGGGTTGGGCTGGCGAGTTCCTT
25 CAGCTGCACGTTCCGCTCCTTGAGCGAGGCGATCTCCTCCTGTTTCTGGGTCAATGTCA
CGT

SEQ ID NO: 28 (see Table 7)

AACGTCTATCACCCAGGGAAAGCTACTCTTGACTCCTTCCACCTATCAAAATTGCCTAAG
30 AAAGGTTGAGTCTGACCAAGGGGCGGCGCAGCTTGCAACTTTCGCCAACTCCGGGA

SEQ ID NO: 29 (see Table 7)

GGTGCATTTGGGATCAGCGACTAGAGACAGCGTCGCTCCAAGAAAAAGCCGGGTTCTG
CTCCCGGGACCGACGCCGCGCCGCCCTGCGCTCTCGCCGCCTGCGCTCGCCCTGCGC
35 TGGCCCGGGTCGCTGTGCTAATC

SEQ ID NO: 30 (see Table 7)

CCGAAAGAAATCCGAGCCAGGGTGAGGGTCTGAGACGCAAGGAGAATCCCAGGCAAG
GCGCTCCTGAGAAAAGATCCCCACGGCGGACGTGGGGCAACAAAACC

5 SEQ ID NO: 31 (see Table 7)

CGAGAGAGGGGAAGGGGCTGGTTGGAACCGGTGGCAAGAGGCTGTGGCGGGACTCA
GGCCTCCCCGAGTCGGCTCCACAATCTGCGCCCCAAGTTCG

SEQ ID NO: 32 (see Table 7)

10 GCCCAAGCCTCACCTCACACAGGAAAGCAGATGTGTTCTGGCCGGAAGTTGAGTGGG
GCCGCGGGGCTGCTGGGAGGTGTTGTCTCGGAAACGTCGCTGGCGCGGAGGGATG
GTTCG

SEQ ID NO: 33 (see Table 7)

15 GGTGCGCTTCACCCAGCATCTCAGAACTGCGCGCGGGATGAACATTGGGTGTTTCC
GGCAGGTGACGCTG

SEQ ID NO: 34 (see Table 7)

CCAGAGGCCCAGGGATCCGTTTCAGGTCAGCGCTGGCGTCCGGGCCTGAGTTTGGAGG
20 TGGCGGGTGCCTTACAAGAATGCTCGCGT

SEQ ID NO: 35 (see Table 7)

GGGAGGAGTGGGCGGCTGAATGGCCAGAGGCCAGGGATCCGTTTCAGGTCAGCGCTG
GCGTCCGGGCCTGAGTTTGGAGGTGGCGGGTGC
25

SEQ ID NO: 36 (see Table 7)

CCCCACGCGTACTCACACCGAAGGCTCAGCCGTCGCGCGTTTCCCTCCCAGGCCCCAG
GAACTAGTAACTAGGGACGCTTCTGGTCTCTAGGCGAGGAGAGGGGGAGAGCGCAATC
TTTGCGCCTGCGCACACTCCTGCTCTTACCCGC
30

SEQ ID NO: 37 (see Table 7)

GTGCGCGTTTTCCCTCCCAGGCCCCAGGAAGTAGTAACTAGGGACGCTTCTGGTCTCT
AGGCGAGGAGAGGGGGAGAGCGCAATCTTTGCGCCTGCGCACACTCCTGCTCTTACCC
GC
35

SEQ ID NO: 38 (see Table 7)

CTGCTCTTACCCGCCGGAACCCTGGGCCACGCCCGGCTCGCGTAATCACGCACTGCG
CAGGCACCGCCCGCTCTGCTCTAAGGTCCCTC

5 SEQ ID NO: 39 (see Table 7)

CTACTGCTAGGTGCTTGCCAAGGTGATTGAGGAATGGCGTTTATTGCGTCGCTGCTCAG
GCAACGCAAACCTACATTATCCAGAAGGACCCTCGCGGTGCCTCAGGGCTGGCCATTGG
CAGCCGAGGAGACAGGCACTTCCGGGCGGAGTGTAAGACGCTGGCCAATCA

10 SEQ ID NO: 40 (see Table 7)

GTGTAAGACGCTGGCCAATCACAGCCTGGCAGCGGGACTTCCGTCGTCGTCCTCGGAC
CATCACTTTGGCATTCTCGATTTTGTCTGCTTCTGAAGGGACCGCGTTGT

SEQ ID NO: 41 (see Table 7)

15 CCGCGTGGTCTGGGCTCTGTAGCGTCCCAGCTGAGCCGGCGATATGCAGCGCACTTGT
GGGGCGGAGGTGGAGGGAATTC

SEQ ID NO: 42 (see Table 7)

CAACGTTAAAGGCAAACACCTTCTGCGGTGTGCTTGGCTCAGCTCAGGCAGGAAGCCC
20 TGCCTGAAAAGGCTGCACCTTCGGCTGTCACTCTGTCCTCATTGCGCC

SEQ ID NO: 43 (see Table 7)

GCCGGTGAGCAGCTTGATGGTGCGGTTCTGGCCGAAGCGCTGCCCGTCCACCTTGTA
AAGACCGGGCCGT
25

SEQ ID NO: 44 (see Table 7)

CGGGAAGGGGCGTTAACAGGGCCACCACTCCGGGGCTCCGCCACTCCCCAGCCGTT
CCTCCTCCGGAGACCTTGCCTGCCAAGA

30 SEQ ID NO: 45 (see Table 13)

CCTCCTACCCATCATCAGCGCCC

SEQ ID NO: 46 (see Table 13)

GGTACCTCCCGCCGCCTCTGTTC

35

SEQ ID NO: 47 (see Table 13)
GCGTGCTGGGTTTAATCTTCACCTCAA

SEQ ID NO: 48 (see Table 13)
5 ACGGCCCGGTCTTTTACAAGGTGG

SEQ ID NO: 49 (see Table 13)
AGGAAGCAAAGTGACCCCTAAGCCT

10 SEQ ID NO: 50 (see Table 13)
GGTCCGCCTCCCCTACACCT

SEQ ID NO: 51 (see Table 13)
CTCTCAGTCCCGCCGGCTTAGGTA
15

SEQ ID NO: 52 (see Table 13)
GCAAGAGACGCGGACCTGGAGC

SEQ ID NO: 53 (see Table 13)
20 CCACTGCGAAGGGAAGGGGCA

SEQ ID NO: 54 (see Table 13)
CCCTGTTAACGCCCTTCCCGGTT

25 SEQ ID NO: 55 (see Table 13)
GCGACCCAGAAAGCCAGCCT

SEQ ID NO: 56 (see Table 13)
AACGGCTGGGGAGTGGCGGA
30

SEQ ID NO: 57 (see Table 13)
CAACGGAACTTCCCGCGCTAC

SEQ ID NO: 58 (see Table 13)
35 CTCGCCGGGGGCTTCGCTAC

- SEQ ID NO: 59 (see Table 13)
CAAAGCGTCTGGGGCGCTAGT
- 5 SEQ ID NO: 60 (see Table 13)
ACTTCTTGCTCCCGGCACCCAGGTC
- SEQ ID NO: 61 (see Table 13)
CGCTCAGCCGCTCTCCTCTTCTCT
- 10 SEQ ID NO: 62 (see Table 13)
AGCGCTAAACACACTGCCAGACCA
- SEQ ID NO: 63 (see Table 13)
- 15 GGGTTCGGAGCGTGCAAAAGGTGA
- SEQ ID NO: 64 (see Table 13)
GAACAGTTCAGTGCATCCCCGCCC
- 20 SEQ ID NO: 65 (see Table 13)
GCGCCCCACTTACATCCAGCACC
- SEQ ID NO: 66 (see Table 13)
ACGTGACATTGACCCAGAAACAGGAGGA
- 25 SEQ ID NO: 67 (see Table 13)
AACGTCTATCACCCAGGGAAAGCT
- SEQ ID NO: 68 (see Table 13)
- 30 TCCCGGAGTTGGCGAAAGTTGCAA
- SEQ ID NO: 69 (see Table 13)
GGTGCATTTGGGATCAGCGACTAGAGAC
- 35 SEQ ID NO: 70 (see Table 13)

GATTAGCACAGCGACCCGGGCCAG

SEQ ID NO: 71 (see Table 13)

CCGAAAGAAATCCGAGCCAGGGTGA

5

SEQ ID NO: 72 (see Table 13)

GGTTTTGTTGCCCCACGTCC

SEQ ID NO: 73 (see Table 13)

10 CGAGAGAGGGGAAGGGGCTGGTTG

SEQ ID NO: 74 (see Table 13)

CGAACTTGGGGCGCAGATTGTGG

15 SEQ ID NO: 75 (see Table 13)

GCCCAAGCCTCACCTCACACAG

SEQ ID NO: 76 (see Table 13)

CGAACCATCCCTCCGCGCCA

20

SEQ ID NO: 77 (see Table 13)

GGTCGCCTTCACCCAGCATCTCAG

SEQ ID NO: 78 (see Table 13)

25 CAGCGTCACCTGCCGAAACACC

SEQ ID NO: 79 (see Table 13)

CCAGAGGCCCAAGGATCCGTTTCAG

30 SEQ ID NO: 80 (see Table 13)

ACGCGAGCATTCTTGTAAAGGCACCC

SEQ ID NO: 81 (see Table 13)

GGGAGGAGTGGGCGGCTGAATGG

35

SEQ ID NO: 82 (see Table 13)
GCACCCGCCACCTCCAAACTCAG

SEQ ID NO: 83 (see Table 13)
5 CCCCACGCGTACTCACACCGAAG

SEQ ID NO: 84 (see Table 13)
GCGGGTAAGAGCAGGAGTGTG

10 SEQ ID NO: 85 (see Table 13)
GTCGCGCGTTTCCCTCCCAG

SEQ ID NO: 86 (see Table 13)
GCGGGTAAGAGCAGGAGTGTG

15
SEQ ID NO: 87 (see Table 13)
CTGCTCTTACCCGCCGGAACCCTG

SEQ ID NO: 88 (see Table 13)
20 GAGGGACCTTAGAGCAGAGCGGGC

SEQ ID NO: 89 (see Table 13)
CTACTGCTAGGTCGTTGCCAAGG

25 SEQ ID NO: 90 (see Table 13)
TGATTGGCCAGCGTCTTACTCCG

SEQ ID NO: 91 (see Table 13)
GTGTAAGACGCTGGCCAATCACA

30
SEQ ID NO: 92 (see Table 13)
ACAACGCGGTCCCTTCAGAAGCAG

SEQ ID NO: 93 (see Table 13)
35 CCGCGTGGTCTGGGCTCTGTAG

SEQ ID NO: 94 (see Table 13)

GAATTCCTCCACCTCCGCCCCAC

5 SEQ ID NO: 95 (see Table 13)

CAACGTTAAAGGCAAACACCTTCTGC

SEQ ID NO: 96 (see Table 13)

GGCCGAATGAGGACAGAGTGACAG

10

SEQ ID NO: 97 (see Table 13)

GCCGGTGAGCAGCTTGATGGT

SEQ ID NO: 98 (see Table 13)

15 ACGGCCCGGTCTTTTACAAGG

SEQ ID NO: 99 (see Table 13)

CGGGAAGGGGCGTTAACAGGGC

20 SEQ ID NO: 100 (see Table 13)

TCTTGGCAGGCAAGGTCTCCGGAG

CLAIMS

1. A in vitro method of screening for colorectal cancer in a subject, the method comprising:
 - a) determining the methylation status of each of the DMR loci in a DMR set as defined in tables 7, 8, 9, 10, 11, 12 or 14, in a subject's DNA sample,
 - 5 b) comparing the data obtained with reference values obtained from healthy individuals,
 - c) diagnosing colorectal cancer in the subject if a hyper-methylation in one or more of the loci is detected, as compared to the reference sample.
- 10 2. The method of claim 1 wherein the methylation status is determined by methylation specific restriction enzyme quantitative polymerase chain reaction (MSRE-qPCR).
3. The method of any one of claim 2 wherein the DMR loci are amplified by the oligonucleotide primer pairs provided in Table 13.
- 15 4. The method of any one of claims 1 to 3, wherein the DNA is isolated from blood, plasma or stool of the human subject.
5. The method of any one of claims 1 to 4, wherein the DNA is cell-free DNA of the human
- 20 subject.
6. The method of any one of claims 1 to 5, wherein the subject was asymptomatic for colorectal cancer at the time of screening.
- 25 7. The method of any one of claims 1 to 18, wherein the subject had been previously screened for colorectal cancer.
8. The method of claim 7, wherein the subject had been screened for colorectal cancer within the last 10 years, within the last 5 years, within the last 4 years, within the last 3 years, within
- 30 the last 2 years, or within the last year.
9. The method of claim 7 or 8, wherein a previous screen for colorectal cancer in the subject had diagnosed the subject as not having colorectal cancer.
- 35 10. The method of claim 9, wherein the previous screen for colorectal cancer that had

diagnosed the subject as not having colorectal cancer was within one year.

11. The method of claim 9 or 10, wherein the previous screen for colorectal cancer that had diagnosed the subject as not having colorectal cancer was a colonoscopy.

5

12. The method of any one of claims 1 to 11, wherein the method includes diagnosis of early stage colorectal cancer, optionally wherein the colorectal cancer is a stage 0, stage I, stage IIA, stage IIB, or stage IIC colorectal cancer.

10 13. The method of any one of claims 1 to 12, wherein the method includes diagnosis of early stage colorectal cancer, wherein the cancer has not metastasized.

14. The method of any one of claims 1 to 13, wherein the method provides a sensitivity for colorectal cancer of at least 0.6.

15

15. The method of claim 14, wherein the method provides a sensitivity for colorectal cancer of at least 0.7.

16. The method of any one of claims 1 to 15, wherein the method provides a specificity of at least 0.7.

20

17. The method of claim 16, wherein the method provides a specificity of at least 0.8.

18. The method of claim 1, wherein methylation status is determined using one or more methods selected from the group consisting of methylation sensitive restriction enzyme quantitative polymerase chain reaction (MSRE-qPCR), Methylation-Specific PCR, Methylation Specific Nuclease-assisted Minor-allele Enrichment PCR, and next-generation sequencing.

25

30 19. A kit comprising:

(a) primer pairs for the detection of the methylation status of each of the DMR loci in one DMR set as defined in tables 7, 8, 9, 10, 11, 12 or 14 wherein the primers pairs are those of table 13,

35

the kit optionally further comprising:

(b) at least one methylation specific restriction enzyme and/or a bisulfite reagent

5 (c) a polymerase

20. The method of any one of claims 1 to 18, the diagnosis further comprising determining the presence or absence of an oncogenic mutation.

10 21. The method of claim 20, wherein the oncogenic mutation is in a gene selected from the group of genes consisting of Kras, NRAS, PIK3CA, PTEN, TP53, BRAF, and APC gene.

22. An in vitro method of screening for colorectal cancer, the method comprising determining the methylation status of one or more colorectal cancer methylation loci selected from:

15 (a) one or more genes according to Table 1; and/or

(b) one or more differentially methylated regions (DMRs) according to Table 7,

in DNA of a human subject, and

diagnosing colorectal cancer in the subject.

20 23. An in vitro method of methylation specific restriction enzyme quantitative polymerase chain reaction (MSRE-qPCR) for colorectal cancer screening, the method comprising:

(a) contacting a sample of DNA of a human subject with one or more methylation specific restriction enzymes; and

25 (b) performing qPCR of enzyme-digested DNA, or amplicons thereof, to determine the methylation status of one or more colorectal cancer methylation loci selected from:

(i) one or more genes according to Table 1; and/or

(ii) one or more differentially methylated regions (DMRs) according to Table 7.

30

24. The method of any one of claims 22 to 23, the method comprising determining whether the one or more methylation loci are hypermethylated as compared to a reference, wherein hypermethylation is indicative of colorectal cancer.

25. The method any one of claims 22 to 24, wherein the method comprises determining the methylation status of at least one DMR of each of ZNF471 and FGF14.
26. The method any one of claims 22 to 24, wherein the method comprises determining the methylation status of at least one DMR of each of ZNF471, FGF14, and PDGFD.
27. The method any one of claims 22 to 24, wherein the method comprises determining the methylation status of at least one DMR of each of ZNF471, FGF14, PDGFD, and ADAMTS2.
28. The method any one of claims 22 to 24, wherein the method comprises determining the methylation status of at least one DMR of each of ZNF471, FGF14, PDGFD, ADAMTS2, ZNF492, ST6GALNAC5.
29. The method any one of claims 22 to 24, wherein the method comprises determining the methylation status of at least one DMR of each of ZNF471, FGF14, PDGFD, ADAMTS2, ZNF492, ST6GALNAC5, ZNF542, LONRF2, ZNF132, CNRIP1, and ALK.
30. The method of any one of claims 22 to 24, wherein the method comprises determining the methylation status of DMRs ZNF471 '558 and FGF14 '577.
31. The method of any one of claims 22 to 24, wherein the method comprises determining the methylation status of DMRs ZNF471 '558, FGF14 '577, and PDGFD '388.
32. The method of any one of claims 22 to 24, wherein the method comprises determining the methylation status of DMRs ZNF471 '558, FGF14 '577, PDGFD '388, ZNF471 '527, and ADAMTS2 '284.
33. The method of any one of claims 22 to 24, wherein the method comprises determining the methylation status of DMRs ZNF471 '558, FGF14 '577, PDGFD '388, ZNF471 '527, ADAMTS2 '284, ADAMTS2 '254, ZNF492 '069, and ST6GALNAC5 '456.
34. The method of any one of claims 22 to 24, wherein the method comprises determining the methylation status of DMRs ZNF471 '558, FGF14 '577, PDGFD '388,

ZNF471 '527, ADAMTS2 '284, ADAMTS2 '254, ZNF492 '069, ST6GALNAC5 '456, ZNF542 '502, LONRF2 '281, ZNF132 '415, PDGFD '921, ZNF132 '268, CNRIP1 '272, and ALK '434.

35. The method of any one of claims 22 to 34, wherein one or more of the DMRs are
5 amplified by oligonucleotide primer pairs provided in Table 13.

36. The method of any one of claims 1 to 15, wherein the DNA is isolated from blood, plasma, urine, saliva, or stool of the human subject.

10 37. The method of any one of claims 22 to 36, wherein the DNA is cell-free DNA of the human subject.

38. The method of any one of claims 22 to 37, wherein the subject was asymptomatic for colorectal cancer at the time of screening.

15

39. The method of any one of claims 22 to 38, wherein the subject had been previously screened for colorectal cancer.

40. The method of claim 39, wherein the subject had been screened for colorectal cancer
20 with in the last 10 years, within the last 5 years, within the last 4 years, within the last 3 years, within the last 2 years, or within the last year.

41. The method of claim 39 or 40, wherein a previous screen for colorectal cancer in the subject had diagnosed the subject as not having colorectal cancer.

25

42. The method of claim 41, wherein the previous screen for colorectal cancer that had diagnosed the subject as not having colorectal cancer was within one year.

43. The method of claim 41 or 42, wherein the previous screen for colorectal cancer that
30 had diagnosed the subject as not having colorectal cancer was a colonoscopy.

44. The method of any one of claims 22 to 43, wherein the method includes diagnosis of early stage colorectal cancer, optionally wherein the colorectal cancer is a stage 0, stage I, stage IIA, stage IIB, or stage IIC colorectal cancer.

35

45. The method of any one of claims 22 to 43, wherein the method includes diagnosis of early stage colorectal cancer, wherein the cancer has not metastasized.
46. The method of any one of claims 22 to 45, wherein the method provides a sensitivity
5 for colorectal cancer of at least 0.6.
47. The method of claim 46, wherein the method provides a sensitivity for colorectal cancer of at least 0.7.
- 10 48. The method of any one of claims 22 to 47, wherein the method provides a specificity for colorectal cancer of at least 0.7.
49. The method of claim 48, wherein the method provides a specificity for colorectal cancer of at least 0.8.
- 15 50. The method of claim 22, wherein methylation status is determined using one or more members selected from the group consisting of methylation sensitive restriction enzyme quantitative polymerase chain reaction (MSRE-qPCR), Methylation-Specific PCR, Methylation Specific Nuclease-assisted Minor-allele Enrichment PCR, and next-generation
20 sequencing.
51. A kit for use in colorectal cancer screening, the kit comprising:
(a) at least one oligonucleotide primer pair of Table 13,
the kit optionally further comprising:
25 (b) at least one methylation specific restriction enzyme and/or a bisulfite reagent.
52. An in vitro diagnostic qPCR reaction for screening of colorectal cancer, the diagnostic qPCR reaction including:
30 (a) a human DNA sample;
(b) a polymerase; and
(c) at least one oligonucleotide primer pair of Table 13,
optionally wherein the human DNA is bisulfite-treated human DNA or methylation
specific restriction enzyme-digested human DNA.
- 35

53. The kit or reaction of claim 51 or 52, wherein the oligonucleotide primer pairs include oligonucleotide primer pairs for amplification of DMRs ZNF471 '558 and FGF14 '577.

54. The kit or reaction of claim 51 or 52, wherein the oligonucleotide primer pairs include
5 oligonucleotide primer pairs for amplification of DMRs ZNF471 '558, FGF14 '577, and PDGFD '388.

55. The kit or reaction of claim 51 or 52, wherein the oligonucleotide primer pairs include oligonucleotide primer pairs for amplification of DMRs ZNF471 '558, FGF14 '577, PDGFD
10 '388, ZNF471 '527, and ADAMTS2 '284.

56. The kit or reaction of claim 51 or 52, wherein the oligonucleotide primer pairs include oligonucleotide primer pairs for amplification of DMRs ZNF471 '558, FGF14 '577, PDGFD '388, ZNF471 '527, ADAMTS2 '284, ADAMTS2 '254, ZNF492 '069, and ST6GALNAC5 '456.
15

57. The kit or reaction of claim 51 or 52, wherein the oligonucleotide primer pairs include oligonucleotide primer pairs for amplification of DMRs ZNF471 '558, FGF14 '577, PDGFD '388, ZNF471 '527, ADAMTS2 '284, ADAMTS2 '254, ZNF492 '069, ST6GALNAC5 '456, ZNF542 '502, LONRF2 '281, ZNF132 '415, PDGFD '921, ZNF132 '268, CNRIP1 '272, and
20 ALK '434.

58. The kit or reaction of any one of claims 22 to 50, further comprising using the determined methylation status of the one or more methylation loci to identify at least one of (i) to (iv) as follows: (i) a presence of colorectal cancer in the human subject; (ii) a
25 predisposition for colorectal cancer in the human subject; (iii) an increased risk of colorectal cancer in the human subject, and (iv) a stage of colorectal cancer in the human subject.

59. The method of any one of claims 22 to 50, the diagnosis further comprising determining the presence or absence of an oncogenic mutation.
30

60. The method of claim 59, wherein the oncogenic mutation is in a gene selected from the group of genes consisting of Kras, NRAS, PIK3CA, PTEN, TP53, BRAF, and APC gene.

61. The method of any one of claims 22 to 50, wherein the method is an in vitro method.
35