

CLAIMS

1. Use of a compound of formula (I), or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for the treatment of cancer,


(I),

wherein:

L represents CH_2SCH_2, CH_2CH_2, $CH_2CH_2CH_2$, CH_2, CH_2S, SCH_2, CH_2NHCH_2, CH=CH, or , wherein any hydrogen atom of a CH or CH_2 unit may be replaced by alkyl or alkoxy, any hydrogen of an NH unit may be replaced by alkyl, and any hydrogen atom of a CH_2 unit of CH_2CH_2, $CH_2CH_2CH_2$ or CH_2 may be replaced by hydroxy;

X, independently for each occurrence, represents S, O or CH=CH, wherein any hydrogen atom of a CH unit may be replaced by alkyl;

Y, independently for each occurrence, represents H or $CH_2O(CO)R_7$;

R_7, independently for each occurrence, represents H or substituted or unsubstituted alkyl, alkoxy, aminoalkyl, alkylaminoalkyl, heterocyclylalkyl, or heterocyclylalkoxy;

Z represents H or $R_3(CO)$;

R_1 and R_2 each independently represent H, alkyl, alkoxy or hydroxy;

R_3, independently for each occurrence, represents substituted or unsubstituted alkyl, hydroxyalkyl, aminoalkyl, acylaminoalkyl, alkenyl, alkoxy, alkoxyalkyl, aryl, arylalkyl, aryloxy, aryloxyalkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, heteroaryloxy, heteroaryloxyalkyl or $C(R_8)(R_9)(R_{10})$, $N(R_4)(R_5)$ or OR_6, wherein any free hydroxyl group may be acylated to form $C(O)R_7$;

R_4 and R_5 each independently represent H or substituted or unsubstituted alkyl, hydroxyalkyl, acyl, aminoalkyl, acylaminoalkyl, alkenyl, alkoxyalkyl, aryl, arylalkyl, aryloxy, aryloxyalkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, heteroaryloxy, or heteroaryloxyalkyl, wherein any free hydroxyl group may be acylated to form $C(O)R_7$;


R_6, independently for each occurrence, represents substituted or unsubstituted alkyl, hydroxyalkyl, aminoalkyl, acylaminoalkyl, alkenyl, alkoxyalkyl, aryl, arylalkyl,

aryloxy, aryloxyalkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, heteroaryloxy, or heteroaryloxyalkyl, wherein any free hydroxyl group may be acylated to form $C(O)R_7$;

R_8, R_9 and R_{10} each independently represent H or substituted or unsubstituted alkyl, hydroxy, hydroxyalkyl, amino, acylamino, aminoalkyl, acylaminoalkyl, alkoxycarbonyl, alkoxycarbonylamino, alkenyl, alkoxy, alkoxyalkyl, aryl, arylalkyl, aryloxy, aryloxyalkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, heteroaryloxy, or heteroaryloxyalkyl, or R_8 and R_9 together with the carbon to which they are attached, form a carbocyclic or heterocyclic ring system, wherein any free hydroxyl group may be acylated to form $C(O)R_7$; and

wherein the medicament is to be orally administered with a meal.

2. Use of a compound of formula (II), or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for the treatment of cancer,



(II),

wherein:

L represents CH_2SCH_2, CH_2CH_2, $CH_2CH_2CH_2$, CH_2, CH_2S, SCH_2, CH_2NHCH_2, CH=CH, or , wherein any hydrogen atom of a CH or CH_2 unit may be replaced by alkyl or alkoxy, any hydrogen of an NH unit may be replaced by alkyl, and any hydrogen atom of a CH_2 unit of CH_2CH_2, $CH_2CH_2CH_2$ or CH_2 may be replaced by hydroxy;

X, independently for each occurrence, represents S, O or CH=CH, wherein any hydrogen atom of a CH unit may be replaced by alkyl;

Y, independently for each occurrence, represents H or $CH_2O(CO)R_7$;

R_7, independently for each occurrence, represents H or substituted or unsubstituted alkyl, alkoxy, aminoalkyl, alkylaminoalkyl, heterocyclylalkyl, arylalkyl, or heterocyclylalkoxy;

Z represents H or $R_3(CO)$;

R_1 and R_2 each independently represent H, alkyl, alkoxy or hydroxy;

R_3 represents substituted or unsubstituted alkyl, hydroxyalkyl, aminoalkyl, acylaminoalkyl, alkenyl, alkoxy, alkoxyalkyl, aryl, arylalkyl, aryloxy, aryloxyalkyl, cycloalkyl,

cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, heteroaryloxy, heteroaryloxyalkyl or $C(R_8)(R_9)(R_{10})$, $N(R_4)(R_5)$ or OR_6, wherein any free hydroxyl group may be acylated to form $C(O)R_7$;

R_4 and R_5 each independently for each occurrence represent H or substituted or unsubstituted alkyl, hydroxyalkyl, acyl, aminoalkyl, acylaminoalkyl, alkenyl, alkoxyalkyl, aryl, arylalkyl, aryloxy, aryloxyalkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, heteroaryloxy, or heteroaryloxyalkyl, wherein any free hydroxyl group may be acylated to form $C(O)R_7$;

R_6 represents substituted or unsubstituted alkyl, hydroxyalkyl, aminoalkyl, acylaminoalkyl, alkenyl, alkoxyalkyl, aryl, arylalkyl, aryloxy, aryloxyalkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, heteroaryloxy, or heteroaryloxyalkyl, wherein any free hydroxyl group may be acylated to form $C(O)R_7$;

R_8, R_9 and R_{10} each independently for each occurrence represent H or substituted or unsubstituted alkyl, hydroxy, hydroxyalkyl, amino, acylamino, aminoalkyl, acylaminoalkyl, alkoxycarbonyl, alkoxycarbonylamino, alkenyl, alkoxy, alkoxyalkyl, aryl, arylalkyl, aryloxy, aryloxyalkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, heteroaryloxy, or heteroaryloxyalkyl, or R_8 and R_9 together with the carbon to which they are attached, form a carbocyclic or heterocyclic ring system, wherein any free hydroxyl group may be acylated to form $C(O)R_7$, and wherein at least two of R_8, R_9 and R_{10} are not H;

R_{11} represents aryl, arylalkyl, aryloxy, aryloxyalkyl, heteroaryl, heteroarylalkyl, heteroaryloxy, or heteroaryloxyalkylor R_{11} represents $C(R_{12})(R_{13})(R_{14})$, $N(R_4)(R_{14})$ or OR_{14}, wherein any free hydroxyl group may be acylated to form $C(O)R_7$;

R_{12} and R_{13} each independently respresent H or substituted or unsubstituted alkyl, hydroxy, hydroxyalkyl, amino, acylamino, aminoalkyl, acylaminoalkyl, alkoxycarbonyl, alkoxycarbonylamino, alkenyl, alkoxy, alkoxyalkyl, aryl, arylalkyl, aryloxy, aryloxyalkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, heteroaryloxy, or heteroaryloxyalkyl, wherein any free hydroxyl group may be acylated to form $C(O)R_7$, and wherein both of R_{12} and R_{13} are not H;

R_{14} represents aryl, arylalkyl, aryloxy, aryloxyalkyl, heteroaryl, heteroarylalkyl, heteroaryloxy, or heteroaryloxyalkyl; and


wherein the medicament is to be orally administered with a meal.

3. The use of claim 2, wherein R_{11} represents aryl, arylalkyl, aryloxy, aryloxyalkyl, heteroaryl, heteroarylalkyl, heteroaryloxy, or heteroaryloxyalkyl, and the aryl or heteroaryl ring is substituted with either $–OCHF_2$ or $–OCF_3$ and is optionally further substituted.

4. The use of claim 2 or 3, wherein R_{14} represents aryl, arylalkyl, aryloxy, aryloxyalkyl, heteroaryl, heteroarylalkyl, heteroaryloxy, or heteroaryloxyalkyl, and the aryl or heteroaryl ring is substituted with either $–OCHF_2$ or $–OCF_3$ and is optionally further substituted.

5. The use of claim 3 or 4, wherein R_{11} represents arylalkyl, wherein the aryl ring is substituted with $–OCF_3$.

6. The use of claim 5, wherein R_{11} represents trifluoromethoxybenzyl.

7. The use of claim 6, wherein R_{11} represents 

.

8. The use of any one of claims 2-7, wherein one or more of (a) – (e) applies:

(a) L represents CH_2SCH_2, CH_2CH_2, CH_2S or SCH_2, preferably CH_2CH_2;

(b) Y represents H;

(c) X represents S or CH=CH, wherein any hydrogen atom of a CH unit may be replaced by alkyl;

(d) Z represents $R_3(CO)$; or

(e) R_1 and R_2 each represent H.

9. The use of claim 8, wherein R_3 and R_{11} are not identical.

10. The use of any one of claims 2-9, wherein Z represents $R_3(CO)$ and R_3 represents substituted or unsubstituted arylalkyl, heteroarylalkyl, cycloalkyl or heterocycloalkyl.

11. The use of claim 10, wherein Z represents $R_3(CO)$ and R_3 represents substituted or unsubstituted heteroarylalkyl.

12. The use of claim 11, wherein Z represents $R_3(CO)$ and R_3 represents substituted or unsubstituted pyridylalkyl.

13. The use of any one of claims 2-7, wherein L represents CH_2SCH_2, CH_2CH_2, CH_2S or SCH_2, each Y represents H, X represents S, Z represents $R_3(CO)$, R_1 and R_2 each represent H, and R_3 represents substituted or unsubstituted arylalkyl, heteroarylalkyl, cycloalkyl or heterocycloalkyl.

14. The use of any one of claims 3-7, wherein L represents CH_2CH_2, each Y represents H, X represents S, Z represents $R_3(CO)$, R_1 and R_2 each represent H, and R_3 represents arylalkyl, heteroarylalkyl, cycloalkyl or heterocycloalkyl.

15. The use of claim 14, wherein R_3 represents heteroarylalkyl.

16. The use of any one of claims 1 to 15, wherein the cancer is selected from breast cancer, colorectal cancer, endocrine cancer, lung cancer, melanoma, mesothelioma, renal cancer, and a B cell malignancy.

17. The use of claim 16, wherein the cancer is breast cancer.

18. The use of claim 17, wherein the breast cancer comprises basal-type breast cancer cells, triple-negative breast cancer cells or claudin-low breast cancer cells.

19. The use of claim 16, wherein the cancer is colorectal cancer.

20. The use of claim 16, wherein the cancer is endocrine cancer.

21. The use of claim 20, wherein the endocrine cancer is selected from adrenal cortex adenoma, adrenal cortex carcicnoma, adrenal gland pheochromocytoma, and parathyroid gland adenoma.

22. The use of claim 16, wherein the cancer is melanoma.

23. The use of claim 16, wherein the cancer is renal cancer.

24. The use of claim 16, wherein the cancer is a B cell malignancy.

25. The use of claim 24, wherein the B cell malignancy is selected from multiple myeloma, leukemia, and lymphoma.

26. The use of claim 25, wherein the B cell malignancy is leukemia, such as acute lymphoblastic leukemia, or chronic lymphoblastic leukemia.

27. The use of claim 25, wherein the B cell malignancy is lymphoma, such as Burkitt's lymphoma, Diffuse large B cell lymphoma, follicular lymphoma, or Hodgkin's lymphoma.

28. The use of any one of the preceding claims, wherein the compound is the compound of formula III and 300 mg to 3000 mg of the medicament is to be administered per day.

29. The use of any one of the preceding claims, wherein the medicament is to be administered two times per day.

30. The use of any one of claims 1 to 28, wherein the medicament is to be administered three times per day.

31. The use of any one of the preceding claim, wherein medicament is to be administered to a human.

32. The use of any one of the preceding claims, wherein the medicament is to be administered with one or more additional chemotherapeutic agents.


33. The use of claim 32, wherein the one or more additional chemotherapeutic agents are selected from ABT-263, aminoglutethimide, amsacrine, anastrozole, asparaginase, azacitidine, AZD5363, Bacillus Calmette–Guérin vaccine (bcg), bicalutamide, bleomycin, bortezomib, buserelin, busulfan, campothecin, capecitabine, carboplatin, carfilzomib, carmustine, chlorambucil, chloroquine, cisplatin, cladribine, clodronate, cobimetinib, colchicine, cyclophosphamide, cyproterone, cytarabine, dacarbazine, dactinomycin, daunorubicin, demethoxyviridin, dexamethasone, dichloroacetate, dienestrol, diethylstilbestrol, docetaxel, doxorubicin, epirubicin, eribulin, erlotinib, estradiol,

estramustine, etoposide, everolimus, exemestane, filgrastim, fludarabine, fludrocortisone, fluorouracil (e.g., 5-fluorouracil), fluoxymesterone, flutamide, gemcitabine, genistein, goserelin, hydroxyurea, idarubicin, ifosfamide, imatinib, interferon, irinotecan, ixabepilone, lenalidomaide, letrozole, leucovorin, leuprolide, levamisole, lomustine, lonidamine, mechlorethamine, medroxyprogesterone, megestrol, melphalan, mercaptopurine, mesna, metformin, methotrexate, miltefosine, mitomycin, mitotane, mitoxantrone, mutamycin, MK-2206, nilutamide, nocodazole, octreotide, oxaliplatin, olaparib, paclitaxel, pamidronate, pazopanib, pentostatin, perifosine, PF-04691502, plicamycin, pomalidomide, porfimer, procarbazine, raltitrexed, rituximab, romidepsin, rucaparib, selumetinib, sorafenib, streptozocin, sunitinib, suramin, talazoparib, tamoxifen, temozolomide, temsirolimus, teniposide, testosterone, thalidomide, thioguanine, thiotepa, titanocene dichloride, topotecan, trametinib, trastuzumab, tretinoin, veliparib, vinblastine, vincristine, vindesine, vinorelbine, and vorinostat (SAHA).

34. The use of claim 33, wherein the one or more additional chemotherapeutic agents are selected from azacitidine, bortezomib, capecitabine, carboplatin, carfilzomib, cyclophosphamide, daunorubicin, dexamethasone, docetaxel, doxorubicin, epirubicin, eribulin, erlotinib, everolimus, fluorouracil, gemcitabine, ixabepilone, lenalidomide, methotrexate, mitoxantrone, mutamycin, paclitaxel, pomalidomide, rituximab, thiotepa, vincristine, and vinorelbine.

35. The use of claim 34, wherein the one or more additional chemotherapeutic agents are selected from azacitidine, dexamethasone, docetaxel, erlotinib, everolimus, paclitaxel and pomalidomide.

36. A compound of formula (III), or a pharmaceutically acceptable salt thereof, for use as a medicament,



(III),

wherein the medicament is to be orally administered with a meal.