

Process of Conducting High Throughput Testing High Performance Liquid Chromatography

TECHNICAL FIELD OF INVENTION

[0001] The invention relates to a process of conducting high throughput testing (HTT) high performance liquid chromatography (HPLC) useful for testing large amounts of samples quickly and accurately. In one embodiment, HTT HPLC is useful for developing process analytical techniques (PAT) for continuous manufacturing of pharmaceutical compositions. In another embodiment, the pharmaceutical compositions are for the treatment of CFTR mediated diseases such as cystic fibrosis and comprise one or more active pharmaceutical ingredient (API).

BACKGROUND

[0002] A common challenge for drugs approved by the FDA is the occasional lack of drug availability for patients in need thereof. Accordingly, a significant unmet need exists for the disclosed processes of preparing drugs in a continuous and controlled manner as opposed to the more traditional batch preparations. To achieve continuous manufacturing, PAT must be developed that accurately monitor properties of the pharmaceutical compositions without interrupting the continuity of the processes. PAT, however, are spectroscopic in nature and must be correlated to references to be of any use. This correlation to references requires running many samples in a timely fashion using HTT HPLC techniques disclosed herein. It is also envisioned that HTT HPLC can be used to test the concentration of API in the final composition as either a back-up to PAT or when PAT is not available.

SUMMARY

[0003] In one embodiment, the present invention features a process of conducting high throughput HPLC comprising a) dropping containers, such as vials, of pre-weighed samples into plastic bottles, such as HDPE bottles; b) adding solution to each set of container and bottle via a bottle top dispenser; c) shaking each set of plastic bottle, container, and solution until sample is dissolved; d) centrifuging each set of plastic bottle, container, and solution; e) loading an aliquot of supernatant from the centrifuge step onto an HPLC column; and f) running the


column with a mobile phase.

[0004] In another embodiment, the process is used to supply correlating values to PAT measurements for continuous manufacturing. In another embodiment, the process is used to measure the concentration of API in the final pharmaceutical composition.

[0005] In another embodiment, the pharmaceutical composition is a tablet. In another embodiment, the tablet is for the treatment of a CFTR mediated disease such as cystic fibrosis (CF).


[0006] In another embodiment, the tablet comprises two API. In another embodiment, one API is a CF corrector. In another embodiment, one API is a CF potentiator. In another embodiment, one API is a CF corrector and the other API is a CF potentiator.

[0007] In another embodiment, one API is 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid (Compound 1), which has the structure below:



Compound 1.

[0008] In another embodiment, one API is N-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide (Compound 2), which has the structure below:



Compound 2.

[0009] In another embodiment, one API is Compound 1 and the other API is Compound 2. In another embodiment, Compound 1 is in Form I, and Compound 2 is the form of a solid dispersion of substantially amorphous Compound 2.

BRIEF DESCRIPTION OF DRAWINGS

[0010] **Figure 1** is a flow chart for the continuous manufacture of a tablet of Compound 1 Form I and a solid dispersion of substantially amorphous Compound 2.

[0011] **Figure 2** is a schematic drawing of a process analytical technique (PAT) enabled continuous manufacturing process where in step 1) feeder/blender one, PAT1 NIR measures material attributes during screening of raw materials; step 2) twin screw granulator, PAT2 NIR measures composition and BU; step 3) fluidized bed dryer, PAT 3a NIR measures granule uniformity, LOD, solid state form and physical attributes of granules, PAT 3b laser diffraction measures particle size distribution; step 4) milling, PAT4 NIR measures composition and BU; step 5) feeder/blender two, PAT 5a Raman measures assay and CU, PAT 5b weight, hardness, thickness; step 6) compression, PAT6 Raman measures coat thickness; and step 7) coating.

[0012] **Figure 3** is a schematic drawing showing a PAT inline Sentronics NIR located after blender one, granule mill, and extra granule blender. Each probe has 7 spots that cycle sequentially to maximize sampling and NIR with multiplexer-NIR ensuring robust and exhaustive sampling by controlled powder flow across the probe optics.

[0013] **Figure 4** is a depiction of NIR in flowing powder.

[0014] **Figure 5** is a Kaiser Raman spectrum of Compound 1 Form I and Compound 1 Form II (Compound 1 Form II is a different polymorph disclosed in US 201131588 incorporated herein in its entirety by reference) taken after tablet pressing. The Kaiser Raman spectrometer is mounted on the Kraemer UTS tablet tester.

[0015] **Figure 6** is a graph showing good correlation between predicted and reference off-line NIR samplings of Compound 2 granules.

[0016] **Figure 7** is a series of NIR spectra measuring water content in samples of Compound 1 granules.

[0017] **Figure 8** is a series of NIR spectra measuring a range of compositions comprising different ratios of Compound 1 Form I and a solid dispersions comprising substantially amorphous Compound 2 on the left, and pretreated spectra on the right depicting Range A for identifying Compound 1 Form I and Range B for identifying amorphous Compound 2.

[0018] **Figure 9** depicts a calibration curve for predicted Compound 1 Form I content versus reference (actual) Compound 1 Form I content using partial least squares (PLS) techniques.

[0019] **Figure 10** depicts actual results of unknown samples comprising different contents of Compound 1 Form I (Y Reference) versus predicted content using the calibration curve calculated from Figure 19 (Y Predicted).

[0020] **Figure 11** depicts the transmission percent of a laser diffraction measurement in response to changes in line rate (flow velocity) for a composition comprising Compound 1 Form I and a solid dispersions comprising substantially amorphous Compound 2 showing the expected reduction in transmission percent as line rate increase.

[0021] **Figure 12** depicts laser diffraction measurements of particles comprising Compound 1 Form I and a solid dispersions comprising substantially amorphous Compound 2 at different line rates showing that the average particle size (Dv(50)) is not affected by line rate.

[0022] **Figure 13** depicts laser diffraction measurements of particles comprising Compound 1 Form I and a solid dispersions comprising substantially amorphous Compound 2 under different processing parameters showing that the particle size measurements are sensitive to such changes.

[0023] **Figure 14** depicts the predictive capabilities of process analytical technology models using Raman spectroscopy, both non-continuously and continuously, for monitoring Compound 1 solid form identity in a tablet.

[0024] **Figure 15** depicts the predictive capabilities of process analytical technology models using Raman spectroscopy, both non-continuously and continuously, for monitoring Compound 2 solid form identity in a tablet.

DETAILED DESCRIPTION

DEFINITIONS

[0025] As used herein, “HTT” stands for high throughput testing and “HPLC” stands for high performance liquid chromatography. The two together as in HTT HPLC refers to a high performance liquid chromatography method that can be used to test a high volume amount of samples quickly and accurately.

[0026] As used herein, the term “active pharmaceutical ingredient” or “API” refers to a biologically active compound.

[0027] As used herein, the term “PAT” stands for process analytical technology.

[0028] As used herein, the term “CU” stands for content uniformity.

[0029] As used herein, “CFTR” stands for cystic fibrosis transmembrane conductance regulator.

[0030] As used herein, a “ Δ F508 mutation” or “F508-del mutation” is a specific mutation within the CFTR protein. The mutation is a deletion of the three nucleotides that comprise the codon for amino acid phenylalanine at position 508, resulting in CFTR protein that lacks this phenylalanine residue.

[0031] As used herein, a patient who is “homozygous” for a particular mutation, e.g. Δ F508, has the same mutation on each allele.

[0032] As used herein, a patient who is “heterozygous” for a particular mutation, e.g. Δ F508, has this mutation on one allele, and a different mutation on the other allele.

[0033] As used herein, the term “CFTR corrector” refers to a compound that increases the amount of functional CFTR protein to the cell surface, resulting in enhanced ion transport.

[0034] As used herein, the term “CFTR potentiator” refers to a compound that increases the channel activity of CFTR protein located at the cell surface, resulting in enhanced ion transport.

[0035] The terms “solid form”, “solid forms” and related terms, when used herein refer to Compound 1 or Compound 2, in a particular solid form e.g. crystals, amorphous states, and the like.

[0036] As used herein, the term “substantially amorphous” refers to a solid material having little or no long range order in the position of its molecules. For example, substantially amorphous materials have less than about 15% crystallinity (e.g., less than about 10% crystallinity or less than about 5% crystallinity). It is also noted that the term ‘substantially amorphous’ includes the descriptor, ‘amorphous’, which refers to materials having no (0%) crystallinity.

[0037] As used herein, the term "substantially crystalline" (as in the phrase substantially crystalline Compound 1 Form I refers to a solid material having predominantly long range order in the position of its molecules. For example, substantially crystalline materials have more than about 85% crystallinity (e.g., more than about 90% crystallinity or more than about 95% crystallinity). It is also noted that the term 'substantially crystalline' includes the descriptor, 'crystalline', which refers to materials having 100% crystallinity.

[0038] The term "crystalline" and related terms used herein, when used to describe a substance, component, product, or form, means that the substance, component or product is substantially crystalline as determined by X-ray diffraction. (See, e.g., Remington: The Science and Practice of Pharmacy, 21st Ed., Lippincott Williams & Wilkins, Baltimore, Md. (2003); The United States Pharmacopeia, 23rd ed., 1843-1844 (1995)).

[0039] The term "tablet" as used herein refers to a physically discrete unit of agent appropriate for the patient to be treated. In general, a compacted mixture has a density greater than that of the mixture prior to compaction. A dosage tablet of the invention can have almost any shape including concave and/or convex faces, rounded or angled corners, and a rounded to rectilinear shape. In some embodiments, the compressed tablets of the invention comprise a rounded tablet having flat faces. The tablets of the invention can be prepared by any compaction and compression method known by persons of ordinary skill in the art of forming compressed solid pharmaceutical dosage forms. In particular embodiments, the formulations provided herein may be prepared using conventional methods known to those skilled in the field of pharmaceutical formulation, as described, e.g., in pertinent textbooks. See, e.g., Remington: The Science and Practice of Pharmacy, 21st Ed., Lippincott Williams & Wilkins, Baltimore, Md. (2003); Ansel et al., Pharmaceutical Dosage Forms And Drug Delivery Systems, 7th Edition, Lippincott Williams & Wilkins, (1999); The Handbook of Pharmaceutical Excipients, 4th edition, Rowe et al., Eds., American Pharmaceuticals Association (2003); Gibson, Pharmaceutical Preformulation And Formulation, CRC Press (2001), these references hereby incorporated herein by reference in their entirety.

[0040] As used herein, an "excipient" includes functional and non-functional ingredients in a pharmaceutical composition.

[0041] An “effective amount” or “therapeutically effective amount” of a compound of the invention may vary according to factors such as the disease state, age, and weight of the subject, and the ability of the compound of the invention to elicit a desired response in the subject. Dosage regimens may be adjusted to provide the optimum therapeutic response. An effective amount is also one in which any toxic or detrimental effects (e.g., side effects) of the compound of the invention are outweighed by the therapeutically beneficial effects.

[0042] As used herein, and unless otherwise specified, the terms “therapeutically effective amount” and “effective amount” of a compound mean an amount sufficient to provide a therapeutic benefit in the treatment or management of a disease or disorder, or to delay or minimize one or more symptoms associated with the disease or disorder. A “therapeutically effective amount” and “effective amount” of a compound mean an amount of therapeutic agent, alone or in combination with one or more other agent(s), which provides a therapeutic benefit in the treatment or management of the disease or disorder. The terms “therapeutically effective amount” and “effective amount” can encompass an amount that improves overall therapy, reduces or avoids symptoms or causes of disease or disorder, or enhances the therapeutic efficacy of another therapeutic agent.

[0043] “Substantially pure” as used in the phrase “substantially pure Compound 1 Form I” means greater than about 90% purity. In another embodiment, substantially pure refers to greater than about 95% purity. In another embodiment, substantially pure refers to greater than about 98% purity. In another embodiment, substantially pure refers to greater than about 99% purity.

[0044] With respect to Compound 1 Form I, or a solid dispersion comprising substantially amorphous Compound 2, the terms “about” and “approximately”, when used in connection with doses, amounts, or weight percent of ingredients of a composition or a dosage form, mean a dose, amount, or weight percent that is recognized by one of ordinary skill in the art to provide a pharmacological effect equivalent to that obtained from the specified dose, amount, or weight percent. Specifically the term “about” or “approximately” means an acceptable error for a particular value as determined by one of ordinary skill in the art, which depends in part on how the value is measured or determined. In certain embodiments, the term “about” or “approximately” means within 1, 2, 3, or 4 standard deviations. In certain embodiments, the term

“about” or “approximately” means within 30%, 25%, 20%, 15%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, 0.5%, 0.1%, or 0.05% of a given value or range.

[0045] Compound 1 Form I is prepared by methods disclosed in US Patent No. 8,507,534 incorporated herein by reference in its entirety. A solid dispersion of substantially amorphous Compound 2 is prepared by methods disclosed in International Published Patent Application No. WO2010/019239 incorporated herein by reference in its entirety. A tablet comprising Compound 1 and Compound 2 may be prepared continuously according to the flow chart of Figure 1.

THERAPEUTIC USES OF THE COMPOSITION

[0046] In one aspect, the invention also provides a method of treating, lessening the severity of, or symptomatically treating a disease in a patient, the method comprising administering an effective amount of the pharmaceutical composition or tablet prepared in a continuous manner using PAT to the patient, preferably a mammal, wherein the disease is selected from cystic fibrosis, asthma, smoke induced COPD, chronic bronchitis, rhinosinusitis, constipation, pancreatitis, pancreatic insufficiency, male infertility caused by congenital bilateral absence of the vas deferens (CBAVD), mild pulmonary disease, idiopathic pancreatitis, allergic bronchopulmonary aspergillosis (ABPA), liver disease, hereditary emphysema, hereditary hemochromatosis, coagulation-fibrinolysis deficiencies, such as protein C deficiency, Type 1 hereditary angioedema, lipid processing deficiencies, such as familial hypercholesterolemia, Type 1 chylomicronemia, abetalipoproteinemia, lysosomal storage diseases, such as I-cell disease/pseudo-Hurler, mucopolysaccharidoses, Sandhof/Tay-Sachs, Crigler-Najjar type II, polyendocrinopathy/hyperinsulemia, Diabetes mellitus, Laron dwarfism, myeloperoxidase deficiency, primary hypoparathyroidism, melanoma, glycanosis CDG type 1, congenital hyperthyroidism, osteogenesis imperfecta, hereditary hypofibrinogenemia, ACT deficiency, Diabetes insipidus (DI), neurophyseal DI, neprogenic DI, Charcot-Marie Tooth syndrome, Perlizaeus-Merzbacher disease, neurodegenerative diseases such as Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, progressive supranuclear palsy, Pick's disease, several polyglutamine neurological disorders such as Huntington's, spinocerebellar ataxia type I, spinal and bulbar muscular atrophy, dentatorubal pallidoluyian, and myotonic dystrophy, as well as spongiform encephalopathies, such as hereditary Creutzfeldt-Jakob disease

(due to prion protein processing defect), Fabry disease, Strausler-Scheinker syndrome, COPD, dry-eye disease, or Sjogren's disease, osteoporosis, osteopenia, bone healing and bone growth (including bone repair, bone regeneration, reducing bone resorption and increasing bone deposition), Gorham's Syndrome, chloride channelopathies such as myotonia congenita (Thomson and Becker forms), Bartter's syndrome type III, Dent's disease, hyperekplexia, epilepsy, lysosomal storage disease, Angelman syndrome, and Primary Ciliary Dyskinesia (PCD), a term for inherited disorders of the structure and/or function of cilia, including PCD with situs inversus (also known as Kartagener syndrome), PCD without situs inversus and ciliary aplasia.

[0047] In one aspect, the invention also provides a method of treating, lessening the severity of, or symptomatically treating a disease in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the disease is selected from generalized epilepsy with febrile seizures plus (GEFS+), general epilepsy with febrile and afebrile seizures, myotonia, paramyotonia congenita, potassium-aggravated myotonia, hyperkalemic periodic paralysis, LQTS, LQTS/Brugada syndrome, autosomal-dominant LQTS with deafness, autosomal-recessive LQTS, LQTS with dysmorphic features, congenital and acquired LQTS, Timothy syndrome, persistent hyperinsulinemic hypoglycemia of infancy, dilated cardiomyopathy, autosomal-dominant LQTS, Dent disease, Osteopetrosis, Bartter syndrome type III, central core disease, malignant hyperthermia, and catecholaminergic polymorphic tachycardia.

[0048] In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation N1303K, ΔI507, or R560T.

[0049] In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation G551D. In another embodiment, the patient is homozygous in G551D. In another embodiment, the patient

is heterozygous in G551D wherein the other CFTR genetic mutation is any one of $\Delta F508$, G542X, N1303K, W1282X, R117H, R553X, 1717-1G->A, 621+1G->T, 2789+5G->A, 3849+10kbC->T, R1162X, G85E, 3120+1G->A, $\Delta I507$, 1898+1G->A, 3659delC, R347P, R560T, R334W, A455E, 2184delA, or 711+1G->T.

[0050] In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation $\Delta F508$. In another embodiment, the patient is homozygous in $\Delta F508$. In another embodiment, the patient is heterozygous in $\Delta F508$ wherein the other CFTR genetic mutation is any one of G551D, G542X, N1303K, W1282X, R117H, R553X, 1717-1G->A, 621+1G->T, 2789+5G->A, 3849+10kbC->T, R1162X, G85E, 3120+1G->A, $\Delta I507$, 1898+1G->A, 3659delC, R347P, R560T, R334W, A455E, 2184delA, or 711+1G->T.

[0051] In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from G178R, G551S, G970R, G1244E, S1255P, G1349D, S549N, S549R, S1251N, E193K, F1052V, G1069R, R117C, D110H, R347H, R352Q, E56K, P67L, L206W, A455E, D579G, S1235R, S945L, R1070W, F1074L, D110E, D1270N, D1152H, 1717-1G->A, 621+1G->T, 3120+1G->A, 1898+1G->A, 711+1G->T, 2622+1G->A, 405+1G->A, 406-1G->A, 4005+1G->A, 1812-1G->A, 1525-1G->A, 712-1G->T, 1248+1G->A, 1341+1G->A, 3121-1G->A, 4374+1G->T, 3850-1G->A, 2789+5G->A, 3849+10kbC->T, 3272-26A->G, 711+5G->A, 3120G->A, 1811+1.6kbA->G, 711+3A->G, 1898+3A->G, 1717-8G->A, 1342-2A->C, 405+3A->C, 1716G/A, 1811+1G->C, 1898+5G->T, 3850-3T->G, IVS14b+5G->A, 1898+1G->T, 4005+2T->C and 621+3A->G.

[0052] In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from G178R, G551S, G970R, G1244E, S1255P, G1349D, S549N, S549R, S1251N, E193K, F1052V

and G1069R. In one embodiment of this aspect, the invention provides a method of treating CFTR comprising administering Compound 1 to a patient possessing a human CFTR mutation selected from G178R, G551S, G970R, G1244E, S1255P, G1349D, S549N, S549R and S1251N. In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from E193K, F1052V and G1069R. In some embodiments of this aspect, the method produces a greater than 10-fold increase in chloride transport relative to baseline chloride transport.

[0053] In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from R117C, D110H, R347H, R352Q, E56K, P67L, L206W, A455E, D579G, S1235R, S945L, R1070W, F1074L, D110E, D1270N and D1152H. In one embodiment of this aspect, the method produces an increase in chloride transport which is greater or equal to 10% above the baseline chloride transport.

[0054] In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from 1717-1G->A, 621+1G->T, 3120+1G->A, 1898+1G->A, 711+1G->T, 2622+1G->A, 405+1G->A, 406-1G->A, 4005+1G->A, 1812-1G->A, 1525-1G->A, 712-1G->T, 1248+1G->A, 1341+1G->A, 3121-1G->A, 4374+1G->T, 3850-1G->A, 2789+5G->A, 3849+10kbC->T, 3272-26A->G, 711+5G->A, 3120G->A, 1811+1.6kbA->G, 711+3A->G, 1898+3A->G, 1717-8G->A, 1342-2A->C, 405+3A->C, 1716G/A, 1811+1G->C, 1898+5G->T, 3850-3T->G, IVS14b+5G->A, 1898+1G->T, 4005+2T->C and 621+3A->G. In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from 1717-1G->A, 1811+1.6kbA->G, 2789+5G->A, 3272-

26A->G and 3849+10kbC->T. In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from 2789+5G->A and 3272-26A->G.

[0055] In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from G178R, G551S, G970R, G1244E, S1255P, G1349D, S549N, S549R, S1251N, E193K, F1052V, G1069R, R117C, D110H, R347H, R352Q, E56K, P67L, L206W, A455E, D579G, S1235R, S945L, R1070W, F1074L, D110E, D1270N, D1152H, 1717-1G->A, 621+1G->T, 3120+1G->A, 1898+1G->A, 711+1G->T, 2622+1G->A, 405+1G->A, 406-1G->A, 4005+1G->A, 1812-1G->A, 1525-1G->A, 712-1G->T, 1248+1G->A, 1341+1G->A, 3121-1G->A, 4374+1G->T, 3850-1G->A, 2789+5G->A, 3849+10kbC->T, 3272-26A->G, 711+5G->A, 3120G->A, 1811+1.6kbA->G, 711+3A->G, 1898+3A->G, 1717-8G->A, 1342-2A->C, 405+3A->C, 1716G/A, 1811+1G->C, 1898+5G->T, 3850-3T->G, IVS14b+5G->A, 1898+1G->T, 4005+2T->C and 621+3A->G, and a human CFTR mutation selected from Δ F508, R117H, and G551D.

[0056] In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from G178R, G551S, G970R, G1244E, S1255P, G1349D, S549N, S549R, S1251N, E193K, F1052V and G1069R, and a human CFTR mutation selected from Δ F508, R117H, and G551D. In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from G178R, G551S, G970R, G1244E, S1255P, G1349D, S549N, S549R and S1251N, and a human CFTR mutation selected from Δ F508, R117H, and G551D. In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic

fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from E193K, F1052V and G1069R, and a human CFTR mutation selected from Δ F508, R117H, and G551D. In some embodiments of this aspect, the method produces a greater than 10-fold increase in chloride transport relative to baseline chloride transport.

[0057] In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from R117C, D110H, R347H, R352Q, E56K, P67L, L206W, A455E, D579G, S1235R, S945L, R1070W, F1074L, D110E, D1270N and D1152H, and a human CFTR mutation selected from Δ F508, R117H, and G551D. In one embodiment of this aspect, the method produces an increase in chloride transport which is greater or equal to 10% above the baseline chloride transport.

[0058] In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from 1717-1G->A, 621+1G->T, 3120+1G->A, 1898+1G->A, 711+1G->T, 2622+1G->A, 405+1G->A, 406-1G->A, 4005+1G->A, 1812-1G->A, 1525-1G->A, 712-1G->T, 1248+1G->A, 1341+1G->A, 3121-1G->A, 4374+1G->T, 3850-1G->A, 2789+5G->A, 3849+10kbC->T, 3272-26A->G, 711+5G->A, 3120G->A, 1811+1.6kbA->G, 711+3A->G, 1898+3A->G, 1717-8G->A, 1342-2A->C, 405+3A->C, 1716G/A, 1811+1G->C, 1898+5G->T, 3850-3T->G, IVS14b+5G->A, 1898+1G->T, 4005+2T->C and 621+3A->G, and a human CFTR mutation selected from Δ F508, R117H, and G551D. In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from 1717-1G->A, 1811+1.6kbA->G, 2789+5G->A, 3272-26A->G and 3849+10kbC->T, and a human CFTR mutation selected from Δ F508, R117H, and G551D. In one aspect, the present invention is directed to a method of treating, lessening the severity of, or

symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from 2789+5G->A and 3272-26A->G, and a human CFTR mutation selected from Δ F508, R117H.

[0059] In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from G178R, G551S, G970R, G1244E, S1255P, G1349D, S549N, S549R, S1251N, E193K, F1052V, G1069R, R117C, D110H, R347H, R352Q, E56K, P67L, L206W, A455E, D579G, S1235R, S945L, R1070W, F1074L, D110E, D1270N, D1152H, 1717-1G->A, 621+1G->T, 3120+1G->A, 1898+1G->A, 711+1G->T, 2622+1G->A, 405+1G->A, 406-1G->A, 4005+1G->A, 1812-1G->A, 1525-1G->A, 712-1G->T, 1248+1G->A, 1341+1G->A, 3121-1G->A, 4374+1G->T, 3850-1G->A, 2789+5G->A, 3849+10kbC->T, 3272-26A->G, 711+5G->A, 3120G->A, 1811+1.6kbA->G, 711+3A->G, 1898+3A->G, 1717-8G->A, 1342-2A->C, 405+3A->C, 1716G/A, 1811+1G->C, 1898+5G->T, 3850-3T->G, IVS14b+5G->A, 1898+1G->T, 4005+2T->C and 621+3A->G, and a human CFTR mutation selected from Δ F508, R117H, and G551D.

[0060] In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from G178R, G551S, G970R, G1244E, S1255P, G1349D, S549N, S549R, S1251N, E193K, F1052V and G1069R. In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from G178R, G551S, G970R, G1244E, S1255P, G1349D, S549N, S549R and S1251N. In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from E193K,

F1052V and G1069R. In some embodiments of this aspect, the method produces a greater than 10-fold increase in chloride transport relative to baseline chloride transport.

[0061] In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from R117C, D110H, R347H, R352Q, E56K, P67L, L206W, A455E, D579G, S1235R, S945L, R1070W, F1074L, D110E, D1270N and D1152H. In one embodiment of this aspect, the method produces an increase in chloride transport which is greater or equal to 10% above the baseline chloride transport.

[0062] In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from 1717-1G->A, 621+1G->T, 3120+1G->A, 1898+1G->A, 711+1G->T, 2622+1G->A, 405+1G->A, 406-1G->A, 4005+1G->A, 1812-1G->A, 1525-1G->A, 712-1G->T, 1248+1G->A, 1341+1G->A, 3121-1G->A, 4374+1G->T, 3850-1G->A, 2789+5G->A, 3849+10kbC->T, 3272-26A->G, 711+5G->A, 3120G->A, 1811+1.6kbA->G, 711+3A->G, 1898+3A->G, 1717-8G->A, 1342-2A->C, 405+3A->C, 1716G/A, 1811+1G->C, 1898+5G->T, 3850-3T->G, IVS14b+5G->A, 1898+1G->T, 4005+2T->C and 621+3A->G. In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from 1717-1G->A, 1811+1.6kbA->G, 2789+5G->A, 3272-26A->G and 3849+10kbC->T. In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from 2789+5G->A and 3272-26A->G.

[0063] In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an

effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from G178R, G551S, G970R, G1244E, S1255P, G1349D, S549N, S549R, S1251N, E193K, F1052V, G1069R, R117C, D110H, R347H, R352Q, E56K, P67L, L206W, A455E, D579G, S1235R, S945L, R1070W, F1074L, D110E, D1270N, D1152H, 1717-1G->A, 621+1G->T, 3120+1G->A, 1898+1G->A, 711+1G->T, 2622+1G->A, 405+1G->A, 406-1G->A, 4005+1G->A, 1812-1G->A, 1525-1G->A, 712-1G->T, 1248+1G->A, 1341+1G->A, 3121-1G->A, 4374+1G->T, 3850-1G->A, 2789+5G->A, 3849+10kbC->T, 3272-26A->G, 711+5G->A, 3120G->A, 1811+1.6kbA->G, 711+3A->G, 1898+3A->G, 1717-8G->A, 1342-2A->C, 405+3A->C, 1716G/A, 1811+1G->C, 1898+5G->T, 3850-3T->G, IVS14b+5G->A, 1898+1G->T, 4005+2T->C and 621+3A->G, and a human CFTR mutation selected from Δ F508, R117H, and G551D, and one or more human CFTR mutations selected from Δ F508, R117H, and G551D.

[0064] In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from G178R, G551S, G970R, G1244E, S1255P, G1349D, S549N, S549R, S1251N, E193K, F1052V and G1069R, and one or more human CFTR mutations selected from Δ F508, R117H, and G551D. In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from G178R, G551S, G970R, G1244E, S1255P, G1349D, S549N, S549R and S1251N, and one or more human CFTR mutations selected from Δ F508, R117H, and G551D. In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from E193K, F1052V and G1069R, and one or more human CFTR mutations selected from Δ F508, R117H, and G551D. In some embodiments of this aspect, the method produces a greater than 10-fold increase in chloride transport relative to baseline chloride transport.

[0065] In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from R117C, D110H, R347H, R352Q, E56K, P67L, L206W, A455E, D579G, S1235R, S945L, R1070W, F1074L, D110E, D1270N and D1152H, and one or more human CFTR mutations selected from Δ F508, R117H, and G551D. In one embodiment of this aspect, the method produces an increase in chloride transport which is greater or equal to 10% above the baseline chloride transport.

[0066] In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from 1717-1G->A, 621+1G->T, 3120+1G->A, 1898+1G->A, 711+1G->T, 2622+1G->A, 405+1G->A, 406-1G->A, 4005+1G->A, 1812-1G->A, 1525-1G->A, 712-1G->T, 1248+1G->A, 1341+1G->A, 3121-1G->A, 4374+1G->T, 3850-1G->A, 2789+5G->A, 3849+10kbC->T, 3272-26A->G, 711+5G->A, 3120G->A, 1811+1.6kbA->G, 711+3A->G, 1898+3A->G, 1717-8G->A, 1342-2A->C, 405+3A->C, 1716G/A, 1811+1G->C, 1898+5G->T, 3850-3T->G, IVS14b+5G->A, 1898+1G->T, 4005+2T->C and 621+3A->G, and one or more human CFTR mutations selected from Δ F508, R117H, and G551D. In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from 1717-1G->A, 1811+1.6kbA->G, 2789+5G->A, 3272-26A->G and 3849+10kbC->T, and one or more human CFTR mutations selected from Δ F508, R117H, and G551D. In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from 2789+5G->A and 3272-26A->G, and one or more human CFTR mutations selected from Δ F508, R117H, and G551D.

[0067] In certain embodiments, the pharmaceutically acceptable composition or tablet of the present invention comprising Compound 1 Form I and a solid dispersion of substantially amorphous Compound 2 are useful for treating, lessening the severity of, or symptomatically treating cystic fibrosis in patients who exhibit residual CFTR activity in the apical membrane of respiratory and non-respiratory epithelia. The presence of residual CFTR activity at the epithelial surface can be readily detected using methods known in the art, e.g., standard electrophysiological, biochemical, or histochemical techniques. Such methods identify CFTR activity using *in vivo* or *ex vivo* electrophysiological techniques, measurement of sweat or salivary Cl^- concentrations, or *ex vivo* biochemical or histochemical techniques to monitor cell surface density. Using such methods, residual CFTR activity can be readily detected in patients heterozygous or homozygous for a variety of different mutations, including patients homozygous or heterozygous for the most common mutation, ΔF508 , as well as other mutations such as the G551D mutation, or the R117H mutation. In certain embodiments, the pharmaceutically acceptable compositions or tablets comprising Compound 1 Form I and a solid dispersion comprising substantially amorphous Compound 2 are useful for treating, lessening the severity of, or symptomatically treating cystic fibrosis in patients who exhibit little to no residual CFTR activity. In certain embodiments, the pharmaceutically acceptable compositions or tablets comprising Compound 1 Form I and a solid dispersion comprising substantially amorphous Compound 2 are useful for treating, lessening the severity of, or symptomatically treating cystic fibrosis in patients who exhibit little to no residual CFTR activity in the apical membrane of respiratory epithelia.

[0068] In another embodiment, the compounds and compositions of the present invention are useful for treating or lessening the severity of cystic fibrosis in patients who have residual CFTR activity induced or augmented using pharmacological methods. In another embodiment, the compounds and compositions of the present invention are useful for treating or lessening the severity of cystic fibrosis in patients who have residual CFTR activity induced or augmented using or gene therapy. Such methods increase the amount of CFTR present at the cell surface, thereby inducing a hitherto absent CFTR activity in a patient or augmenting the existing level of residual CFTR activity in a patient.

[0069] In one embodiment, pharmaceutical compositions and tablets of the present invention comprising Compound 1 Form I and a solid dispersion comprising substantially amorphous Compound 2, as described herein, are useful for treating or lessening the severity of cystic fibrosis in patients within certain genotypes exhibiting residual CFTR activity, e.g., Class I mutations (not synthesized), class II mutation (misfolding), class III mutations (impaired regulation or gating), class IV mutations (altered conductance), or class V mutations (reduced synthesis).

[0070] In one embodiment, pharmaceutical compositions and tablets of the present invention comprising Compound 1 Form I and a solid dispersion comprising substantially amorphous Compound 2, as described herein, are useful for treating, lessening the severity of, or symptomatically treating cystic fibrosis in patients within certain clinical phenotypes, e.g., a moderate to mild clinical phenotype that typically correlates with the amount of residual CFTR activity in the apical membrane of epithelia. Such phenotypes include patients exhibiting pancreatic sufficiency.

[0071] In one embodiment, pharmaceutical compositions and tablets of the present invention comprising Compound 1 Form I and a solid dispersion comprising substantially amorphous Compound 2, as described herein, are useful for treating, lessening the severity of, or symptomatically treating patients diagnosed with pancreatic sufficiency, idiopathic pancreatitis and congenital bilateral absence of the vas deferens, or mild lung disease wherein the patient exhibits residual CFTR activity.

[0072] In one embodiment, pharmaceutical compositions and tablets of the present invention comprising Compound 1 Form I and a solid dispersion comprising substantially amorphous Compound 2, as described herein, are useful for treating, lessening the severity of, or symptomatically treating patients diagnosed with pancreatic sufficiency, idiopathic pancreatitis and congenital bilateral absence of the vas deferens, or mild lung disease wherein the patient has wild type CFTR.

[0073] In addition to cystic fibrosis, modulation of CFTR activity may be beneficial for other diseases not directly caused by mutations in CFTR, such as secretory diseases and other protein folding diseases mediated by CFTR. These include, but are not limited to, chronic obstructive pulmonary disease (COPD), dry eye disease, and Sjögren's Syndrome. COPD is characterized

by airflow limitation that is progressive and not fully reversible. The airflow limitation is due to mucus hypersecretion, emphysema, and bronchiolitis. Activators of mutant or wild-type CFTR offer a potential treatment of mucus hypersecretion and impaired mucociliary clearance that is common in COPD. Specifically, increasing anion secretion across CFTR may facilitate fluid transport into the airway surface liquid to hydrate the mucus and optimized periciliary fluid viscosity. This would lead to enhanced mucociliary clearance and a reduction in the symptoms associated with COPD. Dry eye disease is characterized by a decrease in tear aqueous production and abnormal tear film lipid, protein and mucin profiles. There are many causes of dry eye, some of which include age, Lasik eye surgery, arthritis, medications, chemical/thermal burns, allergies, and diseases, such as cystic fibrosis and Sjögren's syndrome. Increasing anion secretion via CFTR would enhance fluid transport from the corneal endothelial cells and secretory glands surrounding the eye to increase corneal hydration. This would help to alleviate the symptoms associated with dry eye disease. Sjögren's syndrome is an autoimmune disease in which the immune system attacks moisture-producing glands throughout the body, including the eye, mouth, skin, respiratory tissue, liver, vagina, and gut. Symptoms, include, dry eye, mouth, and vagina, as well as lung disease. The disease is also associated with rheumatoid arthritis, systemic lupus, systemic sclerosis, and polymyositis/dermatomyositis. Defective protein trafficking is believed to cause the disease, for which treatment options are limited. Augmenters or inducers of CFTR activity may hydrate the various organs afflicted by the disease and help to alleviate the associated symptoms.

[0074] Anywhere in the present application where a name of a compound may not correctly describe the structure of the compound, the structure supersedes the name and governs.

EXAMPLES

[0075] **Tablet Formation from a Fully Continuous Wet Granulation Process**

[0076] Equipment/Process

[0077] Equipment

[0078] Fully Continuous Development and Launch Rig (DLR) or similar type of equipment.

[0079] Screening

[0080] Compound 1 Form I, the solid dispersion comprising substantially amorphous Compound 2, and excipients may be dispensed in separate intermediate bin containers (IBCs). These materials may be screened using a “bin-to-bin” screening operation. Appropriate screen sizes are mesh 20, mesh 40, or mesh 60.

[0081] Blending

[0082] The IBCs containing the screened Compound 1 Form I, the solid dispersion comprising substantially amorphous Compound 2, and excipients may be docked to the a feeder system, which can feed the materials in a controlled manner, e.g. using volumetric or gravimetric loss in weight feeders, into a continuous blender. The feed rates of the individual components is defined by the formulation composition and the overall line rate. The line rate may be 8 kg/hr to 30 kg/hr. The continuous blender can have different blade configurations to allow appropriate blending and the rotational speed of these blades may be between 80 RPM and 300 RPM.

[0083] Wet Granulation

[0084] A granulation solution may be prepared by dissolving 48 g sodium lauryl sulfate and 159 g polyvinylpyrrolidone in 1,626 g water in a stainless steel container, using an overhead stirrer with a stirring speed of 700 RPM. The granulation solution may be placed in a container from which the solution may be pumped into the twin screw granulator using a peristaltic pump with a mass flow meter and control, using a flow rate that is appropriate for the process. The blend may be granulated using a twin screw granulator such as the granulator that is part of the DLR. The blend may be added to the twin screw granulator using a Loss in Weight feeder, such as the K-Tron feeder on the DLR, with a feed rate of 8 kg/hr to 24 kg/hr. The twin screw granulator may be operated with a barrel temperature of 25 degrees Celsius and a screw speed of 200 to 950 RPM. The granulation process may be performed for three minutes for small batch sizes or several hours for large batch sizes.

[0085] Drying

[0086] The wet granules may be fed directly into a fluid bed dryer, such as the segmented fluid bed dryer on the DLR. The drying end-point may be chosen at a product temperature during discharge ranging from 40 to 55 degrees Celsius at which point the water content of the granules

may be 2.1 %w/w (“Loss on Drying, LOD”) or less. The drying time may be 12 minutes, or shorter or longer, to reach the desired drying endpoint.

[0087] Milling

[0088] The dried granules may be milled to reduce the size of the granules. A cone mill such as the integrated Quadro U10 CoMil may be used for this.

[0089] Blending

[0090] The granules may be blended with extra-granular excipients such as fillers and lubricant using loss in weight feeders and a continuous blender. The blending speed may be 80 – 300 RPM.

[0091] Compression

[0092] The compression blend may be compressed into tablets using a single station or rotary tablet press, such as the Courtoy Modul P press, which is part of the DLR system, using appropriately sized tooling. The weight of the tablets for a dose of 200 mg of Compound 1 Form I and 125 mg of substantially amorphous Compound 2 may be about 500 or 600 mg.

[0093] Film Coating

[0094] Tablets may be film coated using the innovative Omega film coater, which is part of the DLR system. This coater enables fast film coating of sub-batches of 1 to 4 kg to allow continuous manufacturing.

[0095] Printing

[0096] Film coated tablets may be printed with a monogram on one or both tablet faces with, for example, an Ackley ramp printer.

[0097] PAT

[0098] The continuous process described above in one embodiment is enhanced by PAT techniques as described in Table 1. There are 6 PAT positions each of which includes a manual sampling port. In process samples can be obtained for investigational reasons, as needed, and also for PAT model maintenance, transfer, and validation. The PAT systems may be used for real time release testing (RTRT) and may also be employed for in process controls (IPC) and feedback/feed-forward control.

Table 1

Location	Technology	Processing Step	Proposed Purpose	Role
PAT 1	NIR	Dispensing/Charging	Build an NIR raw material library	IPC
PAT 2	NIR	Initial blend	Blend uniformity	IPC
PAT 3	NIR	Wet Granulation	Granule uniformity Moisture	IPC RTRT/IPC
	Laser Diffraction	Wet Granulation	Particle size distribution	RTRT
PAT 4	NIR	Final blend	Blend uniformity Moisture	RTRT RTRT
PAT 5	Raman	Compression	API form Identification	RTRT RTRT
	Tablet Tester	Compression	Weight Thickness Hardness	RTRT/IPC IPC RTRT/IPC
PAT 6	Raman	Coating	Coating thickness	IPC

[0099] Meeting specifications may be done by RTRT as described in Table 2.

Table 2.

Final Product Attribute	PAT Position	In-Process Material	Measurement
Identity	PAT 5a (Raman)	Uncoated Tablet	Confirms spectrum matches the reference standard spectra
Assay	PAT 4 (NIR)	Final Blend	API Concentration
	PAT 5b (Tablet Tester)	Uncoated Tablet	Tablet Weight
CU	PAT 4 (NIR)	Final Blend	Variance in API concentration
	PAT 5b (Tablet Tester)	Uncoated Tablet	Variance in tablet weight
Dissolution	May include:		May include:
	PAT 3b (Laser Diffraction)	Milled granules	Granule Particle Size
	PAT 4 (NIR)	Final Blend	API Concentration

	PAT 5b (Tablet Tester)	Uncoated Tablet	Tablet Weight, Hardness
Moisture	PAT 4	Final Blend	Water Content
Form	PAT 5a (Raman)	Uncoated Tablet	Form I & Absence of Form II

[00100] There is a high probability of detecting non-conforming material. For example, if model classification criterion is set at a minimum of 95% confidence and 800 tablets are tested during batch manufacture, 40 hour run with a sampling rate of 1 tablet every 3 minutes equals 800 tablets. Then, probability of passing a non-conforming batch is extremely low: $<(0.05)^n$, where $n = \#$ of samples, therefore the probability is $< 1.5 \times 10^{-1041}$. Probability of not detecting non-conforming tablets resulting from a short term event (≥ 3 minutes) is as follows: 1 tablet (3 min event) $\rightarrow <0.05$ (probability of detection > 0.95); 2 tablets (6 minute event) $\rightarrow <0.0025$ (probability of detection > 0.9975).

[00101] PAT measurements can serve as surrogates for conventional end-testing directly via combining measurements to express attributes conventionally (i.e. as assay, CU, dissolution, etc.). Validation can be performed using ICH Q2 as guidance. Sequential off-line to on-line method development allows for the assessment of CQAs in a material sparing manner. Ultimately, RTRT will lead to ensuring product quality at a higher confidence level than conventional testing.

[00102] HTT HPLC

[00103] In one embodiment, the continuous process of manufacturing of the present invention utilizes high throughput testing (HTT) HPLC methods to validate samples. High throughput testing HPLC methods achieve 24 hour sample turnaround time for at least 300 samples by improving sample preparation techniques, emphasizing generic analysis methods, using well defined sample workflows, and automating data processing.

[00104] Sample preparation takes the majority of an FTE's time and is the source of most errors. It is often overlooked during method development. In one embodiment, improved sample preparation techniques comprise using wide mouth disposable bottles. In another embodiment, improved sample preparation techniques comprise adding the entire vial of a sample to a disposable bottle, adding diluent, shaking overnight, and centrifuging.

[00105] Generic HPLC methods can be developed and validated for multiple projects. Common HPLC columns and commercial mobile phases can be used. Additional analysis improvements include leveraged standard stability and utilizing injection overlap.

[00106] In another embodiment, HTT HPLC is used in the development of the process analytical techniques as a way of correlating the spectroscopic data collected from the process analytical techniques with an absolute number.

[00107] In one embodiment, the present invention features a process of conducting high throughput HPLC comprising a) dropping containers, such as vials, of pre-weighed samples into plastic bottles, such as HDPE bottles; b) adding solution to each set of container and bottle via a bottle top dispenser; c) shaking the sets of plastic bottles, vials, and solutions until samples are dissolved; d) centrifuging the sets of plastic bottles, vials, and solutions; e) loading an aliquot of supernatant from the centrifuge step onto an HPLC column; and f) running the column with a mobile phase.

[00108] The advantage of HTT HPLC is that it can measure a high volume of samples in a timely, accurate, and cost effective manner. The sample preparation uses plastic bottles as the main vessel which can be placed in large number on a shaker and then transferred directly to a centrifuge. This avoids the more time consuming step of filtering the solution of sample. Additionally, the size of the plastic bottle allows the sample to be added directly by simply dropping the container, such as a vial, of sample into the plastic bottle. Commercially available solution dispensers can then be used to add a fixed amount of solution, thus avoiding another time consuming step of pipetting the solution in.

[00109] Table 3 summarizes the benefits of high throughput testing HPLC compared to traditional HPLC testing methods.

Table 3.

Traditional Method	HTT Method
Samples added to volumetric flasks	Samples added to disposable HDPE bottles
Premixed diluent	Off shelf solvents mixed into sample bottle (no diluent prep necessary)
Diluent added and the QS'd to line	Calibrated bottle top dispensers dispense solvents
For BU: quantitative transfer	For BU: sample bottles rinsed in bottle
Sonication and shaking of samples	Shaking only


Secondary dilution and QS	No dilution (injection volume driven)
Samples filtered	Samples centrifuged
HPLC is project specific (variable MP and column combinations)	HPLC is generic (uses fixed column, fixed mobile phase A and B, and commercially manufactured mobile phases)

OTHER EMBODIMENTS


[00110] All publications and patents referred to in this disclosure are incorporated herein by reference to the same extent as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference. Should the meaning of the terms in any of the patents or publications incorporated by reference conflict with the meaning of the terms used in this disclosure, the meaning of the terms in this disclosure are intended to be controlling. Furthermore, the foregoing discussion discloses and describes merely exemplary embodiments of the invention. One skilled in the art will readily recognize from such discussion and from the accompanying drawings and claims, that various changes, modifications and variations can be made therein without departing from the spirit and scope of the invention as defined in the following claims.

CLAIMS


1. A process of conducting high throughput HPLC comprising:
 - a) dropping containers of pre-weighed samples into plastic bottles;
 - b) adding solution to each set of container and bottle via a bottle top dispenser;
 - c) shaking each set of plastic bottle, container, and solution until sample is dissolved;
 - d) centrifuging each set of plastic bottle, container, and solution;
 - e) loading an aliquot of supernatant from the centrifuge step onto an HPLC column; and
 - f) running the column with a mobile phase.
2. The process of claim 1, wherein the containers of step a) are vials.
3. The process of claim 1, wherein the plastic bottles of step a) are HDPE bottles.
4. The process of claim 1, further comprising correlating the results from the process of claim 1 to process analytical technique (PAT) measurements for continuous manufacturing.
5. The process of claim 4, wherein continuous manufacturing is for a pharmaceutical composition.
6. The process of claim 5, wherein the pharmaceutical composition is a tablet.
7. The process of claim 6, wherein the tablet is for the treatment of a cystic fibrosis transmembrane conductance regulator (CFTR) mediated disease.
8. The process of claim 7, wherein the CFTR mediated disease is cystic fibrosis.
9. The process of claim 6, wherein the tablet comprises two active pharmaceutical ingredients (API).
10. The process of claim 9, wherein one API is a cystic fibrosis (CF) corrector.
11. The process of claim 9, wherein one API is a CF potentiator.
12. The process of claim 9, wherein one API is a CF corrector and the other API is a CF potentiator.
13. The process of claim 9, wherein one API is




14. The process of claim 9, wherein one API is




15. The process of claim 9, wherein one API is




and the other API is



16. The process of claim 15, wherein



is in Form I, and



is substantially amorphous in the form of a solid dispersion.