

- 1 -

SPIRO-LACTAM NMDA RECEPTOR MODULATORS AND USES THEREOF**CROSS REFERENCE TO RELATED APPLICATIONS**

[0001] This application claims the benefit of United States Provisional Application No. 61/757,920, filed on January 29, 2013, which is incorporated by reference in its entirety.

BACKGROUND

[0002] An N-methyl-d-aspartate (NMDA) receptor is a postsynaptic, ionotropic receptor that is responsive to, *inter alia*, the excitatory amino acids glutamate and glycine and the synthetic compound NMDA. The NMDA receptor controls the flow of both divalent and monovalent ions into the postsynaptic neural cell through a receptor associated channel (Foster *et al.*, Nature 1987, 329:395-396; Mayer *et al.*, Trends in Pharmacol. Sci. 1990, 11:254-260). The NMDA receptor has been implicated during development in specifying neuronal architecture and synaptic connectivity, and may be involved in experience-dependent synaptic modifications. In addition, NMDA receptors are also thought to be involved in long term potentiation and central nervous system disorders.

[0003] The NMDA receptor plays a major role in the synaptic plasticity that underlies many higher cognitive functions, such as memory acquisition, retention and learning, as well as in certain cognitive pathways and in the perception of pain (Collingridge *et al.*, The NMDA Receptor, Oxford University Press, 1994). In addition, certain properties of NMDA receptors suggest that they may be involved in the information-processing in the brain that underlies consciousness itself.

[0004] The NMDA receptor has drawn particular interest since it appears to be involved in a broad spectrum of CNS disorders. For instance, during brain ischemia caused by stroke or traumatic injury, excessive amounts of the excitatory amino acid glutamate are released from damaged or oxygen deprived neurons. This excess glutamate binds to the NMDA receptors which opens their ligand-gated ion channels; in turn the calcium influx produces a high level of intracellular calcium which activates a biochemical cascade resulting in protein degradation and cell death. This phenomenon, known as excitotoxicity, is also thought to be responsible for

- 2 -

the neurological damage associated with other disorders ranging from hypoglycemia and cardiac arrest to epilepsy. In addition, there are preliminary reports indicating similar involvement in the chronic neurodegeneration of Huntington's, Parkinson's, and Alzheimer's diseases. Activation of the NMDA receptor has been shown to be responsible for post-stroke convulsions, and, in certain models of epilepsy, activation of the NMDA receptor has been shown to be necessary for the generation of seizures. Neuropsychiatric involvement of the NMDA receptor has also been recognized since blockage of the NMDA receptor Ca^{++} channel by the animal anesthetic PCP (phencyclidine) produces a psychotic state in humans similar to schizophrenia (reviewed in Johnson, K. and Jones, S., 1990). Further, NMDA receptors have also been implicated in certain types of spatial learning.

[0005] The NMDA receptor is believed to consist of several protein chains embedded in the postsynaptic membrane. The first two types of subunits discovered so far form a large extracellular region, which probably contains most of the allosteric binding sites, several transmembrane regions looped and folded so as to form a pore or channel, which is permeable to Ca^{++} , and a carboxyl terminal region. The opening and closing of the channel is regulated by the binding of various ligands to domains (allosteric sites) of the protein residing on the extracellular surface. The binding of the ligands is thought to affect a conformational change in the overall structure of the protein which is ultimately reflected in the channel opening, partially opening, partially closing, or closing.


[0006] NMDA receptor compounds may exert dual (agonist/antagonist) effect on the NMDA receptor through the allosteric sites. These compounds are typically termed "partial agonists". In the presence of the principal site ligand, a partial agonist will displace some of the ligand and thus decrease Ca^{++} flow through the receptor. In the absence of or lowered level of the principal site ligand, the partial agonist acts to increase Ca^{++} flow through the receptor channel.

[0007] A need continues to exist in the art for novel and more specific/potent compounds that are capable of binding the glycine binding site of NMDA receptors, and provide pharmaceutical benefits. In addition, a need continues to exist in the medical arts for orally deliverable forms of such compounds.

- 3 -

SUMMARY

[0008] Provided herein, at least in part, are compounds that are NMDA modulators, for example, partial agonists of NMDA. For example, disclosed herein are compounds represented by the formula:



5 and pharmaceutically acceptable salts, stereoisomers, and N-oxides thereof, wherein

R_b is selected from the group consisting of H, halogen, hydroxyl, cyano or C_1 - C_6 alkyl;

R_1 is H or C_1 - C_6 alkyl;

R_2 is H or C_1 - C_6 alkyl;

10 R_3 is selected from the group consisting of H, C_1 - C_6 alkyl and a nitrogen protecting group;

R_4 is H or C_1 - C_6 alkyl;

R_5 is X or $-C_1$ - C_6 alkylene-X, wherein X is selected from the group consisting of phenyl, a 4- to 6-membered heteroaryl ring having 1, 2, or 3 heteroatoms selected from O, S, or N, or a 4- to 6-membered heterocyclyl ring having 1, 2, or 3 heteroatoms selected from O, S, or

15 N, and wherein X is optionally substituted with ; and

R_6 is selected from the group consisting of H, halogen, hydroxyl, cyano, $-O-C(O)-C_1$ - C_6 alkyl, C_1 - C_6 alkyl, or C_1 - C_6 alkoxy;

or, in other embodiments, the variables set forth in formula (I) are defined as follows:

R_b is selected from the group consisting of H, halogen, hydroxyl, cyano or C_1 - C_6 alkyl;

20 R_1 is H or C_1 - C_6 alkyl;

R_2 is H or C_1 - C_6 alkyl;

R_3 is selected from the group consisting of H, C_1 - C_6 alkyl and a nitrogen protecting group;

R_4 is H or C_1 - C_6 alkyl;


- 4 -

R₅ is X or -C₁-C₆ alkylene-X, wherein X is selected from the group consisting of:

(i) phenyl;


(ii) heteroaryl including from 5 to 6 ring atoms wherein 1, 2, or 3 of the ring atoms are independently selected from the group consisting of N, NH, N(C₁-C₃ alkyl), O, and S; and

5 (iii) heterocyclyl including from 3 to 6 ring atoms wherein 1, 2, or 3 of the ring atoms are independently selected from the group consisting of N, NH, N(C₁-C₃ alkyl), O, and S;

wherein R₅ is optionally substituted with ; and

R₆ is selected from the group consisting of H, halogen, hydroxyl, cyano, -O-C(O)-C₁-C₆ alkyl, C₁-C₆ alkyl, or C₁-C₆ alkoxy.

10 **[0009]** Also disclosed herein are compounds represented by the formula:




and pharmaceutically acceptable salts, stereoisomers, and N-oxides thereof, wherein

R₃ is selected from the group consisting of H, C₁-C₆alkyl and a nitrogen protecting group; and

15 R₇ is a 4- to 6-membered heteroaryl ring having 1, 2, or 3 heteroatoms selected from O, S, or N, optionally substituted on a free carbon by a substituent selected from the group consisting of: halogen, C₁-C₆alkyl, hydroxyl, cyano, and phenyl.

[0010] Also provided herein are compounds represented by the formula:



and pharmaceutically acceptable salts, stereoisomers, and N-oxides thereof, wherein

20 R_b is selected from the group consisting of H, halogen, hydroxyl, cyano or C₁-C₆ alkyl;

R₁ is H or C₁-C₆ alkyl;

R₂ is H or C₁-C₆ alkyl;

R₃ is selected from the group consisting of H, C₁-C₆ alkyl and a nitrogen protecting group;

- 5 -

R₄ is H or C₁-C₆ alkyl;

X is selected from the group consisting of phenyl, a 4- to 6-membered heteroaryl ring having 1, 2, or 3 heteroatoms selected from O, S, or N, or a 4- to 6-membered heterocyclyl ring having 1,

2, or 3 heteroatoms selected from O, S, or N, wherein  is present on a free carbon of X;

5 and

R₆ is selected from the group consisting of H, halogen, hydroxyl, cyano, -O-C(O)-C₁-C₆ alkyl, C₁-C₆ alkyl, or C₁-C₆ alkoxy.

[0011] Also provided herein are pharmaceutically acceptable compositions comprising a disclosed compound, and a pharmaceutically acceptable excipient. For example, such
10 compositions may be suitable for oral or intravenous administration to a patient.

[0012] In another aspect, a method of treating a condition selected from the group consisting of autism, anxiety, depression, bipolar disorder, attention deficit disorder, attention deficit hyperactivity disorder (ADHD), schizophrenia, a psychotic disorder, a psychotic symptom, social withdrawal, obsessive-compulsive disorder, phobia, post-traumatic stress
15 syndrome, a behavior disorder, an impulse control disorder, a substance abuse disorder, a sleep disorder, a memory disorder, a learning disorder, urinary incontinence, multiple system atrophy, progressive supra-nuclear palsy, Friedrich's ataxia, Down's syndrome, fragile X syndrome, tuberous sclerosis, olivio-ponto-cerebellar atrophy, cerebral palsy, drug-induced optic neuritis, ischemic retinopathy, diabetic retinopathy, glaucoma, dementia, AIDS dementia,
20 Alzheimer's disease, Huntington's chorea, spasticity, myoclonus, muscle spasm, Tourette's syndrome, epilepsy, cerebral ischemia, stroke, a brain tumor, traumatic brain injury, cardiac arrest, myelopathy, spinal cord injury, peripheral neuropathy, acute neuropathic pain, and chronic neuropathic, in a patient in need thereof is provided. Such methods may comprise administering to the patient a pharmaceutically effective amount of a disclosed compound or
25 pharmaceutically acceptable salts, stereoisomers, N-oxides, and hydrates thereof.

[0013] In some embodiments, a contemplated method includes treating depression. For example, depression may include one or more of major depressive disorder, dysthymic disorder, psychotic depression, postpartum depression, seasonal affective disorder, bipolar disorder, mood disorder, or depression caused by a chronic medical condition. In other
30 embodiments, a contemplated method may treat schizophrenia. Such schizophrenia may be,

- 6 -

for example, paranoid type schizophrenia, disorganized type schizophrenia, catatonic type schizophrenia, undifferentiated type schizophrenia, residual type schizophrenia, post-schizophrenic depression, or simple schizophrenia.

BRIEF DESCRIPTION OF THE DRAWINGS

5 **[0014]** **Figure 1** shows the potentiation of [³H]MK-801 binding in the presence of Compound Y.

DETAILED DESCRIPTION

[0015] This disclosure is generally directed to compounds that are capable of modulating NMDA, e.g., NMDA antagonists or partial agonists, and compositions and/or methods of using the disclosed compounds.

Definitions

10 **[0016]** “Treating” includes any effect, e.g., lessening, reducing, modulating, or eliminating, that results in the improvement of the condition, disease, disorder and the like.

15 **[0017]** The term “alkenyl” as used herein refers to an unsaturated straight or branched hydrocarbon having at least one carbon-carbon double bond, such as a straight or branched group of 2-6 or 3-4 carbon atoms, referred to herein for example as C₂-C₆ alkenyl, and C₃-C₄ alkenyl, respectively. Exemplary alkenyl groups include, but are not limited to, vinyl, allyl, butenyl, pentenyl, etc.

20 **[0018]** The term “alkoxy” as used herein refers to a straight or branched alkyl group attached to an oxygen (alkyl-O-). Exemplary alkoxy groups include, but are not limited to, alkoxys of 1-6 or 2-6 carbon atoms, referred to herein as C₁-C₆ alkoxy, and C₂-C₆ alkoxy, respectively. Exemplary alkoxy groups include, but are not limited to methoxy, ethoxy, isopropoxy, etc.

25 **[0019]** The term “alkenyloxy” used herein refers to a straight or branched alkenyl group attached to an oxygen (alkenyl-O). Exemplary alkenoxy group include, but are not limited to, groups with an alkenyl group of 3-6 carbon atoms, (also e.g. referred to as C₃-C₆ alkenyloxy). Exemplary “alkenoxo” groups include, but are not limited to allyloxy, butenyloxy, etc.

[0020] The term “alkynyloxy” used herein refers to a straight or branched alkynyl group attached to an oxygen (alkynyl-O)). Exemplary alkynyloxy groups include, but are not limited to, C₃-C₆ alkynyloxy, e.g., propynyloxy.

- 7 -

[0021] The term “alkyl” as used herein refers to a saturated straight or branched hydrocarbon, such as a straight or branched group of 1-6, 1-4, or 1-3 carbon atoms, referred to herein as C₁-C₆ alkyl, C₁-C₄ alkyl, and C₁-C₃ alkyl, respectively. Exemplary alkyl groups include, but are not limited to, methyl, ethyl, propyl, isopropyl, 2-methyl-1-propyl, 2-methyl-2-propyl, 2-methyl-1-butyl, 3-methyl-1-butyl, 3-methyl-2-butyl, 2,2-dimethyl-1-propyl, 2-methyl-1-pentyl, 3-methyl-1-pentyl, 4-methyl-1-pentyl, 2-methyl-2-pentyl, 3-methyl-2-pentyl, 4-methyl-2-pentyl, 2,2-dimethyl-1-butyl, 3,3-dimethyl-1-butyl, 2-ethyl-1-butyl, butyl, isobutyl, t-butyl, pentyl, isopentyl, neopentyl, hexyl, etc. The term “haloalkyl” as used herein refers to a saturated straight or branched alkyl groups, in which one or more hydrogen atoms of the alkyl group are replaced with one or more independently selected halogens. The term “haloalkyl” encompasses alkyl groups in which all of hydrogen atoms of the alkyl group are replaced independently selected halogens (sometimes referred to as “perhalo” alkyl groups. Exemplary haloalkyl groups include, but are not limited to, CH₂F, CH₂CH₂Cl, CF₃, CHFCH₂Cl.

[0022] The term “alkynyl” as used herein refers to an unsaturated straight or branched hydrocarbon having at least one carbon-carbon triple bond, such as a straight or branched group of 2-6, or 3-6 carbon atoms, referred to herein as C₂-C₆ alkynyl, and C₃-C₆ alkynyl, respectively. Exemplary alkynyl groups include, but are not limited to, ethynyl, propynyl, butynyl, pentynyl, hexynyl, methylpropynyl, etc.

[0023] The term “bridged cycloalkyl”, as used herein, is defined as a monocyclic 4- to 7-membered cycloalkyl group in which two non-adjacent atoms are linked by a CH₂ or CH₂CH₂ group. A “bridged cycloalkyl” may be fused to one or more phenyl, partially unsaturated, or saturated rings. Examples of bridged carbocyclic groups include but are not limited to bicyclo[2.2.1]heptane, bicyclo[2.2.2]octane, bicyclo[2.2.2]octene etc.

[0024] The term “carbonyl” as used herein refers to the radical -C(O)-. The term “cyano” as used herein refers to the radical -CN. The term “nitro” refers to the radical -NO₂. The term “H” refers to hydrogen.

[0025] The term “cycloalkoxy” as used herein refers to a cycloalkyl group attached to an oxygen (cycloalkyl-O-).

[0026] The term “cycloalkyl” as used herein refers to a monocyclic saturated or partially unsaturated hydrocarbon group of for example 3-6, or 4-6 carbons, referred to herein, e.g., as “C₃₋₆ cycloalkyl” or “C₄₋₆ cycloalkyl,” and derived from a cycloalkane. Exemplary cycloalkyl

- 8 -

groups include, but are not limited to, cyclohexyl, cyclohexenyl, cyclopentyl, cyclobutyl, cyclopropyl or cyclopentyl.

[0027] The terms “halo” or “halogen” as used herein refer to F, Cl, Br, or I.

[0028] The terms “heteroaryl” as used herein refers to a monocyclic aromatic 4-6
5 membered ring system containing one or more heteroatoms, for example one to three heteroatoms, such as nitrogen, oxygen, and sulfur. Where possible, said heteroaryl ring may be linked to the adjacent radical through carbon or nitrogen. Examples of heteroaryl rings include but are not limited to furyl, thienyl, pyrrolyl, thiazolyl, oxazolyl, isothiazolyl, isoxazolyl, imidazolyl, pyrazolyl, triazolyl, oxadiazolyl (e.g., 1,2,4- oxadiazolyl or 1,3,4- oxadiazolyl),
10 pyridyl, and pyrimidinyl.

[0029] The terms “heterocyclyl” or “heterocyclic group” are art-recognized and refer to saturated or partially unsaturated 4- to 7-membered ring structures, whose ring structures include one to three heteroatoms, such as nitrogen, oxygen, and sulfur. A heterocycle may be fused to one or more phenyl, partially unsaturated, or saturated rings. Examples of heterocyclyl
15 groups include but are not limited to pyrrolidinyl, piperidinyl, morpholino, thiomorpholino, and piperazinyl.

[0030] The term “heterocyclalkoxy” as used herein refers to a heterocyclyl- alkyl-O- group.

[0031] The term “heterocycloxyalkyl” refers to a heterocyclyl-O-alkyl- group.

20 [0032] The term “heterocycloxy” refers to a heterocyclyl-O- group. The term “cycloalkyloxy” refers to a cycloalkyl-O- group.

[0033] The term “heteroaryloxy” refers to a heteroaryl-O- group.

[0034] The terms “hydroxy” and “hydroxyl” as used herein refers to the radical -OH.

[0035] The term “oxo” as used herein refers to the radical =O.

25 [0036] The term “nitrogen protecting group” or “amino protecting group” is art-recognized and as used herein refers to a chemical moiety that is covalently linked to a nitrogen atom of an amino (primary or secondary) group and that temporarily blocks the reactivity of the amino group during a synthetic step and is selectively removed once the synthetic step is complete. Nitrogen protecting groups include, for example, 9-Fluorenylmethyloxycarbonyl (Fmoc), tert-
30 butoxycarbonyl (Boc), carbobenzyloxycarbonyl (Cbz), p-methoxybenzyloxycarbonyl, acetyl,

- 9 -

trifluoroacetyl, benzoyl, phthalimido, benzyl (Bn), p-methoxybenzyl, p-methoxyphenyl, 3,4-dimethoxybenzyl, triphenylmethyl, benzyldiene, and p-toluenesulfonyl (Ts). In some embodiments, the nitrogen protecting group can have one of the following formulas: –

C(O)OR₃₁ or –C(O)R₃₂ as defined herein. In certain embodiments, R₃₁ is selected from the

- 5 group consisting of: C₁-C₆ alkyl; C₁-C₆ haloalkyl; C₂-C₆ alkenyl; C₂-C₆ alkynyl; C₃-C₁₀ cycloalkyl, wherein the C₃-C₁₀ cycloalkyl is optionally substituted with from 1-3 independently selected C₁-C₃ alkyl; –CH₂-C₃-C₁₀ cycloalkyl wherein the C₃-C₁₀ cycloalkyl is optionally substituted with from 1-3 independently selected C₁-C₃ alkyl; –CH₂-phenyl, wherein the phenyl is optionally substituted with from 1-2 substituents independently selected from C₁-C₃ alkyl, C₁-C₃ haloalkyl, C₁-C₃ alkoxy, C₁-C₃ haloalkoxy, nitro, halo, SO₂Me, cyano, and –OC(O)CH₃; and –CH₂-pyridyl. In certain embodiments, R₃₂ is selected from the group consisting of: H; C₁-C₆ alkyl; C₁-C₆ haloalkyl; phenyl, wherein the phenyl is optionally substituted with from 1-2 substituents independently selected from C₁-C₃ alkyl, C₁-C₃ haloalkyl, C₁-C₃ alkoxy, C₁-C₃ haloalkoxy, nitro, halo, SO₂Me, cyano, and –OC(O)CH₃; and pyridyl.

- 15 **[0037]** As used in the present disclosure, the term “partial NMDA receptor agonist” generally refers to a compound that is capable of binding to a glycine binding site of an NMDA receptor; at low concentrations a NMDA receptor agonist acts substantially as agonist and at high concentrations it acts substantially as an antagonist. These concentrations are experimentally determined for each and every “partial agonist.

- 20 **[0038]** “Pharmaceutically or pharmacologically acceptable” include molecular entities and compositions that do not produce an adverse, allergic or other untoward reaction when administered to an animal, or a human, as appropriate. For human administration, preparations should meet sterility, pyrogenicity, general safety and purity standards as required by FDA Office of Biologics standards.

- 25 **[0039]** The term “pharmaceutically acceptable carrier” or “pharmaceutically acceptable excipient” as used herein refers to any and all solvents, dispersion media, coatings, isotonic and absorption delaying agents, and the like, that are compatible with pharmaceutical administration. The use of such media and agents for pharmaceutically active substances is well known in the art. The compositions may also contain other active compounds providing
30 supplemental, additional, or enhanced therapeutic functions.

[0040] The term “pharmaceutical composition” as used herein refers to a composition comprising at least one compound as disclosed herein formulated together with one or more pharmaceutically acceptable carriers.

[0041] “Individual,” “patient,” or “subject” are used interchangeably and include any
5 animal, including mammals, preferably mice, rats, other rodents, rabbits, dogs, cats, swine, cattle, sheep, horses, or primates, and most preferably humans. The compounds of the invention can be administered to a mammal, such as a human, but can also be administered to other mammals such as an animal in need of veterinary treatment, *e.g.*, domestic animals (*e.g.*, dogs, cats, and the like), farm animals (*e.g.*, cows, sheep, pigs, horses, and the like) and
10 laboratory animals (*e.g.*, rats, mice, guinea pigs, and the like). The mammal treated in the methods of the invention is desirably a mammal in which treatment *e.g.*, of pain or depression is desired. “Modulation” includes antagonism (*e.g.*, inhibition), agonism, partial antagonism and/or partial agonism.

[0042] In the present specification, the term “therapeutically effective amount” means the
15 amount of the subject compound that will elicit the biological or medical response of a tissue, system, animal or human that is being sought by the researcher, veterinarian, medical doctor or other clinician. The compounds of the invention are administered in therapeutically effective amounts to treat a disease. Alternatively, a therapeutically effective amount of a compound is the quantity required to achieve a desired therapeutic and/or prophylactic effect, such as an
20 amount which results in lessening a symptom of depression.

[0043] The term “pharmaceutically acceptable salt(s)” as used herein refers to salts of acidic or basic groups that may be present in compounds used in the present compositions. Compounds included in the present compositions that are basic in nature are capable of forming a wide variety of salts with various inorganic and organic acids. The acids that may be used to
25 prepare pharmaceutically acceptable acid addition salts of such basic compounds are those that form non-toxic acid addition salts, *i.e.*, salts containing pharmacologically acceptable anions, including but not limited to malate, oxalate, chloride, bromide, iodide, nitrate, sulfate, bisulfate, phosphate, acid phosphate, isonicotinate, acetate, lactate, salicylate, citrate, tartrate, oleate, tannate, pantothenate, bitartrate, ascorbate, succinate, maleate, gentisinate, fumarate, gluconate,
30 glucuronate, saccharate, formate, benzoate, glutamate, methanesulfonate, ethanesulfonate, benzenesulfonate, *p*-toluenesulfonate and pamoate (*i.e.*, 1,1'-methylene-bis-(2-hydroxy-3-naphthoate)) salts. Compounds included in the present compositions that are acidic in nature

- 11 -

are capable of forming base salts with various pharmacologically acceptable cations. Examples of such salts include alkali metal or alkaline earth metal salts and, particularly, calcium, magnesium, sodium, lithium, zinc, potassium, and iron salts. Compounds included in the present compositions that include a basic or acidic moiety may also form pharmaceutically acceptable salts with various amino acids. The compounds of the disclosure may contain both acidic and basic groups; for example, one amino and one carboxylic acid group. In such a case, the compound can exist as an acid addition salt, a zwitterion, or a base salt.

[0044] The compounds of the disclosure may contain one or more chiral centers and/or double bonds and, therefore, exist as stereoisomers, such as geometric isomers, enantiomers or diastereomers. The term “stereoisomers” when used herein consist of all geometric isomers, enantiomers or diastereomers. These compounds may be designated by the symbols “R” or “S,” depending on the configuration of substituents around the stereogenic carbon atom. The present invention encompasses various stereoisomers of these compounds and mixtures thereof. Stereoisomers include enantiomers and diastereomers. Mixtures of enantiomers or diastereomers may be designated “(±)” in nomenclature, but the skilled artisan will recognize that a structure may denote a chiral center implicitly.

[0045] The compounds of the disclosure may contain one or more chiral centers and/or double bonds and, therefore, exist as geometric isomers, enantiomers or diastereomers. The enantiomer and diastereomers may be designated by the symbols “(+)”, “(-)”, “R” or “S,” depending on the configuration of substituents around the stereogenic carbon atom, but the skilled artisan will recognize that a structure may denote a chiral center implicitly. Geometric isomers, resulting from the arrangement of substituents around a carbon-carbon double bond or arrangement of substituents around a cycloalkyl or heterocyclic ring, can also exist in the compounds of the present invention. The symbol  denotes a bond that may be a single, double or triple bond as described herein. Substituents around a carbon-carbon double bond are designated as being in the “Z” or “E” configuration wherein the terms “Z” and “E” are used in accordance with IUPAC standards. Unless otherwise specified, structures depicting double bonds encompass both the “E” and “Z” isomers. Substituents around a carbon-carbon double bond alternatively can be referred to as “cis” or “trans,” where “cis” represents substituents on the same side of the double bond and “trans” represents substituents on opposite sides of the double bond. The arrangement of substituents around a carbocyclic ring can also be designated as “cis” or “trans.” The term “cis” represents substituents on the same side of the plane of the ring and the term “trans” represents substituents on opposite sides of the plane of the ring.

- 12 -

Mixtures of compounds wherein the substituents are disposed on both the same and opposite sides of plane of the ring are designated "cis/trans."

[0046] The term "stereoisomers" when used herein consist of all geometric isomers, enantiomers or diastereomers. The present invention encompasses various stereoisomers of these compounds and mixtures thereof.

[0047] Individual enantiomers and diastereomers of compounds of the present invention can be prepared synthetically from commercially available starting materials that contain asymmetric or stereogenic centers, or by preparation of racemic mixtures followed by resolution methods well known to those of ordinary skill in the art. These methods of resolution are exemplified by (1) attachment of a mixture of enantiomers to a chiral auxiliary, separation of the resulting mixture of diastereomers by recrystallization or chromatography and liberation of the optically pure product from the auxiliary, (2) salt formation employing an optically active resolving agent, (3) direct separation of the mixture of optical enantiomers on chiral liquid chromatographic columns or (4) kinetic resolution using stereoselective chemical or enzymatic reagents. Racemic mixtures can also be resolved into their component enantiomers by well-known methods, such as chiral-phase gas chromatography or crystallizing the compound in a chiral solvent. Stereoselective syntheses, a chemical or enzymatic reaction in which a single reactant forms an unequal mixture of stereoisomers during the creation of a new stereocenter or during the transformation of a pre-existing one, are well known in the art. Stereoselective syntheses encompass both enantio- and diastereoselective transformations. For examples, see Carreira and Kvaerno, *Classics in Stereoselective Synthesis*, Wiley-VCH: Weinheim, 2009.

[0048] The compounds disclosed herein can exist in solvated as well as unsolvated forms with pharmaceutically acceptable solvents such as water, ethanol, and the like, and it is intended that the invention embrace both solvated and unsolvated forms. In one embodiment, the compound is amorphous. In one embodiment, the compound is a single polymorph. In another embodiment, the compound is a mixture of polymorphs. In another embodiment, the compound is in a crystalline form.

[0049] The invention also embraces isotopically labeled compounds of the invention which are identical to those recited herein, except that one or more atoms are replaced by an atom having an atomic mass or mass number different from the atomic mass or mass number usually found in nature. Examples of isotopes that can be incorporated into compounds of the

invention include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorus, fluorine and chlorine, such as ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F , and ^{36}Cl , respectively. For example, a compound of the invention may have one or more H atom replaced with deuterium.

[0050] Certain isotopically-labeled disclosed compounds (*e.g.*, those labeled with ^3H and ^{14}C) are useful in compound and/or substrate tissue distribution assays. Tritiated (*i.e.*, ^3H) and carbon-14 (*i.e.*, ^{14}C) isotopes are particularly preferred for their ease of preparation and detectability. Further, substitution with heavier isotopes such as deuterium (*i.e.*, ^2H) may afford certain therapeutic advantages resulting from greater metabolic stability (*e.g.*, increased *in vivo* half-life or reduced dosage requirements) and hence may be preferred in some circumstances.

10 Isotopically labeled compounds of the invention can generally be prepared by following procedures analogous to those disclosed in the *e.g.*, Examples herein by substituting an isotopically labeled reagent for a non-isotopically labeled reagent.

[0051] The term “prodrug” refers to compounds that are transformed *in vivo* to yield a disclosed compound or a pharmaceutically acceptable salt, hydrate or solvate of the compound.

15 The transformation may occur by various mechanisms (such as by esterase, amidase, phosphatase, oxidative and or reductive metabolism) in various locations (such as in the intestinal lumen or upon transit of the intestine, blood or liver). Prodrugs are well known in the art (for example, see Rautio, Kumpulainen, *et al*, Nature Reviews Drug Discovery 2008, 7, 255). For example, if a compound of the invention or a pharmaceutically acceptable salt,

20 hydrate or solvate of the compound contains a carboxylic acid functional group, a prodrug can comprise an ester formed by the replacement of the hydrogen atom of the acid group with a group such as (C₁-C₈)alkyl, (C₂-C₁₂)alkanoyloxymethyl, 1-(alkanoyloxy)ethyl having from 4 to 9 carbon atoms, 1-methyl-1-(alkanoyloxy)-ethyl having from 5 to 10 carbon atoms, alkoxycarbonyloxymethyl having from 3 to 6 carbon atoms, 1-(alkoxycarbonyloxy)ethyl

25 having from 4 to 7 carbon atoms, 1-methyl-1-(alkoxycarbonyloxy)ethyl having from 5 to 8 carbon atoms, N-(alkoxycarbonyl)aminomethyl having from 3 to 9 carbon atoms, 1-(N-(alkoxycarbonyl)amino)ethyl having from 4 to 10 carbon atoms, 3-phthalidyl, 4-crotonolactonyl, gamma-butyrolacton-4-yl, di-N,N-(C₁-C₂)alkylamino(C₂-C₃)alkyl (such as β-dimethylaminoethyl), carbamoyl-(C₁-C₂)alkyl, N,N-di(C₁-C₂)alkylcarbamoyl-(C₁-C₂)alkyl

30 and piperidino-, pyrrolidino- or morpholino(C₂-C₃)alkyl.

[0052] Similarly, if a compound of the invention contains an alcohol functional group, a prodrug can be formed by the replacement of the hydrogen atom of the alcohol group with a


- 14 -

group such as (C₁-C₆)alkanoyloxymethyl, 1-((C₁-C₆)alkanoyloxy)ethyl, 1-methyl-1-((C₁-C₆)alkanoyloxy)ethyl (C₁-C₆)alkoxycarbonyloxymethyl, N-(C₁-C₆)alkoxycarbonylaminomethyl, succinoyl, (C₁-C₆)alkanoyl, α -amino(C₁-C₄)alkanoyl, arylacyl and α -aminoacyl, or α -aminoacyl- α -aminoacyl, where each α -aminoacyl group is independently selected from the naturally occurring L-amino acids, P(O)(OH)₂, -P(O)(O(C₁-C₆)alkyl)₂ or glycosyl (the radical resulting from the removal of a hydroxyl group of the hemiacetal form of a carbohydrate).

[0053] If a compound of the invention incorporates an amine functional group, a prodrug can be formed, for example, by creation of an amide or carbamate, an N-acyloxyakyl derivative, an (oxodioxolenyl)methyl derivative, an N-Mannich base, imine or enamine. In addition, a secondary amine can be metabolically cleaved to generate a bioactive primary amine, or a tertiary amine can be metabolically cleaved to generate a bioactive primary or secondary amine. For examples, see Simplicio, *et al.*, *Molecules* 2008, 13, 519 and references therein.

15 Compounds

[0054] Disclosed compounds include those represented by formula I:



and pharmaceutically acceptable salts, stereoisomers, and N-oxides thereof, wherein

R_b is selected from the group consisting of H, halogen, hydroxyl, cyano or C₁-C₆ alkyl;

20 R₁ is H or C₁-C₆ alkyl;

R₂ is H or C₁-C₆ alkyl;

R₃ is selected from the group consisting of H, C₁-C₆ alkyl and a nitrogen protecting group;

R₄ is H or C₁-C₆ alkyl;

25 R₅ is X or -C₁-C₆ alkylene-X, wherein X is selected from the group consisting of phenyl, a 4- to 6-membered heteroaryl ring having 1, 2, or 3 heteroatoms selected from

- 15 -

O, S, or N, or a 4- to 6-membered heterocyclcyl ring having 1, 2, or 3 heteroatoms

selected from O, S, or N, and wherein R₅ is optionally substituted with ; and

R₆ is selected from the group consisting of H, halogen, hydroxyl, cyano, -O-C(O)-C₁-C₆ alkyl, C₁-C₆ alkyl, or C₁-C₆ alkoxy;

5 or, in other embodiments, the variables set forth in formula (I) are defined as follows:

R_b is selected from the group consisting of H, halogen, hydroxyl, cyano or C₁-C₆ alkyl (e.g., H);

R₁ is H or C₁-C₆ alkyl;

R₂ is H or C₁-C₆ alkyl;

10 R₃ is selected from the group consisting of H, C₁-C₆ alkyl and a nitrogen protecting group;

R₄ is H or C₁-C₆ alkyl;

R₅ is X or -C₁-C₆ alkylene-X, wherein X is selected from the group consisting of:

(i) phenyl;

15 (ii) heteroaryl including from 5 to 6 ring atoms wherein 1, 2, or 3 of the ring atoms are independently selected from the group consisting of N, NH, N(C₁-C₃ alkyl), O, and S; and

(iii) heterocyclcyl including from 3 to 6 ring atoms wherein 1, 2, or 3 of the ring atoms are independently selected from the group consisting of N, NH, N(C₁-C₃ alkyl), O, and

20 S; wherein R₅ is optionally substituted with ; and

R₆ is selected from the group consisting of H, halogen, hydroxyl, cyano, -O-C(O)-C₁-C₆ alkyl, C₁-C₆ alkyl, or C₁-C₆ alkoxy.

[0055] In certain embodiments, R₁ is H.

[0056] In certain embodiments, R₂ is H.

25 [0057] In certain embodiments, R₃ is H.

[0058] In other embodiments, R₃ is a nitrogen protecting group. In certain embodiments, R₃ has formula -C(O)OR₃₁, wherein R₃₁ is selected from the group consisting of: C₁-C₆ alkyl;



- 16 -

C₁-C₆ haloalkyl; C₂-C₆ alkenyl; C₂-C₆ alkynyl; C₃-C₁₀ cycloalkyl, wherein the C₃-C₁₀ cycloalkyl is optionally substituted with from 1-3 independently selected C₁-C₃ alkyl; -CH₂-C₃-C₁₀ cycloalkyl wherein the C₃-C₁₀ cycloalkyl is optionally substituted with from 1-3 independently selected C₁-C₃ alkyl; -CH₂-phenyl, wherein the phenyl is optionally substituted with from 1-2 substituents independently selected from C₁-C₃ alkyl; C₁-C₃ haloalkyl; C₁-C₃ alkoxy; C₁-C₃ haloalkoxy; nitro; halo; SO₂Me, cyano; and -OC(O)CH₃; and -CH₂-pyridyl. In certain embodiments, R₃₁ is C₁-C₆ alkyl (e.g., *tert*-butyl). In other embodiments, R₃ has formula -C(O)R₃₂, wherein R₃₂ is selected from the group consisting of: H; C₁-C₆ alkyl; C₁-C₆ haloalkyl; phenyl, wherein the phenyl is optionally substituted with from 1-2 substituents independently selected from C₁-C₃ alkyl; C₁-C₃ haloalkyl; C₁-C₃ alkoxy; C₁-C₃ haloalkoxy; nitro; halo; SO₂Me, cyano; and -OC(O)CH₃; and pyridyl. In certain embodiments, R₃₂ is C₁-C₆ alkyl (e.g., -CH₃ or *iso*-propyl).

[0059] In certain embodiments, R₄ is C₁-C₆ alkyl. In some embodiments, R₄ is methyl.

[0060] In certain embodiments, R_b is H.

[0061] In certain embodiments, R₅ is X, and wherein X is a 5- to 6-membered heteroaryl ring selected from the group consisting of azetidine, pyrrolidine, pyrazolidine, pyridine, pyrimidine, isooxazolidine, imidazolidine, oxazolidine, thiazolidine, and isothiazolidine. In

other embodiments, R₅ is  or . In certain other embodiments, R₅ is X, and

wherein X is optionally substituted with  (sometimes referred to herein as “-CH(R₄)(R₆)”).


[0062] In some embodiments, R₅ is X. In certain embodiments, X is heteroaryl including from 5 to 6 ring atoms wherein 1, 2, or 3 of the ring atoms are independently selected from the group consisting of N, NH, N(C₁-C₃ alkyl), O, and S. For example, X can be selected from the group consisting of 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl, pyridyl, and pyrimidinyl. In certain embodiments, R₅ (here, X as defined anywhere herein) is substituted with -CH(R₄)(R₆). In certain embodiments, R₄ is C₁-C₆ alkyl (e.g., R₄ is methyl). In certain embodiments, R₆ is selected from the group consisting of hydroxyl, -O-C(O)-C₁-C₆ alkyl, and C₁-C₆ alkoxy (e.g., R₆ is hydroxyl). Embodiments in which R₅ is X can include one or more of the following features: R₁ is H or methyl; R₂ is H or methyl; R₃ is H, -C(O)OR₃₁, or -C(O)R₃₂; when present, R₄ is C₁-C₆ alkyl (e.g., R₄ is methyl), and R₆ is selected from the group consisting of hydroxyl, -O-C(O)-C₁-C₆ alkyl, and C₁-C₆ alkoxy (e.g., R₆ is hydroxyl); R_b is H.

- 17 -


[0063] In some embodiments, R_5 is $-C_1-C_6$ alkylene-X (e.g., $-C_1-C_2$ alkylene-X or $-C_1$ alkylene-X). In certain embodiments, X is heteroaryl including from 5 to 6 ring atoms wherein 1, 2, or 3 of the ring atoms are independently selected from the group consisting of N, NH, N(C_1-C_3 alkyl), O, and S. For example, X can be selected from the group consisting of 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl, pyridyl, and pyrimidinyl. In certain embodiments, R_5 is substituted on either the X portion or the alkylene chain portion with $-\text{CH}(R_4)(R_6)$. In certain of these embodiments, R_5 is substituted on the alkylene portion with $-\text{CH}(R_4)(R_6)$, and R_5 can have, for example, the formula $-\text{CH}(\text{CHR}_4\text{R}_6)\text{-X}$. In certain embodiments, R_4 is C_1-C_6 alkyl (e.g., R_4 is methyl). In certain embodiments, R_6 is selected from the group consisting of hydroxyl, $-\text{O}-\text{C}(\text{O})-C_1-C_6$ alkyl, and C_1-C_6 alkoxy (e.g., R_6 is hydroxyl). Embodiments in which R_5 is $-C_1-C_6$ alkylene-X can include one or more of the following features: R_1 is H or methyl; R_2 is H or methyl; R_3 is H, $-\text{C}(\text{O})\text{OR}_{31}$, or $-\text{C}(\text{O})\text{R}_{32}$; when present, R_4 is C_1-C_6 alkyl (e.g., R_4 is methyl), and R_6 is selected from the group consisting of hydroxyl, $-\text{O}-\text{C}(\text{O})-C_1-C_6$ alkyl, and C_1-C_6 alkoxy (e.g., R_6 is hydroxyl); R_b is H.

[0064] In certain embodiments, R_6 is hydroxyl.

[0065] In some embodiments, a disclosed compound includes those delineated in Table 1 and/or the Examples, e.g., one having the formula:



[0066] Disclosed compounds also include compounds represented by formula II:




and pharmaceutically acceptable salts, stereoisomers, and N-oxides thereof, wherein

R_3 is selected from the group consisting of H, C_1-C_6 alkyl and a nitrogen protecting group; and

R_7 is a 4- to 6-membered heteroaryl ring having 1, 2, or 3 heteroatoms selected from O, S, or N, optionally substituted on a free carbon by a substituent selected from the group consisting of: halogen, C_1-C_6 alkyl, hydroxyl, cyano, and phenyl.

- 18 -

[0067] Also disclosed are compounds represented by formula III:



and pharmaceutically acceptable salts, stereoisomers, and N-oxides thereof, wherein

R_b is selected from the group consisting of H, halogen, hydroxyl, cyano or C₁-C₆ alkyl;


5 R₁ is H or C₁-C₆ alkyl;

R₂ is H or C₁-C₆ alkyl;

R₃ is selected from the group consisting of H, C₁-C₆ alkyl and a nitrogen protecting group;

R₄ is H or C₁-C₆ alkyl;

10 X is selected from the group consisting of phenyl, a 4- to 6-membered heteroaryl ring having 1, 2, or 3 heteroatoms selected from O, S, or N, or a 4- to 6-membered heterocyclyl ring having 1, 2, or 3 heteroatoms selected from O, S, or N, wherein



R₆ is selected from the group consisting of H, halogen, hydroxyl, cyano, -O-C(O)-C₁-C₆ alkyl, C₁-C₆ alkyl, or C₁-C₆ alkoxy.

15


[0068] In certain embodiments, R₁ is H.

[0069] In certain embodiments, R₂ is H.

[0070] In some embodiments, R₃ is H.

[0071] In certain embodiments, R₄ is C₁-C₆ alkyl. In other embodiments, R₄ is methyl.

20 [0072] In some embodiments, R_b is H.



[0073] In certain embodiments, X is

[0074] In some embodiments, R₆ is hydroxyl.

[0075] The compounds of the present disclosure and formulations thereof may have a plurality of chiral centers. Each chiral center may be independently *R*, *S*, or any mixture of *R* and *S*. For example, in some embodiments, a chiral center may have an *R*:*S* ratio of between

25

about 100:0 and about 50:50, between about 100:0 and about 75:25, between about 100:0 and about 85:15, between about 100:0 and about 90:10, between about 100:0 and about 95:5, between about 100:0 and about 98:2, between about 100:0 and about 99:1, between about 0:100 and 50:50, between about 0:100 and about 25:75, between about 0:100 and about 15:85, between about 0:100 and about 10:90, between about 0:100 and about 5:95, between about 0:100 and about 2:98, between about 0:100 and about 1:99, between about 75:25 and 25:75, and about 50:50. Formulations of the disclosed compounds comprising a greater ratio of one or more isomers (i.e., *R* and/or *S*) may possess enhanced therapeutic characteristic relative to racemic formulations of a disclosed compounds or mixture of compounds. In some instances, chemical formulas contain the descriptor “-(*R*)-“ or “-(*S*)-“ that is further attached to solid wedge or dashed wedge. This descriptor is intended to show a methine carbon (CH) that is attached to three other substituents and has either the indicated *R* or *S* configuration (see, e.g., Table 1).

[0076] Disclosed compounds may provide for efficient cation channel opening at the NMDA receptor, e.g. may bind or associate with the glutamate site of the NMDA receptor to assist in opening the cation channel. The disclosed compounds may be used to regulate (turn on or turn off) the NMDA receptor through action as an agonist.

[0077] The compounds as described herein may be glycine site NMDA receptor partial agonists. A partial agonist as used in this context will be understood to mean that at a low concentration, the analog acts as an agonist and at a high concentration, the analog acts as an antagonist. Glycine binding is not inhibited by glutamate or by competitive inhibitors of glutamate, and also does not bind at the same site as glutamate on the NMDA receptor. A second and separate binding site for glycine exists at the NMDA receptor. The ligand-gated ion channel of the NMDA receptor is, thus, under the control of at least these two distinct allosteric sites. Disclosed compounds may be capable of binding or associating with the glycine binding site of the NMDA receptor. In some embodiments, disclosed compounds may possess a potency that is 10-fold or greater than the activity of existing NMDA receptor glycine site partial agonists.

[0078] The disclosed compounds may exhibit a high therapeutic index. The therapeutic index, as used herein, refers to the ratio of the dose that produces a toxicity in 50% of the population (i.e., TD_{50}) to the minimum effective dose for 50% of the population (i.e., ED_{50}). Thus, the therapeutic index = $(TD_{50})/(ED_{50})$. In some embodiments, a disclosed compound

- 20 -

may have a therapeutic index of at least about 10:1, at least about 50:1, at least about 100:1, at least about 200:1, at least about 500:1, or at least about 1000:1.

Compositions

5 [0079] In other aspects, formulations and compositions comprising the disclosed compounds and optionally a pharmaceutically acceptable excipient are provided. In some embodiments, a contemplated formulation comprises a racemic mixture of one or more of the disclosed compounds.

[0080] Contemplated formulations may be prepared in any of a variety of forms for use. By way of example, and not limitation, the compounds may be prepared in a formulation
10 suitable for oral administration, subcutaneous injection, or other methods for administering an active agent to an animal known in the pharmaceutical arts.

[0081] Amounts of a disclosed compound as described herein in a formulation may vary according to factors such as the disease state, age, sex, and weight of the individual. Dosage regimens may be adjusted to provide the optimum therapeutic response. For example, a single
15 bolus may be administered, several divided doses may be administered over time or the dose may be proportionally reduced or increased as indicated by the exigencies of the therapeutic situation. It is especially advantageous to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the mammalian subjects to be
20 treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier.

[0082] The specification for the dosage unit forms of the invention are dictated by and directly dependent on (a) the unique characteristics of the compound selected and the particular therapeutic effect to be achieved, and (b) the limitations inherent in the art of compounding
25 such an active compound for the treatment of sensitivity in individuals.

[0083] Therapeutic compositions typically must be sterile and stable under the conditions of manufacture and storage. The composition can be formulated as a solution, microemulsion, liposome, or other ordered structure suitable to high drug concentration. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example,
30 glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. In many cases, it will be preferable to include isotonic agents, for example,

- 21 -

sugars, polyalcohols such as mannitol, sorbitol, or sodium chloride in the composition.

Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, monostearate salts and gelatin.

[0084] The compounds can be administered in a time release formulation, for example in a

5 composition which includes a slow release polymer. The compounds can be prepared with carriers that will protect the compound against rapid release, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, polylactic acid and polylactic, polyglycolic
10 copolymers (PLG). Many methods for the preparation of such formulations are generally known to those skilled in the art.

[0085] Sterile injectable solutions can be prepared by incorporating the compound in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by
15 incorporating the active compound into a sterile vehicle which contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying which yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

20 **[0086]** In accordance with an alternative aspect of the invention, a compound may be formulated with one or more additional compounds that enhance the solubility of the compound.

Methods

[0087] Methods for treating a condition in a patient in need thereof by administering a
25 therapeutically effective dose of a compound described herein are provided. In some embodiments, the condition may be a mental condition. For example, a mental illness may be treated. In another aspect, a nervous system condition may be treated. For example, a condition that affects the central nervous system, the peripheral nervous system, and/or the eye may be treated. In some embodiments, neurodegenerative diseases may be treated.

30 **[0088]** In some embodiments, the methods include administering a compound to treat patients suffering from autism, anxiety, depression, bipolar disorder, attention deficit disorder, attention deficit hyperactivity disorder (ADHD), schizophrenia, a psychotic disorder, a psychotic symptom, social withdrawal, obsessive-compulsive disorder (OCD), phobia, post-

traumatic stress syndrome, a behavior disorder, an impulse control disorder, a substance abuse disorder (e.g., a withdrawal symptom, opiate addiction, nicotine addiction, and ethanol addition), a sleep disorder, a memory disorder (e.g., a deficit, loss, or reduced ability to make new memories), a learning disorder, urinary incontinence, multiple system atrophy, progressive supra-nuclear palsy, Friedrich's ataxia, Down's syndrome, fragile X syndrome, tuberous sclerosis, olivio-ponto-cerebellar atrophy, cerebral palsy, drug-induced optic neuritis, ischemic retinopathy, diabetic retinopathy, glaucoma, dementia, AIDS dementia, Alzheimer's disease, Huntington's chorea, spasticity, myoclonus, muscle spasm, Tourette's syndrome, epilepsy, cerebral ischemia, stroke, a brain tumor, traumatic brain injury, cardiac arrest, myelopathy, spinal cord injury, peripheral neuropathy, acute neuropathic pain, and chronic neuropathic pain.

[0089] In some embodiments, methods of treating a memory disorder associated with aging, schizophrenia, special learning disorders, seizures, post-stroke convulsions, brain ischemia, hypoglycemia, cardiac arrest, epilepsy, migraine, AIDS dementia, Huntington's chorea, Parkinson's disease, early stage Alzheimer's disease, and Alzheimer's disease are contemplated.

[0090] In certain embodiments, methods for treating schizophrenia are provided. For example, paranoid type schizophrenia, disorganized type schizophrenia (i.e., hebephrenic schizophrenia), catatonic type schizophrenia, undifferentiated type schizophrenia, residual type schizophrenia, post-schizophrenic depression, and simple schizophrenia may be treated using the methods and compositions contemplated herein. Psychotic disorders such as schizoaffective disorders, delusional disorders, brief psychotic disorders, shared psychotic disorders, and psychotic disorders with delusions or hallucinations may also be treated using the compositions contemplated herein.

[0091] Paranoid schizophrenia may be characterized where delusions or auditory hallucinations are present, but thought disorder, disorganized behavior, or affective flattening are not. Delusions may be persecutory and/or grandiose, but in addition to these, other themes such as jealousy, religiosity, or somatization may also be present. Disorganized type schizophrenia may be characterized where thought disorder and flat affect are present together. Catatonic type schizophrenia may be characterized where the patient may be almost immobile or exhibit agitated, purposeless movement. Symptoms can include catatonic stupor and waxy flexibility. Undifferentiated type schizophrenia may be characterized where psychotic symptoms are present but the criteria for paranoid, disorganized, or catatonic types have not been met. Residual type schizophrenia may be characterized where positive symptoms are

present at a low intensity only. Post-schizophrenic depression may be characterized where a depressive episode arises in the aftermath of a schizophrenic illness where some low-level schizophrenic symptoms may still be present. Simple schizophrenia may be characterized by insidious and progressive development of prominent negative symptoms with no history of psychotic episodes.

[0092] In some embodiments, methods are provided for treating psychotic symptoms that may be present in other mental disorders, including, but not limited to, bipolar disorder, borderline personality disorder, drug intoxication, and drug-induced psychosis. In another embodiment, methods for treating delusions (e.g., "non-bizarre") that may be present in, for example, delusional disorder are provided.

[0093] Also provided are methods for treating social withdrawal in conditions including, but not limited to, social anxiety disorder, avoidant personality disorder, and schizotypal personality disorder.

[0094] In some embodiments, methods are provided for treating neuropathic pain. The neuropathic pain may be acute or chronic. In some cases, the neuropathic pain may be associated with a condition such as herpes, HIV, traumatic nerve injury, stroke, post-ischemia, fibromyalgia, reflex sympathetic dystrophy, complex regional pain syndrome, spinal cord injury, sciatica, phantom limb pain, diabetic neuropathy, and cancer chemotherapeutic-induced neuropathic pain. Methods for enhancing pain relief and for providing analgesia to a patient are also contemplated.

[0095] Further contemplated methods include a method of treating autism and/or an autism spectrum disorder in a patient need thereof, comprising administering an effective amount of a compound to the patient. In an embodiment, a method for reducing the symptoms of autism in a patient in need thereof is contemplated, comprising administering an effective amount of a disclosed compound to the patient. For example, upon administration, the compound may decrease the incidence of one or more symptoms of autism such as eye contact avoidance, failure to socialize, attention deficit, poor mood, hyperactivity, abnormal sound sensitivity, inappropriate speech, disrupted sleep, and perseveration. Such decreased incidence may be measured relative to the incidence in the untreated individual or an untreated individual(s).

[0096] Also provided herein is a method of modulating an autism target gene expression in a cell comprising contacting a cell with an effective amount of a compound described herein. The autism gene expression may be for example, selected from ABAT, APOE, CHRNA4, GABRA5, GFAP, GRIN2A, PDYN, and PENK. In another embodiment, a method of

- 24 -

modulating synaptic plasticity in a patient suffering from a synaptic plasticity related disorder is provided, comprising administering to the patient an effective amount of a compound.

[0097] In another embodiment, a method of treating Alzheimer's disease, or e.g., treatment of memory loss that e.g., accompanies early stage Alzheimer's disease, in a patient in need thereof is provided, comprising administering a compound. Also provided herein is a method of modulating an Alzheimer's amyloid protein (e.g., beta amyloid peptide, e.g. the isoform A β ₁₋₄₂), in-vitro or in-vivo (e.g. in a cell) comprising contacting the protein with an effective amount of a compound is disclosed. For example, in some embodiments, a compound may block the ability of such amyloid protein to inhibit long-term potentiation in hippocampal slices as well as apoptotic neuronal cell death. In some embodiments, a disclosed compound may provide neuroprotective properties to a Alzheimer's patient in need thereof, for example, may provide a therapeutic effect on later stage Alzheimer's -associated neuronal cell death.

[0098] In a further embodiment, a method of treating depression comprising administering a compound described herein is provided. In some embodiments, the treatment may relieve depression or a symptom of depression without affecting behavior or motor coordination and without inducing or promoting seizure activity. Exemplary depression conditions that are expected to be treated according to this aspect of the invention include, but are not limited to, major depressive disorder, dysthymic disorder, psychotic depression, postpartum depression, premenstrual syndrome, premenstrual dysphoric disorder, seasonal affective disorder (SAD), bipolar disorder (or manic depressive disorder), mood disorder, and depressions caused by chronic medical conditions such as cancer or chronic pain, chemotherapy, chronic stress, and post traumatic stress disorders. In addition, patients suffering from any form of depression often experience anxiety. Various symptoms associated with anxiety include fear, panic, heart palpitations, shortness of breath, fatigue, nausea, and headaches among others. Anxiety or any of the symptoms thereof may be treated by administering a compound as described herein.

[0099] Also provided herein are methods of treating a condition in treatment-resistant patients, e.g., patients suffering from a mental or central nervous system condition that does not, and/or has not, responded to adequate courses of at least one, or at least two, other compounds or therapeutics. For example, provided herein is a method of treating depression in a treatment resistant patient, comprising a) optionally identifying the patient as treatment resistant and b) administering an effective dose of a compound to said patient.

- 25 -

[00100] In some embodiments, a compound described herein may be used for acute care of a patient. For example, a compound may be administered to a patient to treat a particular episode (e.g., a severe episode) of a condition contemplated herein.


[00101] Also contemplated herein are combination therapies comprising a compound in
 5 combination with one or more other active agents. For example, a compound may be combined with one or more antidepressants, such as tricyclic antidepressants, MAO-I's, SSRI's, and double and triple uptake inhibitors and/or anxiolytic drugs. Exemplary drugs that may be used in combination with a compound include Anafranil, Adapin, Aventyl, Elavil, Norpramin, Pamelor, Pertofrane, Sinequan, Surmontil, Tofranil, Vivactil, Parnate, Nardil, Marplan, Celexa,
 10 Lexapro, Luvox, Paxil, Prozac, Zoloft, Wellbutrin, Effexor, Remeron, Cymbalta, Desyrel (trazodone), and Ludiomill. In another example, a compound may be combined with an antipsychotic medication. Non-limiting examples of antipsychotics include butyrophenones, phenothiazines, thioxanthenes, clozapine, olanzapine, risperidone, quetiapine, ziprasidone, amisulpride, asenapine, paliperidone, iloperidone, zotepine, sertindole, lurasidone, and
 15 aripiprazole. It should be understood that combinations of a compound and one or more of the above therapeutics may be used for treatment of any suitable condition and are not limited to use as antidepressants or antipsychotics.

EXAMPLES






[00102] The following examples are provided for illustrative purposes only, and are not intended to limit the scope of the disclosure.

20 **[00103]** Table 1 below shows some exemplary compounds of the disclosure and provides physiochemical characteristics of the compounds.





Table 1.





Compound	Structure	Molecular Weight (Da)	cLogP	tPSA
Compound X		252	-1.14	91.5

- 26 -






Compound Y		262	-0.57	78.35
Compound Z		248	-0.15	78.35
C-16		252.2697	-1.1382	91.49
C-4		262.3076	-0.57075	78.35
C-15		352.3856	0.144989	109


- 27 -

C-20		276.3342	- 0.160256	78.35
C-36		276.3342	- 0.160256	78.35
C-33		290.3608	0.250239	78.35
C-10		252.2697	-1.83479	91.49

C-9		352.3856	- 0.551597	109
C-3		362.4234	0.672806	95.86
C-23		366.4121	- 0.135022	109
C-19		376.45	1.0833	95.86


- 29 -

C-26		366.4121	- 0.135022	109
C-29		376.45	1.0833	95.86
C-32		390.4766	1.4938	95.86
C-11		322.3596	- 0.981326	99.77
C-5		332.3975	0.335914	86.63

C-37		346.424	0.746408	86.63
------	---	---------	----------	-------

Example 1 – Synthesis of Compound X

Scheme 1.



5

Synthesis of 2-((tert-butoxycarbonyl) amino)-3-hydroxybutanoic acid (A)

[00104] To a stirring solution of 2-amino-3-hydroxybutanoic acid (SM2) (10 g, 83.9 mmol) in 1,4-dioxane/water (100 mL, 1: 1)) was added NaHCO₃ (21.1 g, 0.25 mol) followed by Boc-anhydride (21.9 mL, 0.101 mol) at 0°C. The reaction mixture was stirred at RT for 16 h. After consumption of the starting material (by TLC), the reaction mixture was diluted with water and washed with EtOAc. The aqueous layer was acidified using citric acid solution (pH~3-4) and then extracted with CH₂Cl₂ (2 x 150 mL). The separated organic extracts were dried over

- 31 -

anhydrous Na₂SO₄, filtered and concentrated under vacuum to afford **A** (15 g, crude). This material was directly used for the next step without further purification.

Synthesis of 3-(*tert*-butoxycarbonyl)-2, 2, 5-trimethyloxazolidine-4-carboxylic acid (B)

[00105] To a stirring solution of **A** (15 g, 59.28 mmol) in THF (150 mL) was added PPTS (1.47 g, 5.92 mmol) followed by 2,2-dimethoxy propane (21.79 mL, 0.17 mol) at 0°C under N₂ atmosphere. The reaction mixture was stirred at RT for 16 h. The reaction mixture was again heated to reflux for 6 h. The reaction mixture was diluted with aqueous NaHCO₃ solution and washed with EtOAc. The aqueous layer was acidified using citric acid solution (pH~2) and extracted with CH₂Cl₂ (2x 100 mL). The organic layer was washed with brine, dried over anhydrous Na₂SO₄ and concentrated under vacuum to afford **B** (18 g, crude).

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 13.25 (br s, 1H), 4.11-4.05 (m, 1H), 3.79 (d, 1H), 1.50 (s, 3H), 1.67 (s, 3H), 1.45 (s, 9H), 1.29 (d, 3H).

Synthesis of *tert*-butyl 4-carbamoyl-2, 2, 5-trimethyloxazolidine-3-carboxylate (C)

[00106] To a stirring solution of **B** (18 g, 69.4 mmol) in CH₂Cl₂ (180 mL) was added HOBt (14.16 g, 0.104 mol), EDCL.HCl (19.88 g, 0.104 mol) followed by NH₄Cl (5.56 g, 0.104 mol) and DIPEA (31.9 mL, 0.173 mol) at 0°C. The reaction mixture was stirred at RT for 16 h. After consumption of the starting material (by TLC), the reaction mixture was washed with aqueous citric acid, NaHCO₃ followed by brine. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The resulting crude material was purified by silica gel column chromatography eluting with 2% MeOH/CH₂Cl₂ to afford **C** (13 g, 72.5%).

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 7.51 (br s, 1H), 7.14 (br s, 1H), 3.97-3.95 (m, 1H), 3.71 (d, 1H), 1.51 (d, 6H), 1.34 (s, 9H), 1.24 (d, 3H).

LCMS (ESI): 159.1[(M⁺+1)-Boc]

Synthesis of (Z)-*tert*-butyl 4-(((dimethylamino) methylene) carbamoyl)-2, 2, 5-trimethyloxazolidine-3-carboxylate (D)

[00107] A solution of **C** (13 g, 50.3 mmol) in DMF.DMA (130 mL) was stirred at reflux temperature for 3 h under N₂ atmosphere. After consumption of the starting material (by TLC), the reaction mixture was concentrated under reduced pressure to afford **D** (15.7 g, crude). This crude material was directly taken for the next step without further purification.

Synthesis of *tert*-butyl 2, 2, 5-trimethyl-4-(1, 2, 4-oxadiazol-5-yl) oxazolidine-3-carboxylate (E)

- 32 -

[00108] To a stirring solution of **D** (15.7 g, 50.09 mmol) in ethanol (157 mL) was added hydroxylamine hydrochloride (6.96 g, 0.10 mol) under N₂ atmosphere. The reaction mixture was heated to reflux and stirred for 2 h. After consumption of the starting material (by TLC), acetic acid (28.6 mL, 0.50 mol) was added to the reaction mixture and then refluxed for 16 h.

5 Solvents from the reaction mixture were evaporated under vacuum. The resulting crude material was purified by silica gel column chromatography eluting with 10% EtOAc/Hexane to afford **E** (4.5 g, 32%).

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 6.35 (s, 2H), 4.61 (d, 1H), 4.22-4.15 (m, 1H), 1.55 (s, 6H), 1.37 (s, 2H), 1.25 (d, 3H), 1.21 (s, 6H).

10 LCMS (ESI): 284 [M⁺+1]

Mass (m/z): 283 [M⁺]

Synthesis of 1-amino-1-(1, 2, 4-oxadiazol-5-yl) propan-2-ol (Int-F)

[00109] To a stirring solution of **E** (5 g, 17.6 mmol) in water (25 mL) was added trifluoroacetic acid (25 mL). The reaction mixture was stirred at RT for 5 h. After consumption
15 of the starting material (by TLC), the reaction mixture was concentrated under vacuum. The residue was dissolved in water and neutralized with aqueous NaHCO₃. The solvents from the reaction mixture were evaporated under vacuum and extracted with 5% MeOH/CH₂Cl₂ (4x 150 mL). The organic layer was concentrated under reduced pressure to afford **Int-F** (2.5 g, crude).

¹H-NMR: (400 MHz, D₂O): δ 8.84 (s, 1H), 4.05 (d, 1H), 3.98-3.95 (m, 1H), 3.67 (s, 1H), 3.58
20 (d, 1H), 1.15 (d, 3H), 1.12 (d, 3H).

LCMS (ESI): 144.1 [M⁺+1]

Synthesis of methyl pyrrolidine-2-carboxylate (1)

[00110] To a stirring solution of pyrrolidine-2-carboxylic acid (**SM1**) (100 g, 0.87 mol) in methanol (800 mL) was added thionyl chloride (76.9 mL, 1.04 mol) slowly drop wise at 0°C.

25 The reaction mixture was heated to reflux for 12 h. After consumption of the starting material (by TLC), the reaction was concentrated under vacuum. Obtained residue was washed with *n*-Hexane and distilled off the solvent to afford **1** (143.9 g, HCl salt).

¹H-NMR: (400 MHz, CDCl₃) (Rotamers): δ 3.89 (s, 3H), 3.68-3.62 (m, 2H), 3.59-3.47 (m, 2H), 2.49-2.37 (m, 1H), 2.27-2.05 (m, 3H).

30 LCMS (ESI): 166 [M⁺+1]

Synthesis of 1-tert-butyl 2-methyl pyrrolidine-1,2-dicarboxylate (2)

- 33 -

[00111] To a stirring solution of **1** (35 g, 0.22 mol) in CH₂Cl₂ (175 mL) were added Et₃N (90 mL, 0.65 mol) followed by Boc-anhydride (56.9 mL, 0.26 mol) at 0°C. The reaction mixture was stirred at RT for 16 h. After consumption of the starting material (by TLC), the reaction was diluted with water (100 mL) and extracted with CH₂Cl₂ (2x 100 mL). The organic layer was washed with water, brine, dried over Na₂SO₄ and concentrated. Obtained crude material was purified by silica gel column chromatography eluting with 30% EtOAc/Hexane to afford **2** (41 g, 95%).

¹H-NMR: (400 MHz, CDCl₃) (Rotamers): δ 4.25-4.21 (m, 1H), 3.75 (s, 3H), 3.57-3.26 (m, 2H), 2.29-2.10 (m, 1H), 1.99-1.75 (m, 3H), 1.45 (s, 9H).

LCMS (ESI): 130 [(M⁺+1)-Boc]

Synthesis of 1-tert-butyl 2-methyl 2-((benzyloxy) methyl) pyrrolidine-1, 2-dicarboxylate (3)

[00112] To a stirring solution of **2** (100 g, 0.43 mol) in THF (800 mL) was added LiHMDS (873 mL, 0.87 mol) at -78°C and stirred for 1 h. To this BOM-chloride (93.2 mL, 0.65 mol) was added drop wise at -78°C and stirred for 2 h at -20 °C. After consumption of the starting material (by TLC), the reaction was quenched with NH₄Cl at 0 °C. The separated organic layer was washed with water, dried over Na₂SO₄ and concentrated to afford **3** (180 g, crude). This material was directly taken for the next step without further purification.

LCMS (ESI): 250 [(M⁺+1)-Boc]

Synthesis of 1-tert-butyl 2-methyl 2-(hydroxymethyl)pyrrolidine-1, 2-dicarboxylate (4)

[00113] To a stirring solution of **3** (74 g, 0.21 mol) in methanol (740 mL) was added 10% Pd/C (50% wet, 14.8 g) under N₂ atmosphere and stirred for 6 h under H₂ atmosphere (balloon pressure). The reaction mixture was filtered through celite pad and concentrated under reduced pressure to afford **4** (45 g, 82%) as crude.

Synthesis of 1-tert-butyl 2-methyl 2-formylpyrrolidine-1, 2-dicarboxylate (5)

[00114] To a stirring solution of **4** (10 g, 38.5 mmol) in CH₂Cl₂ (100 mL) was added Dess-martin periodinane (19.6 g, 46.27 mmol) at 0°C under N₂ atmosphere and stirred for 3 h. After consumption of the starting material (determined by TLC), the reaction was quenched with aqueous NaHCO₃ solution and extracted with CH₂Cl₂ (2x 100 mL). The organic layer was dried over anhydrous Na₂SO₄ and concentrated under vacuum. The resulting crude material

- 34 -

was purified by column chromatography eluting with 10% EtOAc/Hexane to afford **5** (7 g, 70.5%).

Synthesis of 1-*tert*-butyl 2-methyl 2-((((1*S*, 2*R*)-2-hydroxy-1-(1, 2, 4-oxadiazol-5-yl) propyl) amino) methyl) pyrrolidine-1, 2-dicarboxylate (6**)**

- 5 **[00115]** To a stirring solution of **5** (3 g, 11.6 mmol) in MeOH (30 mL) was added sodium acetate (1.91 g, 23.3 mmol) followed by **Int-F** (3.6 g, 13.9 mmol). The reaction mixture was heated to reflux for 1 h. The reaction mixture was slowly cooled to RT-0°C, to this sodium cyanoborohydride (1.465 g, 23.3 mmol) was added, and stirring was continued for another 6 h at RT. After consumption of the starting material (determined by TLC), methanol from the
- 10 reaction was evaporated under reduced pressure and the residue was diluted with water and extracted with EtOAc (2x 50 mL). The separated organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The resulting crude material was purified by silica gel column chromatography eluting with 40% EtOAc/Hexane to afford **6** (2.5 g, 56%).
- LCMS *m/z*: 385 [M⁺+1]

15 **Synthesis of *tert*-butyl 2-((1*S*, 2*R*)-2-hydroxy-1-(1, 2, 4-oxadiazol-5-yl) propyl)-1-oxo-2, 5-diazaspiro[3.4] octane-5-carboxylate (**7**)**

- [00116]** To a stirring solution of **6** (1.5 g, 3.90 mmol) in THF (30 mL) was cooled to 0 °C and added *t*-BuMgCl (1M in THF, 15.6 mL, 15.6 mmol) and stirred for 15 min. After consumption of the starting material (by TLC), the reaction mixture was quenched with
- 20 aqueous NH₄Cl solution and diluted with water. Aqueous layer was extracted with EtOAc (2x 25 mL). The separated organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The resulting crude material was purified by silica gel column chromatography eluting with 150% EtOAc/CH₂Cl₂ to afford **7** (0.15 g, 11%).
- ¹H-NMR: (400 MHz, DMSO-d₆): δ 9.02 (s, 1H), 5.15 (s, 1H), 4.37-4.32 (m, 1H), 3.95 (d, 1H), 3.66-3.60 (m, 1H), 3.36-3.30 (m, 1H), 2.29-2.09 (m, 2H), 1.87-1.82 (m, 2H), 1.55 (s, 9H), 1.27 (d, 3H).

LCMS (ESI) *m/z*: 351 [M⁺-1]

UPLC Purity: 96%

30 **Synthesis of 2-((1*S*, 2*R*)-2-hydroxy-1-(1, 2, 4-oxadiazol-5-yl) propyl)-2, 5-diazaspiro [3.4] octan-1-one (Compound X)**

[00117] To a stirring solution of **7** (0.4 g, 1.13 mmol) in CH₂Cl₂ (4 mL) was added TFA (0.43 mL) at 0°C and stirred at RT for 30 min. The reaction mixture was concentrated under vacuum. The resulting crude material was purified by prep-HPLC to afford **Compound X** (65 mg) as TFA salt.


- 5 ¹H-NMR: (400 MHz, DMSO-d₆): δ 9.89 (br s, 1H), 9.08 (s, 1H), 5.46 (d, 1H), 5.31 (s, 1H), 4.37-4.35 (m, 1H), 3.99 (d, 1H), 3.81 (d, 1H), 3.42-3.35 (m, 2H), 2.35-2.18 (m, 2H), 2.10-2.03 (m, 2H), 1.24 (d, 3H).

LCMS (ESI) *m/z*: 253.4 [M⁺+1]

Prep-HPLC Purity: 95%

10 Example 2 - Synthesis of Compound Y

Scheme 2.



Synthesis of (S)-methyl pyrrolidine-2-carboxylate (A)

- 36 -

[00118] To a stirring solution of L-proline (**SM1**) (100 g, 0.87 mol) in methanol (800 mL) was slowly added thionyl chloride (76.9 mL, 1.04 mol) at 0°C. The reaction mixture was slowly warmed to RT and heated to reflux for 12 h. After consumption of the starting material (by TLC), the reaction was concentrated under reduced pressure to afford **A** (143.9 g, HCl salt).

5 ¹H-NMR: (400 MHz, CDCl₃): δ 3.89 (s, 3H), 3.68-3.62 (m, 2H), 3.59-3.47 (m, 2H), 2.49-2.37 (m, 1H), 2.27-2.05 (m, 3H).

LCMS (m/z): 166 [M⁺+1]

Synthesis of (S)-1-tert-butyl 2-methyl pyrrolidine-1,2-dicarboxylate (B)

[00119] To a stirring solution of **A** (35 g, 0.22 mol) in CH₂Cl₂ (175 mL) were added Et₃N (90 mL, 0.65 mol) followed by Boc-anhydride (56.9 mL, 0.26 mol) at 0°C. The reaction mixture was stirred at RT for 16 h. After consumption of the starting material (by TLC), the reaction was diluted with water (100 mL) and extracted with CH₂Cl₂ (2x 100 mL). The organic layer was washed with water, brine, dried over Na₂SO₄ and concentrated. The resulting crude material was purified by silica gel column chromatography eluting with 30% EtOAc/Hexane to

10 afford **B** (41 g, 95%).

¹H-NMR: (400 MHz, CDCl₃): δ 4.25-4.21 (m, 1H), 3.75 (s, 3H), 3.57-3.26 (m, 2H), 2.29-2.10 (m, 1H), 1.99-1.75 (m, 3H), 1.45 (s, 9H).

LCMS (m/z): 130 [(M⁺+1)-Boc]

20 **Synthesis of 1-tert-butyl 2-methyl 2-((benzyloxy) methyl) pyrrolidine-1, 2-dicarboxylate (C)**

[00120] To a stirring solution of **B** (100 g, 0.43 mol) in THF (800 mL) was added LiHMDS (873 mL, 0.87 mol) at -78°C and stirred for 1 h. To this BOM-chloride (93.2 mL, 0.65 mol) was added drop wise at -78°C and stirred for 2 h at -20°C. After consumption of the starting material (by TLC), the reaction was quenched with sat.NH₄Cl solution at 0°C. The separated

25 organic layer was washed with water, dried over Na₂SO₄ and concentrated to afford **C** (180 g, crude). This material was directly taken for the next step without further purification.

LCMS (m/z): 250 [(M⁺+1)-Boc]

Synthesis of 2-((benzyloxy) methyl)-1-(tert-butoxycarbonyl) pyrrolidine-2-carboxylic acid (D)

30 [00121] To a stirring solution of **C** (100 g, 0.28 mol) in methanol (200 mL) was added 2N NaOH solution (300 mL) at RT. The reaction mixture was heated to reflux for 4 h. After

- 37 -

consumption of the starting material (by TLC), the solvent from the reaction was evaporated under reduced pressure and diluted with EtOAc (100 mL). The aqueous layer was acidified using citric acid solution and extracted with CH₂Cl₂ (2x 250 mL). The separated organic layer was washed with water, dried over Na₂SO₄ and concentrated to afford **D** (60 g, 63%).

- 5 ¹H-NMR: (400 MHz, CDCl₃): δ 7.37-7.32 (m, 5H), 4.61 (s, 2H), 4.05-3.88 (m, 2H), 3.65-3.42 (m, 2H), 2.54-2.46 (m, 2H), 1.95 (br s, 2H), 1.57 (s, 9H).

LCMS (m/z): 334 [M⁺-1]

Synthesis of 1-(tert-butoxycarbonyl)-2-(hydroxymethyl)pyrrolidine-2-carboxylic acid (Int-E)

- 10 **[00122]** To a stirring solution of **D** (10 g, 29.81 mmol) in methanol (300 mL) was added 50% wet 10% Pd/C (5 g) at RT and stirred for 24 h under H₂ atmosphere (balloon pressure). After consumption of the starting material (by TLC), the reaction mixture was filtered through a pad of celite and the pad was washed with methanol. The filtrate was concentrated under reduced pressure to afford **Int-E** (6 g, 82%).
- 15 ¹H-NMR: (400 MHz, DMSO-d₆): δ 12.55 (br m, 1H), 3.99 (d, 1H), 3.88 (d, 1H), 7.65-7.60 (m, 1H), 3.51-3.45 (m, 1H), 3.39-3.34 (m, 1H), 2.32-2.14 (m, 1H), 1.98-1.69 (m, 3H), 1.39 (s, 9H).

Synthesis of 1-(pyrimidin-2-yl)propan-1-one (2)

- [00123]** To a stirring solution of **1** (15 g, 0.14 mol) in THF (150 mL) was added ethylmagnesium bromide (1M in THF) (171.4 mL, 0.17 mol) at 0°C under nitrogen atmosphere
- 20 slowly over a period of 15 min. After being stirred for 30 min, the reaction mixture was quenched with aqueous NH₄Cl solution and the aqueous layer was extracted with EtOAc. The separated organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to obtain crude product, which was purified by silica gel column chromatography eluting with 50% EtOAc/hexane to afford **2** (8.5 g, 44%).
- 25 ¹H-NMR: (400 MHz, CDCl₃): δ 8.95 (d, 2H), 7.47 (t, 1H), 3.26 (q, 2H), 1.25 (t, 3H).

Synthesis of (Z)-2-(1-((tert-butyldimethylsilyl)oxy)prop-1-en-1-yl)pyrimidine (3)

- [00124]** To a stirring solution of **2** (8.5 g, 62.5 mmol) in THF (170 mL) was added LiHMDS (1M in THF) (93.7 mL, 93.7 mmol) and stirred for 1 h. To this was added TES-Cl (12.5 mL, 75 mmol) at 0°C under N₂ atmosphere and further stirred for 1 h. After consumption of the starting
- 30 material (by TLC), the reaction was quenched with aqueous NH₄Cl solution and extracted with EtOAc. The separated organic layer was dried over anhydrous Na₂SO₄, filtered and

concentrated under reduced pressure to afford **3** (10 g, 64%). The resulting crude material was directly used for the next step without further purification.

Synthesis of 2-bromo-1-(pyrimidin-2-yl) propan-1-one (4)

[00125] To a stirring solution of **3** (10 g, 0.04 mol) in THF: H₂O (100 mL, 4:1) was added NBS (5.6 g, 0.05 mol) at RT and stirred for 1 h. After consumption of the starting material (by TLC), the reaction mixture was concentrated under reduced pressure and obtained residue was diluted with EtOAc and washed with water. The separated organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to obtain crude product, which was purified by silica gel column chromatography eluting with 30% EtOAc/hexane to afford **4** (6.5 g, 75.5%).

¹H-NMR: (400 MHz, CDCl₃): δ 8.97 (d, 2H), 7.52 (t, 1H), 5.92 (q, 1H), 2.79 (s, 1H), 1.99 (d, 3H).

Synthesis of 2-hydroxy-1-(pyrimidin-2-yl) propan-1-one (5)

[00126] To a stirring solution of **4** (6.5 g, 0.03 mol) in MeOH (65 mL) was added sodium formate (10.28 g, 0.15 mol) and heated to reflux for 8 h. After consumption of the starting material (by TLC), the reaction mixture was cooled to RT and solvent was removed under reduced pressure. The resulting crude material was purified by silica gel column chromatography eluting with 2% MeOH/CH₂Cl₂ to afford **5** (3.5 g, 76%).

¹H-NMR: (400 MHz, CDCl₃): δ 9.01 (d, 2H), 7.52 (t, 1H), 5.92 (q, 1H), 3.73 (br s, 1H), 3.41 (s, 1H), 2.78 (s, 2H), 1.55 (d, 3H).

Synthesis of 2-((tert-butyldimethylsilyl)oxy)-1-(pyrimidin-2-yl)propan-1-one (6)

[00127] To a stirring solution of **5** (3.5 g, 23 mmol) in CH₂Cl₂ (70 mL) was added imidazole (3.91 g, 57 mmol) followed by DMAP (281 mg, 2.30 mmol) at RT. The reaction mixture was cooled to 0°C, to which TBS-Cl (5.18 g, 0.03 mol) was added and stirred at RT for 4 h. After consumption of the starting material (by TLC), the reaction mixture was diluted with CH₂Cl₂ and washed with water. The separated organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to obtain a crude product, which was purified by silica gel column chromatography eluting with 20% EtOAc/hexane to afford **6** (3 g, 49%).

¹H-NMR: (400 MHz, CDCl₃): δ 8.93 (d, 2H), 7.45 (t, 1H), 5.59 (q, 1H), 1.52 (d, 3H), 0.89 (s, 9H), 0.09 (s, 6H).

Synthesis of 2-((tert-butyldimethylsilyl)oxy)-1-(pyrimidin-2-yl)propan-1-amine (7)

[00128] To a stirring solution of **6** (3.0 g, 11.27 mmol) in MeOH (60 mL) was added sodium acetate (1.84 g, 22.55 mmol) followed by ammonium carbonate (8.85 g, 56.3 mmol) and acetic acid (0.64 mL, 11.27 mmol) at RT. The reaction mixture was heated to reflux for 1 h, added
5 NaCNBH₃ (1.41 g, 22.5 mmol) and continued reflux for another 6 h. After consumption of the starting material (by TLC), the reaction mixture was cooled to RT and volatiles were evaporated. Obtained residue was diluted with water and extracted with EtOAc. The separated organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to get crude product, which was purified by silica gel column chromatography eluting
10 with 2% MeOH/CH₂Cl₂ to afford **7** (2 g, 66.4%).

¹H-NMR: (400 MHz, DMSO-d₆): δ 8.91 (d, 1H), 8.85 (d, 1H), 7.48 (t, 1H), 5.71 (br m, 2H), 4.24 (t, 1H), 4.05 (d, 1H), 1.29 (d, 1H), 1.12 (d, 2H), 0.74 (s, 9H), 0.03 (s, 3H), 0.02 (s, 2H), 0.01 (s, 1H).

LCMS (m/z): 268 [M⁺+1]

15 **Synthesis of tert-butyl 2-((2-((tert-butyldimethylsilyl)oxy)-1-(pyrimidin-2-yl)propyl)carbamoyl)-2-(hydroxymethyl)pyrrolidine-1-carboxylate (8)**

[00129] To a stirring solution of **Int-E** (2 g, 8.16 mmol) in CH₂Cl₂ (60 mL) was added compound **7** (2.39 g, 8.97 mmol), EDCI. HCl (2.33 g, 12.2 mmol) followed by HOBt (1.66 g, 12.24 mmol) and DIPEA (4.5 mL, 24.4 mmol) at 0°C. The reaction mixture was warmed to RT
20 and stirred for 16 h. After consumption of the starting material (by TLC), the reaction mixture was diluted with CH₂Cl₂. The separated organic layer was washed with aqueous NaHCO₃ solution followed by aqueous NH₄Cl. The organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to get crude product, which was purified by silica gel column chromatography eluting with 70% EtOAc/hexane to afford **8** (2.3 g, 57.5%).

25 ¹H-NMR: (400 MHz, DMSO-d₆): δ 8.91 (d, 1H), 8.68 (d, 2H), 7.41 (br s, 1H), 5.74 (br t, 1H), 5.07-4.89 (m, 1H), 4.15-4.10 (m, 1H), 3.97-3.92 (m, 1H), 3.45-3.41 (m, 1H), 1.79-1.74 (m, 2H), 1.43-1.39 (m, 4H), 1.29-1.21 (m, 6H), 1.12 (d, 5H), 0.71 (s, 9H), 0.12 (t, 1H), 0.09 (s, 2H), 0.08 (s, 1H), 0.04 (s, 2H).

LCMS (m/z): 495.5 [M⁺+1]

30 **Mass:** 495.5 [M⁺+1]

Synthesis of *tert*-butyl 2-(2-((*tert*-butyldimethylsilyl)oxy)-1-(pyrimidin-2-yl)propyl)-1-oxo-2,5-diazaspiro[3.4]octane-5-carboxylate (9)

[00130] To a stirring solution of **8** (2.3 g, 4.65 mmol) in THF (23 mL) was added TPP (1.34 g, 5.12 mmol) followed by DTAD (1.6 g, 6.98 mmol) at 0°C. The reaction mixture was warmed to RT and stirred for 16 h. After consumption of the starting material (by TLC), the reaction mixture was concentrated under reduced pressure to get crude product, which was purified by silica gel column chromatography eluting with 25% EtOAc/hexane to afford **9** (1.2 g, 54.2%).

LCMS (m/z): 477.4 [M⁺+1]

¹H-NMR: (400 MHz, DMSO-d₆): δ 8.82 (d, 2H), 7.49 (t, 1H), 4.72 (d, 1H), 4.31 (q, 1H), 3.62 (br s, 2H), 3.25-3.19 (m, 1H), 2.24-2.05 (m, 2H), 1.85-1.81 (m, 2H), 1.42 (br s, 1H), 1.25 (t, 3H), 0.92 (s, 8H), 0.75 (s, 9H), 0.02 (s, 3H).

Synthesis of *tert*-butyl 2-(2-hydroxy-1-(pyrimidin-2-yl)propyl)-1-oxo-2,5-diazaspiro[3.4]octane-5-carboxylate (10)

[00131] To a stirring solution of **9** (1.0 g, 2.10 mmol) in THF (20 mL) was added TBAF (1M in THF) (6.3 mL, 6.30 mmol) at 0°C under N₂ atmosphere and stirred for 1 h. After consumption of the starting material (by TLC), the reaction mixture was quenched with ice water and extracted with EtOAc. The separated organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to obtain crude product, which was purified by silica gel column chromatography eluting with 5% MeOH/CH₂Cl₂ to afford **10** (0.35 g, 46%).

¹H-NMR: (400 MHz, DMSO-d₆): δ 8.81 (d, 2H), 7.49 (t, 1H), 4.81 (d, 1H), 4.65 (d, 1H), 4.25-4.20 (m, 1H), 3.64-3.51 (m, 2H), 3.34 (s, 1H), 3.25-3.20 (m, 1H), 2.25-2.20 (m, 2H), 1.87-1.82 (m, 2H), 1.19 (d, 3H), 0.97 (s, 9H).

LCMS (m/z): 363.3 [M⁺+1]

Synthesis of 2-(2-hydroxy-1-(pyrimidin-2-yl)propyl)-2,5-diazaspiro[3.4]octan-1-one (Compound Y)

[00132] To a stirring solution of **10** (0.3 g, 0.82 mmol) in CH₂Cl₂ (6 mL) was added molecular sieves (0.3 g) followed by BF₃-etherate (0.31 mL, 2.48 mmol) at 0 °C and stirred at RT for 1 h. After consumption of the starting material (by TLC), the reaction mixture was filtered and obtained residue was dissolved in MeOH and washed with CH₂Cl₂. The volatiles

- 41 -


were evaporated under reduced pressure to obtain crude product, which was purified by silica gel column chromatography eluting with 8% MeOH/CH₂Cl₂ to afford **Compound Y** (0.12 g, 55%).

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 8.81 (d, 2H), 7.49 (t, 1H), 4.81 (d, 1H), 4.65 (d, 1H), 4.25-4.20 (m, 1H), 3.64-3.51 (m, 2H), 3.12-3.01 (m, 2H), 2.15-2.10 (m, 2H), 1.87-1.82 (m, 2H), 1.19 (d, 3H).

LCMS (m/z): 263.1 [M⁺+1]

Preparation of key intermediates, schemes I-1 to I-7:

Scheme I-1



Synthesis of 1-(pyrimidin-2-yl) propan-1-one, A:

[00133] Referring to Scheme I-1, to a stirring solution of pyrimidine-2-carbonitrile (25 g, 238 mmol) in THF (180 mL) was added ethyl magnesium bromide (1M in THF, 37.9 g, 285 mmol) at 0 °C and stirred for 2 h at RT. After completion of starting material (by TLC), the reaction mixture was diluted with saturated ammonium chloride solution and EtOAc (150 mL). The separated organic layer was washed with brine solution (2 x 100 mL). The extracted organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to afford crude material which was purified by column chromatography eluting 30% EtOAc/hexane to afford **A** (18.3 g, 57%) as an off-white solid.

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 9.00 (d, *J* = 5.2 Hz, 2H), 7.70 (t, *J* = 4.8 Hz, 1H), 3.20-3.15 (m, 2H), 1.09 (t, *J* = 7.2 Hz, 3H).

LCMS m/z: 137 [M⁺+1].

Synthesis of (Z)-2-(1-((triethylsilyl) oxy) prop-1-en-1-yl) pyrimidine (B):

- 42 -

[00134] To a stirring solution of **A** (14 g, 103 mmol) in dry THF (140 mL) was added LiHMDS (1M in THF, 206 mL, 206 mmol) slowly at 0 °C and stirred for 30 min. After added chloro triethylsilane (24.8 g, 165 mmol) in THF (50 mL) dropwise at 0 °C and stirred 1 h. After completion of starting material (by TLC), the reaction mixture was diluted with saturated ammonium chloride solution (100 mL) and EtOAc (150 mL). The separated organic layer was washed with brine solution (2 x 100 mL). The organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to afford crude material which was purified by column chromatography eluting 20% EtOAc/hexane to afford **B** (23 g, 89.4%) as yellow thick syrup.

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 8.75 (d, *J* = 4.8 Hz, 2H), 7.32 (t, *J* = 4.8 Hz, 1H), 6.36-6.31 (m, 1H), 1.77 (d, *J* = 7.2 Hz, 3H), 0.95-0.87 (m, 9H), 0.71-0.65 (m, 6H).

Synthesis of 2-bromo-1-(pyrimidin-2-yl) propan-1-one (C):

[00135] To a stirring solution of **B** (23 g, 92 mmol) in THF/H₂O (184 mL/46 mL) were added N-bromosuccinamide (18 g, 101 mmol) slowly at 0 °C and stirred for 3 h at RT. After completion of starting material (by TLC), the reaction mixture was diluted with H₂O and EtOAc (100 ml/150 mL). The separated organic layer was washed with hypo solution (2 x 100 mL) followed by brine solution (2 x 100 mL). The organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to afford crude material which was purified by column chromatography eluting 15% EtOAc/hexane to afford **C** (11.5 g, 58%) as yellow thick syrup.

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 9.06 (d, *J* = 4.8 Hz, 2H), 7.75 (t, *J* = 4.8 Hz, 1H), 5.97-5.92 (m, 1H), 1.83 (d, *J* = 6.4 Hz, 3H).

Synthesis of 2-hydroxy-1-(pyrimidin-2-yl) propan-1-one (D):

[00136] To a stirring solution of **C** (11.5 g, 53.4 mmol) in MeOH (80 mL) was added sodium formate (14.5 g, 214 mmol) and stirred the reaction mass at 80 °C for 6 h. After completion of reaction (by TLC), the reaction mixture was evaporated under reduced pressure to give crude product, which was purified by column chromatography eluting 50% EtOAc/*n*-hexane to afford **D** (5.0 g, 61.6%) as colorless liquid.

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 8.73 (d, *J* = 5.2 Hz, 2H), 7.55 (t, *J* = 4.8 Hz, 1H), 5.28-5.26 (m, 1H), 1.24 (d, *J* = 6.4 Hz, 1H), 0.99 (d, *J* = 6.4 Hz, 3H)

Synthesis of (2-((tert-butyldimethylsilyl) oxy)-1-(pyrimidin-2-yl) propan-1-one (E):

- [00137] To a stirring solution of **D** (5 g, 32.8 mmol) in DCM (50 mL) were added imidazole (5.5 g, 82.2 mmol), DMAP (800 mg, 0.65 mmol) at 0 °C and stirred for 10 min.
- 5 After added TBDMS-Cl (7.38 g, 49.2 mmol) at 0 °C and stirred at RT for 12 h. After completion of starting material (by TLC), diluted the reaction mass with H₂O (50 mL). The separated organic layer was washed with brine solution (2 x 50 mL). The organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to afford crude material which was purified by column chromatography eluting 20% EtOAc/hexane to afford **E**
- 10 (6 g, 68.8%) as an off-white solid.


¹H-NMR: (400 MHz, CDCl₃): δ 9.00 (d, *J* = 5.2 Hz, 2H), 7.71 (t, *J* = 4.8 Hz, 1H), 5.47-5.42 (m, 1H), 1.35 (d, *J* = 6.8 Hz, 3H), 0.79 (s, 9H), 0.05 (s, 6H).

Synthesis of (2-((tert-butyldimethylsilyl) oxy)-1-(pyrimidin-2-yl) propan-1-amine (F):

- 15 [00138] To a stirring solution of **E** (6 g, 22.5 mmol) in MeOH (50 mL) were added sodium acetate (3.69 g, 45.1 mmol), ammonium carbonate (21 g, 135 mmol), AcOH (1.28 mL, 22.5 mmol) at RT and stirred the reaction mixture at 90 °C for 2 h. The temperature of the reaction was cooled to RT and added sodium cyanoborohydride (2.84 g, 45.1 mmol) slowly and stirred at 90 °C for 6 h. After completion of reaction (by TLC), evaporated MeOH under
- 20 reduced pressure. The crude residue was diluted with water (20 mL) and extracted with DCM (2 x 50 mL). The organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to afford crude material was purified by column chromatography eluting 5% MeOH/DCM to afford **F** (4.2 g, 69.6%) as semi solid.

- 25 ¹H-NMR: (400 MHz, CDCl₃): δ 8.83 (d, *J* = 4.8 Hz, 2H), 7.40 (t, *J* = 5.2 Hz, 1H), 4.13 (t, *J* = 6.4 Hz, 2H), 3.90 (d, *J* = 6.4 Hz, 2H), 1.12 (d, *J* = 6.4 Hz, 3H), 0.70 (s, 9H), 0.02 (s, 6H).

Scheme I-2

**Synthesis of 1-tert-butyl 2-methyl 2-((benzyloxy) methyl) pyrrolidine-1, 2-dicarboxylate**5 **(G):**

[00139] To a stirring solution of t-BOC proline methyl ester (25 g, 109 mmol) in THF (250 mL) was added LiHMDS (240 mL, 240 mmol) at -20 °C and stirred for 2 h. To this BOM-chloride (23 mL, 163 mmol) was added drop wise at -30 °C and stirred for 2 h. After consumption of the starting material (by TLC), the reaction was quenched with aqueous NH₄Cl solution (100 mL) and extracted with EtOAc (2 x 200 mL). The combined organic layer was washed with water (2 x 150 mL) followed by brine solution (2 x 100 mL). The organic layer was dried over Na₂SO₄ and concentrated to obtain crude compound which was purified by column chromatography by eluting 10% EtOAc/*n*-hexane to afford compound **G** (30 g, 79%) as thick syrup.

15 **¹H-NMR:** (500 MHz, DMSO-*d*₆): δ 7.36-7.22 (m, 5H), 4.59-4.48 (m, 2H), 4.02-3.88 (m, 1H), 3.63 (s, 3H), 3.49-3.35 (m, 2H), 3.34-3.30 (m, 1H), 2.31-2.23 (m, 1H), 2.04-1.89 (m, 2H), 1.82-1.78 (m, 1H).

LCMS (*m/z*): 249.4 [(M⁺+1)-Boc].

Synthesis of 2-((benzyloxy) methyl)-1-(tert-butoxycarbonyl) pyrrolidine-2-carboxylic acid20 **(H):**

[00140] To a stirring solution of compound **G** (30 g, 86 mmol) in methanol (70 mL) was added NaOH solution (6.88 g in 70 mL H₂O) at RT. The reaction mixture was heated to 70 °C for 16 h. After consumption of the starting material (by TLC), the solvent from the reaction was evaporated under reduced pressure and diluted with EtOAc (2 x 200 mL). The separated

- 45 -

aqueous layer was acidified using citric acid solution (pH~3) and extracted with EtOAc (2 x 250 mL). The combined organic layer was dried over Na₂SO₄ and concentrated to afford crude which was triturated with n-hexane to obtain compound **H** (25 g, 86.8%) as off-white solid.

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 12.35 (br s, 1H), 7.37-7.29 (m, 5H), 4.56-4.48 (m, 2H), 4.06-4.00 (m, 1H), 3.92-3.89 (m, 1H), 3.66-3.45 (m, 1H), 3.37-3.28 (m, 1H), 2.31-2.20 (m, 1H), 2.05-1.97 (m, 1H), 1.87-1.75 (m, 2H), 1.38 (s, 9H).


LCMS (*m/z*): 335.3 [*M*⁺+1].

Synthesis of 1-(tert-butoxycarbonyl)-2-(hydroxymethyl) pyrrolidine-2-carboxylic acid (I):

[00141] To a stirring solution of compound **H** (25 g, 74 mmol) in methanol (150 mL) was added 50% wet 10% Pd/C (7 g) at RT and stirred for 10 h under H₂ atmosphere. After consumption of the starting material (by TLC), the reaction mixture was filtered through a pad of celite and the pad was washed with methanol (100 mL). Obtained filtrate was concentrated under reduced pressure to afford compound **I** (15 g, 82.8%) as white solid.

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 4.66 (br s, 1H), 3.96-3.83 (m, 1H), 3.63-3.59 (m, 1H), 3.49-3.41 (m, 1H), 3.34-3.25 (m, 1H), 2.30-2.17 (m, 1H), 1.95-1.72 (m, 3H), 1.38 (s, 9H).

Scheme I-3



Synthesis of (2*S*, 3*R*)-methyl 2-(((benzyloxy) carbonyl) amino)-3-((tert-butyldimethylsilyl) oxy) butanoate (K):

[00142] Referring to Scheme I-3, to a stirring solution of **J** (50 g, 187 mmol) in DMF (400 mL) were added DIPEA (86 mL, 468 mmol) TBDMS-Cl (33.66 mL, 224 mmol) at 0 °C and stirred at RT for 12 h. After completion of starting material (by TLC), diluted the reaction

- 46 -

mass with EtOAc (500 ml). The separated organic layer was washed with (2x200 mL) of Water followed by brine solution (2x100 mL). The organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to afford crude material was purified by column chromatography eluting 10% EtOAc/hexane to afford **K** (50 g, 70.1%) as colorless syrup.

¹H-NMR: (400 MHz, CDCl₃): δ 7.39-7.32 (m, 5H), 5.43 (d, *J* = 9.6 Hz, 1H), 5.14 (s, 2H), 4.45-4.43 (m, 1H), 4.29-4.26 (m, 1H), 3.72 (s, 3H), 1.21 (d, *J* = 6.0 Hz, 3H), 0.83 (s, 9H), 0.09 (s, 6H).

LCMS *m/z*: 382.2[M⁺+1].

Synthesis of benzyl ((2*S*, 3*R*)-3-((tert-butyldimethylsilyl) oxy)-1-hydrazinyl-1-oxobutan-2-yl) carbamate (L):

[00143] A solution of **K** (50 g, 131 mmol) in EtOH (400 mL) was added hydrazine hydrate (32.8 g, 656 mmol), at RT and after stirred at 90 °C for 24 h. After completion of starting material (by TLC), evaporated ethanol under reduced pressure. The crude residue was diluted with water (100 mL) and EtOAc (500 mL). After the separated organic layer was washed with (2x100 mL) of water followed by brine solution (1x100 mL). Dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. Obtained crude material was purified by column chromatography by eluting with 20% EtOAc/hexane to afford **L** (25 g, 50%) as colorless thick syrup.

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 9.10 (s, 1H), 7.36-7.30 (m, 5H), 6.83 (d, *J* = 9.6 Hz, 1H), 5.02 (s, 2H), 4.19 (s, 2H), 4.05-4.02 (m, 1H), 3.97-3.93 (m, 1H), 1.05 (d, *J* = 6.0 Hz, 3H), 0.81 (s, 9H), 0.01 (s, 6H).

Synthesis of benzyl ((1*S*, 2*R*)-2-((tert-butyldimethylsilyl) oxy)-1-(1, 3, 4-oxadiazol-2-yl) propyl) carbamate (M):

[00144] A solution of **L** (25 g, 65.6 mmol) in triethyl orthoformate (250 mL) was added *p*-TSA (catalytic, 250 mg) at RT and after stirred at 80 °C for 3 h. After completion of starting material (by TLC), evaporated triethyl orthoformate under reduced pressure. The crude residue was purified by column chromatography eluting 10% EtOAc/hexane to afford **M** (15 g, 58%) as thick syrup.

¹H-NMR: (500 MHz, DMSO-*d*₆): δ 9.22 (s, 1H), 7.85 (d, *J* = 9.5 Hz, 1H), 7.36-7.31 (m, 5H), 5.05 (s, 2H), 4.96-4.93 (m, 1H), 4.25 (t, *J* = 6.0 Hz, 1H), 1.23 (d, *J* = 6.0 Hz, 3H), 0.80 (s, 9H), 0.10 (s, 6H).

- 47 -

LCMS m/z: 392.4[M⁺+1].


Synthesis of (1*S*, 2*R*)-2-((tert-butyldimethylsilyl) oxy)-1-(1, 3, 4-oxadiazol-2-yl) propan-1-amine (N):

5 [00145] To a stirring solution of **M** (15 g, 38.3 mmol) in methanol (200 mL) was added 50% wet 10% Pd/C (5 g) and stirred under H₂ atmosphere (balloon pressure) for 4 h at RT. The reaction mixture was filtered through a pad of celite and triturated with methanol (100 mL). The filtrate was concentrated under reduced pressure to afford **N** (10 g, crude) as thick syrup.

¹H-NMR: (500 MHz, DMSO-*d*₆): δ 9.15 (s, 1H), 4.11 (t, *J* = 5.0 Hz, 1H), 4.03 (d, *J* = 2.0 Hz, 1H), 2.05 (br s, 2H), 1.17 (d, *J* = 6.0 Hz, 3H), 0.76 (s, 9H), 0.02 (s, 6H).

LCMS m/z: 258.3 [M⁺+1].

Scheme I-4



Synthesis of 3-(tert-butoxycarbonyl)-2, 2, 5-trimethyloxazolidine-4-carboxylic acid (O)

20 [00146] To a stirring solution of N-BOC threonine (15 g, 59.28 mmol) in THF (150 mL) was added PPTS (1.47 g, 5.92 mmol) followed by 2,2-dimethoxy propane (21.79 mL, 0.17 mol) at 0 °C under N₂ atmosphere. The reaction mixture was stirred at RT for 16 h. The reaction mixture was again heated to reflux for 6 h. The reaction mixture was diluted with aqueous NaHCO₃ solution and washed with EtOAc (1x100 mL). Aqueous layer was acidified using citric acid solution (pH~2) and extracted with CH₂Cl₂ (2x100 mL). The organic layer was

- 48 -

washed with brine, dried over anhydrous Na₂SO₄ and concentrated under vacuum to afford compound **O** (18 g, crude).

¹H-NMR: (400 MHz, DMSO-d₆): δ 13.25 (br s, 1H), 4.11-4.05 (m, 1H), 3.79 (d, 1H), 1.50 (s, 3H), 1.67 (s, 3H), 1.45 (s, 9H), 1.29 (d, 3H).

5 **Synthesis of *tert*-butyl 4-carbamoyl-2, 2, 5-trimethyloxazolidine-3-carboxylate (P):**

[00147] To a stirring solution of compound **O** (18 g, 69.4 mmol) in CH₂Cl₂ (180 mL) was added HOBt (14.16 g, 0.104 mol), EDCL.HCl (19.88 g, 0.104 mol) followed by NH₄Cl (5.56 g, 0.104 mol) and DIPEA (31.9 mL, 0.173 mol) at 0 °C. The reaction mixture was stirred at RT for 16 h. After consumption of the starting material (by TLC), the reaction mixture was
10 washed with aqueous citric acid, NaHCO₃ followed by brine. Organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to give crude; which was purified by silica gel column chromatography eluting with 2% MeOH/CH₂Cl₂ to afford compound **P** (13 g, 72.5%).

¹H-NMR: (400 MHz, DMSO-d₆): δ 7.51 (br s, 1H), 7.14 (br s, 1H), 3.97-3.95 (m, 1H), 3.71 (d, 1H), 1.51 (d, 6H), 1.34 (s, 9H), 1.24 (d, 3H).

15 LCMS (ESI): 159.1[(M⁺+1)-Boc]

Synthesis of (Z)-*tert*-butyl 4-(((dimethylamino) methylene) carbamoyl)-2, 2, 5-trimethyloxazolidine-3-carboxylate (Q):

20 [00148] A solution of compound **P** (13 g, 50.3 mmol) in DMF.DMA (130 mL) was stirred at reflux temperature for 3 h under N₂ atmosphere. After consumption of the starting material (by TLC), the reaction mixture was concentrated under reduced pressure to afford compound **Q** (15.7 g, crude). This crude material was directly taken for the next step without further purification.

25 **Synthesis of *tert*-butyl 2, 2, 5-trimethyl-4-(1, 2, 4-oxadiazol-5-yl) oxazolidine-3-carboxylate (R):**

[00149] To a stirring solution of compound **Q** (15.7 g, 50.09 mmol) in ethanol (157 mL) was added hydroxylamine hydrochloride (6.96 g, 0.10 mol) under N₂ atmosphere. The reaction
30 mixture was heated to reflux and stirred for 2 h. After consumption of the starting material (by

- 49 -

TLC), acetic acid (28.6 mL, 0.50 mol) was added to the reaction mixture and then refluxed for 16 h. The solvents from the reaction mixture was evaporated under vacuum to give crude; which was purified by silica gel column chromatography eluting with 10% EtOAc/Hexane to afford compound **R** (4.5 g, 32%).

5 ¹H-NMR: (400 MHz, DMSO-*d*₆): δ 6.35 (s, 2H), 4.61 (d, 1H), 4.22-4.15 (m, 1H), 1.55 (s, 6H), 1.37 (s, 2H), 1.25 (d, 3H), 1.21 (s, 6H).

LCMS (ESI): 284 [M⁺+1]

Mass (m/z): 283 [M⁺]


Synthesis of 1-amino-1-(1, 2, 4-oxadiazol-5-yl) propan-2-ol (S):

10 [00150] To a stirring solution of compound **R** (5 g, 17.6 mmol) in water (25 mL) was added trifluoroacetic acid (25 mL). The reaction mixture was stirred at RT for 5 h. After consumption of the starting material (by TLC), the reaction mixture was concentrated under vacuum. The residue was dissolved in water and neutralized with aqueous NaHCO₃. The solvent from the reaction mixture was evaporated under vacuum and extracted with 5%
15 MeOH/CH₂Cl₂ (3x 100 mL). The organic layer was concentrated under reduced pressure to afford compound **S** (2.5 g, crude).

¹H-NMR: (400 MHz, D₂O): δ 8.84 (s, 1H), 4.05 (d, 1H), 3.98-3.95 (m, 1H), 3.67 (s, 1H), 3.58 (d, 1H), 1.15 (d, 3H), 1.12 (d, 3H).

LCMS (ESI): 144.1 [M⁺+1]

Scheme I-5

**Synthesis of ethyl 2-((tert-butoxycarbonyl) amino)-5-oxohexanoate (T):**

- 5 **[00151]** To a stirring solution of 1-*tert*-butyl 2-ethyl 5-oxopyrrolidine-1,2-dicarboxylate (12 g, 46.6 mmol) in THF (120 mL) under inert atmosphere was added MeMgBr (3M in ether) (20.2 mL, 60.6 mmol) at 0 °C and stirred for 2 h. After consumption of the starting material (by TLC), the reaction mixture was quenched with aqueous NH₄Cl solution and the aqueous layer was extracted with EtOAc (2 x 200 mL). The combined organic extracts were dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude residue obtained was purified by silica gel column chromatography eluting with 20% EtOAc/Hexane to afford compound T (10 g, 79%).

¹H-NMR: (400 MHz, CDCl₃): δ 5.14 (br s, 1H), 4.23 (q, 2H), 2.62-2.47 (m, 2H), 2.17 (s, 4H), 1.91-1.82 (m, 1H), 1.45 (s, 10H), 1.26 (t, 3H).

15 **Synthesis of ethyl 5-methylpyrrolidine-2-carboxylate (V):**

- [00152]** To a stirring solution of compound T (10 g, 36.7 mmol) in CH₂Cl₂ (50 mL) was added TFA (14.89 mL, 194.6 mmol) at 0 °C. After being stirred for 2 h at RT, the reaction mixture was concentrated under reduced pressure to get compound U. Obtained material was dissolved in ethanol (100 mL) and 10% Pd/C (50% wet, 3 g) under N₂ atmosphere. The reaction mixture was stirred under H₂ atmosphere (balloon pressure) for 16 h. The reaction

- 51 -

mixture was filtered through a pad of celite and filtrate was concentrated under reduced pressure to afford compound **V** (15 g, crude). This material was directly taken for the next step without further purification.

¹H-NMR: (500 MHz, DMSO-*d*₆): 4.4 (m, 1H), 4.2 (m, 2H), 3.6 (m, 1H), 2.3 (m, 1H), 2.1 (m, 2H), 1.6 (m, 1H), 1.3 (d, 3H), 1.2 (t, 3H).

LCMS *m/z*: 158.1 [M^+ +1].

Synthesis of 1-tert-butyl 2-ethyl 5-methylpyrrolidine-1, 2-dicarboxylate (W):

[00153] To a stirring solution of compound **V** (30 g, 191 mmol) in CH₂Cl₂ (150 mL) was added DMAP (23.3 g, 191 mmol) followed by Et₃N (79.8 mL, 573 mmol) and Boc-anhydride (104 mL, 477 mmol) at 0 °C. The reaction mixture was stirred at RT for 16 h. The reaction mixture was diluted with CH₂Cl₂ (50 mL) and washed with water (2x150 mL) followed by brine. The separated organic layer was dried over anhydrous Na₂SO₄ and concentrated under vacuum. Obtained crude material was purified by column chromatography eluting with 6% EtOAc/Hexane to afford compound **W** (12 g, 82%) as pale yellow liquid.

¹H-NMR: (500 MHz, DMSO-*d*₆): δ 4.13-3.86 (m, 4H), 2.15 (d, *J* = 3.5 Hz, 1H), 1.99-1.82 (m, 2H), 1.52 (t, *J* = 4.5 Hz, 1H), 1.38 (s, 9H), 1.24 (t, *J* = 5.5 Hz, 3H), 1.16 (d, *J* = 6.5 Hz, 3H)
LCMS (*m/z*): 258 [M^+ +1].

Synthesis of 1-tert-butyl 2-ethyl 2-((benzyloxy) methyl)-5-methylpyrrolidine-1, 2-dicarboxylate (X):

[00154] To a stirring solution of compound **W** (8.0 g, 31.12 mmol) in THF (70 mL) was added LiHMDS (59 mL, 41.72 mmol) at -78 °C and stirred for 2 h. To this BOM-chloride (6.56 mL, 41.72 mmol) was added drop wise and stirred for 2 h at -30 °C. After consumption of the starting material (by TLC), the reaction was quenched with aqueous NH₄Cl solution (20 mL) and extracted with DCM (30 mL). The separated organic layer was dried over Na₂SO₄ and concentrated to afford crude material was purified by column chromatography eluting with 10% EtOAc/Hexane to afford compound **X** (11 g, 94.2%) as pale yellow liquid.

¹H-NMR: (500 MHz, DMSO-*d*₆): δ 7.33-7.25 (m, 5H), 4.38 (d, *J* = 10.5 Hz, 2H), 4.08-3.98 (m, 1H), 3.88 (d, *J* = 9.5 Hz, 2H), 2.20-2.08 (m, 2H), 1.38 (s, 9H), 1.37-1.29 (m, 4H), 1.19 (t, *J* = 7.5 Hz, 3H), 1.14-1.10 (m, 3H).

LCMS (*m/z*): 378 (M^+ +1).

Synthesis of 2-((benzyloxy) methyl)-1-(tert-butoxycarbonyl)-5-methylpyrrolidine-2-carboxylic acid (Y):

[00155] To a stirring solution of compound **X** (11 g, 29.17 mmol) in CH₃OH/THF (22 mL/20 mL) were added 2N NaOH solution (33 mL) at RT. The reaction mixture was heated to 65 °C for 8 h. After consumption of the starting material (by TLC), the solvent from the reaction was evaporated under reduced pressure and diluted with EtOAc (50 mL). The aqueous layer was acidified using citric acid solution and extracted with CH₂Cl₂ (2 x 100 mL). The separated organic layer was washed with water (1 x 50 mL), dried over Na₂SO₄ and concentrated to afford compound **Y** (8 g, 80%).

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 12.58 (s, 1H), 7.34-7.28 (m, 5H), 4.54-4.47 (m, 2H), 4.05-3.87 (m, 2H), 3.70-3.62 (m, 1H), 2.28-2.08 (m, 3H), 1.46-1.37 (m, 1H), 1.28 (s, 9H).


LCMS (m/z): 350 [M⁺+1].

Synthesis of 1-(tert-butoxycarbonyl)-2-(hydroxymethyl)-5-methylpyrrolidine-2-carboxylic acid (Z):

[00156] To a stirring solution of compound **Y** (8 g, 1.45 mmol) in methanol (40 mL) was added 50% wet Pd/C (4 g) under N₂ atmosphere. The reaction mixture was stirred under H₂ atmosphere (balloon pressure) at RT for 16 h. After consumption of the starting material (by TLC), the reaction mixture was filtered through a pad of celite and the pad was washed with methanol. Obtained filtrate was concentrated under reduced pressure to afford crude compound which was triturated with n-pentane to obtained compound **Z** (4.5 g, 75.2%) as white solid.

¹H-NMR: (500 MHz, DMSO-*d*₆): δ 12.37 (br s, 1H), 4.61 (br s, 1H), 3.95-3.85 (m, 3H), 2.18-2.06 (m, 3H), 1.44-1.41 (m, 1H), 1.38 (s, 9H), 1.09 (d, *J* = 6.0 Hz, 3H); LCMS (ESI): 260 [M⁺+1]

Scheme I-6



Synthesis of 1-(tert-butyl) 2-methyl 2-(1-hydroxyethyl) pyrrolidine-1, 2-dicarboxylate (AA):

[00157] To a stirring solution of N-BOC proline methyl ester (30 g, 131 mmol) in THF (100 mL) was added LiHMDS (323 mL, 327 mmol) at -20 °C and stirred for 30 min. To this
5 acetaldehyde (12.3 mL, 196 mmol) was added drop wise at -20 °C and stirred for 2 h. After consumption of the starting material (by TLC), the reaction was quenched with aqueous NH₄Cl solution (100 mL) and extracted with EtOAc (2 x 200 mL). The combined organic layer was washed with brine solution (2 x 150 mL). The organic layer was dried over Na₂SO₄ and concentrated to obtain crude compound which was purified by column chromatography by
10 eluting 10% EtOAc/*n*-hexane to afford compound **AA** (30 g, 83.8%) as thick syrup.

¹H-NMR: (400 MHz, CDCl₃): δ 5.82 (d, *J* = 10.0 Hz, 1H), 4.71-4.11 (m, 1H), 3.76 (s, 3H), 3.66-3.57 (m, 1H), 3.45-3.36 (m, 1H), 2.51-2.46 (m, 1H), 2.33-2.24 (m, 1H), 2.04-1.97 (m, 1H), 1.95-1.83 (m, 1H), 1.49 (s, 9H), 1.20-1.11 (m, 3H).

LCMS (*m/z*): 274.3 [M⁺+1].


15 **Synthesis of 1-(tert-butoxycarbonyl)-2-(1-hydroxyethyl) pyrrolidine-2-carboxylic acid (BB):**

[00158] To a stirring solution of compound **AA** (30 g, 109 mmol) in methanol (50 mL) was added NaOH solution (8.72 g in 10 mL H₂O, 218 mmol) at RT. The reaction mixture was heated to 70 °C for 12 h. After consumption of the starting material (by TLC), the solvent from
20 the reaction was evaporated under reduced pressure and diluted with DCM (200 mL). The separated aqueous layer was acidified using citric acid solution (pH~3) and extracted with DCM (2 x 250 mL). The combined organic layer was washed with brine solution (1 x 100 mL). The organic layer was dried over Na₂SO₄ and concentrated to afford compound **BB** (9.5 g, 33.6%) as brown solid.

25 ¹H-NMR: (400 MHz, DMSO-*d*₆): δ 4.53-4.45 (m, 1H), 3.59-3.48 (m, 1H), 3.25-3.19 (m, 1H), 2.33-2.16 (m, 1H), 1.90-1.78 (m, 3H), 1.38 (s, 9H), 0.95 (d, *J* = 6.4 Hz, 3H).

LCMS (*m/z*): 258.2 [M⁻-1].

Scheme I-7



5 Synthesis of 1-*tert*-butyl 2-ethyl 2-(1-hydroxyethyl)-5-methylpyrrolidine-1, 2-
dicarboxylate (CC):

[00159] To a stirring solution of compound **W** (18.0 g, 70 mmol) in THF (200 mL) was added LiHMDS (84 mL, 84 mmol) drop wise at -20 °C and stirred for 30 min. To this acetaldehyde (4.2 mL, 77 mmol) was added drop wise and stirred for 45 min at -20 °C. After consumption of the starting material (by TLC), the reaction was quenched with aqueous NH₄Cl solution (100 mL) and extracted with EtOAc (2 x 150 mL). The separated organic layer was dried over Na₂SO₄ and concentrated to afford crude material was purified by column chromatography eluting with 30% EtOAc/Hexane to afford compound **CC** (15 g, 71.4%) as colorless syrup.

¹H-NMR: (500 MHz, DMSO-*d*₆):δ 7.33-7.25 (m, 5H), 4.38 (d, *J* = 10.5 Hz, 2H), 4.08-3.98 (m, 1H), 3.88 (d, *J* = 9.5 Hz, 2H), 2.20-2.08 (m, 2H), 1.38 (s, 9H), 1.37-1.29 (m, 4H), 1.19 (t, *J* = 7.5 Hz, 3H), 1.14-1.10 (m, 3H);

LCMS (m/z):378 (M^++1)


Synthesis of 1-(*tert*-butoxycarbonyl)-2-(1-hydroxyethyl)-5-methylpyrrolidine-2-carboxylic acid (DD):

[00160] To a stirring solution of compound **CC** (15 g, 49 mmol) in CH₃OH/THF (10 mL/40 mL) were added NaOH (3.98 g, 99 mmol) in water (10 mL) at RT. The reaction mixture was heated to 70 °C for 16 h. After consumption of the starting material (by TLC), the solvent from the reaction was evaporated under reduced pressure and acidified by using citric acid (pH~4). The aqueous layer was extracted with EtOAc (2 x 200 mL). The combined organic layer was dried over Na₂SO₄ and concentrated to obtained crude compound, which was purified by column chromatography eluting 40% EtOAc to afford compound **DD** (4 g, 29.4%) as brown syrup.

¹H-NMR: (500 MHz, DMSO-*d*₆): δ 12.15 (br s, 2H), 4.54-4.50 (m, 1H), 4.03-4.02 (m, 1H), 2.17-1.77 (m, 3H), 1.41(s, 9H), 1.39-1.09 (m, 3H), 0.99-0.94 (m, 3H).

LCMS (*m/z*): 272.4 [*M*-1].

5 Scheme 1



Synthesis of *tert*-butyl2-((2-((*tert*-butyldimethylsilyl)oxy)-1-(pyrimidin-2-yl)propyl) carbamoyl)-2-(hydroxymethyl) pyrrolidine-1-carboxylate(C-1):

[00161] Referring to Scheme 1, to a stirring solution of I (2 g, 8.16 mmol) in CH₂Cl₂ (60 mL) was added intermediate F (2.39 g, 8.97 mmol), EDCI. HCl (2.33 g, 12.2 mmol) followed by HOBt (1.66 g, 12.24 mmol) and DIPEA (4.5 mL, 24.4 mmol) at 0°C. The reaction mixture was warmed to RT and stirred for 16 h. After consumption of the starting material (by TLC), the reaction mixture was diluted with CH₂Cl₂. The separated organic layer was washed with aqueous NaHCO₃ solution followed by aqueous NH₄Cl. The organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to get crude product, which was purified by silica gel column chromatography eluting with 70% EtOAc/hexane to afford compound **C-1** (2.3 g, 57.5%).

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 8.91 (d, 1H), 8.68 (d, 2H), 7.41 (br s, 1H), 5.74 (br t, 1H), 5.07-4.89 (m, 1H), 4.15-4.10 (m, 1H), 3.97-3.92 (m, 1H), 3.45-3.41 (m, 1H), 1.79-1.74 (m, 2H), 1.43-1.39 (m, 4H), 1.29-1.21 (m, 6H), 1.12 (d, 5H), 0.71 (s, 9H), 0.12 (t, 1H), 0.09 (s, 2H), 0.08 (s, 1H), 0.04 (s, 2H).

LCMS (*m/z*): 495.5 [*M*⁺+1].

Synthesis of *tert*-butyl 2-(2-((*tert*-butyldimethylsilyl)oxy)-1-(pyrimidin-2-yl)propyl)-1-oxo-2,5-diazaspiro[3.4]octane-5-carboxylate(C-2):

[00162] To a stirring solution of compound C-1 (2.3 g, 4.65mmol) in THF (23 mL) was added TPP (1.34 g, 5.12 mmol) followed by DTAD (1.6 g, 6.98 mmol) at 0 °C. The reaction mixture was warmed to RT and stirred for 16 h. After consumption of the starting material (by TLC), the reaction mixture was concentrated under reduced pressure to get crude product, which was purified by silica gel column chromatography eluting with 25% EtOAc/hexane to afford compound C-2 (1.2 g, 54.2%)

10 ¹H-NMR: (400 MHz, DMSO-d₆): δ8.82 (d, 2H), 7.49 (t, 1H), 4.72 (d, 1H), 4.31 (q, 1H), 3.62 (br s, 2H), 3.25-3.19 (m, 1H), 2.24-2.05 (m, 2H), 1.85-1.81 (m, 2H), 1.42 (br s, 1H), 1.25 (t, 3H), 0.92 (s, 8H), 0.75 (s, 9H), 0.02 (s, 3H).

LCMS (*m/z*): 477.4 [M⁺+1].

Synthesis of *tert*-butyl 2-(2-hydroxy-1-(pyrimidin-2-yl)propyl)-1-oxo-2,5 diazaspiro [3.4]octane-5-carboxylate (C-3):

15 [00163] To a stirring solution of compound C-2 (1.0 g, 2.10 mmol) in THF (20 mL) was added TBAF (1M in THF) (6.3 mL, 6.30 mmol) at 0 °C under N₂ atmosphere and stirred for 1 h. After consumption of the starting material (by TLC), the reaction mixture was quenched with ice water and extracted with EtOAc. The separated organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to obtain crude product, which was purified by silica gel column chromatography eluting with 5% MeOH/CH₂Cl₂ to afford compound C-3 (0.35 g, 46%).

20 ¹H-NMR: (400 MHz, DMSO-d₆): δ8.81 (d, 2H), 7.49 (t, 1H), 4.81 (d, 1H), 4.65 (d, 1H), 4.25-4.20 (m, 1H), 3.64-3.51 (m, 2H), 3.34 (s, 1H), 3.25-3.20 (m, 1H), 2.25-2.20 (m, 2H), 1.87-1.82 (m, 2H), 1.19 (d, 3H), 0.97 (s, 9H).

25 LCMS (*m/z*): 363.3 [M⁺+1].

Synthesis of *tert*-butyl 2-(2-hydroxy-1-(pyrimidin-2-yl) propyl)-1-oxo-2, 5-diazaspiro [3.4]octane-5-carboxylate (C-4)

30 [00164] To a stirring solution of compound C-3 (200 mg, 0.42 mmol) in THF (5 mL) was added TBAF (1M in THF) (0.84 mL, 0.84 mmol) at 0 °C under N₂ atmosphere and stirred for 2 h. After consumption of the starting material (by TLC), the reaction mixture was

- 57 -

concentrated under reduced pressure to obtain crude product, which was purified by silica gel column chromatography eluting with 2% MeOH/CH₂Cl₂ to afford **C-4** (125 mg, 82.2%).

¹H-NMR: (400 MHz, CD₃OD): 8.78 (t, *J* = 3.2 Hz, 2H), 7.42-7.38 (m, 1H), 4.92-4.87 (m, 1H), 4.79-4.56 (m, 1H), 4.43-4.38 (m, 1H), 4.33-3.97 (m, 1H), 3.88-3.80 (m, 1H), 3.48-3.44 (m, 1H), 3.40-3.30 (m, 1H), 2.31-2.12 (m, 2H), 1.96-1.84 (m, 2H), 1.42 (s, 9H), 1.32-1.28 (m, 3H);

LCMS (*m/z*): 363.3 [*M*⁺+1].

HPLC: 94.5%.

Synthesis of 2-(2-hydroxy-1-(pyrimidin-2-yl)propyl)-2,5-diazaspiro[3.4]octan-1-one (C-5):

10 **[00165]** To a stirring solution of compound **C-4** (0.3 g, 0.82 mmol) in CH₂Cl₂ (6 mL) was added molecular sieves (0.3 g) followed by BF₃-etherate (0.31 mL, 2.48 mmol) at 0 °C and stirred at RT for 1 h. After consumption of the starting material (by TLC), the reaction mixture was filtered and obtained residue was dissolved in MeOH and washed with CH₂Cl₂. The volatiles were evaporated under reduced pressure to obtain crude product, which was purified

15 by silica gel column chromatography eluting with 8% MeOH/CH₂Cl₂ to afford **C-5** (0.12 g, 55%).

¹H-NMR: (400 MHz, DMSO-*d*₆): δ8.81 (d, 2H), 7.49 (t, 1H), 4.81 (d, 1H), 4.65 (d, 1H), 4.25-4.20 (m, 1H), 3.64-3.51 (m, 2H), 3.12-3.01 (m, 2H), 2.15-2.10 (m, 2H), 1.87-1.82 (m, 2H), 1.19 (d, 3H).

20 **LCMS** (*m/z*): 263.1 [*M*⁺+1].

Synthesis of 2-(2-hydroxy-1-(pyrimidin-2-yl) propyl)-5-isobutyryl-2, 5-diazaspiro [3.4] octan-1-one (C-6):


[00166] To a stirring solution of compound **C-5** (500 mg, 1.9 mmol) in DCM (10 mL)

25 was added TEA (0.79 mL, 5.72 mmol) followed by isobutyryl chloride (241 mg, 2.28 mmol) at 0 °C under N₂ atmosphere and stirred for 2 h at RT. After consumption of the starting material (by TLC), the reaction mixture was diluted with water (10 mL) and extracted with CH₂Cl₂ (2 x 15 mL). The combined organic layer was washed with citric acid solution (1 x 20 mL). The separated organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated under

30 reduced pressure. Obtained crude material was purified by silica gel column chromatography eluting 2% MeOH/DCM to afford **C-6** (85 mg, 13.4%) as white solid.

¹H-NMR: (400 MHz, CD₃OD): δ 4.91-4.75 (m, 1H), 4.46-4.40 (m, 1H), 3.99-3.85 (m, 1H), 3.70-3.64 (m, 1H), 3.60-3.45 (m, 3H), 2.80-2.69 (m, 1H), 2.29-2.22 (m, 2H), 2.07-1.96 (m, 2H), 1.45-1.33 (m, 3H), 1.16-1.10 (m, 3H), 1.06-1.01 (m, 3H) .

5 Scheme 2



10

Synthesis of tert-butyl 2-(((1R, 2S)-2-((tert-butyldimethylsilyl) oxy)-1-(1, 3, 4-oxadiazol-2-yl) propyl) carbamoyl)-2-(hydroxymethyl) pyrrolidine-1-carboxylate (C-7):

[00167] To a stirring solution of compound **I** (2.1 g, 8.57 mmol) in CH₂Cl₂ (20 mL) were added DIPEA (3.81 mL, 21.4 mmol), intermediate **F** (2.64 g, 10.28 mmol), HATU (3.90 g, 10.28 mmol) at 0 °C and stirred to RT for 16 h. After consumption of the starting material (by TLC), the reaction mixture was diluted with water (30 mL). The organic layer was washed with citric acid (1 x 50 mL) followed by brine solution (1 x 50 mL). The organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. Obtained crude material was purified by silica gel column chromatography eluting with 50% EtOAc/*n*-hexane to afford compound **C-7** (3.5 g, 85.15%) as yellow thick syrup.

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 9.24 (s, 1H), 8.17 (d, *J* = 8.4 Hz, 1H), 5.82-5.54 (m, 1H), 4.36-4.28 (m, 1H), 4.12-4.00 (m, 1H), 3.93-3.66 (m, 1H), 3.43-3.39 (m, 1H), 2.69-2.66 (m, 2H), 2.02-1.94 (m, 2H), 1.78-1.68 (m, 2H), 1.40 (s, 9H), 1.23-1.15 (m, 3H), 0.75 (s, 9H), -0.08 (s, 6H);

Mass (ESI): m/z 484.7 [$M^+ + 1$]

Synthesis of tert-butyl 2-(((1R, 2S)-2-((tert-butyldimethylsilyl) oxy)-1-(1, 3, 4-oxadiazol-2-yl) propyl) carbamoyl)-2-(hydroxymethyl) pyrrolidine-1-carboxylate (C-8):

[00168] To a stirring solution of triphenylphosphine (4.05 g, 15.4 mmol) in THF (30 mL) was added DIAD (3.03 g, 15.4 mmol) at RT and stirred for 15 min. After added compound C-7 (3 g, 6.19 mmol) in (20 mL) THF slowly and reaction mixture was stirred at RT for 3 h. After consumption of the starting material (by TLC), the reaction was concentrated under reduced pressure. The crude material was triturated with 30% Ether/*n*-hexane (2 x 50 mL). The filtered organic solvent was concentrated and crude residue was purified by column chromatography by eluting 30% EtOAc/*n*-hexane to afford compound C-8 (2 g, 69.4%) as thick syrup.

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 9.28 (s, 1H), 5.30-5.21 (m, 1H), 4.84-4.81 (m, 1H), 4.49-4.29 (m, 1H), 4.05-3.91 (m, 1H), 3.65-3.61 (m, 1H), 3.39-3.24 (m, 1H), 2.23-2.12 (m, 2H), 1.81-1.76 (m, 2H), 1.41 (s, 9H), 1.20-1.14 (m, 3H), 0.75 (s, 9H), -0.22 (s, 6H);

Mass (ESI): m/z 467.6 [$M^+ + 1$]

Synthesis of tert-butyl 2-((1R, 2S)-2-hydroxy-1-(1, 3, 4-oxadiazol-2-yl) propyl)-1-oxo-2, 5-diazaspiro [3.4] octane-5-carboxylate (C-9):

[00169] To a stirring solution of compound C-8 (2 g, 4.29 mmol) in THF (20 mL) was added TBAF (4.3 mL, 6.43 mmol) slowly at 0° C and stirred at RT for 30 min. After completion of reaction (by TLC), the reaction mixture was evaporated under reduced pressure. The crude residue was purified by column chromatography by eluting 70% EtOAc/*n*-hexane to afford C-9 (400 mg, 26.5%) as white solid.

¹H-NMR: (400 MHz, DMSO-*d*₆): 9.29 (s, 1H), 5.30-5.21 (m, 1H), 5.08-4.95 (m, 1H), 4.08-4.03 (m, 1H), 3.91-3.71 (m, 1H), 3.75-3.71 (m, 1H), 3.65-3.52 (m, 1H), 3.36-3.22 (m, 1H), 2.22-2.09 (m, 2H), 1.83-1.77 (m, 2H), 1.41 (s, 9H), 1.10-1.04 (m, 3H);

LCMS: 353.4;

HPLC: 99.82%.

Synthesis of 2-((1R, 2S)-2-hydroxy-1-(1, 3, 4-oxadiazol-2-yl)propyl)-2, 5-diazaspiro [3.4] octan-1-one (C-10):

[00170] To a stirring solution of compound C-9 (500 mg, 1.07 mmol) in DCM (5 mL) was added BF₃(OEt)₂ (0.26 mL, 2.14 mmol) slowly at 0 °C and stirred at RT for 30 min. After completion of reaction (by TLC), the reaction mixture was evaporated under reduced pressure.

- 60 -

The crude residue was triturated with di ethylether/*n*-pentane (5 mL/5 mL) to obtain solid, **C-10**, which was taken into the next step without further purification.

Synthesis of 2-((1*R*, 2*S*)-2-hydroxy-1-(1, 3, 4-oxadiazol-2-yl) propyl)-5-isobutyryl-2, 5-diazaspiro [3.4] octan-1-one (C-11):


- 5 **[00171]** To a stirring solution of compound **C-10** (700 mg, 2.77 mmol) in DCM (10 mL) was added TEA (0.94 mL, 6.92 mmol), SM-3 (440 mg, 4.15 mmol) at 0 °C and stirred to RT for 1 h. After completion of reaction (by TLC), diluted with water (20 mL). The organic layer was washed with brine solution (1 x 50 mL). The organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The crude residue was purified by
- 10 column chromatography followed by preparative HPLC purification to afford **C-11** (150 mg, 16.8%) as white solid.

¹H-NMR: (400 MHz, DMSO-*d*₆): 9.26 (s, 1H), 5.24-5.04 (m, 1H), 5.03-4.93 (m, 1H), 4.27-3.91 (m, 1H), 3.90-3.64 (m, 1H), 3.49-3.45 (m, 3H), 2.73-2.61 (m, 1H), 2.17-2.14 (m, 2H), 2.10-2.06 (m, 2H), 1.25-1.07 (m, 3H), 0.95-0.93 (m, 6H).

- 15 **LCMS** *m/z*: 323.3.

HPLC: 97.89%.

Scheme 3



- 20 **Synthesis of 1-*tert*-butyl 2-methyl 2-(hydroxymethyl)pyrrolidine-1,2-dicarboxylate(C-12):**

[00172] To a stirring solution of compound **G** (74 g, 0.21 mol) in methanol (740 mL) was added 10% Pd/C (50% wet, 14.8 g) under N₂ atmosphere and stirred for 6 h under H₂

- 61 -

atmosphere (balloon pressure). The reaction mixture was filtered through celite pad and concentrated under reduced pressure to afford compound **C-12** (45 g, 82%) as crude.

Synthesis of 1-tert-butyl 2-methyl 2-formylpyrrolidine-1,2-dicarboxylate(C-13):

- 5 [00173] To a stirring solution of compound **C-12** (10 g, 38.5 mmol) in CH₂Cl₂ (100 mL) was added Dess-Martin periodinane (19.6 g, 46.27 mmol) at 0 °C under N₂ atmosphere and stirred for 3 h. After consumption of the starting material (by TLC), the reaction was quenched with aqueous NaHCO₃ solution and extracted with CH₂Cl₂ (2x 100 mL). The organic layer was dried over anhydrous Na₂SO₄ and concentrated under vacuum. The crude was purified by
10 column chromatography eluting with 10% EtOAc/Hexane to afford compound **C-13** (7 g, 70.5%).

Synthesis of 1-tert-butyl 2-methyl 2-((((1S,2R)-2-hydroxy-1-(1,2,4-oxadiazol-5-yl) propyl) amino) methyl)pyrrolidine-1,2-dicarboxylate (C-14):

- 15 [00174] To a stirring solution of compound **C-13** (3 g, 11.6 mmol) in MeOH (30 mL) was added sodium acetate (1.91 g, 23.3 mmol) followed by intermediate **S** (3.6 g, 13.9 mmol). The reaction mixture was heated to reflux for 1 h. The reaction mixture was slowly cooled to RT-0 °C, to this sodium cyanoborohydride (1.465 g, 23.3 mmol) and stirring was continued for another 6 h at RT. After consumption of the starting material (by TLC), methanol from the
20 reaction was evaporated under reduced pressure and the residue was diluted with water and extracted with EtOAc (2x 50 mL). The separated organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to give crude; which was purified by silica gel column chromatography eluting with 40% EtOAc/Hexane to afford compound **C-14** (2.5 g, 56%).
25 LCMS *m/z*: 385 [M⁺+1].

Synthesis of tert-butyl 2-((1S,2R)-2-hydroxy-1-(1,2,4-oxadiazol-5-yl) propyl)-1-oxo-2,5-diazaspiro [3.4] octane-5-carboxylate (C-15):

- 30 [00175] To a stirring solution of compound **C-14** (1.5 g, 3.90 mmol) in THF (30 mL) was cooled to 0 °C and added *t*-BuMgCl (1M in THF, 15.6 mL, 15.6 mmol) and stirred for 15 min. After consumption of the starting material (by TLC), the reaction mixture was quenched with aqueous NH₄Cl solution and diluted with water. Aqueous layer was extracted with EtOAc (2x 25 mL). The separated organic layer was dried over anhydrous Na₂SO₄ and concentrated

- 62 -

under reduced pressure. Obtained crude material was purified by silica gel column chromatography eluting with 150% EtOAc/CH₂Cl₂ to afford **C-15** (0.15 g, 11%).

¹H-NMR: (400 MHz, DMSO-d₆): δ 9.02 (s, 1H), 5.15 (s, 1H), 4.37-4.32 (m, 1H), 3.95 (d, 1H), 3.66-3.60 (m, 1H), 3.36-3.30 (m, 1H), 2.29-2.09 (m, 2H), 1.87-1.82 (m, 2H), 1.55 (s, 9H), 1.27 (d, 3H).

LCMS (ESI) *m/z*: 351 [M⁺-1].

HPLC Purity: 96%.

Synthesis of 2-((1*S*,2*R*)-2-hydroxy-1-(1,2,4-oxadiazol-5-yl)propyl)-2,5-diazaspiro[3.4]octan-1-one (C-16):


[00176] To a stirring solution of C-15 (0.4 g, 1.13 mmol) in CH₂Cl₂ (4 mL) was added TFA (0.43 mL) at 0 °C and stirred at RT for 30 min. The reaction mixture was concentrated under vacuum. Obtained crude material was purified by prep-HPLC to afford C-16 (65mg) as TFA salt.

¹H-NMR: (400 MHz, DMSO-d₆): δ 9.89 (br s, 1H), 9.08 (s, 1H), 5.46 (d, 1H), 5.31 (s, 1H), 4.37-4.35 (m, 1H), 3.99 (d, 1H), 3.81 (d, 1H), 3.42-3.35 (m, 2H), 2.35-2.18 (m, 2H), 2.10-2.03 (m, 2H), 1.24 (d, 3H).

LCMS (ESI) *m/z*: 253.4 [M⁺+1].

HPLC Purity: 95%.

Scheme 4



Synthesis of *tert*-butyl 2-((2-((*tert*-butyldimethylsilyl) oxy)-1-(pyrimidin-2-yl) propyl) carbamoyl)-2-(hydroxymethyl)-5-methylpyrrolidine-1-carboxylate (C-17):

- 63 -

[00177] To a stirring solution of compound **Z** (500 mg, 1.93 mmol) in DCM (10 mL) were added *N,N*-diisopropylethylamine (1.0 mL, 5.79 mmol), intermediate **F** (566mg, 2.12mmol), followed by EDCI (737mg, 3.86 mmol), HOBT (521mg, 3.86mmol) at 0 °C and stirred at RT for 16 h. After consumption of the starting material (by TLC), the reaction mixture was diluted with water (20 mL). The separated organic layer was washed with saturated brine solution (1x30 mL). The separated organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford crude compound which was purified by column chromatography to obtained compound **C-17** (300 mg, 30.6%) as pale yellow liquid. ¹H-NMR: (500 MHz, CDCl₃): δ 8.73-8.68 (m, 3H), 5.30 (s, 2H), 4.40-4.15 (m, 1H), 3.76-3.61 (m, 1H), 3.18-3.13 (m, 1H), 2.18-2.13 (m, 4H), 1.46 (s, 9H), 1.32 (d, *J* = 5.5 Hz, 3H), 1.24 (d, *J* = 6.5 Hz, 3H), 1.08 (d, *J* = 6.5 Hz, 1H), 0.79 (s, 9H), 0.74 (t, *J* = 10.5 Hz, 1H), 0.12 (s, 6H)

Synthesis of *tert*-butyl 2-(2-((*tert*-butyldimethylsilyl) oxy)-1-(pyrimidin-2-yl) propyl)-6-methyl-1-oxo-2, 5-diazaspiro [3.4] octane-5-carboxylate (C-18):

[00178] To a stirring solution of triphenylphosphine (380 mg, 1.47 mmol) in dry THF (10 mL) was added DTAD (339 mg, 1.47 mmol) at RT and stirred for 10 min. After added compound **C-17** (300 mg, 0.59 mmol) and the reaction mixture was stirred at RT for 16 h. After consumption of the starting material (by TLC), the reaction mixture was concentrated under reduced pressure. The crude material was purified by silica gel column chromatography eluting 30% EtOAc/hexane to afford compound **C-18** (80 mg, 27.5%) as yellow liquid. ¹H-NMR: (400 MHz, CDCl₃): δ 7.73-7.64 (m, 3H), 5.26 (s, 2H), 4.30 (dd, *J* = 4.4 Hz, 4.4 Hz, 1H), 4.17-4.12 (m, 1H), 2.76 (t, *J* = 6.8 Hz, 1H), 2.32-2.28 (m, 2H), 2.05-2.00 (m, 2H), 1.49 (s, 9H), 1.45 (d, *J* = 5.2 Hz, 3H), 1.42 (d, *J* = 4.8 Hz, 3H), 0.89 (s, 9H), 0.15 (s, 6H).

Synthesis of *tert*-butyl 2-(2-hydroxy-1-(pyrimidin-2-yl) propyl)-6-methyl-1-oxo-2, 5-diazaspiro [3.4] octane-5-carboxylate (C-19):

[00179] To a stirring solution of compound **C-18** (700 mg, 1.42 mmol) in dry THF (10 mL) was added TBAF (744 mg, 2.85mmol) at 0 °C. The reaction mixture was stirred at RT for 3 h. After consumption of the starting material (by TLC), the reaction mixture was concentrated under reduced pressure. The crude material was purified by silica gel column chromatography eluting 80% EtOAc/hexane to afford **C-19** (0.16 g, 30%) as yellow thick syrup.

- 64 -

¹H-NMR: (500 MHz, DMSO-*d*₆): δ 8.77 (d, *J* = 5.0 Hz, 2H), 7.42(t, *J* = 4.5 Hz, 1H), 4.76-4.63 (m, 2H), 4.14-4.11 (m, 1H), 3.85-3.83 (m, 1H), 3.54-3.34 (m, 1H), 2.25-2.21 (m, 2H), 2.08-1.93 (m, 1H), 1.53-1.47 (m, 1H), 1.39 (s, 9H), 1.15(d, *J* = 6.0 Hz, 3H), 1.03-0.80 (m, 4H).

LCMS (ESI): 377.4 [M⁺+1].

5 HPLC: 94.6% (both isomers).

Synthesis of 2-(2-hydroxy-1-(pyrimidin-2-yl) propyl)-6-methyl-2, 5-diazaspiro [3.4] octan-1-one (C-20):


[00180] To a stirring solution of compound **C-19** (260 mg, 0.69 mmol) in DCM (10 mL) was added 4A molecular sieves (70 mg), BF₃·OEt₂ (0.08 mL) at 0° C under N₂ atmosphere. The reaction mixture was stirred at RT for 30 min. After consumption of the starting material (by TLC), the reaction mixture was concentrated under reduced pressure to afford crude, which was purified by preparative HPLC method to afford **C-20** (65 mg, 34%) as white solid.

15 ¹H-NMR: (400 MHz, D₂O): δ 8.88-8.86 (m, 2H), 7.63-7.57 (m, 1H), 4.92-4.89 (m, 1H), 4.61-4.56 (m, 1H), 4.03-3.91 (m, 3H), 2.61-2.41 (m, 3H), 1.92-1.84 (m, 1H), 1.46 (d, *J* = 6.8 Hz, 3H), 1.37 (d, *J* = 4.8 Hz, 3H);

LCMS (ESI): 295.3 [M⁺+H₂O].

HPLC: 91.2% (both enantiomers).

20 **Scheme 5**



Synthesis of tert-butyl 2-((2-((tert-butyldimethylsilyl)oxy)-1-(1,3,4-oxadiazol-2-yl)propyl)carbamoyl)-2-(hydroxymethyl)-5-methylpyrrolidine-1-carboxylate (C-21):

[00181] To a stirring solution of compound Z (4 g, 15.4 mmol) in DCM (30 mL) were added *N,N*-diisopropylethylamine (8 mL, 46.33mmol), intermediate N (4.74 g, 18.48 mmol) followed by HATU (7 g, 18.48 mmol) at 0 °C and stirred at RT for 12 h. After consumption of the starting material (by TLC), the reaction mixture was diluted with water (20 mL). The separated organic layer was washed with 10% citric acid solution (1 x 50 mL) followed by saturated brine solution (1x50 mL). The separated organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford crude compound which was purified by column chromatography eluting 30% EtOAc/*n*-hexane to obtained compound **C-21** (5.5g, 71.8%) as pale yellow liquid.

¹H-NMR: (500 MHz, DMSO-*d*₆): δ 9.25 (s, 1H), 5.50-5.41 (m, 1H), 5.19-5.06 (m, 2H), 4.38-4.36 (m, 1H), 4.05-3.90 (m, 3H), 2.69-1.98 (m, 2H), 1.39(s, 9H), 1.37-1.26 (m, 2H), 1.23-1.16 (m, 6H), 0.76 (s, 9H), -0.01 (s, 6H).

Synthesis of tert-butyl 2-(2-((tert-butyldimethylsilyl)oxy)-1-(1,3,4-oxadiazol-2-yl)propyl)-6-methyl-1-oxo-2,5-diazaspiro[3.4]octane-5-carboxylate (C-22):

[00182] To a stirring solution of triphenylphosphine (5.5 g, 11.04 mmol) in THF (55 mL) was added DTAD (5.07 g, 22.05 mmol) at RT and stirred for 10 min. After added compound C-21 (5.5g, 11.04mmol) and the reaction mixture was stirred at RT for 16 h. After consumption of the starting material (by TLC), the reaction mixture was concentrated under reduced pressure. The crude material was purified by silica gel column chromatography eluting 30% EtOAc/hexane to afford compound **C-22** (4 g, 75.4%) as yellow liquid.

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 9.36 (s, 1H), 5.30-4.48 (m, 1H), 4.47-4.31 (m, 1H), 4.03-3.88 (m, 2H), 3.62-3.58(m, 1H), 2.38-1.98 (m, 3H), 1.57-1.53 (m, 1H), 1.41 (s, 9H), 1.38-1.15 (m, 6H), 0.69 (s, 9H), -0.01 (s, 6H).

Synthesis tert-butyl 2-(2-hydroxy-1-(1,3,4-oxadiazol-2-yl)propyl)-6-methyl-1-oxo-2,5-diazaspiro [3.4] octane-5-carboxylate (C-23):

[00183] To a stirring solution of compound **C-22** (3.5 g, 7.29 mmol) in THF (35 mL) was added TBAF (1M THF) (14.5 mL, 14.58 mmol) at 0 °C. The reaction mixture was stirred at RT for 1 h. After consumption of the starting material (by TLC), the reaction mixture was

- 66 -


concentrated under reduced pressure. The crude material was purified by silica gel column chromatography eluting 60% EtOAc/hexane to afford **C-23** (800 mg, 30.7%) as an off-white solid.

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 9.30 (s, 1H), 5.30-5.21 (m, 1H), 5.13-5.02 (m, 1H), 4.13-4.03 (m, 1H), 3.90-3.83 (m, 1H), 3.74-3.68 (m, 1H), 3.62-3.49 (m, 1H), 2.29 (m, 3H), 1.56-1.52 (m, 1H), 1.40 (s, 9H), 1.25-1.14 (m, 6H).

LCMS (ESI) (*m/z*): 377.4 [*M*⁺+1].

HPLC: 95.38%.

10 **Scheme 6**



Synthesis of tert-butyl 2-(((1*R*, 2*S*)-2-((tert-butyldimethylsilyl)oxy)-1-(1,3,4-oxadiazol-2-yl)propyl) carbamoyl)-2-(1-hydroxyethyl)pyrrolidine-1-carboxylate (**C-24**):

15 **[00185]** To a stirring solution of compound **BB** (3 g, 11.5 mmol) in CH₂Cl₂ (30 mL) were added DIPEA (6 mL, 34.5 mmol), intermediate N (2.95 g, 11.5 mmol), HATU (5.24 g, 13.8 mmol) at 0 °C and stirred to RT for 12 h. After consumption of the starting material (by TLC), the reaction mixture was diluted with water (30 mL) and extracted with CH₂Cl₂ (50 mL). The combined organic layer was washed with citric acid (1 x 50 mL) followed by brine solution (1 x 50 mL). The organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. Obtained crude material was purified by silica gel column chromatography eluting with 35% EtOAc/*n*-hexane to afford compound **C-24** (3.6 g, 62.5%) as yellow thick syrup.

- 67 -

¹H-NMR: (500 MHz, DMSO-*d*₆): δ 9.26 (s, 1H), 8.74 (d, *J* = 8.5 Hz, 2H), 5.30-5.12 (m, 1H), 4.61-4.58 (m, 1H), 4.38-4.35 (m, 2H), 3.57-3.54 (m, 1H), 2.68-1.98 (m, 2H), 1.97-1.69 (m, 2H), 1.40 (s, 9H), 1.19-1.00 (m, 6H), 0.85 (s, 6H), -0.25 (s, 9H).

Mass (ESI): *m/z* 497.6[M⁻-1].

5 **Synthesis of tert-butyl 2-((1*R*, 2*S*)-2-((tert-butyldimethylsilyl) oxy)-1-(1, 3, 4-oxadiazol-2-yl) propyl)-1-methyl-3-oxo-2, 5-diazaspiro [3.4] octane-5-carboxylate (C-25):**

[00186] To a stirring solution of triphenylphosphine (3.6 g, 14.05 mmol) in THF (20 mL) was added DIAD (2.8 g, 14.05 mmol) at RT and stirred for 30 min. After added compound C-24 (3.5 g, 7.02 mmol) in (15 mL) THF slowly and reaction mixture was stirred at RT for 6 h.

10 After consumption of the starting material (by TLC), the reaction was concentrated under reduced pressure. The crude material was purified by column chromatography by eluting 30% EtOAc/*n*-hexane to afford compound C-25 (3 g, 89.2%) as thick syrup.

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 9.33 (s, 1H), 5.33-5.29 (m, 2H), 4.79-4.75 (m, 1H), 4.50-4.45 (m, 1H), 4.06-3.96 (m, 1H), 2.16-2.08 (m, 1H), 2.01-1.89 (m, 1H), 1.81-1.76 (m, 2H),
15 1.40 (s, 9H), 1.24-1.15 (s, 6H), 0.80 (s, 9H), -0.08 (s, 6H).

Mass (ESI): *m/z* 319.3 [M⁺+1].

Synthesis of tert-butyl 2-((1*R*, 2*S*)-2-hydroxy-1-(1, 3, 4-oxadiazol-2-yl) propyl)-1-methyl-3-oxo-2, 5-diazaspiro [3.4] octane-5-carboxylate (C-26):

[00187] To a stirring solution of compound C-25 (3 g, 6.25 mmol) in THF (25 mL) was
20 added TBAF (12.5 mL, 12.5 mmol) at 0 °C and stirred for 30 min. After consumption of the starting material (by TLC), the reaction mixture was concentrated under reduced pressure. Obtained crude material was purified by silica gel column chromatography eluting with 60% EtOAc/*n*-hexane to afford C-26 (0.8 g, 35%) as white solid.


¹H-NMR: (400 MHz, DMSO-*d*₆): δ 9.29 (s, 1H), 5.28-5.15 (m, 1H), 5.08-4.89 (m, 1H), 4.43-4.35 (m, 1H), 3.97-3.79 (m, 1H), 3.65-3.40 (m, 1H), 3.31-3.25 (m, 1H), 2.15-2.00 (m, 2H),
25 1.93-1.86 (m, 1H), 1.80-1.68 (m, 1H), 1.40 (s, 9H), 1.18-1.04 (m, 6H);

Mass (ESI): *m/z* 367.4 [M⁺+1].

HPLC: 95.85% (both isomers).

- 68 -

Scheme 7



Synthesis of tert-butyl 2-((2-((tert-butyldimethylsilyl) oxy)-1-(pyrimidin-2-yl) propyl) carbamoyl)-2-(1-hydroxyethyl) pyrrolidine-1-carboxylate (C-27):

[00188] To a stirring solution of compound **BB** (200 mg, 0.77 mmol) in CH₂Cl₂ (20 mL) were added DIPEA (298 mg, 2.31 mmol), EDCI (221 mg, 1.15 mmol), HOBT (177 mg, 1.15 mmol) followed by intermediate **F** (206 mg, 0.77 mmol) at 0 °C and stirred for 16 h at RT. After consumption of the starting material (by TLC), the reaction mixture was concentrated under reduced pressure. Obtained crude material was purified by silica gel column chromatography eluting with 4% MeOH/DCM to afford compound **C-27** (210 mg, 53.6%) as pale yellow thick syrup.

¹H-NMR: (500 MHz, DMSO-*d*₆): δ 8.77 (d, *J* = 4.5 Hz, 2H), 7.40 (t, *J* = 5.0 Hz, 1H), 4.95-4.92 (m, 2H), 4.21-4.05 (m, 2H), 3.34-3.25 (m, 2H), 3.20-3.12 (m, 1H), 1.81-1.63 (m, 4H), 1.48 (s, 9H), 1.12-1.06 (m, 3H), 0.99-0.85 (m, 3H), 0.70 (s, 9H), -0.06 (s, 6H);

LCMS (*m/z*): 509.4 [M⁺+1]

Synthesis of tert-butyl 2-(2-((tert-butyldimethylsilyl) oxy)-1-(pyrimidin-2-yl) propyl)-1-methyl-3-oxo-2, 5-diazaspiro [3.4] octane-5-carboxylate (C-28):

[00189] To a stirring solution of triphenylphosphine (206 mg, 0.78 mmol) in THF (6 mL) was added DIAD (179 mg, 0.78 mmol) at RT and stirred for 30 min. To this added compound **C-27** (200 mg, 0.39 mmol) in (5 mL) THF slowly and reaction mixture was stirred at RT for 2 h. After consumption of the starting material (by TLC), the reaction was concentrated under reduced pressure. The crude material was triturated with diethylether/*n*-

- 69 -

pentane (10 mL/10 mL). The filtered solvent was concentrated and purified by silica gel column chromatography eluting 30% EtOAc/*n*-hexane to afford compound **C-28** (80 mg, 41.8%) as pale yellow liquid.

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 8.81-8.77 (m, 2H), 7.72 (d, *J* = 5.5 Hz, 1H), 4.92-4.80 (m, 1H), 4.70-4.65 (m, 2H), 4.46-4.20 (m, 1H), 3.30-3.16 (m, 1H), 1.86-1.65 (m, 2H), 1.64-1.43 (m, 2H), 1.40 (s, 9H), 1.20-1.01 (m, 3H), 0.92-0.85 (m, 3H), 0.74 (s, 9H), -0.06 (s, 6H);

LCMS (*m/z*): 491.4 [*M*⁺+1].

Synthesis of tert-butyl 2-(2-hydroxy-1-(pyrimidin-2-yl) propyl)-1-methyl-3-oxo-2, 5-diazaspiro [3.4] octane-5-carboxylate (C-29):

10 **[00190]** To a stirring solution of compound **C-28** (450 mg, 0.91 mmol) in THF (10 mL) was added TBAF (1.8 mL, 1.83 mmol) in 5 mL THF at 0 °C and stirred to RT for 4 h. After consumption of the starting material (by TLC), the reaction was concentrated under reduced pressure. The crude material was purified by silica gel column chromatography eluting 2% MeOH/DCM followed by preparative HPLC purification to afford **C-29** (100 mg, 29%) as a


15 semi solid.

¹H-NMR: (400 MHz, CD₃OD): δ 8.82-8.79 (m, 2H), 8.78-7.43 (m, 1H), 4.94-4.89 (m, 1H), 4.63-4.44 (m, 1H), 3.99-3.73 (m, 1H), 3.56-3.36 (m, 2H), 2.29-2.22 (m, 2H), 1.99-1.73 (m, 2H), 1.40 (s, 9H), 1.24-1.20 (m, 3H), 1.13-1.05 (m, 3H).

LCMS(*m/s*): 377.5 [*M*⁺+1].

20 **HPLC:** 94.4% (both isomers).

Scheme 9



Synthesis of tert-butyl 2-((2-((tert-butyldimethylsilyl) oxy)-1-(pyrimidin-2-yl) propyl) carbamoyl)-2-(1-hydroxyethyl)-5-methylpyrrolidine-1-carboxylate (C-30):

[00191] To a stirring solution of compound DD (500 mg, 1.83 mmol) in DCM (10 mL) were added *N,N*-diisopropylethylamine (0.84 mL, 4.57 mmol), intermediate F (586 mg, 2.19 mmol), followed by EDCI (296 mg, 2.19 mmol), HOBT (338 mg, 2.19 mmol) at 0 °C and stirred at RT for 16 h. After consumption of the starting material (by TLC), the reaction mixture was diluted with water (20 mL). The separated organic layer was washed with brine solution (1 x 20 mL). The separated organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford crude compound which was purified by column chromatography by eluting 2% MeOH/DCM to obtained compound C-30 (800 mg, 83.6%) as pale yellow liquid.

¹H-NMR: (500 MHz, DMSO-d₆): δ 8.78-8.75 (m, 2H), 7.42-7.39 (m, 1H), 5.01-4.92 (m, 1H), 4.21-4.09 (m, 1H), 4.05-4.02 (m, 2H), 1.98 (s, 2H), 1.44-1.31 (m, 7H), 1.29-1.11 (m, 10H), 1.02-0.96 (m, 4H), 0.70-0.63 (m, 9H), -0.25 (s, 3H), -0.32 (s, 3H).

LCMS (ESI) : 523.6 [M⁺+1].

Synthesis of tert-butyl 2-(2-((tert-butyldimethylsilyl) oxy)-1-(pyrimidin-2-yl) propyl)-1, 6-dimethyl-3-oxo-2, 5-diazaspiro [3.4] octane-5-carboxylate (C-31):

[00192] To a stirring solution of triphenylphosphine (1.23 g, 4.87 mmol) in dry THF (10 mL) was added DTAD (1.09 g, 4.77 mmol) as portion-wise and stirred for 15 min at RT. To this precipitated solution, added compound C-30 (1.0 g, 1.91 mmol) in dry THF (10 mL) slowly at RT and stirred for 16 h. After consumption of the starting material (by TLC), the reaction mixture was concentrated under reduced pressure. The crude material was triturated with *n*-pentane (20 mL) and filtered solid (TPPO). The filtrate was concentrated under reduced pressure to obtained crude compound was purified by silica gel column chromatography eluting 30% EtOAc/hexane to afford compound C-31 (400 mg, 41.4%) as yellow liquid.

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 8.82-8.79 (m, 2H), 7.44-7.42 (m, 1H), 3.88-3.70 (m, 2H), 1.82-1.78 (m, 2H), 1.49-1.41 (m, 4H), 1.40 (s, 9H), 1.21-1.08 (m, 9H), 0.57 (s, 6H), - 0.05 (s, 9H); LCMS (ESI): 505.5 [M⁺+1].

Synthesis of tert-butyl 2-(2-hydroxy-1-(pyrimidin-2-yl) propyl)-1, 6-dimethyl-3-oxo-2, 5-diazaspiro [3.4] octane-5-carboxylate (C-32):

- 71 -

[00193] To a stirring solution of compound **C-31** (220 mg, 0.43 mmol) in dry THF (3 mL) was added TBAF (227 mg) slowly at 0 °C and stirred at RT for 16 h. After consumption of the starting material (by TLC), the reaction mixture was concentrated under reduced pressure to afford crude, which was purified by silica gel column chromatography eluting 80% EtOAc/hexane to afford compound **C-32** (150 mg, 88.2%) as yellow liquid.

¹H-NMR: (500 MHz, DMSO-*d*₆): δ 8.81-8.79 (m, 2H), 7.43-7.41 (m, 1H), 4.36-4.33 (m, 1H), 4.27-4.24 (m, 1H), 3.86-3.80 (m, 2H), 2.15-1.86 (m, 4H), 1.53-1.50 (m, 1H), 1.43 (s, 9H), 1.19-1.03 (m, 9H).


10 **Synthesis of 2-(2-hydroxy-1-(pyrimidin-2-yl) propyl)-3, 6-dimethyl-2, 5-diazaspiro [3.4] octan-1-one (C-33):**

[00194] To a stirring solution of compound **C-32** (120 mg, 0.25 mmol) in DCM (10 mL) was added molecular sieves (100 mg), BF₃(OEt)₂ (72 mg, 0.51 mmol) at 0 °C under N₂ atmosphere. The reaction mixture was stirred at RT for 10 min. After consumption of the starting material (by TLC), the obtained precipitate was triturated with *n*-pentane/diethylether (5 mL/5 mL) and the filtered solid was dried on vacuum to afford **C-33** (80 mg, 89.8%) as white solid.

¹H-NMR: (400 MHz, D₂O): δ 8.90 (t, *J* = 4.8 Hz, 2H), 7.64-7.60 (m, 1H), 4.72-4.62 (m, 2H), 4.24-4.15 (m, 1H), 4.08-4.00 (m, 1H), 2.60-2.32 (m, 3H), 2.02-1.91 (m, 1H), 1.55-1.51 (m, 3H), 1.48-1.45 (m, 3H), 1.39-1.25 (m, 3H).

LCMS (ESI): 581.3 [2M⁺+1].

Scheme 10



Synthesis of tert-butyl 2-((2-((tert-butyldimethylsilyl) oxy)-1-(pyrimidin-2-yl) propyl) carbamoyl)-2-(1-hydroxyethyl) pyrrolidine-1-carboxylate (C-34):

[00195] To a stirring solution of compound **BB** (2.5 g, 9.65 mmol) in CH₂Cl₂ (25 mL) were added DIPEA (5 mL, 28.95 mmol), intermediate **F** (2.57 g, 9.65 mmol) followed by HATU (4.4 g, 11.58 mmol) at 0 °C and stirred for 12 h at RT. After consumption of the starting material (by TLC), the reaction mixture was the reaction was diluted with water (20 mL). The separated organic layer was washed with citric acid solution (1 x 50 mL) followed by brine solution (1 x 50 mL). The organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. Obtained crude material was purified by silica gel column chromatography eluting 50% EtOAc/*n*-hexane to afford compound **C-34** (3.1 g, 63.2%) as an off-white solid.

¹H-NMR: (500 MHz, DMSO-d₆): δ 8.76 (d, *J* = 13.5 Hz, 2H), 8.42 (d, *J* = 9.0 Hz, 1H), 7.41 (d, *J* = 9.0 Hz, 1H), 4.60-4.41 (m, 1H), 4.23-4.11 (m, 1H), 3.58-3.40 (m, 3H), 2.37-1.62 (m, 4H), 1.39 (s, 9H), 1.28-1.19 (m, 3H), 1.18-1.15 (m, 3H), 1.09 (s, 9H), -0.04 (s, 6H);

LCMS (ESI): *m/z* 509.7 [M⁺+1]

Synthesis of tert-butyl 2-(2-((tert-butyldimethylsilyl) oxy)-1-(pyrimidin-2-yl) propyl)-1-methyl-3-oxo-2, 5-diazaspiro [3.4] octane-5-carboxylate (C-35):

[00196] To a stirring solution of triphenylphosphine (1.6 g, 6.29 mmol) in THF (15 mL) was added DIAD (1.44 g, 6.29 mmol) at RT and stirred for 30 min. To this added compound **C-34** (1.6 g, 3.14 mmol) in (5 mL) THF slowly and reaction mixture was stirred at RT for 3 h. After consumption of the starting material (by TLC), the reaction was concentrated under reduced pressure. The crude material was triturated with diethyl ether/*n*-pentane (10 mL/10 mL). The filtered solvent was concentrated and purified by silica gel column chromatography eluting 20% EtOAc/*n*-hexane to afford compound **C-35** (1 g, 65.3%) as pale yellow liquid.

¹H-NMR: (500 MHz, DMSO-d₆): δ 8.80 (d, *J* = 14.5 Hz, 2H), 7.44 (d, *J* = 9.0 Hz, 1H), 4.72-4.46 (m, 2H), 3.36-3.23 (m, 3H), 2.06-1.61 (m, 4H), 1.40 (s, 9H), 1.35-1.23 (m, 3H), 1.16-1.12 (m, 3H), 1.00 (s, 9H), -0.01 (s, 6H);

LCMS (ESI): *m/z* 491.4 [M⁺+1]

Synthesis of 2-(2-hydroxy-1-(pyrimidin-2-yl) propyl)-3-methyl-2, 5-diazaspiro [3.4] octan-1-one (C-36):

- 73 -

[00197] To a stirring solution of compound C-35 (1 g, 2.04 mmol) in DCM (5 mL) was added $\text{BF}_3\text{O}(\text{C}_2\text{H}_5)_2$ (0.05 mL, 4.08 mmol) followed by molecular sieves (50 mg) at 0 °C and stirred to RT for 3 h. After consumption of the starting material (by TLC), the reaction was diluted with *n*-pentane and filtered the obtained solid. The filtered solid was purified by silica gel column chromatography eluting 5% MeOH/DCM to afford **C-36** (180 mg, 32%) as an off-white solid.

¹H-NMR: (400 MHz, D₂O): δ 8.89 (d, *J* = 4.8 Hz, 2H), 7.61 (t, *J* = 4.8 Hz, 1H), 4.85-4.73 (m, 2H), 4.18-4.13 (m, 1H), 3.54-3.41 (m, 2H), 2.43-2.36 (m, 1H), 2.31-2.12 (m, 3H), 1.43-1.37 (m, 6H)

10 LCMS (ESI): *m/z* 277.3 [*M*⁺+1];

HPLC: 98.85% .

Synthesis of 2-(2-hydroxy-1-(pyrimidin-2-yl) propyl)-5-isobutyryl-3-methyl-2, 5-diazaspiro [3.4] octan-1-one (C-37):

15 [00198] To a stirring solution of compound **C-36** (600 mg, 2.17 mmol) in CH₂Cl₂ (10 mL) was added TEA (1.05 mL, 7.59 mmol) at 0 °C. After added isobutyryl chloride (253 mg, 2.39 mmol) slowly and stirred for 1 h at RT. After consumption of the starting material (by TLC), the reaction mixture was diluted with water (10 mL). The separated organic layer was washed with brine solution (1 x 50 mL). The organic layer was dried over anhydrous Na₂SO₄,
20 filtered and concentrated under reduced pressure. Obtained crude material was purified by silica gel column chromatography eluting 3% MeOH/DCM followed by preparative HPLC purification to afford **C-37** (120 mg, 16%) as an off-white solid.

¹H-NMR: (500 MHz, DMSO-*d*₆): δ 8.81 (d, *J* = 7.5 Hz, 2H), 7.43 (d, *J* = 11.5 Hz, 1H), 5.05-4.34 (m, 2H), 3.64-3.44 (m, 3H), 2.01-1.81 (m, 5H), 1.35-1.13 (m, 6H), 1.10-1.04 (m, 6H);


25 LCMS (ESI): *m/z* 347.4 [*M*⁺+1]

HPLC: 91.1%

Preparation of key intermediates, schemes I-8 to I-10:

Scheme I-8

- 74 -



Synthesis of ethyl 2-((benzyloxy) methyl)-5-methylpyrrolidine-2-carboxylate (EE):

[00199] To a stirring solution of **X** (8.5 g, 22.51 mmol) in DCM (50 mL) was added TFA (8.6 mL, 112.58 mmol) at 0 °C. The reaction mixture was stirred at RT for 4 h. After consumption of the starting material (by TLC), the reaction mixture was concentrated under reduced pressure to afford **EE** (14 g, crude) as pink liquid (TFA salt).

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 9.71 (br s, 1H), 7.45-7.30 (m, 5H), 4.62 (s, 2H), 4.27-4.22 (m, 2H), 3.90-3.73 (m, 3H), 2.26-2.08 (m, 2H), 2.01-1.95 (m, 1H), 1.59-1.51 (m, 1H), 1.36-1.31 (m, 3H), 1.22-1.18 (m, 3H);

LCMS (ESI): *m/z* 278.36 [*M*⁺+1]

Synthesis of ethyl 2-(hydroxymethyl)-5-methylpyrrolidine-2-carboxylate (FF):

[00200] To a stirring solution of **EE** (14 g (crude), 35.80 mmol) in methanol (100 mL) was added (50% wet) 10% Pd/C (5 g) under N₂ atmosphere. The reaction mixture was stirred under H₂ atmosphere (balloon pressure) at RT for 16 h. After consumption of the starting material (by TLC), the reaction mixture was filtered through a pad of celite and the pad was washed with methanol (100 mL). Obtained filtrate was concentrated under reduced pressure to afford **FF** (8 g, crude) as yellow syrup.

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 9.52 (s, 1H), 8.74 (s, 1H), 4.28-4.20 (m, 2H), 3.94-3.69 (m, 3H), 2.22-2.16 (m, 2H), 2.13-1.92 (m, 1H), 1.60-1.53 (m, 1H), 1.32-1.18 (m, 6H);

LCMS (ESI): *m/z* 188.24 [*M*⁺+1]

Synthesis of ethyl 1-benzyl-2-(hydroxymethyl)-5-methylpyrrolidine-2-carboxylate (GG):

[00201] To a stirring solution of **FF** (8 g, 26.57 mmol) in CH₃CN (50 mL) was added K₂CO₃ (11.02 g, 79.73 mmol) followed by benzyl bromide (3.78 mL, 31.89 mmol) at RT and stirred for 16 h. After consumption of the starting material, filtered the reaction mass through celite bed and filtrate was washed with EtOAc (250 mL). The filtrate was concentrated under

- 75 -

reduced pressure to afford crude. Obtained crude material was purified by column chromatography eluting with 10% EtOAc/Hexane to afford **GG** (5 g, 68%) as brown syrup.

¹H-NMR: (500 MHz, DMSO-*d*₆): δ 7.33-7.15 (m, 5H), 5.17 (t, *J* = 5.5 Hz, 1H), 4.75 (d, *J* = 5.5 Hz, 1H), 4.50 (d, *J* = 6.0 Hz, 1H), 4.14-3.93 (m, 2H), 3.93 (s, 2H), 3.81-3.60 (m, 1H), 3.13-3.09 (m, 1H), 2.20-2.17 (m, 1H), 1.99-1.88 (m, 2H), 1.40-1.36 (m, 3H), 1.14 (t, *J* = 7.0 Hz, 3H)

LCMS (ESI): *m/z* 278.36 [*M*⁺+1]


Synthesis of 1-benzyl-2-(hydroxymethyl)-5-methylpyrrolidine-2-carboxylic acid (HH):

[00202] To a stirring solution of **GG** (5 g, 18.05 mmol) in EtOH/H₂O (25 mL/25 mL) were added NaOH (1.44 g, 36.1 mg) at RT. The reaction mixture was heated to 100 °C for 2 h. After consumption of the starting material (by TLC), the solvent from the reaction was evaporated under reduced pressure and extracted with di ethylether (2 x 75 mL). The aqueous layer was acidified by using 1*N* HCl and extracted with 20% MeOH/CH₂Cl₂ (2 x 100 mL). The organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to afford **HH** (3.5 g, 78%) as brown syrup.

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 11.17 (s, 1H), 7.49-7.27 (m, 5H), 4.55 (d, *J* = 14.0 Hz, 1H), 4.06 (d, *J* = 14.4 Hz, 1H), 3.83 (s, 2H), 3.51-3.41 (m, 2H), 2.18-1.90 (m, 3H), 1.45-1.39 (m, 1H), 0.61-0.45 (m, 3H)

LCMS (ESI): *m/z* 250.13 [*M*⁺+1]

20 Scheme I-9



Synthesis of methyl pyrrolidine-2-carboxylate (II):

[00203] To a stirring solution of L-Proline (50 g, 434 mmol) in methanol was added thionyl chloride (37.5 ml, 521 mmol) at 0 °C and heated to 70 °C for 16 h. The reaction

- 76 -

mixture was brought to RT and concentrated under vacuum to afford II as (70 g, 99 %) as thick syrup (hydrochloride salt).

¹H-NMR: (500 MHz, DMSO-*d*₆): δ 4.15-4.13 (m, 1H), 3.65 (s, 3H), 3.35-3.30 (m, 2H), 2.23-2.15 (m, 1H), 1.86-1.78 (m, 3H), 1.41 (s, 9H)

5 **LCMS (ESI):** *m/z* 129 [*M*⁺+1]

Synthesis of methylbenzylproline (JJ):

[00204] To a stirring solution of compound II (18 g, 108 mmol) in DCM (200 mL) was added TEA (45.35 mL, 326 mmol) followed by benzyl bromide (15.5 mL, 130 mmol) at 0 °C and stirred at RT for 24 h. After completion of the reaction (by TLC) was diluted with water (75 mL) and EtOAc (500 mL). The organic layer was washed with water (2 x 100 mL), brine solution (2 x 50 mL). The organic layer was dried over anhydrous Na₂SO₄ and concentrated under vacuum. Obtained crude material was purified by column chromatography eluting with 10% EtOAc/Hexane to afford JJ (5 g, 21%) as oily liquid.

10 **¹H-NMR:** (400 MHz, DMSO-*d*₆): δ 7.32-7.22 (m, 5H), 3.86-3.80 (m, 1H), 3.58 (s, 3H), 3.48-3.24 (m, 2H), 2.85-2.80 (m, 1H), 2.38-2.32 (m, 1H), 2.10-1.98 (m, 1H), 1.85-1.69 (m, 3H)

15 **LCMS (ESI):** *m/z* 220.28 [*M*⁺+1]

Synthesis of methyl 1-benzyl-2-(1-hydroxyethyl) pyrrolidine-2-carboxylate (KK):

[00205] To a stirring solution of JJ (5 g, 22.80 mmol) in THF (50 mL) was added LiHMDS (46 mL, 45.60 mmol) dropwise at -20 °C and stirred for 45 min. To this acetaldehyde (2.33 mL, 45.60 mmol) was added drop wise and stirred for 3 h at -20 °C. After consumption of the starting material (by TLC), the reaction was quenched with aqueous NH₄Cl solution (100 mL) and extracted with EtOAc (2 x 150 mL). The separated organic layer was washed with water (200 mL) and brine solution (200 mL). The separated organic layer was dried over Na₂SO₄ and concentrated to afford crude which was purified by column chromatography eluting with 10% EtOAc/Hexane to afford KK (3.5 g, 58%) as yellow syrup.

25 **LCMS (ESI):** *m/z* 264.3 [*M*⁺+1]

Synthesis of 1-benzyl-2-(1-hydroxyethyl) pyrrolidine-2-carboxylic acid (LL):

[00206] To a stirring solution of KK (4.5 g, 17.11 mmol) in MeOH/THF/H₂O (10 mL/10 mL/10 mL) were added NaOH (1.02 g, 25.66 mmol) at RT. The reaction mixture was heated to 95 °C for 1 h. After consumption of the starting material (by TLC), the solvent was evaporated

30


under reduced pressure. The aqueous layer was washed with EtOAc (100 mL). The separated aqueous layer was acidified by using 1N HCl (pH~3). The aqueous layer was extracted with 20% MeOH/DCM (2 x 100 mL). The combined organic layer was dried over Na₂SO₄ and concentrated under reduced pressure to afford **LL** (2.7 g, 63%) as an off-white solid.

- 5 ¹H-NMR: (400 MHz, DMSO-*d*₆): δ 7.31-7.13 (m, 5H), 3.87-3.78 (m, 2H), 3.61-3.57 (m, 1H), 2.75-2.64 (m, 2H), 1.90-1.78 (m, 2H), 1.69-1.65 (m, 1H), 1.53-1.43 (m, 1H), 1.06 (d, *J* = 6.4 Hz, 3H);

LCMS (ESI): *m/z* 250.4 [M⁺-1]

Scheme I-10

10



Synthesis of methyl L-serinate (MM):

- 15 **[00207]** To a stirring solution of L-serine (12 g, 114 mmol) in CH₃OH (100 mL) was added thionyl chloride (10 mL, 137 mmol) at 0 °C and stirred at 80 °C for 16 h. After completion of starting material (by TLC), the reaction mixture was concentrated under reduced pressure to afford **MM** (16 g, crude, HCl salt) as white solid. This material was directly used for the next step without further purification.

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 8.61 (s, 2H), 4.50 (s, 1H), 4.20-4.16 (m, 1H), 3.82 (d, *J* = 3.6 Hz, 2H), 3.73 (s, 3H)

- 20 **[00208]** Synthesis of methyl ((benzyloxy)carbonyl)-L-serinate (NN):

- [00208]** To a stirring solution of **MM** (16 g, 103 mmol) in water/1, 4 dioxane (130 mL/65 mL) were added Na₂CO₃ (27.3 g, 257 mmol) at RT. After added Cbz-Cl (17.6 mL, 123 mmol) was added at 0 °C drop wise and stirred for 16 h at RT. After completion of starting material (by TLC), diluted the reaction mass with EtOAc (300 ml). The separated organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated under reduced
- 25

- 78 -

pressure to afford crude material which was purified by column chromatography to afford **NN** (23 g, 88 %) as thick syrup.

¹H-NMR: (500 MHz, DMSO-*d*₆): δ 7.50 (d, *J* = 8.0 Hz, 1H), 7.38-7.31 (m, 5H), 5.04 (s, 2H), 4.95-4.92 (m, 1H), 4.17-4.13 (m, 1H), 3.66 (s, 3H), 3.65-3.56 (m, 2H).

5 **LCMS (ESI):** *m/z* 254.2[M⁺+1]

Synthesis of methyl N-((benzyloxy)carbonyl)-O-(tert-butyldimethylsilyl)-L-serinate (OO):

[00209] To a stirring solution of **NN** (23 g, 91 mmol) in DCM (700 mL) were added imidazole (12.37 g, 182 mmol), DMAP (2.22 g, 18.2 mmol) followed by TBDMS-Cl (20.4 g, 136 mmol) at 0 °C and stirred at RT for 16 h. After completion of starting material (by TLC),
 10 diluted the reaction mass with water (200 ml). The separated organic layer was washed with brine solution (2 x 200 mL). The organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to afford crude material was purified by column chromatography eluting 20% EtOAc/hexane to afford **OO** (32 g, 97%) as thick syrup.

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 7.56 (d, *J* = 8.0 Hz, 1H), 7.38-7.30 (m, 5H), 5.07 (s, 2H),
 15 4.24-4.19 (m, 1H), 3.81 (d, *J* = 5.2 Hz, 2H), 3.63 (s, 3H), 0.84 (s, 9H), -0.04 (s, 6H)

LCMS (ESI): *m/z* 368.5 [M⁺+1]

Synthesis of benzyl (S)-(3-((tert-butyldimethylsilyl)oxy)-1-hydrazinyl-1-oxopropan-2-yl)carbamate (PP):

[00210] To a stirred solution of **OO** (20 g, 54.5 mmol) in methanol (120 mL) was added
 20 hydrazine hydrate (27 g, 545 mmol) at 0 °C and after stirred at 80 °C for 2 h. After completion of starting material (by TLC), ethanol was evaporated under reduced pressure. The crude residue was triturated with *n*-pentane (100 mL) to afford **PP** (17 g, 85%) as an off-white solid.

¹H-NMR: (500 MHz, DMSO-*d*₆): δ 9.17 (s, 1H), 9.10 (s, 1H), 7.35-7.14 (m, 5H), 5.02 (s, 2H), 4.48 (s, 2H), 4.16-4.07 (m, 2H), 3.73-3.58 (m, 1H), 0.81 (s, 9H), -0.04 (s, 6H).

25 **Synthesis of benzyl ((1S, 2R)-2-((tert-butyldimethylsilyl)oxy)-1-(1, 3, 4-oxadiazol-2-yl)propyl) carbamate (QQ):**

[00211] A solution of **PP** (17 g, 46.3 mmol) in triethylorthoformate (68.5 g, 463 mmol) was added *p*-TSA (8.8 g, 4.63 mmol) at RT and after stirred at 80 °C for 2 h. After completion of starting material (by TLC), triethylorthoformate was evaporated under reduced pressure. The
 30 crude residue was purified by column chromatography eluting 10% EtOAc/hexane to afford **QQ** (5 g, 29%) as thick syrup.

- 79 -

¹H-NMR: (500 MHz, DMSO-*d*₆): δ 9.22 (s, 1H), 8.05 (d, *J* = 7.5 Hz, 1H), 7.36-7.30 (m, 5H), 5.08 (s, 2H), 5.05-4.97 (m, 1H), 4.03-3.91 (m, 2H), 0.84 (s, 9H), 0.03 (s, 6H);

LCMS (ESI): *m/z* 378.5 [*M*⁺+1]


Synthesis of (S)-2-((*tert*-butyldimethylsilyl) oxy)-1-(1, 3, 4-oxadiazol-2-yl) ethan-1-amine (RR):

[00212] To a stirring solution of **QQ** (5 g, 13.2 mmol) in methanol (30 mL) was added 50% wet 10% Pd/C (1.2 g) and stirred under H₂ atmosphere (balloon pressure) for 2 h at RT. After completion of reaction, the reaction mixture was filtered through a pad of celite and triturated with methanol (20 mL). The filtrate was concentrated under reduced pressure to afford **RR** (2 g, 63%) as thick syrup.

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 9.16 (s, 1H), 4.19-4.16 (m, 1H), 3.85-3.75 (m, 2H), 2.11 (s, 2H), 0.77 (s, 9H), -0.04 (s, 6H)

LCMS (ESI): *m/z* 244.3 [*M*⁺+1]

Scheme 11



Synthesis of 2-(2-hydroxy-1-(pyrimidin-2-yl) propyl)-5-isobutyryl-6-methyl-2, 5-diazaspiro [3.4] octan-1-one (C-38):

[00213] To a stirring solution of **C-20** (500 mg, 1.81 mmol) in DCM (10 mL) was added DIPEA (0.94 mL, 5.43 mmol) at 0 °C. After added iso butyrylchloride (578 mg, 5.43 mmol) at 0 °C and stirred at RT for 4 h. After consumption of the starting material (by TLC), the reaction mixture was diluted with water (10 mL). The separated organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford crude compound which was purified by column chromatography by eluting 2% MeOH/DCM to obtained **C-38** (150 mg, 24%) as pale yellow solid.


¹H-NMR: (400 MHz, CD₃OD): δ 8.82-8.77 (m, 2H), 7.39 (t, *J* = 5.2 Hz, 1H), 4.76 (d, *J* = 8.0 Hz, 1H), 4.47-4.44 (m, 1H), 4.20-4.14 (m, 1H), 3.82-3.74 (m, 1H), 3.51-3.40 (m, 1H), 2.78-2.72 (m, 1H), 2.46-2.37 (m, 1H), 2.18-2.12 (m, 2H), 1.77-1.72 (m, 1H), 1.34-1.29 (m, 3H), 1.26-1.22 (m, 3H), 1.06-1.02 (m, 6H)

- 80 -

LCMS (ESI): m/z 347.2 [$M^+ + 1$];

HPLC: 99.7%

Scheme 12



5 **Synthesis of 5-(3,3-dimethylbutanoyl)-2-(2-hydroxy-1-(pyrimidin-2-yl)propyl)-6-methyl-2,5-diazaspiro[3.4]octan-1-one (C-39):**


[00214] To a stirring solution of **C-20** (150 mg, 0.54 mmol) in DCM (10 mL) was added TEA (137 mg, 1.35 mmol) at 0 °C. After added 3, 3-dimethylbutanoyl chloride (87 mg, 0.65 mmol) at 0 °C and stirred at RT for 1 h. After consumption of the starting material (by TLC),
 10 the reaction mixture was diluted with water (10 mL). The separated organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford crude compound which was purified by column chromatography by eluting 2% MeOH/DCM to obtained **C-39** (150 mg, 24%) as sticky solid.

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 8.80-8.75 (m, 2H), 7.43-7.37 (m, 1H), 4.73-4.65 (m, 2H),
 15 4.24-4.17 (m, 2H), 3.78-3.59 (m, 1H), 3.23-3.16 (m, 1H), 2.36 (s, 2H), 2.32-2.08 (m, 3H), 1.58-1.50 (m, 1H), 1.20-1.08 (m, 6H), 0.86 (s, 9H)

LCMS (ESI): m/z 375.49 [$M^+ + 1$];

HPLC: 96.9%

20 Scheme 13



Synthesis of cyclopentyl 2-(2-hydroxy-1-(pyrimidin-2-yl) propyl)-6-methyl-1-oxo-2, 5-diazaspiro [3.4] octane-5-carboxylate (C-40):

[00215] To a stirring solution of **C-20** (150 mg, 0.54 mmol) in DCM (5 mL) was added
 25 TEA (0.23 mL, 1.63 mmol) at 0 °C. After added cyclopentyl chloroformate (88 mg, 0.59

- 81 -


mmol) at 0 °C and stirred at RT for 1 h. After consumption of the starting material (by TLC), the reaction mixture was diluted with water (10 mL). The separated organic layer was washed with saturated brine solution (20 mL). The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford crude compound which was purified by column chromatography by eluting 3% MeOH/DCM to obtained **C-40** (110mg, 52%) as sticky solid.

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 8.81-8.76 (m, 2H), 7.45-7.42 (m, 1H), 5.20-4.91 (m, 1H), 4.77-4.64 (m, 2H), 4.30-3.63 (m, 1H), 3.57-3.31 (m, 3H), 2.16-1.86 (m, 4H), 1.77-1.32 (m, 8H), 1.18-1.08 (m, 6H)

LCMS (ESI): *m/z* 389.4 [M⁺+1];

UPLC: 97.6%

Scheme 14



Synthesis of cyclohexyl 2-(2-hydroxy-1-(pyrimidin-2-yl) propyl)-6-methyl-1-oxo-2, 5-diazaspiro [3.4] octane-5-carboxylate (C-41):

[00216] To a stirring solution of **C-20** (150 mg, 0.54 mmol) in DCM (5 mL) was added TEA (0.2 mL, 1.35 mmol) at 0 °C. After added cyclohexyl chloroformate (105 mg, 0.65 mmol) at 0 °C and stirred at RT for 1 h. After consumption of the starting material (by TLC), the reaction mixture was diluted with water (10 mL). The separated organic layer was washed with citric acid solution (20 mL), saturated brine solution (20 mL). The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford crude compound which was purified by column chromatography by eluting 2% MeOH/DCM to obtained **C-41** (90 mg, 41%) as sticky solid.


¹H-NMR: (400 MHz, DMSO-*d*₆): δ 8.82-8.78 (m, 2H), 7.48-7.41 (m, 1H), 4.78-4.61 (m, 2H), 4.24-4.13 (m, 2H), 3.94-3.87 (m, 1H), 3.76-3.53 (m, 2H), 2.27-1.99 (m, 3H), 1.97-1.92 (m, 1H), 1.56-1.32 (m, 10H), 1.29-1.10 (m, 2H)

LCMS (ESI): *m/z* 403.5 [M⁺+1];

UPLC: 98.06%

Scheme 15

- 82 -



Synthesis of 1-benzyl-N-((1*S*, 2*R*)-2-((*tert*-butyldimethylsilyl) oxy)-1-(1, 3, 4-oxadiazol-2-yl) propyl)-2-(hydroxymethyl)-5-methylpyrrolidine-2-carboxamide (C-42):

[00217] To a stirring solution of HH (2.5 g, 10.04 mmol) in DMF (15 mL) were added
 5 *N*, *N*-diisopropylethylamine (5.4 mL, 30.12 mmol), Int N (2.58 g, 10.04 mmol) followed by
 HATU (4.57 g, 12.04 mmol) at 0 °C and stirred at RT for 16 h. After consumption of the
 starting material (by TLC), the reaction mixture was diluted with water (100 mL) and EtOAc
 (100 mL). The organic layer was washed with water (100 mL), saturated sodium bicarbonate
 solution (50 mL) followed by brine solution (50 mL). The organic layer was dried over
 10 anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to afford crude compound
 which was purified by column chromatography by eluting 20% EtOAc/*n*-hexane to obtained C-
 42 (3.2 g, 65%) as brown syrup.

¹H-NMR: (500 MHz, DMSO-*d*₆): δ 9.26 (s, 1H), 8.78 (d, *J* = 10.0 Hz, 1H), 7.30-7.21 (m, 5H),
 5.26 (d, *J* = 10.0 Hz, 1H), 4.43 (s, 2H), 4.05-4.02 (m, 2H), 3.87-3.72 (m, 1H), 3.15-3.10 (m,
 15 1H), 2.81-2.65 (m, 1H), 1.39-1.26 (m, 4H), 1.25-1.16 (m, 6H), 0.05 (s, 6H)

LCMS (ESI): *m/z* 489.70 [M⁺+1]

Synthesis of 5-benzyl-2-((1*S*, 2*R*)-2-((*tert*-butyldimethylsilyl) oxy)-1-(1, 3, 4-oxadiazol-2-yl) propyl)-6-methyl-2, 5-diazaspiro [3.4] octan-1-one (C-43):

[00218] To a stirring solution of triphenylphosphine (4.02 g, 15.36 mmol) in dry THF
 20 (30 mL) was added DIAD (2.48 g, 12.29 mmol) at RT and stirred for 15 min. After added C-42
 (3 g, 6.14 mmol) in THF (30 mL) was added dropwise and the reaction mixture was stirred at
 RT for 4 h. After consumption of the starting material (by TLC), the reaction mixture was
 concentrated under reduced pressure. The crude material was purified by column
 chromatography by eluting 20% EtOAc/*n*-hexane to afford C-43 (1.3 g, 45%) as pale green
 25 syrup.

- 83 -

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 9.28 (s, 1H), 7.33-7.19 (m, 5H), 5.24-5.19 (m, 1H), 4.73 (s, 2H), 4.47-4.41 (m, 1H), 4.02 (t, *J* = 7.2 Hz, 1H), 3.68 (s, 2H), 2.49-2.28 (m, 1H), 2.27-2.10 (m, 1H), 1.98-1.94 (m, 1H), 1.47-1.40 (m, 1H), 1.23-1.15 (m, 6H), 0.87 (s, 9H), 0.02 (s, 6H)

LCMS (ESI): *m/z* 491.6 [*M*⁺+1]

5 **Synthesis of 5-benzyl-2-((1*S*, 2*R*)-2-hydroxy-1-(1, 3, 4-oxadiazol-2-yl) propyl)-6-methyl-2, 5-diazaspiro [3.4] octan-1-one (C-44):**

[00219] To a stirring solution of C-43 (1.3 g, 2.76 mmol) in dry THF (30 mL) was added TBAF (1M in THF) (4.13 mL, 4.14 mmol) at 0 °C and stirred for 2 h at RT. After consumption of the starting material (by TLC), the reaction mixture was concentrated under reduced
10 pressure. The crude material was diluted with water (100 mL) and EtOAc (100 mL). The organic layer was washed with water (100 mL), brine solution (50 mL). The organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to afford crude compound which was purified by column chromatography by eluting 2% MeOH/CH₂Cl₂ to obtained C-44 (350 mg, 35%) as thick syrup.

15 ¹H-NMR: (400 MHz, DMSO-*d*₆): δ 9.18 (s, 1H), 7.34-7.16 (m, 5H), 5.21 (t, *J* = 5.6 Hz, 1H), 5.01-4.83 (m, 1H), 4.22-4.18 (m, 1H), 3.88 (s, 2H), 3.84-3.67 (m, 1H), 3.54-3.35 (m, 1H), 2.98-2.89 (m, 1H), 2.31-1.89 (m, 3H), 1.45-1.41 (m, 1H), 1.39-1.14 (m, 3H), 0.90-0.81 (s, 3H)

LCMS (ESI): *m/z* 357.43 [*M*⁺+1]

HPLC: 92.46%

20 **Synthesis of 5-benzyl-2-((1*S*, 2*R*)-2-hydroxy-1-(1, 3, 4-oxadiazol-2-yl) propyl)-6-methyl-2, 5-diazaspiro [3.4] octan-1-one (C-45):**

[00220] To a stirring solution of C-44 (1.3 g, 2.76 mmol) in dry THF (30 mL) was added TBAF (1M in THF) (4.13 mL, 4.14 mmol) at 0 °C and stirred for 2 h at RT. After consumption of the starting material (by TLC), the reaction mixture was concentrated under reduced
25 pressure. The crude material was diluted with water (100 mL) and EtOAc (100 mL). The organic layer was washed with water (100 mL), brine solution (50 mL). The organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to afford crude compound which was purified by column chromatography by eluting 2% MeOH/CH₂Cl₂ to obtained C-45 (350 mg, 35%) as thick syrup.


- 84 -

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 9.18 (s, 1H), 7.34-7.16 (m, 5H), 5.21 (t, *J* = 5.6 Hz, 1H), 5.01-4.83 (m, 1H), 4.22-4.18 (m, 1H), 3.88 (s, 2H), 3.84-3.67 (m, 1H), 3.54-3.35 (m, 1H), 2.98-2.89 (m, 1H), 2.31-1.89 (m, 3H), 1.45-1.41 (m, 1H), 1.39-1.14 (m, 3H), 0.90-0.81 (s, 3H)

LCMS (ESI): *m/z* 357.43 [*M*⁺+1]

5 HPLC: 92.46%

Scheme 16



Synthesis of 2-((1*S*,2*R*)-2-hydroxy-1-(1,3,4-oxadiazol-2-yl)propyl)-3-methyl-2,5-

diazaspiro[3.4]octan-1-one (C-46):

[00221] To a stirring solution of **C-26** (800 mg, 2.18 mmol) in DCM (10 mL) were added BF₃·OEt₂ (620 mg, 4.37 mmol), molecular sieves (150 mg) at 0 °C and stirred for 3 h at RT. After consumption of the starting material (by TLC), the reaction mixture was diluted with *n*-pentane (5 mL). The precipitated solid was filtered and filtrate was concentrated under reduced pressure. Obtained crude material was purified by silica gel column chromatography eluting with 3% MeOH/DCM to afford **C-46** (700 mg, crude) as yellow liquid.

Mass (ESI): *m/z* 267.3 [*M*⁺+1]

Synthesis of 2-((1*S*,2*R*)-2-hydroxy-1-(1,3,4-oxadiazol-2-yl)propyl)-5-isobutyryl-3-methyl-2,5-diazaspiro[3.4]octan-1-one (C-47):

[00222] To a stirring solution of **C-46** (700 mg (crude), 2.63 mmol) in DCM (10 mL) was added TEA (0.53 mL, 3.94 mmol) at 0 °C. After 10 min added isobutyryl chloride (0.41 mL, 3.94 mmol) at 0 °C and stirred for 2 h at RT. After consumption of the starting material (by TLC), the reaction mixture was diluted with water (5 mL). The separated organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. Obtained crude material was purified by silica gel column chromatography eluting with 3% MeOH/DCM followed by preparative HPLC purification to afford **C-47** (150 mg, 17%) as thick syrup.


- 85 -

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 9.27 (s, 1H), 5.20-5.17 (m, 1H), 4.87-4.85 (m, 1H), 4.30-4.25 (m, 1H), 3.84-3.79 (m, 1H), 3.64-3.50 (m, 2H), 2.72-2.66 (m, 1H), 2.13-2.09 (m, 1H), 2.01-1.79 (m, 3H), 1.37-1.19 (m, 3H), 1.10-0.94 (m, 9H)

Mass (ESI): *m/z* 337.4 [*M*⁺+1]

5 **HPLC:** 99.5%

Scheme 17



Synthesis of 1-benzyl-*N*-((1*S*, 2*R*)-2-((*tert*-butyldimethylsilyl) oxy)-1-(1, 3, 4-oxadiazol-2-yl) propyl)-2-(1-hydroxyethyl) pyrrolidine-2-carboxamide (C-48):

10 **[00223]** To a stirring solution of **LL** (3.9 g, 15.66 mmol) in DMF (10 mL) were added *N*, *N*-diisopropylethylamine (8.16 mL, 46.98 mmol), **Int-N** (4.03 g, 15.66 mmol) followed by HATU (7.14 g, 18.79 mmol) at 0 °C and stirred at RT for 16 h. After consumption of the starting material (by TLC), the reaction mixture was diluted with water (100 mL) and EtOAc (150 mL). The separated organic layer was washed with water (100 mL) followed by brine solution (100 mL). The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford crude compound which was purified by column chromatography by eluting 30% EtOAc/n-hexane to obtained **C-48** (1.8 g, 24%) as yellow syrup.

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 9.25 (s, 1H), 8.74 (d, *J* = 8.8 Hz, 1H), 7.34-7.21 (m, 5H), 5.47-5.27 (m, 1H), 5.26-5.23 (m, 1H), 4.42-4.40 (m, 1H), 4.13-4.01 (m, 2H), 3.82-3.77 (m, 1H), 2.70-2.60 (m, 2H), 1.99-1.82 (m, 2H), 1.69-1.64 (m, 2H), 1.29-1.21 (m, 6H), 0.72 (s, 9H), -0.01 (s, 3H), -0.02 (s, 3H);

LCMS (ESI): *m/z* 489.6 [*M*⁺+1]

Synthesis of 5-benzyl-2-((1*S*, 2*R*)-2-((*tert*-butyldimethylsilyl) oxy)-1-(1, 3, 4-oxadiazol-2-yl) propyl)-3-methyl-2, 5-diazaspiro [3.4] octan-1-one (C-49):

25 **[00224]** To a stirring solution of triphenylphosphine (2.41 g, 9.22 mmol) in dry THF (20 mL) was added DIAD (1.49 g, 7.37 mmol) as portionwise and stirred for 15 min at RT. To this

- 86 -

precipitated solution added **C-48** (1.8 g, 3.68 mmol) in dry THF (15 mL) dropwise at RT and stirred for 16 h. After consumption of the starting material (by TLC), the reaction mixture was concentrated under reduced pressure. The obtained solid was triturated with 20% di ethylether/*n*-hexane (2 x 100 mL). The filtrate was concentrated under reduced pressure to
5 obtained crude compound which was purified by silica gel column chromatography eluting 20% EtOAc/hexane to afford **C-49** (400 mg, crude) as pale green syrup.

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 9.30 (s, 1H), 7.40-7.24 (m, 5H), 5.26-5.22 (m, 1H), 4.80-4.73 (m, 2H), 4.62-4.48 (m, 1H), 4.15-4.00 (m, 1H), 3.81-3.75 (m, 1H), 2.85 (t, *J* = 8.0 Hz, 1H), 2.49-2.39 (m, 3H), 2.26-2.11 (m, 1H), 1.71-1.64 (m, 3H), 1.25-1.15 (m, 3H), 0.78 (s, 9H),
10 0.02 (s, 6H);

LCMS (ESI): *m/z* 471.6 [*M*⁺+1]

Synthesis of 5-benzyl-2-((1*S*, 2*R*)-2-hydroxy-1-(1, 3, 4-oxadiazol-2-yl) propyl)-3-methyl-2, 5-diazaspiro [3.4] octan-1-one (C-50):

[00225] To a stirring solution of **C-49** (1.5 g, 3.19 mmol) in dry THF (30 mL) was added
15 TBAF (1M in THF) (4.78 mL, 1.5 mmol) at 0 °C and stirred at RT for 1 h. After consumption of the starting material (by TLC), the reaction mixture was concentrated under reduced pressure. The crude material was diluted with water (75 mL) and EtOAc (100 mL). The organic layer was washed with water (2 x 50 mL), brine solution (2 x 50 mL). The organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to afford crude
20 compound which was purified by column chromatography by eluting 30% EtOAc/*n*-hexane to obtained **C-50** (500 mg, 44%) as white solid.

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 9.26 (s, 1H), 7.37-7.22 (m, 5H), 5.29-5.21 (m, 1H), 4.96-4.75 (m, 1H), 4.33-4.02 (m, 1H), 4.00-3.94 (m, 2H), 3.77-3.70 (m, 2H), 2.91-2.80 (m, 1H), 2.42-2.38 (m, 1H), 2.16-2.08 (m, 2H), 1.80-1.63 (m, 1H), 1.30 (d, *J* = 6.4 Hz, 3H), 1.12 (d, *J* =
25 6.0 Hz, 3H);

LCMS (ESI): *m/z* 357.43 [*M*⁺+1]

HPLC: 99.61%

Synthesis of 2-((1*S*,2*R*)-2-hydroxy-1-(1,3,4-oxadiazol-2-yl)propyl)-3-methyl-2,5-diazaspiro[3.4]octan-1-one (C-51):

[00226] To a stirring solution of compound **C-50** (420 mg, 1.17 mmol) in methanol (10 mL) was added 10%Pd/C (400 mg) under N₂ atmosphere. The reaction mixture was stirred under H₂ atmosphere at RT for 3 h. After consumption of the starting material (by TLC), the

- 87 -


reaction mixture was filtered through a pad of celite and the pad was washed with methanol (10 mL). Obtained filtrate was concentrated under reduced pressure to afford **C-51** (220 mg, 70%) as an off-white solid.

¹H-NMR: (500 MHz, DMSO-*d*₆): δ 9.26 (s, 1H), 5.25-5.17 (m, 1H), 4.89-4.69 (m, 1H), 4.28-4.21 (m, 1H), 3.84-3.56 (m, 2H), 2.95-2.79 (m, 2H), 1.99-1.84 (m, 2H), 1.77-1.59 (m, 2H), 1.24-1.10 (m, 6H);

LCMS (ESI): *m/z* 260 [*M*⁺+1]

HPLC: 97.94%

Scheme 18



Synthesis of 2-(2-hydroxy-1-(pyrimidin-2-yl) propyl)-3, 6-dimethyl-2, 5-diazaspiro [3.4] octan-1-one (C-52):

[00227] To a stirring solution of **C-33** (500 mg, 1.72 mmol) in dry DCM (10 mL) was added DIPEA (556 mg, 4.31 mmol) followed by isobutyryl chloride (273 mg, 2.58 mmol) slowly at 0 °C and stirred at RT for 3 h. After consumption of the starting material (by TLC), the reaction mixture was concentrated under reduced pressure. The crude residue was diluted with water (10 mL). The separated organic layer was washed with brine solution (20 mL). The separated organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford crude compound which was purified by column chromatography by eluting 1% MeOH/DCM to obtained **C-52** (90 mg, 14.5%) as pale brown syrup.


¹H-NMR: (400 MHz, CD₃OD): δ 8.81-8.77 (m, 2H), 7.41-7.37 (m, 1H), 4.63-4.53 (m, 2H), 4.24-4.18 (m, 2H), 2.87-2.72 (m, 1H), 2.33-2.27 (m, 1H), 2.16-2.08 (m, 2H), 1.73-1.69 (m, 1H), 1.40-1.35 (m, 3H), 1.28-1.20 (m, 3H), 1.15-1.02 (m, 9H);

LCMS (ESI): *m/z* 361.3 [*M*⁺+1];

HPLC: 94.85%

Scheme 19

- 88 -



Synthesis of ethyl 2-(1-hydroxyethyl)-5-methylpyrrolidine-2-carboxylate (C-53):

[00228] To a stirring solution of CC (5 g, 16.61 mmol) in CH_2Cl_2 (50 mL) was added
 5 TFA (6.34 mL, 83.05 mmol) at 0 °C. After being stirred at RT for 4 h, the reaction mixture was concentrated under reduced pressure to afford crude compound which was triturated with n-pentane (50 mL) to obtained C-53 (5 g, crude) as black syrup was directly taken for the next step without further purification.

¹H-NMR: (500 MHz, DMSO-*d*₆): δ 8.71 (br s, 1H), 4.29-4.25 (m, 1H), 4.25-4.21 (m, 3H),
 10 4.14-4.03 (m, 1H), 2.31-2.10 (m, 4H), 1.36-1.18 (m, 6H), 1.18-1.10 (m, 3H)

LCMS (ESI): *m/z* 202.38 [*M*⁺+1]

Synthesis of ethyl 1-benzyl-2-(1-hydroxyethyl)-5-methylpyrrolidine-2-carboxylate (C-54: C-54-A & C-54-B) :

[00229] To a stirring solution of C-53 (5 g (crude), 15.87 mmol) in acetonitrile (50 mL)
 15 was added K_2CO_3 (6.57 g, 47.61 mmol) followed by benzyl bromide (2.82 mL, 23.80 mmol) at RT and stirred for 16 h. After completion of the reaction, diluted the reaction mixture with EtOAc (150 mL) and water (75 mL). The organic layer was washed with brine solution (2 x 150 mL). The organic layer was dried over anhydrous Na_2SO_4 and concentrated under vacuum. Obtained crude material was purified by column chromatography eluting with 10%
 20 EtOAc/Hexane to afford 1 g of C-54-A and 1.2 g of C-54-B (separated isomers of C-54) (48%) as yellow syrups.

¹H-NMR (C-54-A isomer): (500 MHz, DMSO-*d*₆): δ 7.35-7.14 (m, 5H), 4.55 (s, 2H), 4.24-4.19 (m, 1H), 4.13-4.02 (m, 2H), 3.96-3.91 (m, 2H), 2.14-1.99 (m, 1H), 1.95-1.82 (m, 3H), 1.25-1.18 (m, 3H), 1.10-1.06 (m, 3H), 0.75 (d, *J* = 6.0 Hz, 3H)

LCMS (ESI): m/z 292.3 [$M^+ + 1$]

$^1\text{H-NMR}$ (C-54-B isomer): (500 MHz, DMSO- d_6): δ 7.34-7.18 (m, 5H), 5.15 (t, J = 5.5 Hz, 2H), 4.63-4.49 (m, 1H), 4.24-4.22 (m, 1H), 4.10-4.06 (m, 2H), 3.18-3.10 (m, 1H), 2.39-2.33 (m, 1H), 1.89-1.83 (m, 3H), 1.42-1.39 (m, 3H), 1.20-1.13 (m, 3H), 0.76 (d, J = 6.0 Hz, 3H)

5 LCMS (ESI): m/z 292.3 [$M^+ + 1$]

Synthesis of 1-benzyl-2-(1-hydroxyethyl)-5-methylpyrrolidine-2-carboxylic acid (C-55-A):

[00230] To a stirring solution of C-54-A (1 g, 3.43 mmol) in MeOH/THF/H₂O (5 mL/5 mL/5 mL) were added NaOH (206 mg, 5.14 mmol) at 0 °C. The reaction mixture was heated to 90 °C for 6 h. After consumption of the starting material (by TLC), the solvent was evaporated under reduced pressure. The aqueous layer was washed with di ethylether (50 mL). The separated aqueous layer was acidified by using 2N HCl (pH~3). The aqueous layer was extracted with 10% MeOH/DCM (2 x 100 mL). The combined organic layer was dried over Na₂SO₄ and concentrated under reduced pressure to afford compound C-55-A (1 g, crude) as yellow solid.

15 $^1\text{H-NMR}$: (400 MHz, DMSO- d_6): δ 9.84 (s, 1H), 7.60 (d, J = 6.8 Hz, 2H), 7.50-7.30 (m, 3H), 5.12 (br s, 1H), 4.48-4.35 (m, 2H), 4.09-3.81 (m, 1H), 2.39-2.22 (m, 2H), 2.16-2.08 (m, 2H), 1.66-1.57 (m, 1H), 1.16-0.86 (m, 6H);

LCMS (ESI): m/z 264.3 [$M^+ + 1$]

Synthesis of 1-benzyl-2-(1-hydroxyethyl)-5-methylpyrrolidine-2-carboxylic acid (C-55-B):

20 [00231] To a stirring solution of C-54-B (1.2 g, 4.12 mmol) in MeOH/THF/H₂O (5 mL/5 mL/5 mL) were added NaOH (206 mg, 5.14 mmol) at 0 °C. The reaction mixture was heated to 90 °C for 6 h. After consumption of the starting material (by TLC), the solvent was evaporated under reduced pressure. The aqueous layer was washed with di ethylether (50 mL). The separated aqueous layer was acidified by using 2N HCl (pH~3). The aqueous layer was
25 extracted with 10% MeOH/DCM (2 x 100 mL). The combined organic layer was dried over Na₂SO₄ and concentrated under reduced pressure to afford compound C-55-B (780 mg) as yellow solid.

$^1\text{H-NMR}$ (C-55-B isomer): (400 MHz, DMSO- d_6): δ 9.14 (s, 1H), 7.64 (d, J = 6.0 Hz, 2H), 7.49-7.44 (m, 3H), 4.99 (d, J = 12.8 Hz, 1H), 4.48 (s, 2H), 4.45-4.41 (m, 1H), 3.98-3.93 (m, 1H), 2.32-2.13 (m, 4H), 1.40 (d, J = 6.4 Hz, 3H), 0.76 (d, J = 6.4 Hz, 3H);

30 LCMS (ESI): m/z 272.4 [$M^+ - 1$]

Synthesis of 1-benzyl-N-((1S, 2R)-2-((tert-butyldimethylsilyl)oxy)-1-(1, 3, 4-oxadiazol-2-yl) propyl)-2-(1-hydroxyethyl)-5-methylpyrrolidine-2-carboxamide (C-56):

[00232] To a stirring solution of **C-55** (1.78 g, 6.75 mmol) (mixture of two isomers **C-55-A** & **C-55-B**) in DMF (10 mL) were added *N, N*-diisopropylethylamine (3.51 mL, 20.27 mmol), **Int N** (1.74 g, 6.75 mmol) followed by HATU (3.08 g, 8.11 mmol) at 0 °C and stirred at RT for 16 h. After consumption of the starting material (by TLC), the reaction mixture was diluted with water (100 mL) and EtOAc (150 mL). The separated organic layer was washed with water (100 mL) followed by brine solution (100 mL). The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford crude compound which was purified by column chromatography by eluting 30% EtOAc/n-hexane to obtained **C-56** (1.2 g, 35%) as brown syrup.

¹H-NMR: (500 MHz, DMSO-*d*₆): δ 9.31 (s, 1H), 8.42 (s, 1H), 7.44-7.19 (m, 5H), 5.26-4.95 (m, 1H), 4.36 (s, 2H), 4.33-4.18 (m, 2H), 4.09-3.94 (m, 1H), 2.32-2.22 (m, 1H), 2.15-1.86 (m, 4H), 1.37-1.05 (m, 9H), 0.85 (s, 9H), 0.04 (s, 6H);

LCMS (ESI): *m/z* 503.7 [M⁺+1]

Synthesis of 1-benzyl-N-((1S, 2R)-2-((tert-butyldimethylsilyl) oxy)-1-(1, 3, 4-oxadiazol-2-yl) propyl)-2-(1-hydroxyethyl)-5-methylpyrrolidine-2-carboxamide (C-57):

[00233] To a stirring solution of triphenylphosphine (1.56 g, 5.97 mmol) in dry THF (15 mL) was added DIAD (967 mg, 4.78 mmol) as portionwise and stirred for 15 min at RT. To this precipitated solution added **C-56** (1.2 g, 2.39 mmol) in dry THF (15 mL) slowly at RT and stirred for 4 h. After consumption of the starting material (by TLC), the reaction mixture was concentrated under reduced pressure. The obtained solid was triturated with 20% diethylether/n-hexane (2 x 100 mL). The filtrate was concentrated under reduced pressure to obtained crude compound which was purified by silica gel column chromatography eluting 20% EtOAc/hexane to afford **C-57** (400 mg, crude) as brown thick syrup.

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 9.27 (s, 1H), 7.43-7.18 (m, 5H), 4.91-4.73 (m, 2H), 4.42-4.02 (m, 2H), 3.96-3.75 (m, 2H), 2.11-1.84 (m, 3H), 1.42-1.37 (m, 1H), 1.35 (d, *J* = 6.0 Hz, 3H), 1.23-1.11 (m, 6H), 0.70 (s, 9H), 0.02 (s, 6H);

LCMS (ESI): *m/z* 485.7 [M⁺+1]

Synthesis of 5-benzyl-2-((1S, 2R)-2-hydroxy-1-(1, 3, 4-oxadiazol-2-yl) propyl)-3, 6-dimethyl-2, 5-diazaspiro [3.4] octan-1-one (C-58):


[00234] To a stirring solution of **C-57** (400 mg, 0.82 mmol) in dry THF (5 mL) was added TBAF (1M in THF) (1.23 mL, 1.5 mmol) at 0 °C and stirred for 1 h at RT. After consumption of the starting material (by TLC), the reaction mixture was concentrated under reduced pressure. The crude material was diluted with water (30 mL) and EtOAc (50 mL). The organic layer was washed with water (2 x 20 mL), brine solution (2 x 20 mL). The organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to afford crude compound which was purified by column chromatography by eluting 30% EtOAc/*n*-hexane to obtained **C-58** (90 mg, 29%) as thick syrup.

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 9.25 (s, 1H), 7.41-7.18 (m, 5H), 5.28-5.17 (m, 1H), 4.97-4.62 (m, 1H), 4.43-4.20 (m, 1H), 4.02-3.89 (m, 1H), 3.77-3.64 (m, 2H), 3.20-3.01 (m, 1H), 2.09-1.80 (m, 3H), 1.44-1.40 (m, 1H), 1.38-1.30 (m, 1H), 1.28-1.21 (m, 4H), 1.19-1.06 (m, 1H), 0.90-0.85 (m, 3H)

LCMS (ESI):*m/z* 371.45 [M⁺+1]

HPLC: 94.53%

Scheme 20



Synthesis of *tert*-butyl 2-(((*S*)-2-((*tert*-butyldimethylsilyl)oxy)-1-(1,3,4-oxadiazol-2-yl)ethyl)carbamoyl)-2-(hydroxymethyl)-5-methylpyrrolidine-1-carboxylate (**C-59**):

[00235] To a stirring solution of **Z** (2 g, 7.72 mmol) in DCM (25 mL) were added *N,N*-diisopropylethylamine (4 mL, 23.16 mmol), Int **RR** (2.06 g, 8.49 mmol), followed by HATU (3.52 g, 9.26 mmol) at 0 °C and stirred at RT for 16 h. After consumption of the starting material (by TLC), the reaction mixture was diluted with water (50 mL). The separated organic layer was washed with citric acid solution (1 x 50 mL), saturated brine solution (1 x 50 mL). The separated organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford crude compound which was purified by column chromatography by eluting 50% EtOAc/*n*-hexane to obtained **C-59** (2.5 g, 67%) as thick syrup.

¹H-NMR: (500 MHz, DMSO-*d*₆): δ 9.25 (s, 1H), 7.99 (d, *J* = 3.5 Hz, 1H), 5.29-5.26 (m, 2H), 4.05-3.99 (m, 3H), 3.57-3.54 (m, 2H), 2.28-1.91 (m, 3H), 1.42 (s, 9H), 1.38-1.31 (m, 1H), 1.18-1.11 (m, 3H), 0.84 (s, 9H), -0.01 (s, 6H)

Synthesis of tert-butyl 2-((S)-2-((tert-butyldimethylsilyl)oxy)-1-(1,3,4-oxadiazol-2-yl)ethyl)-6-methyl-1-oxo-2,5-diazaspiro[3.4]octane-5-carboxylate (C-60):

[00236] To a stirring solution of triphenylphosphine (2.7 g, 10.33 mmol) in dry THF (20 mL) was added DTAD (2.37 g, 10.33 mmol) at RT and stirred for 15 min. After added **C-59** (2.5 g, 5.16 mmol) and the reaction mixture was stirred at RT for 3 h. After consumption of the starting material (by TLC), the reaction mixture was concentrated under reduced pressure. The crude material was triturated with di ethylether/*n*-pentane and obtained solid was filtered. The filtrate was evaporated, purified by silica gel column chromatography eluting 30% EtOAc/hexane to afford **C-60** (2.1 g, 87.5%) as yellow liquid.

¹H-NMR: (500 MHz, DMSO-*d*₆): δ 9.32 (s, 1H), 5.19-5.08 (m, 1H), 4.12-3.87 (m, 3H), 3.58-3.45 (m, 2H), 1.56-1.42 (m, 3H), 1.40 (s, 9H), 1.38-1.20 (m, 1H), 1.18-1.12 (m, 3H), 0.80 (s, 9H), 0.02 (s, 6H)

Synthesis of tert-butyl 2-(2-hydroxy-1-(pyrimidin-2-yl) propyl)-6-methyl-1-oxo-2, 5-diazaspiro [3.4] octane-5-carboxylate (C-61):


[00237] To a stirring solution of **C-60** (2.1 g, 4.5 mmol) in dry THF (10 mL) was added TBAF (2.34 g, 9 mmol) at 0 °C. The reaction mixture was stirred at RT for 3 h. After consumption of the starting material (by TLC), the reaction mixture was concentrated under reduced pressure. The crude material was purified by silica gel column chromatography eluting 2% MeOH/DCM to afford **C-61** (800 mg, 51%) as thick syrup.

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 9.25 (s, 1H), 5.33-5.08 (m, 2H), 3.94-3.75 (m, 3H), 3.67-3.51 (m, 1H), 3.39-3.31 (m, 1H), 2.13-1.93 (m, 3H), 1.55-1.50 (m, 1H), 1.40 (s, 9H), 1.15-1.08 (m, 3H),

LCMS (ESI): *m/z* 353.3 [M⁺+1];

HPLC: 96.6%

Scheme 21



Synthesis of tert-butyl 2-(((S)-2-((tert-butyldimethylsilyl) oxy)-1-(1, 3, 4-oxadiazol-2-yl) ethyl) carbamoyl)-2-(1-hydroxyethyl) pyrrolidine-1-carboxylate (C-62):

- 93 -

[00238] To a stirring solution of **BB** (3.5 g, 13.51 mmol) in CH₂Cl₂ (40 mL) were added DIPEA (7.0 mL, 40.53 mmol), **RR** (3.2 g, 13.51 mmol), HATU (5.6 g, 14.85 mmol) at 0 °C and stirred for 12 h. After consumption of the starting material (by TLC), the reaction mixture was diluted with water (30 mL) and extracted with CH₂Cl₂ (2 x 100 mL). The separated organic layer was washed with brine, dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to afford **C-62** (3.5 g, crude) as colorless liquid.

Mass (ESI): m/z 485.67 [M⁺+1]

Synthesis of *tert*-butyl 2-((*S*)-2-((*tert*-butyldimethylsilyl) oxy)-1-(1, 3, 4-oxadiazol-2-yl) ethyl)-1-methyl-3-oxo-2, 5-diazaspiro [3.4] octane-5-carboxylate (C-63):

[00239] To a stirring solution of triphenylphosphine (2.03 g, 7.75 mmol) in THF (25 mL) was added DIAD (1.59 g, 7.75 mmol) at 0 °C and stirred for 30 min. **C-62** (1.5 g, 3.10 mmol) in THF (10 mL) was added dropwise and the reaction mixture was stirred at RT for 16 h. After consumption of the starting material (by TLC), the reaction was concentrated under reduced pressure. The crude material was purified by silica gel column chromatography eluting 30% EtOAc/hexane to afford **C-63** (600 mg, 43%) as pale yellow liquid.

¹H-NMR: (500 MHz, DMSO-*d*₆): δ 9.24 (s, 1H), 5.93-4.75 (m, 2H), 4.16-3.91 (m, 2H), 3.78-3.70 (m, 2H), 3.91 (d, *J* = 7.0 Hz, 1H), 3.77 (d, *J* = 7.0 Hz, 2H), 3.44-3.34 (m, 4H), 2.01-1.91 (m, 2H), 1.85-1.68 (m, 6H), 2.11-1.68 (m, 4H), 1.40 (s, 9H), 1.38-1.18 (m, 3H), 0.80 (s, 9H), -0.02 (s, 6H).

Mass (ESI): m/z 467.6 [M⁺+1]

Synthesis of *tert*-butyl 2-((*S*)-2-hydroxy-1-(1, 3, 4-oxadiazol-2-yl) ethyl)-1-methyl-3-oxo-2, 5-diazaspiro [3.4] octane-5-carboxylate (C-64):


[00240] To a stirring solution of **C-63** (0.5 g, 1.07 mmol) in dry THF (10 mL) was added TBAF (1M in THF) (1.07 mL, 1.91 mmol) at 0 °C and stirred at RT for 3 h. After consumption of the starting material (by TLC), the reaction mixture was concentrated under reduced pressure. The crude material was diluted with water (15 mL) and EtOAc (30 mL). The organic layer was washed with water (2 x 15 mL), brine solution (2 x 10 mL). The organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to afford crude compound which was purified by column chromatography by eluting 70% EtOAc/*n*-hexane to obtain **C-64** (95 mg, 25%) as white solid.

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 9.26 (s, 1H), 5.32-5.04 (m, 2H), 3.98-3.72 (m, 4H), 2.14-1.66 (m, 4H), 1.39 (s, 1H), 1.37-1.09 (m, 3H)

LCMS (ESI): m/z 353.3 [M⁺+1]

HPLC: 88.9%

Scheme 22



Synthesis of tert-butyl 2-(((S)-2-((tert-butyldimethylsilyl)oxy)-1-(1,3,4-oxadiazol-2-yl)ethyl)carbamoyl)-2-(1-hydroxyethyl)-5-methylpyrrolidine-1-carboxylate (C-65):

[00241] To a stirring solution of DD (1 g, 3.66 mmol) in DCM (20 mL) were added *N*, *N*-diisopropylethylamine (1.9 mL, 10.98 mmol), Int RR (978 mg, 4.02 mmol), EDCI (1.04 g, 5.49 mmol) followed by HOBt (741 mg, 5.49 mmol) at 0 °C and stirred at RT for 16 h. After consumption of the starting material (by TLC), the reaction mixture was diluted with water (20 mL). The separated organic layer was washed with brine solution (30 mL). The separated organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford crude compound which was purified by column chromatography by eluting 40% EtOAc/*n*-hexane to obtained C-65 (700 mg, 38.4%) as thick syrup.

¹H-NMR: (500 MHz, DMSO-*d*₆): δ 9.26 (s, 1H), 5.22-5.18 (m, 1H), 4.58-4.50 (m, 2H), 4.04-3.92 (m, 3H), 1.47-1.41 (m, 1H), 1.38 (s, 9H), 1.29-1.16 (m, 6H), 1.14-1.11 (m, 3H), 0.83 (s, 9H), -0.02 (s, 6H);

LCMS (ESI): *m/z* 523.6 [M⁺+1]

Synthesis of tert-butyl 2-(((S)-2-((tert-butyldimethylsilyl)oxy)-1-(1,3,4-oxadiazol-2-yl)ethyl)-1,6-dimethyl-3-oxo-2,5-diazaspiro[3.4]octane-5-carboxylate (C-66):

[00242] To a stirring solution of triphenylphosphine (736 mg, 2.81 mmol) in dry THF (10 mL) was added DTAD (646 mg, 2.81 mmol) as portionwise and stirred for 15 min at RT. To this precipitated solution added C-65 (700 mg, 1.4 mmol) in dry THF (15 mL) slowly at RT and stirred for 6 h. After consumption of the starting material (by TLC), the reaction mixture was concentrated under reduced pressure. The crude material was triturated with *n*-pentane (20 mL) /di ethylether (20 mL). The filtrate was concentrated under reduced pressure to obtained crude compound which was purified by silica gel column chromatography eluting 25% EtOAc/hexane to afford C-66 (430 mg, 64%) as thick syrup.

- 95 -

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 9.25 (s, 1H), 5.20-5.16 (m, 1H), 4.16-4.00 (m, 1H), 3.88-3.70 (m, 3H), 2.17-2.08 (m, 1H), 1.96-1.85 (m, 1H), 1.40 (s, 9H), 1.26-1.07 (m, 8H), 0.82 (s, 9H), -0.02 (s, 6H);

LCMS (ESI): *m/z* 505.5 [*M*⁺+1]

5 **Synthesis of tert-butyl 2-((*S*)-2-hydroxy-1-(1, 3, 4-oxadiazol-2-yl) ethyl)-1, 6-dimethyl-3-oxo-2, 5-diazaspiro [3.4] octane-5-carboxylate (C-67):**

[00243] To a stirring solution of **C-66** (430 mg, 0.89 mmol) in dry THF (10 mL) was added TBAF (1.8 mL, 1.79 mmol) slowly at 0 °C and stirred at RT for 6 h. After consumption of the starting material (by TLC), the reaction mixture was concentrated under reduced
10 pressure. The crude residue was purified with column chromatography by eluting 90% EtOAc/n-hexane to afford **C-67** (80 mg, 24.5%) as thick syrup.

¹H-NMR: (400 MHz, DMSO-*d*₆): δ 9.25 (s, 1H), 5.20-5.03 (m, 2H), 3.98-3.72 (m, 3H), 2.50-1.85 (m, 3H), 1.56-1.50 (m, 1H), 1.40 (s, 9H), 1.27-1.11 (m, 6H)

LCMS (ESI): *m/z* 291.3 [*M*⁺+1];

15 **HPLC:** 91.4%

Example 3 - [³H] MK-801 binding assay

Methods

[00244] Assays were conducted as described in Moskal et al. (Moskal, J.R., Kuo, A.G.,
20 Weiss, C., Wood, P.L., O'Connor Hanson, A., Kelso, S., Harris, R.B., Disterhoft, J.F., 2005. GLYX-13: a monoclonal antibody-derived peptide that acts as an N-methyl-D-aspartate receptor modulator. *Neuropharmacology*. 49, 1077-87) The potentiation of [³H]MK-801 binding (5 nM; 22.5 Ci / mmol) to well washed rat cortical membranes (200 µg) was measured under non-equilibrium conditions (15 min @ 25 °C) in the presence of increasing
25 concentrations of test compounds and 50µM glutamate. Zero levels were determined in the absence of any glycine ligand and in the presence of 30µM 5,7 DCKA. Maximal stimulation was measured in the presence of 1 mM glycine, and 50µM glutamate was present in all samples. The facilitation of [³H]MK-801 binding by tests compounds was calculated by using a 3 parameter log agonist vs. response equation (Graph pad Prism, USA) and potency (EC₅₀,
30 expressed in pM) and maximal activity (% maximal stimulation) were calculated for the test compound.

Results

[00245] The potency and maximal activity for Compound Y is 288 pM, 14%.

Table 2. Additional Biological Data

Compound	Unified Activity Data: LTP Augmentation (Percent)	Unified Activity Data: LTP Concentration (uM)	Unified Activity Data: LTP, Significant (S) or Non-significant (NS)	Unified Activity Data: Porsolt Floating Time Inhibition (Percent)	Unified Activity Data: Porsolt Dose (mg/kg)	Unified Activity Data: Porsolt Dose, route	Unified Activity Data: Porsolt Time Post Dose (Hours)	MK-801 Glycine Site Binding Assay: Rat Cortex EC50 (M)
C-4	140	1	S	92	3	IV	1	2.879E-10
C-15	120	1	S	86	1	PO	1	

5 **Table 3.** Additional Biological Data

Compound	MK-801 Glycine Site Binding Assay: Rat Cortex EC50 (M)	LTP: LTP Augmentation (%)	LTP: LTP Concentration (uM)	LTP: LTP Significance, S or NS
C-33	7.129E-12	30	1	NS
C-19	1.096E-10	110	1	S
C-26	7.23E-13	65	1	NS
C-29	1.353E-10	125	1	S

EQUIVALENTS

[00246] Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention

10 described herein. Such equivalents are intended to be encompassed by the following claims.


INCORPORATION BY REFERENCE

[00247] The entire contents of all patents, published patent applications, websites, and other references cited herein are hereby expressly incorporated herein in their entireties by reference.

- 97 -

What is claimed is:

1. A compound represented by formula I:



- or a pharmaceutically acceptable salt thereof, wherein

R_b is selected from the group consisting of H, halogen, hydroxyl, cyano or C₁-C₆ alkyl;

R₁ is H or C₁-C₆ alkyl;

R₂ is H or C₁-C₆ alkyl;

R₃ is selected from the group consisting of H, C₁-C₆ alkyl and a nitrogen protecting group;

R₄ is H or C₁-C₆ alkyl;

R₅ is X or -C₁-C₆ alkylene-X, wherein X is selected from the group consisting of:

(i) phenyl;

(ii) heteroaryl including from 5 to 6 ring atoms wherein 1, 2, or 3 of the ring atoms are independently selected from the group consisting of N, NH, N(C₁-C₃ alkyl), O, and S; and

(iii) heterocyclyl including from 3 to 6 ring atoms wherein 1, 2, or 3 of the ring atoms are independently selected from the group consisting of N, NH, N(C₁-C₃ alkyl), O,

and S; wherein R₅ is optionally substituted with ; and

R₆ is selected from the group consisting of H, halogen, hydroxyl, cyano, -O-C(O)-C₁-C₆ alkyl, C₁-C₆ alkyl, or C₁-C₆ alkoxy.

2. The compound of claim 1, wherein R₁ is H.

3. The compound of claim 1 or 2, wherein R₂ is H.

4. The compound of any one of claims 1-3, wherein R₃ is H.

5. The compound of any one of claims 1-3, wherein R₃ is a nitrogen protecting group.


- 1 6. The compound of claim 5, wherein R_3 has formula $-C(O)OR_{31}$, wherein R_{31} is selected
2 from the group consisting of: C_1 - C_6 alkyl; C_1 - C_6 haloalkyl; C_2 - C_6 alkenyl; C_2 - C_6 alkynyl; C_3 -
3 C_{10} cycloalkyl, wherein the C_3 - C_{10} cycloalkyl is optionally substituted with from 1-3
4 independently selected C_1 - C_3 alkyl; $-CH_2$ - C_3 - C_{10} cycloalkyl wherein the C_3 - C_{10} cycloalkyl is
5 optionally substituted with from 1-3 independently selected C_1 - C_3 alkyl; $-CH_2$ -phenyl, wherein
6 the phenyl is optionally substituted with from 1-2 substituents independently selected from C_1 -
7 C_3 alkyl; C_1 - C_3 haloalkyl; C_1 - C_3 alkoxy; C_1 - C_3 haloalkoxy; nitro; halo; SO_2Me , cyano; and -
8 $OC(O)CH_3$; and $-CH_2$ -pyridyl.
- 1 7. The compound of claim 6, wherein R_{31} is C_1 - C_6 alkyl.
- 1 8. The compound of claim 7, wherein R_{31} is *tert*-butyl.
- 1 9. The compound of claim 5, wherein R_3 has formula $-C(O)R_{32}$, wherein R_{32} is selected
2 from the group consisting of: H; C_1 - C_6 alkyl; C_1 - C_6 haloalkyl; phenyl, wherein the phenyl is
3 optionally substituted with from 1-2 substituents independently selected from C_1 - C_3 alkyl; C_1 -
4 C_3 haloalkyl; C_1 - C_3 alkoxy; C_1 - C_3 haloalkoxy; nitro; halo; SO_2Me , cyano; and $-OC(O)CH_3$;
5 and pyridyl.
- 1 10. The compound of claim 9, wherein R_{32} is C_1 - C_6 alkyl.
- 1 11. The compound of claim 10, wherein R_{32} is $-CH_3$ or *iso*-propyl.
- 1 12. The compound of any one of claims 1-11, wherein R_5 is X.
- 1 13. The compound of claim 10, wherein X is heteroaryl including from 5 to 6 ring atoms
2 wherein 1, 2, or 3 of the ring atoms are independently selected from the group consisting of N,
3 NH, $N(C_1$ - C_3 alkyl), O, and S.
- 1 14. The compound of claim 13, wherein X is selected from the group consisting of 1,2,4-
2 oxadiazolyl, 1,3,4- oxadiazolyl, pyridyl, and pyrimidinyl.
- 1 15. The compound of any one of claims 12-14, wherein R_5 is substituted with $-CH(R_4)(R_6)$.
- 1 16. The compound of claim 15, wherein R_4 is C_1 - C_6 alkyl.
- 1 17. The compound of claim 15 or 16, wherein R_4 is methyl.
- 1 18. The compound of any one of claims 15-17, wherein R_6 is selected from the group
2 consisting of hydroxyl, $-O-C(O)-C_1$ - C_6 alkyl, and C_1 - C_6 alkoxy.
- 1 19. The compound of claim 18, wherein R_6 is hydroxyl.

- 1 20. The compound of any one of claims 1-11, wherein R_5 is $-C_1-C_6$ alkylene-X.
- 1 21. The compound of any one of claims 1-11, wherein R_5 is $-C_1-C_2$ alkylene-X.
- 1 22. The compound of claim 21 or 22, wherein R_5 is substituted with $-\text{CH}(R_4)(R_6)$.
- 1 23. The compound of claim 22, wherein R_5 is substituted on the alkylene portion with –
2 $\text{CH}(R_4)(R_6)$.
- 1 24. The compound of claim 23, wherein R_5 has the formula $-\text{CH}(\text{CHR}_4R_6)-\text{X}$.
- 1 25. The compound of any one of claims 22-24, wherein R_4 is C_1-C_6 alkyl.
- 1 26. The compound of any one of claims 22-25, wherein R_4 is methyl.
- 1 27. The compound of any one of claims 22-26, wherein R_6 is selected from the group
2 consisting of hydroxyl, $-\text{O}-\text{C}(\text{O})-C_1-C_6$ alkyl, and C_1-C_6 alkoxy.
- 1 28. The compound of claim 27, wherein R_6 is hydroxyl.
- 1 29. The compound of claim 20 or 21, wherein X is heteroaryl including from 5 to 6 ring
2 atoms wherein 1, 2, or 3 of the ring atoms are independently selected from the group consisting
3 of N, NH, $\text{N}(C_1-C_3 \text{ alkyl})$, O, and S.
- 1 30. The compound of any one of claims 20-22, wherein X is selected from the group
2 consisting of 1,2,4- oxadiazolyl, 1,3,4- oxadiazolyl, pyridyl, and pyrimidinyl.
- 1 31. The compound of any one of claims 1-30, wherein R_6 is H.
- 1 32. The compound of claim 1, wherein the compound is selected from the compounds
2 delineated in Table 1.
- 1 33. A pharmaceutical composition comprising a compound of any one of claims 1-32, and a
2 pharmaceutically acceptable excipient.
- 1 34. The pharmaceutical composition of claim 33, suitable for oral administration.
- 1 35. The pharmaceutical composition of claim 33, suitable for intravenous administration.
- 1 36. A method of treating of treating depression, Alzheimer's disease, attention deficit
2 disorder, schizophrenia, or anxiety, in a patient in need thereof, comprising administering to
3 said patient:
4 a pharmaceutically effective amount of a compound of any one of claims 1-32.

- 100 -

6

1 37. A compound represented by formula I:



2

3 and pharmaceutically acceptable salts, stereoisomers, and N-oxides thereof, wherein

4 R_b is selected from the group consisting of H, halogen, hydroxyl, cyano or C₁-C₆ alkyl;

5 R₁ is H or C₁-C₆ alkyl;

6 R₂ is H or C₁-C₆ alkyl;

7 R₃ is selected from the group consisting of H, C₁-C₆ alkyl and a nitrogen protecting
8 group;

9 R₄ is H or C₁-C₆ alkyl;

10 R₅ is X or -C₁-C₆ alkylene-X, wherein X is selected from the group consisting of
11 phenyl, a 4- to 6-membered heteroaryl ring having 1, 2, or 3 heteroatoms selected from
12 O, S, or N, or a 4- to 6-membered heterocyclyl ring having 1, 2, or 3 heteroatoms

13 selected from O, S, or N, and wherein R₅ is optionally substituted with ; and

14 R₆ is selected from the group consisting of H, halogen, hydroxyl, cyano, -O-C(O)-C₁-C₆
15 alkyl, C₁-C₆ alkyl, or C₁-C₆ alkoxy.

1 38. The compound of claim 37, wherein R₁ is H.

1 39. The compound of claim 37 or 38, wherein R₂ is H.

1 40. The compound of any one of claims 37-39, wherein R₃ is H.

1 41. The compound of any one of claims 37-40, wherein R₄ is C₁-C₆ alkyl.



1 42. The compound of any one of claims 37-40, wherein R₄ is methyl.

1 43. The compound of any one of claims 37-42, wherein R_b is H.


1 44. The compound of any one of claims 37-43, wherein R₅ is X, and wherein X is a 5- to 6-
2 membered heteroaryl ring selected from the group consisting of azetidine, pyrrolidine,

- 101 -

pyrazolidine, pyridine, pyrimidine, isooxazolidine, imidazolidine, oxazolidine, thiazolidine, and isothiazolidine.


45. The compound of any one of claims 37-43, wherein R_5 is  or .

46. The compound of any one of claims 37-43, wherein R_5 is X, and wherein X is


optionally substituted with .

47. The compound of any one of claims 37-46, wherein R_6 is hydroxyl.

48. The compound of claim 37, wherein the compound is



49. A compound represented by formula II:




and pharmaceutically acceptable salts, stereoisomers, and N-oxides thereof, wherein

R_3 is selected from the group consisting of H, C_1 - C_6 alkyl and a nitrogen protecting group; and

R_7 is a 4- to 6-membered heteroaryl ring having 1, 2, or 3 heteroatoms selected from O, S, or N, optionally substituted on a free carbon by a substituent selected from the group consisting of: halogen, C_1 - C_6 alkyl, hydroxyl, cyano, and phenyl.

50. A compound represented by formula III:



and pharmaceutically acceptable salts, stereoisomers, and N-oxides thereof, wherein

- 102 -

4 R_b is selected from the group consisting of H, halogen, hydroxyl, cyano or C_1-C_6 alkyl;

5 R_1 is H or C_1-C_6 alkyl;

6 R_2 is H or C_1-C_6 alkyl;

7 R_3 is selected from the group consisting of H, C_1-C_6 alkyl and a nitrogen protecting
8 group;

9 R_4 is H or C_1-C_6 alkyl;

10 X is selected from the group consisting of phenyl, a 4- to 6-membered heteroaryl ring
11 having 1, 2, or 3 heteroatoms selected from O, S, or N, or a 4- to 6-membered
12 heterocyclyl ring having 1, 2, or 3 heteroatoms selected from O, S, or N, wherein

13  is present on a free carbon of X; and

14 R_6 is selected from the group consisting of H, halogen, hydroxyl, cyano, $-O-C(O)-C_1-C_6$
15 alkyl, C_1-C_6 alkyl, or C_1-C_6 alkoxy.

1 51. The compound of claim 50, wherein R_1 is H.

1 52. The compound of claim 50 or 51, wherein R_2 is H.


1 53. The compound of any one of claims 50-52, wherein R_3 is H.

1 54. The compound of any one of claims 50-53, wherein R_4 is C_1-C_6 alkyl.

1 55. The compound of any one of claims 50-53, wherein R_4 is methyl.


1 56. The compound of any one of claims 50-55, wherein R_b is H.

1 57. The compound of any one of claims 50-56, wherein X is



1 58. The compound of any one of claims 50-57, wherein R_6 is hydroxyl.

1 59. The compound of claim 50, wherein the compound is



1 60. A pharmaceutical composition comprising a compound of any one of claims 37-59, and
2 a pharmaceutically acceptable excipient.

- 103 -

- 1 61. The pharmaceutical composition of claim 60, suitable for oral administration.
- 1 62. The pharmaceutical composition of claim 60, suitable or intravenous administration.
- 1 63. A method of treating of treating depression, Alzheimer's disease, attention deficit
2 disorder, schizophrenia, or anxiety, in a patient in need thereof, comprising administering to
3 said patient:
- 4 a pharmaceutically effective amount of a compound of any one of claims 37-59.