

DIFFERENTIATION OF HUMAN EMBRYONIC STEM CELLS TO THE PANCREATIC ENDOCRINE LINEAGE

- [0001] The present invention claims priority to application serial number 61/110,287, filed October 31, 2008.

FIELD OF THE INVENTION

- [0002] The present invention provides methods to promote the differentiation of pluripotent stem cells. In particular, the present invention provides a method to increase the expression of MAFA in cells expressing markers characteristic of the pancreatic endocrine lineage.

BACKGROUND

- [0003] Advances in cell-replacement therapy for Type I diabetes mellitus and a shortage of transplantable islets of Langerhans have focused interest on developing sources of insulin-producing cells, or β cells, appropriate for engraftment. One approach is the generation of functional β cells from pluripotent stem cells, such as, for example, embryonic stem cells.
- [0004] In vertebrate embryonic development, a pluripotent cell gives rise to a group of cells comprising three germ layers (ectoderm, mesoderm, and endoderm) in a process known as gastrulation. Tissues such as, for example, thyroid, thymus, pancreas, gut, and liver, will develop from the endoderm, via an intermediate stage. The intermediate stage in this process is the formation of definitive endoderm. Definitive endoderm cells express a number of markers, such as, HNF-3 beta, GATA4, MIXL1, CXCR4 and SOX17.
- [0005] Formation of the pancreas arises from the differentiation of definitive endoderm into pancreatic endoderm. Cells of the pancreatic endoderm express the pancreatic-duodenal homeobox gene, PDX1. In the absence of PDX1, the pancreas fails to develop beyond the formation of ventral and dorsal buds. Thus, PDX1 expression marks a critical step in pancreatic organogenesis. The mature pancreas contains, among other cell types, exocrine tissue and endocrine tissue. Exocrine and endocrine tissues arise from the differentiation of pancreatic endoderm.

- [0006] Cells bearing the features of islet cells have reportedly been derived from embryonic cells of the mouse. For example, Lumelsky *et al.* (Science 292:1389, 2001) report differentiation of mouse embryonic stem cells to insulin-secreting structures similar to pancreatic islets. Soria *et al.* (Diabetes 49:157, 2000) report that insulin-secreting cells derived from mouse embryonic stem cells normalize glycemia in streptozotocin-induced diabetic mice.
- [0007] In one example, Hori *et al.* (PNAS 99: 16105, 2002) disclose that treatment of mouse embryonic stem cells with inhibitors of phosphoinositide 3-kinase (LY294002) produced cells that resembled β cells.
- [0008] In another example, Blyszczuk *et al.* (PNAS 100:998, 2003) reports the generation of insulin-producing cells from mouse embryonic stem cells constitutively expressing Pax4.
- [0009] Micallef *et al.* reports that retinoic acid can regulate the commitment of embryonic stem cells to form Pdx1 positive pancreatic endoderm. Retinoic acid is most effective at inducing Pdx1 expression when added to cultures at day 4 of embryonic stem cell differentiation during a period corresponding to the end of gastrulation in the embryo (Diabetes 54:301, 2005).
- [0010] Miyazaki *et al.* reports a mouse embryonic stem cell line over-expressing Pdx1. Their results show that exogenous Pdx1 expression clearly enhanced the expression of insulin, somatostatin, glucokinase, neurogenin3, P48, Pax6, and HNF6 genes in the resulting differentiated cells (Diabetes 53: 1030, 2004).
- [0011] Skoudy *et al.* reports that activin A (a member of the TGF- β superfamily) upregulates the expression of exocrine pancreatic genes (p48 and amylase) and endocrine genes (Pdx1, insulin, and glucagon) in mouse embryonic stem cells. The maximal effect was observed using 1nM activin A. They also observed that the expression level of insulin and Pdx1 mRNA was not affected by retinoic acid; however, 3nM FGF7 treatment resulted in an increased level of the transcript for Pdx1 (Biochem. J. 379: 749, 2004).
- [0012] Shiraki *et al.* studied the effects of growth factors that specifically enhance differentiation of embryonic stem cells into Pdx1 positive cells. They observed that

TGF- β 2 reproducibly yielded a higher proportion of Pdx1 positive cells (Genes Cells. 2005 Jun; 10(6): 503-16.).

- [0013] Gordon *et al.* demonstrated the induction of brachyury⁺/HNF-3beta⁺ endoderm cells from mouse embryonic stem cells in the absence of serum and in the presence of activin along with an inhibitor of Wnt signaling (US 2006/0003446A1).
- [0014] Gordon *et al.* (PNAS, Vol 103, page 16806, 2006) states “Wnt and TGF-beta/ nodal/ activin signaling simultaneously were required for the generation of the anterior primitive streak”.
- [0015] However, the mouse model of embryonic stem cell development may not exactly mimic the developmental program in higher mammals, such as, for example, humans.
- [0016] Thomson *et al.* isolated embryonic stem cells from human blastocysts (Science 282:114, 1998). Concurrently, Gearhart and coworkers derived human embryonic germ (hEG) cell lines from fetal gonadal tissue (Shamblott *et al.*, Proc. Natl. Acad. Sci. USA 95:13726, 1998). Unlike mouse embryonic stem cells, which can be prevented from differentiating simply by culturing with Leukemia Inhibitory Factor (LIF), human embryonic stem cells must be maintained under very special conditions (U.S. Pat. No. 6,200,806; WO 99/20741; WO 01/51616).
- [0017] D’Amour *et al.* describes the production of enriched cultures of human embryonic stem cell-derived definitive endoderm in the presence of a high concentration of activin and low serum (Nature Biotechnology 2005). Transplanting these cells under the kidney capsule of mice resulted in differentiation into more mature cells with characteristics of some endodermal organs. Human embryonic stem cell-derived definitive endoderm cells can be further differentiated into Pdx1 positive cells after addition of FGF-10 (US 2005/0266554A1).
- [0018] D’Amour *et al.* (Nature Biotechnology - 24, 1392 - 1401 (2006)) states: “We have developed a differentiation process that converts human embryonic stem (hES) cells to endocrine cells capable of synthesizing the pancreatic hormones insulin, glucagon, somatostatin, pancreatic polypeptide and ghrelin. This process mimics in vivo pancreatic organogenesis by directing cells through stages resembling definitive

endoderm, gut-tube endoderm, pancreatic endoderm and endocrine precursor en route to cells that express endocrine hormones”.

- [0019] In another example, Fisk *et al.* reports a system for producing pancreatic islet cells from human embryonic stem cells (US2006/0040387A1). In this case, the differentiation pathway was divided into three stages. Human embryonic stem cells were first differentiated to endoderm using a combination of sodium butyrate and activin A. The cells were then cultured with TGF- β antagonists such as Noggin in combination with EGF or betacellulin to generate Pdx1 positive cells. The terminal differentiation was induced by nicotinamide.
- [0020] In one example, Benvenisty *et al.* states: “We conclude that over-expression of Pdx1 enhanced expression of pancreatic enriched genes, induction of insulin expression may require additional signals that are only present *in vivo*” (Benvenisty *et al.*, Stem Cells 2006; 24:1923–1930).
- [0021] Cyclins have been implicated in beta cell function. For example, Lilja *et al.* report that Cdk5 is present in the insulin-secreting pancreatic β -cell (J. Biol. Chem., Vol. 276, Issue 36, 34199-34205, September 7, 2001). Lilja *et al.* states “Cdk5 is present in β -cells and acts as a positive regulator of insulin exocytosis.”
- [0022] In another example, Marzo *et al.* states “Cdk4 knockin mice have significantly increased beta cell mass and are physiologically functional, indicating that Cdk4 is a potential target for pancreatic beta cell mass regeneration in Type 1 diabetes” (Diabetologia, Vol. 47, Number 4, 686-694, April 1, 2004.)
- [0023] In another example, Ubeda *et al.* report that inhibition of cyclin-dependant kinase 5 activity protects pancreatic beta cells from glucotoxicity (J. Biol. Chem., Vol. 281, Issue 39, 28858-28864, September 29, 2006).
- [0024] In another example, Wei *et al.* report Cdk5-dependent regulation of glucose-stimulated insulin secretion (Nature Medicine 11, 1104-1108 (1 October 2005).)
- [0025] In another example, Vanderford *et al.* state “MafA is a basic leucine zipper transcription factor expressed within the beta cells of the pancreas and is required to maintain normal glucose homeostasis as it is involved in various aspects of beta cell

biology. MafA protein levels are known to increase in response to high glucose through mechanisms that have yet to be fully characterized. We investigated whether discrete intracellular signaling events control mafA expression. We found that the general kinase inhibitor staurosporine induces mafA expression without altering the stability of the protein. Inhibition of the MAP-kinase JNK mimics the effects of staurosporine on the expression of mafA. Calmodulin kinase and calcium signaling are also important in stimulating mafA expression by high glucose. However, staurosporine, JNK, and calmodulin kinase have different effects on the induction of insulin expression. These data reveal that MafA levels are tightly controlled by the coordinated action of multiple kinase pathways.” (Archives of Biochemistry and Biophysics (2008), doi: 10.1016/j.abb.2008.10.001).

[0026] Therefore, there still remains a significant need to develop methods for differentiating pluripotent stem cells into pancreatic endocrine cells, pancreatic hormone expressing cells, or pancreatic hormone secreting cells. The present invention provides methods to increase the expression of MAFA in cells expressing markers characteristic of the pancreatic endocrine lineage.

SUMMARY

[0027] In one embodiment, the present invention provides a method for increasing the expression of MAFA in cells expressing markers characteristic of the pancreatic endocrine lineage comprising the steps of culturing the cells expressing markers characteristic of the pancreatic endocrine lineage in medium comprising a sufficient amount of a cyclin-dependant kinase inhibitor to cause an increase in expression of MAFA.

BRIEF DESCRIPTION OF THE DRAWINGS

[0028] Figure 1, panel a shows the effect of compounds from the EMD Calbiochem kinase inhibitor library on the ratio of insulin to glucagon expression in cells expressing markers characteristic of the pancreatic endocrine lineage, as determined by real-time PCR. The alphanumeric label corresponds to the compound identity as shown in Table 1. Panel b shows the effect of compounds from the EMD Calbiochem kinase inhibitor library on the ratio of MAFA to ARX4 expression in cells expressing markers characteristic of the pancreatic endocrine lineage, as determined by real-time

PCR. The alphanumeric label corresponds to the compound identity as shown in Table 1.

- [0029] Figure 2 A) shows a 4x micrograph of cells treated according to the methods described in Example 1, at day 4 of the stage 6 treatment. B) shows a 4x micrograph of cells treated with 0.5 μ M of the compound PubChemID# 5330812 at day 4 of treatment. C) shows a 4x micrograph of cells treated with 1 μ M of the compound PubChemID# 5330812 at day 4 of treatment. D) shows a 20x micrograph of cells treated according to the methods described in Example 1, at day 6 of the stage 6 treatment. E) shows a 20x micrograph of cells treated with 0.5 μ M of the compound PubChemID# 5330812 at day 6 of treatment. F) shows a 20x micrograph of cells treated with 1 μ M of the compound PubChemID# 5330812 at day 6 of treatment.
- [0030] Figure 3 shows the expression of the 23 genes indicated, in cells expressing markers characteristic of the pancreatic endocrine lineage following a five-day treatment of 0.5 μ M (dark bars) or 1.0 μ M (light bars) of the compound PubChem ID#5330812. Expression levels were determined at day 0, day 2 and day 5.
- [0031] Figure 4 shows the effect of CDK inhibitor III treatment on the expression of markers characteristic of the pancreatic endocrine lineage in cells treated with Stage 7 of the differentiation protocol described in Example 4.
- [0032] Figure 5 shows the effect of CDK inhibitor III treatment on the dithazone staining of islet-like clusters.
- [0033] Figure 6 shows the expression of insulin, synaptophysin and glucagon in insulin-producing cells produced according to the methods described in Example 5. Expression of the proteins indicated was determined by FACS.
- [0034] Figure 7 shows the expression of insulin, synaptophysin and glucagon in insulin-producing cells produced according to the methods described in Example 5. Expression of the proteins indicated was determined by FACS.
- [0035] Figure 8 shows the expression of MAFA (Panel a) and insulin (Panel b), in insulin-producing cells, produced by the methods of the present invention. Samples of cells were taken for PCR analysis at days 1, 2, 3, and 4. Following 4 days of treatment

with CDK inhibitor, the CDK inhibitor was removed from culture and the cells were cultured additional 4 days in DMEM-F12 + 1% B27 + 20 ng/ml of activin A. At the end of the four days, samples were collected in triplicate for PCR analysis.

[0036] Figure 9 shows the effect of compounds from the EMD Calbiochem kinase inhibitor library I on the expression of MAFA in cells expressing markers characteristic of the pancreatic endocrine lineage, as determined by real-time PCR.

[0037] Figure 10 shows the effect of genestein on the mRNA expression of insulin, glucagon, somatostatin and MAFA in cells expressing markers characteristic of the pancreatic endocrine lineage, as determined by real-time PCR.

DETAILED DESCRIPTION

[0038] For clarity of disclosure, and not by way of limitation, the detailed description of the invention is divided into the following subsections that describe or illustrate certain features, embodiments or applications of the present invention.

Definitions

[0039] Stem cells are undifferentiated cells defined by their ability at the single cell level to both self-renew and differentiate to produce progeny cells, including self-renewing progenitors, non-renewing progenitors, and terminally differentiated cells. Stem cells are also characterized by their ability to differentiate *in vitro* into functional cells of various cell lineages from multiple germ layers (endoderm, mesoderm and ectoderm), as well as to give rise to tissues of multiple germ layers following transplantation and to contribute substantially to most, if not all, tissues following injection into blastocysts.

[0040] Stem cells are classified by their developmental potential as: (1) totipotent, meaning able to give rise to all embryonic and extraembryonic cell types; (2) pluripotent, meaning able to give rise to all embryonic cell types; (3) multipotent, meaning able to give rise to a subset of cell lineages but all within a particular tissue, organ, or physiological system (for example, hematopoietic stem cells (HSC) can produce progeny that include HSC (self-renewal), blood cell restricted oligopotent progenitors, and all cell types and elements (e.g., platelets) that are normal

components of the blood); (4) oligopotential, meaning able to give rise to a more restricted subset of cell lineages than multipotent stem cells; and (5) unipotent, meaning able to give rise to a single cell lineage (e.g., spermatogenic stem cells).

[0041] Differentiation is the process by which an unspecialized ("uncommitted") or less specialized cell acquires the features of a specialized cell such as, for example, a nerve cell or a muscle cell. A differentiated or differentiation-induced cell is one that has taken on a more specialized ("committed") position within the lineage of a cell. The term "committed", when applied to the process of differentiation, refers to a cell that has proceeded in the differentiation pathway to a point where, under normal circumstances, it will continue to differentiate into a specific cell type or subset of cell types, and cannot, under normal circumstances, differentiate into a different cell type or revert to a less differentiated cell type. De-differentiation refers to the process by which a cell reverts to a less specialized (or committed) position within the lineage of a cell. As used herein, the lineage of a cell defines the heredity of the cell, i.e., which cells it came from and what cells it can give rise to. The lineage of a cell places the cell within a hereditary scheme of development and differentiation. A lineage-specific marker refers to a characteristic specifically associated with the phenotype of cells of a lineage of interest and can be used to assess the differentiation of an uncommitted cell to the lineage of interest.

[0042] "β-cell lineage" refers to cells with positive gene expression for the transcription factor PDX-1 and at least one of the following transcription factors: NGN3, NKX2.2, NKX6.1, NEUROD, ISL1, HNF3 beta, MAFA, PAX4, or PAX6. Cells expressing markers characteristic of the β cell lineage include β cells.

[0043] "Cells expressing markers characteristic of the definitive endoderm lineage", as used herein, refers to cells expressing at least one of the following markers: SOX17, GATA4, HNF3 beta, GSC, CER1, Nodal, FGF8, Brachyury, Mix-like homeobox protein, FGF4 CD48, eomesodermin (EOMES), DKK4, FGF17, GATA6, CXCR4, C-Kit, CD99, or OTX2. Cells expressing markers characteristic of the definitive endoderm lineage include primitive streak precursor cells, primitive streak cells, mesendoderm cells and definitive endoderm cells.

- [0044] “Cells expressing markers characteristic of the pancreatic endoderm lineage”, as used herein, refers to cells expressing at least one of the following markers: PDX1, HNF1 beta, PTF1 alpha, HNF-, or HB9. Cells expressing markers characteristic of the pancreatic endoderm lineage include pancreatic endoderm cells, primitive gut tube cells, and posterior foregut cells.
- [0045] “Cells expressing markers characteristic of the pancreatic endocrine lineage”, as used herein, refers to cells expressing at least one of the following markers: NGN3, NEUROD, ISL1, PDX1, NKX6.1, PAX4, NGN3, or PTF1 alpha. Cells expressing markers characteristic of the pancreatic endocrine lineage include pancreatic endocrine cells, pancreatic hormone expressing cells, and pancreatic hormone secreting cells, and cells of the β -cell lineage.
- [0046] “Definitive endoderm”, as used herein, refers to cells which bear the characteristics of cells arising from the epiblast during gastrulation and which form the gastrointestinal tract and its derivatives. Definitive endoderm cells express the following markers: HNF3 beta, GATA4, SOX17, Cerberus, OTX2, goosecoid, C-Kit, CD99, or MIXL1.
- [0047] “Extraembryonic endoderm”, as used herein, refers to a population of cells expressing at least one of the following markers: SOX7, AFP, or SPARC.
- [0048] “Markers”, as used herein, are nucleic acid or polypeptide molecules that are differentially expressed in a cell of interest. In this context, differential expression means an increased level for a positive marker and a decreased level for a negative marker. The detectable level of the marker nucleic acid or polypeptide is sufficiently higher or lower in the cells of interest compared to other cells, such that the cell of interest can be identified and distinguished from other cells using any of a variety of methods known in the art.
- [0049] “Mesendoderm cell”, as used herein, refers to a cell expressing at least one of the following markers: CD48, eomesodermin (EOMES), SOX17, DKK4, HNF3 beta, GSC, FGF17, or GATA6.
- [0050] “Pancreatic endocrine cell”, or “pancreatic hormone expressing cell”, as used herein, refers to a cell capable of expressing at least one of the following hormones: insulin, glucagon, somatostatin, or pancreatic polypeptide.

- [0051] “Pancreatic endoderm cell”, as used herein, refers to a cell capable of expressing at least one of the following markers: NGN3, NEUROD, ISL1, PDX1, PAX4, or NKX2.2.
- [0052] “Pancreatic hormone producing cell”, as used herein, refers to a cell capable of producing at least one of the following hormones: insulin, glucagon, somatostatin, or pancreatic polypeptide.
- [0053] “Pancreatic hormone secreting cell” as used herein, refers to a cell capable of secreting at least one of the following hormones: insulin, glucagon, somatostatin, and pancreatic polypeptide.
- [0054] “Posterior foregut cell”, as used herein, refers to a cell capable of secreting at least one of the following markers: PDX1, HNF1, PTF1 alpha, HNF6, HB9, or PROX1.
- [0055] “Pre-primitive streak cell”, as used herein, refers to a cell expressing at least one of the following markers: Nodal, or FGF8.
- [0056] “Primitive gut tube cell”, as used herein, refers to a cell capable of secreting at least one of the following markers: HNF1, or HNF4A.
- [0057] “Primitive streak cell”, as used herein, refers to a cell expressing at least one of the following markers: Brachyury, Mix-like homeobox protein, or FGF4.

Isolation, Expansion and Culture of Pluripotent Stem Cells

Characterization of Pluripotent Stem Cells

- [0058] Pluripotent stem cells may express one or more of the stage-specific embryonic antigens (SSEA) 3 and 4, and markers detectable using antibodies designated Tra-1-60 and Tra-1-81 (Thomson *et al.*, Science 282:1145, 1998). Differentiation of pluripotent stem cells *in vitro* results in the loss of SSEA-4, Tra-1-60, and Tra-1-81 expression (if present) and increased expression of SSEA-1. Undifferentiated pluripotent stem cells typically have alkaline phosphatase activity, which can be detected by fixing the cells with 4% paraformaldehyde, and then developing with Vector Red as a substrate, as described by the manufacturer (Vector Laboratories,

Burlingame Calif.) Undifferentiated pluripotent stem cells also typically express Oct-4 and TERT, as detected by RT-PCR.

[0059] Another desirable phenotype of propagated pluripotent stem cells is a potential to differentiate into cells of all three germinal layers: endoderm, mesoderm, and ectoderm tissues. Pluripotency of pluripotent stem cells can be confirmed, for example, by injecting cells into severe combined immunodeficient (SCID) mice, fixing the teratomas that form using 4% paraformaldehyde, and then examining them histologically for evidence of cell types from the three germ layers. Alternatively, pluripotency may be determined by the creation of embryoid bodies and assessing the embryoid bodies for the presence of markers associated with the three germinal layers.

[0060] Propagated pluripotent stem cell lines may be karyotyped using a standard G-banding technique and compared to published karyotypes of the corresponding primate species. It is desirable to obtain cells that have a "normal karyotype," which means that the cells are euploid, wherein all human chromosomes are present and not noticeably altered.

Sources of Pluripotent Stem Cells

[0061] The types of pluripotent stem cells that may be used include established lines of pluripotent cells derived from tissue formed after gestation, including pre-embryonic tissue (such as, for example, a blastocyst), embryonic tissue, or fetal tissue taken any time during gestation, typically but not necessarily before approximately 10-12 weeks gestation. Non-limiting examples are established lines of human embryonic stem cells or human embryonic germ cells, such as, for example the human embryonic stem cell lines H1, H7, and H9 (WiCell). Also contemplated is use of the compositions of this disclosure during the initial establishment or stabilization of such cells, in which case the source cells would be primary pluripotent cells taken directly from the source tissues. Also suitable are cells taken from a pluripotent stem cell population already cultured in the absence of feeder cells. Also suitable are mutant human embryonic stem cell lines, such as, for example, BG01v (BresaGen, Athens, GA).

- [0062] In one embodiment, human embryonic stem cells are prepared as described by Thomson *et al.* (U.S. Pat. No. 5,843,780; Science 282:1145, 1998; Curr. Top. Dev. Biol. 38:133 ff., 1998; Proc. Natl. Acad. Sci. U.S.A. 92:7844, 1995).

Culture of Pluripotent Stem Cells

- [0063] In one embodiment, pluripotent stem cells are typically cultured on a layer of feeder cells that support the pluripotent stem cells in various ways. Alternatively, pluripotent stem cells are cultured in a culture system that is essentially free of feeder cells, but nonetheless supports proliferation of pluripotent stem cells without undergoing substantial differentiation. The growth of pluripotent stem cells in feeder-free culture without differentiation is supported using a medium conditioned by culturing previously with another cell type. Alternatively, the growth of pluripotent stem cells in feeder-free culture without differentiation is supported using a chemically defined medium.
- [0064] For example, Reubinoff *et al.* (Nature Biotechnology 18: 399 - 404 (2000)) and Thompson *et al.* (Science 6 November 1998: Vol. 282. no. 5391, pp. 1145 – 1147) disclose the culture of pluripotent stem cell lines from human blastocysts using a mouse embryonic fibroblast feeder cell layer.
- [0065] Richards *et al.*, (Stem Cells 21: 546-556, 2003) evaluated a panel of 11 different human adult, fetal and neonatal feeder cell layers for their ability to support human pluripotent stem cell culture. Richards *et al.*, states: “human embryonic stem cell lines cultured on adult skin fibroblast feeders retain human embryonic stem cell morphology and remain pluripotent”.
- [0066] US20020072117 discloses cell lines that produce media that support the growth of primate pluripotent stem cells in feeder-free culture. The cell lines employed are mesenchymal and fibroblast-like cell lines obtained from embryonic tissue or differentiated from embryonic stem cells. US20020072117 also discloses the use of the cell lines as a primary feeder cell layer.
- [0067] In another example, Wang *et al.* (Stem Cells 23: 1221-1227, 2005) discloses methods for the long-term growth of human pluripotent stem cells on feeder cell layers derived from human embryonic stem cells.

- [0068] In another example, Stojkovic *et al* (Stem Cells 2005 23: 306-314, 2005) disclose a feeder cell system derived from the spontaneous differentiation of human embryonic stem cells.
- [0069] In a further example, Miyamoto *et al* (Stem Cells 22: 433-440, 2004) disclose a source of feeder cells obtained from human placenta.
- [0070] Amit *et al* (Biol. Reprod 68: 2150-2156, 2003) discloses a feeder cell layer derived from human foreskin.
- [0071] In another example, Inzunza *et al* (Stem Cells 23: 544-549, 2005) disclose a feeder cell layer from human postnatal foreskin fibroblasts.
- [0072] US6642048 discloses media that support the growth of primate pluripotent stem (pPS) cells in feeder-free culture, and cell lines useful for production of such media. US6642048 states: "This invention includes mesenchymal and fibroblast-like cell lines obtained from embryonic tissue or differentiated from embryonic stem cells. Methods for deriving such cell lines, processing media, and growing stem cells using the conditioned media are described and illustrated in this disclosure."
- [0073] In another example, WO2005014799 discloses conditioned medium for the maintenance, proliferation and differentiation of mammalian cells. WO2005014799 states: "The culture medium produced in accordance with the present invention is conditioned by the cell secretion activity of murine cells, in particular, those differentiated and immortalized transgenic hepatocytes, named MMH (Met Murine Hepatocyte)."
- [0074] In another example, Xu *et al* (Stem Cells 22: 972-980, 2004) discloses conditioned medium obtained from human embryonic stem cell derivatives that have been genetically modified to over express human telomerase reverse transcriptase.
- [0075] In another example, US20070010011 discloses a chemically defined culture medium for the maintenance of pluripotent stem cells.
- [0076] An alternative culture system employs serum-free medium supplemented with growth factors capable of promoting the proliferation of embryonic stem cells. For example, Cheon *et al* (BioReprod DOI:10.1095/biolreprod.105.046870, October 19, 2005)

disclose a feeder-free, serum-free culture system in which embryonic stem cells are maintained in unconditioned serum replacement (SR) medium supplemented with different growth factors capable of triggering embryonic stem cell self-renewal.

- [0077] In another example, Levenstein *et al* (Stem Cells 24: 568-574, 2006) disclose methods for the long-term culture of human embryonic stem cells in the absence of fibroblasts or conditioned medium, using media supplemented with bFGF.
- [0078] In another example, US20050148070 discloses a method of culturing human embryonic stem cells in defined media without serum and without fibroblast feeder cells, the method comprising: culturing the stem cells in a culture medium containing albumin, amino acids, vitamins, minerals, at least one transferrin or transferrin substitute, at least one insulin or insulin substitute, the culture medium essentially free of mammalian fetal serum and containing at least about 100 ng/ml of a fibroblast growth factor capable of activating a fibroblast growth factor signaling receptor, wherein the growth factor is supplied from a source other than just a fibroblast feeder layer, the medium supported the proliferation of stem cells in an undifferentiated state without feeder cells or conditioned medium.
- [0079] In another example, US20050233446 discloses a defined media useful in culturing stem cells, including undifferentiated primate primordial stem cells. In solution, the media is substantially isotonic as compared to the stem cells being cultured. In a given culture, the particular medium comprises a base medium and an amount of each of bFGF, insulin, and ascorbic acid necessary to support substantially undifferentiated growth of the primordial stem cells.
- [0080] In another example, US6800480 states “In one embodiment, a cell culture medium for growing primate-derived primordial stem cells in a substantially undifferentiated state is provided which includes a low osmotic pressure, low endotoxin basic medium that is effective to support the growth of primate-derived primordial stem cells. The basic medium is combined with a nutrient serum effective to support the growth of primate-derived primordial stem cells and a substrate selected from the group consisting of feeder cells and an extracellular matrix component derived from feeder cells. The medium further includes non-essential amino acids, an anti-oxidant, and a first growth factor selected from the group consisting of nucleosides and a pyruvate salt.”

- [0081] In another example, US20050244962 states: "In one aspect the invention provides a method of culturing primate embryonic stem cells. One cultures the stem cells in a culture essentially free of mammalian fetal serum (preferably also essentially free of any animal serum) and in the presence of fibroblast growth factor that is supplied from a source other than just a fibroblast feeder layer. In a preferred form, the fibroblast feeder layer, previously required to sustain a stem cell culture, is rendered unnecessary by the addition of sufficient fibroblast growth factor."
- [0082] In a further example, WO2005065354 discloses a defined, isotonic culture medium that is essentially feeder-free and serum-free, comprising: a. a basal medium; b. an amount of bFGF sufficient to support growth of substantially undifferentiated mammalian stem cells; c. an amount of insulin sufficient to support growth of substantially undifferentiated mammalian stem cells; and d. an amount of ascorbic acid sufficient to support growth of substantially undifferentiated mammalian stem cells.
- [0083] In another example, WO2005086845 discloses a method for maintenance of an undifferentiated stem cell, said method comprising exposing a stem cell to a member of the transforming growth factor-beta (TGF- β) family of proteins, a member of the fibroblast growth factor (FGF) family of proteins, or nicotinamide (NIC) in an amount sufficient to maintain the cell in an undifferentiated state for a sufficient amount of time to achieve a desired result.
- [0084] The pluripotent stem cells may be plated onto a suitable culture substrate. In one embodiment, the suitable culture substrate is an extracellular matrix component, such as, for example, those derived from basement membrane or that may form part of adhesion molecule receptor-ligand couplings. In one embodiment, a the suitable culture substrate is MATRIGEL® (Becton Dickenson). MATRIGEL® is a soluble preparation from Engelbreth-Holm Swarm tumor cells that gels at room temperature to form a reconstituted basement membrane.
- [0085] Other extracellular matrix components and component mixtures are suitable as an alternative. Depending on the cell type being proliferated, this may include laminin, fibronectin, proteoglycan, entactin, heparan sulfate, and the like, alone or in various combinations.

[0086] The pluripotent stem cells may be plated onto the substrate in a suitable distribution and in the presence of a medium that promotes cell survival, propagation, and retention of the desirable characteristics. All these characteristics benefit from careful attention to the seeding distribution and can readily be determined by one of skill in the art.

[0087] Suitable culture media may be made from the following components, such as, for example, Dulbecco's modified Eagle's medium (DMEM), Gibco # 11965-092; Knockout Dulbecco's modified Eagle's medium (KO DMEM), Gibco #10829-018; Ham's F12/50% DMEM basal medium; 200 mM L-glutamine, Gibco # 15039-027; non-essential amino acid solution, Gibco 11140-050; β -mercaptoethanol, Sigma # M7522; human recombinant basic fibroblast growth factor (bFGF), Gibco # 13256-029.

Formation of Pancreatic Hormone Producing Cells from Pluripotent Stem Cells

[0088] In one embodiment, the present invention provides a method for producing pancreatic hormone producing cells from pluripotent stem cells, comprising the steps of:

- a. Culturing pluripotent stem cells,
- b. Differentiating the pluripotent stem cells into cells expressing markers characteristic of the definitive endoderm lineage,
- c. Differentiating the cells expressing markers characteristic of the definitive endoderm lineage into cells expressing markers characteristic of the pancreatic endoderm lineage, and
- d. Differentiating the cells expressing markers characteristic of the pancreatic endoderm lineage into cells expressing markers characteristic of the pancreatic endocrine lineage.

[0089] Pluripotent stem cells suitable for use in the present invention include, for example, the human embryonic stem cell line H9 (NIH code: WA09), the human embryonic stem cell line H1 (NIH code: WA01), the human embryonic stem cell line H7 (NIH code: WA07), and the human embryonic stem cell line SA002 (Cellartis, Sweden). Also suitable for use in the present invention are cells that express at least one of the

following markers characteristic of pluripotent cells: ABCG2, cripto, CD9, FOXD3, Connexin43, Connexin45, OCT4, SOX2, NANOG, hTERT, UTF1, ZFP42, SSEA3, SSEA4, Tra1-60, or Tra1-81.

[0090] Markers characteristic of the definitive endoderm lineage are selected from the group consisting of SOX17, GATA4, HNF3 beta, GSC, CER1, NODAL, FGF8, Brachyury, Mix-like homeobox protein, FGF4 CD48, comesodermin (EOMES), DKK4, FGF17, GATA6, CXCR4, C-Kit, CD99, and OTX2. Suitable for use in the present invention is a cell that expresses at least one of the markers characteristic of the definitive endoderm lineage. In one aspect of the present invention, a cell expressing markers characteristic of the definitive endoderm lineage is a primitive streak precursor cell. In an alternate aspect, a cell expressing markers characteristic of the definitive endoderm lineage is a mesendoderm cell. In an alternate aspect, a cell expressing markers characteristic of the definitive endoderm lineage is a definitive endoderm cell.

[0091] Markers characteristic of the pancreatic endoderm lineage are selected from the group consisting of PDX1, HNF1 beta, PTF1 alpha, HNF6, HB9 and PROX1. Suitable for use in the present invention is a cell that expresses at least one of the markers characteristic of the pancreatic endoderm lineage. In one aspect of the present invention, a cell expressing markers characteristic of the pancreatic endoderm lineage is a pancreatic endoderm cell.

[0092] Markers characteristic of the pancreatic endocrine lineage are selected from the group consisting of NGN3, NEUROD, ISL1, PDX1, NKX6.1, PAX4, NGN3, and PTF1 alpha. In one embodiment, a pancreatic endocrine cell is capable of expressing at least one of the following hormones: insulin, glucagon, somatostatin, and pancreatic polypeptide. Suitable for use in the present invention is a cell that expresses at least one of the markers characteristic of the pancreatic endocrine lineage. In one aspect of the present invention, a cell expressing markers characteristic of the pancreatic endocrine lineage is a pancreatic endocrine cell. The pancreatic endocrine cell may be a pancreatic hormone expressing cell. Alternatively, the pancreatic endocrine cell may be a pancreatic hormone secreting cell.

- [0093] In one aspect of the present invention, the pancreatic endocrine cell is a cell expressing markers characteristic of the β cell lineage. A cell expressing markers characteristic of the β cell lineage expresses Pdx1 and at least one of the following transcription factors: NGN3, NKX2.2, NKX6.1, NEUROD, ISL1, HNF3 beta, MAFA, PAX4, or PAX6. In one aspect of the present invention, a cell expressing markers characteristic of the β cell lineage is a β cell.

**Formation of Cells Expressing Markers Characteristic of the Definitive
Endoderm Lineage**

- [0094] Pluripotent stem cells may be differentiated into cells expressing markers characteristic of the definitive endoderm lineage by any method in the art or by any method proposed in this invention.
- [0095] For example, pluripotent stem cells may be differentiated into cells expressing markers characteristic of the definitive endoderm lineage according to the methods disclosed in D'Amour *et al*, Nature Biotechnology 23, 1534 – 1541 (2005).
- [0096] For example, pluripotent stem cells may be differentiated into cells expressing markers characteristic of the definitive endoderm lineage according to the methods disclosed in Shinozaki *et al*, Development 131, 1651 - 1662 (2004).
- [0097] For example, pluripotent stem cells may be differentiated into cells expressing markers characteristic of the definitive endoderm lineage according to the methods disclosed in McLean *et al*, Stem Cells 25, 29 - 38 (2007).
- [0098] For example, pluripotent stem cells may be differentiated into cells expressing markers characteristic of the definitive endoderm lineage according to the methods disclosed in D'Amour *et al*, Nature Biotechnology 24, 1392 – 1401 (2006).
- [0099] For example, pluripotent stem cells may be differentiated into cells expressing markers characteristic of the definitive endoderm lineage by culturing the pluripotent stem cells in medium containing activin A in the absence of serum, then culturing the cells with activin A and serum, and then culturing the cells with activin A and serum of a different concentration. An example of this method is disclosed in Nature Biotechnology 23, 1534 - 1541 (2005).

- [0100] For example, pluripotent stem cells may be differentiated into cells expressing markers characteristic of the definitive endoderm lineage by culturing the pluripotent stem cells in medium containing activin A in the absence of serum, then culturing the cells with activin A with serum of another concentration. An example of this method is disclosed in D' Amour *et al*, Nature Biotechnology, 2005.
- [0101] For example, pluripotent stem cells may be differentiated into cells expressing markers characteristic of the definitive endoderm lineage by culturing the pluripotent stem cells in medium containing activin A and a Wnt ligand in the absence of serum, then removing the Wnt ligand and culturing the cells with activin A with serum. An example of this method is disclosed in Nature Biotechnology 24, 1392 - 1401 (2006).
- [0102] For example, pluripotent stem cells may be differentiated into cells expressing markers characteristic of the definitive endoderm lineage by treating the pluripotent stem cells according to the methods disclosed in US patent application Ser. No. 11/736,908, assigned to LifeScan, Inc.
- [0103] For example, pluripotent stem cells may be differentiated into cells expressing markers characteristic of the definitive endoderm lineage by treating the pluripotent stem cells according to the methods disclosed in US patent application Ser. No. 11/779,311, assigned to LifeScan, Inc.
- [0104] For example, pluripotent stem cells may be differentiated into cells expressing markers characteristic of the definitive endoderm lineage by treating the pluripotent stem cells according to the methods disclosed in US patent application Ser. No. 60/990,529.
- [0105] For example, pluripotent stem cells may be differentiated into cells expressing markers characteristic of the definitive endoderm lineage by treating the pluripotent stem cells according to the methods disclosed in US patent application Ser. No. 61/076,889.
- [0106] For example, pluripotent stem cells may be differentiated into cells expressing markers characteristic of the definitive endoderm lineage by treating the pluripotent stem cells according to the methods disclosed in US patent application Ser. No. 61/076,900.

[0107] For example, pluripotent stem cells may be differentiated into cells expressing markers characteristic of the definitive endoderm lineage by treating the pluripotent stem cells according to the methods disclosed in US patent application Ser. No. 61/076,908.

[0108] For example, pluripotent stem cells may be differentiated into cells expressing markers characteristic of the definitive endoderm lineage by treating the pluripotent stem cells according to the methods disclosed in US patent application Ser. No. 61/076,915.

*Differentiation of Cells Expressing Markers Characteristic of the Definitive
Endoderm Lineage*

[0109] Formation of cells expressing markers characteristic of the definitive endoderm lineage may be determined by testing for the presence of the markers before and after following a particular protocol. Pluripotent stem cells typically do not express such markers. Thus, differentiation of pluripotent cells is detected when cells begin to express them.

[0110] The efficiency of differentiation may be determined by exposing a treated cell population to an agent (such as an antibody) that specifically recognizes a protein marker expressed by cells expressing markers characteristic of the definitive endoderm lineage.

[0111] Methods for assessing expression of protein and nucleic acid markers in cultured or isolated cells are standard in the art. These include quantitative reverse transcriptase polymerase chain reaction (RT-PCR), Northern blots, *in situ* hybridization (see, e.g., Current Protocols in Molecular Biology (Ausubel *et al.*, eds. 2001 supplement)), and immunoassays such as immunohistochemical analysis of sectioned material, Western blotting, and for markers that are accessible in intact cells, flow cytometry analysis (FACS) (see, e.g., Harlow and Lane, Using Antibodies: A Laboratory Manual, New York: Cold Spring Harbor Laboratory Press (1998)).

[0112] Characteristics of pluripotent stem cells are well known to those skilled in the art, and additional characteristics of pluripotent stem cells continue to be identified. Pluripotent stem cell markers include, for example, the expression of one or more of

the following: ABCG2, *cripto*, FOXD3, Connexin43, Connexin45, OCT4, SOX2, NANOG, hTERT, UTF1, ZFP42, SSEA3, SSEA4, Tra1-60, or Tra1-81.

- [0113] After treating pluripotent stem cells with the methods of the present invention, the differentiated cells may be purified by exposing a treated cell population to an agent (such as an antibody) that specifically recognizes a protein marker, such as CXCR4, expressed by cells expressing markers characteristic of the definitive endoderm lineage.

Formation of Cells Expressing Markers Characteristic of the Pancreatic Endoderm Lineage

- [0114] Cells expressing markers characteristic of the definitive endoderm lineage may be differentiated into cells expressing markers characteristic of the pancreatic endoderm lineage by any method in the art or by any method proposed in this invention.
- [0115] For example, cells expressing markers characteristic of the definitive endoderm lineage may be differentiated into cells expressing markers characteristic of the pancreatic endoderm lineage according to the methods disclosed in D'Amour *et al*, Nature Biotechnology 24, 1392 - 1401 (2006).
- [0116] For example, cells expressing markers characteristic of the definitive endoderm lineage are further differentiated into cells expressing markers characteristic of the pancreatic endoderm lineage, by treating the cells expressing markers characteristic of the definitive endoderm lineage with a fibroblast growth factor and the hedgehog signaling pathway inhibitor KAAD-cyclopamine, then removing the medium containing the fibroblast growth factor and KAAD-cyclopamine and subsequently culturing the cells in medium containing retinoic acid, a fibroblast growth factor and KAAD-cyclopamine. An example of this method is disclosed in Nature Biotechnology 24, 1392 - 1401 (2006).
- [0117] In one aspect of the present invention, cells expressing markers characteristic of the definitive endoderm lineage are further differentiated into cells expressing markers characteristic of the pancreatic endoderm lineage, by treating the cells expressing markers characteristic of the definitive endoderm lineage with retinoic acid and at

least one fibroblast growth factor for a period of time, according to the methods disclosed in US patent application Ser. No. 11/736,908, assigned to LifeScan, Inc.

[0118] In one aspect of the present invention, cells expressing markers characteristic of the definitive endoderm lineage are further differentiated into cells expressing markers characteristic of the pancreatic endoderm lineage, by treating the cells expressing markers characteristic of the definitive endoderm lineage with retinoic acid and at least one fibroblast growth factor for a period of time, according to the methods disclosed in US patent application Ser. No. 11/779,311, assigned to LifeScan, Inc.

[0119] In one aspect of the present invention, cells expressing markers characteristic of the definitive endoderm lineage are further differentiated into cells expressing markers characteristic of the pancreatic endoderm lineage, by treating the cells expressing markers characteristic of the definitive endoderm lineage according to the methods disclosed in US patent application Ser. No. 60/990,529.

Detection of Cells Expressing Markers Characteristic of the Pancreatic Endoderm Lineage

[0120] Markers characteristic of the pancreatic endoderm lineage are well known to those skilled in the art, and additional markers characteristic of the pancreatic endoderm lineage continue to be identified. These markers can be used to confirm that the cells treated in accordance with the present invention have differentiated to acquire the properties characteristic of the pancreatic endoderm lineage. Pancreatic endoderm lineage specific markers include the expression of one or more transcription factors such as, for example, HLXB9, PTF1 alpha, PDX1, HNF6, or HNF1 beta.

[0121] The efficiency of differentiation may be determined by exposing a treated cell population to an agent (such as an antibody) that specifically recognizes a protein marker expressed by cells expressing markers characteristic of the pancreatic endoderm lineage.

[0122] Methods for assessing expression of protein and nucleic acid markers in cultured or isolated cells are standard in the art. These include quantitative reverse transcriptase polymerase chain reaction (RT-PCR), Northern blots, *in situ* hybridization (see, e.g., Current Protocols in Molecular Biology (Ausubel *et al.*, eds. 2001 supplement)), and

immunoassays such as immunohistochemical analysis of sectioned material, Western blotting, and for markers that are accessible in intact cells, flow cytometry analysis (FACS) (see, e.g., Harlow and Lane, *Using Antibodies: A Laboratory Manual*, New York: Cold Spring Harbor Laboratory Press (1998)).

Formation of Cells Expressing Markers Characteristic of the Pancreatic Endocrine Lineage

- [0123] Cells expressing markers characteristic of the pancreatic endoderm lineage may be differentiated into cells expressing markers characteristic of the pancreatic endocrine lineage by any method in the art or by any method disclosed in this invention.
- [0124] For example, cells expressing markers characteristic of the pancreatic endoderm lineage may be differentiated into cells expressing markers characteristic of the pancreatic endocrine lineage according to the methods disclosed in D'Amour *et al*, *Nature Biotechnology* 24, 1392 - 1401 (2006).
- [0125] For example, cells expressing markers characteristic of the pancreatic endoderm lineage are further differentiated into cells expressing markers characteristic of the pancreatic endocrine lineage, by culturing the cells expressing markers characteristic of the pancreatic endoderm lineage in medium containing DAPT and exendin 4, then removing the medium containing DAPT and exendin 4 and subsequently culturing the cells in medium containing exendin 1, IGF-1 and HGF. An example of this method is disclosed in *Nature Biotechnology* 24, 1392 - 1401 (2006).
- [0126] For example, cells expressing markers characteristic of the pancreatic endoderm lineage are further differentiated into cells expressing markers characteristic of the pancreatic endocrine lineage, by culturing the cells expressing markers characteristic of the pancreatic endoderm lineage in medium containing exendin 4, then removing the medium containing exendin 4 and subsequently culturing the cells in medium containing exendin 1, IGF-1 and HGF. An example of this method is disclosed in D'Amour *et al*, *Nature Biotechnology*, 2006.
- [0127] For example, cells expressing markers characteristic of the pancreatic endoderm lineage are further differentiated into cells expressing markers characteristic of the pancreatic endocrine lineage, by culturing the cells expressing markers characteristic

of the pancreatic endoderm lineage in medium containing DAPT and exendin 4. An example of this method is disclosed in D' Amour *et al*, Nature Biotechnology, 2006.

- [0128] For example, cells expressing markers characteristic of the pancreatic endoderm lineage are further differentiated into cells expressing markers characteristic of the pancreatic endocrine lineage, by culturing the cells expressing markers characteristic of the pancreatic endoderm lineage in medium containing exendin 4. An example of this method is disclosed in D' Amour *et al*, Nature Biotechnology, 2006.
- [0129] In one aspect of the present invention, cells expressing markers characteristic of the pancreatic endoderm lineage are further differentiated into cells expressing markers characteristic of the pancreatic endocrine lineage, by treating the cells expressing markers characteristic of the pancreatic endoderm lineage with a factor that inhibits the Notch signaling pathway, according to the methods disclosed in US patent application Ser. No. 11/736,908, assigned to LifeScan, Inc.
- [0130] In one aspect of the present invention, cells expressing markers characteristic of the pancreatic endoderm lineage are further differentiated into cells expressing markers characteristic of the pancreatic endocrine lineage, by treating the cells expressing markers characteristic of the pancreatic endoderm lineage with a factor that inhibits the Notch signaling pathway, according to the methods disclosed in US patent application Ser. No. 11/779,311, assigned to LifeScan, Inc.
- [0131] In one aspect of the present invention, cells expressing markers characteristic of the pancreatic endoderm lineage are further differentiated into cells expressing markers characteristic of the pancreatic endocrine lineage, by treating the cells expressing markers characteristic of the pancreatic endoderm lineage with a factor that inhibits the Notch signaling pathway, according to the methods disclosed in US patent application Ser. No. 60/953,178, assigned to LifeScan, Inc.
- [0132] In one aspect of the present invention, cells expressing markers characteristic of the pancreatic endoderm lineage are further differentiated into cells expressing markers characteristic of the pancreatic endocrine lineage, by treating the cells expressing markers characteristic of the pancreatic endoderm lineage according to the methods disclosed in US patent application Ser. No. 60/990,529.

- [0133] In one aspect of the present invention, the present invention provides a method for increasing the expression of markers associated with the pancreatic endocrine lineage comprising treating cells expressing markers characteristic of the pancreatic endocrine lineage with medium comprising a sufficient amount of a TGF- β receptor agonist to cause an increase in expression of markers associated with the pancreatic endocrine lineage according to the methods disclosed in US patent application Ser. No. 61/110,278.

Detection of Cells Expressing Markers Characteristic of the Pancreatic Endocrine Lineage

- [0134] Markers characteristic of cells of the pancreatic endocrine lineage are well known to those skilled in the art, and additional markers characteristic of the pancreatic endocrine lineage continue to be identified. These markers can be used to confirm that the cells treated in accordance with the present invention have differentiated to acquire the properties characteristic of the pancreatic endocrine lineage. Pancreatic endocrine lineage specific markers include the expression of one or more transcription factors such as, for example, NGN3, NEUROD, or ISL1.
- [0135] Markers characteristic of cells of the β cell lineage are well known to those skilled in the art, and additional markers characteristic of the β cell lineage continue to be identified. These markers can be used to confirm that the cells treated in accordance with the present invention have differentiated to acquire the properties characteristic of the β -cell lineage. β cell lineage specific characteristics include the expression of one or more transcription factors such as, for example, PDX1, NKX2.2, NKX6.1, ISL1, PAX6, PAX4, NEUROD, HNF1 beta, HNF6, HNF3 beta, or MAFA, among others. These transcription factors are well established in the art for identification of endocrine cells. See, e.g., Edlund (Nature Reviews Genetics 3: 524-632 (2002)).
- [0136] The efficiency of differentiation may be determined by exposing a treated cell population to an agent (such as an antibody) that specifically recognizes a protein marker expressed by cells expressing markers characteristic of the pancreatic endocrine lineage. Alternatively, the efficiency of differentiation may be determined by exposing a treated cell population to an agent (such as an antibody) that

specifically recognizes a protein marker expressed by cells expressing markers characteristic of the β cell lineage.

- [0137] Methods for assessing expression of protein and nucleic acid markers in cultured or isolated cells are standard in the art. These include quantitative reverse transcriptase polymerase chain reaction (RT-PCR), Northern blots, *in situ* hybridization (see, e.g., Current Protocols in Molecular Biology (Ausubel *et al.*, eds. 2001 supplement)), and immunoassays such as immunohistochemical analysis of sectioned material, Western blotting, and for markers that are accessible in intact cells, flow cytometry analysis (FACS) (see, e.g., Harlow and Lane, Using Antibodies: A Laboratory Manual, New York: Cold Spring Harbor Laboratory Press (1998)).
- [0138] In one aspect of the present invention, the efficiency of differentiation is determined by measuring the percentage of insulin positive cells in a given cell culture following treatment. In one embodiment, the methods of the present invention produce about 100% insulin positive cells in a given culture. In an alternate embodiment, the methods of the present invention produce about 90% insulin positive cells in a given culture. In an alternate embodiment, the methods of the present invention produce about 80% insulin positive cells in a given culture. In an alternate embodiment, the methods of the present invention produce about 70% insulin positive cells in a given culture. In an alternate embodiment, the methods of the present invention produce about 60% insulin positive cells in a given culture. In an alternate embodiment, the methods of the present invention produce about 50% insulin positive cells in a given culture. In an alternate embodiment, the methods of the present invention produce about 40% insulin positive cells in a given culture. In an alternate embodiment, the methods of the present invention produce about 30% insulin positive cells in a given culture. In an alternate embodiment, the methods of the present invention produce about 20% insulin positive cells in a given culture. In an alternate embodiment, the methods of the present invention produce about 10% insulin positive cells in a given culture. In an alternate embodiment, the methods of the present invention produce about 5% insulin positive cells in a given culture.
- [0139] In one aspect of the present invention, the efficiency of differentiation is determined by measuring glucose-stimulated insulin secretion, as detected by measuring the amount of C-peptide released by the cells. In one embodiment, cells produced by the

methods of the present invention produce about 1000ng C-peptide/pg DNA. In an alternate embodiment, cells produced by the methods of the present invention produce about 900ng C-peptide/pg DNA. In an alternate embodiment, cells produced by the methods of the present invention produce about 800ng C-peptide/pg DNA. In an alternate embodiment, cells produced by the methods of the present invention produce about 700ng C-peptide/pg DNA. In an alternate embodiment, cells produced by the methods of the present invention produce about 600ng C-peptide/pg DNA. In an alternate embodiment, cells produced by the methods of the present invention produce about 500ng C-peptide/pg DNA. In an alternate embodiment, cells produced by the methods of the present invention produce about 400ng C-peptide/pg DNA. In an alternate embodiment, cells produced by the methods of the present invention produce about 500ng C-peptide/pg DNA. In an alternate embodiment, cells produced by the methods of the present invention produce about 400ng C-peptide/pg DNA. In an alternate embodiment, cells produced by the methods of the present invention produce about 300ng C-peptide/pg DNA. In an alternate embodiment, cells produced by the methods of the present invention produce about 200ng C-peptide/pg DNA. In an alternate embodiment, cells produced by the methods of the present invention produce about 100ng C-peptide/pg DNA. In an alternate embodiment, cells produced by the methods of the present invention produce about 90ng C-peptide/pg DNA. In an alternate embodiment, cells produced by the methods of the present invention produce about 80ng C-peptide/pg DNA. In an alternate embodiment, cells produced by the methods of the present invention produce about 70ng C-peptide/pg DNA. In an alternate embodiment, cells produced by the methods of the present invention produce about 60ng C-peptide/pg DNA. In an alternate embodiment, cells produced by the methods of the present invention produce about 50ng C-peptide/pg DNA. In an alternate embodiment, cells produced by the methods of the present invention produce about 40ng C-peptide/pg DNA. In an alternate embodiment, cells produced by the methods of the present invention produce about 30ng C-peptide/pg DNA. In an alternate embodiment, cells produced by the methods of the present invention produce about 20ng C-peptide/pg DNA. In an alternate embodiment, cells produced by the methods of the present invention produce about 10ng C-peptide/pg DNA.

Increasing Expression of MAFA in Cells Expressing Markers Characteristic of the Pancreatic Endocrine Lineage

- [0140]** In one embodiment, the present invention provides a method for increasing the expression of MAFA in cells expressing markers characteristic of the pancreatic endocrine lineage comprising the steps of culturing the cells expressing markers characteristic of the pancreatic endocrine lineage in medium comprising a sufficient amount of a cyclin-dependant kinase inhibitor to cause an increase in expression of MAFA.
- [0141]** The cyclin dependant kinase inhibitor may inhibit cyclin dependant kinase 1. Alternatively, the cyclin dependant kinase inhibitor may inhibit cyclin dependant kinase 2. Alternatively, the cyclin dependant kinase inhibitor may inhibit cyclin dependant kinase 4. Alternatively, the cyclin dependant kinase inhibitor may inhibit cyclin dependant kinase 5. Alternatively, the cyclin dependant kinase inhibitor may inhibit cyclin dependant kinase 9. Alternatively, the cyclin dependant kinase inhibitor may inhibit multiple isoforms of cyclin dependant kinase, in any combination thereof.
- [0142]** The cyclin dependant kinase inhibitor may be a protein. Alternatively, the cyclin dependant kinase inhibitor may be a peptide. Alternatively, the cyclin dependant kinase inhibitor may be a small molecule. In one embodiment, the small molecule cyclin dependant kinase inhibitor is selected from the group consisting of 7-n-Butyl-6-(4-hydroxyphenyl)[5H]pyrrolo[2,3-b]pyrazine, 9-Nitro-7,12-dihydroindolo[3,2-d][1]benzazepin-6(5H)-one, 3-(6-Oxo-9-nitro-5,6,7,12-tetrahydroindolo[3,2-d][1]benzazepin-2-yl)propionitrile, (2R)-2-(((3-Amino-5-chlorophenyl)amino)-9-(1-methylethyl)-9H-purin-2-yl)amino)-3-methyl-1-butanol, Arcyriaflavin A, [6-Benzylamino-2-(3-hydroxypropylamino)-9-isopropylpurine, Butyrolactone I, (Z)-1-(3-Ethyl-5-methoxy-2,3-dihydrobenzothiazol-2-ylidene)propan-2-one, 2-(3-Hydroxypropylamino)-6-(o-hydroxybenzylamino)-9-isopropylpurine, 1-(2,6-Dichlorophenyl)-1,5-dihydro-6-((4-(2-hydroxyethoxy)phenyl)methyl)-3-(1-methylethyl)-4H-pyrazolo[3,4-d]pyrimidin-4-one, Cdk/ Cyclin Inhibitory Peptide III, 3-(2-Chloro-3-indolylmethylene)-1,3-dihydroindol-2-one, Ethyl-(6-hydroxy-4-phenylbenzo[4,5]furo[2,3-b])pyridine-3-carboxylate, RO-3306, N-(cis-2-Aminocyclohexyl)-N-(3-chlorophenyl)-9-ethyl-9H-purine-2,6-diamine, 6-Cyclohexylmethoxy-2-(4'-sulfamoylanilino)purine, 5-Amino-3-((4-

(aminosulfonyl)phenyl)amino)-N-(2,6-difluorophenyl)-1H-1,2,4-triazole-1-carbothioamide, 3-Amino-1H-pyrazolo[3,4-b]quinoxaline, Cdk2 Inhibitor I, Cdk2 Inhibitor II, 2(bis-(Hydroxyethyl)amino)-6-(4-methoxybenzylamino)-9-isopropylpurine, 4-(6-Cyclohexylmethoxy-9H-purin-2-ylamino)-N,N-diethylbenzamide, N4-(6-Aminopyrimidin-4-yl)-sulfanilamide, (4-(2-Amino-4-methylthiazol-5-yl)pyrimidin-2-yl)-(3-nitrophenyl)amine, 2-Bromo-12,13-dihydro-5H-indolo[2,3-a]pyrrolo[3,4-c]carbazole-5,7(6H)-dione, 1,4-Dimethoxyacridine-9(10H)-thione, 5-(N-(4-Methylphenyl)amino)-2-methyl-4,7-dioxobenzothiazole, 4-(3,5-Diamino-1Hpyrazol-4-ylazo)-phenol, 2-(2-Hydroxyethylamino)-6-(3-chloroanilino)-9-isopropylpurine, Fascaplysin, Indirubin-3'-monoxime, Indirubin-3'-monoxime, 5-Iodo-, Indirubin-3'-monoxime-5-sulphonic Acid, Isogranulatimide, 2-(2-Hydroxyethylamino)-6-benzylamino-9-methylpurine, 6-(2-Hydroxybenzylamino)-2-((1R)-(hydroxymethyl)propyl)amino)-9-isopropylpurine, 5-Bromo-3-(2-(4-Fluorophenyl)-2-oxoethylidene)-1,3-dihydroindol-2-one, N6,N6-Dimethyladenine, 2-(1R-Isopropyl-2-hydroxyethylamino)-6-(3-chloroanilino)-9-isopropyl-purine, rapamycin, 2-(R)-(1-Ethyl-2-hydroxyethylamino)-6-benzylamino-9-isopropylpurine, Scytonemin, 3-[1-(3H-Imidazol-4-yl)-meth-(Z)-ylidene]-5-methoxy-1,3-dihydroindol-2-one, and 4-(3'-Hydroxyphenyl)amino-6,7-dimethoxyquinazoline.

- [0143] In one embodiment, the cyclin dependant kinase is ethyl-(6-hydroxy-4-phenylbenzo[4,5]furo[2,3-b])pyridine-3-carboxylate. In one embodiment, ethyl-(6-hydroxy-4-phenylbenzo[4,5]furo[2,3-b])pyridine-3-carboxylate is added to cells expressing markers characteristic of the endocrine lineage at a concentration from about 0.1 μ M to about 10 μ M for about one to seven days.
- [0144] In one embodiment, cells expressing markers characteristic of the endocrine lineage are treated with ethyl-(6-hydroxy-4-phenylbenzo[4,5]furo[2,3-b])pyridine-3-carboxylate for about one to about seven days.
- [0145] The present invention is further illustrated, but not limited by, the following examples.

EXAMPLES

Example 1

**Differentiation of Human Embryonic Stem Cells of the Cell Line H1 to
Pancreatic Endocrine Cells in the Absence of Fetal Bovine Serum**

- [0146]** Cells of the human embryonic stem cells line H1 at passage 52 were cultured on MATRIGEL[®]-coated dishes (1:30 dilution) and exposed to the following differentiation protocol, in order to differentiate the cells to cells expressing markers characteristic of the pancreatic endocrine lineage.
- a. RPMI medium supplemented with 2% BSA (Catalog# 152401, MP Biomedical, Ohio), and 100 ng/ml activin A (R&D Systems, MN) plus 20 ng/ml WNT-3a (Catalog# 1324-WN-002, R&D Systems, MN) plus 8 ng/ml of bFGF (Catalog# 100-18B, PeproTech, NJ), for one day followed by treatment with RPMI media supplemented with 2% BSA and 100 ng/ml activin A plus 8 ng/ml of bFGF for an additional two days (Stage 1), then
 - b. DMEM/F12 + 2% BSA + 50 ng/ml FGF7 + 0.25 μ M Cyclopamine- KAAD (#239804, Calbiochem, CA) for two days (Stage 2), then
 - c. DMEM/F12 + 1% B27 (Invitrogen, CA) + 50 ng/ml FGF7 + 0.25 μ M Cyclopamine- KAAD + 2 μ M Retinoic acid (RA) (Sigma, MO) + 100 ng/ml of Noggin (R & D Systems, MN) for four days (Stage 3), then
 - d. DMEM/F12 + 1% B27 (Invitrogen, CA) + 100 ng/ml Noggin + 1 μ M DAPT (a gamma-secretase inhibitor) (Catalog# 565784, Calbiochem, CA) + 1 μ M ALK5 inhibitor II (Catalog# 616452, Calbiochem, Ca) + 100 ng/ml of Netrin-4 (R&D Systems, MN) for three days (Stage 4), then
 - e. DMEM/F12 + 1% B27 (Invitrogen, CA) + 1 μ M ALK5 inhibitor II (Calbiochem, Ca) for seven days (Stage 5).
- [0147]** Medium was changed daily. At each stage the cell number was calculated using a hemocytometer and RNA was collected for PCR analysis. All samples were collected in triplicate.

Example 2

Screening of the Effects of Compounds from the EMD Kinase Inhibitor Library II on Cells that have been Treated according to the Differentiation Protocol Outlined in Example 1

- [0148]** Cells of the human embryonic stem cell line H1 at passage 44 were seeded onto MATRIGEL™ coated 24-well dishes (1:30 dilution), and differentiated according to the methods described in Example 1 up to stage 5. Following this, the cells were treated for four days in DMEM/F12 + 1% B27 containing a compound from an EMD Calbiochem compound library (Catalog# 539745, Calbiochem, San Diego, Ca) at a final concentration of 1μM. Wells containing vehicle were included as a control. Throughout the protocol media was changed daily. All samples were treated in duplicate. At the completion of this treatment RNA was collected for PCR analysis. Samples were analyzed by real-time PCR for expression of insulin, glucagon, MAFA, and Arx4. Results are expressed as a ratio insulin/glucagon (Figure 1, panel a), or MAFA versus ARX4 (Figure 1, panel b) of the treated samples relative to the untreated control, as measured by real-time PCR. The corresponding PubChem Compound ID# for each well # is listed in Table 1.
- [0149]** Treatment of cells expressing markers characteristic of the pancreatic endocrine lineage with compounds A6, B7, B8, or C2 at a 1μM concentration resulted in an insulin/ glucagon expression ratio of approximately 3.0 or higher (see Figure 1, panel a).
- [0150]** We next examined the effect of these compounds on the ratio of MAFA/ARX4, and we observed that treatment of cells expressing markers characteristic of the pancreatic endocrine lineage with several of the compounds resulted in a much greater change in the ratio of MAFA to ARX4 than other compounds tested in the library: Cells treated with compound C2 showed a ratio of MAFA/ARX4 of approximately 1000. Treatment of cells expressing markers characteristic of the pancreatic endocrine lineage with compound C2 resulted in a MAFA/ARX4 ratio of approximately 100. (See Figure 1, panel b).

Example 3

The Effects of Cyclin Dependant Kinase Inhibitor Treatment on Insulin and MAFA Expression in Cells that have been Treated according to the Differentiation Protocol Outlined in Example 1

- [0151]** Several of the compounds that increased the ratio of insulin to glucagon expression, or MAFA to ARX4 expression in Example 2 were cyclin dependant kinase inhibitors. One such compound was PubChem Compound ID# 5330797 (5-Amino-3-((4-(aminosulfonyl)phenyl)amino)-N-(2,6-difluorophenyl)-1H-1,2,4-triazole-1-carbothioamide) (Catalogue # 217714; Calbiochem, San Diego, Ca). To confirm these observations, cells of the human embryonic stem cell line H1 at passage 42 were cultured in 10cm² MATRIGEL[®]-coated dishes and treated according to the methods described in Example 1 up to stage 5. After stage 5, the cells were treated with DMEM/F12 containing 1% B27 containing 1µM PubChem Compound ID# 5330797 for six days. Medium was changed every other day. Samples of cells were taken for real-time PCR prior to treatment with the compound, and at days two and five of compound treatment.
- [0152]** Characteristic micrographs of the cells at day 4 or day 6 of compound treatment versus untreated controls are shown in Figure 2. Untreated cells are highly packed (Figure 3, panels a and d) and it is difficult to distinguish individual cells. However, after treatment with 0.5µM or 1µM of PubChem Compound ID#5330797 for six days, individual nuclei became visible (Figures 2, panels e and f) as compared to the untreated control (Figure 2, panel d), indicating that there was differentiation occurring in the cell population. This was also accompanied by some cell death, which can be seen by gaps in the layer of cells as shown in Figure 2, panels b and c.
- [0153]** Treatment of cells with PubChem Compound ID# 5330797 resulted in the increase in expression of insulin, glucagon, MAFA, MAFB and somatostatin, albeit to differing degrees. The relative induction of gene expression per treatment as compared to day 0 (pretreatment) cultures is shown in Figures 3, panels a-v. Cells expressing markers characteristic of the pancreatic endocrine lineage that were treated with 1µM PubChem Compound ID# 5330797 resulted in an approximately 1.5 fold increase in glucagon expression at 48 hrs of treatment. This expression declined to below

pretreatment levels after 5 days of treatment. No increase in glucagon expression was observed with treatment of 0.5 μ M PubChem Compound ID# 5330797. (See Figure 3, panel a).

[0154] Cells expressing markers characteristic of the pancreatic endocrine lineage that were treated with 1 μ M PubChem Compound ID# 5330797 for five days resulted in an approximately 1.5 fold increase in insulin expression. (See Figure 3, panel b).

[0155] Cells expressing markers characteristic of the pancreatic endocrine lineage that were treated with 1 μ M PubChem Compound ID# 5330797 for five days resulted in an approximately 200 fold increase in MAFA expression. (See Figure 3, panel d).

[0156] Cells expressing markers characteristic of the pancreatic endocrine lineage that were treated with 0.5 μ M PubChem Compound ID# 5330797 for five days resulted in an approximately 1.5 fold increase in MAFB expression. (See Figure 3, panel c). A dose-dependent increase in the expression of somatostatin was observed (Figure 3, panel e).

[0157] No change in the expression of amylase was observed in cells expressing markers characteristic of the pancreatic endocrine lineage that were treated with PubChem Compound ID# 5330797 for five days. (See Figure 3, panel f). However, decreases in the level of expression of PAX4 (Figure 3, panel h), NKX6.1 (Figure 3, panel k), PDX1 (Figure 3, panel l), NEUROD (Figure 3, panel o), and BRN4 (Figure 3, panel q) was observed.

Example 4

Cyclin Dependant Kinase Inhibitor Treatment Increased the Expression of MAFA in Islet-Like Clusters.

[0158] Cells of the human embryonic stem cell line H1 at passage 52 were cultured on MATRIGEL[®] coated dishes (1:30 dilution) and differentiated according to the methods described in Example 1. An additional stage (Stage 6) was added, in order to further mature the cells expressing markers characteristic of the pancreatic endocrine lineage. Stage 6 in this example consisted of a seven day treatment in DMEM/F12 + 1% B27 (Invitrogen, CA). The medium was changed daily.

- [0159] After stage 6, the cells were treated for 5 mins at room temperature with 1X accutase (Sigma, MO). The accutase was removed, and DMEM/12 + 1% B27 was added to the cells. The attached cells were removed using a cell scarper and gently resuspended and passed through a 40 µm cell strainer. The cells retained on the strainer were removed by rinsing in basal media and cultured in suspension on Ultra-Low culture plates (Catalogue# 3471, Corning, Ma). The cells were then treated as follows: The cells were cultured in DMEM/F12 + 1 % B27, containing 20 ng/ml of activin A (AA), 1 µm of CDK inhibitor III (Catalog# 217714, Calbiochem, Ca) for 10 days (Stage 7). Cells treated with vehicle were included as controls. Samples were collected at days 7 through 10 for PCR analysis and dithizone staining. The cells cultured in suspension according to the methods outlined in this example assumed a morphology similar to pancreatic islet clusters. Treatment with CDK inhibitor III did not appear to affect the morphology of the islet like clusters.
- [0160] Figure 4, panels a-i shows the effect of CDK inhibitor III treatment on gene expression profile of the cell clusters. Treatment with of CDK inhibitor III increased the expression of markers associated with the pancreatic endocrine lineage and in particular increased the expression of the pro-insulin transcription factor, MAFA.
- [0161] Figure 5, panels a-b shows the effect of CDK inhibitor III on dithazone (DTZ) staining of clusters. Cell clusters treated with CDK inhibitor and stained with DTZ, showed a more reddish staining pattern as compared to clusters not treated with the CDK inhibitor III.

Example 5

FACS Analysis of Insulin Producing Cells Produced by the Methods of the Present Invention.

- [0162] Cells of the human embryonic stem cell line H1 at passage 42 were cultured on MATRIGEL[®]-coated plates, and differentiated into insulin producing cells using the following protocol:
- a. RPMI medium supplemented with 2% BSA (Catalog# 152401, MP Biomedical, Ohio), and 100 ng/ml activin A (R&D Systems, MN) plus 20 ng/ml WNT-3a (Catalog# 1324-WN-002, R&D Systems, MN) plus 8 ng/ml of

bFGF (Catalog# 100-18B, PeproTech, NJ), for one day followed by treatment with RPMI media supplemented with 2% BSA and 100 ng/ml activin A plus 8 ng/ml of bFGF for an additional two days (Stage 1), then

- b. DMEM/F12 + 2% BSA + 50 ng/ml FGF7 + 0.25 μ M Cyclopamine- KAAD (#239804, Calbiochem, CA) for two days (Stage 2), then
- c. DMEM/F12 + 1% B27 (Invitrogen, CA) + 50 ng/ml FGF7 + 0.25 μ M Cyclopamine- KAAD + 2 μ M Retinoic acid (RA) (Sigma, MO) + 100 ng/ml of Noggin (R & D Systems, MN) for four days (Stage 3), then
- d. DMEM/F12 + 1% B27 (Invitrogen, CA) + 100 ng/ml Noggin + 1 μ M DAPT (a gamma-secretase inhibitor) (Catalog# 565784, Calbiochem, CA) + 1 μ M ALK5 inhibitor II (Catalog# 616452, Calbiochem, Ca) + 100 ng/ml of Netrin-4 (R&D Systems, MN) for three days (Stage 4), then
- e. DMEM/F12 + 1% B27 (Invitrogen, CA) + 1 μ M ALK5 inhibitor II (Calbiochem, Ca) for seven days (Stage 5), then
- f. DMEM/F12 + 1% B27 for seven days (Stage 6), then
- g. Treatment with Accutase for 5 minutes, followed by scraping to remove any remaining attached cells. The cell suspension was then passed through a 40 μ m cell strainer. The cells retained on the strainer were removed by rinsing in basal media and cultured in suspension on Ultra-Low culture plates in DMEM-High Glucose (Catalogue# 11995-073, Invitrogen, Ca) + 1 % B27 + 20 ng/ml of activin A (AA) 1 μ m of CDK inhibitor III (Catalog# 217714, Calbiochem, Ca) for 5 days (Stage 7).

[0163] Islet-like clusters were dispersed into single cells using TrypLE Express (Invitrogen, Carlsbad, CA) and washed in cold PBS. For fixation, the cells were resuspended in 200-300 μ l Cytofix/Cytoperm Buffer (BD 554722, BD, Ca) and incubated for 30 min at 4°C. Cells were washed two times in 1 ml Perm/Wash Buffer Solution (BD 554723) and resuspended in 100 μ l staining/blocking solution containing 2% normal goat serum in Perm/Wash buffer. For flow cytometric analysis, cells were stained with the following primary antibodies: Anti-Insulin (Rabbit mAb, Cell Signaling No.

C27C9; 1:100 dilution); Anti-Glucagon (Mouse Mab, Sigma No. G2654, 1:100); Anti-Synaptophysin (Rabbit Polyclonal antibody, DakoCytomation No A0010, 1:50). Cells were incubated for 30 min at 4°C followed by two washes in Perm/Wash buffer and a further 30 min incubation in appropriate secondary antibodies as follows: Goat anti-Rabbit Alexa 647 (Invitrogen No. A21246) or Goat anti-Mouse 647 (Invitrogen No. A21235); Goat anti-Rabbit R-PE (BioSource No. ALI4407). All secondary antibodies were used at a 1:200 dilution. Cells were washed at least once in Perm/Wash buffer and analyzed using BD FACSArray. At least 10,000 events were acquired for analysis. Controls included undifferentiated H1 cells and the β -TC (CRL-11506™ ATCC, VA) cell line.

- [0164] Figure 6, panels a-c show the percentage insulin positive, synaptophysin positive, and glucagon positive cells in cells following treatment with Stage 7, in medium containing vehicle. Figure 7, panels a-c shows the percentage insulin positive, synaptophysin positive, and glucagon positive cells following treatment with Stage 7 in medium containing 1 μ M CDK inhibitor III for 5 days. The number of single hormonal insulin positive cells increased from 3% to 8% following treatment with the CDK inhibitor. Additionally, the percentage of poly hormonal (insulin and glucagon positive) cells decreased following treatment with the CDK inhibitor.

Example 6

Kinetics of CDK Inhibitor-Induced MAFA Expression.

- [0165] Cells of the human embryonic stem cell line H1 at passage 42 were cultured on MATRIGEL®-coated plates, and differentiated into insulin producing cells using the following protocol:
- a. RPMI medium supplemented with 2% BSA (Catalog# 152401, MP Biomedical, Ohio), and 100 ng/ml activin A (R&D Systems, MN) plus 20 ng/ml WNT-3a (Catalog# 1324-WN-002, R&D Systems, MN) plus 8 ng/ml of bFGF (Catalog# 100-18B, PeproTech, NJ), for one day followed by treatment with RPMI media supplemented with 2% BSA and 100 ng/ml activin A plus 8 ng/ml of bFGF for an additional two days (Stage 1), then

- b. DMEM/F12 + 2% BSA + 50 ng/ml FGF7 + 0.25 μ M Cyclopamine- KAAD (#239804, Calbiochem, CA) for two days (Stage 2), then
- c. DMEM/F12 + 1% B27 (Invitrogen, CA) + 50 ng/ml FGF7 + 0.25 μ M Cyclopamine- KAAD + 2 μ M Retinoic acid (RA) (Sigma, MO) + 100 ng/ml of Noggin (R & D Systems, MN) for four days (Stage 3), then
- d. DMEM/F12 + 1% B27 (Invitrogen, CA) + 100 ng/ml Noggin + 1 μ M DAPT (a gamma-secretase inhibitor) (Catalog# 565784, Calbiochem, CA) + 1 μ M ALK5 inhibitor II (Catalog# 616452, Calbiochem, Ca) + 100 ng/ml of Netrin-4 (R&D Systems, MN) for three days (Stage 4), then
- e. DMEM/F12 + 1% B27 (Invitrogen, CA) + 1 μ M ALK5 inhibitor II (Calbiochem, Ca) for seven days (Stage 5), then
- f. DMEM/F12 + 1% B27 for seven days (Stage 6), then
- g. Treatment with Accutase for 5 minutes, followed by scraping to remove any remaining attached cells. The cell suspension was then passed through a 40 μ m cell strainer. The cells retained on the strainer were removed by rinsing in basal media and cultured in suspension on Ultra-Low culture plates in DMEM-High Glucose (Catalogue# 11995-073, Invitrogen, Ca) + 1 % B27 + 20 ng/ml of activin A (AA) 2 μ m of CDK inhibitor III (Catalog# 217714, Calbiochem, Ca) for 1-8 days (Stage 7).

[0166] Samples were collected for PCR analysis at days 1, 2, 3, and 4. Following 4 days of treatment with CDK inhibitor, the CDK inhibitor was removed from culture and the cells were cultured additional 4 days in DMEM-F12 + 1% B27 + 20 ng/ml of activin A. At the end of the four days, samples were collected in triplicate for PCR analysis.

[0167] Figure 8, panels a-b show expression pattern of MAFA and insulin at various time points of stage 7. CDK inhibitor treatment resulted in significant increase in MAFA and insulin expression which increased as a function of time. However, removal of CDK inhibitor resulted in a significant drop to both MAFA and insulin expression, in samples obtained four days after removal of the compound.

Example 7

Screening of the Effects of Compounds from the BIOMOL™ Kinase Inhibitor Library on Cells that have been Treated According to the Differentiation Protocol Outlined in Example 1.

- [0168] Cells of the human embryonic stem cell line H1 at passage 51 were seeded onto MATRIGEL[®]-coated 24-well dishes (1:30 dilution), and differentiated according to the methods described in Example 1 up to stage 5. Following this, the cells were grown for one day in DMEM/F12 + 1% B27 and then treated for six days in DMEM/F12 + 1% B27 containing a compound from a BIOMOL™ compound library (Catalog# 2832, BIOMOL, Plymouth Meeting, Pa) at a final concentration of 4μM. Wells containing vehicle were included as a control. Throughout the treatment protocol media containing vehicle or compound was changed every other day. All samples were treated in duplicate. At the completion of this treatment RNA was collected for PCR analysis. Samples were analyzed by real-time PCR for expression of insulin, glucagon, MAFA, and ARX4. Results are expressed as a ratio insulin/glucagon (Table 2), or MAFA versus Arx4 (Table 2) of the treated samples relative to the untreated control, as measured by real-time PCR. The corresponding catalog#, CAS#, and compound name or ID number for each alpha numeric well# is listed in Table 3.
- [0169] Treatment of cells expressing markers characteristic of the pancreatic endocrine lineage with compounds C8 or F1 at a 4μM concentration resulted in an insulin/ glucagon expression ratio of approximately 10.0 or higher. Cells treated with D9 had an insulin/ glucagon expression ratio of approximately 1840.0 (Table 2).
- [0170] We next examined the effect of these compounds on the ratio of MAFA/ARX4, and we observed that treatment of cells expressing markers characteristic of the pancreatic endocrine lineage with several of the compounds resulted in a much greater change in the ratio of MAFA to ARX4 than other compounds tested in the library: Cells treated with compound B6 or F1 showed a ratio of MAFA/ARX4 of approximately greater than 10. Treatment of cells expressing markers characteristic of the pancreatic endocrine lineage with compound C8 resulted in a MAFA/ ARX4 ratio of

approximately 84, while cells treated with D9 had a MAFA/ ARX4 ratio of approximately 212. (Table 2).

Example 8

The Effect of Cyclin Dependant Kinase Inhibitors on Insulin and MAFA Expression in Cells Treated according to the Differentiation Protocol Outlined in Example 1.

- [0171] Cells of the human embryonic stem cell line H1 at passage 51 were seeded onto MATRIGEL™ coated 24-well dishes (1:30 dilution), and differentiated according to the methods described in Example 1 up to stage 5. Following this, the cells were grown for eight days in DMEM/F12 + 1% B27 and then treated for four days in DMEM/F12 + 1% B27 containing a cyclin dependent kinase inhibitor at a final concentration of 0.6125, 1.25, or 5.0 µM. We tested 6 inhibitors: PubChem ID# 5330812 (EMD cat# 217714), PubChem ID# 4566 (EMD cat#217713), PubChem ID# 5330797 (EMD cat# 219476), PubChem ID# 73292 (EMD cat#341251), PubChem ID# 4592 (EMD cat#495620), and PubChem ID# 160355 (EMD cat #557360). Wells containing vehicle were included as a control. Throughout the treatment protocol media containing vehicle or compound was changed every other day. All samples were treated in duplicate. At the completion of this treatment RNA was collected for PCR analysis. Samples were analyzed by real-time PCR for expression of insulin, glucagon, MAFA, and ARX4. Results are expressed as the fold change relative to the vehicle treated control, as measured by real-time PCR.
- [0172] We observed that the compounds PubChem ID# 5330812, PubChem ID# 4566, PubChem ID# 5330797, and PubChem ID# 73292 all stimulated MAFA expression at the concentrations tested (Table 4). PubChem ID# 4592 and PubChem ID# 160355 did not stimulate MAFA at the concentrations tested (Table 4). The compounds PubChem ID# 5330812, PubChem ID# 4566, PubChem ID# 5330797, PubChem ID# 4592 and PubChem ID# 160355 all appeared to stimulate insulin expression (Table 4). The compound PubChem ID# 5330797 reduced both glucagon and Arx4 expression (Table 4) while stimulating MAFA expression.

Example 9**Differentiation of Human Embryonic Stem Cells of the Cell Line H1 to Pancreatic Endocrine Cells with DMEM containing 25mM glucose (DMEM-HG), Lacking Fetal Bovine Serum**

- [0173]** Cells of the human embryonic stem cells line H1 were cultured on MATRIGEL[®]-coated dishes (1:30 dilution) and differentiated into cells expressing markers characteristic of the pancreatic endocrine lineage using the following protocol:
- a. RPMI medium supplemented with 2% BSA (Catalog# 152401, MP Biomedical, Ohio), and 100 ng/ml activin A (R&D Systems, MN) plus 20 ng/ml WNT-3a (Catalog# 1324-WN-002, R&D Systems, MN) plus 8 ng/ml of bFGF (Catalog# 100-18B, PeproTech, NJ), for one day followed by treatment with RPMI media supplemented with 2% BSA and 100 ng/ml activin A plus 8 ng/ml of bFGF for an additional two days (Stage 1), then
 - b. RPMI medium supplemented with 2% BSA + 50 ng/ml FGF7 + 0.25 μ M Cyclopamine- KAAD (#239804, Calbiochem, CA) for two days (Stage 2), then
 - c. DMEM-HG + 1% B27 (Invitrogen, CA) + 50 ng/ml FGF7 + 0.25 μ M Cyclopamine- KAAD + 2 μ M Retinoic acid (RA) (Sigma, MO) + 100 ng/ml of Noggin (R & D Systems, MN) for six days (Stage 3), then
 - d. DMEM-HG + 1% B27 (Invitrogen, CA) + 100 ng/ml Noggin + 1 μ M ALK5 inhibitor II (Catalog# 616452, Calbiochem, Ca) for three days (Stage 4), then
 - e. DMEM-HG + 1% B27 (Invitrogen, CA) + 1 μ M ALK5 inhibitor II (Calbiochem, Ca) for seven days (Stage 5).
- [0174]** Medium was changed daily. At each stage the cell number was calculated using a hemocytometer and RNA was collected for PCR analysis. All samples were collected in triplicate.

Example 10**Screening of the Effects of Compounds from the EMD Kinase Inhibitor Library I on Cells Treated according to the Differentiation Protocol Outlined in Example****9**

- [0175]** Cells of the human embryonic stem cell line H1 at passage 45 were seeded onto MATRIGEL[®]-coated 24-well dishes (1:30 dilution), and differentiated according to the methods described in Example 9 up to stage 5. Following this, the cells were fed and treated on day 1, 3, and 5 of stage 5 with media comprising DMEM-HG, 1% B27 (Invitrogen, CA), 1 μ M ALK5 inhibitor II (Calbiochem, Ca) and a compound from an EMD Calbiochem compound library 1 solubilized in DMSO (Catalog# 539744, Calbiochem, San Diego, Ca) and treated at a final concentration of 2.5 μ M. Wells containing vehicle were included as a control. Throughout the protocol media was changed daily except at stage 5 when media was changed every other day. All samples were treated in duplicate.
- [0176]** At the completion of this treatment RNA was collected for PCR analysis. Samples were analyzed by real-time PCR for expression of MAFA. Results are expressed as the fold increase in MAFA expression versus untreated H1 human embryonic stem cells (Table 5), as measured by real-time PCR.
- [0177]** Treatment of cells expressing markers characteristic of the pancreatic endocrine lineage with compounds A4 (Cat#, 124001, Akt Inhibitor IV), E8 (Cat# 527450, PKR Inhibitor), and F9 (Cat#539648, Staurosporine, N-benzoyl-) at a 2.5 μ M concentration resulted in an increase in MAFA expression at least 4 fold higher than vehicle treated controls (Table 5). Treatment with the compound E6 (Cat# 521233, PDGF Receptor Tyrosine Kinase Inhibitor IV) at a 2.5 μ M concentration resulted in an increase in MAFA expression at least 2.5 fold higher than vehicle treated controls (Table 5).

Example 11**Screening of the Effects of Compounds from the EMD Kinase Inhibitor Library
II on Cells Treated according to the Differentiation Protocol Outlined in
Example 9**

- [0178]** Cells of the human embryonic stem cell line H1 at passage 46 were seeded onto MATRIGEL[®]-coated 24-well dishes (1:30 dilution), and differentiated according to the methods described in Example 9 up to stage 5. Following this, the cells were fed and treated on day 1, 3, and 5 of stage 5 with media comprising DMEM-HG, 1% B27 (Invitrogen, CA), 1 μ M ALK5 inhibitor II (Calbiochem, Ca) (Stage 5) and a compound from an EMD Calbiochem compound library II solubilized in DMSO (Tables 1 and 6, Calbiochem, San Diego, Ca) and treated at a final concentration of 2.5 μ M. Wells containing vehicle were included as a control. Throughout the protocol media was changed daily except at stage 5 when media was changed every other day. All samples were treated in duplicate.
- [0179]** At the completion of this treatment RNA was collected for PCR analysis. Samples were analyzed by real-time PCR for expression of MAFA. Results for compounds that stimulated the expression of MAFA are shown and expressed as the fold increase in MAFA expression versus control samples (Figure 9), as measured by real-time PCR.
- [0180]** Treatment of cells expressing markers characteristic of the pancreatic endocrine lineage with either: Alsterpaullone, 2-Cyanoethyl; SU9516; Alsterpaullone; Cdk1/2 Inhibitor III; Casein Kinase I Inhibitor, D4476; or MEK1/2 Inhibitor at a 2.5 μ M concentration resulted in a 4.5 fold increase in MAFA expression versus untreated controls (Table 7).

Example 12**Inhibiting Cell Cycle Progression in Cells Expressing Markers Characteristic of the Pancreatic Endocrine Lineage with Small Molecule Inhibitors Promotes MAFA Expression in Cells Expressing Markers Characteristic of the Pancreatic Endocrine Lineage**

- [0181] Cell growth resulting from cell cycle progression can be activated and maintained by stimulating cells with extracellular growth factors. Growth factors bind to the extracellular domains of growth factor receptors, inducing a conformational switch in the receptor's intracellular domain. This shift initiates receptor dimerization and activation of tyrosine kinases located on the intracellular domain of the receptor leading to phosphorylation and activation of multiple serine/threonine kinases downstream, ultimately resulting in cell cycle progression and cell proliferation.
- [0182] Under normal physiologic conditions mature pancreatic beta cells, characterized by expression of insulin and the transcription factor MAFA, are quiescent and tend to remain in G0 of the cell cycle. Yet, in order to generate enough cells to form a functional organ and meet the needs of a mature animal, the cells expressing markers characteristic of the pancreatic endocrine lineage of the present invention must be cell cycling. Consequently, at some point in embryonic development, the cells expressing markers characteristic of the pancreatic endocrine lineage of the present invention differentiate to beta cells and transition from an actively cell-cycling proliferating cell, to a quiescent cell.
- [0183] Our data indicate that by inhibiting cell cycle progression by blocking signaling cascades with small molecule kinase inhibitors, we can induce the cells expressing markers characteristic of the pancreatic endocrine lineage to express MAFA, a marker of mature pancreatic beta cells. Kinase inhibitors targeted to a growth factor receptor, (PDGF Receptor Tyrosine Kinase Inhibitor IV), or inhibitors which disrupt kinases downstream of tyrosine kinase receptors (MEK1/2 Inhibitor, PKR Inhibitor, or Akt Inhibitor IV) disrupt proliferative growth factor/kinase based signaling resulting in cell cycle arrest and induction of MAFA expression. Use of a broad spectrum inhibitor like staurosporine, can effectively induce MAFA, however it is also cytotoxic at effective concentrations. More directed compounds like cyclin dependent

kinase inhibitors (Alsterpaullone, 2-Cyanoethyl; SU9516; Alsterpaullone; or Cdk1/2 Inhibitor III) induce MAFA with less toxicity than a broad spectrum inhibitor like staurosporine.

- [0184] In order to determine if a broad spectrum kinase inhibitor could induce MAFA expression and a more mature phenotype in the cells expressing markers characteristic of the pancreatic endocrine lineage of the present invention we differentiated H1 human ES cells according to the methods described in example 9, and treated them on days 1, 3, and 5 of stage 5 with the protein-tyrosine kinase inhibitor, Genistein, which has been shown to induce G2 phase arrest in human and murine cell lines and inhibit multiple kinases. At doses of 10 and 30ng/ml the endocrine hormones insulin, somatostatin, and the transcription factor MAFA, all showed increased expression versus untreated controls, while at 10ng/ml the endocrine hormone glucagon had increased expression (Figure 10). We observed significant toxicity at a 100ng/ml dose of genistein that correlated with loss of insulin, glucagon, and somatostatin expression.
- [0185] These data indicate that by inhibiting cell cycle progression by blocking signaling cascades with small molecule kinase inhibitors targeted to inhibit signal transduction from a growth factor receptor tyrosine kinase through intracellular signaling kinases to the nucleus and cyclin dependent kinases, we can induce the cells expressing markers characteristic of the pancreatic endocrine lineage of the present invention to express MAFA, a marker of mature pancreatic beta cells.
- [0186] Publications cited throughout this document are hereby incorporated by reference in their entirety. Although the various aspects of the invention have been illustrated above by reference to examples and preferred embodiments, it will be appreciated that the scope of the invention is defined not by the foregoing description but by the following claims properly construed under principles of patent law.

Well	PubChemID#	Well	PubChemID#	Well	PubChemID#	Well	PubChemID#
A2	<u>16760529</u>	B2	<u>16760303</u>	C2	<u>5330797</u>	D2	<u>9797929</u>
A3	<u>5278396</u>	B3	<u>5326739</u>	C3	<u>3004085</u>	D3	<u>11493598</u>
A4	<u>6605258</u>	B4	<u>5353431</u>	C4	<u>481747</u>	D4	<u>16760417</u>
A5	<u>5005498</u>	B5	<u>2422</u>	C5	<u>1893668</u>	D5	<u>73292</u>
A6	<u>16760286</u>	B6	<u>5472558</u>	C6	<u>9969021</u>	D6	<u>4124851</u>
A7	<u>5326843</u>	B7	<u>2794188</u>	C7	<u>2856</u>	D7	<u>6539732</u>
A8	<u>3641059</u>	B8	<u>5330812</u>	C8	<u>6918386</u>	D8	<u>448014</u>
A9	<u>6604931</u>	B9	<u>438981</u>	C9	<u>10202471</u>	D9	<u>5287844</u>
A10	<u>11524144</u>	B10	<u>6419753</u>	C10	<u>9549301</u>	D10	<u>6538818</u>
A11	<u>9549303</u>	B11	<u>16760346</u>	C11	<u>5339183</u>	D11	<u>10020713</u>
E2	<u>5312137</u>	F2	<u>11382492</u>	G2	<u>6419739</u>	H2	<u>176155</u>
E3	<u>6419766</u>	F3	<u>11624601</u>	G3	<u>4665</u>	H3	<u>16219471</u>
E4	<u>6419741</u>	F4	<u>490561</u>	G4	<u>4713</u>	H4	<u>3387354</u>
E5	<u>3674</u>	F5	<u>3820</u>	G5	<u>4712</u>	H5	<u>5174</u>
E6	<u>9903786</u>	F6	<u>5312122</u>	G6	<u>5164</u>	H6	<u>5228</u>
E7	<u>6419764</u>	F7	<u>9951490</u>	G7	<u>4987</u>	H7	<u>16760659</u>
E8	<u>8515</u>	F8	<u>389898</u>	G8	<u>9549289</u>	H8	<u>451705</u>
E9	<u>11665831</u>	F9	<u>9549284</u>	G9	<u>5702541</u>	H9	<u>16760660</u>
E10	<u>11422035</u>	F10	<u>11644425</u>	G10	<u>5162</u>	H10	<u>5289419</u>
E11	<u>16760525</u>	F11	<u>509554</u>	G11	<u>5353940</u>	H11	<u>9549300</u>

**Table 1. AlphaNumeric Well
Label and the Corresponding
PubChem ID# for EMD
Calbiochem® Kinase Inhibitor
II compound library**

Table 2. The Effect of compounds of the BIOMOL Inhibitor compound library on the Ratio of Insulin/glucagon and MAFA/Arx4 expression as determined by real-time PCR in Cells Expressing Markers Characteristic of the Pancreatic Endocrine Lineage. The AlphaNumeric well# corresponds to the compound identity in Table 3.

Well #	Ratio vs. Control	
	Insulin to Glucagon	MAFA to Arx4
B1	1.6	0.9
B2	2.4	1.3
B3	2.9	2.9
B4	1.1	2.0
B5	1.3	1.3
B6	1.6	16.3
B7	1.3	0.5
B8	1.6	0.5
B9	1.1	1.5
B10	1.2	1.5
B11	1.1	2.1
B12	1.0	2.0
C1	0.7	0.8
C2	0.9	1.0
C3	1.3	0.9
C4	1.2	1.7
C5	1.0	1.1
C6	1.6	1.2
C7	4.3	0.2
C8	40.2	84.3
C9	0.8	0.5
C10	2.3	1.9
C11	1.1	0.4
C12	1.0	0.4
D1	2.7	1.2

D2	3.4	1.3
D3	1.7	2.1
D4	5.8	6.4
D5	1.4	1.1
D6	1.8	3.9
D7	1.7	0.6
D8	2.8	5.1
D9	1842.5	212.6
D10	0.9	1.3
D11	1.0	0.7
D12	1.1	2.5
E1	1.1	0.9
E2	0.8	0.8
E3	1.2	1.0
E4	2.1	1.3
E5	1.3	1.1
E6	2.0	1.5
E7	4.8	0.2
E8	3.7	0.0
E9	1.0	0.8
E10	0.6	0.2
E11	1.0	0.3
E12	0.8	0.2
F1	10.3	9.5
F2	2.9	1.9
F3	2.6	2.5
F4	1.5	2.7
F5	1.9	1.4
F6	1.6	1.2
F7	1.9	0.6
F8	1.5	0.6
F9	1.0	1.4
F10	5.4	3.4

F11	0.8	1.9
F12	1.0	1.4
G1	0.8	1.2
G2	0.6	1.1
G3	2.0	1.6
G4	1.3	1.6
G5	1.7	1.5
G6	1.5	1.3
G7	4.6	0.2
G8	3.9	0.4
G9	1.0	0.7
G10	1.3	0.7
G11	1.9	0.6
G12	1.4	0.8
H1	3.1	0.6
H2	1.8	3.5
H3	1.8	3.9
H4	1.2	4.0
H5	1.8	2.0
H6	1.4	2.9
H7	1.5	0.6
H8	2.1	0.8
vehicle control	1.0	1.0

Table 3. AlphaNumeric Well Label and the Corresponding Catalog#, CAS#, and Compound Name or ID Number for the BIOMOL Kinase Inhibitor compound library

PLATE LOCATION	CATALOG #	CAS #	COMPOUND NAME OR ID NUMBER
B1	EI-360	167869-21-8	PD-98059
B2	EI-282	109511-58-2	U-0126
B3	EI-286	152121-47-6	SB-203580
B4	EI-148	84477-87-2	H-7
B5	EI-195	84468-17-7	H-9
B6	EI-156	62996-74-1	Staurosporine
B7	EI-228	133550-35-5	AG-494
B8	EI-267		AG-825
B9	EI-185	125697-92-9	Lavendustin A
B10	EI-253	136831-49-7	RG-14620
B11	EI-191	118409-57-7	Tyrphostin 23
B12	EI-187	118409-58-8	Tyrphostin 25
C1	EI-257	122520-85-8	Tyrphostin 46
C2	EI-188	122520-86-9	Tyrphostin 47
C3	EI-189	122520-90-5	Tyrphostin 51
C4	EI-190	2826-26-8	Tyrphostin 1
C5	EI-335	116313-73-6	Tyrphostin AG 1288
C6	EI-277	63177-57-1	Tyrphostin AG 1478
C7	AC-1133	71897-07-9	Tyrphostin AG 1295
C8	EI-215	10537-47-0	Tyrphostin 9
C9	EI-247		HNMPA (Hydroxy-2-naphthalenylmethylphosphonic acid)
C10	EI-370	120685-11-2	PKC-412
C11	EI-271	10083-24-6	Piceatannol
C12	EI-275	172889-26-8	PP1
D1	EI-272	133550-35-3	AG-490
D2	EI-263		AG-126

D3	EI-229		AG-370
D4	EI-258		AG-879
D5	ST-420	154447-36-6	LY 294002
D6	ST-415	19545-26-7	Wortmannin
D7	EI-246	133052-90-1	GF 109203X
D8	EI-226	548-04-9	Hypericin
D9	EI-283	138489-18-6	Ro 31-8220
D10	EI-155	123-78-4	Sphingosine
D11	EI-196	127243-85-0	H-89
D12	EI-158	84478-11-5	H-8
E1	EI-184	91742-10-8	HA-1004
E2	EI-233	103745-39-7	HA-1077
E3	EI-232		HDBA (2-Hydroxy-5-(2,5-dihydroxybenzylamino)benzoic acid)
E4	EI-230	127191-97-3	KN-62
E5	EI-268		KN-93
E6	EI-197	109376-83-2	ML-7
E7	EI-153	105637-50-1	ML-9
E8	CC-100	452-06-2	2-Aminopurine
E9	CC-202	158982-15-1	N9-Isopropyl-olomoucine
E10	CC-200	101622-51-9	Olomoucine
E11	CC-201	101622-50-8	iso-Olomoucine
E12	CC-205	186692-46-6	Roscovitine
F1	EI-293	24386-93-4	5-Iodotubercidin
F2	EI-295	62004-35-7	LFM-A13
F3	EI-294	152121-30-7	SB-202190
F4	EI-297	172889-27-9	PP2
F5	EI-298	208260-29-1	ZM 336372
F6	EI-306	5812-07-7	SU 4312
F7	EI-303	146535-11-7	AG-1296
F8	EI-307	220904-83-6	GW 5074
F9	AC-1121	6865-14-1	Palmitoyl-DL-carnitine Cl
F10	EI-270	82-08-6	Rottlerin

F11	EI-147	446-72-0	Genistein
F12	ST-110	486-66-8	Daidzein
G1	EI-146	63177-57-1	Erbstatin analog
G2	AC-1142	6151-25-3	Quercetin dihydrate
G3	AC-1293		SU1498
G4	EI-357	4452-06-6	ZM 449829
G5	EI-278	195462-67-7	BAY 11-7082
G6	EI-231	53-85-0	DRB (5,6-Dichloro-1- β -D- ribofuranosylbenzimidazole)
G7	EI-273		HBDDE (2,2',3,3',4,4'-Hexahydroxy- 1,1'-biphenyl-6,6'-dimethanol dimethyl ether)
G8	EI-305	129-56-6	SP 600125
G9	CC-206	479-41-4	Indirubin
G10	CC-207	160807-49-8	Indirubin-3'-monoxime
G11	EI-299	146986-50-7	Y-27632
G12	EI-310	142273-20-9	Kenpaullone
H1	EI-328	121-40-4	Terreic acid
H2	EI-332	35943-35-2	Triciribine
H3	EI-336		BML-257
H4	EI-343		SC-514
H5	EI-344		BML-259
H6	EI-345	520-36-5	Apigenin
H7	EI-346		BML-265 (Erlotinib analog)
H8	A-275	53123-88-9	Rapamycin

Table 4. The Effect of compounds of the BIOMOL Inhibitor compound library on the Expression of Insulin, glucagon, MAFA and Arx4 in Cells Expressing Markers Characteristic of the Pancreatic Endocrine Lineage

Concentration and PubChem ID#	MAFA	Insulin	Glucagon	Arx4
0.61μM 5330812	46.3	0.9	0.26	0.68
1.25μM 5330812	209.2	1.3	0.31	0.66
5.0μM 5330812	2909.9	66.3	4.71	0.92
0.61μM 4566	1.0	1.0	0.77	0.78
1.25μM 4566	0.8	1.1	0.90	0.78
5.0μM 4566	1.0	1.1	0.96	0.69
0.61μM 5330797	0.7	0.6	0.34	0.36
1.25μM 5330797	1.5	0.8	0.25	0.37
5.0μM 5330797	6.3	1.3	0.04	0.16
0.61μM 73292	0.7	0.7	0.29	0.38
1.25μM 73292	1.3	1.0	0.25	0.42
5.0μM 73292	3.1	0.8	0.13	0.33
0.61μM 4592	0.9	0.9	0.81	0.61
1.25μM 4592	1.0	1.0	0.70	0.54
5.0μM 4592	0.6	1.3	1.08	0.77
0.61μM 160355	0.9	0.9	0.76	0.77
1.25μM 160355	0.7	1.0	0.61	0.65
5.0μM 160355	0.8	1.1	0.59	0.86
Vehicle Treated	1.0	1.0	1.00	1.00

Table 5. AlphaNumeric Well Label and the Corresponding Catalog#, and Compound Name or ID Number for the EMD Calbiochem Kinase Inhibitor compound library I

Plate Location	Catalog#	Compound Name	Gene Induction vs. H1: MAFa
A10	197221	Bcr-abl Inhibitor	1.5
A11	203290	Bisindolylmaleimide I	0.8
A12	DMSO	Control	1.5
A2	121767	AG 1024	0.8
A3	121790	AGL 2043	0.8
A4	124011	Akt Inhibitor IV	45.7
A5	124012	Akt Inhibitor V, Triciribine	0.9
A6	124018	Akt Inhibitor VIII, Isozyme-Selective, Akti-1/2	1.6
A7	124020	Akt Inhibitor X	1.4
A8	521275	PDK1/Akt/Flt Dual Pathway Inhibitor	2.1
A9	189404	Aurora Kinase Inhibitor II	1.3
B10	317200	DMBI	1.9
B11	324673	EGFR/ErbB-2 Inhibitor	2.4
B12	DMSO	Control	1.9
B2	203297	Bisindolylmaleimide IV	1.9
B3	203696	BPIQ-I	1.6
B4	220285	Chelerythrine Chloride	2.3
B5	234505	Compound 56	1.8
B6	260961	DNA-PK Inhibitor II	2.0
B7	260962	DNA-PK Inhibitor III	2.2
B8	528100	PI-103	1.9
B9	266788	Diacylglycerol Kinase Inhibitor II	1.5
C10	375670	Herbimycin A, Streptomyces sp.	1.3
C11	343022	Flt-3 Inhibitor III	1.1
C12	DMSO	Control	1.1
C2	324674	EGFR Inhibitor	3.5
C3	324840	EGFR/ErbB-2/ErbB-4 Inhibitor	0.9
C4	343020	Flt-3 Inhibitor	0.6
C5	343021	Flt-3 Inhibitor II	0.5
C6	344036	cFMS Receptor Tyrosine Kinase Inhibitor	2.2
C7	365250	Gö 6976	1.9
C8	365251	Gö 6983	1.0
C9	371806	GTP-14564	0.7
D10	440203	LY 303511	1.7
D11	448101	Met Kinase Inhibitor	2.1
D12	BLANK		1.7

D2	407248	IGF-1R Inhibitor II	1.6
D3	407601	IRAK-1/4 Inhibitor	2.2
D4	420099	JAK Inhibitor I	1.4
D5	420104	JAK3 Inhibitor II	2.0
D6	420121	JAK3 Inhibitor IV	1.7
D7	420126	JAK3 Inhibitor VI	1.9
D8	428205	Lck Inhibitor	1.9
D9	440202	LY 294002	2.3
E10	528106	PI 3-Kg Inhibitor	1.6
E11	528108	PI 3-KbInhibitor II	1.4
E12	BLANK		1.6
E2	513035	PD 158780	0.8
E3	513040	PD 174265	1.0
E4	521231	PDGF Receptor Tyrosine Kinase Inhibitor II	0.8
E5	521232	PDGF Receptor Tyrosine Kinase Inhibitor III	1.7
E6	521233	PDGF Receptor Tyrosine Kinase Inhibitor IV	5.5
E7	521234	PDGF RTK Inhibitor	1.9
E8	527450	PKR Inhibitor	24.6
E9	527455	PKR Inhibitor, Negative Control	1.5
F10	567805	Src Kinase Inhibitor I	2.3
F11	572660	SU11652	1.7
F12	DMSO	Control	2.1
F2	529574	PP3	1.3
F3	529581	PP1 Analog II, 1NM-PP1	2.3
F4	539652	PKCbII/EGFR Inhibitor	1.8
F5	539654	PKCb Inhibitor	1.6
F6	553210	Rapamycin	1.2
F7	555553	Rho Kinase Inhibitor III, Rockout	1.7
F8	555554	Rho Kinase Inhibitor IV	2.3
F9	539648	Staurosporine, N-benzoyl-	11.7
G10	658550	AG 1295	1.4
G11	658551	AG 1296	1.1
G12	DMSO	Control	1.2
G2	574711	Syk Inhibitor	1.2
G3	574712	Syk Inhibitor II	0.8
G4	574713	Syk Inhibitor III	1.1
G5	616451	TGF-b RI Kinase Inhibitor	1.1
G6	616453	TGF-b RI Inhibitor III	1.6
G7	658390	AG 9	1.5
G8	658401	AG 490	1.4
G9	658440	AG 112	1.4
H10	189405	Aurora Kinase Inhibitor III	2.0
H11	569397	Staurosporine, Streptomyces sp.	0.0*
H12	DMSO	Control	1.5
H2	658552	AG 1478	2.7
H3	676480	VEGF Receptor 2 Kinase Inhibitor I	1.7
H4	676481	VEGF Receptor Tyrosine Kinase Inhibitor II	1.0
H5	676482	VEGF Receptor Tyrosine Kinase Inhibitor III, KRN633	1.6
H6	676485	VEGF Receptor 2 Kinase Inhibitor II	1.0
H7	676487	VEGF Receptor 2 Kinase Inhibitor III	1.1

H8	676489	VEGF Receptor 2 Kinase Inhibitor IV	2.1
H9	260964	DNA-PK Inhibitor V	1.5

Table 6. AlphaNumeric Well Label, Corresponding Catalog#, and Compound # for EMD Calbiochem® Kinase Inhibitor II compound library

Well #	Catalog #	Compound #
A1	DMSO	
A2	422706	KN-62
A3	118500	ATM Kinase Inhibitor
A4	118501	ATM/ATR Kinase Inhibitor
A5	126870	Alsterpaullone
A6	126871	Alsterpaullone, 2-Cyanoethyl
A7	128125	Aloisine A, RP107
A8	128135	Aloisine, RP106
A9	164640	Aminopurvalanol A
A10	171260	AMPK Inhibitor, Compound C
A11	189405	Aurora Kinase Inhibitor III
A12	BLANK	
B1	DMSO	
B2	189406	Aurora Kinase/Cdk Inhibitor
B3	402085	Indirubin-3'-monoxime
B4	196870	BAY 11-7082
B5	203600	Bohemine
B6	217695	Cdk1 Inhibitor
B7	217696	Cdk1 Inhibitor, CGP74514A
B8	217714	Cdk1/2 Inhibitor III
B9	217720	Cdk1/5 Inhibitor
B10	218696	Casein Kinase I Inhibitor, D4476
B11	218710	Casein Kinase II Inhibitor III, TBCA
B12	BLANK	
C1	DMSO	
C2	219476	Cdk4 Inhibitor
C3	219477	Cdk4 Inhibitor II, NSC 625987
C4	219478	Cdk4 Inhibitor III
C5	219479	Cdc2-Like Kinase Inhibitor, TG003
C6	220486	Chk2 Inhibitor II
C7	234503	Compound 52
C8	238803	Cdk2 Inhibitor III
C9	238804	Cdk2 Inhibitor IV, NU6140
C10	219491	Cdk/Crk Inhibitor
C11	328009	ERK Inhibitor III
C12	BLANK	
D1	DMSO	
D2	688000	ROCK Inhibitor, Y-27632
D3	328007	ERK Inhibitor II, FR180204
D4	328008	ERK Inhibitor II, Negative control
D5	341251	Fascaplysin, Synthetic
D6	361540	GSK-3b Inhibitor I
D7	361541	GSK-3b Inhibitor II
D8	361549	GSK-3b Inhibitor VIII

D9	361550	GSK-3 Inhibitor IX	
D10	361551	GSK-3 Inhibitor X	
D11	361553	GSK-3b Inhibitor XI	
D12	BLANK		
E1	DMSO		
E2	572635	SU6656	
E3	361555	GSK-3 Inhibitor XIII	
E4	371957	Isogranulatimide	
E5	400090	IC261	
E6	401481	IKK-2 Inhibitor IV	
E7	402081	Indirubin Derivative E804	
E8	420119	JNK Inhibitor II	
E9	420123	JNK Inhibitor,	Negative Control
E10	420129	JNK Inhibitor V	
E11	420136	JNK Inhibitor IX	
E12	BLANK		
F1	DMSO		
F2	475863	MK2a Inhibitor	
F3	420135	JNK Inhibitor VIII	
F4	420298	K-252a, Nocardiosis sp.	
F5	422000	Kenpaullone	
F6	422708	KN-93	
F7	444937	MEK Inhibitor I	
F8	444938	MEK Inhibitor II	
F9	444939	MEK1/2 Inhibitor	
F10	454861	MNK1 Inhibitor	
F11	481406	NF- κ B Activation Inhibitor	
F12	BLANK		
G1	DMSO		
G2	506121	p38 MAP Kinase Inhibitor III	
G3	506126	p38 MAP Kinase Inhibitor	
G4	513000	PD 98059	
G5	513030	PD 169316	
G6	559396	SB220025	
G7	540500	Purvalanol A	
G8	361554	GSK3b Inhibitor XII, TWS119	
G9	371963	H-89, Dihydrochloride	
G10	559387	SB 202474, Negative control for p38 MAPK inhibition studies	
G11	559388	SB 202190	
G12	BLANK		
H1	DMSO		
H2	559389	SB 203580	
H3	371970	HA 1077, Dihydrochloride	Fasudil
H4	559402	SB 218078	
H5	565625	SC-68376	
H6	567305	SKF-86002	
H7	567731	Sphingosine Kinase Inhibitor	
H8	569397	Staurosporine, Streptomyces sp.	
H9	570250	STO-609	
H10	572650	SU9516	
H11	616373	Tpl2 Kinase Inhibitor	
H12	BLANK		

Table 7. Fold Induction of MAFA expression by several Compounds from the EMD Kinase Inhibitor Library II on Cells Treated according to the Differentiation Protocol Outlined in Example 9

<u>well#</u>	<u>fold vs. control</u>	<u>cat#</u>	<u>drug name</u>
A6	24.8	126871	Alsterpaullone, 2-Cyanoethyl
H10	18.0	572650	SU9516
A5	15.3	126870	Alsterpaullone
B8	8.2	217714	Cdk1/2 Inhibitor III
B10	5.7	218696	Casein Kinase I Inhibitor, D4476
F9	4.92	444939	MEK1/2 Inhibitor

What is claimed is:

1. A method for increasing the expression of MAFA in cells expressing markers characteristic of the pancreatic endocrine lineage comprising the steps of culturing the cells expressing markers characteristic of the pancreatic endocrine lineage in medium comprising a sufficient amount of a cyclin-dependant kinase inhibitor to cause an increase in expression of MAFA.