

Method of peptide synthesis

The present invention relates to a method of solid-phase peptide synthesis for a peptide that is unusually difficult to synthesize, and to respective peptide-solid phase conjugates.

- 5 The challenge of solid-phase peptide synthesis is that despite well-established routine methods exist, successful and efficient synthesis is highly dependent on the very peptide sequence to be synthesized.

Some peptide sequences, drawn from natural peptide or protein segments, simply do much
10 worse than others on the same resins materials used highly successfully with a lot of 'normal' peptides. In the literature, oftenly careful fine-tuning of solvent systems has been proposed in view of preventing inter-chain aggregation on the resin by virtue of beta-sheet formation. Still then, solvent engineering using dichloromethane, N-methyl-pyrrolidone or dimethylformamide, eventually supplemented with helical structure-stabilising particular polar solvents such
15 trifluoro-ethanol, is far from providing a general remedy for difficult peptides.

Thymosin α_1 (international non-proprietary pharmaceutical name: thymalfasin) is an N-acetylated 28-mer peptide hormone that occurs in the human thymus gland and is used a drug to treat chronic hepatitis B.

20

Its linear full-length solid-phase synthesis has been described early on but suffered from poor yields up to now. Unfortunately, the Thymosin α_1 amino acid sequence does not offer a glycine or proline which could be used as a junctional residue in a classical segment coupling approach in a stepwise approach to synthesis.

25

Echner et al. (Liebigs Ann. Chem. 1988, 1095-1097) describe linear solid-phase synthesis on a Wang-polystyrene resin. Whilst coupling efficiency is allegedly checked after each coupling with BOP reagent in DMF by means of Kaiser Test, only after every fifth coupling capping of uncoupled peptide by means of acetylation is done. Cleavage of the protected and terminally
30 acetylated peptide from the resin is achieved with concentrated trifluoro acetic acid; thereafter, resin and liquid phase are separated by filtration and the peptide product is collected by precipitation with diethylether. The precipitate collected in this way is quantified and is

designated 'crude product', yield is calculated as to amount of 76%. – Only thereafter, a first chromatographic purification of the crude product is undertaken, followed by reverse phase HPLC. No yield is indicated for the pure product thus obtained.


A problem with Echnér's approach is that incomplete chain coupling, leading to shortened undesirable sequence variants during synthesis, is not accounted for in the yield calculation. 'Crude product' amounts to every peptidic species obtained during synthesis, including incompletely deprotected/alkylated peptide. However, purity peptide obtained from linear synthesis is problem when dealing with difficult peptides; interestingly, Echnér does not give any indication of yield of purified product thus obtained, nor is any HPLC chromatogram given in the publication as is routinely done in the art for proving quality of synthesis. In our hands, the synthetic scheme of Echnér gives only very low yields of correct product, a large array of improper peptide product making up for most of the crude product.

Another problem with Echnér's approach which the author raises himself (p.1095, left column, 3rd para) is loading of the Wang resin with the C-terminal amino acid of Thymosin α_1 which is an asparagine; the loading Echnér achieves is 0.15 mmol/g, which is extremely low. Fmoc-Asn(4,4'-dimethoxydiphenylmethyl)-OH amino acid must be used for loading of the resin without running risk of degrading side-reactions.

It is the aim of the embodiments of present invention to devise an improved solid-phase synthesis for Thymosin α_1 . This aim is solved by the method of claim 5.

Summary

In one aspect, there is provided a peptide conjugate of formula I



I

Wherein each amino acid residue is individually protected or unprotected, R3 is a PEG resin solid phase which is comprising an integral linker, R2 is a grafted linker with x being 0 or 1 and R1 is hydrogen, a protection group or a peptidyl radical, preferably a peptidyl radical counting less than 50, preferably less than 25 amino acid residues, and wherein, if R1 is a peptidyl radical, the individual amino acid residues of said radical are individually protected or unprotected and the N-terminus of said radical is free, is acetylated or is protected with a protection group Y.

In another aspect, there is provided a method of synthesizing Thymosin α_1 , a C-terminally truncated version of Thymosin α_1 comprising residues 1-27, an N-terminally truncated version of Thymosin α_1 comprising at least residues 19-28 or a truncated version comprising at least residues 19-27 of mature Thymosin α_1 on a solid phase comprising the steps of a. coupling a suitably protected FMOC-Glu, FMOC-Asp or FMOC-Asn residue onto a PEG resin solid phase which PEG resin is preferably selected from the group comprising polystyrene-PEG mixed resins, PEG polyether-polyester mixed resins or PEG polyether-polyamide mixed resins and essentially pure PEG resins b. elongating the peptide chain by FMOC synthesis wherein the side-chains of individual amino acids may be derivatized with suitable protection groups c. optionally acetylating the last, N-terminal residue and d. cleaving the peptide from the resin.

Detailed Disclosure of Embodiments

According to the present invention, a method of synthesizing on a solid phase Thymosin α_1 or a C-terminally truncated version comprising residues 1-27 or an N-terminally truncated version comprising at least residues 19-28 or an truncated version comprising at least residues 19-27 of mature Thymosin α_1 is devised, comprising the steps of

- a. coupling a suitably protected FMOC-Glu or FMOC-Asn residue onto a PEG resin which PEG resin is preferably selected from the group comprising polystyrene-PEG mixed resins, PEG polyether-polyester mixed resins or PEG polyether-polyamide mixed resins and essentially pure PEG resins

- b. elongating the peptide chain by Fmoc synthesis wherein the side-chains of individual amino acids may be derivatized with suitable protection groups
- c. optionally acetylating the last, N-terminal residue and
- d. cleaving the peptide from the resin

5

The present method allows of synthesizing the thymalfasin peptide in unprecedented purity and hence high yield in a single solid-phase linear synthetic approach. According to the present invention, it may also be feasible though to synthesize only a C-terminal fragment on a solid phase and to constitute any full-length peptide or an at least longer version of the peptide by segment coupling, using in one preferred embodiment coupling of an N-terminal fragment

10 having a C-terminal Ser or Thr which is protected to racemization by the pseudoproline dipeptide method (Wöhr et al., J. Am. Chem. Soc. 118, 9218).

Preferably, the peptide to be synthesized is Thymosin α_1 (also referred to as mature, active

15 Thymosin α_1 as to distinguish from precursor peptides) having the sequence

1-Ser-Asp-Ala-Ala-Val-Asp-Thr-Ser-Ser-Glu-Ile-Thr-Thr-Lys-Asp-Leu-Lys-Glu-Lys-Lys-Glu-Val-Val-Glu-Glu-Ala-Glu-Asn-28

wherein the individual amino acid chains are unprotected or are suitably protected, as need may

20 be for a given individual amino acid residue as is common knowledge in the art (see Bodansky, *infra*, for instance). More preferably, such peptide is N-terminally acetylated in a step subsequent to linear synthesis but prior to cleavage from resin. Natural thymalfasin which is also used as a pharmaceutical is N-terminally acetylated.

25 Notably, the present inventors have found that it is mainly the C-terminal half of the thymalfasin peptide, it is about residues 19-28, that incur most of the impurities by chain deletions and hence losses in final product yield generated during linear phase synthesis. This part of the peptide has been found to be extremely difficult to synthesize.

30 Vice versa, during linear synthesis, it is also feasible to use such oxazolidine dipeptides according to Wöhr et al., *supra*. A benefit of using such dipeptide instead of individual protected, preferably Fmoc-, amino acid can be inferred from the thymalfasin sequence in

particular for the dipeptide Asp-Thr; use of oxazolidine dipeptide derivative would help here to avoid aspartimide formation during synthesis.

Similarly, prevention of glutarimide formation may require use of special protection groups.

5

Whilst the present invention strongly prefers Fmoc chemistry for carrying out the coupling reactions, Boc chemistry may likewise be used in another embodiment to carry out the present invention.

- 10 Coupling reagents for peptide synthesis are well-known in the art (see Bodansky, M., Principles of Peptide Synthesis, 2nd ed. Springer Verlag Berlin/Heidelberg, 1993; also see discussion of role of coupling additives or auxiliaries therein). Coupling reagents may be mixed anhydrides (e.g. T3P: propane phosphonic acid anhydride) or other acylating agents such as activated esters or acid halogenides (e.g. ICBF, isobutyl-chloroformiate), or they may be carbodiimides (e.g. 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide), activated benzotriazin-derivatives (DEPBT: 3-(diethoxyphosphoryloxy)-1,2,3-benzotriazin-4(3H)-one) or uronium or phosphonium salt derivatives of benzotriazol.
- 15

- In view of best yield, short reaction time and protection against racemization during chain elongation, more preferred is that the coupling reagent is selected from the group consisting of uronium salts and phosphonium salts of the benzotriazol capable of activating a free carboxylic acid function along with that the reaction is carried out in the presence of a base. Suitable and likewise preferred examples of such uronium or phosphonium coupling salts are e.g. HBTU (O-(1H-benzotriazole-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate), BOP (benzotriazole-1-yl-oxy-tris-(dimethylamino)-phosphonium hexafluorophosphate), PyBOP (Benzotriazol-1-yl-oxy-tripyrrolidinophosphonium hexafluorophosphate), PyAOP, HCTU (O-(1H-6-chloro-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate), TCTU (O-(1H-6-chlorobenzotriazole-1-yl)-1,1,3,3-tetramethyluronium tetrafluoroborate), HATU (O-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate), TATU (O-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium tetrafluoroborate), TOTU (O-[cyano(ethoxycarbonyl)methyleneamino]-N,N,N',N''-tetramethyluronium tetrafluoroborate), HAPyU (O-(benzotriazol-1-yl)oxybis-(pyrrolidino)-uronium hexafluorophosphate).
- 20
- 25
- 30

Preferably, when using DEPBT or the like, uronium or phosphonium salt reagents, a further or second weak base reagent is needed for carrying out the coupling step. This is matched by base whose conjugated acid has a pKa value of from pKa 7.5 to 15, more preferably of from pKa 7.5 to 10, with the exclusion of an α -amino function of a peptide or amino acid or amino acid derivative, and which base preferably is a tertiary, sterically hindered amine. Examples of such and further preferred are Hünig-base (N,N-diisopropylethylamine), N,N'-dialkylaniline, 2,4,6-trialkylpyridine or N-alkyl-morpholine with the alkyl being straight or branched C1-C4 alkyl, more preferably it is N-methylmorpholine or collidine (2,4,6-trimethylpyridine), most preferably it is collidine.

The use of coupling additives, in particular of coupling additives of the benzotriazol type, is also known (see Bodansky, supra). Their use is particularly preferred when using the highly activating, afore said uronium or phosphonium salt coupling reagents. Hence it is further preferred that the coupling reagent additive is a nucleophilic hydroxy compound capable of forming activated esters, more preferably having an acidic, nucleophilic N-hydroxy function wherein N is imide or is N-acyl or N-aryl substituted triazeno, most preferably the coupling additive is a N-hydroxy-benzotriazol derivative (or 1-hydroxy-benzotriazol derivative) or is an N-hydroxy-benzotriazine derivative. Such coupling additive N-hydroxy compounds have been described in large and wide in WO 94/07910 and EP-410 182 and whose respective disclosure is incorporated by reference hereto. Examples are e.g. N-hydroxy-succinimide, N-hydroxy-3,4-dihydro-4-oxo-1,2,3-benzotriazine (HOObt), 1-hydroxy-7-azabenzotriazole (HOAt) and N-hydroxy-benzotriazole (HObt). N-hydroxy-benzotriazine derivatives, first and foremost HOObt, are particularly preferred.


Ammonium salt compounds of coupling additives are known and their use in coupling chemistry has been described, for instance in US4806641.

In a further particularly preferred embodiment, the uronium or phosphonium salt coupling reagent is an uronium salt reagent, preferably it is HCTU, TCTU or HBTU, more preferably it is HCTU or TCTU, and most preferably it is used in the reaction in combination with N-hydroxy-3,4-dihydro-4-oxo-1,2,3-benzotriazine or a salt thereof.

In the context of the present invention, it is to be noted that HCTU and TCTU are defined as to be encompassed by the term 'uronium salt reagent' despite that these compounds and possible analogues have been shown to comprise an isonitroso moiety rather than an uronium moiety by

means of crystal structure analysis (O. Marder, Y. Shvo, and F. Albericio "*HCTU and TCTU: New Coupling Reagents: Development and Industrial Applications*", Presentation Gordon Conference Feb. 2002 & *Chimica Oggi* 2002, 20:37-41), an N-amidino substituent on the heterocyclic core giving rise to a guanidium structure instead. In the present context, such class
 5 of compounds is termed 'guanidium-type subclass' of uronium salt reagents according to the present invention.

Deprotection of the base labile Nu may be carried out as routinely done in the art, e.g. with 20%
 10 piperidine in N-methyl morpholine in case of Fmoc chemistry. The last amino acid to be added, e.g. typically the N-terminal Serine of mature tymalfasin, may of course carry an N-terminal protection group other than Fmoc, e.g. it may carry Boc or readily an acetyl-protection group, though the latter may more conveniently introduced by post-coupling acetylation of the deprotected N-terminal amino acid of the peptide usually. Preferably, one may use an
 15 orthogonal protection group other than Fmoc for protection of the last, N-terminal residue. Examples are Alloc protection groups (e.g. for protecting primary amines), both being selectively removable from peptide by Pd(0)catalyzed transacylation (Gomez-Martinez, N^α-Alloc temporary protection in solid-phase peptide synthesis, use of amine-borane complexes as allyl group scavengers, *J.Chem. Soc. , Perkin Trans. 1*, 1999, 2871-2874), the Dde group
 20 (Bycroft et al., A novel lysine protecting procedure for SPPS of branched peptides, *J. Chem. Soc. Chem. Commun.* 1993, p. 778-779), a dimedon derivative that is removable by hydrazinolysis, and its functional homologues such as e.g. Nde group (N-1-(4-nitro-1,3-dioxindan-2-ylidene)-ethyl group, Kellam et al., *Tetrahedron* 54, 1998, p. 6817-6832; N-1-(4,4-dimethyl-2,6-dioxocyclohexylidene)-3-methylbutyl, Chan 1995). Dde, its derivatives and
 25 homologues are collectively addressed as Dde-type protection groups sharing in their free, unreacted state a functional dioxoalkylidene moiety of formula III



wherein R is substituted or unsubstituted alkyl and wherein preferably the two carbonyl
 30 functions are forming a cyclic structure connected by a $-\text{CH}_2-\text{CR}'\text{R}''-\text{CH}_2-$, $-\text{NR}'-\text{CO}-\text{NR}''-$ or $-\text{CR}'=\text{CR}''-$ backbone. Preferably, R', R'' are then alkyl or, taken together, aryl.

It is also readily feasible to use modified or tagged orthogonal protection groups, which bear the additional benefit of allowing of simple purification by subsequent affinity chromatography or affinity-improved 'standard' RP HPLC using such reversible affinity tag. In this case, the peptide synthesized is cleaved off the resin suitably under mild condition avoiding global deprotection, purification is taking place, the N-terminal probe is selectively removed allowing of subsequent N-terminal acetylation and finally global deprotection of peptide. An example of such are the Fmoc derivatives described in Ball et al., *Int. J. Peptide Protein Res.* 1992, 40:370-379 and Ball et al., *J. Chromatography* 1994, 686:73-88.

Particularly preferred for the present invention is to carry out removal of the Fmoc group with 20% piperidine-DMF (v/v), e.g. (2 × 1 min, 2 × 10 min, 1 × 5 min). Likewise preferred, couplings of Fmoc-aa-OH (5 equiv) are carried out with the above mentioned coupling reagents in DMF or similar solvent for 60-120 min. For optimizing synthesis and in particular coupling times for individual sequence positions, after the coupling ninhydrin tests were carried out and if it was positive the coupling was repeated under the same conditions, otherwise the process was continued. If after second coupling the ninhydrin test was still positive an acetylation step was carried out with Ac₂O and DIEA or similar method for capping the deletion peptide chains. Such enhanced coupling and/or capping procedure was then applied to repeated synthesis under the same conditions for the very given sequence position. Notably, choice of resin was found to strongly influence sequence-position specific coupling problems as is exemplified in the experimental section.

According to the present invention, the solid-phase comprises a PEG resin. According to the present inventions, they allow of synthesizing tymalfasin or the difficult core peptide section of tymalfasin in exceptional yield and purity. In the present context, the definition of 'solid-phase' comprises apart from the inert resin matrix the presence of an integral linker on the inert matrix material, for linkage of peptide or for linking a further, grafted linker or handle.

Such resins have been described e.g. in US2003078372 A1; by virtue of the PEG incorporated into the solid-phase, such resins gain amphiphilic properties which makes them different from traditional resin materials. They are known to adopt a 'gel-like' behaviour instead of the solid-like behaviour of traditional e.g. Merrifield resin after swelling. This particular feature is said to offer benefits of reduced cycling times in view of diffusion/liquid exchange e.g. during

washing, coupling, deprotection steps etc.. Notably, according to the present invention, such PEG resin is also found to improve efficiency of difficult, peptide-specific coupling steps for a peptide, drastically and in an unexpected, synergistic fashion enhancing purity and yield. The term 'PEG resin' is to be construed strictly *generically* in the present context as to relate to a
5 polymeric resin comprising a polyether copolymeric or block copolymer share at least – a PEG-type polyether or PEG-polyether resin for short. The term polyether as used here relates in its conventional sense to polyoxyalkenyl polymers such as e.g. polyoxyethylene or polyethyleneglycol (PEG) respectively, polyoxypropylene, polyoxybutylene or mixed polymers thereof. Different embodiments of such type of resin exist, polymerisation of carbon monomers
10 (including monomer units other than oxiran) not being restricted to polyether chemistry only. Accordingly, the resin matrix of the solid-phase may be constituted e.g. by an amphiphilic polystyrene-PEG mixed resin (e.g. Tentagel, see US4908405) or PEG-polyamide or PEG-polyester mixed resin, e.g. Kempe et al., J. Am. Chem. Soc. 1996, 118, 7083; also cp. US5,910,554 A. Usually, resin loading is usually less efficient and/or chemical stability in
15 particular in acidic media is oftenly not comparable to that of traditional resins, e.g. classic Merrifield polystyrene-DVB. - Polystyrene-PEG mixed resins are reaching higher loadings, but lower amphiphilicity is obtained because the PEG contents is decreased.

It is also possible to use a PEG resin that has been obtained by grafting of a non-PEG resin
20 matrix with a layer of a suitable PEG resin material, e.g. cp. Kates et al., High-load polyethylene glycol-polystyrene (PEG-PS) graft supports for solid-phase synthesis, Biopolymers 1998, 47:365-380.


In a strongly preferred embodiment according to the present invention, the PEG resin is a
25 substantially pure PEG-polyether resin that is more preferably devoid of any, eventually further substituted, polystyrene co-polymeric or block-copolymeric share and, again more preferably, further does also not comprise any internal ester or amide functional groups but only polyether functional groups. A substantially pure PEG-polyether resin may also be termed an essentially pure PEG-polyether resin or a resin consisting essentially of PEG-polyether share as defined
30 above. Most preferably, it is a pure PEG resin or PEG-polyether resin. Preferably, the meaning of 'substantially pure' is that the above structural definition of pure PEG resin amounts to at least 70% of the dry weight of the solid-phase material, allowing of a minor share of polymeric additions of other type. Such ideally pure PEG resin material offers optimal chemical stability

along with good compatibility and ease of handling with standard solvents during normal Fmoc synthesis. Notably, in the latter definition the terminal ester or amide bonds bridging the resin directly or via the integral and/or grafted linker to the peptide are not accounted for, since they are not internal structural, crosslinking features and accordingly do not affect the definition of 'pure PEG' in this regard. Such 'pure', highly amphiphilic PEG resins are offered by different companies, e.g. Matrix Innovations Inc. from Quebec, Canada ('ChemMatrix' brand) or Versamatrix A.S. from Denmark and are described e.g. in WO 02/40559. They may offer loading capacity of more than 0.5 mmol/g, see ChemMatrix brand resin described in afore said WO'559 for instance. For linkage of peptide, such resin may have terminal integral linkers such as hydroxymethyl radicals or derived thereof 'quasi-native' amino-methyl, carboxyl or bromomethyl or iodomethyl radicals for instance, or may be derivatized by known grafted linkers or handles for solid phase linkage such as e.g. Wang, PAL, 4-alkoxy-benzaldehyde or Rink or Sieber amide linkers.

Notably, as a further element of the surprising properties of PEG resin according to the present invention, coupling of the second amino acid No. 27, Glu, was suddenly found to become a critically difficult step, in particular when using side-chain anchoring of the terminal Asp/Asn residue as compared to traditional resins (e.g. PS-DVB or CTC-PS-DVB), whilst the coupling efficiency of the other critical residues in the sequence stretch 19-28 was found to be improved, though not rendered trivial. Hence extended and/or repeated coupling of residue No. 27 to the resin-anchored Asn moiety was required when using PEG resin, and especially pure or substantially pure PEG-polyether resin. 'Extended' relates to using more equivalents of an Fmoc amino acid or longer coupling times than for the average residue.

Further aims of the embodiments of the present invention are the various peptide-solid phase conjugates obtained in this way. The definitions and explanations given in the foregoing apply likewise to these aims.

Such aim is a peptide conjugate of formula I



I

5 wherein each amino acid residue may be individually protected or unprotected, R3 is a PEG-resin solid phase which is comprising an integral linker, R2 is a grafted linker with x being 0 or 1 and R1 is hydrogen, a protection group or a peptidyl radical, preferably a peptidyl radical counting less than 50, preferably less than 25 amino acid residues, and wherein, if R1 is a peptidyl radical, the individual amino acid residues of said radical are individually protected or
 10 unprotected and the N-terminus of said peptidyl radical is free, is acetylated or is protected with a protection group Y.

Accordingly, preferably R1 is a peptidyl radical which is Y-Ser-Asp-Ala-Ala-Val-Asp-Thr-Ser-Ser-Glu-Ile-Thr-Thr-Lys-Asp-Leu-Lys-Glu-, Ac-Ser-Asp-Ala-Ala-Val-Asp-Thr-Ser-Ser-
 15 Glu-Ile-Thr-Thr-Lys-Asp-Leu-Lys-Glu or H-Ser-Asp-Ala-Ala-Val-Asp-Thr-Ser-Ser-Glu-Ile-Thr-Thr-Lys-Asp-Leu-Lys-Glu-Leu-Lys-Glu with Y having the meaning named before. Ac: Acetyl.

20 Linkers as defined by Guillier, Orain, and Bradley, Chem. Rev. 100, 2091-2157, 2000 are considered simply as immobilized protecting groups and are classified into one of two types: (a) *Integral linkers* in which part of the solid phase core material forms or is inseparably part of the linker, both together constituting R3, and (b) *Nonintegral or grafted linkers R2* in which the linker R2 is additionally attached to the solid support, on top of an integral linker/solid phase
 25 R3, e.g. for allowing of more mild cleavage conditions. Typical examples of integral linker-solid phases are *p*-methylbenzhydrylamine (MBHA), 2-chlorotrityl (CTC), amino-methyl, etc... Integral linkers are mandatory whilst grafted linkers R2 are optional. Typical examples are 4-hydroxymethylphenoxyacetyl- or 4-hydroxymethylbenzoic acid.

30 Conventionally, the peptide is conjugated through its C-terminal residue via its 1-carboxylic acid function to the solid phase or the grafted linker, respectively.

Preferably, according to the present invention, the synthesis of tymalfasin or of its aforementioned fragments and variants is achieved by side chain anchoring of the second last-Glu or of a C-terminal Asp or Asn residue. 'Second last-Glu' may mean, in the present context, a Glu-Asn protected dipeptide that is anchored to a solid phase or it may mean that synthesis of a variant of tymalfasin that is devoid of the C-terminal Asn of natural tymalfasin starts with said Glu residue. - Unexpectedly, side chain anchoring has been found to result in more efficient synthesis. A terminal Asp residue that is anchored via its beta-carboxylic function to the solid phase or a grafted linker may be post-synthetically amidated, after cleavage from resin. More preferably though, an Fmoc-Asp which is suitably protected at its C carboxylic function, preferably by means of a tert.butyl protection group, is side-chain anchored to an amide generating linker, preferably a grafted amide-generating linker; with regard to the final product after cleavage from resin, such conjugate may be considered and termed a 'side-chain anchored Asn' as well due to the use of an amide-generating linker. Examples and strongly preferred embodiments of such amide generating linkers are e.g. 'Rink amide' 4-(2',4'-dimethoxybenzyl-aminomethyl)-phenoxy- resin, Sieber resin (Tetrahedron Lett. 1987, 28, 2107-2110) or similar 9-amino-xanthenyl-type resins, PAL resins (Albericio et al., 1987, Int. J. Pept. Protein Research 30, 206-216). Another, less preferred example for amide-generating linkers are the specially substituted trityl-amine derivatives according to Meisenbach et al., 1997, Chem. Letters, p. 1265 f.). These are examples of (Fmoc-chemistry compatible) linker groups from which a C-carboxamid is generated or liberated upon cleavage of the peptide from the resin. It goes without saying that use of such amide linkers is of course dependent on the type of solid phase synthesis carried out, ie. whether is traditional Boc or now customary, orthogonal Fmoc protection chemistry that is used for coupling; a Boc-specific amide resin linker is PAM, for instance. Accordingly, solid phases comprising such linker groups are termed 'amide-producing solid phases' in the present context.

Incorporation of Fmoc-Asp-OtBu onto a solid-phase, containing one or two linkers, using coupling reagents (Albericio, van Abel, Barany, *Int. J. Peptide Protein Res.*, 35, 284-286, 1990), has been described and may be applied in the context of the present invention. Usually, uronium type coupling reagents are preferred for achieving this.


Advantages of this side-chain anchoring strategy are:

(i) Incorporation of the first amino acid into the solid support is carried out with rather quantitative yields through an amide bond and therefore the same reagents used for the formation of peptide bond can be used. In addition, the incorporation of asparagine derivatives into a Wang type resin takes place with low yields, with risk of racemization and with formation of β -cyanoalanine if side-chain amide is not protected (Katsoyannis et al., 1958, Am. chem. soc. 80:2558). Furthermore, side-chain unprotected asparagines could be incorporated to halogen resins through both the 1-carboxylic and the carboxamide function, requiring use of e.g. suitably protected Fmoc-Asn(Mbh)-OH amino acids, creating further problems at the stage of deprotecting the peptide finally.

(ii) Side-chain anchoring strategy involving just acid-labile tBu-based protecting groups (tBu esters for the 1-carboxylic acid of the first residue and ω -carboxylic acids of Asp and Glu residues, tBu ethers for Ser and Thr side-chain) and acid-labile Boc for Lys side-chain will facilitate the final global deprotection, because simple scavengers such as H₂O can be used. In contrast, if Asn is incorporated with trityl, xantyl, or 2,4,6-trimethoxybenzyl protecting groups, it will require the presence of additional scavengers.

(iii) The growing backbone peptide chain into the side-chain anchoring strategy will have more flexibility than in the 1-carboxylic anchoring strategy, because the anchoring is through the side-chain.

Scheme I below recapitulates the most preferred mode of synthesis on a PEG resin according to the present invention for tymalfasin (Zadaxine), most preferably on a pure or substantially pure PEG resin, wherein linker 1 is an amide-generating linker and linker 2 is an integral linker:

Scheme I

- 5 (i) Incorporation of Fmoc-Asp-OtBu onto a solid support (SS), containing one or two linkers, using coupling reagents (Albericio, van Abel, Barany, *Int. J. Peptide Protein Res.*, **35**, 284-286, 1990) .
- (ii) elongating the peptidic chain with the rest of Fmoc-protected amino acid (tBu for Asp, Glu, Ser, and Thr; and Boc for Lys).
- 10 (iii) acetylation

- (iv) cleavage of the peptide from the solid support and removal of the protecting groups, which can be carried out in one (high acid concentration solution) or two steps (low acid concentration solution followed by high acid concentration solution).

EXAMPLES

General Procedures. Fmoc-Sieber-PS-resin was from NovaBiochem (Läufelfingen, Switzerland), ChemMatrix resin from Matrix Innovation (Quebec, Canada), 2-Cl-TrtCl-resin
5 from CBL (Patras, Greece), HCTU from Luxembourg Industries Ltd. (Tel Aviv, Israel), Protected Fmoc-amino acid derivatives from IRIS Biotech (Marktredwitz, Germany).

Solid-phase syntheses were carried out in either glass funnels fitted with filters or polypropylene syringes (50 mL) fitted with a polyethylene porous disc. Solvents and soluble reagents were removed by suction. Removal of the Fmoc group was carried out with piperidine–
10 DMF (2:8, v/v) (2 × 1 min, 2 × 10 min, 1 × 5 min). Washings between deprotection, coupling, and, again, deprotection steps were carried out with DMF (5 × 0.5 min) and CH₂Cl₂ (5 × 0.5 min) using each time 10 mL solvent/g resin. Peptide synthesis transformations and washes were performed at 25 °C. Syntheses carried out on solid-phase were controlled by HPLC of the intermediates obtained after cleaving with TFA–H₂O (95:5) for 1 h an aliquot (aprox. 2 mg) of
15 the peptidyl-resin. HPLC reversed phase columns Symmetry™ C₁₈ 4,6 × 150 mm, 5 μm (column A) and were from Waters (Ireland). Analytical HPLC was carried out on a Waters instrument comprising two solvent delivery pumps (Waters 1525), and automatic injector (Waters 717 autosampler), dual wavelength detector (Waters 2487). UV detection was at 215 or 220 nm, and linear gradients of CH₃CN (+0.036% TFA) into H₂O (+0.045% TFA), from 30%
20 to 100% in 15 min.

MALDI-TOF and ES-MS analysis of peptide samples were performed in a PerSeptive Biosystems Voyager DE RP, using ACH matrix.

25

Example 1

Ac-Ser(tBu)-Asp(OtBu)-Ala-Ala-Val-Asp(OtBu)-Thr(tBu)-Ser(tBu)-Ser(tBu)-Glu(OtBu)-
Ile-Thr(tBu)-Thr(tBu)-Lys(Boc)-Asp(OtBu)-Leu-Lys(Boc)-Glu(OtBu)-Lys(Boc)-Lys(Boc)-
30 Glu(OtBu)-Val-Val-Glu(OtBu)-Glu(OtBu)-Ala-Glu(OtBu)-Asp(Rink-ChemMatrix-resin)-
OtBu

Step 1

Side chain anchoring of Fmoc-Asp(Rink-ChemMatrix-resin)-OtBu

Fmoc-Rink-ChemMatrix-resin (0.45 mmol/g) (5 g, 2.25 mmol) was placed in a glass funnel fitted with a porous disk. The resin was subjected to the following washings/treatments with CH₂Cl₂ (3 × 0.5 min), DMF (3 × 0.5 min), and piperidine as indicated in General Procedures, and DMF (5 × 0.5 min). Then, Fmoc-Asp-OtBu (4.11 g, 5 equiv) and DIEA (3.4 mL, 10 equiv) in DMF (9 mL) were added, followed by HCTU (3.96 g, 4.8 equiv) in DMF (5 mL). The mixture was left to stir mechanically for 2 h and the resin was washed with DMF (3 × 0.5 min) and the ninhydrin test was negative. Ac₂O (0.5 mL, 5 mmol) and DIEA (1.7 mL, 5 mmol) in DMF (14 mL) were added and left to stir mechanically for 15 min, where the resin was filtered and washed with DMF (3 × 0.5 min). Successful loading was determined as to amount to 0.4 mmol/g.

Step 2

H-Ser(tBu)-Asp(OtBu)-Ala-Ala-Val-Asp(OtBu)-Thr(tBu)-Ser(tBu)-Ser(tBu)-Glu(OtBu)-Ile-Thr(tBu)-Thr(tBu)-Lys(Boc)-Asp(OtBu)-Leu-Lys(Boc)-Glu(OtBu)-Lys(Boc)-Lys(Boc)-Glu(OtBu)-Val-Val-Glu(OtBu)-Glu(OtBu)-Ala-Glu(OtBu)-Asp(Rink-ChemMatrix-resin)-OtBu

The Fmoc group was removed and Fmoc-aa-OH's (5-10 equiv) were sequentially added to the above peptidyl-resin (step 1) together with DIEA (3.4 mL, 10 equiv) in DMF (9 mL), by HCTU (3.96 g, 4.8 equiv) in DMF (5 mL). The mixture was left to stir mechanically for 1-2 h, the resin was washed with DMF (3 × 0.5 min) and the ninhydrin test was taken. If test resulted positive (blue), the coupling was repeated, while if the test was slightly (brown), an acetylation step was carried out with Ac₂O (0.5 mL, 5 mmol) and DIEA (1.7 mL, 5 mmol) in DMF (14 mL) for 15 min with mechanical stirring. Aliquots of resins were taken after piperidine treatments. At positions E27, K19, for each position a double coupling with at least 35 min. reaction time per single and using 10 eq. of Fmoc amino acids was performed.

Step 3

*Ac-Ser(tBu)-Asp(OtBu)-Ala-Ala-Val-Asp(OtBu)-Thr(tBu)-Ser(tBu)-Ser(tBu)-Glu(OtBu)-Ile-
Thr(tBu)-Thr(tBu)-Lys(Boc)-Asp(OtBu)-Leu-Lys(Boc)-Glu(OtBu)-Lys(Boc)-Lys(Boc)-Glu(OtBu)-
5 Val-Val-Glu(OtBu)-Glu(OtBu)-Ala-Glu(OtBu)-Asp(Rink-ChemMatrix-resin)-OtBu*

The final acetylation was carried out with Ac₂O (0.5 mL, 5 mmol) and DIEA (1.7 mL, 5 mmol) in DMF (14 mL) for 15 min with mechanical stirring, where ninhydrin was negative. The resin was dried and 16.5 g were obtained.

10

Step 4

*Ac-Ser-Asp-Ala-Ala-Val-Asp-Thr-Ser-Ser-Glu-Ile-Thr-Thr-Lys-Asp-Leu-Lys-Glu-Lys-
15 Lys-Glu-Val-Val-Glu-Glu-Ala-Glu-Asn-OH (Zadaxin, tymalfasin)*

One step cleavage from resin and global deprotection was performed. Portions of 5 g of resin from Step 3 were placed in the glass funnel fitted with a porous disk were treated with cold TFA-H₂O (95:5) (35 mL) for 2 h, then the solution is filtered off and the resin is further washed
20 with TFA-H₂O (95:5) (10 mL). The combined TFA solution is poured on cold *t*-butyl methyl ether (350 mL) and the mixture is centrifugated. The solid is isolated by decantation.

MALDI-TOF-MS, calcd. 3106.50. Found: *m/z* 3107.36 [M+H]⁺.

RP-HPLC analysis of the crude peptide showed that the product was predominantly pure, amounting to 90% yield (Fig.1). Two minor contaminating peaks could be easily removed by
25 RP-HPLC on C18 Hypersil with 10-30% acetonitril/water over 45 min., giving pure product with almost complete recovery of the correct product peak.(Fig. 2).

30

Example 2: Comparative example

*Ac-Ser-Asp-Ala-Ala-Val-Asp-Thr-Ser-Ser-Glu-Ile-Thr-Thr-Lys-Asp-Leu-Lys-Glu-Lys-
Lys-Glu-Val-Val-Glu-Glu-Ala-Glu-Asn-OH (Zadaxin, tymalfasin)*

35

Experimental procedures as described in Examples 1 and 2, except that Fmoc-Sieber-PS-resin (0.59 mmol/g) (3.33 g, 2 mmol) was used instead of Fmoc-Rink-ChemMatrix-resin.

MALDI-TOF-MS, calcd. 3106.50. Found: m/z 3107.24 $[M+H]^+$. The purity and yield of the curde product mix analyzed by RP-HPLC as determined by peak area amounted to 40% yield only (Fig. 3), with a broad spectrum of impurities.

Example 3: Comparative Example

Ac-Ser(tBu)-Asp(OtBu)-Ala-Ala-Val-Asp(OtBu)-Thr(tBu)-Ser(tBu)-Ser(tBu)-Glu(OtBu)-Ile-Thr(tBu)-Thr(tBu)-Lys(Boc)-Asp(OtBu)-Leu-Lys(Boc)-Glu(OtBu)-Lys(Boc)-Lys(Boc)-Glu(OtBu)-Val-Val-Glu(OtBu)-Glu(OtBu)-Ala-Glu(OtBu)-Asn(Trt)-O-(2-Cl-Trt)-resin

Step 1

Ca-terminal anchoring on CTC resin, giving Fmoc-Asn(Trt)-O-TrtCl-resin

2-Cl-TrtCl-resin (0.2 g, 1.64 mmol/g; Ambersynth™ CTC from Rohm & Haas, Paris, France with matrix made from polystyrene-1%divinylbenzene) was placed in a 10 mL polypropylene syringe fitted with a polyethylene filter disk. The resin was then washed with CH₂Cl₂ (5 × 0.5 min), and a solution of Fmoc-Asn(Trt)-OH (0.7 equiv) and DIEA (241 μL) in CH₂Cl₂ (2.5 mL) was added, and the mixture was stirred for 15 min, when extra DIEA (121 μL, total 7 equiv) and the mixture stirred for 45 min. The reaction was terminated by addition of MeOH (160 μL), after a stirring of 10 min. The Fmoc-Asn(Trt)-O-TrtCl-resin was subjected to the following washings with CH₂Cl₂ (3 × 0.5 min) and DMF (3 × 0.5 min). The loading calculated by Fmoc determination was 0.8 mmol/g.

Step 2

H-Ser(tBu)-Asp(OtBu)-Ala-Ala-Val-Asp(OtBu)-Thr(tBu)-Ser(tBu)-Ser(tBu)-Glu(OtBu)-Ile-Thr(tBu)-Thr(tBu)-Lys(Boc)-Asp(OtBu)-Leu-Lys(Boc)-Glu(OtBu)-Lys(Boc)-Lys(Boc)-Glu(OtBu)-Val-Val-Glu(OtBu)-Glu(OtBu)-Ala-Glu(OtBu)-Asn(Trt)-ClTrt-PS-resin

The Fmoc group was removed and Fmoc-aa-OH's (10 equiv) were sequentially added to the above peptidyl-resin (step 1) together with DIEA (20 equiv) in DMF (9 mL), by HCTU (9.6 equiv) in DMF an ABI automatic synthesizer 433A with 60 min coupling time.

5

Step 3

Ac-Ser(tBu)-Asp(OtBu)-Ala-Ala-Val-Asp(OtBu)-Thr(tBu)-Ser(tBu)-Ser(tBu)-Glu(OtBu)-Ile-Thr(tBu)-Thr(tBu)-Lys(Boc)-Asp(OtBu)-Leu-Lys(Boc)-Glu(OtBu)-Lys(Boc)-Lys(Boc)-Glu(OtBu)-Val-Val-Glu(OtBu)-Glu(OtBu)-Ala-Glu(OtBu)-Asn(Trt)-ClTrt-PS-resin

10

The final acetylation was carried out with with Ac₂O (0.5 mmol) and DIEA (5 mmol) in DMF (2 mL) for 15 min with sporadic manual stirring, where ninhydrin was negative.

Step 4

15

Ac-Ser(tBu)-Asp(OtBu)-Ala-Ala-Val-Asp(OtBu)-Thr(tBu)-Ser(tBu)-Ser(tBu)-Glu(OtBu)-Ile-Thr(tBu)-Thr(tBu)-Lys(Boc)-Asp(OtBu)-Leu-Lys(Boc)-Glu(OtBu)-Lys(Boc)-Lys(Boc)-Glu(OtBu)-Val-Val-Glu(OtBu)-Glu(OtBu)-Ala-Glu(OtBu)-Asn(Trt)-OH

20

The protected peptide from step 3 above was cleaved from the resin by TFA-Et₃SiH-CH₂Cl₂ (1:1:98) (5 × 30 sec). Filtrate was collected on H₂O (4 mL) and the H₂O was partially removed under reduced pressure. Acetonitril (ACN) was then added to dissolve solid that appeared during the H₂O removal, and the solution was lyophilized.

Step 5

25

Ac-Ser-Asp-Ala-Ala-Val-Asp-Thr-Ser-Ser-Glu-Ile-Thr-Thr-Lys-Asp-Leu-Lys-Glu-Lys-Lys-Glu-Val-Val-Glu-Glu-Ala-Glu-Asn-OH (Zadaxin, tymalfasin)

30

The protected peptide from Step 4 was dissolved in TFA-H₂O (95:5, 5 mL) and the mixture was allowed to stir for 1 h. Then the solution is filtered off and the resin is further washed with TFA-H₂O (95:5) (1 mL). The combined TFA solution is poured on cold butyl methyl ether (25 mL) and the mixture is centrifugated. The solid is isolated by decantation. Then, H₂O (5 mL) was added and lyophilized.

MALDI-TOF-MS, calcd. 3106.50. Found: *m/z* [M+H]⁺ 3107.98.

The crude purity and yield, as determined by RP-HPLC peak area, amounted to dissapointing 20% proper product with a strong background smear of multiple improper chain variants (Fig. 4). – Careful repetition of the solid-phase synthesis of a shortened fragement (Lys19 up to Asn28) with manual coupling and ninhydrin testing after every step showed that chain synthesis often effectively terminated after incorporation of V22; three consecutive couplings E21, K20 and K19 proved to be highly inefficient, requiring capping of unreacted chains. Double coupling, elongated coupling times and using more eq. of Fmoc amino acid did not satisfactorily overcome the problem, not allowing to establish significantly more efficient synthesis this way (Fig. 5).

Example 4: Comparative example

Ac-Ser-Asp-Ala-Ala-Val-Asp-Thr-Ser-Ser-Glu-Ile-Thr-Thr-Lys-Asp-Leu-Lys-Glu-Lys-Lys-Glu-Val-Val-Glu-Glu-Ala-Glu-Asn-OH (Zadaxin, tymalfasin)

Experimental procedures essentially as as described in Examples 2, except that Fmoc-Rink handle was incorporated onto MBHA resin (4-methylbenzhydramine resin LL on polystyrene-1%divinylbenzene basis, Novabiochem™, Merck/Germany), Fmoc-Asp-OtBu was side-chain anchored as described in Step1, Example 1 onto Rink aminomethyl grafted linker after removal of Fmoc group of the Rink moiety, loading was determined to amount to 0.66 mmol/g. Couplings were carried out with HCTU (10 eq.) for 35-min. with double couplings at positions D28 (i.e.N28), E27,E21,K20,K19. One step cleavage and deprotection was carried out with TFA-H2O (95:5) after acetylation has been carried out as described above. The crude purity and yielded obtained, as determined by RP-HPLC, was 27% (Fig. 6).

Example 5

NH2-Lys-Lys-Glu-Val-Val-Glu-Glu-Ala-Glu-Asn-OH (fragment 19-28 of tymalfasin)

Side chain anchoring and synthesis was performed on Rink-Chemmatrix resin essentially as described in example 1, except that no double coupling was performed at the sequence positions indicated in example 1 but a uniform, single coupling time of 35 min. was applied. Deletion of the Glu residue No. 27 could was observed as the only but dominant chain termination

byproduct in the HPLC diagram (Fig 7). —As set forth in example 1, this byproduct could be successfully suppressed by extended, repeated coupling at E27.


Example 6

5 NH₂-Lys-Lys-Glu-Val-Val-Glu-Glu-Ala-Glu-Asn-OH (fragment 19-28 of tymalfasin)

Side chain anchoring and synthesis was performed essentially as described in example 1, except that a PEG-MBHA resin from Millipore-Waters (MeO[PEG]2000-CO-Orn-MBHA-PS-resin, similarly described in Kates et al., High-load PEG-polystyrene graft supports for solid-phase
10 synthesis, supra) that was grafted with a Rink aminomethyl linker was employed. The loading of the Rink-PEG-MBHA resin was determined with 0.55 mmol/g. Specially, double couplings were performed at the sequence positions E27,E21,K20,K19. The crude purity and yield determined by RP-HPLC amounted to 92% (Fig. 8).

Claims

- 5 1. Peptide conjugate of formula I



I

10 Wherein each amino acid residue is individually protected or unprotected, R3 is a PEG resin solid phase which is comprising an integral linker, R2 is a grafted linker with x being 0 or 1 and R1 is hydrogen, a protection group or a peptidyl radical, preferably a peptidyl radical counting less than 50, preferably less than 25 amino acid residues, and wherein, if R1 is a peptidyl radical, the individual amino acid residues of said radical are individually protected or unprotected and the N-terminus of said radical is free, is acetylated or is protected with a protection group Y.

- 20 2. Peptide conjugate according to the preceding claim, characterised in that the protection group Y is an Fmoc group, a Dde-type group, a Boc group or a derivative of said protection groups, preferably a derivative of a Fmoc or Dde-type group, that comprises a moiety that allows of reversible affinity binding of the thus protected peptide to a suitable chromatography solid matrix material.
- 25 3. Peptide conjugate according to claim 1, characterised in that the protection group Y allows of reversible affinity binding to an immobilised-metal affinity chromatography solid matrix material.
- 30 4. Peptide conjugate according to one of the preceding claims, characterised in that R1 is Y-Ser-Asp-Ala-Ala-Val-Asp-Thr-Ser-Ser-Glu-Ile-Thr-Thr-Lys-Asp- Leu-Lys-Glu- , Ac-Ser-Asp-Ala-Ala-Val-Asp-Thr-Ser-Ser-Glu-Ile-Thr-Thr-Lys-Asp- Leu-Lys-Glu or H- Ser-Asp-Ala-Ala-Val-Asp-Thr-Ser-Ser-Glu-Ile-Thr-Thr-Lys-Asp- Leu-Lys-Glu-Leu-Lys-Glu with Y having the meaning named before.

5 5. Method of synthesizing Thymosin α_1 , a C-terminally truncated version of Thymosin α_1 comprising residues 1-27, an N-terminally truncated version of Thymosin α_1 comprising at least residues 19-28 or a truncated version comprising at least residues 19-27 of mature Thymosin α_1 on a solid phase comprising the steps of

- 10 a. coupling a suitably protected Fmoc-Glu, Fmoc-Asp or Fmoc-Asn residue onto a PEG resin solid phase which PEG resin is preferably selected from the group comprising polystyrene-PEG mixed resins, PEG polyether-polyester mixed resins or PEG polyether-polyamide mixed resins and essentially pure PEG resins
- b. elongating the peptide chain by Fmoc synthesis wherein the side-chains of individual amino acids may be derivatized with suitable protection groups
- c. optionally acetylating the last, N-terminal residue and
- 15 d. cleaving the peptide from the resin

20 6. Method according to claim 5, characterised in that the peptide synthesized is Thymosin α_1 having the sequence

1-Ser-Asp-Ala-Ala-Val-Asp-Thr-Ser-Ser-Glu-Ile-Thr-Thr-Lys-Asp-Leu-Lys-Glu-Lys-Lys-Glu-Val-Val-Glu-Glu-Ala-Glu-Asn-28

wherein the individual amino acid chains are unprotected or are suitably protected.

25 7. Method according to claim 5, characterised in that the PEG resin is an amphiphilic resin.

8. Method according to claim 7, characterised in that the loading capacity of the PEG resin is at least 0.5 mmol/g and which PEG resin preferably is essentially devoid of polystyrene co-polymeric share.

9. Method according to claim 7, characterised in that the PEG-resin with the exception of the integral and/or grafted linkers is a substantially pure polyether-PEG resin, preferably that it is a pure PEG-polyether resin.