Method and markers for the diagnosis of renal diseases

The present invention relates to the diagnosis, particularly differential diagnosis, of renal diseases.

The number of patients presenting with renal diseases has been increasing in the recent years. Thus, renal diseases present an increasing problem to the health system. Many renal diseases are irreversible, therefore an early diagnosis and/or a differential diagnosis of renal diseases is important. Early diagnosis and a therapy precisely tailored to each particular disease could reduce the number of patients requiring dialysis and could also reduce the high cardiovascular risk of the patients.

Currently, precise diagnosis and/or differential diagnosis relies mostly on kidney biopsies. Although biopsies serve as the current "gold standard" in renal diagnostics, biopsies have the disadvantage of being invasive and therefore being conducted only on selected patients.

Urine analysis is a different approach to diagnose renal diseases. However, currently only few parameters of urine are routinely measured, for example creatinin, urea, albumin, blood cells (such as leukocytes and erythrocytes), bacteria, sugar, urobilinogen, bilirubin and pH value. The diagnostic value of these analyses is limited, as they lack sufficient sensitivity and/or selectivity, particularly for differential diagnosis.

Several attempts have been made to analyze the proteins contained in urine.

V. Thongboonkerd et al. have used two-dimensional polyacrylamide gel electrophoresis (2D-PAGE) in combination with matrix-assisted laser desorption ionization-time-of-flight (MALDI-TOF) mass spectrometry followed by mass fingerprinting to investigate normal human urinary proteins. A total of 67 protein forms of 47 unique proteins was identified (V. Thongboonkerd et al. (2001). Proteomic analysis of normal human urinary proteins

isolated by acetone precipitation or ultracentrifugation. Kidney International, vol. 62, p. 1461-1469).

C.S. Spahr et al. have digested the proteins contained in urine samples with trypsin and identified 751 peptides from 124 proteins by means of liquid chromatography-tandem mass spectrometry (C.S. Spahr et al. (2001). Towards defining the urinary proteome using liquid chromatography-tandem mass spectrometry. I. Profiling an unfractionated tryptic digest. Proteomics vol. 1, p. 93-107).

These studies relate only to healthy individuals. The studies have not addressed the question whether alterations in presence of urinary polypeptides can be used for diagnosis or differential diagnosis of renal diseases.

It has been proposed to use the presence or absence of polypeptides in urine for the diagnosis of membranous glomerulonephritis (MGN) (von Neuhoff et al. (2004). Mass Spectrometry for the Detection of Differentially Expressed Proteins: A Comparison of Surface-Enhanced Laser Desorption/Ionization and Capillary Electrophoresis/Mass Spectrometry. Rapid Communications in Mass Spectrometry, vol. 18: 149-156). However, samples of only 8 patients were used in the study, which was mainly concerned with the comparison of different analysis methods. The actual diagnostic value of the markers has remained unclear.

Consequently, there is need for a fast and simple methods and means for diagnosis, particularly differential diagnosis, of renal diseases.

Accordingly, the object the present invention is to provide methods and means for the diagnosis of renal diseases, particularly for differential diagnosis of renal diseases. It is an particular object of the present invention to provide methods and means for the diagnosis and/or differential diagnosis of IgA-nephropathy, which is the most common glomerulopathy.

According to a first aspect of the present invention, the problem is solved by the use of the presence of at least one polypeptide marker in a urine sample for the diagnosis, preferably the differential diagnosis, of a renal disease, wherein the polypeptide marker is selected from the group of polypeptide markers as shown in table 1 to 22.

In the context of the present invention, it has been found that with the help of the polypeptide markers as shown in table 1 to 22 it is possible to reliably diagnose or differentially diagnose, respectively, different renal diseases.

The present invention has numerous advantages compared to the state of the art. First, the presence of the polypeptide markers according to the invention can be determined in urine samples. Therefore, there is no need to take biopsies. Thus, the present invention allows a simplified and fast diagnosis of renal diseases, allowing to screen patients regularly for the presence of renal diseases and to diagnose renal diseases at early stages. Furthermore, the polypeptide markers according to the invention can be used for differential diagnosis between different renal diseases. The high number of markers identified according to the present invention allows to increase both specificity and sensitivity of diagnosis as compared to the use of only a single or a small number of markers. Also, the present invention provides methods which allow to measure said polypeptide markes without the use of specific ligands such as antibodies or aptamers.

The polypeptide markers as shown in the tables have been identified by a method named capillary electrophoresis-mass spectrometry (CE-MS), which will be described further below. Furthermore, the method has been described in detail in von Neuhoff et al. (2004) (Mass Spectrometry for the Detection of Differentially Expressed Proteins: A Comparison of Surface-Enhanced Laser Desorption/Ionization and Capillary Electrophoresis/Mass Spectrometry. Rapid Communications in Mass Spectrometry, vol. 18: 149-156). Starting from the parameters defining the polypeptide markers, it is possible by methods known in the art to identify the sequence of the corresponding polypeptides and then to synthesize or produce the corresponding polypeptides, e.g. with the help of protein synthesis or expression of the corresponding gene in appropriate cells.

The markers are defined by there mass and their migration time in capillary electrophoresis (CE), particularly mass and their migration time obtained according to Example 1. It is known that CE migration times can vary, typically in the range of 5 min, more typically in the range of 3 minutes. However, the sequence of markers being eluted is typically the same or very similar for each CE system applied. The system can be calibrated by use of polypeptides which are present in almost any urine sample, e.g. by the polypeptides given in tables 23 or 24. Furthermore, the polypeptides given in SEQ ID NO: 1 to SEQ ID NO: 5 can serve for calibration.

Variation of the masses between measurements or between different mass spectrometers is relatively small, typically it is in the range of plus or minus 0.05%.

In table 1, polypeptide markers are listed which are preferred for the discrimination between healthy individuals and individuals suffering from a renal disease, particularly from a glomerulonephritis or glomerulopathy.

In table 2, polypeptide markers are listed, which are preferred for a discrimination between FSGS and the healthy condition.

In table 3, polypeptide markers are listed, which can be used for differential diagnosis between FSGS and MCD.

In table 4, polypeptide markers are listed, which are preferred for a differential diagnosis of FSGS and MGN.

In table 5, polypeptide markers are listed, which are preferred for a differential diagnosis between FSGS on the one hand, and MCD or MGN on the other hand.

In table 6, polypeptide markers are listed, which are preferred for diagnosis of MCD as compared to the healthy condition.

In table 7, polypeptide markers are listed, which are preferred for differential diagnosis between MCD and MGN.

In table 8, polypeptide markers are listed, which are preferred for differential diagnosis between MCD on the one hand, and FSGS or MGN on the other hand.

In table 9, polypeptide markers are listed, which are preferred for diagnosis of MGN as compared to the healthy condition.

In table 10, polypeptide markers are listed, which are preferred for differential diagnosis between MGN on the one hand, and FSGS or MCD on the other hand.

In table 11, polypeptide markers are listed, which are preferred for diagnosis of IgA-nephropathy or MGN on the one hand as compared to the healthy condition.

In table 12, polypeptide markers are listed, which are preferred for diagnosis of IgA-nephropathy as compared to the healthy condition.

In table 13, polypeptide markers are listed, which are preferred for differential diagnosis between IgA-nephropathy and MGN.

In table 14, polypeptides are listed with their respective frequency in healthy, FSGS, MCD, and MGN patients.

In table 15, polypeptides are listed which have been used for differential diagnosis between healthy individuals and renal patients using support vector machines according to Example 1.

In table 16, polypeptides are listed which have been used for differential diagnosis between healthy, FSGS, MCD, and MGN patients using random forest analysis according to Example 1.

In table 17, polypeptides are listed which have been used for differential diagnosis between MCD and MGN patients using using support vector machines according to Example 1.

In table 18, polypeptides are listed which have been used for differential diagnosis between MCD and FSGS patients using using support vector machines according to Example 1.

In table 19, polypeptides are listed which have been used for differential diagnosis between MGN and FSGS patients using using support vector machines according to Example 1.

In table 20 and 21, polypeptides are listed which have been identified in von Neuhoff et al. (2004), which has been cited above.

In table 22, polypeptides are listed which can be used for diagnosis of diabetes and/or diabetic nephropathy.

In table 23, polypeptides are listed, which are preferred as internal standards to standardize the CE-time.

In table 24, polypeptides are listed, which are preferred as internal standards to standardize the CE-time if the pressure method (0.3 to 1 psi) according to Example 1 is used. These standards are e.g. preferred as internal standards in diagnosis of IgA-nephropathy.

In table 25, clinical data of renal patients are listed whose samples were used for identification of polypeptide markers according to Example 1. Abbreviations: CsA, Cyclosporin A; PS , prednisolone; +, frequent relapse; -, currently no immunosuppression; *, clinically unclear whether MCD or FSGS.

The polypeptide markers used according to the present invention can be identified and their presence can be measured in urine samples. Urine samples can be taken as known in the state of the art. Preferably, midstream urine is used in the context of the present invention.

The polypeptide markers used according to the present invention can be gene expression products such as proteins, peptides, and fragments or other degradation products of proteins or peptides. They can be modified by posttranslational modifications, e.g. by glycosylation, phoshorylation, alkylation or disulfide bond. It is known that fragments and degradation products can have a different diagnostic value and/or physiological role than the protein or peptide they have been derived from. For example, in different diseases, different proteolytic degradation products or fragments can be found. It is also considered to be within the scope of the present invention if the urine sample is pretreated to chemically modify the polypeptide markers contained in the urine and to measure these chemically modified polypeptide markers. The polypeptide markers according to the present invention have a molecular mass between 400 and 20 000 Da, particularly between 700 and 14000 Da, more particularly between 800 and 11000 Da.

Preferred polypeptide markers according to the present invention are listed in tables 1 to 22, particularly in tables 1 to 21, more particularly in tables 1 to 13.
Preferred polypeptides for use as internal standards are listed in tables 23 to 24.
Preferred are also polypeptide markers which are listed in table 1, but not in table 14 and/or 15 and/or 16 and/or 17 and/or 18 and/or 19 and/or 20 and/or 21 and/or 22.
Preferred are also polypeptide markers which are listed in table 2, but not in table 14 and/or 15 and/or 16 and/or 18.
Preferred are also polypeptide markers which are listed in table 3, but not in table 14 and/or 16 and/or 18.

Preferred are also polypeptide markers which are listed in table 4, but not in table 14 and/or 16 and/or 19.
Preferred are also polypeptide markers which are listed in table 5, but not in table 14 and/or 16 and/or 18 and/or 19.
Preferred are also polypeptide markers which are listed in table 6, but not in table 14 and/or 16.
Preferred are also polypeptide markers which are listed in table 7, but not in table 14 and/or 16 and/or 17.
Preferred are also polypeptide markers which are listed in table 8, but not in table 14 and/or 16.
Preferred are also polypeptide markers which are listed in table 9, but not in table 14 and/or 16 and/or 20 and/or 21.
Preferred are also polypeptide markers which are listed in table 10, but not in table 14 and/or 16.
Preferred are also polypeptide markers which are listed in table 11, but not in table 14 and/or 16.

Renal disease according to the present invention relates to any kind of renal disease or kidney dysfunction known to the person skilled in the art, for example IgA-nephropathy, MGN (membranous glomerulonephritis), MCD (minimal-change disease), FSGS (focal-segmental glomerulosclerosis), or diabetic nephropathy. Particularly, renal disease relates to a glomerulopathy such as IgA-nephropathy, MGN, MCD, or FSGS. Even more particularly renal disease relates to IgA-nephropathy, MCD, or FSGS. Most particularly, renal disease relates to IgA-nephropathy

The glomerulopathies are a subgroup of renal diseases. Glomerulopathies comprise a several diseases of different etiology. Glomerulopathies are characterized by pathomorphological changes in malpighian corpuscles, glomerulus, and Bowman's capsule. As a consequence of these changes, further pathomorphological changes may appear in other parts of the nephron and interstice.

IgA-nephropathy is also known as Berger-Nephritis. IgA-nephropathy is the most common glomerulopathy. It may be a specific, kidney-limited, form of purpura Schoenlein-Henoch (also known as anaphylactoid purpura) with increased plasma concentration of IgA. The histopathology includes all forms of glomerular lesions and deposits of IgA in the mesangium. Clinically, IgA nephropathy presents as micro- and macro-hematouria. Therapy may be attempted with ACE inhibitors and omega-3 fatty acids. Progression of

the disease occurs over the course of several years and includes transition into progressive renal insufficiency.

MGN is characterized by thickening of the basal membrane and granular subepithelial IgG deposits. MGN becomes frequently manifest in the between the age of 40 and 50. It is frequently caused by medicaments, e.g. gold, D-penicillamine, or ACE inhibitors. Therapy of MGN may be attempted with glucocorticoids or cyclophosphamide. MGN is a nephrotic syndrome, a transition into progressive renal insufficiency may take several years.

MCD is also known as lipoid nephrosis. MCD is the most common cause of a nephrotic syndrome in children. The etiology of the disease is unknown. Histologically, no or only very discrete changes can be found. Therapy of MCD may include treatment with glucocorticoids, cyclosporin A, or cyclophosphamide. In children, the disease spontaneously heals in 90% of the cases, in adults in 50% of the cases. A transition into FSGS is possible.

FSGS is also known as IgM-nephropathy. FSGS is typically characterized by deposits of IgM and C3 in the mesangium. Clinically, it becomes manifest as a nephrotic syndrome. Therapy of FSGS may include treatment with glucocorticoids, cyclosporin A, or cyclophosphamide. Prognosis is poor and includes transition into progressive renal insufficiency.

Diabetic nephropathy is also known as diabetic glomerulosclerosis. Diabetic nephropathy is the most common cause for requirement of dialysis treatment.

In summary, it is evident that renal diseases include a variety of diseases which may show quite similar histology. However, etiology, treatment, and prognosis can be quite different for each disease. For example, IgA-nephropathy requires different treatment from any other glomerulopathy described above: In IgA-nephropathy, treatment with ACE inhibitors may be attempted, which would not be recommendable in the case of MGN. Therefore, fast and reliable diagnosis is of great importance for treatment.

In the context of the present invention, diagnosing or diagnosis means that, for an individual patient, the probability of having the respective disease is determined.

Diagnosis may also include confirming a preliminary diagnosis, particularly a preliminary diagnosis established by a different method.

Furthermore, in a preferred embodiment, diagnosis according to the present invention particularly relates to "differential diagnosis". The term "differential diagnosis" relates to distinguishing between two different diseases, i.e. to determining for an individual patient the probability of having a certain first disease as compared to having a certain second disease. More particularly, differential diagnosis according to the present invention relates to distinguishing between at least two renal diseases chosen from the group consisting of IgA-nephropathy, MGN, MCD, FSGS, and diabetic nephropathy.

In another embodiment, the present invention relates to a method for the differential diagnosis of a renal disease, the method comprising:

a) measuring the presence or the absence of a polypeptide marker in a urine sample, wherein the polypeptide marker is selected from the group of polypeptide markers shown in table 1 to 22, and

b) comparing the probability of the presence of this marker in a disease patient to the probability of the presence of this marker in a control patient, wherein

c1) if the probability of the presence of this marker in a disease patient is higher than the probability of the presence of this marker in a control patient, the presence of this marker is indicative for a higher probability of having the disease rather than the control condition, or

c2) if the probability of the presence of this marker in a disease patient is lower than the probability of the presence of this marker in a control patient, the absence of the marker is indicative for a higher probability of having the disease rather than the control condition.

Preferably, the individual probabilities according to step b) are as indicated in the tables.

The term "measuring" according to the present invention relates to determining the presence of a polypeptide or other substance of interest.

The decision whether a polypeptide marker is present or absent may depend on definition of a suitable threshold value. The threshold value can either be defined through the sensitivity of the method of measurement, or it can be defined at will. The threshold in the

context of the present invention is 25 fmol/µl in a sample which has been injected into a mass spectrometer according to Example 1. However, this threshold may be the same when other methods are used. This threshold coincides with the detection threshold of a typical mass spectrometer. This threshold corresponds approximately to a concentration of the polypeptide marker in the urine sample of 50-5000 pmol/l. If different thresholds are to be used (e.g. when using another detection method), the corresponding probabilities may differ, but can easily be established by the person skilled in the art.

The "disease patient" according to the present invention is suffering from a renal disease. Particularly, the disease is at least one from the group consisting of IgA-nephropathy, MGN, MCD, FSGS, and diabetic nephropathy.

The "control patient" can either be healthy or suffering from a disease different from the one the disease patient is suffering from, i.e. the control patient can either represent the healthy condition or a disease or group of diseases. Particularly, the represented disease is at least one from the group consisting of IgA-nephropathy, MGN, MCD, FSGS, and diabetic nephropathy.

Tables 1 to 14, 16, 20, 21, and 22 list the probability (also designated as "frequency") of a given polypeptide marker being present in a urine sample of a healthy control patient or a control patient suffering from a certain disease. The discrimination factor indicates the difference between the probability of presence in the disease as compared to a given control condition. The discrimination factor can easily be calculated from the respective probabilities. The higher the discrimination factor, the better is the potential of the given marker to distinguish between the disease and the control condition. An absolute value of the discrimination factor of 0.40 or higher is preferred.

The person skilled in the art is able to establish similar tables for the polypeptide markers by himself and/or to refine the data contained in the tables, e.g. based on further patient data and/or according to different thresholds for the presence of the polypeptide marker.

For diagnosis, the probability of the presence of the polypeptide marker in a disease patient is compared to the probability of the presence of this marker in a control patient, wherein the individual probabilities are as indicated in the tables. If the probability of the presence of this marker in a disease patient is higher than the probability of the presence of this marker in a control patient, then the presence of this marker in the sample is indicative that the patient from whom the sample originates has a higher probability of having the disease

rather than the control condition. If the probability of the presence of this marker in a disease patient is lower than the probability of the presence of this marker in a control patient, then the absence of this marker in the sample is indicative that the patient from whom the sample originates has a higher probability of having the disease rather than the control condition.

For example, a given marker may have a probability of 73% of being present in a control representing IgA-nephropathy but a probability of 0% of being present in a control representing the healthy condition. If this marker is present in the sample, then the individual is diagnosed as having a 73% probability of suffering from IgA-nephropathy as compared to being healthy. If this marker is not present in the sample, then the individual is diagnosed as having a 73% probability of being healthy instead of suffering from IgA-nephropathy.

Thus, diagnosis can be established according to statistical methods familiar to the person skilled in the art.

The invention can be carried out using only one of the polypeptide markers or using a plurality of the polypeptide markers. Preferably, presence of a plurality of polypeptide markers is measured. Preferably at least 3 of the markers, more preferably at least 10 of the markers, even more preferably at least 20, most preferred at least 50 of the markers according to the present invention are measured.

An advantage of the present invention is that it provides a multitude of suitable markers. Measuring a plurality of markes can increase both sensitivity and selectivity of diagnosis. Therefore, also markers which show low discrimination factors between the disease and control can be used for diagnosis if they are combined with other markers.

If a plurality of polypeptide markers is used, a "pattern" is be generated which contains the information about the presence for each marker measured. This pattern can then be compared to the pattern of probabilities of presence of the polypeptide markers in a disease or control patient. Each table represents a pattern of probabilities of finding given polypeptide markers in certain disease and control patients.

Therefore, in a preferred embodiment, the present invention relates to a method for the differential diagnosis of a renal disease, the method comprising:

a) establishing a pattern of presence or absence for a plurality of polypeptide markers in a urine sample, wherein at least one polypeptide marker is selected from the group of polypeptide markers shown in table 1 to 22, and

b) comparing the probability of finding this pattern in a disease patient to the probability of finding this pattern in a control patient, wherein

c1) if the probability of finding the pattern in a disease patient is higher than the probability of the finding the pattern in a control patient, finding this pattern is indicative for a higher probability of having the disease rather than the control condition, or

c2) if the probability of finding the pattern in a disease patient is lower than the probability of the finding the pattern in a control patient, finding this pattern is indicative for a lower probability of having the disease rather than the control condition, or

Preferably, the individual probability for the at least one polypeptide marker according to step b) is as indicated in the tables.

Comparison of the found pattern with the probability of finding the pattern in a disease or control patient can be performed according to statistical methods known in the art. Preferably, automated methods are employed, e.g. CART-analysis, random forest analysis, and support vector machines (SVM, see e.g. Xiong. M., et al. (2001). Biomarker identification by feature wrappers. Genome Research vol. 11, p. 1878-1887). Comparison can also be performed simultaneously for several different patterns and the probability of finding them.

Thus, the measured pattern is typically compared to the probability of finding the pattern in at least two different conditions. An example for diagnosis and differential diagnosis of renal diseases according to this method is shown in Fig. 3.

If necessary, the urine samples may be pre-treated before measurement of the polypeptide marker. Particularly, lipids, nucleic acids or polypeptides may be purified from the sample according to methods known in the art, including filtration, centrifugation, or extraction methods such as chloroform/phenol extraction.

Measuring the presence of a polypeptide marker can be done by any method known in the art.

Preferred methods include gas phase ion spectrometry, such as laser desorption/ionization mass spectrometry, surface enhanced laser desorption/ionization time-of flight mass spectrometry (SELDI-TOF MS) and CE-MS. These spectrometry methods allow to measure the polypeptide markers without the need for ligands such as antibodies or aptamers.

Urine sample generally are highly complex, i.e. they contain numerous polypeptides. In case of high complexity, a spectrometric analysis becomes difficult. To reduce the complexity of the sample, the polypeptides contained in the sample may be separated by any suitable means, e.g. by electrophoretic separation, affinity-based separation, or separation based on ion exchange chromatography. Particular examples include gel electrophoresis, two-dimensional polyacrylamide gel electrophoresis (2D-PAGE), capillary electrophoresis, metal-affinity chromatography, immobilized metal-affinity chromatography (IMAC), affinity chromatography based on lectins, liquid chromatography, high pressure liquid chromatography (HPLC), and reversed-phase HPLC, cation exchange chromatography, and selectively binding surfaces (such as the surfaces used in SELDI-TOF, see below).

2D-PAGE is commonly used for polypeptide separation and can be combined with mass spectrometry (MS) yielding identification of individual polypeptides. Over 1000 protein spots can be discerned with 2D-PAGE. However, each single spot must be analysed separately by MS/MS for identification.

SELDI (surface enhanced laser desorption/ionization) time-of-flight mass spectrometry is currently applied in many fields of biomedical sciences.

In the SELDI system, the ProteinChip Arrays are the most important component. They are narrow metal strips carrying 8 or 16 spots in a row on the surface. Samples to be analyzed are directly applied to the spots, either as a standing drop or in volumes up to 500 µl, by using sample holders called "bioprocessors" as supporting units. They are placed onto the arrays during incubation and washing steps and removed again afterwards. The different types of arrays belong to two main series: chromatographic arrays, presenting hydrophobic, hydrophilic, cation-exchanging, anion- exchanging or immobilized metal ion

affinity-surfaces, and preactivated arrays with chemical groups to allow the covalent coupling of proteins. Preferably, a chip with cation-exchange surfaces is used. As the ProteinChip Arrays do not only support the sample but specifically interact with the biomolecules, the composition of the analyte depends on the array type used and the washing conditions applied. This explains why the SELDI-process can be defined as a further development of the traditional MALDI (matrix assisted laser desorption / ionization)-technique. In the SELDI-process, only on those polypeptides are measured that actually bind to the chip surface.

After binding of sample proteins, the energy absorbing matrix is applied to each spot. The matrix rapidly crystallizes and the analysis can start immediately.

The ProteinChip Arrays are placed into the ProteinChip Reader for analysis. The reader is a TOF (time-of-flight) mass spectrometer in which the proteins are desorbed and ionized with the help of a laser beam. As the crystallized proteins are equally distributed on the spot surface, the ionizing laser beam always hits a representative average of the molecules in the analyte, allowing quantitative calculations. After ionization, the proteins are accelerated by an electric field to fly down the flight tube, before reaching the detector. The flight time between the laser striking the array surface and the molecules reaching the detector at the end of the flight tube enables the system to accurately determine the mass of the protein species present in the sample (for more detailed information on the method see the following review: Merchant M and Weinberger SR (2000). Recent advancements in surface-enhanced laser desorption / ionization – time of flight mass spectrometry. Electrophoresis vol. 212, p. 1164-1177).

However, the most preferred method is CE-MS, in which capillary electrophoresis (CE) is coupled to mass spectrometry (MS). CE-MS has been described in detail elsewhere (see e.g. German patent application DE 100 21 737, and Kaiser, T., et al., Capillary Electrophoresis coupled mass spectrometry to establish polypeptide patterns in dialysis fluids. J Chromatogr A, vol. 1013, p. 157-171(2003)).

CE is known to the person skilled in the art. In brief, the sample is loaded onto an electrophoresis capillary and a voltage of up to 50 kV, typically up to 30 kV, is applied. Typical capillaries are fused silica capillaries, i.e. glas capillaries comprising an outer sheath as mechanical support and to improve mechanical flexibility, e.g. a sheath made of thermoplastic material. Typically, the capillary is untreated, i.e. it shows hydroxy-groups on its inside. However, the capillary may also be coated on the inside. E.g., hydrophobic coating can be used to improve discriminatory power. In addition to the voltage, also pressure may be applied, which is typically in the range of 0 to 1 psi. The pressure can also be applied or increased during the run.

To improve discriminatory power, also a stacking protocol can be applied when loading the sample: Before loading of the sample, a base is loaded, then the sample is loaded, then an acid. The principle is to capture the analyte ions between a base and an acid. If voltage is applied, the positively charges analyte ions move towards the base. There, they get negatively charged and move into the opposite direction towards the acid, where they get positively charged. This stacking repeats itself until acid and base are neutralized. Then, the separation starts from a well concentrated sample.

The sample is contained in an appropriate buffer in which polypeptides are soluble, e.g. phosphate buffer. For CE-MS coupling, it is preferred to use volatile solvents and to work under mostly salt-free conditions to avoid contamination of the MS. Examples comprise acetonitrile, isopropanol, methanol, and the like. The solvents can also be combined with water and a weak acid (e.g. 0.1% formic acid), the latter to protonate the analyte. The polypeptides in the sample are separated according to size and charge, which determine the run-time in the capillary. CE is characterized by high separating power and short time of analysis.

For subsequent MS analysis, either fractions collected from the CE can be analyzed as separate batches or, preferably, the CE system can be coupled via a suitable interface to the mass spectrometer to allow continous flow analysis. Alternatively, the flow from the CE may be used to generate continuous "separation tracks", which can be analyzed separately.

In the mass spectrometer, ions generated from the sample are analyzed according to the mass/charge (m/z) quotient. Using mass spectrometry, it is possible to routinely analyze 10 fmol (i.e. 0.1. ng of a 10 kDa polypeptide) with a precision of ± 0.01%. Experimentally, it is possible to analyze even less than 0.1 fmol.

Any type of mass spectrometer can be used. In mass spectrometers, an ion-generating device is coupled with an suitable analyzer. For example, the electrospray ionization (ESI) interfaces are most commonly used to produce ions from liquid samples, whereas MALDI is most commonly used to produce ions from individually processed samples. Different kinds of analyzers are available, e.g. ion trap analyzers or time-of-flight (TOF) analyzers. Both ESI and MALDI can be combined with essentially all types of mass spectrometers, although ESI has usually been combined with ion traps, whereas MALDI has usually been combined with TOF.

A preferred CE-MS method according to the present invention includes capillary electrophoresis coupled online via ESI to a TOF analyzer.

The CE-MS technique permits to measure the presence of several hundred polypeptide markers simultaneously in a short time in a small volume with high sensitivity. Once the presence of the polypeptide markers has been measured, a pattern of the measured polypeptide markers is generated and can be compared to a disease pattern by any of the methods described further above. However, in many cases it will be sufficient for diagnosis to measure only one or a limited number of the markers.

The polypeptide sequences can be determined according to methods well-known to the person skilled in the art (see e.g. C.S. Spahr et al. (2001). Towards defining the urinary proteome using liquid chromatography-tandem mass spectrometry. I. Profiling an unfractionated tryptic digest. Proteomics vol. 1, p. 93-107).

Depending on the type of polypeptide marker, it is possible to measure its presence or absence by further means. For example, if the polypeptide is biologically active, its presence may be determined by cellular or enzymatic assays.

Presence of a polypeptide can also be determined by use of ligands binding to the polypeptide of interest. Binding according to the present invention includes both covalent and non-covalent binding.

A ligand according to the present invention can be any peptide, polypeptide, nucleic acid, or other substance binding to the polypeptide of interest. It is well known that polypeptides, if obtained or purified from the human or animal body, can be modified, e.g. by glycosylation. A suitable ligand according to the present invention may bind the polypeptide also via such sites.

Preferred ligands include antibodies, nucleic acids, peptides or polypeptides, and aptamers, e.g. nucleic acid or peptide aptamers. For many polypeptides, suitable ligands are commercially available. Furthermore, methods to generate suitable ligands are well-known in the art. For example, identification and production of suitable antibodies or aptamers is also offered by commercial suppliers.

The term "antibody" as used herein includes both polyclonal and monoclonal antibodies, as well as fragments thereof, such as Fv, Fab and $F(ab)_2$ fragments that are capable of binding antigen or hapten.

Preferably, the ligand should bind specifically to the polypeptide to be measured. "Specific binding" according to the present invention means that the ligand should not bind substantially to ("cross-react" with) another polypeptide or substance present in the sample investigated. Preferably, the specifically bound protein or isoform should be bound with at least 3 times higher, more preferably at least 10 times higher and even more preferably at least 50 times higher affinity than any other relevant polypeptide.

Non-specific binding may be tolerable, particularly if the investigated peptide or polypeptide can still be distinguished and measured unequivocally, e.g. according to its size on a Western Blot, or by its relatively higher abundance in the sample.

A method for measuring the presence of a polypeptide of interest may comprise the steps of (a) contacting a polypeptide with a specifically binding ligand, (b) (optionally) removing non-bound ligand, (c) measuring the presence or amount of bound ligand.

Binding of the ligand can be measured by any method known in the art. First, binding of a ligand may be measured directly, e.g. by NMR or surface plasmon resonance. Second, the ligand also serves as a substrate of an enzymatic activity of the peptide or polypeptide of interest, an enzymatic reaction product may be measured (e.g. the presence of a protease can be measured by measuring the amount of cleaved substrate, e.g. by Western Blot). Third, the ligand may be coupled covalently or non-covalently to a label allowing detection and measurement of the ligand.

Labeling may be done by direct or indirect methods. Direct labeling involves coupling of the label directly (covalently or non-covalently) to the ligand. Indirect labeling involves binding (covalently or non-covalently) of a secondary ligand to the first ligand. The secondary ligand should specifically bind to the first ligand. Said secondary ligand may be coupled with a suitable label and/or be the target (receptor) of tertiary ligand binding to the secondary ligand. The use of secondary, tertiary or even higher order ligands is often used to increase the signal. Suitable secondary and higher order ligands may include antibodies, secondary antibodies, and the well-known streptavidin-biotin system (Vector Laboratories, Inc.)

The ligand or substrate may also be "tagged" with one or more tags as known in the art. Such tags may then be targets for higher order ligands. Suitable tags include biotin, digoxygenin, His-Tag, Glutathion-S-Transferase, FLAG, GFP, myc-tag, influenza A virus haemagglutinin (HA), maltose binding protein, and the like. In the case of a peptide or polypeptide, the tag is preferably at the N-terminus and/or C-terminus.

Suitable labels are any labels detectable by an appropriate detection method. Typical labels include gold particles, latex beads, acridan ester, luminol, ruthenium, enzymatically active labels, radioactive labels, magnetic labels ("e.g. magnetic beads", including paramagnetic and superparamagnetic labels), and fluorescent labels.

Enzymatically active labels include e.g. horseradish peroxidase, alkaline phosphatase, beta-Galactosidase, Luciferase, and derivatives thereof. Suitable substrates for detection

include di-amino-benzidine (DAB), 3,3'-5,5'-tetramethylbenzidine, NBT-BCIP (4-nitro blue tetrazolium chloride and 5-bromo-4-chloro-3-indolyl-phosphate, available as ready-made stock solution from Roche Diagnostics), CDP-Star™ (Amersham Biosciences), ECF™ (Amersham Biosciences). A suitable enzyme-substrate combination may result in a colored reaction product, fluorescence or chemoluminescence, which can be measured according to methods known in the art.

Typical fluorescent labels include fluorescent proteins (such as GFP and its derivatives), Cy3, Cy5, Texas Red, Fluorescein, the Alexa dyes (e.g. Alexa 568), and quantum dots.

Typical radioactive labels include ^{35}S, ^{125}I, ^{32}P, ^{33}P, and the like.

Thus, suitable measurement methods according the present invention also include precipitation (particularly immunoprecipitation), electrochemiluminescence (electro-generated chemiluminescence), RIA (radioimmunoassay), ELISA (enzyme-linked immunosorbent assay), sandwich enzyme immune tests, electrochemiluminescence sandwich immunoassays (ECLIA), dissociation-enhanced lanthanide fluoro immuno assay (DELFIA), scintillation proximity assay (SPA), turbidimetry, nephelometry, latex-enhanced turbidimetry or nephelometry, or solid phase immune tests. Further methods known in the art (such as gel electrophoresis, 2D gel electrophoresis, SDS polyacrylamide gel electrophoresis (SDS-PAGE), Western Blotting), can be used alone or in combination with labeling or other detection methods as described above.

The ligand may also be present on an array. Said array contains at least one additional ligand, which may be directed against a peptide, polypeptide or a nucleic acid of interest. Said additional ligand may also be directed against a peptide, polypeptide or a nucleic acid of no particular interest in the context of the present invention. Preferably, ligands for at least five, more preferably at least 10, even more preferably at least 20 polypeptide markers according to the present invention are contained on the array.

According to the present invention, the term "array" refers to a solid-phase or gel-like carrier upon which at least two compounds are attached or bound in one-, two- or three-

dimensional arrangement. Such arrays (including "gene chips", "protein chips", antibody arrays and the like) are generally known to the person skilled in the art and typically generated on glass microscope slides, specially coated glass slides such as polycation-, nitrocellulose- or biotin-coated slides, cover slips, and membranes such as, for example, membranes based on nitrocellulose or nylon.

The array may include a bound ligand or at least two cells expressing each at least one ligand.

It is also contemplated to use "suspension arrays" as arrays according to the present invention (Nolan JP, Sklar LA. (2002). Suspension array technology: evolution of the flat-array paradigm. Trends Biotechnol. vol. 20(1), p. 9-12). In such suspension arrays, the carrier, e.g. a microbead or microsphere, is present in suspension. The array consists of different microbeads or microspheres, possibly labeled, carrying different ligands.

The invention further relates to a method of producing arrays as defined above, wherein at least one ligand is bound to the carrier material in addition to other ligands.

Methods of producing such arrays, for example based on solid-phase chemistry and photo-labile protective groups, are generally known (US 5,744,305). Such arrays can also be brought into contact with substances or substance libraries and tested for interaction, for example for binding or change of conformation. Therefore, arrays comprising a polypeptide marker according to the present invention may be used for identifying ligands binding specifically to said peptides or polypeptides.

To determine the sequence of a polypeptide, it should be purified to the highest level achievable. However, the polypeptide does not need to be completely isolated. For example, it is enough to have the polypeptide detectable as a coomassie-stained band in a polyacrylamide gel. The corresponding gel piece can then be cut out and used for the next identification steps. After purification of the polypeptide, it can be enzymatically digested with trypsin and the molecular weights of the resulting fragments determined using any suitable method, for example mass spectrometry. Using mass spectrometry, each polypeptide displays a characteristic "fingerprint" of fragments allowing its identification

by database searches. In case that the polypeptide to be identified is not present in the database or if the researcher wants to have a closer characterization for any reasons, the polypeptide fragments can also be sequenced according to methods known in the art.

CE-MS allows particularly easy determination of the polypeptide sequences. The capillary electrophoresis elution time for each marker is listed in the tables. Thus, it is possible to collect the fraction containing the polypeptide at relatively high purity. If a single fraction contains insufficient material, fractions of more than one experiment may be pooled.

Sequences of some of the polypeptide markers are listed as SEQ ID NO: 1 to 5. Their masses as measured by CE-MS and their respective sequences are as follows:

SEQ ID NO:	mass [Da]	sequence	description
1	8765,9	FTFHADICTLSEKERQIKKQTALVEL VKHKPKATKEQLKAVMDDFAAFV EKCCKADDKETCFAEEGKKLVAAS QAALGL	fragment of human albumin, C-terminus amino acids 531-609.
2	10046,3	TYVPKEFNAETFTFHADICTLSEKER QIKKQTALVELVKHKPKATKEQLKA VMDDFAAFVEKCCKADDKETCFAE EGKKLVAASQAALGL	fragment of human albumin, C-terminus, amino acids 520 to 609.
3	950.0	GGRPSRPPQ	fragment of Salivary proline-rich protein
4	1292.5	GFRHRHPDEAA	fragment of alpha fibrinogen
5	1448.8	GLITLIGINPSLHT	fragment of olfactory receptor 8B4

Figure legends:

Fig. 1 Depiction of the information from a crude CE-MS analysis (A) as a three dimensional contour plot (left side). Here a contour plot of urine from a healthy volunteer is shown, mass per charge on the Y-axis against the retention time in min (X-axis), signal intensity colour coded. Next, the signal to noise is calculated and the noise removed, thus leaving only actual signals (B). The software calculates the actual mass (C) based on both isotopic distribution and conjugated masses. This

results in a table of up to 1500 polypeptides defined via their mass and retention time. As an example, bottom right shows 17 polypeptides found in the sample. CE-t, CE-time (migration time); int., intensity; m.p.c., mass per charge, cal. m., calculated mass.

Fig. 2 Contour plots of polypeptides (actual masses) for healthy subjects (NC) and for patients with focal-segmental glomerulosclerosis (FSGS), minimal-change disease (MCD) and membranous glomerulonephritis (MGN) are shown. The upper mass limit for each plot (i.e. the maximum value along the X-axis) is indicated on the top left of each plot. As evident, the contour plots differ significantly between the healthy subjects and the renal disease groups.

Fig. 3 Flow sheet for diagnosis and differential diagnosis of renal diseases (example). Samp., sample; MS-dat., MS-data; Disea., disease; Y, yes; N, no; n.d., no disease; d.n., diabetic nephropathy, FSGS, FSGS; MGN, MGN; MCD, MCD; IgA, IgA-nephropathy, diff., differential diagnosis

The invention is further illustrated by the following examples:

Example 1

Participants:

After local Ethics Committee approval, informed consent was obtained from all participants. We examined a group of 57 healthy individuals with normal renal function in order to establish normal urinary protein patterns with CE-MS. In addition, we studied 44 patients with biopsy-proven minimal-change disease (n = 16; MCD), membranous glomerulonephritis (n = 18; MGN), and focal-segmental glomerulosclerosis (n = 10; FSGS) (Table 1).

CE-MS Analysis:

Spot urine samples were collected from all participants in the morning after voiding the first urine. Samples were prepared as described in detail elsewhere (Wittke S, Fliser D, Haubitz M, et al: Determination of peptides and proteins in human urine with CE-MS – suitable tool for the establishment of new diagnostic markers. *J Chromatogr A* 1013:173-

181, 2003). The CE-MS analysis was established as described previously (Kaiser T, Hermann A, Kielstein JT, et al: Capillary Electrophoresis coupled mass spectrometry to establish polypeptide patterns in dialysis fluids. *J Chromatogr A* 1013: 157-171, 2003), using a Beckman Coulter PAC/E system coupled to a Mariner TOF mass spectrometer (ABI). CE capillaries were from Beckman, ID/OD 75/360 µm and 90 cm in length. The mobile phase used contained 30% methanol and 0.5% formic acid in water. The same liquid was used for the sheath flow, which was applied at 2 µl/min. Sample injection was performed with pressure: 1 psi for 20 sec. Under these conditions about 100 nl of sample could be injected. For sample stacking, the following protocol was applied: injection of 1M NH_3 for 7 sec., injection of sample, injection of 2M formic acid for 5 sec. The subsequent CE-MS run was performed at +30 kV with the sequence of the following pressures: 40 min at 0 psi, 2 min at 0.1 psi, 2 min at 0.2 psi, 2 min at 0.3 psi, 2 min at 0.4 psi, 80 min at 0.5 psi. For diagnosis of IgA-nephropathy, the following pressure sequence was used: 40 min at 0.3 psi, 2 min at 0.4 psi, 2 min at 0.6 psi, 2 min at 0.8 psi, 80 min at 1 psi. After each run, the CE capillary was rinsed for 5 min with 0.1 M NaOH, followed by 5 min with water and 5 min with running buffer.

Statistical analysis:

For discrimination between healthy subjects and different groups of patients with renal diseases we used the method of Random Forests and the corresponding S-Plus program version 6/2002 Breiman L: Random Forests. (http://oz.berkeley.edu/users/breiman/randomforest2001.pdf). In this procedure, a series of PP subsets of fixed size is selected randomly from all candidate PP. For each subset, a classification tree as described in the Classification and Regression Tree (CART) analysis is generated (Steinberg D, Colla P: CART – Classification and Regression trees. San Diego, CA, Salford Systems 1997), resulting in a classification rule. The forest prediction is the unweight plurality of class votes of the series of classification rules. Over-fitting is not generated due to large numbers of subset selections. The estimated generalisation error is unbiased due to the method of "out of bag" (oob) estimation: each tree is grown on a bootstrap sample of cases of the learning sample and the validation is estimated on the basis of those cases not selected in the bootstrap sample.

Further, discrimination between groups was also performed using support vector machines. This tool has the advantage of discriminating data in high dimensional parameter space. Its fast and stable algorithms showed good performance in the evaluation of clinical markers (Dieterle F, Muller-Hagedorn S, Liebich HM, Gauglitz G: Urinary nucleosides as potential

tumor markers evaluated by learning vector quantization. *Artif Intell Med* 28:265-279, 2003) and different areas of biological analyses like DNA arrays (Brown MP, Grundy WN, Lin D, et al: Knowledge-based analysis of microarray gene expression data by using support vector machines. *Proc Natl Acad Sci USA* 97:262-267, 2000).

Normal urinary polypeptide pattern analysed with CE-MS:

A graphical depiction (contour plot) of a typical sample is presented in Fig. 1 (raw data). In one individual sample, between 900 and 2500 PP with molecular weights from 800 up to 30.000 Dalton were detected. Under the conditions used for CE polypeptides with higher molecular weights tend to precipitate. Thus larger proteins in general cannot be detected, although some (e.g. albumin) can be visualised. A list of polypeptides present with high probability that were chosen as internal standards to assure sample comparability is shown in table 23. For analysis of protein-rich samples, such as samples from suspected IgA-nephropathy patients, higher pressure was applied and the polypeptides according to table 24 were preferred as internal standards. Repeated analyses of identical samples did not reveal any significant differences under identical conditions of the CE-MS run for an individual sample.

The subsequent electronic data manipulation for one example is summarised in Fig. 1. Each run results in the crude spectrum depicted in the upper part of Fig. 1 and is composed of single spectra (blow up Fig. 1) generated every 3 seconds. CE-MS peaks were identified in the first data analysis run (Fig. 1A). Next, the charge of each peak was ascertained utilising both isotopic distribution and conjugated peaks (Fig. 1B). As a result, conjugated peaks were summarised in one single peak and the real mass was calculated, as shown in Fig. 1C. Initially, the samples were spiked with external standards of known mass. This allowed subsequent definition of internal standards of PP present with high probability in the urine samples. Thus the CE-time could be normalized to the internal standards. By applying this technique on an average urine sample, roughly 1000 PP can be detected and described/identified by the two parameters mass and CE-migration time.

The examination of urine obtained from healthy subjects led to the establishment of peaks defined by actual mass and CE-time of the PP detected, so-called peak lists, and contour plots for each individual. The individual peak lists were deposited in an MS-Access database and the probability of each of the PP to appear in a single sample was calculated. One-hundred seventy-three PP were present in over 90% of the control samples examined. In addition, 156 PP were present in more than 75% of the samples, while additional 361 PP

were found in over 50% of samples from the healthy individuals. These 690 PP were found in more than 50% of all samples obtained from healthy subjects and were used to establish a "normal PP pattern".

Urine from patients with renal diseases analysed with CE-MS:

Data from the individual runs of 44 patients were sub-grouped in the three disease groups and analysed. The values from these databases, representing typical PP patterns, were subsequently compared. Significant homology of the protein patterns present in urine samples from each patient group was found within the groups. Typical examples of urinary PP patterns from patients with MCD, FSGS, and MGN are shown in Fig. 2. Each disease presents a typical protein contour plot, revealing more than 500 PP. Subsequently, the data from the three groups were compared with those obtained in healthy subjects. Table 16 shows 124 PP found in the urine of more than 95% of healthy subjects and reveals the differences to patients with MCD, FSGS, and MGN.

Statistical analysis for discrimination of healthy individuals and patients with renal disease using CE-MS data was applied. A list of 800 PP, present with more than 50% probability in either disease group was chosen for Random Forest analysis. The correct classification rate for the discrimination between healthy subjects and renal patients was 96.5 %, as shown in the following list:

Class	Healthy subjects (n=57)	Renal patients (n=44)	Classification error [%]
classified as healthy	56	2	3.5
classified as patients	1	42	2.3

After cross-validation a sensitivity of 81.3% and a specificity of 94.3% could be obtained. Discrimination of the disease groups was achieved in the learning sample. However, most likely due to the small number of FSGS patients, these could not be discriminated from MCD when applying cross-validation. Hence, FSGS and MCD were combined into one group. For the discrimination between healthy subjects, MCD/FSGS and MGN, four PP were selected by CART from the list to build a classification tree with five terminal nodes (table 15).The correct classification rate in the learning sample is 94.1%. After cross-

validation it reduces to 84.3% (93.8% for healthy controls, 71.4% for MCD/FSGS and 92.9% for MGN).

Alternatively, statistical analysis was performed using support vector machines on the same data; table 16 shows PP that were employed in this analysis. Using these PP, the correct classification was 98.0% after complete cross-validation. Table 17 depicts PP that were used to discriminate between MCD and MGN. Here the correct classification was 94.1 % after complete cross-validation. Further, it was possible to separate patients with MCD and FSGS and patients with MGN and FSGS with (cross-validated) classification rates of 92.3 % and 89.3%, respectively (tables 18 and 19). These results can be valued as a first approach using support vector machines to classify a limited number of patients. With increasing patients data the classification will further improve and become more stable. The results also indicate that for stable classification the number of applicable variables (polypeptides) depends on the number of cases (patients), hence an increase in patients will allow to use even more PP for classification.

Table 1:

molecular weight [Da]	migration time [min]	discrimination factor	frequency [%]	
			glomerulo-nephritis	healthy
830,5	25,3	0,42	55	12
836,5	35	0,64	80	16
862,4	48,7	0,45	57	12
866,4	37,9	0,62	77	16
870,4	33,9	0,45	52	7
874,5	29,7	0,46	75	29
876,4	48,9	-0,41	57	98
882,6	36,5	-0,4	55	95
903,4	46,8	-0,42	14	55
909,4	40,3	0,67	70	3
925,4	50,7	0,63	77	14
926,5	36,1	-0,44	32	76
943,5	30	0,52	80	28
946,5	46,8	-0,44	41	84
950,5	36	0,61	66	5
952,5	32	0,52	64	12
956,4	49,5	-0,54	7	60
978,5	35	0,53	55	2
980,6	34,6	-0,45	43	88
982,5	31,9	0,53	64	10
983,5	35,1	0,41	55	14
988,6	49,9	-0,48	9	57
990,6	32	0,42	61	19
991,4	37,7	0,41	55	14
994,5	33,3	0,51	75	24
995,6	36,4	0,46	68	22
1000,5	34	-0,58	36	95
1006,4	35,7	-0,48	30	78
1008,5	34,4	-0,5	43	93
1010,6	30,6	-0,58	25	83
1013,4	39,3	-0,43	9	52
1015,6	38,2	0,47	52	5
1028,6	37,8	-0,55	43	98
1033,6	39,1	0,62	70	9
1038,6	34,4	0,43	57	14
1046,6	38,6	-0,76	20	97
1047,6	30,4	-0,44	41	84
1073,6	34,7	0,59	75	16
1075,6	29	-0,43	20	64
1102,6	32,9	0,4	52	12
1108,6	29,8	0,47	86	40

1110,4	46,9	-0,5	20	71
1121,6	42,3	-0,46	16	62
1122,5	50,2	-0,41	25	66
1135,6	42,7	-0,61	5	66
1138,6	39,3	0,55	73	17
1139,6	32,2	0,52	59	7
1141,6	38	-0,6	7	67
1157,6	28,5	0,5	80	29
1159,6	39	-0,65	16	81
1163,7	38,1	0,47	50	3
1171,6	32,8	0,64	66	2
1182,6	47,2	-0,45	20	66
1191,6	50,5	-0,41	50	91
1191,8	18,3	0,56	66	10
1198,8	29,2	0,44	75	31
1203,7	24,7	-0,52	5	57
1209,6	50,5	-0,42	48	90
1211,6	31,3	0,5	66	16
1212,7	30,6	0,57	66	9
1219,6	37,3	0,57	77	21
1220,6	30,2	0,42	75	33
1223,5	51,6	-0,6	30	90
1224,7	33,6	-0,41	59	100
1225,7	41,3	0,45	50	5
1235,6	41,4	-0,41	59	100
1237,7	41,6	-0,49	18	67
1246,7	30,5	-0,47	18	66
1256,6	53,4	0,41	55	14
1264,7	26,7	0,44	52	9
1268,6	53,7	-0,47	5	52
1269,7	39,8	0,45	64	19
1270,5	52,5	-0,51	5	55
1279,7	38,3	0,48	64	16
1280,6	51,9	-0,53	9	62
1286	30,7	0,41	84	43
1292,5	53	-0,59	27	86
1297,6	38,7	0,45	86	41
1302,7	31,8	0,54	86	33
1303,6	40,7	-0,44	11	55
1311,8	31,5	0,58	77	19
1319,9	34,8	0,4	59	19
1324,2	40,5	0,5	59	9
1325,5	35,2	0,54	80	26
1333,8	38,8	0,65	82	17
1335,7	39,2	0,57	80	22
1338,7	29,6	0,46	57	10
1338,7	47,2	0,8	82	2
1350,7	50,3	-0,48	2	50
1353,7	39,3	-0,44	45	90
1354,8	45,6	0,55	93	38
1371,7	39,9	0,6	64	3
1371,8	19,3	0,63	89	26
1389,8	19,5	0,5	84	34
1390,7	41,1	0,45	64	19

1398,9	30,5	0,59	73	14
1401,8	46,2	-0,53	9	62
1405,9	17,3	0,49	59	10
1408,9	26,8	0,42	52	10
1414,6	38,1	0,62	86	24
1415,7	33,3	0,45	50	5
1419,8	39,7	0,48	77	29
1424,9	35,4	-0,46	25	71
1442,8	33,3	0,63	84	21
1444,6	37,8	0,53	82	29
1448,8	30,3	0,42	75	33
1465,9	28,8	0,59	66	7
1472,1	31,2	0,57	70	14
1474,9	16,9	0,65	77	12
1482	30,4	0,57	84	28
1484	30,4	0,58	89	31
1486,5	30,6	0,45	66	21
1498,7	34,9	0,52	66	14
1499,9	30,6	0,56	91	34
1502,8	28,8	0,44	75	31
1502,9	16,8	0,66	68	2
1508,9	16,8	0,48	57	9
1511,7	38,4	0,55	80	24
1518	26,8	0,45	93	48
1520,7	27,9	0,45	64	19
1527,9	34,7	0,43	73	29
1529,7	54,1	-0,48	36	84
1535	28,3	0,61	73	12
1537,9	31,5	0,43	70	28
1540,7	29,8	0,5	66	16
1542,5	27,2	0,4	52	12
1548,3	31,1	0,46	89	43
1556,8	33,7	0,59	89	29
1567	31,9	0,45	86	41
1567,6	53,9	-0,53	2	55
1568,6	34,3	0,41	70	29
1573,8	40,4	0,44	89	45
1574,8	33,9	0,41	61	21
1582,9	27,8	0,51	61	10
1588,4	47,9	-0,61	11	72
1596,9	34	0,64	86	22
1604,3	21,6	0,5	64	14
1604,7	38,1	0,42	73	31
1605,7	53,3	-0,4	34	74
1611,7	53,2	-0,48	36	84
1612,8	36,8	0,58	91	33
1622	19,2	0,51	91	40
1633,8	24,6	0,42	75	33
1644	18,8	0,46	55	9
1652,3	28,6	0,44	84	40
1669,9	33,4	0,54	75	21
1676	25,3	0,52	64	12
1681,6	40	0,51	82	31
1686,8	38,2	0,67	91	24

1690,8	25,5	0,44	75	31
1692,5	44,2	-0,41	39	79
1699,1	41,9	0,62	86	24
1711	43,3	-0,45	20	66
1718,5	22,6	0,57	66	9
1726	36,3	0,62	70	9
1729,2	26	0,57	70	14
1732	51,6	-0,41	36	78
1739,8	35,7	0,45	86	41
1746,2	46,2	-0,59	25	84
1747,7	50,8	-0,52	5	57
1752,9	39,9	0,46	68	22
1763	24,4	0,57	80	22
1770,4	45,4	0,4	89	48
1777,6	28,6	0,53	70	17
1793,6	28,3	0,49	55	5
1804,7	34	0,45	100	55
1808,1	45,6	0,49	55	5
1810,1	31,8	0,45	64	19
1811,3	31,3	0,55	93	38
1813,4	54,7	-0,47	7	53
1815,2	27,7	0,5	66	16
1819,9	24,1	0,5	64	14
1820,1	31,8	0,48	91	43
1821,2	18,2	0,52	57	5
1822,9	40,7	-0,6	23	83
1824,3	37	-0,52	27	79
1826,1	21,8	0,45	59	14
1831,9	41,5	0,5	59	9
1847,8	57	-0,66	27	93
1851,2	31,6	0,48	86	38
1853	31,2	0,59	82	22
1853,6	46,7	0,47	50	3
1854,2	28,8	0,45	52	7
1856,8	56,3	-0,51	18	69
1857,1	39	0,46	75	29
1864,6	28,6	0,65	82	17
1867	31,8	0,6	91	31
1883	29,1	-0,49	32	81
1885,7	57,5	-0,4	55	95
1889,2	30,1	0,41	55	14
1889,8	46,4	-0,58	39	97
1891,6	32,3	0,55	77	22
1894,9	22	0,67	86	19
1896,8	53,3	-0,44	11	55
1898,7	26,5	0,45	59	14
1904	27,5	0,48	50	2
1913,4	30,1	0,44	55	10
1916,8	44,7	-0,52	14	66
1920,7	30,6	0,46	91	45
1933,9	32,8	-0,55	36	91
1934,2	16,1	0,71	73	2
1936,5	46,6	-0,42	25	67
1936,7	32,8	0,5	80	29

1944,2	47	-0,68	18	86
1951,1	53	-0,48	16	64
1966,3	25,1	0,71	82	10
1973,7	57,1	-0,5	9	59
1977,4	42,9	-0,47	41	88
1982,9	32,2	0,58	82	24
1989,3	43,7	0,69	84	16
1990,8	47,3	-0,71	11	83
2011,5	42	0,41	70	29
2022,6	34,6	0,46	68	22
2025	24,2	0,41	50	9
2028,4	29,9	0,55	80	24
2030,4	31,7	-0,49	32	81
2030,8	46,5	-0,61	25	86
2033,5	27,5	0,46	55	9
2042	26,4	0,47	95	48
2042,5	40,7	0,5	70	21
2045,9	25,3	0,43	84	41
2047	45,4	-0,46	52	98
2050,8	38,2	0,47	82	34
2065,3	20,9	0,49	52	3
2092	26,7	0,52	66	14
2092,5	41,3	0,59	75	16
2099,2	36,9	0,58	84	26
2103,6	26,7	0,46	89	43
2105,4	32,5	0,52	66	14
2109,3	27,9	0,47	68	21
2117,1	57,1	-0,57	20	78
2127,2	39,6	0,46	75	29
2129,5	35,1	-0,58	39	97
2140,1	26,8	0,52	64	12
2144,3	22	0,47	75	28
2146,3	25,8	0,74	77	3
2147,2	38,4	0,43	64	21
2152,7	29,5	0,51	70	19
2157,2	24,4	0,41	50	9
2174,4	24,6	0,46	68	22
2178,5	21,4	0,48	57	9
2182,5	27,6	0,55	80	24
2207,2	41,9	0,41	61	21
2210,7	25,7	0,64	86	22
2217,7	41,9	0,5	84	34
2221,1	40,7	-0,5	14	64
2223,5	22,6	0,6	68	9
2228,1	25,9	0,51	91	40
2233	31,1	-0,4	55	95
2241,1	22,7	0,49	80	31
2290,7	36,2	0,47	52	5
2291,1	21,9	0,45	50	5
2308,9	26,2	0,41	61	21
2312,5	22,9	0,45	57	12
2322,5	47,1	0,47	52	5
2356,3	24	0,41	57	16
2364,4	38,9	0,49	73	24

2370,7	27,3	0,4	52	12
2391,2	24,3	0,58	70	12
2406,4	31,8	0,43	84	41
2409,1	41,9	0,43	84	41
2421	28,7	0,41	70	29
2423,1	27,4	0,41	68	28
2426,5	38,5	0,58	89	31
2427,4	24	0,53	84	31
2432,2	38,3	0,66	80	14
2464	47,2	-0,55	7	62
2465	22,8	0,7	75	5
2473,4	41,9	0,44	52	9
2490,7	26,7	0,43	70	28
2493,6	24,6	0,63	77	14
2522,9	24,4	0,47	68	21
2529,2	41,4	-0,47	14	60
2535	37,7	0,42	82	40
2540,5	25,5	0,65	75	10
2548,2	35,1	-0,46	36	83
2566,4	22,2	0,5	57	7
2568,9	26,9	0,41	70	29
2573,7	16,3	0,57	66	9
2584	43,8	-0,56	41	97
2593,4	25	0,41	57	16
2614,1	22,5	0,42	59	17
2619,7	22,9	0,47	50	3
2621,4	25,8	0,56	68	12
2644,1	32,5	-0,48	45	93
2660,8	27,1	0,4	59	19
2665,3	39,4	0,46	57	10
2677,6	23,6	0,58	68	10
2698,4	32,1	-0,47	32	79
2713,2	41,3	-0,51	11	62
2719,9	20,2	0,49	55	5
2752,8	25,3	0,56	82	26
2780,4	28,3	0,52	66	14
2790,3	26,8	0,46	61	16
2793,7	36,3	0,64	80	16
2809,1	37,2	-0,48	30	78
2812,5	32,8	0,46	61	16
2830,9	33,2	0,49	68	19
2937,1	26,6	0,46	55	9
2973,7	34,9	-0,58	30	88
2978	26,3	-0,49	34	83
2990,4	33,6	-0,47	20	67
3007,5	30,5	-0,45	23	67
3017,7	46,8	-0,42	18	60
3057,1	56,4	-0,41	43	84
3058,8	35,5	-0,41	45	86
3121,4	42,5	-0,43	57	100
3137	37	-0,42	41	83
3139,4	43,7	-0,53	25	78
3152,6	38,2	-0,45	55	100
3177,4	22,3	-0,45	27	72

3187,7	48,6	-0,4	41	81
3209,2	34,3	-0,47	50	97
3219,5	20,2	0,48	61	14
3255,8	42,9	-0,48	36	84
3262	31,5	-0,52	34	86
3281	36,8	-0,66	32	98
3282	49,4	-0,4	55	95
3290,9	36,9	-0,57	36	93
3295,8	38,4	-0,55	43	98
3303,2	38,6	-0,57	27	84
3308,6	21,3	0,53	57	3
3309,7	43,6	-0,41	32	72
3319,3	46,2	-0,45	50	95
3333,4	23,3	-0,56	32	88
3334,6	41,7	-0,54	32	86
3337,4	36,2	-0,45	43	88
3343,8	43,8	-0,46	45	91
3405,7	37,8	-0,6	39	98
3422,5	38,7	-0,58	32	90
3436	26,4	-0,5	20	71
3479,3	48,5	-0,5	50	100
3503,3	23,2	-0,43	16	59
3530,9	36,8	-0,54	27	81
3583,3	25,2	-0,67	23	90
3589,5	39,1	-0,65	25	90
3617,4	44,8	-0,4	18	59
3631,2	33,1	-0,55	16	71
3634,9	42,6	-0,41	39	79
3682,4	42,8	-0,47	27	74
3686,1	32,6	-0,71	11	83
3697,4	38,8	-0,42	11	53
3701,8	43,4	-0,63	9	72
3707	31,9	-0,69	7	76
3719,6	44,7	-0,42	41	83
3723,3	32,5	-0,65	32	97
3735,8	43,9	-0,5	30	79
3760,8	25,9	-0,51	18	69
3802,7	46,2	-0,43	14	57
3816,7	32,2	-0,52	14	66
3852,2	36,9	-0,45	36	81
3871,7	42,9	-0,41	23	64
3946,9	33,1	-0,54	39	93
3969,6	31,3	-0,52	34	86
3987	30,5	-0,42	55	97
4026,2	30,5	-0,42	11	53
4044,7	31,2	-0,57	30	86
4055,2	24,1	0,41	57	16
4154,2	23,7	0,63	77	14
4170,6	46,1	-0,45	7	52
4183,7	26,6	0,42	52	10
4241,2	24,4	0,75	89	14
4283,1	24,3	0,55	64	9
4290,8	41,1	-0,45	43	88
4654,8	38,8	-0,4	11	52

4713,7	26,9	0,63	68	5
4748,5	25,4	-0,56	39	95
4772,1	28,9	-0,43	9	52
4801,2	37,5	-0,49	39	88
4827,1	27,3	0,51	52	2
4863,7	39,2	-0,53	18	71
5213,8	36,8	-0,43	7	50
5229,1	39,9	-0,43	9	52
5575,8	35,7	-0,48	14	62
6171,5	39,6	-0,5	43	93
6212,4	30,6	-0,5	9	59
6400,9	23,4	0,51	52	2
7409,9	26,2	0,49	61	12
7556,6	26,2	0,59	75	16
7572,8	25,7	0,42	55	12
8054,8	16,7	0,63	82	19
8341,2	16,6	0,59	66	7
8653,1	17,2	0,4	52	12
8765,9	17,6	0,56	89	33
9060,7	23	0,46	57	10
9076	23	0,58	68	10
9182	17,1	0,55	64	9
9223,1	22,8	0,64	70	7
9335,5	17,5	0,47	50	3
9868,8	29,5	-0,57	20	78
9933,5	18,4	0,47	50	3
10046,3	18,1	0,7	89	19
10390,1	20,2	0,58	70	12
10518,8	20,9	0,56	75	19

Table 2

molecular weight [Da]	migration time [min]	discrimination factor	frequency [%]	
			FSGS	healthy
803,4	35,2	0,44	60	16
807,3	35,4	0,41	50	9
809,3	25,2	0,5	50	0
817,2	25,6	0,47	50	3
830,5	25,3	0,68	80	12
836,5	35	0,54	70	16
866,4	37,9	0,64	80	16
870,4	33,9	0,43	50	7
874,4	49,7	0,57	60	3
907,4	27,5	0,57	60	3
909,4	40,3	0,57	60	3
915,4	35	0,48	50	2
925,4	50,7	0,56	70	14
926,5	36,1	-0,46	30	76
939,6	33,1	0,61	70	9
943,5	30	0,42	70	28
946,5	46,8	-0,54	30	84
950,5	36	0,75	80	5
952,5	32	0,58	70	12

953,6	34,5	-0,52	20	72
956,4	49,5	-0,5	10	60
958,5	32,5	0,48	70	22
974,5	27,6	0,41	60	19
978,5	35	0,48	50	2
980,6	34,6	-0,58	30	88
981,6	37,4	-0,47	50	97
982,5	31,9	0,7	80	10
988,6	49,9	-0,47	10	57
990,6	32	0,41	60	19
994,5	33,3	0,46	70	24
995,6	36,4	0,48	70	22
1000,5	34	-0,55	40	95
1002,6	38,5	0,53	60	7
1005,5	35	0,54	70	16
1006,4	35,7	-0,58	20	78
1008,5	34,4	-0,63	30	93
1010,6	30,6	-0,53	30	83
1010,6	50,8	-0,64	0	64
1013,4	39,3	-0,52	0	52
1015,6	38,2	0,55	60	5
1021,5	32,1	0,45	50	5
1028,6	37,8	-0,58	40	98
1033,6	39,1	0,71	80	9
1041,5	51,6	-0,4	20	60
1046,6	38,6	-0,77	20	97
1049,5	39,9	0,45	50	5
1055,6	36,4	0,51	60	9
1071,5	38,7	0,48	50	2
1073,6	34,7	0,64	80	16
1075,6	29	-0,44	20	64
1090,5	36,2	0,42	70	28
1106,6	19,6	0,41	50	9
1108,6	29,8	0,4	80	40
1110,4	46,9	-0,41	30	71
1121,6	42,3	-0,62	0	62
1126,5	42,7	0,51	60	9
1135,6	42,7	-0,66	0	66
1138,6	39,3	0,63	80	17
1139,6	32,2	0,53	60	7
1141,6	38	-0,57	10	67
1157,6	28,5	0,51	80	29
1157,6	16,9	0,5	50	0
1159,6	39	-0,61	20	81
1164,6	38,7	0,46	60	14
1171,6	32,8	0,48	50	2
1182,6	47,2	-0,66	0	66
1186,6	32	0,47	80	33
1191,6	50,5	-0,71	20	91
1191,8	18,3	0,5	60	10
1192,6	40,9	0,46	70	24
1203,7	24,7	-0,47	10	57
1209,6	50,5	-0,5	40	90
1211,6	31,3	0,64	80	16

1212,7	30,6	0,61	70	9
1219,6	37,3	0,59	80	21
1220,6	30,2	0,57	90	33
1224,7	33,6	-0,5	50	100
1225,7	41,3	0,55	60	5
1235,6	41,4	-0,4	60	100
1236,7	34,8	0,44	70	26
1237,7	41,6	-0,47	20	67
1246,7	30,5	-0,46	20	66
1258,7	20,9	0,5	60	10
1263,7	38,6	0,65	70	5
1268,6	53,7	-0,52	0	52
1269,7	39,8	0,51	70	19
1270,5	52,5	-0,55	0	55
1270,6	25,7	0,55	60	5
1274,6	38	0,56	70	14
1279,7	38,3	0,44	60	16
1280,6	51,9	-0,52	10	62
1288,6	46,1	-0,44	40	84
1292,5	53	-0,66	20	86
1294,6	54,4	0,56	70	14
1302,7	31,8	0,57	90	33
1303,6	40,7	-0,45	10	55
1304,8	24,6	0,5	50	0
1305,9	33,4	0,6	70	10
1308,6	53,6	-0,54	30	84
1309,8	38,9	0,41	50	9
1310,7	52,5	0,5	50	0
1311,8	31,5	0,51	70	19
1325,5	35,2	0,44	70	26
1333,8	38,8	0,63	80	17
1335,7	39,2	0,68	90	22
1338,7	47,2	0,78	80	2
1338,7	29,6	0,7	80	10
1350,7	50,3	-0,5	0	50
1353,7	39,3	-0,5	40	90
1354,8	45,6	0,52	90	38
1367,7	26,2	0,41	50	9
1371,7	39,9	0,67	70	3
1371,8	19,3	0,64	90	26
1377,7	25,4	0,67	70	3
1383,7	52,7	-0,53	0	53
1386	24,4	0,41	50	9
1389,8	19,5	0,56	90	34
1390,7	41,1	0,71	90	19
1394,3	48,1	0,48	60	12
1398,9	30,5	0,56	70	14
1401,8	46,2	-0,62	0	62
1403,8	34,4	0,57	60	3
1405,9	17,3	0,6	70	10
1408,9	26,8	0,5	60	10
1411,2	45,4	0,41	70	29
1414,6	38,1	0,76	100	24
1414,7	26,6	0,48	50	2

1419,8	39,7	0,41	70	29
1424,9	35,4	-0,61	10	71
1426,8	38,7	0,41	50	9
1434,8	41,2	0,48	50	2
1442,8	33,3	0,69	90	21
1444,6	37,8	0,61	90	29
1465,9	28,8	0,53	60	7
1472,1	31,2	0,46	60	14
1474,9	16,9	0,58	70	12
1482	30,4	0,42	70	28
1484	30,4	0,49	80	31
1487,7	41,4	-0,41	50	91
1490,9	53,4	-0,41	30	71
1493,7	33,7	0,41	70	29
1498,7	34,9	0,46	60	14
1499,9	30,6	0,56	90	34
1501,1	29,5	0,45	50	5
1502,9	16,8	0,68	70	2
1508,9	16,8	0,51	60	9
1511,7	38,4	0,56	80	24
1517,6	54,8	0,48	60	12
1518	26,8	0,52	100	48
1518,9	42,5	0,48	60	12
1520,7	27,9	0,41	60	19
1527,9	34,7	0,41	70	29
1529,7	54,1	-0,44	40	84
1535	28,3	0,68	80	12
1536,6	35	0,46	60	14
1537,9	31,5	0,52	80	28
1540,7	29,8	0,44	60	16
1547	38,3	0,5	90	40
1548,3	31,1	0,47	90	43
1553,2	54,4	0,48	60	12
1556,8	33,7	0,51	80	29
1567	31,9	0,49	90	41
1567,6	53,9	-0,55	0	55
1588,4	47,9	-0,62	10	72
1589,7	54,3	-0,44	40	84
1591,6	32,6	0,53	60	7
1596,9	34	0,58	80	22
1598,7	28,5	0,48	60	12
1604,3	21,6	0,56	70	14
1604,7	38,1	0,49	80	31
1605,7	53,3	-0,44	30	74
1607,7	41	0,48	60	12
1611,7	53,2	-0,54	30	84
1612,8	36,8	0,47	80	33
1619,7	53,9	-0,52	10	62
1622	19,2	0,6	100	40
1644	18,8	0,41	50	9
1669,9	33,4	0,59	80	21
1671,3	25,4	0,42	80	38
1681,6	40	0,59	90	31
1686,8	38,2	0,66	90	24

1692,5	44,2	-0,59	20	79
1695,1	23,6	0,48	60	12
1699,1	41,9	0,46	70	24
1711	43,3	-0,46	20	66
1713,4	24,6	0,41	60	19
1718,5	22,6	0,41	50	9
1723,3	26,5	0,43	60	17
1726	36,3	0,61	70	9
1729,1	37,7	0,41	50	9
1729,2	26	0,66	80	14
1732	51,6	-0,48	30	78
1739,8	35,7	0,49	90	41
1746,2	46,2	-0,54	30	84
1747,7	50,8	-0,47	10	57
1751,4	40,8	0,42	80	38
1752,9	39,9	0,48	70	22
1763	24,4	0,68	90	22
1770,4	45,4	0,52	100	48
1772,6	28,5	0,44	60	16
1776,1	43,6	-0,43	40	83
1777,6	28,6	0,73	90	17
1783,4	29,2	0,51	70	19
1784,9	31,4	0,47	50	3
1786,9	35,9	0,53	70	17
1788,6	31	0,49	90	41
1792,4	42,3	0,59	80	21
1794,9	40,4	0,4	80	40
1804,7	34	0,45	100	55
1808,1	45,6	0,55	60	5
1810,1	31,8	0,41	60	19
1811,3	31,3	0,52	90	38
1813,4	54,7	-0,43	10	53
1815,1	39,3	0,41	60	19
1815,2	27,7	0,44	60	16
1820,1	31,8	0,57	100	43
1821,2	18,2	0,45	50	5
1821,5	42,1	0,46	70	24
1822,9	40,7	-0,73	10	83
1824,3	37	-0,59	20	79
1831,7	25,2	0,43	50	7
1831,9	41,5	0,61	70	9
1844,2	34,6	0,62	90	28
1847,8	57	-0,73	20	93
1849,6	37,2	-0,41	40	81
1851,2	31,6	0,52	90	38
1853	31,2	0,58	80	22
1853,4	18,3	0,5	50	0
1853,6	46,7	0,57	60	3
1854,2	28,8	0,43	50	7
1856,8	56,3	-0,59	10	69
1857,1	39	0,51	80	29
1864,6	28,6	0,43	60	17
1867	31,8	0,49	80	31
1870,5	16,1	0,45	50	5

1871,7	43,2	0,65	70	5
1881,4	34,3	0,47	80	33
1889,2	30,1	0,56	70	14
1889,8	46,4	-0,67	30	97
1891,6	32,3	0,58	80	22
1894,9	22	0,71	90	19
1894,9	56	-0,46	20	66
1898,7	26,5	0,46	60	14
1900,7	30,4	-0,63	20	83
1902,8	33,1	0,53	70	17
1904	27,5	0,48	50	2
1904,3	43,2	0,41	50	9
1916,8	44,7	-0,46	20	66
1920,5	46,1	0,44	60	16
1920,7	30,6	0,45	90	45
1925,3	52,5	0,57	90	33
1928,4	33,2	0,48	70	22
1933,9	32,8	-0,41	50	91
1934,2	16,1	0,78	80	2
1936,5	46,6	-0,47	20	67
1936,7	32,8	0,51	80	29
1944,2	47	-0,66	20	86
1950,9	34,5	0,59	90	31
1951,1	53	-0,54	10	64
1966,3	25,1	0,6	70	10
1973,7	57,1	-0,49	10	59
1977,4	42,9	-0,48	40	88
1982,9	32,2	0,46	70	24
1989,3	43,7	0,54	70	16
1990,8	47,3	-0,83	0	83
1999,4	35,6	0,4	80	40
2005,3	39,6	0,49	80	31
2013,8	45,3	-0,47	20	67
2022,6	34,6	0,48	70	22
2028,4	29,9	0,66	90	24
2030,8	46,5	-0,56	30	86
2033,5	27,5	0,51	60	9
2042	26,4	0,52	100	48
2042,5	40,7	0,49	70	21
2047	45,4	-0,48	50	98
2048,2	33,1	-0,6	40	100
2057,2	36,3	0,43	100	57
2063	24,3	0,6	60	0
2065,3	20,9	0,57	60	3
2077,3	35,8	-0,42	10	52
2092,5	41,3	0,44	60	16
2099,2	36,9	0,54	80	26
2105,4	32,5	0,56	70	14
2114,9	42	0,48	70	22
2117,1	57,1	-0,68	10	78
2121	26,9	0,61	80	19
2127,2	39,6	0,61	90	29
2140,1	26,8	0,48	60	12
2144,3	22	0,42	70	28

2146,3	25,8	0,57	60	3
2147,2	38,4	0,69	90	21
2152,7	29,5	0,51	70	19
2163,4	27,6	0,6	70	10
2174,4	24,6	0,58	80	22
2178,5	21,4	0,41	50	9
2178,7	47,5	-0,4	20	60
2182,5	27,6	0,46	70	24
2200,3	47	0,5	50	0
2210,7	25,7	0,58	80	22
2212,9	46,3	-0,44	40	84
2221,1	40,7	-0,44	20	64
2223,5	22,6	0,41	50	9
2228,1	25,9	0,5	90	40
2233	31,1	-0,65	30	95
2257,2	46,6	-0,4	60	100
2258,9	33,6	0,62	100	38
2290,7	36,2	0,45	50	5
2291,1	21,9	0,45	50	5
2322,5	47,1	0,55	60	5
2334,2	41,2	0,45	50	5
2338,2	40,4	0,53	60	7
2367,7	43,2	-0,43	40	83
2375,2	36,5	0,42	80	38
2391,2	24,3	0,48	60	12
2409,1	41,9	0,49	90	41
2423,1	27,4	0,42	70	28
2426,5	38,5	0,49	80	31
2427,4	24	0,59	90	31
2429,9	39,3	-0,46	30	76
2432,2	38,3	0,76	90	14
2435	21,6	0,48	50	2
2438,3	52,6	0,46	60	14
2443,4	31,9	-0,44	40	84
2464	47,2	-0,52	10	62
2465	22,8	0,65	70	5
2473,4	41,9	0,51	60	9
2490,5	43	0,43	60	17
2490,7	26,7	0,42	70	28
2493,6	24,6	0,66	80	14
2529,2	41,4	-0,5	10	60
2536,6	24,8	0,51	60	9
2540,5	25,5	0,5	60	10
2542,6	42,1	0,41	50	9
2568,9	26,9	0,41	70	29
2570,5	57,1	-0,58	20	78
2573,7	16,3	0,51	60	9
2584	43,8	-0,67	30	97
2591,5	37,7	-0,4	10	50
2592,5	56,6	-0,42	20	62
2593,4	25	0,44	60	16
2608,6	37,6	0,43	90	47
2614,1	22,5	0,53	70	17
2619,7	22,9	0,47	50	3

2621,4	25,8	0,48	60	12
2627,4	44,8	-0,41	50	91
2630,6	41,7	0,41	60	19
2646,7	21,9	0,53	60	7
2660,8	27,1	0,51	70	19
2677,6	23,6	0,5	60	10
2679,5	35	-0,5	50	100
2687,4	41,9	-0,4	60	100
2690,3	24,8	0,45	50	5
2713,2	41,3	-0,52	10	62
2720,6	39,5	0,47	50	3
2752,8	25,3	0,44	70	26
2767,4	31,4	-0,55	40	95
2780,4	28,3	0,46	60	14
2793,7	36,3	0,64	80	16
2825,4	36,5	-0,48	50	98
2830,9	33,2	0,51	70	19
2841,6	37,1	-0,42	30	72
2854,4	43,8	-0,47	50	97
2883,6	28,9	0,63	80	17
2892,2	32,1	0,51	80	29
2898,5	42,3	-0,44	40	84
2902,9	42,1	0,48	60	12
2911,7	36,8	-0,54	30	84
2918	42,2	-0,54	20	74
2937,1	26,6	0,51	60	9
2945,1	22,6	0,41	50	9
2973,7	34,9	-0,58	30	88
2978	26,3	-0,53	30	83
2986,9	47,3	-0,41	40	81
2990,4	33,6	-0,57	10	67
3012,1	39,4	-0,4	60	100
3017,7	46,8	-0,5	10	60
3022,8	33,8	-0,57	40	97
3038,1	33,7	-0,51	30	81
3047,7	35,9	-0,49	30	79
3057,1	56,4	-0,54	30	84
3058,8	35,5	-0,46	40	86
3080,2	31,7	-0,4	20	60
3098,8	42,6	-0,4	60	100
3108,8	44,7	-0,45	50	95
3121,4	42,5	-0,5	50	100
3137	37	-0,53	30	83
3139,4	43,7	-0,58	20	78
3152,6	38,2	-0,4	60	100
3158,8	43,3	-0,51	40	91
3166,2	41,2	-0,43	50	93
3177,4	22,3	-0,52	20	72
3187,7	48,6	-0,41	40	81
3209,2	34,3	-0,67	30	97
3255,8	42,9	-0,64	20	84
3262	31,5	-0,46	40	86
3281	36,8	-0,78	20	98
3290,9	36,9	-0,53	40	93

3293,2	54,2	-0,47	50	97
3295,8	38,4	-0,48	50	98
3303,2	38,6	-0,54	30	84
3308,6	21,3	0,57	60	3
3315,1	54,1	-0,61	10	71
3319,3	46,2	-0,45	50	95
3322,8	27,3	-0,48	30	78
3333,4	23,3	-0,68	20	88
3334,6	41,7	-0,56	30	86
3337,4	36,2	-0,58	30	88
3343,8	43,8	-0,61	30	91
3360,1	44,3	-0,4	60	100
3376,2	45,2	-0,47	50	97
3402,4	33,8	-0,48	50	98
3405,7	37,8	-0,58	40	98
3421,8	21,1	0,47	50	3
3422,5	38,7	-0,6	30	90
3436	26,4	-0,61	10	71
3503,3	23,2	-0,49	10	59
3530,9	36,8	-0,51	30	81
3547,3	38,5	-0,4	10	50
3583,3	25,2	-0,9	0	90
3589,5	39,1	-0,8	10	90
3631,2	33,1	-0,51	20	71
3634,9	42,6	-0,49	30	79
3682,4	42,8	-0,44	30	74
3686,1	32,6	-0,73	10	83
3697,4	38,8	-0,43	10	53
3701,8	43,4	-0,62	10	72
3707	31,9	-0,76	0	76
3719,6	44,7	-0,43	40	83
3723,3	32,5	-0,77	20	97
3760,8	25,9	-0,49	20	69
3802,7	46,2	-0,47	10	57
3816,7	32,2	-0,56	10	66
3852,2	36,9	-0,41	40	81
3871,7	42,9	-0,54	10	64
3946,9	33,1	-0,63	30	93
3955,9	23,6	0,41	50	9
3969,6	31,3	-0,46	40	86
3987	30,5	-0,47	50	97
4026,2	30,5	-0,53	0	53
4044,7	31,2	-0,56	30	86
4055,2	24,1	0,54	70	16
4154,2	23,7	0,66	80	14
4170,6	46,1	-0,42	10	52
4241,2	24,4	0,66	80	14
4283,1	24,3	0,41	50	9
4306,5	41,4	-0,4	20	60
4335,8	27,1	0,43	50	7
4527,7	26	0,48	50	2
4594,6	20,6	0,45	50	5
4654,8	38,8	-0,42	10	52
4713,7	26,9	0,65	70	5

4748,5	25,4	-0,45	50	95
4772,1	28,9	-0,42	10	52
4863,7	39,2	-0,51	20	71
5213,8	36,8	-0,4	10	50
5229,1	39,9	-0,52	0	52
5428,4	33,5	-0,44	30	74
5575,8	35,7	-0,52	10	62
5845,8	21,8	0,5	50	0
6171,5	39,6	-0,63	30	93
6212,4	30,6	-0,49	10	59
6238,6	30,9	-0,56	20	76
7556,6	26,2	0,44	60	16
7885,4	20,9	0,45	50	5
8054,8	16,7	0,61	80	19
8341,2	16,6	0,53	60	7
8765,9	17,6	0,47	80	33
9076	23	0,5	60	10
9223,1	22,8	0,53	60	7
9465,1	23,3	0,5	50	0
9868,8	29,5	-0,68	10	78
9933,5	18,4	0,47	50	3
10046,3	18,1	0,61	80	19
10518,8	20,9	0,51	70	19

Table 3:

molecular	migration	discrimination	frequency FSGS	frequency MCD
830,5	25,3	0,49	80	31
865,4	35,5	0,43	80	38
907,4	27,5	0,41	60	19
1005,5	35	0,45	70	25
1008,5	34,4	-0,45	30	75
1015,6	38,2	0,47	60	13
1026,5	33,2	-0,4	10	50
1041,5	51,6	-0,42	20	63
1055,6	36,4	0,41	60	19
1085,6	50,8	-0,42	20	63
1088,6	37,4	0,49	80	31
1107,5	40,2	-0,45	30	75
1128,5	44,3	0,41	60	19
1138,6	22,9	-0,4	10	50
1160,6	48,8	-0,44	50	94
1191,6	50,5	-0,49	20	69
1199,6	31	-0,63	0	63
1207,7	36,6	0,41	60	19
1208,6	38,6	0,41	60	19
1211,6	31,3	0,43	80	38
1224,7	33,6	-0,44	50	94
1270,6	25,7	0,41	60	19
1274,6	50,7	-0,44	50	94

1282,7	38,4	0,43	80	38
1294,6	54,4	0,45	70	25
1304,8	24,6	0,44	50	6
1305,9	33,4	0,51	70	19
1308,6	53,6	-0,45	30	75
1377,7	25,4	0,45	70	25
1390,7	41,1	0,4	90	50
1404,9	29,4	0,43	80	38
1493,7	33,7	0,51	70	19
1518,9	42,5	0,41	60	19
1581	37,8	-0,44	50	94
1594,8	54,8	-0,4	60	100
1607,7	41	0,41	60	19
1650,7	25,4	-0,46	10	56
1695,7	54,7	-0,4	10	50
1766,6	44,9	-0,41	40	81
1826,9	50,8	-0,59	10	69
1880,3	57,4	-0,42	20	63
1887,8	33,8	0,4	90	50
1900,7	30,4	-0,61	20	81
1925,3	52,5	0,59	90	31
1950,9	34,5	0,59	90	31
1992,9	48,5	0,44	50	6
2005,3	39,6	0,43	80	38
2011,5	42	-0,64	30	94
2048,2	33,1	-0,41	40	81
2063	24,3	0,41	60	19
2077,3	35,8	-0,59	10	69
2121	26,9	0,43	80	38
2163,4	27,6	0,51	70	19
2174,4	24,6	0,43	80	38
2258,9	33,6	0,75	100	25
2412,3	42,7	-0,42	20	63
2453,2	49,7	-0,42	20	63
2487,9	38	0,41	60	19
2570,5	57,1	-0,42	20	63
2679,5	35	-0,44	50	94
2690,3	24,8	0,5	50	0
2819,4	32,2	0,44	50	6
2864,7	29,1	-0,49	20	69
2883,6	28,9	0,55	80	25
2889,2	20,2	0,41	60	19
2918	42,2	-0,68	20	88
2986,9	47,3	-0,41	40	81
3209,2	34,3	-0,51	30	81
3255,8	42,9	-0,42	20	63
3315,1	54,1	-0,4	10	50
3402,4	33,8	-0,44	50	94
3583,3	25,2	-0,5	0	50
4335,8	27,1	0,44	50	6
9182	17,1	-0,42	20	63

Table 4:

molecular weight [Da]	migration time [min]	discrimination factor	frequency [%]	
			FSGS	MGN
819,5	35,7	-0,47	20	67
909,4	40,3	-0,4	60	100
939,6	33,1	0,59	70	11
978,5	23,9	-0,4	10	50
1017,4	36,6	-0,43	40	83
1081,7	29,6	0,49	60	11
1201,5	51,6	0,43	60	17
1282,7	38,4	0,47	80	33
1284,8	55,1	-0,4	10	50
1305,9	33,4	0,42	70	28
1338,7	29,6	0,41	80	39
1341,8	33,1	-0,43	40	83
1359,5	47,4	0,53	70	17
1394,3	48,1	0,43	60	17
1403,8	34,4	0,43	60	17
1423,6	22,3	-0,47	20	67
1490,7	33,7	-0,47	20	67
1493,5	57	-0,46	10	56
1497,9	26,7	-0,48	30	78
1504,4	30,5	-0,42	30	72
1517,6	54,8	0,43	60	17
1526,4	39,4	-0,42	30	72
1529,5	32,4	0,43	60	17
1576,4	42,5	0,48	70	22
1591,7	51,2	0,44	100	56
1595,4	31	-0,48	30	78
1692,4	30,4	-0,41	20	61
1734,6	27,2	-0,48	30	78
1768,9	44,7	0,63	80	17
1790,8	38,8	0,4	90	50
1802,5	25,6	-0,42	30	72
1844,2	34,6	0,4	90	50
1883	29,1	0,44	50	6
1885,7	57,5	0,42	70	28
1900,7	30,4	-0,47	20	67
1920,5	46,1	0,49	60	11
1925,3	52,5	0,57	90	33
1933,9	32,8	0,44	50	6
1971,5	18,9	-0,47	20	67
1986,6	35,8	-0,42	30	72
2011,5	42	-0,42	30	72
2015,1	49,6	-0,42	30	72
2079,7	21,8	-0,51	10	61
2121	26,9	0,47	80	33
2129,5	35,1	0,43	60	17
2146,3	25,8	-0,4	60	100
2274	37	-0,4	10	50
2292,4	35,3	-0,48	30	78

2312,5	22,9	-0,63	20	83
2338,2	40,4	0,43	60	17
2338,6	26	-0,49	40	89
2356,3	24	-0,43	40	83
2421	28,7	-0,44	50	94
2449,3	28,3	-0,53	30	83
2451,7	35,5	-0,43	40	83
2453,6	32	-0,53	30	83
2469,3	32,5	-0,51	10	61
2471,7	23,8	-0,42	30	72
2525,5	35,6	0,68	90	22
2566,4	22,2	-0,42	30	72
2591,5	37,7	-0,4	10	50
2607	47,6	0,48	70	22
2639,6	45,2	-0,46	10	56
2665,3	39,4	-0,49	40	89
2712,9	22,6	-0,42	30	72
2758,5	40,9	0,42	70	28
2912,9	57,5	-0,4	10	50
3041,2	45	0,41	80	39
3107,2	26,4	-0,4	10	50
3182,9	34,3	0,44	50	6
3313,8	31,6	-0,48	30	78
3479,3	48,5	0,53	70	17
4827,1	27,3	-0,43	40	83
5829,7	20,8	-0,41	20	61
8216,9	16,8	-0,4	10	50
8371,2	15,8	-0,41	20	61
8466,3	18	-0,51	10	61
8518,7	15,7	-0,48	30	78
8578,4	17	-0,47	20	67
9182	17,1	-0,69	20	89

Table 5

molecular weight [Da]	migration time [min]	discrimination factor	frequency [%]	
			FSGS	MCD + MGN
939,6	33,1	0,49	70	21
1282,7	38,4	0,45	80	35
1305,9	33,4	0,46	70	24
1359,5	47,4	0,44	70	26
1493,7	33,7	0,41	70	29
1650,7	25,4	-0,4	10	50
1734,6	27,2	-0,41	30	71
1900,7	30,4	-0,54	20	74
1925,3	52,5	0,58	90	32
2011,5	42	-0,52	30	82
2121	26,9	0,45	80	35
2258,9	33,6	0,41	100	59

2312,5	22,9	-0,48	20	68
2449,3	28,3	-0,41	30	71
2525,5	35,6	0,49	90	41
2607	47,6	0,41	70	29
2690,3	24,8	0,41	50	9
2918	42,2	-0,51	20	71
9182	17,1	-0,56	20	76

Table 6:

molecular weight [Da]	Migration time [min]	discrimination factor	frequency [%]	
			MCD	control
836,5	35	0,59	75	16
862,4	48,7	0,44	56	12
866,4	37,9	0,41	56	16
866,5	23,1	0,55	56	2
870,4	33,9	0,43	50	7
876,4	48,9	-0,61	38	98
881,5	25,7	0,46	75	29
882,6	36,5	-0,64	31	95
888,6	29,9	-0,47	6	53
903,4	46,8	-0,49	6	55
914,5	34,3	0,47	50	3
925,4	50,7	0,61	75	14
943,5	30	0,47	75	28
950,5	36	0,51	56	5
956,4	49,5	-0,48	13	60
958,5	32,5	0,46	69	22
974,5	37,9	0,41	50	9
988,5	33,9	0,43	50	7
990,6	32	0,44	63	19
991,4	37,7	0,42	56	14
1000,5	34	-0,51	44	95
1006,4	35,7	-0,4	38	78
1010,6	50,8	-0,45	19	64
1010,6	30,6	-0,52	31	83
1033,6	39,1	0,41	50	9
1034,5	31,4	0,47	50	3
1046,6	38,6	-0,59	38	97
1047,6	30,4	-0,47	38	84
1085,6	50,8	0,49	63	14
1102,6	32,9	0,44	56	12
1104,6	43,3	-0,45	6	52
1108,6	29,8	0,42	81	40
1110,4	46,9	-0,46	25	71
1122,5	50,2	-0,41	25	66
1135,6	42,7	-0,53	13	66
1138,6	39,3	0,7	88	17
1138,6	22,9	0,48	50	2
1139,6	32,2	0,49	56	7
1141,6	38	-0,61	6	67
1159,6	39	-0,56	25	81
1171,6	32,8	0,8	81	2
1182,6	47,2	-0,41	25	66
1191,8	18,3	0,46	56	10

1199,6	31	0,61	63	2
1203,7	24,7	-0,51	6	57
1219,6	37,3	0,48	69	21
1223,5	51,6	-0,65	25	90
1233,7	49,6	0,45	50	5
1237,7	41,6	-0,42	25	67
1246,7	30,5	-0,41	25	66
1256,6	53,4	0,49	63	14
1264,7	26,7	0,66	75	9
1268,6	53,7	-0,45	6	52
1269,7	39,8	0,44	63	19
1270,5	52,5	-0,49	6	55
1274,6	50,7	0,49	94	45
1280,6	51,9	-0,5	13	62
1292,5	53	-0,49	38	86
1296,6	53,8	0,53	56	3
1302,7	31,8	0,55	88	33
1310,7	36,8	0,41	56	16
1311,8	31,5	0,44	63	19
1324,2	40,5	0,6	69	9
1324,5	54,3	0,45	63	17
1325,5	35,2	0,55	81	26
1333,8	38,8	0,52	69	17
1338,7	47,2	0,86	88	2
1338,7	29,6	0,52	63	10
1350,7	50,3	-0,44	6	50
1354,8	45,6	0,62	100	38
1365	22,3	0,49	63	14
1371,8	19,3	0,49	75	26
1389,8	19,5	0,41	75	34
1401,8	46,2	-0,5	13	62
1414,6	38,1	0,57	81	24
1415,7	33,3	0,51	56	5
1424,9	35,4	-0,52	19	71
1442,8	33,3	0,61	81	21
1444,6	37,8	0,46	75	29
1448,8	30,3	0,42	75	33
1472,1	31,2	0,42	56	14
1474,9	16,9	0,63	75	12
1482	30,4	0,47	75	28
1484	30,4	0,5	81	31
1486,5	30,6	0,61	81	21
1499,9	30,6	0,53	88	34
1502,8	28,8	0,5	81	31
1502,9	16,8	0,55	56	2
1508,9	16,8	0,54	63	9
1511,7	38,4	0,7	94	24
1535	28,3	0,57	69	12
1548,3	31,1	0,44	88	43
1556,8	33,7	0,52	81	29
1561,9	28,1	0,46	88	41
1567,6	53,9	-0,55	0	55
1573,8	40,4	0,43	88	45
1574,3	53,4	-0,41	13	53
1588,4	47,9	-0,6	13	72
1591,6	32,6	0,49	56	7
1596,9	34	0,53	75	22
1604,3	21,6	0,55	69	14
1611,7	53,2	-0,41	44	84

1612,8	36,8	0,55	88	33
1622	19,2	0,42	81	40
1629,6	49,6	0,47	50	3
1635,2	27,8	0,41	75	34
1644	18,8	0,41	50	9
1658,4	39	0,44	88	43
1669,9	33,4	0,48	69	21
1671,3	42,6	0,42	56	14
1676	25,3	0,44	56	12
1681,6	40	0,56	88	31
1686,8	38,2	0,63	88	24
1692,4	30,4	0,41	56	16
1699,1	41,9	0,63	88	24
1718,5	22,6	0,48	56	9
1746,2	46,2	-0,53	31	84
1747,7	50,8	-0,51	6	57
1751,4	40,8	0,43	81	38
1752,9	39,9	0,46	69	22
1766,6	44,9	0,52	81	29
1776,1	43,6	-0,45	38	83
1777,6	28,6	0,58	75	17
1804,7	34	0,45	100	55
1811,3	31,3	0,5	88	38
1813,4	54,7	-0,53	0	53
1815,2	27,7	0,41	56	16
1820,1	31,8	0,44	88	43
1821,2	18,2	0,45	50	5
1822,9	40,7	-0,52	31	83
1824,3	37	-0,42	38	79
1831,9	41,5	0,54	63	9
1847,8	57	-0,62	31	93
1851,2	31,6	0,43	81	38
1853	31,2	0,4	63	22
1854,9	53,6	-0,44	6	50
1856,8	56,3	-0,44	25	69
1864,6	28,6	0,64	81	17
1867	31,8	0,56	88	31
1889,8	46,4	-0,53	44	97
1894,9	22	0,56	75	19
1896,8	53,3	-0,43	13	55
1909,7	47,9	0,49	63	14
1913,4	30,1	0,46	56	10
1916,8	44,7	-0,59	6	66
1934,2	16,1	0,48	50	2
1944,2	47	-0,61	25	86
1951,1	53	-0,45	19	64
1955,3	48,4	0,44	63	19
1966,3	25,1	0,65	75	10
1973,7	57,1	-0,46	13	59
1982,9	32,2	0,57	81	24
1989,3	43,7	0,66	81	16
1990,8	47,3	-0,7	13	83
2011,5	42	0,64	94	29
2017,6	33,2	0,45	81	36
2030,4	31,7	-0,44	38	81
2030,8	46,5	-0,61	25	86
2047	45,4	-0,42	56	98
2050,8	38,2	0,47	81	34
2092,5	41,3	0,66	81	16

2098,3	52	0,5	69	19
2099,2	36,9	0,49	75	26
2103,6	26,7	0,44	88	43
2106,1	46,1	0,41	56	16
2117,1	57,1	-0,4	38	78
2129,5	35,1	-0,47	50	97
2130,3	18,4	0,42	56	14
2139,3	36,9	0,41	56	16
2146,3	25,8	0,59	63	3
2151,6	42,6	0,44	56	12
2157,2	24,4	0,54	63	9
2182,5	27,6	0,51	75	24
2189,1	40,9	-0,48	19	67
2207,2	41,9	0,48	69	21
2210,7	25,7	0,59	81	22
2217,7	41,9	0,53	88	34
2223,5	22,6	0,54	63	9
2228,1	25,9	0,48	88	40
2238,4	46,3	-0,41	44	84
2281,7	45,6	-0,5	31	81
2426,5	38,5	0,56	88	31
2432,2	38,3	0,55	69	14
2464	47,2	-0,5	13	62
2465	22,8	0,51	56	5
2522,9	24,4	0,42	63	21
2529,2	41,4	-0,42	19	60
2535	37,7	0,42	81	40
2540,5	25,5	0,58	69	10
2548,2	35,1	-0,45	38	83
2566,4	22,2	0,49	56	7
2593,4	25	0,41	56	16
2621,4	25,8	0,5	63	12
2644,1	32,5	-0,43	50	93
2698,4	32,1	-0,48	31	79
2713,2	41,3	-0,43	19	62
2752,8	25,3	0,49	75	26
2790,3	26,8	0,41	56	16
2793,7	36,3	0,66	81	16
2809,1	37,2	-0,46	31	78
2921,4	30,4	0,43	69	26
2933,8	39,4	-0,47	6	53
2973,7	34,9	-0,5	38	88
3007,5	30,5	-0,48	19	67
3017,7	46,8	-0,42	19	60
3139,4	43,7	-0,4	38	78
3179,2	44,3	0,41	75	34
3262	31,5	-0,49	38	86
3281	36,8	-0,48	50	98
3282	49,4	-0,45	50	95
3290,9	36,9	-0,56	38	93
3295,8	38,4	-0,48	50	98
3333,4	23,3	-0,5	38	88
3334,6	41,7	-0,42	44	86
3343,8	43,8	-0,41	50	91
3433,3	44,5	-0,42	56	98
3530,9	36,8	-0,5	31	81
3589,5	39,1	-0,58	31	90
3631,2	33,1	-0,52	19	71
3686,1	32,6	-0,7	13	83

3697,4	38,8	-0,41	13	53
3701,8	43,4	-0,54	19	72
3707	31,9	-0,63	13	76
3723,3	32,5	-0,47	50	97
3760,8	25,9	-0,5	19	69
3816,7	32,2	-0,47	19	66
4154,2	23,7	0,42	56	14
4170,6	46,1	-0,45	6	52
4241,2	24,4	0,67	81	14
4283,1	24,3	0,54	63	9
4707,5	20,5	0,42	56	14
4713,7	26,9	0,45	50	5
4748,5	25,4	-0,57	38	95
4772,1	28,9	-0,45	6	52
5213,8	36,8	-0,44	6	50
7409,9	26,2	0,44	56	12
7556,6	26,2	0,53	69	16
8054,8	16,7	0,5	69	19
8765,9	17,6	0,48	81	33
9076	23	0,58	69	10
9182	17,1	0,54	63	9
9223,1	22,8	0,56	63	7
9868,8	29,5	-0,53	25	78
10046,3	18,1	0,62	81	19
10390,1	20,2	0,5	63	12
10518,8	20,9	0,62	81	19

Table 7:

molecular weight [Da]	migration time [min]	discrimination factor	frequency [%]	
			MCD	MGN
814,5	28,8	-0,46	38	83
819,5	35,7	-0,48	19	67
856,5	28,9	0,42	75	33
863,4	28,8	-0,54	13	67
864,5	37,3	-0,48	19	67
879,6	26,9	0,72	94	22
882,6	36,5	-0,41	31	72
909,4	40,3	-0,56	44	100
928,4	49,4	0,53	81	28
934,5	33,9	-0,41	31	72
935,6	36,6	0,46	63	17
946,5	46,8	0,47	69	22
952,5	32	-0,46	38	83
1005,5	35	-0,42	25	67
1008,5	34,4	0,53	75	22
1015,6	38,2	-0,71	13	83
1017,4	36,6	-0,52	31	83
1022,5	39,1	-0,52	31	83
1028,6	37,8	0,47	69	22
1073,6	34,7	-0,56	44	100
1085,6	50,8	0,51	63	11
1108,5	50,1	0,45	56	11
1113,6	33,7	-0,48	19	67
1138,6	22,9	0,44	50	6
1147,6	49,7	0,46	63	17
1152,5	40,7	-0,65	19	83
1208,6	38,6	-0,42	19	61

1211,6	31,3	-0,46	38	83
1213,6	50	-0,43	13	56
1224,7	33,6	0,6	94	33
1225,7	41,3	-0,42	25	67
1270,6	25,7	-0,48	19	67
1277,6	50	0,46	63	17
1279,7	38,3	-0,63	31	94
1283,9	28,9	-0,48	19	67
1301,7	34	-0,43	13	56
1319,9	34,8	-0,46	38	83
1329,8	37,5	-0,47	25	72
1337,6	52	0,47	69	22
1341,8	33,1	-0,58	25	83
1350,8	26,8	-0,42	19	61
1365	22,3	0,4	63	22
1381,1	32,3	-0,63	31	94
1398,9	30,5	-0,44	50	94
1404,9	29,4	-0,63	38	100
1423,6	22,3	-0,48	19	67
1426,8	38,7	-0,42	25	67
1433	33,7	-0,53	19	72
1465,9	28,8	-0,45	44	89
1482,8	36,3	-0,46	38	83
1487,7	41,4	0,54	88	33
1490,7	33,7	-0,48	19	67
1512,8	35,9	0,41	69	28
1527,9	34,7	-0,44	50	94
1543,8	34,9	-0,52	31	83
1558,1	23,4	-0,42	19	61
1560,5	39,5	-0,46	38	83
1569,8	48,3	0,5	50	0
1574,8	33,9	-0,58	31	89
1593,8	36,7	0,41	69	28
1595,4	31	-0,4	38	78
1602,8	58,1	0,51	63	11
1605,9	23,7	-0,65	13	78
1607,7	41	-0,42	19	61
1612,8	26,3	-0,43	13	56
1623,3	41,4	-0,47	31	78
1671,3	42,6	0,45	56	11
1726	36,3	-0,51	44	94
1729,2	26	-0,45	44	89
1744,1	34,3	-0,63	38	100
1768,9	44,7	0,52	69	17
1774,6	36,5	-0,47	31	78
1786,9	35,9	-0,41	31	72
1799	28,8	-0,47	25	72
1802,5	25,6	-0,53	19	72
1826,9	50,8	0,52	69	17
1839,1	35,5	-0,44	50	94
1857,1	39	-0,44	50	94
1859,4	22,8	-0,42	25	67
1863,8	57,5	0,42	81	39
1876,2	40,1	-0,51	38	89
1878,7	49,9	0,47	75	28
1880,3	57,4	0,51	63	11
1883	29,1	0,44	50	6
1885,7	57,5	0,47	75	28
1887,8	33,8	-0,5	50	100

1898,7	26,5	-0,52	31	83
1924,2	32,9	-0,64	25	89
1933,9	32,8	0,57	63	6
1936,7	32,8	-0,44	56	100
1949,1	38,5	0,43	88	44
1950,9	34,5	-0,58	31	89
1971,5	18,9	-0,48	19	67
1977,4	42,9	0,4	63	22
1988,9	28,8	-0,4	38	78
2005,3	39,6	-0,46	38	83
2011,3	29	-0,43	13	56
2033,5	27,5	-0,53	25	78
2035,6	30,9	-0,54	13	67
2065,3	20,9	-0,47	25	72
2077,3	35,8	0,41	69	28
2109,3	27,9	-0,51	44	94
2140,1	26,8	-0,51	38	89
2152,7	29,5	-0,51	44	94
2160,4	27,9	-0,49	6	56
2163,4	27,6	-0,48	19	67
2167,3	27,8	-0,41	31	72
2174,4	24,6	-0,51	38	89
2178,5	21,4	-0,4	38	78
2189,1	40,9	-0,48	19	67
2258,9	33,6	-0,64	25	89
2274	37	-0,44	6	50
2288,8	41,4	-0,65	13	78
2291,1	21,9	-0,47	25	72
2292,4	35,3	-0,4	38	78
2308,9	26,2	-0,46	38	83
2332,4	35,4	-0,54	13	67
2341,2	26,3	-0,49	6	56
2356,3	24	-0,46	38	83
2367,7	43,2	0,58	75	17
2380	39,6	-0,51	44	94
2391,2	24,3	-0,44	50	94
2423,1	27,4	-0,45	44	89
2434,4	34,7	-0,44	6	50
2446,2	24,7	-0,42	19	61
2451,7	35,5	-0,46	38	83
2453,6	32	-0,52	31	83
2453,8	20,4	-0,49	6	56
2455,6	27,7	-0,41	31	72
2461,1	40,5	-0,47	25	72
2469,3	32,5	-0,42	19	61
2471,7	23,8	-0,41	31	72
2475,5	22,3	-0,42	19	61
2480,2	47,2	0,4	63	22
2483,8	19,6	-0,47	25	72
2493,6	24,6	-0,5	50	100
2500,3	30,4	0,53	75	22
2518,7	38,9	-0,46	38	83
2521,3	48,3	-0,49	13	61
2525,5	35,6	0,4	63	22
2527,3	40,8	-0,53	19	72
2553,7	24,7	-0,42	19	61
2573,7	16,3	-0,45	44	89
2579,5	15,2	-0,48	19	67
2608,6	57,7	0,45	56	11

2614,1	22,5	-0,47	31	78
2619,6	38,3	-0,42	19	61
2642,4	40,9	0,46	63	17
2660,8	27,1	-0,47	31	78
2665,3	39,4	-0,58	31	89
2666	23	-0,43	13	56
2677,6	23,6	-0,51	44	94
2701	34,8	0,4	63	22
2784,3	45,2	-0,59	19	78
2825,4	36,5	0,49	88	39
2830,9	33,2	-0,57	38	94
2864,7	29,1	0,52	69	17
2889,2	20,2	-0,42	19	61
2902,9	42,1	-0,42	25	67
2912,9	57,5	-0,44	6	50
2921,4	30,4	0,52	69	17
2940,5	40,4	-0,4	38	78
3041,2	45	0,42	81	39
3044,8	48,6	0,4	63	22
3082,3	43,1	0,42	75	33
3169	37,5	0,42	75	33
3205,8	28,3	0,53	75	22
3209,2	34,3	0,48	81	33
3255,8	42,9	0,4	63	22
3256,3	23,1	-0,48	19	67
3303,2	38,6	0,44	50	6
3308,6	21,3	-0,47	31	78
3313,8	31,6	-0,53	25	78
3325,5	43,5	0,44	50	6
3336,8	53,8	0,51	56	6
3405,7	37,8	0,46	63	17
3422,5	38,7	0,45	56	11
3479,3	48,5	0,58	75	17
3578,2	32,5	-0,53	19	72
3881,9	26,2	-0,42	19	61
3969,6	31,3	0,45	56	11
4183,7	26,6	-0,47	31	78
4290,8	41,1	0,4	63	22
4527,7	26	-0,53	19	72
4565,8	25,1	-0,44	6	50
4719,5	39,3	-0,44	6	50
4827,1	27,3	-0,58	25	83
5112,9	33,1	-0,5	0	50
5829,7	20,8	-0,49	13	61
6106,5	27	-0,56	0	56
7885,4	20,9	-0,49	13	61
8341,2	16,6	-0,57	38	94
8371,2	15,8	-0,49	13	61
8466,3	18	-0,42	19	61
8518,7	15,7	-0,53	25	78
8578,4	17	-0,48	19	67
9335,5	17,5	-0,41	31	72
9465,1	23,3	-0,49	13	61
9944,2	16,7	-0,48	19	67
10949,7	26,3	-0,56	0	56

Table 8:

molecular weight [Da]	migration time [min]	discrimination factor	frequency [%]	
			MCD	FSGS + MGN
863,4	28,8	-0,45	13	57
879,6	26,9	0,58	94	36
909,4	40,3	-0,42	44	86
928,4	49,4	0,46	81	36
935,6	36,6	0,41	63	21
946,5	46,8	0,44	69	25
952,5	32	-0,41	38	79
1005,5	35	-0,43	25	68
1008,5	34,4	0,5	75	25
1015,6	38,2	-0,63	13	75
1022,5	39,1	-0,44	31	75
1028,6	37,8	0,4	69	29
1073,6	34,7	-0,49	44	93
1085,6	50,8	0,48	63	14
1138,6	22,9	0,43	50	7
1147,6	49,7	0,41	63	21
1152,5	40,7	-0,53	19	71
1199,6	31	0,41	63	21
1208,6	38,6	-0,42	19	61
1211,6	31,3	-0,45	38	82
1224,7	33,6	0,54	94	39
1270,6	25,7	-0,46	19	64
1279,7	38,3	-0,51	31	82
1341,8	33,1	-0,43	25	68
1381,1	32,3	-0,54	31	86
1404,9	29,4	-0,55	38	93
1433	33,7	-0,46	19	64
1487,7	41,4	0,48	88	39
1543,8	34,9	-0,44	31	75
1560,5	39,5	-0,41	38	79
1569,8	48,3	0,43	50	7
1574,8	33,9	-0,47	31	79
1602,8	58,1	0,41	63	21
1605,9	23,7	-0,52	13	64
1607,7	41	-0,42	19	61
1623,3	41,4	-0,44	31	75
1726	36,3	-0,42	44	86
1729,2	26	-0,42	44	86
1744,1	34,3	-0,52	38	89
1786,9	35,9	-0,4	31	71
1826,9	50,8	0,54	69	14
1876,2	40,1	-0,41	38	79
1880,3	57,4	0,48	63	14
1887,8	33,8	-0,46	50	96
1898,7	26,5	-0,44	31	75
1924,2	32,9	-0,54	25	79
1933,9	32,8	0,41	63	21
1950,9	34,5	-0,58	31	89
2005,3	39,6	-0,45	38	82
2033,5	27,5	-0,46	25	71
2035,6	30,9	-0,48	13	61
2065,3	20,9	-0,43	25	68
2077,3	35,8	0,47	69	21
2140,1	26,8	-0,41	38	79
2152,7	29,5	-0,42	44	86
2163,4	27,6	-0,49	19	68

2174,4	24,6	-0,48	38	86
2189,1	40,9	-0,42	19	61
2258,9	33,6	-0,68	25	93
2288,8	41,4	-0,55	13	68
2332,4	35,4	-0,45	13	57
2367,7	43,2	0,5	75	25
2493,6	24,6	-0,43	50	93
2500,3	30,4	0,43	75	32
2527,3	40,8	-0,42	19	61
2614,1	22,5	-0,44	31	75
2660,8	27,1	-0,44	31	75
2665,3	39,4	-0,4	31	71
2784,3	45,2	-0,46	19	64
2825,4	36,5	0,45	88	43
2830,9	33,2	-0,48	38	86
2864,7	29,1	0,51	69	18
2883,6	28,9	-0,43	25	68
2889,2	20,2	-0,42	19	61
2918	42,2	0,45	88	43
2921,4	30,4	0,47	69	21
3205,8	28,3	0,43	75	32
3209,2	34,3	0,49	81	32
3255,8	42,9	0,41	63	21
3308,6	21,3	-0,4	31	71
3402,4	33,8	0,4	94	54
3578,2	32,5	-0,46	19	64
3583,3	25,2	0,43	50	7
4527,7	26	-0,46	19	64
4827,1	27,3	-0,43	25	68
7885,4	20,9	-0,45	13	57
8341,2	16,6	-0,45	38	82
9465,1	23,3	-0,45	13	57

Table 9:

molecular weight [Da]	Migration time [min]	discrimination factor	frequency [%]	
			MGN	control
803,4	35,2	0,4	56	16
814,5	28,8	0,59	83	24
815,5	30,9	-0,45	17	62
819,5	35,7	0,63	67	3
830,5	25,3	0,49	61	12
836,5	35	0,73	89	16
844,5	30,9	0,61	61	0
847,5	35,9	0,44	61	17
862,4	48,7	0,55	67	12
863,4	28,8	0,67	67	0
864,5	37,3	0,65	67	2
866,4	37,9	0,79	94	16
870,4	33,9	0,49	56	7
873,5	38,3	0,45	50	5
874,4	49,7	0,58	61	3
874,5	29,7	0,6	89	29

879,6	26,9	-0,74	22	97
881,5	25,7	0,48	78	29
903,4	46,8	-0,44	11	55
907,4	27,5	0,47	50	3
909,4	40,3	0,97	100	3
925,4	50,7	0,7	83	14
926,5	36,1	-0,48	28	76
928,4	49,4	-0,57	28	84
929,5	39,9	-0,46	33	79
934,5	33,9	0,52	72	21
937,5	41,7	0,48	100	52
943,5	30	0,61	89	28
946,5	46,8	-0,62	22	84
950,5	36	0,61	67	5
952,5	32	0,71	83	12
956,4	49,5	-0,6	0	60
968,6	30,5	0,43	56	12
978,5	35	0,7	72	2
978,5	23,9	0,48	50	2
980,6	34,6	-0,43	44	88
981,6	37,4	-0,41	56	97
982,5	31,9	0,56	67	10
983,5	35,1	0,58	72	14
986,5	30,3	0,54	61	7
987,4	45,1	0,54	56	2
988,5	33,9	0,49	56	7
988,6	49,9	-0,57	0	57
990,6	32	0,42	61	19
991,4	37,7	0,47	61	14
994,5	33,3	0,7	94	24
995,6	36,4	0,61	83	22
998,5	35,8	0,42	56	14
1000,5	34	-0,67	28	95
1005,5	35	0,51	67	16
1006,4	35,7	-0,5	28	78
1008,5	34,4	-0,71	22	93
1010,6	50,8	-0,58	6	64
1010,6	30,6	-0,66	17	83
1013,4	39,3	-0,52	0	52
1015,6	38,2	0,78	83	5
1017,4	36,6	0,63	83	21
1022,5	39,1	0,61	83	22
1028,6	37,8	-0,76	22	98
1033,6	39,1	0,75	83	9
1038,6	34,4	0,58	72	14
1046,6	38,6	-0,91	6	97
1047,6	30,4	-0,46	39	84
1049,5	39,9	0,56	61	5
1051,5	36,2	-0,47	17	64
1055,6	36,4	0,41	50	9
1058,6	21,5	0,56	56	0
1060,6	32	0,55	94	40

1071,5	38,7	0,54	56	2
1073,6	34,7	0,84	100	16
1075,6	29	-0,47	17	64
1081,7	29,6	-0,41	11	52
1090,5	36,2	0,56	83	28
1106,5	37,2	0,44	89	45
1108,6	29,8	0,55	94	40
1109,6	34,9	-0,54	11	66
1110,4	46,9	-0,6	11	71
1113,6	33,7	0,55	67	12
1114,5	37,4	-0,48	6	53
1121,6	42,3	-0,51	11	62
1122,5	50,2	-0,43	22	66
1131,7	34,9	0,41	67	26
1132,6	36,7	-0,45	44	90
1134,7	16,9	0,48	50	2
1135,6	42,7	-0,66	0	66
1136,6	31,6	0,43	56	12
1139,6	32,2	0,54	61	7
1141,6	38	-0,62	6	67
1150,6	35,8	0,46	94	48
1152,5	40,7	0,7	83	14
1157,6	28,5	0,6	89	29
1159,6	39	-0,75	6	81
1163,7	38,1	0,58	61	3
1171,6	32,8	0,59	61	2
1181,6	37,8	0,48	89	41
1186,6	32	0,51	83	33
1191,6	50,5	-0,41	50	91
1191,8	18,3	0,67	78	10
1198,8	29,2	0,63	94	31
1199,3	49,9	-0,46	6	52
1203,7	24,7	-0,57	0	57
1211,6	31,3	0,68	83	16
1212,7	30,6	0,75	83	9
1219,6	37,3	0,63	83	21
1220,6	30,2	0,51	83	33
1223,5	51,6	-0,67	22	90
1224,7	33,6	-0,67	33	100
1225,7	41,3	0,61	67	5
1226,7	41,6	0,47	50	3
1235,6	41,4	-0,5	50	100
1236,7	34,8	0,41	67	26
1237,7	41,6	-0,56	11	67
1246,7	30,5	-0,54	11	66
1254,8	52,4	-0,44	33	78
1264,6	45,9	0,52	61	9
1268,6	53,7	-0,46	6	52
1269,7	39,8	0,42	61	19
1270,5	52,5	-0,5	6	55
1270,6	25,7	0,61	67	5
1273,8	24,6	0,54	61	7

1274,6	38	0,42	56	14
1277,6	50	-0,45	17	62
1279,7	38,3	0,79	94	16
1280,6	51,9	-0,57	6	62
1283,9	28,9	0,61	67	5
1286	30,7	0,57	100	43
1292,5	53	-0,64	22	86
1297,6	38,7	0,53	94	41
1302,7	31,8	0,51	83	33
1303,6	40,7	-0,55	0	55
1308,6	53,6	-0,4	44	84
1311,8	31,5	0,75	94	19
1319,9	34,8	0,64	83	19
1321,9	41,1	-0,5	50	100
1324,2	40,5	0,52	61	9
1325,5	35,2	0,57	83	26
1331,7	35,4	-0,48	11	59
1333,8	38,8	0,77	94	17
1335,4	53,4	-0,52	0	52
1335,7	39,2	0,72	94	22
1338,7	47,2	0,76	78	2
1341,8	33,1	0,68	83	16
1350,7	50,3	-0,5	0	50
1350,8	26,8	0,46	61	16
1353,7	39,3	-0,51	39	90
1354,8	45,6	0,51	89	38
1355,7	36,3	0,4	56	16
1367,7	26,2	0,41	50	9
1370,8	33	0,41	50	9
1371,7	39,9	0,74	78	3
1371,8	19,3	0,74	100	26
1374,8	42,1	0,43	56	12
1377,7	25,4	0,52	56	3
1378,5	45,4	-0,41	56	97
1381,1	32,3	0,58	94	36
1386	24,4	0,47	56	9
1389,8	19,5	0,54	89	34
1390,7	41,1	0,42	61	19
1395,5	25,4	0,43	50	7
1397,8	36,1	0,45	56	10
1398,9	30,5	0,81	94	14
1401,8	46,2	-0,51	11	62
1404,9	29,4	0,53	100	47
1405,9	17,3	0,62	72	10
1408,9	26,8	0,56	67	10
1414,6	38,1	0,59	83	24
1415,7	33,3	0,45	50	5
1419,8	39,7	0,6	89	29
1423,6	22,3	0,61	67	5
1424,9	18,5	0,55	67	12
1426,8	38,7	0,58	67	9
1433	33,7	0,46	72	26

1439,6	25,4	0,41	50	9
1439,7	38,1	0,41	89	48
1442,8	33,3	0,63	83	21
1444,6	37,8	0,54	83	29
1448,8	30,3	0,45	78	33
1453,1	27,1	0,51	61	10
1462,7	53,6	-0,48	50	98
1465,9	28,8	0,82	89	7
1472,1	31,2	0,75	89	14
1473,6	30,3	0,41	67	26
1474,9	16,9	0,71	83	12
1482	30,4	0,72	100	28
1482,8	36,3	0,47	83	36
1484	30,4	0,69	100	31
1487,7	41,4	-0,58	33	91
1493,5	57	0,56	56	0
1497,9	26,7	0,57	78	21
1498,7	34,9	0,7	83	14
1499,9	30,6	0,6	94	34
1501,1	29,5	0,5	56	5
1502,8	28,8	0,41	72	31
1502,9	16,8	0,76	78	2
1504,4	30,5	0,41	72	31
1508,9	16,8	0,41	50	9
1510,1	39,5	0,48	89	41
1511,7	38,4	0,43	67	24
1518	26,8	0,46	94	48
1520,7	27,9	0,64	83	19
1522,5	26,4	0,42	61	19
1527,9	34,7	0,65	94	29
1529,7	54,1	-0,62	22	84
1535	28,3	0,6	72	12
1537,9	31,5	0,56	83	28
1539,4	28,7	0,49	89	40
1540,7	29,8	0,68	83	16
1542,5	27,2	0,49	61	12
1543,8	34,9	0,54	83	29
1548,3	31,1	0,46	89	43
1552,3	35,5	0,41	100	59
1556,8	33,7	0,71	100	29
1558,1	23,4	0,52	61	9
1567	31,9	0,59	100	41
1567,6	53,9	-0,5	6	55
1568,6	34,3	0,54	83	29
1573,8	40,4	0,5	94	45
1574,3	53,4	-0,42	11	53
1574,8	33,9	0,68	89	21
1576,4	42,5	-0,62	22	84
1578	52,5	-0,48	50	98
1582,9	27,8	0,67	78	10
1588,4	47,9	-0,61	11	72
1589,7	54,3	-0,46	39	84

1595,4	31	0,61	78	17
1596,9	34	0,78	100	22
1604,3	21,6	0,42	56	14
1604,7	38,1	0,41	72	31
1605,7	53,3	-0,46	28	74
1605,9	23,7	0,74	78	3
1607,7	41	0,49	61	12
1611,7	53,2	-0,51	33	84
1612,8	36,8	0,67	100	33
1612,8	26,3	0,52	56	3
1613,9	36,3	-0,46	39	84
1617,9	44,8	-0,4	44	84
1619,7	53,9	-0,45	17	62
1622	19,2	0,55	94	40
1629,5	32	0,42	61	19
1633,8	24,6	0,62	94	33
1635,2	27,8	0,43	78	34
1644	18,8	0,52	61	9
1652,3	28,6	0,6	100	40
1665	27,5	0,45	50	5
1666,3	23,3	0,48	89	41
1669,9	33,4	0,57	78	21
1676	25,3	0,66	78	12
1681,6	40	0,41	72	31
1686,8	38,2	0,7	94	24
1690,8	25,5	0,63	94	31
1692,4	30,4	0,46	61	16
1695,1	23,6	0,43	56	12
1699,1	41,9	0,7	94	24
1702,9	24,5	0,42	56	14
1706,8	21,5	0,51	67	16
1711	43,3	-0,54	11	66
1713,4	24,6	0,42	61	19
1718,5	22,6	0,75	83	9
1723,3	26,5	0,49	67	17
1726	36,3	0,86	94	9
1729,2	26	0,75	89	14
1732	51,6	-0,44	33	78
1734,6	27,2	0,47	78	31
1739,8	35,7	0,48	89	41
1744,1	34,3	0,67	100	33
1746,2	46,2	-0,68	17	84
1747,7	50,8	-0,57	0	57
1752,9	39,9	0,44	67	22
1763	24,4	0,72	94	22
1768,9	44,7	-0,63	17	79
1770,4	45,4	0,46	94	48
1772,6	28,5	0,4	56	16
1774,6	36,5	0,61	78	17
1782,1	33	0,43	94	52
1786,9	35,9	0,55	72	17
1788,6	31	0,42	83	41

1791	25	0,58	67	9
1792	25,1	0,4	56	16
1793,6	28,3	0,67	72	5
1802,5	25,6	0,53	72	19
1804,7	34	0,45	100	55
1808,1	45,6	0,56	61	5
1810,1	31,8	0,64	83	19
1811,3	31,3	0,62	100	38
1813,4	54,7	-0,42	11	53
1815,2	27,7	0,62	78	16
1819,9	24,1	0,7	83	14
1820,1	31,8	0,46	89	43
1821,2	18,2	0,61	67	5
1822,9	40,7	-0,61	22	83
1824,3	37	-0,57	22	79
1826,1	21,8	0,64	78	14
1831,9	41,5	0,41	50	9
1833,5	20,4	0,48	50	2
1837,6	38,2	0,41	72	31
1839,1	35,5	0,58	94	36
1847,8	57	-0,65	28	93
1849,6	37,2	-0,48	33	81
1851,2	31,6	0,51	89	38
1853	31,2	0,78	100	22
1853,6	46,7	0,47	50	3
1854,2	28,8	0,54	61	7
1856,8	56,3	-0,52	17	69
1857,1	39	0,65	94	29
1859,4	22,8	0,63	67	3
1860,3	25,9	0,4	56	16
1863,8	57,5	-0,54	39	93
1864,6	28,6	0,77	94	17
1867	31,8	0,69	100	31
1870,4	30,4	0,46	94	48
1870,5	16,1	0,45	50	5
1876,2	40,1	0,65	89	24
1878,7	49,9	-0,52	28	79
1878,9	30,2	0,41	72	31
1880,3	57,4	-0,44	11	55
1883	29,1	-0,75	6	81
1885,7	57,5	-0,67	28	95
1887,8	33,8	0,48	100	52
1889,8	46,4	-0,58	39	97
1891,6	32,3	0,72	94	22
1894,9	22	0,75	94	19
1894,9	56	-0,43	22	66
1896,8	53,3	-0,5	6	55
1898,7	26,5	0,7	83	14
1904	27,5	0,59	61	2
1913,4	30,1	0,51	61	10
1913,9	53,9	-0,4	39	79
1916,8	44,7	-0,49	17	66

1920,7	30,6	0,55	100	45
1924,2	32,9	0,51	89	38
1931,4	26,6	0,43	56	12
1933,9	32,8	-0,86	6	91
1934,2	16,1	0,87	89	2
1936,5	46,6	-0,45	22	67
1936,7	32,8	0,71	100	29
1944,1	32,2	0,47	83	36
1944,2	47	-0,75	11	86
1949,1	38,5	-0,42	44	86
1950,9	34,5	0,58	89	31
1951,1	53	-0,47	17	64
1966,3	25,1	0,84	94	10
1971,3	35,1	0,48	89	41
1971,5	18,9	0,61	67	5
1973,7	57,1	-0,53	6	59
1977	25,5	0,46	94	48
1977,4	42,9	-0,66	22	88
1982,9	32,2	0,65	89	24
1986,6	35,8	0,46	72	26
1988,9	28,8	0,61	78	17
1989,3	43,7	0,79	94	16
1990,8	47,3	-0,66	17	83
2005,3	39,6	0,52	83	31
2011,3	29	0,47	56	9
2011,5	42	0,43	72	29
2013,8	45,3	-0,51	17	67
2022,6	34,6	0,55	78	22
2025	24,2	0,58	67	9
2028,4	29,9	0,7	94	24
2030,4	31,7	-0,64	17	81
2030,8	46,5	-0,64	22	86
2032,1	30,6	0,42	56	14
2033,5	27,5	0,69	78	9
2035,6	30,9	0,41	67	26
2042	26,4	0,52	100	48
2042,5	40,7	0,63	83	21
2045,9	25,3	0,53	94	41
2047	45,4	-0,48	50	98
2050,8	38,2	0,54	89	34
2052,5	38,7	0,41	67	26
2057,2	36,3	0,43	100	57
2065,3	20,9	0,69	72	3
2079,7	21,8	0,54	61	7
2092	26,7	0,75	89	14
2092,5	41,3	0,62	78	16
2093,1	25,3	0,47	56	9
2095,3	33,7	0,46	89	43
2099,2	36,9	0,69	94	26
2103,6	26,7	0,51	94	43
2105,4	32,5	0,64	78	14
2109,3	27,9	0,74	94	21

2116,3	20,3	0,51	67	16
2117,1	57,1	-0,66	11	78
2121,1	43,1	0,4	61	21
2127,2	39,6	0,48	78	29
2129,5	35,1	-0,8	17	97
2140,1	26,8	0,77	89	12
2144,3	22	0,67	94	28
2146,3	25,8	0,97	100	3
2152,7	29,5	0,75	94	19
2157,2	24,4	0,41	50	9
2163,4	27,6	0,56	67	10
2167,3	27,8	0,45	72	28
2172,5	36,7	0,46	94	48
2174,4	24,6	0,66	89	22
2178,5	21,4	0,69	78	9
2182,5	27,6	0,65	89	24
2197,9	29	0,48	67	19
2200,2	33,6	0,47	78	31
2205,6	23	0,42	56	14
2207,2	41,9	0,46	67	21
2210,7	25,7	0,72	94	22
2212,9	46,3	-0,46	39	84
2217,7	41,9	0,6	94	34
2221,1	40,7	-0,64	0	64
2223,5	22,6	0,75	83	9
2228,1	25,9	0,55	94	40
2230,1	22,8	0,52	56	3
2241	41,1	0,47	78	31
2241,1	22,7	0,63	94	31
2246,6	39,1	0,41	94	53
2253,1	22,4	0,54	56	2
2258,9	33,6	0,51	89	38
2264,4	34,8	-0,44	17	60
2266	18,5	0,5	50	0
2273,5	22,4	0,42	56	14
2279,1	47,2	-0,5	44	95
2279,5	34,8	0,46	94	48
2288,8	41,4	0,45	78	33
2288,9	27	0,48	67	19
2290,7	36,2	0,61	67	5
2291,1	21,9	0,67	72	5
2302,9	36,7	0,52	100	48
2308,9	26,2	0,63	83	21
2312,5	22,9	0,71	83	12
2322,5	47,1	0,61	67	5
2325,5	19,5	0,5	50	0
2334,2	41,2	0,45	50	5
2352,4	24,7	0,42	61	19
2356,3	24	0,68	83	16
2364,4	38,9	0,65	89	24
2367,7	43,2	-0,66	17	83
2370,7	27,3	0,49	61	12

2380	39,6	0,62	94	33
2383,9	21,1	0,52	56	3
2389,2	34,4	0,41	89	48
2391,2	24,3	0,82	94	12
2396,5	34,6	0,49	61	12
2406,4	31,8	0,59	100	41
2409,1	41,9	0,48	89	41
2414,5	40,8	0,4	61	21
2421	28,7	0,65	94	29
2423,1	27,4	0,61	89	28
2426,5	38,5	0,63	94	31
2427,4	24	0,63	94	31
2429,9	39,3	-0,48	28	76
2432,2	38,3	0,7	83	14
2435	21,6	0,54	56	2
2443,4	31,9	-0,46	39	84
2446,2	24,7	0,44	61	17
2449,3	28,3	0,45	83	38
2451,7	35,5	0,44	83	40
2453,6	32	0,7	83	14
2453,8	20,4	0,56	56	0
2455,6	27,7	0,5	72	22
2461,1	40,5	0,58	72	14
2464	47,2	-0,62	0	62
2465	22,8	0,89	94	5
2469,3	32,5	0,49	61	12
2471,7	23,8	0,53	72	19
2473,4	41,9	0,52	61	9
2475,5	22,3	0,49	61	12
2480,2	47,2	-0,5	22	72
2483,8	19,6	0,55	72	17
2485,9	47,5	0,41	72	31
2490,7	26,7	0,56	83	28
2493,4	46	-0,48	33	81
2493,6	24,6	0,86	100	14
2500,3	30,4	-0,57	22	79
2502,9	33	0,46	61	16
2507,3	17,2	0,48	50	2
2518,7	38,9	0,59	83	24
2521,3	48,3	0,56	61	5
2522,9	24,4	0,63	83	21
2525,5	35,6	-0,64	22	86
2527,3	40,8	0,6	72	12
2529,2	41,4	-0,49	11	60
2535	37,7	0,55	94	40
2536,6	24,8	0,47	56	9
2540,5	25,5	0,79	89	10
2548,2	35,1	-0,55	28	83
2553,7	24,7	0,44	61	17
2561,3	21,6	0,48	50	2
2566,4	22,2	0,65	72	7
2568,9	26,9	0,48	78	29

2570,5	57,1	-0,44	33	78
2573,7	16,3	0,8	89	9
2576,2	25,4	0,46	67	21
2579,5	15,2	0,63	67	3
2584	43,8	-0,69	28	97
2592,5	56,6	-0,45	17	62
2593,4	25	0,4	56	16
2601,6	23,2	0,45	56	10
2607	47,6	-0,52	22	74
2608,6	37,6	0,42	89	47
2614,1	22,5	0,61	78	17
2619,6	38,3	0,4	61	21
2619,7	22,9	0,63	67	3
2621,4	25,8	0,66	78	12
2627,4	44,8	-0,41	50	91
2630,6	41,7	0,53	72	19
2639,6	45,2	0,5	56	5
2642,4	40,9	-0,61	17	78
2644,1	32,5	-0,6	33	93
2646,7	21,9	0,43	50	7
2654,3	37,2	-0,54	11	66
2658,5	24,7	-0,42	44	86
2660,8	27,1	0,59	78	19
2665,3	39,4	0,79	89	10
2666	23	0,45	56	10
2677,6	23,6	0,84	94	10
2687,9	28,2	0,5	56	5
2697,3	42,4	-0,48	6	53
2698,4	32,1	-0,52	28	79
2706,7	18,5	0,48	50	2
2707,2	34,1	0,45	72	28
2712,9	22,6	0,52	72	21
2713,2	41,3	-0,57	6	62
2719,9	20,2	0,67	72	5
2733,4	34,6	-0,42	44	86
2752,8	25,3	0,69	94	26
2758,5	40,9	-0,43	28	71
2775,5	26,3	0,43	50	7
2780,4	28,3	0,7	83	14
2784,3	45,2	0,61	78	17
2790,3	26,8	0,62	78	16
2793,7	36,3	0,62	78	16
2809,1	37,2	-0,61	17	78
2812,5	32,8	0,62	78	16
2823,3	39,9	-0,43	44	88
2825,4	36,5	-0,59	39	98
2830,9	33,2	0,75	94	19
2834,1	38,2	-0,45	28	72
2841,6	37,1	-0,45	28	72
2848,8	36,3	-0,5	44	95
2854,4	43,8	-0,41	56	97
2875,1	59,1	0,5	50	0

2883,6	28,9	0,44	61	17
2898,5	42,3	-0,4	44	84
2902,9	42,1	0,55	67	12
2908,2	49,2	-0,42	22	64
2912,9	57,5	0,5	50	0
2937,1	26,6	0,52	61	9
2945,1	22,6	0,52	61	9
2972,2	25,6	0,48	67	19
2973,7	34,9	-0,66	22	88
2978	26,3	-0,66	17	83
2978,3	41,7	-0,52	28	79
2990,4	33,6	-0,51	17	67
3007,5	30,5	-0,45	22	67
3041,2	45	-0,61	39	100
3044,8	48,6	-0,52	22	74
3057,1	56,4	-0,51	33	84
3058,8	35,5	-0,42	44	86
3061,9	30,4	0,54	89	34
3077	28,4	-0,55	28	83
3082,3	43,1	-0,46	33	79
3098,8	42,6	-0,44	56	100
3114,9	44,5	-0,53	33	86
3121,4	42,5	-0,56	44	100
3133,8	43,9	-0,45	17	62
3139,4	43,7	-0,61	17	78
3149,7	41,6	-0,48	50	98
3152,6	38,2	-0,61	39	100
3169	37,5	-0,41	33	74
3177,4	22,3	-0,5	22	72
3187,7	48,6	-0,53	28	81
3190,9	28,8	-0,42	22	64
3193,1	35,5	-0,46	6	52
3205,8	28,3	-0,64	22	86
3209,2	34,3	-0,63	33	97
3219,5	20,2	0,7	83	14
3232,5	35,7	-0,46	11	57
3255,8	42,9	-0,62	22	84
3256,3	23,1	0,65	67	2
3258,6	36,3	-0,52	17	69
3260,9	57,3	-0,48	6	53
3262	31,5	-0,58	28	86
3281	36,8	-0,76	22	98
3290,9	36,9	-0,6	33	93
3293,2	54,2	-0,47	50	97
3295,8	38,4	-0,65	33	98
3300,3	44,5	0,41	50	9
3303,2	38,6	-0,79	6	84
3308,6	21,3	0,74	78	3
3309,7	43,6	-0,56	17	72
3319,3	46,2	-0,56	39	95
3320	26,7	0,42	56	14
3333,4	23,3	-0,55	33	88

3334,6	41,7	-0,64	22	86
3336,8	53,8	-0,58	6	64
3337,4	36,2	-0,49	39	88
3343,8	43,8	-0,41	50	91
3372,2	32,5	0,51	83	33
3381,9	43,9	-0,42	11	53
3398,9	44,5	-0,41	50	91
3402,4	33,8	-0,43	56	98
3405,7	37,8	-0,82	17	98
3422,5	38,7	-0,79	11	90
3433,3	44,5	-0,43	56	98
3436	26,4	-0,54	17	71
3442,8	42,5	-0,44	56	100
3451,5	32,6	-0,43	22	66
3479,3	48,5	-0,83	17	100
3503,3	23,2	-0,48	11	59
3530,9	36,8	-0,59	22	81
3552	38,8	-0,48	6	53
3578,2	32,5	0,58	72	14
3583,3	25,2	-0,79	11	90
3589,5	39,1	-0,62	28	90
3617,4	44,8	-0,53	6	59
3631,2	33,1	-0,6	11	71
3634,9	42,6	-0,52	28	79
3669,7	36,7	-0,47	17	64
3682,4	42,8	-0,57	17	74
3686,1	32,6	-0,72	11	83
3697,4	38,8	-0,42	11	53
3701,8	43,4	-0,72	0	72
3707	31,9	-0,7	6	76
3719,6	44,7	-0,55	28	83
3723,3	32,5	-0,74	22	97
3735,8	43,9	-0,68	11	79
3739,7	47,7	-0,44	33	78
3760,8	25,9	-0,52	17	69
3802,7	46,2	-0,51	6	57
3816,7	32,2	-0,54	11	66
3852,2	36,9	-0,59	22	81
3871,7	42,9	-0,42	22	64
3881,9	26,2	0,51	61	10
3946,9	33,1	-0,65	28	93
3955,9	23,6	0,41	50	9
3969,6	31,3	-0,75	11	86
3987	30,5	-0,52	44	97
4006,7	45,9	-0,43	22	66
4026,2	30,5	-0,48	6	53
4044,7	31,2	-0,75	11	86
4055,2	24,1	0,46	61	16
4102,1	41,9	-0,5	0	50
4154,2	23,7	0,81	94	14
4170,6	46,1	-0,46	6	52
4183,7	26,6	0,67	78	10

4241,2	24,4	0,86	100	14
4283,1	24,3	0,64	72	9
4290,8	41,1	-0,66	22	88
4364,4	23,9	-0,42	11	53
4369,9	27,1	0,53	94	41
4527,7	26	0,7	72	2
4565,8	25,1	0,41	50	9
4626,4	27,2	0,61	67	5
4654,8	38,8	-0,41	11	52
4713,7	26,9	0,78	83	5
4719,5	39,3	0,5	50	0
4748,5	25,4	-0,61	33	95
4772,1	28,9	-0,41	11	52
4801,2	37,5	-0,71	17	88
4827,1	27,3	0,82	83	2
4863,7	39,2	-0,65	6	71
5112,9	33,1	0,5	50	0
5213,8	36,8	-0,44	6	50
5229,1	39,9	-0,41	11	52
5575,8	35,7	-0,57	6	62
5829,7	20,8	0,59	61	2
5845,8	21,8	0,5	50	0
6106,5	27	0,56	56	0
6171,5	39,6	-0,54	39	93
6212,4	30,6	-0,59	0	59
6238,6	30,9	-0,43	33	76
6400,9	23,4	0,7	72	2
7190,3	26,5	0,43	50	7
7284,9	40,7	0,5	50	0
7409,9	26,2	0,6	72	12
7556,6	26,2	0,73	89	16
7572,8	25,7	0,55	67	12
7885,4	20,9	0,56	61	5
8054,8	16,7	0,75	94	19
8216,9	16,8	0,45	50	5
8341,2	16,6	0,88	94	7
8371,2	15,8	0,61	61	0
8466,3	18	0,56	61	5
8518,7	15,7	0,76	78	2
8578,4	17	0,61	67	5
8653,1	17,2	0,55	67	12
8765,9	17,6	0,67	100	33
8803,1	16,2	0,47	50	3
8928,6	17	0,49	56	7
9060,7	23	0,62	72	10
9076	23	0,62	72	10
9182	17,1	0,8	89	9
9208	22,9	0,61	67	5
9223,1	22,8	0,76	83	7
9335,5	17,5	0,69	72	3
9465,1	23,3	0,61	61	0
9480,1	23,6	0,56	56	0

9621,9	19,2	0,58	67	9
9868,8	29,5	-0,55	22	78
9933,5	18,4	0,52	56	3
9944,2	16,7	0,63	67	3
10046,3	18,1	0,81	100	19
10390,1	20,2	0,77	89	12
10518,8	20,9	0,53	72	19
10949,7	26,3	0,5	56	5

Table 10:

molecular weight [Da]	migration time [min]	discrimination factor	frequency [%]	
			MGN	FSGS + MCD
814,5	28,8	0,41	83	42
819,5	35,7	0,47	67	19
863,4	28,8	0,44	67	23
864,5	37,3	0,44	67	23
879,6	26,9	-0,59	22	81
909,4	40,3	0,5	100	50
928,4	49,4	-0,41	28	69
1015,6	38,2	0,53	83	31
1017,4	36,6	0,49	83	35
1022,5	39,1	0,41	83	42
1073,6	34,7	0,42	100	58
1152,5	40,7	0,53	83	31
1224,7	33,6	-0,44	33	77
1279,7	38,3	0,52	94	42
1283,9	28,9	0,44	67	23
1301,7	34	0,4	56	15
1319,9	34,8	0,41	83	42
1329,8	37,5	0,41	72	31
1341,8	33,1	0,53	83	31
1381,1	32,3	0,48	94	46
1404,9	29,4	0,46	100	54
1423,6	22,3	0,47	67	19
1433	33,7	0,41	72	31
1490,7	33,7	0,47	67	19
1543,8	34,9	0,41	83	42
1574,8	33,9	0,47	89	42
1595,4	31	0,43	78	35
1602,8	58,1	-0,43	11	54
1605,9	23,7	0,55	78	23
1612,8	26,3	0,4	56	15
1726	36,3	0,41	94	54
1744,1	34,3	0,5	100	50
1768,9	44,7	-0,56	17	73
1774,6	36,5	0,43	78	35
1799	28,8	0,41	72	31
1802,5	25,6	0,49	72	23
1839,1	35,5	0,41	94	54
1876,2	40,1	0,43	89	46
1883	29,1	-0,44	6	50
1885,7	57,5	-0,45	28	73
1898,7	26,5	0,41	83	42
1924,2	32,9	0,5	89	38

1933,9	32,8	-0,52	6	58
1971,5	18,9	0,47	67	19
2011,3	29	0,4	56	15
2079,7	21,8	0,42	61	19
2109,3	27,9	0,44	94	50
2140,1	26,8	0,43	89	46
2152,7	29,5	0,41	94	54
2160,4	27,9	0,4	56	15
2274	37	0,42	50	8
2288,8	41,4	0,51	78	27
2292,4	35,3	0,43	78	35
2312,5	22,9	0,45	83	38
2332,4	35,4	0,44	67	23
2338,6	26	0,43	89	46
2341,2	26,3	0,44	56	12
2356,3	24	0,45	83	38
2367,7	43,2	-0,45	17	62
2380	39,6	0,44	94	50
2391,2	24,3	0,41	94	54
2421	28,7	0,41	94	54
2451,7	35,5	0,45	83	38
2453,6	32	0,53	83	31
2453,8	20,4	0,44	56	12
2461,1	40,5	0,41	72	31
2469,3	32,5	0,46	61	15
2471,7	23,8	0,41	72	31
2500,3	30,4	-0,43	22	65
2521,3	48,3	0,42	61	19
2525,5	35,6	-0,51	22	73
2527,3	40,8	0,45	72	27
2639,6	45,2	0,4	56	15
2642,4	40,9	-0,41	17	58
2665,3	39,4	0,54	89	35
2677,6	23,6	0,44	94	50
2784,3	45,2	0,51	78	27
2830,9	33,2	0,44	94	50
2912,9	57,5	0,42	50	8
3041,2	45	-0,42	39	81
3205,8	28,3	-0,43	22	65
3256,3	23,1	0,44	67	23
3313,8	31,6	0,51	78	27
3336,8	53,8	-0,44	6	50
3479,3	48,5	-0,56	17	73
3578,2	32,5	0,41	72	31
4183,7	26,6	0,43	78	35
4527,7	26	0,41	72	31
4827,1	27,3	0,53	83	31
5112,9	33,1	0,42	50	8
5829,7	20,8	0,46	61	15
6106,5	27	0,48	56	8
8341,2	16,6	0,48	94	46
8371,2	15,8	0,46	61	15
8466,3	18	0,46	61	15
8518,7	15,7	0,51	78	27
8578,4	17	0,47	67	19
9182	17,1	0,43	89	46
9944,2	16,7	0,44	67	23
10949,7	26,3	0,48	56	8

Table 11:

molecular weight [Da]	migration time [min]	discrimination factor	frequency [%]	
			IgA + MGN	control
800,5	22,5	-0,41	14	55
866,5	22	0,52	52	0
874,7	19,2	0,45	52	7
879,5	15,6	-0,84	9	93
937,5	23	0,43	57	14
981,6	21,3	-0,42	34	76
995,5	21,4	0,5	54	3
1010,6	16,3	-0,57	5	62
1028,6	21,2	-0,56	13	69
1046,5	21,5	-0,74	9	83
1060,7	17,4	0,47	61	14
1134,6	20,4	-0,59	41	100
1169,5	45,7	-0,52	4	55
1194,7	22,9	-0,68	32	100
1219,7	23,7	0,43	61	17
1224,7	18,1	-0,58	11	69
1235,3	23,2	-0,59	27	86
1250,6	23	-0,52	48	100
1265,6	23	-0,77	23	100
1285,8	18	0,54	57	3
1297,6	22,1	0,61	75	14
1321,7	23,6	-0,46	20	66
1333,7	22,8	0,54	54	0
1335,5	21,1	0,45	63	17
1368,8	17	0,65	75	10
1386,9	17,2	0,75	79	3
1404,8	16	0,61	75	14
1424,4	32,1	-0,57	43	100
1438,5	23,6	-0,43	54	97
1448,6	17,5	0,57	64	7
1460,9	19,5	0,54	54	0
1482	17,7	0,65	71	7
1493,8	20	0,48	52	3
1499,9	17,6	0,77	88	10
1539,7	34,1	-0,72	25	97
1567	19,2	0,59	63	3
1579,5	22,9	-0,56	30	86
1585	18,1	0,77	91	14
1621,8	13,4	0,47	75	28
1644,4	20,8	0,63	63	0
1651,9	17,2	0,64	68	3
1698,1	18,4	0,72	96	24
1760	17,7	0,5	57	7
1805,1	19,8	0,5	50	0
1811,2	17,9	0,83	96	14
1819,7	20,4	0,52	59	7
1829,1	18	0,5	98	48
1851,1	17,8	0,77	84	7
1863,8	37	-0,52	38	90
1867,2	18,6	0,82	86	3
1872,9	18,6	0,65	71	7
1878,5	17,3	0,52	59	7

1880,1	37,5	-0,48	7	55
1895,1	16,2	0,63	77	14
1924,5	20,4	0,52	52	0
1943	19,5	0,79	86	7
1955	19,9	0,41	52	10
1977	12,7	-0,57	5	62
2039,1	18,6	-0,49	27	76
2042,1	17,7	0,68	75	7
2048	19,9	-0,54	46	100
2057,3	19,1	0,71	71	0
2133,3	21,5	0,5	64	14
2147	19,5	0,54	68	14
2174,9	27,8	-0,81	9	90
2233,1	18,1	-0,48	11	59
2246,2	22,1	0,68	68	0
2249,1	18,7	-0,42	23	66
2258,6	18,9	0,5	50	0
2279,5	22,5	0,5	54	3
2377,4	18,4	-0,67	13	79
2389,2	18,6	0,59	66	7
2405,6	17,8	0,59	59	0
2427,1	16,4	0,84	95	10
2502,2	19,2	0,5	61	10
2518,7	18,8	0,61	64	3
2540,4	16	0,68	75	7
2562,9	19,1	-0,52	38	90
2566,7	13,9	0,66	70	3
2608,3	21,8	0,54	75	21
2621,5	16,5	0,68	71	3
2649,6	28,9	-0,5	13	62
2695,4	19,7	-0,54	43	97
2742,3	23,7	-0,61	36	97
2752,4	15,5	0,83	93	10
2755,3	23,6	0,68	71	3
2790,6	16,4	0,64	64	0
2799,7	20,4	-0,45	48	93
2825	20,8	-0,63	38	100
2838,7	20,3	-0,57	36	93
2914,8	17	0,43	50	7
2936,8	16	0,7	70	0
3011,5	24,5	-0,68	32	100
3013,3	16,8	-0,42	20	62
3040,9	25,5	-0,72	25	97
3098,3	24,8	-0,51	11	62
3205,9	15,7	-0,53	9	62
3209,4	19	-0,67	16	83
3265,5	24	-0,66	23	90
3281,1	27,3	-0,56	38	93
3287,4	25,2	-0,54	39	93
3303,5	28,7	-0,49	23	72
3333,2	14,1	-0,64	5	69
3359,6	26,2	-0,72	4	76
3375,6	25,3	-0,5	9	59
3385,7	21,2	-0,61	32	93
3402,3	17,8	-0,56	13	69
3405,3	21,6	-0,75	18	93
3416,7	26,4	-0,46	9	55
3432,5	25,6	-0,69	7	76
3441,6	25,4	-0,45	55	100

3457,8	25,4	-0,45	55	100
3502,8	14,8	-0,48	7	55
3582,9	14,2	-0,72	7	79
3969,1	18,1	-0,43	13	55
3987,1	17,3	-0,63	27	90
4044,5	18	-0,44	14	59
4054,8	15,5	0,45	59	14
4098,2	20,5	-0,56	23	79
4153,7	14,7	0,58	71	14
4240,6	14,8	0,63	84	21
4290,3	23,4	-0,65	7	72
4306,5	23,5	-0,57	2	59
4369,2	15,5	0,44	75	31
4626	15,7	0,52	55	3
4712,9	15,8	0,72	79	7
4748	14,7	-0,67	20	86
6171,4	23	-0,7	16	86
6186,7	23,3	-0,67	20	86
8764,7	12,8	0,46	88	41
9868,4	17,2	-0,63	9	72
10044,3	13,1	0,65	89	24
10516,9	13,9	0,45	52	7
12719	22,6	-0,44	25	69

Table 12:

molecular weight [Da]	migration time [min]	discrimination factor	frequency [%]	
			IgA	control
800,5	22,5	-0,44	12	55
852,6	19,7	0,51	51	0
862,4	27,2	0,56	56	0
866,5	22	0,58	58	0
879,5	15,6	-0,91	2	93
937,5	23	0,49	63	14
981,6	21,3	-0,48	28	76
995,5	21,4	0,55	58	3
1008,5	20	-0,44	12	55
1010,6	16,3	-0,6	2	62
1028,6	21,2	-0,57	12	69
1033,7	21,6	0,53	53	0
1046,5	21,5	-0,8	2	83
1134,6	20,4	-0,63	37	100
1144,7	25,8	0,48	51	3
1169,5	45,7	-0,51	5	55
1181,6	23,7	0,48	65	17
1194,7	22,9	-0,65	35	100
1219,7	23,7	0,55	72	17
1224,7	18,1	-0,64	5	69
1235,3	23,2	-0,61	26	86
1250,6	23	-0,49	51	100
1265,6	23	-0,79	21	100
1297,6	22,1	0,63	77	14
1321,7	23,6	-0,52	14	66
1333,7	22,8	0,63	63	0
1335,5	21,1	0,55	72	17
1368,8	17	0,64	74	10
1386,9	17,2	0,71	74	3

1404,8	16	0,58	72	14
1424,4	32,1	-0,49	51	100
1448,6	17,5	0,54	60	7
1482	17,7	0,61	67	7
1493,8	20	0,55	58	3
1499,9	17,6	0,76	86	10
1539,7	34,1	-0,66	30	97
1567	19,2	0,5	53	3
1573,8	27,8	0,49	77	28
1579,5	22,9	-0,65	21	86
1585	18,1	0,79	93	14
1621,8	13,4	0,47	74	28
1644,4	20,8	0,63	63	0
1651,9	17,2	0,62	65	3
1689,7	32,2	0,47	53	7
1698,1	18,4	0,74	98	24
1811,2	17,9	0,84	98	14
1819,7	20,4	0,47	53	7
1829,1	18	0,49	98	48
1851,1	17,8	0,74	81	7
1863,8	37	-0,43	47	90
1867,2	18,6	0,8	84	3
1872,9	18,6	0,63	70	7
1878,5	17,3	0,49	56	7
1880,1	37,5	-0,48	7	55
1895,1	16,2	0,58	72	14
1924,5	20,4	0,51	51	0
1943	19,5	0,79	86	7
1955	19,9	0,43	53	10
1977	12,7	-0,57	5	62
2039,1	18,6	-0,53	23	76
2042,1	17,7	0,7	77	7
2048	19,9	-0,6	40	100
2057,3	19,1	0,67	67	0
2133,3	21,5	0,56	70	14
2147	19,5	0,44	58	14
2174,9	27,8	-0,8	9	90
2233,1	18,1	-0,52	7	59
2246,2	22,1	0,79	79	0
2249,1	18,7	-0,49	16	66
2279,5	22,5	0,5	53	3
2377,4	18,4	-0,79	0	79
2389,2	18,6	0,63	70	7
2405,6	17,8	0,58	58	0
2427,1	16,4	0,83	93	10
2483,4	21,6	-0,47	12	59
2502,2	19,2	0,57	67	10
2518,7	18,8	0,52	56	3
2540,4	16	0,68	74	7
2562,9	19,1	-0,52	37	90
2566,7	13,9	0,71	74	3
2608,3	21,8	0,65	86	21
2621,5	16,5	0,69	72	3
2649,6	28,9	-0,5	12	62
2695,4	19,7	-0,59	37	97
2742,3	23,7	-0,62	35	97
2752,4	15,5	0,85	95	10
2755,3	23,6	0,8	84	3
2761,3	17,7	-0,4	12	52
2790,6	16,4	0,6	60	0
2799,7	20,4	-0,42	51	93

2825	20,8	-0,67	33	100
2838,7	20,3	-0,58	35	93
2936,8	16	0,67	67	0
3011,5	24,5	-0,63	37	100
3013,3	16,8	-0,48	14	62
3040,9	25,5	-0,69	28	97
3098,3	24,8	-0,57	5	62
3205,9	15,7	-0,57	5	62
3209,4	19	-0,73	9	83
3265,5	24	-0,69	21	90
3281,1	27,3	-0,56	37	93
3287,4	25,2	-0,47	47	93
3303,5	28,7	-0,49	23	72
3333,2	14,1	-0,67	2	69
3359,6	26,2	-0,71	5	76
3375,6	25,3	-0,54	5	59
3385,7	21,2	-0,56	37	93
3402,3	17,8	-0,57	12	69
3405,3	21,6	-0,79	14	93
3416,7	26,4	-0,44	12	55
3432,5	25,6	-0,67	9	76
3502,8	14,8	-0,53	2	55
3582,9	14,2	-0,72	7	79
3841,4	14,9	-0,44	42	86
3969,1	18,1	-0,41	14	55
3987,1	17,3	-0,62	28	90
4044,5	18	-0,42	16	59
4098,2	20,5	-0,51	28	79
4153,7	14,7	0,51	65	14
4240,6	14,8	0,61	81	21
4290,3	23,4	-0,63	9	72
4306,5	23,5	-0,56	2	59
4712,9	15,8	0,68	74	7
4748	14,7	-0,68	19	86
6171,4	23	-0,68	19	86
6186,7	23,3	-0,65	21	86
8764,7	12,8	0,45	86	41
9868,4	17,2	-0,63	9	72
10044,3	13,1	0,64	88	24

Table 13:

molecular weight [Da]	migration time [min]	discrimination factor	frequency [%]	
			IgA	MGN
816,7	16	-0,47	7	54
852,6	19,7	0,51	51	0
862,4	27,2	0,56	56	0
874,7	19,2	-0,43	42	85
943,6	17,5	-0,59	26	85
977,6	18	-0,5	12	62
994,5	18,3	-0,52	2	54
1004,6	22,4	-0,49	5	54
1040,4	19,5	-0,59	2	62
1060,7	17,4	-0,41	51	92
1099,8	17,5	-0,57	28	85
1108,6	17,3	-0,5	42	92
1143,6	30,2	0,45	60	15

1157,5	31,6	0,47	63	15
1179,6	31,6	0,49	72	23
1186,7	19,7	-0,49	5	54
1195,6	31,4	0,64	79	15
1198,8	17,7	-0,49	28	77
1203,7	19,3	0,45	60	15
1217,6	30,1	0,78	93	15
1219,7	23,7	0,49	72	23
1239,6	30,6	0,75	91	15
1252,7	20,5	-0,81	12	92
1255,6	29,4	0,49	72	23
1261,5	30,1	0,51	74	23
1268,4	23	-0,51	26	77
1270,7	16,6	-0,47	7	54
1285,8	18	-0,46	47	92
1287,6	18,8	-0,55	14	69
1311,8	17,9	-0,46	23	69
1335,5	21,1	0,41	72	31
1340,6	16,7	-0,52	9	62
1343,1	18,5	-0,48	21	69
1350,8	19	-0,48	14	62
1365,7	21,2	-0,62	7	69
1377,8	17,1	-0,47	7	54
1381	19,8	-0,62	7	69
1402,4	18,7	-0,52	2	54
1405,8	31,3	0,54	70	15
1424,8	14,7	-0,5	12	62
1446,7	32,7	0,44	67	23
1455,8	20,7	-0,7	7	77
1464,1	21,3	-0,55	7	62
1465,7	19,4	-0,67	2	69
1470,8	17,3	-0,47	7	54
1472	21,3	-0,53	16	69
1484,5	17,5	-0,73	12	85
1486,4	19,5	-0,58	12	69
1514,6	19,4	-0,49	5	54
1519,1	13,4	-0,5	12	62
1523,7	33	0,65	95	31
1539,6	21,8	-0,62	30	92
1545,8	33,5	0,61	84	23
1556,7	19,6	-0,85	0	85
1561,8	33,3	0,55	93	38
1561,9	19,8	-0,57	5	62
1573,8	27,8	0,61	77	15
1579,5	22,9	-0,41	21	62
1596,9	18,3	-0,6	9	69
1603	19,1	-0,55	7	62
1611,6	32,9	0,51	51	0
1627,6	19,6	-0,52	2	54
1637,8	19,3	-0,6	9	69
1651,8	34	0,59	74	15
1707,6	19,7	-0,42	12	54
1726	20,1	-0,6	9	69
1775,9	20,8	-0,42	12	54
1782,3	18,9	-0,42	35	77
1787,7	19,3	-0,48	14	62
1791,3	19,3	-0,56	21	77
1793,1	23	-0,61	16	77
1799,3	18,7	-0,54	0	54
1845,3	20,1	-0,68	9	77
1876,8	19,2	-0,52	33	85

1887,6	21,5	-0,45	9	54
1891,2	18,7	-0,59	2	62
1936,5	21,6	-0,46	23	69
1968,5	27,3	-0,5	12	62
1971,3	22,2	-0,5	12	62
1988,9	18,9	-0,42	12	54
2014,9	18,1	-0,69	23	92
2025,9	21,7	-0,49	5	54
2063,8	17,7	-0,49	51	100
2085,9	21	-0,45	9	54
2113	18,6	-0,42	12	54
2129,1	16,6	-0,57	5	62
2135,3	18,1	-0,68	9	77
2147	19,5	-0,42	58	100
2154,1	21,2	-0,44	33	77
2159	23,1	-0,47	37	85
2166,6	20,1	-0,48	14	62
2171,7	21,1	-0,41	21	62
2178,4	14,9	-0,52	9	62
2183,5	16,3	-0,47	7	54
2229,5	20,2	-0,7	7	77
2246,2	22,1	0,48	79	31
2290,5	20,1	-0,41	21	62
2292,2	20,6	-0,47	7	54
2309,7	16	-0,52	2	54
2325,2	18,2	-0,53	16	69
2361,6	17,3	-0,49	5	54
2377,4	18,4	-0,54	0	54
2407,5	19,3	-0,45	9	54
2421,1	16,1	-0,62	0	62
2464,4	19,8	-0,69	23	92
2466,6	17	-0,48	21	69
2475,8	18,8	-0,64	21	85
2483,4	21,6	-0,73	12	85
2493,3	14,9	-0,57	28	85
2522,4	16,3	-0,41	28	69
2547,1	17,9	-0,55	14	69
2553,7	17,7	-0,72	5	77
2573,5	20,3	-0,45	16	62
2586,7	18,5	-0,59	2	62
2599,2	21	-0,68	9	77
2608,3	21,8	0,48	86	38
2669,1	18,4	-0,52	2	54
2676	20,6	-0,55	14	69
2681,5	20,1	-0,42	12	54
2684,2	19,4	-0,55	7	62
2687,5	19,2	-0,59	2	62
2755,3	23,6	0,53	84	31
2807,6	17,3	-0,47	7	54
2810,5	18,1	-0,5	12	62
2831,2	19	-0,43	19	62
2847,7	16,5	-0,5	35	85
2914,8	17	-0,45	40	85
2959,7	16,6	-0,47	7	54
3030	16,7	-0,44	26	69
3062,1	17,9	-0,54	23	77
3441,6	25,4	0,62	70	8
3478,6	27,6	0,78	86	8
3495,5	25,8	0,71	79	8
3841,4	14,9	-0,43	42	85
4183,6	16,3	-0,45	9	54

4479,4	14,8	-0,45	9	54
4483,2	16,2	-0,47	7	54
4527,6	16,3	-0,5	12	62
4566,6	16,6	-0,52	2	54
4594,4	14,2	-0,45	9	54
8053,8	12,9	-0,44	26	69

Table 14:

mass	CE_t	frequency [%]				mass	CE_t	frequency [%]			
[Da]	[min]	control	FSGS	MCD	MGN	[Da	[min]	control	FSGS	MCD	MGN
3012,09	39,45	100	60	94	72	1028,57	37,79	98	40	69	22
1539,67	50,20	100	90	100	89	876,41	48,89	98	70	38	67
2249,19	33,82	100	80	100	83	2205,03	36,94	98	70	88	83
3152,55	38,22	100	60	69	39	3402,40	33,83	98	50	94	56
3360,09	44,33	100	60	81	72	2377,60	32,06	98	80	94	83
3001,97	48,35	100	100	100	89	2175,03	44,27	98	80	100	100
2257,19	46,60	100	60	81	67	2385,45	45,47	98	80	81	94
2563,42	32,22	100	90	75	72	2046,99	45,39	98	50	56	50
2158,98	46,70	100	90	88	89	2409,90	32,54	98	100	94	94
3287,97	43,92	100	90	100	89	3433,28	44,46	98	70	56	56
3385,76	36,60	100	100	81	72	1545,75	54,72	98	90	100	89
3271,80	44,05	100	80	94	61	2736,31	32,30	97	100	94	89
2007,69	33,06	100	90	88	89	3723,33	32,48	97	20	50	22
1194,61	39,46	100	70	88	83	1737,76	41,29	97	90	88	89
1265,62	40,37	100	70	94	72	1378,54	45,45	97	60	81	56
1435,69	39,91	100	90	94	89	2854,41	43,80	97	50	75	56
1261,53	49,63	100	80	88	61	2068,54	41,06	97	80	81	61
1438,56	37,63	100	100	100	100	2663,36	36,38	97	80	88	83
1446,50	52,53	100	80	94	67	2085,50	39,21	97	100	94	94
3265,77	42,28	100	80	75	78	2682,49	35,03	97	90	75	89
3121,36	42,46	100	50	75	44	1046,57	38,63	97	20	38	6
1911,14	37,40	100	90	100	94	2994,61	40,63	97	90	88	72
1321,91	41,10	100	70	69	50	2583,98	43,75	97	30	62	28
2695,49	35,27	100	70	88	78	2129,48	35,14	97	60	50	17
1235,59	41,42	100	60	69	50	981,56	37,39	97	50	81	56
2799,94	37,08	100	80	88	83	879,55	26,95	97	60	94	22
2169,75	39,56	100	100	100	94	2394,29	36,32	97	100	100	94
1224,74	33,57	100	50	94	33	3986,98	30,46	97	50	69	44
1451,67	41,11	100	100	100	83	2483,58	38,74	97	90	81	78
3479,32	48,53	100	70	75	17	1523,73	54,29	97	90	100	89
2649,94	45,91	100	90	75	78	1889,76	46,38	97	30	44	39
2687,36	41,87	100	60	75	61	1507,64	54,43	97	90	100	89
3458,52	44,64	100	90	94	72	3209,22	34,27	97	30	81	33
3442,84	42,54	100	80	94	56	3022,82	33,82	97	40	62	61
2048,19	33,07	100	40	81	61	1765,17	35,50	97	90	88	94
2679,46	34,99	100	50	94	67	1367,67	53,27	97	90	94	94
2227,34	38,28	100	70	62	83	2726,38	40,38	97	100	88	100
1239,52	50,23	100	100	100	78	1179,57	52,17	97	80	100	83
3098,80	42,63	100	60	81	56	1651,86	55,15	97	90	100	78

2839,07	35,41	100	80	88	83	3376,24	45,17	97	50	88	67
3417,12	45,12	100	70	81	61	3293,15	54,21	97	50	75	50
3426,20	42,48	100	70	88	72	1579,50	39,43	97	100	94	94
3041,16	45,04	100	80	81	39	3474,27	43,37	95	70	75	67
1508,70	41,26	98	90	94	61	2848,83	36,33	95	70	75	44
1462,67	53,58	98	70	88	50	3319,28	46,22	95	50	62	39
3280,96	36,76	98	20	50	22	1000,52	33,96	95	40	44	28
1877,33	29,62	98	100	100	100	3281,97	49,44	95	60	50	56
2742,25	42,25	98	90	75	89	1885,74	57,47	95	70	75	28
3092,71	43,86	98	90	81	78	3556,92	34,85	95	100	75	78
2196,66	45,45	98	80	100	89	1609,17	42,60	95	100	94	89
6187,55	39,78	98	60	94	67	2767,41	31,39	95	40	69	56
2825,42	36,54	98	50	88	39	3108,81	44,70	95	50	75	56
1255,55	49,81	98	100	100	78	2233,00	31,06	95	30	69	56
2717,56	34,43	98	60	69	61	882,55	36,55	95	60	31	72
3149,67	41,62	98	70	88	50	1680,16	37,32	95	100	94	100
1195,53	51,76	98	70	94	83	1673,80	54,59	95	80	94	61
3496,02	43,85	98	90	94	61	2336,78	42,47	95	70	94	89
1561,69	54,17	98	90	100	83	1217,64	48,54	95	100	94	83
1250,63	41,97	98	70	100	89	1489,49	42,21	95	90	94	72
3295,77	38,36	98	50	50	33	2442,06	46,85	95	70	81	67
3405,68	37,84	98	40	62	17	2279,06	47,16	95	70	69	44
1578,01	52,53	98	70	75	50	4748,51	25,38	95	50	38	33
1134,58	37,11	98	90	94	94	1766,84	35,15	95	100	100	100

Table 15:

Mass [Da]	CE-time [min]	Frequency Healthy [%]	Frequency FSGS [%]	Frequency MCD [%]	Frequency MGN [%]
1435,69	32,7	94	86	100	7
1282,39	29,3	69	29	29	0
3531,01	26,9	69	0	0	0
5801,94	13,3	69	0	7	7

Table 16:

healthy vs. renal patients	
mass [Da]	CE time [min]
909,4	40,3
1159,6	39,0
1338,7	47,2
1686,8	38,2
1847,8	57,0
1966,3	25,1
1990,8	47,3

2146,3	25,8
2432,2	38,3
2465,0	22,8
3707,0	31,9

Table 17:

MGN vs. MCD	
Mass	CE_t
879,6	26,9
1279,7	38,3
1341,8	33,1
1404,9	29,4
1569,8	48,3
1574,8	33,9
1605,9	23,7
2527,3	40,8
5112,9	33,1

Table 18:

MCD vs. FSGS	
Mass	CE_t
1199,6	31,0
1826,9	50,8
2077,3	35,8
2258,9	33,6
2918,0	42,2

Table 19:

MGN vs. FSGS	
Mass	CE_t
2312,5	22,9
2453,6	32,0
2639,6	45,2
9182,0	17,1

Table 20:

mass [Da]	CE time [min]	%healthy	% MGN	mass [Da]	CE time [min]	%healthy	% MGN
4098,2	40,1	100	0	1933,02	41,5	100	12
3685,9	35,9	100	0	1889,82	49,3	100	12
3531,3	42,9	100	0	1636,69	46,8	100	12
3359,7	48,4	100	0	1579,76	47,3	100	12
3287,4	47,4	100	0	1438,66	45,4	100	12

3265,3	51,6	100	0	1321,59	45,8	100	12
3098,5	46,9	100	0	1255,53	53	100	12
3041,3	46,5	100	0	1200,53	53,4	100	12
3011,3	46,5	100	0	2427,43	27	12	100
2742,2	45,8	100	0	1829,09	33,8	12	100
2563,2	34,5	100	0	4627,01	28,7	0	88
2483,5	44,5	100	0	2621,42	29	0	88
2385,2	50,4	100	0	1942,57	34,8	0	88
1893,1	42,5	100	0	1867,06	33,8	0	88
1639,9	47,5	100	0	1759,92	32,5	0	88
1609,7	47,3	100	0	1460,83	39,8	0	88
1580,9	41,3	100	0	3013,36	36,6	88	12
1508,7	46,6	100	0	2838,9	39,8	88	12
1489,6	46,2	100	0	2710,31	52,8	88	12
1424,7	56	100	0	2395,04	40,4	88	12
1407,6	54,6	100	0	1876,91	37,2	88	12
1160,6	52,7	100	0	1863,86	59,2	88	12
981,53	41	100	0	1651,81	56,5	88	12
980,54	38	100	0	1561,56	56,1	88	12
876,4	52,2	100	0	1523,72	56,1	88	12
2752,9	29,3	0	100	1473,66	46,4	88	12
6171,1	43,4	88	0	1261,49	53,2	88	12
3851,9	41,2	88	0	1195,5	54	88	12
3706,8	35,2	88	0	10047	22,3	12	88
3634,2	43,6	88	0	4713,94	28,8	12	88
3631,3	36,3	88	0	4241,41	26,7	12	88
3478,9	47,9	88	0	1811,13	34,6	12	88
3376,3	48,5	88	0	1753,98	32,7	12	88
3338,2	38,6	88	0	1698,06	34,1	12	88
3292,7	56,7	88	0	1584,91	32,7	12	88
3280,6	42,2	88	0	4353,62	33,6	75	0
3271,5	47,3	88	0	4102,45	45,2	75	0
3248,5	47,2	88	0	4044,58	34,1	75	0
2849,2	39,4	88	0	3987,48	34,8	75	0
2736,4	39,3	88	0	3947,22	36	75	0
2682,1	37,3	88	0	3589,65	41,3	75	0
2642,6	44,7	88	0	3433,12	48,6	75	0
2584,3	51,9	88	0	3416,92	48,6	75	0
2257,1	50,3	88	0	3295,53	42	75	0
2204,9	44	88	0	3261,55	35,6	75	0
2196,9	49,9	88	0	3258,52	37,8	75	0
2039,2	35,8	88	0	3193,48	37,3	75	0
1680,8	47	88	0	3152,6	40,3	75	0
1635,8	56,8	88	0	3092,08	47,6	75	0
1539,7	46,3	88	0	2863,25	40,6	75	0
1423,7	54,4	88	0	2854,55	52,4	75	0
1422,5	55	88	0	2698,37	37,2	75	0
1353,7	43,2	88	0	2548,42	37,7	75	0
1046,6	42,6	88	0	2464,07	50,8	75	0

3969,5	34,4	100	12	2406,98	50,6	75	0
3496,1	47,1	100	12	2279	50,2	75	0
3442,2	47,9	100	12	2233,02	35,7	75	0
3405,4	42,4	100	12	2226,97	43	75	0
3385,6	41,5	100	12	2019,97	41,1	75	0
3281,7	53	100	12	1991,95	36,2	75	0
3209,4	37,1	100	12	1849,85	41,1	75	0
2799,9	42,4	100	12	1768,95	48,7	75	0
2378	38,8	100	12	1755,02	48,2	75	0
2170	42,6	100	12	1737,78	48,2	75	0
2008	37	100	12	1462,63	56,1	75	0
1949	41,5	100	12	1446,6	56	75	0
				1425,78	41,6	75	0

Table 21:

mass [Da]	CE time [min]	%healthy	% MGN
1405,6	55,7	75	0
1389,6	55,3	75	0
1322,6	45,4	75	0
1262,6	56,4	75	0
1246,6	55,6	75	0
1224,8	35,4	75	0
1141,7	41,6	75	0
1028,6	41,9	75	0
946,43	50,5	75	0
3723,1	35,5	100	25
3458,2	48,2	100	25
3001,8	51,8	100	25
2825,3	40,8	100	25
2695,3	39,1	100	25
2679,2	39,2	100	25
2410	39,6	100	25
2394	39,3	100	25
2048	35,9	100	25
1911,1	41,6	100	25
1545,7	57,3	100	25
1507,7	57,3	100	25
1467,8	41	100	25
1451,7	46,4	100	25
1435,7	46,3	100	25
1265,6	44,6	100	25
1250,6	45,7	100	25
1239,4	53,7	100	25
1235,6	44	100	25
1217,6	53,3	100	25
1194,6	44,1	100	25

1179,5	55	100	25
1716	32,1	25	100
4827,2	29,3	0	75
2937,4	29,6	0	75
2057,4	37	0	75
1851,1	33,8	0	75
1680,1	33,6	0	75
1517,9	30,2	0	75
1483,9	32,5	0	75
1481,9	33,8	0	75
1404,8	29	0	75
1398,8	34,1	0	75
1367,6	56,1	88	25
1157,6	54,9	88	25
3474,3	47,9	75	12
3402,5	37	75	12
2761,4	34,7	75	12
2644,1	33,5	75	12
2587,2	34,9	75	12
2579,7	50,5	75	12
2579,7	41,4	75	12
2175	50	75	12
2069,1	49,5	75	12
2047	49,5	75	12
1170,6	46	75	12
1386,8	32,3	25	88
8766,7	21,6	12	75
4154,4	26,4	12	75
3842,8	25,7	12	75
1873	33,9	12	75
1566,9	33,2	12	75
1499,9	33,7	12	75
1368,8	31,6	12	75
1285,7	31,1	12	75
1108,6	32,2	12	75
1099,6	31,2	12	75
1060,6	31,6	12	75

Table 22:

time [min]	mass [Da]	healthy, frequency	disease, frequency	type
22,9 ± 3,05	834,5 ± 0,10	3%	54%	Diabetes pos.
22,9 ± 3,03	869,4 ± 0,17	14%	63%	Diabetes pos.
24,2 ± 1,89	874,5 ± 0,09	28%	66%	Diabetes pos.
22,2 ± 2,19	907,5 ± 0,13	0%	41%	Diabetes pos.
29,0 ± 2,35	910,5 ± 0,09	15%	47%	Diabetes pos.

22,9 ± 3,18	947,6 ± 0,22	17%	51%	Diabetes pos.
26,8 ± 2,98	950,5 ± 0,12	0%	24%	Diabetes pos.
23,2 ± 4,87	995,6 ± 0,14	23%	50%	Diabetes pos.
27,4 ± 3,59	1082,6 ± 0,16	0%	44%	Diabetes pos.
32,3 ± 1,99	1096,5 ± 0,14	10%	51%	Diabetes pos.
26,8 ± 3,85	1176,6 ± 0,13	21%	59%	Diabetes pos.
22,3 ± 3,45	1222,8 ± 0,22	17%	56%	Diabetes pos.
30,6 ± 3,31	1236,6 ± 0,11	24%	59%	Diabetes pos.
52,6 ± 4,80	1285,0 ± 0,09	14%	54%	Diabetes pos.
28,8 ± 3,98	1332,7 ± 0,20	23%	55%	Diabetes pos.
49,8 ± 4,72	1332,8 ± 0,16	8%	38%	Diabetes pos.
26,7 ± 2,79	1355,8 ± 0,15	17%	56%	Diabetes pos.
24,6 ± 2,84	1386,8 ± 0,14	53%	77%	Diabetes pos.
26,8 ± 3,26	1403,7 ± 0,21	8%	46%	Diabetes pos.
17,8 ± 4,12	1405,9 ± 0,15	14%	56%	Diabetes pos.
31,5 ± 3,71	1442,7 ± 0,27	15%	55%	Diabetes pos.
32,1 ± 3,38	1449,8 ± 0,14	41%	85%	Diabetes pos.
31,3 ± 5,27	1592,4 ± 0,38	3%	46%	Diabetes pos.
43,4 ± 4,41	1783,4 ± 0,30	33%	63%	Diabetes pos.
29,4 ± 3,08	1789,2 ± 0,39	28%	75%	Diabetes pos.
38,4 ± 1,09	1818,9 ± 0,21	28%	67%	Diabetes pos.
37,7 ± 1,04	1821,4 ± 0,39	14%	56%	Diabetes pos.
24,4 ± 2,55	1829,2 ± 0,23	45%	81%	Diabetes pos.
51,1 ± 4,11	1854,7 ± 0,41	14%	54%	Diabetes pos.
37,6 ± 3,30	1856,8 ± 0,48	33%	56%	Diabetes pos.
24,7 ± 2,63	1872,9 ± 0,35	43%	72%	Diabetes pos.
28,3 ± 3,47	1949,5 ± 0,32	17%	73%	Diabetes pos.
31,6 ± 2,90	1955,1 ± 0,32	55%	79%	Diabetes pos.
31,3 ± 3,00	1971,0 ± 0,45	20%	54%	Diabetes pos.
37,8 ± 2,40	2032,0 ± 0,30	25%	60%	Diabetes pos.
30,9 ± 4,69	2061,4 ± 0,58	10%	38%	Diabetes pos.
33,8 ± 3,76	2092,2 ± 0,46	18%	45%	Diabetes pos.
27,7 ± 4,43	2185,6 ± 0,46	10%	36%	Diabetes pos.
32,9 ± 1,48	2189,4 ± 0,34	14%	54%	Diabetes pos.
39,6 ± 5,31	2229,4 ± 0,48	5%	39%	Diabetes pos.
24,5 ± 5,14	2229,9 ± 0,33	25%	63%	Diabetes pos.
28,3 ± 3,30	2502,9 ± 0,56	20%	48%	Diabetes pos.
24,9 ± 4,84	2621,6 ± 0,97	20%	45%	Diabetes pos.
37,5 ± 4,52	2669,8 ± 0,39	23%	67%	Diabetes pos.
20,8 ± 4,47	2752,2 ± 0,76	35%	64%	Diabetes pos.
24,9 ± 4,31	2795,7 ± 0,96	13%	40%	Diabetes pos.
48,2 ± 3,61	3246,1 ± 0,43	0%	30%	Diabetes pos.
20,9 ± 3,33	3844,0 ± 0,52	3%	54%	Diabetes pos.
21,9 ± 2,62	4961,5 ± 0,89	10%	40%	Diabetes pos.
18,6 ± 2,91	5497,0 ± 0,66	18%	42%	Diabetes pos.
20,4 ± 2,20	808,4 ± 0,10	58%	9%	Diabetes neg.
45,3 ± 2,03	897,5 ± 0,09	48%	7%	Diabetes neg.
31,4 ± 1,08	929,5 ± 0,11	98%	46%	Diabetes neg.
41,2 ± 1,41	946,4 ± 0,10	85%	36%	Diabetes neg.

28,0 ± 1,04	980,5 ± 0,07	85%	31%	Diabetes neg.
26,7 ± 2,26	1000,5 ± 0,09	83%	41%	Diabetes neg.
27,8 ± 1,51	1008,5 ± 0,10	95%	41%	Diabetes neg.
29,3 ± 2,55	1012,5 ± 0,10	63%	17%	Diabetes neg.
43,6 ± 2,03	1047,5 ± 0,11	90%	26%	Diabetes neg.
25,0 ± 3,91	1052,6 ± 0,08	45%	4%	Diabetes neg.
37,4 ± 5,63	1066,5 ± 0,14	58%	13%	Diabetes neg.
22,8 ± 1,78	1075,5 ± 0,13	68%	26%	Diabetes neg.
28,9 ± 3,89	1088,6 ± 0,15	65%	21%	Diabetes neg.
44,4 ± 2,06	1106,5 ± 0,11	80%	18%	Diabetes neg.
34,1 ± 1,80	1107,5 ± 0,10	88%	35%	Diabetes neg.
42,8 ± 3,26	1120,5 ± 0,06	60%	14%	Diabetes neg.
29,1 ± 2,26	1134,6 ± 0,10	95%	49%	Diabetes neg.
28,2 ± 3,00	1137,7 ± 0,11	70%	24%	Diabetes neg.
45,5 ± 2,34	1139,5 ± 0,20	83%	22%	Diabetes neg.
32,9 ± 1,25	1159,6 ± 0,11	80%	27%	Diabetes neg.
23,3 ± 4,17	1180,5 ± 0,16	50%	9%	Diabetes neg.
43,8 ± 2,08	1200,6 ± 0,11	95%	50%	Diabetes neg.
27,2 ± 3,22	1204,6 ± 0,17	60%	17%	Diabetes neg.
44,9 ± 2,53	1209,5 ± 0,09	83%	17%	Diabetes neg.
47,8 ± 2,73	1224,6 ± 0,12	75%	19%	Diabetes neg.
25,6 ± 2,43	1246,7 ± 0,15	73%	30%	Diabetes neg.
47,9 ± 2,66	1268,6 ± 0,09	68%	25%	Diabetes neg.
43,9 ± 1,80	1277,5 ± 0,10	70%	28%	Diabetes neg.
46,0 ± 2,69	1278,5 ± 0,09	58%	10%	Diabetes neg.
33,1 ± 1,82	1282,6 ± 0,13	62%	7%	Diabetes neg.
29,3 ± 3,88	1331,7 ± 0,18	65%	12%	Diabetes neg.
45,9 ± 4,78	1405,5 ± 0,33	93%	45%	Diabetes neg.
44,4 ± 3,90	1423,6 ± 0,16	60%	20%	Diabetes neg.
19,2 ± 3,40	1484,8 ± 0,19	68%	13%	Diabetes neg.
36,9 ± 2,02	1609,6 ± 0,13	85%	13%	Diabetes neg.
38,9 ± 3,78	1639,7 ± 0,27	63%	19%	Diabetes neg.
33,2 ± 3,34	1662,9 ± 0,21	62%	5%	Diabetes neg.
35,8 ± 2,19	1664,6 ± 0,29	66%	10%	Diabetes neg.
36,2 ± 4,78	1666,6 ± 0,34	75%	29%	Diabetes neg.
35,9 ± 2,98	1678,1 ± 0,44	60%	18%	Diabetes neg.
37,3 ± 2,99	1716,8 ± 0,23	73%	19%	Diabetes neg.
46,5 ± 4,38	1717,5 ± 0,37	79%	15%	Diabetes neg.
37,9 ± 4,18	1746,0 ± 0,33	83%	34%	Diabetes neg.
25,1 ± 2,25	1817,6 ± 0,27	65%	8%	Diabetes neg.
34,2 ± 3,95	1823,4 ± 0,47	73%	30%	Diabetes neg.
29,1 ± 3,59	1849,8 ± 0,30	100%	56%	Diabetes neg.
49,3 ± 4,49	1914,1 ± 0,36	88%	38%	Diabetes neg.
44,2 ± 4,23	1916,7 ± 0,33	69%	10%	Diabetes neg.
39,8 ± 2,19	2030,8 ± 0,35	93%	38%	Diabetes neg.
31,9 ± 1,61	2118,9 ± 0,21	73%	14%	Diabetes neg.
41,2 ± 2,45	2179,3 ± 0,42	58%	17%	Diabetes neg.
20,1 ± 2,78	2219,0 ± 0,26	53%	13%	Diabetes neg.
25,8 ± 2,70	2256,9 ± 0,47	85%	26%	Diabetes neg.

45,1 ± 5,23	2273,4 ± 0,42	79%	22%	Diabetes neg.
40,7 ± 1,90	2279,0 ± 0,33	90%	20%	Diabetes neg.
26,8 ± 3,73	2320,2 ± 0,55	78%	34%	Diabetes neg.
23,6 ± 3,10	2332,2 ± 0,35	53%	11%	Diabetes neg.
44,5 ± 3,08	2345,6 ± 0,46	75%	34%	Diabetes neg.
25,7 ± 5,16	2384,5 ± 0,63	65%	21%	Diabetes neg.
38,5 ± 3,62	2423,9 ± 0,41	88%	29%	Diabetes neg.
34,2 ± 2,92	2429,9 ± 0,51	65%	18%	Diabetes neg.
23,3 ± 2,54	2443,3 ± 0,46	66%	5%	Diabetes neg.
41,7 ± 3,72	2548,1 ± 0,57	69%	15%	Diabetes neg.
27,3 ± 4,77	2548,3 ± 0,66	83%	35%	Diabetes neg.
43,6 ± 2,08	2548,3 ± 0,23	95%	41%	Diabetes neg.
24,0 ± 3,11	2581,5 ± 0,47	60%	13%	Diabetes neg.
24,0 ± 2,70	2587,4 ± 0,40	80%	26%	Diabetes neg.
41,7 ± 3,06	2606,8 ± 0,55	78%	35%	Diabetes neg.
31,3 ± 4,92	2636,4 ± 0,48	72%	12%	Diabetes neg.
25,5 ± 3,62	2644,2 ± 0,41	88%	33%	Diabetes neg.
29,2 ± 1,07	2654,0 ± 0,37	66%	0%	Diabetes neg.
29,8 ± 3,50	2698,2 ± 0,63	90%	29%	Diabetes neg.
43,0 ± 2,26	2710,5 ± 0,37	79%	5%	Diabetes neg.
25,1 ± 1,64	2761,3 ± 0,35	88%	44%	Diabetes neg.
31,3 ± 2,79	2808,5 ± 0,56	79%	22%	Diabetes neg.
42,0 ± 3,22	2876,5 ± 0,48	62%	7%	Diabetes neg.
33,7 ± 3,34	2898,7 ± 0,50	85%	43%	Diabetes neg.
42,2 ± 2,68	2908,1 ± 0,53	72%	17%	Diabetes neg.
35,4 ± 2,63	2917,6 ± 0,58	72%	12%	Diabetes neg.
35,4 ± 0,77	2978,1 ± 0,49	85%	35%	Diabetes neg.
36,1 ± 1,42	2994,6 ± 0,80	83%	24%	Diabetes neg.
43,5 ± 2,99	3023,4 ± 0,65	93%	34%	Diabetes neg.
44,4 ± 3,35	3045,2 ± 0,61	69%	12%	Diabetes neg.
22,9 ± 3,47	3076,4 ± 0,96	66%	7%	Diabetes neg.
35,7 ± 1,99	3082,3 ± 0,43	73%	22%	Diabetes neg.
33,6 ± 3,53	3136,8 ± 0,61	95%	47%	Diabetes neg.
21,7 ± 3,14	3154,8 ± 0,44	55%	10%	Diabetes neg.
26,5 ± 1,92	3193,7 ± 0,53	78%	32%	Diabetes neg.
24,4 ± 3,02	3206,3 ± 0,72	66%	7%	Diabetes neg.
28,2 ± 2,80	3250,9 ± 0,71	63%	18%	Diabetes neg.
48,2 ± 3,46	3293,2 ± 0,74	93%	39%	Diabetes neg.
31,4 ± 1,60	3295,7 ± 0,33	95%	40%	Diabetes neg.
27,2 ± 3,58	3338,4 ± 0,79	80%	34%	Diabetes neg.
37,3 ± 2,11	3381,6 ± 0,63	78%	26%	Diabetes neg.
27,6 ± 2,49	3452,1 ± 0,49	58%	15%	Diabetes neg.
37,3 ± 1,50	3463,0 ± 0,83	72%	15%	Diabetes neg.
19,6 ± 2,89	3583,4 ± 0,75	79%	20%	Diabetes neg.
34,0 ± 2,55	3634,4 ± 0,74	86%	29%	Diabetes neg.
37,7 ± 2,61	3681,8 ± 1,38	55%	14%	Diabetes neg.
25,5 ± 2,25	3686,2 ± 0,60	86%	20%	Diabetes neg.
36,0 ± 3,89	3735,7 ± 0,57	70%	28%	Diabetes neg.
30,3 ± 1,58	3852,3 ± 0,56	83%	41%	Diabetes neg.

29,6 ± 1,46	4098,4 ± 0,59	93%	20%	Diabetes neg.
28,8 ± 1,18	5428,8 ± 0,67	70%	19%	Diabetes neg.
33,1 ± 0,69	6187,5 ± 1,13	83%	10%	Diabetes neg.
26,0 ± 4,82	6212,0 ± 1,41	75%	26%	Diabetes neg.
23,3 ± 2,19	9868,8 ± 1,33	66%	0%	Diabetes neg.
21,7 ± 5,12	830,5 ± 0,11	4%	40%	Nephropathy pos.
32,4 ± 1,83	866,4 ± 0,11	0%	40%	Nephropathy pos.
30,6 ± 3,07	909,5 ± 0,13	11%	40%	Nephropathy pos.
32,8 ± 3,14	937,5 ± 0,11	14%	73%	Nephropathy pos.
24,9 ± 2,97	952,5 ± 0,16	11%	40%	Nephropathy pos.
32,1 ± 2,44	1033,5 ± 0,11	5%	40%	Nephropathy pos.
24,4 ± 2,87	1060,6 ± 0,16	17%	68%	Nephropathy pos.
27,5 ± 2,86	1131,6 ± 0,16	20%	68%	Nephropathy pos.
33,4 ± 3,48	1181,6 ± 0,15	22%	73%	Nephropathy pos.
33,0 ± 2,52	1203,6 ± 0,14	9%	50%	Nephropathy pos.
26,5 ± 3,68	1211,6 ± 0,14	14%	40%	Nephropathy pos.
33,1 ± 0,91	1219,6 ± 0,15	18%	40%	Nephropathy pos.
32,8 ± 3,30	1225,6 ± 0,13	12%	40%	Nephropathy pos.
30,7 ± 3,18	1297,7 ± 0,20	31%	82%	Nephropathy pos.
34,1 ± 2,05	1333,7 ± 0,23	9%	40%	Nephropathy pos.
44,7 ± 4,06	1337,5 ± 0,20	19%	59%	Nephropathy pos.
27,9 ± 4,19	1398,8 ± 0,36	29%	77%	Nephropathy pos.
21,3 ± 5,08	1423,7 ± 0,49	6%	50%	Nephropathy pos.
28,1 ± 4,95	1439,8 ± 0,19	19%	68%	Nephropathy pos.
24,5 ± 2,42	1466,0 ± 0,27	9%	77%	Nephropathy pos.
27,5 ± 4,93	1482,0 ± 0,42	33%	40%	Nephropathy pos.
29,8 ± 4,43	1482,9 ± 0,28	18%	40%	Nephropathy pos.
24,3 ± 2,65	1483,7 ± 0,28	26%	91%	Nephropathy pos.
24,6 ± 1,98	1500,0 ± 0,20	38%	86%	Nephropathy pos.
24,6 ± 2,90	1553,1 ± 0,28	14%	64%	Nephropathy pos.
29,0 ± 4,83	1556,7 ± 0,45	26%	73%	Nephropathy pos.
24,2 ± 2,48	1567,0 ± 0,22	26%	86%	Nephropathy pos.
28,8 ± 4,53	1596,9 ± 0,31	21%	86%	Nephropathy pos.
24,5 ± 2,43	1652,8 ± 0,25	14%	59%	Nephropathy pos.
26,3 ± 2,63	1669,8 ± 0,37	20%	64%	Nephropathy pos.
33,1 ± 3,22	1729,2 ± 0,36	6%	45%	Nephropathy pos.
30,5 ± 4,11	1744,4 ± 0,46	16%	59%	Nephropathy pos.
25,1 ± 3,42	1754,4 ± 0,41	53%	95%	Nephropathy pos.
24,2 ± 1,56	1776,0 ± 0,27	9%	50%	Nephropathy pos.
18,5 ± 3,55	1791,0 ± 0,38	7%	40%	Nephropathy pos.
32,2 ± 5,38	1792,9 ± 0,31	28%	40%	Nephropathy pos.
9,7 ± 2,54	1799,8 ± 0,29	0%	40%	Nephropathy pos.
25,3 ± 2,89	1810,9 ± 0,38	43%	91%	Nephropathy pos.
24,6 ± 2,34	1851,1 ± 0,21	43%	95%	Nephropathy pos.
27,2 ± 4,46	1867,3 ± 0,42	38%	91%	Nephropathy pos.
25,0 ± 3,97	1966,0 ± 0,53	16%	40%	Nephropathy pos.
28,7 ± 3,08	1982,8 ± 0,57	11%	40%	Nephropathy pos.
29,5 ± 5,53	1986,3 ± 0,36	15%	64%	Nephropathy pos.
23,3 ± 4,46	2045,9 ± 0,32	32%	40%	Nephropathy pos.

33,7 ± 3,16	2115,1 ± 0,53	30%	40%	Nephropathy pos.
20,5 ± 2,78	2177,1 ± 0,37	9%	40%	Nephropathy pos.
18,1 ± 4,24	2241,6 ± 0,41	9%	59%	Nephropathy pos.
21,2 ± 2,49	2250,7 ± 0,38	23%	64%	Nephropathy pos.
27,5 ± 2,53	2258,7 ± 0,49	9%	59%	Nephropathy pos.
20,0 ± 3,30	2356,4 ± 0,41	13%	59%	Nephropathy pos.
28,1 ± 3,95	2391,4 ± 0,42	13%	64%	Nephropathy pos.
25,7 ± 4,85	2406,1 ± 0,57	20%	77%	Nephropathy pos.
22,8 ± 4,28	2423,2 ± 0,53	14%	64%	Nephropathy pos.
21,9 ± 4,45	2427,3 ± 0,40	31%	91%	Nephropathy pos.
19,2 ± 4,24	2465,1 ± 0,62	9%	77%	Nephropathy pos.
25,4 ± 5,25	2493,0 ± 0,38	9%	50%	Nephropathy pos.
19,5 ± 4,66	2494,0 ± 0,66	12%	77%	Nephropathy pos.
23,7 ± 4,27	2494,9 ± 0,49	7%	40%	Nephropathy pos.
24,4 ± 5,51	2522,0 ± 0,67	17%	82%	Nephropathy pos.
20,1 ± 3,61	2540,5 ± 0,54	14%	68%	Nephropathy pos.
22,3 ± 4,72	2593,5 ± 0,30	7%	55%	Nephropathy pos.
20,0 ± 4,87	2613,9 ± 0,83	14%	55%	Nephropathy pos.
35,1 ± 1,62	2726,5 ± 0,67	61%	20%	Nephropathy pos.
25,0 ± 4,39	2775,1 ± 0,56	12%	40%	Nephropathy pos.
21,8 ± 3,78	2790,7 ± 0,55	19%	86%	Nephropathy pos.
25,9 ± 3,30	2892,2 ± 0,50	9%	50%	Nephropathy pos.
16,8 ± 2,72	2919,0 ± 0,26	2%	50%	Nephropathy pos.
21,9 ± 3,23	2937,0 ± 0,49	13%	86%	Nephropathy pos.
20,0 ± 4,81	2958,8 ± 0,80	5%	59%	Nephropathy pos.
34,4 ± 2,72	2962,0 ± 0,54	12%	20%	Nephropathy pos.
28,9 ± 3,56	3059,7 ± 0,78	30%	40%	Nephropathy pos.
28,3 ± 5,96	3088,0 ± 0,79	7%	20%	Nephropathy pos.
26,1 ± 2,72	3369,2 ± 0,73	21%	40%	Nephropathy pos.
26,0 ± 2,89	3483,4 ± 0,95	30%	40%	Nephropathy pos.
24,5 ± 3,92	4183,3 ± 1,44	4%	40%	Nephropathy pos.
21,0 ± 5,35	4241,0 ± 0,62	29%	73%	Nephropathy pos.
23,4 ± 4,09	4370,2 ± 1,01	11%	40%	Nephropathy pos.
22,8 ± 2,94	4527,6 ± 0,67	1%	45%	Nephropathy pos.
21,7 ± 3,00	4713,6 ± 0,44	7%	64%	Nephropathy pos.
24,6 ± 3,73	7556,6 ± 1,55	2%	40%	Nephropathy pos.
16,7 ± 5,54	8055,1 ± 2,10	12%	40%	Nephropathy pos.
13,2 ± 5,19	8765,8 ± 0,96	37%	82%	Nephropathy pos.
15,3 ± 4,97	9181,0 ± 1,28	10%	64%	Nephropathy pos.
14,0 ± 4,20	10046,1 ± 0,96	21%	77%	Nephropathy pos.
18,7 ± 5,50	10208,0 ± 1,24	2%	40%	Nephropathy pos.
17,4 ± 4,02	10518,2 ± 1,10	23%	64%	Nephropathy pos.
35,3 ± 5,04	924,5 ± 0,12	50%	0%	Nephropathy neg.
43,1 ± 2,61	928,4 ± 0,08	65%	14%	Nephropathy neg.
45,7 ± 2,25	955,5 ± 0,14	60%	5%	Nephropathy neg.
23,8 ± 2,94	1010,6 ± 0,09	67%	5%	Nephropathy neg.
31,2 ± 1,53	1028,5 ± 0,09	84%	32%	Nephropathy neg.
45,9 ± 2,27	1041,4 ± 0,10	57%	0%	Nephropathy neg.
31,5 ± 1,98	1046,5 ± 0,09	87%	32%	Nephropathy neg.

43,4 ± 2,24	1047,5 ± 0,12	68%	0%	Nephropathy neg.
18,1 ± 4,34	1050,7 ± 0,12	60%	0%	Nephropathy neg.
32,9 ± 3,03	1084,4 ± 0,11	69%	18%	Nephropathy neg.
46,7 ± 2,63	1125,5 ± 0,12	63%	9%	Nephropathy neg.
46,3 ± 2,70	1157,5 ± 0,10	83%	32%	Nephropathy neg.
43,7 ± 1,70	1160,5 ± 0,07	72%	18%	Nephropathy neg.
44,5 ± 3,67	1179,5 ± 0,09	97%	36%	Nephropathy neg.
45,0 ± 2,24	1191,6 ± 0,09	60%	9%	Nephropathy neg.
46,2 ± 2,59	1195,5 ± 0,10	98%	32%	Nephropathy neg.
44,2 ± 1,83	1200,6 ± 0,13	86%	0%	Nephropathy neg.
45,9 ± 2,04	1223,5 ± 0,10	80%	9%	Nephropathy neg.
44,5 ± 2,15	1239,6 ± 0,08	89%	0%	Nephropathy neg.
47,8 ± 3,08	1246,6 ± 0,11	60%	5%	Nephropathy neg.
46,8 ± 2,20	1254,7 ± 0,19	56%	5%	Nephropathy neg.
43,2 ± 2,90	1261,5 ± 0,16	91%	36%	Nephropathy neg.
48,6 ± 2,90	1262,5 ± 0,09	65%	0%	Nephropathy neg.
43,9 ± 2,16	1277,6 ± 0,11	67%	0%	Nephropathy neg.
36,7 ± 3,04	1288,7 ± 0,18	72%	23%	Nephropathy neg.
47,2 ± 3,17	1292,5 ± 0,14	67%	18%	Nephropathy neg.
47,8 ± 2,58	1308,5 ± 0,09	66%	0%	Nephropathy neg.
48,2 ± 2,67	1321,6 ± 0,11	53%	0%	Nephropathy neg.
34,8 ± 1,81	1321,7 ± 0,23	98%	41%	Nephropathy neg.
46,0 ± 4,93	1351,7 ± 0,15	63%	9%	Nephropathy neg.
47,7 ± 2,99	1367,6 ± 0,14	97%	23%	Nephropathy neg.
37,8 ± 2,93	1378,6 ± 0,16	87%	36%	Nephropathy neg.
47,5 ± 2,59	1389,7 ± 0,15	86%	18%	Nephropathy neg.
46,5 ± 2,28	1407,8 ± 0,20	79%	9%	Nephropathy neg.
44,6 ± 4,84	1422,1 ± 0,33	70%	0%	Nephropathy neg.
45,4 ± 3,62	1423,8 ± 0,19	75%	0%	Nephropathy neg.
48,0 ± 2,97	1424,7 ± 0,16	95%	18%	Nephropathy neg.
47,6 ± 3,40	1446,7 ± 0,16	92%	23%	Nephropathy neg.
46,5 ± 2,95	1450,4 ± 0,25	62%	9%	Nephropathy neg.
48,0 ± 2,95	1462,6 ± 0,17	97%	9%	Nephropathy neg.
35,7 ± 1,90	1487,7 ± 0,15	70%	18%	Nephropathy neg.
47,8 ± 2,35	1490,6 ± 0,12	72%	9%	Nephropathy neg.
49,2 ± 2,77	1491,7 ± 0,12	81%	14%	Nephropathy neg.
49,0 ± 3,14	1507,8 ± 0,17	99%	32%	Nephropathy neg.
49,2 ± 2,86	1523,7 ± 0,11	97%	18%	Nephropathy neg.
48,6 ± 2,70	1529,7 ± 0,19	83%	9%	Nephropathy neg.
49,2 ± 3,26	1539,7 ± 0,19	98%	23%	Nephropathy neg.
49,0 ± 3,19	1545,7 ± 0,13	99%	23%	Nephropathy neg.
49,8 ± 2,76	1561,6 ± 0,19	90%	18%	Nephropathy neg.
48,4 ± 3,12	1567,7 ± 0,20	65%	9%	Nephropathy neg.
48,1 ± 2,66	1573,7 ± 0,27	63%	5%	Nephropathy neg.
48,5 ± 4,03	1577,8 ± 0,35	94%	9%	Nephropathy neg.
50,6 ± 3,40	1587,1 ± 0,34	65%	0%	Nephropathy neg.
48,6 ± 2,68	1589,7 ± 0,14	86%	18%	Nephropathy neg.
45,9 ± 3,83	1591,7 ± 0,30	79%	18%	Nephropathy neg.
49,3 ± 3,22	1594,8 ± 0,14	88%	14%	Nephropathy neg.

48,8 ± 2,78	1605,7 ± 0,13	73%	18%	Nephropathy neg.
48,5 ± 2,81	1611,7 ± 0,14	73%	5%	Nephropathy neg.
46,3 ± 5,12	1636,4 ± 0,39	79%	23%	Nephropathy neg.
49,5 ± 3,37	1651,8 ± 0,19	99%	23%	Nephropathy neg.
45,2 ± 5,96	1657,7 ± 0,23	60%	5%	Nephropathy neg.
49,5 ± 3,33	1673,8 ± 0,14	95%	23%	Nephropathy neg.
49,6 ± 3,05	1689,8 ± 0,18	86%	0%	Nephropathy neg.
26,9 ± 3,18	1706,8 ± 0,30	78%	27%	Nephropathy neg.
49,4 ± 2,84	1734,4 ± 0,40	65%	5%	Nephropathy neg.
49,2 ± 3,17	1739,7 ± 0,22	59%	5%	Nephropathy neg.
45,1 ± 4,21	1748,0 ± 0,28	55%	5%	Nephropathy neg.
44,2 ± 4,71	1813,6 ± 0,38	58%	5%	Nephropathy neg.
39,1 ± 3,48	1817,0 ± 0,29	85%	18%	Nephropathy neg.
51,7 ± 3,48	1841,0 ± 0,23	59%	9%	Nephropathy neg.
50,4 ± 4,56	1848,2 ± 0,43	58%	0%	Nephropathy neg.
51,5 ± 2,94	1856,8 ± 0,24	59%	5%	Nephropathy neg.
52,7 ± 4,24	1863,8 ± 0,31	88%	14%	Nephropathy neg.
52,7 ± 3,92	1885,8 ± 0,20	70%	5%	Nephropathy neg.
47,7 ± 4,69	1902,1 ± 0,33	75%	0%	Nephropathy neg.
50,6 ± 3,95	1924,0 ± 0,48	68%	0%	Nephropathy neg.
26,6 ± 1,76	2048,5 ± 0,44	86%	20%	Nephropathy neg.
25,8 ± 1,39	2085,9 ± 0,24	83%	32%	Nephropathy neg.
39,9 ± 1,45	2087,8 ± 0,34	72%	23%	Nephropathy neg.
52,8 ± 4,09	2117,1 ± 0,17	78%	9%	Nephropathy neg.
28,3 ± 3,90	2129,7 ± 0,42	63%	0%	Nephropathy neg.
40,4 ± 1,53	2158,9 ± 0,26	86%	32%	Nephropathy neg.
39,7 ± 1,71	2174,9 ± 0,36	97%	45%	Nephropathy neg.
32,6 ± 1,79	2227,1 ± 0,41	81%	23%	Nephropathy neg.
29,3 ± 3,50	2249,0 ± 0,41	92%	41%	Nephropathy neg.
40,6 ± 1,25	2257,1 ± 0,35	94%	45%	Nephropathy neg.
46,2 ± 5,11	2273,5 ± 0,38	71%	18%	Nephropathy neg.
40,8 ± 2,66	2296,0 ± 0,40	63%	20%	Nephropathy neg.
40,9 ± 3,32	2327,6 ± 0,52	85%	36%	Nephropathy neg.
41,8 ± 2,45	2343,3 ± 0,43	77%	27%	Nephropathy neg.
40,8 ± 1,31	2385,3 ± 0,32	95%	45%	Nephropathy neg.
40,9 ± 2,68	2471,5 ± 0,52	69%	14%	Nephropathy neg.
41,5 ± 2,64	2493,5 ± 0,48	74%	18%	Nephropathy neg.
52,9 ± 3,98	2570,4 ± 0,27	71%	5%	Nephropathy neg.
34,1 ± 0,72	2642,8 ± 0,40	86%	36%	Nephropathy neg.
36,1 ± 2,56	2687,1 ± 0,49	84%	23%	Nephropathy neg.
42,8 ± 2,33	2710,6 ± 0,46	88%	18%	Nephropathy neg.
50,6 ± 4,73	2748,6 ± 0,36	64%	0%	Nephropathy neg.
37,8 ± 1,92	2986,6 ± 0,55	74%	23%	Nephropathy neg.
23,3 ± 2,07	3007,4 ± 0,50	65%	9%	Nephropathy neg.
25,9 ± 2,35	3038,3 ± 0,70	46%	0%	Nephropathy neg.
46,0 ± 2,91	3045,4 ± 0,36	59%	5%	Nephropathy neg.
53,3 ± 4,05	3057,2 ± 0,64	76%	9%	Nephropathy neg.
38,9 ± 2,57	3109,0 ± 0,57	88%	14%	Nephropathy neg.
41,9 ± 3,55	3187,6 ± 0,47	71%	14%	Nephropathy neg.

26,6 ± 1,15	3193,6 ± 0,41	61%	0%	Nephropathy neg.
48,3 ± 3,69	3223,8 ± 0,41	88%	18%	Nephropathy neg.
31,7 ± 3,65	3265,1 ± 0,64	93%	41%	Nephropathy neg.
29,5 ± 1,76	3291,0 ± 0,52	81%	23%	Nephropathy neg.
49,2 ± 3,70	3293,1 ± 0,43	91%	14%	Nephropathy neg.
49,9 ± 3,57	3315,0 ± 0,45	67%	5%	Nephropathy neg.
43,3 ± 2,04	3319,9 ± 0,66	86%	23%	Nephropathy neg.
49,1 ± 3,35	3336,7 ± 0,38	63%	9%	Nephropathy neg.
38,5 ± 2,05	3359,9 ± 0,42	98%	41%	Nephropathy neg.
38,5 ± 1,92	3360,1 ± 0,65	98%	20%	Nephropathy neg.
38,5 ± 2,03	3417,1 ± 0,48	95%	45%	Nephropathy neg.
38,5 ± 1,09	3433,3 ± 0,43	92%	41%	Nephropathy neg.
51,6 ± 3,50	3478,9 ± 0,48	74%	5%	Nephropathy neg.
31,7 ± 2,29	3589,7 ± 0,48	73%	18%	Nephropathy neg.
33,2 ± 3,71	3633,4 ± 0,95	80%	18%	Nephropathy neg.
36,0 ± 3,18	3636,6 ± 0,73	58%	0%	Nephropathy neg.
37,9 ± 2,69	3719,5 ± 0,61	67%	9%	Nephropathy neg.
42,0 ± 3,21	3739,7 ± 0,99	73%	14%	Nephropathy neg.
25,8 ± 1,20	3947,3 ± 0,67	92%	32%	Nephropathy neg.
39,4 ± 1,13	4006,6 ± 0,49	62%	5%	Nephropathy neg.
26,0 ± 3,97	4044,9 ± 0,56	78%	14%	Nephropathy neg.
30,5 ± 2,17	4070,4 ± 0,48	57%	5%	Nephropathy neg.
29,5 ± 0,93	4098,6 ± 0,52	86%	32%	Nephropathy neg.
34,3 ± 2,08	4102,5 ± 0,50	77%	14%	Nephropathy neg.
34,7 ± 0,63	4290,7 ± 0,52	76%	18%	Nephropathy neg.
23,5 ± 1,61	4405,8 ± 0,54	51%	0%	Nephropathy neg.
30,4 ± 1,31	4801,5 ± 1,06	65%	0%	Nephropathy neg.
32,4 ± 1,31	4863,8 ± 0,64	67%	5%	Nephropathy neg.
29,5 ± 2,25	5214,0 ± 1,29	51%	0%	Nephropathy neg.
33,0 ± 0,99	6172,0 ± 1,57	65%	0%	Nephropathy neg.
33,2 ± 0,75	6187,8 ± 0,75	95%	45%	Nephropathy neg.
23,8 ± 1,86	9869,7 ± 1,06	69%	14%	Nephropathy neg.

Table 23:

migration time [min]	dt [min]	mass [Da]
15,490396	0,158804	8054,473633
15,803237	0,155143	8765,233398
16,034266	0,174906	1621,9104
16,185061	0,147871	9180,99707
16,645294	0,198704	10045,20703
17,663696	0,165531	10388,81348
17,980883	0,178564	10518,18457
19,917442	0,234131	9220,939453
20,34516	0,170572	1877,789429
20,479975	0,221246	3842,693604
20,519386	0,265078	4747,932617
21,465685	0,217493	4154,003906
21,480436	0,362197	2427,251709
21,804012	0,271715	4240,856445

22,221563	0,191069	4282,796387
22,777784	0,245503	3840,540527
24,304148	0,319715	7556,177734
24,579231	0,291986	879,519653
24,813087	0,224198	1867,731689
25,283239	0,22054	2266,040771
26,177101	0,289898	2172,188721
26,773794	0,352887	2914,05542
26,81407	0,297343	962,591919
28,254925	0,581783	4353,585938
28,825331	0,258778	1250,62439
29,308136	0,852391	1060,239014
29,822325	0,595913	1682,720947
30,75272	0,175961	943,492859
30,762201	0,263861	1108,647949
30,926645	0,138075	1368,781738
31,305229	0,301605	3987,548828
31,433071	0,515308	1099,419434
32,165497	0,198377	3122,730713
32,222111	0,226858	1829,089966
33,427856	0,151562	2767,015625
34,053886	0,252424	1302,691772
34,15913	0,233032	3722,875977
34,557327	0,186137	2039,143433
34,681156	0,20976	3685,918213
35,30254	0,207782	2389,097168
35,502213	0,388916	3209,800293
36,314056	0,183495	980,526123
36,404907	0,145751	1008,513733
36,424831	0,150486	1000,48761
36,720509	0,128397	2717,472656
36,777012	0,164648	2663,246826
37,557594	0,165628	3556,580566
37,572525	0,185484	1743,890381
37,680653	0,160958	1134,580566
37,700241	0,171622	4097,981934
38,050472	0,156383	3152,361572
38,155159	0,217341	2825,309082
38,17057	0,432096	882,532654
38,281631	0,20781	996,190369
38,57658	0,370648	1425,324829
38,687305	0,15052	3385,513916
38,830559	0,056085	1352,824097
38,921108	0,150325	5000,982422
39,241917	0,178206	3775,720459
39,433277	0,235333	3405,60791
39,484215	0,140887	1046,52771
39,513248	0,093703	2154,053955
39,936756	0,195951	6171,129395
40,533363	0,158628	1194,581543
40,537457	0,221485	2205,064941
40,607231	0,426674	1235,384888

40,686531	0,122381	1265,634888
40,83009	0,191972	2642,264893
41,506096	0,161887	4159,304199
41,604115	0,217324	1250,585449
41,818069	0,163642	2742,253418
42,079609	0,266392	1463,643311
42,105633	0,172054	1489,658936
42,131275	0,184863	1473,643555
42,144161	0,162716	1451,710938
42,573879	0,234118	3098,450928
42,636433	0,041732	1487,660034
42,811199	0,246696	1579,670776
42,940624	0,1884	3121,243164
43,093792	0,106392	3271,523438
43,115334	0,607341	1834,878052
43,46143	0,193155	3442,135498
43,494144	0,20218	3495,841797
43,549488	0,217899	3473,905029
43,740391	0,12795	3108,919434
44,191006	0,18629	3359,583496
44,230297	0,233319	3416,526611
44,934914	0,127421	1991,917114
45,538418	0,214716	2197,337158
45,675098	0,12333	1889,864502
46,313114	0,259721	2385,597168
47,216648	0,168651	2649,602539
47,279705	0,127824	2343,072998
47,526871	0,19233	2584,635986
48,441795	0,239347	1160,526001
48,804813	0,251244	1261,53125
49,519478	0,243133	1274,625244
51,416531	0,33207	1195,518677
51,492035	0,213235	1211,559204
51,657627	0,822884	1223,348633
53,168346	0,293424	1351,643433
53,240913	0,216809	1367,655151
53,259499	0,15916	1770,30481
54,59832	0,234281	1507,742432
55,038143	0,329349	1594,211426
57,475471	0,325805	1840,810547
58,191887	0,129	2021,900879
58,898354	0,484288	2608,239746
60,082333	0,507699	1863,939453

Table 24:

migration time [min]	dt [min]	mass [Da]
12,295616	0,092835	8053,516
12,331619	0,12201	1621,946
12,508785	0,139706	8765,729
12,696615	0,122507	9181,114

12,906662	0,115952	10046,58
13,103853	0,041984	2427,001
14,332394	0,144029	4153,814
14,426023	0,131007	4240,702
14,496774	0,385605	3841,615
14,585806	0,105399	4282,281
15,094264	0,132582	879,5324
15,123884	0,069059	1868,033
15,236325	0,136994	7555,679
15,641728	0,159929	962,6218
16,194395	0,167525	1060,664
16,280394	0,28676	4353,476
16,363562	0,082856	1682,889
16,427116	0,09725	1743,982
16,50071	0,133981	1108,646
16,904119	0,128321	1829,115
17,017418	0,23895	3987,366
17,409172	0,158398	2767,263
17,716999	0,1571	1302,722
17,891594	0,202698	3722,962
18,049681	0,191037	2039,257
18,140236	0,176836	3686,508
18,528196	0,076336	3209,884
19,106394	0,148381	1008,572
19,118612	0,156251	1000,564
19,173443	0,122128	980,5635
19,335644	0,098819	2663,262
19,367334	0,112794	2718,314
20,023649	0,199337	3556,408
20,041323	0,195353	1134,629
20,063593	0,224531	4098,26
20,300522	0,113124	3152,333
20,347666	0,200969	882,5596
20,470793	0,208903	2825,334
20,889994	0,26629	3385,819
20,93943	0,057322	1425,772
21,519066	0,226397	5000,98
21,655712	0,298068	3775,697
21,755213	0,316991	1046,586
21,850452	0,518151	3405,871
22,747589	0,302407	1235,601
22,763943	0,277557	1194,603
22,997269	0,34528	1265,661
23,013165	0,225846	2642,188
23,017294	0,551478	6171,03
23,838888	0,406385	1250,653
24,025209	0,261109	2742,267
24,137253	0,135523	1463,693
24,14039	0,158709	1473,664

24,220345	0,206216	1489,686
24,355286	0,409203	1451,684
24,686199	0,240303	3098,376
24,915867	0,332783	1579,718
25,093962	0,214003	3121,259
25,181305	0,33936	3272,276
25,634459	0,407648	3441,958
25,648405	0,344555	3495,801
25,928818	0,283113	3108,66
26,411203	0,355909	3359,75
26,493782	0,341234	3416,324
27,775286	0,346393	2196,686
28,415859	0,219954	2385,565
29,471397	0,2699	2649,791
29,74654	0,131224	2584,214
30,499264	0,32736	1160,556
30,832899	0,269278	1261,477
32,240211	0,415696	1195,543
32,240601	0,406316	1223,53
32,29216	0,268596	1212,024
33,24297	0,403599	1367,633
34,039223	0,467469	1507,75
34,274136	0,432896	1594,746
35,978645	0,326975	1841,202
37,237282	0,110906	2608,186
37,342949	0,6411	1863,833

Table 25:

Sex	Age	Diagnosis	S-creatinine	Proteinuria	Immunosuppression
M	63	FSGS	95	0.02	PS
M	18	FSGS	99	0.05	CsA
M	63	FSGS	93	0.05	PS
F	49	FSGS	80	0.05	CsA+PS
F	23	FSGS	69	0.54	CsA
F	26	FSGS	16	0,7	CsA
F	56	FSGS	80	0.8	-
M	62	FSGS	150	1,9	-
M	26	FSGS	144	4,9	-
F	26	FSGS	150	11.0	CsA+PS
M	69	MGN	128	0.02	CsA
M	62	MGN	91	0.17	-
M	23	MGN	150	0.3	-
M	37	MGN	73	0.33	-
M	43	MGN	82	0.7	PS
M	48	MGN	100	1.0	CsA+PS
F	68	MGN	150	1.0	-
F	21	MGN	80	1.0	CsA+PS
M	44	MGN	118	1.0	CsA
M	45	MGN	93	1.3	-

M	48	MGN	133	2.4	-
M	37	MGN	93	2.6	-
M	78	MGN	99	3.3	-
M	47	MGN	93	3.5	PS
F	34	MGN	80	3.5	CsA+PS
M	66	MGN	132	3.6	-
M	38	MGN	100	4.0	CsA+PS
M	43	MGN	85	5,1	-
F	43	MCD	114	0.01	CsA
M	45	MCD +	93	0.01	-
F	52	MCD +	118	0.01	-
M	52	MCD	93	0.01	-
F	44	MCD +	80	0.02	CsA
M	39	MCD *	93	0.02	-
M	51	MCD	93	0.05	-
M	18	MCD	77	0.05	CsA+PS
F	70	MCD *	95	0.08	-
M	69	MCD	93	0,08	-
F	29	MCD +	160	0.1	-
M	62	MCD +	93	0.1	-
M	21	MCD	57	0.12	-
F	43	MCD	114	0,01	CSA
F	25	MCD	80	1,2	-
M	52	MCD	93	0.4	PS
F	80	MCD *	145	7.9	-

Claims

1. Use of the presence of at least one polypeptide marker in a urine sample for the diagnosis of a renal disease, wherein the polypeptide marker is selected from the group of polypeptide markers as shown in tables 1 to 22.

2. The use according to claim 1, wherein the renal disease is chosen from the group consisting of IgA-nephropathy, MGN, MCD, FSGS, and diabetic nephropathy.

3. The use according to claim 1, wherein the renal disease is IgA-nephropathy.

4. The use according to claim 1, wherein diagnosis relates to differential diagnosis between at least two diseases chosen from the group consisting of IgA-nephropathy, MGN, MCD, FSGS, and diabetic nephropathy.

5. The use according to claim 1, wherein the polypeptide marker is selected from the group of polypeptide markers as shown in tables 1 to 13.

6. The use according to claim 3, wherein the polypeptide marker is selected from the group of polypeptide markers as shown in tables 11 to 13.

7. A method for the diagnosis of a renal disease, the method comprising:

   a) measuring the presence or the absence of a polypeptide marker in a urine sample, wherein the polypeptide marker is selected from the group of polypeptide markers shown in tables 1 to 22, and
   b) comparing the probability of the presence of this marker in a disease patient to the probability of the presence of this marker in a control patient, wherein
   c1) if the probability of the presence of this marker in a disease patient is higher than the probability of the presence of this marker in a control patient, the presence of this marker is indicative for a higher probability of having the disease rather than the control condition, or
   c2) if the probability of the presence of this marker in a disease patient is lower than the probability of the presence of this marker in a control patient, the absence of the marker is indicative for a higher probability of having the disease rather than the con-

trol condition.

8. The method according to claim 7, wherein the individual probabilities in step b) are as indicated in the tables.

9. The method according to claim 7, wherein the control represents the healthy condition.

10. The method according to claim 7, wherein the control represents a renal disease, particularly chosen from the group consisting of IgA-nephropathy, MGN, MCD, FSGS, and diabetic nephropathy.

11. The method according to claim 7, wherein the polypeptide marker is selected from the group of polypeptide markers shown in tables 11 to 13.

12. The method according to claim 7, wherein the method comprises detecting a plurality of the polypeptide markers, preferably at least 3, more preferably at least 10, most preferably at least 50 of the polypeptide markers.

13. The method according to claim 7, wherein ELISA, quantitative Western Blot, radio-immunoassay, surface plasmon resonance, array, gel electrophoresis, capillary electrophoresis, gas phase ion spectrometry, or mass spectrometry is used for detecting the presence of the marker or markers.

14. The method according to claim 7, wherein the polypeptide markers in the sample are separated by capillary electrophoresis before measurement.

15. The method according to claim 14, wherein mass spectrometry is used for detecting the presence of the marker or markers.

16. Use of capillary electrophoresis-mass spectrometry for the diagnosis, particularly differential diagnosis, of a renal disease in vitro.

17. The use according to claim 16, wherein the renal disease is selected from the group consisting of IgA-nephropathy, MGN, MCD, FSGS, and diabetic nephropathy.

18. Use of capillary electrophoresis-mass spectrometry for the differential diagnosis between at least two renal diseases selected from the group consisting of IgA-nephropathy, MGN, MCD, FSGS, and diabetic nephropathy.