

HETEROARYL DERIVATIVE OR PHARMACEUTICALLY
ACCEPTABLE SALT THEREOF, PREPARATION METHOD THEREFOR,
AND PHARMACEUTICAL COMPOSITION FOR PREVENTING OR
TREATING DISEASES ASSOCIATED WITH PI3 KINASES,
5 CONTAINING SAME AS ACTIVE INGREDIENT

BACKGROUND OF THE INVENTION

1. Field of the Invention

10 The present invention relates to a heteroaryl
derivatives or a pharmaceutically acceptable salt
thereof, a preparation method thereof, and a
pharmaceutical composition comprising the same as an
active ingredient for the prevention or treatment of
15 PI3 kinase related diseases.

2. Description of the Related Art

As indicated in the fluid mosaic model, the
eukaryotic cell membrane is not even and floated or
20 anchored to form a specialized compartment, which is
called a lipid raft. This lipid raft is rich in
cholesterol so as to prevent lysis of the membrane by
a detergent. Some proteins prefer lipid attachment
than being hydrophobic transmembrane anchored.
25 Phosphatidyl inositol is an intracellular protein

found in the lipid raft on cell membrane, which is attached on cell membrane by fatty acid or prenyl link. The lipid raft is very dynamic and can coagulate proteins to make them very active.

5

The protein phosphorylation mediated by kinase is an important way for a cell to regulate physiological activities. The activity of many enzymes is regulated by the phosphorylation mediated by kinase. Another
10 important role of the kinase-mediated phosphorylation is to provide a protein binding site. That is, other proteins are gathered together in the binding site of the phosphorylated area and bound together there without changing the inherent characteristics of the
15 phosphorylated protein. Many kinases involved in signal transduction are found in the lipid raft on the intracellular surface of cell membrane. When a membrane-associated protein is phosphorylated by the activation of a cell surface receptor, this
20 phosphorylated area becomes the protein binding site for those proteins floating alone. These target proteins are not active when they are floating alone without being attached in cytoplasm but become phosphorylated and active when they are gathered

together in the binding site to increase their concentration there.

Phosphatidylinositol 3-kinase (PI3 kinase; PI3K) is the lipid kinase that is responsible for the phosphorylation of a lipid molecule instead of a protein, which plays an important role in cell survival, signal transduction, and control of membrane trafficking. Once there is a problem in any of these regulations, a disease such as cancer, inflammatory disease, and autoimmune disease is developed.

Signal transduction in the cell through 3'-phosphorylated phosphoinositide is associated with various cellular processes such as malignant transformation, growth factor signaling, inflammation, and immunity. PI3 kinase playing a role in producing the phosphorylated signal transduction product was first identified in the course of investigation of the interaction between the viral oncoprotein that induced the phosphorylation of phosphatidylinositol (PI) and its phosphorylated derivative at 3-OH of inositol ring and the growth factor receptor tyrosin kinase.

Phosphatidylinositol-3,4,5-triphosphate (PIP3), the primary product of the PI3 kinase activation, is

up-regulated by treating cells with various stimuli which are exemplified by growth factor and inflammatory stimuli, hormone, neurotransmitter, and antigen receptor mediated signal transduction, etc.

5 So, the activation of PI3 kinase, even though it is not dominant, presents one of the cell surface receptor activation associated signal transductions in mammals. Therefore, the PI3 kinase activation is believed to be involved in a variety of cell responses

10 including cell growth, migration, differentiation, and apoptosis.

PI3 kinase is the enzyme to phosphorylate the 3rd position (3-OH) of the inositol ring moiety of phosphatidylinositol by using ATP (adenosine

15 triphosphate). Precisely, PI3 kinase phosphorylates 3'-OH of inositol ring of phosphatidylinositide to change PIP2 into PIP3. This PIP3 is then functioning as a binding site for the protein kinases containing pleckstrin homology. These protein kinases regulate

20 important cell functions one another. Among the PIP3 binding protein kinases, the most important one is the serine/threonine kinase that is AKT or PKB (protein kinase B), which regulates cell growth, cell survival, and cell division via its downstream mTOR, GSK3 β , Foxo

25 3a, p70S6K, and NF- κ B.

It was confirmed by the primary purification and molecular cloning of PI3 kinase that PI3 kinase is a heterodimer composed of p85 and p110 subunits. Considering sequence homology and substrate
5 specificity, it belongs to class I and class I is also classified into class IA and class IB.

Class IA includes PI3K α , PI3K β , and PI3K δ , and class IA is the downstream of RTK (receptor tyrosine kinase). Class IB represents PI3K γ , and is the
10 downstream of G protein coupled receptor. Each class is composed of the 110 kDa catalytic subunit and the control subunit.

More particularly, three catalytic subunits, p110 α , p110 β , and p110 δ , contain ATP binding domain
15 and interact with the control subunit p85 equally and also are activated by receptor tyrosine kinase. In the meantime, PI3K γ interacts with another control subunit p101 and is activated by the heterotrimer G-protein. Control domain includes the domain for
20 anchoring on the cell surface receptor.

When ATP binding is inhibited, PIP2 phosphorylation is suppressed and therefore PIP3 is not generated. As a result, an important regulatory protein such as AKT cannot be functioning for
25 anchoring on the cell membrane. So, it is an

important target of drug development to inhibit such a catalytic subunit and its ATP binding site.

As explained hereinafter, the expression pattern
5 of each PI3K differs from each other in human cells and tissues. PI3K α and PI3K β display a wide tissue distribution, while PI3K γ is mainly found in white blood cells and also found in skeleton muscle, liver, pancreas, and heart. PI3K δ is only expressed in
10 spleen, thymus, and peripheral blood lymphocytes. The expression pattern above indicates that PI3K α and PI3K β are deeply involved in cancer, while PI3K γ and PI3K δ are rather associated with adaptive immune system such as rheumatoid arthritis (RA), systemic
15 lupus erythematosus (SLE), and hematological malignance.

Particularly, p110 α mutation has been found in some solid tumors. For example, α amplification has been confirmed to be associated with 50% of ovarian
20 cancer, cervical cancer, lung cancer, and breast cancer, and hyperactivation has been confirmed to be associated with at least 50% of colon cancer and at least 25% of breast cancer. P110 β is confirmed to be involved in thrombopoiesis and the p110 γ associated
25 compounds have been developed as an immunosuppressant

for autoimmune disease. The autoimmune disease herein includes rheumatoid arthritis and systemic lupus erythematosus.

Pl10 δ plays a key role in B and T cell
5 activation. Further, δ is also partly involved in
neutrophil migration and sudden increase of
respiration. It is also confirmed that δ can
partially interrupt the antigen-IgE mediated mast cell
degranulation. Pl10 δ draws our attention as an
10 important mediator for not only autoimmune disease and
allergic reaction but also multiple major inflammatory
reactions including abnormal inflammatory diseases.
Target evaluation data for pl10 δ have been filed up
from the studies using the genetic tools and
15 pharmacological agonists, which support the
confirmation above. The inhibition of δ results in
the significant improvement of inflammation and the
related disease, confirmed in the ovalbumin induced
airway inflammation murine asthma model. Rituximab
20 and Belimumab, the PI3K δ monoclonal antibodies, are
effective in RA and SLE.

It is also disclosed recently that PI3K is
involved in the inflammation in the lung and the ear.
25 The mechanism involved in there is not completely

explained yet the over-expressed p110 δ -AKT-mTOR pathway increases aerobic glycolysis but decreases the function and survival of lymphocytes, leading to the decrease of immune response.

5 Chronic inflammation is not unique in autoimmune disease, but PI3K δ and phosphorylated-AKT are characteristically up-regulated in chronic obstructive pulmonary disease (COPD). The over-expression of PI3K δ and phosphorylated-AKT is not only associated
10 with immune disease but also associated with inflammation.

Therefore, it is suggested that the inhibition of PI3K δ can be effective not only in treating autoimmune disease such as rheumatoid arthritis (RA) and systemic
15 lupus erythematosus (SLE) but also in treating chronic non-autoimmune disease such as chronic obstructive pulmonary disease (COPD).

It has been recently reported that a novel
20 compound with a novel structure to inhibit selectively PI3 kinase has been developed. Precisely, patent reference 1 presents a compound useful for cancer treatment which displays the PI3K enzyme inhibiting activity. Patent reference 2 describes that 4-

morpholino-substituted bicyclic heteroaryl compound
has the PI3K activity inhibiting effect.

Thus, the present inventors tried to develop a
5 novel compound with a novel structure that is
effective in inhibiting PI3 kinase selectively. In
the course of the study, the inventors confirmed that
the heteroaryl derivative with a specific structure
had the activity of inhibiting PI3K α , β , δ , and γ
10 selectively and particularly was more excellent in
inhibiting PI3K δ and γ . Therefore, the inventors
confirmed that the heteroaryl derivative can be
effectively used as a pharmaceutical composition for
the prevention and treatment of PI3 kinase related
15 diseases, leading to the completion of this invention.

【PRIOR ART REFERENCE】

【PATENT REFERENCE】

(Patent Reference 1) International Patent
20 Publication No. 2004/048365

(Patent Reference 2) European Patent No.
1,277,738

SUMMARY OF THE INVENTION

25

It is an object of the present invention to provide a heteroaryl derivative, an optical isomer thereof, or a pharmaceutically acceptable salt of the same.

5 It is another object of the present invention to provide a method for preparing the said heteroaryl derivative, the optical isomer thereof, or the pharmaceutically acceptable salt of the same.


It is also an object of the present invention to
10 provide a pharmaceutical composition for the prevention or treatment of PI3 kinase related diseases which comprises the said heteroaryl derivative, the optical isomer thereof, or the pharmaceutically acceptable salt of the same as an active ingredient.

15 It is further an object of the present invention to provide a health food composition for the prevention or improvement of PI3 kinase related diseases which comprises the said heteroaryl derivative, the optical isomer thereof, or the
20 pharmaceutically acceptable salt of the same as an active ingredient.

To achieve the above objects, the present invention provides the compound represented by formula

1, the optical isomer thereof, or the pharmaceutically acceptable salt of the same:

[Formula 1]



5 In the formula 1,

----- is single bond or double bond;


A is carbon (C) or nitrogen (N);

R¹ is hydrogen (H), -NH₂, or C₁₋₅ straight or branched alkylthio;

10 R² is H, -CN, C₁₋₅ straight or branched alkyl, unsubstituted C₃₋₇ cycloalkyl or halogen;

R³ and R⁴ are independently H or C₁₋₅ straight or branched alkyl; or

R³ and R⁴ can form 5 ~ 7 membered unsubstituted
15 heterocycloalkyl containing one or more hetero atoms of N along with the atoms which are conjugated to the same; and



R⁶ is unsubstituted or substituted C₆₋₁₀ aryl or
 5 unsubstituted or substituted 5 ~ 10 membered
 heteroaryl containing one or more hetero atoms
 selected from the group consisting of N, O, and S. In
 the said substituted aryl and the substituted
 heteroaryl, one or more substituents selected from the
 10 group consisting of halogen, C₁₋₅ straight or branched
 alkyl, and C₁₋₅ straight or branched alkylsulfonyl can
 be substituted,

R⁷ and R⁸ are independently H, halogen, -CN, -OH,
 unsubstituted or substituted C₆₋₁₀ aryl, unsubstituted
 15 or substituted 5 ~ 10 membered heteroaryl containing
 one or more hetero atoms selected from the group
 consisting of N, O, and S, C₁₋₅ straight or branched
 alkyl, C₁₋₅ straight or branched alkoxy, C₁₋₅ straight
 or branched alkyloxyalkyl, C₁₋₅ straight or branched
 20 alkylsulfonyl, C₁₋₅ straight or branched alkylthio, or

-NR⁹R¹⁰, wherein R⁹ and R¹⁰ are independently H, C₁₋₅ straight or branched alkyl, C₁₋₅ straight or branched alkylamino, unsubstituted or substituted C₆₋₁₀ aryl, unsubstituted or substituted 5 ~ 10 membered
5 heteroaryl containing one or more hetero atoms selected from the group consisting of N, O, and S, or unsubstituted or substituted 3 ~ 8 membered heterocycloalkyl containing one or more hetero atoms selected from the group consisting of N, O, and S. In
10 the said substituted C₆₋₁₀ aryl, the substituted 5 ~ 10 membered heteroaryl, and the substituted 3 ~ 8 membered heterocycloalkyl, one or more substituents selected from the group consisting of halogen and C₁₋₅ straight or branched alkyl can be substituted.

15

The present invention also provides a method for preparing the compound represented by formula 1 comprising the following steps as shown in the below reaction formula 1:

20 preparing the compound represented by formula 2A by reacting the compound represented by formula 2 and the compound represented by formula 3 (step 1);

preparing the compound represented by formula 5 by reacting the compound represented by formula 2A

prepared in step 1 and the compound represented by formula 4 (step 2);

preparing the compound represented by formula 7 by reacting the compound represented by formula 5 prepared in step 2 and the compound represented by formula 6 (step 3);

preparing the compound represented by formula 8 by reacting the compound represented by formula 7 prepared in step 3 and the compound represented by formula 2B under basic condition (step 4);


preparing the compound represented by formula 10 by reacting the compound represented by formula 8 prepared in step 4 and the compound represented by formula 9 (step 5);

preparing the compound represented by formula 11 by reacting the compound represented by formula 10 prepared in step 5 under acidic condition (step 6);

preparing the compound represented by formula 12 by reacting the compound represented by formula 11 prepared in step 6 and the compound represented by formula 2C (step 7); and

preparing the compound represented by formula 1a by eliminating the amine protecting group from the compound represented by formula 12 prepared in step 7 under acidic condition (step 8):

[Reaction Formula 1]



In the reaction formula 1,

PG is amine protecting group;

the compound represented by formula 1a is a derivative of the compound represented by formula 1, in which --- is double bond and A is carbon, and R^1 , R^2 , R^3 , R^4 , and R^5 are as defined in formula 1.

The present invention also provides a method for preparing the compound represented by formula 1 comprising the following steps as shown in the below reaction formula 2:

preparing the compound represented by formula 15
by reacting the compound represented by formula 2A and
the compound represented by formula 14 (step 1);

preparing the compound represented by formula 17
5 by reacting the compound represented by formula 15
prepared in step 1 and the compound represented by
formula 16 (step 2);


preparing the compound represented by formula 18
by reacting the compound represented by formula 17
10 prepared in step 2 and the compound represented by
formula 2B (step 3);

preparing the compound represented by formula 20
by reacting the compound represented by formula 18
prepared in step 3 and the compound represented by
15 formula 19 (step 4);

preparing the compound represented by formula 21
by reacting the compound represented by formula 20
prepared in step 4 and the compound represented by
formula 2C under basic condition (step 5); and

20 preparing the compound represented by formula 1b
by eliminating the amine protecting group from the
compound represented by formula 21 prepared in step 5
under acidic condition (step 6):

[Reaction Formula 2]



In the reaction formula 2,

PG is amine protecting group;

the compound represented by formula 1b is a
 5 derivative of the compound represented by formula 1,
 in which --- is single bond and A is nitrogen, and R^1 ,
 R^2 , R^3 , R^4 , and R^5 are as defined in formula 1.

The present invention also provides a method for
 10 preparing the compound represented by formula 1
 comprising the following steps as shown in the below
 reaction formula 3:


preparing the compound represented by formula 15
 by reacting the compound represented by formula 2A and
 15 the compound represented by formula 14 (step 1);

preparing the compound represented by formula 23
 by reacting the compound represented by formula 15

prepared in step 1 and the compound represented by formula 22 (step 2); and

preparing the compound represented by formula 1c by reacting the compound represented by formula 23 prepared in step 2 and the compound represented by formula 2C under basic condition (step 3):

[Reaction Formula 3]



In the reaction formula 3,

PG is amine protecting group;

the compound represented by formula 1c is a derivative of the compound represented by formula 1, in which ~~----~~ is double bond and A is nitrogen, and R^1 , R^2 , R^3 , R^4 , and R^5 are as defined in formula 1.

15

The present invention also provides a pharmaceutical composition for the prevention or treatment of PI3 kinase related diseases which comprises the said heteroaryl derivative, the optical isomer thereof, or the pharmaceutically acceptable salt of the same as an active ingredient.

In addition, the present invention provides a health food composition for the prevention or improvement of PI3 kinase related diseases which comprises the said heteroaryl derivative, the optical
5 isomer thereof, or the pharmaceutically acceptable salt of the same as an active ingredient.

ADVANTAGEOUS EFFECT

The heteroaryl derivative of the present invention is excellent in inhibiting PI3 kinase selectively, so that it can be effectively used for the prevention or treatment of PI3 kinase related diseases including cancer such as hematological malignance, ovarian cancer, cervical cancer, breast cancer, colorectal cancer, liver cancer, stomach cancer, pancreatic cancer, colon cancer, peritoneal metastasis, skin cancer, bladder cancer, prostate cancer, thyroid cancer, lung cancer, osteosarcoma, fibrous tumor, and brain tumor; autoimmune disease such as rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis, type 1 diabetes, hyperthyroidism, myasthenia, Crohn's disease, ankylosing spondylitis, psoriasis, autoimmune pernicious anemia, and Sjogren's syndrome; and respiratory disease such as chronic obstructive pulmonary disease(COPD), rhinitis, asthma, chronic bronchitis, chronic inflammatory lung disease, silicosis, pulmonary sarcoidosis, pleurisy, alveolitis, vasculitis, emphysema, pneumonia, and bronchiectasis.


DESCRIPTION OF THE PREFERRED EMBODIMENTS

Hereinafter, the present invention is described in detail.

5

The present invention provides the compound represented by formula 1, the optical isomer thereof, or the pharmaceutically acceptable salt of the same:

[Formula 1]



10

In the formula 1,

— is single bond or double bond;

A is carbon (C) or nitrogen (N);


R¹ is hydrogen (H), -NH₂, or C₁₋₅ straight or
15 branched alkylthio;

R² is H, -CN, C₁₋₅ straight or branched alkyl, unsubstituted C₃₋₇ cycloalkyl or halogen;

when — is double bond and A is N, R² is not existed,

20 R³ and R⁴ are independently H or C₁₋₅ straight or branched alkyl; or

R^3 and R^4 can form 5 ~ 7 membered unsubstituted heterocycloalkyl containing one or more hetero atoms of N along with the atoms which are conjugated to the same; and



R^6 is unsubstituted or substituted C_{6-10} aryl or unsubstituted or substituted 5 ~ 10 membered heteroaryl containing one or more hetero atoms selected from the group consisting of N, O, and S. In the said substituted aryl and the substituted heteroaryl, one or more substituents selected from the group consisting of halogen, C_{1-5} straight or branched alkyl, and C_{1-5} straight or branched alkylsulfonyl can be substituted,

R^7 and R^8 are independently H, halogen, -CN, -OH, unsubstituted or substituted C_{6-10} aryl, unsubstituted or substituted 5 ~ 10 membered heteroaryl containing one or more hetero atoms selected from the group

consisting of N, O, and S, C₁₋₅ straight or branched alkyl, C₁₋₅ straight or branched alkoxy, C₁₋₅ straight or branched alkyloxyalkyl, C₁₋₅ straight or branched alkylsulfonyl, C₁₋₅ straight or branched alkylthio, or
5 -NR⁹R¹⁰, wherein R⁹ and R¹⁰ are independently H, C₁₋₅ straight or branched alkyl, C₁₋₅ straight or branched alkylamino, unsubstituted or substituted C₆₋₁₀ aryl, unsubstituted or substituted 5 ~ 10 membered heteroaryl containing one or more hetero atoms
10 selected from the group consisting of N, O, and S, or unsubstituted or substituted 3 ~ 8 membered heterocycloalkyl containing one or more hetero atoms selected from the group consisting of N, O, and S. In the said substituted C₆₋₁₀ aryl, the substituted 5 ~ 10
15 membered heteroaryl, and the substituted 3 ~ 8 membered heterocycloalkyl, one or more substituents selected from the group consisting of halogen and C₁₋₅ straight or branched alkyl can be substituted.

20 Preferably,

In the formula 1,

~~----~~ is single bond or double bond;


A is carbon (C) or nitrogen (N);

R¹ is H, -NH₂, or methylthio;

R^2 is H, -CN, C_{1-3} straight or branched alkyl, unsubstituted C_{3-5} cycloalkyl or halogen;

R^3 and R^4 are independently H or C_{1-5} straight or branched alkyl; or

5 R^3 and R^4 can form 5 ~ 7 membered unsubstituted heterocycloalkyl containing one or more hetero atoms of N along with the atoms which are conjugated to the same; and



wherein, n is an integer of 0 or 1,

R^6 is unsubstituted or substituted C_{6-10} aryl or unsubstituted or substituted 5 ~ 10 membered heteroaryl containing one or more hetero atoms
 15 selected from the group consisting of N, O, and S. In the said substituted aryl and the substituted heteroaryl, one or more substituents selected from the group consisting of halogen and C_{1-5} straight or branched alkyl can be substituted,

R⁷ is H, halogen, unsubstituted or substituted C₆-
10 aryl, or unsubstituted or substituted 5 ~ 7
membered heteroaryl containing one or more hetero
atoms selected from the group consisting of N, O, and
5 S. In the said substituted aryl and the substituted
heteroaryl, one or more substituents selected from the
group consisting of halogen and C₁₋₅ straight or
branched alkyl can be substituted, and

R⁸ is H, halogen, C₁₋₃ straight or branched alkyl,
10 or C₁₋₃ straight or branched alkoxy.

More preferably,

In the formula 1,

~~----~~ is single bond or double bond;

15 A is carbon (C) or nitrogen (N);


R¹ is H, -NH₂, or methylthio;

R² is H, -CN, C₁₋₃ straight or branched alkyl,
unsubstituted C₃₋₅ cycloalkyl or halogen;

R³ is H;

20 R⁴ is H or C₁₋₃ straight or branched alkyl; or

R³ and R⁴ can form 5 ~ 7 membered unsubstituted
heterocycloalkyl containing one atom of N along with
the atoms which are conjugated to the same; and



More preferably,

In the formula 1,

--- is single bond or double bond;

A is carbon (C) or nitrogen (N);


R¹ is H or -NH₂;

5 R² is H, -F, -Cl, -CN, methyl, ethyl, propyl, isopropyl, cyclopropyl, or cyclopentyl;

R³ is H;

R⁴ is H or methyl; or

R³ and R⁴ can form pyrrolidine along with the
10 atoms which are conjugated to the same; and



wherein, n is an integer of 0 or 1,

R⁶ is unsubstituted or substituted phenyl or
15 pyridinyl,


in the said substituted phenyl and pyridinyl, one or more substituents selected from the group consisting of -F, -Cl, and methyl can be substituted;

R⁷ is H, -F, -Cl, or pyridinyl; and

20 R⁸ is H, -F, or -Cl.

The compound represented by formula 1 herein can be the compound represented by formula 1A, the optical isomer thereof, or the pharmaceutically acceptable
 5 salt of the same.


[Formula 1A]







In the formula 1A,

 , A, R¹, R², R³, R⁴, and R⁵ are as defined in
 10 formula 1.

In the compound represented by formula 1 of the present invention, the ring containing A and R² is







5

The compound represented by formula 1 of the present invention can be exemplified by the following compounds:

<1> 4-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<2> 4-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<3> 4-((1-(5-chloro-4-oxo-3-(pyridine-2-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<4> 4-((1-(5-chloro-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<5> 4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

10 <6> 4-((1-(2-phenylquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<7> 4-((1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<8> 4-((1-(7-fluoro-2-(3-fluorophenyl)quinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

15 <9> 4-(1-(7-fluoro-2-(pyridine-2-yl)quinoline-3-yl)ethylamino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<10> 4-((1-(4,8-dichloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

20 <11> 4-((1-(8-chloro-4-fluoro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<12> 4-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<13> 4-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrido[2,3-d]pyrimidine-5(8H)-one;

<14> 4-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)pyrido[2,3-d]pyrimidine-5(8H)-one;

10 <15> 2-amino-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<16> 2-amino-4-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)pyrido[2,3-d]pyrimidine-5(8H)-one;

15 <17> 4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-6-methylpyrido[2,3-d]pyrimidine-5(8H)-one;

<18> 2-amino-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-6-methylpyrido[2,3-d]pyrimidine-5(8H)-one;

20 <19> 4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-5-oxo-5,8-dihydropyrido[2,3-d]pyrimidine-6-carbonitrile;

- <20> 4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-6-fluoropyrido[2,3-d]pyrimidine-5(8H)-one;
- <21> 4-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)-6-fluoropyrido[2,3-d]pyrimidine-5(8H)-one;
- <22> 6-chloro-4-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
- 10 <23> 6-chloro-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
- <24> 6-chloro-4-((1-(4,8-dichloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
- 15 <25> 2-amino-4-((1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
- <26> 4-((1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
- 20 <27> 4-((1-(6-fluoro-3,4-di(pyridine-2-yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<28> 4-((1-(6-fluoro-3-phenyl-4-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<29> 4-((1-(6-fluoro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<30> 4-((1-(6-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

10 <31> 4-((1-(6-fluoro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<32> 4-((1-(5-chloro-3-(2-chlorobenzyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

15 <33> 4-((1-(6-fluoro-4-oxo-3-(pyridine-2-ylmethyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<34> 4-((1-(5-chloro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

20 <35> 5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<36> 5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<37> 5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<38> 5-((1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

10 <39> 5-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<40> 3-methyl-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

15 <41> 3-methyl-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<42> 5-((1-(4,8-dichloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

20 <43> 5-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<44> 5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<45> 5-(2-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<46> 5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

10 <47> 5-(2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<48> 5-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

15 <49> 5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

20 <50> 7-amino-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<51> 7-amino-5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

25

<52> 7-amino-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<53> 7-amino-5-((1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<54> 7-amino-5-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

10 <55> 7-amino-3-methyl-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<56> 7-amino-5-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

15 <57> 7-amino-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<58> 7-amino-5-(2-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

20 <59> 7-amino-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<60> 7-amino-5-(2-(5-chloro-4-oxo-3-(m-tolyl)-
 3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-
 methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<61> 7-amino-5-(2-(8-chloro-1-oxo-2-phenyl-1,2-
 5 dihydroisoquinoline-3-yl)pyrrolidine-1-yl)-3-methyl-
 2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<62> 5-(1-(8-chloro-1-oxo-2-phenyl-1,2-
 dihydroisoquinoline-3-yl)ethylamino)-3-ethyl-2,3-
 dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

10 <63> 5-(1-(8-chloro-1-oxo-2-phenyl-1,2-
 dihydroisoquinoline-3-yl)ethylamino)-3-propyl-2,3-
 dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<64> 5-(1-(8-chloro-1-oxo-2-phenyl-1,2-
 dihydroisoquinoline-3-yl)ethylamino)-3-cyclopropyl-
 15 2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<65> 5-(1-(8-chloro-1-oxo-2-phenyl-1,2-
 dihydroisoquinoline-3-yl)ethylamino)-3-cyclopentyl-
 2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<66> 5-(1-(8-chloro-1-oxo-2-phenyl-1,2-
 20 dihydroisoquinoline-3-yl)ethylamino)-3-isopropyl-2,3-
 dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<67> 5-(1-(5-fluoro-4-oxo-3-phenyl-3,4-
 dihydroquinazoline-2-yl)propylamino)-3-isopropyl-2,3-
 dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<68> 5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<69> 5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<70> 5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

10 <71> 5-((1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<72> 5-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

15 <73> 5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<74> 5-((1-(2-phenylquinoline-3-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

20 <75> 5-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<76> 5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<77> 5-(2-(8-chloro-1-oxo-2-(pyridine-3-yl)-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<78> 5-(2-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

10 <79> 5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<80> 5-(2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

15 <81> 5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<82> 5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

20 <83> 7-amino-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<84> 7-amino-5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;
 <85> 7-amino-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;
 <86> 7-amino-5-((1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;
 10 <87> 7-amino-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)pyrimido[4,5-d]pyrimidine-4(1H)-one;
 <88> 7-amino-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;
 15 <89> 7-amino-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;
 <90> 7-amino-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;
 20 <91> 7-amino-5-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<92> 7-amino-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<93> 7-amino-5-(2-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<94> 7-amino-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

10 <95> 7-amino-5-(2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<96> 7-amino-5-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<97> 7-amino-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<98> 7-amino-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<99> 4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-7,8-dihydropyrido[2,3-d]pyrimidine-5(6H)-one;

<100> 4-((1-(4,8-dichloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-7,8-dihydropyrido[2,3-d]pyrimidine-5(6H)-one.

5 The compound represented by formula 1 of the present invention can be preferably exemplified by the following optical isomer compounds:

<1> (S)-4-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<2> (S)-4-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<3> (S)-4-((1-(5-chloro-4-oxo-3-(pyridine-2-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<4> (S)-4-((1-(5-chloro-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

20 <5> (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<6> (S)-4-((1-(2-phenylquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<7> (S)-4-((1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<8> (S)-4-((1-(7-fluoro-2-(3-fluorophenyl)quinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<9> (S)-4-(1-(7-fluoro-2-(pyridine-2-yl)quinoline-3-yl)ethylamino)pyrido[2,3-d]pyrimidine-5(8H)-one;

10 <10> (S)-4-((1-(4,8-dichloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<11> (S)-4-((1-(8-chloro-4-fluoro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

15 <12> (S)-4-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<13> (S)-4-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrido[2,3-d]pyrimidine-5(8H)-one;

20 <14> (S)-4-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)pyrido[2,3-d]pyrimidine-5(8H)-one;

- <15> (S)-2-amino-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
- <16> (S)-2-amino-4-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)pyrido[2,3-d]pyrimidine-5(8H)-one;
- <17> (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-6-methylpyrido[2,3-d]pyrimidine-5(8H)-one;
- 10 <18> (S)-2-amino-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-6-methylpyrido[2,3-d]pyrimidine-5(8H)-one;
- <19> (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-5-oxo-5,8-dihydropyrido[2,3-d]pyrimidine-6-carbonitrile;
- 15 <20> (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-6-fluoropyrido[2,3-d]pyrimidine-5(8H)-one;
- <21> (S)-4-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)-6-fluoropyrido[2,3-d]pyrimidine-5(8H)-one;
- 20 <22> (S)-6-chloro-4-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<23> (S)-6-chloro-4-((1-(8-chloro-1-oxo-2-phenyl-
 1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-
 d]pyrimidine-5(8H)-one;

<24> (S)-6-chloro-4-((1-(4,8-dichloro-1-oxo-2-
 5 phenyl-1,2-dihydroisoquinoline-3-
 yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<25> (S)-2-amino-4-((1-(6-fluoro-3-(pyridine-2-
 yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-
 5(8H)-one;

10 <26> (S)-4-((1-(6-fluoro-3-(pyridine-2-
 yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-
 5(8H)-one;

<27> (S)-4-((1-(6-fluoro-3,4-di(pyridine-2-
 yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-
 15 5(8H)-one;

<28> (S)-4-((1-(6-fluoro-3-phenyl-4-(pyridine-2-
 yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-
 5(8H)-one;

<29> (S)-4-((1-(6-fluoro-4-oxo-3-(pyridine-3-yl)-
 20 3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-
 d]pyrimidine-5(8H)-one;

<30> (S)-4-((1-(6-fluoro-4-oxo-3-phenyl-3,4-
 dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-
 d]pyrimidine-5(8H)-one;

<31> (S)-4-((1-(6-fluoro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
 <32> (S)-4-((1-(5-chloro-3-(2-chlorobenzyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
 <33> (S)-4-((1-(6-fluoro-4-oxo-3-(pyridine-2-ylmethyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
 10 <34> (S)-4-((1-(5-chloro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
 <35> (S)-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;
 15 <36> (S)-5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;
 <37> (S)-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;
 20 <38> (S)-5-((1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<39> (S)-5-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<40> (S)-3-methyl-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<41> (S)-3-methyl-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

10 <42> (S)-5-((1-(4,8-dichloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<43> (S)-5-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

15 <44> (S)-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<45> (S)-5-(2-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

20 <46> (S)-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<47> (S)-5-(2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<48> (S)-5-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<49> (S)-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<50> (S)-7-amino-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<51> (S)-7-amino-5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<52> (S)-7-amino-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<53> (S)-7-amino-5-((1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<54> (S)-7-amino-5-((1-(8-chloro-1-oxo-2-phenyl-
 1,2-dihydroisoquinoline-3-yl)ethyl)amino)-3-methyl-
 2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<55> (S)-7-amino-3-methyl-5-((1-(2-
 5 phenylquinoline-3-yl)ethyl)amino)-2,3-
 dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<56> (S)-7-amino-5-((1-(5-fluoro-4-oxo-3-phenyl-
 3,4-dihydroquinazoline-2-yl)propyl)amino)-3-methyl-
 2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

10 <57> (S)-7-amino-5-(2-(5-chloro-4-oxo-3-phenyl-
 3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-
 methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<58> (S)-7-amino-5-(2-(5-chloro-4-oxo-3-
 (pyridine-3-yl)-3,4-dihydroquinazoline-2-
 15 yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-
 d]pyrimidine-4(1H)-one;

<59> (S)-7-amino-5-(2-(5-chloro-3-(3-
 fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-
 yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-
 20 d]pyrimidine-4(1H)-one;

<60> (S)-7-amino-5-(2-(5-chloro-4-oxo-3-(m-
 tolyl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-
 3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-
 one;

<61> (S)-7-amino-5-(2-(8-chloro-1-oxo-2-phenyl-
 1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)-3-
 methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<62> (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-
 5 dihydroisoquinoline-3-yl)ethylamino)-3-ethyl-2,3-
 dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<63> (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-
 dihydroisoquinoline-3-yl)ethylamino)-3-propyl-2,3-
 dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

10 <64> (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-
 dihydroisoquinoline-3-yl)ethylamino)-3-cyclopropyl-
 2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<65> (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-
 dihydroisoquinoline-3-yl)ethylamino)-3-cyclopentyl-
 15 2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<66> (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-
 dihydroisoquinoline-3-yl)ethylamino)-3-isopropyl-2,3-
 dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<67> (S)-5-(1-(5-fluoro-4-oxo-3-phenyl-3,4-
 20 dihydroquinazoline-2-yl)propylamino)-3-isopropyl-2,3-
 dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<68> (S)-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-
 dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-
 d]pyrimidine-4(1H)-one;

<69> (S)-5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-
 3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-
 d]pyrimidine-4(1H)-one;

<70> (S)-5-((1-(5-chloro-3-(3-fluorophenyl)-4-
 5 oxo-3,4-dihydroquinazoline-2-
 yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<71> (S)-5-((1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-
 dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-
 d]pyrimidine-4(1H)-one;

10 <72> (S)-5-((1-(8-chloro-1-oxo-2-phenyl-1,2-
 dihydroisoquinoline-3-yl)ethyl)amino)pyrimido[4,5-
 d]pyrimidine-4(1H)-one;

<73> (S)-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-
 dihydropyrrolo[2,1-f][1,2,4]triazine-2-
 15 yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<74> (S)-5-((1-(2-phenylquinoline-3-
 yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<75> (S)-5-((1-(5-fluoro-4-oxo-3-phenyl-3,4-
 dihydroquinazoline-2-yl)propyl)amino)pyrimido[4,5-
 20 d]pyrimidine-4(1H)-one;

<76> (S)-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-
 dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-
 d]pyrimidine-4(1H)-one;

<77> (S)-5-(2-(8-chloro-1-oxo-2-(pyridine-3-yl)-
 1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-
 yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<78> (S)-5-(2-(5-chloro-4-oxo-3-(pyridine-3-yl)-
 5 3,4-dihydroquinazoline-2-yl)pyrrolidine-1-
 yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<79> (S)-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-
 3,4-dihydroquinazoline-2-yl)pyrrolidine-1-
 yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

10 <80> (S)-5-(2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-
 dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-
 d]pyrimidine-4(1H)-one;

<81> (S)-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-
 dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)pyrrolidine-
 15 1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<82> (S)-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-
 3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-
 yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-
 one;

20 <83> (S)-7-amino-5-((1-(5-chloro-4-oxo-3-phenyl-
 3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-
 d]pyrimidine-4(1H)-one;

<84> (S)-7-amino-5-((1-(5-chloro-4-oxo-3-
 (pyridine-3-yl)-3,4-dihydroquinazoline-2-
 25 yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<85> (S)-7-amino-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<86> (S)-7-amino-5-((1-(5-chloro-4-oxo-3-(m-
 5 tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<87> (S)-7-amino-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

10 <88> (S)-7-amino-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<89> (S)-7-amino-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydropyrrolo[2,1-
 15 f][1,2,4]triazine-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<90> (S)-7-amino-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<91> (S)-7-amino-5-((1-(5-fluoro-4-oxo-3-phenyl-
 20 3,4-dihydroquinazoline-2-yl)propyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<92> (S)-7-amino-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<93> (S)-7-amino-5-(2-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

5 <94> (S)-7-amino-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<95> (S)-7-amino-5-(2-(5-chloro-4-oxo-3-(m-
 10 tolyl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<96> (S)-7-amino-5-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

15 <97> (S)-7-amino-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;


<98> (S)-7-amino-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;
 20

<99> (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-7,8-dihydropyrido[2,3-d]pyrimidine-5(6H)-one;
 25

<100> (S)-4-((1-(4,8-dichloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-7,8-dihydropyrido[2,3-d]pyrimidine-5(6H)-one.

5 The present invention also provides the intermediate compound represented by formula 1B of the compound represented by formula 1, or the optical isomer of the same.

[Formula 1B]



10

In the formula 1B,

---, A, R¹, R², R³, R⁴, and R⁵ are as defined in formula 1; and

PG is an amine protecting group selected from the
 15 group consisting of t-butyloxycarbonyl (Boc), carbobenzyloxy (Cbz), 9-fluorenylmethyloxycarbonyl (Fmoc), acetyl (Ac), benzoyl (Bz), benzyl (Bn), p-methoxybenzyl (PMB), 3,4-dimethoxybenzyl (DMPM), p-methoxyphenyl (PMP), tosyl (Ts), 2,2,2-
 20 trichloroethoxycarbonyl (Troc), 2-

trimethylsilylethoxycarbonyl (Teoc), and
aryloxycarbonyl (Alloc).

The compound represented by formula 1B is an
intermediate of the compound represented by formula 1
5 which can be prepared by eliminating the amine
protecting group of PG.

The compound represented by formula 1 of the
present invention can be used as a form of a
10 pharmaceutically acceptable salt, in which the salt is
preferably acid addition salt formed by
pharmaceutically acceptable free acids. The acid
addition salt herein can be obtained from inorganic
acids such as hydrochloric acid, nitric acid,
15 phosphoric acid, sulfuric acid, hydrobromic acid,
hydriodic acid, nitrous acid, and phosphorous acid;
non-toxic organic acids such as aliphatic
mono/dicarboxylate, phenyl-substituted alkanoate,
hydroxy alkanoate, alkandioate, aromatic acids, and
20 aliphatic/aromatic sulfonic acids; or organic acids
such as acetic acid, benzoic acid, citric acid, lactic
acid, maleic acid, gluconic acid, methanesulfonic
acid, 4-toluenesulfonic acid, tartaric acid, and
fumaric acid. The pharmaceutically non-toxic salts
25 are exemplified by sulfate, pyrosulfate, bisulfate,

sulphite, bisulphite, nitrate, phosphate, monohydrogen
phosphate, dihydrogen phosphate, metaphosphate,
pyrophosphate, chloride, bromide, iodide, fluoride,
acetate, propionate, decanoate, caprylate, acrylate,
5 formate, isobutylate, caprate, heptanoate, propiolate,
oxalate, malonate, succinate, suberate, sebacate,
fumarate, maleate, butyne-1,4-dioate, hexane-1,6-
dioate, benzoate, chlorobenzoate, methylbenzoate,
dinitrobenzoate, hydroxybenzoate, methoxybenzoate,
10 phthalate, terephthalate, benzenesulfonate,
toluenesulfonate, chlorobenzenesulfonate,
xylenesulfonate, phenylacetate, phenylpropionate,
phenylbutylate, citrate, lactate, hydroxybutylate,
glycolate, malate, tartrate, methanesulfonate,
15 propanesulfonate, naphthalene-1-sulfonate,
naphthalene-2-sulfonate, and mandelate.

The acid addition salt in this invention can be
prepared by the conventional method known to those in
the art. For example, the compound represented by
20 formula 1 is dissolved in an organic solvent such as
methanol, ethanol, acetone, methylenechloride, or
acetonitrile, to which organic acid or inorganic acid
is added to induce precipitation. Then, the
precipitate is filtered and dried to give the salt.
25 Or the solvent and the excessive acid are distilled

under reduced pressure, and dried to give the salt.
Or the precipitate is crystallized in an organic
solvent to give the same.

A pharmaceutically acceptable metal salt can be
5 prepared by using a base. Alkali metal or alkali
earth metal salt is obtained by the following
processes: dissolving the compound in excessive alkali
metal hydroxide or alkali earth metal hydroxide
solution; filtering non-soluble compound salt;
10 evaporating the remaining solution and drying thereof.
At this time, the metal salt is preferably prepared in
the pharmaceutically suitable form of sodium,
potassium, or calcium salt. And the corresponding
silver salt is prepared by the reaction of alkali
15 metal or alkali earth metal salt with proper silver
salt (ex; silver nitrate).

The present invention includes not only the
compound represented by formula 1 but also a
pharmaceutically acceptable salt thereof, and a
20 solvate, an optical isomer, or a hydrate possibly
produced from the same.

The present invention also provides a method for
preparing the compound represented by formula 1

comprising the following steps as shown in the below reaction formula 1:

preparing the compound represented by formula 2A by reacting the compound represented by formula 2 and the compound represented by formula 3 (step 1);

preparing the compound represented by formula 5 by reacting the compound represented by formula 2A prepared in step 1 and the compound represented by formula 4 (step 2);

10 preparing the compound represented by formula 7 by reacting the compound represented by formula 5 prepared in step 2 and the compound represented by formula 6 (step 3);

preparing the compound represented by formula 8 by reacting the compound represented by formula 7 prepared in step 3 and the compound represented by formula 2B under basic condition (step 4);


preparing the compound represented by formula 10 by reacting the compound represented by formula 8 prepared in step 4 and the compound represented by formula 9 (step 5);

preparing the compound represented by formula 11 by reacting the compound represented by formula 10 prepared in step 5 under acidic condition (step 6);

preparing the compound represented by formula 12 by reacting the compound represented by formula 11 prepared in step 6 and the compound represented by formula 2C (step 7); and

5 preparing the compound represented by formula 1a by eliminating the amine protecting group from the compound represented by formula 12 prepared in step 7 under acidic condition (step 8):

[Reaction Formula 1]



In the reaction formula 1,

PG is amine protecting group;

the compound represented by formula 1a is a derivative of the compound represented by formula 1,

in which --- is double bond and A is carbon, and R^1 , R^2 , R^3 , R^4 , and R^5 are as defined in formula 1.

Hereinafter, the method for preparing the
5 compound represented by formula 1 of the present invention is described in more detail.

In the method for preparing the compound represented by formula 1 of the present invention, step 1 is to give the compound represented by formula
10 2A by reacting the compound represented by formula 2 with the compound represented by formula 3.

At this time, the compound represented by formula 3 is exemplified by ZnCl_2 , SnCl_2 , SnCl_4 , FeCl_2 , FeCl_3 , and POCl_3 , which can be used as an equivalent or an
15 excess. Among these, POCl_3 is more preferred.

In step 1, a compound capable of providing carbon to form aldehyde of the compound represented by formula 2A is used. The compound that can provide carbon herein is not limited but dimethylformamide is
20 preferred.

In the method for preparing the compound represented by formula 1 of the present invention, step 2 is to give the compound represented by formula 5 by reacting the compound represented by formula 2

prepared in step 1 with the compound represented by formula 4, the Grignard reagent.

In the method for preparing the compound represented by formula 1 of the present invention, 5 step 3 is to give the compound represented by formula 7 by reacting the compound represented by formula 5 prepared in step 2 with the compound represented by formula 6.

Particularly, step 3 is to prepare the aldehyde 10 compound represented by formula 7 by reacting the alcohol compound represented by formula 5 with the oxidant represented by formula 6. At this time, the oxidant represented by formula 6 is exemplified by PCC (pyridinium chlorochromate), PDC (pyridinium 15 dichromate), and CrO_3 , which can be used as an equivalent or an excess. Among these, CrO_3 is more preferred.

In the method for preparing the compound represented by formula 1 of the present invention, 20 step 4 is to give the compound represented by formula 8 by reacting the compound represented by formula 7 prepared in step 3 with the compound represented by formula 2B.

At this time, PG in the compound represented by 25 formula 2B is the amine protecting group and the amine

protecting group is exemplified by t-butyloxycarbonyl (Boc), carbobenzyloxy (Cbz), 9-fluorenylmethyloxycarbonyl (Fmoc), acetyl (Ac), benzoyl (Bz), benzyl (Bn), p-methoxybenzyl (PMB), 3,4-dimethoxybenzyl (DMPM), p-methoxyphenyl (PMP), tosyl (Ts), 2,2,2-trichloroethoxycarbonyl (Troc), 2-trimethylsilylethoxycarbonyl (Teoc), or aryloxycarbonyl (Alloc), and p-methoxybenzyl (PMB) is more preferred.

10 In the method for preparing the compound represented by formula 1 of the present invention, step 5 is to give the compound represented by formula 10 by reacting the compound represented by formula 8 prepared in step 4 with DMF-DMA (dimethylformamide-dimethylacetal), the compound represented by formula 15 9.

In the method for preparing the compound represented by formula 1 of the present invention, step 6 is to give the compound represented by formula 11 by reacting the compound represented by formula 10 20 11 by reacting the compound represented by formula 10 prepared in step 5 under acidic condition.

At this time, the acid is exemplified by HCl, H₂SO₄, bromic acid, and acetic acid, which can be used as an equivalent or an excess. Among these, acetic 25 acid is more preferred.

In the method for preparing the compound represented by formula 1 of the present invention, step 7 is to give the compound represented by formula 12 by reacting the compound represented by formula 11 prepared in step 6 with the compound represented by formula 2C.

Particularly, step 3 is to prepare the compound represented by formula 1 by dehydration-condensation of the compound represented by formula 11 and the compound represented by formula 2C in the presence of (benzotriazole-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate (BOP) and a base.

At this time, the base herein is exemplified by an organic base such as pyridine, triethylamine, N,N-diisopropylethylamine (DIPEA), and 1,8-diazabicyclo[5.4.0]-7-undecene (DBU) or an inorganic base such as sodiumhydroxide, sodiumcarbonate, potassiumcarbonate, cesiumcarbonate, and sodiumhydride, which can be used as an equivalent or an excess independently or mixed together. Among these, 1,8-diazabicyclo[5.4.0]-7-undecene (DBU) is more preferred.

The stereo-selectivity of the compound represented by formula 1 is determined by the stereo-selectivity of the compound represented by formula 2C

used in step 7. Thus, the optical isomer of the compound represented by formula 1 can be prepared by using the optical isomer of the compound represented by formula 2C.

5 In the method for preparing the compound represented by formula 1 of the present invention, step 8 is to give the compound represented by formula 1a by eliminating the amine protecting group of the compound represented by formula 12 prepared in step 7
10 under acidic condition.

At this time, the acid is exemplified by HCl, H₂SO₄, acetic acid, and trifluoroacetic acid, which can be used as an equivalent or an excess. Among these, trifluoroacetic acid is more preferred.

15

In the method for preparing the compound represented by formula 1 of the present invention, each step of the reaction formula 1 can be executed by the conventional method known to those in the art and
20 at this time the usable base is exemplified by such organic bases as pyridine, triethylamine, N,N-diisopropylethylamine (DIPEA), and 1,8-diazabicyclo[5.4.0]-7-undecene (DBU); or such inorganic bases as sodium hydroxide, sodium carbonate,
25 potassium carbonate, cesium carbonate, and sodium

hydride, which can be used as an equivalent or an excess independently or as mixed. The acceptable reaction solvent is exemplified by tetrahydrofuran (TFH); dioxane; ether solvents such as ethylether and 1,2-dimethoxyethane; lower alcohols such as methanol, ethanol, propanol, and butanol; dimethylformamide (DMF), dimethylsulfoxide (DMSO), dichloromethane (DCM), dichloroethane, water, acetone, sulfonates, toluenesulfonate, chlorobenzenesulfonate, xylenesulfonate, ethylacetate, phenylacetate, phenylpropionate, phenylbutylate, citrate, lactate, hydroxybutylate, glycolate, malate, tartrate, methanesulfonate, propanesulfonate, naphthalene-1-sulfonate, naphthalene-2-sulfonate, and mandelate.

The reaction solvent can be added independently or as a mixture of those.

As explained hereinbefore, the preparation method represented by the reaction formula 1 of the invention is not only a novel method for preparing easily the compound represented by formula 11, one of the intermediates of the compound represented by formula 1 but also a useful method for preparing various pyridopyrimidine derivatives from the compound represented by formula 1 by reacting the compound represented by

formula 11 with the compound reacting to hydroxyl group (-OH), the substituent for the compound represented by formula 11.


5 In the preparation method represented by the reaction formula 1 of the invention, the compound represented by formula 1a can be prepared by the method comprising the following steps as shown in reaction formula 1-a:

10 preparing the compound represented by formula 13 by reacting the compound represented by formula 8 prepared in step 4 of reaction formula 1 and the compound represented by formula 2C (step 1);

preparing the compound represented by formula 12
15 by reacting the compound represented by formula 13 prepared in step 1 and the compound represented by formula 9 (step 2); and

preparing the compound represented by formula 1a by eliminating the amine protecting group from the
20 compound represented by formula 12 prepared in step 2 under acidic condition (step 3):

[Reaction Formula 1-a]



In the reaction formula 1-a,


the compound represented by formula 1a is a derivative of the compound represented by formula 1, in which ---- is double bond and A is carbon, and R¹, R², R³, R⁴, and R⁵ are as defined in formula 1.

Further, in the preparation method represented by the reaction formula 1, the following steps can be added to the method in order to prepare the compound having the substitution of R¹ (-SCH₃ → -NH₂), as shown in reaction formula 1-b:

preparing the compound represented by formula 1a" by reacting the compound represented by formula 1a' with mCPBA (3-chlorobenzoic acid) (step 1); and

preparing the compound represented by formula 1a wherein R¹ is -NH₂ by reacting the compound represented by formula 1a" prepared in step 1 in the presence of NH₄OH (step 2):

[Reaction Formula 1-b]



In the reaction formula 1-b,

the compound represented by formula 1a is a derivative of the compound represented by formula 1, in which ~~----~~ is double bond and A is carbon, and R², R³, R⁴, and R⁵ are as defined in formula 1. m is 1 or 2.

The present invention also provides a method for preparing the compound represented by formula 1 comprising the following steps as shown in the below reaction formula 2:

preparing the compound represented by formula 15 by reacting the compound represented by formula 2A and the compound represented by formula 14 (step 1);

preparing the compound represented by formula 17 by reacting the compound represented by formula 15 prepared in step 1 and the compound represented by formula 16 (step 2);

preparing the compound represented by formula 18 by reacting the compound represented by formula 17


prepared in step 2 and the compound represented by formula 2B (step 3);

preparing the compound represented by formula 20 by reacting the compound represented by formula 18 prepared in step 3 and the compound represented by formula 19 (step 4);

preparing the compound represented by formula 21 by reacting the compound represented by formula 20 prepared in step 4 and the compound represented by formula 2C under basic condition (step 5); and

preparing the compound represented by formula 1b by eliminating the amine protecting group from the compound represented by formula 21 prepared in step 5 under acidic condition (step 6):

[Reaction Formula 2]



In the reaction formula 2,

PG is amine protecting group;

the compound represented by formula 1b is a derivative of the compound represented by formula 1, in which ~~----~~ is single bond and A is nitrogen, and R¹, R², R³, R⁴, and R⁵ are as defined in formula 1.

5

Each step of the preparation method represented by the reaction formula 2 is executed by the same or similar method to the method of reaction formula 1 above or the conventional method known to those in the art.

Therefore, the preparation method represented by the reaction formula 2 of the invention is not only a novel method for preparing easily the compound represented by formula 20, one of the intermediates of the compound represented by formula 1 but also a useful method for preparing diverse dihydro pyrimido-pyrimidine derivatives from the compound represented by formula 1 by reacting the compound represented by formula 20 with the compound reacting to chloride (-Cl), the substituent of the compound.


The present invention also provides a method for preparing the compound represented by formula 1 comprising the following steps as shown in the below reaction formula 3:

preparing the compound represented by formula 15 by reacting the compound represented by formula 2A and the compound represented by formula 14 (step 1);

preparing the compound represented by formula 23 by reacting the compound represented by formula 15 prepared in step 1 and the compound represented by formula 22 (step 2); and

preparing the compound represented by formula 1c by reacting the compound represented by formula 23 prepared in step 2 and the compound represented by formula 2C under basic condition (step 3):

[Reaction Formula 3]



In the reaction formula 3,

the compound represented by formula 1c is a derivative of the compound represented by formula 1, in which is double bond and A is nitrogen, and R¹, R², R³, R⁴, and R⁵ are as defined in formula 1.

Each step of the preparation method represented by the reaction formula 3 is executed by the same or similar method to the method of reaction formula 1

above or the conventional method known to those in the art.

Therefore, the preparation method represented by the reaction formula 3 of the invention is not only a novel method for preparing easily the compound represented by formula 23, one of the intermediates of the compound represented by formula 1 but also a useful method for preparing diverse pyrimido-pyrimidine derivatives from the compound represented by formula 1 by reacting the compound represented by formula 23 with the compound reacting to chloride (-Cl), the substituent of the compound.

In the preparation method represented by the reaction formula 3, the compound represented by formula 1c can be prepared by the method comprising the following steps as shown in reaction formula 3-a:

preparing the compound represented by formula 25 by reacting the compound represented by formula 15 prepared in step 1 of reaction formula 3 and the compound represented by formula 24 (step 1);


preparing the compound represented by formula 26 by reacting the compound represented by formula 25 prepared in step 1 and the compound represented by formula 2B (step 2);

preparing the compound represented by formula 29 by reacting the compound represented by formula 26 prepared in step 2 and the compound represented by formula 27 (step 3);

5 preparing the compound represented by formula 30 by reacting the compound represented by formula 29 prepared in step 3 and the compound represented by formula 2C (step 4); and

preparing the compound represented by formula 1b by eliminating the amine protecting group from the
10 compound represented by formula 30 prepared in step 4 under acidic condition (step 5):

[Reaction Formula 3-a]



15 In the reaction formula 3-a,
PG is amine protecting group;

the compound represented by formula 1c is a derivative of the compound represented by formula 1, in which ~~----~~ is double bond and A is nitrogen, and R¹, R², R³, R⁴, and R⁵ are as defined in formula 1.

5

The present invention also provides a pharmaceutical composition for the prevention or treatment of PI3 kinase related diseases which comprises the said heteroaryl derivative, the optical
10 isomer thereof, or the pharmaceutically acceptable salt of the same as an active ingredient.

The heteroaryl derivative, the optical isomer thereof, or the pharmaceutically acceptable salt thereof of the invention is characterized by the
15 selective inhibition of PI3 kinase selected from the group consisting of PI3K α , PI3K β , PI3K δ , and PI3K γ .

Particularly, the PI3 kinase related disease includes cancer, autoimmune disease, and respiratory disease.

20 The cancer herein is exemplified by hematological malignance such as myeloid metaplasia, chronic myelomonocytic leukemia, acute lymphoblastic leukemia, acute erythroid leukemia, Hodgikin`s/non-Hodgikin`s disease, B-cell lymphoma, acute T-cell leukemia,
25 myelodysplastic syndrome, plasma cell dysfunction,

hairy cell leukemia, Kaposi's sarcoma, and lymphoma, ovarian cancer, cervical cancer, breast cancer, colorectal cancer, liver cancer, stomach cancer, pancreatic cancer, colon cancer, peritoneal
5 metastasis, skin cancer, bladder cancer, prostate cancer, thyroid cancer, lung cancer, osteosarcoma, fibrous tumor, and brain tumor.

The autoimmune disease herein includes rheumatoid arthritis, systemic lupus erythematosus, multiple
10 sclerosis, type 1 diabetes, hyperthyroidism, myasthenia, Crohn's disease, ankylosing spondylitis, psoriasis, autoimmune pernicious anemia, and Sjogren's syndrome.

The respiratory disease herein includes chronic
15 obstructive pulmonary disease(COPD), rhinitis, asthma, chronic bronchitis, chronic inflammatory lung disease, silicosis, pulmonary sarcoidosis, pleurisy, alveolitis, vasculitis, emphysema, pneumonia, and bronchiectasis.

20

The present inventors investigated the inhibitory effect of the compound represented by formula 1 of the present invention on PI3K α , β , γ , and δ . As a result, it was confirmed that the compound of the
25 invention was excellent in inhibiting PI3K α , β , γ ,

and δ . In particular, the inhibitory effect on PI3 kinase γ or δ was more peculiar even at a low concentration (see Experimental Examples 1 ~ 4).

Therefore, the compound of the present invention
5 plays a role as a PI3 kinase inhibitor, so that it can be effectively used for the prevention or treatment of PI3 kinase related diseases including cancer such as hematological malignance, ovarian cancer, cervical cancer, breast cancer, colorectal cancer, liver
10 cancer, stomach cancer, pancreatic cancer, colon cancer, peritoneal metastasis, skin cancer, bladder cancer, prostate cancer, thyroid cancer, lung cancer, osteosarcoma, fibrous tumor, and brain tumor; autoimmune disease such as rheumatoid arthritis,
15 systemic lupus erythematosus, multiple sclerosis, type 1 diabetes, hyperthyroidism, myasthenia, Crohn's disease, ankylosing spondylitis, psoriasis, autoimmune pernicious anemia, and Sjogren's syndrome; and respiratory disease such as chronic obstructive
20 pulmonary disease(COPD), rhinitis, asthma, chronic bronchitis, chronic inflammatory lung disease, silicosis, pulmonary sarcoidosis, pleurisy, alveolitis, vasculitis, emphysema, pneumonia, and bronchiectasis.

25

The compound represented by formula 1, the optical isomer thereof, or the pharmaceutically acceptable salt thereof of the present invention can be prepared for oral or parenteral administration by
5 mixing with generally used diluents or excipients such as fillers, extenders, binders, wetting agents, disintegrating agents and surfactants.

The formulations for oral administration are exemplified by tablets, pills, hard/soft capsules,
10 solutions, suspensions, emulsions, syrups, granules, and elixirs, etc. These formulations can include diluents (for example, lactose, dextrose, sucrose, mannitol, sorbitol, cellulose, and/or glycine) and lubricants (for example, silica, talc, stearate and
15 its magnesium or calcium salt, and/or polyethylene glycol) in addition to the active ingredient. Tablets can include binding agents such as magnesium aluminum silicate, starch paste, gelatin, methylcellulose, sodium carboxymethylcellulose and/or
20 polyvinylpyrrolidone, and if necessary disintegrating agents such as starch, agarose, alginic acid or its sodium salt or azeotropic mixtures and/or absorbents, coloring agents, flavours, and sweeteners can be additionally included thereto.

The pharmaceutical composition comprising the compound represented by formula 1, the optical isomer thereof, or the pharmaceutically acceptable salt thereof as an active ingredient can be administered by
5 parenterally and the parenteral administration includes subcutaneous injection, intravenous injection, intramuscular injection, or intrathoracic injection.

To prepare the composition as a formulation for
10 parenteral administration, the compound represented by formula 1 or the pharmaceutically acceptable salt thereof is mixed with a stabilizer or a buffering agent in water to produce a solution or suspension, which is then formulated as ampoules or vials. The
15 composition herein can be sterilized and additionally contains preservatives, stabilizers, wettable powders or emulsifiers, salts and/or buffers for the regulation of osmotic pressure, and other therapeutically useful materials, and the composition
20 can be formulated by the conventional mixing, granulating or coating method.

The effective dosage of the pharmaceutical composition comprising the compound represented by formula 1, the optical isomer thereof, or the
25 pharmaceutically acceptable salt thereof as an active

ingredient of the present invention can be determined according to age, weight, gender, administration method, health condition, and severity of disease. The dosage is 0.1 ~ 1000 mg/day for an adult patient
5 (70 Kg), preferably 1 ~ 500 mg/day, which can be administered several times a day or preferably once a day or a couple of times a day.

The pharmaceutical composition comprising the compound represented by formula 1, the optical isomer
10 thereof, or the pharmaceutically acceptable salt thereof as an active ingredient of the present invention can be administered alone or treated together with surgical operation, hormone therapy, chemo-therapy and biological regulators.

15

In addition, the present invention provides a health food composition for the prevention or improvement of PI3 kinase related diseases which comprises the said heteroaryl derivative, the optical
20 isomer thereof, or the pharmaceutically acceptable salt of the same as an active ingredient.

The PI3 kinase related disease herein includes cancer such as hematological malignance, ovarian cancer, cervical cancer, breast cancer, colorectal
25 cancer, liver cancer, stomach cancer, pancreatic

cancer, colon cancer, peritoneal metastasis, skin cancer, bladder cancer, prostate cancer, thyroid cancer, lung cancer, osteosarcoma, fibrous tumor, and brain tumor, autoimmune disease such as rheumatoid
5 arthritis, systemic lupus erythematosus, multiple sclerosis, type 1 diabetes, hyperthyroidism, myasthenia, Crohn's disease, ankylosing spondylitis, psoriasis, autoimmune pernicious anemia, and Sjogren's syndrome, and respiratory disease such as chronic
10 obstructive pulmonary disease(COPD), rhinitis, asthma, chronic bronchitis, chronic inflammatory lung disease, silicosis, pulmonary sarcoidosis, pleurisy, alveolitis, vasculitis, emphysema, pneumonia, and bronchiectasis.

15 The compound represented by formula 1 of the present invention acts as a PI3 kinase inhibitor, so that it can be added to a health functional supplement including food and beverages as a health food composition for the prevention or improvement of PI3
20 kinase related diseases.

The compound represented by formula 1 of the present invention can be used as a food additive. In that case, the compound of the present invention can be added as it is or as mixed with other food
25 components according to the conventional method. The

mixing ratio of active ingredients can be regulated according to the purpose of use (prevention or improvement). In general, to produce health food or beverages, the compound of the present invention is
5 added preferably by 0.1 ~ 90 weight part. However, if long term administration is required for health and hygiene or regulating health condition, the content can be lower than the above but higher content can be accepted as well since the compound of the present
10 invention has been proved to be very safe.

The composition for health beverages of the present invention can additionally include various flavors or natural carbohydrates, etc, like other beverages. The natural carbohydrates above can be one
15 of monosaccharides such as glucose and fructose, disaccharides such as maltose and sucrose, polysaccharides such as dextrin and cyclodextrin, and glucose alcohols such as xylitol, sorbitol and erythritol. Besides, natural sweetening agents
20 (thaumatin, stevia extract, for example rebaudioside A, glycyrrhizin, etc.) and synthetic sweetening agents (saccharin, aspartame, etc.) can be included as a sweetening agent. The content of the natural carbohydrate is preferably 1 ~ 20 g and more
25 preferably 5 ~ 12 g in 100 g of the composition.


In addition to the ingredients mentioned above, the compound represented by formula 1 of the present invention can include in variety of nutrients, vitamins, minerals (electrolytes), flavors including
5 natural flavors and synthetic flavors, coloring agents and extenders (cheese, chocolate, etc.), pectic acid and its salts, alginic acid and its salts, organic acid, protective colloidal viscosifiers, pH regulators, stabilizers, antiseptics, glycerin,
10 alcohols, carbonators which used to be added to soda, etc. The compound represented by formula 1 of the present invention can also include fruit flesh addable to natural fruit juice, fruit beverages, and vegetable beverages.

15

Practical and presently preferred embodiments of the present invention are illustrative as shown in the following Examples.

However, it will be appreciated that those
20 skilled in the art, on consideration of this disclosure, may make modifications and improvements within the spirit and scope of the present invention.

Preparative Example 1: Preparation of (S)-2-(1-aminoethyl)-5-chloro-3-phenylquinazoline-4(3H)-one
25



Step 1: Preparation of 2-amino-6-chlorobenzoic acid

The reaction mixture composed of 5 g of 2-amino-6-chlorobenzonitrile (32.77 mmol), 30% potassium hydroxide (50 mL), and 30% hydrogen peroxide aqueous solution (3 mL) was heat-refluxed for 12 hours, which was then cooled down at room temperature. The aqueous layer was separated by using diethyl ether and then acidized with 12 N HCl (pH: 3-4) to separate an organic layer. The organic layer was washed with saturated brine and then separated, dried (Na₂SO₄), filtered, and concentrated under reduced pressure. As a result, 5.31 g of the target compound 2-amino-6-chlorobenzoic acid was obtained as a yellow solid (30.95 mmol, yield: 94%).

¹H NMR (300 MHz, DMSO-d₆) δ 8.24 (s, 2H), 7.00-7.06 (t, J = 7.5 Hz, 1H), 6.64 (d, J = 8.4 Hz, 1H), 6.56 (d, J = 7.8 Hz, 1H).

20

Step 2: Preparation of tert-butyl (S)-(1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)carbamate

1.00 g (5.89 mmol) of 2-amino-6-chlorobenzoic acid obtained in step 1 above was mixed with N-Boc-L-alanine (1 equivalent), triphenyl phosphite (1.2 equivalent), and anhydrous pyridine (5 mL). The
5 reaction mixture was stirred at 55°C for 12 hours, to which aniline (1 equivalent) was added. The mixture was heated for 6 hours, and then cooled down at room temperature. The mixture was concentrated under reduced pressure, followed by acidization with 1N HCl
10 (pH: 5-6). The reaction mixture was extracted by using ethyl acetate to separate an organic layer. The organic layer was washed with saturated brine, separated, dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was separated by
15 column chromatography (SiO₂, eluent: hexane/ethyl acetate, 10/1 -> hexane/ethyl acetate, 1/1) to give 1.63 g of the target compound tert-butyl (S)-(1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)carbamate as a yellow solid (4.09 mmol,
20 yield: 69%).


¹H NMR (300 MHz, CDCl₃) δ 7.61-7.63(m, 2H), 7.46-7.57(m, 4H), 7.36-7.39(m, 1H), 7.29(s, 1H), 5.59(s, 1H), 4.50(s, 1H), 1.37-1.46(m, 9H), 1.25(d, J = 6.5Hz, 3H).

Step 3: Preparation of (S)-2-(1-aminoethyl)-5-chloro-3-phenylquinazoline-4(3H)-one

1.634 g (4.09 mmol) of tert-butyl (S)-(1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)carbamate prepared in step 2 was dissolved in dichloromethane (15 mL), to which trifluoroacetic acid (TFA, 5 mL) was added. After the reflux at 40°C for 3 hours, the mixture was cooled down at room temperature, to which saturated NaHCO₃ aqueous solution was slowly added to neutralize the mixture. The organic layer was extracted by using ethyl acetate, which was washed with saturated brine, separated, dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was separated by column chromatography (SiO₂, eluent: dichloromethane/methanol, 20/1 -> dichloromethane/methanol, 5/1) to give 1.046 g of the target compound (S)-2-(1-aminoethyl)-5-chloro-3-phenylquinazoline-4(3H)-one as a white solid (3.49 mmol, yield: 85%).

¹H NMR (300 MHz, CDCl₃) δ 7.60-7.64 (m, 2H), 7.51-7.59 (m, 3H), 7.44-7.48 (m, 1H), 7.27-7.29 (m, 2H), 3.63-3.70 (m, 1H), 1.83 (s, 2H), 1.27 (d, J = 6.5 Hz, 3H).

Preparative Example 2: Preparation of (S)-5-chloro-3-phenyl-2-(pyrrolidine-2-yl)quinazoline-4(3H)-one



5

Step 1: Preparation of tert-butyl (S)-2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-carboxylate

5.51 g of tert-butyl (S)-2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-carboxylate was prepared as a beige solid by using 3.76 g (17.47 mmol) of (tert-butoxycarbonyl)-L-proline according to the same manner as described in step 2 of Preparative Example 1 (12.94 mmol, yield: 74%).


^1H NMR (300 MHz, $CDCl_3$) δ 7.42-7.61 (m, 6H), 7.32-7.34 (m, 1H), 7.20-7.24 (m, 1H), 4.40-4.43 (m, 1H), 3.41-3.52 (m, 2H), 1.86-2.06 (m, 3H), 1.70-1.76 (m, 1H), 1.30 (s, 9H).

Step 2: Preparation of (S)-5-chloro-3-phenyl-2-(pyrrolidine-2-yl)quinazoline-4(3H)-one

3.3 g of (S)-5-chloro-3-phenyl-2-(pyrrolidine-2-yl)quinazoline-4(3H)-one was prepared as a beige solid by using 5.53 g (12.99 mmol) of tert-butyl (S)-2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-carboxylate prepared in step 1 according to the same manner as described in step 3 of Preparative Example 1 (10.13 mmol, yield: 78%).

^1H NMR (300 MHz, CDCl_3) δ 7.60-7.62(m, 2H), 7.47-7.54(m, 4H), 7.27-7.29(m, 2H), 3.75-3.79(m, 1H), 3.22-3.26(m, 1H), 3.73-3.76(m, 1H), 1.69-1.77(m, 4H).

Preparative Example 3: Preparation of (S)-2-(1-aminoethyl)-5-chloro-3-(pyridine-3-yl)quinazoline-4(3H)-one



Step 1: Preparation of tert-butyl (S)-2-(1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)carbamate

4.06 g of tert-butyl (S)-2-(1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)carbamate was prepared as an ivory solid by using 1.59 g (16.90 mmol) of 3-aminopyridine according

to the same manner as described in step 2 of Preparative Example 1 (10.14 mmol, yield: 60%).

¹H NMR (300 MHz, CDCl₃) δ 8.77 (s, 1H), 8.57 (s, 1H), 7.45-7.82 (m, 5H), 5.47 (s, 1H), 4.35-4.38 (m, 1H),
5 1.41 (s, 9H), 1.26-1.31 (m, 3H).


Step 2: Preparation of (S)-2-(1-aminoethyl)-5-chloro-3-(pyridine-3-yl)quinazoline-4(3H)-one

2.6 g of (S)-2-(1-aminoethyl)-5-chloro-3-
10 (pyridine-3-yl)quinazoline-4(3H)-one was prepared as a white solid by using 4.08 g (10.18 mmol) of tert-butyl (S)-(1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)carbamate prepared in
step 1 according to the same manner as described in
15 step 3 of Preparative Example 1 (8.65 mmol, yield: 85%).

¹H NMR (300 MHz, CDCl₃) δ 8.78 (s, 1H), 8.58 (s, 1H), 7.65-7.71 (m, 3H), 7.52-7.56 (m, 2H), 3.57-3.64 (m, 1H), 1.29 (dd, J = 22.7, 5.9 Hz, 3H).

20

Preparative Example 4: Preparation of (S)-5-chloro-3-(pyridine-3-yl)-2-(pyrrolidine-2-yl)quinazoline-4(3H)-one



Step 1: Preparation of tert-butyl (S)-2-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-carboxylate

3.82 g of tert-butyl (S)-2-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-carboxylate was prepared as a white solid by using 1.65 g (17.55 mmol) of 3-aminopyridine according to the same manner as described in step 2 of Preparative Example 1 (8.95 mmol, yield: 51%).

¹H NMR (300 MHz, CDCl₃) δ 8.74-8.77(m, 1H), 8.51 (s, 1H), 7.45-7.70(m, 5H), 4.27-4.41(m, 1H), 3.70-3.83(m, 1H), 3.45-3.60(m, 1H), 1.92-1.99(m, 2H), 1.77-1.87(m, 2H), 1.31(d, J = 11.3 Hz, 9H).

15


Step 2: Preparation of (S)-5-chloro-3-(pyridine-3-yl)-2-(pyrrolidine-2-yl)quinazoline-4(3H)-one

2.6 g of (S)-5-chloro-3-(pyridine-3-yl)-2-(pyrrolidine-2-yl)quinazoline-4(3H)-one was prepared as a white solid by using 3.83 g (8.97 mmol) of tert-butyl (S)-2-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-carboxylate

prepared in step 1 according to the same manner as described in step 3 of Preparative Example 1 (7.80 mmol, yield: 87%).

¹H NMR (300 MHz, CDCl₃) δ 8.75 (s, 1H), 8.55-8.59 (d, J = 12.2 Hz, 1H), 7.63-7.70 (m, 3H), 7.50-7.52 (m, 2H), 3.63-3.81 (m, 1H), 3.20-3.27 (m, 1H), 2.74-2.79 (m, 1H), 1.65-1.78 (m, 4H).

Preparative Example 5: Preparation of (S)-2-(1-aminoethyl)-5-chloro-3-(m-tolyl)quinazoline-4(3H)-one



Step 1: Preparation of tert-butyl (S)-(1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)carbamate

5.1 g of tert-butyl (S)-(1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)carbamate was prepared as a white solid by using 1.88 g (17.56 mmol) of m-toluidine according to the same manner as described in step 2 of Preparative Example 1 (12.29 mmol, yield: 70%).

¹H NMR (300 MHz, CDCl₃) δ 7.62 (s, 2H), 7.39-7.47 (m, 2H), 7.31-7.33 (m, 1H), 7.15 (s, 1H), 7.08 (s, 1H),


5.61(s, 1H), 4.50-4.53(m, 1H), 2.42 (s, 3H), 1.42(s, 9H), 1.27(s, 3H).

Step 2: Preparation of (S)-2-(1-aminoethyl)-5-chloro-3-(m-tolyl)quinazoline-4(3H)-one

3.0 g of (S)-2-(1-aminoethyl)-5-chloro-3-(m-tolyl)quinazoline-4(3H)-one was prepared as a white solid by using 5.0 g (12.10 mmol) of tert-butyl (S)-(1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)carbamate prepared in step 1 according to the same manner as described in step 3 of Preparative Example 1 (9.56 mmol, yield: 79%).

^1H NMR (300 MHz, CDCl_3) δ 7.61-7.63(m, 2H), 7.41-7.48(m, 2H), 7.30-7.33(d, J = 7.7 Hz, 1H), 7.05-7.08(m, 2H), 3.66-3.73(q, J = 13.0, 6.5 Hz, 1H), 2.42(s, 3H), 1.27-1.29(d, J = 6.5 Hz, 3H).

Preparative Example 6: Preparation of (S)-5-chloro-2-(pyrrolidine-2-yl)-3-(m-tolyl)quinazoline-4(3H)-one



Step 1: Preparation of (S)-tert-butyl 2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-carboxylate

5.67 g of tert-butyl (S)-2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-carboxylate was prepared as a yellow solid by using 1.87 g (17.41 mmol) of m-toluidine according to the same manner as described in step 2 of Preparative Example 1 (12.88 mmol, yield: 74%).

¹H NMR (300 MHz, CDCl₃) δ 7.53-7.58 (m, 2H), 7.45-7.51 (m, 2H), 7.30-7.33 (m, 1H), 7.14 (s, 1H), 7.02 (s, 1H), 4.43-4.51 (m, 1H), 3.63-3.74 (m, 1H), 3.42-3.50 (m, 1H), 2.42 (s, 3H), 1.93-2.04 (m, 3H), 1.73-1.79 (m, 1H), 1.23-1.37 (m, 9H).

15


Step 2: Preparation of (S)-5-chloro-2-(pyrrolidine-2-yl)-3-(m-tolyl)quinazoline-4(3H)-one

4.0 g of (S)-5-chloro-2-(pyrrolidine-2-yl)-3-(m-tolyl)quinazoline-4(3H)-one was prepared as a white solid by using 5.69 g (12.93 mmol) of tert-butyl (S)-2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-carboxylate prepared in step 1 according to the same manner as described in step 3 of Preparative Example 1 (11.77 mmol, yield: 91%).

^1H NMR (300 MHz, CDCl_3) δ 7.42-7.58 (m, 4H), 7.35 (d, $J = 8.1$ Hz, 1H), 7.20-7.24 (m, 1H), 7.05 (s, 1H), 4.44-4.51 (m, 1H), 3.42-3.50 (m, 1H), 3.18-3.24 (m, 1H), 2.44 (d, $J = 11.7$ Hz, 3H), 1.76-1.93 (m, 4H).

5

Preparative Example 7: Preparation of (S)-2-(1-aminoethyl)-5-chloro-3-(3-fluorophenyl)quinazoline-4(3H)-one



10 Step 1: Preparation of tert-butyl (S)-2-(1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)carbamate

4.88 g of tert-butyl (S)-2-(1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)carbamate was prepared as a yellow solid by using 1.94 g (17.43 mmol) of 3-fluoroaniline according to the same manner as described in step 2 of Preparative Example 1 (11.68 mmol, yield: 67%).


20 ^1H NMR (300 MHz, CDCl_3) δ 7.42-7.63 (m, 4H), 7.14-7.23 (m, 1H), 7.03-7.17 (m, 2H), 5.44-5.55 (m, 1H), 4.48-4.52 (m, 1H), 1.42 (s, 9H), 1.18-1.31 (m, 3H).

Step 2: Preparation of (S)-2-(1-aminoethyl)-5-chloro-3-(3-fluorophenyl)quinazoline-4(3H)-one

2.4 g of (S)-2-(1-aminoethyl)-5-chloro-3-(3-fluorophenyl)quinazoline-4(3H)-one was prepared as a white solid by using 4.88 g (11.80 mmol) of tert-butyl (S)-(1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)carbamate prepared in step 1 according to the same manner as described in step 3 of Preparative Example 1 (7.55 mmol, yield: 64%).

^1H NMR (300 MHz, CDCl_3) δ 1.64 (s, 1H), 7.63 (d, J = 1.5 Hz, 1H), 7.46-7.60 (m, 2H), 7.22-7.25 (m, 1H), 7.04-7.10 (m, 2H), 3.65-3.71 (m, 1H), 1.31 (dd, J = 6.5, 1.3 Hz, 3H).

Preparative Example 8: Preparation of (S)-5-chloro-3-(3-fluorophenyl)-2-(pyrrolidine-2-yl)quinazoline-4(3H)-one



Step 1: Preparation of tert-butyl (S)-2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-carboxylate

6.33 g of tert-butyl (S)-2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-carboxylate was prepared as a yellow solid by using 1.93 g (17.39 mmol) of 3-fluoroaniline according to the same manner as described in step 2 of Preparative Example 1 (14.26 mmol, yield: 82%).

¹H NMR(300 MHz, CDCl₃) δ 7.52-7.65(m, 3H), 7.34-7.49(m, 2H), 6.97-7.23(m, 2H), 4.42-4.51(m, 1H), 3.65-3.77(m, 1H), 3.42-3.54(m, 1H), 1.91-2.11(m, 3H), 1.79-1.88(m, 1H), 1.26-1.37(m, 9H).


Step 2: Preparation of (S)-5-chloro-3-(3-fluorophenyl)-2-(pyrrolidine-2-yl)quinazoline-4(3H)-one

3.82 g of (S)-5-chloro-3-(3-fluorophenyl)-2-(pyrrolidine-2-yl)quinazoline-4(3H)-one was prepared as a white solid by using 6.49 g (14.62 mmol) of tert-butyl (S)-2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-carboxylate prepared in step 1 according to the same manner as described in step 3 of Preparative Example 1 (11.11 mmol, yield: 76%).

¹H NMR(300 MHz, CDCl₃) δ 7.62(s, 1H), 7.61(s, 1H), 7.44-7.58(m, 2H), 7.20-7.25(m, 1H), 7.02-7.12(m, 2H),

3.78-3.81 (m, 1H), 3.24-3.28 (m, 1H), 3.00 (s, 1H), 2.77-2.80 (m, 1H), 1.72-1.82 (m, 4H).

Preparative Example 9: Preparation of (S)-2-(1-aminopropyl)-5-fluoro-3-phenylquinazoline-4(3H)-one



Step 1: Preparation of tert-butyl (S)-2-(1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)carbamate

4.42 g of tert-butyl (S)-2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-carboxylate was prepared as a white solid by using 2.97g (14.61 mmol) of (S)-2-((tert-butoxycarbonyl)amino)butanoic acid according to the same manner as described in step 2 of Preparative Example 1 (11.11 mmol, yield: 76%).

¹H NMR (300 MHz, DMSO-d₆) δ 7.39-7.43 (m, 1H), 7.07-7.19 (m, 3H), 6.99-7.02 (m, 2H), 6.72-6.88 (m, 2H), 3.51-3.56 (m, 1H), 2.92 (s, 1H), 1.26-1.31 (m, 1H), 1.10-1.17 (m, 1H), 0.92 (s, 9H), 0.21 (t, J = 6.7 Hz, 3H).


Step 2: Preparation of (S)-2-(1-aminopropyl)-5-fluoro-3-phenylquinazoline-4(3H)-one

32.43 g of (S)-2-(1-aminopropyl)-5-fluoro-3-phenylquinazoline-4(3H)-one was prepared as a white solid by using 4.16 g (10.47 mmol) of tert-butyl (S)-2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-carboxylate prepared in step 1 according to the same manner as described in step 3 of Preparative Example 1 (8.17 mmol, yield: 78%).

^1H NMR (300 MHz, CDCl_3) δ 7.66-7.73(m, 1H), 7.50-7.56(m, 4H), 7.27-7.28(m, 3H), 7.11(t, J = 5.1 Hz, 1H), 3.40-3.44(m, 1H), 1.75-1.84(m, 1H), 1.46-1.55(m, 1H), 0.79(t, J = 7.4 Hz, 3H).

15

Preparative Example 10: Preparation of (S)-3-(1-aminoethyl)-4,8-dichloro-2-phenylisoquinoline-1(2H)-one



Step 1: Preparation of 2-chloro-6-methylbenzoyl chloride

10.073 g (59.04 mmol) of 2-chloro-6-methylbenzoic acid was mixed with anhydrous dichloromethane (150 mL), to which 10.3 ml (118.09 mmol, 2 equivalents) of oxalylchloride was added. Dimethylformamide was dropped thereto 1-2 drops, and the mixture was stirred at room temperature for 4 hours. The mixture was concentrated under reduced pressure. As a result, 11.479 g of the target compound brown liquid target compound 2-chloro-6-methylbenzoyl chloride was obtained as a brown liquid (59.04 mmol, yield: 100%).

Step 2: Preparation of 2-chloro-6-methyl-N-phenylbenzamide

5.8 mL (63.76 mmol, 1.05 equivalent) of aniline and 14.8 mL (106.26 mmol, 1.75 equivalent) of triethylamine were dissolved in anhydrous dichloromethane (150 mL), to which 11.48 g (60.7 mmol, 1.0 equivalent) of 2-chloro-6-methylbenzoyl chloride prepared in step 1 that had been dissolved in anhydrous dichloromethane (20 mL) was slowly dropped at 0°C for 10 minutes, followed by stirring for 5 hours. Then, the mixture was washed with 1N HCl, water, and saturated sodiumbicarbonate solution stepwise. The organic layer was separated, dried (Na₂SO₄), filtered, and distilled under reduced pressure. The obtained solid was recrystallized with hexane/ethyl acetate to give 13.0 g of the target compound 2-chloro-6-methyl-N-phenylbenzamide as a white solid (52.9 mmol, yield: 87%).

¹H NMR (300 MHz, DMSO-d₆) δ 10.56(s, 1H), 7.69-7.72(d, J = 7.7 Hz, 2H), 7.27-7.37(m, 5H), 7.10(t, J = 7.3 Hz, 1H), 2.31(s, 3H).

Step 3: Preparation of tert-butyl (S)-(4-(3-chloro-2-(phenylcarbamoyl)phenyl)-3-oxobutane-2-yl)carbamate

6 g (24.42 mmol) of 2-chloro-6-methyl-N-phenylbenzamide prepared in step 2 was dissolved in anhydrous THF (50 mL), to which 24.42 ml (61.05 mmol, 2.5 equivalent) of n-BuLi was slowly added at -30°C. 5 The mixture was stirred for 1 hour. 8.5 g (36.63 mmol, 1.5 equivalent) of tert-butyl (S)-(1-(methoxy(methyl)amino)-1-oxopropane-2-yl)carbamate was dissolved in anhydrous THF (50 mL), to which 56.35 ml (73.26 mmol, 3.0 equivalent) of isopropyl 10 magnesiumchloride was slowly added at -30°C. The reaction mixture was stirred for 1 hour, which was slowly added to the above mixture by using a cannula, followed by stirring at -15°C for 2 hours. The temperature was maintained at -15°C ~ -10°C, while the 15 reaction mixture was added with water and 1N HCl stepwise. PH of the reaction mixture was regulated to be 5 and the mixture was heated at room temperature. The organic layer was extracted by using ethyl acetate, which was washed with saturated brine, 20 separated, dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was separated by column chromatography (SiO₂, eluent: CH₂Cl₂/MeOH, 30/1 -> CH₂Cl₂/MeOH, 10/1) to give 8.8 g of the target compound tert-butyl (S)-(4-(3-chloro-2-

(phenylcarbamoyl)phenyl)-3-oxobutane-2-yl)carbamate as a white solid (21.11 mmol, yield: 86%).

¹H NMR (300 MHz, CDCl₃) δ 7.90 (s, 1H), 7.59 (d, J = 7.6 Hz, 2H), 7.29-7.35 (m, 4H), 7.13-7.18 (m, 2H), 5.01 (s, 1H), 4.33-4.37 (m, 1H), 3.91-4.06 (m, 2H), 1.40 (s, 9H), 1.24 (d, J = 7.3 Hz, 3H).

Step 4: Preparation of (S)-3-(1-aminoethyl)-8-chloro-2-phenylisoquinoline-1(2H)-one

8.8 g (21.11 mmol) of tert-butyl (S)-(4-(3-chloro-2-(phenylcarbamoyl)phenyl)-3-oxobutane-2-yl)carbamate prepared in step 3 was dissolved in IPA/12 N HCl (5/3, 160 mL), followed by stirring at 65°C for 2 hours. The mixture was concentrated under reduced pressure, to which saturated sodiumbicarbonate aqueous solution was added. The organic layer was extracted by using dichloromethane, which was separated, dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was separated by column chromatography (SiO₂, eluent: CH₂Cl₂/MeOH, 10/1 -> CH₂Cl₂/MeOH, 5/1) to give 4.871 g of the target compound (S)-3-(1-aminoethyl)-8-chloro-2-phenylisoquinoline-1(2H)-one as a white solid (16.30 mmol, yield: 77%).

Step 5: Preparation of (S)-3-(1-aminoethyl)-8-chloro-2-phenylisoquinoline-1(2H)-one

4.871 g (16.30 mmol) of (S)-3-(1-aminoethyl)-8-chloro-2-phenylisoquinoline-1(2H)-one prepared in step 4 was dissolved in methanol (100 mL), to which 2.45 g (16.30 mmol, 1.0 equivalent) of (D)-tartaric acid was added. The mixture was stirred at room temperature for 30 minutes, followed by reflux for 90 minutes. The reaction mixture was stirred at room temperature for 12 hours and then the generated white solid was filtered. Water was added to the white solid and pH was adjusted to be 8 with saturated sodium bicarbonate aqueous solution, followed by stirring at room temperature for 30 minutes. The white solid was filtered and dried. As a result, 3.74 g of the target compound (S)-3-(1-aminoethyl)-8-chloro-2-phenylisoquinoline-1(2H)-one was obtained as a white solid (12.50 mmol, yield: 77%).

^1H NMR (300 MHz, CDCl_3) δ 7.41-7.56(m, 7H), 7.28(s, 1H), 6.71(s, 1H), 3.68-3.74(q, J = 6.5 Hz, 1H), 1.31(s, 2H), 1.25(d, J = 6.5 Hz, 3H).

Step 6: Preparation of (S)-N-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)-2,2,2-trifluoroacetamide

2.99 g (10.00 mmol) of (S)-3-(1-aminoethyl)-8-chloro-2-phenylisoquinoline-1(2H)-one prepared in step 5 and anhydrous pyridine (3 equivalent) were added to anhydrous CH_2Cl_2 (15 mL), to which trifluoroacetic anhydride $[(\text{CF}_3\text{CO})_2\text{O}]$ (1.2 equivalent) was added at 0°C . 30 minutes later, the reaction mixture was warmed up to room temperature, followed by stirring for 2 hours. The organic layer was extracted by using water and ethyl acetate, which was washed with saturated brine, separated, dried (Na_2SO_4), filtered, and concentrated under reduced pressure. The residue was separated by column chromatography (SiO_2 , eluent: hexane/ethyl acetate, 10/1 $\rightarrow$ hexane/ethyl acetate, 2/1) to give 3.83 g of the target compound (S)-N-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)-2,2,2-trifluoroacetamide as a white solid (9.70 mmol, yield: 97%).

^1H NMR (300 MHz, CDCl_3) δ 7.20-7.60 (m, 8H), 6.52 (s, 1H), 6.38 (br d, 1H), 4.64-4.74 (m, 1H), 1.43 (d, $J = 6.9\text{Hz}$, 3H).

Step 7: Preparation of (S)-N-(1-(4,8-dichloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)-2,2,2-trifluoroacetamide

3.55 g (9.00 mmol) of (S)-N-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)-2,2,2-trifluoroacetamide prepared in step 6, N-chlorosuccinimide (NCS, 1.2 equivalent), and anhydrous acetonitrile (25 mL) were mixed together. The reaction mixture was refluxed for 4 hours and then cooled down the temperature to room temperature. Saturated sodiumthiosulfate (Na₂S₂O₃) solution (2 mL) and water were added thereto, followed by extraction using ethyl acetate. The organic layer was separated and washed with saturated brine. The organic layer was separated, dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was separated by column chromatography (SiO₂, eluent: hexane/ethyl acetate, 10/1 -> hexane/ethyl acetate, 3/1) to give 3.79 g of the target compound (S)-N-(1-(4,8-dichloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)-2,2,2-trifluoroacetamide as a white solid (8.82 mmol, yield: 98%).


¹H NMR(300 MHz, CDCl₃) δ 7.98 (m, 1H), 7.51-7.69 (m, 6H), 7.15-7.20 (m, 1H), 7.03 (br s, 1H), 4.85-5.00 (m, 1H), 1.58 (d, J = 7.2Hz, 3H).

Step 8: Preparation of (S)-3-(1-aminoethyl)-4,8-dichloro-2-phenylisoquinoline-1(2H)-one

3.78 g (8.8 mmol) of (S)-N-(1-(4,8-dichloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)-2,2,2-trifluoroacetamide prepared in step 7, K₂CO₃ (5 equivalent), and MeOH/H₂O (10/1, 20 mL) were mixed and
5 the reaction mixture was refluxed for 12 hours. The mixture was cooled down to room temperature. The solvent was eliminated under reduced pressure. Water and ethyl acetate were added thereto, followed by extraction. The organic layer was separated and
10 washed with saturated brine. The organic layer was separated, dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was separated by column chromatography (SiO₂, eluent: CH₂Cl₂/MeOH, 20/1
-> hexane CH₂Cl₂/MeOH, 10/1) to give 2.90 g of the
15 target compound (S)-3-(1-aminoethyl)-4,8-dichloro-2-phenylisoquinoline-1(2H)-one as a white solid (8.7 mmol, yield: 99%).

¹H NMR(300 MHz, CDCl₃) δ 7.98-8.03 (m, 1H), 7.45-7.65 (m, 5H), 7.17-7.30 (m, 2H), 3.87-4.00 (m, 1H),
20 1.80 (br s, 2H), 1.46 (d, J = 7.1Hz, 3H).

Preparative Example 11: Preparation of (S)-8-chloro-2-phenyl-3-(pyrrolidine-2-yl)isoquinoline-1(2H)-one



Step 1: Preparation of tert-butyl (S)-2-(2-(3-chloro-2-(phenylcarbamoyl)phenyl)acetyl)pyrrolidine-1-carboxylate

5 6 g (24.42 mmol) of 2-chloro-6-methyl-N-phenylbenzamide prepared in step 2 of Preparative Example 13 was dissolved in anhydrous THF (50 mL). 24.42 ml (61.05 mmol, 2.5 equivalent) of n-BuLi was slowly added thereto at -30°C , followed by stirring
10 for 1 hour. 9.46 g (36.63 mmol, 1.5 equivalent) of tert-butyl (S)-2-(methoxy(methyl)carbamoyl)pyrrolidine-1-carboxylate was dissolved in anhydrous THF (50 mL), which was slowly added to the reaction mixture above. Then,
15 5.05 g of the target compound tert-butyl (S)-2-(2-(3-chloro-2-(phenylcarbamoyl)phenyl)acetyl)pyrrolidine-1-carboxylate was obtained as a white solid by the same manner as described in step 3 of Preparative Example 10 (11.41 mmol, yield: 88%).

20 ^1H NMR (300 MHz, CDCl_3) δ 8.03(s, 1H), 7.60(d, J = 6.9 Hz, 2H), 7.32-7.38(m, 3H), 7.16(d, J = 8.6 Hz, 2H), 7.05(d, J = 4.1 Hz, 1H), 4.28-4.39(m, 1H),


3.98(s, 1H), 3.41-3.52(m, 2H), 1.69-1.78(m, 4H),
1.38(d, J = 12.1 Hz, 9H).

Step 2: Preparation of (S)-8-chloro-2-phenyl-3-
5 (pyrrolidine-2-yl)isoquinoline-1(2H)-one

2.3 g of (S)-8-chloro-2-phenyl-3-(pyrrolidine-2-yl)isoquinoline-1(2H)-one was prepared as a white solid by using 3.34g (7.53 mmol) of tert-butyl (S)-2-(2-(3-chloro-2-(phenylcarbamoyl)phenyl)acetyl)pyrrolidine-1-
10 carboxylate prepared in step 1 according to the same manner as described in step 4 of Preparative Example 10 (7.08 mmol, yield: 94%).

¹H NMR(300 MHz, CDCl₃) δ 7.40-7.54(m, 6H), 7.27-
15 7.30(m, 1H), 7.21-7.23(m, 1H), 6.87(s, 1H), 3.78(t, J = 7.0 Hz, 1H), 3.05-3.12(m, 1H), 2.82-2.90(m, 1H), 1.75-1.84(m, 1H), 1.75(s, 1H), 1.54-1.66(m, 3H).

Preparative Example 12: Preparation of (S)-1-(2-
20 phenylquinoline-3-yl)ethaneamine



Step 1: Preparation of 2-phenylquinoline-3-carbaldehyde

10 g (52.19 mmol, 1.0 equivalent) of 2-chloro-3-quinolinecarbaldehyde was dissolved in toluene/water (4/1, 150 mL), to which 7 g (57.41 mmol, 1.1 equivalent) of phenylboronic acid, 12.17 g (114.82 mmol, 2.2 equivalent) of Na_2CO_3 , 1.5 g (1.30 mmol, 2.5 mol%) of $\text{Pd}(\text{PPh}_3)_4$, and 7 ~ 8 drops of Aliquat 336 were slowly added stepwise. The mixture was refluxed for 12 hours under argon atmosphere. The mixture was then cooled down to room temperature. Water was added thereto, followed by extraction using ethyl acetate. The organic layer was separated, dried (N_2SO_4), filtered, and concentrated under reduced pressure. The residue was separated by column chromatography (SiO_2 , eluent: hexane/dichloromethane, 10/1 $\rightarrow$ hexane/dichloromethane, 3/1) to give 12.156 g of the

target compound 2-phenylquinoline-3-carbaldehyde as a white solid (52.11 mmol, yield: 94%).

¹H NMR (300 MHz, CDCl₃) δ 10.19(s, 1H), 8.86(s, 1H), 8.22(d, J = 8.4 Hz, 1H), 8.03(d, J = 7.9 Hz, 1H),
5 7.88(t, J = 7.7 Hz, 1H), 7.64-7.71(m, 3H), 7.55-7.61(m, 3H).

Step 2: Preparation of (S,E)-2-methyl-N-((2-phenylquinoline-3-yl)methylene)propane-2-sulfinamide

10 3 g (12.89 mmol, 1.1 equivalent) of 2-phenylquinoline-3-carbaldehyde prepared in step 1 was dissolved in THF (100 mL), to which 5 ml (23.43 mmol, 2 equivalent) of Ti(OEt)₄ and 1.42 g (11.72 mmol, 1.0 equivalent) of (R)-(+)-2-methyl-2-propanesulfinamide
15 were added, followed by reflux for 12 hours. The reaction mixture was cooled down to room temperature, to which saturated sodiumbicarbonate aqueous solution was added, followed by stirring for 1 hour. The mixture was filtered with celite pad, followed by
20 extraction using ethyl acetate. The organic layer was washed with saturated brine, dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was separated by column chromatography (SiO₂, eluent: hexane/ethyl acetate, 5/1 -> hexane/ethyl acetate,
25 1/1) to give 3.96 g of the target compound (S,E)-2-

methyl-N-((2-phenylquinoline-3-yl)methylene)propane-2-sulfinamide as a yellow solid (11.77 mmol, yield: 91%).

¹H NMR (300 MHz, CDCl₃) δ 8.90 (s, 1H), 8.80 (s, 1H),
5 8.19 (d, J = 8.2 Hz, 1H), 7.97 (d, J = 7.8 Hz, 1H),
7.81 (s, 1H), 7.50-7.61 (m, 6H), 1.31 (s, 9H).

Step 3: Preparation of (R)-2-methyl-N-((S)-1-(2-phenylquinoline-3-yl)ethyl)propane-2-sulfinamide

10 3.96 g (11.76 mmol, 1.0 equivalent) of (S,E)-2-methyl-N-((2-phenylquinoline-3-yl)methylene)propane-2-sulfinamide prepared in step 2 was dissolved in anhydrous dichloromethane (70 mL), to which 11.76 mL (23.53 mmol, 3 equivalent) of 2 M MeMgBr was slowly
15 added at -78°C, followed by stirring for 3 hours. The mixture was further stirred at room temperature for 12 hours, to which saturated NH₄Cl aqueous solution was added. The organic layer was separated, dried (Na₂SO₄), filtered, and concentrated under reduced
20 pressure. The residue was separated by column chromatography (SiO₂, eluent: hexane/ethyl acetate, 3/1 → hexane/ethyl acetate, 1/2) to give 2.52 g of the target compound (R)-2-methyl-N-((S)-1-(2-phenylquinoline-3-yl)ethyl)propane-2-sulfinamide as a
25 white solid (7.15 mmol, yield: 61%).

¹H NMR (300 MHz, CDCl₃) δ 8.32 (s, 1H), 8.16 (d, J = 8.5 Hz, 1H), 7.84 (d, J = 8.2 Hz, 1H), 7.71 (t, J = 7.6 Hz, 1H), 7.41-7.58 (m, 6H), 4.90-4.98 (m, 1H), 3.42 (d, J = 3.1 Hz, 1H), 1.47 (d, J = 6.6 Hz, 3H), 1.20 (s, 9H).

5


Step 4: Preparation of (S)-1-(2-phenylquinoline-3-yl)ethane-1-amine

2.42 g (7.15 mmol, 1.0 equivalent) of (R)-2-methyl-N-((S)-1-(2-phenylquinoline-3-yl)ethyl)propane-
10 2-sulfonamide prepared in step 3 was dissolved in methanol (50 mL), to which 4 M HCl dioxan solution (15 mL) was added at room temperature. The mixture was stirred at room temperature for 2 hours. Saturated sodium bicarbonate aqueous solution was added thereto,
15 followed by extraction using ethyl acetate. The organic layer was separated, dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was separated by column chromatography (SiO₂, eluent: dichloromethane/methanol, 20/1 -
20 > dichloromethane/methanol, 5/1) to give 1.65 g of the target compound (S)-1-(2-phenylquinoline-3-yl)ethane-1-amine as a pale yellow solid (6.64 mmol, yield: 93%).

¹H NMR (300 MHz, CDCl₃) δ 8.43 (s, 1H), 8.13 (d, J = 8.4 Hz, 1H), 7.86 (d, J = 8.1 Hz, 1H), 7.68 (t, J = 7.6

Hz, 1H), 7.44-7.55(m, 6H), 4.42-4.48(q, J = 6.5 Hz, 1H), 1.58(s, 2H), 1.34(d, J = 6.5 Hz, 3H).

Preparative Example 13: Preparation of tert-butyl (S)-(1-(methoxy(methyl)amino)-1-oxopropane-2-yl)carbamate




10 g (52.85 mmol, 1.0 equivalent) of (S)-2-((tert-butoxycarbonyl)amino)propanoic acid was dissolved in anhydrous dichloromethane (250 ml), to which 29.5 ml (211.40 mmol, 4.0 equivalent) of triethylamine and 7.14 g (52.85 mmol, 1.0 equivalent) of hydroxybenzotriazole (HOBt) were added at 0°C. 20.3 g (105.70 mmol, 2.0 equivalent) of EDCI·HCl was added thereto, followed by stirring at room temperature for 30 minutes. 5.7 g (58.14 mmol, 1.1 equivalent) of N,O-dimethylhydroxylamine was added thereto, followed by stirring at room temperature for 12 hours. Water was added to the reaction mixture, followed by extraction using ethyl acetate. The organic layer was separated, dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The obtained solid was recrystallized with hexane/ethyl acetate to

give 11.7 g of the target compound tert-butyl (S)-(1-(methoxy(methyl)amino)-1-oxopropane-2-yl)carbamate as a white solid (50.37 mmol, yield: 95%).

¹H NMR (300 MHz, CDCl₃) δ 5.23 (s, 1H), 4.68-4.70 (m, 1H), 3.77 (s, 3H), 3.12 (s, 3H), 1.44 (s, 9H), 1.31 (d, J = 3.5 Hz, 3H).

Preparative Example 14: Preparation of 2-((tert-butoxycarbonyl)amino)butanoic acid



10 g (96.97 mmol, 1.0 eq) of 2-aminobutanoic acid was dissolved in 65 ml of methanol, to which 97 ml of 1 M sodiumhydroxide (NaOH) and 25.4 g (116.37 mmol, 1.2 equivalent) of di-tert-butyl dicarbonate (Boc₂O) were added at 0°C. The mixture was stirred at room temperature for 48 hours, followed by concentration under reduced pressure. The reaction mixture was acidized with 1N HCl (pH 2-3), followed by extraction using ethyl acetate. The organic layer was separated, dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was separated by column chromatography (SiO₂, eluent: hexane/ethyl acetate, 10/1 → hexane/ethyl acetate, 3/1) to give 18.5 g of

15

20

¹H NMR (300 MHz, CDCl₃) δ 6.24 (s, 1H), 5.00-5.03 (d, J = 7.6 Hz, 1H), 4.27-4.29 (m, 1H), 1.87-1.94 (m, 1H), 1.66-1.78 (m, 1H), 1.45 (s, 9H), 0.96-1.01 (t, J = 7.3 Hz, 3H).

116

5-methyl-3,4-dihydro-2H-pyrrole(**1**) (4 g, 0.05 mol) was dissolved in THF (120 ml), to which N-chlorosuccinimide (51.4 g, 0.39 mol) was slowly added at 0°C. The mixture was stirred for 15 minutes, followed by reflux for 2.5 hours. THF was eliminated under reduced pressure. Extraction was performed with dichloromethane. The organic layer was washed with saturated brine, separated, dried (anhydrous MgSO₄), filtered, and concentrated under reduced pressure.

10 The obtained compound 4,4-dichloro-5-(trichloromethyl)-3,4-dihydro-2H-pyrrole(**2**) was used for the next reaction without purification. 4,4-dichloro-5-(trichloromethyl)-3,4-dihydro-2H-pyrrole(**2**) (12 g, 0.05 mol) was dissolved in methanol (100 ml),

15 to which sodium methoxide (NaOMe) (28 wt% methanol solution) (16 g, 0.29 mol) was slowly added at 0°C, followed by reaction at room temperature for 2 hours. Extraction was performed with ethyl acetate. The organic layer was washed with saturated brine,

20 separated, dried (MgSO₄), filtered, and concentrated under reduced pressure. The residue was separated by column chromatography (SiO₂, eluent: hexane/ethyl acetate, 5/1) to give 6.5 g of the target compound methyl 3-chloro-1H-pyrrole-2-carboxylate(**3**) as a brown

25 solid (0.04 mmol, yield: 77%).

¹H NMR (300 MHz, CDCl₃) δ 9.11 (br s, 1H, NH), 6.87 (t, J = 2.7 Hz, 1H), 6.26 (t, J = 2.7 Hz, 1H), 3.90 (s, 3H).

5 Step 3: Preparation of 3-chloro-1H-pyrrole-2-carboxylic acid(4)

Methyl 3-chloro-1H-pyrrole-2-carboxylate(3) (5 g, 0.03 mol) was dissolved in methanol/water (2/1) (30 ml), to which LiOH·H₂O (5.3 g, 0.13 mol) was added, 10 followed by reflux at room temperature for 1.5 hours. 12 N HCl (13 ml) was slowly added thereto at 0°C. The reaction mixture was extracted by using ethyl acetate. The extract was washed with saturated brine, separated, dried (anhydrous Na₂SO₄), filtered, and 15 concentrated under reduced pressure. The obtained solid compound was washed with hexane to give the target compound 3-chloro-1H-pyrrole-2-carboxylic acid(4).

¹H NMR (300 MHz, DMSO-d₆) δ 12.58 (br s, 1H), 20 11.92 (br s, 1H), 6.94 (t, J = 2.7 Hz, 1H), 6.19 (t, J = 2.7 Hz, 1H).

Step 4-1: Preparation of 3-chloro-N-phenyl-1H-pyrrole-2-carboxamide(5a)

3-chloro-1H-pyrrole-2-carboxylic acid(**4**) (1 g, 6.87 mmol) was dissolved in anhydrous dichloromethane (25 ml), to which oxalyl chloride (1.3 g, 10.31 mmol) and dimethylformamide (2 drops) were slowly added at room temperature. The reaction mixture was refluxed for 1 hour, followed by concentration under reduced pressure. The obtained solid compound was dissolved in anhydrous 1,4-dioxane (8 ml), to which aniline (0.8 g, 8.25 mmol) and N,N-diisopropylethylamine (DIPEA) (2.7 g, 20.61 mmol) were slowly added at 0°C. The mixture was reacted at 60°C for 1 hour, followed by extraction with ethyl acetate. The extract was washed with saturated brine, separated, dried (anhydrous Na₂SO₄), filtered, and concentrated under reduced pressure. The obtained solid compound was washed with hexane to give the target compound 3-chloro-N-phenyl-1H-pyrrole-2-carboxamide(**5a**) as a dark brown solid.

¹H NMR (300 MHz, CDCl₃) δ 10.35 (br s, 1H), 8.60 (br s, 1H), 7.64 (d, J = 7.2 Hz, 2H), 7.37 (t, J = 7.8 Hz, 2H), 7.15 (t, J = 7.2 Hz, 1H), 6.91 (s, 1H), 6.27 (s, 1H).

Step 4-2: Preparation of 3-chloro-N-(3-fluorophenyl)-1H-pyrrole-2-carboxamide(**5b**)

2.6 g of 3-chloro-N-(3-fluorophenyl)-1H-pyrrole-2-carboxamide(**5b**) was prepared as a pale brown solid by using 3-chloro-1H-pyrrole-2-carboxylic acid(**4**) (2 g, 13.75 mmol) and 3-fluoroaniline (1.9 g, 17.19 mmol) by the same method to prepare the compound **5a** (10.85 mmol, yield: 67%).

¹H NMR (300 MHz, CDCl₃) δ 9.81 (br s, 1H), 8.61 (br s, 1H), 7.61 (d, J = 11.1 Hz, 1H), 7.34-7.21 (m, 2H), 6.93 (t, J = 3.0 Hz, 1H), 6.81-6.87 (m, 1H), 6.29 (t, J = 3.0 Hz, 1H).

Step 5-1: Preparation of 1-amino-3-chloro-N-phenyl-1H-pyrrole-2-carboxamide(**6a**)

NH₄Cl (2.1 g, 39 mmol), NaOH (28 wt%) aqueous solution (5.2 g, 130 mmol), NH₄OH (ammoniumhydroxide) (28 wt%) (2.3 g, 65 mmol), and Aliquat 336 (0.3 g, 0.65 mmol) were mixed to prepare a mixed solution. 3-chloro-N-phenyl-1H-pyrrole-2-carboxamide(**5a**) (1.4 g, 6.50 mmol) was dissolved in t-butylmethyl ether/diethyl ether (1:1) (80 ml), which was slowly added to the mixed solution at 0°C. NaOCl (sodiumhypochloride) aqueous solution (10 wt%) was slowly added thereto at the same temperature, followed by reaction at room temperature for 4 hours. Extraction was performed with ethyl acetate. The

organic layer was washed with saturated brine, separated, dried (MgSO₄), filtered, and concentrated under reduced pressure. The residue was separated by column chromatography (SiO₂, eluent: hexane/ethyl acetate, 5/1) to give 1.1 g of the target compound 1-amino-3-chloro-N-phenyl-1H-pyrrole-2-carboxamide (**6a**) as a white solid (4.56 mmol, yield: 70%).

¹H NMR (300 MHz, CDCl₃) δ 8.53 (br s, 1H), 7.59 (d, J = 8.4 Hz, 2H), 7.37 (t, J = 7.8 Hz, 2H), 7.16 (t, J = 7.5 Hz, 1H), 6.91 (d, J = 2.7 Hz, 1H), 6.08 (d, J = 3.0 Hz, 1H), 5.91 (s, 2H).

Step 5-2: Preparation of 1-amino-3-chloro-N-(3-fluorophenyl)-1H-pyrrole-2-carboxamide (**6b**)

1.7 g of 1-amino-3-chloro-N-(3-fluorophenyl)-1H-pyrrole-2-carboxamide (**6b**) was prepared as a white solid by using 3-chloro-N-(3-fluorophenyl)-1H-pyrrole-2-carboxamide (**5b**) (3.9 g, 0.02 mol) by the same method to prepare the compound **6a** (6.78 mmol, yield: 63%).

¹H NMR (300 MHz, DMSO-d₆) δ 11.03 (br s, 1H), 7.68 (d, J = 12 Hz, 1H), 7.36 (s, 2H), 6.98 (d, J = 2.7 Hz, 1H), 6.90-6.94 (m, 1H), 6.54 (s, 2H), 6.12 (d, J = 3.0 Hz, 1H).

Step 6-1-1: Preparation of tert-butyl (S)-2-((3-chloro-2-(phenylcarbamoyl)-1H-pyrrole-1-yl)carbamoyl)pyrrolidine-1-carboxylate(7a)

1-amino-3-chloro-N-phenyl-1H-pyrrole-2-carboxamide(**6a**) (150 mg, 0.64 mmol), N-(tert-butoxycarbonyl)-L-proline (192 mg, 0.89 mmol), and EDC·HCl (171 mg, 0.89 mmol) were dissolved in anhydrous THF (1 ml), followed by reaction at room temperature for 20 hours. The reaction mixture was
10 extracted by using ethyl acetate. The organic layer was washed with saturated brine, separated, dried (MgSO₄), filtered, and concentrated under reduced pressure. The residue was separated by column chromatography (SiO₂, eluent: hexane/ethyl acetate,
15 5/1) to give 193 mg of the target compound tert-butyl (S)-2-((3-chloro-2-(phenylcarbamoyl)-1H-pyrrole-1-yl)carbamoyl)pyrrolidine-1-carboxylate(**7a**) as a white solid (0.45 mmol, yield: 70%).

¹H NMR (300 MHz, CDCl₃) δ 10.61 (br s, 1H), 8.32
20 (brs, 1H), 7.57 (d, J = 7.8 Hz, 2H), 7.34 (t, J = 7.8 Hz, 2H), 7.01-7.15 (m, 2H), 6.20 (s, 1H), 4.30-4.56 (m, 1H), 3.30-3.70 (m, 2H), 2.14-2.44 (m, 2H), 1.82-2.08 (m, 2H), 1.49 (s, 9H).

Step 6-1-2: Preparation of tert-butyl (S)-2-((3-chloro-2-((3-fluorophenyl)carbamoyl)-1H-pyrrole-1-yl)carbamoyl)pyrrolidine-1-carboxylate (**7b**)

Tert-butyl (S)-2-((3-chloro-2-((3-fluorophenyl)carbamoyl)-1H-pyrrole-1-yl)carbamoyl)pyrrolidine-1-carboxylate (**7b**) was prepared by using 1-amino-3-chloro-N-(3-fluorophenyl)-1H-pyrrole-2-carboxamide (**6b**) (0.7 g, 2.76 mmol) by the same method to prepare the compound **7a**.

¹H NMR (300 MHz, CDCl₃) δ 10.61 (br s, 1H), 8.38 (br s, 1H), 7.64 (br s, 1H), 7.23-7.31 (m, 2H), 7.13 (br s, 1H), 6.99 (br s, 1H), 6.79-6.85 (m, 1H), 6.21 (s, 1H), 4.50 (br s, 1H), 3.51 (br s, 1H), 3.42 (br s, 1H), 1.84-2.39 (m, 4H), 1.50 (s, 9H).

Step 6-2-1: Preparation of tert-butyl (S)-(1-((3-chloro-2-(phenylcarbamoyl)-1H-pyrrole-1-yl)amino)-1-oxopropane-2-yl)carbamate (**7a'**)

3.5 g of tert-butyl (S)-(1-((3-chloro-2-(phenylcarbamoyl)-1H-pyrrole-1-yl)amino)-1-oxopropane-2-yl)carbamate (**7a'**) was prepared as a white solid by using 1-amino-3-chloro-N-phenyl-1H-pyrrole-2-carboxamide (**6a**) (2.3 g, 9.76 mmol) and N-(tert-butoxycarbonyl)-L-aniline (2.6 g, 13.66 mmol) by the

same method to prepare the compound **7a** (8.56 mmol, yield: 88%).

¹H NMR (300 MHz, CDCl₃) δ 10.25 (s, 1H), 8.37 (s, 1H), 7.56 (d, J = 8.1 Hz, 2H), 7.34 (t, J = 7.8 Hz, 2H), 7.14 (t, J = 7.5 Hz, 1H), 7.03 (d, J = 2.7 Hz, 1H), 6.21 (d, J = 2.7 Hz, 1H), 5.06 (d, J = 7.2 Hz, 1H), 4.40 (br s, 1H), 1.47 (s, 9H), 1.44 (d, J = 7.5 Hz, 3H).

10 Step 6-2-2: Preparation of tert-butyl (S)-(1-((3-chloro-2-((3-fluorophenyl)carbamoyl)-1H-pyrrole-1-yl)amino)-1-oxopropane-2-yl)carbamate (**7b'**)

Tert-butyl (S)-(1-((3-chloro-2-((3-fluorophenyl)carbamoyl)-1H-pyrrole-1-yl)amino)-1-oxopropane-2-yl)carbamate (**7b'**) was prepared as a white solid by using 1-amino-3-chloro-N-(3-fluorophenyl)-1H-pyrrole-2-carboxamide (**6b**) (3.1 g, 12.26 mmol) and N-(tert-butoxycarbonyl)-L-alanine (3.3 g, 17.16 mmol) by the same method to prepare the
20 compound **7a** (8.56 mmol, yield: 88%).

¹H NMR (300 MHz, CDCl₃) δ 10.16 (s, 1H), 8.42 (s, 1H), 7.61 (d, J = 11.1 Hz, 1H), 7.24-7.31 (m, 2H), 7.14-7.11 (m, 1H), 7.02 (d, J = 3.3 Hz, 1H), 6.86-6.80 (m, 1H), 6.22 (d, J = 3.3 Hz, 1H), 5.00 (d, J = 7.8

Hz, 1H) 4.35-4.42 (m, 1H), 1.48 (s, 9H), 1.44 (d, J = 7.2 Hz, 3H).

Step 7-1-1: Preparation of tert-butyl (S)-2-(5-chloro-4-oxo-3-phenyl-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)pyrrolidine-1-carboxylate (**8a**)

Triphenylphosphine (303 mg, 1.16 mmol) was dissolved in dichloromethane (1 ml), to which Br₂ (184 mg, 1.16 mmol) was slowly added at 0°C, followed by stirring at room temperature for 10 minutes. Tert-butyl (S)-2-((3-chloro-2-(phenylcarbamoyl)-1H-pyrrole-1-yl)carbamoyl)pyrrolidine-1-carboxylate (**7a**) (250 mg, 0.58 mmol) was dissolved in dichloromethane (1 ml), which was slowly added to the mixture above at 0°C. Triethylamine (146 mg, 1.44 mmol) was also added thereto at the same temperature. The reaction mixture was stirred at 0°C for 10 minutes, followed by extraction using dichloromethane. The organic layer was washed with saturated brine, separated, dried (MgSO₄), filtered, and concentrated under reduced pressure. The residue was separated by column chromatography (SiO₂, eluent: hexane/ethyl acetate, 5/1) to give 82 mg of the target compound tert-butyl (S)-2-(5-chloro-4-oxo-3-phenyl-3,4-dihydropyrrolo[2,1-

f][1,2,4]triazine-2-yl)pyrrolidine-1-carboxylate (**8a**)
as a white solid (0.20 mmol, yield: 34%).

¹H NMR (300 MHz, CDCl₃) δ 7.29-7.36 (m, 2H), 7.05-7.13 (m, 3H), 6.36-6.40 (m, 1H), 4.46-4.51 (m, 0.5H),
5 4.36-4.40 (m, 0.5H), 3.09-3.41 (m, 2H), 2.12-2.25 (m, 1H), 1.86-2.00 (m, 1H), 1.71-1.79 (m, 2H), 1.45 (s, 5H), 1.35 (s, 4H).

Step 7-1-2: Preparation of tert-butyl (S)-2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-
10 dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)pyrrolidine-
1-carboxylate (8b)

45 mg of tert-butyl (S)-2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydropyrrolo[2,1-
15 f][1,2,4]triazine-2-yl)pyrrolidine-1-carboxylate (**8b**)
was prepared as a white solid by using tert-butyl (S)-2-((3-chloro-2-((3-fluorophenyl)carbamoyl)-1H-pyrrole-1-yl)carbamoyl)pyrrolidine-1-carboxylate (**7b**) (100 mg, 0.22 mmol) by the same method to prepare the compound
20 **8a** (0.10 mmol, yield: 47%).

¹H NMR (300 MHz, CDCl₃) δ 7.43-7.57 (m, 1H), 7.18-7.37 (m, 2H), 6.99-7.13 (m, 1H), 6.48 (dd, 1H, J = 2.7 Hz, J = 12.9 Hz), 4.46-4.53 (m, 0.5H), 4.41 (br s, 0.5H), 3.32-3.70 (m, 2H), 1.80-2.11 (m, 4H), 1.45 (s,
25 4H), 1.38 (s, 5H).

Step 7-2-1: Preparation of tert-butyl (S)-(1-(5-chloro-4-oxo-3-phenyl-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)ethyl)carbamate(8a')

5 105 mg of tert-butyl (S)-(1-(5-chloro-4-oxo-3-phenyl-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)ethyl)carbamate(**8a'**) was prepared as a white solid by using tert-butyl (S)-(1-((3-chloro-2-(phenylcarbamoyl)-1H-pyrrole-1-yl)amino)-1-oxopropane-2-yl)carbamate(**7a'**) (500 mg, 1.23 mmol) by
10 the same method to prepare the compound **8a** (0.27 mmol, yield: 22%).

¹H NMR (300 MHz, CDCl₃) δ 7.48-7.60 (m, 3H), 7.39-7.41 (m, 1H), 7.28 (brs, 2H), 6.50 (d, J = 2.1 Hz, 1H), 5.09 (brs, 1H), 4.48 (br s, 1H), 1.42 (s, 9H),
15 1.26 (d, J = 6.3 Hz, 3H).

Step 7-2-2: Preparation of tert-butyl (S)-(1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)ethyl)carbamate(8b')

20 140 mg of tert-butyl (S)-(1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)ethyl)carbamate(**8b'**) was
25 prepared as a white solid by using tert-butyl (S)-(1-

((3-chloro-2-((3-fluorophenyl)carbamoyl)-1H-pyrrole-1-yl)amino)-1-oxopropane-2-yl)carbamate(**7b'**) (500 mg, 1.18 mmol) by the same method to prepare the compound **8a** (0.34 mmol, yield: 29%).

5 ^1H NMR (300 MHz, CDCl_3) δ 7.47-7.58 (m, 1H), 7.15-7.30 (m, 3H), 7.02-7.09 (m, 1H), 6.51 (d, $J = 2.1$ Hz, 1H), 4.99-5.10 (m, 1H), 4.48 (br s, 1H), 1.41 (s, 9H), 1.24-1.31 (m, 3H).

10 Step 8-1-1: Preparation of (S)-5-chloro-3-phenyl-2-(pyrrolidine-2-yl)pyrrolo[2,1-f][1,2,4]triazine-4(3H)-one (**9a**)

Tert-butyl (S)-2-(5-chloro-4-oxo-3-phenyl-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)pyrrolidine-15 1-carboxylate(**8a**) (130mg, 0.31 mmol) was dissolved in trifluoroacetic acid (50 wt% dichloromethane) (2 ml) at 0°C , followed by stirring at room temperature for 30 minutes. The reaction mixture was neutralized with NaHCO_3 at 0°C , followed by extraction using 20 dichloromethane. The organic layer was washed with saturated brine, separated, dried (MgSO_4), filtered, and concentrated under reduced pressure. As a result, 96 mg of the target compound (S)-5-chloro-3-phenyl-2-(pyrrolidine-2-yl)pyrrolo[2,1-f][1,2,4]triazine-4(3H)-

one(**9a**) was obtained as a white solid (0.30 mmol, yield: 97%).

¹H NMR (300 MHz, CDCl₃) δ 7.47-7.55 (m, 3H), 7.26-7.30 (m, 3H), 6.49 (d, J = 2.7 Hz, 1H), 3.81 (t, J = 5.7 Hz, 1H), 3.12-3.19 (m, 1H), 2.74-2.81 (m, 1H), 2.02 (br s, 1H), 1.77-1.82 (m, 2H), 1.61-1.73 (m, 2H).

Step 8-1-2: Preparation of (S)-5-chloro-3-(3-fluorophenyl)-2-(pyrrolidine-2-yl)pyrrolo[2,1-f][1,2,4]triazine-4(3H)-one HCl salt(**9b**)

Conc. HCl (15 wt% methanol) (10 ml) was added to tert-butyl (S)-2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)pyrrolidine-1-carboxylate(**8b**) (40 mg, 0.09 mmol) at 0°C, followed by stirring for 1 hour. The solvent was eliminated from the reaction mixture under reduced pressure. As a result, the target compound (S)-5-chloro-3-(3-fluorophenyl)-2-(pyrrolidine-2-yl)pyrrolo[2,1-f][1,2,4]triazine-4(3H)-one hydrochloride salt (**9b**) was obtained as a white solid.

¹H NMR (300 MHz, DMSO-d₆) δ 9.86 (brs, 1H), 9.08 (brs, 1H), 7.63-7.69 (m, 2H), 7.40-7.54 (m, 3H), 6.78 (s, 1H), 4.23 (br s, 1H), 3.17 (br s, 1H), 2.09-2.14 (m, 1H), 1.90-1.98 (m, 1H), 1.69-1.87 (m, 2H).

Step 8-2-1: Preparation of (S)-2-(1-aminoethyl)-5-chloro-3-phenylpyrrolo[2,1-f][1,2,4]triazine-4(3H)-one (9a')

69 mg of (S)-2-(1-aminoethyl)-5-chloro-3-phenylpyrrolo[2,1-f][1,2,4]triazine-4(3H)-one (**9a'**) was prepared as a white solid by using tert-butyl (S)-(1-(5-chloro-4-oxo-3-phenyl-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)ethyl)carbamate (**8a'**) (105 mg, 0.27 mmol) by the same method to prepare the compound **9a** (0.24 mmol, yield: 88%).

¹H NMR (300 MHz, CDCl₃) δ 7.48-7.57 (m, 3H), 7.26-7.30 (m, 3H), 6.50 (d, J = 2.4 Hz, 1H), 3.66 (q, J = 6.6 Hz, J = 13.2 Hz, 1H), 1.29 (d, J = 6.6 Hz, 3H).


Step 8-2-2: Preparation of (S)-2-(1-aminoethyl)-5-chloro-3-(3-fluorophenyl)pyrrolo[2,1-f][1,2,4]triazine-4(3H)-one (9b')

103 mg of (S)-2-(1-aminoethyl)-5-chloro-3-(3-fluorophenyl)pyrrolo[2,1-f][1,2,4]triazine-4(3H)-one (**9b'**) was prepared as a white solid by using tert-butyl (S)-(1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)ethyl)carbamate (**8b'**) (140 mg, 0.34 mmol) by the same method to prepare the compound **9a** (0.33 mmol, yield: 97%).

^1H NMR (300 MHz, CDCl_3) δ 7.48–7.56 (m, 1H), 7.22–7.29 (m, 2H), 7.02–7.14 (m, 2H), 6.50 (d, J = 2.7 Hz, 1H), 3.76 (q, J = 6.3 Hz, J = 12.8 Hz, 1H), 2.22 (br s, 2H), 1.34 (d, J = 6.6 Hz, 3H).

5

Preparative Example 16: Preparation of (S)-1-(7-fluoro-2-(3-fluorophenyl)quinoline-3-yl)ethane-1-amine



Step 1: Preparation of 7-fluoro-2-(3-fluorophenyl)quinoline-3-carbaldehyde

2.48 g of 7-fluoro-2-(3-fluorophenyl)quinoline-3-carbaldehyde was prepared as a pale yellow solid by using 2.10g (10.0 mmol) of 2-chloro-7-fluoroquinoline-3-carbaldehyde by the same manner as described in step 1 of Preparative Example 12 (9.2 mmol, yield: 92%).

MS[m/z; $(M + 1)^+$]: 270.

Step 2: Preparation of (R,E)-N-((7-fluoro-2-(3-fluorophenyl)quinoline-3-yl)methylene)-2-methylpropane-2-sulfinamide

1.3 g of (R,E)-N-((7-fluoro-2-(3-fluorophenyl)quinoline-3-yl)methylene)-2-methylpropane-2-sulfinamide was prepared as a yellow solid by using 1.0 g (3.71 mmol, 1.0 equivalent) of 7-fluoro-2-(3-fluorophenyl)quinoline-3-carbaldehyde prepared in step 1 by the same manner as described in step 2 of Preparative Example 12 (3.49 mmol, yield: 94%).

^1H NMR (300 MHz, CDCl_3) δ 1.31 (s, 9H), 7.18-7.54 (m, 5H), 7.79-7.83 (m, 1H), 7.98-8.03 (m, 1H), 8.76 (s, 1H), 8.91 (s, 1H).

15

Step 3: Preparation of (R)-N-((S)-1-(7-fluoro-2-(3-fluorophenyl)quinoline-3-yl)ethyl)-2-methylpropane-2-sulfinamide

1.30 g of (R)-N-((S)-1-(7-fluoro-2-(3-fluorophenyl)quinoline-3-yl)ethyl)-2-methylpropane-2-sulfinamide was prepared as a pale yellow solid by using 1.3 g (3.49 mmol) of (R,E)-N-((7-fluoro-2-(3-fluorophenyl)quinoline-3-yl)methylene)-2-methylpropane-2-sulfinamide prepared in step 2 by the

same manner as described in step 3 of Preparative Example 12 (3.35 mmol, yield: 96%)


¹H NMR (500 MHz, CDCl₃) δ 1.23 (s, 9H), 1.51-1.53 (d, J = 10.0, 3H), 3.38-3.39 (d, J = 5.0, 1H), 4.92-4.94 (m, 1H), 7.17-7.21 (m, 1H), 7.29-7.32 (m, 1H), 7.38-7.41 (m, 2H), 7.49-7.53 (m, 1H), 7.78-7.80 (m, 1H), 7.85-7.88 (m, 1H), 8.35 (s, 1H).

Step 4: Preparation of (S)-1-(7-fluoro-2-(3-fluorophenyl)quinoline-3-yl)ethane-1-amine

0.37 g of (S)-1-(7-fluoro-2-(3-fluorophenyl)quinoline-3-yl)ethane-1-amine was prepared as a pale yellow solid by using 0.52 g (1.34 mmol) of (R)-N-((S)-1-(7-fluoro-2-(3-fluorophenyl)quinoline-3-yl)ethyl)-2-methylpropane-2-sulfinamide prepared in step 3 by the same manner as described in step 4 of Preparative Example 12 (1.30 mmol, yield: 97%)

¹H NMR (300 MHz, CDCl₃) δ 1.35 (d, J = 9.0, 3H), 1.53 (br s, 2H), 4.43 (t, J = 6.0, 1H), 4.92-4.94 (m, 1H), 7.16 (t, J = 9.0, 1H), 7.30-7.36 (m, 3H), 7.42-7.49 (m, 1H), 7.73 (d, J = 12.0, 1H), 7.82-7.87 (m, 1H), 8.47 (s, 1H).

Preparative Example 17: Preparation of (S)-1-(7-fluoro-2-(pyridine-2-yl)quinoline-3-yl)ethane-1-amine



Step 1: Preparation of 1-(2-chloro-7-fluoroquinoline-3-yl)ethane-1-ol

2.5 g (11.927 mmol) of 2-chloro-7-fluoroquinoline-3-carbaldehyde was dissolved in anhydrous THF (30 mL), to which 4.77 mL (14.312 mmol) of 3 M MeMgBr (Et₂O) solution was added at -78°C, followed by stirring at -78°C ~ -10°C for 2 hours. The temperature was adjusted at -20°C. After adding saturated NH₄Cl aqueous solution, the reaction mixture was heated at room temperature, followed by extraction

using ethyl acetate. The organic layer was separated, dried (Na_2SO_4), filtered, and concentrated under reduced pressure. The residue was separated by column chromatography (SiO_2 , eluent: hexane/ethyl acetate, 3/1) to give 2.4 g of the target compound 1-(2-chloro-7-fluoroquinoline-3-yl)ethane-1-ol as a yellow solid (10.636 mmol, yield: 89%).

^1H NMR (300 MHz, CDCl_3) δ 8.38(s, 1H), 7.79~7.87(m, 1H), 7.63(dd, J = 9.6, 2.2 Hz, 1H), 7.35(td, J = 8.6, 2.4 Hz, 1H), 5.32~5.41(m, 1H), 2.31(d, J = 2.9 Hz, 1H), 1.61(d, J = 7.1 Hz, 3H).

Step 2: Preparation of 1-(2-chloro-7-fluoroquinoline-3-yl)ethane-1-one

2.4 g (10.636 mmol) of 1-(2-chloro-7-fluoroquinoline-3-yl)ethane-1-ol prepared in step 1 was dissolved in 30 mL of anhydrous toluene, to which 9.2 g (106.36 mmol) of manganese dioxide (MnO_2) was added, followed by reflux for 10 hours. The reaction mixture was cooled down at room temperature, filtered with celite pad, concentrated under reduced pressure. The residue was separated by column chromatography (SiO_2 , eluent: hexane/ethyl acetate, 3/1) to give 1.8 g of the target compound 1-(2-chloro-7-

fluoroquinoline-3-yl)ethane-1-one as a yellow solid (8.049 mmol, yield: 76%).

^1H NMR (300 MHz, CDCl_3) δ 8.42 (s, 1H), 7.88~7.95 (m, 1H), 7.68 (dd, $J = 9.8, 2.2$ Hz, 1H), 7.41 (td, $J = 8.4, 2.4$ Hz, 1H), 2.79 (s, 3H).

Step 3: Preparation of (R)-1-(2-chloro-7-fluoroquinoline-3-yl)ethane-1-ol

5 g (15.588 mmol) of B-chlorodiisopinocampheolborane ((+)DIP-Cl) was dissolved in anhydrous THF (10 ml), which was frozen at -47°C . 1.8 g (8.049 mmol) of 1-(2-chloro-7-fluoroquinoline-3-yl)ethane-1-one prepared in step 2 was dissolved in anhydrous THF (20 ml), which was added to the mixture above, followed by stirring at room temperature for 12 hours. The reaction mixture was cooled down at 0°C , to which 1 ml of acetone and 1 ml of 10 % Na_2CO_3 were added, followed by stirring at room temperature for 1 hour. Ethyl acetate and water were added to the reaction mixture. The separated organic layer was dried (Na_2SO_4), filtered, and concentrated under reduced pressure. The residue was separated by column chromatography (SiO_2 , eluent: hexane/ethyl acetate, 4/1) to give 1.4 g of the target compound (R)-1-(2-chloro-7-fluoroquinoline-3-

yl)ethane-1-ol as a white solid (6.204 mmol, yield: 77%).

¹H NMR (300 MHz, CDCl₃) δ 8.39 (s, 1H), 7.81~7.88 (m, 1H), 7.65 (dd, J = 9.9, 2.6 Hz, 1H), 7.36 (td, J = 8.9, 2.8 Hz, 1H), 5.31~5.41 (m, 1H), 2.15 (d, J = 3.8 Hz, 1H), 1.61 (d, J = 6.4 Hz, 3H).

Step 4: Preparation of (S)-2-(1-(2-chloro-7-fluoroquinoline-3-yl)ethyl)isoindoline-1,3-dione

10 1.4 g (6.204 mmol) of (R)-1-(2-chloro-7-fluoroquinoline-3-yl)ethane-1-ol prepared in step 3 was dissolved in anhydrous THF (30 mL), to which 1.95 g (7.445 mmol) of triphenylphosphine (PPh₃) and 1.1 g (7.445 mmol) of phthalimide were added. The mixture
15 was cooled down at 0°C, to which 1.47 mL (7.445 mmol) of diisopropyl azodicarboxylate (DIAD) was added, followed by stirring at room temperature for 15 hours. Water and ethyl acetate were added to the reaction mixture, followed by extraction. The separated
20 organic layer was dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was separated by column chromatography (SiO₂, eluent: hexane/ethyl acetate, 4/1) to give 2 g of the target compound (S)-2-(1-(2-chloro-7-fluoroquinoline-3-

yl)ethyl)isoindoline-1,3-dione as a white solid (5.637 mmol, yield: 91%).

¹H NMR (300 MHz, CDCl₃) δ 8.56(s, 1H), 7.87~7.94(m, 1H), 7.77~7.83(m, 2H), 7.68~7.74(m, 2H), 7.61(dd, J = 9.7, 2.2 Hz, 1H), 7.37(td, J = 8.4, 2.4 Hz, 1H), 5.95(q, J = 7.1, 6.9 Hz, 1H), 1.97(d, J = 7.1 Hz, 3H).

Step 5: Preparation of (S)-2-(1-(7-fluoro-2-(pyridine-2-yl)quinoline-3-yl)ethyl)isoindoline-1,3-dione

1 g (2.819 mmol) of (S)-2-(1-(2-chloro-7-fluoroquinoline-3-yl)ethyl)isoindoline-1,3-dione prepared in step 4 was dissolved in 1,4-dioxane (5 mL), to which 163 mg (0.141 mmol) of Pd(PPh₃)₄ and 1.25 g (3.383 mmol) of 2-(tributylstanyl)-pyridine were added, followed by reflux at 100°C for 3 days under argon atmosphere. Water and ethyl acetate were added to the reaction mixture, followed by extraction. The extracted organic layer was dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was separated by column chromatography (SiO₂, eluent: hexane/ethyl acetate, 4/1) to give 500 mg of the target compound (S)-2-(1-(7-fluoro-2-(pyridine-2-yl)quinoline-3-yl)ethyl)isoindoline-1,3-dione as a white solid (1.258 mmol, yield: 91%).

¹H NMR (300 MHz, CDCl₃) δ 8.69 (s, 1H), 8.65 (d, J = 5.0 Hz, 1H), 7.90~7.97 (m, 1H), 7.60~7.76 (m, 7H), 7.28~7.42 (m, 2H), 6.31 (q, J = 7.4, 7.1 Hz, 1H), 1.98 (d, J = 7.5 Hz, 3H).

5


Step 6: Preparation of (S)-1-(7-fluoro-2-(pyridine-2-yl)quinoline-3-yl)ethane-1-amine

500 mg (1.258 mmol) of (S)-2-(1-(7-fluoro-2-(pyridine-2-yl)quinoline-3-yl)ethyl)isoindoline-1,3-dione prepared in step 5 was dissolved in ethanol (20 mL), to which 612 μL (12.58 mmol) of hydrazine hydrate was added, followed by reflux for 2 hours. The reaction mixture was cooled down at room temperature, and then filtered. The filtrate was added with ethyl acetate and water, followed by extraction. The extracted organic layer was dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was separated by column chromatography (SiO₂, eluent: dichloromethane/methanol, 20/1 -> dichloromethane/methanol, 10/1) to give 312 mg of the target compound (S)-1-(7-fluoro-2-(pyridine-2-yl)quinoline-3-yl)ethane-1-amine as a yellow liquid (1.167 mmol, yield: 93%).

¹H NMR (300 MHz, CDCl₃) δ 8.70 (d, J = 4.6 Hz, 1H), 8.43 (s, 1H), 7.82~7.95 (m, 3H), 7.75 (dd, J = 9.7, 2.4

Hz, 1H), 7.31~7.41(m, 2H), 4.63(q, J = 6.7, 6.7 Hz, 1H), 2.01(br s, 2H), 1.43(d, J = 6.8 Hz, 3H).

Preparative Example 18: Preparation of 1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethane-1-amine



Step 1: Preparation of tert-butyl (S)-(3-oxo-4-(pyridine-2-yl)butane-2-yl)carbamate

5 g (21.526 mmol) of tert-butyl (S)-(1-(methoxy(methyl)amino)-1-oxopropyl)carbamate was dissolved in anhydrous THF (40 mL), to which 16.6 mL (21.526 mmol) of isopropylmagnesium chloride lithium chloride solution was added at -40°C , followed by stirring at -30°C for 30 minutes. The reaction mixture was cooled down to -40°C . 2.6 g (27.984 mmol) of 2-picoline was dissolved in anhydrous THF (20 mL), to which 11 mL (27.984 mmol) of 2.5 M n-BuLi was added

at -40°C, followed by stirring at -20°C for 1 hour. This solution was added to the reaction mixture above, followed by stirring at -20°C ~ -10°C for 3 hours. The reaction mixture was frozen at -78°C, to which
5 saturated ammoniumchloride aqueous solution was added. Ethyl acetate and water were added thereto, followed by extraction. The extracted organic layer was dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was separated by column
10 chromatography (SiO₂, eluent: hexane/ethyl acetate, 4/1) to give 5 g of the target compound tert-butyl (S)-(3-oxo-4-(pyridine-2-yl)butane-2-yl)carbamate as a yellow liquid (18.916 mmol, yield: 99%).

¹H NMR (300 MHz, CDCl₃) δ 8.55(d, J = 4.0 Hz, 1H),
15 7.66(td, J = 7.8, 1.8 Hz, 1H), 7.16~7.24(m, 2H), 5.37(br s, 1H), 4.39~4.49(m, 1H), 3.95~4.11(m, 2H), 1.45(s, 9H), 1.37(d, J = 7.2 Hz, 3H).

Step 2: Preparation of tert-butyl(1-(6-fluoro-3-
20 (pyridine-2-yl)quinoline-2-yl)ethyl)carbamate

254 mg (0.916 mmol) of tert-butyl (S)-(3-oxo-4-(pyridine-2-yl)butane-2-yl)carbamate prepared in step 1, 134 mg (0.961mmol) of 2-amino-5-fluorobenzaldehyde, and 398 mg (2.883 mmol) of potassiumcarbonate (K₂CO₃)
25 were dissolved in ethanol (3 mL), followed by stirring

at 90°C for 2 hours. The reaction mixture was added with ethyl acetate and water, followed by extraction. The extracted organic layer was dried (Na₂SO₄), filtered, and concentrated under reduced pressure.

5 The residue was separated by column chromatography (SiO₂, eluent: hexane/ethyl acetate, 5/1) to give 250 mg of the target compound tert-butyl(1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)carbamate as a yellow solid (0.680 mmol, yield: 71%).

10 ¹H NMR (300 MHz, CDCl₃) δ 8.76(d, J = 4.5 Hz, 1H), 8.08~8.15(m, 2H), 7.84(td, J = 7.9, 1.9 Hz, 1H), 7.41~7.60(m, 3H), 7.35(t, J = 4.5 Hz, 1H), 6.34(d, J = 7.6 Hz, 1H), 5.37~5.48(m, 1H), 1.45(s, 9H), 1.33(d, J = 6.3 Hz, 3H).

15


Step 3: Preparation of 1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethane-1-amine

250 mg (0.680 mmol) of tert-butyl(1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)carbamate prepared
20 in step 2 was dissolved in dichloromethane (3 mL), to which 1 mL of TFA was added, followed by stirring at room temperature for 3 hours. The reaction mixture was filtered under reduced pressure and neutralized with NaHCO₃ aqueous solution. The reaction mixture
25 was added with dichloromethane and water, followed by

extraction. The extracted organic layer was separated, dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was separated by column chromatography (SiO₂, eluent: hexane/ethyl acetate, 5/1) to give 120 mg of the target compound 1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethane-1-amine as a yellow oil (0.449 mmol, yield: 66%).

¹H NMR (500 MHz, CDCl₃) δ 8.75(d, J = 4.1 Hz, 1H), 8.01~8.15(m, 2H), 7.84(td, J = 7.8, 1.5 Hz, 1H), 7.40~7.50(m, 3H), 7.33~7.39(m, 1H), 4.49(br s, 1H), 2.11(br s, 2H), 1.39(d, J = 5.6 Hz, 3H).

Preparative Example 19: Preparation of (S)-3-(1-aminoethyl)-8-chloro-4-fluoro-2-phenylisoquinoline-1(2H)-one



Step 1: Preparation of (S)-N-(1-(8-chloro-4-fluoro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)-2,2,2-trifluoroacetamide

1.97 g (5.0 mmol, 1 equivalent) of (S)-N-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)-2,2,2-trifluoroacetamide prepared in step 6

of Preparative Example 10, Selectfluor (1.5
equivalent), and anhydrous CH₃CN (30 mL) were mixed,
which was refluxed for 12 hours. The reaction mixture
was cooled down at room temperature. Water and ethyl
5 acetate were added thereto, followed by extraction.
The extracted organic layer was washed with saturated
NaHCO₃ aqueous solution, separated, dried (Na₂SO₄),
filtered, and concentrated under reduced pressure.
The residue was separated by column chromatography
10 (SiO₂, eluent: hexane/ethyl acetate, 10/1 ->
hexane/ethyl acetate, 3/1) to give 1.61 g of the
target compound (S)-N-(1-(8-chloro-4-fluoro-1-oxo-2-
phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)-2,2,2-
trifluoroacetamide as a white solid (3.9 mmol, yield:
15 78%).

¹H NMR(500 MHz, CDCl₃) δ 10.99 (br d, J = 5.4Hz,
1H), 7.77-7.85 (m, 2H), 7.64-7.71 (m, 1H), 7.50-7.61
(m, 3H), 7.42-7.46 (m, 2H), 4.17-4.24 (m, 1H), 1.47
(d, J = 7.1Hz, 3H).

20


Step 2: Preparation of (S)-3-(1-aminoethyl)-8-
chloro-4-fluoro-2-phenylisoquinoline-1(2H)-one

1.20 g of (S)-3-(1-aminoethyl)-8-chloro-4-fluoro-
2-phenylisoquinoline-1(2H)-one was prepared as a white
25 solid by using 1.65 g (4.0 mmol) of (S)-N-(1-(8-

chloro-4-fluoro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)-2,2,2-trifluoroacetamide prepared in step 1 by the same manner as described in step 8 of Preparative Example 5 10 (3.8 mmol, yield: 95%).

^1H NMR (500 MHz, CDCl_3) δ 7.74 (d, j = 8.0Hz, 1H), 7.64 (t, J = 7.9Hz, 1H), 7.54-7.60 (m, 3H), 7.47-7.53 (m, 1H), 7.32-7.35 (m, 1H), 7.22-7.25 (m, 1H), 3.57-3.64 (m, 1H), 1.85 (br s, 2H), 1.46 (d, J = 6.9Hz, 10 3H).

Preparative Example 20: Preparation of 4-chloro-8-(4-methoxybenzyl)-7,8-dihydropyrido[2,3-d]pyrimidine-5(6H)-one



Step 1: Preparation of 1-(4,6-dichloropyrimidine-5-yl)propene-2-en-1-ol

200 mg (2.8 mmol) of 4,6-dichloropyrimidine-5-carbaldehyde was dissolved in anhydrous toluene (15 20 mL), to which 2.1 mL (1.2 equivalent) of vinylmagnesium chloride (1.6 M in THF) was slowly added at -20°C , followed by stirring for 1 hour. Saturated NH_4Cl aqueous solution (10 mL) was added

thereto. The reaction mixture was extracted by using ethyl acetate. The extracted organic layer was washed with saturated brine, separated, dried (Na_2SO_4), filtered, and concentrated under reduced pressure.

5 The residue was separated by column chromatography (SiO_2 , eluent: hexane/ethyl acetate, 10/1) to give 475 mg of the target compound 1-(4,6-dichloropyrimidine-5-yl)propene-2-en-1-ol as a yellow oil (2.3 mmol, yield: 82%).

10 ^1H NMR (300 MHz, CDCl_3) δ 8.72 (s, 1H), 6.23-6.12 (m, 1H), 5.90 (s, -OH), 5.43-5.34 (m, 2H).

Step 2: Preparation of 1-(4,6-dichloropyrimidine-5-yl)propene-2-en-1-one

15 323 mg of 1-(4,6-dichloropyrimidine-5-yl)propene-2-en-1-one was prepared as a colorless oil by using 394 mg (1.9 mmol) of 1-(4,6-dichloropyrimidine-5-yl)propene-2-en-1-ol prepared in step 1 by the same manner as described in step 3 of Example 1 (1.57 mmol,

20 yield: 83%).

^1H NMR (300 MHz, CDCl_3) δ 8.87 (s, 1H), 6.69-6.59 (m, 1H), 6.31 (d, $J = 10.6$ Hz, 1H), 6.08 (d, $J = 17.9$ Hz, 1H).


Step 3: Preparation of 4-chloro-8-(4-methoxybenzyl)-7,8-dihydropyrido[2,3-d]pyrimidine-5(6H)-one

4-methoxybenzylamine (1.1 equivalent) was dissolved in anhydrous CH_2Cl_2 (5 mL), which was slowly added to the reaction mixture comprising 300 mg (1.48 mmol) of 1-(4,6-dichloropyrimidine-5-yl)propene-2-en-1-one prepared in step 2, DIPEA (1.1 equivalent), and anhydrous CH_2Cl_2 (15 mL) at 0°C . The mixture was heated at room temperature, followed by stirring for 1 hour. The reaction mixture was added with 1N HCl (5 mL), followed by extraction with ethyl acetate. The extracted organic layer was washed with saturated NaHCO_3 aqueous solution, separated, dried (Na_2SO_4), filtered, and concentrated under reduced pressure. The residue was separated by column chromatography (SiO_2 , eluent: hexane/ethyl acetate, 2/1) to give 413 mg of the target compound 4-chloro-8-(4-methoxybenzyl)-7,8-dihydropyrido[2,3-d]pyrimidine-5(6H)-one as a pale yellow solid (1.36 mmol, yield: 92%).

^1H NMR (300 MHz, CDCl_3) δ 8.72 (s, 1H), 6.23-6.12 (m, 1H), 5.90

^1H NMR (300 MHz, CDCl_3) δ 8.40 (s, 1H), 7.22 (d, J = 8.5 Hz, 2H), 6.87 (d, J = 8.5 Hz, 2H), 4.92 (s, 2H), 3.80 (s, 3H), 3.56 (m, 2H), 2.96 (m, 2H).

5 Preparative Example 21: Preparation of (S)-1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethane-1-amine



Step 1: Preparation of 1-(3-(benzyloxy)-6-fluoroquinoline-2-yl)ethane-1-one

10 20.52 g (100.0 mmol) of 1-(6-fluoro-3-hydroxyquinoline-2-yl)ethane-1-one [reference: WO 2010-151740], BnBr (1.1 equivalent), and K_2CO_3 (3 equivalent) were dissolved in anhydrous DMF (150 mL), followed by stirring at room temperature for 6 hours.

15 The reaction solvent was eliminated under reduced pressure. Water and ethyl acetate were added to the reaction mixture, followed by extraction. The

extracted organic layer was washed with saturated brine, separated, dried (Na_2SO_4), filtered, and concentrated under reduced pressure. The residue was separated by column chromatography (SiO_2 , eluent: 5 hexane/ethyl acetate, 10/1 $\rightarrow$ hexane/ethyl acetate, 3/1) to give 29.53 g of the target compound 1-(3-(benzyloxy)-6-fluoroquinoline-2-yl)ethane-1-one as a pale brown oil (100.0 mmol, yield: 100%).

^1H NMR (300 MHz, CDCl_3) δ 8.00-8.10 (m, 1H), 7.28-10 7.57 (m, 8H), 5.26 (s, 2H), 2.76 (s, 3H).

Step 2: Preparation of 3-(benzyloxy)-6-fluoroquinoline-2-carboxylic acid

28.06 g (95.0 mmol) of 1-(3-(benzyloxy)-6-15 fluoroquinoline-2-yl)ethane-1-one prepared in step 1 was dissolved in dioxane/ H_2O (4/1, 300 mL), to which NaOCl aqueous solution (12%, 5 equivalent) was slowly added at room temperature for 30 minutes, followed by stirring for 5 hours. PH of the reaction mixture was 20 adjusted (pH=4) with saturated 2 N HCl solution. The reaction mixture was extracted by using ethyl acetate. The extracted organic layer was washed with saturated brine, separated, dried (Na_2SO_4), filtered, and concentrated under reduced pressure. The residue was 25 separated by column chromatography (SiO_2 , eluent:

hexane/ethyl acetate, 1/1 -> ethylacetate) to give 27.96 g of the target compound 3-(benzyloxy)-6-fluoroquinoline-2-carboxylic acid as a pale yellow oil (94.1 mmol, yield: 99%).

5 ^1H NMR (300 MHz, CDCl_3) δ 8.05-8.12 (m, 1H), 7.67 (s, 1H), 7.28-7.60 (m, 7H), 5.37 (s, 2H).

Step 3: Preparation of methyl 3-(benzyloxy)-6-fluoroquinoline-2-carboxylate

10 26.76 g (90.0 mmol) of 3-(benzyloxy)-6-fluoroquinoline-2-carboxylic acid prepared in step 2, anhydrous MeOH (200 mL), $\text{CH}(\text{OMe})_3$ (50 mL), and conc H_2SO_4 (2 mL) were mixed, followed by heating at 45°C for 12 hours. The reaction mixture was slowly added
15 to cold saturated NaHCO_3 aqueous solution, followed by extraction with ethyl acetate. The extracted organic layer was washed with saturated brine, separated, dried (Na_2SO_4), filtered, and concentrated under reduced pressure. The residue was separated by column
20 chromatography (SiO_2 , eluent: hexane/ethyl acetate, 10/1 -> hexane/ethylacetate, 3/1) to give 27.46 g of the target compound methyl 3-(benzyloxy)-6-fluoroquinoline-2-carboxylate as a pale yellow solid (88.2 mmol, yield: 98%).

¹H NMR (300 MHz, CDCl₃) δ 8.07-8.14 (m, 1H), 7.50 (s, 1H), 7.28-7.49 (m, 7H), 5.28 (s, 2H), 4.04 (s, 3H).

5 Step 4: Preparation of (3-(benzyloxy)-6-fluoroquinoline-2-yl)methanol

14.17 g (50.0 mmol) of methyl 3-(benzyloxy)-6-fluoroquinoline-2-carboxylate prepared in step 3 was dissolved in anhydrous THF (200 mL). The reaction
10 mixture was cooled down to 0°C, to which LiAlH₄ (1 equivalent) was slowly added for 10 minutes, followed by stirring for 1 hour. The reaction mixture was heated at room temperature, followed by stirring for 5 hours. The reaction mixture was added with
15 diethylether (200 mL) and distilled water (10 mL) slowly to degrade LiAlH₄, followed by stirring for 1 hour. The mixture was added with anhydrous MgSO₄, dried, filtered, and concentrated. The obtained compound was separated by column chromatography (SiO₂,
20 eluent: hexane/ethyl acetate, 10/1 -> hexane/ethylacetate, 1/1) to give 12.04 g of the target compound (3-(benzyloxy)-6-fluoroquinoline-2-yl)methanol as a pale yellow solid (42.5 mmol, yield: 85%).

¹H NMR (300 MHz, CDCl₃) δ 7.97-8.05 (m, 1H), 7.27-7.46 (m, 8H), 5.22 (s, 2H), 4.93 (d, J = 4.6Hz, 2H), 4.54 (t, J = 4.6Hz, 1H, OH).

5 Step 5: Preparation of 3-(benzyloxy)-6-fluoroquinoline-2-carbaldehyde

11.91 g of 3-(benzyloxy)-6-fluoroquinoline-2-carbaldehyde was prepared as a pale yellow solid by Swern oxidation using 12.00 g (42.4 mmol) of (3-
10 (benzyloxy)-6-fluoroquinoline-2-yl)methanol prepared in step 4 (42.3 mmol, yield: 100%).

¹H NMR (300 MHz, CDCl₃) δ 10.53 (s, 1H), 8.20 (m, 1H), 7.59 (s, 1H), 7.25-7.58 (m, 7H), 5.32 (s, 2H).

15 Step 6: Preparation of (S)-(E)-((3-(benzyloxy)-6-fluoroquinoline-2-yl)methylene)-2-methylpropane-2-sulfinamide

5.63 g (20.00 mmol) of 3-(benzyloxy)-6-fluoroquinoline-2-carbaldehyde prepared in step 5,
20 (S)-(-)-2-methyl-2-propanesulfinamide (1.1 equivalent), and Cs₂CO₃ (1.2 equivalent) were dissolved in anhydrous CH₂Cl₂ (30 mL), followed by stirring at room temperature for 12 hours. The reaction mixture was filtered and concentrated. The obtained compound
25 was separated by column chromatography (SiO₂, eluent:

hexane/ethyl acetate, 7/1 -> hexane/ethylacetate, 3/1)
to give 7.30 g of the target compound (S)-(E)-((3-(benzyloxy)-6-fluoroquinoline-2-yl)methylene)-2-methylpropane-2-sulfinamide as a pale yellow solid
5 (18.99 mmol, yield: 95%).

¹H NMR (300 MHz, CDCl₃) δ 9.22 (s, 1H), 8.14-8.20 (m, 1H), 7.53 (s, 1H), 7.28-7.50 (m, 7H), 5.30 (s, 2H), 1.28 (s, 9H).

10 Step 7: Preparation of (S)-N-((S)-1-(3-(benzyloxy)-6-fluoroquinoline-2-yl)ethyl)-2-methylpropane-2-sulfinamide

3.85 g (10.00 mmol) of (S)-(E)-((3-(benzyloxy)-6-fluoroquinoline-2-yl)methylene)-2-methylpropane-2-sulfinamide prepared in step 6 was dissolved in
15 anhydrous CH₂Cl₂ (50 mL), which was frozen at -78°C. MeMgBr (3 M diethylether solution, 3 equivalent) was slowly added thereto for 10 minutes. 2 hours later, the reaction mixture was slowly heated to -20°C,
20 followed by stirring for 1 hour. Saturated NH₄Cl aqueous solution (50 mL) was added thereto. The reaction mixture was heated at room temperature, followed by extraction with ethyl acetate. The extracted organic layer was washed with saturated
25 brine, separated, dried (Na₂SO₄), filtered, and

concentrated under reduced pressure. The residue was separated by column chromatography (SiO₂, eluent: hexane/ethyl acetate, 3/1 → hexane/ethylacetate, 1/2) to give 3.00 g of the target compound (S)-N-((S)-1-(3-(benzyloxy)-6-fluoroquinoline-2-yl)ethyl)-2-methylpropane-2-sulfinamide as a pale yellow solid (7.40 mmol, yield: 74%).

¹H NMR (500 MHz, CDCl₃) δ 7.97-8.01 (m, 1H), 7.45-7.52 (m, 4H), 7.38-7.44 (m, 2H), 7.28-7.33 (m, 2H), 5.60 (d, J = 6.5 Hz, 1H), 5.24 (s, 2H), 5.08-5.13 (m, 1H), 1.53 (d, J = 6.7 Hz, 3H), 1.32 (s, 9H).

Step 8: Preparation of (S)-1-(3-(benzyloxy)-6-fluoroquinoline-2-yl)ethane)-1-amine

2.81 g (7.02 mmol) of (S)-N-((S)-1-(3-(benzyloxy)-6-fluoroquinoline-2-yl)ethyl)-2-methylpropane-2-sulfinamide prepared in step 7 was dissolved in anhydrous MeOH (10 mL), to which 4M HCl (dioxane solution) was added at room temperature, followed by stirring for 1 hour. The solvent was eliminated under reduced pressure. The mixture was added slowly with saturated NaHCO₃ aqueous solution, followed by extraction with ethyl acetate. The extracted organic layer was washed with saturated brine, separated, dried (Na₂SO₄), filtered, and

concentrated under reduced pressure. The obtained compound was separated by column chromatography (SiO₂, eluent: MeOH/CH₂Cl₂, 1/20 -> MeOH/CH₂Cl₂, 1/10) to give 2.00 g of the target compound (S)-1-(3-(benzyloxy)-6-fluoroquinoline-2-yl)ethane)-1-amine as a pale white solid (6.75 mmol, yield: 96%).

¹H NMR (300 MHz, CDCl₃) δ 7.96-8.02 (m, 1H), 7.25-7.50 (m, 8H), 5.21 (s, 2H), 4.60-4.70 (m, 1H), 2.04 (br s, 2H), 1.47 (d, J = 6.6 Hz, 3H).

10

Step 9: Preparation of tert-butyl (S)-1-(3-(benzyloxy)-6-fluoroquinoline-2-yl)ethyl)carbamate

1.90 g (6.41 mmol) of (S)-1-(3-(benzyloxy)-6-fluoroquinoline-2-yl)ethane)-1-amine prepared in step 8 was dissolved in anhydrous CH₂Cl₂ (15 mL), to which Boc₂O (1.3 equivalent) was added at room temperature, followed by stirring for 2 hours. The reaction mixture was concentrated under reduced pressure. The obtained compound was separated by column chromatography (SiO₂, eluent: hexane/ethyl acetate, 10/1 -> hexane/ethylacetate, 3/1) to give 2.52 g of the target compound tert-butyl (S)-1-(3-(benzyloxy)-6-fluoroquinoline-2-yl)ethyl)carbamate as a white solid (6.36 mmol, yield: 99%).

¹H NMR(300 MHz, CDCl₃) δ 8.00-8.04 (m, 1H), 7.30-7.52 (m, 8H), 6.44 (br d, J = 7.6Hz, 1H), 5.38-5.45 (m, 1H), 5.20-5.27 (m, 2H), 1.47-1.56 (m, 12H).

5 Step 10: Preparation of tert-butyl (S)-(1-(6-fluoro-3-hydroxyquinoline-2-yl)ethyl)carbamate

1.98 g (4.99 mmol) of tert-butyl (S)-1-(3-(benzyloxy)-6-fluoroquinoline-2-yl)ethylcarbamate prepared in step 9 was dissolved in MeOH (20 mL), to
10 which 200 mg of 10% Pd/C was added, followed by hydrogenation at room temperature for 1 hour under 1 atm H₂. The reaction mixture was filtered with celite pad and concentrated under reduced pressure. The obtained compound was separated by column
15 chromatography (SiO₂, eluent: hexane/ethyl acetate, 3/1 -> hexane/ethylacetate, 1/1) to give 1.53 g of the target compound tert-butyl (S)-(1-(6-fluoro-3-hydroxyquinoline-2-yl)ethyl)carbamate as a white solid (4.99 mmol, yield: 100%).

20 ¹H NMR(300 MHz, CDCl₃) δ 9.47 (br s, 1H), 7.87-7.90 (m, 1H), 7.18-7.23 (m, 1H), 7.00-7.04 (m, 1H), 6.84-6.89 (m, 1H), 5.52 (br s, 1H), 5.29-5.36 (m, 1H), 1.62 (d, J = 6.8Hz, 3H), 1.53 (s, 9H).

Step 11: Preparation of (S)-2-(1-((tert-butoxycarbonyl)amino)ethyl)-6-fluoroquinoline-3-yl trifluoromethanesulfonate

1.50 g (4.90 mmol) of tert-butyl (S)-(1-(6-fluoro-3-hydroxyquinoline-2-yl)ethyl)carbamate prepared in step 10 and anhydrous Et₃N (3 equivalent) were dissolved in anhydrous CH₂Cl₂ (15 mL), to which Tf₂O (1.2 equivalent) was slowly added at 0°C for 5 minutes, followed by stirring for 2 hours. The solvent was eliminated under reduced pressure. The reaction mixture was added with water, followed by extraction with ethyl acetate. The extracted organic layer was washed with saturated brine, separated, dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The obtained compound was separated by column chromatography (SiO₂, eluent: hexane/ethyl acetate, 10/1 → hexane/ethylacetate, 5/1) to give 2.15 g of the target compound (S)-2-(1-((tert-butoxycarbonyl)amino)ethyl)-6-fluoroquinoline-3-yl trifluoromethanesulfonate as a colorless oil (4.90 mmol, yield: 100%).

¹H NMR (300 MHz, CDCl₃) δ 8.11-8.17 (m, 1H), 8.08 (s, 1H), 7.53-7.62 (m, 1H), 7.47-7.52 (m, 1H), 7.98 (br s, 1H), 5.30-5.40 (m, 1H), 1.54 (d, J = 6.7 Hz, 3H), 1.48 (s, 9H).

Step 12: Preparation of tert-butyl (S)-(1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)carbamate

438 mg (1.00 mmol) of (S)-2-(1-((tert-butoxycarbonyl)amino)ethyl)-6-fluoroquinoline-3-yl trifluoromethanesulfonate prepared in step 11, 2-(tributylstanyl)pyridine (2.0 equivalent), LiCl (3 equivalent), Pd(Ph₃P)₄ (0.1 equivalent), and anhydrous dioxane (13 mL) were mixed, which was heated at 100°C for 24 hours under argon atmosphere. The reaction mixture was cooled down at room temperature and filtered with celite pad. The filtrate was added with water, followed by extraction with ethyl acetate. The extracted organic layer was washed with saturated brine, separated, dried (Na₂SO₄), and concentrated under reduced pressure. The obtained compound was separated by column chromatography (SiO₂, eluent: hexane/ethyl acetate, 10/1 → hexane/ethylacetate, 3/1) to give 286 mg of the target compound tert-butyl (S)-(1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)carbamate as a pale yellow solid (0.78 mmol, yield: 78%).


¹H NMR (300 MHz, CDCl₃) δ 8.76 (d, J = 4.5 Hz, 1H), 8.08-8.15 (m, 2H), 7.84 (td, J = 7.9, 1.9 Hz, 1H), 7.41-7.60 (m, 3H), 7.35 (t, J = 4.5 Hz, 1H), 6.34 (d, J

= 7.6Hz, 1H), 5.37-5.48 (m, 1H), 1.45 (s, 9H), 1.33 (d, J = 6.3Hz, 3H).

Step 13: Preparation of (S)-1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethane-1-amine

(S)-1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethane-1-amine was prepared by using tert-butyl (S)-1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethylcarbamate prepared in step 12 by the same
10 manner as described in step 3 of Preparative Example 18.

Preparative Example 22: Preparation of 1-(6-fluoro-3,4-di(pyridine-2-yl)quinoline-2-yl)ethane-1-
15 amine



Step 1: Preparation of (5-fluoro-2-nitrophenyl)(pyridine-2-yl)methanol

1.90 g (12.0 mmol) of 2-bromopyridine was dissolved in anhydrous THF (20 mL), which was cooled down to 0°C. Isopropylmagnesium chloride lithium chloride complex solution (1.3M THF solution, 1.2
5 equivalent) was slowly added thereto for 5 minutes, followed by stirring for 1 hour. 1.69 g (10.0 mmol) of 3-fluoro-6-nitrobenzaldehyde was dissolved in anhydrous THF (10 mL), which was slowly added to the mixture above for 10 minutes, followed by stirring for
10 1 hour. The reaction mixture was heated to room temperature, followed by stirring for 2 hours. Saturated NH₄Cl solution (20 mL) was added thereto, followed by extraction with ethyl acetate. The extracted organic layer was washed with saturated
15 brine, separated, dried (Na₂SO₄), and concentrated under reduced pressure. The obtained compound was separated by column chromatography (SiO₂, eluent: hexane/ethyl acetate, 10/1 -> hexane/ethylacetate, 5/1) to give 1.61 g of the target compound (5-fluoro-
20 2-nitrophenyl)(pyridine-2-yl)methanol as a pale yellow solid (6.5 mmol, yield: 65%).

¹H NMR (300 MHz, CDCl₃) δ 8.56-8.60 (m, 1H), 8.01-8.06 (m, 1H), 7.65-7.72 (m, 1H), 7.32-7.41 (m, 2H), 7.23-7.30 (m, 1H), 7.05-7.13 (m, 1H), 6.51 (s, 1H),
25 5.44 (br s, 1H).

Step 2: Preparation of (5-fluoro-2-nitrophenyl)(pyridine-2-yl)methanone

1.50 g (6.04 mmol) of (5-fluoro-2-nitrophenyl)(pyridine-2-yl)methanol prepared in step 1 was dissolved in anhydrous CH_2Cl_2 (30 mL), to which MnO_2 (6 equivalent) was added, followed by stirring at room temperature for 5 hours. The reaction mixture was filtered with celite pad and concentrated under reduced pressure. The obtained compound was separated by column chromatography (SiO_2 , eluent: hexane/ethyl acetate, 10/1 $\rightarrow$ hexane/ethylacetate, 4/1) to give 1.48 g of the target compound (5-fluoro-2-nitrophenyl)(pyridine-2-yl)methanone as a pale brown solid (6.01 mmol, yield: 100%).

^1H NMR (300 MHz, CDCl_3) δ 8.49-8.52 (m, 1H), 8.22-8.29 (m, 2H), 7.88-7.96 (m, 1H), 7.43-7.48 (m, 1H), 7.23-7.36 (m, 2H).

Step 3: Preparation of (2-amino-5-fluorophenyl)(pyridine-2-yl)methanone

1.40 g (5.69 mmol) of (5-fluoro-2-nitrophenyl)(pyridine-2-yl)methanone and Fe (5 equivalent) were dissolved in EtOH/ H_2O (4/1, 30 mL), to which 2~3 drops of conc HCl were added. The

reaction mixture was heated at 85°C for 30 minutes and cooled down to room temperature. The reaction mixture was filtered with celite pad, followed by extraction with ethyl acetate. The extracted organic layer was washed with saturated brine, separated, dried (Na₂SO₄), and concentrated under reduced pressure. The obtained compound was separated by column chromatography (SiO₂, eluent: hexane/ethyl acetate, 10/1 -> hexane/ethylacetate, 4/1) to give 1.23 g of the target compound (2-amino-5-fluorophenyl)(pyridine-2-yl)methanone as a pale yellow solid (5.69 mmol, yield: 100%).

¹H NMR (300 MHz, CDCl₃) δ 8.71 (d, J = 4.7 Hz, 1H), 7.92-7.80 (m, 2H), 7.52-7.43 (m, 2H), 7.12-7.05 (m, 1H), 6.66-6.72 (m, 1H), 6.13 (br s, 2H).

Step 4: Preparation of tert-butyl (1-(6-fluoro-3,4-di(pyridine-2-yl)quinoline-2-yl)ethyl)carbamate

1.08 g (5.0 mmol) of (2-amino-5-fluorophenyl)(pyridine-2-yl)methanone prepared in step 3, tert-butyl (S)-3-oxo-4-(pyridine-2-yl)carbamate (1.0 equivalent), and InCl₃ (0.2 equivalent) were added to anhydrous CH₃CN (10 mL), which was heated at 80°C for 15 minutes and cooled down to room temperature. The reaction mixture was filtered with

celite pad, followed by extraction with ethyl acetate. The extracted organic layer was washed with saturated brine, separated, dried (Na₂SO₄), and concentrated under reduced pressure. The obtained compound was
5 separated by column chromatography (SiO₂, eluent: hexane/ethyl acetate, 10/1 -> hexane/ethylacetate, 3/1) to give 2.22 g of the target compound tert-butyl (1-(6-fluoro-3,4-di(pyridine-2-yl)quinoline-2-yl)ethyl)carbamate as a white solid (4.99 mmol, yield:
10 100%).

¹H NMR(300 MHz, CDCl₃) δ 8.60-8.67 (m, 2H), 8.15-8.21 (m, 1H), 7.45-7.55 (m, 3H), 7.10-7.21 (m, 3H), 7.05 (br t, J = 6.8Hz, 2H), 6.35 (br s, 1H), 5.03 (br s, 1H), 1.44 (s, 9H).

15


Step 5: Preparation of 1-(6-fluoro-3,4-di(pyridine-2-yl)quinoline-2-yl)ethane-1-amine

133 mg (0.3 mmol) of tert-butyl (1-(6-fluoro-3,4-di(pyridine-2-yl)quinoline-2-yl)ethyl)carbamate
20 prepared in step 4 was dissolved in CH₂Cl₂ (10 mL), to which TFA (1.0 mL) was added, followed by stirring at room temperature for 1 hour. The reaction mixture was added with saturated NaHCO₃ aqueous solution (25 mL), followed by extraction with ethyl acetate. The
25 extracted organic layer was washed with saturated

brine, separated, dried (Na_2SO_4), and concentrated under reduced pressure. The obtained compound was separated by column chromatography (SiO_2 , eluent: $\text{MeOH}/\text{CH}_2\text{Cl}_2$, 1/20 $\rightarrow$ $\text{MeOH}/\text{CH}_2\text{Cl}_2$, 1/10) to give 117 mg of the target compound 1-(6-fluoro-3,4-di(pyridine-2-yl)quinoline-2-yl)ethane-1-amine as a white solid (0.3 mmol, yield: 100%).

^1H NMR (300 MHz, CDCl_3) δ 8.66 (dd, $J = 12.0$, 4.1 Hz, 2H), 8.11 (dd, $J = 9.0$, 5.5 Hz, 1H), 7.61-7.46 (m, 3H), 7.24-7.16 (m, 3H), 7.04-6.95 (m, 2H), 4.78 (br s, 1H), 2.54 (br s, 2H), 1.51 (d, $J = 6.1$ Hz, 3H).

Preparative Example 23: Preparation of (S)-1-(6-fluoro-3-phenyl-4-(pyridine-2-yl)quinoline-2-yl)ethane-1-amine



Step 1: Preparation of 1,1-diethoxy-3-phenylpropane-2-one

5.29 g (30.0 mmol) of ethyl diethoxyacetate was dissolved in anhydrous THF (50 mL), which was frozen at -78°C. PhMgCl (2M THF solution, 1.5 equivalent) was added slowly thereto for 5 minutes, followed by stirring for 12 hours. While cooling the reaction mixture with ice water, saturated NH₄Cl aqueous solution (50 mL) was slowly added thereto. The reaction mixture was extracted by using ethyl acetate. The extracted organic layer was washed with saturated brine, separated, dried (Na₂SO₄), and concentrated under reduced pressure. The obtained compound was separated by column chromatography (SiO₂, eluent: hexane/ethyl acetate, 20/1 -> hexane/ethylacetate, 5/1) to give 6.21 g of the target compound 1,1-diethoxy-3-phenylpropane-2-one as a colorless oil (27.9 mmol, yield: 93%).

¹H NMR (300 MHz, CDCl₃) δ 7.18-7.35 (m, 5H), 4.63 (s, 1H), 3.89 (s, 2H), 3.63-3.71 (m, 2H), 3.47-3.61 (m, 2H), 1.19-1.27 (m, 6H).

20

Step 2: Preparation of 2-(diethoxymethyl)-6-fluoro-3-phenyl-4-(pyridine-2-yl)quinoline

3.33 g (15.0 mmol) of 1,1-diethoxy-3-phenylpropane-2-one prepared in step 1, 1.62 g (7.49 mmol) of (2-amino-5-fluorophenyl)(pyridine-2-

yl)methanone, and Cs_2CO_3 (30.0 mmol) were added to EtOH (40 mL), followed by reflux for 12 hours. The reaction mixture was cooled down to room temperature. The reaction mixture was added with water, followed by
5 extraction with ethyl acetate. The extracted organic layer was washed with saturated brine, separated, dried (Na_2SO_4), and concentrated under reduced pressure. The obtained compound was separated by column chromatography (SiO_2 , eluent: hexane/ethyl
10 acetate, 10/1 $\rightarrow$ hexane/ethylacetate, 3/1) to give 3.02 g of the target compound 2-(diethoxymethyl)-6-fluoro-3-phenyl-4-(pyridine-2-yl)quinoline as a white solid (7.50 mmol, yield: 100%).

^1H NMR (300 MHz, CDCl_3) δ 8.64-8.67 (m, 1H), 8.32-
15 8.38 (m, 1H), 7.44-7.54 (m, 2H), 7.14-7.26 (m, 6H), 7.03-7.08 (m, 1H), 6.93-6.97 (m, 1H), 5.37 (s, 1H), 3.55-3.80 (m, 2H), 3.39-3.46 (m, 2H), 1.17 (t, J = 7.0Hz, 6H).

20 Step 3: Preparation of 6-fluoro-3-phenyl-4-(pyridine-2-yl)quinoline-2-carbaldehyde

2.01 g (4.99 mmol) of 2-(diethoxymethyl)-6-fluoro-3-phenyl-4-(pyridine-2-yl)quinoline prepared in step 2 was dissolved in THF (20 mL), to which 2N HCl
25 aqueous solution (20 mL) was added, followed by

stirring at room temperature for 3 hours. Saturated NaHCO₃ aqueous solution was slowly added thereto, followed by extraction with ethyl acetate. The extracted organic layer was washed with saturated brine, separated, dried (Na₂SO₄), and concentrated under reduced pressure. The obtained compound was separated by column chromatography (SiO₂, eluent: hexane/ethyl acetate, 10/1 -> hexane/ethylacetate, 3/1) to give 1.64 g of the target compound 6-fluoro-3-phenyl-4-(pyridine-2-yl)quinoline-2-carbaldehyde as a pale yellow solid (4.99 mmol, yield: 100%).

¹H NMR(300 MHz, CDCl₃) δ 10.14 (s, 1H), 8.69-8.72 (m, 1H), 8.38-8.44 (m, 1H), 7.53-7.64 (m, 2H), 7.11-7.27 (m, 7H), 6.96-7.00 (m, 1H).

15

Step 4: Preparation of (S,E)-N-((6-fluoro-3-phenyl-4-(pyridine-2-yl)quinoline-2-yl)methylene)-2-methylpropane-2-sulfinamide

328 mg (1.0 mmol) of 6-fluoro-3-phenyl-4-(pyridine-2-yl)quinoline-2-carbaldehyde prepared in step 3 was reacted by the same manner as described in step 6 of Preparative Example 21. The obtained compound was separated by column chromatography (SiO₂, eluent: hexane/ethyl acetate, 4/1 -> hexane/ethylacetate, 1/1) to give 418 mg of the target

compound (S,E)-N-((6-fluoro-3-phenyl-4-(pyridine-2-yl)quinoline-2-yl)methylene)-2-methylpropane-2-sulfinamide as a pale yellow solid (0.97 mmol, yield: 97%).

5 ^1H NMR (500 MHz, CDCl_3) δ 8.70-8.72 (m, 2H), 8.38-8.40 (m, 1H), 7.56-7.58 (m, 2H), 6.95-7.28 (m, 8H), 1.18 (s, 9H).

Step 5: Preparation of (S)-N-((S)-1-(6-fluoro-3-phenyl-4-(pyridine-2-yl)quinoline-2-yl)ethyl)-2-methylpropane-2-sulfinamide

388 mg (0.90 mmol) of (S,E)-N-((6-fluoro-3-phenyl-4-(pyridine-2-yl)quinoline-2-yl)methylene)-2-methylpropane-2-sulfinamide prepared in step 4 was
15 reacted by the same manner as described in step 7 of Preparative Example 21. The obtained compound was separated by column chromatography (SiO_2 , eluent: CH_2Cl_2 /ethyl acetate, 4/1 $\rightarrow$ CH_2Cl_2 /ethylacetate, 1/1) to give 306 mg of the target compound (S)-N-((S)-1-(6-
20 fluoro-3-phenyl-4-(pyridine-2-yl)quinoline-2-yl)ethyl)-2-methylpropane-2-sulfinamide as a pale white solid (0.68 mmol, yield: 76%).

^1H NMR (300 MHz, CDCl_3) δ 8.62-8.67 (m, 1H), 8.08-8.15 (m, 1H), 7.35-7.56 (m, 4H), 7.11-7.25 (m, 3H),
25 7.01-7.07 (m, 1H), 6.90-7.00 (m, 2H), 5.61-5.92 (m,


1H), 4.66-4.80 (m, 1H), 1.29 (s, 9H), 1.20 (d, J = 6.6Hz, 3H).

Step 6: Preparation of (S)-1-(6-fluoro-3-phenyl-4-(pyridine-2-yl)quinoline-2-yl)ethane-1-amine

224 mg (0.5 mmol) of (S)-N-((S)-1-(6-fluoro-3-phenyl-4-(pyridine-2-yl)quinoline-2-yl)ethyl)-2-methylpropane-2-sulfinamide prepared in step 5 was reacted by the same manner as described in step 8 of Preparative Example 21. The obtained compound was separated by column chromatography (SiO₂, eluent: MeOH/CH₂Cl₂, 1/20 → MeOH/CH₂Cl₂, 1/10) to give 163 mg of the target compound (S)-1-(6-fluoro-3-phenyl-4-(pyridine-2-yl)quinoline-2-yl)ethane-1-amine as a yellow oil (0.47 mmol, yield: 95%).

¹H NMR (300 MHz, CDCl₃) δ 8.62-8.67 (m, 1H), 8.10-8.17 (m, 1H), 7.41-7.55 (m, 2H), 6.91-7.32 (m, 8H), 4.40-4.50 (m, 1H), 3.50 (br s, 2H), 1.23-1.30 (m, 3H).

Preparative Example 24: Preparation of (S)-2-(1-aminoethyl)-6-fluoro-3-(pyridine-3-yl)quinazoline-4(3H)-one



Step 1: Preparation of tert-butyl (S)-(1-(6-fluoro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)carbamate

Tert-butyl (S)-(1-(6-fluoro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)carbamate was prepared by using 2-amino-5-fluorobenzoic acid and 3-aminopyridine according to the same manner as described in step 2 of Preparative Example 1.

¹H NMR (300 MHz, CDCl₃) δ 8.79 (t, J = 3.2 Hz, 2H), 7.92-7.72 (m, 2H), 7.60-7.48 (m, 2H), 5.46 (d, J = 8.3 Hz, 2H), 4.45-4.37 (m, 1H), 1.41 (m, 9H), 1.28 (t, J = 6.8 Hz, 3H).


Step 2: Preparation of (S)-2-(1-aminoethyl)-6-fluoro-3-(pyridine-3-yl)quinazoline-4(3H)-one

(S)-2-(1-aminoethyl)-6-fluoro-3-(pyridine-3-yl)quinazoline-4(3H)-one was prepared by using the compound prepared in step 1 according to the same manner as described in step 3 of Preparative Example 1.

¹H NMR (300 MHz, CDCl₃) δ 8.79 (d, J = 4.0 Hz, 1H), 7.90 (d, J = 8.1 Hz, 1H), 7.74 (d, J = 6.5 Hz, 1H), 7.53 (d, J = 7.7 Hz, 2H), 7.31-7.11 (m, 2H), 3.80-3.73 (m, 1H), 2.80 (s, 2H), 1.32 (dd, J = 21, 6.3 Hz, 3H).

25

Preparative Example 25: Preparation of (S)-2-(1-aminoethyl)-6-fluoro-3-phenylquinazoline-4(3H)-one



Step 1: Preparation of tert-butyl (S)-1-(6-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)carbamate

The target compound was prepared by using 2-amino-5-fluorobenzoic acid and aniline according to the same manner as described in step 2 of Preparative
 10 Example 1.

^1H NMR (300 MHz, $CDCl_3$) δ 7.88 (d, J = 8.0 Hz, 1H), 7.70-7.74 (m, 1H), 7.60-7.39 (m, 3H), 7.28 (d, J = 6.8 Hz, 1H), 5.66 (d, J = 6.7 Hz, 1H), 4.54 (t, J = 6.3 Hz, 1H), 1.41 (s, 9H), 1.26 (d, J = 6.3 Hz, 3H).


15

Step 2: Preparation of (S)-2-(1-aminoethyl)-6-fluoro-3-phenylquinazoline-4(3H)-one

The target compound was prepared by using the compound prepared in step 1 according to the same
 20 manner as described in step 3 of Preparative Example 1.

^1H NMR (300 MHz, CDCl_3) δ 7.87 (d, J = 7.6 Hz, 1H), 7.66 (dd, J = 8.5, 5.0 Hz, 1H), 7.57-7.29 (m, 2H), 4.03-4.09 (m, 1H), 1.36 (d, J = 6.4 Hz, 3H).

5 Preparative Example 26: Preparation of (S)-2-(1-aminoethyl)-6-fluoro-3-(3-fluorophenyl)quinazoline-4(3H)-one



10 Step 1: Preparation of tert-butyl (S)-(1-(6-fluoro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)carbamate

The target compound was prepared by using 2-amino-5-fluorobenzoic acid and 3-fluoroaniline according to the same manner as described in step 2 of Preparative Example 1.


^1H NMR (300 MHz, CDCl_3) δ 7.88 (d, J = 8.0 Hz, 1H), 7.73 (dd, J = 8.4, 4.6 Hz, 1H), 7.62-7.47 (m, 2H), 7.28-7.03 (m, 3H), 5.55-5.50 (m, 1H), 4.56-4.49 (m, 1H), 1.41 (s, 9H), 1.30 (t, J = 4.4 Hz, 3H).

20 Step 2: Preparation of (S)-2-(1-aminoethyl)-6-fluoro-3-(3-fluorophenyl)quinazoline-4(3H)-one

The target compound was prepared by using the compound prepared in step 1 according to the same manner as described in step 3 of Preparative Example 1.

¹H NMR (300 MHz, CDCl₃) δ 7.89(d, J = 8.3 Hz, 1H), 7.74(dd, J = 8.5, 4.4 Hz, 1H), 7.59-7.48(m, 2H), 7.29-7.24(m, 1H), 7.12-7.03(m, 2H), 3.75-3.67(m, 1H), 1.85(s, 2H), 1.31(d, J = 6.4 Hz, 3H).

Preparative Example 27: Preparation of (S)-2-(1-aminoethyl)-5-chloro-3-(2-chlorobenzyl)quinazoline-4(3H)-one



Step 1: Preparation of tert-butyl (S)-((1-(5-chloro-3-(2-chlorobenzyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)carbamate

The target compound was prepared by using 2-amino-6-chlorobenzoic acid and 2-chlorobenzylamine according to the same manner as described in step 2 of Preparative Example 1.

¹H NMR (500 MHz, CDCl₃) δ 7.60-7.67 (m, 2H), 7.48-7.54 (m, 1H), 7.42-7.46 (m, 1H), 7.14-7.27 (m, 2H),

6.82 (d, $J = 7.3\text{Hz}$, 1H), 5.63 (s, 2H), 5.39-5.44 (m, 1H), 4.82-4.85 (m, 1H), 1.49 (s, 9H), 1.37 (d, $J = 6.6\text{Hz}$, 3H).


5 Step 2: Preparation of (S)-2-(1-aminoethyl)-5-chloro-3-(2-chlorobenzyl)quinazoline-4(3H)-one

The target compound was prepared by using the compound prepared in step 1 according to the same manner as described in step 3 of Preparative Example 10 1.

^1H NMR (300 MHz, CDCl_3) δ 7.57-7.65 (m, 2H), 7.39-7.52 (m, 2H), 7.13-7.26 (m, 2H), 6.75-6.85 (m, 1H), 5.75 (d, $J = 17.1\text{ Hz}$, 1H), 5.27 (d, $J = 17.1\text{Hz}$, 1H), 3.83-3.91 (m, 1H), 1.40 (d, $J = 6.5\text{Hz}$, 3H).

15

Preparative Example 28: Preparation of (S)-2-(1-aminoethyl)-6-fluoro-3-(pyridine-2-ylmethyl)quinazoline-4(3H)-one



20 Step 1: Preparation of tert-butyl (S)-(1-(6-fluoro-4-oxo-3-(pyridine-2-ylmethyl)-3,4-dihydroquinazoline-2-yl)ethyl)carbamate

The target compound was prepared by using 2-amino-6-fluorobenzoic acid and pyridine-2-ylmethaneamine according to the same manner as described in step 2 of Preparative Example 1.

5 ^1H NMR (300 MHz, CDCl_3) δ 8.48(d, J = 4.0 Hz, 1H), 7.89(dd, J = 8.5, 3.0 Hz, 1H), 7.71-7.62(m, 2H), 7.46(td, J = 8.6, 2.9 Hz, 1H), 7.19-7.15(m, 2H), 5.61(s, 2H), 5.22-5.13(m, 1H), 1.44(s, 3H), 1.41(s, 9H).


10

Step 2: Preparation of (S)-2-(1-aminoethyl)-6-fluoro-3-(pyridine-2-ylmethyl)quinazoline-4(3H)-one

The target compound was prepared by using the compound prepared in step 1 according to the same
15 manner as described in step 3 of Preparative Example 1.

^1H NMR (300 MHz, CDCl_3) δ 8.48(d, J = 4.5 Hz, 1H), 7.83(dd, J = 8.4, 2.9 Hz, 1H), 7.71-7.63(m, 2H), 7.42-7.33(m, 2H), 7.19(dd, J = 7.1, 5.1 Hz, 1H), 8.48(d, J = 4.5 Hz, 1H), 5.73(d, J = 15.8 Hz, 1H), 5.21(d, J = 15.8 Hz, 1H), 4.72(q, J = 6.5 Hz, 1H), 4.45(s, 2H),
20 1.53(d, J = 6.6 Hz, 3H)

Preparative Example 29: Preparation of 1-(5-chloro-3-(pyridine-2-yl)quinoline-2-yl)ethane-1-amine
25



Step 1: Preparation of (2-amino-6-chlorophenyl)methanol

3.43 g (20.0 mmol) of 2-amino-6-chlorobenzoic acid was dissolved in anhydrous THF (30 mL), to which LiAlH₄ (1.5 equivalent) was slowly added at room temperature for 10 minutes, followed by stirring for 12 hours. Diethylether (40 mL) and water (5 mL) were added thereto. The reaction mixture was dried (MgSO₄) and concentrated under reduced pressure. The residue was separated by column chromatography (SiO₂, eluent: CH₂Cl₂/ethyl acetate, 5/1 → CH₂Cl₂/ethylacetate, 2/1) to give 2.36 g of the target compound (2-amino-6-chlorophenyl)methanol as a pale yellow solid (15.0 mmol, yield: 75%).

¹H NMR (300 MHz, CDCl₃) δ 7.01 (t, J = 8.0 Hz, 1H), 6.76 (d, J = 7.9 Hz, 1H), 6.58 (d, J = 8.0 Hz, 1H), 4.89 (s, 2H), 4.30 (br s, 2H), 1.66 (br s, 1H).

Step 2: Preparation of 2-amino-6-chlorobenzaldehyde

2.30 g (14.6 mmol) of (2-amino-6-chlorophenyl)methanol prepared in step 1, MnO₂ (10 equivalent), and diethylether (50 mL) were mixed together, which was stirred at room temperature for 6 hours. The reaction mixture was filtered with celite pad, and concentrated under reduced pressure. The residue was separated by column chromatography (SiO₂, eluent: hexane/CH₂Cl₂, 5/1 → CH₂Cl₂) to give 2.27 g of the target compound 2-amino-6-chlorobenzaldehyde as a yellow solid (14.6 mmol, yield: 10%).

¹H NMR (300 MHz, CDCl₃) δ 10.48 (s, 1H), 7.17 (t, J = 8.2 Hz, 1H), 6.67 (d, J = 7.7 Hz, 1H), 6.54 (d, J = 8.4 Hz, 1H), 6.48 (br s, 2H).

Step 3: Preparation of tert-butyl (1-(5-chloro-3-(pyridine-2-yl)quinoline-2-yl)ethylcarbamate

793 mg (3.0 mmol) of 2-amino-6-chlorobenzaldehyde (1.2 equivalent) prepared in step 2, tert-butyl (S)-(3-oxo-4-(pyridine-2-yl)butane-2-yl)carbamate, K₂CO₃ (3 equivalent), and ethanol (15 mL) were mixed together, which was refluxed for 6 hours. The reaction mixture was cooled down to room temperature. Water was added

thereto, followed by extraction with ethyl acetate. The extracted organic layer was washed with saturated brine, separated, dried (Na₂SO₄), and concentrated under reduced pressure. The obtained compound was
5 separated by column chromatography (SiO₂, eluent: hexane/ethyl acetate, 8/1 → hexane/ethylacetate, 3/1) to give 1.16 g of the target compound tert-butyl (1-(5-chloro-3-(pyridine-2-yl)quinoline-2-yl)ethylcarbamate as a white solid (3.0 mmol, yield:
10 100%).

¹H NMR (300 MHz, CDCl₃) δ 8.77 (br d, J = 4.1 Hz, 1H), 8.53 (s, 1H), 8.05 (d, J = 8.6 Hz, 1H), 7.83-7.89 (m, 1H), 7.59-7.68 (m, 3H), 7.34-7.39 (m, 1H), 6.33 (br d, J = 6.6 Hz, 1H), 5.43-5.52 (m, 1H), 1.45 (s,
15 9H), 1.34 (d, J = 6.5 Hz, 3H).


Step 4: Preparation of 1-(5-chloro-3-(pyridine-2-yl)quinoline-2-yl)ethane-1-amine

739 mg of 1-(5-chloro-3-(pyridine-2-yl)quinoline-2-yl)ethane-1-amine was prepared by using 1.0 g (2.60
20 mmol) of tert-butyl (1-(5-chloro-3-(pyridine-2-yl)quinoline-2-yl)ethylcarbamate prepared in step 3 according to the same manner as described in step 3 of Preparative Example 1 (2.60 mmol, yield: 100%).


¹H NMR (300 MHz, CDCl₃) δ 8.64 (d, J = 2.7 Hz, 1H),
8.60 (s, 1H), 8.05-7.99 (m, 1H), 8.05-7.99 (m, 1H),
7.88 (td, J = 7.7, 1.3 Hz, 1H), 7.70-7.63 (m, 3H), 7.36-
7.32 (m, 1H), 5.18-5.24 (m, 1H), 1.58 (d, J = 6.6 Hz,
5 3H).

The following examples 1 ~ 33 were performed by
the method represented by the reaction formula 1A.

[Reaction Formula 1A]



Example 1: Preparation of (S)-4-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one



Step 1: Preparation of 4,6-dichloropyrimidine-5-carbaldehyde

30 mL of phosphorylchloride (POCl_3) was cooled
 5 down to 0°C , to which 9.6 mL of anhydrous
 dimethylformamide (DMF) was slowly added. 1 hour
 later, 7.85 g (70.0 mmol) of 4,6-dihydroxypyrimidine
 was added thereto. The reaction mixture was heated at
 room temperature, followed by stirring at room
 10 temperature for 30 minutes. The reaction mixture was
 refluxed for 3 hours. The mixture was cooled down to
 room temperature. The reaction mixture was slowly
 added to ice water, followed by extraction with
 ethylacetate. The extracted organic layer was dried

over Na₂SO₄, filtered, and concentrated under reduced pressure. The obtained solid was washed with hexane/diethyl ether (5/1, v/v) to give 10.5 g of 4,6-dichloropyrimidine-5-carbaldehyde as a white solid
5 (5.95 mmol, yield: 85%).

¹H NMR (300 MHz, CDCl₃) δ 10.47 (s, 1H), 8.90 (s, 1H).

Step 2: preparation of 1-(4,6-dichloropyrimidine-5-yl)ethane-1-ol
10

1.2 g (6.8 mmol) of 4,6-dichloropyrimidine-5-carbaldehyde was dissolved in THF (25 mL), to which 8.14 mL (8.14 mmol, 1.2 equivalent) of methylmagnesium bromide (18% in THF) was slowly added at 0°C.
15 Saturated ammoniumchloride aqueous solution (10 mL) was slowly added thereto, followed by extraction with ethyl acetate. The extracted organic layer was dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The residue was separated by column
20 chromatography (SiO₂, eluent: hexane/ethylacetate, 4/1) to give 1.1 g of the target compound 1-(4,6-dichloropyrimidine-5-yl)ethane-1-ol as a white solid (5.7 mmol, yield: 80%).

¹H NMR(300 MHz, CDCl₃) δ 8.69 (s, 1H), 5.57-5.47 (m, 1H), 2.64 (d, J = 9.3 Hz, 1H), 1.68 (d, J = 6.8 Hz, 3H).

5 Step 3: Preparation of 1-(4,6-dichloropyrimidine-5-yl)ethane-1-one

980 mg (5.08 mmol) of 1-(4,6-dichloropyrimidine-5-yl)ethane-1-ol prepared in step 2 was dissolved in 30 mL of acetone, to which 1.0 g (10.2 mmol, 2.0 eq) of chromium trioxide was slowly added, followed by
10 stirring at room temperature for 2 hours. 2 mL of Isopropylalcohol was added thereto, followed by stirring for 10 minutes. 20 mL of saturated sodiumbicarbonate aqueous solution was added to the
15 reaction mixture, followed by extraction with ethylacetate. The extracted organic layer was dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The residue was separated by column chromatography (SiO₂, eluent: hexane/ethylacetate,
20 6/1) to give 823 mg of the target compound 1-(4,6-dichloropyrimidine-5-yl)ethane-1-one as a white solid (4.3 mmol, yield: 85%).

¹H NMR(300 MHz, CDCl₃) δ 8.84 (s, 1H), 2.63 (s, 3H).

Step 4: Preparation of 1-(4-chloro-6-((4-methoxybenzyl)amino)pyrimidine-5-yl)ethane-1-one

3.82 g (20.0 mmol) of 1-(4,6-dichloropyrimidine-5-yl)ethane-1-one was dissolved in 30 mL of
5 dichloromethane, which was cooled down to 0°C, to which 3.88 g (30.0 mmol) of diisopropylethylamine and 3.29 g (24.0 mmol) of p-methoxybenzylamine (PMBNH₂) were added stepwise. 1 hour later, the reaction mixture was heated to room temperature, followed by
10 stirring at room temperature for 6 hours. Water and ethylacetate were added to the reaction mixture, followed by extraction. The extracted organic layer was dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The residue was separated by
15 column chromatography (SiO₂, eluent: hexane/ethylacetate, 3/1) to give 5.54 g of the target compound 1-(4-chloro-6-((4-methoxybenzyl)amino)pyrimidine-5-yl)ethane-1-one as a white solid (19.0 mmol, yield: 95%).

20 ¹H NMR (300 MHz, CDCl₃) δ 9.07 (br s, 1H, NH), 8.38 (s, 1H), 7.25 (d, J = 8.1 Hz, 2H), 6.88 (d, J = 8.1 Hz, 2H), 4.67 (d, J = 4.8 Hz, 2H), 3.81 (s, 3H), 2.74 (s, 3H).

Steps 5 and 6: Preparation of 4-hydroxy-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one

5.83 g (20.0 mmol) of 1-(4-chloro-6-((4-methoxybenzyl)amino)pyrimidine-5-yl)ethane-1-one

5 prepared in step 4 and 3.57 g (30.0 mmol) of N,N-dimethylformamide dimethyl acetal (DMF-DMA) were dissolved in 30 mL of anhydrous toluene, which was heated at 100°C for 3 hours. The reaction mixture was cooled down to room temperature. The solvent and DMF-DMA were eliminated under reduced pressure. The obtained intermediate compound was added with 100 mL of acetic acid and 20 mL of water, followed by reflux for 4 days. The reaction mixture was cooled down to room temperature. The solvent was eliminated under reduced pressure. The obtained yellow product was washed with water/isopropanol (IPA) (1/1) to give 4.53 g of the target compound 4-hydroxy-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one as a white solid (16.0 mmol, yield: 80%).

¹H NMR (300 MHz, DMSO-d₆) δ 8.78 (br s, 1H, NH), 7.76 (d, J = 4.7 Hz, 1H), 7.28 (d, J = 5.2 Hz, 2H), 6.93 (d, J = 5.2 Hz, 2H), 6.47 (d, J = 4.7 Hz, 1H), 5.48 (s, 3H), 3.83 (s, 3H).

Step 7: Preparation of (S)-4-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one
 50 mg (0.176 mmol) of 4-hydroxy-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one prepared in step 5 and step 6 was dissolved in 2 mL of anhydrous acetonitrile, to which 101 mg (0.229 mmol) of (benzotriazole-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate (BOP) and 39 μ L (0.264 mmol) of 1,8-diazabicyclo[5.4.0]-7-undecene (DBU) were added, followed by stirring for 30 minutes. 58 mg (0.194 mmol) of (S)-2-(1-aminoethyl)-5-chloro-3-phenylquinazoline-4(3H)-one was added thereto, followed by stirring at 60°C for 12 hours. The reaction mixture was filtered under reduced pressure. Ethyl acetate and water were added thereto, followed by extraction. The extracted organic layer was dried (Na_2SO_4), filtered, and concentrated under reduced pressure. The residue was separated by column chromatography (SiO_2 , eluent: hexane/ethylacetate, 6/1 $\rightarrow$ hexane/ethylacetate, 1/1) to give 72 mg of the target compound (S)-4-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)-8-(4-

methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one as a yellow solid (0.127 mmol, yield: 50%).

¹H NMR (300 MHz, CDCl₃) δ 11.02 (d, J = 7.7 Hz, 1H), 8.28 (s, 1H), 7.7-7.71 (d, J = 8.1 Hz, 1H), 7.54-7.62 (m, 2H), 7.40-7.53 (m, 5H), 7.33 (d, J = 7.9 Hz, 1H), 7.20 (d, J = 8.6 Hz, 2H), 6.85 (d, J = 8.6 Hz, 2H), 6.30 (d, J = 7.9 Hz, 1H), 5.33 (s, 2H), 5.01 (q, J = 6.9 Hz, 6.9 Hz, 1H), 3.78 (s, 3H), 1.49 (d, J = 6.7 Hz, 3H).

10


Step 8: Preparation of (S)-4-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one

72 mg (0.127 mmol) of (S)-4-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one prepared in step 7 was dissolved in 1 mL of dichloromethane, to which 1 mL of trifluoroacetic acid (TFA) and 0.5 mL of methanesulfonic acid were added, followed by stirring at 70°C for 10 hours. Saturated sodiumbicarbonate aqueous solution was added thereto, followed by neutralization. Dichloromethane and water were added thereto, followed by extraction. The extracted organic layer was dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue

was separated by column chromatography (SiO₂, eluent: dichloromethane/methanol, 30/1) to give 51 mg of the target compound (S)-4-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one as a pale yellow solid (0.115 mmol, yield: 90%).

¹H NMR (500 MHz, CDCl₃) δ 11.06 (d, J = 7.0 Hz, 1H), 8.24 (s, 1H), 7.73 (d, J = 8.2 Hz, 1H), 7.58-7.64 (m, 2H), 7.51-7.57 (m, 3H), 7.46-7.51 (m, 2H), 7.37 (d, J = 7.6 Hz, 1H), 6.38 (d, J = 7.6 Hz, 1H), 5.12 (q, J = 6.8 Hz, 6.8 Hz, 1H), 1.53 (d, J = 6.4 Hz, 3H).

Example 2: Preparation of (S)-4-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one



Step 1: Preparation of (S)-4-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one

30 mg of (S)-4-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a yellow solid by using 58 mg (0.194 mmol) of (S)-2-(1-aminoethyl)-5-chloro-3-(pyridine-3-yl)quinazoline-4(3H)-one according to the same manner as described in step 7 of Example 1 (0.053 mmol, yield: 30%).

MS[m/z; (M + 1)⁺]: 567.

10


Step 2: Preparation of (S)-4-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one

14 mg of (S)-4-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a pale yellow solid by using 30 mg (0.053 mmol) of (S)-4-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one prepared in step 1 according to the same manner as described in step 8 of Example 1 (0.031 mmol, yield: 59%).

¹H NMR (300 MHz, CDCl₃) δ 10.78 (t, J = 6.5 Hz, 1H), 8.76 (s, 1H), 8.20 (d, J = 4.6 Hz, 1H), 7.44-7.77

(m, 6H), 6.33 (d, J = 7.4 Hz, 1H), 4.93-4.50 (m, 1H), 1.49-1.60 (m, 3H).

Example 3: Preparation of (S)-4-((1-(5-chloro-4-oxo-3-(pyridine-2-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one



Step 1: Preparation of (S)-4-((1-(5-chloro-4-oxo-3-(pyridine-2-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one

50 mg of (S)-4-((1-(5-chloro-4-oxo-3-(pyridine-2-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a pale yellow solid by using 58 mg (0.194 mmol) of (S)-2-(1-aminoethyl)-5-chloro-3-(pyridine-2-yl)quinazoline-4(3H)-one according to the same manner as described in step 7 of Example 1 (0.112 mmol, yield: 64%).

¹H NMR (300 MHz, CDCl₃) δ 10.97 (d, J = 4.6 Hz, 1H), 8.68 (d, J = 4.6 Hz, 1H), 8.23 (s, 1H), 7.85 (t, J = 7.9 Hz, 1H), 7.72 (d, J = 7.9 Hz, 1H), 7.59 (t, J = 7.9 Hz, 1H), 7.38-7.53 (m, 4H), 7.19 (d, J = 8.5 Hz, 2H), 6.85 (d, J = 8.6 Hz, 2H), 6.27 (d, J = 7.9 Hz, 1H), 5.32 (s, 2H), 4.20-5.03 (m, 1H), 3.78 (s, 3H), 1.60 (d, J = 6.6 Hz, 3H).


Step 2: Preparation of (S)-4-((1-(5-chloro-4-oxo-3-(pyridine-2-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one

14 mg of (S)-4-((1-(5-chloro-4-oxo-3-(pyridine-2-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a pale yellow solid by using 50 mg (0.112 mmol) of (S)-4-((1-(5-chloro-4-oxo-3-(pyridine-2-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one prepared in step 1 according to the same manner as described in step 8 of Example 1 (0.031 mmol, yield: 28%).

¹H NMR (300 MHz, CDCl₃) δ 11.20 (brs, 1H), 10.92 (d, J = 6.1 Hz, 1H), 8.70 (d, J = 4.7 Hz, 1H), 8.16 (s, 1H), 7.90 (t, J = 7.7 Hz, 1H), 7.72 (d, J = 7.7 Hz, 1H), 7.61 (t, J = 8.3 Hz, 1H), 7.41-7.55 (m, 4H),

6.33 (d, J = 7.7 Hz, 1H), 4.92-5.03 (m, 1H), 1.60 (d, J = 7.0 Hz, 3H).

Example 4: Preparation of (S)-4-((1-(5-chloro-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one



Step 1: Preparation of (S)-4-((1-(5-chloro-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one

50 mg of (S)-4-((1-(5-chloro-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a white solid by using 65 mg (0.194 mmol) of (S)-2-(1-aminoethyl)-5-chloro-3-(3,5-difluorophenyl)quinazoline-4(3H)-one according to the same manner as described in step 7 of Example 1 (0.083 mmol, yield: 47%).

¹H NMR (500 MHz, CDCl₃) δ 10.94 (d, J = 7.2 Hz, 1H), 8.33 (s, 1H), 7.70 (d, J = 8.1 Hz, 1H), 7.63 (t, J = 8.1 Hz, 1H), 7.51 (t, J = 8.0 Hz, 2H), 7.23 (d, J = 8.5 Hz, 2H), 7.11 (d, J = 7.1 Hz, 1H), 6.02-7.00 (m, 2H), 6.89 (d, J = 8.5 Hz, 2H), 6.33 (d, J = 8.0 Hz, 1H), 5.37 (d, J = 3.1 Hz, 2H), 5.09-5.15 (m, 1H), 3.82 (s, 3H), 1.58 (d, J = 7.2 Hz, 3H).


Step 2: Preparation of (S)-4-((1-(5-chloro-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one

35 mg of (S)-4-((1-(5-chloro-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a pale yellow solid by using 50 mg (0.083 mmol) of (S)-4-((1-(5-chloro-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one prepared in step 1 according to the same manner as described in step 8 of Example 1 (0.073 mmol, yield: 87%).

¹H NMR (300 MHz, CDCl₃) δ 10.84 (br s, 1H), 10.81 (d, J = 6.3 Hz, 1H), 8.24 (s, 1H), 7.58-7.71 (m, 2H), 7.46-7.56 (m, 2H), 7.07-7.13 (m, 1H), 6.89-7.03 (m,

2H), 6.36 (d, $J = 8.0$ Hz, 1H), 5.09 (q, $J = 5.5$ Hz, 6.8 Hz, 1H), 1.56 (t, $J = 6.6$ Hz, 3H).

Example 5: Preparation of (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one



Step 1: Preparation of (S)-3-(1-((5-acetyl-6-((4-methoxybenzyl)amino)pyrimidine-4-yl)amino)ethyl)-8-chloro-2-phenylisoquinoline-1(2H)-one

292 mg (1.0 mmol, 1 equivalent) of 1-(4-chloro-6-((4-methoxybenzyl)amino)pyrimidine-5-yl)ethane-1-one prepared in step 4 of Example 1 and (S)-3-(1-aminoethyl)-8-chloro-2-phenylisoquinoline-1(2H)-one (1.1 equivalent) were dissolved in 10 mL of anhydrous dimethylsulfoxide (DMSO), to which diisopropylethylamine (DIPEA) (3 equivalent) was added, followed by stirring at 80°C for 10 hours. Ethyl acetate and water was added to the reaction mixture, followed by extraction. The extracted organic layer was dried (Na_2SO_4), filtered, and

concentrated under reduced pressure. The residue was separated by column chromatography (SiO₂, eluent: hexane/ethyl acetate, 3/1 → hexane/ethyl acetate, 1/1) to give 482 mg of the target compound (S)-3-(1-
5 ((5-acetyl-6-((4-methoxybenzyl)amino)pyrimidine-4-yl)amino)ethyl)-8-chloro-2-phenylisoquinoline-1(2H)-one as a white solid (0.87 mmol, yield: 87%).

¹H NMR (300 MHz, CDCl₃) δ 8.56 (br d, J = 6.3 Hz, 1H), 8.02 (s, 1H), 7.20-7.55 (m, 10H), 6.87 (d, J =
10 8.7 Hz, 2H), 6.48 (s, 1H), 6.44 (br t, 1H), 4.84-4.95 (m, 1H), 4.66 (d, J = 4.8 Hz, 2H), 3.79 (s, 3H), 2.52 (s, 3H), 1.38 (d, J = 6.6 Hz, 3H).

Step 2: Preparation of (S)-4-((1-(8-chloro-1-oxo-
15 2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one

200 mg (0.361 mmol) of (S)-3-(1-((5-acetyl-6-((4-methoxybenzyl)amino)pyrimidine-4-yl)amino)ethyl)-8-chloro-2-phenylisoquinoline-1(2H)-one prepared in step
20 1 was dissolved in 3 mL of anhydrous dimethylformamide (DMF), to which 0.24 mL (1.805 mmol) of N,N-dimethylformamide dimethyl acetal was added, followed by stirring at 130°C for 15 hours. Ethyl acetate and water was added to the reaction mixture, followed by
25 extraction. The extracted organic layer was dried

(Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was separated by column chromatography (SiO₂, eluent: dichloromethane/ethyl acetate, 10/1 → dichloromethane/ethyl acetate, 1/1) to give 90 mg of the target compound (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one as a white solid (0.160 mmol, yield: 44%).

¹H NMR (300 MHz, CDCl₃) δ 10.71 (d, J = 6.9 Hz, 1H), 8.30 (s, 1H), 7.47-7.57 (m, 3H), 7.29-7.47 (m, 6H), 7.23 (d, J = 8.3 Hz, 2H), 6.87 (d, J = 8.3 Hz, 2H), 6.59 (s, 1H), 6.30 (d, J = 8.3 Hz, 1H), 5.35 (s, 2H), 4.93 (t, J = 6.9 Hz, 1H), 3.78 (s, 3H), 1.45 (d, J = 6.9 Hz, 3H).


Step 3: Preparation of (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one

67 mg of (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a white solid by using 90 mg (0.160 mmol) of (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-

d]pyrimidine-5(8H)-one prepared in step 2 according to the same manner as described in step 8 of Example 1 (0.151 mmol, yield: 95%).

¹H NMR (300 MHz, DMSO-d₆) δ 12.14 (s, 1H), 10.48 (d, J = 7.2 Hz, 1H), 8.19 (s, 1H), 7.75-7.82 (m, 1H), 7.58-7.69 (m, 2H), 7.46-7.54 (m, 2H), 7.28-7.45 (m, 4H), 6.77 (s, 1H), 6.15 (d, J = 8.6 Hz, 1H), 4.71 (t, J = 7.1 Hz, 1H), 1.40 (d, J = 8.6 Hz, 3H).

10 Example 6: Preparation of (S)-4-((1-(2-phenylquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one



15 Step 1: Preparation of (S)-8-(4-methoxybenzyl)-4-((1-(2-phenylquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one

30 mg of (S)-8-(4-methoxybenzyl)-4-((1-(2-phenylquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a pale yellow
20 solid by using 29 mg (0.117 mmol) of (S)-1-(2-

(
phenylquinoline-3-yl)ethane-1-amine according to the same manner as described in step 7 of Example 1 (0.058 mmol, yield: 55%).

¹H NMR (300 MHz, CDCl₃) δ 10.94 (d, J = 6.5 Hz, 1H), 8.42-8.56 (m, 1H), 8.28 (d, J = 12.7 Hz, 2H), 7.38-8.17 (m, 9H), 7.21 (d, J = 8.6 Hz, 2H), 6.85 (d, J = 8.6 Hz, 2H), 6.30 (d, J = 7.6 Hz, 1H), 5.64-5.75 (m, 1H), 5.34 (d, J = 7.6 Hz, 2H), 3.78 (s, 3H), 1.49 (d, J = 7.1 Hz, 3H).

10


Step 2: Preparation of (S)-4-((1-(2-phenylquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one

20 mg of (S)-4-((1-(2-phenylquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a pale yellow solid by using 30 mg (0.058 mmol) of (S)-8-(4-methoxybenzyl)-4-((1-(2-phenylquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one prepared in step 1 according to the same manner as described in step 8 of Example 1 (0.051 mmol, yield: 87%).

¹H NMR (300 MHz, CDCl₃) δ 10.78 (d, J = 8.0 Hz, 1H), 10.55 (brs, 1H), 8.24 (d, J = 6.7 Hz, 2H), 8.14 (d, J = 6.7 Hz, 1H), 7.64-7.84 (m, 4H), 7.41-7.55 (m,

5H), 6.34 (d, J = 7.4 Hz, 1H), 5.71 (q, J = 5.3 Hz, 6.6 Hz, 1H), 1.50 (d, J = 7.4 Hz, 3H).

Example 7: Preparation of 4-((1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one



Step 1: Preparation of 4-((1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one

30 mg of 4-((1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a pale yellow solid by using 52 mg (0.194 mmol) of 1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethane-1-amine according to the same manner as described in step 7 of Example 1 (0.031 mmol, yield: 59%).

¹H NMR (300 MHz, CDCl₃) δ 11.61 (d, J = 6.7 Hz, 1H), 8.79 (d, J = 3.6 Hz, 1H), 8.27-8.36 (m, 2H), 8.10 (s, 1H), 7.82 (t, J = 6.5 Hz, 1H), 7.40-7.62 (m, 4H),

7.32-7.39 (m, 1H), 7.19 (d, J = 7.9 Hz, 2H), 6.85 (d, J = 8.6 Hz, 2H), 6.29 (d, J = 7.8 Hz, 1H), 6.03-6.14 (m, 1H), 5.33 (s, 2H), 3.78 (s, 3H), 1.57 (d, J = 6.5 Hz, 3H).

5


Step 2: 4-((1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one

22 mg of 4-((1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a pale yellow solid by using 30 mg (0.056 mmol) of 4-((1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one prepared in step 1 according to the same manner as described in step 8 of Example 1 (0.053 mmol, yield: 95%).

¹H NMR (500 MHz, CDCl₃) δ 11.54 (d, J = 7.1 Hz, 1H), 8.79 (d, J = 4.7 Hz, 1H), 8.30-8.35 (m, 1H), 8.21 (s, 1H), 8.12 (s, 1H), 7.86 (t, J = 8.7 Hz, 1H), 7.59 (d, J = 8.1 Hz, 1H), 7.53 (t, J = 8.1 Hz, 1H), 7.46 (d, J = 7.6 Hz, 2H), 7.39 (t, J = 6.4 Hz, 1H), 6.32 (d, J = 7.6 Hz, 1H), 6.02-6.08 (m, 1H), 1.54 (d, J = 6.9 Hz, 3H).

25


Example 7-1: Preparation of (S)-4-((1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one



5 The target compound (S)-4-((1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one was prepared by using tert-butyl (S)-1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethane-1-amine prepared in Preparative Example 21
 10 according to the same manner as described in step 7 and step 8 of Example 1.

^1H NMR (500 MHz, CDCl_3) δ 11.54 (d, $J = 7.1$ Hz, 1H), 8.79 (d, $J = 4.7$ Hz, 1H), 8.30-8.35 (m, 1H), 8.21 (s, 1H), 8.12 (s, 1H), 7.86 (t, $J = 8.7$ Hz, 1H), 7.59 (d, $J = 8.1$ Hz, 1H), 7.53 (t, $J = 8.1$ Hz, 1H), 7.46 (d, $J = 7.6$ Hz, 2H), 7.39 (t, $J = 6.4$ Hz, 1H), 6.32 (d, $J = 7.6$ Hz, 1H), 6.02-6.08 (m, 1H), 1.54 (d, $J = 6.9$ Hz, 3H).

Example 8: Preparation of (S)-4-((1-(7-fluoro-2-(3-fluorophenyl)quinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one



Step 1: Preparation of (S)-4-((1-(7-fluoro-2-(3-fluorophenyl)quinoline-3-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one

40 mg of (S)-4-((1-(7-fluoro-2-(3-fluorophenyl)quinoline-3-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a yellow solid by using 55 mg (0.194 mmol) of (S)-1-(7-fluoro-2-(3-fluorophenyl)quinoline-3-yl)ethane-1-amine according to the same manner as described in step 7 of Example 1 (0.073 mmol, yield: 40%).

^1H NMR (300 MHz, CDCl_3) δ 10.94 (d, J = 5.3 Hz, 1H), 8.32 (s, 1H), 8.27 (s, 1H), 7.71-7.84 (m, 2H), 7.40-7.55 (m, 4H), 7.27-7.36 (m, 1H), 7.21 (d, J = 8.7 Hz, 2H), 7.08-7.17 (m, 1H), 6.85 (d, J = 8.5 Hz, 2H), 6.31 (d, J = 7.5 Hz, 1H), 5.61-5.73 (m, 1H), 5.34 (d,


J = 6.8 Hz, 2H), 3.78 (s, 3H), 1.49 (d, J = 6.9 Hz, 3H).

5 Step 2: Preparation of (S)-4-((1-(7-fluoro-2-(3-fluorophenyl)quinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one

25 mg of (S)-4-((1-(7-fluoro-2-(3-fluorophenyl)quinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a white solid
10 by using 40 mg (0.073 mmol) of (S)-4-((1-(7-fluoro-2-(3-fluorophenyl)quinoline-3-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one prepared in step 1 according to the same manner as described in step 8 of Example 1 (0.058 mmol, yield:
15 80%).

¹H NMR (300 MHz, CDCl₃) δ 11.25(brs, 1H), 10.84 (d, J = 7.2 Hz, 1H), 8.25 (d, J = 3.3 Hz, 2H), 7.71-7.84 (m, 2H), 7.42-7.56 (m, 4H), 7.28-7.36 (m, 1H), 7.11-7.19 (m, 1H), 6.35 (d, J = 7.2 Hz, 1H), 5.61-5.72
20 (m, 1H), 1.51 (d, J = 6.9 Hz, 3H).

Example 9: Preparation of (S)-4-(1-(7-fluoro-2-(pyridine-2-yl)quinoline-3-yl)ethylamino)pyrido[2,3-d]pyrimidine-5(8H)-one



Step 1: Preparation of (S)-4-(1-(7-fluoro-2-(pyridine-2-yl)quinoline-3-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one

5 245 mg of (S)-4-(1-(7-fluoro-2-(pyridine-2-yl)quinoline-3-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a pale yellow solid by using 212 mg (0.793 mmol) of (S)-1-(7-fluoro-2-(pyridine-2-yl)quinoline-3-yl)ethane-1-amine according to the same manner as described in step 7 of Example 1 (0.460 mmol, yield: 58%).

¹H NMR(300 MHz, CDCl₃) δ 10.99(d, J = 7.3 Hz, 1H), 8.75(d, J = 4.7 Hz, 1H), 8.33(s, 1H), 8.22(s, 1H), 7.99(d, J = 7.6 Hz, 1H), 7.72~7.88(m, 3H), 7.46(d, J = 7.6 Hz, 1H), 7.27~7.37(m, 2H), 7.18(d, J = 8.5 Hz, 2H), 6.83(d, J = 8.8 Hz, 2H), 6.27(d, J = 8.0 Hz, 1H), 6.07(q, J = 7.2 Hz, 6.4 Hz, 1H), 5.31(q, J = 14.8 Hz, 5.9 Hz, 2H), 3.76(s, 3H), 1.66(d, J = 6.7 Hz, 3H).


20

Step 2: Preparation of (S)-4-(1-(7-fluoro-2-(pyridine-2-yl)quinoline-3-yl)ethylamino)pyrido[2,3-d]pyrimidine-5(8H)-one

187 mg of (S)-4-(1-(7-fluoro-2-(pyridine-2-yl)quinoline-3-yl)ethylamino)pyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a pale yellow solid by using 245 mg (0.460 mmol) of (S)-4-(1-(7-fluoro-2-(pyridine-2-yl)quinoline-3-yl)ethylamino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one according to the same manner as described in step 8 of Example 1 (0.453 mmol, yield: 99%).

¹H NMR (500 MHz, CDCl₃) δ 11.26 (br s, 1H), 10.93 (d, J = 7.0 Hz, 1H), 8.82 (d, J = 4.6 Hz, 1H), 8.35 (s, 1H), 8.17 (s, 1H), 8.03 (d, J = 7.7 Hz, 1H), 7.86~7.93 (m, 1H), 7.83~7.87 (m, 1H), 7.77~7.81 (m, 1H), 7.44 (d, J = 7.7 Hz, 1H), 7.33~7.42 (m, 2H), 6.31 (d, J = 7.0 Hz, 1H), 6.12 (q, J = 7.0 Hz, 7.0 Hz, 1H), 1.68 (d, J = 6.8 Hz, 3H).

20 Example 10: Preparation of (S)-4-((1-(4,8-dichloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one




50 mg (0.113 mmol) of (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one prepared in Example 5 was dissolved in 2 mL of acetic acid, to which 17 mg (0.124 mmol) of N-chlorosuccinimide (NCS) was added, followed by stirring at 50°C for 15 hours. The reaction mixture was filtered under reduced pressure. Saturated sodiumbicarbonate aqueous solution was added thereto, followed by neutralization. Dichloromethane and water were added thereto, followed by extraction. The extracted organic layer was dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was separated by column chromatography (SiO₂, eluent: dichloromethane/methanol, 30/1 -> dichloromethane/methanol, 10/1) to give 25 mg of the target compound (S)-4-((1-(4,8-dichloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-

yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one as a pale yellow solid (0.052 mmol, yield: 46%).

¹H NMR (300 MHz, CDCl₃) δ 10.99 (d, J = 4.8 Hz, 1H), 8.25 (s, 1H), 7.95 (dd, J = 1.9 Hz, J = 7.5 Hz, 1H), 7.75 (d, J = 7.8 Hz, 1H), 7.46-7.62 (m, 6H), 7.20 (d, J = 6.7 Hz, 1H), 6.3 (d, J = 7.5 Hz, 1H), 5.04 (t, J = 67.2 Hz, 1H), 1.67 (d, J = 7.2 Hz, 3H).

Example 11: Preparation of (S)-4-((1-(8-chloro-4-fluoro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one



Step 1: Preparation of (S)-4-((1-(8-chloro-4-fluoro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one

5 mg of (S)-4-((1-(8-chloro-4-fluoro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a white solid by using 25 mg (0.079 mmol) of (S)-3-(1-aminoethyl)-8-chloro-4-fluoro-2-

phenylisoquinoline-1(2H)-one according to the same manner as described in step 7 of Example 1 (0.009 mmol, yield: 11%).

¹H NMR (300 MHz, CDCl₃) δ 10.96 (d, J = 7.8 Hz, 1H), 8.32 (s, 1H), 7.7.68 (t, J = 6.5 Hz, 2H), 7.43-7.62 (m, 7H), 7.19 (d, J = 8.8 Hz, 2H), 6.84 (d, J = 9.0 Hz, 2H), 6.27 (d, J = 7.8 Hz, 1H), 5.33 (q, J = 12.6 Hz, J = 9.1 Hz, 2H), 4.95 (q, J = 5.2 Hz, 6.5 Hz, 1H), 3.77 (s, 3H), 1.60 (d, J = 6.5 Hz, 3H).

10

Step 2: Preparation of (S)-4-((1-(8-chloro-4-fluoro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one

2 mg of (S)-4-((1-(8-chloro-4-fluoro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-

yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a white solid by using 5 mg (0.009 mmol) of (S)-4-((1-(8-chloro-4-fluoro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-8-(4-


20 methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one

prepared in step 1 according to the same manner as described in step 8 of Example 1 (0.004 mmol, yield: 50%).

¹H NMR (500 MHz, CDCl₃) δ 10.82 (d, J = 6.2 Hz, 1H), 8.22 (s, 1H), 7.71 (d, J = 6.2 Hz, 1H), 7.46-7.66

(m, 7H), 7.21-7.24 (m, 1H), 6.31 (d, J = 7.3 Hz, 1H), 4.96 (q, J = 4.9 Hz, 6.2 Hz, 1H), 1.61 (d, J = 7.3 Hz, 3H).

5 Example 12: Preparation of (S)-4-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one



10 Step 1: Preparation of (S)-4-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one

60 mg of (S)-4-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a white solid by using 58 mg (0.194 mmol) of (S)-2-(1-aminopropyl)-5-fluoro-3-phenylquinazoline-4(3H)-one according to the same manner as described in step 7 of Example 1 (0.107 mmol, yield: 60%).

¹H NMR (300 MHz, CDCl₃) δ 10.96 (d, J = 5.9 Hz, 1H), 8.28 (s, 1H), 7.43-7.73 (m, 8H), 7.32 (d, J = 6.5

Hz, 1H), 7.21 (d, J = 8.3 Hz, 2H), 7.15 (d, J = 9.7 Hz, 1H), 6.85 (d, J = 7.6 Hz, 2H), 6.23-6.37 (m, 1H), 5.34 (s, 2H), 4.96-5.07 (m, 1H), 3.77 (s, 3H), 1.75-1.99 (m, 2H), 0.88 (t, J = 7.0 Hz, 3H).


5

Step 2: Preparation of (S)-4-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one

33 mg of (S)-4-((1-(5-fluoro-4-oxo-3-phenyl-3,4-
10 dihydroquinazoline-2-yl)propyl)amino)pyrido[2,3-
d]pyrimidine-5(8H)-one was prepared as a pale brown solid by using 60 mg (0.107 mmol) of (S)-4-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-
15 d]pyrimidine-5(8H)-one prepared in step 1 according to the same manner as described in step 8 of Example 1 (0.075 mmol, yield: 88%).

¹H NMR (300 MHz, CDCl₃) δ 11.88(brs, 1H), 10.90 (d, J = 7.3 Hz, 1H), 8.20 (s, 1H), 7.41-7.70 (m, 7H),
20 7.33 (d, J = 7.0 Hz, 1H), 7.09 (t, J = 8.2 Hz, 1H), 6.34 (d, J = 7.6 Hz, 1H), 4.99-5.09 (m, 1H), 1.78-2.00 (m, 2H), 0.89 (t, J = 7.6 Hz, 3H).

Example 13: Preparation of (S)-4-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrido[2,3-d]pyrimidine-5(8H)-one



5 Step 1: Preparation of (S)-4-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one

60 mg of (S)-4-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a pale yellow solid by using 63 mg (0.194 mmol) of (S)-5-chloro-3-phenyl-2-(pyrrolidine-2-yl)quinazoline-4(3H)-one according to the same manner as described in step 7 of Example 1 (0.102 mmol, yield: 58%).

¹H NMR (300 MHz, CDCl₃) δ 8.21 (s, 1H), 7.71-7.78 (m, 1H), 7.45-7.65 (m, 8H), 7.37-7.44 (m, 3H), 7.19 (d, J = 6.8 Hz, 2H), 6.83 (d, J = 8.3 Hz, 2H), 6.23 (d, J = 7.5 Hz, 1H), 5.39 (d, J = 15.0 Hz, 1H), 5.23


(d, $J = 14.3$ Hz, 1H), 4.74-4.83 (m, 1H), 3.82-3.95 (m, 1H), 3.63-3.74 (m, 1H), 2.23-2.36 (m, 1H), 2.06-2.16 (m, 2H), 1.71-1.86 (m, 1H).

5 Step 2: Preparation of (S)-4-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrido[2,3-d]pyrimidine-5(8H)-one

36 mg of (S)-4-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a pale yellow solid by using 60 mg (0.102 mmol) of (S)-4-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one prepared in step 1 according to
10
15 the same manner as described in step 8 of Example 1 (0.076 mmol, yield: 75%).

^1H NMR (500 MHz, CDCl_3) δ 10.28 (br s, 1H), 8.15 (s, 1H), 7.75 (d, $J = 8.1$ Hz, 1H), 7.49-7.65 (m, 5H), 7.41-7.47 (m, 3H), 6.28 (d, $J = 7.8$ Hz, 1H), 4.87 (t, $J = 6.5$ Hz, 1H), 3.90-3.97 (m, 1H), 3.81-3.87 (m, 1H),
20 2.32-2.41 (m, 1H), 2.09-2.16 (m, 2H), 1.86-1.92 (m, 1H).

Example 14: Preparation of (S)-4-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)pyrido[2,3-d]pyrimidine-5(8H)-one



5 Step 1: Preparation of (S)-3-(1-(5-acetyl-6-((4-methoxybenzyl)amino)pyrimidine-4-yl)pyrrolidine-2-yl)-8-chloro-2-phenylisoquinoline-1(2H)-one

180 mg of (S)-3-(1-(5-acetyl-6-((4-methoxybenzyl)amino)pyrimidine-4-yl)pyrrolidine-2-yl)-
 10 8-chloro-2-phenylisoquinoline-1(2H)-one was prepared as a pale yellow solid by using 100 mg (0.343 mmol) of 1-(4-chloro-6-((4-methoxybenzyl)amino)pyrimidine-5-yl)ethane-1-one prepared in step 4 of Example 1 and
 122 mg (0.377 mmol) of (S)-8-chloro-2-phenyl-3-(pyrrolidine-2-yl)isoquinoline-1(2H)-one according to
 15 the same manner as described in step 1 of Example 5 (0.310 mmol, yield: 91%).

¹H NMR (300 MHz, CDCl₃) δ 8.61 (t, J = 3.4 Hz, 1H), 8.10 (s, 1H), 7.72 (d, J = 5.2 Hz, 1H), 7.38-7.63
 20 (m, 5H), 7.27-7.33 (m, 2H), 7.23 (d, J = 8.3 Hz, 2H), 6.84 (d, J = 8.3 Hz, 2H), 6.49 (s, 1H), 4.82-4.90 (m,

1H), 4.63-4.72 (m, 1H), 4.46-4.55 (m, 1H), 3.78 (s, 3H), 3.70-3.76 (m, 1H), 3.25 (t, J = 9.0 Hz, 1H), 2.53 (s, 3H), 2.03-2.14 (m, 1H), 1.87-2.01 (m, 1H), 1.75-1.87 (m, 1H), 1.57 (m, 1H).

5

Step 2: Preparation of (S)-4-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one

10 70 mg of (S)-4-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a pale yellow solid by using 180 mg (0.310 mmol) of (S)-3-(1-(5-acetyl-6-(4-methoxybenzyl)amino)pyrimidine-4-yl)pyrrolidine-2-yl)-8-chloro-2-phenylisoquinoline-1(2H)-one prepared in step 1 according to the same manner as described in step 2 of Example 5 (0.119 mmol, yield: 38%).

¹H NMR (300 MHz, CDCl₃) δ 8.36 (s, 1H), 7.71 (t, J = 6.6 Hz, 1H), 7.43-7.65 (m, 6H), 7.30-7.42 (m, 4H), 7.23 (d, J = 7.3 Hz, 2H), 6.71 (brs, 1H), 6.25 (d, J = 7.3 Hz, 1H), 5.26-5.42 (m, 2H), 4.97 (t, J = 7.3 Hz, 1H), 4.30-4.43 (m, 1H), 2.94-3.06 (m, 1H), 1.82-2.12 (m, 4H).


25

Step 3: Preparation of (S)-4-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)pyrido[2,3-d]pyrimidine-5(8H)-one

48 mg of (S)-4-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)pyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a pale yellow solid by using 70 mg (0.119 mmol) of (S)-4-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one prepared in step 2 according to the same manner as described in step 8 of Example 1 (0.102 mmol, yield: 86%).

¹H NMR (300 MHz, CDCl₃) δ 11.17 (brs, 1H), 8.29 (s, 1H), 7.43-7.73 (m, 5H), 7.29-7.42 (m, 4H), 6.66 (s, 1H), 6.29 (d, J = 8.6 Hz, 1H), 4.98 (t, J = 7.3 Hz, 1H), 4.34-4.49 (m, 1H), 3.05-3.18 (m, 1H), 1.82-2.15 (m, 3H), 1.74 (brs, 1H).

Example 15: Preparation of (S)-2-amino-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one



Step 1: Preparation of (S)-3-(1-((5-acetyl-6-((4-methoxybenzyl)amino)-2-(methylthio)pyrimidine-4-yl)amino)ethyl)-8-chloro-2-phenylisoquinoline-1(2H)-

5 one

1.5 g of (S)-3-(1-((5-acetyl-6-((4-methoxybenzyl)amino)-2-(methylthio)pyrimidine-4-yl)amino)ethyl)-8-chloro-2-phenylisoquinoline-1(2H)-one was prepared as a pale yellow solid by using 920
 10 mg (2.723 mmol) of 1-(4-chloro-6-((4-methoxybenzyl)amino)-2-(methylthio)pyrimidine-5-yl)ethane-1-one and 895 mg (2.996 mmol) of (S)-3-(1-aminoethyl)-8-chloro-2-phenylisoquinoline-1(2H)-one according to the same manner as described in step 1 of
 15 Example 5 (2.199 mmol, yield: 92%).

¹H NMR (300 MHz, CDCl₃) δ 8.71 (d, J = 6.7 Hz, 1H), 7.56-7.63 (m, 1H), 7.38-7.55 (m, 6H), 7.27-7.38 (m, 2H), 7.20-7.26 (m, 2H), 6.87 (d, J = 8.9 Hz, 2H), 6.53 (brs, 1H), 6.47 (s, 1H), 4.91 (t, J = 7.7 Hz, 1H), 4.67 (d, J = 4.5 Hz, 2H), 3.80 (s, 3H), 3.51 (s, 3H), 3.37 (s, 3H), 1.35 (d, J = 6.7 Hz, 3H).

Step 2: Preparation of (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-8-(4-methoxybenzyl)-2-(methylthio)pyrido[2,3-d]pyrimidine-5(8H)-one

90 mg of (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a white solid by using 700 mg (1.166 mmol) of (S)-3-(1-((5-acetyl-6-((4-methoxybenzyl)amino)-2-(methylthio)pyrimidine-4-yl)amino)ethyl)-8-chloro-2-phenylisoquinoline-1(2H)-one prepared in step 1 according to the same manner as described in step 2 of Example 5 (0.160 mmol, yield: 44%).

¹H NMR (300 MHz, CDCl₃) δ 10.71 (d, J = 6.9 Hz, 1H), 8.30 (s, 1H), 7.47-7.57 (m, 3H), 7.29-7.47 (m, 6H), 7.23 (d, J = 8.3 Hz, 2H), 6.87 (d, J = 8.3 Hz, 2H), 6.59 (s, 1H), 6.30 (d, J = 8.3 Hz, 1H), 5.35 (s,

2H), 4.93 (t, J = 6.9 Hz, 1H), 3.78 (s, 3H), 1.45 (d, J = 6.9 Hz, 3H).

Step 3: Preparation of (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-2-(methylthio)pyrido[2,3-d]pyrimidine-5(8H)-one

195 mg of (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-2-(methylthio)pyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a white solid by using 249 mg (0.408 mmol) of (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-8-(4-methoxybenzyl)-2-(methylthio)pyrido[2,3-d]pyrimidine-5(8H)-one prepared in step 2 according to the same manner as described in step 8 of Example 1 (0.398 mmol, yield: 98%).

¹H NMR (300 MHz, DMSO-d₆) δ 10.56 (d, J = 6.6 Hz, 1H), 7.58-7.72 (m, 3H), 7.47-7.57 (m, 2H), 7.34-7.46 (m, 2H), 7.29 (d, J = 3.2 Hz, 2H), 6.77 (s, 1H), 6.07 (d, J = 6.6 Hz, 1H), 4.74 (t, J = 7.7 Hz, 1H), 2.34 (s, 3H), 1.39 (d, J = 6.6 Hz, 3H).


Step 4: Preparation of (S)-2-amino-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one

100 mg (0.204 mmol) of (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-2-(methylthio)pyrido[2,3-d]pyrimidine-5(8H)-one prepared in step 3 was dissolved in 7 mL of dichloromethane:methanol (2:5), to which 70 mg (0.408 mmol) of 3-chloroperoxybenzoic acid (mCPBA) was added at 0°C, followed by stirring at room temperature for 30 minutes. Water was added to the reaction mixture, followed by extraction with ethyl acetate. The extract was washed with saturated sodiumbicarbonate solution. The organic layer was separated, dried (Na₂SO₄), and concentrated. The obtained compound was dissolved in 5 mL of tetrahydrofuran:isopropanol (1:1), to which 2 mL of 28% ammonia water was added, followed by stirring at 50°C for 10 hours. The reaction mixture was cooled down to room temperature. Water was added to the reaction mixture, followed by extraction with ethyl acetate. The extracted organic layer was dried (Na₂SO₄) and concentrated. The obtained compound was separated by column chromatography (SiO₂, eluent: 2% methanol dichloromethane/methanol, 50/1 -> dichloromethane/methanol, 20/1) to give 49 mg of the target compound (S)-2-amino-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-

yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one as a white solid (0.107 mmol, yield: 52%).

¹H NMR (300 MHz, CDCl₃) δ 10.48 (d, J = 5.9 Hz, 1H), 7.28-7.55 (m, 8H), 7.22 (d, J = 7.3 Hz, 1H), 6.60 (s, 1H), 6.07 (d, J = 7.3 Hz, 1H), 4.99 (brs, 2H), 4.83 (t, J = 7.3 Hz, 1H), 1.40 (d, J = 7.3 Hz, 3H).

Example 16: Preparation of (S)-2-amino-4-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)pyrido[2,3-d]pyrimidine-5(8H)-one



Step 1: Preparation of (S)-4-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)-2-(methylthio)pyrido[2,3-d]pyrimidine-5(8H)-one

42 mg of (S)-4-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)-2-(methylthio)pyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a white solid by using 25 mg (0.11 mmol, 1.1 equivalent) of 4-chloro-2-(methylthio)pyrido[2,3-d]pyrimidine-5(8H)-one and 33 mg (0.10 mmol) of (S)-8-

chloro-2-phenyl-3-(pyrrolidine-2-yl)isoquinoline-
1(2H)-one according to the same manner as described in
step 1 of Example 15 (0.081 mmol, yield: 81%).


¹H NMR(300 MHz, CDCl₃) δ 11.55 (s, -NH), 8.08 (s,
5 1H), 7.85-7.83 (m, 1H), 7.69-7.64 (m, 1H), 7.69-7.33
(m, 7H), 6.63 (s, 1H), 5.02-4.96 (m, 1H), 4.40-4.31
(m, 1H), 3.18-3.12 (m, 1H), 2.57 (s, 3H), 2.12-1.98
(m, 2H), 1.87-1.81 (m, 1H), 1.64-1.55 (m, 1H).

10 Step 2: Preparation of (S)-2-amino-4-(2-(8-
chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-
yl)pyrrolidine-1-yl)pyrido[2,3-d]pyrimidine-5(8H)-one

23 mg of (S)-2-amino-4-(2-(8-chloro-1-oxo-2-
phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-
15 yl)pyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a
white solid by using 35 mg (0.068 mmol) of (S)-4-(2-
(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-
yl)pyrrolidine-1-yl)-2-(methylthio)pyrido[2,3-
d]pyrimidine-5(8H)-one according to the same manner as
20 described in step 4 of Example 15 (0.047 mmol, yield:
70%).

¹H NMR(300 MHz, CDCl₃) δ 8.76 (br s, 1H), 7.81-
7.20 (m, 9H), 6.73 (s, 1H), 6.19 (d, J = 7.5Hz, 1H),
5.02-4.95 (m, 1H), 4.75 (br s, 2H), 4.44-4.31 (m, 1H),
25 3.20-3.10 (m, 1H), 2.57 (s, 3H), 2.10-1.40 (m, 4H).

Example 17: Preparation of (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-6-methylpyrido[2,3-d]pyrimidine-5(8H)-one



Step 1: Preparation of (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-8-(4-methoxybenzyl)-6-methylpyrido[2,3-d]pyrimidine-5(8H)-one

168 mg of (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-8-(4-methoxybenzyl)-6-methylpyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a white solid by using 300 mg (0.528 mmol) of (S)-8-chloro-3-(1-((4-methoxybenzyl)amino)-5-propionylpyrimidine-4-yl)amino)-2-phenylisoquinoline-1(2H)-one according to the same manner as described in step 2 of Example 5 (0.291 mmol, yield: 55%).

¹H NMR (300 MHz, CDCl₃) δ 10.87 (d, J = 7.3 Hz, 1H), 8.28 (s, 1H), 7.29-7.57 (m, 9H), 7.22 (d, J = 8.3

Hz, 2H), 6.86 (d, J = 8.3 Hz, 2H), 6.60 (s, 1H), 5.36 (s, 2H), 4.93 (t, J = 6.3 Hz, 1H), 3.79 (s, 3H), 2.06 (s, 3H), 1.46 (d, J = 7.3 Hz, 3H).


5 Step 2: Preparation of (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-6-methylpyrido[2,3-d]pyrimidine-5(8H)-one

120 mg of (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-6-methylpyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a white solid by using 168 mg (0.291 mmol) of (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-8-(4-methoxybenzyl)-6-methylpyrido[2,3-d]pyrimidine-5(8H)-one prepared in step 1 according to the same manner as described in step 8 of Example 1 (0.262 mmol, yield: 90%).

¹H NMR (300 MHz, DMSO-d₆) δ 10.69 (d, J = 5.0 Hz, 1H), 8.15 (s, 1H), 7.77 (s, 1H), 7.57-7.69 (m, 2H), 7.30-7.56 (m, 7H), 6.76 (s, 1H), 4.64-4.73 (m, 1H), 1.92 (s, 3H), 1.40 (d, J = 6.0 Hz, 3H).

Example 18: Preparation of (S)-2-amino-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-

yl)ethyl)amino)-6-methylpyrido[2,3-d]pyrimidine-5(8H)-
one



Step 1: Preparation of (S)-4-((1-(8-chloro-1-oxo-

5 2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-6-
methyl-2-(methylthio)pyrido[2,3-d]pyrimidine-5(8H)-one

110 mg of (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-
dihydroisoquinoline-3-yl)ethyl)amino)-6-methyl-2-
(methylthio)pyrido[2,3-d]pyrimidine-5(8H)-one was
10 prepared as a white solid by using 150 mg (0.240 mmol)
of (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-
dihydroisoquinoline-3-yl)ethyl)amino)-8-(4-
methoxybenzyl)-6-methyl-2-(methylthio)pyrido[2,3-
d]pyrimidine-5(8H)-one according to the same manner as
15 described in step 8 of Example 1 (0.218 mmol, yield:
91%).


¹H NMR (300 MHz, CDCl₃) δ 10.97(brs, 1H), 10.82
(d, J = 7.4 Hz, 1H), 7.28-7.60 (m, 9H), 6.67 (s, 1H),
5.11 (t, J = 7.4 Hz, 1H), 2.38 (s, 3H), 2.06 (s, 3H),
20 1.48 (d, J = 7.4 Hz, 3H).

Step 2: Preparation of (S)-2-amino-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-6-methylpyrido[2,3-d]pyrimidine-5(8H)-one

5 84 mg of (S)-2-amino-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-6-methylpyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a white solid by using 110 mg (0.218 mmol) of (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-6-methyl-2-(methylthio)pyrido[2,3-d]pyrimidine-5(8H)-one prepared in step 1 according to the same manner as described in step 4 of Example 15 (0.178 mmol, yield: 81%).

15 ¹H NMR (300 MHz, CDCl₃) δ 10.67 (d, J = 5.9 Hz, 1H), 7.29-7.59 (m, 8H), 7.20 (s, 1H), 6.61 (s, 1H), 4.78-4.93 (m, 3H), 1.99 (s, 3H), 1.41 (d, J = 7.3 Hz, 3H).

Example 19: Preparation of (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-5-oxo-5,8-dihydropyrido[2,3-d]pyrimidine-6-carbonitrile



Step 1: Preparation of (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-8-(4-methoxybenzyl)-5-oxo-5,8-dihydropyrido[2,3-d]pyrimidine-6-carbonitrile

(S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-8-(4-methoxybenzyl)-5-oxo-5,8-dihydropyrido[2,3-d]pyrimidine-6-carbonitrile was prepared by using 4-hydroxy-8-(4-methoxybenzyl)-5-oxo-5,8-dihydropyrido[2,3-d]pyrimidine-6-carbonitrile and (S)-3-(1-aminoethyl)-8-chloro-2-phenylisoquinoline-1(2H)-one according to the same manner as described in step 7 of Example 1.

MS[m/z; (M + 1)+]: 590.

Step 2: Preparation of (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-5-oxo-5,8-dihydropyrido[2,3-d]pyrimidine-6-carbonitrile

(S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-5-oxo-5,8-

2.8 g (9.598 mmol) of 1-(4-chloro-6-((4-methoxybenzyl)amino)pyrimidine-5-yl)ethane-1-one prepared in step 4 of Example 1 was dissolved in 15 mL anhydrous dichloromethane, to which 2 mL (14.397 mmol) of Et₃N was added, followed by stirring at room temperature for 30 minutes. 3.09 mL (13.437 mmol) of TBS-OTf was added thereto, followed by stirring at room temperature for 12 hours. Then, the reaction mixture was concentrated under reduced pressure. The residue was separated by column chromatography (SiO₂, eluent: hexane/ethyl acetate, 10/1) to give 3.8 g of the target compound 5-(1-((tert-butyl)dimethylsilyl)oxy)vinyl)-6-chloro-N-(4-methoxybenzyl)pyrimidine-4-amine as a white liquid (9.360 mmol, yield: 98%).

¹H NMR (300 MHz, CDCl₃) δ 8.34 (s, 1H), 7.35 (d, J = 8.6 Hz, 2H), 6.99 (d, J = 8.4 Hz, 2H), 5.93 (br s, 1H), 4.97 (d, J = 1.4 Hz, 1H), 4.81 (d, J = 1.4 Hz, 1H), 4.73 (d, J = 5.4 Hz, 2H), 3.83 (s, 3H), 0.91 (s, 9H), 0.12 (m, 6H).

Step 2: Preparation of 1-(4-chloro-6-((4-methoxybenzyl)amino)pyrimidine-5-yl)-2-fluoroethane-1-one

3.8 g (9.360 mmol) of 5-(1-((tert-butyl)dimethylsilyl)oxy)vinyl)-6-chloro-N-(4-methoxybenzyl)pyrimidine-4-amine prepared in step 1 was dissolved in 40 mL of anhydrous acetonitrile, to which 3.65 g (10.300 mmol) of 1-chloromethyl-4-fluoro-1,4-diazoniabicyclo[2.2.2]octanebis(tetrafluoroborate) (selectfluor) was added, followed by stirring at room temperature for 15 hours. Ethyl acetate and water was added to the reaction mixture, followed by extraction. The extracted organic layer was dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was separated by column chromatography (SiO₂, eluent: hexane/ethylacetate, 4/1) to give 2.5 g of the target compound 1-(4-chloro-6-((4-methoxybenzyl)amino)pyrimidine-5-yl)-2-fluoroethane-1-one as a white liquid (8.072 mmol, yield: 86%).

¹H NMR (300 MHz, CDCl₃) δ 9.55 (br s, 1H), 8.44 (s, 1H), 7.28 (d, J = 7.9 Hz, 2H), 6.91 (d, J = 8.4 Hz, 2H), 5.60 (s, 1H), 5.51 (s, 1H), 4.73 (d, J = 5.0 Hz, 2H), 3.83 (s, 3H).

Step 3: Preparation of 3-(dimethylamino)-2-fluoro-1-(4-methoxy-6-((4-methoxybenzyl)amino)pyrimidine-5-yl)prop-2-en-1-one

2.5 g (8.072 mmol) of 1-(4-chloro-6-((4-methoxybenzyl)amino)pyrimidine-5-yl)-2-fluoroethane-1-one prepared in step 2 was dissolved in 50 mL of anhydrous toluene, to which 10.76 mL (80.720 mmol) of N,N-dimethylformamide dimethyl acetal (DMF-DMA) was added, followed by stirring at 90°C for 1 hour. The reaction mixture was cooled down to room temperature. Ethyl acetate and water was added to the reaction mixture, followed by extraction. The extracted organic layer was dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was separated by column chromatography (SiO₂, eluent: hexane/ethyl acetate, 3/1, -> hexane/ethyl acetate, 1/1) to give 2.1 g of the target compound 3-(dimethylamino)-2-fluoro-1-(4-methoxy-6-((4-methoxybenzyl)amino)pyrimidine-5-yl)prop-2-en-1-one as a yellow liquid (5.827 mmol, yield: 72%).

¹H NMR (500 MHz, CDCl₃) δ 8.19 (s, 1H), 7.28 (d, J = 8.4 Hz, 2H), 6.86 (d, J = 8.4 Hz, 2H), 4.64 (d, J = 5.5 Hz, 2H), 3.91 (s, 3H), 3.80 (s, 3H), 3.12 (s, 6H).

Step 4: Preparation of 6-fluoro-4-hydroxy-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one

2.1 g (5.827 mmol) of (Z)-3-(dimethylamino)-2-fluoro-1-(4-methoxy-6-((4-

methoxybenzyl)amino)pyrimidine-5-yl)prop-2-en-1-one prepared in step 3 was dissolved in 120 mL of acetic acid:water (5:1), which was stirred at 90 ~ 150°C for 2 days. The reaction mixture was cooled down to room temperature and filtered under reduced pressure. Isopropanol and ether were added thereto, followed by filtration to give 1.5 g of the target compound 6-fluoro-4-hydroxy-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one as a pale yellow solid (4.978 mmol, yield: 85%).

¹H NMR (500 MHz, DMSO-d₆) δ 8.39 (d, J = 7.7 Hz, 1H), 8.27 (s, 1H), 7.28 (d, J = 8.5 Hz, 2H), 6.90 (d, J = 8.8 Hz, 2H), 5.34 (s, 2H), 3.73 (s, 3H).

Step 5: Preparation of 4-chloro-6-fluoro-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one

50 mg (0.166 mmol) of 6-fluoro-4-hydroxy-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one prepared in step 4 was dissolved in 2 mL of anhydrous toluene, to which 131 mg (0.498 mmol) of triphenylphosphine (PPh₃) and 50 µL (0.498 mmol) of trichloroacetonitrile (CCl₃CN) were added, followed by stirring at 120°C for 4 hours. The reaction mixture was cooled down to room temperature. Ethyl acetate and water were added thereto, followed by extraction.

The extracted organic layer was dried (Na_2SO_4), filtered, and concentrated under reduced pressure. The residue was separated by column chromatography (SiO_2 , eluent: hexane/ethyl acetate, 5/1 -> 5 hexane/ethyl acetate, 3/1) to give 4 mg of the target compound 4-chloro-6-fluoro-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one as a pale yellow solid (0.013 mmol, yield: 8%).

^1H NMR (300 MHz, CDCl_3) δ 8.87 (s, 1H), 7.71 (d, J = 7.1 Hz, 1H), 7.24 (d, J = 8.8 Hz, 2H), 6.90 (d, J = 8.8 Hz, 2H), 5.47 (s, 2H), 3.81 (s, 3H).

Step 6: Preparation of (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-6-fluoro-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one

4 mg (0.013 mmol) of 4-chloro-6-fluoro-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one prepared in step 5 and 4 mg (0.014 mmol) of (S)-3-(1-aminoethyl)-8-chloro-2-phenylisoquinoline-1(2H)-one were dissolved in 1 mL of anhydrous dimethylsulfoxide (DMSO), to which 6.6 μL (0.039 mmol) of diisopropylethylamine (DIPEA) was added, followed by stirring at 70°C for 5 hours. The reaction mixture was cooled down to room temperature. Ethyl acetate

and water were added thereto, followed by extraction. The extracted organic layer was dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was separated by column chromatography
5 (SiO₂, eluent: hexane/ethyl acetate, 4/1 -> hexane/ethyl acetate, 1/1) to give 6 mg of the target compound (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-6-fluoro-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one as a
10 pale yellow solid (0.010 mmol, yield: 82%).

¹H NMR (300 MHz, CDCl₃) δ 10.53 (d, J = 6.6 Hz, 1H), 8.31 (s, 1H), 7.61 (d, J = 6.9 Hz, 1H), 7.47-7.57 (m, 2H), 7.30-7.47 (m, 6H), 7.22 (d, J = 8.2 Hz, 2H), 7.88 (d, J = 8.2 Hz, 2H), 6.58 (s, 1H), 5.38 (s, 2H),
15 4.95 (q, J = 4.2 Hz, 5.2 Hz, 1H), 3.79 (s, 3H), 1.46 (d, J = 6.9 Hz, 3H).

Step 7: Preparation of (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-6-fluoropyrido[2,3-d]pyrimidine-5(8H)-one
20


3 mg of (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-6-fluoropyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a pale yellow solid by using 6 mg (0.010 mmol) of
25 (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-

dihydroisoquinoline-3-yl)ethyl)amino)-6-fluoro-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one prepared in step 6 according to the same manner as described in step 8 of Example 1 (0.006 mmol, yield: 5 63%).

¹H NMR (500 MHz, CD₃OD) δ 8.19 (s, 1H), 8.07 (d, J = 6.5 Hz, 1H), 7.62 (t, J = 4.7 Hz, 2H), 7.53-7.60 (m, 2H), 7.39-7.45 (m, 3H), 7.34 (t, J = 7.9 Hz, 1H), 6.90 (s, 1H), 4.99-5.05 (m, 1H), 1.56 (d, J = 6.6 Hz, 3H).

10

Example 21: Preparation of (S)-4-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)-6-fluoropyrido[2,3-d]pyrimidine-5(8H)-one



15

Step 1: Preparation of (S)-4-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)-6-fluoro-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one

50 mg (0.166 mmol) of 6-fluoro-4-hydroxy-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one was dissolved in 2 mL of anhydrous dimethylformamide, to 20

which 95 mg (0.216 mmol) of (benzotriazole-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate (BOP) and 81 mg (0.249 mmol) of cesiumcarbonate (Cs_2CO_3) were added, followed by
5 stirring at room temperature for 30 minutes. 55 mg (0.183 mmol) of (S)-2-(1-aminoethyl)-5-chloro-3-phenylquinazoline-4(3H)-one was added thereto, followed by stirring at 60 ~ 80°C for 2 hours. Ethyl acetate and water were added to the reaction mixture,
10 followed by extraction. The extracted organic layer was dried (Na_2SO_4), filtered, and concentrated under reduced pressure. The residue was separated by column chromatography (SiO_2 , eluent: hexane/ethyl acetate, 5/1 -> hexane/ethyl acetate, 1/1) to give 10 mg of the
15 target compound (S)-4-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)-6-fluoro-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one as a pale yellow solid (0.017 mmol, yield: 10%).


^1H NMR (300 MHz, CDCl_3) δ 10.87 (d, J = 7.0 Hz, 1H), 8.29 (s, 1H), 7.72 (d, J = 7.8 Hz, 1H), 7.44-7.62 (m, 7H), 7.33 (d, J = 7.3 Hz, 1H), 7.21 (d, J = 8.6 Hz, 2H), 6.87 (d, J = 8.6 Hz, 2H), 5.32 (s, 2H), 5.11 (q, J = 5.4 Hz, 6.8 Hz, 1H), 3.79 (s, 3H), 1.50 (d, J = 7.1 Hz, 3H).

Step 2: Preparation of (S)-4-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)-6-fluoropyrido[2,3-d]pyrimidine-5(8H)-one

6 mg of (S)-4-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)-6-fluoropyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a pale yellow solid by using 10 mg (0.017 mmol) of (S)-4-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)-6-fluoro-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one prepared in step 1 according to the same manner as described in step 8 of Example 1 (0.013 mmol, yield: 76%).

¹H NMR (300 MHz, CDCl₃) δ 10.84 (d, J = 8.1 Hz, 1H), 8.22 (s, 1H), 7.72 (d, J = 7.9 Hz, 1H), 7.52-7.64 (m, 6H), 7.42-7.49 (m, 2H), 7.35 (d, J = 7.3 Hz, 1H), 5.12 (q, J = 5.5 Hz, 6.7 Hz, 1H), 1.51 (d, J = 6.7 Hz, 3H).

Example 22: Preparation of (S)-6-chloro-4-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one




10 mg (0.022 mmol) of (S)-4-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one prepared in Example 1 was dissolved in 1 mL of acetic acid, to which 3.3 mg (0.025 mmol) of N-chlorosuccinimide (NCS) was added, followed by stirring at 50 ~ 60°C for 12 hours. The reaction mixture was cooled down to room temperature and filtered under reduced pressure. Saturated sodiumbicarbonate aqueous solution was added thereto, followed by neutralization. Dichloromethane and water were added thereto, followed by extraction. The extracted organic layer was dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was separated by column chromatography (SiO₂, eluent: dichloromethane/methanol, 50/1 -> dichloromethane/methanol, 20/1) to give 3 mg of the target compound (S)-6-chloro-4-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-

yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one as a pale yellow solid (0.006 mmol, yield: 28%).

¹H NMR (300 MHz, CDCl₃) δ 9.62 (d, J = 8.1 Hz, 1H), 8.07 (s, 1H), 7.74 (d, J = 7.3 Hz, 1H), 7.46-7.67 (m, 6H), 7.29-7.40 (m, 2H), 5.08-5.18 (m, 1H), 1.45 (d, J = 5.7 Hz, 3H).

Example 23: Preparation of (S)-6-chloro-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one




10 mg (0.023 mmol) of (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one prepared in Example 5 was dissolved in 2 mL of methanol:water (1:1), to which 34 µL (0.068 mmol) of 2N sodiumhydroxide and 34 µL of 12% sodiumhypochloride at 0°C, followed by stirring at room temperature for 1 hour. The reaction mixture was filtered under reduced pressure, followed by neutralization with 1N HCl.

Dichloromethane and water were added thereto, followed by extraction. The extracted organic layer was dried (Na_2SO_4), filtered, and concentrated under reduced pressure. The residue was separated by column chromatography (SiO_2 , eluent: dichloromethane/methanol, 50/1 -> dichloromethane/methanol, 20/1) to give 2 mg of the target compound (S)-6-chloro-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one as a yellow solid (0.004 mmol, yield: 19%).

^1H NMR (300 MHz, CDCl_3) δ 10.53 (d, J = 5.6 Hz, 1H), 8.25 (s, 1H), 7.84 (s, 1H), 7.49-7.57 (m, 2H), 7.40-7.47 (m, 4H), 7.30-7.39 (m, 2H), 6.58 (s, 1H), 4.95 (t, J = 6.7 Hz, 1H), 1.47 (d, J = 6.6 Hz, 3H).


Example 24: Preparation of (S)-6-chloro-4-((1-(4,8-dichloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one



50 mg (0.113 mmol) of (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one prepared in Example 5 was dissolved in 2 mL of acetic acid, to which 17 mg (0.124 mmol) of N-chlorosuccinimide (NCS) was added, followed by stirring at 50°C for 15 hours. The reaction mixture was filtered under reduced pressure. Saturated sodiumbicarbonate aqueous solution was added thereto, followed by neutralization. Dichloromethane and water were added thereto, followed by extraction. The extracted organic layer was dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was separated by column chromatography (SiO₂, eluent: dichloromethane/methanol, 50/1 -> dichloromethane/methanol, 20/1) to give 17 mg of the target compound (S)-6-chloro-4-((1-(4,8-dichloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one as a pale yellow solid (0.036 mmol, yield: 29%).

¹H NMR (300 MHz, CDCl₃) δ 10.92 (d, J = 4.4 Hz, 1H), 8.28 (s, 1H), 7.91-7.98 (m, 1H), 7.70-7.79 (m, 2H), 7.48-7.64 (m, 5H), 7.20 (d, J = 6.2 Hz, 1H), 5.04 (t, J = 7.12 Hz, 1H), 1.68 (d, J = 7.1 Hz, 3H).

Example 25: Preparation of (S)-2-amino-4-((1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one



5 Step 1: Preparation of 4,6-dichloro-2-methylmercaptopyrimidine-5-carbaldehyde

8.36 g of 4,6-dichloro-2-methylmercaptopyrimidine-5-carbaldehyde was prepared as a white solid by using 7.91g (50.0 mmol) of 4,6-dihydroxy-2-methylmercaptopyrimidine according to the same manner as described in step 1 of Example 1 (37.5 mmol, yield: 75%).

^1H NMR (300 MHz, CDCl_3) δ 10.38 (s, 1H), 2.64 (s, 3H).

Step 2: Preparation of 1-(4,6-dichloro-2-methylmercaptopyrimidine-5-yl)ethane-1-ol

2.27 g of 1-(4,6-dichloro-2-methylmercaptopyrimidine-5-yl)ethane-1-ol was prepared as a white solid by using 2.23 g (10.0 mmol) of 4,6-dichloro-2-methylmercaptopyrimidine-5-carbaldehyde prepared in step 1 according to the same manner as described in step 2 of Example 1 (9.5 mmol, yield: 95%).

¹H NMR (300 MHz, CDCl₃) δ 5.40-5.47 (m, 1H), 2.57 (s, 3H), 2.52 (d, J = 9.2 Hz, 1H), 1.64 (d, J = 6.8 Hz, 3H).

Step 3: Preparation of 1-(4,6-dichloro-2-methylmercaptopyrimidine-5-yl)ethane-1-one

1.09 g of 1-(4,6-dichloro-2-methylmercaptopyrimidine-5-yl)ethane-1-one was prepared as a white solid by using 1.20 g (5.0 mmol) of 1-(4,6-dichloro-2-methylmercaptopyrimidine-5-yl)ethane-1-ol prepared in step 2 according to the same manner as described in step 3 of Example 1 (4.6 mmol, yield: 92%).

¹H NMR (300 MHz, CDCl₃) δ 2.62 (s, 3H), 2.61 (s, 3H).

Step 4: Preparation of 1-(4-chloro-2-methylmercapto-6-((4-methoxybenzyl)amino)pyrimidine-5-yl)ethane-1-one

1.014 g of 1-(4-chloro-2-methylmercapto-6-((4-methoxybenzyl)amino)pyrimidine-5-yl)ethane-1-one was prepared as a colorless oil by using 712 mg (3.0 mmol) of 1-(4,6-dichloro-2-methylmercaptopyrimidine-5-yl)ethane-1-one prepared in step 3 according to the same manner as described in step 4 of Example 1 (3.0 mmol, yield: 100%).

¹H NMR (300 MHz, CDCl₃) δ 9.53 (br s, 1H, NH), 7.25 (d, J = 8.7 Hz, 2H), 6.87 (d, J = 8.7 Hz, 2H), 4.66 (d, J = 5.6 Hz, 2H), 3.80 (s, 3H), 2.71 (s, 3H), 2.50 (s, 3H).

15

Step 5: Preparation of (S)-1-(4-((1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)-6-((4-methoxybenzyl)amino)-2-(methylmercapto)pyrimidine-5-yl)ethane-1-one

376 mg of (S)-1-(4-((1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)-6-((4-methoxybenzyl)amino)-2-(methylmercapto)pyrimidine-5-yl)ethane-1-one was prepared as a colorless oil by using 338 mg (1.0 mmol) of 1-(4-chloro-2-methylmercapto-6-((4-methoxybenzyl)amino)pyrimidine-5-

25

yl)ethane-1-one prepared in step 4 and 267 mg (1.0 mmol) of (S)-1-(6-fluoro-3-(pyridine-2-yl)ethane-1-amine according to the same manner as described in step 1 of Example 5 (0.66 mmol, yield: 66%).

5 ^1H NMR (300 MHz, CDCl_3) δ 9.71 (br t, $J = 5.4$ Hz, 1H, NH), 7.79 (d, $J = 4.7$ Hz, 1H), 8.16 (s, 1H), 8.04 (m, 1H), 7.86-7.93 (m, 2H), 7.61-7.64 (m, 1H), 7.30-7.60 (m, 3H), 7.24 (m, 2H), 6.84 (m, 2H), 6.28 (m, 1H), 4.64 (d, $J = 5.4$ Hz, 2H), 3.78 (s, 3H), 2.77 (s, 10 3H), 2.37 (s, 3H), 1.27 (d, $J = 7.2$ Hz, 3H).

Step 6: Preparation of (S)-4-((1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)-8-(4-methoxybenzyl)-2-(methylmercapto)pyrido[2,3-
15 d]pyrimidine-5(8H)-one

168 mg of (S)-4-((1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)-8-(4-methoxybenzyl)-2-(methylmercapto)pyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a white solid by using 285 mg (0.5 mmol) of (S)-1-(4-((1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)-6-((4-methoxybenzyl)amino)-2-(methylmercapto)pyrimidine-5-yl)ethane-1-one prepared in step 5 according to the same manner as described in step 2 of Example 5 (0.29 mmol, yield: 58%).

¹H NMR (300 MHz, CDCl₃) δ 11.42 (br d, J = 7.6 Hz, 1H, NH), 8.77 (br d, J = 4.8 Hz, 1H), 8.27 (m, 1H), 8.11 (s, 1H), 7.82 (m, 1H), 7.61 (d, J = 7.8 Hz, 1H), 7.20-7.60 (m, 4H), 7.18 (d, J = 8.6 Hz, 2H), 6.84 (d, J = 8.6 Hz, 2H), 6.21 (d, J = 7.9 Hz, 1H), 6.14 (m, 1H), 5.26 (s, 2H), 3.78 (s, 3H), 2.36 (s, 3H), 1.56 (d, J = 6.6 Hz, 3H).

Steps 7 and 8: Preparation of (S)-2-amino-4-((1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one

145 mg (0.25 mmol) of (S)-4-((1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)-8-(4-methoxybenzyl)-2-(methylmercapto)pyrido[2,3-d]pyrimidine-5(8H)-one prepared in step 6 was dissolved in 5 mL of dichloromethane, to which 3-chloroperoxybenzoic acid (mCPBA) (2 equivalent) was added, followed by stirring at room temperature for 30 minutes. Water was added to the reaction mixture, followed by extraction with ethyl acetate. The extracted organic layer was washed with saturated sodium bicarbonate solution, separated, dried (Na₂SO₄), and concentrated. The obtained compound was dissolved in 5 mL of tetrahydrofuran:isopropanol (1:1), to which

2 mL of 28% ammonia water was added, followed by stirring at 50°C for 10 hours. The reaction mixture was cooled down to room temperature. Water was added thereto, followed by extraction with ethyl acetate. 5 The extracted organic layer was separated, dried (Na₂SO₄), and concentrated. The obtained compound was separated by column chromatography (SiO₂, eluent: hexane/ethyl acetate, 2/1 → ethyl acetate) to give 78 mg of the target compound (S)-2-amino-4-((1-(6-fluoro-10 3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one as a white solid (0.14 mmol, yield: 57%).

¹H NMR (300 MHz, CDCl₃) δ 11.20 (br d, J = 7.2 Hz, 1H, NH), 8.79 (m, 1H), 8.25 (m, 1H), 8.08 (s, 1H), 15 7.80 (m, 1H), 7.30-7.55 (m, 4H), 7.10-7.26 (m, 3H), 6.83 (m, 2H), 6.08 (d, J = 7.9 Hz, 1H), 5.85 (m, 1H), 5.16 (s, 2H), 4.84 (s, 2H), 3.77 (s, 3H), 1.62 (d, J = 6.6 Hz, 3H).


20 Step 9: Preparation of (S)-2-amino-4-((1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one

54 mg of (S)-2-amino-4-((1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-25 d]pyrimidine-5(8H)-one was prepared as a white solid

by using 70 mg (0.128 mmol) of (S)-2-amino-4-((1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one prepared in step 8 according to the same manner as described in step 8 of Example 1 (0.126 mmol, yield: 99%).

¹H NMR (300 MHz, CDCl₃) δ 11.39 (s, 1H), 8.77 (d, J = 4.5 Hz, 1H), 8.22 (dd, J = 9.0, 5.5 Hz, 1H), 8.03 (s, 1H), 7.86 (td, J = 1.5, 7.6 Hz, 1H), 7.54-7.36 (m, 4H), 7.19 (d, J = 7.6 Hz, 1H), 6.08 (d, J = 7.6 Hz, 1H), 5.93-5.84 (m, 1H), 5.59 (br s, 2H), 1.55 (d, J = 6.5 Hz, 3H).

Example 26: Preparation of 4-((1-(6-fluoro-3,4-di(pyridine-2-yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one



Step 1: Preparation of 4-((1-(6-fluoro-3,4-di(pyridine-2-yl)quinoline-2-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one

99 mg of 4-((1-(6-fluoro-3,4-di(pyridine-2-yl)quinoline-2-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a beige solid by using 100 mg (0.29 mmol)
5 of 1-(6-fluoro-3,4-di(pyridine-2-yl)quinoline-2-yl)ethane-1-amine prepared in Preparative Example 22 according to the same manner as described in step 7 of Example 1 (0.16 mmol, yield: 56%).

¹H NMR (300 MHz, CDCl₃) δ 8.69 (s, 2H), 8.39-8.34
10 (m, 2H), 7.65-7.42 (m, 4H), 7.24-7.19 (m, 3H), 7.13-7.05 (m, 2H), 6.91-6.84 (m, 2H), 6.32 (s, 1H), 5.61 (s, 1H), 5.34 (s, 2H), 3.77 (s, 3H), 3.57-3.50 (m, 1H), 3.46 (s, 1H), 1.59 (d, J = 2.1 Hz, 3H).


15 Step 2: Preparation of 4-((1-(6-fluoro-3,4-di(pyridine-2-yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one

54 mg of 4-((1-(6-fluoro-3,4-di(pyridine-2-yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-
20 5(8H)-one was prepared as a beige solid by using 90 mg (0.15 mmol) of 4-((1-(6-fluoro-3,4-di(pyridine-2-yl)quinoline-2-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one prepared in step 1 according to the same manner as

described in step 8 of Example 1 (0.11 mmol, yield: 75%).

^1H NMR (300 MHz, CDCl_3) δ 11.48 (br s, 1H), 10.69 (s, 1H), 8.67 (t, $J = 5.7\text{Hz}$, 2H), 8.67 (t, $J = 5.7\text{Hz}$, 2H), 8.37 (dd, $J = 9.2, 5.6\text{Hz}$, 1H), 8.17 (s, 2H), 7.54-7.44 (m, 3H), 7.34 (d, $J = 7.7\text{Hz}$, 1H), 7.21-7.13 (m, 3H), 7.08-7.02 (m, 2H), 6.29 (d, $J = 7.6\text{Hz}$, 1H), 5.70 (br s, 1H), 1.54 (d, $J = 2.3\text{Hz}$, 3H).

10 Example 27: Preparation of (S)-4-((1-(6-fluoro-3-phenyl-4-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one



15 Step 1: Preparation of (S)-4-((1-(6-fluoro-3-phenyl-4-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one

(S)-4-((1-(6-fluoro-3-phenyl-4-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a pale beige solid by using 120 mg (0.35

20

mmol) of (S)-1-(6-fluoro-3-phenyl-4-(pyridine-2-yl)quinoline-2-yl)ethane-1-amine prepared in Preparative Example 23 according to the same manner as described in step 7 of Example 1.

5 ^1H NMR(300 MHz, CDCl_3) δ 8.64 (br d, $J = 4.9\text{Hz}$, 1H), 8.29-8.35 (m, 2H), 7.42-7.55 (m, 3H), 7.13-7.34 (m, 7H), 6.91-7.09 (m, 3H), 6.82-6.89 (m, 2H), 6.30-6.32 (m, 1H), 5.65-5.69 (m, 1H), 5.30 (s, 2H), 3.78 (s, 3H), 1.47 (d, $J = 6.3\text{Hz}$, 3H).

10


Step 2: Preparation of (S)-4-((1-(6-fluoro-3-phenyl-4-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one

(S)-4-((1-(6-fluoro-3-phenyl-4-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a pale beige solid by using 95 mg (0.16 mmol) of (S)-4-((1-(6-fluoro-3-phenyl-4-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one
15 prepared in step 1 according to the same manner as described in step 8 of Example 1.
20

^1H NMR(300 MHz, CDCl_3) δ 11.58 (br s, 1H), 11.32 (br s, 1H), 8.65 (br d, $J = 4.2\text{Hz}$, 1H), 8.32-8.38 (m, 1H), 8.23 (s, 1H), 7.43-7.54 (m, 3H), 7.13-7.36 (m,

6H), 6.94-7.10 (m, 3H), 6.36 (d, J = 7.6Hz, 1H), 5.65-6.75 (m, 1H), 1.45 (d, J = 6.3Hz, 3H).

Example 28: Preparation of (S)-4-((1-(6-fluoro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one



Step 1: Preparation of (S)-4-((1-(6-fluoro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one

The target compound was prepared by using the compound prepared in Preparative Example 24 according to the same manner as described in step 7 of Example 1.


¹H NMR (300 MHz, CDCl₃) δ 10.91 (s, 1H), 8.75-8.62 (m, 1H), 8.37-8.26 (m, 1H), 7.89-7.71 (m, 2H), 7.89-7.71 (m, 2H), 7.57-7.40 (m, 3H), 7.21 (d, J = 7.2Hz, 2H), 6.31 (d, J = 7.7Hz, 2H), 5.41-5.26 (m, 2H), 5.13-4.99 (m, 1H), 3.78 (s, 3H), 1.53 (dd, J = 11.3, 6.7Hz, 3H).

Step 2: Preparation of (S)-4-((1-(6-fluoro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one

5 The target compound was prepared by using the compound prepared in step 1 according to the same manner as described in step 8 of Example 1.

¹H NMR(300 MHz, CDCl₃) δ 10.89 (d, J = 6.7 Hz, 1H), 8.77 (s, 1H), 8.68 (d, J = 7.4Hz, 1H), 8.17 (d, J = 3.2Hz, 1H), 7.90-7.73 (m, 3H), 7.59-7.47 (m, 3H), 10
6.35 (t, J = 4.9Hz, 2H), 5.13-5.03 (m, 1H), 1.55 (t, J = 8.0Hz, 3H).

Example 29: Preparation of (S)-4-((1-(6-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one



Step 1: Preparation of (S)-4-((1-(6-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one

The target compound was prepared by using the compound prepared in Preparative Example 25 according to the same manner as described in step 7 of Example 1.

5 ¹H NMR(300 MHz, CDCl₃) δ 11.01(d, J = 6.8 Hz, 1H),
8.27(s, 1H), 7.90-7.88(m, 2H), 7.90-7.88(m, 2H), 7.58-
7.44(m, 6H), 7.33(d, J = 6.8 Hz, 1H), 7.20(d, J = 7.8
Hz, 1H), 6.85(d, J = 8.0 Hz, 2H), 6.32(d, J = 7.8 Hz,
1H), 5.34(s, 1H), 5.18-5.09(m, 1H), 3.78(s, 3H),
10 1.50(d, J = 6.5 Hz, 3H).


Step 2: Preparation of (S)-4-((1-(6-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one

15 The target compound was prepared by using the compound prepared in step 1 according to the same manner as described in step 8 of Example 1.

¹H NMR(300 MHz, CDCl₃) δ 11.05(d, J = 6.9 Hz, 1H),
8.17(s, 1H), 7.90(d, J = 8.3 Hz, 1H), 7.85-7.80(m,
20 1H), 7.60-7.41(m, 6H), 6.36(d, J = 7.6 Hz, 1H), 5.20-
5.11(m, 1H), 1.51(d, J = 6.4 Hz, 3H).

Example 30: Preparation of (S)-4-((1-(6-fluoro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one

25



Step 1: Preparation of (S)-4-((1-(6-fluoro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one

The target compound was prepared by using the compound prepared in Preparative Example 26 according to the same manner as described in step 7 of Example 1.

¹H NMR (300 MHz, CDCl₃) δ 10.97-10.91(m, 1H), 8.28(s, 1H), 7.90-7.72(m, 3H), 7.59-7.50(m, 4H), 7.24-7.20(m, 3H), 6.86(d, J = 7.8Hz 2H), 6.32(d, J = 7.8 Hz, 1H), 5.35(s, 2H), 5.17-5.10(m, 1H), 3.78(s, 3H), 1.53(d, J = 6.5 Hz, 3H).

15

Step 2: Preparation of (S)-4-((1-(6-fluoro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one


The target compound was prepared by using the compound prepared in step 1 according to the same manner as described in step 8 of Example 1.

20

¹H NMR (300 MHz, CDCl₃) δ 10.84-10.78 (m, 1H), 8.20 (d, J = 5.4 Hz, 1H), 7.91-7.77 (m, 2H), 7.59-7.46 (m, 3H), 7.22-7.08 (m, 2H), 6.34 (d, J = 7.4 Hz, 1H), 5.18-5.09 (m, 1H), 1.53 (d, J = 6.5 Hz, 3H).

5

Example 31: Preparation of (S)-4-((1-(5-chloro-3-(2-chlorobenzyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one



10 단계 1: (S)-4-((1-(5-chloro-3-(2-chlorobenzyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one의 제조

The target compound was prepared by using the compound prepared in Preparative Example 27 according to the same manner as described in step 7 of Example 1.

¹H NMR (300 MHz, CDCl₃) δ 10.84 (d, J = 7.8 Hz, 1H), 8.32 (s, 1H), 7.71-7.58 (m, 2H), 7.47 (t, J = 8.3 Hz, 2H), 7.26-7.18 (m, 3H), 7.01 (s, 2H), 6.87 (d, J = 7.7 Hz, 2H), 6.79 (d, J = 5.2 Hz, 1H), 6.23 (d, J = 7.8 Hz,

20


1H), 5.79-5.73(m, 1H), 5.63-5.54(m, 1H), 5.46-5.36(m, 3H), 3.80(s, 3H), 1.56(d, J = 6.2 Hz, 3H).

Step 2: Preparation of ((S)-4-((1-(5-chloro-3-(2-chlorobenzyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one

The target compound was prepared by using the compound prepared in step 1 according to the same manner as described in step 8 of Example 1.

¹H NMR(300 MHz, CDCl₃) δ 10.84(d, J = 7.9 Hz, 1H), 8.22(s, 1H), 7.72(dd, J = 7.9, 0.8 Hz, 1H), 7.62(t, J = 7.8 Hz, 2H), 7.51-7.46(m, 2H), 7.33-7.30(m, 1H), 7.13-7.04(m, 2H), 6.83-6.80(m, 1H), 6.29(d, J = 7.7 Hz, 1H), 5.78(d, J = 17.1 Hz, 1H), 5.61-5.52(m, 1H), 5.49-5.41(m, 1H), 1.57(d, J = 6.6 Hz, 1H). (m, 1H), 5.46-5.36(m, 3H), 3.80(s, 3H), 1.56(d, J = 6.2 Hz, 3H).

Example 32: Preparation of (S)-4-((1-(6-fluoro-4-oxo-3-(pyridine-2-ylmethyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one



Step 1: Preparation of (S)-4-((1-(6-fluoro-4-oxo-3-(pyridine-2-ylmethyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one

The target compound was prepared by using the compound prepared in Preparative Example 28 according to the same manner as described in step 7 of Example 1.


¹H NMR (300 MHz, CDCl₃) δ 11.02 (d, J = 7.7 Hz, 1H), 8.41 (d, J = 4.4 Hz, 1H), 8.40 (s, 1H), 7.89 (dd, J = 8.5, 2.9 Hz, 1H), 7.78 (dd, J = 9.0, 4.9 Hz, 1H), 7.81-7.39 (m, 2H), 7.21 (d, J = 8.6 Hz, 2H), 7.09 (dd, J = 7.0, 5.3 Hz, 1H), 6.86 (d, J = 8.6 Hz, 2H), 6.29 (d, J = 7.9 Hz, 1H), 5.89-5.80 (m, 2H), 5.59 (d, J = 16.2 Hz, 1H), 5.33 (dd, J = 14.6, 16.1 Hz, 2H), 3.78 (s, 3H), 1.59 (d, J = 6.5 Hz, 3H).

Step 2: Preparation of (S)-4-((1-(6-fluoro-4-oxo-3-(pyridine-2-ylmethyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one

The target compound was prepared by using the compound prepared in step 1 according to the same manner as described in step 8 of Example 1.

¹H NMR (300 MHz, CDCl₃) δ 11.19(d, J = 7.6 Hz, 1H), 8.46(d, J = 4.3 Hz, 1H), 8.23(s, 1H), 7.90(dd, J = 8.5, 3.0 Hz, 1H), 7.81(dd, J = 9.0, 4.9 Hz, 1H), 7.63(td, J = 1.7, 7.7 Hz, 1H), 7.51-7.43(m, 2H), 7.31(d, J = 7.9 Hz, 1H), 7.15(dd, J = 7.0, 5.0 Hz, 1H), 6.34(d, J = 7.7 Hz, 1H), 5.95-5.81(m 2H), 5.54(d, J = 16.1 Hz, 1H), 1.61(d, J = 6.5 Hz, 3H).

Example 33: Preparation of 4-((1-(5-chloro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one



Step 1: Preparation of 4-((1-(5-chloro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one

4-((1-(5-chloro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a pale yellow solid by using 1-(5-chloro-3-(pyridine-2-yl)quinoline-2-yl)ethane-1-amine prepared in Preparative Example 29 according to the same manner as described in step 7 of Example 1.

^1H NMR (300 MHz, CDCl_3) δ 11.6(d, J = 7.4 Hz, 1H), 8.81(d, J = 4.8 Hz, 1H), 8.51(s, 1H), 8.32(s, 1H), 8.81(d, J = 4.8 Hz, 1H), 8.23(d, J = 8.0 Hz, 1H), 7.87-7.82(m, 1H), 7.67-7.60(m, 3H), 7.44(d, J = 7.9 Hz, 1H), 7.40-7.35(m, 1H), 7.19(d, J = 8.6 Hz, 2H), 6.85(d, J = 8.6 Hz, 2H), 6.31(d, J = 7.9 Hz, 1H), 6.16-6.07(m, 1H), 5.33(s, 2H), 3.78(s, 3H), 1.56(d, J = 6.6 Hz, 2H).

Step 2: Preparation of 4-((1-(5-chloro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one


4-((1-(5-chloro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one was prepared as a white solid by using 4-((1-(5-chloro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)-8-(4-methoxybenzyl)pyrido[2,3-d]pyrimidine-5(8H)-one

prepared in step 1 according to the same manner as described in step 8 of Example 1.


^1H NMR (300 MHz, CDCl_3) δ 11.54 (d, $J = 7.1$ Hz, 1H), 10.86 (s, 1H), 8.83 (d, $J = 4.7$ Hz, 1H), 8.54 (s, 1H), 8.26 (s, 1H), 8.24 (s, 1H), 7.87 (t, $J = 7.7$ Hz, 1H), 7.69-7.61 (m, 3H), 7.44-7.37 (m, 2H), 6.32 (d, $J = 6.8$ Hz, 1H), 6.20-6.12 (m, 1H), 1.57 (d, $J = 6.3$ Hz, 3H).

The following examples 34 ~ 65 were performed by the method represented by the reaction formula 2.

[Reaction Formula 2]



Example 34: Preparation of (S)-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one



Steps 1 and 2: Preparation of 4,6-dichloro-N-methylpyrimidine-5-carboxamide

1.00 g (5.65 mmol) of 4,6-dichloropyrimidine-5-carboxaldehyde was dissolved in 15 mL of CCl_4 , to which 0.78 mL (9.61 mmol) of sulfuryl chloride and 46 mg (0.28 mmol) of 2,2-azobis(2-methyl propionitrile) were added, followed by stirring at 80°C for 3 hours. The reaction mixture was cooled down to room temperature and filtered under reduced pressure. 5 mL of anhydrous toluene was added thereto, followed by filtration under reduced pressure. The resultant product was dissolved in 15 mL of anhydrous tetrahydrofuran, to which 4.73 mL of 2.0 M methylamine/tetrahydrofuran solution was added at -20°C , followed by stirring at -20°C for 2 hours. The reaction mixture was added with 1 N HCl and filtered

under reduced pressure. Ethyl acetate and water were added thereto, followed by extraction. The extracted organic layer was separated, dried (Na_2SO_4), filtered, and concentrated under reduced pressure. The residue
5 was separated by column chromatography (SiO_2 , eluent: dichloromethane/methanol, 20/1) to give 0.8 g of the target compound 4,6-dichloro-N-methylpyrimidine-5-carboxamide as a pale yellow solid (3.88 mmol, yield: 69%).
10 ^1H NMR (300 MHz, CDCl_3) δ 8.81 (s, 1H), 5.87 (brs, 1H), 3.08 (d, $J = 2.6$ Hz, 3H).

Step 3: Preparation of 4-chloro-6-((4-methoxybenzyl)amino)-N-methylpyrimidine-5-carboxamide
15 3.0 g (14.5 mmol) of 4,6-dichloro-N-methylpyrimidine-5-carboxamide prepared in step 1 and step 2 was dissolved in 80 mL of anhydrous tetrahydrofuran, to which 1.8 mL (15.2 mmol, 1.05 eq) of p-methoxybenzylamine and 2.8 mL (16.0 mmol, 1.1 eq)
20 of diisopropylethylamine (DIPEA) were added, followed by stirring for 6 hours. The reaction mixture was distilled under reduced pressure. Water was added thereto, followed by extraction with ethyl acetate. The extracted organic layer was separated, dried
25 (Na_2SO_4), filtered, and concentrated under reduced

pressure. The residue was separated by column chromatography (SiO₂, eluent: hexane/ethylacetate, 4/1) to give 3.73 g of the target compound 4-chloro-6-((4-methoxybenzyl)amino)-N-methylpyrimidine-5-carboxamide as a transparent oil (12.1 mmol, yield: 84%).

¹H NMR (300 MHz, CDCl₃) δ 8.31 (s, 1H), 8.24 (s, 1H), 7.25 (d, J = 8.9 Hz, 2H), 6.86 (d, J = 8.5 Hz, 2H), 6.70 (s, 1H), 4.62 (d, J = 5.4 Hz, 2H), 3.79 (s, 3H), 2.97 (d, J = 4.7 Hz, 2H).

Step 4: Preparation of 5-chloro-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

3.7 g (12 mmol) of 4-chloro-6-((4-methoxybenzyl)amino)-N-methyl-pyrimidine-5-carboxamide prepared in step 3, 3.6 g (120 mmol, >10 eq) of paraformaldehyde, and 228 mg (1.2 mmol, 0.1 eq) of p-toluenesulfonic acid were dissolved in 100 mL of toluene, which was stirred at 130°C for 12 hours in dean-stark trap. The reaction mixture was cooled down to room temperature and distilled under reduced pressure. The resultant product was separated by column chromatography (SiO₂, eluent:

hexane/ethylacetate, 3/1) to give 3.0 g of the target compound as a white solid (9.4 mmol, yield: 78%).

¹H NMR (300 MHz, CDCl₃) δ 8.43 (s, 1H), 7.22 (d, J = 8.9 Hz, 2H), 6.88 (d, J = 8.4 Hz, 2H), 4.86 (s, 2H),
5 4.55 (s, 2H), 3.81 (s, 3H), 2.98 (s, 3H).

Step 5: Preparation of (S)-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-
10 d]pyrimidine-4(1H)-one

50 mg (0.16 mmol) of 5-chloro-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one prepared in step 4 was dissolved in 2 mL of anhydrous dimethylsulfoxide
15 (DMSO), to which 57 mg (0.19 mmol, 1.2 equivalent) of (S)-2-(1-aminoethyl)-5-chloro-3-phenylquinazoline-4(3H)-one and 0.06 mL (0.35 mmol, 2.2 equivalent) of diisopropylethylamine (DIPEA) were added, followed by stirring at 70°C for 12 hours. The reaction mixture
20 was cooled down to room temperature. Water was added thereto, followed by extraction with ethyl acetate. The extracted organic layer was separated, dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was separated by column
25 chromatography (SiO₂, eluent: hexane/ethylacetate,

2/1) to give 88 mg of the target compound (S)-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one as a white
5 solid (0.15 mmol, yield: 94%).


¹H NMR (300 MHz, CDCl₃) δ 9.49 (d, J = 7.0 Hz, -NH), 8.03 (s, 1H), 7.70-7.68 (m, 1H), 7.61-7.44 (m, 7H), 7.31-7.28 (m, 1H), 7.21 (d, J = 8.6 Hz, 2H), 6.86 (d, J = 8.6 Hz, 2H), 5.06-5.02 (m, 1H), 4.74 (s, 2H),
10 4.45 (s, 2H), 3.79 (s, 3H), 2.89 (s, 3H), 1.44 (d, J = 6.6 Hz, 3H).

Step 6: Preparation of (S)-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-
15 methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

57 mg of (S)-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid by using 70 mg (0.13 mmol)
20 of (S)-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one prepared in step 5 according to the same manner as described in step 8 of Example 1 (0.12 mmol, yield:
25 99%).

¹H NMR (300 MHz, CDCl₃) δ 9.43 (d, J = 7.9 Hz, -1H), 7.88 (s, 1H), 7.69-7.67 (m, 1H), 7.61-7.43 (m, 6H), 7.32-7.29 (m, 1H), 6.97 (s, -NH), 5.02-4.97 (m, 1H), 4.69 (s, 2H), 2.97 (s, 3H), 1.43 (d, J = 5.9 Hz, 3H).

Example 35: Preparation of (S)-5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one



Step 1: Preparation of (S)-5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

83 mg of (S)-5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 5 of Example 34 except that (S)-2-(1-aminoethyl)-5-chloro-

3-(pyridine-3-yl)quinazoline-4(3H)-one was used (0.14 mmol, yield: 94%).

¹H NMR (300 MHz, CDCl₃) δ 9.39-9.35 (m, 1H), 8.76-8.71 (m, 1H), 8.03 (d, J = 8.1 Hz, 2H), 7.70-7.46 (m, 3H), 7.20 (d, J = 8.6 Hz, 2H), 6.86 (d, J = 8.6 Hz, 2H), 5.02-4.89 (m, 1H), 4.8-4.65 (m, 2H), 4.46 (s, 2H), 3.79 (s, 3H), 2.90 (s, 3H), 1.50-1.44 (m, 3H).


Step 2: Preparation of (S)-5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

56 mg of (S)-5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 6 of Example 34 except that (S)-5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was used (0.12 mmol, yield: 99%).

¹H NMR (300 MHz, CDCl₃) δ 9.30-9.25 (m, 1H), 8.74-8.71 (m, 1H), 8.57 (s, 1H), 7.91 (d, J = 8.9 Hz, 2H), 7.91 (d, J = 7.2 Hz, 1H), 7.70-7.59 (m, 2H), 7.53-7.44

(m, 2H), 5.88 (s, 1H), 5.73 (s, 1H), 5.00-4.86 (m, 1H), 4.72 (s, 2H), 2.99 (s, 3H), 1.50-1.43 (m, 3H).

Example 36: Preparation of (S)-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one



Step 1: Preparation of (S)-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

87 mg of (S)-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 5 of Example 34 except that (S)-2-(1-aminoethyl)-5-chloro-3-(3-fluorophenyl)quinazoline-4(3H)-one was used (0.14 mmol, yield: 91%).


¹H NMR (300 MHz, CDCl₃) δ 9.45-9.36 (m, 1H), 7.62-7.50 (m, 4H), 7.42-7.39 (m, 1H), 7.27-7.14 (m, 4H), 7.07-6.99 (m, 1H), 6.84 (d, J = 8.0 Hz, 2H), 4.84-4.69 (m, 3H), 4.73 (s, 2H), 3.89-3.80 (m, 1H), 3.79 (s, 3H), 3.57-3.48 (m, 1H), 2.90 (s, 3H), 2.33-2.24 (m, 1H), 2.13-2.06 (m, 2H), 1.50-1.44 (m, 3H).

Step 2: Preparation of (S)-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

61 mg of (S)-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 6 of Example 34 except that (S)-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was used (0.12 mmol, yield: 99%).

¹H NMR (300 MHz, CDCl₃) δ 9.40-9.31 (m, 1H), 7.89 (d, J = 4.65 Hz, 1H), 7.69-7.44 (m, 4H), 7.25-7.04 (m, 2H), 6.89-6.80 (m, 1H), 5.04-4.95 (m, 1H), 4.71 (s, 2H), 2.98 (m, 2H), 1.46 (m, J = 5.98, 3H).

Example 37: Preparation of (S)-5-((1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one



Step 1: Preparation of (S)-5-((1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

85 mg of (S)-5-((1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 5 of Example 34 except that (S)-2-(1-aminoethyl)-5-chloro-3-m-tolylquinazoline-4(3H)-one was used (0.14 mmol, yield: 89%).

¹H NMR (300 MHz, CDCl₃) δ 9.52-9.43 (m, 1H), 8.03 (s, 1H), 7.70-7.67 (m, 2H), 7.60-54 (m, 1H), 7.44-7.40 (m, 2H), 7.29-7.18 (m, 3H), 7.10 (s, 1H), 6.84 (d, J =


8.8 Hz, 2H), 5.10-04 (m, 1H), 4.73 (s, 2H), 4.44 (s, 2H), 3.78 (s, 3H), 2.89 (s, 3H), 2.35 (s, 3H), 1.47-1.43 (m, 3H).

5 Step 2: Preparation of (S)-5-((1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

57 mg of (S)-5-((1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 6 of Example 34 except that (S)-5-((1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was used (0.12 mmol, yield: 99%).

¹H NMR (300 MHz, CDCl₃) δ 9.44-9.36 (m, 1H), 7.90 (d, J = 4.9 Hz, 1H), 7.69 (d, J = 6.8 Hz, 1H), 7.61-7.56 (m, 1H), 7.46-7.37 (m, 2H), 7.31-7.28 (m, 1H), 7.23 (s, 1H), 7.07 (s, 1H), 6.11 (m, 1H), 5.08-5.02 (m, 1H), 4.70 (s, 2H), 2.99 (s, 3H), 2.35 (s, 3H), 1.45-1.43 (m, 3H).

Example 38: Preparation of (S)-5-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one



5

Step 1: Preparation of (S)-5-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

10 30 mg of (S)-5-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a pale yellow solid by using 20 mg (0.063 mmol) of 5-chloro-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one prepared in step 4 of Example 34 and 19 mg (0.063 mmol) of (S)-3-(1-aminoethyl)-8-chloro-2-phenylisoquinoline-1(2H)-one according to the same manner as described in step 5 of Example 34

15

20 (0.052 mmol, yield: 82%).


¹H NMR (300 MHz, CDCl₃) δ 9.16 (d, J = 7.8 Hz, 1H), 8.06 (s, 1H), 7.43-7.54 (m, 8H), 7.32 (d, J = 8.6 Hz, 2H), 7.22 (d, J = 8.4 Hz, 2H), 6.56 (s, 1H), 4.87 (t, J = 7.1 Hz, 1H), 4.75 (s, 2H), 4.47 (s, 2H), 3.80 (s, 3H), 2.91 (s, 3H), 1.38 (d, J = 6.8 Hz, 3H).

Step 2: Preparation of (S)-5-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

22 mg of (S)-5-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a yellow solid by using 30 mg (0.052 mmol) of (S)-5-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one prepared in step 1 according to the same manner as described in step 8 of Example 1 (0.048 mmol, yield: 92%).

¹H NMR (300 MHz, CDCl₃) δ 9.07 (d, J = 6.4 Hz, 1H), 7.92 (s, 1H), 7.36-7.51 (m, 7H), 7.29-7.32 (m, 1H), 6.55 (s, 1H), 6.11 (brs, 1H), 4.85 (t, J = 7.9 Hz, 1H), 4.73 (s, 2H), 2.99 (s, 3H), 1.38 (d, J = 7.2 Hz, 3H).

Example 39: Preparation of 3-methyl-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one



5 Step 1: Preparation of 1-(4-methoxybenzyl)-3-methyl-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

10 mg of 1-(4-methoxybenzyl)-3-methyl-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a yellow solid by using 8 mg (0.025 mmol) of 5-chloro-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one prepared in step 4 of Example 34 and 7.5 mg (0.030 mmol) of 1-(2-phenylquinoline-3-yl)ethane-1-amine according to the same manner as described in step 5 of Example 34 (0.019 mmol, yield: 75%).

¹H NMR (300 MHz, CDCl₃) δ 9.39 (d, J = 7.2 Hz, 1H), 8.25 (s, 1H), 8.15 (d, J = 9.3 Hz, 1H), 8.01 (s, 1H), 7.83 (d, J = 8.6 Hz, 1H), 7.76 (d, J = 7.2 Hz, 2H),

7.68-7.70 (m, 1H), 7.47-7.54 (m, 5H), 7.22 (d, J =8.6 Hz, 1H), 6.85 (d, J =8.6 Hz, 2H) 6.64 (t, J =7.2 Hz, 1H), 4.75 (d, J =6.5 Hz, 2H), 4.47 (s, 2H), 3.79 (s, 3H), 2.93 (s, 3H), 1.47 (m, 3H).

5


Step 2: Preparation of 3-methyl-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

6 mg of 3-methyl-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a yellow solid by using 10 mg (0.019 mmol) of 1-(4-methoxybenzyl)-3-methyl-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one prepared in
15 step 1 according to the same manner as described in step 8 of Example 1 (0.015 mmol, yield: 78%).

¹H NMR (300 MHz, CDCl₃) δ 9.52 (d, J =6.1 Hz, 1H), 8.21 (s, 1H), 8.14 (d, J =8.8 Hz, 1H), 7.92 (brs, 1H), 7.83 (d, J =7.5 Hz, 1H), 7.69 (d, J =8.1 Hz, 3H),
20 7.48-7.59 (m, 5H), 5.65 (t, J =6.8 Hz, 1H), 4.78 (brs, 2H) 2.99 (s, 3H), 1.44 (d, J =6.8 Hz, 3H).

Example 39-1: Preparation of (S)-3-methyl-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

25



Step 1: Preparation of (S)-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidin-4(1H)-

5 one

84 mg of (S)-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidin-4(1H)-one was prepared as a white solid according to the same manner as described in step 5 of Example 34 except that (S)-1-(2-phenylquinoline-3-yl)ethaneamine was used (0.15 mmol, yield: 99%).

¹H NMR (300 MHz, CDCl₃) δ 9.39 (d, J = 7.8 Hz, 1H), 8.23 (s, 1H), 8.44 (d, J = 8.1 Hz, 1H), 8.08 (s, 1H), 7.82 (d, J = 7.8 Hz, 1H), 7.75 (d, J = 7.8 Hz, 2H), 7.69-7.64 (m, 1H), 7.53-7.43 (m, 4H), 7.21 (d, J = 8.4 Hz, 1H), 6.85 (d, J = 9.0 Hz, 2H), 5.67-5.62 (m, 1H), 4.74 (d, J = 8.4 Hz, 2H), 4.47 (s, 2H), 3.79 (s, 3H), 2.92 (s, 3H), 1.40 (d, J = 6.5 Hz, 3H).


20

Step 2: Preparation of (S)-3-methyl-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

52 mg of (S)-3-methyl-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 6 of Example 34 except that (S)-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was used (0.12 mmol, yield: 99%).

¹H NMR (300 MHz, CDCl₃) δ 9.31 (d, J = 6.9 Hz, -1H), 8.22 (s, 1H), 8.13 (d, J = 7.7 Hz, -1H), 7.93 (s, 1H), 7.82 (d, J = 7.7 Hz, -1H), 7.75-7.64 (m, 3H), 7.53-7.43 (m, 4H), 6.71 (s, -NH), 5.66-5.56 (m, 1H), 4.69 (s, 2H), 2.98 (s, 3H), 1.41 (d, J = 7.4 Hz, 3H).

Example 40: Preparation of (S)-5-((1-(4,8-dichloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one



Step 1: Preparation of (S)-5-((1-(4,8-dichloro-1-oxo-2-phenylquinoline-3-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

(S)-5-((1-(4,8-dichloro-1-oxo-2-phenylquinoline-3-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared according to the same manner as described in step 5 of Example 34 except that (S)-3-(1-aminoethyl)-4,8-dihydro-2-phenylisoquinoline-1(2H)-one was used.

^1H NMR (300 MHz, CDCl_3) δ 9.54 (br d, $J = 6.6\text{Hz}$, 1H), 8.09 (s, 1H), 7.97 (dd, $J = 7.7, 1.5\text{Hz}$, 1H), 7.75 (br d, $J = 7.7\text{Hz}$, 1H), 7.46-7.61 (m, 6H), 7.15-7.21 (m, 3H), 6.80-6.87 (m, 2H), 4.97-5.05 (m, 1H), 4.72 (s, 2H), 4.39-4.47 (m, 2H), 3.78 (s, 3H), 2.88 (s, 3H), 1.60 (d, $J = 7.2\text{Hz}$, 3H).


Step 2: Preparation of (S)-5-((1-(4,8-dichloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-

yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

(S)-5-((1-(4,8-dichloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared according to the same manner as described in step 6 of Example 34 except that (S)-5-((1-(4,8-dichloro-1-oxo-2-phenylquinoline-3-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was used.

¹H NMR (500 MHz, CDCl₃) δ 9.48 (br s, 1H), 7.98 (dd, J = 8.1, 0.9 Hz, 1H), 7.96 (s, 1H), 7.74 (d, J = 7.8 Hz, 1H), 7.49-7.61 (m, 5H), 7.18-7.21 (m, 1H), 6.70 (br s, 1H), 4.96-4.50 (m, 1H), 4.67-4.72 (m, 2H), 2.98 (s, 3H), 1.62 (d, J = 7.2 Hz, 3H).

Example 41: Preparation of (S)-5-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one




Step 1: Preparation of (S)-5-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

prepared as a white solid according to the same manner as described in step 6 of Example 34 except that (S)-5-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was used (0.12 mmol, yield: 99%).

^1H NMR (300 MHz, CDCl_3) δ 9.35 (d, $J = 8.2$ Hz, 1H), 7.88 (s, 1H), 7.69-7.62 (m, 1H), 7.57-7.50 (m, 4H), 7.44-7.41 (m, 1H), 7.30-7.28 (m, 1H), 7.11-7.05 (m, 1H), 6.28 (s, 1H), 4.83-4.91 (m, 1H), 4.70 (s, 2H), 2.98 (s, 3H), 1.92-1.75 (m, 2H), 0.87-0.82 (m, 3H).

Example 42: Preparation of (S)-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one



Step 1: Preparation of (S)-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-

yl)-1-(4-methoxybenzyl)-3-methyl-2,3-
dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

102 mg of (S)-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-
dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-
5 methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-
d]pyrimidine-4(1H)-one was prepared as a white solid
according to the same manner as described in step 5 of
Example 34 except that (S)-5-chloro-3-phenyl-2-
(pyrrolidine-2-yl)quinazoline-4(3H)-one was used (0.16
10 mmol, yield: 99%).

¹H NMR (300 MHz, CDCl₃) δ 8.03 (s, 1H), 7.75-7.73
(m, 1H), 7.60-7.50 (m, 5H), 7.41-7.38 (m, 1H), 7.16
(d, J = 7.6 Hz, 2H), 6.84 (d, J = 8.6 Hz, 2H), 4.90-
4.64 (m, 1H), 3.79 (s, 3H), 3.72-3.65 (m, 1H), 2.89
15 (s, 3H), 2.38-2.28 (m, 1H), 1.86-1.76 (m, 4H).


Step 2: Preparation of (S)-5-(2-(5-chloro-4-oxo-
3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-
yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-
20 4(1H)-one

58 mg of (S)-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-
dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-
2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was
prepared as a white solid according to the same manner
25 as described in step 6 of Example 34 except that (S)-

5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was used (0.12 mmol, yield: 99%).

¹H NMR (300 MHz, CDCl₃) δ 7.91 (s, 1H), 7.71-7.69 (m, 1H), 7.58-7.48 (m, 5H), 7.40 (d, J = 7.5 Hz, 2H), 7.25-7.23 (m, 1H), 4.72-4.68 (m, 1H), 4.56 (s, 2H), 3.90-3.81 (m, 1H), 3.62-3.53 (m, 1H), 3.32-3.25 (m, 2H), 3.05 (s, 3H), 2.31-2.24 (m, 1H), 2.10-2.09 (m, 2H), 1.84-1.75 (m, 1H).

Example 43: Preparation of (S)-5-(2-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one



Step 1: Preparation of (S)-5-(2-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

106 mg of (S)-5-(2-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 5 of Example 34 except that (S)-5-chloro-3-(pyridine-3-yl)-2-(pyrrolidine-2-yl)quinazoline-4(3H)-one was used (0.17 mmol, yield: 99%).

¹H NMR (300 MHz, CDCl₃) δ 9.04 (s, 1H), 8.76 (d, J = 3.5 Hz, 1H), 8.54 (s, 1H), 8.12-8.00 (m, 3H), 7.61-7.40 (m, 7H), 7.17-7.14 (m, 2H), 6.84 (d, J = 8.5 Hz, 2H), 4.90-4.64 (m, 1H), 3.87-3.83 (m, 1H), 3.78 (s, 5H), 3.60-3.53 (m, 1H), 2.89 (s, 3H), 2.37-2.30 (m, 2H), 1.87-1.79 (m, 2H).

15


Step 2: Preparation of (S)-5-(2-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

59 mg of (S)-5-(2-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 6 of Example 34 except that (S)-5-(2-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-

dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was used (0.12 mmol, yield: 99%).

5 ^1H NMR (300 MHz, CDCl_3) δ 9.04 (s, 1H), 8.77-8.74 (m, 2H), 8.53 (s, 1H), 8.12-8.06 (m, 2H), 8.00 (s, 1H), 7.63-7.50 (m, 7H), 7.46-7.40 (m, 2H), 7.19-7.12 (m, 2H), 6.84 (d, $J = 8.5$ Hz, 2H), 4.87-4.67 (m, 2H), 4.36 (s, 2H), 3.60-3.53 (m, 1H), 2.89 (s, 3H), 2.39-
10 2.27 (m, 2H), 1.87-1.79 (m, 2H).

Example 44: Preparation of (S)-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one



Step 1: Preparation of (S)-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

102 mg of (S)-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 5 of Example 34 except that (S)-5-chloro-3-(3-fluorophenyl)-2-(pyrrolidine-2-yl)quinazoline-4(3H)-one was used (0.16 mmol, yield: 99%).

¹H NMR (300 MHz, CDCl₃) δ 8.06-8.03 (m, 1H), 7.61-7.40 (m, 4H), 7.42-7.39 (m, 1H), 7.27-7.14 (m, 3H), 7.07-6.98 (m, 1H), 6.84 (d, J = 8.4 Hz, 2H), 4.85-4.69 (m, 2H), 4.37 (s, 2H), 3.87-3.83 (m, 1H), 3.78 (s, 3H), 3.56-3.48 (m, 1H), 2.90 (s, 3H), 2.32-2.24 (m, 1H), 2.12-2.07 (m, 1H), 1.87-1.76 (m, 2H).

15


Step 2: Preparation of (S)-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

20 60 mg of (S)-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 6 of Example 34 except
25 that (S)-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-

dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was used (0.12 mmol, yield: 99%).

¹H NMR (300 MHz, CDCl₃) δ 7.95-7.91 (m, 1H), 7.57-7.38 (m, 5H), 7.24-7.19 (m, 1H), 7.07-6.98 (m, 1H), 6.13-6.10 (m, 1H), 6.02-6.00 (m, 1H), 4.79-4.37 (m, 1H), 4.66-4.53 (m, 2H), 3.79-3.70 (m, 1H), 3.07 (s, 3H), 2.38-2.32 (m, 1H), 2.13-2.01 (m, 2H), 1.95-1.82 (m, 2H).

Example 45: Preparation of (S)-5-(2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one



Step 1: Preparation of (S)-5-(2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

95 mg of (S)-5-(2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 5 of Example 34 except that (S)-5-chloro-2-(pyrrolidine-2-yl)-3-m-tolylquinazoline-4(3H)-one was used (0.15 mmol, yield: 95%).

¹H NMR (300 MHz, CDCl₃) δ 8.04 (d, J = 6.3 Hz, 1H), 7.56-7.44 (m, 4H), 7.41-7.38 (m, 2H), 7.30 (d, J = 7.6 Hz, 1H), 7.16 (d, J = 8.5 Hz, 2H), 7.04-7.02 (m, 1H), 6.84 (d, J = 8.5 Hz, 2H), 4.81-4.71 (m, 3H), 4.37 (s, 2H), 3.86-3.81 (m, 1H), 3.78 (s, 3H), 3.56-3.51 (m, 1H), 2.89 (s, 3H), 2.76-2.72 (m, 1H), 2.42 (s, 3H), 2.33-2.26 (m, 1H), 2.13-2.07 (m, 2H), 1.84-1.76 (m, 1H).


Step 2: Preparation of (S)-5-(2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

60 mg of (S)-5-(2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same

manner as described in step 6 of Example 34 except that (S)-5-(2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was used (0.12 mmol, yield: 99%).

¹H NMR (300 MHz, CDCl₃) δ 7.88 (s, 1H), 7.49-7.44 (m, 4H), 7.41-7.37 (m, 2H), 7.30-7.27 (m, 1H), 7.03-7.01 (m, 1H), 4.78-4.75 (m, 1H), 4.61-4.53 (m, 2H), 3.73 (s, 2H), 3.05 (s, 3H), 2.41 (s, 3H), 2.35-2.31 (m, 1H), 1.88-1.81 (m, 2H).

Example 46: Preparation of (S)-5-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one



Step 1: Preparation of (S)-5-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-

yl)-1-(4-methoxybenzyl)-3-methyl-2,3-
dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

97 mg of (S)-5-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 5 of Example 34 except that (S)-8-chloro-2-phenyl-3-(pyrrolidine-2-yl)isoquinoline-1(2H)-one was used
10 (0.16 mmol, yield: 99%).

¹H NMR (300 MHz, CDCl₃) δ 8.16 (s, 1H), 7.70-7.47 (m, 6H), 7.38 (s, 2H), 7.33-7.30 (m, 2H), 7.19 (d, J = 8.6 Hz, 2H), 6.86 (d, J = 8.6 Hz, 2H), 6.72 (s, 2H), 5.03-4.91 (m, 2H), 4.72-4.53 (m, 2H), 4.25-4.12 (m, 2H), 3.79 (s, 3H), 3.12-3.04 (m, 1H), 2.96 (s, 3H),
15 2.05-1.95 (m, 2H), 1.87-1.79 (m, 1H).


Step 2: Preparation of (S)-5-(2-(8-chloro-1-oxo-
2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-
20 yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-
4(1H)-one

58 mg of (S)-5-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was
25 prepared as a white solid according to the same manner

as described in step 6 of Example 34 except that (S)-5-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was used (0.12 mmol, yield: 99%).

¹H NMR (300 MHz, CDCl₃) δ 8.05 (s, 1H), 7.65-7.46 (m, 4H), 7.37-7.29 (m, 4H), 6.65 (s, 1H), 6.03 (s, 1H), 5.01-4.98 (m, 1H), 4.77-4.73 (m, 1H), 4.48-4.44 (m, 1H), 4.20-4.10 (m, 1H), 3.10 (s, 3H), 2.05-1.96 (m, 2H), 83-1.60 (m, 2H).

Example 47: Preparation of (S)-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one



Step 1: Preparation of (S)-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

35 mg of (S)-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid by using 20 mg (0.064 mmol) of 5-chloro-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one prepared in step 4 of Example 34 and 20 mg (0.064 mmol) of (S)-5-chloro-3-phenyl-2-(pyrrolidine-2-yl)pyrrolo[2,1-f][1,2,4]triazine-4(3H)-one according to the same manner as described in step 5 of Example 34 (0.059 mmol, yield: 93%).

¹H NMR (500 MHz, CDCl₃) δ 8.14 (s, 1H), 7.74 (d, J = 7.7 Hz, 1H), 7.49-7.61 (m, 3H), 7.17-7.28 (m, 4H), 6.87 (d, J = 8.5 Hz, 2H), 6.44 (d, J = 2.8 Hz, 1H), 4.73-4.91 (m, 3H), 4.35-4.44 (m, 2H), 3.82 (s, 3H), 3.72 (brs, 1H), 3.58 (brs, 1H), 2.90 (s, 3H), 2.26 (brs, 1H), 2.09 (s, 1H), 2.02-2.08 (m, 1H), 1.79-1.87 (m, 1H).


20

Step 2: Preparation of (S)-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

11 mg of (S)-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a yellow solid by using 35
 5 mg (0.059 mmol) of (S)-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one according to the same manner as described in step 6 of Example
 10 34 (0.023 mmol, yield: 39%).

¹H NMR (500 MHz, CDCl₃) δ 8.03 (s, 1H), 7.70 (d, J = 7.2 Hz, 1H), 7.48-7.60 (m, 3H), 7.25-7.27 (m, 1H), 7.20 (d, J = 2.6 Hz, 1H), 6.43 (d, J = 3.0 Hz, 1H), 6.16 (brs, 1H), 4.76-4.81 (m, 1H), 4.58-4.67 (m, 2H),
 15 3.64-3.76 (m, 2H), 3.09 (s, 3H), 2.27 (brs, 1H), 2.01-2.11 (m, 2H), 1.83-1.90 (m, 2H).

Example 48: Preparation of (S)-7-amino-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one



Step 1: Preparation of (S)-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-7-(methylthio)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

5 65 mg of (S)-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-7-(methylthio)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid by using 40 mg (0.11 mmol, 10 1.0 equivalent) of 5-chloro-1-(4-methoxybenzyl)-3-methyl-7-(methylthio)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one and 40 mg (0.13 mmol, 1.2 equivalent) of (S)-2-(1-aminoethyl)-5-chloro-3-phenylquinazoline-4(3H)-one according to the same 15 manner as described in step 5 of Example 34 (0.10 mmol, yield: 94%).

¹H NMR (300 MHz, CDCl₃) δ 9.44-9.47 (d, J = 4.5 Hz, 1H), 7.69-7.72 (d, J = 8.0 Hz, 1H), 7.55-7.61 (m, 5H), 7.46 (s, 1H), 7.27-7.31 (m, 1H), 7.19-7.22 (d, J = 4.5 Hz, 2H), 6.84-6.87 (d, J = 8.6 Hz, 2H), 5.07-5.12 (m, 1H), 4.74 (s, 2H), 4.41 (s, 2H), 3.79 (s, 3H), 2.89 (s, 3H), 2.29 (s, 3H), 1.39-1.42 (d, J = 6.7 Hz, 3H).

Step 2: Preparation of (S)-7-amino-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

5 52 mg of (S)-7-amino-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid by using 65 mg (0.10 mmol) of (S)-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-7-(methylthio)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one prepared in step 1 according to the same manner as described in step 4 of Example 15 (0.09 mmol, yield: 73%).

15 ¹H NMR (300 MHz, CDCl₃) δ 9.46-9.48 (d, J = 7.4 Hz, 1H), 7.70-7.72 (d, J = 8.2 Hz, 1H), 7.42-7.61 (m, 5H), 7.29-7.31 (d, J = 7.8 Hz, 1H), 7.18-7.21 (d, J = 8.1 Hz, 2H), 6.83-6.86 (d, J = 7.8 Hz, 2H), 5.05-5.10 (m, 1H), 4.66 (s, 2H), 4.61 (s, 2H), 4.33 (s, 3H),
20 3.79 (s, 3H), 2.86 (s, 3H), 1.40-1.42 (d, J = 6.3 Hz, 3H).


Step 3: Preparation of (S)-7-amino-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-

yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

25 mg of (S)-7-amino-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid by using 52 mg (0.09 mmol) of (S)-7-amino-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one prepared in step 2 according to the same manner as described in step 8 of Example 1 (0.05 mmol, yield: 60%).

¹H NMR (300 MHz, CDCl₃) δ 9.41-9.43 (d, J = 7.9 Hz, 1H), 7.70-7.73 (d, J = 7.9 Hz, 1H), 7.44-7.62 (m, 5H), 7.29-7.36 (m, 2H), 5.01-5.06 (m, 1H), 4.74 (s, 2H), 4.60 (s, 2H), 2.94 (s, 3H), 1.40-1.42 (d, J = 6.6 Hz, 3H).

Example 49: Preparation of (S)-7-amino-5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one



Step 1: Preparation of (S)-5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-7-(methylthio)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

77 mg of (S)-5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-7-(methylthio)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 1 of Example 48 except that (S)-2-(1-aminoethyl)-5-chloro-3-(pyridine-3-yl)quinazoline-4(3H)-one was used (0.12 mmol, yield: 90%).

^1H NMR (300 MHz, CDCl_3) δ 9.32-9.34 (m, 1H), 8.72-8.74 (m, 1H), 7.98-8.01 (d, $J = 4.5$ Hz, 1H), 7.59-7.68 (m, 3H), 7.46-7.52 (m, 2H), 7.19-7.22 (d, $J = 7.4$ Hz, 2H), 6.84-6.87 (d, $J = 7.9$ Hz, 2H), 4.92-4.98 (m, 1H), 4.74 (s, 2H), 4.42 (s, 2H), 3.79 (s, 3H), 2.88 (s, 3H), 2.35 (s, 3H), 1.45-1.48 (d, $J = 6.4$ Hz, 3H).

Step 2: Preparation of (S)-7-amino-5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

5 47 mg of (S)-7-amino-5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 2 of Example 48 except that (S)-
10 5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-7-(methylthio)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was used
15 (0.08 mmol, yield: 64%).

¹H NMR (300 MHz, CDCl₃) δ 9.40-9.43 (d, J = 8.8 Hz, 1H), 8.75-8.77 (m, 1H), 7.58-7.72 (m, 3H), 6.45-7.55 (m, 2H), 7.18-7.21 (d, J = 2.6 Hz, 2H), 6.83-6.87 (t, J = 6.8 Hz, 3H), 4.93-5.15 (m, 2H), 4.61-4.73 (m,
20 2H), 4.33 (s, 2H), 3.79 (s, 3H), 2.86 (s, 3H), 1.41-1.49 (dd, J = 17.4, 6.5 Hz, 3H).


Step 3: Preparation of (S)-7-amino-5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-

2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

19 mg of (S)-7-amino-5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 8 of Example 1 except that (S)-7-amino-5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was used (0.04 mmol, yield: 51%).

¹H NMR (300 MHz, DMSO-d₆O) δ 9.19-9.27 (m, 1H), 8.70-8.77 (m, 1H), 8.05-8.10 (m, 1H), 7.75-7.80 (m, 1H), 7.57-7.67 (m, 3H), 7.20-7.21 (d, J = 5.0 Hz, 1H), 6.14 (s, 2H), 4.61-4.65 (m, 1H), 4.44 (s, 2H), 2.78 (s, 3H), 1.28-1.31 (d, J = 6.4 Hz, 3H).

Example 50: Preparation of (S)-7-amino-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one



Step 1: Preparation of (S)-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-7-(methylthio)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

75 mg of (S)-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-7-(methylthio)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 1 of Example 48 except that (S)-2-(1-aminoethyl)-5-chloro-3-(3-fluorophenyl)quinazoline-4(3H)-one was used (0.12 mmol, yield: 94%).

¹H NMR (300 MHz, CDCl₃) δ 9.43-9.30 (m, 1H), 7.68-7.70 (d, J = 7.2 Hz, 1H), 7.44-7.62 (m, 4H), 7.30-7.38 (m, 1H), 7.19-7.21 (d, J = 8.0 Hz, 2H), 7.03-7.10 (m, 1H), 6.84-6.87 (d, J = 7.8 Hz, 2H), 5.06-5.13 (m, 1H), 4.74 (s, 2H), 4.41 (s, 3H), 3.79 (s, 3H), 2.88 (s, 3H), 2.32 (s, 3H), 1.43-1.45 (d, J = 6.1 Hz, 3H).

Step 2: Preparation of (S)-7-amino-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

56 mg of (S)-7-amino-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 2 of Example 48 except that (S)-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-7-(methylthio)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was used (0.09 mmol, yield: 78%).

¹H NMR (300 MHz, CDCl₃) δ 9.39-9.46 (m, 1H), 7.70-7.72 (d, J = 7.3 Hz, 1H), 7.58-7.62 (m, 2H), 7.44-7.47 (m, 2H), 7.07-7.21 (m, 4H), 6.84-6.86 (d, J = 8.5 Hz, 2H), 5.07-5.14 (m, 1H), 4.63-4.76 (m, 4H), 4.34 (s, 2H), 3.79 (s, 3H), 2.86 (s, 3H), 1.42-1.44 (d, J = 3.0 Hz, 3H).


Step 3: Preparation of (S)-7-amino-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-

dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-
dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

26 mg of (S)-7-amino-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 8 of Example 1 except that (S)-7-amino-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was used (0.05 mmol, yield: 58%).

¹H NMR (300 MHz, CDCl₃) δ 9.30-9.37 (t, J = 9.1 Hz, 1H), 7.69-7.72 (m, 1H), 7.40-7.62 (m, 3H), 7.04-7.23 (m, 3H), 5.50 (s, 1H), 5.00-5.11 (m, 1H), 4.62-4.75 (d, J = 22.7 Hz, 2H), 4.60 (s, 2H), 2.94 (s, 3H), 1.40-1.45 (t, J = 7.0 Hz, 3H).

Example 51: Preparation of (S)-7-amino-5-((1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one



Step 1: Preparation of (S)-5-((1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-7-(methylthio)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

80 mg of (S)-5-((1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-7-(methylthio)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 1 of Example 48 except that (S)-2-(1-aminoethyl)-5-chloro-3-m-tolylquinazolin-4(3H)-one was used (0.12 mmol, yield: 88%).

¹H NMR (300 MHz, CDCl₃) δ 9.38-9.46 (m, 1H), 7.69-7.71 (m, 1H), 7.55-7.60 (t, J = 7.5 Hz, 1H), 7.42-7.46 (m, 2H), 7.27-7.31 (m, 2H), 7.19-7.22 (d, J = 7.9 Hz, 2H), 7.06-7.09 (m, 1H), 6.84-6.86 (d, J = 7.8 Hz, 2H), 5.10-5.16 (m, 1H), 4.74 (s, 2H), 4.40 (s, 2H), 3.79 (s, 3H), 2.89 (s, 3H), 2.39-2.43 (d, J = 11.3 Hz, 3H), 2.29 (s, 3H), 1.65 (s, 4H), 1.41-1.44 (d, J = 7.9 Hz, 3H).

Step 2: Preparation of (S)-7-amino-5-((1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

5 49 mg of (S)-7-amino-5-((1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 2 of
10 Example 48 except that (S)-5-((1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-7-(methylthio)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was used (0.12 mmol, yield: 64%).

15 ¹H NMR (300 MHz, CDCl₃) δ 9.42-9.48 (m, 1H), 7.69-7.72 (m, 1H), 7.58-7.60 (m, 1H), 7.42-7.46 (m, 2H), 7.27-7.32 (m, 2H), 7.18-7.20 (m, 2H), 7.09-7.11 (m, 1H), 6.84-6.87 (m, 2H), 5.08-5.12 (m, 1H), 4.62-4.67 (m, 3H), 4.34 (s, 3H), 3.79 (s, 3H), 2.86 (s, 3H),
20 2.34-2.44 (d, J = 30.0 Hz, 3H), 1.42-1.44 (s, J = 3.0 Hz, 3H).

Step 3: Preparation of (S)-7-amino-5-((1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-


yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

22 mg of (S)-7-amino-5-((1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 8 of Example 1 except that (S)-7-amino-5-((1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was used (0.04 mmol, yield: 56%).

¹H NMR (300 MHz, CDCl₃) δ 9.34-9.44 (dd, J = 20.1, 8.4 Hz, 1H), 7.70-7.72 (d, J = 7.2 Hz, 1H), 7.56-7.62 (t, J = 7.9 Hz, 1H), 7.37-7.43 (m, 2H), 7.29-7.31 (m, 1H), 7.08-7.16 (m, 2H), 5.77 (s, 3H), 5.04-5.16 (m, 1H), 4.72-4.75 (d, J = 10.0 Hz, 1H), 4.60 (s, 2H), 2.94 (s, 3H), 2.35-2.44 (d, J = 28.7 Hz, 3H), 1.40-1.43 (dd, J = 6.3, 3.3 Hz, 3H).

20

Example 52 Preparation of (S)-7-amino-5-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one



Step 1: Preparation of (S)-5-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-7-(methylthio)-2,3-

5 dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

66 mg of (S)-5-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-7-(methylthio)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 1 of Example 48 except that (S)-2-(1-aminoethyl)-5-chloro-3-(S)-3-(1-aminoethyl)-8-chloro-2-phenylisoquinoline-1(2H)-one was used (0.10 mmol, yield: 100%).

15 ^1H NMR (300 MHz, CDCl_3) δ 9.17-9.19 (d, J = 4.6 Hz, 1H), 7.59-7.62 (m, 1H), 7.40-7.50 (m, 6H), 7.30-7.32 (m, 1H), 7.21-7.23 (d, J = 5.7 Hz, 2H), 6.85-6.87 (d, J = 5.1 Hz, 2H), 6.56 (s, 1H), 4.85-4.91 (m, 1H), 4.70-4.81 (m, 2H), 4.44 (s, 2H), 3.80 (s, 3H), 2.89
20 (s, 3H), 2.37 (s, 3H), 1.35-1.36 (d, J = 3.8 Hz, 3H).

Step 2: Preparation of (S)-7-amino-5-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

5 51 mg of (S)-7-amino-5-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 2 of
10 Example 48 except that (S)-5-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-7-(methylthio)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was used (0.09 mmol, yield: 83%).

15 ¹H NMR (300 MHz, CDCl₃) δ 9.22-9.24 (d, J = 7.8 Hz, 1H), 7.31-7.53 (m, 8H), 7.19-7.22 (d, J = 7.8 Hz, 2H), 6.84-6.87 (d, J = 8.3 Hz, 2H), 6.58 (s, 1H), 4.79-4.83 (m, 1H), 4.67 (s, 2H), 4.36 (s, 2H), 3.79 (s, 3H), 2.87 (s, 3H), 1.32-1.34 (d, J = 6.5 Hz, 3H).


20

Step 3: Preparation of (S)-7-amino-5-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

26 mg of (S)-7-amino-5-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 8 of Example 1 except that (S)-7-amino-5-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was used (0.05 mmol, yield: 64%).

^1H NMR (300 MHz, DMSO- d_6) δ 9.07-9.10 (d, J = 6.7 Hz, 1H), 7.39-7.61 (m, 8H), 7.22 (s, 1H), 6.58 (s, 1H), 6.18 (s, 2H), 4.47-4.53 (m, 3H), 2.80 (s, 3H), 1.20-1.22 (d, J = 3.0 Hz, 3H).

Example 53: Preparation of (S)-7-amino-3-methyl-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one



Step 1: Preparation of (S)-1-(4-methoxybenzyl)-3-methyl-7-(methylthio)-5-((1-(2-phenylquinoline-3-

yl)ethyl)amino)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

60 mg of (S)-1-(4-methoxybenzyl)-3-methyl-7-(methylthio)-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 1 of Example 48 except that (S)-1-(2-phenylquinoline-3-yl)ethaneamine was used (0.10 mmol, yield: 100%).

¹H NMR (300 MHz, CDCl₃) δ 9.44-9.46 (d, J = 6.7 Hz, 1H), 8.24 (s, 1H), 8.81-8.84 (d, J = 8.1 Hz, 1H), 7.75-8.77 (d, J = 7.9 Hz, 2H), 7.64-8.69 (t, J = 6.8 Hz, 1H), 7.44-7.53 (m, 4H), 7.20-7.22 (d, J = 8.0 Hz, 2H), 6.83-6.86 (d, J = 8.6 Hz, 2H), 5.67-5.71 (m, 1H), 4.67-4.80 (q, J = 15.3, 10.8 Hz, 2H), 4.42 (s, 2H), 3.79 (s, 3H), 2.91 (s, 3H), 2.29 (s, 3H), 1.38-1.40 (d, J = 6.7 Hz, 3H).

Step 2: Preparation of (S)-7-amino-1-(4-methoxybenzyl)-3-methyl-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

47 mg of (S)-7-amino-1-(4-methoxybenzyl)-3-methyl-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was

prepared as a white solid according to the same manner as described in step 2 of Example 48 except that (S)-1-(4-methoxybenzyl)-3-methyl-7-(methylthio)-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)-2,3-

5 dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was used (0.09 mmol, yield: 83%).

¹H NMR (300 MHz, CDCl₃) δ 9.44-9.46 (d, J = 8.4 Hz, 1H), 8.27 (s, 1H), 8.11-8.14 (d, J = 7.9 Hz, 1H), 7.82-7.84 (d, J = 7.8 Hz, 1H), 7.75-7.78 (d, J = 7.2
10 Hz, 2H), 7.63-7.68 (t, J = 5.7 Hz, 1H), 7.47-7.54 (m, 4H), 7.18-7.20 (d, J = 7.4 Hz, 2H), 6.82-6.85 (d, J = 8.1 Hz, 2H), 5.56-5.62 (m, 1H), 4.57-4.72 (m, 4H), 4.34 (s, 2H), 3.78 (s, 3H), 2.88 (s, 3H), 1.42-1.44 (d, J = 6.2 Hz, 3H).

15


Step 3: Preparation of (S)-7-amino-3-methyl-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

18 mg of (S)-7-amino-3-methyl-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)-2,3-
20 dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 8 of Example 1 except that (S)-7-amino-1-(4-methoxybenzyl)-3-methyl-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)-2,3-
25

dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was used (0.04 mmol, yield: 49%).

¹H NMR (300 MHz, DMSO-d₆) δ 9.27-9.29 (d, J = 6.3 Hz, 1H), 8.30 (s, 1H), 7.97-7.80 (d, J = 7.8 Hz, 2H), 7.69-7.75 (m, 2H), 7.50-7.60 (m, 4H), 7.20 (s, 1H), 6.06 (s, 2H), 5.40-5.44 (m, 1H), 4.47 (s, 2H), 2.80 (s, 3H), 1.25-1.27 (d, J = 6.7 Hz, 1H).

Example 54: Preparation of (S)-7-amino-5-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one



Step 1: Preparation of (S)-5-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)-1-(4-methoxybenzyl)-3-methyl-7-(methylthio)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

64 mg of (S)-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 1 of

Example 48 except that (S)-2-(1-aminopropyl)-5-fluoro-3-phenylquinazoline-4(3H)-one was used (0.10 mmol, yield: 100%).

¹H NMR (300 MHz, CDCl₃) δ 9.43-9.46 (d, J = 8.5 Hz, 1H), 7.47-7.69 (m, 6H), 7.27-7.31 (m, 1H), 7.19-7.22 (d, J = 8.3 Hz, 2H), 7.05-7.11 (m, 1H), 6.84-6.87 (d, J = 8.2 Hz, 2H), 5.01-5.08 (m, 1H), 4.73 (s, 2H), 4.41 (s, 2H), 3.79 (s, 3H), 2.89 (s, 3H), 2.26 (s, 3H), 1.73-1.81 (m, 2H), 0.80-0.85 (t, J = 7.2 Hz, 3H).

10

Step 2: Preparation of (S)-7-amino-5-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

15 43 mg of (S)-7-amino-5-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 2 of
20 Example 48 except that (S)-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was used (0.07 mmol, yield: 71%).

¹H NMR (300 MHz, CDCl₃) δ 9.39-9.41 (d, J = 8.5 Hz, 1H), 7.52-7.64 (m, 5H), 7.37-7.39 (m, 1H), 7.28-7.31 (m, 1H), 7.19-7.21 (d, J = 7.5 Hz, 2H), 7.05-7.11 (t, J = 8.5 Hz, 1H), 6.84-6.86 (d, J = 7.0 Hz, 2),
5 4.99-5.01 (m, 1H), 4.66 (s, 2H), 4.57 (s, 2H), 4.33 (s, 2H), 3.79 (s, 3H), 2.86 (s, 3H), 1.77-1.79 (m, 2H), 0.82-0.87 (t, J = 7.5 Hz, 3H).


Step 3: Preparation of (S)-7-amino-5-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

16 mg of (S)-7-amino-5-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one
15 was prepared as a white solid according to the same manner as described in step 8 of Example 1 except that (S)-7-amino-5-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was used (0.03 mmol, yield: 47%).

¹H NMR (300 MHz, DMSO-d₆) δ 9.22-9.25 (d, J = 5.4 Hz, 1H), 7.79-7.81 (m, 1H), 7.45-7.61 (m, 5H), 7.24-7.30 (t, J = 4.5 Hz, 1H), 7.18 (s, 1H), 6.06 (s, 2H),
25

4.54-4.61 (m, 1H), 4.45 (s, 2H), 2.79 (s, 3H), 1.49-1.53 (m, 2H), 0.65-0.70 (t, J = 7.9 Hz, 3H).

Example 55: Preparation of (S)-7-amino-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one



Step 1: Preparation of (S)-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-7-(methylthio)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

82 mg of (S)-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-7-(methylthio)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 1 of Example 48 except that (S)-5-chloro-3-phenyl-2-(pyrrolidine-2-yl)quinazoline-4(3H)-one was used (0.13 mmol, yield: 88%).

^1H NMR (300 MHz, CDCl_3) δ 8.04-8.07 (d, J = 4.5 Hz, 1H), 7.62-7.65 (m, 2H), 7.49-7.54 (m, 3H), 7.39-

7.41 (m, 1H), 7.15-7.22 (m, 3H), 6.83-6.86 (d, J = 8.5 Hz, 2H), 4.77-4.80 (d, J = 7.7 Hz, 2H), 4.66-4.72 (m, 1H), 4.30-4.40 (q, J = 6.0, 9.2 Hz, 2H), 3.91-3.97 (m, 1H), 3.79 (s, 3H), 3.42-3.47 (m, 1H), 2.90 (s, 3H),
5 2.38 (s, H), 2.15-2.23 (m, 2H), 1.63-1.75 (m, 2H).

Step 2: Preparation of (S)-7-amino-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

10

55 mg of (S)-7-amino-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid
15 according to the same manner as described in step 2 of Example 48 except that (S)-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-7-(methylthio)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was used
20 (0.09 mmol, yield: 71%).

¹H NMR (300 MHz, CDCl₃) δ 7.48-7.64 (m, 6H), 7.37-7.39 (d, J = 7.6 Hz, 1H), 7.20-7.23 (d, J = 6.9 Hz, 1H), 7.11-7.14 (d, J = 7.7 Hz, 2H), 6.81-6.84 (d, J = 7.9 Hz, 2H), 4.66-4.70 (m, 2H), 4.23-4.29 (m, 2H),
25 3.84-3.90 (m, 1H), 3.78 (s, 3H), 3.50-3.53 (m, 1H),

2.85 (s, 3H), 2.06-2.13 (m, 1H), 1.98-2.01 (m, 2H),
1.68-1.71 (m, 1H).


Step 3: Preparation of (S)-7-amino-5-(2-(5-
5 chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-
yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-
d]pyrimidine-4(1H)-one

33 mg of (S)-7-amino-5-(2-(5-chloro-4-oxo-3-
phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-
10 3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-
one was prepared as a white solid according to the
same manner as described in step 8 of Example 1 except
that (S)-7-amino-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-
dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-
15 methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-
d]pyrimidine-4(1H)-one was used (0.07 mmol, yield:
74%).

¹H NMR (300 MHz, DMSO-d₆) δ 7.87 (s, 1H), 7.65-
7.70 (m, 1H), 7.40-7.54 (m, 7H), 5.94 (s, 2H), 4.79
20 (s, 1H), 4.35 (s, 2H), 3.57 (s, 1H), 2.86 (s, 3H),
1.98-2.04 (m, 2H), 1.78-1.83 (m, 1H), 1.63-1.69 (m,
1H).

Example 56: Preparation of (S)-7-amino-5-(2-(5-
25 chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-

2-yl)pyrrolidine-1-yl)-3-methyl-2,3-
dihydropyrimido[4,5-d]pyrimidine-4(1H)-one



Step 1: Preparation of (S)-5-(2-(5-chloro-4-oxo-

5 3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-
yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-7-
(methylthio)-2,3-dihydropyrimido[4,5-d]pyrimidine-
4(1H)-one

79 mg of (S)-5-(2-(5-chloro-4-oxo-3-(pyridine-3-
 10 yl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-7-(methylthio)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 1 of Example 48 except that (S)-
 15 5-chloro-3-(pyridine-3-yl)-2-(pyrrolidine-2-yl)quinazoline-4(3H)-one was used (0.12 mmol, yield: 91%).

¹H NMR (300 MHz, CDCl₃) δ 8.74-8.76 (d, J = 5.9 Hz, 1H), 8.47-8.50 (m, 1H), 7.58-7.66 (m, 3H), 7.42-
 20 7.44 (d, J = 7.6 Hz, 2H), 7.15-7.18 (d, J = 8.4 Hz, 2H), 6.83-6.86 (d, J = 8.0 Hz, 2H), 4.78-4.79 (d, J = 4.4 Hz, 2H), 4.31-4.37 (m, 2H), 3.85-4.04 (m, 1H),

3.79 (s, 3H), 3.55-3.35 (m, 1H), 2.90 (s, 3H), 2.39 (s, 3H), 2.08-2.13 (m, 2H), 1.60-1.83 (m, 2H).

Step 2: Preparation of (S)-7-amino-5-(2-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

49 mg of (S)-7-amino-5-(2-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 2 of Example 48 except that (S)-5-(2-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-7-(methylthio)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was used (0.08 mmol, yield: 65%).


¹H NMR (300 MHz, CDCl₃) δ 8.75 (s, 7H), 7.61-7.65 (m, 2H), 7.52-7.57 (m, 3H), 7.40-7.43 (d, J = 6.8 Hz, 1H), 7.13-7.17 (m, 2H), 6.81-6.84 (d, J = 8.4 Hz, 2H), 4.81 (s, 2H), 4.61-4.76 (m, 2H), 4.19-4.33 (m, 2H), 3.94-4.05 (m, 1H), 3.78 (s, 3H), 3.47-3.57 (m, 1H), 2.86 (s, 3H), 2.15-2.24 (m, 1H), 1.96-2.09 (m, 2H), 1.72-1.79 (m, 1H).

Step 3: Preparation of (S)-7-amino-5-(2-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

29 mg of (S)-7-amino-5-(2-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 8 of Example 1 except that (S)-7-amino-5-(2-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was used (0.06 mmol, yield: 73%).

¹H NMR (300 MHz, DMSO-d₆O) δ 8.72 (s, 2H), 8.00-8.03 (d, J = 9.0 Hz, 1H), 7.61-7.73 (m, 2H), 7.43-7.52 (m, 2H), 6.04 (s, 1H), 5.31-5.43 (m, 1H) 4.56-4.65 (m, 1H), 4.36 (s, 2H), 3.97-4.05 (m, 1H), 2.86 (s, 3H), 1.95-2.04 (m, 2H), 1.63-1.82 (m, 2H).

Example 57: Preparation of (S)-7-amino-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one



Step 1: Preparation of (S)-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-7-(methylthio)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

84 mg of (S)-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-7-(methylthio)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 1 of Example 48 except that (S)-5-chloro-3-(3-fluorophenyl)-2-(pyrrolidine-2-yl)quinazoline-4(3H)-one was used (0.13 mmol, yield: 88%).

^1H NMR (300 MHz, CDCl_3) δ 7.89-7.91 (d, J = 6.1 Hz, 1H), 7.50-7.66 (m, J = 38.5, 9.8 Hz, 3H), 7.40-7.43 (m, 1H), 7.11-7.24 (m, 3H), 6.91-7.06 (m, 1H), 6.83-6.86 (d, J = 7.4 Hz, 2H), 4.73-4.78 (d, J = 4.9 Hz, 1H), 4.62-4.72 (m, 1H), 4.30-3.41 (m, 2H), 3.86-4.03 (m, 1H), 3.79 (s, 3H), 3.35-3.42 (m, 1H), 2.90

(s, 3H), 2.38 (s, 3H), 2.05-2.22 (m, 3H), 1.72-1.82 (m, 1H).

Step 2: Preparation of (S)-7-amino-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

54 mg of (S)-7-amino-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 2 of Example 48 except that (S)-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-7-(methylthio)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was used (0.08 mmol, yield: 67%).


¹H NMR (300 MHz, CDCl₃) δ 7.64-7.66 (m, 1H), 7.38-7.58 (m, 4H), 6.98-7.15 (m, 4H), 6.81-6.84 (d, J = 8.0 Hz, 2H), 4.62-4.74 (m, 3H), 4.25-4.30 (m, 2H), 3.84-3.96 (m, 1H), 3.78 (s, 3H), 3.48-3.56 (m, 1H), 2.86 (s, 3H), 2.14-2.18 (m, 1H), 2.02-2.05 (m, 2H), 1.72-1.79 (m, 1H).

Step 3: Preparation of (S)-7-amino-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

30 mg of (S)-7-amino-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 8 of Example 1 except that (S)-7-amino-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was used (0.06 mmol, yield: 68%).

¹H NMR (300 MHz, DMSO-d₆) δ 7.85 (s, 1H), 7.63-7.71 (m, 2H), 7.36-7.51 (m, 5H), 5.97-6.05 (ss, 2H), 4.49-4.61 (m, 1H), 4.35 (s, 2H), 3.52-3.59 (m, 1H), 2.86 (s, 3H), 1.98-2.06 (m, 2H), 1.81-1.85 (m, 1H), 1.66-1.70 (m, 1H).

Example 58: Preparation of (S)-7-amino-5-(2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one



Step 1: Preparation of (S)-5-(2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-7-(methylthio)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

76 mg of (S)-5-(2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-7-(methylthio)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 1 of Example 48 except that (S)-5-chloro-2-(pyrrolidine-2-yl)-3-m-tolylquinazoline-4(3H)-one was used (0.11 mmol, yield: 96%).

^1H NMR (300 MHz, CDCl_3) δ 7.83-7.88 (m, 1H), 7.60-7.63 (m, 1H), 7.38-7.58 (m, 3H), 7.29-7.31 (m, 1H), 7.16-7.19 (d, J = 7.4 Hz, 2H), 6.99-7.02 (m, 1H), 6.84-6.86 (d, J = 5.7 Hz, 2H), 4.73-4.81 (m, 3H), 4.35 (s, 2H), 3.85-3.98 (m, 1H), 3.79 (s, 3H), 3.47-3.58 (m, 1H), 2.89 (s, 3H), 2.41-2.47 (d, J = 9.0 Hz, 3H), 2.30-2.41 (m, 3H), 2.15 (s, 3H), 1.70-1.77 (m, 1H).

Step 2: Preparation of (S)-7-amino-5-(2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

5 57 mg of (S)-7-amino-5-(2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 2 of
10 Example 48 except that (S)-5-(2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-7-(methylthio)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was used (0.09 mmol, yield: 79%).

15 ¹H NMR (300 MHz, CDCl₃) δ 7.62-7.65 (m, 1H), 7.41-7.55 (m, 4H), 7.29-7.32 (m, 1H), 7.13-7.16 (d, J = 6.0 Hz, 2H), 7.03 (s, 1H), 6.82-6.85 (d, J = 8.9 Hz, 2H), 4.68-4.76 (m, 3H), 4.59 (s, 2H), 4.27 (s, 2H), 3.85-3.91 (m, 1H), 3.78 (s, 3H), 3.52-3.59 (m, 1H), 2.85
20 (s, 3H), 2.43 (s, 3H), 2.14-2.22 (m, 1H), 2.00-2.04 (m, 2H), 1.72-1.77 (m, 1H).

Step 3: Preparation of (S)-7-amino-5-(2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-


yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

31 mg of (S)-7-amino-5-(2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-
5 3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 8 of Example 1 except that (S)-7-amino-5-(2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was used (0.06 mmol, yield: 67%).

¹H NMR (300 MHz, CDCl₃) δ 7.63-7.65 (m, 1H), 7.51-7.56 (t, J = 7.7 Hz, 1H), 7.39-7.45 (m, 3H), 7.30-7.32
15 (d, J = 7.4 Hz, 1H), 7.01-7.03 (m, 1H), 6.56 (s, 9H), 4.71-4.76 (m, 2H), 4.42-4.58 (m, 1H), 4.94-4.00 (m, 1H), 3.79-3.85 (m, 2H), 3.52-3.60 (m, 1H), 3.02 (s, 3H), 2.42 (s, 3H), 2.12-2.25 (m, 1H), 1.98-2.06 (m, 2H), 1.69-1.76 (m, 1H).

20

Example 59: Preparation of (S)-7-amino-5-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one



Step 1: Preparation of (S)-5-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-7-(methylthio)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

81 mg of (S)-5-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-7-(methylthio)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 1 of Example 48 except that (S)-8-chloro-2-phenyl-3-(pyrrolidine-2-yl)isoquinoline-1(2H)-one was used (0.12 mmol, yield: 88%).

^1H NMR (300 MHz, CDCl_3) δ 7.76-7.78 (d, J = 5.8 Hz, 1H), 7.59-7.64 (t, J = 6.4 Hz, 1H), 7.42-7.52 (m, 2H), 7.36-7.41 (m, 3H), 7.30-7.32 (d, J = 6.4 Hz, 1H), 7.18-7.20 (m, 1H), 6.84-6.86 (d, J = 8.1 Hz, 2H), 6.77 (s, 1H), 5.05-5.09 (t, J = 6.1 Hz, 1H), 4.89-4.94 (d, J = 15.3 Hz, 1H), 4.69-4.74 (d, J = 15.0 Hz, 1H), 4.52-4.56 (d, J = 10.5 Hz, 1H), 4.18-4.22 (d, J = 10.9 Hz, 1H), 4.11-4.16 (m, 1H), 3.79 (s, 3H), 3.07-3.13

(m, 1H), 2.94 (s, 3H), 2.44 (s, 3H), 1.93-1.99 (m, 2H), 1.61-1.81 (m, 2H).

Step 2: Preparation of (S)-7-amino-5-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

38 mg of (S)-7-amino-5-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 2 of Example 48 except that (S)-5-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-7-(methylthio)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was used (0.06 mmol, yield: 49%).

¹H NMR (300 MHz, CDCl₃) δ 7.48-7.55 (m, 4H), 7.31-7.38 (m, 4H), 7.16-7.18 (d, J = 7.9 Hz, 2H), 6.83-6.86 (m, 3H), 5.01-5.07 (m, 1H), 4.84-4.89 (d, J = 15.4 Hz, 1H), 4.69 (s, 2H), 4.48-4.59 (m, 2H), 4.10-4.13 (d, J = 10.3 Hz, 2H), 3.78 (s, 3H), 3.11-3.19 (m, 1H), 2.91 (s, 3H), 1.89-1.93 (m, 2H), 1.73-1.79 (m, 1H), 1.55-1.62 (m, 1H).


Step 3: Preparation of (S)-7-amino-5-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

5 13 mg of (S)-7-amino-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 8 of
10 Example 1 except that (S)-7-amino-5-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)-1-(4-methoxybenzyl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was used (0.03 mmol, yield: 42%).

15 ¹H NMR (300 MHz, DMSO-d₆) δ 7.49-7.63 (m, 7H), 7.23-7.33 (m, 1H), 6.82 (s, 1H), 6.19 (s, 2H), 6.12 (s, 1H), 5.47-5.58 (m, 1H), 4.42 (s, 2H), 3.98-4.04 (m, 1H), 3.06-3.09 (m, 1H), 2.76 (s, 3H), 1.90-1.98 (m, 2H), 1.46-1.57 (m, 2H).

20

Example 60: Preparation of (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-3-ethyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one



Step 1: Preparation of 5-chloro-3-ethyl-1-(4-methoxybenzyl)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

400 mg of 5-chloro-3-ethyl-1-(4-methoxybenzyl)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid by using 600 mg 1.87 mmol) of 4-chloro-N-ethyl-6-(4-methoxybenzylamino)pyrimidine-5-carboxamide according to the same manner as described in step 4 of Example 34 (1.2 mmol, yield: 64%).

¹H NMR (300 MHz, CDCl₃) δ 8.44 (s, 1H), 7.22 (d, J = 8.4 Hz, 2H), 6.89 (d, J = 8.6 Hz, 2H), 4.85 (s, 2H), 4.54 (s, 2H), 3.82 (s, 3H), 3.46 (q, J = 7.2 Hz, 2H), 1.08 (t, J = 7.2 Hz, 3H).

Step 2: Preparation of (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-3-ethyl-1-(4-methoxybenzyl)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

60 mg of (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-3-ethyl-1-(4-methoxybenzyl)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a yellow solid by using 40
5 mg (0.12 mmol) of 5-chloro-3-ethyl-1-(4-methoxybenzyl)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one prepared in step 1 according to the same manner as described in step 5 of Example 34 (0.1 mmol, yield: 84%).

10 ¹H NMR (300 MHz, CDCl₃) δ 9.15 (s, -NH), 7.99 (s, 1H), 7.51-7.30 (m, 8H), 7.23 (d, J = 8.7 Hz, 2H), 6.87 (d, J = 8.1 Hz, 2H), 6.58 (s, 1H), 4.86 (s, 1H), 4.75 (s, 2H), 4.47 (s, 2H), 3.80 (s, 3H), 3.44-3.39 (m, 2H), 1.39 (d, J = 6.8 Hz, 3H), 1.11 (t, J = 7.2 Hz,
15 3H).

Step 3: Preparation of (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-3-ethyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one


20 47 mg of (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-3-ethyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid by using 60 mg (0.1 mmol) of (S)-5-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydro-
25 isoquinoline-3-yl)ethyl)amino)-3-ethyl-1-(4-

methoxybenzyl)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one prepared in step 2 according to the same manner as described in step 8 of Example 1 (0.1 mmol, yield: 99%).

¹H NMR (300 MHz, CDCl₃) δ 9.06 (d, J = 6.9 Hz, -NH), 7.94 (s, 1H), 7.49-7.39 (m, 8H), 6.56 (s, 1H), 5.80 (s, 1H), 4.87-4.82 (m, 1H), 4.74 (s, 2H), 3.52-3.45 (m, 2H), 1.39 (d, J = 6.7 Hz, 3H), 1.23 (t, J = 7.2 Hz, 3H).

10

Example 61: Preparation of (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-3-propyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one



15

Step 1: Preparation of 5-chloro-1-(4-methoxybenzyl)-3-propyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

289 mg of 5-chloro-1-(4-methoxybenzyl)-3-propyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid by using 653 mg (1.95 mmol) of 4-chloro-6-(4-methoxybenzylamino)-N-

propylpyrimidine-5-carboxamide according to the same manner as described in step 4 of Example 34 (1.12 mmol, yield: 57%).

¹H NMR (300 MHz, CDCl₃) δ 8.44 (s, 1H), 7.22 (d, J = 8.5 Hz, 2H), 6.89 (d, J = 8.5 Hz, 2H), 4.85 (s, 2H), 4.52 (s, 2H), 3.82 (s, 3H), 3.37 (t, J = 7.4 Hz, 2H), 1.52-1.45 (m, 2H), 0.86 (t, J = 7.4 Hz, 3H).

Step 2: Preparation of (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-1-(4-methoxybenzyl)-3-propyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

56 mg of (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-1-(4-methoxybenzyl)-3-propyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a yellow solid by using 41 mg (0.12 mmol) of 5-chloro-1-(4-methoxybenzyl)-3-propyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one prepared in step 1 according to the same manner as described in step 5 of Example 34 (0.09 mmol, yield: 73%).

¹H NMR (300 MHz, CDCl₃) δ 9.17-9.15 (m, -NH), 7.99 (s, 1H), 7.51-7.29 (m, 8H), 7.21-7.07 (m, 1H), 6.87 (d, J = 8.0 Hz, 2H), 6.52 (s, 1H), 4.96-4.87 (s, 1H), 4.75 (s, 2H), 4.47 (s, 2H), 3.80 (s, 3H), 3.44-3.39

(m, 2H), 1.68-1.63 (m, 2H), 1.39 (d, J = 6.9 Hz, 3H), 1.01-0.96 (m, 3H).


Step 3: Preparation of (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-3-propyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

45 mg of (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-3-propyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid by using 56 mg (0.09 mmol) of (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-1-(4-methoxybenzyl)-3-propyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one prepared in step 2 according to the same manner as described in step 8 of Example 1 (0.09 mmol, yield: 99%).

¹H NMR (300 MHz, CDCl₃) δ 9.04 (s, -NH), 7.95 (s, 1H), 7.50-7.31 (m, 8H), 6.57 (s, 1H), 5.28 (s, 1H), 4.91-4.84 (m, 1H), 4.73 (s, 2H), 3.41-3.36 (m, 2H), 1.68-1.63 (m, 2H), 1.39 (d, J = 6.9 Hz, 3H), 1.01-0.96 (m, 3H).

Example 62: Preparation of (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-

3-cyclopropyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one



Step 1: Preparation of 5-chloro-3-cyclopropyl-1-(4-methoxybenzyl)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

224 mg of 5-chloro-3-cyclopropyl-1-(4-methoxybenzyl)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid by using 360 mg (1.08 mmol) of 4-chloro-N-cyclopropyl-6-(4-methoxybenzylamino)pyrimidine-5-carboxamide according to the same manner as described in step 4 of Example 34 (0.65 mmol, yield: 60%).

¹H NMR (300 MHz, CDCl₃) δ 8.43 (s, 1H), 7.21 (d, J = 7.8 Hz, 2H), 6.90 (d, J = 8.5 Hz, 2H), 4.84 (s, 2H), 4.52 (s, 2H), 3.82 (s, 3H), 2.60-2.56 (m, 1H), 0.84-0.82 (m, 2H), 0.52-0.50 (m, 2H).

Step 2: Preparation of (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-3-

cyclopropyl-1-(4-methoxybenzyl)-2,3-
dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

43 mg of (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-3-cyclopropyl-1-(4-methoxybenzyl)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid by using 40 mg (0.12 mmol) of 5-chloro-3-cyclopropyl-1-(4-methoxybenzyl)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one prepared in step 1 according to the same manner as described in step 5 of Example 34 (0.07 mmol, yield: 59%).

¹H NMR (300 MHz, CDCl₃) δ 9.19-9.15 (m, -NH), 8.05 (s, 1H), 7.53-7.32 (m, 4H), 7.23-7.18 (m, 1H), 6.87 (d, J = 9.4 Hz, 2H), 6.56 (s, 1H), 4.84 (s, 1H), 4.43 (s, 1H), 3.79 (s, 2H), 2.03 (s, 1H), 1.38 (d, J = 6.2 Hz, 3H), 0.86-0.80 (m, 2H), 0.56-0.52 (m, 2H).


Step 3: Preparation of (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-3-cyclopropyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

34 mg of (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-3-cyclopropyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid by using 43 mg (0.07 mmol)

of (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-3-cyclopropyl-1-(4-methoxybenzyl)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one prepared in step 2 according to the same manner as described in step 8 of Example 1 (0.07 mmol, yield: 99%).

¹H NMR (300 MHz, CDCl₃) δ 9.09 (d, J = 6.8 Hz, 1H), 7.94 (s, 1H), 7.51-7.32 (m, 8H), 6.56 (s, 1H), 5.66 (s, 1H), 4.88-4.84 (m, 1H), 4.71 (s, 2H), 2.59-2.53 (m, 1H), 1.39 (d, J = 6.7 Hz, 3H), 0.95-0.92 (m, 2H), 0.75-0.73 (m, 2H).

Example 63: Preparation of (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-3-cyclopentyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one



Step 1: Preparation of 5-chloro-3-cyclopentyl-1-(4-methoxybenzyl)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

405 mg of 5-chloro-3-cyclopentyl-1-(4-methoxybenzyl)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid by using 620 mg (1.72 mmol) of 4-chloro-N-cyclopentyl-6-(4-methoxybenzylamino)pyrimidine-5-carboxamide according to the same manner as described in step 4 of Example 34 (1.08 mmol, yield: 63%).

¹H NMR (300 MHz, CDCl₃) δ 8.44 (s, 1H), 7.22 (d, J = 8.6 Hz, 2H), 6.90 (d, J = 8.7 Hz, 2H), 5.31 (s, 2H), 4.97-4.88 (m, 1H), 4.83 (s, 2H), 4.41 (s, 2H), 3.82 (s, 3H), 1.86-1.77 (m, 2H), 1.57-1.53 (m, 2H), 1.26-1.16 (m, 2H).

Step 2: Preparation of (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-3-cyclopentyl-1-(4-methoxybenzyl)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

45 mg of (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-3-cyclopentyl-1-(4-methoxybenzyl)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a yellow solid by using 40 mg (0.1 mmol) of 5-chloro-3-cyclopentyl-1-(4-methoxybenzyl)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one prepared in step 1 according to

the same manner as described in step 5 of Example 34 (0.07 mmol, yield: 70%).

¹H NMR (300 MHz, CDCl₃) δ 9.20 (d, J = 6.8 Hz, 1H), 8.06 (s, 1H), 7.56-7.30 (m, 8H), 7.23 (d, J = 8.4 Hz, 2H), 6.87 (d, J = 8.4 Hz, 2H), 6.59 (s, 1H), 4.91-4.82 (m, 2H), 4.74 (s, 2H), 4.36 (d, J = 2.3 Hz, 1H), 1.86-1.74 (m, 2H), 1.38 (d, J = 6.8 Hz, 3H).


Step 3: Preparation of (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-3-cyclopentyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

37 mg of (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-3-cyclopentyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid by using 45 mg (0.07 mmol) of (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-3-cyclopentyl-1-(4-methoxybenzyl)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one prepared in step 2 according to the same manner as described in step 8 of Example 1 (0.07 mmol, yield: 99%).

¹H NMR (300 MHz, CDCl₃) δ 9.11 (d, J = 6.7 Hz, 1H), 7.96 (s, 1H), 7.51-7.32 (m, 8H), 6.57 (s, 1H), 5.56 (s, 1H), 4.95-4.65 (m, 2H), 4.65 (s, 2H), 1.99-

1.94 (m, 2H), 1.75-1.65 (m, 4H), 1.57-1.49 (m, 2H),
1.38 (d, J = 6.6 Hz, 3H).

Example 64: Preparation of (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-3-isopropyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one



Step 1: Preparation of (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-3-isopropyl-1-(4-methoxybenzyl)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

Quantitative yield of (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-3-isopropyl-1-(4-methoxybenzyl)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was obtained as a white solid by using 69 mg (0.20 mmol) of 5-chloro-3-isopropyl-1-(4-methoxybenzyl)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one according to the same manner as described in step 5 of Example 34.


¹H NMR (300MHz, CDCl₃) δ 1.02-1.08 (m, 6H), 1.37-1.40 (d, J = 9.0, 3H), 3.80 (s, 3H), 4.32-4.40 (m, 2H), 4.76-4.89 (m, 4H), 6.59 (s, 1H), 6.85-6.88 (m, 2H), 7.21-7.53 (m, 10H), 8.05 (s, 1H), 9.17-9.19 (d, J = 6.0, 1H).

Step 2: Preparation of (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-3-isopropyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

14 mg of (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-3-isopropyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid by using 25 mg 0.041 mmol) of (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-3-isopropyl-1-(4-methoxybenzyl)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one according to the same manner as described in step 8 of Example 1 (yield: 70%).

¹H NMR (300MHz, CDCl₃) δ 1.19-1.25 (m, 6H), 1.37-1.40 (d, J = 9.0, 3H), 4.61-4.68 (m, 2H), 4.80-4.87 (m, 2H), 5.73(br s, 1H), 6.57 (s, 1H), 7.26-7.50 (m, 8H), 7.94 (s, 1H), 9.10-9.13 (d, J = 9.0, 1H).

Example 65: Preparation of (S)-5-(1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propylamino)-3-isopropyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one



5

Step 1: Preparation of (S)-5-(1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propylamino)-3-isopropyl-1-(4-methoxybenzyl)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one

10 97 mg of (S)-5-(1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propylamino)-3-isopropyl-1-(4-methoxybenzyl)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 5 of Example 34

15 except that 69 mg (0.20 mmol) of 5-chloro-3-isopropyl-1-(4-methoxybenzyl)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one and (S)-2-(1-aminopropyl)-5-fluoro-3-phenylquinazoline-4(3H)-one (1.1 equivalent) were used (0.20 mmol, yield: 100%).

¹H NMR (300MHz, CDCl₃) δ 0.84-0.88 (t, J = 6.0, 3H), 1.02-1.04 (d, J = 6.0, 6H), 1.74-1.94 (m, 2H), 3.79 (s, 3H), 4.34 (s, 2H), 4.73 (s, 2H), 4.79-4.86 (m, 1H), 4.96-5.03 (m, 1H), 6.84-6.87 (m, 2H), 7.05-7.30 (m, 4H), 7.45-7.69 (m, 6H), 8.02 (s, 1H), 9.44-9.46 (d, J = 6.0, 1H).

Step 2: Preparation of (S)-5-(1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propylamino)-3-isopropyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one


14 mg of (S)-5-(1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propylamino)-3-isopropyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid by using 30 mg (0.049 mmol, 1.0 eq) of (S)-5-(1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propylamino)-3-isopropyl-1-(4-methoxybenzyl)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one prepared in step 1 according to the same manner as described in step 8 of Example 1 (yield: 75%).

¹H NMR (300MHz, CDCl₃) δ 0.83-0.87 (t, J = 6.0, 3H), 1.19-1.21 (d, J = 6.0, 6H), 1.73-1.95 (m, 2H), 4.61 (s, 2H), 4.84-4.99 (m, 2H), 6.33 (br s, 1H), 7.05-


7.12 (m, 1H), 7.26-7.30 (m, 1H), 7.44-7.69 (m, 6H),
7.90 (s, 1H), 9.39-9.41 (d, J = 6.0, 1H).

The following examples 66 ~ 98 were performed by
5 the method represented by the reaction formula 3A.

[Reaction Formula 3A]



Example 66: Preparation of (S)-5-((1-(5-chloro-4-
10 oxo-3-phenyl-3,4-dihydroquinazoline-2-
yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one



Step 1: Preparation of 4,6-dichloropyrimidine-5-carbonyl chloride

1 g (5.65 mmol) of 4,6-dichloropyrimidine-5-carboxaldehyde was dissolved in 15 mL of CCl_4 , to which 0.78 mL (9.61 mmol) of sulfonyl chloride and 46 mg (0.28 mmol) of 2,2-azobis(2-methyl propionitrile) were added, followed by stirring at 80°C for 3 hours. The reaction mixture was filtered under reduced pressure. 5 mL of anhydrous toluene was added thereto, followed by filtration under reduced pressure. As a result, 4,6-dichloropyrimidine-5-carbonyl chloride was obtained.

Step 2: Preparation of 5-chloropyrimido[4,5-d]pyrimidine-4(1H)-one

4,6-dichloropyrimidine-5-carbonyl chloride (1.0 eq) prepared in step 1 was dissolved in toluene, to which excessive thionyl chloride (SOCl_2) was added, followed by stirring at 115°C for 12 hours. The reaction mixture was cooled down to room temperature.

The reaction solvent was concentrated under reduced pressure and dried to give acid chloride. Formamidine hydrochloride (1.1 equivalent) was dissolved in tetrahydrofuran at 0°C, to which triethylamine (4.0
5 equivalent) was added. The prepared acid chloride was dissolved in 5.0 mL of anhydrous tetrahydrofuran, which was slowly added to the mixture above. The reaction mixture was heated at room temperature, followed by stirring for 4 hours. Water was added
10 thereto, followed by extraction with diethyl ether. The water layer was extracted by using ethylacetate:tetrahydrofuran (1:1). Then, the organic layer was separated, dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was
15 separated by column chromatography (SiO₂, eluent: dichloromethane/methanol, 15/1 -> dichloromethane/methanol, 10/1) to give the target compound 5-chloropyrimido[4,5-d]pyrimidine-4(1H)-one.

20 Step 3: Preparation of (S)-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one


25 25 mg of (S)-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid

by using 15 mg (0.08 mmol, 1.0 eq) of 5-chloropyrimido[4,5-d]pyrimidine-4(1H)-one prepared in step 2 and 30 mg (0.10 mmol, 1.2 equivalent) of (S)-2-(1-aminoethyl)-5-chloro-3-phenylquinazoline-4(3H)-one according to the same manner as described in step 5 of Example 34 (0.06 mmol, yield: 68%).

¹H NMR (300 MHz, CDCl₃) δ 9.77 (s, 1H), 8.82 (s, 1H), 8.67 (s, 1H), 7.49-7.58 (m, 7H), 7.37 (s, 1H), 5.11-5.16 (m, 1H), 1.51-1.53 (d, J = 3.0 Hz, 3H).

10

Example 67: Preparation of (S)-5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one




15 18.8 mg of (S)-5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 3 of Example 66 except that (S)-
20 2-(1-aminoethyl)-5-chloro-3-(pyridine-3-

yl)quinazoline-4(3H)-one was used (0.04 mmol, yield: 77%).

¹H NMR (300 MHz, DMSO-d₆) δ 9.69 (s, 1H), 8.66-8.69 (m, 2H), 8.40 (s, 1H), 8.31 (s, 1H), 8.00-8.11 (m, 1H), 7.74-7.78 (m, 1H), 7.56-7.65 (m, 3H), 4.82-4.86 (m, 1H), 1.34-1.39 (m, 3H).


Example 68: Preparation of (S)-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one



13.8 mg of (S)-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 3 of Example 66 except that (S)-2-(1-aminoethyl)-5-chloro-3-(3-fluorophenyl)quinazoline-4(3H)-one was used (0.03 mmol, yield: 55%).

¹H NMR (300 MHz, CDCl₃) δ 9.53-9.62 (m, 1H), 8.63-8.65 (d, J = 3.0 Hz, 1H), 8.43-8.44 (m, 1H), 7.63-7.69 (m, 3H), 7.58-7.62 (m, 2H), 7.50-7.55 (m, 1H), 7.28-7.29 (m, 1H), 7.12-7.19 (m, 1H), 5.12-5.18 (m, 1H),
 5 1.47-1.56 (m, 3H).

Example 69: Preparation of (S)-5-((1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one



10


10.5 mg of (S)-5-((1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 3 of
 15 Example 66 except that (S)-2-(1-aminoethyl)-5-chloro-3-(m-tolyl)quinazoline-4(3H)-one was used (0.02 mmol, yield: 42%).

¹H NMR (300 MHz, CDCl₃) δ 9.79-9.81 (d, J = 4.5 Hz, 1H), 8.77-8.79 (d, J = 3.0 Hz, 1H), 8.66-8.68 (d,

$J = 3.0$ Hz, 1H), 7.61-7.68 (m, 2H), 7.46-7.51 (m, 3H), 7.35-7.38 (m, 1H), 7.17-7.18 (m, 1H), 5.18-5.22 (m, 1H), 2.41-2.49 (d, $J = 12.0$ Hz, 3H), 1.53-1.55 (m, 3H).

5

Example 70: Preparation of (S)-5-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one



10 Step 1: Preparation of 5-chloro-1-(4-methoxybenzyl)pyrimido[4,5-d]pyrimidine-4(1H)-one

50 mg (0.171 mmol) of 4-chloro-6-((4-methoxybenzyl)amino)pyrimidine-5-carboxamide, 1 mL of triethyl orthoformate, and 10 μ L of methanesulfonic acid were mixed together, which was stirred at 50°C for 2 hours. The reaction mixture was cooled down to room temperature. Ethyl acetate and water were added thereto, followed by extraction. The extracted organic layer was separated, dried (Na_2SO_4), filtered, and concentrated under reduced pressure. The residue was separated by column chromatography (SiO_2 , eluent:

15

20

hexane/ethyl acetate, 4/1 -> hexane/ethyl acetate, 1/1) to give 10 mg of the target compound 5-chloro-1-(4-methoxybenzyl)pyrimido[4,5-d]pyrimidine-4(1H)-one as a pale yellow liquid (0.033 mmol, yield: 19%).

5 ¹H NMR (300 MHz, CDCl₃) δ 8.96 (s, 1H), 8.44 (s, 1H), 7.29 (d, J = 8.9 Hz, 2H), 6.89 (d, J = 8.9 Hz, 2H), 5.37 (s, 2H), 3.80 (s, 3H).

Step 2: Preparation of (S)-5-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-1-(4-methoxybenzyl)pyrimido[4,5-d]pyrimidine-4(1H)-one

18 mg of (S)-5-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-1-(4-methoxybenzyl)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a pale yellow solid by using 10 mg (0.033 mmol) of 5-chloro-1-(4-methoxybenzyl)pyrimido[4,5-d]pyrimidine-4(1H)-one prepared in step 1 and 9 mg (0.036 mmol) of (S)-3-(1-aminoethyl)-8-chloro-2-phenylisoquinoline-1(2H)-one according to the same manner as described in step 5 of Example 34 (0.032 mmol, yield: 96%).

¹H NMR (300 MHz, CDCl₃) δ 9.89 (d, J = 7.1 Hz, 1H), 8.36 (d, J = 9.4 Hz, 2H), 7.38-7.54 (m, 5H), 7.29-7.37 (m, 5H), 6.87 (d, J = 8.6 Hz, 2H), 6.56 (s,


1H), 5.29 (s, 2H), 4.95 (t, J = 6.3 Hz, 1H), 3.77 (s, 3H), 1.44 (d, J = 6.3 Hz, 3H).

Step 3: Preparation of (S)-5-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one

13 mg of (S)-5-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a pale yellow solid in 25 ml round-bottom flask by using 18 mg (0.032 mmol) of (S)-5-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-1-(4-methoxybenzyl)pyrimido[4,5-d]pyrimidine-4(1H)-one prepared in step 1 according to the same manner as described in step 8 of Example 1 (0.029 mmol, yield: 92%).

¹H NMR (300 MHz, CDCl₃) δ 9.24 (d, J = 5.5 Hz, 1H), 8.57 (s, 1H), 7.47-7.52 (m, 5H), 7.32-7.38 (m, 4H), 6.60 (s, 1H), 5.03 (t, J = 7.0 Hz, 1H), 1.50 (d, J = 6.5 Hz, 3H).


Example 71: Preparation of (S)-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one



20 mg of (S)-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid by using 15 mg (0.08 mmol, 1.0 equivalent) of 5-chloropyrimido[4,5-d]pyrimidine-4(1H)-one and 28 mg (0.10 mmol, 1.2 equivalent) of (S)-2-(1-aminoethyl)-5-chloro-3-phenylpyrrolo[2,1-f][1,2,4]triazine-4(3H)-one according to the same manner as described in step 5 of Example 34 (0.05 mmol, yield: 58%).

^1H NMR (300 MHz, CDCl_3) δ 9.44 (s, 1H), 8.81 (s, 1H), 8.64 (s, 1H), 7.49-7.56 (m, 5H), 7.27-7.36 (m, 2H), 7.50 (s, 1H), 5.10-5.14 (m, 1H), 1.51-1.53 (d, J = 3.2 Hz, 3H).


Example 72: Preparation of (S)-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one



6 mg of (S)-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 5 of Example 34 except that (S)-1-(2-phenylquinoline-3-yl)ethanamine was used (0.015 mmol, yield: 28%).

¹H NMR (300 MHz, DMSO-d₆) δ 12.86 (s, 1H), 9.42-9.44 (m, 1H), 8.46 (s, 1H), 8.39 (s, 1H), 8.31 (s, 1H), 9.58-7.74 (m, 5H), 7.42-7.45 (m, 3H), 5.48-5.52 (m, 1H), 1.46-1.48 (d, J = 3.0 Hz, 3H).


Example 73: Preparation of (S)-5-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one



10.7 mg of (S)-5-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 5 of Example 34 except that (S)-2-(1-aminopropyl)-5-fluoro-3-(phenylquinazoline-4(3H)-one was used (0.02 mmol, yield: 44%).

¹H NMR (300 MHz, CDCl₃) δ 9.69-9.72 (d, J = 4.5 Hz, 1H), 8.84 (s, 1H), 8.68 (s, 1H), 7.51-7.69 (m, 7H), 7.34-7.36 (m, 1H), 7.09-7.15 (t, J = 9.1 Hz, 1H), 5.07-5.11 (m, 1H), 1.86-1.96 (m, 2H), 0.87-0.92 (t, J = 7.1 Hz, 6H).


Example 74: Preparation of (S)-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one



24.7 mg of (S)-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 5 of Example 34 except that (S)-5-chloro-3-phenyl-2-(pyrrolidine-2-yl)quinazoline-4(3H)-one was used (0.05 mmol, yield: 95%).

¹H NMR (300 MHz, CDCl₃) δ 9.56 (s, 1H), 8.53 (s, 1H), 7.74-7.61 (m, 1H), 7.42-7.63 (m, 6H), 7.29 (s, 1H), 4.84-4.88 (m, 1H), 4.00-4.02 (m, 1H), 3.70-3.74 (m, 1H), 2.04-2.29 (m, 2H), 1.83-1.87 (m, 2H).


Example 75: Preparation of (S)-5-(2-(8-chloro-1-oxo-2-(pyridine-3-yl)-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one



10.6 mg of (S)-5-(2-(8-chloro-1-oxo-2-(pyridine-3-yl)-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 5 of Example 34 except that (S)-8-chloro-2-(pyridine-3-yl)-3-(pyrrolidine-2-yl)isoquinoline-1(2H)-one was used (0.02 mmol, yield: 41%).

¹H NMR (300 MHz, DMSO-d₆) δ 8.49 (s, 1H), 8.25 (s, 1H), 7.48-7.61 (m, 6H), 7.35-7.41 (m, 3H), 6.46 (s, 1H), 4.67-4.71 (m, 1H), 4.09-4.15 (m, 1H), 3.09-3.16 (m, 1H), 1.88-1.97 (m, 3H), 1.49-1.53 (m, 1H).


Example 76: Preparation of (S)-5-(2-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one



15.8 mg of (S)-5-(2-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as
 5 a white solid according to the same manner as described in step 5 of Example 34 except that (S)-5-chloro-3-(pyridine-3-yl)-2-(pyrrolidine-2-yl)quinazoline-4(3H)-one was used (0.03 mmol, yield: 61%).

10 ^1H NMR (300 MHz, DMSO- d_6) δ 12.41 (s, 1H), 8.44 (s, 1H), 8.01-8.22 (m, 3H), 7.39-7.63 (m, 4H), 4.41-4.47 (m, 1H), 3.77-3.83 (m, 1H), 3.45-3.52 (m, 1H), 1.99-2.15 (m, 3H), 1.69-1.75 (m, 1H).


15 Example 77: Preparation of (S)-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one



4.1 mg of (S)-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 5 of Example 34 except that (S)-5-chloro-3-(3-fluorophenyl)-2-(pyrrolidine-2-yl)isoquinoline-4(3H)-one was used (0.01 mmol, yield: 15%).

¹H NMR (300 MHz, CDCl₃) δ 9.56 (s, 1H), 8.54 (s, 1H), 7.52-7.64 (m, 4H), 7.44-7.47 (m, 2H), 7.07-7.14 (m, 1H), 4.86-4.90 (q, J = 3.0, 1.5 Hz, 1H), 4.03-4.08 (m, 1H), 3.74-3.76 (m, 1H), 2.32-2.37 (m, 1H), 2.11-2.23 (m, 2H), 1.87-1.91 (m, 1H).


Example 78: Preparation of (S)-5-(2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one



10.7 mg of (S)-5-(2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as
 5 a white solid according to the same manner as described in step 5 of Example 34 except that (S)-5-chloro-2-(pyrrolidine-2-yl)-3-m-tolylquinazoline-4(3H)-one was used (0.02 mmol, yield: 40%).

¹H NMR (300 MHz, CDCl₃) δ 8.53 (s, 1H), 8.47-8.48 (d, J = 1.5 Hz, 1H), 7.42-7.57 (m, 6H), 7.35-7.36 (m, 1H), 7.09-7.11 (m, 1H), 4.86-4.92 (m, 1H), 4.04-4.07 (m, 1H), 3.68-3.72 (m, 1H), 2.46-2.49 (d, J = 4.5 Hz, 3H), 2.31-2.34 (m, 1H), 2.15-2.19 (m, 3H).

15 Example 79: Preparation of (S)-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one




10 mg of (S)-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared
 5 as a white solid by using 13 mg (0.07 mmol, 1.0 equivalent) of 5-chloropyrimido[4,5-d]pyrimidine-4(1H)-one and 30 mg (0.09 mmol, 1.3 eq) of (S)-5-chloro-3-phenyl-2-(pyrrolidine-2-yl)pyrrolo[2,1-f][1,2,4]triazine-4(3H)-one according to the same
 10 manner as described in step 5 of Example 34 (0.02 mmol, yield: 31%).

¹H NMR (300 MHz, DMSO-d₆) δ 8.55 (s, 1H), 8.27 (s, 1H), 7.72-7.77 (m, 1H), 7.50-7.62 (m, 5H), 6.61 (s, 1H), 4.52 (s, 1H), 3.78-3.83 (m, 1H), 3.40-3.46 (m,
 15 1H), 1.96-2.15 (m, 3H), 1.70-1.76 (m, 1H).

Example 80: Preparation of (S)-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydropyrrolo[2,1-


f][1,2,4]triazine-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one



20 mg of (S)-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid by using 16 mg (0.09 mmol, 1.0 equivalent) of 5-chloropyrimido[4,5-d]pyrimidine-4(1H)-one and 30 mg (0.09 mmol, 1.2 equivalent) of (S)-5-chloro-3-(3-fluorophenyl)-2-(pyrrolidine-2-yl)pyrrolo[2,1-f][1,2,4]triazine-4(3H)-one according to the same manner as described in step 5 of Example 34 (0.04 mmol, yield: 47%).

¹H NMR (300 MHz, CDCl₃) δ 8.56-8.63 (m, 2H), 7.51-7.60 (m, 2H), 7.28-7.31 (m, 1H), 7.03-7.14 (m, 2H), 6.42 (s, 1H), 4.87 (s, 1H), 3.94-4.02 (m, 1H), 3.65-3.75 (m, 1H), 3.26-3.33 (m, 1H), 1.86-2.28 (m, 4H).

Example 81: Preparation of (S)-7-amino-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one



5 Step 1: Preparation of 5-chloro-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one

3.0 g (12.55 mmol, 1.0 eq) of 4,6-dichloro-2-(methylthio)pyrimidine-5-carboxylic acid was dissolved in 40 mL of anhydrous toluene, to which 15 mL of
 10 thionyl chloride (SOCl₂) was added, followed by stirring at 115°C for 12 hours. The reaction solvent was concentrated under reduced pressure and dried to give acid chloride. Then, 1.6 g of 5-chloro-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one was
 15 prepared as a white solid according to the same manner as described in step 2 of Example 66 (2 step yield: 56%).

¹H NMR (300 MHz, DMSO-d₆) δ 2.56-2.58 (d, J = 6.0, 3H), 8.42-8.44 (d, J = 6.0, 1H), 12.88 (br s, 1H).

Step 2: Preparation of (S)-5-(1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethylamino)-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one

22 mg of (S)-5-(1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethylamino)-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid by using 10 mg (0.055 mmol) of 5-chloro-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one prepared in step 1 and 20 mg (0.066 mmol, 1.2 equivalent) of (S)-2-(1-aminoethyl)-5-chloro-2-phenylquinazoline-4(3H)-one according to the same manner as described in step 5 of Example 34 (0.044 mmol, yield: 80%).

¹H NMR (300 MHz, CDCl₃) δ 9.42 (s, 1H), 8.21 (s, 1H), 7.64-7.33 (m, 7H), 7.36-7.33 (m, 1H), 5.19-5.13 (m, 1H), 2.43 (s, 3H), 1.47 (d, J = 6.5 Hz, 3H).

Step 3: Preparation of (S)-7-amino-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one

16 mg of (S)-7-amino-5-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid by using 19.6 mg (0.04 mmol) of (S)-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-

dihydroquinazoline-2-yl)ethyl)amino)-7-


(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one

prepared in step 2 according to the same manner as described in step 4 of Example 15 (0.034 mmol, yield:

5 85%).

¹H NMR (300 MHz, DMSO-d₆) δ 9.61-9.58 (m, -NH), 8.06 (s, 1H), 7.77-7.72 (m, 1H), 7.61-7.54 (m, 6H), 4.77-4.72 (m, 1H), 1.31 (d, J = 6.5 Hz, 3H).

10 Example 82: Preparation of (S)-7-amino-5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one



Step 1: Preparation of (S)-5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one

30 mg of (S)-5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-7-

20 (methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner

as described in step 5 of Example 34 except that (S)-2-(1-aminoethyl)-5-chloro-3-(pyridine-3-yl)quinazoline-4(3H)-one was used (0.06 mmol, yield: 60%).


5 ¹H NMR (300 MHz, CDCl₃) δ 11.77 (s, 1H), 11.32 (s, 1H), 9.28-9.20 (m, 1H), 8.89 (s, 1H), 8.81-8.79 (m, 1H), 8.72-8.70 (m, 1H), 8.63-8.62 (m, 1H), 8.24 (s, 1H), 8.10-8.05 (m, 1H), 7.77-7.75 (m, 1H), 7.66-7.48 (m, 5H), 5.18-5.13 (m, 1H), 4.95-4.90 (m, 1H), 2.49
10 (d, J = 11.7 Hz, 3H), 1.56-1.52 (m, 3H).

Step 2: Preparation of (S)-7-amino-5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one

15 11 mg of (S)-7-amino-5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a pale yellow solid by using 24 mg (0.049 mmol) of (S)-5-(1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroisoquinoline-2-yl)ethylamino)-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one prepared in step 1 according to the same manner as described in step 4 of Example 15 (0.024 mmol, yield: 49%).

¹H NMR (300 MHz, DMSO-d₆) δ 12.14 (brs, 1H), 9.40 (brs, 1H), 8.74-8.81 (m, 1H), 8.64-8.73 (m, 1H), 8.03-8.14 (m, 2H), 7.72-7.84 (m, 1H), 7.52-7.67 (m, 3H), 6.80 (brs, 2H), 4.62-4.75 (m, 1H), 1.33 (d, J = 5 6.6 Hz, 3H).

Example 83: Preparation of (S)-7-amino-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one



Step 1: Preparation of (S)-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one

38 mg of (S)-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 5 of Example 34 except that (S)-2-(1-aminoethyl)-5-chloro-3-(3-

fluorophenyl)quinazoline-4(3H)-one was used (0.075 mmol, yield: 75%).


¹H NMR (300 MHz, CDCl₃) δ 9.40 (s, -NH), 8.26 (s, 1H), 7.65-7.47 (m, 5H), 7.37-7.29 (m, 1H), 7.16-7.09 (m, 1H), 5.19-5.12 (m, 1H), 4.09-4.02 (m, 1H), 2.46 (s, 3H), 1.50 (d, J = 6.5 Hz, 1H).

Step 2: Preparation of (S)-7-amino-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one

27 mg of (S)-7-amino-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 4 of Example 15 except that (S)-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one was used (0.056 mmol, yield: 81%).

¹H NMR (300 MHz, DMSO-d₆) δ 12.13 (s, -NH), 9.46 (s, -NH), 8.06 (s, 1H), 7.78-7.73 (m, 1H), 7.62-7.31 (m, 6H), 6.87 (s, -NH), 6.64 (s, -NH), 4.80-4.72 (m, 1H), 1.35-1.33 (m, 3H).

Example 84: Preparation of (S)-7-amino-5-((1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one



5 Step 1: Preparation of (S)-5-((1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one

45 mg of (S)-5-((1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-7-

10 (methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 5 of Example 34 except that (S)-2-(1-aminoethyl)-5-chloro-3-m-tolylquinazoline-4(3H)-one was used (0.089 mmol, yield: 89%).


15 ^1H NMR (300 MHz, CDCl_3) δ 9.46 (s, 1H), 8.23 (s, 1H), 7.65-7.56 (m, 2H), 7.47-7.41 (m, 2H), 7.30-7.27 (m, 1H), 7.14 (s, 1H), 5.22-5.15 (m, 1H), 2.46-2.39 (m, 6H), 2.48-47 (m, 3H).

Step 2: Preparation of (S)-7-amino-5-((1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one

35 mg of (S)-7-amino-5-((1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 4 of Example 15 except that (S)-5-((1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one was used (0.073 mmol, yield: 82%).

¹H NMR (300 MHz, DMSO-d₆) δ 12.13 (s, -NH), 9.60-9.48 (s, -NH), 8.06 (s, 1H), 7.78-7.72 (m, 1H), 7.63-7.53 (m, 2H), 7.45-7.27 (m, 4H), 6.80-6.71 (m, -NH₂), 4.84-4.77 (m, 1H), 2.47 (s, 3H), 1.35-1.30 (m, 3H).

Example 85: Preparation of (S)-7-amino-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)pyrimido[4,5-d]pyrimidine-4(1H)-one



Step 1: Preparation of (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one

35 mg of (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid by using 17 mg (0.0743 mmol, 1.0 equivalent) of 5-chloro-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one prepared in step 1 of Example 81 and 27 mg (0.0892 mmol, 1.2 equivalent) of (S)-3-(1-aminoethyl)-8-chloro-2-phenylisoquinoline-1(2H)-one according to the same manner as described in step 5 of Example 34 (yield: 96%).

¹H NMR (300 MHz, DMSO-d₆) δ 1.40-1.42 (d, J = 6.0, 3H), 2.34 (s, 3H), 4.75-4.80 (m, 1H), 6.82 (s, 1H), 7.27-7.36 (m, 3H), 7.42-7.44 (m, 1H), 7.49-7.54 (m, 2H), 7.60-7.67 (m, 2H), 8.28 (s, 1H), 9.13-9.15 (d, J = 6.0, 1H), 12.78 (br s, 1H).

Step 2: Preparation of (S)-7-amino-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)pyrimido[4,5-d]pyrimidine-4(1H)-one


60 mg of (S)-7-amino-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-

yl)ethylamino)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid by using 170 mg (0.346 mmol) of (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-7-

5 (methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one prepared in step 2 according to the same manner as described in step 4 of Example 15 (yield: 38%).

¹H NMR (300 MHz, DMSO-d₆) δ 1.29-1.31 (d, J = 6.0, 3H), 4.58-4.62 (m, 1H), 6.57-6.94 (m, 3H), 7.42-7.64
10 (m, 8H), 8.08 (s, 1H), 9.05-9.06 (d, J = 3.0, 1H), 12.24 (s, 1H).

Example 86: Preparation of (S)-7-amino-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)ethyl)amino)pyrimido[4,5-
15 d]pyrimidine-4(1H)-one



Step 1: Preparation of (S)-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-
20 yl)ethyl)amino)-7-(methylthio)pyrimido[4,5-
d]pyrimidine-4(1H)-one

102 mg of (S)-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)ethyl)amino)-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid
5 by using 58 mg (0.20 mmol) of (S)-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)ethyl)amino)-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one according to the same manner as described in step 5 of Example 34 (yield: 94%).

10 ^1H NMR (300 MHz, CDCl_3) δ 9.15(br s, 1H), 8.32(s, 1H), 7.30-7.65(m, 7H), 6.52(s, 1H), 5.10-5.25(m, 1H), 2.47(s, 3H), 1.50(d, $J = 6.4\text{Hz}$, 3H).


Step 2: Preparation of (S)-7-amino-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one
15

14 mg of (S)-7-amino-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid by using 43 mg (0.089 mmol) of (S)-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)ethyl)amino)-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one according to the same manner as
25

described in step 4 of Example 15 (0.031 mmol, yield: 35%).

¹H NMR (500 MHz, DMSO-d₆) δ 12.18 (brs, 1H), 9.11 (brs, 1H), 8.10 (s, 1H), 7.70-7.73 (m, 1H), 7.54-7.62 (m, 1H), 7.42-7.52 (m, 2H), 7.24-7.35 (m, 1H), 6.93 (brs, 1H), 6.70 (dd, J = 1.0 Hz, J = 3.1 Hz, 1H), 6.62 (brs, 1H), 4.75-4.83 (m, 1H), 1.35-1.41 (m, 3H).

Example 87: Preparation of (S)-7-amino-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one




20 mg of (S)-7-amino-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid by using 86 mg (0.172 mmol) of (S)-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)ethyl)amino)-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one

according to the same manner as described in step 4 of Example 15 (0.043 mmol, yield: 25%).

¹H NMR (500 MHz, DMSO-d₆) δ 9.19 (brs, 1H), 8.09 (s, 1H), 7.68 (s, 1H), 7.53-7.63 (m, 3H), 7.48 (s, 2H), 6.95 (brs, 1H), 6.68 (s, 1H), 6.60 (brs, 1H), 7.71-7.78 (m, 1H), 1.32-1.38 (m, 3H).

Example 88: Preparation of (S)-7-amino-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one



Step 1: Preparation of (S)-7-(methylthio)-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one

23 mg of (S)-7-(methylthio)-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 5 of Example 34 except that (S)-1-(2-phenylquinoline-3-yl)ethan-1-amine was used (0.052 mmol, yield: 52%).

¹H NMR (300 MHz, CDCl₃) δ 12.32 (s, -NH), 9.31-9.29 (m, 1H), 8.26 (s, 1H), 8.20 (d, J = 8.5 Hz, 1H), 8.04 (s, 1H), 7.83 (d, J = 8.2 Hz, 1H), 7.74-7.68 (m, 3H), 7.57-7.43 (m, 4H), 5.78-5.68 (m, 1H), 1.51 (d, J = 6.6 Hz, 3H).


Step 2: Preparation of (S)-7-amino-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one

10 16 mg of (S)-7-amino-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 4 of Example 15 except that (S)-7-(methylthio)-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one was
15 used (0.039 mmol, yield: 81%).

¹H NMR (300 MHz, DMSO-d₆) δ 12.16 (s, -NH), 9.36 (s, -NH), 8.36 (s, 1H), 8.07 (s, 1H), 8.01-7.96 (m, 2H), 7.75-7.44 (m, 7H), 7.63-7.53 (m, 2H), 7.45-7.27 (m, 4H), 6.86 (m, -NH), 6.75 (m, -NH), 5.48-5.44 (m, 1H), 1.33 (d, J = 6.1 Hz, 3H).

Example 89: Preparation of (S)-7-amino-5-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one

25



Step 1: Preparation of (S)-5-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one

45 mg of (S)-5-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 5 of Example 34 except that (S)-2-(1-aminopropyl)-5-fluoro-3-(pyridine-3-yl)quinazoline-4(3H)-one was used (0.089 mmol, yield: 89%).

¹H NMR (300 MHz, CDCl₃) δ 9.49-9.45 (m, 1H), 8.21 (s, 1H), 7.72-7.47 (m, 5H), 7.35-7.31 (m, 1H), 7.15-7.08 (m, 1H), 5.13-5.06 (m, 1H), 2.38 (s, 3H), 1.98-1.76 (m, 2H), 0.87-0.78 (m, 3H).


Step 2: Preparation of (S)-7-amino-5-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one

34 mg of (S)-7-amino-5-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 4 of Example 15 except that (S)-5-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one was used (0.075 mmol, yield: 88%).

¹H NMR (300 MHz, DMSO-d₆) δ 12.14 (s, -NH), 9.45 (s, -NH), 8.06 (s, 1H), 7.82-7.74 (m, 1H), 7.61-7.51 (m, 5H), 7.46-7.43 (m, 1H), 7.30-7.23 (m, 1H), 6.85 (s, -NH), 6.50 (s, -NH), 4.67 (s, 1H), 1.90-1.82 (m, 1H), 1.60-1.50 (m, 1H), 0.71-0.66 (m, 3H).

15

Example 90: Preparation of (S)-7-amino-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one



20

Step 1: Preparation of (S)-5-(2-(5-chloro-4--oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one

23 mg of (S)-5-(2-(5-chloro-4--oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 5 of Example 34 except that (S)-5-chloro-3-phenyl-2-(pyrrolidine-2-yl)quinazoline-4(3H)-one was used (0.044 mmol, yield: 80%).

¹H NMR(300 MHz, CDCl₃) δ 8.17 (s, 1H), 8.03-8.00 (m, 1H), 7.69-7.40 (m, 8H), 4.85-4.80 (m, 1H), 4.06-4.00 (m, 1H), 3.68-3.60 (m, 1H), 2.53 (s, 3H), 2.29-2.10 (2H), 1.83-1.76 (m, 2H).

15

Step 2: Preparation of (S)-7-amino-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one


20 20 mg of (S)-7-amino-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 4 of Example 15 except that (S)-5-(2-(5-chloro-4--oxo-3-phenyl-3,4-dihydroquinazoline-2-

25

yl)pyrrolidine-1-yl)-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one was used (0.043 mmol, yield: 86%).

¹H NMR (300 MHz, CDCl₃) δ 11.77 (s, 1H), 9.10 (s, 1H), 8.72-8.70 (m, 2H), 8.37 (s, 1H), 8.00 (s, 1H), 7.69-7.47 (m, 4H), 4.83-4.80 (m, 1H), 4.49-4.41 (m, 1H), 3.87-3.73 (m, 2H), 1.28-1.24 (m, 4H).

Example 91: Preparation of (S)-7-amino-5-(2-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one



Step 1: Preparation of (S)-5-(2-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one

30 mg of (S)-5-(2-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner

as described in step 5 of Example 34 except that (S)-5-chloro-3-(pyridine-3-yl)-2-(pyrrolidine-2-yl)quinazoline-4(3H)-one was used (0.057 mmol, yield: 57%).

5 ^1H NMR (300 MHz, CDCl_3) δ 10.96 (s, 1H), 9.31 (s, 1H), 8.82-8.78 (m, 1H), 8.56 (s, 1H), 8.45-8.42 (m, 1H), 8.24 (s, 1H), 7.56-7.43 (m, 4H), 4.83-4.80 (m, 1H), 4.69-4.64 (m, 1H), 3.71-3.60 (m, 1H), 2.53 (s, 3H), 1.28-1.24 (m, 1H).

10


Step 2: Preparation of (S)-7-amino-5-(2-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one

15 20 mg of (S)-7-amino-5-(2-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 4 of Example 15
20 except that (S)-5-(2-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one was used (0.043 mmol, yield: 86%).

25 ^1H NMR (300 MHz, CDCl_3) δ 11.77 (s, 1H), 9.10 (s, 1H), 8.72-8.70 (m, 2H), 8.37 (s, 1H), 8.00 (s, 1H),

7.69-7.47 (m, 4H), 4.83-4.80 (m, 1H), 4.49-4.41 (m, 1H), 3.87-3.73 (m, 2H), 1.28-1.24 (m, 4H).

Example 92: Preparation of (S)-7-amino-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one



Step 1: Preparation of (S)-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one

38 mg of (S)-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 5 of Example 34 except that (S)-5-chloro-3-(3-fluorophenyl)-2-(pyrrolidine-2-yl)quinazoline-4(3H)-one was used (0.071 mmol, yield: 71%).

¹H NMR (300 MHz, CDCl₃) δ 11.62 (s, -NH), 8.29 (s, 1H), 7.86-8.72 (m, 1H), 7.68-7.41 (m, 5H), 7.29-7.24 (m, 1H), 7.09-7.02 (m, 1H), 4.84-4.78 (m, 1H), 4.09-4.02 (m, 1H), 3.65-3.60 (m, 1H), 2.52 (s, 3H),
5 2.25-2.08 (m, 4H), 1.84-1.77 (m, 1H).


Step 2: Preparation of (S)-7-amino-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one
10

(S)-7-amino-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared according to the same manner as described in step 4 of
15 Example 15 except that (S)-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one was used.

¹H NMR (300 MHz, DMSO-d₆) δ 11.76 (br s, 1H), 8.00 (s, 1H), 7.95-7.30 (m, 8H), 6.62 (br s, 2H), 4.60-4.40 (m, 1H), 3.90-3.70 (m, 1H), 3.65-1.77 (m, 6H).
20

Example 93: Preparation of (S)-7-amino-5-(2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-

yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one



Step 1: Preparation of (S)-5-(2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one

45 mg of (S)-5-(2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 5 of Example 34 except that (S)-5-chloro-2-(pyrrolidine-2-yl)-3-m-tolylquinazoline-4(3H)-one was used (0.085 mmol, yield: 85%).

¹H NMR (300 MHz, CDCl₃) δ 11.72 (s -NH), 8.26 (s, 1H), 7.84 (s, 1H), 7.55-7.41 (m, 5H), 7.07 (s, 1H), 4.87-4.81 (m, 1H), 4.13-4.07 (m, 1H), 3.58-3.51 (m, 1H), 2.50 (s, 3H), 2.20-2.04 (m, 4H), 1.79-1.73 (m, 2H).


Step 2: Preparation of (S)-7-amino-5-(2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-

yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one

31 mg of (S)-7-amino-5-(2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 4 of Example 15 except that (S)-5-(2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one was used (0.061 mmol, yield: 77%).

^1H NMR (300 MHz, DMSO- d_6) δ 11.75 (br s, 1H), 8.00 (s, 1H), 7.80-7.20 (m, 7H), 6.53 (br s, 2H), 4.63-4.53 (m, 1H), 3.84-3.74 (m, 1H), 3.60-3.40 (m, 1H), 2.37 (s, 3H), 2.10-1.27 (m, 4H).

Example 94: Preparation of (S)-7-amino-5-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one



Step 1: Preparation of (S)-5-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one

42 mg of (S)-5-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared as a white solid according to the same manner as described in step 5 of Example 34 except that (S)-8-chloro-2-phenyl-3-(pyrrolidine-2-yl)isoquinoline-4(3H)-one was used (0.081 mmol, yield: 81%).

¹H NMR (300 MHz, CDCl₃) δ 11.55 (s, -NH), 8.08 (s, 1H), 7.85-7.83 (m, 1H), 7.69-7.64 (m, 1H), 7.69-7.33 (m, 7H), 6.63 (s, 1H), 5.02-4.96 (m, 1H), 4.40-4.31 (m, 1H), 3.18-3.12 (m, 1H), 2.57 (s, 3H), 2.12-1.98 (m, 2H), 1.87-1.81 (m, 1H), 1.64-1.55 (m, 1H).


Step 2: Preparation of (S)-7-amino-5-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one

(S)-7-amino-5-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared according to the same manner as described in step 4 of Example 15 except that (S)-5-(2-(8-chloro-1-oxo-2-

phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)-
7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one was
used.

¹H NMR (300 MHz, DMSO-d₆) δ 11.86 (br s, 1H),
5 8.04 (s, 1H), 7.85-7.83 (m, 1H), 7.69-7.33 (m, 8H),
6.65 (br s, 2H), 6.48 (s, 1H), 4.96-4.80 (m, 1H), 4.20-
4.00 (m, 1H), 3.18-3.00 (m, 1H), 1.90-1.30 (m, 4H).

Example 95: Preparation of (S)-7-amino-5-(2-(5-
10 chloro-4-oxo-3-phenyl-3,4-dihydropyrrolo[2,1-
f][1,2,4]triazine-2-yl)pyrrolidine-1-yl)pyrimido[4,5-
d]pyrimidine-4(1H)-one



Step 1: Preparation of (S)-5-(2-(5-chloro-4-oxo-
15 3-phenyl-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-
yl)pyrrolidine-1-yl)-7-(methylthio)pyrimido[4,5-
d]pyrimidine-4(1H)-one

30 mg of (S)-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-
dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)pyrrolidine-
20 1-yl)-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-
one was prepared as a white solid according to the

same manner as described in step 5 of Example 34 except that (S)-5-chloro-3-phenyl-2-(pyrrolidine-2-yl)pyrrolo[1,2-f][1,2,4]triazine-4(3H)-one hydrochloride was used (0.06 mmol, yield: 66%).

5 ¹H NMR (300 MHz, CDCl₃) δ 8.10 (s, 1H), 7.96-7.93 (m, 1H), 7.65-7.61 (m, 1H), 7.55-7.53 (m, 2H), 7.35 (s, 1H), 7.35-7.34 (m, 1H), 7.28-7.26 (m, 1H), 6.45-6.44 (m, 1H), 4.85-4.81 (m, 1H), 4.16-4.09 (m, 1H), 4.03-3.93 (m, 1H), 2.56 (s, 3H), 2.21-2.12 (m, 1H),
10 2.08-2.05 (m, 2H), 1.82-1.72 (s, 1H).


Step 2: Preparation of (S)-7-amino-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one

15

(S)-7-amino-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared according to the same manner as described in step 4 of
20 Example 15 except that (S)-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)pyrrolidine-1-yl)-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one was used.

^1H NMR (300 MHz, CDCl_3) δ 8.45 (s, 1H), 7.60-7.05 (m, 7H), 6.43 (s, 1H), 4.85-4.81 (m, 1H), 4.16-4.09 (m, 1H), 3.80-3.60 (m, 1H), 2.45-1.65 (m, 4H).

5 Example 96: Preparation of (S)-7-amino-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-
dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)pyrrolidine-
1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one



10 Step 1: Preparation of (S)-5-(2-(5-chloro-3-(3-
fluorophenyl)4-oxo-3,4-dihydropyrrolo[2,1-
f][1,2,4]triazine-2-yl)pyrrolidine-1-yl)-7-
(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one

60 mg of (S)-5-(2-(5-chloro-3-(3-fluorophenyl)4-
 15 oxo-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-
 yl)pyrrolidine-1-yl)-7-(methylthio)pyrimido[4,5-
 d]pyrimidine-4(1H)-one was prepared as a white solid
 according to the same manner as described in step 5 of
 Example 34 except that (S)-5-chloro-3-(3-
 20 fluorophenyl)-2-(pyrrolidine-2-yl)pyrrolo[1,2-

f][1,2,4]triazine-4(3H)-one hydrochloride was used (0.011 mmol, yield: 79%).


¹H NMR(300 MHz, CDCl₃) δ 9.94 (s, -NH), 8.16 (s, 1H), 7.81-7.74 (m, 1H), 7.62-7.48 (m, 2H), 7.14 (s, 1H), 7.09-7.01 (m, 1H), 6.43 (s, 1H), 4.84 (s, 1H), 4.02-3.96 (m, 1H), 3.69-3.58 (m, 1H), 2.58 (s, 3H), 2.21-2.04 (m, 2H), 1.85-1.77 (m, 2H).

Step 2: Preparation of (S)-7-amino-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one

(S)-7-amino-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one was prepared according to the same manner as described in step 4 of Example 15 except that (S)-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydropyrrolo[2,1-f][1,2,4]triazine-2-yl)pyrrolidine-1-yl)-7-(methylthio)pyrimido[4,5-d]pyrimidine-4(1H)-one was used.

¹H NMR(300 MHz, CDCl₃) δ 8.58 (br s, 1H), 7.65-6.80 (m, 6H), 6.44 (s, 1H), 6.10 (s, 1H), 4.90 (s, 1H), 4.25-4.10 (m, 1H), 3.75-3.50 (m, 1H), 2.25-1.10 (m, 4H).

Example 97: Preparation of (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-7,8-dihydropyrido[2,3-d]pyrimidine-5(6H)-one



Step 1: Preparation of (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-8-(4-methoxybenzyl)-7,8-dihydropyrido[2,3-d]pyrimidine-5(6H)-one

326 mg of (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-8-(4-methoxybenzyl)-7,8-dihydropyrido[2,3-d]pyrimidine-5(6H)-one was prepared as a white solid by using 200 mg (0.658 mmol) of 4-chloro-8-(4-methoxybenzyl)-7,8-dihydropyrido[2,3-d]pyrimidine-5(6H)-one and 136 mg (0.790 mmol) of (S)-3-(1-aminoethyl)-8-chloro-2-phenylisoquinoline-1(2H)-one according to the same manner as described in step 5 of Example 34 (0.576 mmol, yield: 87%).


¹H NMR (300 MHz, CDCl₃) δ 9.52 (d, J = 6.4 Hz, 1H), 8.04 (s, 1H), 7.35-7.56 (m, 7H), 7.31 (d, J = 6.4 Hz, 1H), 7.21 (d, J = 8.3 Hz, 2H), 6.86 (d, J = 8.3 Hz, 2H), 6.52 (s, 1H), 4.93 (t, J = 7.4 Hz, 1H), 4.83 (s, 2H), 3.79 (s, 3H), 3.45 (t, J = 6.4 Hz, 2H), 2.58-2.67 (m, 2H), 1.40 (d, J = 6.4 Hz, 3H).

Step 2: Preparation of (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-7,8-dihydropyrido[2,3-d]pyrimidine-5(6H)-one

36 mg of (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-7,8-dihydropyrido[2,3-d]pyrimidine-5(6H)-one was prepared as a white solid by using 50 mg (0.088 mmol) of (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-8-(4-methoxybenzyl)-7,8-dihydropyrido[2,3-d]pyrimidine-5(6H)-one according to the same manner as described in step 8 of Example 1 (0.081 mmol, yield: 91%).

¹H NMR (300 MHz, CDCl₃) δ 9.35 (d, J = 7.2 Hz, 1H), 7.93 (s, 1H), 7.34-7.55 (m, 7H), 7.30 (d, J = 8.4 Hz, 1H), 6.51 (s, 1H), 6.11 (brs, 1H), 4.91 (t, J = 6.1 Hz, 1H), 3.58-3.67 (m, 2H), 2.67-2.75 (m, 2H), 1.40 (d, J = 6.9 Hz, 3H).

Example 98: Preparation of (S)-4-((1-(4,8-dichloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-7,8-dihydropyrido[2,3-d]pyrimidine-5(6H)-one



Step 1: Preparation of (S)-4-((4,8-dichloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-8-(4-methoxybenzyl)-7,8-dihydropyrido[2,3-d]pyrimidine-5(6H)-one

150 mg of (S)-4-((4,8-dichloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-8-(4-methoxybenzyl)-7,8-dihydropyrido[2,3-d]pyrimidine-5(6H)-one was prepared as a white solid by using 100 mg (0.30 mmol) of (S)-3-(1-aminoethyl)-4,8-dichloro-2-phenylisoquinoline-1(2H)-one prepared in step 8 of Preparative Example 10 according to the same manner as described in step 5 of Example 34 (0.25 mmol, yield: 83%).

¹H NMR (300 MHz, CDCl₃) δ 9.97 (br s, 1H), 8.11 (s, 1H), 8.00 (d, J = 8.2Hz, 1H), 7.76 (d, J = 7.7Hz,

1H), 7.51-7.64 (m, 5H), 7.19-7.24 (m, 3H), 6.86 (d, J = 8.6Hz, 2H), 5.07-5.11 (m, 1H), 4.80-4.87 (m, 2H), 3.81 (s, 3H), 3.42-3.46 (m, 2H), 2.59-2.63 (m, 2H), 1.64 (d, J = 7.2Hz, 3H).

5







Step 2: Preparation of (S)-4-((4,8-dichloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-7,8-dihydropyrido[2,3-d]pyrimidine-5(6H)-one








10 78 mg of (S)-4-((4,8-dichloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-7,8-dihydropyrido[2,3-d]pyrimidine-5(6H)-one was prepared as a white solid by using 100 mg (0.17 mmol) of (S)-4-((4,8-dichloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-8-(4-methoxybenzyl)-7,8-dihydropyrido[2,3-d]pyrimidine-5(6H)-one prepared in step 1 according to the same manner as described in step 8 of Example 1 (0.16 mmol, yield: 98%).

15 ¹H NMR (300 MHz, CDCl₃) δ 9.82 (br s, 1H), 7.99 (d, J = 6.7Hz, 1H), 7.98 (s, 1H), 7.68-7.73 (m, 1H), 9.48-7.63 (m, 5H), 7.19-7.21 (m, 1H), 5.80 (s, 1H), 5.03-5.11 (m, 2H), 3.58-3.65 (m, 2H), 2.65-2.72 (m, 2H), 1.64 (d, J = 7.2Hz, 3H).








The formulae of the compounds prepared in Examples 1 ~ 98 are shown in Table 1 and Table 2.









【Table 1】









Example	Formula	Example	Formula
1		2	
3		4	
5		6	







7		7-1	
8			
9		10	
11		12	

13		14	
15		16	
17		18	
19		20	



21		22	
23		24	
25			
26		27	









28		29	
30		31	
32		33	
34		35	

36		37	
38		39	
39-1		40	
41		42	









43		44	
45		46	
47		48	









【Table 2】









Example	Formula	Example	Formula
49		50	

51		52	
53		54	
55		56	
57		58	

59		60	
61		62	
63		64	
65		66	

67		68	
69		70	
71		72	
73		74	

75		76	
77		78	
79		80	
81		82	

83		84	
85		86	
87		88	
89		90	

91		92	
93		94	
95		96	
97		98	

Experimental Example 1: Inhibition of phosphatidylinositol 3-kinase alpha (PI3K α)

The inhibition of phosphatidylinositol 3 kinase alpha (PI3K α) by those compounds prepared in Examples 1 ~ 98 was investigated by the following experiment. The inhibition activity of each compound was measured in the following conditions: ATP = 10 uM, and sample concentration = 100 nM.

Step 1: Human breast cancer cells (MDA-MB-453) cultured in DMEM (Dulbecco's Modified Eagle Medium, Hyclon, SH30243.01) supplemented with 10% fetal bovine serum (Hyclon, USA) were distributed in a 12-well plate at the density of 1,000,000 cells/well. The cells were stabilized in a 37°C CO₂ for 24 hours and then treated with each compound for 1 and half hours. The cells were treated with 10 ng/mL of EGF (Epidermal Growth Factor, 10 µg/mL; R&D, 2150-C5) to increase the intracellular activity of PI3K alpha. 5 minutes later, the medium was discarded and the cells were washed with cold PBS (phosphate buffered saline, Gibco, 14190-250). PBS was then eliminated completely by using a pipette. Western blotting was performed as described in step 2 to evaluate the intracellular PI3K alpha activity.

Step 2: Western blotting

The stimulated cells were transferred in a 1.5 mL tube, followed by centrifugation at 3000 rpm for 1 minute. 100 μ L of RIPA buffer (radioimmunoprecipitation assay buffer, 50 mM Tris-HCl, 5 mM EDTA, 150 mM NaCl, 1% NP-40, 1 mM PMSF, pH 8.0; ELPIS, Korea) was added thereto, which was stored in a 4°C refrigerator for 12 hours. Centrifugation was performed at 4°C at 14,000 rpm for 20 minutes. The supernatant was transferred into a new 1.5 mL tube. Protein was quantified by BCA (bicinchoninic acid) method and samples were prepared. The composition of the sample was as follows: sample buffer (ELPIS, EBA-1052) 5X, protein 10 μ g, and 1X sample buffer to make the total volume 20 μ L. The sample was heated at 100°C for 5 minutes. The evaporated water vapor was condensed in a refrigerator. The liquid stained on the wall was fallen down by centrifugation for a few seconds. The sample proceeded to 10% SDS (Sodium Dodecyl Sulfate) acrylamide gel for separation. The separated protein was transferred onto PVDF (poly-vinyl difluoride) membrane, followed by reaction with pAkt (phospho protein kinase B) antibody (Ser473 or Thr308; Cell signaling, 9271s or 13038s) for 12 hours at 4°C. The membrane was washed with TBST (Tris-Buffered Saline

with Tween 20, 10 mM Tris-HCl, pH 7.5, 150 mM NaCl, 0.1% Tween-20) three times for 5 minutes each, to which the secondary antibody (rabbit antibody, Santacruz, sc-2004) was added, followed by reaction at room temperature for 2 hours. The membrane was washed with TBST three times for 10 minutes each, to which ECL (enhanced chemiluminescence, Thermo, NCI34095KR) was sprayed. Bands were investigated by using LAS-3000.

10

Experimental Example 2: Inhibition of phosphatidylinositol 3-kinase beta (PI3K β)

The inhibition of phosphatidylinositol 3 kinase beta (PI3K β) by those compounds prepared in Examples 1 ~ 98 was investigated by the following experiment. The inhibition activity of each compound was measured in the following conditions: ATP = 10 uM, and sample concentration = 100 nM.

Human prostate cancer cells (PC3 cells) cultured in DMEM (Dulbecco's Modified Eagle Medium, Hyclon, SH30243.01) supplemented with 10% fetal bovine serum (Hyclon, USA) were distributed in a 12-well plate at the density of 1,000,000 cells/well. The cells were stabilized in a 37°C CO₂ for 24 hours and then treated with each compound for 1 and half hours. The cells

were treated with 10 ng/mL of LPA (lysophosphatidic acid, 10 µg/mL; R&D, 2150-C5) to increase the intracellular activity of PI3K beta. 5 minutes later, the medium was discarded and the cells were washed
5 with cold PBS (phosphate buffered saline, Gibco, 14190-250). PBS was then eliminated completely by using a pipette. Western blotting was performed according to the same manner as described in step 2 of Experimental Example 1 to evaluate the intracellular
10 PI3K beta activity.

Experimental Example 3: Inhibition of
phosphatidylinositol 3-kinase gamma (PI3K γ)

The inhibition of phosphatidylinositol 3 kinase
15 gamma (PI3K γ) by those compounds prepared in Examples 1 ~ 98 was investigated by the following experiment. The inhibition activity of each compound was measured in the following conditions: ATP = 10 uM, and sample concentration = 100 nM.

20 Macrophages (RAW264.7) cultured in DMEM (Dulbecco's Modified Eagle Medium, Hyclon, SH30243.01) supplemented with 10% fetal bovine serum (Hyclon, USA) were distributed in a 12-well plate at the density of 1,000,000 cells/well. The cells were stabilized in a
25 37°C CO₂ for 24 hours and then treated with each

compound for 1 and half hours. The cells were treated with 10 ng/mL of C5a (complement component 5a, 10 µg/mL; R&D, 2150-C5) to increase the intracellular activity of PI3K gamma. 5 minutes later, the medium was discarded and the cells were washed with cold PBS (phosphate buffered saline, Gibco, 14190-250). PBS was then eliminated completely by using a pipette. Western blotting was performed according to the same manner as described in step 2 of Experimental Example 1 to evaluate the intracellular PI3K gamma activity.

Experimental Example 4: Inhibition of phosphatidylinositol 3-kinase delta (PI3K δ)

The inhibition of phosphatidylinositol 3 kinase delta (PI3K δ) by those compounds prepared in Examples 1 ~ 98 was investigated by the following experiment. The inhibition activity of each compound was measured in the following conditions: ATP = 10 uM, and sample concentration = 100 nM.

Raji cells cultured in RPMI1640 (Hyclone, SH30027.02) supplemented with 10% fetal bovine serum (Hyclon, USA) were distributed in a 12-well plate at the density of 1,000,000 cells/well. The cells were stabilized in a 37°C CO₂ for 24 hours and then treated with each compound for 1 and half hours. The cells

were treated with 0.25 µg/mL of IgM (immunoglobulin M, Southern Biotech, USA) to increase the intracellular activity of PI3K delta. 30 minutes later, the medium was discarded and the cells were washed with cold PBS (phosphate buffered saline, Gibco, 14190-250). PBS was then eliminated completely by using a pipette. Western blotting was performed according to the same manner as described in step 2 of Experimental Example 1 to evaluate the intracellular PI3K delta activity.

10

The inhibition of phosphatidylinositol 3-kinase alpha, beta, gamma, and delta (PI3K α , β , γ , and δ) by those compounds prepared in Examples 1 ~ 98 was investigated and the results obtained in Experimental Examples 1 ~ 4 are shown in Table 3.

【Table 3】

Example	PI3 Kinase (p110 α /p85 α) (h)	PI3 Kinase (p110 β /p86 α) (h)	PI3 Kinase (p120 γ) (h)	PI3 Kinase (p110 δ /p85 α) (h)
1	+	++	++	+++
2	+	+	++	+++
3	+	+	++	+++
4	+	+	++	+++
5	+	++	++	+++
6	+	+	++	+++
7	+	+	+	+++
7-1	+	+	+	++
8	+	+	++	+++
9	+	+	++	+++

10	+	+	++	+++
11	+	+	++	+++
12	+	+	++	+++
13	+	+	++	+++
14	+	+	++	++
15	++	++	++	+++
16	+	+	++	++
17	+	+	++	++
18	++	++	++	+++
19	+	+	+	+
20	+	+	++	+++
21	++	++	++	+++
22	+	+	++	+++
23	+	+	++	+++
24	+	+	++	++
25	+	+	+++	+++
26	+	+	+	++
27	+	+	++	++
28	+	+	+	++
29	+	+	++	+++
30	+	+	++	+++
31	+	+	+	++
32				+
33	+	+	++	+++
34			++	+++
35			++	++
36			++	+++
37			++	+++
38	++	+	+++	+++
39			++	++
39-1			+++	++
40	+	+	++	+++
41			++	++
42			++	++
43			+	+
44			++	+++
45			++	++
46			++	+++
47			++	++
48			+++	+++

49			++	+++
50			++	+++
51			++	+++
52			+++	+++
53			++	++
54			++	++
55			++	++
56			+	+
57			++	+
58			+	+
59			+	+
60	+	+	++	++
61	+	+	+	+
62	+	+	+	+
63	+	+	+	+
64			+	+
65			+	+
66			+	++
67			+	+
68			+	+
69			+	++
70			++	++
71			+	++
72			+	+
73			+	+
74			+	+
75			+	++
76			+	+
77			+	+
78			+	+
79	+	+	+	+
80			+	+
81			++	++
82			++	++
83			+	+
84			++	++
85	+	+	++	++
86			++	++
87			++	+++
88	++	++	++	++

89			++	++
90			+	+
91			+	+
92			+	+
93			+	+
94			++	++
95	+	+	++	++
96			+	+
97			++	+++
98	+	++	++	+++

(In Table 3,

+ indicates 'over 500 nM';

++ indicates 'over 10 nM ~ up to 500 nM'; and

+++ indicates 'up to 10 uM'.

5

As shown in Table 3, as a result of the investigation of the inhibition of PI3K α , β , γ , and δ by the compounds represented by formula 1 of the present invention, the compounds of the invention were
10 excellent in inhibiting the activity of PI3K α , β , γ , and δ . In particular, the compounds inhibited the activity of PI3 kinase γ or δ even at a very low concentration.

15 Therefore, the compounds of the present invention can act as a PI3 kinase inhibitor, so that they can be effectively used for the prevention or treatment of PI3 kinase related diseases including cancer such as hematological malignance, ovarian cancer, cervical

cancer, breast cancer, colorectal cancer, liver cancer, stomach cancer, pancreatic cancer, colon cancer, peritoneal metastasis, skin cancer, bladder cancer, prostate cancer, thyroid cancer, lung cancer, 5 osteosarcoma, fibrous tumor, and brain tumor; autoimmune disease such as rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis, type 1 diabetes, hyperthyroidism, myasthenia, Crohn's disease, ankylosing spondylitis, psoriasis, autoimmune 10 pernicious anemia, and Sjogren's syndrome; and respiratory disease such as chronic obstructive pulmonary disease (COPD), rhinitis, asthma, chronic bronchitis, chronic inflammatory lung disease, silicosis, pulmonary sarcoidosis, pleurisy, 15 alveolitis, vasculitis, emphysema, pneumonia, and bronchiectasis.

The preparation methods of the invention represented by reaction formulae 1 ~ 3 are novel 20 methods not only facilitating the preparation of those compounds represented by formulae 11, 20, and 23, which are the intermediates of the compound represented by formula 1, but also providing various steps to prepare the compound represented by formula 1 25 by reacting the intermediate compounds represented by

formulae 11, 20, and 23 with the compound that can react to the substituents of the said compounds.

Manufacturing Example 1: Preparation of powders

5 The compound represented by formula 1 2 g
 Lactose 1 g

 Powders were prepared by mixing all the above components, which were filled in airtight packs according to the conventional method for preparing
10 powders.

Manufacturing Example 2: Preparation of tablets

 The compound represented by formula 1 100 mg
 Corn starch 100 mg
15 Lactose 100 mg
 Magnesium stearate 2 mg

 Tablets were prepared by mixing all the above components by the conventional method for preparing tablets.

20

Manufacturing Example 3: Preparation of capsules

 The compound represented by formula 1 100 mg
 Corn starch 100 mg
 Lactose 100 mg
25 Magnesium stearate 2 mg

Capsules were prepared by mixing all the above components, which were filled in gelatin capsules according to the conventional method for preparing capsules.

5

Manufacturing Example 4: Preparation of injectable solution

	The compound represented by formula 1	100 mg
	Mannitol	180 mg
10	$\text{Na}_2\text{HPO}_4 \cdot 2\text{H}_2\text{O}$	26 mg
	Distilled water	2974 mg

Injectable solutions were prepared by mixing all the above components by the conventional method for preparing injectable solutions.

15

Manufacturing Example 5: Preparation of health food

	The compound represented by formula 1	500 ng
	Vitamin A acetate	70 mg
20	Vitamin E	1.0 mg
	Vitamin B1	0.13 mg
	Vitamin B2	0.15 mg
	Vitamin B6	0.5 mg
	Vitamin B12	0.2 μg
25	Vitamin C	10 mg

	Biotin	10 μ g
	Nicotinic acid amide	1.7 mg
	Folic acid	50 μ g
	Calcium pantothenate	0.5 mg
5	Minerals	proper amount
	Ferrous sulfate	1.75 mg
	Zinc oxide	0.82 mg
	Magnesium carbonate	25.3 mg
	Potassium phosphate monobasic	15 mg
10	Potassium phosphate dibasic	55 mg
	Potassium citrate	90 mg
	Calcium carbonate	100 mg
	Magnesium chloride	24.8 mg

Vitamins and minerals were mixed according to the
 15 preferable composition rate for health food. However,
 the composition rate can be adjusted. The
 constituents were mixed according to the conventional
 method for preparing health food and then the
 composition for health food was prepared according to
 20 the conventional method.


Those skilled in the art will appreciate that
 the conceptions and specific embodiments disclosed in
 the foregoing description may be readily utilized as
 25 a basis for modifying or designing other embodiments

for carrying out the same purposes of the present invention. Those skilled in the art will also appreciate that such equivalent embodiments do not depart from the spirit and scope of the invention as
5 set forth in the appended Claims.

CLAIMS:

1. A compound represented by formula 1, an optical isomer thereof, or a pharmaceutically acceptable salt thereof:

[Formula 1]



In the formula 1,

---- is single bond or double bond;


10 A is carbon (C) or nitrogen (N);

R¹ is hydrogen (H) or -NH₂;

R² is H, -CN, C₁₋₅ straight or branched alkyl, unsubstituted C₃₋₇ cycloalkyl or halogen;

R³ and R⁴ are independently H or C₁₋₅ straight or
15 branched alkyl; or

R³ and R⁴ can form 5 ~ 7 membered unsubstituted heterocycloalkyl containing one hetero atoms of N along with the atoms which are conjugated to the same; and



wherein, n is an integer of 0 or 1,

R^6 is unsubstituted or substituted C_{6-8} aryl or
 5 unsubstituted or substituted 5 ~ 6 membered heteroaryl
 containing one or more hetero atoms selected from the
 group consisting of N, O, and S, wherein the
 substituted aryl and the substituted heteroaryl, can
 be independently substituted with one or more
 10 substituents selected from the group consisting of
 halogen, C_{1-5} straight or branched alkyl, and C_{1-5}
 straight or branched alkylsulfonyl,

R^7 and R^8 are independently H, halogen, -CN,
 unsubstituted or substituted C_{6-8} aryl, unsubstituted
 15 or substituted 5 ~ 6 membered heteroaryl containing
 one or more hetero atoms selected from the group
 consisting of N, O, and S, C_{1-5} straight or branched
 alkyl, C_{1-5} straight or branched alkoxy, or $-NR^9R^{10}$,
 wherein R^9 and R^{10} are independently H, C_{1-5} straight or
 20 branched alkyl, C_{1-5} straight or branched alkylamino,

unsubstituted or substituted C₆₋₈ aryl, unsubstituted or substituted 5 ~ 6 membered heteroaryl containing one or more hetero atoms selected from the group consisting of N, O, and S, or unsubstituted or substituted 3 ~ 6 membered heterocycloalkyl containing one or more hetero atoms selected from the group consisting of N, O, and S, wherein the substituted C₆₋₈ aryl, the substituted 5 ~ 6 membered heteroaryl, and the substituted 3 ~ 6 membered heterocycloalkyl can be independently substituted with one or more substituents selected from the group consisting of halogen and C₁₋₅ straight or branched alkyl.

2. The compound represented by formula 1, the optical isomer thereof, or the pharmaceutically acceptable salt thereof according to claim 1, wherein:

--- is single bond or double bond;

A is carbon (C) or nitrogen (N);


R¹ is H or -NH₂;

R² is H, -CN, C₁₋₃ straight or branched alkyl, unsubstituted C₃₋₅ cycloalkyl or halogen;

R³ and R⁴ are independently H or C₁₋₅ straight or branched alkyl; or

R³ and R⁴ can form 5 ~ 7 membered unsubstituted heterocycloalkyl containing one or more hetero atoms

of N along with the atoms which are conjugated to the same; and



5 wherein, n is an integer of 0 or 1,

R⁶ is unsubstituted or substituted C₆₋₈ aryl or unsubstituted or substituted 5 ~ 6 membered heteroaryl containing one or more hetero atoms selected from the group consisting of N, O, and S, wherein the substituted aryl and the substituted heteroaryl can be independently substituted with one or more substituents selected from the group consisting of halogen and C₁₋₅ straight or branched alkyl,

R⁷ is H, halogen, unsubstituted or substituted C₆₋₈ aryl, or unsubstituted or substituted 5 ~ 6 membered heteroaryl containing one or more hetero atoms selected from the group consisting of N, O, and S, wherein the substituted aryl and the substituted heteroaryl can be independently substituted with one or more substituents selected from the group

consisting of halogen and C₁₋₅ straight or branched alkyl, and

R⁸ is H, halogen, C₁₋₃ straight or branched alkyl, or C₁₋₃ straight or branched alkoxy.

5

3. The compound represented by formula 1, the optical isomer thereof, or the pharmaceutically acceptable salt thereof according to claim 1, wherein:

--- is single bond or double bond;

10 A is carbon (C) or nitrogen (N);



R¹ is H or -NH₂;

R² is H, -CN, C₁₋₃ straight or branched alkyl, C₃₋₅ cycloalkyl or halogen;

R³ is H;

15 R⁴ is H or C₁₋₃ straight or branched alkyl; or

R³ and R⁴ can form 5 ~ 7 membered unsubstituted heterocycloalkyl containing one atom of N along with the atoms which are conjugated to the same; and



wherein, n is an integer of 0 or 1,

R⁶ is unsubstituted or substituted phenyl or pyridinyl,

the substituted phenyl and pyridinyl can be
5 independently substituted with one or more
substituents selected from the group consisting of
halogen and C₁₋₃ straight or branched alkyl;

R⁷ is H, halogen, or unsubstituted or substituted
5 ~ 6 membered heteroaryl containing one or more
10 hetero atoms of N, wherein the substituted aryl and
the substituted heteroaryl can be independently
substituted with one or more substituents selected
from the group consisting of halogen and C₁₋₃ straight
or branched alkyl, and

15 R⁸ is H, halogen, or C₁₋₃ straight or branched
alkyl.

4. The compound represented by formula 1, the
optical isomer thereof, or the pharmaceutically
20 acceptable salt thereof according to claim 1, wherein:

--- is single bond or double bond;

A is carbon (C) or nitrogen (N);


R¹ is H or -NH₂;

R² is H, -F, -Cl, -CN, methyl, ethyl, propyl,
25 isopropyl, cyclopropyl, or cyclopentyl;

R^3 is H;

R^4 is H or methyl; or

R^3 and R^4 can form pyrrolidine along with the atoms which are conjugated to the same; and



wherein, n is an integer of 0 or 1,


R^6 is unsubstituted or substituted phenyl or pyridinyl,

10 the substituted phenyl and pyridinyl can be independently substituted with one or more substituents selected from the group consisting of -F, -Cl, and methyl;


R^7 is H, -F, -Cl, or pyridinyl; and


15 R^8 is H, -F, or -Cl.

5. The compound represented by formula 1, the optical isomer thereof, or the pharmaceutically acceptable salt thereof according to claim 1, wherein
20 the ring containing A and R^2 is exemplified



is exemplified by






6. The compound represented by formula 1, the optical isomer thereof, or the pharmaceutically acceptable salt thereof according to claim 1, wherein
 10 the compound represented by formula 1 is

characteristically the compound represented by formula 1A:

[Formula 1A]



5 In the formula 1A,

---, A, R¹, R², R³, R⁴, and R⁵ are as defined in formula 1.

7. The compound represented by formula 1, the
10 optical isomer thereof, or the pharmaceutically acceptable salt thereof according to claim 1, wherein the compound represented by formula 1 is selected from the group consisting of the following compounds:

<1> 4-((1-(5-chloro-4-oxo-3-phenyl-3,4-
15 dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-
d]pyrimidine-5(8H)-one;

<2> 4-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-
dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-
d]pyrimidine-5(8H)-one;

- <3> 4-((1-(5-chloro-4-oxo-3-(pyridine-2-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
- <4> 4-((1-(5-chloro-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
- <5> 4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
- 10 <6> 4-((1-(2-phenylquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
- <7> 4-((1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
- <8> 4-((1-(7-fluoro-2-(3-fluorophenyl)quinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
- 15 <9> 4-(1-(7-fluoro-2-(pyridine-2-yl)quinoline-3-yl)ethylamino)pyrido[2,3-d]pyrimidine-5(8H)-one;
- <10> 4-((1-(4,8-dichloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
- 20 <11> 4-((1-(8-chloro-4-fluoro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

- <12> 4-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
- <13> 4-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrido[2,3-d]pyrimidine-5(8H)-one;
- <14> 4-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)pyrido[2,3-d]pyrimidine-5(8H)-one;
- 10 <15> 2-amino-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
- <16> 2-amino-4-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)pyrido[2,3-d]pyrimidine-5(8H)-one;
- 15 <17> 4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-6-methylpyrido[2,3-d]pyrimidine-5(8H)-one;
- <18> 2-amino-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-6-methylpyrido[2,3-d]pyrimidine-5(8H)-one;
- 20 <19> 4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-5-oxo-5,8-dihydropyrido[2,3-d]pyrimidine-6-carbonitrile;

- <20> 4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-6-fluoropyrido[2,3-d]pyrimidine-5(8H)-one;
- <21> 4-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)-6-fluoropyrido[2,3-d]pyrimidine-5(8H)-one;
- 5 <22> 6-chloro-4-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
- 10 <23> 6-chloro-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
- <24> 6-chloro-4-((1-(4,8-dichloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
- 15 <25> 2-amino-4-((1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
- <26> 4-((1-(6-fluoro-3,4-di(pyridine-2-yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
- 20 <27> 4-((1-(6-fluoro-3-phenyl-4-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<28> 4-((1-(6-fluoro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<29> 4-((1-(6-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<30> 4-((1-(6-fluoro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

10 <31> 4-((1-(5-chloro-3-(2-chlorobenzyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<32> 4-((1-(6-fluoro-4-oxo-3-(pyridine-2-ylmethyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<33> 4-((1-(5-chloro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<34> 5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<35> 5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<36> 5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<37> 5-((1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<38> 5-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

10 <39> 3-methyl-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<40> 5-((1-(4,8-dichloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<41> 5-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<42> 5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<43> 5-(2-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<44> 5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<45> 5-(2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<46> 5-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

10 <48> 7-amino-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<49> 7-amino-5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

15 <50> 7-amino-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<51> 7-amino-5-((1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

20 <52> 7-amino-5-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<53> 7-amino-3-methyl-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<54> 7-amino-5-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<55> 7-amino-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

10 <56> 7-amino-5-(2-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<57> 7-amino-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

15 <58> 7-amino-5-(2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<59> 7-amino-5-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

20 <60> 5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-3-ethyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<61> 5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-3-propyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<62> 5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-3-cyclopropyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<63> 5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-3-cyclopentyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

10 <64> 5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-3-isopropyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<65> 5-(1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propylamino)-3-isopropyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

15 <66> 5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<67> 5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

20 <68> 5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

- <69> 5-((1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;
- <70> 5-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;
- <72> 5-((1-(2-phenylquinoline-3-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;
- <73> 5-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;
- <74> 5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;
- <75> 5-(2-(8-chloro-1-oxo-2-(pyridine-3-yl)-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;
- <76> 5-(2-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;
- <77> 5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<78> 5-(2-(5-chloro-4-oxo-3-(m-tolyl))-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<81> 7-amino-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<82> 7-amino-5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl))-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

10 <83> 7-amino-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<84> 7-amino-5-((1-(5-chloro-4-oxo-3-(m-tolyl))-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<85> 7-amino-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<88> 7-amino-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<89> 7-amino-5-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<90> 7-amino-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<91> 7-amino-5-(2-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<92> 7-amino-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

10 <93> 7-amino-5-(2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<94> 7-amino-5-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

15 <97> 4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-7,8-dihydropyrido[2,3-d]pyrimidine-5(6H)-one; and

<98> 4-((1-(4,8-dichloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-7,8-dihydropyrido[2,3-d]pyrimidine-5(6H)-one.

8. The compound represented by formula 1, the optical isomer thereof, or the pharmaceutically acceptable salt thereof according to claim 1, wherein

the compound represented by formula 1 is selected from the group consisting of the following compounds:

- <1> (S)-4-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
- 5 <2> (S)-4-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
- <3> (S)-4-((1-(5-chloro-4-oxo-3-(pyridine-2-yl)-10 3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
- <4> (S)-4-((1-(5-chloro-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
- 15 <5> (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
- <6> (S)-4-((1-(2-phenylquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
- 20 <7> (S)-4-((1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
- <8> (S)-4-((1-(7-fluoro-2-(3-fluorophenyl)quinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
- 25

- <9> (S)-4-(1-(7-fluoro-2-(pyridine-2-yl)quinoline-3-yl)ethylamino)pyrido[2,3-d]pyrimidine-5(8H)-one;
- <10> (S)-4-((1-(4,8-dichloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
- <11> (S)-4-((1-(8-chloro-4-fluoro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
- 10 <12> (S)-4-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
- <13> (S)-4-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrido[2,3-d]pyrimidine-5(8H)-one;
- 15 <14> (S)-4-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)pyrido[2,3-d]pyrimidine-5(8H)-one;
- <15> (S)-2-amino-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
- 20 <16> (S)-2-amino-4-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)pyrido[2,3-d]pyrimidine-5(8H)-one;

- <17> (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-6-methylpyrido[2,3-d]pyrimidine-5(8H)-one;
- <18> (S)-2-amino-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-6-methylpyrido[2,3-d]pyrimidine-5(8H)-one;
- <19> (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-5-oxo-5,8-dihydropyrido[2,3-d]pyrimidine-6-carbonitrile;
- 10 <20> (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-6-fluoropyrido[2,3-d]pyrimidine-5(8H)-one;
- <21> (S)-4-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)-6-fluoropyrido[2,3-d]pyrimidine-5(8H)-one;
- 15 <22> (S)-6-chloro-4-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
- <23> (S)-6-chloro-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;
- 20 <24> (S)-6-chloro-4-((1-(4,8-dichloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<25> (S)-2-amino-4-((1-(6-fluoro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<26> (S)-4-((1-(6-fluoro-3,4-di(pyridine-2-yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<27> (S)-4-((1-(6-fluoro-3-phenyl-4-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

10 <28> (S)-4-((1-(6-fluoro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<29> (S)-4-((1-(6-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

15 <30> (S)-4-((1-(6-fluoro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<31> (S)-4-((1-(5-chloro-3-(2-chlorobenzyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

20 <32> (S)-4-((1-(6-fluoro-4-oxo-3-(pyridine-2-ylmethyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<33> (S)-4-((1-(5-chloro-3-(pyridine-2-yl)quinoline-2-yl)ethyl)amino)pyrido[2,3-d]pyrimidine-5(8H)-one;

<34> (S)-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<35> (S)-5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

10 <36> (S)-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<37> (S)-5-((1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<38> (S)-5-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<39> (S)-3-methyl-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<40> (S)-5-((1-(4,8-dichloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

- <41> (S)-5-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;
- <42> (S)-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;
- <43> (S)-5-(2-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;
- 10 <44> (S)-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;
- <45> (S)-5-(2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;
- 15 <46> (S)-5-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;
- <48> (S)-7-amino-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;
- 20 <49> (S)-7-amino-5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;
- 25

- <50> (S)-7-amino-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;
- 5 <51> (S)-7-amino-5-((1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;
- <52> (S)-7-amino-5-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-3-methyl-
10 2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;
- <53> (S)-7-amino-3-methyl-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;
- <54> (S)-7-amino-5-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)-3-methyl-
15 2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;
- <55> (S)-7-amino-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;
- 20 <56> (S)-7-amino-5-(2-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;
- <57> (S)-7-amino-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-
- 25

yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<58> (S)-7-amino-5-(2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<59> (S)-7-amino-5-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)-3-methyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

10 <60> (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-3-ethyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<61> (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-3-propyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<62> (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-3-cyclopropyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

20 <63> (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-3-cyclopentyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

<64> (S)-5-(1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethylamino)-3-isopropyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;

- <65> (S)-5-(1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propylamino)-3-isopropyl-2,3-dihydropyrimido[4,5-d]pyrimidine-4(1H)-one;
- <66> (S)-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;
- <67> (S)-5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;
- 10 <68> (S)-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;
- <69> (S)-5-((1-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;
- 15 <70> (S)-5-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;
- <72> (S)-5-((1-(2-phenylquinoline-3-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;
- 20 <73> (S)-5-((1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)propyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<74> (S)-5-(2-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<75> (S)-5-(2-(8-chloro-1-oxo-2-(pyridine-3-yl)-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<76> (S)-5-(2-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

10 <77> (S)-5-(2-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<78> (S)-5-(2-(5-chloro-4-oxo-3-(m-tolyl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

15 <81> (S)-7-amino-5-((1-(5-chloro-4-oxo-3-phenyl-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<82> (S)-7-amino-5-((1-(5-chloro-4-oxo-3-(pyridine-3-yl)-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

20 <83> (S)-7-amino-5-((1-(5-chloro-3-(3-fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<84> (S)-7-amino-5-((1-(5-chloro-4-oxo-3-(m-
tolyl)-3,4-dihydroquinazoline-2-
yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<85> (S)-7-amino-5-(1-(8-chloro-1-oxo-2-phenyl-
5 1,2-dihydroisoquinoline-3-yl)ethylamino)pyrimido[4,5-
d]pyrimidine-4(1H)-one;

<88> (S)-7-amino-5-((1-(2-phenylquinoline-3-
yl)ethyl)amino)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<89> (S)-7-amino-5-((1-(5-fluoro-4-oxo-3-phenyl-
10 3,4-dihydroquinazoline-2-yl)propyl)amino)pyrimido[4,5-
d]pyrimidine-4(1H)-one;

<90> (S)-7-amino-5-(2-(5-chloro-4-oxo-3-phenyl-
3,4-dihydroquinazoline-2-yl)pyrrolidine-1-
yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<91> (S)-7-amino-5-(2-(5-chloro-4-oxo-3-
15 (pyridine-3-yl)-3,4-dihydroquinazoline-2-
yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-
one;

<92> (S)-7-amino-5-(2-(5-chloro-3-(3-
20 fluorophenyl)-4-oxo-3,4-dihydroquinazoline-2-
yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-
one;

<93> (S)-7-amino-5-(2-(5-chloro-4-oxo-3-(m-
tolyl)-3,4-dihydroquinazoline-2-yl)pyrrolidine-1-
25 yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;

<94> (S)-7-amino-5-(2-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)pyrrolidine-1-yl)pyrimido[4,5-d]pyrimidine-4(1H)-one;


<97> (S)-4-((1-(8-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-7,8-dihydropyrido[2,3-d]pyrimidine-5(6H)-one; and

<98> (S)-4-((1-(4,8-dichloro-1-oxo-2-phenyl-1,2-dihydroisoquinoline-3-yl)ethyl)amino)-7,8-dihydropyrido[2,3-d]pyrimidine-5(6H)-one.

10

9. An intermediate compound represented by formula 1B below or an optical isomer of the same:

[Formula 1B]



15 In the formula 1B,

---, A, R¹, R², R³, R⁴, and R⁵ are as defined in formula 1; and

PG is an amine protecting group selected from the group consisting of t-butyloxycarbonyl (Boc), carbobenzyloxy (Cbz), 9-fluorenylmethyloxycarbonyl (Fmoc), acetyl (Ac), benzoyl (Bz), benzyl (Bn), p-

methoxybenzyl (PMB), 3,4-dimethoxybenzyl (DMPM), p-
methoxyphenyl (PMP), tosyl (Ts), 2,2,2-
trichloroethoxycarbonyl (Troc), 2-
trimethylsilylethoxycarbonyl (Teoc), and
5 aryloxycarbonyl (Alloc).

10. A method for preparing the compound
represented by formula 1 comprising the following
steps as shown in the below reaction formula 1:

10 preparing the compound represented by formula 2A
by reacting the compound represented by formula 2 and
the compound represented by formula 3 (step 1);

preparing the compound represented by formula 5
by reacting the compound represented by formula 2A
15 prepared in step 1 and the compound represented by
formula 4 (step 2);

preparing the compound represented by formula 7
by reacting the compound represented by formula 5
prepared in step 2 and the compound represented by
20 formula 6 (step 3);

preparing the compound represented by formula 8
by reacting the compound represented by formula 7
prepared in step 3 and the compound represented by
formula 2B under basic condition (step 4);


preparing the compound represented by formula 10 by reacting the compound represented by formula 8 prepared in step 4 and the compound represented by formula 9 (step 5);

5 preparing the compound represented by formula 11 by reacting the compound represented by formula 10 prepared in step 5 under acidic condition (step 6);

 preparing the compound represented by formula 12 by reacting the compound represented by formula 11
10 prepared in step 6 and the compound represented by formula 2C (step 7); and

 preparing the compound represented by formula 1a by eliminating the amine protecting group from the compound represented by formula 12 prepared in step 7
15 under acidic condition (step 8):

[Reaction Formula 1]



In the reaction formula 1,

PG is amine protecting group;

the compound represented by formula 1a is a derivative of the compound represented by formula 1, in which --- is double bond and A is carbon, and R^1 , R^2 , R^3 , R^4 , and R^5 are as defined in formula 1.

11. A method for preparing the compound represented by formula 1 comprising the following steps as shown in the below reaction formula 2:

preparing the compound represented by formula 15 by reacting the compound represented by formula 2A and the compound represented by formula 14 (step 1);

preparing the compound represented by formula 17
by reacting the compound represented by formula 15
prepared in step 1 and the compound represented by
formula 16 (step 2);


5 preparing the compound represented by formula 18
by reacting the compound represented by formula 17
prepared in step 2 and the compound represented by
formula 2B (step 3);

preparing the compound represented by formula 20
10 by reacting the compound represented by formula 18
prepared in step 3 and the compound represented by
formula 19 (step 4);

preparing the compound represented by formula 21
by reacting the compound represented by formula 20
15 prepared in step 4 and the compound represented by
formula 2C under basic condition (step 5); and

preparing the compound represented by formula 1b
by eliminating the amine protecting group from the
compound represented by formula 21 prepared in step 5
20 under acidic condition (step 6):

[Reaction Formula 2]



In the reaction formula 2,

PG is amine protecting group;

the compound represented by formula 1b is a derivative of the compound represented by formula 1, in which --- is single bond and A is nitrogen, and R¹, R², R³, R⁴, and R⁵ are as defined in formula 1.


12. A method for preparing the compound represented by formula 1 comprising the following steps as shown in the below reaction formula 3:

preparing the compound represented by formula 15 by reacting the compound represented by formula 2A and the compound represented by formula 14 (step 1);

preparing the compound represented by formula 23 by reacting the compound represented by formula 15 prepared in step 1 and the compound represented by formula 22 (step 2); and

preparing the compound represented by formula 1c by reacting the compound represented by formula 23 prepared in step 2 and the compound represented by formula 2C under basic condition (step 3):

5 [Reaction Formula 3]



In the reaction formula 3,

the compound represented by formula 1c is a
 10 derivative of the compound represented by formula 1,
 in which --- is double bond and A is nitrogen, and R^1 ,
 R^2 , R^3 , R^4 , and R^5 are as defined in formula 1.

13. A pharmaceutical composition for the
 15 prevention or treatment of PI3 kinase related disease,
 which comprises the compound represented by formula 1,
 the optical isomer thereof, or the pharmaceutically
 acceptable salt thereof of claim 1 as an active
 ingredient, wherein the PI3 kinase related disease is
 20 selected from the group consisting of cancer;
 autoimmune disease; and respiratory disease selected
 from the group consisting of chronic obstructive

pulmonary disease(COPD), rhinitis, asthma, chronic
bronchitis, chronic inflammatory lung disease,
silicosis, pulmonary sarcoidosis, pleurisy,
alveolitis, vasculitis, emphysema, pneumonia, and
5 bronchiectasis.

14. The pharmaceutical composition for the
prevention or treatment of PI3 kinase related disease
according to claim 13, wherein the cancer is selected
10 from the group consisting of hematological malignance,
ovarian cancer, cervical cancer, breast cancer,
colorectal cancer, liver cancer, stomach cancer,
pancreatic cancer, colon cancer, peritoneal
metastasis, skin cancer, bladder cancer, prostate
15 cancer, thyroid cancer, lung cancer, osteosarcoma,
fibrous tumor, and brain tumor.

15. The pharmaceutical composition for the
prevention or treatment of PI3 kinase related disease
20 according to claim 13, wherein the autoimmune disease
is selected from the group consisting of rheumatoid
arthritis, systemic lupus erythematosus, multiple
sclerosis, type 1 diabetes, hyperthyroidism,
myasthenia, Crohn's disease, ankylosing spondylitis,

psoriasis, autoimmune pernicious anemia, and Sjogren's syndrome.