

NOVEL KINASE INHIBITORS

The present invention relates to the use of novel pyrimidine and pyridine derivatives which inhibit Btk and are useful for the treatment of auto-immune and inflammatory diseases caused by aberrant B-cell activation. The novel pyrimidine and pyridine derivatives described herein are useful for the treatment of arthritis.

5 Protein kinases constitute one of the largest families of human enzymes and regulate many different signaling processes by adding phosphate groups to proteins (Hunter, *Cell* 50:823-829 (1987)). Specifically, tyrosine kinases phosphorylate proteins on the phenolic moiety of tyrosine residues. The tyrosine kinase family includes members that control cell growth, migration, and differentiation. Abnormal kinase activity has been implicated in a variety of human diseases
10 including cancers, autoimmune and inflammatory diseases. Since protein kinases are among the key regulators of cell signaling they provide a target to modulate cellular function with small molecular kinase inhibitors and thus make good drug design targets. In addition to treatment of kinase-mediated disease processes, selective and efficacious inhibitors of kinase activity are also useful for investigation of cell signaling processes and identification of other cellular targets of
15 therapeutic interest.

There is good evidence that B-cells play a key role in the pathogenesis of autoimmune and/or inflammatory disease. Protein-based therapeutics that deplete B cells such as Rituxan are effective against autoantibody-driven inflammatory diseases such as rheumatoid arthritis [Rastetter *et al. Annu Rev Med* 55:477 (2004)]. Therefore inhibitors of the protein kinases that
20 play a role in B-cell activation should be useful therapeutics for B-cell mediated disease pathology such as autoantibody production.

Signaling through the B-cell receptor (BCR) controls a range of B-cell responses including proliferation and differentiation into mature antibody producing cells. The BCR is a key regulatory point for B-cell activity and aberrant signaling can cause deregulated B-cell
25 proliferation and formation of pathogenic autoantibodies that lead to multiple autoimmune and/or inflammatory diseases. Bruton's Tyrosine Kinase (Btk) is a non-BCR associated kinase that is membrane proximal and immediately downstream from BCR. Lack of Btk has been


shown to block BCR signaling and therefore inhibition of Btk could be a useful therapeutic approach to block B-cell mediated disease processes.

Btk is a member of the Tec family of tyrosine kinases, and has been shown to be a critical regulator of early B-cell development and mature B-cell activation and survival (Khan *et al.* 5 *Immunity* 3:283 (1995); Ellmeier *et al. J. Exp. Med.* 192:1611 (2000). Mutation of Btk in humans leads to the condition X-linked agammaglobulinemia (XLA) (reviewed in Rosen *et al. New Eng. J. Med.* 333:431 (1995) and Lindvall *et al. Immunol. Rev.* 203:200 (2005). These patients are immunocompromised and show impaired maturation of B-cells, decreased immunoglobulin and peripheral B-cell levels, diminished T-cell independent immune responses 10 as well as attenuated calcium mobilization following BCR stimulation.

Evidence for a role for Btk in autoimmune and inflammatory diseases has also been provided by Btk-deficient mouse models. In preclinical murine models of systemic lupus erythematosus (SLE), Btk-deficient mice show marked amelioration of disease progression. In addition, Btk-deficient mice are resistant to collagen-induced arthritis (Jansson and Holmdahl *Clin. Exp.* 15 *Immunol.* 94:459 (1993). A selective Btk inhibitor has been demonstrated dose-dependent efficacy in a mouse arthritis model (Pan *et al., Chem. Med Chem.* 2:58-61 (2007).

Btk is also expressed by cells other than B-cells that may be involved in disease processes. For example, Btk is expressed by mast cells and Btk-deficient bone marrow derived mast cells demonstrate impaired antigen induced degranulation (Iwaki *et al. J. Biol. Chem.* 280:40261 20 (2005). This shows Btk could be useful to treat pathological mast cells responses such as allergy and asthma. Also monocytes from XLA patients, in which Btk activity is absent, show decreased TNF alpha production following stimulation (Horwood *et al. J Exp Med* 197:1603 (2003). Therefore TNF alpha mediated inflammation could be modulated by small molecular Btk inhibitors. Also, Btk has been reported to play a role in apoptosis (Islam and Smith *Immunol.* 25 *Rev.* 178:49 (2000) and thus Btk inhibitors would be useful for the treatment of certain B-cell lymphomas and leukemias (Feldhahn *et al. J. Exp. Med.* 201:1837 (2005).

In a first aspect, there is provided a compound of formula I wherein:



(I)

wherein

X^1 is N;

R^1 is $C(=O)NHR^6$, phenyl or heteroaryl wherein said heteroaryl is selected from the group consisting of pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl; pyrrolyl, furanyl, thiophenyl, imidazolyl, pyrazolyl, oxazolyl, thiazolyl, isoxazolyl, isothiazolyl, 2- C_{1-6} alkyl-pyridazin-3-on-6-yl, 2-benzyl-pyridazin-3-on-6-yl, 1,4,5,6-tetrahydro-pyrimidin-2-yl and 4,5-dihydro-1H-imidazol-2-yl; said phenyl and said heteroaryl optionally independently substituted with one to three groups selected from the group consisting of halogen, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} heteroalkyl, C_{1-6} heteroalkoxy, C_{1-6} alkylsulfanyl, C_{1-6} alkylsulfonyl, C_{1-6} heteroalkylsulfonyl and heterocyclyl- C_{1-6} alkoxy wherein said heterocyclyl is azetindinyl, pyrrolidinyl or piperidinyl, $CONR^aR^b$, CO_2R^g , $SO_2NR^aR^b$, NR^aR^b , $NHSO_2R^7$ or $NHCOR^7$; wherein

R^6 is hydrogen, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{1-6} heteroalkyl, C_{3-6} cycloalkyl, pyridinyl- C_{1-3} alkyl, phenyl or phenyl- C_{1-3} alkyl said phenyl optionally substituted with C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} heteroalkoxy, halogen, $CONR^aR^b$ or CO_2R^g ;

R^a and R^b (i) selected independently are hydrogen, C_{1-6} alkyl, C_{1-6} heteroalkyl, C_{1-3} alkoxy- C_{1-3} alkyl, carboxy- C_{1-3} alkyl, C_{3-6} cycloalkyl or heterocyclyl; (ii) together are $(CH_2)_mX^2(CH_2)_2$ wherein m is 2 or 3 and X^2 is O, $S(O)_n$ or NR^8 and n is zero to two, and R^8 is hydrogen, C_{1-6} alkyl, C_{1-6} hydroxyalkyl or C_{1-3} acyl; or (iii) together with the nitrogen to which they are attached are piperidinyl or pyrrolidinyl said piperidinyl or said pyrrolidinyl optionally substituted with hydroxy, C_{1-3} alkoxy, C_{1-3} hydroxyalkyl;

R^g is hydrogen or C_{1-6} alkyl;

R^7 is C_{1-6} alkyl or C_{1-6} heteroalkyl;

R^2 is hydrogen or C_{1-6} alkyl;




R^3 and R^4 are independently hydrogen, halogen, amino, C_{1-6} hydroxyalkyl, C_{1-6} alkyl or cyclopropyl;

R^5 is (i) phenyl,

(ii) heteroaryl selected from the group consisting of pyridinyl, benzo[b]thiophen-2-yl, 4,5,6,7-tetrahydro-benzo[b]thiophen-2-yl, thiophenyl, 1',2',3',4',5',6'-hexahydro-[2,4']bipyridin-5-yl and C_{1-3} alkyl-indolyl optionally substituted with one or two C_{1-6} alkyl or halogen;

(iii) azetidin-1-yl, pyrrolidin-2-yl, piperidin-1-yl, azepan-1-yl or 2,3-dihydro-1H-isoindolin-2-yl, or 6,7-dihydro-5H-pyrrolo[3,4-b]pyridine;

(iv) NR^cR^d wherein R^c and R^d together are $(CH_2)_oX^2(CH_2)_p$ wherein o and p are independently 1 or 2, and X^2 is as defined above, or R^c and R^d independently are hydrogen C_{1-10} alkyl or C_{1-10} hydroxyalkyl;



wherein R^9 is hydrogen, halogen or C_{1-6} alkyl;

wherein said phenyl, heteroaryl, azetidin-1-yl, pyrrolidin-2-yl, piperidin-1-yl, azepan-1-yl or 2,3-dihydro-1H-isoindolin-2-yl ring is optionally substituted with one to three groups independently selected from the group consisting of (a) halogen, (b) C_{1-6} alkyl, (b) C_{2-6} alkenyl, (c) C_{2-6} alkynyl, (d) C_{1-6} haloalkyl, (e) C_{1-6} heteroalkyl, (f) C_{3-7} cycloalkyl optionally

substituted with one to three groups selected from C_{1-6} alkyl, C_{1-6} haloalkyl or halogen, (g) C_{1-6} alkoxyalkyl, (h) hydroxy, (i) NR^eR^f , wherein R^e and R^f are (i) independently hydrogen, C_{1-6} alkyl or (ii) together are $(CH_2)_mX^3(CH_2)_2$ wherein m is 2 or 3 and X^3 is CH_2 , O, $S(O)_n$ or NR^8 and n is zero to two and R^8 is as defined above; (j) C_{1-6} alkoxy, (k) trialkylsilyl, (l) C_{1-6} cyanoalkyl and (m) SF_5 ;


A is $-NHC(=O)-$, $C(=O)NH-$, $NHC(=O)NH$, $CH_2C(=O)$, CH_2SO_2 or if R^5 is (v) or (vi) A is absent;

or a pharmaceutically acceptable salt thereof.

In a second aspect, there is provided a pharmaceutical composition comprising a compound as defined above admixed with at least one pharmaceutically acceptable carrier, excipient or
5 diluent.

In a third aspect, there is provided the use of a compound of formula I as defined above for the preparation of a medicament for the treatment of an inflammatory and/or autoimmune condition.

10 The present invention relates to compounds of formula I



(I)

wherein

X¹ is CH or N;

R¹ is C(=O)NHR⁶, phenyl or heteroaryl wherein said heteroaryl is selected from the group consisting of pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl; pyrrolyl, furanyl, thiophenyl, imidazolyl, pyrazolyl, oxazolyl, thiazolyl, isoxazolyl, isothiazolyl, 2-C₁₋₆alkyl-pyridazin-3-on-6-yl, 2-benzyl-pyridazin-3-on-6-yl, 1,4,5,6-tetrahydro-pyrimidin-2-yl and 4,5-dihydro-1H-imidazol-2-yl, with the proviso that when X¹ is CH, R¹ is C(=O)NHR⁶ or R³ is CH₂OH;

said phenyl and said heteroaryl optionally independently substituted with one to three

groups selected from the group consisting of halogen, C₁₋₆alkyl, C₁₋₆alkoxy, C₁₋₆heteroalkyl, C₁₋₆heteroalkoxy, C₁₋₆alkylsulfanyl, C₁₋₆alkylsulfonyl, C₁₋₆heteroalkylsulfonyl and heterocyclyl-C₁₋₆alkoxy wherein said heterocyclyl is azetindinyl, pyrrolidinyl or piperidinyl, CONR^aR^b, CO₂R^g, SO₂NR^aR^b, NR^aR^b, NHSO₂R⁷ or NHCOR⁷; wherein

R⁶ is hydrogen, C₁₋₆alkyl, C₁₋₆haloalkyl, C₁₋₆heteroalkyl, C₃₋₆cycloalkyl, pyridinyl-

C₁₋₃alkyl, phenyl or phenyl-C₁₋₃alkyl said phenyl optionally substituted with C₁₋₆alkyl,

C₁₋₆alkoxy, C₁₋₆heteroalkoxy, halogen, CONR^aR^b or CO₂R^g;

R^a and R^b (i) selected independently are hydrogen, C₁₋₆alkyl, C₁₋₆heteroalkyl, C₁₋₃alkoxy-

C₁₋₃alkyl, carboxy-C₁₋₃alkyl, C₃₋₆cycloalkyl or heterocyclyl; (ii) together are

(CH₂)_mX²(CH₂)₂ wherein m is 2 or 3 and X² is O, S(O)_n or NR⁸ and n is zero to two,

and R⁸ is hydrogen, C₁₋₆alkyl, C₁₋₆hydroxyalkyl or C₁₋₃acyl; or (iii) together with the

nitrogen to which they are attached are piperidinyl or pyrrolidinyl said piperidinyl or

said pyrrolidinyl optionally substituted with hydroxy, C₁₋₃alkoxy, C₁₋₃hydroxyalkyl;

R^g is hydrogen or C₁₋₆alkyl;

R⁷ is C₁₋₆alkyl or C₁₋₆heteroalkyl;

R² is hydrogen or C₁₋₆alkyl;




R³ and R⁴ are independently hydrogen, halogen, amino, C₁₋₆hydroxyalkyl, C₁₋₆alkyl or cyclopropyl;

R⁵ is (i) phenyl,

(ii) heteroaryl selected from the group consisting of pyridinyl, benzo[b]thiophen-2-yl, 4,5,6,7-tetrahydro-benzo[b]thiophen-2-yl, thiophenyl, 1',2',3',4',5',6'-hexahydro-[2,4']bipyridin-5-yl and C₁₋₃alkyl-indolyl optionally substituted with one or two C₁₋₆alkyl or halogen;

(iii) azetidin-1-yl, pyrrolidin-2-yl, piperidin-1-yl, azepan-1-yl or 2,3-dihydro-1H-isoindolin-2-yl, or 6,7-dihydro-5H-pyrrolo[3,4-b]pyridine;

(iv) NR^cR^d wherein R^c and R^d together are (CH₂)_oX²(CH₂)_p wherein o and p are independently 1 or 2, and X² is as defined above, or R^c and R^d independently are hydrogen C₁₋₁₀alkyl or C₁₋₁₀hydroxyalkyl;



wherein R⁹ is hydrogen, halogen or C₁₋₆alkyl;

wherein said phenyl, heteroaryl, azetidin-1-yl, pyrrolidin-2-yl, piperidin-1-yl, azepan-1-yl or 2,3-dihydro-1H-isoindolin-2-yl ring is optionally substituted with one to three groups independently selected from the group consisting of (a) halogen, (b) C₁₋₆alkyl, (b) C₂₋₆alkenyl, (c) C₂₋₆alkynyl, (d) C₁₋₆haloalkyl, (e) C₁₋₆heteroalkyl, (f) C₃₋₇cycloalkyl optionally substituted with one to three groups selected from C₁₋₆alkyl, C₁₋₆haloalkyl or halogen, (g) C₁₋₆alkoxyalkyl, (h) hydroxy, (i) NR^eR^f, wherein R^e and R^f are (i) independently hydrogen, C₁₋₆alkyl or (ii) together are (CH₂)_mX³(CH₂)₂ wherein m is 2 or 3 and X³ is CH₂, O, S(O)_n or NR⁸ and n is zero to two and R⁸ is as defined above; (j) C₁₋₆alkoxy, (k) trialkylsilyl, (l) C₁₋₆cyanoalkyl and (m) SF₅;

A is -NHC(=O)-, C(=O)NH-, NHC(=O)NH-, CH₂C(=O)-, CH₂SO₂ or if R⁵ is (v) or (vi) A is absent;

or a pharmaceutically acceptable salt thereof.

In one embodiment the present invention relates to compounds of formula I wherein

X^1 is CH or N;

R^1 is $C(=O)NHR^6$, phenyl or heteroaryl wherein said heteroaryl is selected from the group consisting of pyridin-2-yl, pyridazinyl, pyrimidinyl, pyrazinyl, pyrrolyl, furanyl, thiophenyl, imidazolyl, pyrazolyl, oxazolyl, thiazolyl, isoxazolyl, isothazolyl, 1,4,5,6-tetrahydro-pyrimidin-2-yl and 4,5-dihydro-1H-imidazol-2-yl, with the proviso that when X^1 is CH, R^1 is not phenyl; said heteroaryl and said phenyl optionally independently substituted with one to three groups selected from the group consisting of halogen, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} heteroalkyl, C_{1-6} heteroalkoxy, C_{1-6} alkylsulfanyl, C_{1-6} alkylsulfonyl, C_{1-6} heteroalkylsulfonyl, heterocyclyl- C_{1-6} alkoxy wherein said heterocycle is azetindinyl, pyrrolidinyl or piperidinyl, $CONR^aR^b$, CO_2R^g , $SO_2NR^aR^b$, NR^aR^b , $NHSO_2R^7$ or $NHCOR^7$;

R^2 and R^g are independently hydrogen or C_{1-6} alkyl;



R^3 and R^4 are independently hydrogen, halogen, amino, C_{1-6} hydroxyalkyl, C_{1-6} alkyl or cyclopropyl;

R^5 is (i) phenyl,

(ii) heteroaryl selected from the group consisting of pyridinyl, benzo[b]thiophen-2-yl, 4,5,6,7-tetrahydro-benzo[b]thiophen-2-yl and thiophenyl;

(iii) azetidin-1-yl, pyrrolidin-2-yl, piperidin-1-yl, azepan-1-yl or 2,3-dihydro-1H-isoindolin-2-yl;

(iv) NR^cR^d wherein R^c and R^d together are $(CH_2)_oX^3(CH_2)_p$ wherein o and p are independently is 1 or 2, and X^3 is O, $S(O)_n$, NR^8 and n is 0 to 2, or R^c and R^d independently are hydrogen C_{1-10} alkyl or C_{1-10} hydroxyalkyl;



wherein said phenyl, heteroaryl, azetidin-1-yl, pyrrolidin-2-yl, piperidin-1-yl or azepan-1-yl ring is optionally substituted with one to three groups independently selected from the group consisting of halogen, C_{1-6} alkyl, C_{1-6} heteroalkyl, hydroxy, NR^eR^f and C_{1-6} alkoxy; wherein R^e and R^f are (i) independently hydrogen, C_{1-6} alkyl or (ii) together are $(CH_2)_mX^3(CH_2)_2$ wherein m is 2 or 3 and X^3 is O, $S(O)_n$, NR^8 and n is zero to two;

R^6 is hydrogen, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{1-6} heteroalkyl, C_{3-6} cycloalkyl, phenyl or benzyl said benzyl or phenyl optionally substituted with C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} heteroalkoxy, halogen, $CONR^aR^b$, CO_2R^c ;

A is $-NHC(=O)-$, $C(=O)NH-$, $NHC(=O)NH-$, $CH_2C(=O)-$, CH_2SO_2- or if R^5 is (v) or (vi) A is absent;

R^a and R^b (i) selected independently are hydrogen, C_{1-6} alkyl, C_{1-6} heteroalkyl, C_{1-3} alkoxy C_{1-3} alkyl, carboxy C_{1-3} alkyl, C_{3-6} cycloalkyl or heterocyclyl (ii) together are $(CH_2)_mX^3(CH_2)_2$ wherein m is 2 or 3 and X^3 is O, $S(O)_n$, NR^8 and n is zero to two, or (iii) together with the nitrogen to which they are attached are piperidine or pyrrolidine said piperidine or said pyrrolidine optionally substituted with hydroxy, C_{1-3} alkoxy, C_{1-3} hydroxyalkyl;

R^7 is C_{1-6} alkyl or C_{1-6} heteroalkyl;

R^8 is hydrogen, C_{1-6} alkyl, C_{1-6} hydroxyalkyl or C_{1-3} acyl;

R^9 is hydrogen, halogen or C_{1-6} alkyl; or

pharmaceutically acceptable salts thereof.

The compounds of formula I inhibit Bruton's tyrosine kinase (Btk). Activation of Btk by upstream kinases results in activation of phospholipase-C γ which, in turn, stimulates release of pro-inflammatory mediators. Compounds of formula I are useful in the treatment of arthritis and other anti-inflammatory and auto-immune diseases. Compounds according to formula I are, accordingly, useful for the treatment of arthritis. Compounds of formula I are useful for inhibiting Btk in cells and for modulating B-cell development. The present invention further comprises pharmaceutical compositions containing compounds of formula I admixed with pharmaceutically acceptable carriers, excipients or diluents.

The phrase "a" or "an" entity as used herein refers to one or more of that entity; e.g., a compound refers to one or more compounds or at least one compound. As such, the terms "a" (or "an"), "one or more", and "at least one" can be used interchangeably herein.

The phrase "as defined herein above" refers to the broadest definition for each group as provided herein or the broadest claim. In all other embodiments provided below, substituents which can be present in each embodiment and which are not explicitly defined retain the broadest definition provided herein.

As used in this specification, whether in a transitional phrase or in the body of the claim, the terms "comprise(s)" and "comprising" are to be interpreted as having an open-ended meaning.


That is, the terms are to be interpreted synonymously with the phrases "having at least" or "including at least". When used in the context of a process, the term "comprising" means that the process includes at least the recited steps, but may include additional steps. When used in the context of a compound or composition, the term "comprising" means that the compound or composition includes at least the recited features or components, but may also include additional features or components.

As used herein, unless specifically indicated otherwise, the word "or" is used in the "inclusive" sense of "and/or" and not the "exclusive" sense of "either/or".

The term "independently" is used herein to indicate that a variable is applied in any one instance without regard to the presence or absence of a variable having that same or a different definition within the same compound. Thus, in a compound in which R" appears twice and is defined as "independently carbon or nitrogen", both R"s can be carbon, both R"s can be nitrogen, or one R" can be carbon and the other nitrogen.

When any variable (e.g., R¹, R^{4a}, Ar, X¹ or Het) occurs more than one time in any moiety or formula depicting and describing compounds employed or claimed in the present invention, its definition on each occurrence is independent of its definition at every other occurrence. Also, combinations of substituents and/or variables are permissible only if such compounds result in stable compounds.

The symbols "*" at the end of a bond or "-----" drawn through a bond each refer to the point of attachment of a functional group or other chemical moiety to the rest of the molecule of which it is a part. Thus, e.g.:



A bond drawn into ring system (as opposed to connected at a distinct vertex) indicates that the bond may be attached to any of the suitable ring atoms.

The term "optional" or "optionally" as used herein means that a subsequently described event or circumstance may, but need not, occur, and that the description includes instances where the event or circumstance occurs and instances in which it does not. For example, "optionally substituted" means that the optionally substituted moiety may incorporate a hydrogen or a substituent.

The phrase "optional bond" means that the bond may or may not be present, and that the description includes single, double, or triple bonds. If a substituent is designated to be a "bond" or "absent", the atoms linked to the substituents are then directly connected.

The term "about" is used herein to mean approximately, in the region of, roughly, or around.

- 5 When the term "about" is used in conjunction with a numerical range, it modifies that range by extending the boundaries above and below the numerical values set forth. In general, the term "about" is used herein to modify a numerical value above and below the stated value by a variance of 20%.

- The definitions described herein may be appended to form chemically-relevant combinations, such as "heteroalkylaryl", "haloalkylheteroaryl", "arylalkylheterocyclyl", "alkylcarbonyl", "alkoxyalkyl" and the like. When the term "alkyl" is used as a suffix following another term, as in "phenylalkyl" or "hydroxyalkyl" this is intended to refer to an alkyl group, as defined above, being substituted with one to two substituents selected from the other specifically-named group. Thus, e.g., "phenylalkyl" refers to an alkyl group having one to two phenyl substituents, and thus includes benzyl, phenylethyl, and biphenyl. An "alkylaminoalkyl" is an alkyl group having one to two alkylamino substituents. "Hydroxyalkyl" includes 2-hydroxyethyl, 2-hydroxypropyl, 1-(hydroxymethyl)-2-methylpropyl, 2-hydroxybutyl, 2,3-dihydroxybutyl, 2-(hydroxymethyl), 3-hydroxypropyl, and so forth. Accordingly, as used herein, the term "hydroxyalkyl" is used to define a subset of heteroalkyl groups defined below. The term -(ar)alkyl refers to either an unsubstituted alkyl or an aralkyl group. The term (hetero)aryl or (het)aryl refers to either an aryl or a heteroaryl group.
- 10
15
20

- Compounds of formula I exhibit tautomerism. Tautomeric compounds can exist as two or more interconvertible species. Prototropic tautomers result from the migration of a covalently bonded hydrogen atom between two atoms. Tautomers generally exist in equilibrium and attempts to isolate an individual tautomer usually produce a mixture whose chemical and physical properties are consistent with a mixture of compounds. The position of the equilibrium is dependent on chemical features within the molecule. For example, in many aliphatic aldehydes and ketones, such as acetaldehyde, the keto form predominates while; in phenols, the enol form predominates. Common prototropic tautomers include keto/enol ($-C(=O)-CH- \leftrightarrow -C(OH)=CH-$), amide/imidic acid ($-C(=O)-NH- \leftrightarrow -C(OH)=N-$) and amidine ($-C(=NR)-NH- \leftrightarrow -C(NHR)=N-$) tautomers. The latter two are particularly common in heteroaryl and heterocyclic rings and the present invention encompasses all tautomeric forms of the compounds.
- 25
30

Technical and scientific terms used herein have the meaning commonly understood by one of skill in the art to which the present invention pertains, unless otherwise defined. Reference is made herein to various methodologies and materials known to those of skill in the art. Standard reference works setting forth the general principles of pharmacology include Goodman and
5 Gilman's *The Pharmacological Basis of Therapeutics*, 10th Ed., McGraw Hill Companies Inc., New York (2001). Any suitable materials and/or methods known to those of skill can be utilized in carrying out the present invention. However, preferred materials and methods are described. Materials, reagents and the like to which reference are made in the following description and examples are obtainable from commercial sources, unless otherwise noted.

- 10 In one embodiment of the present invention there is provided a compound according to formula I wherein R^1 , R^2 , R^3 , R^4 , R^5 , X^1 and A are as defined herein above.

In a second embodiment of the present invention there is provided a compound according to formula I wherein X^1 is N and R^1 , R^2 , R^3 , R^4 , R^5 and A are as defined herein above.

- 15 In a third embodiment of the present invention there is provided a compound according to formula I wherein X^1 is N; R^1 is phenyl, pyridin-2-yl or pyrimidin-4-yl; R^2 is C_{1-3} alkyl; R^3 is hydrogen or C_{1-3} alkyl; A is $NHC(=O)$; and R^4 and R^5 are as defined herein above.

In another embodiment of the present invention there is provided a compound according to formula I wherein X^1 is N; R^1 is phenyl, pyridin-2-yl or pyrimidin-4-yl; R^2 is C_{1-3} alkyl; R^3 is C_{1-6} hydroxyalkyl; A is $NHC(=O)$; and R^4 and R^5 are as defined herein above.

- 20 In a fourth embodiment of the present invention there is provided a compound according to formula I wherein X^1 is N; R^1 is phenyl, pyridin-2-yl or pyrimidin-4-yl; R^2 is C_{1-3} alkyl; R^3 is hydroxymethyl; A is $NHC(=O)$; and R^4 and R^5 are as defined herein above.

- In a fifth embodiment of the present invention there is provided a compound according to formula I wherein X^1 is N; R^1 is phenyl, pyridin-2-yl or pyrimidin-4-yl; R^2 is C_{1-3} alkyl; R^3 is
25 hydroxymethyl; A is $NHC(=O)$; R^5 is optionally substituted phenyl, optionally substituted pyridinyl, benzo[b]thiophen-2-yl or 4,5,6,7-tetrahydro-benzo[b]thiophen-2-yl; and R^4 is as defined herein above.

- In a sixth embodiment of the present invention there is provided a compound according to formula I wherein X^1 is N; R^1 is pyridin-2-yl optionally substituted with $CONR^aR^b$, NR^aR^b , C_{1-6} alkoxy, C_{1-6} heteroalkoxy, C_{1-6} alkylsulfanyl or C_{1-6} alkylsulfonyl; R^2 is C_{1-3} alkyl; R^3 is
30

hydroxymethyl; A is NHC(=O); R⁵ is optionally substituted phenyl, optionally substituted pyridinyl, benzo[b]thiophen-2-yl or 4,5,6,7-tetrahydro-benzo[b]thiophen-2-yl; and R⁴ is as defined herein above.

In another embodiment of the present invention there is provided a compound according to
5 formula I wherein X¹ is N; R¹ is pyridin-2-yl optionally substituted with CONR^aR^b, NR^aR^b, C₁₋₆alkoxy, C₁₋₆heteroalkoxy, C₁₋₆alkylsulfanyl or C₁₋₆alkylsulfonyl; R² is C₁₋₃alkyl; R³ is hydrogen or C₁₋₃alkyl; A is NHC(=O); R⁵ is optionally substituted phenyl, optionally substituted pyridinyl, benzo[b]thiophen-2-yl or 4,5,6,7-tetrahydro-benzo[b]thiophen-2-yl; and R⁴ is as defined herein above.

10 In a seventh embodiment of the present invention there is provided a compound according to formula I wherein X¹ is N; R¹ is pyrimidin-4-yl optionally substituted with CONR^aR^b, NR^aR^b, C₁₋₆alkoxy, C₁₋₆heteroalkoxy, C₁₋₆alkylsulfanyl or C₁₋₆alkylsulfonyl; R² is C₁₋₃alkyl; R³ is hydroxymethyl; A is NHC(=O); R⁵ is optionally substituted phenyl, optionally substituted pyridinyl, benzo[b]thiophen-2-yl or 4,5,6,7-tetrahydro-benzo[b]thiophen-2-yl; and R⁴ is as
15 defined herein above.

In another embodiment of the present invention there is provided a compound according to formula I wherein X¹ is N; R¹ is pyrimidin-4-yl optionally substituted with CONR^aR^b, NR^aR^b, C₁₋₆alkoxy, C₁₋₆heteroalkoxy, C₁₋₆alkylsulfanyl or C₁₋₆alkylsulfonyl; R² is C₁₋₃alkyl; R³ is hydrogen or C₁₋₃alkyl; A is NHC(=O); R⁵ is optionally substituted phenyl, optionally substituted
20 pyridinyl, benzo[b]thiophen-2-yl or 4,5,6,7-tetrahydro-benzo[b]thiophen-2-yl; and R⁴ is as defined herein above.

In an eighth embodiment of the present invention there is provided a compound according to formula I wherein X¹ is N; R¹ is phenyl optionally substituted with CONR^aR^b, NR^aR^b, C₁₋₆alkoxy, C₁₋₆heteroalkoxy, C₁₋₆alkylsulfanyl or C₁₋₆alkylsulfonyl; R² is C₁₋₃alkyl; R³ is
25 hydroxymethyl; A is NHC(=O); R⁵ is optionally substituted phenyl, optionally substituted pyridinyl, benzo[b]thiophen-2-yl or 4,5,6,7-tetrahydro-benzo[b]thiophen-2-yl; and R⁴ is as defined herein above.

In a another embodiment of the present invention there is provided a compound according to formula I wherein X¹ is N; R¹ is phenyl optionally substituted with CONR^aR^b, NR^aR^b,
30 C₁₋₆alkoxy, C₁₋₆heteroalkoxy, C₁₋₆alkylsulfanyl or C₁₋₆alkylsulfonyl; R² is C₁₋₃alkyl; R³ is hydrogen or C₁₋₃alkyl; A is NHC(=O); R⁵ is optionally substituted phenyl, optionally substituted

pyridinyl, benzo[b]thiophen-2-yl or 4,5,6,7-tetrahydro-benzo[b]thiophen-2-yl; and R⁴ is as defined herein above.

In a ninth embodiment of the present invention there is provided a compound according to formula I wherein X¹ is N; R¹ is C(=O)NHR⁶; R⁶ is C₁₋₆alkyl; R³ is hydrogen, C₁₋₃alkyl or
5 C₁₋₆heteroalkyl; A is NHC(=O) and R², R⁴ and R⁵ are as defined herein above.

In a tenth embodiment of the present invention there is provided a compound according to formula I wherein X¹ is N; R¹ is C(=O)NHR⁶; R⁶ is C₁₋₆alkyl; R³ is hydroxymethyl; A is NHC(=O) and R², R⁴ and R⁵ are as defined herein above.

In an eleventh embodiment of the present invention there is provided a compound according to
10 formula I wherein X¹ is CH and R¹, R², R³, R⁴, R⁵ and A are as defined herein above.

In a twelfth embodiment of the present invention there is provided a compound according to formula I wherein X¹ is CH; R¹ is pyridin-2-yl or pyrimidin-4-yl; R² is C₁₋₃alkyl; R³ is hydroxymethyl; A is NHC(=O); and R⁴ and R⁵ are as defined herein above.

In a thirteenth embodiment of the present invention there is provided a compound according to
15 formula I wherein X¹ is CH; R¹ is pyridin-2-yl or pyrimidin-4-yl; R² is C₁₋₃alkyl; R³ is hydroxymethyl; R⁵ is optionally substituted phenyl, optionally substituted pyridinyl, benzo[b]thiophen-2-yl or 4,5,6,7-tetrahydro-benzo[b]thiophen-2-yl; A is NHC(=O); and R⁴ is as defined herein above.

In a fourteenth embodiment of the present invention there is provided a compound according to
20 formula I wherein X¹ is CH; R¹ is pyridin-2-yl optionally substituted with CONR^aR^b, NR^aR^b, C₁₋₆alkoxy, C₁₋₆heteroalkoxy, C₁₋₆alkylsulfanyl or C₁₋₆alkylsulfonyl; R³ is hydroxymethyl; R⁵ is optionally substituted phenyl, optionally substituted pyridinyl, benzo[b]thiophen-2-yl or 4,5,6,7-tetrahydro-benzo[b]thiophen-2-yl; A is NHC(=O); and R⁴ is as defined herein above.

In a fifteenth embodiment of the present invention there is provided a compound according to
25 formula I wherein X¹ is CH; R¹ is pyrimidin-4-yl optionally substituted with CONR^aR^b, NR^aR^b, C₁₋₆alkoxy, C₁₋₆heteroalkoxy, C₁₋₆alkylsulfanyl or C₁₋₆alkylsulfonyl; R³ is hydroxymethyl; R⁵ is optionally substituted phenyl, optionally substituted pyridinyl, benzo[b]thiophen-2-yl or 4,5,6,7-tetrahydro-benzo[b]thiophen-2-yl; A is NHC(=O); and R⁴ is as defined herein above.

In a sixteenth embodiment of the present invention there is provided a compound according to formula I wherein X^1 is CH and R^1 is $C(=O)NHR^6$, R^6 is C_{1-6} alkyl, R^3 is hydrogen, C_{1-3} alkyl or C_{1-6} heteroalkyl; A is $NHC(=O)$, and R^2 , R^4 , R^5 are as defined herein above.

In a seventeenth embodiment of the present invention there is provided a compound according to
 5 formula I wherein X^1 is CH and R^1 is $C(=O)NHR^6$, R^6 is C_{1-6} alkyl, R^3 is hydroxymethyl; A is $NHC(=O)$, and R^2 , R^4 , R^5 are as defined herein above.

In an eighteenth embodiment of the present invention there is provided a compound according to formula I wherein R^5 is (v) or (vi); A is absent; and R^1 , R^2 , R^3 , R^4 and X^1 are as defined herein above.

10 In a nineteenth embodiment of the present invention there is provided a compound according to formula I wherein A is $CH_2C(=O)$ and R^5 is (i), (ii), (iii) or (iv); and R^1 , R^2 , R^3 , R^4 and X^1 are as defined herein above.

In a twentieth embodiment of the present invention there is provided a compound selected from compounds I-1 to I-46 of TABLE I or from II-4 to II-19, II-21 to II-23, II-27 to II-31, II-45 to II-
 15 47 and II-49 to II-87 of TABLE II.

In another embodiment of the present invention there is provided a compound of formula I selected from the group consisting of:

4-*tert*-butyl-*N*-(2-methyl-3-{1-methyl-5-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-6-oxo-1,6-dihydro-pyridazin-3-yl}-phenyl)-benzamide;

20 4-*tert*-butyl-*N*-(2-methyl-3-{1-methyl-5-[5-(4-methyl-piperazine-1-carbonyl)-pyridin-2-ylamino]-6-oxo-1,6-dihydro-pyridazin-3-yl}-phenyl)-benzamide;

4-*tert*-butyl-*N*-(2-methyl-3-[1-methyl-6-oxo-5-(pyridin-2-ylamino)-1,6-dihydro-pyridazin-3-yl]-phenyl)-benzamide;

25 4-*tert*-butyl-*N*-(2-methyl-3-[1-methyl-6-oxo-5-(pyrimidin-4-ylamino)-1,6-dihydro-pyridazin-3-yl]-phenyl)-benzamide;

4-(1-hydroxy-1-methyl-ethyl)-*N*-(2-methyl-3-{1-methyl-5-[5-(4-methyl-piperazine-1-carbonyl)-pyridin-2-ylamino]-6-oxo-1,6-dihydro-pyridazin-3-yl}-phenyl)-benzamide;

4-*tert*-butyl-piperazine-1-carboxylic acid (2-methyl-3-{1-methyl-5-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-6-oxo-1,6-dihydro-pyridazin-3-yl}-phenyl)-amide;

30 4-*tert*-butyl-2-methoxy-*N*-(2-methyl-3-{1-methyl-5-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-6-oxo-1,6-dihydro-pyridazin-3-yl}-phenyl)-benzamide;

- 7-*tert*-butyl-3-(2-methyl-3-{1-methyl-5-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-6-oxo-1,6-dihydro-pyridazin-3-yl}-phenyl)-3*H*-quinazolin-4-one;
- 6-{6-[3-(4-*tert*-butyl-benzoylamino)-2-methyl-phenyl]-2-methyl-3-oxo-2,3-dihydro-pyridazin-4-ylamino}-nicotinic acid methyl ester;
- 5 3-*tert*-butoxy-azetidine-1-carboxylic acid (2-methyl-3-{1-methyl-5-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-6-oxo-1,6-dihydro-pyridazin-3-yl}-phenyl)-amide;
- 4-*tert*-butyl-*N*-(2-methyl-3-{1-methyl-5-[2-(4-methyl-piperazin-1-yl)-pyrimidin-4-ylamino]-6-oxo-1,6-dihydro-pyridazin-3-yl}-phenyl)-benzamide;
- 4-*tert*-butyl-*N*-(2-methyl-3-[1-methyl-5-(2-methylsulfanyl-pyrimidin-4-ylamino)-6-oxo-1,6-dihydro-pyridazin-3-yl]-phenyl)-benzamide
- 10 4-*tert*-butyl-*N*-(3-[5-(2-methanesulfonyl-pyrimidin-4-ylamino)-1-methyl-6-oxo-1,6-dihydro-pyridazin-3-yl]-2-methyl-phenyl)-benzamide;
- 4-*tert*-butyl-*N*-(2-methyl-3-[1-methyl-5-(2-morpholin-4-yl-pyrimidin-4-ylamino)-6-oxo-1,6-dihydro-pyridazin-3-yl]-phenyl)-benzamide;
- 15 4-(1-hydroxy-1-methyl-ethyl)-*N*-(2-methyl-3-{1-methyl-5-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-6-oxo-1,6-dihydro-pyridazin-3-yl}-phenyl)-benzamide;
- 4-(1-hydroxy-1-methyl-ethyl)-*N*-(3-{1-methyl-5-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-6-oxo-1,6-dihydro-pyridazin-3-yl}-phenyl)-benzamide;
- 4-*tert*-butyl-*N*-(3-[5-(2-methoxy-pyrimidin-4-ylamino)-1-methyl-6-oxo-1,6-dihydro-pyridazin-3-yl]-2-methyl-phenyl)-benzamide;
- 20 4-*tert*-butyl-*N*-(3-{5-[2-(2-dimethylamino-ethoxy)-pyrimidin-4-ylamino]-1-methyl-6-oxo-1,6-dihydro-pyridazin-3-yl}-2-methyl-phenyl)-benzamide;
- 4-*tert*-butyl-*N*-(2-methyl-3-{1-methyl-6-oxo-5-[2-(pyrrolidin-3-ylmethoxy)-pyrimidin-4-ylamino]-1,6-dihydro-pyridazin-3-yl}-phenyl)-benzamide;
- 25 4-*tert*-butyl-*N*-(3-{5-[2-(3-hydroxymethyl-pyrrolidin-1-yl)-pyrimidin-4-ylamino]-1-methyl-6-oxo-1,6-dihydro-pyridazin-3-yl}-2-methyl-phenyl)-benzamide;
- 4-*tert*-butyl-*N*-(2-methyl-3-[1-methyl-6-oxo-5-(2-pyrrolidin-1-yl-pyrimidin-4-ylamino)-1,6-dihydro-pyridazin-3-yl]-phenyl)-benzamide;
- 4-*tert*-butyl-*N*-(3-{5-[2-(3-hydroxy-pyrrolidin-1-yl)-pyrimidin-4-ylamino]-1-methyl-6-oxo-1,6-dihydro-pyridazin-3-yl}-2-methyl-phenyl)-benzamide;
- 30 4-*tert*-butyl-*N*-(2-methyl-3-[1-methyl-5-(3-methyl-ureido)-6-oxo-1,6-dihydro-pyridazin-3-yl]-phenyl)-benzamide;
- 4-*tert*-butyl-*N*-(2-methyl-3-{1-methyl-5-[4-(morpholine-4-carbonyl)-phenylamino]-6-oxo-1,6-dihydro-pyridazin-3-yl}-phenyl)-benzamide;

- 4-*tert*-butyl-*N*-(3-{1-methyl-5-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-6-oxo-1,6-dihydro-pyridazin-3-yl}-phenyl)-benzamide;
- 4-*tert*-butyl-*N*-(3-{5-[4-(4-hydroxy-piperidine-1-carbonyl)-phenylamino]-1-methyl-6-oxo-1,6-dihydro-pyridazin-3-yl}-2-methyl-phenyl)-benzamide;
- 5 4-*tert*-butyl-*N*-(3-[5-(3-ethyl-ureido)-1-methyl-6-oxo-1,6-dihydro-pyridazin-3-yl]-2-methyl-phenyl)-benzamide;
- 4-dimethylamino-*N*-(3-{5-[5-(4-hydroxy-piperidine-1-carbonyl)-pyridin-2-ylamino]-1-methyl-6-oxo-1,6-dihydro-pyridazin-3-yl}-2-methyl-phenyl)-benzamide;
- N*-(3-Hydroxy-4,4-dimethyl-pentyl)-2-(3-{1-methyl-5-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-6-oxo-1,6-dihydro-pyridazin-3-yl}-phenyl)-acetamide;
- 10 4-*tert*-butyl-2-hydroxy-*N*-(2-methyl-3-{1-methyl-5-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-6-oxo-1,6-dihydro-pyridazin-3-yl}-phenyl)-benzamide;
- 6-{3-[2-(1,3-dihydro-isoindol-2-yl)-2-oxo-ethyl]-phenyl}-4-[5-(4-hydroxy-piperidine-1-carbonyl)-pyridin-2-ylamino]-2-methyl-2*H*-pyridazin-3-one
- 15 *N*-(2-*tert*-butoxy-ethyl)-2-(3-{1-methyl-5-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-6-oxo-1,6-dihydro-pyridazin-3-yl}-phenyl)-acetamide;
- 2-methyl-4-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-6-[3-(2-oxo-2-pyrrolidin-1-yl-ethyl)-phenyl]-2*H*-pyridazin-3-one;
- 7-*tert*-butyl-3-(2-methyl-3-{1-methyl-5-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-6-oxo-1,6-dihydro-pyridazin-3-yl}-phenyl)-2,3-dihydro-1*H*-quinazolin-4-one;
- 20 6-{3-[2-(3-isopropoxy-azetidin-1-yl)-2-oxo-ethyl]-phenyl}-2-methyl-4-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-2*H*-pyridazin-3-one;
- (3-{1-methyl-5-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-6-oxo-1,6-dihydro-pyridazin-3-yl}-phenyl)-acetic acid 2-*tert*-butoxy-ethyl ester;
- 25 6-{3-[2-(4-*tert*-butyl-phenyl)-2-oxo-ethyl]-phenyl}-2-methyl-4-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-2*H*-pyridazin-3-one;
- 6-{3-[2-(4-*tert*-butyl-piperazin-1-yl)-2-oxo-ethyl]-phenyl}-2-methyl-4-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-2*H*-pyridazin-3-one;
- 6-{3-[2-(3-*tert*-butoxy-azetidin-1-yl)-2-oxo-ethyl]-phenyl}-2-methyl-4-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-2*H*-pyridazin-3-one;
- 30 6-{3-[2-(1,3-dihydro-isoindol-2-yl)-2-oxo-ethyl]-phenyl}-2-methyl-4-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-2*H*-pyridazin-3-one;
- 6-{3-[2-(4-isopropyl-piperazin-1-yl)-2-oxo-ethyl]-phenyl}-2-methyl-4-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-2*H*-pyridazin-3-one;

- 6- {3-[2-(4-*tert*-butyl-piperidin-1-yl)-2-oxo-ethyl]-phenyl}-2-methyl-4-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-2*H*-pyridazin-3-one;
- N*-(3,3-dimethyl-butyl)-2-(3-{1-methyl-5-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-6-oxo-1,6-dihydro-pyridazin-3-yl}-phenyl)-acetamide;
- 5 6- {3-[2-(4-acetyl-piperazin-1-yl)-2-oxo-ethyl]-phenyl}-2-methyl-4-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-2*H*-pyridazin-3-one;
- 4-*tert*-butyl-*N*-(2-methyl-3-[1-methyl-5-(3-methyl-ureido)-6-oxo-1,6-dihydro-pyridin-3-yl]-phenyl)-benzamide;
- 4-*tert*-butyl-*N*-(3-[5-(3-ethyl-ureido)-1-methyl-6-oxo-1,6-dihydro-pyridin-3-yl]-2-methyl-phenyl)-benzamide;
- 10 4-*tert*-butyl-*N*-(3-[5-(3-isopropyl-ureido)-1-methyl-6-oxo-1,6-dihydro-pyridin-3-yl]-2-methyl-phenyl)-benzamide;
- 4-*tert*-butyl-*N*-(2-methyl-3-{1-methyl-6-oxo-5-[3-(2,2,2-trifluoro-ethyl)-ureido]-1,6-dihydro-pyridin-3-yl}-phenyl)-benzamide;
- 15 4-*tert*-butyl-*N*-(2-methyl-3-[1-methyl-6-oxo-5-(3-phenyl-ureido)-1,6-dihydro-pyridin-3-yl]-phenyl)-benzamide;
- N*-(3-[5-(3-benzyl-ureido)-1-methyl-6-oxo-1,6-dihydro-pyridin-3-yl]-2-methyl-phenyl)-4-*tert*-butyl-benzamide;
- benzo[*b*]thiophene-2-carboxylic acid {2-methyl-3-[1-methyl-5-(3-methyl-ureido)-6-oxo-1,6-dihydro-pyridin-3-yl]-phenyl}-amide;
- 20 4,5,6,7-tetrahydro-benzo[*b*]thiophene-2-carboxylic acid {2-methyl-3-[1-methyl-5-(3-methyl-ureido)-6-oxo-1,6-dihydro-pyridin-3-yl]-phenyl}-amide;
- 1-{5-[3-(7-*tert*-butyl-4-oxo-4*H*-quinazolin-3-yl)-2-methyl-phenyl]-1-methyl-2-oxo-1,2-dihydro-pyridin-3-yl}-3-methyl-urea;
- 25 4-dimethylamino-*N*-(2-methyl-3-[1-methyl-5-(3-methyl-ureido)-6-oxo-1,6-dihydro-pyridin-3-yl]-phenyl)-benzamide;
- 5-fluoro-1,3-dihydro-isoindole-2-carboxylic acid {2-methyl-3-[1-methyl-5-(3-methyl-ureido)-6-oxo-1,6-dihydro-pyridin-3-yl]-phenyl}-amide;
- 6-dimethylamino-*N*-(2-methyl-3-[1-methyl-5-(3-methyl-ureido)-6-oxo-1,6-dihydro-pyridin-3-yl]-phenyl)-nicotinamide;
- 30 4-*tert*-butyl-*N*-(3-{5-[3-(2-dimethylamino-ethyl)-ureido]-1-methyl-6-oxo-1,6-dihydro-pyridin-3-yl}-2-methyl-phenyl)-benzamide;
- 1,3-dihydro-isoindole-2-carboxylic acid {2-methyl-3-[1-methyl-5-(3-methyl-ureido)-6-oxo-1,6-dihydro-pyridin-3-yl]-phenyl}-amide;

- 4-*tert*-butyl-*N*-(3-{5-[3-(2-hydroxy-ethyl)-ureido]-1-methyl-6-oxo-1,6-dihydro-pyridin-3-yl}-2-methyl-phenyl)-benzamide;
- 3,4,5,6-tetrahydro-2*H*-[1,2']bipyridinyl-5'-carboxylic acid {2-methyl-3-[1-methyl-5-(3-methyl-ureido)-6-oxo-1,6-dihydro-pyridin-3-yl]-phenyl}-amide;
- 5 6-{5-[3-(4-dimethylamino-benzoylamino)-2-methyl-phenyl]-1-methyl-2-oxo-1,2-dihydro-pyridin-3-ylamino}-*N*-(2-dimethylamino-ethyl)-nicotinamide;
- 4-*tert*-butyl-*N*-(3-{5-[3-(3-dimethylamino-propyl)-ureido]-1-methyl-6-oxo-1,6-dihydro-pyridin-3-yl}-2-methyl-phenyl)-benzamide
- 1-methyl-1*H*-indole-5-carboxylic acid {2-methyl-3-[1-methyl-5-(3-methyl-ureido)-6-oxo-1,6-dihydro-pyridin-3-yl]-phenyl}-amide
- 10 4-*tert*-butyl-*N*-(3-[1-methyl-5-(3-methyl-ureido)-6-oxo-1,6-dihydro-pyridin-3-yl]-phenyl)-benzamide;
- 4-(1-hydroxy-1-methyl-ethyl)-*N*-(2-methyl-3-[1-methyl-5-(3-methyl-ureido)-6-oxo-1,6-dihydro-pyridin-3-yl]-phenyl)-benzamide;
- 15 1-(5-{3-[2-(3-*tert*-butoxy-azetidin-1-yl)-2-oxo-ethyl]-phenyl}-1-methyl-2-oxo-1,2-dihydro-pyridin-3-yl)-3-methyl-urea;
- 1-(5-{3-[2-(3-isopropoxy-azetidin-1-yl)-2-oxo-ethyl]-phenyl}-1-methyl-2-oxo-1,2-dihydro-pyridin-3-yl)-3-methyl-urea;
- 1-(5-{3-[2-(5,6-dichloro-1,3-dihydro-isoindol-2-yl)-2-oxo-ethyl]-phenyl}-1-methyl-2-oxo-1,2-dihydro-pyridin-3-yl)-3-methyl-urea;
- 20 4-(cyano-dimethyl-methyl)-*N*-(2-methyl-3-[1-methyl-5-(3-methyl-ureido)-6-oxo-1,6-dihydro-pyridin-3-yl]-phenyl)-benzamide;
- 1-(5-{3-[2-(1,3-dihydro-isoindol-2-yl)-2-oxo-ethyl]-phenyl}-1-methyl-2-oxo-1,2-dihydro-pyridin-3-yl)-3-methyl-urea;
- 25 1-(5-{3-[2-(5-fluoro-1,3-dihydro-isoindol-2-yl)-2-oxo-ethyl]-phenyl}-1-methyl-2-oxo-1,2-dihydro-pyridin-3-yl)-3-methyl-urea; and,
- 1-[5-(3-{2-[3-(2-hydroxy-2-methyl-propyl)-azetidin-1-yl]-2-oxo-ethyl}-phenyl)-1-methyl-2-oxo-1,2-dihydro-pyridin-3-yl]-3-methyl-urea.

In a twenty-first embodiment of the present invention there is provided a method for treating an inflammatory and/or autoimmune condition comprising administering to a patient in need thereof a compound according of formula I wherein R¹, R², R³, R⁴, R⁵, X¹ and A are as defined herein above.

In a twenty-second embodiment of the present invention there is provided a method for treating rheumatoid arthritis comprising administering to a patient in need thereof a compound according to formula I wherein R^1 , R^2 , R^3 , R^4 , R^5 , X^1 and A are as defined herein above.

5 In a twenty-third embodiment of the present invention there is provided a method of inhibiting B-cell proliferation comprising administering to a patient in need thereof a compound according to formula I wherein R^1 , R^2 , R^3 , R^4 , R^5 , X^1 and A are as defined herein above.

10 In a twenty-fourth embodiment of the present invention there is provided a method for inhibiting Btk activity comprising administering to a patient in need thereof a compound according to formula I wherein R^1 , R^2 , R^3 , R^4 , R^5 , X^1 and A are as defined herein above wherein the compound exhibits an IC_{50} of 1 micromolar or less in an *in vitro* biochemical assay of Btk activity.

15 In a twenty-fifth embodiment of the present invention there is provided a method for inhibiting Btk activity comprising administering to a patient in need thereof a compound according to formula I wherein R^1 , R^2 , R^3 , R^4 , R^5 , X^1 and A are as defined herein above wherein the compound exhibits an IC_{50} of 100 nanomolar or less in an *in vitro* biochemical assay of Btk activity.

20 In a twenty-sixth embodiment of the present invention there is provided a method for inhibiting Btk activity comprising co-administering to a patient in need thereof a compound according to formula I wherein R^1 , R^2 , R^3 , R^4 , R^5 , X^1 and A are as defined herein above and a therapeutically effective amount of another anti-inflammatory compound.

25 In a twenty-seventh embodiment of the present invention there is provided a method for treating an inflammatory condition, e.g. rheumatoid arthritis, comprising co-administering to a patient in need thereof a therapeutically effective amount of an anti-inflammatory compound along with a compound according to formula I wherein R^1 , R^2 , R^3 , R^4 , R^5 , X^1 and A are as defined herein above.

In a twenty-eighth embodiment of the present invention there is provided a pharmaceutical composition comprising a compound according to formula I wherein R^1 , R^2 , R^3 , R^4 , R^5 , X^1 and A are as defined herein above admixed with at least one pharmaceutically acceptable carrier, excipient or diluent.

The term "acyl" as used herein denotes a group of formula $-C(=O)R$ wherein R is hydrogen or lower alkyl as defined herein. The term "alkylcarbonyl" as used herein denotes a group of formula $C(=O)R$ wherein R is alkyl as defined herein. The term C_{1-6} acyl refers to a group $-C(=O)R$ wherein R is hydrogen or C_{1-5} alkyl. The term "arylcarbonyl" as used herein means a group of formula $C(=O)R$ wherein R is an aryl group; the term "benzoyl" as used herein an "arylcarbonyl" group wherein R is phenyl.

The term "alkyl" as used herein denotes an unbranched or branched chain, saturated, monovalent hydrocarbon residue containing 1 to 10 carbon atoms. The term "lower alkyl" denotes a straight or branched chain hydrocarbon residue containing 1 to 6 carbon atoms. " C_{1-10} alkyl" as used herein refers to an alkyl composed of 1 to 10 carbons. Examples of alkyl groups include, but are not limited to, methyl, ethyl, propyl, *i*-propyl, *n*-butyl, *i*-butyl, *t*-butyl or pentyl, isopentyl, neopentyl, hexyl, heptyl, and octyl.

When the term "alkyl" is used as a suffix following another term, as in "phenylalkyl," or "hydroxyalkyl," this is intended to refer to an alkyl group, as defined above, being substituted with one to two substituents selected from the other specifically-named group. Thus, e.g., "phenylalkyl" denotes the radical $R'R^-$, wherein R' is a phenyl radical, and R" is an alkylene radical as defined herein with the understanding that the attachment point of the phenylalkyl moiety will be on the alkylene radical. Examples of arylalkyl radicals include, but are not limited to, benzyl, phenylethyl, 3-phenylpropyl. The terms "arylalkyl" or "aralkyl" are interpreted similarly except R' is an aryl radical. The terms "(het)arylalkyl" or "(het)aralkyl" are interpreted similarly except R' is optionally an aryl or a heteroaryl radical.

The term "haloalkyl" as used herein denotes an unbranched or branched chain alkyl group as defined above wherein 1, 2, 3 or more hydrogen atoms are substituted by a halogen. Examples are 1-fluoromethyl, 1-chloromethyl, 1-bromomethyl, 1-iodomethyl, difluoromethyl, trifluoromethyl, trichloromethyl, tribromomethyl, triiodomethyl, 1-fluoroethyl, 1-chloroethyl, 1-bromoethyl, 1-iodoethyl, 2-fluoroethyl, 2-chloroethyl, 2-bromoethyl, 2-iodoethyl, 2,2-dichloroethyl, 3-bromopropyl or 2,2,2-trifluoroethyl.

The term "alkylene" as used herein denotes a divalent saturated linear hydrocarbon radical of 1 to 10 carbon atoms (e.g., $(CH_2)_n$) or a branched saturated divalent hydrocarbon radical of 2 to 10 carbon atoms (e.g., $-CHMe-$ or $-CH_2CH(i-Pr)CH_2-$), unless otherwise indicated. Except in the case of methylene, the open valences of an alkylene group are not attached to the same atom.

Examples of alkylene radicals include, but are not limited to, methylene, ethylene, propylene, 2-methyl-propylene, 1,1-dimethyl-ethylene, butylene, 2-ethylbutylene.

The term "alkoxy" as used herein means an -O-alkyl group, wherein alkyl is as defined above such as methoxy, ethoxy, *n*-propyloxy, *i*-propyloxy, *n*-butyloxy, *i*-butyloxy, *t*-butyloxy, pent-
5 yloxy, hexyloxy, including their isomers. "Lower alkoxy" as used herein denotes an -O-alkyl group wherein alkyl is "lower alkyl" as previously defined. "C₁₋₁₀alkoxy" as used herein refers to an-O-alkyl wherein alkyl is C₁₋₁₀alkyl.

The term "alkoxyalkyl" as used herein refers to the radical R'R"-, wherein R' is an alkoxy radical as defined herein, and R" is an alkylene radical as defined herein with the understanding that the
10 attachment point of the alkoxyalkyl moiety will be on the alkylene radical. C₁₋₆alkoxyalkyl denotes a group wherein the alkyl portion is comprised of 1-6 carbon atoms exclusive of carbon atoms in the alkoxy portion of the group. C₁₋₃alkoxy-C₁₋₆alkyl denotes a group wherein the alkyl portion is comprised of 1-6 carbon atoms and the alkoxy group is 1-3 carbons. Examples are
15 methoxymethyl, methoxyethyl, methoxypropyl, ethoxymethyl, ethoxyethyl, ethoxypropyl, propyloxypropyl, methoxybutyl, ethoxybutyl, propyloxybutyl, butyloxybutyl, *t*-butyloxybutyl, methoxypentyl, ethoxypentyl, propyloxypentyl including their isomers.

The term "heteroalkoxy" as used herein means an -O-(heteroalkyl) group wherein heteroalkyl is as defined herein. "C₁₋₁₀heteroalkoxy" as used herein refers to an -O-(C₁₋₁₀heteroalkyl) wherein
20 C₁₋₁₀heteroalkyl is as defined herein. Representative examples include, but are not limited to, 2-dimethylaminoethoxy and 3-sulfonamido-1-propoxy.

The term "heteroalkyl" as used herein refers to an alkyl radical as defined herein wherein one, two or three hydrogen atoms have been replaced with a substituent independently selected from the group consisting of -OR^a, -NR^bR^c, and -S(O)_nR^d (where n is an integer from 0 to 2), with the
25 understanding that the point of attachment of the heteroalkyl radical is through a carbon atom, and wherein R^a is hydrogen, acyl, alkyl, cycloalkyl, or cycloalkylalkyl; R^b and R^c are independently of each other hydrogen, acyl, alkyl, cycloalkyl, or cycloalkylalkyl; and when n is 0, R^d is hydrogen, alkyl, cycloalkyl, or cycloalkylalkyl, and when n is 1 or 2, R^d is alkyl, cycloalkyl, cycloalkylalkyl, amino, alkylamino, dialkylamino or phenyl. Representative
30 examples include, but are not limited to, 2-hydroxyethyl, 3-hydroxypropyl, 2-hydroxy-1-hydroxymethylethyl, 2,3-dihydroxypropyl, 1-hydroxymethylethyl, 3-hydroxybutyl, 2,3-dihydroxybutyl, 2-hydroxy-1-methylpropyl, 2-aminoethyl, 3-aminopropyl, 2-methylsulfonylethyl,

aminosulfonylmethyl, aminosulfonylethyl, aminosulfonylpropyl, methylaminosulfonylmethyl, methylaminosulfonylethyl, methylaminosulfonylpropyl, and the like.

The term "cyanoalkyl" as used herein refers to an alkyl radical as defined herein wherein one hydrogen atom has been replaced with a cyano substituent.

- 5 The term "hydroxyalkyl" as used herein denotes an alkyl radical as herein defined wherein one to three hydrogen atoms on different carbon atoms is/are replaced by hydroxyl groups.

- The term "alkylthio" or "alkylsulfanyl" refers to an -S-alkyl group, wherein alkyl is as defined above such as methylthio, ethylthio, *n*-propylthio, *i*-propylthio, *n*-butylthio, hexylthio, including their isomers. "Lower alkylthio" as used herein denotes an alkylthio group with a "lower alkyl" group as previously defined. "C₁₋₁₀alkylthio" as used herein refers to an-S-alkyl wherein alkyl is C₁₋₁₀. "Phenylthio" is an "arylthio" moiety wherein aryl is phenyl.
- 10

The terms "alkylcarbonylamino" and "arylcarbonylamino" as used herein refer to a group of formula -NC(=O)R wherein R is alkyl or aryl respectively and alkyl and aryl are as defined herein.

- 15 The terms "alkylsulfinyl" and "arylsulfinyl" as used herein refer to a group of formula -S(=O)R wherein R is alkyl or aryl respectively and alkyl and aryl are as defined herein.

- The terms "alkylsulfonyl" and "arylsulfonyl" as used herein refer to a group of formula -S(=O)₂R wherein R is alkyl or aryl respectively and alkyl and aryl are as defined herein. The term "heteroalkylsulfonyl" as used herein denotes a group of formula -S(=O)₂R wherein R is "heteroalkyl" as defined herein.
- 20

The terms "alkylsulfonylamino" and "arylsulfonylamino" as used herein refer to a group of formula -NR'S(=O)₂R wherein R is alkyl or aryl respectively, R' is hydrogen or C₁₋₃alkyl, and alkyl and aryl are as defined herein.

- The term "cycloalkyl" as used herein refers to a saturated carbocyclic ring containing 3 to 8 carbon atoms, i.e. cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or cyclooctyl. "C₃₋₇cycloalkyl" as used herein refers to a cycloalkyl composed of 3 to 7 carbons in the carbocyclic ring.
- 25


The term carboxy-alkyl as used herein refers to an alkyl moiety wherein one hydrogen atom has been replaced with a carboxyl with the understanding that the point of attachment of the heteroalkyl radical is through a carbon atom. The term carboxyl refers to a $-\text{CO}_2\text{H}$ moiety.

The term "halogen" or "halo" as used herein means fluorine, chlorine, bromine, or iodine.


- 5 The terms azetidine, pyrrolidine, piperidine and azepine refer to a four, five, six and seven membered saturated ring containing one nitrogen atom in the ring. The "1"-position refers to the nitrogen atom.

The terms benzo[b]thiophen-2-yl, 4,5,6,7-tetrahydro-benzo[b]thiophen-2-yl, thiophenyl 2,3-dihydro-1H-isoindolin-2-yl, 1',2',3',4',5',6'-hexahydro-[2,4']bipyridin-5-yl, and 6,7-dihydro-5H-

- 10 pyrrolo[3,4-b]pyridine refer to moieties (i) to (vi) respectively.



The terms "1,4,5,6-tetrahydro-pyrimidin-2-ylamine" and "4,5-dihydro-1H-imidazol-2-ylamine" refer to the moieties (vii) and (viii) respectively.



- 15 The term "heteroaryl" or "heteroaromatic" as used herein means a monocyclic or bicyclic radical of 5 to 12 ring atoms having at least one aromatic ring containing four to eight atoms per ring, incorporating one or more N, O, or S heteroatoms, the remaining ring atoms being carbon, with the understanding that the attachment point of the heteroaryl radical will be on an aromatic ring. As well known to those skilled in the art, heteroaryl rings have less aromatic character than their
- 20 all-carbon counter parts. Thus, for the purposes of the invention, a heteroaryl group need only have some degree of aromatic character. Examples of heteroaryl moieties include monocyclic aromatic heterocycles having 5 to 6 ring atoms and 1 to 3 heteroatoms include, but is not limited to, pyridinyl, pyrimidinyl, pyrazinyl, pyrrolyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, triazolyl, thiadiazolyl and oxadiazolyl which can optionally be
- 25 substituted with one or more, preferably one or two substituents selected from hydroxy, cyano, alkyl, alkoxy, thio, lower haloalkoxy, alkylthio, halo, haloalkyl, alkylsulfinyl, alkylsulfonyl, halogen, amino, alkylamino, dialkylamino, aminoalkyl, alkylaminoalkyl, and dialkylaminoalkyl,

nitro, alkoxycarbonyl and carbamoyl, alkylcarbamoyl, dialkylcarbamoyl, arylcarbamoyl, alkylcarbonylamino and arylcarbonylamino. Examples of bicyclic moieties include, but are not limited to, quinolinyl, isoquinolinyl, benzofuryl, benzothiophenyl, benzoxazolyl, benzisoxazolyl, benzothiazolyl and benzisothiazolyl. Bicyclic moieties can be optionally substituted on either
5 ring; however the point of attachment is on a ring containing a heteroatom.

The term (hetero)aryl as used herein refers to an aryl or a heteroaryl moiety as each is defined herein.

The term "heterocyclyl" or "heterocycle" as used herein denotes a monovalent saturated cyclic radical, consisting of one or more rings, preferably one to two rings, of three to eight atoms per
10 ring, incorporating one or more ring heteroatoms (chosen from N, O or S(O)₀₋₂), and which can optionally be independently substituted with one or more, preferably one or two substituents selected from hydroxy, oxo, cyano, lower alkyl, lower alkoxy, lower haloalkoxy, alkylthio, halo, haloalkyl, hydroxyalkyl, nitro, alkoxycarbonyl, amino, alkylamino, alkylsulfonyl, arylsulfonyl, alkylaminosulfonyl, arylaminosulfonyl, alkylsulfonylamino, arylsulfonylamino,
15 alkylaminocarbonyl, arylaminocarbonyl, alkylcarbonylamino, arylcarbonylamino, unless otherwise indicated. Examples of heterocyclic radicals include, but are not limited to, azetidiny, pyrrolidinyl, hexahydroazepinyl, oxetanyl, tetrahydrofuranyl, tetrahydrothiophenyl, oxazolidinyl, thiazolidinyl, isoxazolidinyl, morpholinyl, piperazinyl, piperidinyl, tetrahydropyranyl, thiomorpholinyl, quinuclidinyl and imidazolinyl.

20 The term "heterocyclyl-C₁₋₆alkoxy" as used herein refers to the radical R'R"-, wherein R' is an alkoxy radical as defined herein, wherein one, two of the hydrogen atoms attached to a carbon has been replaced with a heterocyclic substituent and heterocyclyl-C₁₋₆alkoxy moiety is attached at the oxygen atom of the alkoxy group.

The term "N-alkyl-ureido" as used herein refers to a group RNHC(=O)NH- wherein R is an alkyl
25 group as defined herein.

Commonly used abbreviations include: acetyl (Ac), azo-*bis*-isobutyronitrile (AIBN), atmospheres (Atm), 9-borabicyclo[3.3.1]nonane (9-BBN or BBN), *tert*-butoxycarbonyl (Boc), di-*tert*-butyl pyrocarbonate or boc anhydride (BOC₂O), benzyl (Bn), butyl (Bu), Chemical Abstracts Registration Number (CASRN), benzyloxycarbonyl (CBZ or Z), carbonyl diimidazole (CDI),
30 1,4-diazabicyclo[2.2.2]octane (DABCO), diethylaminosulfur trifluoride (DAST), di-benzylideneacetone (dba), 1,5-diazabicyclo[4.3.0]non-5-ene (DBN), 1,8-diazabicyclo[5.4.0]-

undec-7-ene (DBU), N,N'-dicyclohexylcarbodiimide (DCC), 1,2-dichloroethane (DCE), dichloromethane (DCM), diethyl azodicarboxylate (DEAD), di-*iso*-propylazodicarboxylate (DIAD), di-*iso*-butylaluminumhydride (DIBAL or DIBAL-H), di-*iso*-propylethylamine (DIPEA), N,N-dimethyl acetamide (DMA), 4-N,N-dimethylaminopyridine (DMAP), N,N-di-

5 methylformamide (DMF), dimethyl sulfoxide (DMSO), 1,1'-*bis*-(diphenylphosphino)ethane (dppe), 1,1'-*bis*-(diphenylphosphino)ferrocene (dppf), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDCI), ethyl (Et), ethyl acetate (EtOAc), ethanol (EtOH), 2-ethoxy-2*H*-quinoline-1-carboxylic acid ethyl ester (EEDQ), diethyl ether (Et₂O), O-(7-azabenzotriazole-1-yl)-N, N,N',N'-tetramethyluronium hexafluorophosphate acetic acid (HATU),

10 acetic acid (HOAc), 1-N-hydroxybenzotriazole (HOBT), high pressure liquid chromatography (HPLC), *iso*-propanol (IPA), lithium hexamethyl disilazane (LiHMDS), methanol (MeOH), melting point (mp), MeSO₂- (mesyl or Ms), , methyl (Me), acetonitrile (MeCN), *m*-chloroperbenzoic acid (MCPBA), mass spectrum (ms), methyl *t*-butyl ether (MTBE), N-bromosuccinimide (NBS), N-carboxyanhydride (NCA), N-chlorosuccinimide (NCS), N-methyl-

15 morpholine (NMM), N-methylpyrrolidone (NMP), pyridinium chlorochromate (PCC), pyridinium dichromate (PDC), phenyl (Ph), propyl (Pr), *iso*-propyl (*i*-Pr), pounds per square inch (psi), pyridine (pyr), room temperature (rt or RT), *tert*-butyldimethylsilyl or *t*-BuMe₂Si (TBDMS), triethylamine (TEA or Et₃N), 2,2,6,6-tetramethylpiperidine 1-oxyl (TEMPO), triflate or CF₃SO₂- (Tf), trifluoroacetic acid (TFA), 1,1'-*bis*-2,2,6,6-tetramethylheptane-2,6-dione

20 (TMHD), O-benzotriazol-1-yl-N,N,N',N'-tetramethyluronium tetrafluoroborate (TBTU), thin layer chromatography (TLC), tetrahydrofuran (THF), trimethylsilyl or Me₃Si (TMS), *p*-toluenesulfonic acid monohydrate (TsOH or pTsOH), 4-Me-C₆H₄SO₂- or tosyl (Ts), N-urethane-N-carboxyanhydride (UNCA),. Conventional nomenclature including the prefixes *normal* (*n*), *iso* (*i*-), *secondary* (*sec*-), *tertiary* (*tert*-) and *neo* have their customary meaning when used with

25 an alkyl moiety. (Rigaudy and Klesney, *Nomenclature in Organic Chemistry*, IUPAC 1979 Pergamon Press, Oxford.).





Examples of representative compounds encompassed by the present invention and within the scope of the invention are provided in the following Table. These examples and preparations which follow are provided to enable those skilled in the art to more clearly understand and to

30 practice the present invention. They should not be considered as limiting the scope of the invention, but merely as being illustrative and representative thereof.







In general, the nomenclature used in this Application is based on AUTONOMTM v.4.0, a Beilstein Institute computerized system for the generation of IUPAC systematic nomenclature.






If there is a discrepancy between a depicted structure and a name given that structure, the depicted structure is to be accorded more weight. In addition, if the stereochemistry of a structure or a portion of a structure is not indicated with, e.g., bold or dashed lines, the structure or portion of the structure is to be interpreted as encompassing all stereoisomers of it.







5 TABLE I: examples of pyridazinone compounds according to formula I wherein X¹ is N.







Cpd. #	Structure	mw	ms	mp	IC ₅₀ ^I (μ M)
I-1		580	581	148.0-150.0	<0.01
I-2		593	594	158.0-160.0	0.014
I-3		467	468	248.5-249.3	0.094
I-4		468	469	249.0-250.0	0.208







I-5		595	596	167.5-168.5	0.122
I-6		588	589	166.0-169.0	15.5
I-7		610	611	156.5-157.0	0.272
I-8		605	606	188.5-191.0	0.116
I-9		525	526		0.16
I-10		575	576	171.0-172.0	0.228






I-11		566	567	220.0-222.0	0.078
I-12		514	515	224.0-225.0	0.051
I-13		546	547		1.064
I-14		553	554	211.0-213.0	0.249
I-15		582	583		0.374
I-16		568	569	172.0-173.0	0.12

I-17		498	499	262.0-264.0	0.104
I-18		555	556	188.0-189.0	0.326
I-19		567	568		0.143
I-20		567	568	210.0-211.0	0.125
I-21		537	538	193.0-195.0	0.354

I-22		553	554		0.052
I-23		447	448	287.0-288.0	0.101
I-24		579	580	157.0-158.0	0.11
I-25		566	567	229.0-230.0	0.12
I-26		593	594		0.045
I-27		461	462	278.0-279.0	0.122

I-28		581	582		0.034
I-29		562	563		1.715
I-30		596	597	216.0- 217.0	0.059
I-31		564	565		0.155
I-32		548	549	220.0- 221.0	0.82
I-33		502	503	117.0- 118.0	16.35

I-34		607	608		
I-35		546	547	92.0-93.0	0.418
I-36		549	550	80.0-81.0	2.2
I-37		565	566		0.206
I-38		573	574	234.0-236.0	33.5
I-39		560	561	134.0-135.0	0.138






I-40		550	551	179.0-180.0	0.68
I-41		559	560	209.0-210.0	6.245
I-42		572	573	231.0-232.0	2.345
I-43		532	533	138.0-139.0	6.29
I-44		559	560		61.6






I-45		470	471		
I-46		580	581		<0.01






1. Bruton's Tyrosine Kinase Assay (Example 34)






TABLE II: examples of pyridinone compounds according to formula I wherein X¹ is CH.






Cpd. #	Structure	mw	ms	mp	IC ₅₀ ¹ (μ M)
II-4		446	447	262.0-264.0	
II-5		460	461	215.0-216.0	0.013
II-6		474	475	185.0-186.0	0.308






II-7		514	515	258.0-259.0	0.867
II-8		508	509	219.0-220.0	7.325
II-9		522	523	169.0-170.0	1.24
II-10		446	447	286.0-287.0	0.131
II-11		450	451	254.0-255.0	0.04






II-12		471	472		0.112
II-13		433	434	291.0-292.0	0.096
II-14		449	450	262.0-263.0	1.95
II-15		434	435		2.31
II-16		503	504	164.0-165.0	0.129






II-17		431	432	264.0-265.0	0.666
II-18		476	477	220.0-221.0	0.038
II-19		474	475		0.31
II-21		567	568		0.039
II-22		517	518	194.0-195.0	0.029






II-23		443	444		9.93
II-27		432	433		0.131
II-28		448	449		0.23
II-29		426	427	125.0-126.0	1.405
II-30		412	413	118.0-119.0	4.37






II-31		485	486		14.85
II-42		457	458		0.681
II-45		416	417	251.0-252.0	1.563
II-46		434	435		3.44
II-47		426	427	113.0-114.0	54.65






II-49		582	583	<0.01
II-50		446	447	<0.01
II-51		635	636	
II-52		599	600	<0.01
II-53		597	598	






II-54		490	491		0.068
II-55		607	608		0.085
II-56		476	477		<0.01
II-57		474	475	201.0-203.0	<0.01
II-58		595	596		<0.01

II-59		469	470	0.02
II-60		623	624	<0.01
II-61		463	464	-
II-62		579	580	-
II-63		618	618,620	0.031

II-64		484	485		0.05
II-65		478	479		0.015
II-66		611	612		0.024
II-67		539	540		0.77
II-68		707	708		0.071






II-69		563	564		0.076
II-70		569	570	203.0-204.0	<0.01
II-71		609	610		<0.01
II-72		629	630		<0.01
II-73		649	650		






II-74		656	656,658	0.05
II-75		614	615	<0.01
II-76		611	612	0.04
II-77		460	461	0.04
II-78		537	538	0.214


II-79		537	538	0.16
II-80		517	518	0.043
II-81		648	649	<0.01
II-82		676	677	
II-83		559	560	0.054

II-84		516	517	0.06
II-85		504	505	0.046
II-86		490	491	0.068
II-87		579	580	
II-88		525	526	0.14

II-89		665	666	<0.01
II-90		510	511	0.081
II-91		511	512	0.066
II-92		472	473	0.034
II-93		573	574	0.298

II-94		481	482	0.024
II-95		467	468	0.02
II-96		606	607	<0.01
II-97		622	623	0.114
II-98		467	468	0.09


II-99		563	564	0.011
II-100		585	586	0.031
II-101		549	550	<0.01
II-102		580	581	<0.01
II-103		552	553	0.014

II-104		565	566		
1. Bruton's Tyrosine Kinase Assay (Example 34)					

Compounds of the present invention that are 2-amino(hetero)aryl-4-aryl-pyridazinone (TABLE I) derivatives are prepared utilizing a two step sequence starting from a 4,6-dihalo-2-alkyl-2H-pyridazin-3-one, *e.g.*, A-1b. Displacement of the 2-halo moiety with a heterocyclic amine results in the introduction of the (hetero)aryl amine into the 6-position. The heterocyclic amine is optionally substituted with other groups within the scope of the invention or with a moiety which can be converted into desired substituents. Thus in SCHEME A the pyrimidine is 2-methylsulfanyl-pyrimidin-4-ylamine (A-5) The methylsulfanyl moiety can be oxidized to the corresponding sulfone which is easily displaced by nucleophiles in a subsequent step. The 4-substituted phenyl moiety is introduced by a Suzuki type coupling of a 4-halo-pyridazinone and a boronic acid derivative such as 4-(1,1-dimethylethyl)-N-[2-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]-benzamide (A-6, CASRN 910235-65-3) which affords A-3a. One skilled in the art will appreciate that a wide variety of boronic acids are available including 4,4,5,5-tetramethyl-2-(3-nitro-phenyl)-[1,3,2]dioxaborolane (CASRN 68716-48-3) or 4,4,5,5-tetramethyl-2-(2-methyl-3-nitro-phenyl)-[1,3,2]dioxaborolane (C-1, CASRN 910235-64-2) which can undergo reduction of the nitro group and functionalization of the resulting amine to produce amides, ureas and sulfonamides. Other useful boronic acid derivatives which are described in the examples which follow. Elaboration of the phenyl substituents can occur either prior to or after the Suzuki coupling. Thus, *e.g.*, the Suzuki condensation can be carried out with 2-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenylamine (**14a**, CASRN 882678-96-8) or 3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenylamine (**14b**) which can be acylated after the palladium-catalyzed coupling step (example 5). Coupling of [3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid (CASRN 797755-05-6 allows introduction of the amine moiety after the Suzuki condensation (example 8). Compounds of the present invention with a 3H-quinazolin-4-one moiety (*e.g.*, **I-8**) were prepared from **38** which is prepared by sequential condensation of 4-*tert*-butyl-anthranilic acid with trimethylorthoformate and **14a** to afford **38** which is utilized in the Suzuki coupling. Compounds with a 2,3-dihydro-

1H-quinazolin-4-one moiety (e.g., **I-34**) are prepared by reduction of the corresponding 3H-quinazolin-4-one.

SCHEME A




The Suzuki reaction (Miyama and Suzuki, Chem Rev. 1995 95:2457-2483; Suzuki, J.

- 5 Organomet. Chem. 1999 576:147-168) is a palladium-catalyzed coupling of a boronic acid with an aryl or vinyl halide or triflate. The reaction is postulated to proceed by an oxidative-addition mechanism. Pd(II) compounds used as catalysts are believed to be reduced to the catalytically active Pd(0) species in situ. Typical catalysts bis-(*tri*-*o*-tolylphosphine)-palladium(II) chloride, (dba)₃Pd₂(0)/tris-*o*-tolylphosphine, (dba)₃Pd₂(0)/tris-(2-furyl)phosphine, (dba)₃Pd₂(0)/2,2'-bis-
- 10 (diphenylphosphino)-1,1'-binaphthyl, Pd(0)(PPh₃)₄, Pd(0)Cl₂(dppf) or Pd(II)(OAc)₂/1,3-bis-(triphenylphosphino)-propane. The reaction is often carried out in the presence of a base such as sodium-*tert*-butoxide, bis-(trimethylsilyl)-lithium amide, K₂CO₃, Cs₂CO₃ or TEA. The reaction can be carried out in a variety of organic solvents including toluene, THF, dioxane, DCE, DMF, DMSO and MeCN, aqueous solvents and under biphasic conditions. Reactions are typically run
- 15 from about RT to about 150°C. Additives such as CsF, KF, TIOH, NaOEt and KOH frequently accelerate the coupling. Although there are numerous components in the Suzuki reaction including the palladium source, the ligand, additive solvent, temperature, etc., numerous protocols have been identified. Recently useful general conditions have been disclosed. Litke et al. J. Am. Chem. Soc. 2000 122:4020-4028 disclose conditions for Suzuki cross-coupling with
- 20 arylboronic acids in high yield at RT utilizing Pd₂(dba)₃/P(*tert*-bu)₃ and conditions for cross-coupling of aryl- and vinyl triflates utilizing Pd(OAc)₂/P(C₆H₁₁)₃ at RT. Wolf et al. (J. Am. Chem. Soc. 1999 121:9550-9561) disclose efficient condition for Suzuki cross-coupling utilizing

Pd(OAc)₂/o-(di-tert-butylphosphino)biphenyl or o-(dicyclohexylphosphino)biphenyl. One skilled in the art will be able to identify a satisfactory protocol without undue experimentation. Oxidation of a thiol to a sulfone is typically facile. Sulfur oxidations are commonly carried out with aqueous solution of hydrogen peroxide, NaIO₄, tert-butylhypochlorite, acyl nitrites, sodium perborate potassium hydrogen persulfate and peracids such as peracetic acid and meta-chloroperbenzoic acid. Exposure to two or more equivalents of oxidant results in oxidation to the sulfone. Displacement of the sulfone with oxygen and nitrogen nucleophiles yielded compounds as described in examples 1 and 2.

SCHEME B



Compounds of the present invention wherein the pyridazinone is substituted with a 6-amino-nicotinamide moiety are prepared by contacting A-1 with a 6-amino-nicotinamide derivative in which the amide can be a primary, secondary or tertiary amide which can be prepared by condensation of 6-amino-nicotinic acid with a suitable amine. (SCHEME B)


- 15 The amides are prepared by forming an activated carboxylic acid, such as an acid chloride or a symmetrical or mixed acid anhydride, and reacting the activated derivative with the appropriate amine in a solvent such as DMF, DCM, THF, with or without water as a co-solvent at temperatures between 0° and 60°C generally in the presence of a base such as Na₂CO₃, NaHCO₃, K₂CO₃, DIPEA, TEA or pyridine and the like to afford an amide. Carboxylic acids are
- 20 converted into their acid chlorides using standard reagents well known to someone skilled in the art, such as thionyl chloride, oxalyl chloride, phosphoryl chloride and the like. Those reagents

can be used in the presence of a base such as DIPEA, TEA or pyridine in an inert solvent such as DCM or DMF.

Alternatively a carboxylic acid can be converted *in situ* into an activated acid by different coupling procedures known to those skilled in the art. These activated acids were reacted directly with the amines to afford amides. Said activation with those peptide coupling procedures can involve the use of an activating agent like EDCI, DCC, HOBt, BOP, PyBrOP, HATU or Mukaiyama's reagent (2-fluoro-1-methylpyridinium p-toluenesulphonate) and the like with or without a base such as NMM, TEA or DIPEA in an inert solvent such as DMF or DCM at temperatures between 0° and 60°C. The reaction may alternatively be carried out in the presence of HATU or 1-hydroxy-7-azabenzotriazole (HOAt) and TEA or DIPEA in DMF, DCM or THF. Acylation of amines (March, *supra* pp.417-425; Benz, *Synthesis of Amides and Related Compounds in Comprehensive Organic Synthesis*, Winterfeldt, ed., vol. 6, Pergamon Press, Oxford 1991 pp. 381-411) has been reviewed.

The boronic acid derivatives utilized in the coupling step also contain amide linkages (see *e.g.*, SCHEME D) and these can be introduced by utilizing similar methodology. One skilled in the art will appreciate that the choice of coupling reagent is a matter of convenience.

SCHEME C




Compounds of the present invention substituted at the 3-position by an N-alkyl-ureido moiety are accessible from 4-amino-6-chloro-2-methyl-2H-pyridazin-3-one (CASRN 3366-87-8).

Suzuki coupling of A-6, or alternate dioxaborolanes, and C-1 affords C-2 which is converted to a urea C-3 by sequential treatment with phosgene and an amine or by direct condensation with an isocyanate. Alternatively the amino group of 42b can be converted to a urea prior to the Suzuki coupling step.

Compounds according to formula I wherein X is CH (TABLE II) are prepared analogously from 1-alkyl-3-amino-5-bromo-1H-pyridin-2-one (CASRN 910543-72-5) wherein R¹-NH- is a urea or from 1-alkyl-3,5-dibromo-1H-pyridin-2-one wherein R¹-NH- is a (hetero)aryl-amine.

- Compounds of the present invention with a hydroxy methyl substituent were prepared by
 5 condensation of a boronic acid derivative substituted by a *tert*-butyl-dimethyl-silyloxymethyl) moiety and an appropriately substituted pyridone or pyridazinone D-4. While the present examples utilize the TBDMS protecting group one skilled in the art will appreciate that other alcohol protecting groups could be substituted for the TBDMS group. Suzuki-coupling of 2-(*tert*-butyl-dimethyl-silyloxymethyl)-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-
 10 phenylamine or 2-[2-(*tert*-butyl-dimethyl-silyloxymethyl)-3-nitro-phenyl]-4,4,5,5-tetramethyl-[1,3,2]dioxaborolane permit incorporation of the amide side chain at a later stage in the synthesis.

SCHEME D



R = optionally substituted heteroaryl, phenyl, alkylcarbamoyl or heteroalkylcarbamoyl
 Ar = optionally substituted aryl or heteroaryl

- The pyrimidine and pyridine derivatives described herein are kinase inhibitors, in particular Btk inhibitors. These inhibitors can be useful for treating one or more diseases responsive to kinase
 15 inhibition, including diseases responsive to Btk inhibition and/or inhibition of B-cell proliferation, in mammals. Without wishing to be bound to any particular theory, it is believed that the interaction of the compounds of the invention with Btk results in the inhibition of Btk activity and thus in the pharmaceutical utility of these compounds. Accordingly, the invention includes a method of treating a mammal, for instance a human, having a disease responsive to
 20 inhibition of Btk activity, and/or inhibiting B-cell proliferation, comprising administering to the mammal having such a disease, an effective amount of at least one chemical entity provided herein. An effective concentration may be ascertained experimentally, e.g. by assaying blood concentration of the compound, or theoretically, by calculating bioavailability. Other kinases that

may be affected in addition to Btk include, but are not limited to, other tyrosine kinases and serine/threonine kinases. Example 24 describes an enzyme inhibition assay to measure inhibition of Btk activity by compounds of the present invention. Example 25 describes a functional assay to measure inhibition of Ca^{2+} influx mediated by Btk kinase.

- 5 Kinases play notable roles in signaling pathways controlling fundamental cellular processes such as proliferation, differentiation, and death (apoptosis). Abnormal kinase activity has been implicated in a wide range of diseases, including multiple cancers, autoimmune and/or inflammatory diseases, and acute inflammatory reactions. The multifaceted role of kinases in key cell signaling pathways provides a significant opportunity to identify novel drugs targeting
- 10 kinases and signaling pathways.

An embodiment includes a method of treating a patient having an autoimmune and/or inflammatory disease, or an acute inflammatory reaction responsive to inhibition of Btk activity and/or B-cell proliferation.

- Autoimmune and/or inflammatory diseases that can be affected using compounds and compositions according to the invention include, but are not limited to: psoriasis, allergy, Crohn's disease, irritable bowel syndrome, Sjogren's disease, tissue graft rejection, and hyperacute rejection of transplanted organs, asthma, systemic lupus erythematosus (and associated glomerulonephritis), dermatomyositis, multiple sclerosis, scleroderma, vasculitis (ANCA-associated and other vasculitides), autoimmune hemolytic and thrombocytopenic states, Goodpasture's syndrome (and
- 15 associated glomerulonephritis and pulmonary hemorrhage), atherosclerosis, rheumatoid arthritis, chronic Idiopathic thrombocytopenic purpura (ITP), Addison's disease, Parkinson's disease, Alzheimer's disease, diabetes, septic shock, and myasthenia gravis,
- 20

- Included herein are methods of treatment in which at least one chemical entity provided herein is administered in combination with an anti-inflammatory agent. Anti-inflammatory agents include
- 25 but are not limited to NSAIDs, non-specific and COX-2 specific cyclooxygenase enzyme inhibitors, gold compounds, corticosteroids, tumor necrosis factor (TNF) receptor antagonists, immunosuppressants and methotrexate.

- Examples of NSAIDs include, but are not limited to, ibuprofen, flurbiprofen, naproxen and naproxen sodium, diclofenac, combinations of diclofenac sodium and misoprostol, sulindac,
- 30 oxaprozin, diflunisal, piroxicam, indomethacin, etodolac, fenoprofen calcium, ketoprofen, sodium nabumetone, sulfasalazine, tolmetin sodium, and hydroxychloroquine. Examples of

NSAIDs also include COX-2 specific inhibitors such as celecoxib, valdecoxib, lumiracoxib and/or etoricoxib.

In some embodiments, the anti-inflammatory agent is a salicylate. Salicylates include but are not limited to acetylsalicylic acid or aspirin, sodium salicylate, and choline and magnesium

5 salicylates.

The anti-inflammatory agent may also be a corticosteroid. For example, the corticosteroid may be cortisone, dexamethasone, methylprednisolone, prednisolone, prednisolone sodium phosphate, or prednisone.

In additional embodiments the anti-inflammatory agent is a gold compound such as gold sodium
10 thiomalate or auranofin.

The invention also includes embodiments in which the anti-inflammatory agent is a metabolic inhibitor such as a dihydrofolate reductase inhibitor, such as methotrexate or a dihydroorotate dehydrogenase inhibitor, such as leflunomide.

Other embodiments of the invention pertain to combinations in which at least one anti-in-
15 flammatory compound is an anti-C5 monoclonal antibody (such as eculizumab or pexelizumab), a TNF antagonist, such as entanercept, or infliximab, which is an anti-TNF alpha monoclonal antibody.

Still other embodiments of the invention pertain to combinations in which at least one active agent is an immunosuppressant compound such as an immunosuppressant compound chosen
20 from methotrexate, leflunomide, cyclosporine, tacrolimus, azathioprine, and mycophenolate mofetil.

B-cells and B-cell precursors expressing BTK have been implicated in the pathology of B-cell malignancies, including, but not limited to, B-cell lymphoma, lymphoma (including Hodgkin's and non-Hodgkin's lymphoma), hairy cell lymphoma, multiple myeloma, chronic and acute
25 myelogenous leukemia and chronic and acute lymphocytic leukemia.

BTK has been shown to be an inhibitor of the Fas/APO-1 (CD-95) death inducing signaling complex (DISC) in B-lineage lymphoid cells. The fate of leukemia/lymphoma cells may reside in the balance between the opposing proapoptotic effects of caspases activated by DISC and an

upstream anti-apoptotic regulatory mechanism involving BTK and/or its substrates (Vassilev *et al.*, *J. Biol. Chem.* 1998, 274, 1646-1656).

It has also been discovered that BTK inhibitors are useful as chemosensitizing agents, and, thus, are useful in combination with other chemotherapeutic drugs, in particular, drugs that induce
5 apoptosis. Examples of other chemotherapeutic drugs that can be used in combination with chemosensitizing BTK inhibitors include topoisomerase I inhibitors (camptothecin or topotecan), topoisomerase II inhibitors (e.g. daunomycin and etoposide), alkylating agents (e.g. cyclophosphamide, melphalan and BCNU), tubulin directed agents (e.g. taxol and vinblastine), and biological agents (e.g. antibodies such as anti CD20 antibody, IDEC 8, immunotoxins, and
10 cytokines).

Btk activity has also be associated with some leukemias expressing the *bcr-abl* fusion gene resulting from translocation of parts of chromosome 9 and 22. This abnormality is commonly observed in chronic myelogenous leukemia. Btk is constitutively phosphorylated by the *bcr-abl* kinase which initiates downstream survival signals which circumvents apoptosis in *bcr-abl* cells.
15 (Feldhahn *et al. J. Exp. Med.* 2005 201(11):1837-1852).

The compounds of the present invention may be formulated in a wide variety of oral administration dosage forms and carriers. Oral administration can be in the form of tablets, coated tablets, dragées, hard and soft gelatine capsules, solutions, emulsions, syrups, or suspensions. Compounds of the present invention are efficacious when administered by other routes of administration including continuous (intravenous drip) topical parenteral, intramuscular, intra-
20 venous, subcutaneous, transdermal (which may include a penetration enhancement agent), buccal, nasal, inhalation and suppository administration, among other routes of administration. The preferred manner of administration is generally oral using a convenient daily dosing regimen which can be adjusted according to the degree of affliction and the patient's response to the
25 active ingredient.

A compound or compounds of the present invention, as well as their pharmaceutically useable salts, together with one or more conventional excipients, carriers, or diluents, may be placed into the form of pharmaceutical compositions and unit dosages. The pharmaceutical compositions and unit dosage forms may be comprised of conventional ingredients in conventional proportions,
30 with or without additional active compounds or principles, and the unit dosage forms may contain any suitable effective amount of the active ingredient commensurate with the intended daily dosage range to be employed. The pharmaceutical compositions may be employed as

solids, such as tablets or filled capsules, semisolids, powders, sustained release formulations, or liquids such as solutions, suspensions, emulsions, elixirs, or filled capsules for oral use; or in the form of suppositories for rectal or vaginal administration; or in the form of sterile injectable solutions for parenteral use. A typical preparation will contain from about 5% to about 95%
5 active compound or compounds (w/w). The term "preparation" or "dosage form" is intended to include both solid and liquid formulations of the active compound and one skilled in the art will appreciate that an active ingredient can exist in different preparations depending on the target organ or tissue and on the desired dose and pharmacokinetic parameters.

The term "excipient" as used herein refers to a compound that is useful in preparing a pharmaceutical composition, generally safe, non-toxic and neither biologically nor otherwise
10 undesirable, and includes excipients that are acceptable for veterinary use as well as human pharmaceutical use. The compounds of this invention can be administered alone but will generally be administered in admixture with one or more suitable pharmaceutical excipients, diluents or carriers selected with regard to the intended route of administration and standard
15 pharmaceutical practice.

"Pharmaceutically acceptable" means that which is useful in preparing a pharmaceutical composition that is generally safe, non-toxic, and neither biologically nor otherwise undesirable and includes that which is acceptable for veterinary as well as human pharmaceutical use.

A "pharmaceutically acceptable salt" form of an active ingredient may also initially confer a
20 desirable pharmacokinetic property on the active ingredient which were absent in the non-salt form, and may even positively affect the pharmacodynamics of the active ingredient with respect to its therapeutic activity in the body. The phrase "pharmaceutically acceptable salt" of a compound means a salt that is pharmaceutically acceptable and that possesses the desired pharmacological activity of the parent compound. Such salts include: (1) acid addition salts,
25 formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like; or formed with organic acids such as acetic acid, propionic acid, hexanoic acid, cyclopentanepropionic acid, glycolic acid, pyruvic acid, lactic acid, malonic acid, succinic acid, malic acid, maleic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, 3-(4-hydroxybenzoyl)benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid,
30 ethanesulfonic acid, 1,2-ethane-disulfonic acid, 2-hydroxyethanesulfonic acid, benzenesulfonic acid, 4-chlorobenzenesulfonic acid, 2-naphthalenesulfonic acid, 4-toluenesulfonic acid, camphorsulfonic acid, 4-methylbicyclo[2.2.2]-oct-2-ene-1-carboxylic acid, glucoheptonic acid,

3-phenylpropionic acid, trimethylacetic acid, tertiary butylacetic acid, lauryl sulfuric acid, gluconic acid, glutamic acid, hydroxynaphthoic acid, salicylic acid, stearic acid, muconic acid, and the like; or (2) salts formed when an acidic proton present in the parent compound either is replaced by a metal ion, *e.g.*, an alkali metal ion, an alkaline earth ion, or an aluminum ion; or
5 coordinates with an organic base such as ethanolamine, diethanolamine, triethanolamine, tromethamine, N-methylglucamine, and the like.

Solid form preparations include powders, tablets, pills, capsules, cachets, suppositories, and dispersible granules. A solid carrier may be one or more substances which may also act as diluents, flavoring agents, solubilizers, lubricants, suspending agents, binders, preservatives,
10 tablet disintegrating agents, or an encapsulating material. In powders, the carrier generally is a finely divided solid which is a mixture with the finely divided active component. In tablets, the active component generally is mixed with the carrier having the necessary binding capacity in suitable proportions and compacted in the shape and size desired. Suitable carriers include but are not limited to magnesium carbonate, magnesium stearate, talc, sugar, lactose, pectin, dextrin,
15 starch, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose, a low melting wax, cocoa butter, and the like. Solid form preparations may contain, in addition to the active component, colorants, flavors, stabilizers, buffers, artificial and natural sweeteners, dispersants, thickeners, solubilizing agents, and the like.

Liquid formulations also are suitable for oral administration include liquid formulation including
20 emulsions, syrups, elixirs, aqueous solutions, aqueous suspensions. These include solid form preparations which are intended to be converted to liquid form preparations shortly before use. Emulsions may be prepared in solutions, *e.g.*, in aqueous propylene glycol solutions or may contain emulsifying agents such as lecithin, sorbitan monooleate, or acacia. Aqueous solutions can be prepared by dissolving the active component in water and adding suitable colorants,
25 flavors, stabilizing, and thickening agents. Aqueous suspensions can be prepared by dispersing the finely divided active component in water with viscous material, such as natural or synthetic gums, resins, methylcellulose, sodium carboxymethylcellulose, and other well known suspending agents.

The compounds of the present invention may be formulated for parenteral administration (*e.g.*,
30 by injection, *e.g.* bolus injection or continuous infusion) and may be presented in unit dose form in ampoules, pre-filled syringes, small volume infusion or in multi-dose containers with an added preservative. The compositions may take such forms as suspensions, solutions, or emulsions in

oily or aqueous vehicles, e.g. solutions in aqueous polyethylene glycol. Examples of oily or nonaqueous carriers, diluents, solvents or vehicles include propylene glycol, polyethylene glycol, vegetable oils (*e.g.*, olive oil), and injectable organic esters (*e.g.*, ethyl oleate), and may contain formulatory agents such as preserving, wetting, emulsifying or suspending, stabilizing and/or dispersing agents. Alternatively, the active ingredient may be in powder form, obtained by aseptic isolation of sterile solid or by lyophilisation from solution for constitution before use with a suitable vehicle, *e.g.*, sterile, pyrogen-free water.

The compounds of the present invention may be formulated for topical administration to the epidermis as ointments, creams or lotions, or as a transdermal patch. Ointments and creams may, *e.g.*, be formulated with an aqueous or oily base with the addition of suitable thickening and/or gelling agents. Lotions may be formulated with an aqueous or oily base and will in general also contain one or more emulsifying agents, stabilizing agents, dispersing agents, suspending agents, thickening agents, or coloring agents. Formulations suitable for topical administration in the mouth include lozenges comprising active agents in a flavored base, usually sucrose and acacia or tragacanth; pastilles comprising the active ingredient in an inert base such as gelatin and glycerin or sucrose and acacia; and mouthwashes comprising the active ingredient in a suitable liquid carrier.

The compounds of the present invention may be formulated for administration as suppositories. A low melting wax, such as a mixture of fatty acid glycerides or cocoa butter is first melted and the active component is dispersed homogeneously, *e.g.*, by stirring. The molten homogeneous mixture is then poured into convenient sized molds, allowed to cool, and to solidify.

The compounds of the present invention may be formulated for vaginal administration. Pessaries, tampons, creams, gels, pastes, foams or sprays containing in addition to the active ingredient such carriers as are known in the art to be appropriate.

The compounds of the present invention may be formulated for nasal administration. The solutions or suspensions are applied directly to the nasal cavity by conventional means, *e.g.*, with a dropper, pipette or spray. The formulations may be provided in a single or multidose form. In the latter case of a dropper or pipette, this may be achieved by the patient administering an appropriate, predetermined volume of the solution or suspension. In the case of a spray, this may be achieved *e.g.* by means of a metering atomizing spray pump.

The compounds of the present invention may be formulated for aerosol administration, particularly to the respiratory tract and including intranasal administration. The compound will generally have a small particle size e.g. of the order of five (5) microns or less. Such a particle size may be obtained by means known in the art, e.g. by micronization. The active ingredient is
5 provided in a pressurized pack with a suitable propellant such as a chlorofluorocarbon (CFC), e.g., dichlorodifluoromethane, trichlorofluoromethane, or dichlorotetrafluoroethane, or carbon dioxide or other suitable gas. The aerosol may conveniently also contain a surfactant such as lecithin. The dose of drug may be controlled by a metered valve. Alternatively the active ingredients may be provided in a form of a dry powder, e.g. a powder mix of the compound in a
10 suitable powder base such as lactose, starch, starch derivatives such as hydroxypropylmethyl cellulose and polyvinylpyrrolidone (PVP). The powder carrier will form a gel in the nasal cavity. The powder composition may be presented in unit dose form e.g. in capsules or cartridges of e.g., gelatin or blister packs from which the powder may be administered by means of an inhaler.

When desired, formulations can be prepared with enteric coatings adapted for sustained or
15 controlled release administration of the active ingredient. For example, the compounds of the present invention can be formulated in transdermal or subcutaneous drug delivery devices. These delivery systems are advantageous when sustained release of the compound is necessary and when patient compliance with a treatment regimen is crucial. Compounds in transdermal delivery systems are frequently attached to an skin-adhesive solid support. The compound of
20 interest can also be combined with a penetration enhancer, e.g., Azone (1-dodecylazacycloheptan-2-one). Sustained release delivery systems are inserted subcutaneously into to the subdermal layer by surgery or injection. The subdermal implants encapsulate the compound in a lipid soluble membrane, e.g., silicone rubber, or a biodegradable polymer, e.g., polyactic acid.

Suitable formulations along with pharmaceutical carriers, diluents and excipients are described
25 in *Remington: The Science and Practice of Pharmacy* 1995, edited by Martin, Mack Publishing Company, 19th edition, Easton, Pennsylvania. A skilled formulation scientist may modify the formulations within the teachings of the specification to provide numerous formulations for a particular route of administration without rendering the compositions of the present invention unstable or compromising their therapeutic activity.

30 The modification of the present compounds to render them more soluble in water or other vehicle, e.g., may be easily accomplished by minor modifications (salt formulation, esterification, etc.), which are well within the ordinary skill in the art. It is also well within the ordinary skill of

the art to modify the route of administration and dosage regimen of a particular compound in order to manage the pharmacokinetics of the present compounds for maximum beneficial effect in patients.

The term "therapeutically effective amount" as used herein means an amount required to reduce
5 symptoms of the disease in an individual. The dose will be adjusted to the individual requirements in each particular case. That dosage can vary within wide limits depending upon numerous factors such as the severity of the disease to be treated, the age and general health condition of the patient, other medicaments with which the patient is being treated, the route and form of administration and the preferences and experience of the medical practitioner involved.
10 For oral administration, a daily dosage of between about 0.01 and about 1000 mg/kg body weight per day should be appropriate in monotherapy and/or in combination therapy. A preferred daily dosage is between about 0.1 and about 500 mg/kg body weight, more preferred 0.1 and about 100 mg/kg body weight and most preferred 1.0 and about 10 mg/kg body weight per day. Thus, for administration to a 70 kg person, the dosage range would be about 7 mg to 0.7 g per
15 day. The daily dosage can be administered as a single dosage or in divided dosages, typically between 1 and 5 dosages per day. Generally, treatment is initiated with smaller dosages which are less than the optimum dose of the compound. Thereafter, the dosage is increased by small increments until the optimum effect for the individual patient is reached. One of ordinary skill in treating diseases described herein will be able, without undue experimentation and in reliance on
20 personal knowledge, experience and the disclosures of this application, to ascertain a therapeutically effective amount of the compounds of the present invention for a given disease and patient.

The pharmaceutical preparations are preferably in unit dosage forms. In such form, the preparation is subdivided into unit doses containing appropriate quantities of the active component.
25 The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation, such as packeted tablets, capsules, and powders in vials or ampoules. Also, the unit dosage form can be a capsule, tablet, cachet, or lozenge itself, or it can be the appropriate number of any of these in packaged form.

The following examples illustrate the preparation and biological evaluation of compounds within
30 the scope of the invention. These examples and preparations which follow are provided to enable those skilled in the art to more clearly understand and to practice the present invention.

They should not be considered as limiting the scope of the invention, but merely as being illustrative and representative thereof.

Example 1: 4-tert-Butyl-N-(2-methyl-3-{1-methyl-6-oxo-5-[2-(pyrrolidin-3-ylmethoxy)-pyrimidin-4-ylamino]-1,6-dihydro-pyridazin-3-yl}-phenyl)-benzamide (I-19,

SCHEME A)

step 1 - A solution of 4,6-dichloro-2H-pyridazin-one (**A-1a**, 3.9 g, 23 mmol, CASRN 17285-37-9) in anhydrous DMF (45 mL) was cooled at 0°C under nitrogen. To this solution was added sodium hydride (0.92 g, 23 mmol, 60% dispersion in mineral oil). The mixture was warmed to RT and stirred for 30 min. then cooled to about 10°C and methyl iodide (3.2 g, 23 mmol) was added drop wise over 5 min. The resulting mixture was stirred at RT for about 4 h and then diluted with water (50 mL) and extracted with EtOAc (100 mL). The organic layer was washed three times with brine, dried (Na₂SO₄) and concentrated *in vacuo* to give 4 g (100%) of **A-1b** which used in the next step without further purification.

step 2 - To a solution of 2-methylsulfanyl-pyrimidin-4-ylamine (**A-5**, 1.4 g, 9.8 mmol, CASRN 2183-66-6) and anhydrous DMF (14 mL) cooled at 0°C and under nitrogen atmosphere was added NaH (0.392 g, 609.8 mmol, 60% mineral oil dispersion). The reaction mixture was warmed to RT and stirred for 30 min. To the resulting reaction mixture was added dropwise over 5 min a solution of **A-1b** (0.872 g, 4.9 mmol) in DMF (5 mL). The reaction mixture was stirred at RT for 2 h and then diluted with water (20 mL) and EtOAc (50 mL). The organic layer was washed three times with brine. The desired product which precipitated out of an EtOAc solution, was filtered and dried under high vacuum to afford 1 g (71.4%) of **A-2** which was used in the next step without further purification.

step 3 - To a suspension of **A-2** (0.2 g, 0.7 mmol) and DME (8 mL) in a microwave tube was added a solution of Na₂CO₃ (0.223 g, 2.1 mmol) in water (3 mL) and **A-6** (0.277 g, 0.70 mmol). The mixture was sparged with nitrogen, Pd(0)(PPh₃)₄ (0.041 g, 0.035 mmol) was added and the tube sealed and heated at 175°C in a microwave apparatus for 45 min. After cooling, the combined reaction mixture was partitioned between EtOAc and H₂O. The organic layer was washed with brine, dried (Na₂SO₄), filtered and concentrated *in vacuo*. The crude residue was purified by SiO₂ chromatography eluting with a MeOH/DCM gradient (2% to 4% MeOH) to afford 0.4 g (55%) of **I-12**.

step 4 - To a solution of **I-12** (0.370 g, 0.719 mmol), MeOH (16 mL) and THF (10 mL) was added dropwise over 5 min a solution of oxone (0.884 g, 1.43 mmol) in H₂O (6 mL). The solution was stirred at RT for 4 h then a second portion of oxone (0.6g, 0.97 mmol) in H₂O (4

mL) was added and the mixture stirred for additional 2 h. The reaction mixture was diluted with H₂O (50 mL) and extracted with EtOAc (100 mL). The organic layer was washed with brine, dried (Na₂SO₄), filtered, concentrated and dried *in vacuo* to afford 0.380 g (96%) of **I-13** which was used in the next step without further purification.

- 5 step 5 - To a solution of **I-13** (0.100 g, 0.18 mmol) and DMF (2 mL) was added NaH (0.028 g, 60%, 0.72 mmol) and the resulting mixture warmed to RT and stirred 30 min. To the resulting solution was added pyrrolidin-3-yl- methanol (0.054 g, 0.535 mmol) and the mixture stirred at RT for 4 h. The reaction mixture was next diluted with water and extracted with EtOAc. The organic layer was washed with brine, dried (Na₂SO₄) and concentrated *in vacuo*. The resulting
10 residue was purified on a preparative SiO₂ TLC plate developed with 10% MeOH/DCM to afford 0.025 g (26%) of **I-19**.

I-17 and **I-18** were prepared analogously except pyrrolidin-3-ol was replaced with methanol and N,N-dimethylamino-ethanol, respectively.

- I-45** can be prepared analogously except in step 2, 2-methylsulfanyl-pyrimidin-4-ylamine is
15 replaced with 3-amino-1-methyl-pyrazole, in step 3, **A-6** is replaced with **120** (Ar = 4-cyclopropylphenyl) and steps 3 and 4 are omitted.

Example 2: 4-tert-Butyl-N-(3-{5-[2-(3-hydroxy-pyrrolidin-1-yl)-pyrimidin-4-ylamino]-1-methyl-6-oxo-1,6-dihydro-pyridazin-3-yl}-2-methyl-phenyl)-benzamide (I-22**)**

- To a solution of **I-13** (0.069 g, 0.12 mmol) and DMF (2 mL) was added pyrrolidin-3-ol (0.032g,
20 0.36 mmol). The reaction mixture was stirred for 2 h at RT and then for 2 h at 60°C. The reaction mixture was cooled to RT and partitioned between EtOAc and H₂O. The organic layer was washed with brine, dried (Na₂SO₄), filtered and concentrated *in vacuo*. The recovered residue was purified on a preparative SiO₂ TLC plate developed with 5% MeOH/DCM to afford 0.028 g (42%) of **I-22**.

- 25 The following analogs were prepared analogously except pyrrolidin-3-ol was replaced with the amine or alcohol in parenthesis: **I-11** (N-methyl piperazine), **I-14** (morpholine), **I-20** (pyrrolidin-3-yl- methanol) and **I-21** (pyrrolidine).

Example 3: 4-tert-Butyl-N-(2-methyl-3-{1-methyl-5-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-6-oxo-1,6-dihydro-pyridazin-3-yl}-phenyl)-benzamide (I-1**;**

- 30 **SCHEME B)**

step 1 - To a solution of morpholine (9.00 g, 103 mmol) in EtOH (400 mL) was added EDCI (10.0 g, 52.2 mmol), HOBt (7.00 g 51.8 mmol), and 6-aminonicotinic acid (6.00 g, 43.4 mmol). After stirring for 18 h, the resulting solid was filtered. The solid was triturated with a mixture of

MeOH (100 mL) and DCM (100 mL) to afford 3.08 g of **B-2** (NR^1R^2 together are morpholine): MS (ESI) 208.1 (M+H).

step 2 - To an ice cold suspension of **B-2** (2.921 g, 14.11 mmol, CASRN 827587-90-6) and DMF (50 mL) was added NaH (847 mg, 21.2 mmol, 60% suspension in mineral oil). The mixture was stirred at RT for 30 min then re-cooled to 0°C and **A-1b** (1.256 g, 7.056 mmol) was added. After 2 h, the reaction mixture was partitioned between saturated aqueous NH_4Cl and EtOAc. The EtOAc layer was dried (MgSO_4), filtered and concentrated *in vacuo*. The crude product was purified by SiO_2 chromatography eluting with 2% MeOH/DCM to afford 0.732 g (2.09 mmol) of **B-3**: MS (ESI) 350.1 (M+H)⁺.

step 3 - To a suspension of **B-3** (0.030 g, 0.085 mmol) and **A-6** (0.035 g, 0.089 mmol) and DME (2 mL) in a microwave tube was added a solution of Na_2CO_3 (0.028 g, 0.27 mmol) and H_2O (0.5 mL). After sparging the mixture with nitrogen, $\text{Pd(0)(PPh}_3)_4$ (0.01 g, 0.009 mmol) was added and the tube sealed and heated at 175°C for 30 min in a microwave apparatus. The reaction mixture was partitioned between H_2O and EtOAc. The organic layer was washed with brine, dried (Na_2SO_4) and concentrated *in vacuo*. The crude product was purified on a preparative SiO_2 TLC plate developed with 5% MeOH/DCM to afford 0.034 g (69%) of **I-1**.

I-2 was prepared analogously except in step 1, morpholine was replaced with 1-methyl-piperazine which afforded (6-amino-pyridin-3-yl)-(4-methyl-piperazin-1-yl)-methanone (CASRN 652138-83-5).


I-3, I-4, I-9, and I-24 were prepared analogously except in step 1 was omitted and 2-amino-pyridine, 4-aminopyrimidine, methyl 6-amino-nicotinate (CASRN 36052-24-1) and 4-(4-aminobenzoyl)-morpholine (CASRN 51207-86-4), respectively, were used in place of **B-2** in step 2.

I-25 was prepared analogously except in step 3, **A-6** was replaced with **14b** (CASRN 210907-84-9) which was subsequently acylated with 4-*tert*-butyl-benzoic acid after the Suzuki coupling step.

I-26 was prepared by condensation of methyl 6-aminonicotinate (CASRN 36052-24-1) in place of **B-2** in step 2. The Suzuki coupling proceeded with concomitant hydrolysis of the methyl ester and the resulting carboxylic acid was converted to the amide by BOP-mediated coupling of the acid and 4-hydroxy-piperidine in the presence of TEA and DMF. Step 1 in the example was omitted.

I-46 is prepared analogously except **A-6** is replaced by **120** (Ar = 4-cyclopropyl-phenyl) as described in step 5 of example 25.

Example 4: 4-*tert*-Butyl-piperazine-1-carboxylic acid (2-methyl-3-{1-methyl-5-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-6-oxo-1,6-dihydro-pyridazin-3-yl}-phenyl)-amide (I-6)



5 step 1 - A solution of **B-3** (149 mg, 0.426 mmol), **14a** (105 mg, 0.426 mmol, CASRN 882678-96-8), Pd(0)(PPh₃)₄ (49 mg, 0.042 mmol), and Na₂CO₃ (135 mg, 1.28 mmol) in DME (2 mL) and H₂O (1 mL) was heated at 170°C for 12.5 min in a microwave synthesizer. The resulting mixture was partitioned between EtOAc and water. The EtOAc layer was washed with brine, dried (MgSO₄), filtered and concentrated *in vacuo*. The crude product was purified by SiO₂

10 chromatography eluting with 4 % MeOH/DCM to afford 0.154 g (0.366 mmol) of **12a**: MS (ESI) 421 (M+H)⁺.


step 2 - To a suspension of **12a** (50mg, 0.12mmol) in THF (4 mL) was added a phosgene solution (0.18 mL, 0.36mmol, 1.93 M in toluene). After 20 min, 1-*tert*-butyl-piperazine (85 mg, 0.60 mmol) was added and the resulting solution stirred for 15 min. The resulting mixture was

15 partitioned between EtOAc and dilute aqueous NaHCO₃. The EtOAc layer was washed with brine, dried (MgSO₄), filtered and concentrated *in vacuo*. The crude product was purified on a preparative SiO₂ TLC plate developed with 5% MeOH/DCM to afford 0.050 g (0.085 mmol) of **I-6**: MS (ESI) 589 (M+H)⁺.


I-15 was prepared analogously except in step 2, **12a** was condensed with with 4-(1-hydroxy-1-methyl-ethyl)-benzoic acid (**86**) using BOP-mediated coupling as described in step 2 of example

20 6.

I-7 can be prepared analogously except in step 2, **12** is condensed with 4-*tert*-butyl-2-methoxy benzoic acid using BOP-mediated coupling as described in step 2 of example 6.



16a: X = NMe
16b: X = CH(OH)




18a: X = NMe
18b: X = CH(OH)

I-5 was prepared analogously starting with **18a** which was converted to **16a** and condensed with **86**. The requisite amine **18a** was prepared by condensation of (6-amino-pyridin-3-yl)-(4-methylpiperazin-1-yl)-methanone (CASRN 652138-83-5) and **A-1b**.

I-30 was prepared by the condensation of **12a** and 4-*tert*-butyl-2-methoxy-benzoyl chloride. **I-16** was prepared by BBr₃ mediated demethylation of **I-30**.

Example 5: 3-*tert*-Butoxy-azetidine-1-carboxylic acid (2-methyl-3-{1-methyl-5-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-6-oxo-1,6-dihydro-pyridazin-3-yl}-phenyl)-amide (I-10)



step 1 - A solution of **20a** (CASRN 141699-55-0, 1.313 g, 7.58 mmol), cyclohexane (20 mL), DCM (10 mL), *tert*-butyl trichloroacetimidate (5.4 mL, 30 mmol) and BF₃·Et₂O (0.48 mL, 3.8 mmol) was stirred for 16 h. NaHCO₃ (2.5 g) was added, the mixture was filtered through CELITE®, and the filtrate was concentrated to 2.52 g of an oily white solid. The crude product was purified by SiO₂ chromatography eluting with an EtOAc/hexane gradient (0-50% EtOAc) to afford 0.408 g (23%) of **20b** as a colorless oil.

step 2 - A premixed solution of MeOH (2 mL) and acetyl chloride (0.5 mL) was added to a solution of **20b** (0.251 g, 1.09 mmol) in MeOH (2 mL) at 0-5°C. The colorless solution was


warmed to RT and stirred for 4 h, then concentrated, chasing with diethyl ether, to afford crude **22** as a white solid that was used without further purification.

step 3 - Phosgene (20% in toluene, 0.459 mL, 0.874 mmol) was added to a solution of **14a** (0.1697 g, 0.728 mmol) and DIPEA (0.510 mL, 2.93 mmol) in DCM (5 mL) cooled to 0-5°C.

- 5 The reaction mixture was stirred for 10 min and then transferred *via* Pasteur pipette to a flask containing crude **22**. A total of 5 mL of additional DCM was used in rinsing the flask. The yellow solution was stirred at 0-5°C for 10 min, then 4 g of SiO₂ was added, and the mixture was concentrated to a pale yellow powder. The dry SiO₂ was applied to the top of a SiO₂ column and eluted with a EtOAc/hexane gradient (50-100% EtOAc) to afford 0.226 g (80%) of **24** as a white solid.

The synthesis of **I-10** was completed by coupling of **B-3** and **24** as previously described in step 3 of example 3.

Example 6: N-(2-*tert*-Butoxy-ethyl)-2-(2-methyl-3-{1-methyl-5-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-6-oxo-1,6-dihydro-pyridin-3-yl}-phenyl)-acetamide (**I-32**)




- step 1 - A solution of **B-3** (256 mg, 0.732 mmol), **28** (192 mg, 0.732 mmol), Pd(0)(PPh₃)₄ (84 mg, 0.073 mmol), and Na₂CO₃ (310 mg, 2.93 mmol) in DME (2 mL) and H₂O (1 mL) was heated at 170°C for 12.5 min in a microwave synthesizer. The resulting mixture was partitioned between EtOAc and H₂O. The pH of the aqueous layer was adjusted to *ca.* pH 1 with 6M aqueous HCl and the resulting solid was filtered, washed with water and dried to afford 0.263 g (0.586 mmol) of **26**: MS (ESI) 450.1 (M+H)⁺.

step 2 - To **26** (50 mg, 0.11 mmol) and BOP (55 mg, 0.12 mmol) was added DMF (2 mL), N,N-dimethylaminoethanol (0.017 g, 0.11 mmol) and TEA (56 mg, 0.56 mmol). After 4 h the reaction mixture was partitioned between EtOAc and dilute aqueous NaHCO₃. The EtOAc layer was washed with brine, dried (MgSO₄), filtered and concentrated *in vacuo*. The crude product was purified on a preparative SiO₂ TLC plate developed with 5% MeOH/DCM to afford 0.033 g (0.060 mmol) of **I-32**: MS (ESI) 549 (M+H)⁺.

The following were prepared from **26**: **I-29**, **I-33** and **I-36** were prepared by EDCI mediated coupling of **26**, with 1-amino-4,4-dimethyl-3-hydroxypentane, 3-*tert*-butoxy-cyclobutylamine, pyrrolidine and 2-*tert*-butoxy-ethanol respectively using the EDCI coupling protocol described in step 2 of example 13 except excess DIPEA was added as the base. **I-35** was prepared by BOP-catalyzed coupling of **28** and 3-*iso*-propoxy-cyclobutylamine using the protocol described in step 2 to afford 1-(3-*tert*-butoxy-azetidin-1-yl)-2-[3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-ethanone which was subjected to Suzuki-coupling with **B-3** (NR¹R² = morpholine). **I-31** was prepared analogously from **30** by EDCI mediated coupling with 2,3-dihydro-1H-isoindole. Suzuki coupling of 5-chloro-3-[5-(4-hydroxy-piperidine-1-carbonyl)-pyridin-2-ylamino]-1-methyl-1H-pyridin-2-one and **18** afforded **30**.

Example 7: 6-{3-[2-(4-*tert*-Butyl-phenyl)-2-oxo-ethyl]-phenyl}-2-methyl-4-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-2H-pyridazin-3-one (I-37**)**




step 1 - To **26** (163 mg, 0.362 mmol) and BOP (160 mg, 0.362 mmol) was added DMF (5 mL), TEA (125 mg, 1.23 mmol) and thiophenol (40 mg, 0.36 mmol). After stirring for 18 h, the resulting solution was partitioned between EtOAc and dilute aqueous NaHCO₃. The EtOAc layer was washed with brine, dried (MgSO₄), filtered and concentrated *in vacuo*. The crude product was purified by SiO₂ chromatography eluting with (2%MeOH/DCM) to afford 0.041 g (0.076 mmol) of **32**: MS (ESI) 542.0 (M+H)⁺.

step 2 - To a solution of **32** (40 mg, 0.074 mmol), 4-*tert*-butylphenylboronic acid (16 mg, 0.089 mmol), copper(I)thiophenecarboxylate (17 mg, 0.089 mmol) and Pd₂(dba)₃ (1.6 mg, 0.0017 mmol) and THF (2 mL) was added a solution of triethylphosphite (1.2 mg, 0.072 mmol) and THF (2 mL). The resulting mixture was heated to 30°C under an argon atmosphere for 18 h.

The mixture was partitioned between EtOAc and H₂O. The EtOAc layer was washed with brine, dried (MgSO₄), filtered and concentrated *in vacuo*. The crude product was purified by preparative HPLC to afford 0.004 g (0.007 mmol) of **I-37**: MS (ESI) 566 (M+H)⁺.

I-38 to **I-44** can be prepared analogously except the palladium coupling in step 2 is replaced by an amide coupling procedure of **26** with 1-methyl-piperazine, 3-*tert*-butoxy-azetidine, 2H-isindole, 1-isopropyl-piperazine, 4-*tert*-butyl-piperidine, 3,3-dimethyl-butylamine, 1-acetyl-piperazine, respectively. The coupling can be carried out using BOP-mediated condensation as described in step 2 of example 6.

Example 8: 7-*tert*-Butyl-3-(2-methyl-3-{1-methyl-5-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-6-oxo-1,6-dihydro-pyridazin-3-yl}-phenyl)-3H-quinazolin-4-one (**I-8**)



step 1 - To **34a** (3.715 g, 11.35 mmol), Cl₂(PPh₃)₂Pd(II) (336 mg, 0.478 mmol), CuI (168 mg, 0.882 mmol) was added DMF (45 mL), TEA (1.72 g, 17.0 mmol), and trimethylsilylacetylene (2.229 g, 22.70 mmol). The resulting mixture was heated at 90°C under a nitrogen atmosphere for 15 min. The mixture was partitioned between EtOAc and H₂O. The EtOAc layer was washed with brine, dried (MgSO₄), filtered and concentrated *in vacuo*. The crude product was purified by SiO₂ chromatography eluting with DCM to yield 3.193 g (11.61 mmol) of **34b**.

step 2 - A solution of **34b** (3.167 g, 11.51 mmol), K₂CO₃ (3.18 g, 23.0 mmol) were stirred in MeOH (75 mL) for 20 min. The resulting mixture was partitioned between EtOAc and water.

The EtOAc layer was washed with brine, dried (MgSO₄), filtered and concentrated *in vacuo*.

The crude product was purified by SiO₂ chromatography eluting with an EtOAc/hexane gradient (0 to 5% EtOAc/hexanes) to yield 1.910 g (9.398 mmol) of **34c**.

step 3 – To a solution of **34c** (1.662 g, 8.176 mmol), CCl₄ (20 mL), MeCN (20 mL) and H₂O

5 (40 mL) was added sequentially H₂IO₄ (9.322 g, 40.89 mmol) and Ru(III)Cl₃ (85 mg, 0.41 mmol). After stirring of 90 min, the resulting mixture was diluted with water and extracted with three portions of DCM. The combined DCM extracts were dried (MgSO₄), filtered and concentrated *in vacuo* to afford 1.924 g (19 mmol) of **34d**.

step 4 – A suspension of **34d** (199 mg, 8.91 mmol), 10% Pd/C (40 mg) were stirred in EtOH

10 (6 mL) under a hydrogen atmosphere 3 h. The resulting mixture was filtered and concentrated *in vacuo* to yield 0.177 g (0.916 mmol) of **36**.

step 5 – A solution of **36** (78 mg, 0.41 mmol) and trimethylorthoformate (2 mL) was heated to

105°C for 30 min. The resulting solution was concentrated *in vacuo*. A solution of **A-6** (100 mg, 0.41 mmol) in toluene (2 mL) was added. The resulting mixture was heated at reflux for 1 h,

15 concentrated *in vacuo*, and purified by SiO₂ chromatography eluting with 15% EtOAc/hexanes to afford 0.068 g (0.16 mmol) of **38**: MS (ESI) 419.2 (M+H)⁺.

step 6 – A solution of **B-3** (56 mg, 0.16 mmol, NR¹R² together are morpholine), **38** (68 mg, 0.16 mmol), Pd(0)(PPh₃)₄ (19 mg, 0.016 mmol), and Na₂CO₃ (52 mg, 0.49 mmol) in DME (2 mL) and H₂O (1 mL) was heated at 170°C for 12.5 min in a microwave synthesizer. The resulting


20 mixture was partitioned between EtOAc and H₂O. The EtOAc layer was washed with brine, dried (MgSO₄), filtered and concentrated *in vacuo*. The crude product was purified on a preparative SiO₂ TLC plate developed with 10% MeOH/DCM to afford 0.264 g (0.044 mmol) of **I-8**: MS (ESI) 606 (M+H)⁺.

Example 9: 7-*tert*-Butyl-3-(2-methyl-3-{1-methyl-5-[5-(morpholine-4-carbonyl)-pyridin-
25 2-ylamino]-6-oxo-1,6-dihydro-pyridazin-3-yl}-phenyl)-2,3-dihydro-1H-quinazolin-4-one (**I-34**)

To a mixture of **I-8** (11 mg, 0.018 mmol) and NaBCNH₃ (1.3 mg, 0.019 mmol) in 1 mL MeOH was added 1 drop of 2M HCl in MeOH to bring the pH to *ca.* 3. After stirring for 45 min, the resulting mixture was partitioned between EtOAc and dilute aqueous NaHCO₃. The EtOAc

30 layer was washed with brine, dried (MgSO₄), filtered and concentrated *in vacuo*. The crude product was purified on a SiO₂ preparative TLC plate developed with 5% MeOH/DCM to afford 6.5 mg (0.011 mmol) of **I-34**: MS (ESI) 608 (M+H)⁺.

Example 10: 4-*tert*-Butyl-*N*-{2-methyl-3-[1-methyl-5-(3-methyl-ureido)-6-oxo-1,6-dihydro-pyridazin-3-yl]-phenyl}-benzamide (I-23)



step 1 - A mixture of **40** (0.127 g, 0.774 mmol; CASRN 823-58-5) and 4% aqueous NaOH

5 solution (3 mL) was heated at reflux for 21 h, then cooled to RT. The pH was adjusted to 4 with glacial HOAc, then the mixture was partitioned between H₂O (20 mL) and EtOAc (30 mL). The aqueous layer was extracted with EtOAc (30 mL). The combined organic layers were dried (MgSO₄), filtered, and concentrated to give 0.082 g (73%) of **42a** as a pink solid that was used without further purification.

10 step 2 - A mixture of **42a** (0.082 g, 0.56 mmol) and K₂CO₃ (0.117 g, 0.845 mmol) in DMF (3 mL) was treated with methyl iodide (0.035 mL, 0.56 mmol), stirred for 42 h, then diluted with H₂O (20 mL). The mixture was twice extracted with EtOAc (20 mL each), and the combined organic layers were dried (MgSO₄), filtered, and concentrated. The crude product was purified by SiO₂ chromatography eluting with a EtOAc/hexane gradient (10 to 50% EtOAc) to afford

15 0.035 g (39%) of **42b** as an off-white solid.

step 3 - A mixture of **42b** (0.035 g, 0.22 mmol), **A-6** (0.102 g, 0.26 mmol), Na₂CO₃ (0.070 g,

0.66 mmol), and Pd(0)(PPh₃)₄ (0.025 g, 0.022 mmol) in DME (2 mL) and H₂O (1 mL) was

stirred and heated at 170°C for 30 min in a microwave synthesizer. The mixture was partitioned between EtOAc (30 mL) and H₂O (30 mL). The aqueous layer was extracted with two 30 mL

20 portions of EtOAc. The combined organic layers were dried (MgSO₄), filtered, and concentrated. The crude product was purified by SiO₂ chromatography eluting with a MeOH/DCM gradient (0 to 3% MeOH) to afford 0.079 g (92%) of **44** as a white solid. The material was not completely pure but was used as is in the next step.

step 4 - A solution of **44** (0.039 g, 0.100 mmol) and DIPEA (0.044 mL, 0.25 mmol) in DCM (2.5


25 mL) cooled to 0-5°C was treated with a 20% phosgene in toluene solution (0.25 mL, 0.55 mmol). The mixture was stirred for 10 min, then a 2 M methylamine in tetrahydrofuran solution (1.0 mL, 2.0 mmol) was added. The mixture was stirred for 1 h, treated with methanol, and adsorbed onto SiO₂. The SiO₂ was placed atop a SiO₂ flash column and eluted with a MeOH/EtOAc gradient

(0 to 4% MeOH) to afford 0.039 g (87%) of impure **I-23** as a white solid. This solid was combined with other batches and finally completely purified with a second SiO₂ chromatography eluting with a MeOH/DCM gradient (2 to 3% MeOH).

I-27 was prepared analogously except in step 4, the methylamine/THF solution was replaced

5 with an ethylamine/THF solution.

Example 11: 4-*tert*-Butyl-N-{2-methyl-3-[1-methyl-5-(3-methyl-ureido)-6-oxo-1,6-dihydro-pyridin-3-yl]-phenyl}-benzamide (II-4**)**



step 1 - A mixture of **46** (0.248 g, 1.22 mmol, CASRN 910543-72-5), **A-6** (0.574 g, 1.46 mmol), Na₂CO₃ (0.388 g, 3.66 mmol), and Pd(0)(PPh₃)₄ (0.141 g, 0.122 mmol) in DME (10 mL) and H₂O (5 mL) was stirred and heated at 170°C for 30 min in a microwave synthesizer. The mixture was partitioned between EtOAc (100 mL) and H₂O (100 mL). The aqueous layer was extracted with two 100 mL portions of EtOAc. The combined organic layers were dried (MgSO₄), filtered and concentrated. The crude product was purified by SiO₂ chromatography eluting with a MeOH/DCM gradient (0 to 4% MeOH) to afford 0.341 g (72%) of **48** as a yellow solid. The material was not completely pure but was used as is in the next step.


step 2 - A solution of **48** (0.136 g, 0.348 mmol) and DIPEA (0.15 mL, 0.87 mmol) in DCM (7 mL) at 0-5°C was treated with a 20% phosgene solution in toluene (0.19 mL, 0.42 mmol). The mixture was stirred for 10 min, then a 2 M methylamine THF solution (0.87 mL, 1.8 mmol) was added. The mixture was stirred for 15 min, then adsorbed onto SiO₂. The crude product was purified by two SiO₂ chromatographies eluting first with a MeOH/EtOAc gradient (0 to 5 % MeOH) then (0 to 3% MeOH) to afford 0.101 g (65%) of **II-4** as a white solid.

II-5, II-6, II-7, II-8, II-9, II-16, II-18 and **II-22** were prepared analogously except in step 2 methyl amine was replaced with ethylamine, *iso*-propylamine, 2,2,2-trifluoroethylamine, aniline, benzyl amine, N¹,N¹-dimethyl-ethane-1,2-diamine, 3-dimethylamino-propan-1-ol and N¹,N¹-dimethyl-propane-1,3-diamine respectively.

II-27 was prepared analogously except in step 1, **A-6** was replaced with 4-*tert*-butyl-N-[3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-benzamide (**49**) which was prepared by acylation of 3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenylamine with 4-*tert*-butyl-benzoyl chloride

- 5 **II-13** was prepared analogously except in the Suzuki coupling in step 1, **A-6** was replaced with **56**. **II-21** was prepared analogously except in step 2, methyl amine was replaced with N^1,N^1 -dimethyl-ethane-1,2-diamine.

Example 12: 4,5,6,7-Tetrahydro-benzo[b]thiophene-2-carboxylic acid {2-methyl-3-[1-methyl-5-(3-methyl-ureido)-6-oxo-1,6-dihydro-pyridin-3-yl]-phenyl}-amide
(**II-11**)



II-11 was prepared by Suzuki coupling of **46** and **50** and subsequently converting the amine the urea as described in steps 1 and 2 of example 11. **II-10** and **II-12** (dioxaborolane **38** was described in example 8) were prepared analogously except **50** was replaced with **53** (*infra*) and **38**, respectively.


The requisite dioxaborolanes were prepared by acylation **14a** (CASRN 882678-96-8) with 4,5,6,7-tetrahydrobenzo[b]thiophene-2-carbonyl chloride (CASRN 65361-26-4) and benzo[b]thiophene-2-carbonyl chloride (CASRN 39827-11-7).

Benzo[b]thiophene-2-carboxylic acid [2-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-amide **53**.

To a solution of **14a** (0.396 g, 1.7 mmol) and TEA (0.355 mL, 2.55 mmol) in THF (10 mL) was added a solution of benzo[b]-thiophene-2-carbonyl chloride (0.334 g, 1.7 mmol) in THF (2 mL). The white suspension was stirred for 1 h, then H₂O (5 mL) was added. The mixture was partitioned between EtOAc (30 mL) and H₂O (30 mL), and the aqueous layer was extracted with EtOAc (30 mL). The combined organic extracts were dried (MgSO₄), filtered, and concentrated to a solid. The crude product was purified by SiO₂ chromatography eluting with a EtOAc/hexane gradient (0 to 15% EtOAc) to afford 0.399g (60%) of benzo[b]thiophen-2-

carboxylic acid [2-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-amide (**53**) as a white solid.

Example 13: 4-*tert*-Butyl-N-(2-methyl-3-{1-methyl-5-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-6-oxo-1,6-dihydro-pyridin-3-yl}-phenyl)-benzamide (II-2)



step 1 - To a suspension of **50** (0.80 g, 3.0 mmol, CASRN 14529-54-5), methyl 6-amino-nicotinate (**52**, 0.55g, 3.6 mmol, CASRN 36052-24-1) and toluene (10 mL) was added BINAP (0.140 g, 0.22 mmol), Pd₂(dba)₃ (0.137 g, 0.15 mmol), Cs₂CO₃ (1.37 g, 4.2 mmol). The glass reaction vessel was sealed and heated to 130°C overnight. The reaction mixture was cooled, concentrated *in vacuo* and purified by SiO₂ chromatography eluting with 1%MeOH/DCM to afford 1.80 g of **54a**.

step 2 - To a solution of **54a** (0.75 g, 2.22 mmol) and EtOH (100 mL) was added 3 N NaOH (2.22 mL). The solution was heated at reflux for 2 h, cooled, concentrated *in vacuo* and dried under a high vacuum. The residue was dissolved in DMF (50 mL) and EDCI (1.28 g, 6.65 mmol), HOBt (0.90 g, 6.66 mmol) and morpholine (0.29 mL, 3.33 mmol) were added and the resulting reaction mixture stirred at RT overnight. The solution was diluted with H₂O (50 mL) and stirred overnight. The reaction mixture was extracted with DCM and the organic extracts washed with Na₂CO₃ and brine. The DCM solution was dried (MgSO₄), filtered and evaporated. The crude product was purified by SiO₂ chromatography eluting with 2% MeOH/DCM to afford 0.30 g of **54b**.

step 3 - A solution of **54b** (0.110 g, 0.28 mmol), A-6 (0.132 g, 0.33 mmol), Pd(0)(PPh₃)₄ (0.032 g, 0.03 mmol) in 1 N Na₂CO₃ (1 ml) and DME (1 mL) was heated at 120°C for 40 min in a microwave synthesizer. The solution was cooled and partitioned between EtOAc and H₂O. The organic extract was washed with brine, dried (MgSO₄), filtered and evaporated. The crude

product was purified by SiO₂ chromatography eluting with 4% MeOH/DCM to afford 0.062 g of **II-2**.

II-1 and **II-25** were prepared analogously except in step 2, morpholine was replaced with 1-methyl-piperazine and pyrrolidin-3-ol, respectively.

- 5 **II-3** was prepared analogously except in step 1, methyl 6-amino-nicotinate was replaced with 2-amino-pyridine and step 2 was omitted.

II-36 was prepared analogously except in step 3, **A-6** was replaced with **60** (see example 16). **II-34** was prepared by Suzuki-mediated coupling of **54b** and **14a**. The resulting aryl amine was acylated with 4-*tert*-butyl-3-methoxy-benzoyl chloride to afford **II-34** which was demethylated
10 with BBr₃ to afford **II-32**.

II-48 is prepared analogously except in step 3, **A-6** is replaced with **60** (Ar = 4-dimethylamino-phenyl, see example 16).


II-95 is prepared analogously except in step I, **52** is replaced with pyridazin-3-ylamine and in step 3, **A-6** is replaced with **120** (Ar = 4-cyclopropyl-phenyl).

- 15 **II-94** is prepared analogously except in step I, **52** is replaced with 3-amino-6-methyl-pyridazine and in step 3, **A-6** is replaced with **120** (Ar = 4-cyclopropyl-phenyl).

II-98 is prepared analogously except in step I, **52** is replaced with pyrazin-2-ylamine and in step 3, **A-6** is replaced with **120** (Ar = 4-cyclopropyl-phenyl).

- 20 **II-103** is prepared analogously except in step I, **52** is replaced with 6-morpholin-4-yl-pyridazin-3-ylamine and in step 3, **A-6** is replaced with **120** (Ar = 4-cyclopropyl-phenyl).

Example 14: 4-Dimethylamino-N-(3-{5-[5-(4-hydroxy-piperidine-1-carbonyl)-pyridin-2-ylamino]-1-methyl-6-oxo-1,6-dihydro-pyridin-3-yl}-2-methyl-phenyl)-benzamide (II-20)



step 1 – To a solution of 4-dimethylamino-benzoyl chloride (1.49 g, 8.15 mmol) and TEA (1.2 g, 12 mmol) and DCM (60 mL) under nitrogen was added **14a** (1.9 g, 8.15 mmol). The mixture was stirred at RT for 16 h, then diluted with water and extracted with EtOAc. The organic layer was washed with brine, dried (Na₂SO₄), filtered and concentrated *in vacuo*. The residue was purified by SiO₂ chromatography eluting with a EtOAc/hexane gradient (10% to 50% EtOAc) affording 2 g (66%) of **56**.

step 2 – A suspension of **50** (5 g, 18.7 mmol) and **62** (3.4 g, 22.5 mmol) in toluene (60 mL) was sparged with argon, racemic BINAP (0.931 g, 1.5 mmol) was added followed by Pd₂(dba)₃ (0.514 g, 0.56 mmol) and sodium *tert*-butoxide (2.51 g, 26.18 mmol). The mixture was heated at 120°C for 16 h. After cooling the reaction mixture was diluted with H₂O (200 mL) and twice extracted with EtOAc (150 mL each). The organic layer was concentrated *in vacuo*, the residue triturated with EtOAc and filtered. The solid was washed with water, MeOH and EtOAc and dried to afford 6 g of **58** which used in the next step without further purification.

step 3 – To a suspension of **56** (0.562 g, 1.47 mmol) and **58** (0.5 g, 1.47 mmol) in DME (10 mL) in a microwave tube was added a solution of Na₂CO₃ (0.467 g, 4.41 mmol) in H₂O (3 mL). After sparging with argon, Pd(0)(PPh₃)₄ (0.085 g, 0.0735 mmol) was added, the tube was sealed and the reaction mixture was heated at 175°C for 45 min in a microwave apparatus. The reaction mixture was partitioned between H₂O and EtOAc. The aqueous layer was acidified with 4N HCl to *ca.* pH 5 and thrice extracted with EtOAc to afford 0.5 g (68%) of **60**.

step 4 – To a solution of **60** (0.1 g, 0.2 mmol) and anhydrous DMF (3 mL) maintained under a nitrogen atmosphere was added BOP (0.088 g, 0.2 mmol) followed by TEA (0.060 g, 0.6 mmol) and piperidin-4-ol (0.02 g, 0.2 mmol). The mixture was stirred at RT for 6 h. After this time, the mixture was diluted with water, extracted with EtOAc, washed with brine and concentrated *in*

vacuo. The residue was purified on a preparative TLC plate three times developed with 5% MeOH/DCM to afford 0.024 g (20%) of **II-20** (yield 20%).

II-33 was prepared analogously except in step 4, 1-methyl-piperazine was used in place of piperidin-4-ol. **II-35** was prepared analogously except in step 3, **56** is replaced with **88** (example 21) and in step 4, 1-methyl-piperazine is used in place of piperidin-4-ol. **II-87** is prepared analogously except in step 1 4-dimethylamino-benzoyl chloride is replaced with 4-cyclopropyl-benzoyl chloride and in step 4, 1-ethyl-piperazine is used in place of 1-methyl-piperazine.

II-60 is prepared analogously except in step 3, **56** is replaced with **106** and in step 4, piperidin-4-ol is replaced with 4-dimethylamino-piperidine.

II-99 is prepared analogously except in step 3, **56** is replaced with **106** and in step 4, piperidin-4-ol is replaced with pyrrolidine.

II-100 is prepared analogously except in step 3, **56** is replaced with **106** and in step 4, piperidin-4-ol is replaced with 3,3-difluoroazetidine (CASRN 679431-52-8).

II-101 is prepared analogously except in step 3, **56** is replaced with **106** and in step 4, piperidin-4-ol is replaced with azetidine.

Example 15: 1-Methyl-1H-indole-5-carboxylic acid {2-methyl-3-[1-methyl-5-(3-methyl-ureido)-6-oxo-1,6-dihydro-pyridin-3-yl]-phenyl}-amide (II-23**)**

step 1 - Sodium hydride (1.71 g, 42.80 mmol, 60% in mineral oil) was added at 0°C in portions to a stirred solution of methyl indole-5-carboxylate (5.0 g, 28.53 mmol, CASRN 1670-81-1) in DMF over a 30 min period. While still at 0°C MeI (5.33 mL, 85.61 mmol) was added. The resulting mixture was stirred at 0°C for 60 min then at RT overnight. The mixture was partitioned between H₂O and EtOAc and the aqueous layer was further extracted with EtOAc. The combined organic extracts were washed with water, dried (MgSO₄), filtered and concentrated to afford 4.91 g of 1-methyl-1H-indole-5-carboxylic acid methyl ester (**53**).


step 2 - To the solution of **53** (4.90g, 25.89 mmol) in THF/H₂O was added 1N NaOH and the resulting mixture was heated at reflux for 60 min. The reaction mixture was cooled to 0°C, acidified with 1N HCl to *ca.* pH=3, and the mixture was partitioned between H₂O and EtOAc and the combined organic layers were dried (MgSO₄), filtered and concentrated to afford 4.25 g of 1-methyl-1H-indole-5-carboxylic acid (**55**).

step 3 - To a solution of **55** (0.608 g, 3.474 mmol, CASRN 186129-25-9) and DMF was added **14a** (0.27 g, 1.158 mmol), DIPEA (0.60 mL, 3.474 mmol) and HATU (0.484 g, 1.274 mmol). The reaction mixture was stirred at RT over night. The reaction mixture was partitioned between EtOAc and H₂O, the EtOAc phase was washed with brine, dried (MgSO₄), filtered and

concentrated. The crude product was purified by SiO₂ chromatography eluting with 42% EtOAc/hexane to afford 236 mg of **59**.

The synthesis of **II-23** was completed by Suzuki coupling of **59** and **46** followed by sequential treatment with phosgene and methylamine as described in example 11

5 **Example 16: 4-(Cyano-dimethyl-methyl)-N-{2-methyl-3-[1-methyl-5-(3-methyl-ureido)-6-oxo-1,6-dihydro-pyridin-3-yl]-phenyl}-benzamide (II-42)**




step 1 – To a solution of 4-(cyano-dimethyl-methyl) benzoic acid (0.5 g, 2.64 mmol, CASRN 129488-74-0) and DMF (15 mL) was added BOP (1.16 g; 2.64 mmol), TEA (0.8 g; 7.92 mmol) and **14a** (0.615 g, 2.64 mmol). The mixture was stirred at RT for 16 h, then diluted with EtOAc and washed with brine. The organic layer was dried (Na₂SO₄), filtered and concentrated *in vacuo*. The residue was triturated with EtOAc and the solid was filtered to afford 0.650 g (61%) of **60**. Additional impure product (0.3 g) was recovered from filtrate.

step 2 – To a solution of **46** (0.5 g, 2.46 mmol), DIPEA (0.952 g; 7.38 mmol) and DCM (40 mL) cooled to 0°C under a nitrogen atmosphere was added phosgene (1.8 mL, 3.69 mmol, 20% in toluene). The reaction mixture was stirred for 20 min then methylamine (12 mL, 24.6 mmol, 2M solution in THF) was added slowly. The mixture warmed to RT and stirred for 5 h. The product was filtered, washed with DCM and dried *in vacuo* to afford 0.8 g (99%, 80% pure) of **62** which was used in the following step without further purification.

step 3 – To a suspension of **60** (0.153 g, 0.38 mmol) and **62** (0.125 g, 0.38 mmol) in DME (3 mL) in a microwave tube was added Na₂CO₃ (0.120 g, 1.14 mmol) in H₂O (1.5 mL). After sparging with argon, Pd(0)(PPh₃)₄ (22 mg, 0.019 mmol) was added, the tube was sealed and the mixture heated at 175°C in a microwave apparatus. The reaction mixture was cooled, diluted with water and extracted with EtOAc. The organic layer was concentrated and the precipitate filtered and

washed with H₂O (2 mL), MeOH (2 mL) and DCM (4 mL), then dried, to afford 0.058 g (33%) of II-42.


Example 17: 4-(1-Hydroxy-1-methyl-ethyl)-N-{2-methyl-3-[1-methyl-5-(3-methyl-ureido)-6-oxo-1,6-dihydro-pyridin-3-yl]-phenyl}-benzamide (II-28)



step 1 - A solution of **86** (2.10 g, 11.653 mmol) and **14a** in DMF (30 mL) was stirred at RT and DIPEA (5.53 mL, 31.78 mmol) and HATU (4.43 g, 11.65 mmol) were added and the resulting mixture was stirred at RT over night. The mixture was partitioned between H₂O (150 mL) and EtOAc (150 mL). The aqueous layer was further extracted with 50 mL of EtOAc. The combined organic extracts were thrice washed with H₂O (50 mL each), dried (MgSO₄), filtered and concentrated to afford 2.90 g of **88**.

The Suzuki coupling of **88** and **62** was carried out as described in step 3 of example 16.

Example 18: 5-{3-[2-(1,3-Dihydro-isoindol-2-yl)-2-oxo-ethyl]-phenyl}-1-methyl-3-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-1H-pyridin-2-one (II-24)



(6-chloro-pyridin-3-yl)-morpholin-4-yl-methanone

To a solution of 6-chloronicotinic acid (25.0 g, 142 mmol) and DCM cooled to a 0°C was added morpholine (13.6 g, 156 mmol) and TEA (28.7 g, 284 mmol). After stirring for 2 h at 0°C, the solution was washed with aqueous NaHCO₃, dried (MgSO₄), filtered and concentrated *in vacuo* to afford 29.5g (130mmol) of **66**: MS (ESI) 227.1 (M+H)⁺.


- 5 step 1 - To a 0°C solution of **66** (14.00 g, 61.77 mmol) and **46** (12.54 g, 61.77 mmol) in DMF (160 mL) was added sodium hydride (95%, 3.12 g, 123.5 mmol). This solution was slowly warmed to RT, stirred for 18 h then poured into 1.3 L cold water. EtOAc (3mL) was added and the resulting precipitate was filtered, washed with water, and air dried for 3 d to afford 18.84g (47.91mmol) of **68**: MS (ESI) 393.1 (M+H)⁺.

- 10 The Suzuki coupling of **68** and **28** was carried out as described in step 1 of example 8 to afford **72** which was converted to **II-24** by EDCI coupling of 2,3-dihydro-1H-isoindole using the protocol described in step 2 of example 14.

II-26, **II-40** and **II-41** were prepared analogously except 6,7-dihydro-5H-pyrrolo[3,4-b]pyridine (CASRN 147739-88-6, 3-*tert*-butoxy-cyclobutylamine and 3-*iso*-propoxy-cyclobutylamine

- 15 respectively were used in place of 2,3-dihydro-1H-isoindole.

Example 19: N-(2-*tert*-Butoxy-ethyl)-2-(2-methyl-3-{1-methyl-5-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-6-oxo-1,6-dihydro-pyridin-3-yl}-phenyl)-acetamide (II-39)



step 1 - To a solution of **74a** (1.00 g, 4.65 mmol) in DCM (15 mL) was added oxalyl chloride (649 mg, 5.12 mmol) and 4 drops DMF. After 40 min, the resulting mixture was concentrated *in vacuo*. The residue was dissolved in toluene (20 mL) and (trimethylsilyl)diazomethane solution (6.30 mL, 12.6 mmol, 2.0M in Et₂O) was added. After stirring for 3 d, the solution was

5 partitioned between EtOAc and H₂O. The organic layer was washed with brine, dried (MgSO₄), filtered and concentrated *in vacuo*. The crude product was purified by SiO₂ chromatography and eluted with a EtOAc/hexane gradient (50 to 100% EtOAc) to afford 0.394 g of **74b**: MS (ESI) 238.9 (M-H).

step 2 - A solution of **74b** (115 mg, 0.479 mmol) in benzyl alcohol (0.5 mL) and 2,4,6-collidine (0.5 mL) was heated to 180°C for 7 min in a microwave synthesizer. The resulting solution was

10 partitioned between EtOAc and 1M aq. HCl. The organic layer was washed with water and brine, dried (MgSO₄), filtered and concentrated *in vacuo*. The crude product was purified by SiO₂ chromatography eluting with 5% EtOAc/hexanes to afford 0.072 g of **76**: ¹HNMR (300MHz, CDCl₃) δ 2.36 (s, 3H), 3.73 (s, 2H), 5.14 (s, 2H), 7.00 (t, *J* = 8 Hz, 1H), 7.14 (d, *J* = 8 Hz, 1H), 7.26-7.36 (m, 5H), and 7.49 ppm (d, *J* = 8 Hz, 1H).

step 3 - To a solution of **76** (200 mg, 0.626 mmol), *bis*-(pinacolato)diboron (191 mg, 0.752 mmol), KOAc (184 mg, 1.88 mmol) in DMSO (4mL) was added a solution of [1,1'-*bis*-(di-phenylphosphino)ferrocene]dichloropalladium(II) in DCM (15 mg, 0.018 mmol) and the resulting solution was heated at 80°C under an atmosphere of argon for 2 h. The resulting solution

20 was partitioned between EtOAc and H₂O. The organic layer was washed with brine, dried (MgSO₄), filtered and concentrated *in vacuo*. The crude product was purified by SiO₂ chromatography eluting with 4% EtOAc/hexanes to afford 0.094 g of **78**: MS (ESI) 389.2 (M+Na)⁺.

step 4 - A solution of **54b** (99 mg, 0.25 mmol), **78** (92 mg, 0.25 mmol), Pd(0)(PPh₃)₄ (29 mg, 0.025 mmol), and Na₂CO₃ (80 mg, 0.75 mmol) in DME (3 mL) and H₂O (1 mL) was heated at 170°C for 10 min in a microwave synthesizer. The resulting mixture was partitioned between EtOAc and water. The EtOAc layer was washed with brine, dried (MgSO₄), filtered and concentrated *in vacuo*. The crude product was purified by on a preparative TLC plate developed with 5 % MeOH/DCM) to afford 36 mg of **80a**: MS (ESI) 553.2 (M+H)⁺.


step 5 - A suspension of **80a** (36 mg, 0.065 mmol) and 10% Pd/C (22 mg) and EtOH (15 mL) was stirred under H₂ atmosphere for 2 h. The resulting mixture was filtered and concentrated *in vacuo* to afford 0.020 g of **80b**: MS (ESI) 463.1 (M+H)⁺.

step 6 - To a solution of **80b** (30 mg, 0.065 mmol), BOP (29 mg, 0.066 mmol), and 2-*tert*-butoxy-ethylamine hydrochloride (10 mg, 0.065 mmol, CASRN 335598-67-9) in DMF (2 mL) was

added TEA (33 mg, 0.33 mmol). The solution was stirred at RT for 5 h and the resulting mixture was partitioned between EtOAc and aqueous NaHCO₃. The EtOAc layer was washed with brine, dried MgSO₄, filtered and concentrated *in vacuo*. The crude product was purified on a preparative SiO₂ TLC plate developed with 5% MeOH/DCM) to afford 0.0092 g of **II-39**: MS (ESI) 562 (M+H)⁺.

II-43 was prepared analogously except in step 6, 2-*tert*-butoxy-ethylamine hydrochloride was replaced with 2,3-dihydro-1H-isoinidole

Example 20: 6-Dimethylamino-N-{2-methyl-3-[1-methyl-5-(3-methyl-ureido)-6-oxo-1,6-dihydro-pyridin-3-yl]-phenyl}-nicotinamide (II-15)



step 1 – To a solution of **82a** (1.0 g, 6.34 mmol) and THF (15 mL) was added Me₂NH (6.35 mL, 12.69 mmol; 2.0 M solution in THF) and the mixture was stirred in a microwave and heated to 150°C for 45 min. The solvent was evaporated to afford 1.0 g of **82b** which was used in the next step.

step 2 – A solution of **82b** (0.60 g, 3.61 mmol) and **14a** (0.280g, 1.20 mmol) and DMF (20 mL) was stirred at RT. DIPEA (0.63 mL, 3.61 mmol) and HATU (0.50 g, 1.32 mmol) were added sequentially and the resulting mixture was stirred at RT overnight. The mixture was partitioned between H₂O and EtOAc. The aqueous layer was separated and further extracted with EtOAc (50 mL). The combined organic extracts were thrice washed with H₂O (50 mL each), dried (MgSO₄), filtered and concentrated. The crude product was purified by SiO₂ chromatography eluting with a EtOAc/hexanes gradient (0-50% EtOAc) to afford 0.140 g of **84**.


The Suzuki coupling of **84** and **62** was carried out as described in step 4 of example 17.

II-19 was prepared analogously except in step 1, dimethylamine was replaced with piperidine.

II-50 can be prepared analogously except **84** is replaced with 4-*tert*-butyl-N-[2-methyl-3-

(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-benzamide and **62** is replaced with 1-(5-bromo-2-oxo-1,2-dihydro-pyridin-3-yl)-3-methyl-urea.

Example 21: *N*-(2-Hydroxymethyl-3-{1-methyl-6-oxo-5-[5-(piperazine-1-carbonyl)-pyridin-2-ylamino]-1,6-dihydro-pyridin-3-yl}-phenyl)-4-(2,2,2-trifluoro-1,1-dimethyl-ethyl)-benzamide (**II-81**)



step 1 - To a solution of **144** (2.16 g, 12.27 mmol) and TEA (3.42 mL, 2 eq) in DCM (50 mL) at 0°C was added **146** (2.42 mL, 1 eq) and the resulting mixture was stirred and warmed from 0°C to RT over one h. The DCM was removed under reduced pressure at 55°C and the residue partitioned between EtOAc (200 mL) and water (50 mL). The organic phase was washed sequentially with water (2 x 50 mL), saturated NaHCO₃ (3 x 50 mL) and brine (1 x 50 mL). The organic layer was dried (MgSO₄), filtered, and concentrated to afford 4.4 g of **148** as a viscous colorless oil.

step 2 - Condensation of **148** and **46** can be carried out as described in step 1 of example 18 to afford **150**.


step 3 - Palladium-catalyzed coupling of **120** (Ar = 4-(2,2,2-trifluoro-1,1-dimethyl-ethyl)-phenyl) and **150** can be carried out as described in step 5 of example 25 to afford **152a**.

step 4 - A solution of **152a** (73 mg, 0.093 mmol) in EtOH (40 mL) and THF (10 mL) and was purged with nitrogen gas for five min then 10% Pd/C carbon (50 mg) was carefully added. The resulting mixture was stirred under 1 atm of hydrogen gas (balloon) at RT for about 18 h. The mixture was filtered through a 3 cm bed of SOLKA-FLOC and the filtrate was concentrated.

The crude product was purified on a preparative SiO₂ TLC plate developed with 30% MeOH/DCM to afford 7 mg of **II-81** as an off-white powder.

II-82 is prepared analogously except **152b** is formylated with one equivalent of acetic formic mixed anhydride.

Example 22: 1-(5-{3-[2-(3-*tert*-Butoxy-azetidin-1-yl)-2-oxo-ethyl]-phenyl}-1-methyl-2-oxo-1,2-dihydro-pyridin-3-yl)-3-methyl-urea (**II-29**)



The title compound is prepared by Suzuki coupling (step 2) of **92a** and **62** using the protocol described in step 1 of example 6. The amide **92a** is prepared (step 2) by BOP-mediated coupling of **28** and 3-*tert*-butoxy-azetidine using the protocol described in step of example 6.


II-30 (3-*iso*-propoxy-azetidine) and **II-31** (5,6-dichloro-2,3-dihydro-1H-isoindole) were

prepared analogously except in step 2, 3-*tert*-butoxy-azetidine was replaced by the secondary amine in parenthesis.

II-45 and **II-46** can be prepared analogously except in step 1, **28** is coupled with 2H-isoindole and 5-fluoro-2H-isoindole respectively.

II-47 can be prepared analogously except in step 1, **28** is coupled with azetidin-3-yl-propan-2-ol using HATU as described in example 21.

Example 23: 1,3-Dihydro-isoindole-2-carboxylic acid {2-methyl-3-[1-methyl-5-(3-methyl-ureido)-6-oxo-1,6-dihydro-pyridin-3-yl]-phenyl}-amide (**II-17**)




step 1 - Phosgene (0.270 mL, 0.513 mmol, 20% in toluene) was added to a solution of **14a**

(0.100 g, 0.43 mmol) and TEA (0.150 mL, 1.08 mmol) in DCM (6 mL) cooled to 0-5°C. The

reaction mixture was stirred for 10 min, then 2,3-dihydro-1H-isoindole (0.122 mL, 1.08 mmol) was added. The brown mixture was stirred at 0-5°C for 10 min. To the solution was added SiO₂ (10 g) of silica gel and the mixture was concentrated to a powder. The resulting solid was applied to the top of a SiO₂ column eluting with a EtOAc/hexanes gradient (0 to 40% EtOAc) to afford 0.150 g (92%) of **94** as a dark brown solid that was used without further purification. Suzuki coupling of **14a** and **94** was carried out utilizing the protocol described in step 1 of example 6 to afford **II-17**.

II-14 was prepared analogously except in step 1, 2,3-dihydro-1H-isoindole was replaced by 5-fluoro-2,3-dihydro-1H-isoindole (CASRN 57584-71-1).

Example 24: 4-Dimethylamino-N-(3-{5-[5-(4-hydroxy-piperidine-1-carbonyl)-pyridin-2-ylamino]-1-methyl-6-oxo-1,6-dihydro-pyridazin-3-yl}-2-methyl-phenyl)-benzamide (I-28)




step 1 – To a solution methyl 6-amino-nicotinate (0.234 g, 1.53 mmol) and DMF (4 mL) cooled to 0°C was added NaH (61.2 mg, 60% mineral oil dispersion) and the solution warmed to RT and stirred for 0.5 h. The resulting solution was re-cooled to 0°C and a solution of **A-1b** (0.137 g, 0.76 mmol) and DMF (1 mL) was added and the resulting solution stirred for 4 h at RT. The solution was diluted with EtOAc and brine which resulted in the formation of an insoluble solid which was filtered and dried to afford 0.140 g of **96**.

step 2 – To a suspension of **96** (0.140 g, 0.47 mmol) and **56** (0.1809 g, 0.47 mmol) in a microwave tube containing DME (5 mL) was added a solution of Na₂CO₃ in water (2 mL). The tube was flushed with N₂ and Pd(0)(PPh₃)₄ (0.050, 0.047 mmol) was added and the tube sealed and heated to 175°C for 45 min in a microwave. The resulting mixture was cooled to RT and partition between EtOAc and brine. The ester hydrolyzed during the coupling procedure and the aqueous brine solution was acidified with 2N HCl, extracted with EtOAc. The aqueous solution was extracted with EtOAc and the combined extracts were dried (Na₂SO₄), filtered and evaporated to afford 0.060 g of **98**.

step 3 – To a solution of **98** (0.060 g, 0.12 mmol) and BOP (0.053 g, 0.12 mmol) in DMF (3 mL) was added 4-hydroxy-piperidine (0.012 g, 0.21 mmol) and TEA (0.036 g, 0.36 mmol) and the resulting solution was stirred at RT for 8 h. The solution was partition between EtOAc and brine. The EtOAc phase was dried (Na₂SO₄), filtered and evaporated. The crude product was purified on a preparative SiO₂ plate developed with 5% MeOH/DCM and eluted to afford **I-28**.

Example 25: 4-Dimethylamino-N-(2-hydroxymethyl-3-{1-methyl-5-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-6-oxo-1,6-dihydro-pyridin-3-yl}-phenyl)-benzamide (II-49)



step 1 - To a solution of above **100a** (1.1 g, 4.74 mmol, CASRN 861106-91-4) in DCM (10 mL) were added sequentially ^tBuMe₂SiCl (0.86 g, 5.69 mmol) and TEA (0.8 mL, 5.69 mmol). The resulting mixture was stirred at RT until the reaction was complete. The reaction mixture was partitioned between EtOAc (150 mL) and water (100 mL). The organic layer was dried (MgSO₄), filtered and concentrated. The residue was then purified by SiO₂ chromatography eluting with an EtOAc/hexane gradient (0 to 50% EtOAc) to afford 1.5 g of **100b**.

step 2 - A mixture of the **102b** (1.5 g, 6.46 mmol), iron powder (1.08 g, 19.4 mmol) and ammonium chloride (1.73 g, 32.3 mmol) in EtOH (25 mL) and water (25 mL) was heated at 70°C for 2 h. The mixture was filtered and extracted with EtOAc (100 mL). The organic layer was dried (MgSO₄), filtered and concentrated to afford 0.87 g of **102**.

step 3 - A mixture of the **102** (0.87 g, 2.76 mmol), PdCl₂(dppf) (10 mg, 0.012 mmol), pinacol diborane (0.84 g, 3.31 mmol), KOAc (0.974 g, 9.94 mmol) in DMSO (4 mL) was heated at 70°C overnight. The mixture was then partitioned between EtOAc (100 mL) and water (100 mL). The organic layer was dried (MgSO₄), filtered and concentrated. The residue was purified by SiO₂

chromatography eluting with an EtOAc/hexane gradient (0 to 30% EtOAc) to afford 0.090 g of **104**.

step 4 - To a mixture of **104** (90 mg, 0.248 mmol) and *p*-N,N-dimethylamino-benzoyl chloride (45 mg, 0.248 mmol) in DCM (3 mL), was added dropwise TEA (54 μ L, 0.372 mmol). The resulting solution was stirred at RT for 0.5 h and purified by SiO₂ chromatography eluting a EtOAc/hexane gradient (0 to 50% EtOAc) to afford 100 mg of **106**.

step 5 - A mixture of above **106** (100 mg, 0.196 mmol), **54b** (77 mg, 0.196 mmol), Pd(Ph₃P)₄ (10 mg, 0.009 mmol), and Na₂CO₃ (62 mg, 0.588 mmol) in DME (1.5 mL) and water (1.5 mL) was heated at 170°C for 0.5 h in a microwave. The resulting mixture was partitioned between EtOAc (100 mL) and water (100 mL). The organic layer was then dried (MgSO₄), filtered and concentrated. The residue was purified on a preparative SiO₂ TLC plate developed with 8% MeOH/DCM to afford 0.035 g of **II-49** as white solid: MS: 583 (M+1)⁺.

II-52 can be prepared analogously, except in step 4, *p*-N,N-dimethylamino-benzoyl chloride is replaced with 4-(1-fluoro-1-methyl-ethyl)-benzoyl chloride. The latter is prepared from 4-(1-fluoro-1-methyl-ethyl)-benzoic acid (CASRN 477219-30-0) by standard methodology.

II-53 is prepared analogously, except in step 4, *p*-N,N-dimethylamino-benzoyl chloride is replaced with 4-(1-fluoro-cyclopropyl)-benzoyl chloride. The latter is prepared from 4-(1-fluoro-cyclopropyl)-benzoic acid (CASRN 946118-80-5) by standard methodology.

II-74 is prepared analogously, except in step 4, *p*-N,N-dimethylamino-benzoyl chloride is replaced with 4-trichloromethyl-benzoyl chloride.

II-75 is prepared analogously, except in step 4, *p*-N,N-dimethylamino-benzoyl chloride is replaced with 4-(1-chloro-cyclopropyl)-benzoyl chloride.

II-55 is prepared analogously, except in step 4, *p*-N,N-dimethylamino-benzoyl chloride is replaced with 4-trifluoromethyl-benzoyl chloride (CASRN 329-15-7).

II-58 is prepared analogously, except in step 4, *p*-N,N-dimethylamino-benzoyl chloride is replaced with 4-*tert*-butyl-benzoyl chloride (CASRN 329-15-7).

II-64 is prepared analogously except in step 4, *p*-N,N-dimethylamino-benzoyl chloride is replaced with is replaced with 4-(1-fluoro-cyclopropyl)-benzoyl chloride and in step 5, **54b** is replaced with 5-bromo-1-methyl-3-(pyridin-2-ylamino)-1H-pyridin-2-one.

II-59 can be prepared analogously except in step 5, **54b** is replaced with 5-bromo-1-methyl-3-(pyridin-2-ylamino)-1H-pyridin-2-one which is prepared as described in step 1 of example 13 wherein methyl 6-amino-nicotinate is replaced by 2-amino-pyridine.

II-62 is prepared analogously, except in step 4, *p*-N,N-dimethylamino-benzoyl chloride is replaced with 4-cyclopropyl-benzoyl chloride (CASRN 76274-96-77).

II-63 is prepared analogously, except in step 4, *p*-N,N-dimethylamino-benzoyl chloride is replaced with 4-bromo-benzoyl chloride.

II-66 is prepared analogously, except in step 4, *p*-N,N-dimethylamino-benzoyl chloride is replaced with 4-(1-methoxy-1-methyl-ethyl)-benzoyl chloride. The latter is prepared from 4-(1-methoxy-1-methyl-ethyl)-benzoic acid (CASRN 50604-11-0) by standard methodology.

II-67 is prepared analogously, except in step 4, *p*-N,N-dimethylamino-benzoyl chloride is replaced with benzoyl chloride.

II-68 is prepared analogously, except in step 4, *p*-N,N-dimethylamino-benzoyl chloride is replaced with 4-(1,2,2,2-tetrafluoro-1-trifluoromethyl-ethyl)-benzoyl chloride (CASRN 62480-31-3).

II-69 is prepared analogously, except in step 4, *p*-N,N-dimethylamino-benzoyl chloride is replaced with 4-ethynyl-benzoyl chloride (CASRN 62480-31-3).

II-96 is prepared analogously, except in step 4, *p*-N,N-dimethylamino-benzoyl chloride is replaced with 4-(cyano-dimethyl-methyl)-benzoyl chloride (CASRN 129488-75-1).

II-89 is prepared analogously, except in step 4, *p*-N,N-dimethylamino-benzoyl chloride is replaced with 4-pentafluorothio-benzoic acid and the coupling of the carboxylic acid to **104** is carried out with HATU as described in step 1 of example 21.

II-76 is prepared analogously, except in step 4, *p*-N,N-dimethylamino-benzoyl chloride is replaced with trimethylsilyl-benzoic acid (CASRN 15290-29-6) and the coupling of the

carboxylic acid to **104** is carried out with HATU as described in step 1 of example 21.

II-97 is prepared analogously, except in step 4, *p*-N,N-dimethylamino-benzoyl chloride is replaced with 4-piperidin-1-yl-benzoic acid (CASRN 22090-24-0) and the coupling of the carboxylic acid to **104** is carried out with HATU as described in step 1 of example 21.

II-90 is prepared by palladium-catalyzed coupling of **50** and 2-amino-6-ethoxy-pyridine (CASRN 768-42-3) which can be carried out as described in step 1 of example 13 to afford 5-bromo-3-(6-ethoxy-pyridin-2-ylamino)-1-methyl-1H-pyridin-2-one (**105**). Condensation of **105** with **120** (Ar is 4-cyclopropyl-phenyl) is carried out as described in step 5 of the present example.


II-91 is prepared by palladium-catalyzed coupling of **50** and 6-amino-2-ethyl-2H-pyridazin-3-one (CASRN 50500-52-2) which can be carried out as described in step 1 of example 13 to afford 6-(5-bromo-1-methyl-2-oxo-1,2-dihydro-pyridin-3-ylamino)-2-ethyl-2H-pyridazin-3-one (**107**). Condensation of **107** with **120** (Ar is 4-cyclopropyl-phenyl) is carried out as described in step 5 of the present example.

II-93 is prepared by coupling of **42b** and 2-benzyl-6-chloro-2H-pyridazin-3-one which can be carried out as described in step 1 of example 10 to afford 2-benzyl-6-(5-chloro-1-methyl-2-oxo-

1,2-dihydro-pyridin-3-ylamino)-2H-pyridazin-3-one (**109**). Condensation of **109** with **120** (Ar is 4-cyclopropyl-phenyl) is carried out as described in step 5 of the present example.

II-102 is prepared analogously except in step 5, **106** is replaced with 1,3-dihydro-isindole-2-carboxylic acid [2-(tert-butyl-dimethyl-silanyloxymethyl)-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-amide which can be prepared treating **104** sequentially with phosgene and 2,3-dihydro-1H-isindole according to the procedure in step 1 of example 23.

Example 26: *N*-(2-Hydroxymethyl-3-{1-methyl-5-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-6-oxo-1,6-dihydro-pyridin-3-yl}-phenyl)-4-(2,2,2-trifluoro-1-methyl-ethyl)-benzamide (**II-51**)



step 1 - To a solution of **110** (4 g, 22.4 mmol) in THF (120 mL) was added trimethyl-trifluoromethyl-silane (9.5 g, 67 mmol). The mixture was cooled to 0°C and a solution of tetrabutylammonium fluoride (67 mL, 66.7 mmol, 1M in THF) was added slowly. The reaction mixture was warmed to RT and stirred for 16 h. The mixture was diluted with EtOAc and washed with brine. The organic layer was dried (Na₂SO₄), filtered and concentrated *in vacuo*. The residue was purified by SiO₂ chromatography eluting with a EtOAc/hexane gradient (15% to 35% EtOAc) to afford 5 g (91.7%) of **112**.

step 2 - To a solution of **112** (1 g, 4 mmol) in THF (6 mL) cooled to 0°C was added pyridine (0.9 g, 12 mmol) and SOCl₂ (1.4 g, 12 mmol) followed by DMAP (20 mg). The mixture was warmed to 50°C for 2 h with stirring. The reaction mixture was cooled to RT, diluted with EtOAc, washed with brine, dried (Na₂SO₄), filtered and concentrated *in vacuo*. The residue was purified by SiO₂ chromatography eluting with a EtOAc/hexane gradient (10% to 30% EtOAc) to afford 0.4 g (43.4%) of **114**.


step 3 - A suspension of **114** (0.150 g, 0.65 mmol) in MeOH (50 mL) and 10% Pd/C (20 mg) was stirred under H₂ at RT and atmospheric pressure. After 4 h the Pd was filtered and the solvent removed *in vacuo* to afford 100 mg (66%) of **116a**.

step 4 - To a solution of **116a** (100 mg, 0.43 mmol) in MeOH (4 mL) was added a solution of LiOH monohydrate (65.1 mg, 1.55 mmol) and water (1 mL). The mixture was stirred at RT for 16 h. The reaction mixture was concentrated *in vacuo*. The residue was dissolved in water (3

mL), acidified to a pH of about 2 with 3N HCl and extracted with EtOAc. The organic layer was dried (Na₂SO₄), filtered and concentrated *in vacuo* to afford 93 mg (99%) of **116b**.

II-51 can be prepared analogously to the procedure described in example 25, except in step 4, *p*-N,N-dimethylamino-benzoyl chloride is replaced with **116b** and the coupling of the **116** to **104** is carried out with HATU as described in step 1 of example 21.

Example 27: 4-*tert*-Butyl-N-{3-[5-(3-ethyl-ureido)-1-methyl-6-oxo-1,6-dihydro-pyridin-3-yl]-2-hydroxymethyl-phenyl}-benzamide (II-56)



The urea (**118**) is prepared from **46** by the procedure in step 2 of example 16 except methyl amine was replaced by ethyl amine. The palladium-coupling of **120** and **118** was carried out as described in step 5 of example 25.

II-65 is made analogously except the Suzuki coupling is carried out with **120** (Ar = 4-(1-fluoro-cyclopropyl)-phenyl).

II-54 can be prepared analogously except **118** is replaced with 1-(5-bromo-2-oxo-1,2-dihydro-pyridin-3-yl)-3-((S)-2-hydroxy-propyl)-urea which can be prepared from **46** and (S)-1-amino-propan-2-ol by the procedure described in step 2 of example 16 and the Suzuki coupling is carried out with **120** (Ar = 4-cyclopropyl-phenyl).


II-57 was prepared analogously except **120** is replaced by N-[2-(*tert*-butyl-dimethyl-silanyl-oxymethyl)-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-4-cyclopropyl-benzamide (**120**, Ar = 4-cyclopropyl-phenyl).

II-61 was prepared analogously except **120** is replaced by N-[2-(*tert*-butyl-dimethyl-silanyloxy-methyl)-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-4-dimethylamino-benzamide (**120**, Ar = 4-dimethylamino-phenyl).

II-77 was prepared analogously except **120** is replaced by N-[2-(*tert*-butyl-dimethyl-silanyl-oxymethyl)-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-4-cyclopropyl-benzamide (**120**, Ar = cyclopropyl-phenyl). **II-78**, **II-79**, **II-80**, **II-83**, **II-84**, **I-85** and **II-86** were prepared analogously except the requisite ureas were prepared by substituting 2-pyridin-2-yl-ethylamine (CASRN 2706-56-1), 2-pyridin-3-yl-ethylamine (CASRN 20173-24-4), N¹,N¹-dimethyl-


propane-1,3-diamine (CASRN 109-55-7), 3-morpholin-4-yl-propylamine (CASRN 123-00-2), C-(tetrahydro-furan-2-yl)-methylamine (CASRN 4795-29-3) and 1-amino-butan-2-ol (CASRN 13552-21-1) and 1-amino-propan-2-ol respectively for methylamine in the procedure described in step 2 of example 16.

5 **Example 28:** *N*-{3-[5-(5-Diethylaminomethyl-pyridin-2-ylamino)-1-methyl-6-oxo-1,6-dihydro-pyridin-3-yl]-2-hydroxymethyl-phenyl}-4-(1-fluoro-cyclopropyl)-benzamide (II-70)



Step 1 was carried out by converting **122a** to the corresponding acid chloride which was subsequently contacted with diethyl amine to afford **122b**. Treating 3-amino-5-bromo-1-methyl-1H-pyridin-2-one with sodium hydride in DMF as described in step 1 of example 18 and subsequently adding **122b** afforded **124** which was reduced to **126** by diborane-THF. The palladium-coupling of **126** and **120** (Ar = 1-fluoro-cyclopropyl-phenyl) was carried out as described in step 5 of example 25.

15 **Example 29:** 3,3-Dimethyl-2,3-dihydro-benzofuran-6-carboxylic acid (2-hydroxymethyl-3-{1-methyl-5-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-6-oxo-1,6-dihydro-pyridin-3-yl}-phenyl)-amide (II-71)




step 1 - To a stirred suspension of the **128** (5.01 g, 21.7 mmol) and K₂CO₃ (4.5 g, 32.5 mmol) in DMF (15 mL) was added dropwise 3-bromo-2-methylpropene (2.93 g, 21.7 mmol) and the resulting suspension was maintained at RT with vigorous stirring overnight. The mixture was diluted with 20% EtOAc/hexanes (100 mL), filtered, and concentrated *in vacuo* to a clear oil. The residue was purified with SiO₂ chromatography eluting with a EtOAc/hexane gradient (0 to 8% EtOAc) to afford 5.8 g (94%) of **130** as a clear oil.

step 2 - Solid AIBN (150 mg) was added in one portion to a solution of **130** (2.32 g, 8.14 mmol) and tributyltin hydride (2.84 g, 9.76 mmol) in degassed benzene (75 mL), and the solution was heated at 100°C under a N₂ atmosphere for 32 h. The benzene was removed *in vacuo*, and the residue was dissolved in Et₂O (100 mL). Aqueous KF (10% w/v, 150 mL) was added and the biphasic mixture was vigorously stirred for 3.5 h. The layers were separated and the organic layer was washed sequentially with saturated NaHCO₃ (25 mL) and brine (25 mL), dried (MgSO₄), filtered and concentrated to a yellow oil. The residue was purified with SiO₂ chromatography eluting with a EtOAc/hexane gradient (0 to 8% EtOAc) to afford 1.65 g (99%) of **132a** as a clear oil.

- 10 step 3 - Aqueous LiOH (1 N, 21 mL) was added to a solution of **132a** (1.45 g, 7.03 mmol) in MeOH (42 mL), and the solution was maintained at RT for 4 h. The reaction pH was adjusted to 2.0 by the addition of 2 N HCl, and this suspension was extracted with EtOAc (2 x 50 mL). The combined organics were washed with brine (25 mL), dried (MgSO₄), filtered and concentrated *in vacuo* to afford 1.34 g (99%) of **132b** as a white amorphous solid which was used without
15 further purification.

The preparation of **II-71** is then carried out in analogy to the procedures described in example 25, except in step 4, *p*-N,N-dimethylamino-benzoyl chloride is replaced with **132b** and the coupling to **104** was accomplished with HATU as described in step 1 of example 21.

Example 30: 4-(1-Difluoromethyl-cyclopropyl)-N-(2-hydroxymethyl-3-{1-methyl-5-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-6-oxo-1,6-dihydro-pyridin-3-yl}-phenyl)-benzamide (II-72)



- step 1 - To a solution of **134a** (0.8 g, 3.6 mmol, 124276-67-1) in toluene (8 mL) cooled to -50°C was added dropwise DIBAH (4.4 mL, 4.4 mmol, 1 M solution in toluene) while maintaining
25 temperature below -40°C. After stirring for 1 h at -40°C the reaction was quenched with 6 M HCl (3 mL) and stirred at RT for 30 min. The mixture was diluted with EtOAc, the organic layer was separated and washed sequentially with saturated NaHCO₃ and brine, dried (Na₂SO₄),


filtered and evaporated to afford 0.67 g (83%) of **134b** which was used in the next step without further purification.

step 2 – To a solution of **134b** (0.56 g, 2.5 mmol) and DCM (6 mL) under a nitrogen atmosphere was added dropwise a solution of DAST (0.40 g, 2.5mmol) in DCM (1.5 mL). The reaction mixture was stirred at RT for 2 h, then cooled in an ice bath and carefully quenched with ice water (4 mL). The resulting mixture was diluted with EtOAc and the organic phase washed sequentially with saturated NaHCO₃ and brine, dried (Na₂SO₄), filtered and concentrated *in vacuo*. The residue was purified by SiO₂ chromatography eluting with a EtOAc/hexane gradient (0 to 10% EtOAc) to afford 0.208 g (33% yield) of **134c**.

step 3 - To a solution of *n*-BuLi and pentane (0.5 mL, 1 mmol, 2M solution) was added THF (8 ml) and the solution was cooled to -78°C. A solution of compound **134c** (0.208 g, 0.84 mmol) in THF (1 mL) was added. The mixture was stirred at -78°C for 20 min and then CO₂ gas was bubbled through the solution for 0.5 h. The reaction mixture was warmed to RT, quenched with saturated NH₄Cl (4 mL) and extracted with ether. The combined organic extracts were washed with brine, dried (Na₂SO₄), filtered and concentrated *in vacuo*. The residue was dissolved in saturated NaHCO₃ (10 mL) and extracted with 10% EtOAc in hexane. The aqueous layer was acidified to a pH of about 2 with 3N HCl and the acidified solution extracted with EtOAc. The organic layer was washed with brine, dried (Na₂SO₄), filtered and concentrated *in vacuo* to afford 0.040 g (22%) of **136**.

The preparation of **II-72** is then carried out in analogy to the procedures described in example 25, except in step 4, *p*-N,N-dimethylamino-benzoyl chloride is replaced with **136** and the coupling to **104** was accomplished with HATU as described in step 1 of example 21.

Example 31: N-(2-Hydroxymethyl-3-{1-methyl-5-[5-(morpholine-4-carbonyl)-pyridin-2-ylamino]-6-oxo-1,6-dihydro-pyridin-3-yl}-phenyl)-4-(2,2,2-trifluoro-1,1-dimethyl-ethyl)-benzamide (II-73)



step 1 - To a solution of 1.0 M 4-methoxyphenyl magnesium bromide in THF (50 mL) at 0°C with stirring was added dropwise a solution of 1,1,1-trifluoroacetone (6.11 mL, 1.3 eq) in THF (50 mL). After the addition was completed, the reaction mixture was removed from the ice bath and stirred at RT overnight. The reaction was complete by TLC, and the solution was

5 partitioned between EtOAc (300 mL) and 1M HCl (100 mL) and the mixture was stirred for 10 min. Then the layers were separated and the organic layer was washed with brine (1 x 100 mL). Finally the EtOAc layer was dried (MgSO₄), filtered and concentrated. The crude product was purified by SiO₂ chromatography eluting with 5% EtOAc/hexane to afford 2.181 g of **140a** as a clear colorless oil.

10 step 2 - To a solution of **140a** (2.181g, 9.90 mmol) in DCM (33 mL) cooled to -78°C was added TiCl₄ (2.17 mL, 2 eq) *via* syringe and the resulting mixture was stirred at -78°C for 1.5 h. A solution of dimethylzinc in heptane (39.6 mL, 4 eq, 1.0 M) was added slowly *via* syringe while maintaining the temperature at -78°C and the resulting mixture was warmed to RT over about 3.5 h. Finally the reaction mixture was poured into crushed ice (500 g) and stirred vigorously for

15 about 20 min. DCM (300 mL) then was added and the mixture was partitioned and the layers separated. The aqueous layer was extracted with DCM (2 x 200 mL) and the combined DCM extracts were washed with brine (2 x 200 mL), dried (MgSO₄), filtered and concentrated. The crude product was purified by SiO₂ chromatography eluting with a EtOAc/hexane gradient (1% to 5% EtOAc) to afford 1.002 g of **140b** as a white powder.

20 step 3 - To a solution of **140b** (735mg, 3.37 mmol) and DCM (50 mL) cooled to -78°C was added BBr₃ (0.955 mL, 3 eq) was added with stirring at -78°C and the resulting mixture was stirred at -78°C for 30 min before the dry ice bath was removed and the solution stirred at RT for 3.5 h. TLC analysis indicated that the starting material was entirely consumed. Water (15 mL) was added carefully to the mixture while stirring vigorously followed by 2N NaOH (10 mL) with

25 stirring and then the solution was acidified to pH of about 2 with 4N HCl. The layers were separated and the aqueous layer was further extracted with DCM (2 x 50 mL) and the combined DCM extracts were washed with brine (1 x 50mL), dried (MgSO₄), filtered and concentrated to afford 0.711 g of **142a** as a brownish powder.

step 4 - To a solution of **142a** (702 mg, 3.44 mmol) in pyridine (3.5 mL) cooled to 0°C was

30 added dropwise with stirring triflic anhydride (0.64 mL, 1.1 eq) *via* syringe and the resulting mixture warmed to RT overnight with continued stirring. EtOAc (100 mL) was added and the mixture was washed sequentially with 1M Cu(SO₄)₂ (4 x 20 mL), water (2 x 20 mL) and brine (1 x 25 mL) and the organic layer was dried (MgSO₄), filtered and concentrated. The crude


reaction mixture was purified by SiO₂ chromatography eluting with a EtOAc/hexane gradient (1% to 3% EtOAc) to afford 0.927 g of **142b** as a colorless oil.

step 5 - A 5 mL microwave vessel was charged with **142b** (410 mg, 1.22 mmol), dppf (68 mg, 0.1eq), Pd(II)(OAc)₂ (28 mg, 0.1 eq), pyridine (0.45 mL), molybdenum hexacarbonyl (161 mg,

5 0.5 eq) and water (3.9 mL). The sample was irradiated for 20 min at 150°C then the mixture was partitioned between EtOAc (120 mL) and 4 N HCl. The EtOAc layer was concentrated, 2N NaOH (75 mL) and ether (75 mL) were added, the layers portioned and separated. Finally EtOAc (75 mL) was added and the stirred mixture at 0°C was acidified to pH of about 2 with concentrated HCl. The EtOAc layer was washed with brine (2 x 50 mL), dried (MgSO₄),
10 filtered and concentrated to afford 0.169 g of **142c** as a light tan powder.

The preparation of **II-73** is then carried out in analogy to the procedures described in example 25, except in step 4, *p*-N,N-dimethylamino-benzoyl chloride is replaced with **142c** and the coupling to **104** was accomplished with HATU as described in step 1 of example 21.

Example 32: 4-Cyclopropyl-N-{2-hydroxymethyl-3- [1-methyl-5-(4-methyl-thiazol-2-ylamino)-6-oxo-1,6-dihydro-pyridin-3-yl]-phenyl}-benzamide (II-92)




A sealable vial was charged with **50** (0.05 g, 0.19 mmol), 4-methyl-thiazol-2-ylamine (0.03 g, 1.2 eq), Pd(OAc)₂ (0.004 g, 0.1 eq), Xantphos (0.02 g, 0.2 eq), and K₂CO₃ (0.05 g, 2 eq) and 1,4-dioxane (0.5 mL). The argon purged mixture was sealed and heated to 95°C for 5 h. The cooled
20 reaction was filtered through CELITE, washed with DCM and purified on a preparative SiO₂ TLC plate developed with 3% MeOH/DCM to afford 55 mg of **139** as a dark green solid.

The condensation of **139** and **120** (Ar = 4-cyclopropyl-phenyl) to afford **II-92** was carried out as described in step 5 of example 25.

Example 33: 4-Cyclopropyl-N-(2-hydroxymethyl-3-{1-methyl-5-[6-(4-methyl-piperazin-1-yl)-pyridazin-3-ylamino]-6-oxo-1,6-dihydro-pyridin-3-yl}-phenyl)-benzamide (II-104)

25



step 1 - A mixture of **154** (1.5 g, 15 mmol) and **156** (2.23 g, 15 mmol) in dry toluene (10 mL) was heated in a sealed tube to 110°C for 3 h. The precipitous mixture was cooled to RT dissolved in methanol and dry loaded onto granular SiO₂. The crude reaction was purified by SiO₂ chromatography eluting with 7% MeOH/DCM to afford 1.3 g of **158** as an off-white solid.

step 2 - A heterogenous mixture of **46** (0.1 g, 0.49 mmol), **158** (0.16 g, 1.5 eq), Pd₂(dba)₃ (32 mg, 0.07 eq), xantphos (31 mg, 0.1 eq), and Cs₂CO₃ (0.32 g, 2 eq) in dioxane (1.5 mL) was purged with argon. The mixture was sealed and heated for 18 h in a sand bath at 95°C. The cooled mixture was partitioned between EtOAc and H₂O, concentrated, and purified on a preparative SiO₂ plate developed with 6% MeOH/DCM to afford 39 mg of **160** as a dark green solid. Condensation of **160** with **D-2** (Ar = 4-cyclopropyl-phenyl) was carried out by the procedure in set 5 of example 25.

Example 34: Bruton's tyrosine kinase (Btk) inhibition Assay

The assay is a capture of radioactive ³³P phosphorylated product through filtration. The interactions of Btk, biotinylated SH₂ peptide substrate (Src homology), and ATP lead to phosphorylation of the peptide substrate. Biotinylated product is bound streptavidin sepharose beads. All bound, radiolabeled products are detected by scintillation counter.

Plates assayed are 96-well polypropylene (Greiner) and 96-well 1.2 μm hydrophilic PVDF filter plates (Millipore). Concentrations reported here are final assay concentrations: 10- 100 μM compounds in DMSO (Burdick and Jackson), 5-10 nM Btk enzyme (His-tagged, full-length), 30 μM peptide substrate (Biotin-Aca-AAAEIYGEI-NH₂), 100 μM ATP (Sigma), 8 mM imidazole (Sigma, pH 7.2), 8 mM glycerol-2-phosphate (Sigma), 200 μM EGTA (Roche Diagnostics), 1 mM MnCl₂ (Sigma), 20 mM MgCl₂ (Sigma), 0.1 mg/ ml BSA (Sigma), 2 mM DTT (Sigma), 1 μCi ³³P ATP (Amersham), 20% streptavidin sepharose beads (Amersham), 50 mM EDTA

(Gibco), 2 M NaCl (Gibco), 2 M NaCl w/ 1% phosphoric acid (Gibco), microscint-20 (Perkin Elmer).

IC₅₀ determinations are calculated from 10 data points per compound utilizing data produced from a standard 96-well plate assay template. One control compound and seven unknown inhibitors were tested on each plate and each plate was run twice. Typically, compounds were diluted in half-log starting at 100 µM and ending at 3 nM. The control compound was staurosporine. Background was counted in the absence of peptide substrate. Total activity was determined in the presence of peptide substrate. The following protocol was used to determine Btk inhibition.

- 1) Sample preparation: The test compounds were diluted at half-log increments in assay buffer (imidazole, glycerol-2-phosphate, EGTA, MnCl₂, MgCl₂, BSA).
 - 2) Bead preparation
 - a.) rinse beads by centrifuging at 500 g
 - b.) reconstitute the beads with PBS and EDTA to produce a 20% bead slurry
 - 3) Pre-incubate reaction mix without substrate (assay buffer, DTT, ATP, ³³P ATP) and mix with substrate (assay buffer, DTT, ATP, ³³P ATP, peptide substrate) 30°C for 15 min.
 - 4) To start assay, pre-incubate 10 µL Btk in enzyme buffer (imidazole, glycerol-2-phosphate, BSA) and 10 µL of test compounds for 10 min at RT.
 - 5) Add 30 µL reaction mixture without or with substrate to Btk and compounds.
 - 6) Incubate 50 µL total assay mix for 30 min at 30°C.
 - 7) Transfer 40 µL of assay to 150 µL bead slurry in filter plate to stop reaction.
 - 8) Wash filter plate after 30 min, with following steps
 - a. 3 x 250 µL NaCl
 - b. 3 x 250 µL NaCl containing 1% phosphoric acid
 - c. 1 x 250 µL H₂O
 - 9) Dry plate for 1 h at 65°C or overnight at RT
 - 10) Add 50 µL microscint-20 and count ³³P cpm on scintillation counter.
- Calculate percent activity from raw data in cpm

$$\text{percent activity} = (\text{sample} - \text{bkg}) / (\text{total activity} - \text{bkg}) \times 100$$

- Calculate IC₅₀ from percent activity, using one-site dose response sigmoidal model

$$y = A + ((B - A) / (1 + ((x / C)^D)))$$

$$x = \text{cmpd conc}, y = \% \text{ activity}, A = \text{min}, B = \text{max}, C = \text{IC}_{50}, D = 1 \text{ (hill slope)}$$

Example 35: Inhibition of B-cell Activation - B cell FLIPR assay in Ramos cells

Inhibition of B-cell activation by compounds of the present invention is demonstrated by determining the effect of the test compounds on anti-IgM stimulated B cell responses.

The B cell FLIPR assay is a cell based functional method of determining the effect of potential inhibitors of the intracellular calcium increase from stimulation by an anti-IgM antibody. Ramos cells (human Burkitt's lymphoma cell line. ATCC-No. CRL-1596) were cultivated in Growth Media (described below). One day prior to assay, Ramos cells were resuspended in fresh growth media (same as above) and set at a concentration of $0.5 \times 10^6/\text{mL}$ in tissue culture flasks. On day of assay, cells are counted and set at a concentration of $1 \times 10^6/\text{mL}$ in growth media supplemented with $1\mu\text{M}$ FLUO-3AM (TefLabs Cat-No. 0116, prepared in anhydrous DMSO and 10% Pluronic acid) in a tissue culture flask, and incubated at 37°C ($4\% \text{CO}_2$) for one h. To remove extracellular dye, cells were collected by centrifugation (5min, 1000 rpm), resuspended in FLIPR buffer (described below) at 1×10^6 cells/mL and then dispensed into 96-well poly-D-lysine coated black/clear plates (BD Cat-No. 356692) at 1×10^5 cells per well. Test compounds were added at various concentrations ranging from $100\mu\text{M}$ to $0.03\mu\text{M}$ (7 concentrations, details below), and allowed to incubate with cells for 30 min at RT. Ramos cell Ca^{2+} signaling was stimulated by the addition of $10\mu\text{g/mL}$ anti-IgM (Southern Biotech, Cat-No. 2020-01) and measured on a FLIPR (Molecular Devices, captures images of 96 well plates using a CCD camera with an argon laser at 480nm excitation).

Media/Buffers:

Growth Medium: RPMI 1640 medium with L-glutamine (Invitrogen, Cat-No. 61870-010), 10% Fetal Bovine Serum (FBS, Summit Biotechnology Cat-No. FP-100-05); 1mM Sodium Pyruvate (Invitrogen Cat. No. 11360-070).

FLIPR buffer: HBSS (Invitrogen, Cat-No. 141175-079), 2mM CaCl_2 (Sigma Cat-No. C-4901), HEPES (Invitrogen, Cat-No. 15630-080), 2.5mM Probenecid (Sigma, Cat-No. P-8761), 0.1%

BSA (Sigma, Cat-No.A-7906), 11mM Glucose (Sigma, Cat-No.G-7528)

Compound dilution details:

In order to achieve the highest final assay concentration of $100\mu\text{M}$, $24\mu\text{L}$ of 10 mM compound stock solution (made in DMSO) is added directly to $576\mu\text{L}$ of FLIPR buffer. The test compounds are diluted in FLIPR Buffer (using Biomek 2000 robotic pipettor) resulting in the following dilution scheme: vehicle, 1.00×10^{-4} M, 1.00×10^{-5} , 3.16×10^{-6} , 1.00×10^{-6} , 3.16×10^{-7} , 1.00×10^{-7} , 3.16×10^{-8} .

Assay and Analysis:

Intracellular increases in calcium were reported using a max – min statistic (subtracting the resting baseline from the peak caused by addition of the stimulatory antibody using a Molecular

Devices FLIPR control and statistic exporting software. The IC_{50} was determined using a non-linear curve fit (GraphPad Prism software). Representative results are in Table III.

TABLE III

Compound	Ramos B Cell FLIPR [IC_{50} (μ M)]
I-2	0.0747
I-8	0.205
I-32	0.032
I-17	0.415
II-13	0.552
II-48	0.222
II-2	0.067
II-56	0.004
II-58	0.015

Example 36: Rat Collagen-induced arthritis (rCIA)

- 5 On day 0, rats are injected with an emulsion of Bovine Type II Collagen in Incomplete Freund's adjuvant (IFA), injected intradermally (i.d.) on several locations on the back. A booster injection of collagen emulsion is given around day 7, (i.d.) at the base of the tail or alternative sites on the back. Arthritis is generally observed 12-14 days after the initial collagen injection. Animals may be evaluated for the development of arthritis as described below (Evaluation of arthritis) from
- 10 day 14 onwards. Animals are dosed with candidate therapeutic agents in a preventive fashion starting at the time of secondary challenge and for the prescribed time (typically 2—3 weeks) and dosing frequency, daily (QD) or twice-daily (BID).

In both models, developing inflammation of the paws and limb joints is quantified using a scoring system that involves the assessment of the 4 paws following the criteria described below:

- 15 Scoring: 1= swelling and/or redness of paw or one digit.
 2= swelling in two or more joints.
 3= gross swelling of the paw with more than two joints involved.
 4= severe arthritis of the entire paw and digits.

- Evaluations are made on day 0 for baseline measurement and starting again at the first signs or
- 20 swelling for up to three times per week until the end of the experiment. The arthritic index for each mouse is obtained by adding the four scores of the individual paws, giving a maximum score of 16 per animal.

Example 37: Pharmaceutical compositions**Composition for Oral Administration (A)**

Ingredient	% wt./wt.
Active ingredient	20.0%
Lactose	79.5%
Magnesium stearate	0.5%

The ingredients are mixed and dispensed into capsules containing about 100 mg each; one capsule would approximate a total daily dosage.

5 Composition for Oral Administration (B)

Ingredient	% wt./wt.
Active ingredient	20.0%
Magnesium stearate	0.5%
Crosscarmellose sodium	2.0%
Lactose	76.5%
PVP (polyvinylpyrrolidone)	1.0%

The ingredients are combined and granulated using a solvent such as methanol. The formulation is then dried and formed into tablets (containing about 20 mg of active compound) with an appropriate tablet machine.

10 Composition for Oral Administration (C)

Ingredient	% wt./wt.
Active compound	1.0 g
Fumaric acid	0.5 g
Sodium chloride	2.0 g
Methyl paraben	0.15 g
Propyl paraben	0.05 g
Granulated sugar	25.5 g
Sorbitol (70% solution)	12.85 g
Veegum K (Vanderbilt Co.)	1.0 g
Flavoring	0.035 ml
Colorings	0.5 mg
Distilled water	q.s. to 100 ml

The ingredients are mixed to form a suspension for oral administration.

Parenteral Formulation (D)

Ingredient	% wt./wt.
Active ingredient	0.25 g
Sodium Chloride	qs to make isotonic
Water for injection to	100 ml

- The active ingredient is dissolved in a portion of the water for injection. A sufficient quantity of sodium chloride is then added with stirring to make the solution isotonic. The solution is made up to weight with the remainder of the water for injection, filtered through a 0.2 micron membrane filter and packaged under sterile conditions.

Suppository Formulation (E)

Ingredient	% wt./wt.
Active ingredient	1.0%
Polyethylene glycol 1000	74.5%
Polyethylene glycol 4000	24.5%

- The ingredients are melted together and mixed on a steam bath, and poured into molds containing 2.5 g total weight.

Topical Formulation (F)

Ingredients	grams
Active compound	0.2-2
Span 60	2
Tween 60	2
Mineral oil	5
Petrolatum	10
Methyl paraben	0.15
Propyl paraben	0.05
BHA (butylated hydroxy anisole)	0.01
Water	q.s. 100

The features disclosed in the foregoing description, or the following claims expressed in their specific forms or in terms of a means for performing the disclosed function, or a method or

process for attaining the disclosed result, as appropriate, may, separately, or in any combination of such features, be utilized for realizing the invention in diverse forms thereof.

The foregoing invention has been described in some detail by way of illustration and example, for purposes of clarity and understanding. It will be obvious to one of skill in the art that


5 changes and modifications may be practiced within the scope of the appended claims. Therefore, it is to be understood that the above description is intended to be illustrative and not restrictive.

The scope of the invention should, therefore, be determined not with reference to the above description, but should instead be determined with reference to the following appended claims, along with the full scope of equivalents to which such claims are entitled.

10 All patents, patent applications and publications cited in this application are hereby incorporated by reference in their entirety for all purposes to the same extent as if each individual patent, patent application or publication were so individually denoted.

Claims

1. A compound of formula I wherein:



(I)

wherein

5 X^1 is N;

R^1 is $C(=O)NHR^6$, phenyl or heteroaryl wherein said heteroaryl is selected from the group consisting of pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl; pyrrolyl, furanyl, thiophenyl, imidazolyl, pyrazolyl, oxazolyl, thiazolyl, isoxazolyl, isothiazolyl, 2- C_{1-6} alkyl-pyridazin-3-on-6-yl, 2-benzyl-pyridazin-3-on-6-yl, 1,4,5,6-tetrahydro-pyrimidin-2-yl and 4,5-dihydro-1H-imidazol-2-yl; said phenyl and said heteroaryl optionally independently substituted with one to three groups selected from the group consisting of halogen, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} heteroalkyl, C_{1-6} heteroalkoxy, C_{1-6} alkylsulfanyl, C_{1-6} alkylsulfonyl, C_{1-6} heteroalkylsulfonyl and heterocyclyl- C_{1-6} alkoxy wherein said heterocyclyl is azetindinyl, pyrrolidinyl or piperidinyl, $CONR^aR^b$, CO_2R^g , $SO_2NR^aR^b$, NR^aR^b , $NHSO_2R^7$ or $NHCOR^7$;

15 wherein

R^6 is hydrogen, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{1-6} heteroalkyl, C_{3-6} cycloalkyl, pyridinyl- C_{1-3} alkyl, phenyl or phenyl- C_{1-3} alkyl said phenyl optionally substituted with C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} heteroalkoxy, halogen, $CONR^aR^b$ or CO_2R^g ;

R^a and R^b (i) selected independently are hydrogen, C_{1-6} alkyl, C_{1-6} heteroalkyl, C_{1-3} alkoxy- C_{1-3} alkyl, carboxy- C_{1-3} alkyl, C_{3-6} cycloalkyl or heterocyclyl; (ii) together are $(CH_2)_mX^2(CH_2)_2$ wherein m is 2 or 3 and X^2 is O, $S(O)_n$ or NR^8 and n is zero to two, and R^8 is hydrogen, C_{1-6} alkyl, C_{1-6} hydroxyalkyl or C_{1-3} acyl; or (iii) together with the nitrogen to which they are attached are piperidinyl or pyrrolidinyl said piperidinyl or said pyrrolidinyl optionally substituted with hydroxy, C_{1-3} alkoxy, C_{1-3} hydroxyalkyl;

25 R^g is hydrogen or C_{1-6} alkyl;

R^7 is C_{1-6} alkyl or C_{1-6} heteroalkyl;

R^2 is hydrogen or C_{1-6} alkyl;




R^3 and R^4 are independently hydrogen, halogen, amino, C_{1-6} hydroxyalkyl, C_{1-6} alkyl or cyclopropyl;

R^5 is (i) phenyl,

(ii) heteroaryl selected from the group consisting of pyridinyl, benzo[b]thiophen-2-yl, 4,5,6,7-tetrahydro-benzo[b]thiophen-2-yl, thiophenyl, 1',2',3',4',5',6'-hexahydro-[2,4']bipyridin-5-yl and C_{1-3} alkyl-indolyl optionally substituted with one or two C_{1-6} alkyl or halogen;

(iii) azetidin-1-yl, pyrrolidin-2-yl, piperidin-1-yl, azepan-1-yl or 2,3-dihydro-1H-isoindolin-2-yl, or 6,7-dihydro-5H-pyrrolo[3,4-b]pyridine;

(iv) NR^cR^d wherein R^c and R^d together are $(CH_2)_oX^2(CH_2)_p$ wherein o and p are independently 1 or 2, and X^2 is as defined above, or R^c and R^d independently are hydrogen C_{1-10} alkyl or C_{1-10} hydroxyalkyl;



wherein R^9 is hydrogen, halogen or C_{1-6} alkyl;

wherein said phenyl, heteroaryl, azetidin-1-yl, pyrrolidin-2-yl, piperidin-1-yl, azepan-1-yl or 2,3-dihydro-1H-isoindolin-2-yl ring is optionally substituted with one to three groups independently selected from the group consisting of (a) halogen, (b) C_{1-6} alkyl, (b) C_{2-6} alkenyl, (c) C_{2-6} alkynyl, (d) C_{1-6} haloalkyl, (e) C_{1-6} heteroalkyl, (f) C_{3-7} cycloalkyl optionally

substituted with one to three groups selected from C_{1-6} alkyl, C_{1-6} haloalkyl or halogen, (g) C_{1-6} alkoxyalkyl, (h) hydroxy, (i) NR^eR^f , wherein R^e and R^f are (i) independently hydrogen, C_{1-6} alkyl or (ii) together are $(CH_2)_mX^3(CH_2)_2$ wherein m is 2 or 3 and X^3 is CH_2 , O, $S(O)_n$ or NR^8 and n is zero to two and R^8 is as defined above; (j) C_{1-6} alkoxy, (k) trialkylsilyl, (l) C_{1-6} cyanoalkyl and (m) SF_5 ;

A is $-NHC(=O)-$, $C(=O)NH-$, $NHC(=O)NH$, $CH_2C(=O)$, CH_2SO_2 or if R^5 is (v) or (vi) A is absent;

or a pharmaceutically acceptable salt thereof.

2. A compound according to claim 1 wherein R^1 is phenyl, pyridin-2-yl or pyrimidin-4-yl, R^2 is C_{1-3} alkyl; A is $NHC(=O)$ and R^3 is hydrogen or C_{1-3} alkyl; or hydroxymethyl.
3. A compound according to claim 1 wherein R^1 is $C(=O)NHR^6$, R^6 is C_{1-6} alkyl, R^3 is hydrogen,
5 C_{1-3} alkyl or C_{1-6} heteroalkyl; and, A is $NHC(=O)$.
4. A compound according claim 1 wherein R^5 is (v) or (vi).
5. A compound according claim 1 wherein A is $CH_2C(=O)$ and R^5 is (i), (ii), (iii) or (iv).
6. A pharmaceutical composition comprising a compound according to claim 1 admixed with at least one pharmaceutically acceptable carrier, excipient or diluent.
- 10 7. The use of a compound of formula I according to claim 1 for the preparation of a medicament for the treatment of an inflammatory and/or autoimmune condition.
8. The use according to claim 7, wherein the inflammatory and/or auto-immune condition is rheumatoid arthritis.