

SALMONELLA-BASED VECTORS FOR CANCER IMMUNOTHERAPY TARGETING WILMS' TUMOR GENE WT1

FIELD OF THE INVENTION

[0001] The present invention relates to an attenuated mutant strain of Salmonella comprising a recombinant DNA molecule encoding Wilms' tumor Protein 1. In particular, the present invention relates to the use of said attenuated mutant strain of Salmonella in cancer immunotherapy.

BACKGROUND OF THE INVENTION

[0002] Wilms' tumor gene 1 (WT1) encodes a zinc finger transcription factor involved in cell proliferation and differentiation. It is highly expressed in a wide variety of malignancies including several types of hematological malignancies and various solid tumors. In contrast, normal tissue expression of WT1 in adults is restricted to gonads, uterus, kidney, mesothelium and CD34⁺ progenitor cells in various types of tissues. WT1 was originally proposed as a tumor suppressor gene. However, more recent evidence points to oncogenic functions of this transcription factor; Wt-1 negatively affects differentiation and promotes proliferation of progenitor cells. Furthermore, overexpressed WT1 is immunogenic; WT1 specific T cells as well as IgG anti-WT1 antibodies have been observed in cancer patients. Thus, WT-1 is a promising candidate for the development of cancer vaccines.

[0003] Human clinical trials with WT1 vaccines based on HLA (human leukocyte antigen)-restricted WT1 peptide fragments have been reported. Osada et al., Clin Cancer Res 2009;15:2789-2796, discloses a WT1-encoding Adenovirus Vaccine.

[0004] To the inventor's knowledge, no live bacterial cancer vaccine targeting WT1 has been reported. Furthermore, no oral cancer vaccine targeting WT1 has been described.

[0005] Attenuated derivatives of *Salmonella enterica* are attractive vehicles for the delivery of heterologous antigens to the mammalian immune system, since *S. enterica* strains can potentially be delivered via mucosal routes of immunization, i.e. orally or nasally, which offers advantages of simplicity and safety compared to parenteral administration. Furthermore, *Salmonella* strains elicit strong humoral and cellular immune responses at the level of both systemic and mucosal compartments. Batch preparation costs are relatively low and formulations of live bacterial vaccines are highly stable. Attenuation can be accomplished by deletion of various genes, including virulence, regulatory, and metabolic genes.

[0006] Several *Salmonella typhimurium* strains attenuated by *aro* mutations have been shown to be safe and effective delivery vehicles for heterologous antigens in animal models.

[0007] Approaches of delivering DNA constructs encoding antigens, in particular VEGF receptor proteins, via live attenuated *Salmonella typhimurium* strains into mouse target cells are described in WO 03/073995. Niethammer et al., (*Nature Medicine* 2002, 8(12), 1369) demonstrated that the attenuated *S. typhimurium aroA* strain SL7207 harboring an expression vector encoding the murine vascular endothelial growth factor receptor 2 (VEGFR-2 or FLK-1), which is essential for tumor angiogenesis, is functional as a cancer vaccine.

[0008] There is however only one attenuated *Salmonella enterica* serovar strain, namely *Salmonella enterica* serovar typhi Ty21a (short: *S. typhi* Ty21a), which has been accepted for use in humans and is distributed under the trade name of Vivotif® (Berna Biotech Ltd., a Crucell Company, Switzerland; marketing authorization number PL 15747/0001 dated 16 December 1996).

[0009] This well-tolerated, live oral vaccine against typhoid fever was derived by chemical mutagenesis of the wild-type virulent bacterial isolate *S. typhi* Ty2 and harbors a loss-of-function mutation in the *galE* gene, as well as other less defined mutations. It has been licensed as typhoid vaccine in many countries after it was shown to be efficacious and safe in field trials.

[0010] WT1 is a promising tumor antigen for the development of cancer vaccines. Major limitations of previously available WT1 peptide vaccines are HLA-restriction and parenteral administration. The great need for improved cancer therapy approaches based on targeting WT1 has not been met so far.

OBJECTS OF THE INVENTION

[0011] In view of the prior art, it is an object of the present invention to provide a novel oral WT1 targeting cancer vaccine. Such a WT1 targeting cancer vaccine would offer major advantages for improving the treatment options for cancer patients.

SUMMARY OF THE INVENTION

[0012] In one aspect, the present invention relates to an attenuated mutant strain of *Salmonella* comprising at least one copy of a recombinant DNA molecule comprising an expression cassette encoding Wilms' Tumor Protein (WT1).

[0013] The attenuated *Salmonella* strain of the present invention was shown to exhibit antitumor activity in a mouse model challenged with murine leukemia cells. To the inventor's knowledge, this novel attenuated *Salmonella* strain is the first live bacterial cancer vaccine targeting WT1. In addition, the attenuated *Salmonella* strain of the present invention is the first oral cancer vaccine targeting WT1. Since WT1 is overexpressed in a wide variety of hematological malignancies and solid tumor, the attenuated *Salmonella* strain of the present invention has great potential as universal cancer vaccine.

[0014] In a first study, the vaccine according to the present invention (VXM06) has been demonstrated to efficiently prolong survival of mice bearing intraperitoneally implanted leukemia cells. These results indicate that vaccination with VXM06 may lead to an immune response and the development of an immune memory against tumor cells overexpressing WT1. It is remarkable and surprising that the novel vaccine VXM06 is effective at relatively low doses. The attenuated *Salmonella* mutant strain of the present invention may be applied in monotherapy or in combination with a second attenuated mutant strain of *Salmonella* comprising a DNA molecule encoding a second tumor antigen. Furthermore, the attenuated mutant strain of the present invention may be administered in combination with chemotherapy, radiotherapy or biological cancer therapy. Treatment with VXM06

may also be effective, if the patient has developed a resistance to chemotherapy (chemo-refractory patients). The novel attenuated *Salmonella* strain of the present invention might therefore be useful in novel, greatly improved cancer therapy approaches.

[0015] In particular embodiments, the attenuated mutant strain of *Salmonella* is of the species *Salmonella enterica*. In particular embodiments, the attenuated mutant strain of *Salmonella* is *Salmonella typhi* Ty21a.

[0016] In particular embodiments, the expression cassette is a eukaryotic expression cassette.

[0017] In particular embodiments, WT1 is selected from the group consisting of human WT1 and a protein that shares at least about 80% sequence identity therewith.

[0018] In particular embodiments, WT1 is truncated. In particular embodiments, the zinc finger domain of WT1 is deleted. In particular embodiments, the truncated WT1 has the amino acid sequence as found in SEQ ID NO 1.

[0019] In particular embodiments, the recombinant DNA molecule comprises the kanamycin antibiotic resistance gene, the pMB1 ori, and a eukaryotic expression cassette encoding human WT1 or a protein that shares at least 80% sequence identity therewith, particularly truncated human WT1, under the control of a CMV promoter. In particular embodiments, the recombinant DNA molecule is a plasmid designated pVAX10.hWT1 and has the nucleic acid sequence as found in SEQ ID NO 2.

[0020] In particular embodiments, the attenuated mutant strain of Salmonella is for use as a medicament.

[0021] In particular embodiments, the attenuated mutant strain of Salmonella is for use as a vaccine.

[0022] In particular embodiments, the attenuated mutant strain of Salmonella is for use in cancer immunotherapy.

[0023] In particular embodiments, cancer immunotherapy further comprises administration of one or more further attenuated mutant strain(s) of Salmonella comprising at least one copy of a recombinant DNA molecule comprising an expression cassette encoding a tumor antigen and/or a tumor stroma antigen. In particular embodiments, said one or more further attenuated mutant strain(s) of Salmonella is/are Salmonella typhi Ty21a comprising a eukaryotic expression cassette. In particular embodiments, said one or more further strain(s) of Salmonella comprise(s) an attenuated mutant strain(s) of Salmonella encoding human VEGFR-2.

[0024] In particular embodiments, the attenuated mutant strain of Salmonella is co-administered with said one or more further attenuated mutant strain(s) of Salmonella.

[0025] In particular embodiments, cancer immunotherapy is accompanied by chemotherapy, radiotherapy or biological cancer therapy.

[0026] In particular embodiments, the attenuated mutant strain of Salmonella is administered during the chemotherapy or the radiotherapy treatment cycle or during biological cancer therapy.

[0027] In particular embodiments, the attenuated mutant strain of *Salmonella* is administered before the chemotherapy or the radiotherapy treatment cycle or before biological cancer therapy.

[0028] In particular embodiments, the attenuated mutant strain of *Salmonella* is administered after the chemotherapy or the radiotherapy treatment cycle or after biological cancer therapy.

[0029] In further embodiments the attenuated mutant strain of *Salmonella* is administered before and during at least one of the chemotherapy, the radiotherapy treatment cycle and the biological cancer therapy. In cases where more than one of the chemotherapy, the radiotherapy and the biological cancer therapy are carried out the attenuated mutant strain of *Salmonella* may be administered before or during or before and during at least one of these therapies, particularly during at least two of these therapies.

[0030] In particular embodiments, the attenuated mutant strain of *Salmonella* is administered orally.

[0031] In particular embodiments, the cancer is selected from leukemia, particularly from acute myeloid leukemia (AML) and acute lymphoid leukemia (ALL), and from solid tumors, particularly from lung cancer, breast cancer, esophageal, colon, colorectal, gastric, cholangioductal, pancreatic cancer, glioblastoma, head and neck cancer, synovial sarcoma, angiosarcoma, osteosarcoma, thyroid cancer, cervical, endometrial, ovarian cancer, neuroblastoma, rhabdomyosarcoma, prostate cancer.

[0032] In particular embodiments, the single dose is from about 10^5 to about 10^{11} , particularly from about 10^6 to about 10^{10} , more particularly from about 10^6 to about

10⁹, more particularly from about 10⁶ to about 10⁸, most particularly from about 10⁶ to about 10⁷ colony forming units (CFU).

[0033] In particular embodiments, the attenuated mutant strain of Salmonella is for use in personalized cancer immunotherapy comprising the step of assessing the tumor antigen expression pattern and/or stroma antigen expression pattern of a patient.

[0033a] In one aspect, the present invention relates to an attenuated mutant strain of Salmonella comprising at least one copy of a recombinant DNA molecule comprising a eukaryotic expression cassette encoding Wilms' Tumor Protein (WT1).

[0033b] In one aspect, the present invention relates to the use of an attenuated mutant strain of Salmonella as described herein, in the manufacture of a vaccine.

DETAILED DESCRIPTION OF THE INVENTION

[0034] The present invention may be understood more readily by reference to the following detailed description of the invention and the examples included therein.

[0035] In one aspect, the present invention relates to an attenuated mutant strain of Salmonella comprising at least one copy of a recombinant DNA molecule comprising an expression cassette encoding Wilms' Tumor Protein (WT1).

[0036] According to the invention, the attenuated Salmonella strain functions as the bacterial carrier of the recombinant DNA molecule comprising an expression cassette encoding Wilms' Tumor Protein (WT1) for the delivery of said recombinant DNA molecule into a target cell.

[0037] In the context of the present invention, the term "attenuated" refers to a bacterial strain of reduced virulence compared to the parental bacterial strain, not harboring the attenuating mutation. Attenuated bacterial strains have preferably lost their virulence but retained their ability to induce protective immunity. Attenuation can be accomplished by deletion of various genes, including virulence, regulatory, and metabolic genes. Attenuated bacteria may be found naturally or they may be

produced artificially in the laboratory, for example by adaptation to a new medium or cell culture or they may be produced by recombinant DNA technology.

[0038] In the context of the present invention, the term "mutant strain" refers to a bacterial strain harboring a mutation in its genome. In this context, the term "mutation" refers to a change in a nucleic acid sequence, including point mutations, insertions, deletions, translocations and inversions.

[0039] In the context of the present invention, the term "comprises" or "comprising" means "including, but not limited to". The term is intended to be open-ended, to specify the presence of any stated features, elements, integers, steps or components, but not to preclude the presence or addition of one or more other features, elements, integers, steps, components or groups thereof. The term "comprising" thus includes the more restrictive terms "consisting of" and "essentially consisting of". In one embodiment the term "comprising" as used throughout the application and in particular within the claims may be replaced by the term "consisting of".

[0040] In the context of the present invention, the term "recombinant DNA molecule" refers to an engineered DNA construct, preferably composed of DNA pieces of different origin. The recombinant DNA molecule can be a linear nucleic acid, or preferably, a circular recombinant DNA plasmid generated by introducing an open reading frame encoding WT1 into an expression vector plasmid.

[0041] In the context of the present invention, the term "expression cassette" refers to a nucleic acid unit comprising at least the WT1 gene under the control of regulatory sequences controlling its expression. The expression cassette comprised in the attenuated mutant strain of *Salmonella* can preferably mediate transcription of the

included open reading frame encoding WT1 in a target cell. Expression cassettes typically comprise a promoter, at least one open reading frame and a transcription termination signal.

[0042] The zinc finger transcription factor Wilms' tumor protein 1 is encoded by the WT1 gene. It contains four zinc finger motifs at the C-terminus and a proline/glutamine-rich DNA-binding domain at the N-terminus. Multiple transcript variants, *resulting from alternative splicing at two coding exons, have been well characterized*. WT1 plays an essential role in the development of the urogenital system and is involved in cell proliferation and differentiation. The WT1 gene was isolated as the gene responsible for a childhood renal neoplasm, Wilms' tumor. It is highly expressed in a wide variety of malignancies including several types of hematological malignancies and various solid tumors. In contrast, normal tissue expression of WT1 in adults is restricted to gonads, uterus, kidney, mesothelium and progenitor cells in various types of tissues. Due to its expression profile, its oncogenic functions and its immunogenic potential, the tumor antigen WT1 is a promising candidate for the development of cancer vaccines.

[0043] In particular embodiments, the attenuated mutant strain of Salmonella is of the species *Salmonella enterica*. In particular embodiments, the attenuated mutant strain of Salmonella is *Salmonella typhi* Ty21a. The attenuated *S. typhi* Ty21a strain is the active component of Typhoral L[®], also known as Vivotif[®] (manufactured by Berna Biotech Ltd., a Crucell Company, Switzerland). It is currently the only licensed live oral vaccine against typhoid fever. This vaccine has been extensively tested and has proved to be safe regarding patient toxicity as well as transmission to third parties (Wahdan et al., J. Infectious Diseases 1982, 145:292-295). The vaccine is licensed in more than 40 countries. The Marketing Authorization number of Typhoral L[®] is PL 15747/0001 dated 16 December 1996. One dose of vaccine contains at least 2×10^9

viable *S. typhi* Ty21a colony forming units and at least 5×10^9 non-viable *S. typhi* Ty21a cells.

[0044] One of the biochemical properties of the *Salmonella typhi* Ty21a bacterial strain is its inability to metabolize galactose. The attenuated bacterial strain is also not able to reduce sulfate to sulfide which differentiates it from the wild-type *Salmonella typhi* Ty2 strain. With regard to its serological characteristics, the *Salmonella typhi* Ty21a strain contains the O9-antigen which is a polysaccharide of the outer membrane of the bacteria and lacks the O5-antigen which is in turn a characteristic component of *Salmonella typhimurium*. This serological characteristic supports the rationale for including the respective test in a panel of identity tests for batch release.

[0045] In particular embodiments, the expression cassette is a eukaryotic expression cassette. In the context of the present invention, the term "eukaryotic expression cassette" refers to an expression cassette which allows for expression of the open reading frame in a eukaryotic cell. It has been shown that the amount of heterologous antigen required to induce an adequate immune response may be toxic for the bacterium and result in cell death, over-attenuation or loss of expression of the heterologous antigen. Using a eukaryotic expression cassette that is not expressed in the bacterial vector but only in the target cell may overcome this toxicity problem and the protein expressed may exhibit a eukaryotic glycosylation pattern.

[0046] A eukaryotic expression cassette comprises regulatory sequences that are able to control the expression of an open reading frame in a eukaryotic cell, preferably a promoter and a polyadenylation signal. Promoters and polyadenylation signals included in the recombinant DNA molecules comprised by the attenuated mutant strain of *Salmonella* of the present invention are preferably selected to be

functional within the cells of the subject to be immunized. Examples of suitable promoters, especially for the production of a DNA vaccine for humans, include but are not limited to promoters from Cytomegalovirus (CMV), such as the strong CMV immediate early promoter, Simian Virus 40 (SV40), Mouse Mammary Tumor Virus (MMTV), Human Immunodeficiency Virus (HIV), such as the HIV Long Terminal Repeat (LTR) promoter, Moloney virus, Epstein Barr Virus (EBV), and from Rous Sarcoma Virus (RSV) as well as promoters from human genes such as human actin, human myosin, human hemoglobin, human muscle creatine, and human metallothionein. In a particular embodiment, the eukaryotic expression cassette contains the CMV promoter. In the context of the present invention, the term "CMV promoter" refers to the strong immediate-early cytomegalovirus promoter.

[0047] Examples of suitable polyadenylation signals, especially for the production of a DNA vaccine for humans, include but are not limited to the bovine growth hormone (BGH) polyadenylation site, SV40 polyadenylation signals and LTR polyadenylation signals. In a particular embodiment, the eukaryotic expression cassette included in the recombinant DNA molecule comprised by the attenuated mutant strain of *Salmonella* of the present invention comprises the BGH polyadenylation site.

[0048] In addition to the regulatory elements required for expression of the heterologous WT1 gene, like a promoter and a polyadenylation signal, other elements can also be included in the recombinant DNA molecule. Such additional elements include enhancers. The enhancer can be, for example, the enhancer of human actin, human myosin, human hemoglobin, human muscle creatine and viral enhancers such as those from CMV, RSV and EBV.

[0049] Regulatory sequences and codons are generally species dependent, so in order to maximize protein production, the regulatory sequences and codons are

preferably selected to be effective in the species to be immunized. The person skilled in the art can produce recombinant DNA molecules that are functional in a given subject species.

[0050] In particular embodiments, WT1 is selected from the group consisting of human WT1 and a protein that shares at least about 80% sequence identity therewith.

[0051] In this context, the term "about" or "approximately" means within 80% to 120%, alternatively within 90% to 110%, including within 95% to 105% of a given value or range.

[0052] In the context of the present invention, the term "protein that shares at least about 80% sequence identity with human WT1" refers to a protein that differs in the amino acid sequence and/or the nucleic acid sequence encoding the amino acid sequence of human WT1. The protein may be of natural origin, e.g. a homolog of WT1 of a different species, or an engineered protein, e.g. an engineered WT1 derivative. It is known that the usage of codons is different between species. Thus, when expressing a heterologous protein in a target cell, it may be necessary, or at least helpful, to adapt the nucleic acid sequence to the codon usage of the target cell. Methods for designing and constructing derivatives of a given protein are well known to anyone of ordinary skill in the art.

[0053] The protein that shares at least about 80% sequence identity with human WT1 may contain one or more mutations comprising an addition, a deletion and/or a substitution of one or more amino acids. According to the teaching of the present invention, said deleted, added and/or substituted amino acids may be consecutive amino acids or may be interspersed over the length of the amino acid sequence of

the protein that shares at least about 80% sequence identity with human WT1. According to the teaching of the present invention, any number of amino acids may be added, deleted, and/or substitutes, as long as the sequence identity with human WT1 is at least about 80%. In particular embodiments, the sequence identity with human WT1 is at least about 80%, at least about 85%, at least about 90%, or most particularly at least about 95%. Methods and algorithms for determining sequence identity including the comparison of a parental protein and its derivative having deletions, additions and/or substitutions relative to a parental sequence, are well known to the practitioner of ordinary skill in the art. On the DNA level, the nucleic acid sequences encoding the protein that shares at least about 80% sequence identity with human WT1 may differ to a larger extent due to the degeneracy of the genetic code.

[0054] In particular embodiments, WT1 is truncated. In particular embodiments, the zinc finger domain of WT1 is deleted. In particular embodiments, the truncated WT1 has the amino acid sequence as found in SEQ ID NO 1.

[0055] The zinc finger domain at the C-terminus of WT1 comprises four zinc finger motifs. Truncated WT1 of the amino acid sequence as found in SEQ ID NO 1 represents amino acids 1 to 371 of UniProt ref P19544-7. Deletion of the zinc finger domain minimizes the risk of immunological cross reactivity with other zinc finger containing transcription factors. Furthermore, truncated WT1 lacking the zinc finger domain has greater immunogenic potential than full-length WT1. In addition, deletion of the zinc finger motifs, which are essential for DNA binding, abrogates the oncogenic potential of WT1, thus minimizing the risk of oncogenesis.

[0056] In particular embodiments, the recombinant DNA molecule comprises the kanamycin antibiotic resistance gene, the pMB1 ori, and a eukaryotic expression

cassette encoding human WT1 or a protein that shares at least 80% sequence identity therewith, particularly truncated human WT1, under the control of a CMV promoter. In particular embodiments, the recombinant DNA molecule is a plasmid designated pVAX10.hWT1 and has the nucleic acid sequence as found in SEQ ID NO 2.

[0057] In particular embodiments, the recombinant DNA molecule is derived from commercially available pVAX1TM expression plasmid (Invitrogen, San Diego, California). This expression vector was modified by replacing the high copy pUC origin of replication by the low copy pMB1 origin of replication of pBR322. The low copy modification was made in order to reduce the metabolic burden and to render the construct more stable. The generated expression vector backbone was designated pVAX10. Inserting human, truncated WT1 into this expression vector backbone via *NheI/XhoI* yielded the expression plasmid pVAX10.hWT1 having the nucleic acid sequence as found in SEQ ID NO 2. The expression plasmid pVAX10.hWT1 is schematically depicted in Figure 3.

[0058] In particular embodiments, the attenuated mutant strain of *Salmonella* is for use as a medicament.

[0059] In particular embodiments, the attenuated mutant strain of *Salmonella* is for use as a vaccine.

[0060] In the context of the present invention, the term "vaccine" refers to an agent which is able to induce an immune response in a subject upon administration. A vaccine can preferably prevent, ameliorate or treat a disease. A vaccine in accordance with the present invention comprises an attenuated mutant strain of *Salmonella*, preferably *S. typhi* Ty21a. The vaccine in accordance with the present

invention further comprises at least one copy of a recombinant DNA molecule comprising an expression cassette, preferably a eukaryotic expression cassette, encoding WT1, preferably selected from human WT1 or a protein that shares at least about 80% sequence identity therewith. Preferably, said WT1 is truncated, particularly the zinc finger domain is deleted.

[0061] The live attenuated Salmonella mutant strain according to the present invention comprising a recombinant DNA molecule encoding WT1 can be used as a vehicle for the oral delivery of this recombinant DNA molecule. Such a delivery vector comprising a DNA molecule encoding a heterologous antigen, such as WT1, is termed DNA vaccine.

[0062] Genetic immunization might be advantageous over conventional vaccination. The target DNA can be detected for a considerable period of time thus acting as a depot of the antigen. Sequence motifs in some plasmids, like GpC islands, are immunostimulatory and can function as adjuvants furthered by the immunostimulation due to LPS and other bacterial components.

[0063] In contrast to peptide vaccines that can only mediate immunity against a small fragment of the WT1 protein, genetic vaccination may result in immunity against a wide variety of epitopes present over the whole length of the encoded WT1 protein.

[0064] Apart from that, WT1 peptide vaccine, which have been used in clinical trials for the most part, have limited application due to HLA restriction of the peptides, i.e. their binding capacity to HLA molecules of antigen presenting cells (APCs). In contrast, the DNA vaccine of the present invention is not HLA restricted.

[0065] Live bacterial vectors produce their own immunomodulatory factors such as lipopolysaccharides (LPS) *in situ* which may constitute an advantage over other forms of administration such as microencapsulation. Moreover, the use of the natural route of entry proves to be of benefit since many bacteria, like *Salmonella*, egress from the gut lumen via the M cells of Peyer's patches and migrate eventually into the lymph nodes and spleen, thus allowing targeting of vaccines to inductive sites of the immune system. The vaccine strain of *Salmonella typhi*, Ty21a, has been demonstrated to-date to have an excellent safety profile. Upon exit from the gut lumen via the M cells, the bacteria are taken up by phagocytic cells, such as macrophages and dendritic cells. These cells are activated by the pathogen and start to differentiate, and probably migrate into the lymph nodes and spleen. Due to their attenuating mutations, bacteria of the *S. typhi* Ty21 strain are not able to persist in these phagocytic cells but die at this time point. The recombinant DNA molecules are released and subsequently transferred into the cytosol of the phagocytic immune cells, either via a specific transport system or by endosomal leakage. Finally, the recombinant DNA molecules enter the nucleus, where they are transcribed, leading to WT1 expression in the cytosol of the phagocytic cells. Specific cytotoxic T cells against WT1 are induced by the activated antigen presenting cells.

[0066] There is no data available to-date indicating that *S. typhi* Ty21a is able to enter the bloodstream systemically. The live attenuated *Salmonella typhi* Ty21a vaccine strain thus allows specific targeting of the immune system while exhibiting an excellent safety profile.

[0067] Attenuated derivatives of *Salmonella enterica* are attractive as vehicles for the delivery of heterologous antigens to the mammalian immune system because *S. enterica* strains can potentially be delivered via mucosal routes of immunization, i.e. orally or nasally, which offers advantages of simplicity and safety compared to

parenteral administration. Furthermore, *Salmonella* strains elicit strong humoral and cellular immune responses at the level of both systemic and mucosal compartments.

[0068] In particular embodiments, the attenuated mutant strain of *Salmonella* is for use in cancer immunotherapy.

[0069] In particular embodiments, cancer immunotherapy further comprises administration of one or more further attenuated mutant strain(s) of *Salmonella* comprising at least one copy of a recombinant DNA molecule comprising an expression cassette encoding a tumor antigen and/or a tumor stroma antigen. In particular embodiments, said one or more further mutant strain(s) of *Salmonella* is/are *Salmonella typhi* Ty21a comprising a eukaryotic expression cassette. In particular embodiments, said one or more further strain(s) of *Salmonella* comprise(s) an attenuated mutant strain of *Salmonella* encoding human VEGFR-2.

[0070] Combining the attenuated mutant strain of *Salmonella* of the present invention with a second attenuated mutant strain comprising a DNA molecule encoding a second tumor antigen may have synergistic antitumor effects. In particular, simultaneous targeting of different tumor antigens may minimize the risk of tumor escape. Combining WT1 based cancer immunotherapy with VEGFR-2 based immunotherapy may prove especially effective, since WT1 overexpressing tumor cells and the tumor vasculature are attacked at the same time.

[0071] In particular embodiments, the attenuated mutant strain of *Salmonella* is co-administered with said one or more further attenuated mutant strain(s) of *Salmonella*.

[0072] In the context of the present invention, the term "co-administration" or "co-administer" means administration of two different attenuated mutant strains of

Salmonella within three consecutive days, more particularly within two consecutive days, more particularly on the same day, more particularly within 12 hours. Most particularly, in the context of the present invention, the term "co-administration" refers to simultaneous administration of two different attenuated mutant strains of Salmonella.

[0073] In particular embodiments, cancer immunotherapy is accompanied by chemotherapy, radiotherapy or biological cancer therapy. For cure of cancer, complete eradication of cancer stem cells may be essential. For maximal efficacy, a combination of different therapy approaches may be beneficial.

[0074] In the context of the present invention, the term "biological cancer therapy" or "cancer immunotherapy" refers to the stimulation of the patient's immune system to attack malignant tumor cells or the tumor stroma. Biological cancer therapy approaches include delivery of tumor antigens, delivery of therapeutic antibodies as drugs, administration of immunostimulatory cytokines and administration of immune cells.

[0075] Chemotherapeutic agents that may be used in combination with the attenuated mutant strain of Salmonella of the present invention may be, for example: gemcitabine, amifostine (ethiol), cabazitaxel, cisplatin, dacarbazine (DTIC), dactinomycin, docetaxel, mechlorethamine, streptozocin, cyclophosphamide, carmustine (BCNU), lomustine (CCNU), doxorubicin (adriamycin), doxorubicin lipo (doxil), folinic acid, gemcitabine (gemzar), daunorubicin, daunorubicin lipo (daunoxome), procarbazine, ketokonazole, mitomycin, cytarabine, etoposide, methotrexate, 5-fluorouracil (5-FU), vinblastine, vincristine, bleomycin, paclitaxel (taxol), docetaxel (taxotere), aldesleukin, asparaginase, busulfan, carboplatin, cladribine, camptothecin, CPT-11, 10-hydroxy-7-ethyl-camptothecin (SN38),

dacarbazine, floxuridine, fludarabine, hydroxyurea, ifosfamide, idarubicin, mesna, interferon alpha, interferon beta, irinotecan, mitoxantrone, topotecan, leuprolide, megestrol, melphalan, mercaptopurine, oxaliplatin, plicamycin, mitotane, pegaspargase, pentostatin, pipobroman, plicamycin, streptozocin, tamoxifen, teniposide, testolactone, thioguanine, thiotepa, uracil mustard, vinorelbine, chlorambucil and combinations thereof.

[0076] Most preferred chemotherapeutic agents according to the invention in combination with VXM06 are cabazitaxel, carboplatin, oxaliplatin, cisplatin, cyclophosphamide, docetaxel, gemcitabine, doxorubicin, paclitaxel (taxol), irinotecan, vincristine, vinblastine, vinorelbin, folinic acid, 5-fluorouracil and bleomycin, especially gemcitabine.

[0077] It may be also favorable dependent on the occurrence of possible side effects, to include treatment with antibiotics or anti-inflammatory agents.

[0078] Should adverse events occur that resemble hypersensitivity reactions mediated by histamine, leukotrienes, or cytokines, treatment options for fever, anaphylaxis, blood pressure instability, bronchospasm, and dyspnoea are available. Treatment options in case of unwanted T-cell derived auto-aggression are derived from standard treatment schemes in acute and chronic graft vs. host disease applied after stem cell transplantation. Cyclosporin and glucocorticoids are proposed as treatment options.

[0079] In the unlikely case of systemic *Salmonella typhi* Ty21a type infection, appropriate antibiotic therapy is recommended, for example with fluoroquinolones including ciprofloxacin or ofloxacin. Bacterial infections of the gastrointestinal tract are to be treated with respective agents, such as rifaximin.

[0080] In particular embodiments, the attenuated mutant strain of *Salmonella* is administered during the chemotherapy or the radiotherapy treatment cycle or during biological cancer therapy.

[0081] In particular embodiments, the attenuated mutant strain of *Salmonella* is administered before the chemotherapy or the radiotherapy treatment cycle or before biological cancer therapy. This approach may have the advantage that chemotherapy or radiotherapy can be performed under conditions of enhanced cancer immunity.

[0082] In particular embodiments, the attenuated mutant strain of *Salmonella* is administered after the chemotherapy or the radiotherapy treatment cycle or after biological cancer therapy.

[0083] In particular embodiments, the attenuated mutant strain of *Salmonella* is administered orally. Oral administration is simpler, safer and more comfortable than parenteral administration. WT1 peptide vaccines, which have been used in clinical trials for the most part, are usually administered subcutaneously or intradermally, often resulting in skin erythema and local inflammation reactions. These adverse effects may be overcome by oral administration of the DNA vaccine of the present invention. The attenuated mutant strain of *Salmonella* of the present invention may however also be administered by any other suitable route. Preferably, a therapeutically effective dose is administered to the subject, and this dose depends on the particular application, the type of malignancy, the subject's weight, age, sex and state of health, the manner of administration and the formulation, etc. Administration may be single or multiple, as required.

[0084] The attenuated mutant strain of *Salmonella* of the present invention may be provided in the form of a solution, a suspension, lyophilisate, or any other suitable form. It may be provided in combination with pharmaceutically acceptable carriers, diluents, and/or excipients. Agents for adjusting the pH value, buffers, agents for adjusting toxicity, and the like may also be included. In the context of the present invention, the term "pharmaceutically acceptable" refers to molecular entities and other ingredients of pharmaceutical compositions that are physiologically tolerable and do not typically produce untoward reactions when administered to a mammal (e.g., human). The term "pharmaceutically acceptable" may also mean approved by a regulatory agency of a Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in mammals, and, more particularly, in humans.

[0085] In particular embodiments, the cancer is selected from leukemia, particularly from acute myeloid leukemia (AML) and acute lymphoid leukemia (ALL), and from solid tumors, particularly from lung cancer, breast cancer, esophageal, colon, colorectal, gastric, cholangioductal, pancreatic cancer, glioblastoma, head and neck cancer, synovial sarcoma, angiosarcoma, osteosarcoma, thyroid cancer, cervical, endometrial, ovarian cancer, neuroblastoma, rhabdomyosarcoma, prostate cancer.

[0086] The vaccine of the present invention is surprisingly effective at relatively low doses. In particular embodiments, the single dose is from about 10^5 to about 10^{11} , particularly from about 10^6 to about 10^{10} , more particularly from about 10^6 to about 10^9 , more particularly from about 10^6 to about 10^8 , most particularly from about 10^6 to about 10^7 colony forming units (CFU). Administration of low doses of this live bacterial vaccine minimizes the risk of excretion and thus of transmission to third parties.

[0087] In this context, the term "about" or "approximately" means within a factor of 3, alternatively within a factor of 2, including within a factor of 1.5 of a given value or range.

[0088] In particular embodiments, the attenuated mutant strain of *Salmonella* is for use in individualized cancer immunotherapy comprising the step of assessing the tumor antigen expression pattern and/or stroma antigen expression pattern of a patient.

[0089] VXM06 can be used - either by itself or in combination with other *Salmonella typhi* Ty21a based cancer vaccines comprising eukaryotic expression systems – for the treatment of various cancer types. In particular embodiments, VXM06 may be used for individualized patient specific cancer treatment. For that purpose, the patient's tumor and/or stromal antigen expression pattern may be assessed in a first step for example by companion diagnostics targeting the patient's specific tumor and/or stromal antigen pattern. Depending on the patient's tumor and/or stromal antigen expression pattern VXM06 may be administered either alone or in combination with one or more suitable further *Salmonella typhi* Ty21a based cancer vaccine(s) comprising eukaryotic expression systems. Combinations of VXM06 with one or more further *Salmonella typhi* Ty21a based cancer vaccine(s) may however also be administered as fixed combinations. These cocktails combining two or more *Salmonella typhi* Ty21a based cancer vaccines can be composed from separate off the shelf products. The combinations, either fixed or individualized may contain VXM01 as anti-angiogenic basis therapy.

SHORT DESCRIPTION OF FIGURES AND TABLES

Figure 1: Amino acid sequence of truncated human WT1 encoded by WT1 cDNA contained in plasmid pVAX10.hWT1

Figure 2: Nucleic acid sequence of pVAX10.hWT1

Figure 3: Plasmid map of pVAX10.hWT1

Figure 4: Kaplan-Meier survival curves of mice bearing FBL-3 leukemia treated with VXM0m_empty, VXM06m and VXM06

Figure 5: Mean survival of mice bearing FBL-3 leukemia treated with VXM0m_empty, VXM06m and VXM06

Table 1: *In vitro* gene synthesis: phosphorylation reaction setup

Table 2: *In vitro* gene synthesis: amplification of ligation product - PCR profile

Table 3: Vaccine compositions

Table 4: Assessment of antitumor activity – experimental design

Table 5: Survival of mice bearing FBL-3 leukemia treated with VXM0m_empty, VXM06m and VXM06

Table 6: Number of dead animals after tumor challenge over time

Table 7: Mean and median survival of mice bearing FBL-3 leukemia treated with VXM0m_empty, VXM06m and VXM06

EXAMPLES

Example 1: Preparation of recombinant plasmids pVAX10.mWT1 and pVAX10.hWT1

[0090] Human truncated WT1 (1116 bp, WT1 sequence according to UniProt reference sequence P19544-7, truncated by the zinc finger domain) and murine truncated WT1 (1101 bp, WT1 sequence according to UniProt reference sequence

P22561-5, truncated by the zinc finger domain) were cloned into the pVAX10 backbone derived of pVAX10.VR2-1. WT1 DNA fragments were generated by double-strand gene synthesis, where oligonucleotides were linked together using a thermostable ligase.

Oligo design and synthesis:

[0091] In a first step, the gene sequence of truncated human and truncated murine WT1 (truncated by the zinc finger domain) were subdivided into individual oligonucleotides of 40 to 50 bases using the software "SeqEditor" (Entelechon). The defined oligonucleotides overlapped and corresponded to both DNA strands. After synthesis of the oligonucleotides of both DNA strands, the oligonucleotides were diluted with 10 mM Tris (pH 8.5) to a final concentration of 50 pmol/μl.

Kinase reaction:

[0092] The *in vitro* synthesized oligonucleotides were then phosphorylated by incubation with T4 polynucleotide kinase in order to allow for subsequent ligation of the oligonucleotides. Forward and reverse oligonucleotides were phosphorylated.

[0093] The reaction setup is summarized in the following Table 1:

Volume	Ingredient
10 μl	Primer mix
10 μl	10x T4 Polynucleotide kinase (PNK) buffer
10 μl	ATP (25 mM)
68 μl	Sterile water
2 μl	T4 PNK (10 U/μl)

[0094] The reaction mixture was incubated for 1 hour at 37°C in a water bath. Then the T4 polynucleotide kinase was inactivated by a five-minute heat step at 95°C and afterwards immediately cooled on ice until further treatment.

[0095] 12.5 µL of the kinase mixture was used directly for ligation (20 U Taq DNA ligase, 35 µl reaction volume (New England Biolabs, M0208S).

[0096] The denaturation step at 95°C was followed by progressive cooling (1°C/min). During this process, the complementary oligonucleotides assemble to a double strand DNA. The process was performed in a thermocycler (Personal Cycler, Biometra). By addition of the thermostable Taq DNA ligase, the 5'-PO₄-end of the oligonucleotides were linked with the free 3'-OH-end of the following oligonucleotide. By repeated denaturation and renaturation steps, mismatched oligonucleotides were released. At the end of the program, the mixture was cooled at 4°C.

Amplification of the ligation products by PCR:

[0097] 5 µl of the obtained ligation products (about 1.1 kb) were amplified by PCR in a 50 µl volume with flanking primers as depicted in the following Table 2:

no	step	temperature	time	No of cycles
1	denaturation	95°C	5 min	1
2	denaturation	95°C	30 sec	} 30x
3	annealing	57°C	30 sec	
4	elongation	72°C	90 sec	
5	extra-elongation	72°C	5 min	1
6	cooling	4°C	∞	1

[0098] The PCR mixture contained the following components: 0.2 – 0.5 μ M of each primer, 20 mM Tris-Cl, pH 8.8, 10 mM KCl, 10 mM $(\text{NH}_4)_2\text{SO}_4$, 2 mM MgSO_4 , 0.1% Triton X-100, 25 mM dNTP (each), 2 U Vent_R polymerase.

[0099] The *in vitro* synthesized DNA fragments (human and murine truncated WT1; about 1.1 kb each) were cloned into the pVAX10 backbone via *NheI/XhoI* (the VEGFR-2 coding region of recombinant plasmid pVAX10.VR2-1 was replaced by truncated human or murine WT1). For quality control, the entire plasmids were sequenced and aligned to the respective reference sequence after transformation into *E. coli*. Both sequences proved to be free of errors. The resulting plasmids were designated pVAX10.mWT1 and pVAX10.hWT1.

Example 2: Transformation of attenuated Salmonella strains with the recombinant plasmids:

[00100] *S. typhi* Ty 21a was transformed with plasmid pVAX10.hWT1. *S. typhimurium* SL7207 (aroA⁻) was transformed with plasmid pVAX10.mWT1. The transformation was performed by electroporation.

Preparation of competent Salmonella cells:

[00101] Glycerol cultures of *S. typhi* Ty21a and *S. typhimurium* SL7207 were inoculated on LB plates (animal component free [ACF] soy peptone). The plates were incubated at 37°C overnight. One colony each was used for overnight-liquid-preculture. 3 ml LB medium (ACF soy peptone) inoculated with one colony each was incubated at 37°C and 180 rpm overnight. To prepare competent cells, 2 x 300 ml of LB medium (ACF soy peptone) were inoculated with 3 ml of the overnight culture and incubated at 37°C and 180 rpm up to an OD₆₀₀ of about 0.5. The cultures were then put on ice for 10 minutes. Subsequently, the bacteria were centrifuged for 10 minutes

at 3000xg at 4°C and each pellet was resuspended in 500 mL of ice cold H₂O_{dest.} After a new centrifugation step, the bacterial pellets were washed twice in 10% ice cold glycerol. Both pellets were put together in 2 ml of 10% glycerol and finally frozen in aliquots of 50 µL on dry ice. The used glycerol did not contain any animal ingredients (Sigma Aldrich, G5150).

Transformation of competent Salmonella cells:

[00102] For each transformation reaction, a 50 µl aliquot of competent cells was thawed on ice for 10 minutes. After adding 3-5 µL of plasmid DNA (pVAX10.hWT1 for competent *S. typhi* Ty21a cells and pVAX10.mWT1 for competent *S. typhimurium* SL7207 cells) the mixtures were incubated on ice for five minutes. Subsequently, the mixtures were transferred to pre-cooled cuvettes (1 mm thickness). The electric pulse was carried out at 12.5 kV/cm. Immediately afterwards, 1ml of LB medium (ACF soy peptone) was added to the cells, the cells were transferred into a 2 ml Eppendorf tube and shaken for 1 hour at 37°C. After a short centrifugation step on a bench centrifuge (16600 rcf, 20s), the bacterial pellet was resuspended in 200 µl of LB (ACF soy peptone) antibiotic-free medium. The mixtures were applied with a Drigalski spatula on LB plates (ACF soy peptone) containing kanamycin (concentration = 25 µg/ml or 50 µg/ml). The plates were incubated at 37°C overnight.

Plasmid preparation of recombinant Salmonella clones:

[00103] Three clones of each recombinant Salmonella strain were incubated overnight in 3 ml of LB medium (ACF soy peptone) containing kanamycin (50 µg/ml) at 37°C. The bacterial culture was then pelleted by centrifugation (16600 rcf, 30 s). Plasmid isolation was performed using the NucleoSpin Plasmid Kit from Macherey-

Nagel. The plasmid DNA was eluted from the silica gel columns with 50 µl water. 5 µl of the eluate was used in agarose gel electrophoresis for control.

[00104] For long-term storage, 1ml glycerol cultures of the positive clones were produced. For this purpose, 172 µl glycerol (no animal ingredients) was added to 828 µl medium of a logarithmically growing 3 ml culture in a 1 low ml screw microtube. The samples were stored at -70°C until further use.

Complete sequencing of recombinant plasmid DNA isolated from Salmonella:

[00105] 3ml of liquid LB-Kan medium (ACF soy peptone) were inoculated with one colony of recombinant Salmonella (*S. typhi* Ty21a harboring pVAX10.hWT1 and *S. typhimurium* SL7207 harboring pVAX10.mWT1) and incubated overnight at 37°C and 180 rpm. The overnight culture was pelleted by centrifugation at 1300 rpm for 30 s on a bench centrifuge (Biofuge pico, Heraeus). The plasmid isolation was performed with the NucleoSpin Plasmid Kit from Macherey-Nagel. After alkaline lysis and precipitation of high molecular weight genomic DNA and cellular components, the plasmid DNA was loaded onto columns with a silica membrane. After a washing step, the plasmids were eluted from the column with 50 µl of sterile water and sequenced. The sequences were then compared with the respective reference sequence by clone specific alignments, i.e. the plasmid sequences of each Salmonella clone was one by one aligned with the reference sequence. All sequences were in line with the respective reference sequences. The recombinant Salmonella strains were designated VXM06 (*S. typhi* Ty21a harboring plasmid pVAX10.hWT1) and VXM06m (*S. typhimurium* SL7207 harboring plasmid pVAX10.mWT1).

Example 3: Assessing antitumor activity of VXM06 and VXM06m in syngeneic leukemia mouse model

[00106] The efficacy of VXM06 and VXM06m was assessed in a syngeneic leukemia C57/BL6J mouse model over a period of 43 days (with dose administration on alternate days for 4 occasions followed by leukemia cell inoculation at day 17). Two groups, each comprising 10 male mice (n=10) either received VXM06 (*S. typhi* Ty21a containing pVAX10.hWT1 coding for truncated human WT1) or VXM06m (*S. typhimurium* containing pVAX10.mWT1 coding for truncated murine WT1) at doses of 10^{10} CFU/occasion. One similarly constituted control group received VXM0m_empty (*S. typhimurium* vector control with no expression plasmid) at the same dose as the treated groups. During the study, body weight, mortality, and survival investigations were undertaken. Prior to this main study, a pilot study over 14 days in male C57B16 mice (n=5 per group) without vaccination was performed using 5.0×10^6 and 3.0×10^7 FBL-3 cells, after which the first cell concentration was judged as optimal for the main vaccination study.

DNA vaccines:

[00107] VXM06, VXM06m, and the *S. typhimurium* empty vector (VXM0m_empty) without expression plasmid were stored at $\leq -80^\circ\text{C}$ until use. Vials being used during vaccination were not frozen again but discarded afterwards. The DNA vaccines under investigation are characterized in the following Table 3:

Test item	Batch	Concentration	Quantity
VXM0m_empty	VXM01m-e.1-01/2010	10^{11} CFU/ml	0,7 ml/vial (10 vials)
VXM06m	VXM06m-031.1-01/2012	10^{11} CFU/ml	0,7 ml/vial (10 vials)
VXM06	VXM06-031.1-01/2012	10^{11} CFU/ml	0,7 ml/vial (10 vials)

Route of drug administration:

[00108] 100 µl of VXM0m_empty, VXM06m and VXM06 were applied per animal and application. After thawing of test items, application took place within 30 min. All test substances were administered by oral gavage (per os, P.O) via cannula with an injection volume of 100 µl/mouse.

[00109] Regardless of animal groups, each animal received pre-dose application buffer to neutralize acid medium in the stomach prior to dosing (100 µl/animal/application for all dose groups). This buffer contained 2.6 g sodium hydrogen carbonate, 1.7 g L-ascorbic acid, 0.2 g lactose monohydrate and 100 ml of drinking water. Pre-dose applications were performed up to 30 minutes prior to application of the test items.

Cell culture:

[00110] The WT1 overexpressing mouse leukemia cells FBL-3 required passaging to ensure viability and to attain the required amount of cells.

[00111] The exponentially growing tumor cells were collected, mixed with trypan blue (at the recommended dilution 1:1) for viability determination, and manually counted using a counting chamber under optical microscope. FBL-3 cells were washed and resuspended in serum-free RPMI medium for injections into C57BL/6 mice.

Vaccination and tumor cell inoculation:

[00112] Cell injection conditions for intraperitoneal (I.P.) injection: cell viability $\geq 97\%$; 5.0×10^6 cells/500 µl/mouse.

[00113] Animals (30 C57BL/6 mice, 4-6 weeks, male, $\approx$ 20 g each, Charles River, France) were numbered, given a unique animal identification ear notch mark; the body weight was measured twice a week.

[00114] Ten mice each ($n = 10$ male) were vaccinated with VXM0m_empty, VXM06m, and VMX06. Vaccination was carried out by oral gavage of 10^{10} CFU/application at days 1, 3, 5, and 7. At day 17, FBL-3 tumor cells were inoculated in mice by I.P. route. The experimental design is summarized in the following Table 4:

Group	Animals	Tumor cells	Treatment	Dose (CFU/adm)	Admin. route	Treatment schedule
1	10	5×10^6	VXM0m_empty	10^{10} (in 100 μ l)	PO	Q2Dx4 (D1, D3, D5, D7)
2	10	5×10^6	VXM06m	10^{10} (in 100 μ l)	PO	Q2Dx4 (D1, D3, D5, D7)
3	10	5×10^6	VXM96	10^{10} (in 100 μ l)	PO	Q2Dx4 (D1, D3, D5, D7)

Results:

[00115] Survival data for the three treatment groups are listed in the following Table 5:

	Group 1; VXM0m_empty	Group 2; VXM06m	Group 3; VXM06
Day	Survival	Survival	Survival
-3	10	10	10
1	10	10	10
5	9	9	10
7	9	9	10
12	9	9	10
15	9	9	10
19	9	8	10
22	9	8	10
26	9	8	10
29	5	7	9
33	0	4	4
36	0	1	3
40	0	1	1
43	0	0	0

[00116] A daily clinical examination of all animals was performed: behavior, signs of suffering (cachexia, becoming, and difficulties moving or feeding).

[00117] The number of dead animals after tumor challenge are listed in the following Table 6:

Time	Survival	Dead	treatment
1	9	0	VXM0m_empty
12	6	3	VXM0m_empty
15	2	4	VXM0m_empty
16	0	2	VXM0m_empty
1	8	0	VXM06m
12	7	1	VXM06m
14	6	1	VXM06m
16	4	2	VXM06m
19	1	3	VXM06m
26	0	1	VXM06m
1	10	0	VXM06
12	9	1	VXM06
14	8	1	VXM06
15	6	2	VXM06
16	4	2	VXM06
19	3	1	VXM06
v21	2	1	VXM06
22	1	1	VXM06
26	0	1	VXM06

[00118] Figure 4 depicts Kaplan-Meier survival curves of mice bearing FBL-3 leukemia treated with VXM0m_empty, VXM06m and VXM06. Mice treated with VXM06m and VXM06 survived longer compared to the control mice (up to 26 days). About 40% the mice that received VXM06m and VXM06 survived longer than those treated with VXM0m_empty. Mean and median survival of the three test groups are

depicted in the following Table 7:

Mice	VXM0m_empty 9	VMX06m 8	VXM06 10
	Survival time		
1	12	12	12
2	12	14	14
3	12	16	15
4	15	16	15
5	15	19	16
6	15	19	16
7	15	19	19
8	16	26	21
9	16		22
10			26
Median	15	17.5	16
Mean	14.2	17.6	17.6
SD	1.6	4.0	4.1
SEM	0.5	1.4	1.3
Var.	2.6	15.7	16.6

[00119] Figure 5 depicts the mean survival of mice bearing FBL-3 leukemia treated with VXM0m_empty, VXM06m and VXM06. Mice treated with VXM06m and VXM06 survived longer compared to the control mice.

[00120] This study demonstrated the effectiveness of the constructs VXM06 and VXM06m in targeting WT1 overexpressing leukemia cells in a mouse model. The control mice treated with empty vector survived for up to 16 days after tumor cell challenge (see fig. 4). In contrast, the mice treated with VXM06m or with VXM06 showed prolonged survival compared to the mice vaccinated with the control vector VXM0m_empty (up to 26 days for both test items). About 40% of the mice that received VXM06m or VXM06 survived longer than those in the control group treated with VXM0m_empty.

[00121] In summary, VXM06m and VXM06 showed a pharmacodynamic effect on the survival of test animals in this syngeneic C57B16 leukemia mouse model. A similar pharmacodynamic effect of the two compounds VXM06m and VXM06 compared to empty vector was observed. These results show that these vaccines were able to trigger a response against WT1 in an immune-competent mouse leukemia model resulting in longer survival of animals compared to animals treated with empty vector.

CLAIMS

1. An attenuated mutant strain of *Salmonella typhi* Ty21a comprising at least one copy of a recombinant DNA molecule comprising a eukaryotic expression cassette encoding Wilms' Tumor Protein (WT1).
2. The attenuated mutant strain of *Salmonella typhi* Ty21a of claim 1, wherein WT1 is selected from the group consisting of human WT1 and a protein that shares at least 80% sequence identity therewith, particularly wherein the zinc finger domain of WT1 is deleted, more particularly wherein WT1 has the amino acid sequence as found in SEQ ID NO 1.
3. The attenuated mutant strain of *Salmonella typhi* Ty21a of claim 1 or 2, wherein the recombinant DNA molecule comprises the kanamycin antibiotic resistance gene, the pMB1 ori, and a eukaryotic expression cassette encoding human WT1 or a protein that shares at least 80% sequence identity therewith, particularly wherein the zinc finger domain of WT1 is deleted, under the control of a CMV promoter.
4. The attenuated mutant strain of *Salmonella typhi* Ty21a of any one of claims 1 to 3 for use as a medicament.
5. Use of an attenuated mutant strain of *Salmonella typhi* Ty21a of any one of claims 1 to 3, in the manufacture of a vaccine.
6. The use of claim 5, wherein the vaccine is for cancer immunotherapy.
7. The use of claim 6, wherein cancer immunotherapy further comprises administration of one or more further attenuated mutant strain(s) of *Salmonella* comprising at least one copy of a recombinant DNA molecule comprising an expression cassette encoding a tumor antigen and/or a tumor stroma antigen, particularly wherein said one or more further attenuated mutant strain(s) of *Salmonella* is/are *Salmonella typhi* Ty21a comprising a eukaryotic expression cassette, more particularly wherein said one or more further attenuated mutant strain(s) of *Salmonella* comprise(s) an attenuated mutant strain of *Salmonella* encoding human VEGFR-2, more particularly wherein said one or more further attenuated mutant strain(s) of *Salmonella* comprise(s) the attenuated mutant strain of *Salmonella* designated VXM01.

8. The use of claim 7, wherein the attenuated mutant strain of *Salmonella* is co-administered with said one or more further attenuated mutant strain(s) of *Salmonella typhi* Ty21a.
9. The use of any one of claims 6 to 8, wherein cancer immunotherapy is accompanied by chemotherapy, radiotherapy or biological cancer therapy, particularly wherein the attenuated mutant strain of *Salmonella* is administered before or during the chemotherapy or the radiotherapy treatment cycle or before or during biological cancer therapy, or before and during the chemotherapy or the radiotherapy treatment cycle or the biological cancer therapy.
10. The use of any one of claims 5 to 9, wherein the attenuated mutant strain of *Salmonella typhi* Ty21a is administered orally.
11. The use of any one of claims 6 to 10, wherein the cancer is selected from leukemia, particularly from acute myeloid leukemia (AML) and acute lymphoid leukemia (ALL), and from solid tumors, particularly from lung cancer, breast cancer, esophageal, colon, colorectal, gastric, cholangioductal, pancreatic cancer, glioblastoma, head and neck cancer, synovial sarcoma, angiosarcoma, osteosarcoma, thyroid cancer, cervical, endometrial, ovarian cancer, neuroblastoma, rhabdomyosarcoma, prostate cancer.
12. The attenuated mutant strain of *Salmonella* of claim 4 or use in any one of claims 5 to 11, wherein the attenuated mutant strain of *Salmonella typhi* Ty21a or the vaccine is to be administered in a single dose, wherein the single dose comprises from about 10^5 to about 10^{11} , particularly from about 10^6 to about 10^{10} , more particularly from about 10^6 to about 10^9 , more particularly from about 10^6 to about 10^8 , most particularly from about 10^6 to about 10^7 colony forming units (CFU).
13. The use of any one of claims 6 to 12 for use in individualized cancer immunotherapy comprising the step of assessing the tumor antigen expression pattern and/or stroma antigen expression pattern of a patient.