

Description

Title of Invention: COMPOSITION FOR CLEAVING A TARGET DNA COMPRISING A GUIDE RNA SPECIFIC FOR THE TARGET DNA AND CAS PROTEIN-ENCODING NUCLEIC ACID OR CAS PROTEIN, AND USE THEREOF

Technical Field

- [1] The present invention relates to targeted genome editing in eukaryotic cells or organisms. More particularly, the present invention relates to a composition for cleaving a target DNA in eukaryotic cells or organisms comprising a guide RNA specific for the target DNA and Cas protein-encoding nucleic acid or Cas protein, and use thereof.

[2]

Background Art

- [3] CRISPRs (Clustered Regularly Interspaced Short Palindromic Repeats) are loci containing multiple short direct repeats that are found in the genomes of approximately 40% of sequenced bacteria and 90% of sequenced archaea. CRISPR functions as a prokaryotic immune system, in that it confers resistance to exogenous genetic elements such as plasmids and phages. The CRISPR system provides a form of acquired immunity. Short segments of foreign DNA, called spacers, are incorporated into the genome between CRISPR repeats, and serve as a memory of past exposures. CRISPR spacers are then used to recognize and silence exogenous genetic elements in a manner analogous to RNAi in eukaryotic organisms.
- [4] Cas9, an essential protein component in the Type II CRISPR/Cas system, forms an active endonuclease when complexed with two RNAs termed CRISPR RNA (crRNA) and trans-activating crRNA (tracrRNA), thereby slicing foreign genetic elements in invading phages or plasmids to protect the host cells. crRNA is transcribed from the CRISPR element in the host genome, which was previously captured from such foreign invaders. Recently, Jinek et al. (1) demonstrated that a single-chain chimeric RNA produced by fusing an essential portion of crRNA and tracrRNA could replace the two RNAs in the Cas9/RNA complex to form a functional endonuclease.
- [5] CRISPR/Cas systems offer an advantage to zinc finger and transcription activator-like effector DNA-binding proteins, as the site specificity in nucleotide binding CRISPR-Cas proteins is governed by a RNA molecule instead of the DNA-binding protein, which can be more challenging to design and synthesize.
- [6] However, until now, a genome editing method using the RNA-guided en-

endonuclease(RGEN) based on CRISPR/Cas system has not been developed.

[7]

[8] Meanwhile, Restriction fragment length polymorphism (RFLP) is one of the oldest, most convenient, and least expensive methods of genotyping that is still used widely in molecular biology and genetics but is often limited by the lack of appropriate sites recognized by restriction endonucleases.

[9] Engineered nuclease-induced mutations are detected by various methods, which include mismatch-sensitive T7 endonuclease I (T7E1) or Surveyor nuclease assays, RFLP, capillary electrophoresis of fluorescent PCR products, Dideoxy sequencing, and deep sequencing. The T7E1 and Surveyor assays are widely used but are cumbersome. Furthermore, these enzymes tend to underestimate mutation frequencies because mutant sequences can form homoduplexes with each other and cannot distinguish homozygous bi-allelic mutant clones from wildtype cells. RFLP is free of these limitations and therefore is a method of choice. Indeed, RFLP was one of the first methods to detect engineered nuclease-mediated mutations in cells and animals. Unfortunately, however, RFLP is limited by the availability of appropriate restriction sites. It is possible that no restriction sites are available at the target site of interest.

[10]

Disclosure of Invention

Technical Problem

[11] Until now, a genome editing and genotyping method using the RNA-guided endonuclease(RGEN) based on CRISPR/Cas system has not been developed.

[12] Under these circumstances, the present inventors have made many efforts to develop a genome editing method based on CRISPR/Cas system and finally established a programmable RNA-guided endonuclease that cleave DNA in a targeted manner in eukaryotic cells and organisms.

[13] In addition, the present inventors have made many efforts to develop a novel method of using RNA-guided endonucleases (RGENs) in RFLP analysis. They have used RGENs to genotype recurrent mutations found in cancer and those induced in cells and organisms by engineered nucleases including RGENs themselves, thereby completing the present invention.

[14]

Solution to Problem

[15] It is an object of the present invention to provide a composition for cleaving target DNA in eukaryotic cells or organisms comprising a guide RNA specific for target DNA or DNA that encodes the guide RNA, and Cas protein-encoding nucleic acid or Cas protein.

[16]

[17] It is another object of the present invention to provide a composition for inducing targeted mutagenesis in eukaryotic cells or organisms, comprising a guide RNA specific for target DNA or DNA that encodes the guide RNA, and Cas protein-encoding nucleic acid or Cas protein.

[18]

[19] It is still another object of the present invention to provide a kit for cleaving a target DNA in eukaryotic cells or organisms comprising a guide RNA specific for target DNA or DNA that encodes the guide RNA, and Cas protein-encoding nucleic acid or Cas protein.

[20]

[21] It is still another object of the present invention to provide a kit for inducing targeted mutagenesis in eukaryotic cells or organisms comprising a guide RNA specific for target DNA or DNA that encodes the guide RNA, and Cas protein-encoding nucleic acid or Cas protein.

[22]

[23] It is still another object of the present invention to provide a method for preparing a eukaryotic cell or organism comprising Cas protein and a guide RNA comprising a step of co-transfecting or serial-transfecting the eukaryotic cell or organism with a Cas protein-encoding nucleic acid or Cas protein, and a guide RNA or DNA that encodes the guide RNA.

[24]

[25] It is still another object of the present invention to provide a eukaryotic cell or organism comprising a guide RNA specific for target DNA or DNA that encodes the guide RNA, and Cas protein-encoding nucleic acid or Cas protein.

[26]

[27] It is still another object of the present invention to provide a method for cleaving a target DNA in eukaryotic cells or organisms comprising a step of transfecting the eukaryotic cells or organisms comprising a target DNA with a composition comprising a guide RNA specific for target DNA or DNA that encodes the guide RNA, and Cas protein-encoding nucleic acid or Cas protein.

[28]

[29] It is still another object of the present invention to provide a method for inducing targeted mutagenesis in a eukaryotic cell or organism comprising a step of treating a eukaryotic cell or organism with a composition comprising a guide RNA specific for target DNA or DNA that encodes the guide RNA, and Cas protein-encoding nucleic acid or Cas protein.

[30]

- [31] It is still another object of the present invention to provide an embryo, a genome-modified animal, or genome-modified plant comprising a genome edited by a composition comprising a guide RNA specific for target DNA or DNA that encodes the guide RNA, and Cas protein-encoding nucleic acid or Cas protein.
- [32]
- [33] It is still another object of the present invention to provide a method of preparing a genome-modified animal comprising a step of introducing the composition comprising a guide RNA specific for target DNA or DNA that encodes the guide RNA, and Cas protein-encoding nucleic acid or Cas protein into an embryo of an animal; and a step of transferring the embryo into a oviduct of pseudopregnant foster mother to produce a genome-modified animal.
- [34]
- [35] It is still another object of the present invention to provide a composition for genotyping mutations or variations in an isolated biological sample, comprising a guide RNA specific for the target DNA sequence Cas protein.
- [36]
- [37] It is still another object of the present invention to provide a
- [38] method of using a RNA-guided endonuclease (RGEN) to genotype mutations induced by engineered nucleases in cells or naturally-occurring mutations or variations, wherein the RGEN comprises a guide RNA specific for target DNA and Cas protein.
- [39]
- [40] It is still another object of the present invention to provide a kit for genotyping mutations induced by engineered nucleases in cells or naturally-occurring mutations or variations, comprising a RNA-guided endonuclease (RGEN), wherein the RGEN comprises a guide RNA specific for target DNA and Cas protein.
- [41]
- [42] It is an object of the present invention to provide a composition for cleaving target DNA in eukaryotic cells or organisms comprising a guide RNA specific for target DNA or DNA that encodes the guide RNA, and Cas protein-encoding nucleic acid or Cas protein.
- [43]
- [44] It is another object of the present invention to provide a composition for inducing targeted mutagenesis in eukaryotic cells or organisms, comprising a guide RNA specific for target DNA or DNA that encodes the guide RNA, and Cas protein-encoding nucleic acid or Cas protein.
- [45]
- [46] It is still another object of the present invention to provide a kit for cleaving a target

DNA in eukaryotic cells or organisms comprising a guide RNA specific for target DNA or DNA that encodes the guide RNA, and Cas protein-encoding nucleic acid or Cas protein.

[47]

[48] It is still another object of the present invention to provide a kit for inducing targeted mutagenesis in eukaryotic cells or organisms comprising a guide RNA specific for target DNA or DNA that encodes the guide RNA, and Cas protein-encoding nucleic acid or Cas protein.

[49]

[50] It is still another object of the present invention to provide a method for preparing a eukaryotic cell or organism comprising Cas protein and a guide RNA comprising a step of co-transfecting or serial-transfecting the eukaryotic cell or organism with a Cas protein-encoding nucleic acid or Cas protein, and a guide RNA or DNA that encodes the guide RNA.

[51]

[52] It is still another object of the present invention to provide a eukaryotic cell or organism comprising a guide RNA specific for target DNA or DNA that encodes the guide RNA, and Cas protein-encoding nucleic acid or Cas protein.

[53]

[54] It is still another object of the present invention to provide a method for cleaving a target DNA in eukaryotic cells or organisms comprising a step of transfecting the eukaryotic cells or organisms comprising a target DNA with a composition comprising a guide RNA specific for target DNA or DNA that encodes the guide RNA, and Cas protein-encoding nucleic acid or Cas protein.

[55]

[56] It is still another object of the present invention to provide a method for inducing targeted mutagenesis in a eukaryotic cell or organism comprising a step of treating a eukaryotic cell or organism with a composition comprising a guide RNA specific for target DNA or DNA that encodes the guide RNA, and Cas protein-encoding nucleic acid or Cas protein.

[57]

[58] It is still another object of the present invention to provide an embryo, a genome-modified animal, or genome-modified plant comprising a genome edited by a composition comprising a guide RNA specific for target DNA or DNA that encodes the guide RNA, and Cas protein-encoding nucleic acid or Cas protein.

[59]

[60] It is still another object of the present invention to provide a method of preparing a genome-modified animal comprising a step of introducing the composition comprising

a guide RNA specific for target DNA or DNA that encodes the guide RNA, and Cas protein-encoding nucleic acid or Cas protein into an embryo of an animal; and a step of transferring the embryo into a oviduct of pseudopregnant foster mother to produce a genome-modified animal.

[61]

[62] It is still another object of the present invention to provide a composition for genotyping mutations or variations in an isolated biological sample, comprising a guide RNA specific for the target DNA sequence Cas protein.

[63]

[64] It is still another object of the present invention to provide a composition for genotyping nucleic acid sequences in pathogenic microorganisms in an isolated biological sample, comprising a guide RNA specific for the target DNA sequence and Cas protein.

[65]

[66] It is still another object of the present invention to provide a kit for genotyping mutations or variations in an isolated biological sample, comprising the composition, specifically comprising a RNA-guided endonuclease (RGEN), wherein the RGEN comprises a guide RNA specific for target DNA and Cas protein.

[67]

[68] It is still another object of the present invention to provide a method of genotyping mutations or variations in an isolated biological sample, using the composition, specifically comprising a RNA-guided endonuclease (RGEN), wherein the RGEN comprises a guide RNA specific for target DNA and Cas protein.

Advantageous Effects of Invention

[69] The present composition for cleaving a target DNA or inducing a targeted mutagenesis in eukaryotic cells or organisms, comprising a guide RNA specific for the target DNA and Cas protein-encoding nucleic acid or Cas protein, the kit comprising the composition, and the method for inducing targeted mutagenesis provide a new convenient genome editing tools. In addition, because custom RGENs can be designed to target any DNA sequence, almost any single nucleotide polymorphism or small insertion/deletion (indel) can be analyzed via RGEN-mediated RFLP, therefore, the composition and method of the present invention may be used in detection and cleaving naturally-occurring variations and mutations.

[70]

Brief Description of Drawings

[71] Fig. 1 shows Cas9-catalyzed cleavage of plasmid DNA *in vitro*. (a) Schematic representation of target DNA and chimeric RNA sequences. Red triangles indicate cleavage

sites. The PAM sequence recognized by Cas9 is shown in bold. The sequences in the guide RNA derived from crRNA and tracrRNA are shown in box and underlined, respectively. (b) *In vitro* cleavage of plasmid DNA by Cas9. An intact circular plasmid or ApaLI-digested plasmid was incubated with Cas9 and guide RNA.

[72]

[73] Fig. 2 shows Cas9-induced mutagenesis at an episomal target site. (a) Schematic overview of cell-based assays using a RFP-GFP reporter. GFP is not expressed from this reporter because the GFP sequence is fused to the RFP sequence out-of-frame. The RFP-GFP fusion protein is expressed only when the target site between the two sequences is cleaved by a site-specific nuclease. (b) Flow cytometry of cells transfected with Cas9. The percentage of cells that express the RFP-GFP fusion protein is indicated.

[74]

[75] Fig. 3 shows RGEN-driven mutations at endogenous chromosomal sites. (a) CCR5 locus. (b) C4BPB locus. (Top) The T7E1 assay was used to detect RGEN-driven mutations. Arrows indicate the expected position of DNA bands cleaved by T7E1. Mutation frequencies (Indels (%)) were calculated by measuring the band intensities. (Bottom) DNA sequences of the CCR5 and C4BPB wild-type (WT) and mutant clones. The region of the target sequence complementary to the guide RNA is shown in box. The PAM sequence is shown in bold. Triangles indicate the cleavage site. Bases corresponding to microhomologies are underlined. The column on the right indicates the number of inserted or deleted bases.

[76]

[77] Fig. 4 shows that RGEN-driven off-target mutations are undetectable. (a) On-target and potential off-target sequences. The human genome was searched *in silico* for potential off-target sites. Four sites were identified, each of which carries 3-base mismatches with the CCR5 on-target site. Mismatched bases are underlined. (b) The T7E1 assay was used to investigate whether these sites were mutated in cells transfected with the Cas9/RNA complex. No mutations were detected at these sites. N/A (not applicable), an intergenic site. (c) Cas9 did not induce off-target-associated chromosomal deletions. The CCR5-specific RGEN and ZFN were expressed in human cells. PCR was used to detect the induction of the 15-kb chromosomal deletions in these cells.

[78]

[79] Fig. 5 shows RGEN-induced Foxn1 gene targeting in mice. (a) A schematic diagram depicting a sgRNA specific to exon 2 of the mouse Foxn1 gene. PAM in exon 2 is shown in red and the sequence in the sgRNA that is complementary to exon 2 is underlined. Triangles indicate cleavage sites. (b) Representative T7E1 assays demon-

strating gene-targeting efficiencies of Cas9 mRNA plus Foxn1-specific sgRNA that were delivered via intra-cytoplasmic injection into one-cell stage mouse embryos. Numbers indicate independent founder mice generated from the highest dose. Arrows indicate bands cleaved by T7E1. (c) DNA sequences of mutant alleles observed in three Foxn1 mutant founders identified in b. The number of occurrences is shown in parentheses. (d) PCR genotyping of F1 progenies derived from crossing Foxn1 founder #108 and wild-type FVB/NTac. Note the segregation of the mutant alleles found in Foxn1 founder #108 in the progenies.

[80]

[81] Fig. 6 shows Foxn1 gene targeting in mouse embryos by intra-cytoplasmic injection of Cas9 mRNA and Foxn1-sgRNA. (a) A representative result of a T7E1 assay monitoring the mutation rate after injecting the highest dose. Arrows indicate bands cleaved by T7E1. (b) A summary of T7E1 assay results. Mutant fractions among *in vitro* cultivated embryos obtained after intra-cytoplasmic injection of the indicated RGEN doses are indicated. (c) DNA sequences of Foxn1 mutant alleles identified from a subset of T7E1-positive mutant embryos. The target sequence of the wild-type allele is denoted in box.

[82]

[83] Fig. 7 shows Foxn1 gene targeting in mouse embryos using the recombinant Cas9 protein: Foxn1-sgRNA complex. (a) and (b) are representative T7E1 assays results and their summaries. Embryos were cultivated *in vitro* after they underwent pronuclear (a) or intra-cytoplasmic injection (b). Numbers in red indicate T7E1-positive mutant founder mice. (c) DNA sequences of Foxn1 mutant alleles identified from the *in vitro* cultivated embryos that were obtained by the pronucleus injection of recombinant Cas9 protein: Foxn1-sgRNA complex at the highest dose. The target sequence of the wild-type allele is denoted in box.

[84]

[85] Fig. 8 shows Germ-line transmission of the mutant alleles found in Foxn1 mutant founder #12. (a) fPCR analysis. (b) PCR genotyping of wild-type FVB/NTac, the founder mouse, and their F1 progenies.

[86]

[87] Fig. 9 shows Genotypes of embryos generated by crossing Prkdc mutant founders. Prkdc mutant founders ♂25 and ♀15 were crossed and E13.5 embryos were isolated. (a) fPCR analysis of wild-type, founder ♂25, and founder ♀15. Note that, due to the technical limitations of fPCR analysis, these results showed small differences from the precise sequences of the mutant alleles; e.g., from the sequence analysis, Δ269/Δ61/WT and Δ5+1/+7/+12/WT were identified in founders ♂25 and ♀15, respectively. (b) Genotypes of the generated embryos.

- [88]
- [89] Fig. 10 shows Cas9 protein/sgRNA complex induced targeted mutation.
- [90]
- [91] Fig. 11 shows recombinant Cas9 protein-induced mutations in Arabidopsis protoplasts.
- [92]
- [93] Fig. 12 shows recombinant Cas9 protein-induced mutant sequences in the Arabidopsis BRI1 gene.
- [94]
- [95] Fig. 13 shows T7E1 assay showing endogenous CCR5 gene disruption in 293 cells by treatment of Cas9-mal-9R4L and sgRNA/C9R4LC complex.
- [96]
- [97] Fig. 14 (a, b) shows mutation frequencies at on-target and off-target sites of RGENs reported in Fu et al. (2013). T7E1 assays analyzing genomic DNA from K562 cells (R) transfected serially with 20 µg of Cas9-encoding plasmid and with 60 µg and 120 µg of *in vitro* transcribed GX19 crRNA and tracrRNA, respectively (1 x 10⁶ cells), or (D) co-transfected with 1 µg of Cas9-encoding plasmid and 1 µg of GX₁₉ sgRNA expression plasmid (2 x 10⁵ cells).
- [98]
- [99] Fig. 15 (a, b) shows comparison of guide RNA structure. Mutation frequencies of the RGENs reported in Fu et al. (2013) were measured at on-target and off-target sites using the T7E1 assay. K562 cells were co-transfected with the Cas9-encoding plasmid and the plasmid encoding GX19 sgRNA or GGX20 sgRNA. Off-target sites (OT1-3 etc.) are labeled as in Fu et al. (2013).
- [100]
- [101] Fig. 16 shows that *in vitro* DNA cleavage by Cas9 nickases. (a) Schematic overview of the Cas9 nuclease and the paired Cas9 nickase. The PAM sequences and cleavage sites are shown in box. (b) Target sites in the human AAVS1 locus. The position of each target site is shown in triangle. (c) Schematic overview of DNA cleavage reactions. FAM dyes (shown in box) were linked to both 5' ends of the DNA substrate. (d) DSBs and SSBs analyzed using fluorescent capillary electrophoresis. Fluorescently-labeled DNA substrates were incubated with Cas9 nucleases or nickases before electrophoresis.
- [102]
- [103] Fig. 17 shows comparison of Cas9 nuclease and nickase behavior. (a) On-target mutation frequencies associated with Cas9 nucleases (WT), nickases (D10A), and paired nickases. Paired nickases that would produce 5' overhangs or 3' overhangs are indicated. (b) Analysis of off-target effects of Cas9 nucleases and paired nickases. A

total of seven potential off-target sites for three sgRNAs were analyzed.

[104]

[105] Fig. 18 shows paired Cas9 nickases tested at other endogenous human loci. (a,c) The sgRNA target sites at human CCR5 and BRCA2 loci. PAM sequences are indicated in red. (b,d) Genome editing activities at each target site were detected by the T7E1 assay. The repair of two nicks that would produce 5' overhangs led to the formation of indels much more frequently than did those producing 3' overhangs.

[106]

[107] Fig. 19 shows that paired Cas9 nickases mediate homologous recombination. (a) Strategy to detect homologous recombination. Donor DNA included an XbaI restriction enzyme site between two homology arms, whereas the endogenous target site lacked this site. A PCR assay was used to detect sequences that had undergone homologous recombination. To prevent amplification of contaminating donor DNA, primers specific to genomic DNA were used. (b) Efficiency of homologous recombination. Only amplicons of a region in which homologous recombination had occurred could be digested with XbaI; the intensities of the cleavage bands were used to measure the efficiency of this method.

[108]

[109] Fig. 20 shows DNA splicing induced by paired Cas9 nickases. (a) The target sites of paired nickases in the human AAVS1 locus. The distances between the AS2 site and each of the other sites are shown. Arrows indicate PCR primers. (b) Genomic deletions detected using PCR. Asterisks indicate deletion-specific PCR products. (c) DNA sequences of deletion-specific PCR products obtained using AS2 and L1 sgRNAs. Target site PAM sequences are shown in box and sgRNA-matching sequences are shown in capital letters. Intact sgRNA-matching sequences are underlined. (d) A schematic model of paired Cas9 nickase-mediated chromosomal deletions. Newly-synthesized DNA strands are shown in box.

[110]

[111] Fig. 21 shows that paired Cas9 nickases do not induce translocations. (a) Schematic overview of chromosomal translocations between the on-target and off-target sites. (b) PCR amplification to detect chromosomal translocations. (c) Translocations induced by Cas9 nucleases but not by the nickase pair.

[112]

[113] Fig. 22 shows a conceptual diagram of the T7E1 and RFLP assays. (a) Comparison of assay cleavage reactions in four possible scenarios after engineered nuclease treatment in a diploid cell: (A) wild type, (B) a monoallelic mutation, (C) different biallelic mutations (hetero), and (D) identical biallelic mutations (homo). Black lines represent PCR products derived from each allele; dashed and dotted boxes indicate

insertion/deletion mutations generated by NHEJ. (b) Expected results of T7E1 and RGEN digestion resolved by electrophoresis.

[114]

[115] Fig. 23 shows *in vitro* cleavage assay of a linearized plasmid containing the C4BPB target site bearing indels. DNA sequences of individual plasmid substrates (upper panel). The PAM sequence is underlined. Inserted bases are shown in box. Arrows (bottom panel) indicate expected positions of DNA bands cleaved by the wild-type-specific RGEN after electrophoresis.

[116]

[117] Fig. 24 shows genotyping of mutations induced by engineered nucleases in cells via RGEN-mediated RFLP. (a) Genotype of C4BPB mutant K562 cell clones. (b) Comparison of the mismatch-sensitive T7E1 assay with RGEN-mediated RFLP analysis. Black arrows indicate the cleavage product by treatment of T7E1 enzyme or RGENs.

[118]

[119] Fig. 25 shows genotyping of RGEN-induced mutations via the RGEN-RFLP technique. (a) Analysis of C4BPB-disrupted clones using RGEN-RFLP and T7E1 assays. Arrows indicate expected positions of DNA bands cleaved by RGEN or T7E1. (b) Quantitative comparison of RGEN-RFLP analysis with T7E1 assays. Genomic DNA samples from wild-type and C4BPB-disrupted K562 cells were mixed in various ratios and subjected to PCR amplification. (c) Genotyping of RGEN-induced mutations in the HLA-B gene in HeLa cells with RFLP and T7E1 analyses.

[120]

[121] Fig. 26 shows genotyping of mutations induced by engineered nucleases in organisms via RGEN-mediated RFLP. (a) Genotype of Pibf1 mutant founder mice. (b) Comparison of the mismatch-sensitive T7E1 assay with RGEN-mediated RFLP analysis. Black arrows indicate the cleavage product by treatment of T7E1 enzyme or RGENs.

[122]

[123] Fig. 27 shows RGEN-mediated genotyping of ZFN-induced mutations. The ZFN target site is shown in box. Black arrows indicate DNA bands cleaved by T7E1.

[124]

[125] Fig. 28 shows polymorphic sites in a region of the human HLA-B gene. The sequence, which surrounds the RGEN target site, is that of a PCR amplicon from HeLa cells. Polymorphic positions are shown in box. The RGEN target site and the PAM sequence are shown in dashed and bolded box, respectively. Primer sequences are underlined.

[126]

- [127] Fig. 29 shows genotyping of oncogenic mutations via RGEN-RFLP analysis. (a) A recurrent mutation (c.133-135 deletion of TCT) in the human CTNNB1 gene in HCT116 cells was detected by RGENs. HeLa cells were used as a negative control. (b) Genotyping of the KRAS substitution mutation (c.34 G>A) in the A549 cancer cell line with RGENs that contain mismatched guide RNA. Mismatched nucleotides are shown in box. HeLa cells were used as a negative control. Arrows indicate DNA bands cleaved by RGENs. DNA sequences confirmed by Sanger sequencing are shown.
- [128]
- [129] Fig. 30 shows genotyping of the CCR5 delta32 allele in HEK293T cells via RGEN-RFLP analysis. (a) RGEN-RFLP assays of cell lines. K562, SKBR3, and HeLa cells were used as wild-type controls. Arrows indicate DNA bands cleaved by RGENs. (b) DNA sequence of wild-type and delta32 CCR5 alleles. Both on-target and off-target sites of RGENs used in RFLP analysis are underlined. A single-nucleotide mismatch between the two sites is shown in box. The PAM sequence is underlined. (c) *In vitro* cleavage of plasmids harboring WT or del32 CCR5 alleles using the wild-type-specific RGEN. (d) Confirming the presence of an off-target site of the CCR5-delta32-specific RGEN at the CCR5 locus. *In vitro* cleavage assays of plasmids harboring either on-target or off-target sequences using various amounts of the del32-specific RGEN.
- [130]
- [131] Fig. 31 shows genotyping of a KRAS point mutation (c.34 G>A). (a) RGEN-RFLP analysis of the KRAS mutation (c.34 G>A) in cancer cell lines. PCR products from HeLa cells (used as a wild-type control) or A549 cells, which are homozygous for the point mutation, were digested with RGENs with perfectly matched crRNA specific to the wild-type sequence or the mutant sequence. KRAS genotypes in these cells were confirmed by Sanger sequencing. (b) Plasmids harboring either the wild-type or mutant KRAS sequences were digested using RGENs with perfectly matched crRNAs or attenuated, one-base mismatched crRNAs. Attenuated crRNAs that were chosen for genotyping are labeled in box above the gels.
- [132]
- [133] Fig. 32 shows genotyping of a PIK3CA point mutation (c.3140 A>G). (a) RGEN-RFLP analysis of the PIK3CA mutation (c.3140 A>G) in cancer cell lines. PCR products from HeLa cells (used as a wild-type control) or HCT116 cells that are heterozygous for the point mutation were digested with RGENs with perfectly matched crRNA specific to the wild-type sequence or the mutant sequence. PIK3CA genotypes in these cells were confirmed by Sanger sequencing. (b) Plasmids harboring either the wild-type or mutant PIK3CA sequences were digested using RGENs with perfectly matched crRNAs or attenuated, one-base mismatched crRNAs. Attenuated crRNAs that were chosen for genotyping are labeled in box above the gels.

[134]

[135] Fig. 33 shows genotyping of recurrent point mutations in cancer cell lines. (a) RGEN-RFLP assays of recurrent oncogenic point mutations in IDH (c.394c>T), (b) PIK3CA (c.3140A>G), (c) NRAS (c.181C>A), (d) and BRAF genes (c.1799T>A). Genotypes of each cell line confirmed by Sanger sequencing are shown. Mismatched nucleotides are shown in box. Black arrows indicate DNA bands cleaved by RGENs.

[136]

Best Mode for Carrying out the Invention

[137] In accordance with one aspect of the invention, the present invention provides a composition for cleaving target DNA in eukaryotic cells or organisms comprising a guide RNA specific for target DNA or DNA that encodes the guide RNA, and Cas protein-encoding nucleic acid or Cas protein. In addition, the present invention provides a use of the composition for cleaving target DNA in eukaryotic cells or organisms comprising a guide RNA specific for target DNA or DNA that encodes the guide RNA, and Cas protein-encoding nucleic acid or Cas protein.

[138]

[139] In the present invention, the composition is also referred to as a RNA-guided endonuclease(RGEN) composition.

[140]

[141] ZFNs and TALENs enable targeted mutagenesis in mammalian cells, model organisms, plants, and livestock, but the mutation frequencies obtained with individual nucleases are widely different from each other. Furthermore, some ZFNs and TALENs fail to show any genome editing activities. DNA methylation may limit the binding of these engineered nucleases to target sites. In addition, it is technically challenging and time-consuming to make customized nucleases.

[142]

[143] The present inventors have developed a new RNA-guided endonuclease composition based on Cas protein to overcome the disadvantages of ZFNs and TALENs.

[144]

[145] Prior to the present invention, an endonuclease activity of Cas proteins has been known. However, it has not been known whether the endonuclease activity of Cas protein would function in an eukaryotic cell because of the complexity of the eukaryotic genome. Further, until now, a composition comprising Cas protein or Cas protein-encoding nucleic acid and a guide RNA specific for the target DNA to cleave a target DNA in eukaryotic cells or organisms has not been developed.

[146]

[147] Compared to ZFNs and TALENs, the present RGEN composition based on Cas

protein can be more readily customized because only the synthetic guide RNA component is replaced to make a new genome-editing nuclease. No sub-cloning steps are involved to make customized RNA guided endonucleases. Furthermore, the relatively small size of the Cas gene (for example, 4.2 kbp for Cas9) as compared to a pair of TALEN genes (~6 kbp) provides an advantage for this RNA-guided endonuclease composition in some applications such as virus-mediated gene delivery. Further, this RNA-guided endonuclease does not have off-target effects and thus does not induce unwanted mutations, deletion, inversions, and duplications. These features make the present RNA-guided endonuclease composition a scalable, versatile, and convenient tools for genome engineering in eukaryotic cells and organisms. In addition, RGEN can be designed to target any DNA sequence, almost any single nucleotide polymorphism or small insertion/deletion (indel) can be analyzed via RGEN-mediated RFLP. The specificity of RGENs is determined by the RNA component that hybridizes with a target DNA sequence of up to 20 base pairs (bp) in length and by the Cas9 protein that recognize the protospacer-adjacent motif (PAM). RGENs are readily re-programmed by replacing the RNA component. Therefore, RGENs provide a platform to use simple and robust RFLP analysis for various sequence variations.

[148]

[149] The target DNA may be an endogenous DNA, or artificial DNA, preferably, endogenous DNA.

[150]

[151] As used herein, the term "Cas protein" refers to an essential protein component in the CRISPR/Cas system, forms an active endonuclease or nickase when complexed with two RNAs termed CRISPR RNA (crRNA) and trans-activating crRNA (tracrRNA).

[152] The information on the gene and protein of Cas are available from GenBank of National Center for Biotechnology Information (NCBI), without limitation.

[153] The CRISPR-associated (cas) genes encoding Cas proteins are often associated with CRISPR repeat-spacer arrays. More than forty different Cas protein families have been described. Of these protein families, Cas1 appears to be ubiquitous among different CRISPR/Cas systems. There are three types of CRISPR-Cas system. Among them, Type II CRISPR/Cas system involving Cas9 protein and crRNA and tracrRNA is representative and is well known. Particular combinations of cas genes and repeat structures have been used to define 8 CRISPR subtypes (Ecoli, Ypest, Nmeni, Dvulg, Tneap, Hmari, Aperi, and Mtube).

[154] The Cas protein may be linked to a protein transduction domain. The protein transduction domain may be poly-arginine or a TAT protein derived from HIV, but it is not limited thereto.

[155]

- [156] The present composition may comprise Cas component in the form of a protein or in the form of a nucleic acid encoding Cas protein.
- [157]
- [158] In the present invention, Cas protein may be any Cas protein provided that it has an endonuclease or nickase activity when complexed with a guide RNA.
- [159] Preferably, Cas protein is Cas9 protein or variants thereof.
- [160] The variant of the Cas9 protein may be a mutant form of Cas9 in which the catalytic aspartate residue is changed to any other amino acid. Preferably, the other amino acid may be an alanine, but it is not limited thereto.
- [161] Further, Cas protein may be the one isolated from an organism such as *Streptococcus* sp., preferably *Streptococcus pyogenes* or a recombinant protein, but it is not limited thereto.
- [162] The Cas protein derived from *Streptococcus pyogenes* may recognize NGG trinucleotide. The Cas protein may comprise an amino acid sequence of SEQ ID NO: 109, but it is not limited thereto.
- [163]
- [164] The term "recombinant" when used with reference, e.g., to a cell, nucleic acid, protein, or vector, indicates that the cell, nucleic acid, protein or vector, has been modified by the introduction of a heterologous nucleic acid or protein or the alteration of a native nucleic acid or protein, or that the cell is derived from a cell so modified. Thus, for example, a recombinant Cas protein may be generated by reconstituting Cas protein-encoding sequence using the human codon table.
- [165]
- [166] As for the present invention, Cas protein-encoding nucleic acid may be a form of vector, such as plasmid comprising Cas-encoding sequence under a promoter such as CMV or CAG. When Cas protein is Cas9, Cas9 encoding sequence may be derived from *Streptococcus* sp., and preferably derived from *Streptococcus pyogenes*. For example, Cas9 encoding nucleic acid may comprise the nucleotide sequence of SEQ ID NO: 1. Moreover, Cas9 encoding nucleic acid may comprise the nucleotide sequence having homology of at least 50% to the sequence of SEQ ID NO: 1, preferably at least 60, 70, 80, 90, 95, 97, 98, or 99% to the SEQ ID NO:1, but it is not limited thereto. Cas9 encoding nucleic acid may comprise the nucleotide sequence of SEQ ID NOs. 108, 110, 106, or 107.
- [167]
- [168] As used herein, the term "guide RNA" refers to a RNA which is specific for the target DNA and can form a complex with Cas protein and bring Cas protein to the target DNA.
- [169] In the present invention, the guide RNA may consist of two RNA, i.e., CRISPR

RNA(crRNA) and transactivating crRNA(tracrRNA) or be a single-chain RNA(sgRNA) produced by fusion of an essential portion of crRNA and tracrRNA.

- [170] The guide RNA may be a dualRNA comprising a crRNA and a tracrRNA.
- [171] If the guide RNA comprises the essential portion of crRNA and tracrRNA and a portion complementary to a target, any guide RNA may be used in the present invention.
- [172] The crRNA may hybridize with a target DNA.
- [173] The RGEN may consist of Cas protein, and dualRNA (invariable tracrRNA and target-specific crRNA), or Cas protein and sgRNA (fusion of an essential portion of invariable tracrRNA and target-specific crRNA), and may be readily reprogrammed by replacing crRNA.
- [174] The guide RNA further comprises one or more additional nucleotides at the 5' end of the single-chain guide RNA or the crRNA of the dualRNA.
- [175] Preferably, the guide RNA further comprises 2-additional guanine nucleotides at the 5' end of the single-chain guide RNA or the crRNA of the dualRNA.
- [176] The guide RNA may be transferred into a cell or an organism in the form of RNA or DNA that encodes the guide RNA. The guide RNA may be in the form of an isolated RNA, RNA incorporated into a viral vector, or is encoded in a vector. Preferably, the vector may be a viral vector, plasmid vector, or agrobacterium vector, but it is not limited thereto.
- [177] A DNA that encodes the guide RNA may be a vector comprising a sequence coding for the guide RNA. For example, the guide RNA may be transferred into a cell or organism by transfecting the cell or organism with the isolated guide RNA or plasmid DNA comprising a sequence coding for the guide RNA and a promoter.
- [178] Alternatively, the guide RNA may be transferred into a cell or organism using virus-mediated gene delivery.
- [179] When the guide RNA is transfected in the form of an isolated RNA into a cell or organism, the guide RNA may be prepared by *in vitro* transcription using any *in vitro* transcription system known in the art. The guide RNA is preferably transferred to a cell in the form of isolated RNA rather than in the form of plasmid comprising encoding sequence for a guide RNA. As used herein, the term "isolated RNA" may be interchangeable to "naked RNA". This is cost- and time-saving because it does not require a step of cloning. However, the use of plasmid DNA or virus-mediated gene delivery for transfection of the guide RNA is not excluded.
- [180]
- [181] The present RGEN composition comprising Cas protein or Cas protein-encoding nucleic acid and a guide RNA can specifically cleave a target DNA due to a specificity of the guide RNA for a target and an endonuclease or nickase activity of Cas protein.

[182]

[183] As used herein, the term "cleavage" refers to the breakage of the covalent backbone of a nucleotide molecule.

[184]

[185] In the present invention, a guide RNA may be prepared to be specific for any target which is to be cleaved. Therefore, the present RGEN composition can cleave any target DNA by manipulating or genotyping the target-specific portion of the guide RNA.

[186]

[187] The guide RNA and the Cas protein may function as a pair. As used herein, the term "paired Cas nickase" may refer to the guide RNA and the Cas protein functioning as a pair. The pair comprises two guide RNAs. The guide RNA and Cas protein may function as a pair, and induce two nicks on different DNA strand. The two nicks may be separated by at least 100 bps, but are not limited thereto.

[188]

[189] In the Example, the present inventors confirmed that paired Cas nickase allow targeted mutagenesis and large deletions of up to 1-kbp chromosomal segments in human cells. Importantly, paired nickases did not induce indels at off-target sites at which their corresponding nucleases induce mutations. Furthermore, unlike nucleases, paired nickases did not promote unwanted translocations associated with off-target DNA cleavages. In principle, paired nickases double the specificity of Cas9-mediated mutagenesis and will broaden the utility of RNA-guided enzymes in applications that require precise genome editing such as gene and cell therapy.

[190]

[191] In the present invention, the composition may be used in the genotyping of a genome in the eukaryotic cells or organisms *in vitro*.

[192]

[193] In one specific embodiment, the guide RNA may comprise the nucleotide sequence of Seq ID.No. 1, wherein the portion of nucleotide position 3 ~ 22 is a target-specific portion and thus, the sequence of this portion may be changed depending on a target.

[194]

[195] As used herein, a eukaryotic cell or organism may be yeast, fungus, protozoa, plant, higher plant, and insect, or amphibian cells, or mammalian cells such as CHO, HeLa, HEK293, and COS-1, for example, cultured cells (*in vitro*), graft cells and primary cell culture (*in vitro* and *ex vivo*), and *in vivo* cells, and also mammalian cells including human, which are commonly used in the art, without limitation.

[196]

[197] In one specific embodiment, it was found that Cas9 protein/single-chain guide RNA

could generate site-specific DNA double-strand breaks *in vitro* and in mammalian cells, whose spontaneous repair induced targeted genome mutations at high frequencies.

[198] Moreover, it was found that gene-knockout mice could be induced by the injection of Cas9 protein/guide RNA complexes or Cas9 mRNA/guide RNA into one-cell stage embryo and germ-line transmittable mutations could be generated by Cas9/guide RNA system.

[199] Using Cas protein rather than a nucleic acid encoding Cas protein to induce a targeted mutagenesis is advantageous because exogenous DNA is not introduced into an organism. Thus, the composition comprising Cas protein and a guide RNA may be used to develop therapeutics or value-added crops, livestock, poultry, fish, pets, etc.

[200]

[201] In accordance with another aspect of the invention, the present invention provides a composition for inducing targeted mutagenesis in eukaryotic cells or organisms, comprising a guide RNA specific for target DNA or DNA that encodes the guide RNA, and Cas protein-encoding nucleic acid or Cas protein. In addition, the present invention provides a use of the composition for inducing targeted mutagenesis in eukaryotic cells or organisms, comprising a guide RNA specific for target DNA or DNA that encodes the guide RNA, and Cas protein-encoding nucleic acid or Cas protein.

[202]

[203] A guide RNA, Cas protein-encoding nucleic acid or Cas protein are as described in the above.

[204]

[205] In accordance with another aspect of the invention, the present invention provides a kit for cleaving a target DNA or inducing targeted mutagenesis in eukaryotic cells or organisms comprising a guide RNA specific for target DNA or DNA that encodes the guide RNA, and Cas protein-encoding nucleic acid or Cas protein.

[206]

[207] A guide RNA, Cas protein-encoding nucleic acid or Cas protein are as described in the above.

[208]

[209] The kit may comprise a guide RNA and Cas protein-encoding nucleic acid or Cas protein as separate components or as one composition.

[210]

[211] The present kit may comprise some additional components necessary for transferring the guide RNA and Cas component to a cell or an organism. For example, the kit may comprise an injection buffer such as DEPC-treated injection buffer, and materials necessary for analysis of mutation of a target DNA, but are not limited thereto.

[212]

[213] In accordance with another aspect, the present invention provides a method for preparing a eukaryotic cell or organism comprising Cas protein and a guide RNA comprising a step of co-transfecting or serial-transfecting the eukaryotic cell or organism with a Cas protein-encoding nucleic acid or Cas protein, and a guide RNA or DNA that encodes the guide RNA.

[214]

[215] A guide RNA, Cas protein-encoding nucleic acid or Cas protein are as described in the above.

[216]

[217] In the present invention, a Cas protein-encoding nucleic acid or Cas protein and a guide RNA or DNA that encodes the guide RNA may be transferred into a cell by various methods known in the art, such as microinjection, electroporation, DEAE-dextran treatment, lipofection, nanoparticle-mediated transfection, protein transduction domain mediated transduction, virus-mediated gene delivery, and PEG-mediated transfection in protoplast, and so on, but are not limited thereto. Also, a Cas protein encoding nucleic acid or Cas protein and a guide RNA may be transferred into an organism by various method known in the art to administer a gene or a protein such as injection. A Cas protein-encoding nucleic acid or Cas protein may be transferred into a cell in the form of complex with a guide RNA, or separately. Cas protein fused to a protein transduction domain such as Tat can also be delivered efficiently into cells.

[218]

[219] Preferably, the eukaryotic cell or organisms is co-transfected or serial-transfected with a Cas9 protein and a guide RNA.

[220] The serial-transfection may be performed by transfection with Cas protein-encoding nucleic acid first, followed by second transfection with naked guide RNA. Preferably, the second transfection is after 3, 6, 12, 18, 24 hours, but it is not limited thereto.

[221]

[222] In accordance with another aspect, the present invention provides a eukaryotic cell or organism comprising a guide RNA specific for target DNA or DNA that encodes the guide RNA, and Cas protein-encoding nucleic acid or Cas protein.

[223]

[224] The eukaryotic cells or organisms may be prepared by transferring the composition comprising a guide RNA specific for target DNA or DNA that encodes the guide RNA, and Cas protein-encoding nucleic acid or Cas protein into the cell or organism.

[225]

[226] The eukaryotic cell may be yeast, fungus, protozoa, higher plant, and insect, or amphibian cells, or mammalian cells such as CHO, HeLa, HEK293, and COS-1, for

example, cultured cells (*in vitro*), graft cells and primary cell culture (*in vitro* and *ex vivo*), and *in vivo* cells, and also mammalian cells including human, which are commonly used in the art, without limitation. Further the organism may be yeast, fungus, protozoa, plant, higher plant, insect, amphibian, or mammal.

[227]

[228] In accordance with another aspect of the invention, the present invention provides a method for cleaving a target DNA or inducing targeted mutagenesis in eukaryotic cells or organisms, comprising a step of treating a cell or organism comprising a target DNA with a composition comprising a guide RNA specific for target DNA or DNA that encodes the guide RNA, and Cas protein-encoding nucleic acid or Cas protein.

[229]

[230] The step of treating a cell or organism with the composition may be performed by transferring the present composition comprising a guide RNA specific for target DNA or DNA that encodes the guide RNA, and Cas protein-encoding nucleic acid or Cas protein into the cell or organism.

[231] As described in the above, such transfer may be performed by microinjection, transfection, electroporation, and so on.

[232]

[233] In accordance with another aspect of the invention, the present invention provides an embryo comprising a genome edited by the present RGEN composition comprising a guide RNA specific for target DNA or DNA that encodes the guide RNA, and Cas protein-encoding nucleic acid or Cas protein.

[234]

[235] Any embryo can be used in the present invention, and for the present invention, the embryo may be an embryo of a mouse. The embryo may be produced by injecting PMSG (Pregnant Mare Serum Gonadotropin) and hCG (human Chorionic Gonadotropin) into a female mouse of 4 to 7 weeks and the super-ovulated female mouse may be mated to males, and the fertilized embryos may be collected from oviducts.

[236]

[237] The present RGEN composition introduced into an embryo can cleave a target DNA complementary to the guide RNA by the action of Cas protein and cause a mutation in the target DNA. Thus, the embryo into which the present RGEN composition has been introduced has an edited genome.

[238]

[239] In one specific embodiment, it was found that the present RGEN composition could cause a mutation in a mouse embryo and the mutation could be transmitted to offsprings.

[240]

- [241] A method for introducing the RGEN composition into the embryo may be any method known in the art, such as microinjection, stem cell insertion, retrovirus insertion, and so on. Preferably, a microinjection technique can be used.
- [242]
- [243] In accordance with another aspect, the present invention provides a genome-modified animal obtained by transferring the embryo comprising a genome edited by the present RGEN composition into the oviducts of an animal.
- [244]
- [245] In the present invention, the term “genome-modified animal” refers to an animal of which genome has been modified in the stage of embryo by the present RGEN composition and the type of the animal is not limited.
- [246]
- [247] The genome-modified animal has mutations caused by a targeted mutagenesis based on the present RGEN composition. The mutations may be any one of deletion, insertion, translocation, inversion. The site of mutation depends on the sequence of guide RNA of the RGEN composition.
- [248]
- [249] The genome-modified animal having a mutation of a gene may be used to determine the function of the gene.
- [250]
- [251] In accordance with another aspect of the invention, the present invention provides a method of preparing a genome-modified animal comprising a step of introducing the present RGEN composition comprising a guide RNA specific for the target DNA or DNA that encodes the guide RNA and Cas protein-encoding nucleic acid or Cas protein into an embryo of an animal; and a step of transferring the embryo into a oviduct of pseudopregnant foster mother to produce a genome-modified animal.
- [252]
- [253] The step of introducing the present RGEN composition may be accomplished by any method known in the art such as microinjection, stem cell insertion, retroviral insertion, and so on.
- [254]
- [255] In accordance with another aspect of the invention, the present invention provides a plant regenerated from the genome-modified protoplasts prepared by the method for eukaryotic cells comprising the RGEN composition.
- [256]
- [257] In accordance with another aspect of the invention, the present invention provides a composition for genotyping mutations or variations in an isolated biological sample, comprising a guide RNA specific for the target DNA sequence Cas protein. In addition,

the present invention provides a composition for genotyping nucleic acid sequences in pathogenic microorganisms in an isolated biological sample, comprising a guide RNA specific for the target DNA sequence and Cas protein.

[258]

[259] A guide RNA, Cas protein-encoding nucleic acid or Cas protein are as described in the above.

[260]

[261] As used herein the term "genotyping" refers to the "Restriction fragment length polymorphism (RFLP) assay".

[262] RFLP may be used in 1) the detection of indel in cells or organisms induced by the engineered nucleases, 2) the genotyping naturally-occurring mutations or variations in cells or organisms, or 3) the genotyping the DNA of infected pathogenic microorganisms including virus or bacteria, etc.

[263]

[264] The mutations or variation may be induced by engineered nucleases in cells.

[265] The engineered nuclease may be a Zinc Finger Nuclease (ZFNs), Transcription Activator-Like Effector Nucleases (TALENs), or RGENs, but it is not limited thereto.

[266]

[267] As used herein the term "biological sample" includes samples for analysis, such as tissues, cells, whole blood, serum, plasma, saliva, sputum, cerebrospinal fluid or urine, but is not limited thereto

[268]

[269] The mutations or variation may be a naturally-occurring mutations or variations.

[270]

[271] The mutations or variations are induced by the pathogenic microorganisms. Namely, the mutations or variation occur due to the infection of pathogenic microorganisms, when the pathogenic microorganisms are detected, the biological sample is identified as infected.

[272] The pathogenic microorganisms may be virus or bacteria, but are not limited thereto.

[273]

[274] Engineered nuclease-induced mutations are detected by various methods, which include mismatch-sensitive Surveyor or T7 endonuclease I (T7E1) assays, RFLP analysis, fluorescent PCR, DNA melting analysis, and Sanger and deep sequencing. The T7E1 and Surveyor assays are widely used but often underestimate mutation frequencies because the assays detect heteroduplexes (formed by the hybridization of mutant and wild-type sequences or two different mutant sequences); they fail to detect homoduplexes formed by the hybridization of two identical mutant sequences. Thus, these assays cannot distinguish homozygous biallelic mutant clones from wild-type

cells nor heterozygous biallelic mutants from heterozygous monoallelic mutants (Fig. 22). In addition, sequence polymorphisms near the nuclease target site can produce confounding results because the enzymes can cleave heteroduplexes formed by hybridization of these different wild-type alleles. RFLP analysis is free of these limitations and therefore is a method of choice. Indeed, RFLP analysis was one of the first methods used to detect engineered nuclease-mediated mutations. Unfortunately, however, it is limited by the availability of appropriate restriction sites.

[275]

[276] In accordance with another aspect of the invention, the present invention provides a kit for genotyping mutations or variations in an isolated biological sample, comprising the composition for genotyping mutations or variations in an isolated biological sample. In addition, the present invention provides a kit for genotyping nucleic acid sequences in pathogenic microorganisms in an isolated biological sample, comprising a guide RNA specific for the target DNA sequence and Cas protein.

[277]

[278] A guide RNA, Cas protein-encoding nucleic acid or Cas protein are as described in the above.

[279]

[280] In accordance with another aspect of the invention, the present invention provides a method of genotyping mutations or variations in an isolated biological sample, using the composition for genotyping mutations or variations in an isolated biological sample. In addition, the present invention provides a method of genotyping nucleic acid sequences in pathogenic microorganisms in an isolated biological sample, comprising a guide RNA specific for the target DNA sequence and Cas protein.

[281]

[282] A guide RNA, Cas protein-encoding nucleic acid or Cas protein are as described in the above.

[283]

Mode for the Invention

[284] Hereinafter, the present invention will be described in more detail with reference to Examples. However, these Examples are for illustrative purposes only, and the invention is not intended to be limited by these Examples.

[285]

[286] **Example 1: Genome editing assay**

[287]

[288] **1-1. DNA cleavage activity of Cas9 protein**

[289]

- [290] Firstly, the DNA cleavage activity of Cas9 derived from *Streptococcus pyogenes* in the presence or absence of a chimeric guide RNA *in vitro* was tested.
- [291] To this end, recombinant Cas9 protein that was expressed in and purified from *E. coli* was used to cleave a predigested or circular plasmid DNA that contained the 23-base pair (bp) human CCR5 target sequence. A Cas9 target sequence consists of a 20-bp DNA sequence complementary to crRNA or a chimeric guide RNA and the trinucleotide (5'-NGG-3') protospacer adjacent motif (PAM) recognized by Cas9 itself (Fig. 1A).
- [292] Specifically, the Cas9-coding sequence (4,104 bp), derived from *Streptococcus pyogenes* strain M1 GAS (NC_002737.1), was reconstituted using the human codon usage table and synthesized using oligonucleotides. First, 1-kb DNA segments were assembled using overlapping ~35-mer oligonucleotides and Phusion polymerase (New England Biolabs) and cloned into T-vector (SolGent). A full-length Cas9 sequence was assembled using four 1-kbp DNA segments by overlap PCR. The Cas9-encoding DNA segment was subcloned into p3s, which was derived from pcDNA3.1 (Invitrogen). In this vector, a peptide tag (NH₂-GGSGPPKKRKVYPYDVPDYA-COOH, SEQ ID NO: 2) containing the HA epitope and a nuclear localization signal (NLS) was added to the C-terminus of Cas9. Expression and nuclear localization of the Cas9 protein in HEK 293T cells were confirmed by western blotting using anti-HA antibody (Santa Cruz).
- [293] Then, the Cas9 cassette was subcloned into pET28-b(+) and transformed into BL21(DE3). The expression of Cas9 was induced using 0.5 mM IPTG for 4 h at 25 °C. The Cas9 protein containing the His₆-tag at the C terminus was purified using Ni-NTA agarose resin (Qiagen) and dialyzed against 20 mM HEPES (pH 7.5), 150 mM KCl, 1 mM DTT, and 10% glycerol (1). Purified Cas9 (50 nM) was incubated with supercoiled or pre-digested plasmid DNA (300 ng) and chimeric RNA (50 nM) in a reaction volume of 20 µl in NEB buffer 3 for 1 h at 37 °C. Digested DNA was analyzed by electrophoresis using 0.8% agarose gels.
- [294] Cas9 cleaved the plasmid DNA efficiently at the expected position only in the presence of the synthetic RNA and did not cleave a control plasmid that lacked the target sequence (Fig. 1B).
- [295]
- [296] **1-2. DNA cleavage by Cas9/guide RNA complex in human cells**
- [297]
- [298] A RFP-GFP reporter was used to investigate whether the Cas9/guide RNA complex can cleave the target sequence incorporated between the RFP and GFP sequences in mammalian cells.
- [299] In this reporter, the GFP sequence is fused to the RFP sequence out-of-frame (2). The

active GFP is expressed only when the target sequence is cleaved by site-specific nucleases, which causes frameshifting small insertions or deletions (indels) around the target sequence via error-prone non-homologous end-joining (NHEJ) repair of the double-strand break (DSB) (Fig 2).

- [300] The RFP-GFP reporter plasmids used in this study were constructed as described previously (2). Oligonucleotides corresponding to target sites (Table 1) were synthesized (Macrogen) and annealed. The annealed oligonucleotides were ligated into a reporter vector digested with EcoRI and BamHI.
- [301] HEK 293T cells were co-transfected with Cas9-encoding plasmid (0.8 μ g) and the RFP-GFP reporter plasmid (0.2 μ g) in a 24-well plate using Lipofectamine 2000 (Invitrogen).
- [302] Meanwhile, the *in vitro* transcribed chimeric RNA had been prepared as follows. RNA was *in vitro* transcribed through run-off reactions using the MEGAscript T7 kit (Ambion) according to the manufacturer's manual. Templates for RNA *in vitro* transcription were generated by annealing two complementary single strand DNAs or by PCR amplification (Table 1). Transcribed RNA was resolved on a 8% denaturing urea-PAGE gel. The gel slice containing RNA was cut out and transferred to probe elution buffer. RNA was recovered in nuclease-free water followed by phenol:chloroform extraction, chloroform extraction, and ethanol precipitation. Purified RNAs were quantified by spectrometry.
- [303] At 12h post transfection, chimeric RNA (1 μ g) prepared by *in vitro* transcription was transfected using Lipofectamine 2000.
- [304] At 3d post-transfection, transfected cells were subjected to flow cytometry and cells expressing both RFP and GFP were counted.
- [305] It was found that GFP-expressing cells were obtained only when the cells were transfected first with the Cas9 plasmid and then with the guide RNA 12 h later (Fig. 2), demonstrating that RGENs could recognize and cleave the target DNA sequence in cultured human cells. Thus GFP-expressing cells were obtained by serial-transfection of the Cas9 plasmid and the guide RNA rather than co-transfection.
- [306]
- [307] Table 1

[Table 1]

Gene		sequence (5' to 3')	SEQ ID NO.
Oligonucleotides used for the construction of the reporter plasmid			
CCR5	F	AATTCATGACATCAATTATTATACATCGGAG GAG	3
	R	GATCCTCCTCCGATGTATAATAATTGATGTC ATG	4
Primers used in the T7E1 assay			
CCR5	F1	CTCCATGGTGCTATAGAGCA	5
	F2	GAGCCAAGCTCTCCATCTAGT	6
	R	GCCCTGTCAAGAGTTGACAC	7
C4BPB	F1	TATTTGGCTGGTTGAAAGGG	8
	R1	AAAGTCATGAAATAAACACACCCA	9
	F2	CTGCATTGATATGGTAGTACCATG	10
	R2	GCTGTTCATTGCAATGGAATG	11
Primers used for the amplification of off-target sites			
ADCY5	F1	GCTCCCACCTTAGTGCTCTG	12
	R1	GGTGGCAGGAACCTGTATGT	13
	F2	GTCATTGGCCAGAGATGTGGA	14
	R2	GTCCCATGACAGGCGTGTAT	15
KCNJ6	F	GCCTGGCCAAGTTTCAGTTA	16
	R1	TGGAGCCATTGGTTTGCATC	17
	R2	CCAGAACTAAGCCGTTTCTGAC	18
CNTNAP2	F1	ATCACCGACAACCAGTTTCC	19
	F2	TGCAGTGCAGACTCTTTCCA	20
	R	AAGGACACAGGGCAACTGAA	21
N/A Chr. 5	F1	TGTGGAACGAGTGGTGACAG	22
	R1	GCTGGATTAGGAGGCAGGATTC	23

	F2	GTGCTGAGAACGCTTCATAGAG	24
	R2	GGACCAAACCACATTCTTCTCAC	25
Primers used for the detection of chromosomal deletions			
Deletion	F	CCACATCTCGTTCTCGGTTT	26
	R	TCACAAGCCCACAGATATTT	27

[308]

[309] **1-3. Targeted disruption of endogeneous genes in mammalian cells by RGEN**

[310]

[311] To test whether RGENs could be used for targeted disruption of endogenous genes in mammalian cells, genomic DNA isolated from transfected cells using T7 endonuclease I (T7E1), a mismatch-sensitive endonuclease that specifically recognizes and cleaves heteroduplexes formed by the hybridization of wild-type and mutant DNA sequences was analyzed (3).

[312]

[313] To introduce DSBs in mammalian cells using RGENs, 2×10^6 K562 cells were transfected with 20 μ g of Cas9-encoding plasmid using the 4D-Nucleofector, SF Cell Line 4D-Nucleofector X Kit, Program FF-120 (Lonza) according to the manufacturer's protocol. For this experiment, K562 (ATCC, CCL-243) cells were grown in RPMI-1640 with 10% FBS and the penicillin/streptomycin mix (100 U/ml and 100 μ g/ml, respectively).

[314] After 24h, 10-40 μ g of *in vitro* transcribed chimeric RNA was nucleofected into 1×10^6 K562 cells. The *in vitro* transcribed chimeric RNA had been prepared as described in the Example 1-2.

[315]

[316] Cells were collected two days after RNA transfection and genomic DNA was isolated. The region including the target site was PCR-amplified using the primers described in Table 1. The amplicons were subjected to the T7E1 assay as described previously (3). For sequencing analysis, PCR products corresponding to genomic modifications were purified and cloned into the T-Blunt vector using the T-Blunt PCR Cloning Kit (SolGent). Cloned products were sequenced using the M13 primer.

[317] It was found that mutations were induced only when the cells were transfected serially with Cas9-encoding plasmid and then with guide RNA (Fig. 3). Mutation frequencies (Indels (%)) in Fig. 3A) estimated from the relative DNA band intensities were RNA-dosage dependent, ranging from 1.3% to 5.1%. DNA sequencing analysis of the PCR amplicons corroborated the induction of RGEN-mediated mutations at the endogenous sites. Indels and microhomologies, characteristic of error-prone NHEJ,

were observed at the target site. The mutation frequency measured by direct sequencing was 7.3% (= 7 mutant clones/96 clones), on par with those obtained with zinc finger nucleases (ZFNs) or transcription-activator-like effector nucleases (TALENs).

[318]

[319] Serial-transfection of Cas9 plasmid and guide RNA was required to induce mutations in cells. But when plasmids that encode guide RNA, serial transfection was unnecessary and cells were co-transfected with Cas9 plasmid and guide RNA-encoding plasmid.

[320]

[321] In the meantime, both ZFNs and TALENs have been successfully developed to disrupt the human CCR5 gene (3-6), which encodes a G-protein-coupled chemokine receptor, an essential co-receptor of HIV infection. A CCR5-specific ZFN is now under clinical investigation in the US for the treatment of AIDS (7). These ZFNs and TALENs, however, have off-target effects, inducing both local mutations at sites whose sequences are homologous to the on-target sequence (6, 8-10) and genome rearrangements that arise from the repair of two concurrent DSBs induced at on-target and off-target sites (11-12). The most striking off-target sites associated with these CCR5-specific engineered nucleases reside in the CCR2 locus, a close homolog of CCR5, located 15-kbp upstream of CCR5. To avoid off-target mutations in the CCR2 gene and unwanted deletions, inversions, and duplications of the 15-kbp chromosomal segment between the CCR5 on-target and CCR2 off-target sites, the present inventors intentionally chose the target site of our CCR5-specific RGEN to recognize a region within the CCR5 sequence that has no apparent homology with the CCR2 sequence.

[322] The present inventors investigated whether the CCR5-specific RGEN had off-target effects. To this end, we searched for potential off-target sites in the human genome by identifying sites that are most homologous to the intended 23-bp target sequence. As expected, no such sites were found in the CCR2 gene. Instead, four sites, each of which carries 3-base mismatches with the on-target site, were found (Fig. 4A). The T7E1 assays showed that mutations were not detected at these sites (assay sensitivity, ~0.5%), demonstrating exquisite specificities of RGENs (Fig. 4B). Furthermore, PCR was used to detect the induction of chromosomal deletions in cells separately transfected with plasmids encoding the ZFN and RGEN specific to CCR5. Whereas the ZFN induced deletions, the RGEN did not (Fig. 4C).

[323]

[324] Next, RGENs was reprogrammed by replacing the CCR5-specific guide RNA with a newly-synthesized RNA designed to target the human C4BPB gene, which encodes the beta chain of C4b-binding protein, a transcription factor. This RGEN induced

mutations at the chromosomal target site in K562 cells at high frequencies (Fig. 3B). Mutation frequencies measured by the T7E1 assay and by direct sequencing were 14% and 8.3% (= 4 mutant clones/48 clones), respectively. Out of four mutant sequences, two clones contained a single-base or two-base insertion precisely at the cleavage site, a pattern that was also observed at the CCR5 target site. These results indicate that RGENs cleave chromosomal target DNA at expected positions in cells.

[325]

[326]

Example 2: Proteinaceous RGEN-mediated genome editing

[327]

[328]

RGENs can be delivered into cells in many different forms. RGENs consist of Cas9 protein, crRNA, and tracrRNA. The two RNAs can be fused to form a single-chain guide RNA (sgRNA). A plasmid that encodes Cas9 under a promoter such as CMV or CAG can be transfected into cells. crRNA, tracrRNA, or sgRNA can also be expressed in cells using plasmids that encode these RNAs. Use of plasmids, however, often results in integration of the whole or part of the plasmids in the host genome. The bacterial sequences incorporated in plasmid DNA can cause unwanted immune response *in vivo*. Cells transfected with plasmid for cell therapy or animals and plants derived from DNA-transfected cells must go through a costly and lengthy regulation procedure before market approval in most developed countries. Furthermore, plasmid DNA can persist in cells for several days post-transfection, aggravating off-target effects of RGENs.

[329]

Here, we used recombinant Cas9 protein complexed with *in vitro* transcribed guide RNA to induce targeted disruption of endogenous genes in human cells. Recombinant Cas9 protein fused with the hexa-histidine tag was expressed in and purified from *E. coli* using standard Ni ion affinity chromatography and gel filtration. Purified recombinant Cas9 protein was concentrated in storage buffer (20 mM HEPES pH 7.5, 150 mM KCl, 1 mM DTT, and 10% glycerol). Cas9 protein/sgRNA complex was introduced directly into K562 cells by nucleofection: 1×10^6 K562 cells were transfected with 22.5-225 (1.4-14 μ M) of Cas9 protein mixed with 100ug (29 μ M) of *in vitro* transcribed sgRNA (or crRNA 40ug and tracrRNA 80ug) in 100 μ l solution using the 4D-Nucleofector, SF Cell Line 4D-Nucleofector X Kit, Program FF-120 (Lonza) according to the manufacturer's protocol. After nucleofection, cells were placed in growth media in 6-well plates and incubated for 48hr. When 2×10^5 K562 cells were transfected with 1/5 scale-downed protocol, 4.5-45 μ g of Cas9 protein mixed with 6-60ug of *in vitro* transcribed sgRNA (or crRNA 8 μ g and tracrRNA 16 μ g) were used and nucleofected in 20 μ l solution. Nucleofected cell were then placed in growth media in 48-well plates. After 48hr, cells were collected and genomic DNA was isolated. The genomic DNA region spanning the target site was PCR-amplified and subjected to the

T7E1 assay.

[330] As shown in Fig. 10, Cas9 protein/sgRNA complex induced targeted mutation at the CCR5 locus at frequencies that ranged from 4.8 to 38% in a sgRNA or Cas9 protein dose-dependent manner, on par with the frequency obtained with Cas9 plasmid transfection (45%). Cas9 protein/crRNA/tracrRNA complex was able to induce mutations at a frequency of 9.4%. Cas9 protein alone failed to induce mutations. When 2×10^5 cells were transfected with 1/5 scale-downed doses of Cas9 protein and sgRNA, mutation frequencies at the CCR5 locus ranged from 2.7 to 57 % in a dose-dependent manner, greater than that obtained with co-transfection of Cas9 plasmid and sgRNA plasmid (32%).

[331] We also tested Cas9 protein/sgRNA complex that targets the ABCC11 gene and found that this complex induced indels at a frequency of 35%, demonstrating general utility of this method.

[332]

[333]

[334] Table 2

[Table 2]

Sequences of guide RNA

Target	RNA type	RNA sequence (5' to 3')	Length	SEQ ID NO
CCR5	sgRNA	GGUGACAUCAAUUAAUUAUACAUGUUU UAGAGCUAGAAAUAGCAAGUUAUAAAU AAGGCUAGUCCGUUAUCAACUUGAAAA AGUGGCACCGAGUCGGUGCUUUUUUU	104bp	28
	crRNA	GGUGACAUCAAUUAAUUAUACAUGUUU UAGAGCUAUGCUGUUUUUG	44bp	29
	tracrRNA	GGAACCAUUCAAAACAGCAUAGCAAGU UAAAAUAAGGCUAGUCCGUUAUCAACU UGAAAAAGUGGCACCGAGUCGGUGCUU UUUUU	86bp	30

[335]

[336] **Example 3: RNA-guided genome editing in mice**

[337]

[338] To examine the gene-targeting potential of RGENs in pronuclear (PN)-stage mouse embryos, the forkhead box N1 (Foxn1) gene, which is important for thymus development and keratinocyte differentiation (Nehls et al., 1996), and the protein kinase,

DNA activated, catalytic polypeptide (Prkdc) gene, which encodes an enzyme critical for DNA DSB repair and recombination (Taccioli et al., 1998) were used.

- [339] To evaluate the genome-editing activity of the Foxn1-RGEN, we injected Cas9 mRNA (10-ng/ μ l solution) with various doses of the sgRNA (Fig. 5a) into the cytoplasm of PN-stage mouse embryos, and conducted T7 endonuclease I (T7E1) assays (Kim et al. 2009) using genomic DNAs obtained from *in vitro* cultivated embryos (Fig. 6a).
- [340] Alternatively, we directly injected the RGEN in the form of recombinant Cas9 protein (0.3 to 30 ng/ μ l) complexed with the two-fold molar excess of Foxn1-specific sgRNA (0.14 to 14 ng/ μ l) into the cytoplasm or pronucleus of one-cell mouse embryos, and analyzed mutations in the Foxn1 gene using *in vitro* cultivated embryos (Fig. 7).
- [341]
- [342] Specifically, Cas9 mRNA and sgRNAs were synthesized *in vitro* from linear DNA templates using the mMESSAGE mMACHINE T7 Ultra kit (Ambion) and MEGAscript T7 kit (Ambion), respectively, according to the manufacturers' instructions, and were diluted with appropriate amounts of diethyl pyrocarbonate (DEPC, Sigma)-treated injection buffer (0.25 mM EDTA, 10 mM Tris, pH 7.4). Templates for sgRNA synthesis were generated using oligonucleotides listed in Table 3. Recombinant Cas9 protein was obtained from ToolGen, Inc.
- [343]
- [344] Table 3

[Table 3]

RNA Name	Direction	Sequence (5' to 3')	SEQ ID NO
Foxn1 #1 sgRNA	F	<u>GAAATTAATACGACTCACTATAGG</u> CA GTCTGACGTCACACTTCCGTTTTAGAGC TAGAAATAGCAAGTTAAAATAAGGCT AGTCCG	31
Foxn1 #2 sgRNA	F	<u>GAAATTAATACGACTCACTATAGG</u> ACT TCCAGGCTCCACCCGACGTTTTAGAGC TAGAAATAGCAAGTTAAAATAAGGCT AGTCCG	32
Foxn1 #3 sgRNA	F	<u>GAAATTAATACGACTCACTATAGG</u> CCA GGCTCCACCCGACTGGAGTTTTAGAGC TAGAAATAGCAAGTTAAAATAAGGCT AGTCCG	33
Foxn1 #4 sgRNA	F	<u>GAAATTAATACGACTCACTATAGG</u> ACT GGAGGGCGAACCCCAAGGTTTTAGAGC TAGAAATAGCAAGTTAAAATAAGGCT AGTCCG	34
Foxn1 #5 sgRNA	F	<u>GAAATTAATACGACTCACTATAGG</u> ACC CCAAGGGGACCTCATGCGTTTTAGAGC TAGAAATAGCAAGTTAAAATAAGGCT AGTCCG	35
Prkdc #1 sgRNA	F	<u>GAAATTAATACGACTCACTATAGG</u> TTA GTTTTTCCAGAGACTTGTTTTAGAGCT <u>AGAAATAGCAAGTTAAAATAAGGCTA</u> <u>GTCCG</u>	36
Prkdc #2 sgRNA	F	<u>GAAATTAATACGACTCACTATAGG</u> TTG GTTTGCTTGTTTATCGTTTTAGAGCT <u>AGAAATAGCAAGTTAAAATAAGGCTA</u> <u>GTCCG</u>	37
Prkdc #3 sgRNA	F	<u>GAAATTAATACGACTCACTATAGG</u> CAC AAGCAAACCAAAGTCTCGTTTTAGAGCT AGAAATAGCAAGTTAAAATAAGGCTA GTCCG	38

Prkdc #4 sgRNA	F	GAAATTAATACGACTCACTATAGGCCT CAATGCTAAGCGACTTCGTTTTAGAGCT AGAAATAGCAAGTTAAAATAAGGCTA GTCCG	39
-------------------	---	--	----

[345]

[346]

All animal experiments were performed in accordance with the Korean Food and Drug Administration (KFDA) guidelines. Protocols were reviewed and approved by the Institutional Animal Care and Use Committees (IACUC) of the Laboratory Animal Research Center at Yonsei University (Permit Number: 2013-0099). All mice were maintained in the specific pathogen-free facility of the Yonsei Laboratory Animal Research Center. FVB/NTac (Taconic) and ICR mouse strains were used as embryo donors and foster mothers, respectively. Female FVB/NTac mice (7-8 weeks old) were super-ovulated by intra-peritoneal injections of 5 IU pregnant mare serum gonadotropin (PMSG, Sigma) and 5 IU human chorionic gonadotropin (hCG, Sigma) at 48-hour intervals. The super-ovulated female mice were mated to FVB/NTac stud males, and fertilized embryos were collected from oviducts.

[347]

Cas9 mRNA and sgRNAs in M2 medium (Sigma) were injected into the cytoplasm of fertilized eggs with well-recognized pronuclei using a Piezo-driven micromanipulator (Prime Tech).

[348]

In the case of injection of recombinant Cas9 protein, the recombinant Cas9 protein: Foxn1-sgRNA complex was diluted with DEPC-treated injection buffer (0.25 mM EDTA, 10 mM Tris, pH 7.4) and injected into male pronuclei using a TransferMan NK2 micromanipulator and a FemtoJet microinjector (Eppendorf).

[349]

The manipulated embryos were transferred into the oviducts of pseudopregnant foster mothers to produce live animals, or were cultivated *in vitro* for further analyses.

[350]

[351]

To screen F0 mice and *in vitro* cultivated mouse embryos with RGEN-induced mutations, T7E1 assays were performed as previously described using genomic DNA samples from tail biopsies and lysates of whole embryos (Cho et al., 2013).

[352]

Briefly, the genomic region encompassing the RGEN target site was PCR-amplified, melted, and re-annealed to form heteroduplex DNA, which was treated with T7 endonuclease I (New England Biolabs), and then analyzed by agarose gel electrophoresis. Potential off-target sites were identified by searching with bowtie 0.12.9 and were also similarly monitored by T7E1 assays. The primer pairs used in these assays were listed in Tables 4 and 5.

[353]

[354]

Table 4

[Table 4]

Primers used in the T7E1 assay

Gene	Direction	Sequence(5' to 3')	SEQ ID NO
Foxn1	F1	GTCTGTCTATCATCTCTTCCCTTCTCTCC	40
	F2	TCCCTAATCCGATGGCTAGCTCCAG	41
	R1	ACGAGCAGCTGAAGTTAGCATGC	42
	R2	CTACTCAATGCTCTTAGAGCTACCAGGCTTG C	43
Prkdc	F	GACTGTTGTGGGGAGGGCCG	44
	F2	GGGAGGGCCGAAAGTCTTATTTTG	45
	R1	CCTGAAGACTGAAGTTGGCAGAAGTGAG	46
	R2	CTTTAGGGCTTCTTCTCTACAATCACG	47

[355]

[356] Table 5

[Table 5]

Primers used for amplification of off-target sites

Gene	Notation	Direction	Sequence(5' to 3')	SEQ ID NO
Foxn1	off 1	F	CTCGGTGTGTAGCCCTGAC	48
		R	AGACTGGCCTGGAACTCACA G	49
	off 2	F	CACTAAAGCCTGTCAGGAAG CCG	50
		R	CTGTGGAGAGCACACAGCAG C	51
	off 3	F	GCTGCGACCTGAGACCATG	52
		R	CTTCAATGGCTTCCTGCTTAG GCTAC	53
	off 4	F	GGTTCAGATGAGGCCATCCTT TC	54
		R	CCTGATCTGCAGGCTTAACCC TTG	55

Prkdc	off 1	F	CTCACCTGCACATCACATGTG G	56
		R	GGCATCCACCCTATGGGGTC	57
	off 2	F	GCCTTGACCTAGAGCTTAAAG AGCC	58
		R	GGTCTTGTTAGCAGGAAGGAC ACTG	59
	off 3	F	AAAACCTCTGCTTGATGGGATA TGTGGG	60
		R	CTCTCACTGGTTATCTGTGCT CCTTC	61
	off 4	F	GGATCAATAGGTGGTGGGGG ATG	62
		R	GTGAATGACACAATGTGACA GCTTCAG	63
	off 5	F	CACAAGACAGACCTCTCAAC ATTCAGTC	64
		R	GTGCATGCATATAATCCATTC TGATTGCTCTC	65
	off 6	F1	GGGAGGCAGAGGCAGGT	66
		F2	GGATCTCTGTGAGTTTGAGGC CA	67
		R1	GCTCCAGAACTCACTCTTAGG CTC	68

[357]

[358] Mutant founders identified by the T7E1 assay were further analyzed by fPCR. Appropriate regions of genomic DNA were sequenced as described previously (Sung et al., 2013). For routine PCR genotyping of F1 progenies, the following primer pairs were used for both wild-type and mutant alleles:

5'-CTACTCCCTCCGCAGTCTGA-3' (SEQ ID NO: 69) and

5'-CCAGGCCTAGGTTCCAGGTA-3' (SEQ ID NO: 70) for the Foxn1 gene,

5'-CCCCAGCATTGCAGATTTCC-3' (SEQ ID NO: 71) and

5'-AGGGCTTCTTCTCTACAATCACG-3' (SEQ ID NO: 72) for Prkdc gene.

[359] In the case of injection of Cas9 mRNA, mutant fractions (the number of mutant

embryos/the number of total embryos) were dose-dependent, ranging from 33% (1 ng/ μ l sgRNA) to 91% (100 ng/ μ l) (Fig. 6b). Sequence analysis confirmed mutations in the *Foxn1* gene; most mutations were small deletions (Fig. 6c), reminiscent of those induced by ZFNs and TALENs (Kim et al., 2013).

[360] In the case of injection of Cas9 protein, these injection doses and methods minimally affected the survival and development of mouse embryos *in vitro*: over 70% of RGEN-injected embryos hatched out normally in both experiments. Again, mutant fractions obtained with Cas9 protein injection were dose-dependent, and reached up to 88% at the highest dose via pronucleus injection and to 71% via intra-cytoplasmic injection (Figs. 7a and 7b). Similar to the mutation patterns induced by Cas9 mRNA plus sgRNA (Fig. 6c), those induced by the Cas9 protein-sgRNA complex were mostly small deletions (Fig. 7c). These results clearly demonstrate that RGENs have high gene-targeting activity in mouse embryos.

[361]

[362] Encouraged by the high mutant frequencies and low cytotoxicity induced by RGENs, we produced live animals by transferring the mouse embryos into the oviducts of pseudo-pregnant foster mothers.

[363] Notably, the birth rates were very high, ranging from 58% to 73%, and were not affected by the increasing doses of *Foxn1*-sgRNA (Table 6).

[364]

[365] Table 6

[Table 6]

RGEN-mediated gene-targeting in FVB/NTac mice

Target Gene	Cas9 mRNA + sgRNA (ng/ μ l)	Injected embryos	Transferred embryos (%)	Total newborns (%)	Live newborns* (%)	Founders [†] (%)
<i>Foxn1</i>	10 + 1	76	62 (82)	45 (73)	31 (50)	12 (39)
	10 + 10	104	90 (87)	52 (58)	58 (64)	33 (57)
	10 + 100	100	90 (90)	62 (69)	58 (64)	54 (93)
	Total	280	242 (86)	159 (66)	147 (61)	99 (67)
<i>Prkdc</i>	50 + 50	73	58 (79)	35 (60)	33 (57)	11 (33)
	50 + 100	79	59 (75)	22 (37)	21 (36)	7 (33)
	50 + 250	94	73 (78)	37 (51)	37 (51)	21 (57)
	Total	246	190 (77)	94 (49)	91 (48)	39 (43)

[366]

[367] Out of 147 newborns, we obtained 99 mutant founder mice. Consistent with the results observed in cultivated embryos (Fig. 6b), mutant fractions were proportional to the doses of Foxn1-sgRNA, and reached up to 93% (100 ng/μl Foxn1-sgRNA) (Tables 6 and 7, Fig. 5b).

[368]

[369] Table 7

[Table 7]

DNA sequences of Foxn1 mutant alleles identified from a subset of T7E1-positive mutant founders

ACTTCCAGGCTCCACCCGACTGGAGGGCGAACC CCAAGGGGACCTCATGCAGG	del+ins	#	Founder mice
ACTTCCAGGC-----AACCCCAAGGGGAC CTCATGCAGG	Δ19	1	20
ACTTCCAGGC-----GAACCCCAAGGGGA CCTCATGCAGG	Δ18	1	115
ACTTCCAGGCTCC-----	Δ60	1	19
ACTTCCAGGCTCC-----	Δ44	1	108
ACTTCCAGGCTCC-----CAAGGGGACC TCATGCAGG	Δ21	1	64
ACTTCCAGGCTCC-----TTAGGAGGCGAACCC CAAGGGGACCTCA	Δ12+6	1	126
ACTTCCAGGCTCCACC-----TCAT GCAGG	Δ28	1	5
ACTTCCAGGCTCCACCC-----CCAAGG GACCTCATG	Δ21+4	1	61
ACTTCCAGGCTCCACCC-----AAGGGGAC CTCATGCAGG	Δ18	2	95, 29
ACTTCCAGGCTCCACCC-----CAAGGGGA CCTCATGCAGG	Δ17	7	12, 14, 27, 66, 108, 114, 126
ACTTCCAGGCTCCACCC-----ACCCAAGGG GACCTCATGCAG	Δ15+1	1	32
ACTTCCAGGCTCCACCC-----CACCCAAGG GGACCTCATGCA	Δ15+2	1	124
ACTTCCAGGCTCCACCC-----ACCCAAGGG GACCTCATGCAGG	Δ13	1	32
ACTTCCAGGCTCCACCC-----GGCGAACCCCAAG GGGACCTCATGCAGG	Δ8	1	110
ACTTCCAGGCTCCACCCT-----GGGGAC CTCATGCAGG	Δ20+1	1	29

ACTTCCAGGCTCCACCCG-----AACCCCAAGG GGACCTCATGCAGG	$\Delta 11$	1	111
ACTTCCAGGCTCCACCCGA-----ACCT CATGCAGG	$\Delta 22$	1	79
ACTTCCAGGCTCCACCCGA-----GGGGAC CTCATGCAGG	$\Delta 18$	2	13, 127
ACTTCCAGGCTCCACCCCA-----AGGGGA CCTCATGCAGG	$\Delta 17$	1	24
ACTTCCAGGCTCCACCCGA-----ACCCCAAGG GGACCTCATGCAGG	$\Delta 11$	5	14, 53, 58, 69, 124
ACTTCCAGGCTCCACCCGA-----GACCCCAAGG GGACCTCATGCAGG	$\Delta 10$	1	14
ACTTCCAGGCTCCACCCGA----GGGCGAACCCCA AGGGGACCTCATGCAGG	$\Delta 5$	3	53, 79, 115
ACTTCCAGGCTCCACCCGAC-----CTC ATGCAGG	$\Delta 23$	1	108
ACTTCCAGGCTCCACCCGAC-----CCCCAAGG GGACCTCATGCAGG	$\Delta 11$	1	3
ACTTCCAGGCTCCACCCGAC-----GAAGGGCC CCAAGGGGACCTCA	$\Delta 11+6$	1	66
ACTTCCAGGCTCCACCCGAC-----GAACCCCAAG GGGACCTCATGCAGG	$\Delta 8$	2	3, 66
ACTTCCAGGCTCCACCCGAC----GGCGAACCCCA AGGGGACCTCATGCAGG	$\Delta 5$	1	27
ACTTCCAGGCTCCACCCGAC--GTGCTTGAGGGCG AACCCCAAGGGGACCTCA	$\Delta 2+6$	2	5
ACTTCCAGGCTCCACCCGACT-----CACTATCTTC TGGGCTCCTCCATGTC	$\Delta 6+25$	2	21, 114
ACTTCCAGGCTCCACCCGACT---TGGCGAACCCC AAGGGGACCTCATGCAG	$\Delta 4+1$	1	53
ACTTCCAGGCTCCACCCGACT--TGCAGGGCGAAC CCCAAGGGGACCTCATGC	$\Delta 2+3$	1	126
ACTTCCAGGCTCCACCCGACTTGGAGGGCGAAC CCCAAGGGGACCTCATGCAG	+1	15	3, 5, 12, 19, 29, 55, 56,

			61, 66, 68, 81, 108, 111, 124, 127
ACTTCCAGGCTCCACCCGACTTTGGAGGGCGAAC CCCAAGGGGACCTCATGCA	+2	2	79, 120
ACTTCCAGGCTCCACCCGACTGTTGGAGGGCGA ACCCCAAGGGGACCTCATGC	+3	1	55
ACTTCCAGGCTCCACCCGACTGGAG(+455)GGCG AACCCCAAGGGGACCTCC	+455	1	13

[370]

[371] To generate Prkdc-targeted mice, we applied a 5-fold higher concentration of Cas9 mRNA (50 ng/μl) with increasing doses of Prkdc-sgRNA (50, 100, and 250 ng/μl). Again, the birth rates were very high, ranging from 51% to 60%, enough to produce a sufficient number of newborns for the analysis (Table 6). The mutant fraction was 57% (21 mutant founders among 37 newborns) at the maximum dose of Prkdc-sgRNA. These birth rates obtained with RGENs were approximately 2- to 10-fold higher than those with TALENs reported in our previous study (Sung et al., 2013). These results demonstrate that RGENs are potent gene-targeting reagents with minimal toxicity.

[372] To test the germ-line transmission of the mutant alleles, we crossed the Foxn1 mutant founder #108, a mosaic with four different alleles (Fig. 5c, and Table 8) with wild-type mice, and monitored the genotypes of F1 offspring.

[373]

[374] Table 8

[Table 8]

Genotypes of Foxn1 mutant mice

Founder NO.	sgRNA (ng/ml)	Genotyping Summary	Detected alleles
58*	1	not determined	$\Delta 11$
19	100	bi-allelic	<u>$\Delta 60/+1$</u>
20	100	bi-allelic	<u>$\Delta 67/\Delta 19$</u>
13	100	bi-allelic	<u>$\Delta 18/+455$</u>
32	10	bi-allelic (heterozygote)	<u>$\Delta 13/\Delta 15+1$</u>
115	10	bi-allelic (heterozygote)	<u>$\Delta 18/\Delta 5$</u>
111	10	bi-allelic (heterozygote)	<u>$\Delta 11/+1$</u>
110	10	bi-allelic (homozygote)	<u>$\Delta 8/\Delta 8$</u>
120	10	bi-allelic (homozygote)	<u>$+2/+2$</u>
81	100	heterozygote	<u>$+1/WT$</u>
69	100	homozygote	<u>$\Delta 11/\Delta 11$</u>
55	1	mosaic	<u>$\Delta 18/\Delta 1/+1/+3$</u>
56	1	mosaic	<u>$\Delta 127/\Delta 41/\Delta 2/+1$</u>
127	1	mosaic	<u>$\Delta 18/+1/WT$</u>
53	1	mosaic	<u>$\Delta 11/\Delta 5/\Delta 4+1/WT$</u>
27	10	mosaic	<u>$\Delta 17/\Delta 5/WT$</u>
29	10	mosaic	<u>$\Delta 18/\Delta 20+1/+1$</u>
95	10	mosaic	<u>$\Delta 18/\Delta 14/\Delta 8/\Delta 4$</u>
108	10	mosaic	<u>$+1/\Delta 17/\Delta 23/\Delta 44$</u>
114	10	mosaic	<u>$\Delta 17/\Delta 8/\Delta 6+25$</u>
124	10	mosaic	<u>$\Delta 11/\Delta 15+2/+1$</u>
126	10	mosaic	<u>$\Delta 17/\Delta 2+3/\Delta 12+6$</u>
12	100	mosaic	<u>$\Delta 30/\Delta 28/\Delta 17/+1$</u>
5	100	mosaic	<u>$\Delta 28/\Delta 11/\Delta 2+6/+1$</u>
14	100	mosaic	<u>$\Delta 17/\Delta 11/\Delta 10$</u>
21	100	mosaic	<u>$\Delta 127/\Delta 41/\Delta 2/\Delta 6+25$</u>
24	100	mosaic	<u>$\Delta 17/+1/WT$</u>
64	100	mosaic	<u>$\Delta 31/\Delta 21/+1/WT$</u>

68	100	mosaic	<u>$\Delta 17/\Delta 11/+1/WT$</u>
79	100	mosaic	<u>$\Delta 22/\Delta 5/+2/WT$</u>
61	100	mosaic	<u>$\Delta 21+4/\Delta 6/+1/+9$</u>
66**	100	mosaic	<u>$\Delta 17/\Delta 8/\Delta 11+6/+1/WT$</u>
3	100	mosaic	<u>$\Delta 11/\Delta 8/+1$</u>

[375] Underlined alleles were sequenced.

[376] Alleles in red, detected by sequencing, but not by fPCR.

[377] *only one clone sequenced.

[378] **Not determined by fPCR.

[379]

[380] As expected, all the progenies were heterozygous mutants possessing the wild-type allele and one of the mutant alleles (Fig. 5d). We also confirmed the germ-line transmission in independent founder mice of Foxn1 (Fig. 8) and Prkdc (Fig. 9). To the best of our knowledge, these results provide the first evidence that RGEN-induced mutant alleles are stably transmitted to F1 progenies in animals.

[381]

[382] **Example 4: RNA-guided genome editing in plants**

[383]

[384] **4-1. Production of Cas9 protein**

[385]

[386] The Cas9 coding sequence (4104 bps), derived from Streptococcus pyogenes strain M1 GAS (NC_002737.1), was cloned to pET28-b(+) plasmid. A nuclear targeting sequence (NLS) was included at the protein N terminus to ensure the localization of the protein to the nucleus. pET28-b(+) plasmid containing Cas9 ORF was transformed into BL21(DE3). Cas9 was then induced using 0.2mM IPTG for 16 hrs at 18°C and purified using Ni-NTA agarose beads (Qiagen) following the manufacturer's instructions. Purified Cas9 protein was concentrated using Ultracel - 100K (Millipore).

[387]

[388] **4-2. Production of guide RNA**

[389]

[390] The genomic sequence of the Arabidopsis gene encoding the BRI1 was screened for the presence of a NGG motif, the so called protospacer adjacent motif (PAM), in an exon which is required for Cas9 targeting To disrupt the BRI1 gene in Arabidopsis, we identified two RGEN target sites in an exon that contain the NGG motif. sgRNAs were produced in vitor using template DNA. Each template DNA was generated by extension with two partially overlapped oligonucleotides (Macrogen, Table X1) and

Phusion polymerase (Thermo Scientific) using the following conditions - 98°C 30 sec {98°C 10 sec, 54°C 20 sec, 72°C 2 min}x20, 72°C 5 min.

[391]

[392] Table 9

[Table 9]

Oligonucleotides for the production of the template DNA for *in vitro* transcription

Oligonucleotides	Sequence (5'-3')	SEQ ID NO
BRI1 target 1 (Forward)	GAAATTAATACGACTCACTATAGGTTTGAA AGATGGAAGCGCGGGTTTTAGAGCTAGAA ATAGCAAGTTAAAATAAGGCTAGTCCG	73
BRI1 target 2 (Forward)	GAAATTAATACGACTCACTATAGGTGAAAC TAAACTGGTCCACAGTTTTAGAGCTAGAAA TAGCAAGTTAAAATAAGGCTAGTCCG	74
Universal (Reverse)	AAAAAAGCACCGACTCGGTGCCACTTTTTTC AAGTTGATAACGGACTAGCCTTATTTTAAC TTGC	75

[393]

[394] The extended DNA was purified and used as a template for the *in vitro* production of the guide RNA's using the MEGAscript T7 kit (Life Technologies). Guide RNA were then purified by Phenol/Chloroform extraction and ethanol precipitation. To prepare Cas9/sgrNA complexes, 10 μ l of purified Cas9 protein (12 μ g/ μ l) and 4 μ l each of two sgRNAs (11 μ g/ μ l) were mixed in 20 μ l NEB3 buffer (New England Biolabs) and incubated for 10 min at 37°C.

[395]

[396] 4-3. Transfection of Cas9/sgrNA complex to protoplast

[397]

[398] The leaves of 4-week-old Arabidopsis seedlings grown aseptically in petri dishes were digested in enzyme solution (1% cellulose R10, 0.5% macerozyme R10, 450 mM mannitol, 20mM MES pH 5.7 and CPW salt) for 8~16 hrs at 25°C with 40 rpm shaking in the dark. Enzyme/protoplast solutions were filtered and centrifuged at 100 X g for 3~5 min. Protoplasts were re-suspended in CPW solution after counting cells under the microscope (X100) using a hemacytometer. Finally, protoplasts were re-suspended at 1×10^6 /ml in MMG solution (4mM HEPES pH 5.7, 400 mM mannitol and 15 mM MgCl₂). To transfect the protoplasts with Cas9/sgrNA complex, 200 μ L (200,000 protoplasts) of the protoplast suspension were gently mixed with 3.3 or 10 μ L

of Cas9/sgRNA complex [Cas9 protein (6 $\mu\text{g}/\mu\text{L}$) and two sgRNAs (2.2 $\mu\text{g}/\mu\text{L}$ each)] and 200 μL of 40% polyethylene glycol transfection buffer (40% PEG4000, 200 mM mannitol and 100 mM CaCl_2) in 2 ml tubes. After 5~20 min incubation at room temperature, transfection was stopped by adding wash buffer with W5 solution (2 mM MES pH 5.7, 154 mM NaCl, 125 mM CaCl_2 and 5 mM KCl). Protoplasts were then collected by centrifugation for 5 min at 100 X g, washed with 1 ml of W5 solution, centrifuged for another 5 min at 100 X g. The density of protoplasts was adjusted to 1×10^5 /ml and they were cultured in modified KM 8p liquid medium with 400 mM glucose.

[399]

[400] **4-4. Detection of mutations in Arabidopsis protoplasts and plants**

[401]

[402] After 24 hr or 72 hr post-transfection, protoplasts were collected and genomic DNA was isolated. The genomic DNA region spanning the two target sites was PCR-amplified and subjected to the T7E1 assay. As shown in Figure 11, indels were induced by RGENs at high frequencies that ranged from 50% to 70%. Surprisingly, mutations were induced at 24 hr post-transfection. Apparently, Cas9 protein functions immediately after transfection. PCR products were purified and cloned into T-Blunt PCR Cloning Kit (Solgent). Plasmids were purified and subjected to Sanger sequencing with M13F primer. One mutant sequence had a 7-bp deletion at one site(Figure 12). The other three mutant sequences had deletions of ~220-bp DNA segments between the two RGEN site.

[403]

[404] **Example 5: Cas9 protein transduction using a cell-penetrating peptide or protein transduction domain**

[405]

[406] **5-1. Construction of His-Cas9-encoding plasmid**

[407]

[408] Cas9 with a cysteine at the C-terminal was prepared by PCR amplification using the previously described Cas9 plasmid {Cho, 2013 #166} as the template and cloned into pET28-(a) vector (Novagen, Merk Millipore, Germany) containing His-tag at the N-terminus.

[409]

[410] **5-2. Cell culture**

[411]

[412] 293T (Human embryonic kidney cell line), and HeLa (human ovarian cancer cell line) were grown in DMEM (GIBCO-BRL Rockville) supplemented with 10% FBS and 1% penicillin and streptomycin.

[413]

[414] **5-3. Expression and purification of Cas9 protein**

[415]

[416] To express the Cas9 protein, E. coli BL21 cells were transformed with the pET28-(a) vector encoding Cas9 and plated onto Luria-Bertani (LB) agar medium containing 50 µg/mL kanamycin (Amresco, Solon, OH). Next day, a single colony was picked and cultured in LB broth containing 50 µg/mL kanamycin at 37°C overnight. Following day, this starter culture at 0.1 OD₆₀₀ was inoculated into Luria broth containing 50 µg/mL kanamycin and incubated for 2 hrs at 37°C until OD₆₀₀ reached to 0.6-0.8. To induce Cas9 protein expression, the cells were cultured at 30°C overnight after addition of isopropyl-β-D-thiogalactopyranoside (IPTG) (Promega, Madison, WI) to the final concentration of 0.5mM.

[417] The cells were collected by centrifugation at 4000 rpm for 15-20 mins, resuspended in a lysis buffer (20mM Tris-Cl pH8.0, 300mM NaCl, 20mM imidazole, 1X protease inhibitor cocktail, 1 mg/ml lysozyme), and lysed by sonication (40% duty, 10 sec pulse, 30 sec rest, for 10 mins on ice). The soluble fraction was separated as the supernatant after centrifugation at 15,000 rpm for 20 mins at 4°C. Cas9 protein was purified at 4°C using a column containing Ni-NTA agarose resin (QIAGEN) and AKTA prime instrument (AKTA prime, GE Healthcare, UK). During this chromatography step, soluble protein fractions were loaded onto Ni-NTA agarose resin column (GE Healthcare, UK) at the flow rate of 1 mL/min. The column was washed with a washing buffer (20mM Tris-Cl pH8.0, 300mM NaCl, 20mM imidazole, 1X protease inhibitor cocktail) and the bound protein was eluted at the flow rate of 0.5 mL/min with an elution buffer (20mM Tris-Cl pH8.0, 300mM NaCl, 250mM imidazole, 1X protease inhibitor cocktail). The pooled eluted fraction was concentrated and dialyzed against storage buffer (50 mM Tris-HCl, pH8.0, 200 mM KCl, 0.1 mM EDTA, 1 mM DTT, 0.5 mM PMSF, 20% Glycerol). Protein concentration was quantitated by Bradford assay (Biorad, Hercules, CA) and purity was analyzed by SDS-PAGE using bovine serum albumin as the control.

[418]

[419] **5-4. Conjugation of Cas9 to 9R4L**

[420]

[421] 1mg Cas9 protein diluted in PBS at the concentration of 1mg/mL and 50 µg of maleimide-9R4L peptide in 25µL DW (Peptron, Korea) were gently mixed using a rotor at room temperature for 2 hrs and at 4°C overnight. To remove unconjugated maleimide-9R4L, the samples were dialyzed using 50kDa molecular weight cutoff membrane against DPBS (pH 7.4) at 4°C for 24 hrs. Cas9-9R4L protein was collected from the dialysis membrane and the protein amount was determined using

Bradford assay.

[422]

[423] **5-5. Preparation of sgRNA-9R4L**

[424]

[425] sgRNA (1 µg) was gently added to various amounts of C9R4LC peptide (ranging from 1 to 40 weight ratio) in 100 µl of DPBS (pH 7.4). This mixture was incubated at room temperature for 30 mins and diluted to 10 folds using RNase-free deionized water. The hydrodynamic diameter and z-potential of the formed nanoparticles were measured using dynamic light scattering (Zetasizer-nano analyzer ZS; Malvern instruments, Worcestershire, UK).

[426]

[427] **5-6. Cas9 protein and sgRNA treatments**

[428]

[429] Cas9-9R4L and sgRNA-C9R4LC were treated to the cells as follows: 1 µg of sgRNA and 15 µg of C9R4LC peptide were added to 250 mL of OPTIMEM medium and incubated at room temperature for 30 mins. At 24 hrs after seeding, cells were washed with OPTIMEM medium and treated with sgRNA-C9R4LC complex for 4 hrs at 37°C. Cells were washed again with OPTIMEM medium and treated with Cas9-9R4L for 2 hrs at 37°C. After treatment, culture media was replaced with serum-containing complete medium and incubated at 37°C for 24 hrs before the next treatment. Same procedure was followed for multiple treatments of Cas9 and sgRNA for three consecutive days.

[430]

[431] **5-7. Cas9-9R4L and sgRNA-9R4L can edit endogenous genes in cultured mammalian cells without the use of additional delivery tools**

[432]

[433] To determine whether Cas9-9R4L and sgRNA-9R4L can edit endogenous genes in cultured mammalian cells without the use of additional delivery tools, we treated 293 cells with Cas9-9R4L and sgRNA-9R4L targeting the CCR5 gene and analyzed the genomic DNA. T7E1 assay showed that 9% of CCR5 gene was disrupted in cells treated with both Cas9-9R4L and sgRNA-9R4L and that the CCR5 gene disruption was not observed in control cells including those untreated, treated with either Cas9-9R or sgRNA-9R4L, or treated with both unmodified Cas-9 and sgRNA (Fig 13), suggesting that the treatment with Cas9-9R4L protein and sgRNA conjugated with 9R4L, but not unmodified Cas9 and sgRNA, can lead to efficient genome editing in mammalian cells.

[434]

[435] **Example 6: Control of off-target mutation according to guide RNA structure**

[436]

[437] Recently, three groups reported that RGENs had off-target effects in human cells. To our surprise, RGENs induced mutations efficiently at off-target sites that differ by 3 to 5 nucleotides from on-target sites. We noticed, however, that there were several differences between our RGENs and those used by others. First, we used dualRNA, which is crRNA plus tracrRNA, rather than single-guide RNA (sgRNA) that is composed of essential portions of crRNA and tracrRNA. Second, we transfected K562 cells (but not HeLa cells) with synthetic crRNA rather than plasmids encoding crRNA. HeLa cells were transfected with crRNA-encoding plasmids. Other groups used sgRNA-encoding plasmids. Third, our guide RNA had two additional guanine nucleotides at the 5' end, which are required for efficient transcription by T7 polymerase *in vitro*. No such additional nucleotides were included in the sgRNA used by others. Thus, the RNA sequence of our guide RNA can be shown as 5'-GGX₂₀, whereas 5'-GX₁₉, in which X₂₀ or GX₁₉ corresponds to the 20-bp target sequence, represents the sequence used by others. The first guanine nucleotide is required for transcription by RNA polymerase in cells. To test whether off-target RGEN effects can be attributed to these differences, we chose four RGENs that induced off-target mutations in human cells at high frequencies(13). First, we compared our method of using *in vitro* transcribed dualRNA with the method of transfecting sgRNA-encoding plasmids in K562 cells and measured mutation frequencies at the on-target and off-target sites via the T7E1 assay. Three RGENs showed comparable mutation frequencies at on-target and off-target sites regardless of the composition of guide RNA. Interestingly, one RGEN (VEFGA site 1) did not induce indels at one validated off-target site, which differs by three nucleotides from the on-target site (termed OT1-11, Fig. 14), when synthetic dualRNA was used. But the synthetic dualRNA did not discriminate the other validated off-target site (OT1-3), which differs by two nucleotides from the on-target site.

[438] Next, we tested whether the addition of two guanine nucleotides at the 5' end of sgRNA could make RGENs more specific by comparing 5'-GGX₂₀ (or 5'-GGGX₁₉) sgRNA with 5'-GX₁₉ sgRNA. Four GX₁₉ sgRNAs complexed with Cas9 induced indels equally efficiently at on-target and off-target sites, tolerating up to four nucleotide mismatches. In sharp contrast, GGX₂₀ sgRNAs discriminated off-target sites effectively. In fact, the T7E1 assay barely detected RGEN-induced indels at six out of the seven validated off-target sites when we used the four GGX₂₀ sgRNAs (Fig. 15). We noticed, however, that two GGX₂₀ sgRNAs (VEGFA sites 1 and 3) were less active at on-target sites than were the corresponding GX₁₉ sgRNAs. These results show that the extra nucleotides at the 5' end can affect mutation frequencies at on-target and off-target sites, perhaps by altering guide RNA stability, concentration, or secondary

structure.

[439]

[440] These results suggest that three factors-the use of synthetic guide RNA rather than guide RNA-encoding plasmids, dualRNA rather than sgRNA, and GGX₂₀ sgRNA rather than GX₁₉ sgRNA-have cumulative effects on the discrimination of off-target sites.

[441]

[442] **Example 7: Paired Cas9 nickases**

[443]

[444] In principle, single-strand breaks (SSBs) cannot be repaired by error-prone NHEJ but still trigger high fidelity homology-directed repair (HDR) or base excision repair. But nickase-induced targeted mutagenesis via HDR is much less efficient than is nuclease-induced mutagenesis. We reasoned that paired Cas9 nickases would produce composite DSBs, which trigger DNA repair via NHEJ or HDR, leading to efficient mutagenesis (Fig. 16A). Furthermore, paired nickases would double the specificity of Cas9-based genome editing.

[445]

We first tested several Cas9 nucleases and nickases designed to target sites in the AAVS1 locus (Fig. 16B) *in vitro* via fluorescent capillary electrophoresis. Unlike Cas9 nucleases that cleaved both strands of DNA substrates, Cas9 nickases composed of guide RNA and a mutant form of Cas9 in which a catalytic aspartate residue is changed to an alanine (D10A Cas9) cleaved only one strand, producing site-specific nicks (Fig. 16C,D). Interestingly, however, some nickases (AS1, AS2, AS3, and S6 in Fig. 17A) induced indels at target sites in human cells, suggesting that nicks can be converted to DSBs, albeit inefficiently, *in vivo*. Paired Cas9 nickases producing two adjacent nicks on opposite DNA strands yielded indels at frequencies that ranged from 14% to 91%, comparable to the effects of paired nucleases (Fig. 17A). The repair of two nicks that would produce 5' overhangs led to the formation of indels much more frequently than those producing 3' overhangs at three genomic loci (Fig. 17A and Fig. 18). In addition, paired nickases enabled targeted genome editing via homology-directed repair more efficiently than did single nickases (Fig. 19).

[446]

[447] We next measured mutation frequencies of paired nickases and nucleases at off-target sites using deep sequencing. Cas9 nucleases complexed with three sgRNAs induced off-target mutations at six sites that differ by one or two nucleotides from their corresponding on-target sites with frequencies that ranged from 0.5% to 10% (Fig. 17B). In contrast, paired Cas9 nickases did not produce indels above the detection limit of 0.1% at any of the six off-target sites. The S2 Off-1 site that differs by a single nucleotide at the first position in the PAM (i.e., N in NGG) from its on-target site can be

considered as another on-target site. As expected, the Cas9 nuclease complexed with the S2 sgRNA was equally efficient at this site and the on-target site. In sharp contrast, D10A Cas9 complexed with the S2 and AS2 sgRNAs discriminated this site from the on-target site by a factor of 270 fold. This paired nickase also discriminated the AS2 off-target sites (Off-1 and Off-9 in Fig. 17B) from the on-target site by factors of 160 fold and 990 fold, respectively.

[448]

[449]

Example 8: Chromosomal DNA splicing induced by paired Cas9 nickases

[450]

[451]

Two concurrent DSBs produced by engineered nucleases such as ZFNs and TALENs can promote large deletions of the intervening chromosomal segments has reported. We tested whether two SSBs induced by paired Cas9 nickases can also produce deletions in human cells. We used PCR to detect deletion events and found that seven paired nickases induced deletions of up to 1.1-kbp chromosomal segments as efficiently as paired Cas9 nucleases did (Fig. 20A,B). DNA sequences of the PCR products confirmed the deletion events (Fig. 20C). Interestingly, the sgRNA-matching sequence remained intact in two out of seven deletion-specific PCR amplicons (underlined in Fig. 20C). In contrast, Cas9 nuclease pairs did not produce sequences that contained intact target sites. This finding suggests that two distant nicks were not converted to two separate DSBs to promote deletions of the intervening chromosomal segment. In addition, it is unlikely that two nicks separated by more than a 100 bp can produce a composite DSB with large overhangs under physiological conditions because the melting temperature is very high.

[452]

[453]

We propose that two distant nicks are repaired by strand displacement in a head-to-head direction, resulting in the formation of a DSB in the middle, whose repair via NHEJ causes small deletions (Fig. 20D). Because the two target sites remain intact during this process, nickases can induce SSBs again, triggering the cycle repeatedly until the target sites are deleted. This mechanism explains why two offset nicks producing 5' overhangs but not those producing 3' overhangs induced indels efficiently at three loci.

[454]

[455]

We then investigated whether Cas9 nucleases and nickases can induce unwanted chromosomal translocations that result from NHEJ repair of on-target and off-target DNA cleavages (Fig. 21A). We were able to detect translocations induced by Cas9 nucleases using PCR (Fig. 21B,C). No such PCR products were amplified using genomic DNA isolated from cells transfected with the plasmids encoding the AS2+S3 Cas9 nickase pair. This result is in line with the fact that both AS2 and S3 nickases,

unlike their corresponding nucleases, did not produce indels at off-target sites (Fig. 17B).

[456]

[457] These results suggest that paired Cas9 nickases allow targeted mutagenesis and large deletions of up to 1-kbp chromosomal segments in human cells. Importantly, paired nickases did not induce indels at off-target sites at which their corresponding nucleases induce mutations. Furthermore, unlike nucleases, paired nickases did not promote unwanted translocations associated with off-target DNA cleavages. In principle, paired nickases double the specificity of Cas9-mediated mutagenesis and will broaden the utility of RNA-guided enzymes in applications that require precise genome editing such as gene and cell therapy. One caveat to this approach is that two highly active sgRNAs are needed to make an efficient nickase pair, limiting targetable sites. As shown in this and other studies, not all sgRNAs are equally active. When single clones rather than populations of cells are used for further studies or applications, the choice of guide RNAs that represent unique sequences in the genome and the use of optimized guide RNAs would suffice to avoid off-target mutations associated with Cas9 nucleases. We propose that both Cas9 nucleases and paired nickases are powerful options that will facilitate precision genome editing in cells and organisms.

[458]

[459] **Example 9: Genotyping with CRISPR/Cas-derived RNA-guided endonucleases**

[460]

[461] Next, We reasoned that RGENs can be used in Restriction fragment length polymorphism (RFLP) analysis, replacing conventional restriction enzymes. Engineered nucleases including RGENs induce indels at target sites, when the DSBs caused by the nucleases are repaired by the error-prone non-homologous end-joining (NHEJ) system. RGENs that are designed to recognize the target sequences cannot cleave mutant sequences with indels but will cleave wildtype target sequences efficiently.

[462]

[463] **9-1. RGEN components**

[464]

[465] crRNA and tracrRNA were prepared by *in vitro* transcription using MEGAshortcript T7 kit (Ambion) according to the manufacturer's instruction. Transcribed RNAs were resolved on a 8% denaturing urea-PAGE gel. The gel slice containing RNA was cut out and transferred to elution buffer. RNA was recovered in nuclease-free water followed by phenol:chloroform extraction, chloroform extraction, and ethanol precipitation. Purified RNA was quantified by spectrometry. Templates for crRNA were prepared by annealing an oligonucleotide whose sequence is shown as 5'-GAAATTAATACGACTCACTATAGGX₂₀GTTTtagagctatgctgttttg-3' (

SEQ ID NO: 76), in which X_{20} is the target sequence, and its complementary oligonucleotide. The template for tracrRNA was synthesized by extension of forward and reverse oligonucleotides

(5'-GAAATTAATACGACTCACTATAGGAACCATTCAAAACAGCATAGCAAGTTAAAATAAGGCTAGTCCG-3' (SEQ ID NO: 77) and

5'-AAAAAAAGCACCGACTCGGTGCCACTTTTTCAAGTTGATAACGGACTAGCCTTATTTTAACTTGCTATG-3' (SEQ ID NO: 78)) using Phusion polymerase (New England Biolabs).

[466]

[467] **9-2. Recombinant Cas9 protein purification**

[468]

[469] The Cas9 DNA construct used in our previous Example, which encodes Cas9 fused to the His6-tag at the C terminus, was inserted in the pET-28a expression vector. The recombinant Cas9 protein was expressed in *E. coli* strain BL21(DE3) cultured in LB medium at 25°C for 4 hour after induction with 1 mM IPTG. Cells were harvested and resuspended in buffer containing 20 mM Tris PH 8.0, 500 mM NaCl, 5 mM imidazole, and 1 mM PMSF. Cells were frozen in liquid nitrogen, thawed at 4°C, and sonicated. After centrifugation, the Cas9 protein in the lysate was bound to Ni-NTA agarose resin (Qiagen), washed with buffer containing 20 mM Tris pH 8.0, 500 mM NaCl, and 20 mM imidazole, and eluted with buffer containing 20 mM Tris pH 8.0, 500 mM NaCl, and 250 mM imidazole. Purified Cas9 protein was dialyzed against 20 mM HEPES (pH 7.5), 150 mM KCl, 1 mM DTT, and 10% glycerol and analyzed by SDS-PAGE.

[470]

[471] **9-3. T7 endonuclease I assay**

[472]

[473] The T7E1 assay was performed as following. In brief, PCR products amplified using genomic DNA were denatured at 95°C, reannealed at 16°C, and incubated with 5 units of T7 Endonuclease I (New England BioLabs) for 20 min at 37°C. The reaction products were resolved using 2 to 2.5% agarose gel electrophoresis.

[474]

[475] **9-4. RGEN-RFLP assay**

[476]

[477] PCR products (100-150 ng) were incubated for 60 min at 37°C with optimized concentrations (Table 10) of Cas9 protein, tracrRNA, crRNA in 10 µl NEB buffer 3 (1X). After the cleavage reaction, RNase A (4 µg) was added, and the reaction mixture was incubated for 30 min at 37°C to remove RNA. Reactions were stopped with 6X stop solution buffer containing 30% glycerol, 1.2% SDS, and 100 mM EDTA. Products

were resolved with 1-2.5% agarose gel electrophoresis and visualized with EtBr staining.

[478]

[479] Table 10

[Table 10]

Concentration of RGEN components in RFLP assays

Target Name	Cas9 (ng/μl)	crRNA (ng/μl)	tracrRNA (ng/μl)
<i>C4BPB</i>	100	25	60
<i>PIBF</i> -NGG-RGEN	100	25	60
<i>HLA-B</i>	1.2	0.3	0.7
<i>CCR5</i> -ZFN	100	25	60
<i>CTNNB1</i> Wild type specific	30	10	20
<i>CTNNB1</i> mutant specific	30	10	20
<i>CCR5</i> WT-specific	100	25	60
<i>CCR5</i> Δ32-specific	10	2.5	6
<i>KRAS</i> WT specific(wt)	30	10	20
<i>KRAS</i> mutant specific(m8)	30	10	20
<i>KRAS</i> WT specific (m6)	30	10	20
<i>KRAS</i> mutant specific (m6,8)	30	10	20
<i>PIK3CA</i> WT specific (wt)	100	25	60
<i>PIK3CA</i> mutant specific(m4)	30	10	20
<i>PIK3CA</i> WT specific (m7)	100	25	60
<i>PIK3CA</i> mutant specific(m4,7)	30	10	20
<i>BRAF</i> WT-specific	30	10	20
<i>BRAF</i> mutant-specific	100	25	60
<i>NRAS</i> WT-specific	100	25	60
<i>NRAS</i> mutant-specific	30	10	20
<i>IDH</i> WT-specific	30	10	20
<i>IDH</i> mutant-specific	30	10	20
<i>PIBF</i> -NAG-RGEN	30	10	60

[480]

[481] Table 11

[Table 11]

Primers

Gene(site)	Direction	Sequence(5' to 3')	SEQ ID NO
CCR5(RGEN)	F1	CTCCATGGTGCTATAGAGCA	79
	F2	GAGCCAAGCTCTCCATCTAGT	80
	R	GCCCTGTCAAGAGTTGACAC	81
CCR5(ZFN)	F	GCACAGGGTGAACAAGATGGA	82
	R	GCCAGGTACCTATCGATTGTCAGG	83
CCR5(del32)	F	GAGCCAAGCTCTCCATCTAGT	84
	R	ACTCTGACTG GGTCACCAGC	85
C4BPB	F1	TATTTGGCTGGTTGAAAGGG	86
	R1	AAAGTCATGAAATAAACACACCCA	87
	F2	CTGCATTGATATGGTAGTACCATG	88
	R2	GCTGTTCAATTGCAATGGAATG	89
CTNNB1	F	ATGGAGTTGGACATGGCCATGG	90
	R	ACTCACTATCCACAGTTCAGCATTTAC C	91
KRAS	F	TGGAGATAGCTGTCAGCAACTTT	92
	R	CAACAA AGCAAAG- GTAAAGTTGGTAATAG	93
PIK3CA	F	GGTTTCAGGAGATGTGTTACAAGGC	94
	R	GATTGTGCAATTCCTATGCAATCGGTC	95
NRAS	F	CACTGGGTACTTAATCTGTAGCCTC	96
	R	GGTTCCAAGTCATTCCCAGTAGC	97
IDH1	F	CATCACTGCAGTTGTAGGTTATAACTATC C	98
	R	TTGAAAACCACAGATCTGGTTGAACC	99
BRAF	F	GGAGTGCCAAGAGAATATCTGG	100
	R	CTGAAACTGGTTTCAAAATATTCGTTTTA AGG	101

<i>PIBF</i>	<i>F</i>	<i>GCTCTGTATGCCCTGTAGTAGG</i>	<i>102</i>
	<i>R</i>	<i>TTTGCATCTGACCTTACCTTTG</i>	<i>103</i>

[482]

[483] **9-5. Plasmid cleavage assay**

[484]

[485] Restriction enzyme-treated linearized plasmid (100 ng) was incubated for 60 min at 37 °C with Cas9 protein (0.1 µg), tracrRNA (60 ng), and crRNA (25 ng) in 10 µl NEB 3 buffer (1X). Reactions were stopped with 6X stop solution containing 30% glycerol, 1.2% SDS, and 100 mM EDTA. Products were resolved with 1% agarose gel electrophoresis and visualized with EtBr staining.

[486]

[487] **9-6. Strategy of RFLP**

[488]

[489] New RGENs with desired DNA specificities can be readily created by replacing crRNA; no de novo purification of custom proteins is required once recombinant Cas9 protein is available. Engineered nucleases, including RGENs, induce small insertions or deletions (indels) at target sites when the DSBs caused by the nucleases are repaired by error-prone non-homologous end-joining (NHEJ). RGENs that are designed to recognize the target sequences cleave wild-type sequences efficiently but cannot cleave mutant sequences with indels (Fig. 22).

[490] We first tested whether RGENs can differentially cleave plasmids that contain wild-type or modified C4BPB target sequences that harbor 1- to 3-base indels at the cleavage site. None of the six plasmids with these indels were cleaved by a C4BPB-specific RGEN5 composed of target-specific crRNA, tracrRNA, and recombinant Cas9 protein (Fig. 23). In contrast, the plasmid with the intact target sequence was cleaved efficiently by this RGEN.

[491]

[492] **9-7. Detection of mutations induced by the same RGENs using RGEN-mediated RFLP**

[493]

[494] Next, to test the feasibility of RGEN-mediated RFLP for detection of mutations induced by the same RGENs, we utilized gene-modified K562 human cancer cell clones established using an RGEN targeting C4BPB gene (Table 12).

[495]

[496] Table 12

[Table 12]

Target sequence of RGENs used in this study

Gene	Target sequence	SEQ ID NO
human C4BPB	<u>AATGACCACTACATCCTCAA</u> GGG	104
mouse Pibf1	<u>AGATGATGTCTCATCATCAG</u> AGG	105

[497]

[498] C4BPB mutant clones used in this study have various mutations ranging from 94 bp deletion to 67 bp insertion (Fig. 24A). Importantly, all mutations occurred in mutant clones resulted in the loss of RGEN target site. Among 6 C4BPB clones analyzed, 4 clones have both wildtype and mutant alleles (+/-) and 2 clones have only mutant alleles (-/-).

[499]

[500] The PCR products spanning the RGEN target site amplified from wildtype K562 genomic DNA were digested completely by the RGEN composed of target-specific crRNA, tracrRNA, and recombinant Cas9 protein expressed in and purified from *E. coli* (Fig. 24B/Lane 1). When the C4BPB mutant clones were subjected to RFLP analysis using the RGEN, PCR amplicons of +/- clones that contained both wildtype and mutant alleles were partially digested, and those of -/- clones that did not contain the wildtype allele were not digested at all, yielding no cleavage products corresponding to the wildtype sequence (Fig. 24B). Even a single-base insertion at the target site blocked the digestion (#12 and #28 clones) of amplified mutant alleles by the C4BPB RGEN, showing the high specificity of RGEN-mediated RFLP. We subjected the PCR amplicons to the mismatch-sensitive T7E1 assay in parallel (Fig. 24B). Notably, the T7E1 assay was not able to distinguish -/- clones from +/- clones. To make it matters worse, the T7E1 assay cannot distinguish homozygous mutant clones that contain the same mutant sequence from wildtype clones, because annealing of the same mutant sequence will form a homoduplex. Thus, RGEN-mediated RFLP has a critical advantage over the conventional mismatch-sensitive nuclease assay in the analysis of mutant clones induced by engineered nucleases including ZFNs, TALENs and RGENs.

[501]

[502] 9-8. Quantitative assay for RGEN-RFLP analysis

[503]

[504] We also investigated whether RGEN-RFLP analysis is a quantitative method.

Genomic DNA samples isolated from the C4BPB null clone and the wild-type cells were mixed at various ratios and used for PCR amplifications. The PCR products were subjected to RGEN genotyping and the T7E1 assay in parallel (Fig. 25b). As expected, DNA cleavage by the RGEN was proportional to the wild type to mutant ratio. In contrast, results of the T7E1 assay correlated poorly with mutation frequencies inferred from the ratios and were inaccurate, especially at high mutant %, a situation in which complementary mutant sequences can hybridize with each other to form homoduplexes.

[505]

[506] **9-9. Analysis of mutant mouse founders using a RGEN-mediated RFLP genotyping**

[507]

[508] We also applied RGEN-mediated RFLP genotyping (RGEN genotyping in short) to the analysis of mutant mouse founders that had been established by injection of TALENs into mouse one-cell embryos (Fig. 26A). We designed and used an RGEN that recognized the TALEN target site in the Pibf1 gene (Table 10). Genomic DNA was isolated from a wildtype mouse and mutant mice and subjected to RGEN genotyping after PCR amplification. RGEN genotyping successfully detected various mutations, which ranged from one to 27-bp deletions (Fig. 26B). Unlike the T7E1 assay, RGEN genotyping enabled differential detection of +/- and -/- founder.

[509]

[510] **9-10. Detection of mutations induced in human cells by a CCR5-specific ZFN using RGENs**

[511]

[512] In addition, we used RGENs to detect mutations induced in human cells by a CCR5-specific ZFN, representing yet another class of engineered nucleases (Fig. 27). These results show that RGENs can detect mutations induced by nucleases other than RGENs themselves. In fact, we expect that RGENs can be designed to detect mutations induced by most, if not all, engineered nucleases. The only limitation in the design of an RGEN genotyping assay is the requirement for the GG or AG (CC or CT on the complementary strand) dinucleotide in the PAM sequence recognized by the Cas9 protein, which occurs once per 4 bp on average. Indels induced anywhere within the seed region of several bases in crRNA and the PAM nucleotides are expected to disrupt RGEN-catalyzed DNA cleavage. Indeed, we identified at least one RGEN site in most (98%) of the ZFN and TALEN sites.

[513]

[514] **9-11. Detection of polymorphisms or variations using RGEN**

[515]

[516] Next, we designed and tested a new RGEN that targets a highly polymorphic locus, HLA-B, that encodes Human Leukocyte Antigen B (a.k.a. MHC class I protein) (Fig. 28). HeLa cells were transfected with RGEN plasmids, and the genomic DNA was subjected to T7E1 and RGEN-RFLP analyses in parallel. T7E1 produced false positive bands that resulted from sequence polymorphisms near the target site (Fig. 25c). As expected, however, the same RGEN used for gene disruption cleaved PCR products from wild-type cells completely but those from RGEN-transfected cells partially, indicating the presence of RGEN-induced indels at the target site. This result shows that RGEN-RFLP analysis has a clear advantage over the T7E1 assay, especially when it is not known whether target genes have polymorphisms or variations in cells of interest.

[517]

[518] **9-12. Detection of recurrent mutations found in cancer and naturally-occurring polymorphisms through RGEN-RFLP analysis**

[519]

[520] RGEN-RFLP analysis has applications beyond genotyping of engineered nuclease-induced mutations. We sought to use RGEN genotyping to detect recurrent mutations found in cancer and naturally-occurring polymorphisms. We chose the human colorectal cancer cell line, HCT116, which carries a gain-of-function 3-bp deletion in the oncogenic CTNNB1 gene encoding beta-catenin. PCR products amplified from HCT116 genomic DNA were cleaved partially by both wild-type-specific and mutant-specific RGENs, in line with the heterozygous genotype in HCT116 cells (Fig. 29a). In sharp contrast, PCR products amplified from DNA from HeLa cells harboring only wild-type alleles were digested completely by the wild-type-specific RGEN and were not cleaved at all by the mutation-specific RGEN.

[521]

[522] We also noted that HEK293 cells harbor the 32-bp deletion (del32) in the CCR5 gene, which encodes an essential co-receptor of HIV infection: Homozygous del32 CCR5 carriers are immune to HIV infection. We designed one RGEN specific to the del32 allele and the other to the wild-type allele. As expected, the wild-type-specific RGEN cleaved the PCR products obtained from K562, SKBR3, or HeLa cells (used as wild-type controls) completely but those from HEK293 cells partially (Fig. 30a), confirming the presence of the uncleavable del32 allele in HEK293 cells. Unexpectedly, however, the del32-specific RGEN cleaved the PCR products from wild-type cells as efficiently as those from HEK293 cells. Interestingly, this RGEN had an off-target site with a single-base mismatch immediately downstream of the on-target site (Fig. 30). These results suggest that RGENs can be used to detect naturally-occurring indels but cannot distinguish sequences with single nucleotide polymorphisms or point mutations due to their off-target effects.

[523]

[524] To genotype oncogenic single-nucleotide variations using RGENs, we attenuated RGEN activity by employing a single-base mismatched guide RNA instead of a perfectly-matched RNA. RGENs that contained the perfectly-matched guide RNA specific to the wild-type sequence or mutant sequence cleaved both sequences (Figs. 31a and 32a). In contrast, RGENs that contained a single-base mismatched guide RNA distinguished the two sequences, enabling genotyping of three recurrent oncogenic point mutations in the KRAS, PIK3CA, and IDH1 genes in human cancer cell lines (Fig. 29b and Figs. 33a, b). In addition, we were able to detect point mutations in the BRAF and NRAS genes using RGENs that recognize the NAG PAM sequence (Figs. 33c, d). We believe that we can use RGEN-RFLP to genotype almost any, if not all, mutations or polymorphisms in the human and other genomes.

[525]

[526] The above data proposes RGENs as providing a platform to use simple and robust RFLP analysis for various sequence variations. With high flexibility in reprogramming target sequence, RGENs can be used to detect various genetic variations (single nucleotide variations, small insertion/deletions, structural variations) such as disease-related recurring mutations, genotypes related to drug-response by a patient and also mutations induced by engineered nucleases in cells. Here, we used RGEN genotyping to detect mutations induced by engineered nucleases in cells and animals. In principle, one could also use RGENs that will specifically detect and cleave naturally-occurring variations and mutations.

[527]

[528] Based on the above description, it should be understood by those skilled in the art that various alternatives to the embodiments of the invention described herein may be employed in practicing the invention without departing from the technical idea or essential features of the invention as defined in the following claims. In this regard, the above-described examples are for illustrative purposes only, and the invention is not intended to be limited by these examples. The scope of the present invention should be understood to include all of the modifications or modified form derived from the meaning and scope of the following claims or its equivalent concepts.

[529]

[530] [References]

[531] 1. M. Jinek et al., Science 337, 816 (Aug 17, 2012).

[532] 2. H. Kim, E. Um, S. R. Cho, C. Jung, J. S. Kim, Nat Methods 8, 941 (Nov, 2011).

[533] 3. H. J. Kim, H. J. Lee, H. Kim, S. W. Cho, J. S. Kim, Genome Res 19, 1279 (Jul, 2009).

[534] 4. E. E. Perez et al., Nat Biotechnol 26, 808 (Jul, 2008).

- [535] 5. J. C. Miller et al., Nat Biotechnol 29, 143 (Feb, 2011).
- [536] 6. C. Mussolino et al., Nucleic Acids Res 39, 9283 (Nov, 2011).
- [537] 7. J. Cohen, Science 332, 784 (May 13, 2011).
- [538] 8. V. Pattanayak, C. L. Ramirez, J. K. Joung, D. R. Liu, Nat Methods 8, 765 (Sep, 2011).
- [539] 9. R. Gabriel et al., Nat Biotechnol 29, 816 (Sep, 2011).
- [540] 10. E. Kim et al., Genome Res, (Apr 20, 2012).
- [541] 11. H. J. Lee, J. Kweon, E. Kim, S. Kim, J. S. Kim, Genome Res 22, 539 (Mar, 2012).
- [542] 12. H. J. Lee, E. Kim, J. S. Kim, Genome Res 20, 81 (Jan, 2010).
- [543] 13. Fu Y, Foden JA, Khayter C, Maeder ML, Reyon D, Joung JK, Sander JD. High-frequency off-target mutagenesis induced by CRISPR-Cas nucleases in human cells. Nat Biotech advance online publication (2013)

Claims

1. A composition comprising a Type II Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR)/Cas system for cleavage of a target nucleic acid in a eukaryotic cell, said CRISPR/Cas system comprising (i) a nucleic acid encoding a Cas9 polypeptide and a nuclear localization sequence, and (ii) a nucleic acid encoding a guide RNA that hybridizes to the target nucleic acid in a eukaryotic cell, wherein the guide RNA is a chimeric guide RNA comprising a CRISPR RNA (crRNA) sequence fused to a trans activating crRNA (tracrRNA) sequence, wherein the crRNA sequence is at the 5' end relative to the tracrRNA sequence.
2. The composition of claim 1, wherein said Cas9 polypeptide is a *Streptococcus* Cas9 polypeptide.
3. The composition of claim 2, wherein said Cas9 polypeptide is a *Streptococcus pyogenes* Cas9 polypeptide.
4. The composition of any one of claims 1-3, wherein said nucleic acid encoding a Cas9 polypeptide is codon-optimized for expression in eukaryotic cells.
5. The composition of claim 4, wherein said nucleic acid encoding a Cas9 polypeptide is codon-optimized for expression in mammalian cells.
6. The composition of any one of claims 1-5, wherein said nuclear localization sequence is located at the C terminus or N terminus of the Cas9 polypeptide.
7. The composition of any one of claims 1-6, wherein the target nucleic acid is an endogenous target nucleic acid.
8. The composition of any one of claims 1-7, wherein the guide RNA is in the form of a vector.
9. The composition of any one of claims 1-8, wherein the guide RNA comprises one or more additional nucleic acid residues 5' to the crRNA sequence.
10. The composition of claim 9, wherein the one or more additional nucleic residues are two

nucleic acid residues, and wherein there are no additional nucleic acid residues between the two nucleic acid residues and the crRNA sequence.

11. The composition of claim 10, wherein the two nucleic acid residues at the 5' end are both guanine residues.
12. A composition comprising a Type II Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR)/Cas system for cleavage of a target nucleic acid in a eukaryotic cell, said CRISPR/Cas system comprising (i) a nucleic acid encoding a Cas9 polypeptide and a nuclear localization sequence, wherein said nucleic acid comprises SEQ ID NO:1, and (ii) a nucleic acid encoding a guide RNA that hybridizes to the target nucleic acid in a eukaryotic cell, wherein the guide RNA is a chimeric guide RNA comprising a CRISPR RNA (crRNA) sequence fused to a trans activating crRNA (tracrRNA) sequence, wherein the crRNA sequence is at the 5' end relative to the tracrRNA sequence.
13. The composition of claim 12, wherein the target nucleic acid sequence comprises a sequence having a trinucleotide protospacer adjacent motif (PAM), wherein the PAM is a trinucleotide of 5'-NGG-3' selected from the group consisting of 5'-AGG-3', 5'-GGG-3', 5'-TGG-3' and 5'-CGG-3'.
14. The composition of claim 12 or 13, wherein said nucleic acid encoding a Cas9 polypeptide is codon-optimized for expression in eukaryotic cells.
15. The composition of claim 14, wherein said nucleic acid encoding a Cas9 polypeptide is codon-optimized for expression in mammalian cells.
16. The composition of any one of claims 13-15, wherein said nuclear localization sequence is located at the C terminus or N terminus of the Cas9 polypeptide.
17. The composition of any one of claims 12-16, wherein the target nucleic acid is an endogenous target nucleic acid.
18. The composition of any one of claims 12-17, wherein the guide RNA is in the form of a vector.

19. The composition of any one of claims 12-18, wherein the guide RNA comprises one or more additional nucleic acid residues 5' to the crRNA sequence.
20. The composition of claim 19, wherein the one or more additional nucleic acid residues are two nucleic acid residues, and wherein there are no additional nucleic acid residues between the two nucleic acid residues and the crRNA sequence.
21. The composition of claim 20, wherein the two nucleic acid residues at the 5' end are both guanine residues.
22. An *in vitro* method of introducing a site-specific, double-stranded break at a target nucleic acid sequence in a eukaryotic cell, the method comprising introducing into the eukaryotic cell a Type II Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR)/Cas system, wherein the CRISPR/Cas system comprises:
 - a) a nucleic acid encoding a Cas9 polypeptide and a nuclear localization signal, and
 - b) a nucleic acid encoding a guide RNA that hybridizes to the target nucleic acid, wherein the guide RNA is a chimeric guide RNA comprising a CRISPR RNA (crRNA) sequence fused to a trans activating crRNA (tracrRNA) sequence, wherein the crRNA sequence is at the 5' end relative to the tracrRNA sequence; wherein the target nucleic acid sequence comprises a sequence that binds to the crRNA sequence,and wherein the Cas9 polypeptide and the guide RNA form a Cas9/RNA complex in the eukaryotic cell, whereby a site-specific, double stranded break at the target nucleic acid sequence is introduced.
23. The method of claim 22, wherein the Cas9 polypeptide is a *Streptococcus* Cas9 polypeptide.
24. The method of claim 23, wherein the Cas9 polypeptide is a *Streptococcus pyogenes* Cas9 polypeptide.
25. The method of claim 22 or 23, wherein said nucleic acid encoding a Cas9 polypeptide is

codon-optimized for expression in eukaryotic cells.

26. The method of claim 25, wherein the nucleic acid encoding the Cas 9 polypeptide is codon-optimized for expression in mammalian cells.
27. The method of any one of claims 22-26, wherein the nuclear localization signal is located at the C terminus or N terminus of the Cas9 polypeptide.
28. The method of any one of claims 22-27, wherein the eukaryotic cell is a mammalian cell.
29. The method of claim 28, wherein the mammalian cell is a human cell.
30. The method of any one of claims 22-29, wherein the target nucleic acid sequence is a genomic sequence located at its endogenous site in the genome of the eukaryotic cell.
31. The method of any one of claims 22-30, wherein the nucleic acid encoding the guide RNA is a vector.
32. The method of any one of claims 22-30, wherein the nucleic acid encoding the guide RNA is in vitro transcribed RNA.
33. The method of any one of claims 22-30, wherein the guide RNA comprises one or more additional nucleic acid residues 5' to the crRNA sequence.
34. The method of claim 33, wherein the one or more additional nucleic acid residues are two nucleic acid residues, and wherein there are no additional nucleic acid residues between the two nucleic acid residues and the crRNA sequence.
35. The method of claim 34, wherein the two nucleic acid residues at the 5' end are both guanine residues.
36. The method of any one of claims 22-35, wherein the nucleic acid encoding the Cas9 polypeptide is introduced into the eukaryotic cell before introducing the nucleic acid encoding the guide RNA into the eukaryotic cell.
37. The method of any one of claims 22-35, wherein the nucleic acid encoding the Cas9 polypeptide is co-transfected into the eukaryotic cell with the nucleic acid encoding the

guide RNA.

38. The method of any one of claims 22-37, wherein the eukaryotic cell is a non-human eukaryotic cell.
39. The composition of any one of claims 1-11 or the method of any one of claims 22-38, wherein the target nucleic acid sequence comprises a sequence having a trinucleotide protospacer adjacent motif (PAM), wherein the PAM is a trinucleotide of 5'-NGG-3' selected from the group consisting of 5'-AGG-3', 5'-GGG-3', 5'-TGG-3' and 5'-CGG-3'.
40. The composition of claim 6, wherein said nuclear localization sequence is located at the C terminus of the Cas9 polypeptide.
41. The composition of claim 16, wherein said nuclear localization sequence is located at the C terminus of the Cas9 polypeptide.
42. The method of claim 27, wherein the nuclear localization signal is located at the C terminus of the Cas9 polypeptide.