



EP 0 644 202 A1

The implementation of systematic testing for hepatitis B virus (HBV) has been instrumental in eliminating this virus from the blood supply. Nevertheless, a significant number of post-transfusion hepatitis (PTH) cases still occur. These cases are generally attributable to non-A, non-B hepatitis (NANBH) virus(es), the diagnosis of which is usually made by exclusion of other viral markers.

The etiological agent responsible for a large proportion of these cases has recently been cloned (Choo, Q-L et al. *Science* (1988) 244:359-362) and a first-generation antibody test developed (Kuo, G. et al. *Science* (1989) 244:362-364). The agent has been identified as a positive-stranded RNA virus, and the sequence of its genome has been partially determined. Studies suggest that this virus, referred to subsequently as hepatitis C virus (HCV), may be related to flaviviruses and pestiviruses. A portion of the genome of an HCV isolated from a chimpanzee (HCV_{CDC/CHI}) is disclosed in EPO 88310922.5. The coding sequences disclosed in this document do not include sequences originating from the 5'-end of the viral genome which code for putative structural proteins. Recently however, sequences derived from this region of the HCV genome have been published (Okamoto, H. et al., *Japan J. Exp. Med.* 60:167-177, 1990.). The amino acid sequences encoded by the Japanese clone HC-J1 were combined with the HCV_{CDC/CHI} sequences in a region where the two sequences overlap to generate the composite sequence depicted in Figure 1. Specifically, the two sequences were joined at glycine₄₅₁. It should be emphasized that the numbering system used for the HCV amino acid sequence is not intended to be absolute since the existence of variant HCV strains harboring deletions or insertions is highly probable. Sequences corresponding to the 5' end of the HCV genome have also recently been disclosed in EPO 90302866.0.

In order to detect potential carriers of HCV, it is necessary to have access to large amounts of viral proteins. In the case of HCV, there is currently no known method for culturing the virus, which precludes the use of virus-infected cultures as a source of viral antigens. The current first-generation antibody test makes use of a fusion protein containing a sequence of 363 amino acids encoded by the HCV genome. It was found that antibodies to this protein could be detected in 75 to 85% of chronic NANBH patients. In contrast, only approximately 15% of those patients who were in the acute phase of the disease, had antibodies which recognized this fusion protein (Kuo, G. et al. *Science* (1989) 244:362-364). The absence of suitable confirmatory tests, however, makes it difficult to verify these statistics. The seeming similarity between the HCV genome and that of flaviviruses makes it possible to predict the location of epitopes which are likely to be of diagnostic value. An analysis of the HCV genome reveals the presence of a continuous long open reading frame. Viral RNA is presumably translated into a long polyprotein which is subsequently cleaved by cellular and/or viral proteases. By analogy with, for example, Dengue virus, the viral structural proteins are presumed to be derived from the amino-terminal third of the viral polyprotein. At the present time, the precise sites at which the polyprotein is cleaved can only be surmised. Nevertheless, the structural proteins are likely to contain epitopes which would be useful for diagnostic purposes, both for the detection of antibodies as well as for raising antibodies which could subsequently be used for the detection of viral antigens. Similarly, domains of nonstructural proteins are also expected to contain epitopes of diagnostic value, even though these proteins are not found as structural components of virus particles.

Brief Description of the Drawings

Figure 1 shows the amino acid sequence of the composite HCV_{HC-J1/CDC/CHI}

Figure 2 shows the antibody binding to individual peptides and various mixtures in an ELISA assay

Description of the Specific Embodiments

It is known that RNA viruses frequently exhibit a high rate of spontaneous mutation and, as such, it is to be expected that no two HCV isolates will be completely identical, even when derived from the same individual. For the purpose of this disclosure, a virus is considered to be the same or equivalent to HCV if it exhibits a global homology of 60 percent or more with the HCV_{HC-J1/CDC/CHI} composite sequence at the nucleic acid level and 70 percent at the amino acid level.

Peptides are described which immunologically mimic proteins encoded by HCV. In order to accommodate strain-to-strain variations in sequence, conservative as well as non-conservative amino acid substitutions may be made. These will generally account for less than 35 percent of a specific sequence. It may be desirable in cases where a peptide corresponds to a region in the HCV polypeptide which is highly polymorphic, to vary one or more of the amino acids so as to better mimic the different epitopes of different viral strains.

The peptides of interest will include at least five, sometimes six, sometimes eight, sometimes twelve, usually fewer than about fifty, more usually fewer than about thirty-five, and preferably fewer than about

twenty-five amino acids included within the sequence encoded by the HCV genome. In each instance, the peptide will preferably be as small as possible while still maintaining substantially all of the sensitivity of the larger peptide. It may also be desirable in certain instances to join two or more peptides together in one peptide structure.

5 It should be understood that the peptides described need not be identical to any particular HCV sequence, so long as the subject compounds are capable of providing for immunological competition with at least one strain of HCV. The peptides may therefore be subject to insertions, deletions, and conservative or non-conservative amino acid substitutions where such changes might provide for certain advantages in their use.

10 Substitutions which are considered conservative are those in which the chemical nature of the substitute is similar to that of the original amino acid. Combinations of amino acids which could be considered conservative are Gly, Ala; Asp, Glu; Asn, Gln; Val, Ile, Leu; Ser, Thr; Lys, Arg; and Phe, Tyr.

Furthermore, additional amino acids or chemical groups may be added to the amino- or carboxyl terminus for the purpose of creating a "linker arm" by which the peptide can conveniently be attached to a carrier. The linker arm will be at least one amino acid and may be as many as 60 amino acids but will most frequently be 1 to 10 amino acids. The nature of the attachment to a solid phase or carrier need not be covalent.

Natural amino acids such as cysteine, lysine, tyrosine, glutamic acid, or aspartic acid may be added to either the amino- or carboxyl terminus to provide functional groups for coupling to a solid phase or a carrier. However, other chemical groups such as, for example, biotin and thioglycolic acid, may be added to the termini which will endow the peptides with desired chemical or physical properties. The termini of the peptides may also be modified, for example, by N-terminal acetylation or terminal carboxy-amidation. The peptides of interest are described in relation to the composite amino acid sequence shown in Figure 1. The amino acid sequences are given in the conventional and universally accepted three-letter code. In addition to the amino acids shown, other groups are defined as follows: Y is, for example, NH₂, one or more N-terminal amino acids, or other moieties added to facilitate coupling. Y may itself be modified by, for example, acetylation. Z is a bond, (an) amino acid(s), or (a) chemical group(s) which may be used for linking. X is intended to represent OH, NH₂, or a linkage involving either of these two groups.

Peptide I corresponds to amino acids 1 to 20 and has the following amino acid sequences:

30

(I)

Y-Met-Ser-Thr-Ile-Pro-Lys-Pro-Gln-Arg-Lys-Thr-Lys-Arg-Asn-Thr-Asn-Arg-Arg-Pro-
35 Gln-Z-X. (SEQ ID NO 1)

Peptide II corresponds to amino acids 7 to 26 and has the amino acid sequence:

40

(II)

Y-Pro-Gln-Arg-Lys-Thr-Lys-Arg-Asn-Thr-Asn-Arg-Arg-Pro-Gln-Asp-Val-Lys-Phe-
45 Pro-Gly-Z-X. (SEQ ID NO 2)

Of particular interest is the oligopeptide IIA:

(IIA)

Y-Gln-Arg-Lys-Thr-Lys-Arg-Asn-Thr-Asn-Arg-Arg-Z-X. (SEQ ID NO 3)

50 Peptide III corresponds to amino acids 13 to 32 and has the sequence:

(III)

Y-Arg-Asn-Thr-Asn-Arg-Arg-Pro-Gln-Asp-Val-Lys-Phe-Pro-Gly-Gly-Gly-Gln-Ile-Val-
55 Gly-Z-X. (SEQ ID NO 4)

This Sheet Has Been Replaced



Peptide IV corresponds to amino acid 37 to 56 and has the sequences:

(IV)

5 Y-Leu-Pro-Arg-Arg-Gly-Pro-Arg-Leu-Gly-Val-Arg-Ala-Thr-Arg-Lys-Thr-Ser-Glu-Arg-Ser-Z-X. (SEQ ID NO 5)

10 Peptide V corresponds to amino acids 49 to 68 and has the sequence:

(V)

15 Y-Thr-Arg-Lys-Thr-Ser-Glu-Arg-Ser-Gln-Pro-Arg-Gly-Arg-Arg-Gln-Pro-Ile-Pro-Lys-Val-Z-X. (SEQ ID NO 6)

20 Peptide VI corresponds to amino acid 61 to 80 and has the following sequence:

(VI)

25 Y-Arg-Arg-Gln-Pro-Ile-Pro-Lys-Val-Arg-Arg-Pro-Glu-Gly-Arg-Thr-Trp-Ala-Gln-Pro-Gly-Z-X. (SEQ ID NO 7)

Peptide VII corresponds to amino acids 73 to 92 and has the sequence:

(VII)

30 Y-Gly-Arg-Thr-Trp-Ala-Gln-Pro-Gly-Tyr-Pro-Trp-Pro-Leu-Tyr-Gly-Asn-Glu-Gly-Cys-Gly-Z-X. (SEQ ID NO 8)

35 Peptide VIII corresponds to amino acids 1688 to 1707 and has the sequence:

(VIII)

40 Y-Leu-Ser-Gly-Lys-Pro-Ala-Ile-Ile-Pro-Asp-Arg-Glu-Val-Leu-Tyr-Arg-Glu-Phe-Asp-Glu-Z-X. (SEQ ID NO 9)

45 Peptide IX corresponds to amino acids 1694 to 1713 and has the sequence:

(IX)

50 Y-Ile-Ile-Pro-Asp-Arg-Glu-Val-Leu-Tyr-Arg-Glu-Phe-Asp-Glu-Met-Glu-Glu-Cys-Ser-Gln-Z-X. (SEQ ID NO 10)

55 Peptide X corresponds to amino acids 1706 to 1725 and has the sequence:

This Sheet Has Been Replaced



(X)

Y-Asp-Glu-Met-Glu-Glu-Cys-Ser-Gln-His-Leu-Pro-Tyr-Ile-Glu-Gln-Gly-Met-Met-Leu-
Ala-Z-X. (SEQ ID NO 11)

Peptide XI corresponds to amino acids 1712 to 1731 and has the sequence:

(XI)

Y-Ser-Gln-His-Leu-Pro-Tyr-Ile-Glu-Gln-Gly-Met-Met-Leu-Ala-Glu-Gln-Phe-Lys-Gln-
Lys-Z-X. (SEQ ID NO 12)

Peptide XII corresponds to amino acids 1718 to 1737 and has the sequence:

(XII)

Y-Ile-Glu-Gln-Gly-Met-Met-Leu-Ala-Glu-Gln-Phe-Lys-Gln-Lys-Ala-Leu-Gly-Leu-Leu-
Gln-Z-X. (SEQ ID NO 13)

Peptide XIII corresponds to amino acids 1724 to 1743 and has the sequence:

(XIII)

Y-Leu-Ala-Glu-Gln-Phe-Lys-Gln-Lys-Ala-Leu-Gly-Leu-Leu-Gln-Thr-Ala-Ser-Arg-Gln-
Ala-Z-X. (SEQ ID NO 14)

Peptide XIV corresponds to amino acids 1730 to 1749 and has the sequence:

(XIV)

Y-Gln-Lys-Ala-Leu-Gly-Leu-Leu-Gln-Thr-Ala-Ser-Arg-Gln-Ala-Glu-Val-Ile-Ala-Pro-
Ala-Z-X. (SEQ ID NO 15)

Peptide XV corresponds to amino acids 2263 to 2282 and has the sequence:

(XV)

Y-Glu-Asp-Glu-Arg-Glu-Ile-Ser-Val-Pro-Ala-Glu-Ile-Leu-Arg-Lys-Ser-Arg-Arg-Phe-
Ala-Z-X. (SEQ ID NO 16)

Peptide XVI corresponds to amino acids 2275 to 2294 and has the sequence:

This Sheet Has Been Replaced





(XVI)

Y-Leu-Arg-Lys-Ser-Arg-Arg-Phe-Ala-Gln-Ala-Leu-Pro-Val-Trp-Ala-Arg-Pro-Asp-Tyr-
Asn-Z-X. (SEQ ID NO 17)

Peptide XVII corresponds to amino acids 2287 to 2306 and has the sequence:

(XVII)

Y-Val-Trp-Ala-Arg-Pro-Asp-Tyr-Asn-Pro-Pro-Leu-Val-Glu-Thr-Trp-Lys-Lys-Pro-Asp-
Tyr-Z-X. (SEQ ID NO 18)

Peptide XVIII corresponds to amino acids 2299 to 2318 and has the sequence:

(XVIII)

Y-Glu-Thr-Trp-Lys-Lys-Pro-Asp-Tyr-Glu-Pro-Pro-Val-Val-His-Gly-Cys-Pro-Leu-Pro-
Pro-Z-X. (SEQ ID NO 19)

Peptide XIX corresponds to amino acids 2311 to 2330 and has the sequence:

(XIX)

Y-Val-His-Gly-Cys-Pro-Leu-Pro-Pro-Pro-Lys-Ser-Pro-Pro-Val-Pro-Pro-Pro-Arg-Lys-
Lys-Z-X. (SEQ ID NO 20)

Of particular interest is the use of the mercapto-group of cysteines or thioglycolic acids used for acylating terminal amino groups for cyclizing the peptides or coupling two peptides together. The cyclization or coupling may occur via a single bond or may be accomplished using thiol-specific reagents to form a molecular bridge.

The peptides may be coupled to a soluble carrier for the purpose of either raising antibodies or facilitating the adsorption of the peptides to a solid phase. The nature of the carrier should be such that it has a molecular weight greater than 5000 and should not be recognized by antibodies in human serum. Generally, the carrier will be a protein. Proteins which are frequently used as carriers are keyhole limpet hemocyanin, bovine gamma globulin, bovine serum albumin, and poly-L-lysine.

There are many well described techniques for coupling peptides to carriers. The linkage may occur at the N-terminus, C-terminus or at an internal site in the peptide. The peptide may also be derivatized for coupling. Detailed descriptions of a wide variety of coupling procedures are given, for example, in Van Regenmortel, M.H.V., Briand, J.P., Muller, S., and Plaué, S., Laboratory Techniques in Biochemistry and Molecular Biology, Vol. 19, Synthetic Polypeptides as Antigens, Elsevier Press, Amsterdam, New York, Oxford, 1988.

The peptides may also be synthesized directly on an oligo-lysine core in which both the alpha as well as the epsilon-amino groups of lysines are used as growth points for the peptides. The number of lysines comprising the core is preferably 3 or 7. Additionally, a cysteine may be included near or at the C-terminus of the complex to facilitate the formation of homo- or heterodimers. The use of this technique has been amply illustrated for hepatitis B antigens (Tam, J.P., and Lu, Y.-A., Proc. Natl. Acad. Sci. USA (1989) 86:9084-9088) as well as for a variety of other antigens (see Tam, J.P., Multiple Antigen Peptide System: A Novel Design for Synthetic Peptide Vaccine and Immunoassay, in Synthetic Peptides, Approaches to Biological Problems, Tam, J.P., and Kaiser, E.T., ed. Alan R. Liss Inc., New York, 1989).

Depending on their intended use, the peptides may be either labeled or unlabeled. Labels which may be employed may be of any type, such as enzymatic, chemical, fluorescent, luminescent, or radioactive. In addition, the peptides may be modified for binding to surfaces or solid phases, such as, for example, microtiter plates, nylon membranes, glass or plastic beads, and chromatographic supports such as cellulose, silica, or agarose. The methods by which peptides can be attached or bound to solid support or surface are well known to those versed in the art.

Of particular interest is the use of mixtures of peptides for the detection of antibodies specific for hepatitis C virus. Mixtures of peptides which are considered particularly advantageous are:

- A. II (SEQ ID NO 2), III (SEQ ID NO 4), V (SEQ ID NO 6), IX (SEQ ID NO 10), and XVIII (SEQ ID NO 19),
- B. I (SEQ ID NO 1), II (SEQ ID NO 2), V (SEQ ID NO 6), IX (SEQ ID NO 10), XI (SEQ ID NO 12), XVI (SEQ ID NO 17), and XVIII (SEQ ID NO 19),
- C. II (SEQ ID NO 2), III (SEQ ID NO 4), IV (SEQ ID NO 5), V (SEQ ID NO 6), VIII (SEQ ID NO 9), XI (SEQ ID NO 12), XVI (SEQ ID NO 17), and XVIII (SEQ ID NO 19),
- D. II (SEQ ID NO 2), IX (SEQ ID NO 10), and XVIII (SEQ ID NO 19),
- E. II (SEQ ID NO 2), III (SEQ ID NO 4), IV (SEQ ID NO 5), and V (SEQ ID NO 6),
- F. VIII (SEQ ID NO 9), IX (SEQ ID NO 10), XI (SEQ ID NO 12), XIII (SEQ ID NO 14), and XIX (SEQ ID NO 15),
- G. XV (SEQ ID NO 16), XVI (SEQ ID NO 17), XVII (SEQ ID NO 18), XVIII (SEQ ID NO 19), and XIX (SEQ ID NO 20).

Antibodies which recognize the peptides can be detected in a variety of ways. A preferred method of detection is the enzyme-linked immunosorbent assay (ELISA) in which a peptide or mixture of peptides is bound to a solid support. In most cases, this will be a microtiter plate but may in principle be any sort of insoluble solid phase. A suitable dilution or dilutions of serum or other body fluid to be tested is brought into contact with the solid phase to which the peptide is bound. The incubation is carried out for a time necessary to allow the binding reaction to occur. Subsequently, unbound components are removed by washing the solid phase. The detection of immune complexes is achieved using antibodies which specifically bind to human immunoglobulins, and which have been labeled with an enzyme, preferably but not limited to either horseradish peroxidase, alkaline phosphatase, or beta-galactosidase, which is capable of converting a colorless or nearly colorless substrate or co-substrate into a highly colored product or a product capable of forming a colored complex with a chromogen. Alternatively, the detection system may employ an enzyme which, in the presence of the proper substrate(s), emits light. The amount of product formed is detected either visually, spectrophotometrically, electrochemically, or luminometrically, and is compared to a similarly treated control. The detection system may also employ radioactively labeled antibodies, in which case the amount of immune complex is quantified by scintillation counting or gamma counting.

Other detection systems which may be used include those based on the use of protein A derived from Staphylococcus aureus Cowan strain I, protein G from group C Staphylococcus sp. (strain 26RP66), or systems which make use of the high affinity biotin-avidin or streptavidin binding reaction.

Antibodies raised to carrier-bound peptides can also be used in conjunction with labeled peptides for the detection of antibodies present in serum or other body fluids by competition assay. In this case, antibodies raised to carrier-bound peptides are attached to a solid support which may be, for example, a plastic bead or a plastic tube. Labeled peptide is then mixed with suitable dilutions of the fluid to be tested and this mixture is subsequently brought into contact with the antibody bound to the solid support. After a suitable incubation period, the solid support is washed and the amount of labeled peptide is quantified. A reduction in the amount of label bound to the solid support is indicative of the presence of antibodies in the original sample. By the same token, the peptide may also be bound to the solid support. Labeled antibody may then be allowed to compete with antibody present in the sample under conditions in which the amount of peptide is limiting. As in the previous example, a reduction in the measured signal is indicative of the presence of antibodies in the sample tested.

Another preferred method of antibody detection is the homogeneous immunoassay. There are many possible variations in the design of such assays. By way of example, numerous possible configurations for homogeneous enzyme immunoassays and methods by which they may be performed are given in Tijssen, P., Practice and Theory of Enzyme Immunoassays, Elsevier Press, Amersham, Oxford, New York, 1985. Detection systems which may be employed include those based on enzyme channeling, bioluminescence, allosteric activation and allosteric inhibition. Methods employing liposome-entrapped enzymes or coenzymes may also be used (see Pinnaduwege, P. and Huang, L., Clin. Chem. (1988) 34/2: 268-272, and Ullman, E.F. et al., Clin. Chem. (1987) 33/9: 1579-1584 for examples).

This Sheet Has Been Replaced





The synthesis of the peptides can be achieved in solution or on a solid support. Synthesis protocols generally employ the use t-butyloxycarbonyl- or 9-fluorenylmethoxy-carbonyl-protected activated amino acids. The procedures for carrying out the syntheses, the types of side-chain protection, and the cleavage methods are amply described in, for example, Stewart and Young, Solid Phase Peptide Synthesis, 2nd Edition, Pierce Chemical Company, 1984; and Atherton and Sheppard, Solid Phase Peptide Synthesis, IRL Press, 1989.

Experimental

I. Peptide Synthesis

All of the peptides described were synthesized on Pepsyn K polyamide-Kieselguhr resin (Milligen, Novato, California) which had been functionalized with ethylenediamine and onto which the acid-labile linker 4-(alpha-Fmoc-amino-2',4'-dimethoxybenzyl) phenoxyacetic acid had been coupled (Rink, Tetrahedron Lett. (1987)28:3787). t-Butyl-based side-chain protection and Fmoc alpha-amino-protection was used. The guanidino-group of arginine was protected by the 2,2,5,7,8-pentamethylchroman-6-sulfonyl moiety. The imidazole group of histidine was protected by either t-Boc or trityl and the sulfhydryl group of cysteine was protected by a trityl group. Couplings were carried out using performed O-pentafluorophenyl esters except in the case of arginine where diisopropylcarbodiimide-mediated hydroxybenzotriazole ester formation was employed. Except for peptide I, all peptides were N-acetylated using acetic anhydride. All syntheses were carried out on a Milligen 9050 PepSynthesizer (Novato, California) using continuous flow procedures. Following cleavage with trifluoroacetic acid in the presence of scavengers and extraction with diethylether, all peptides were analyzed by C₁₈ -reverse phase chromatography.

II. Detection of Antibodies to Hepatitis C Virus

A. Use of peptides bound to a nylon membrane.

Peptides were dissolved in a suitable buffer to make a concentrated stock solution which was then further diluted in phosphate-buffered saline (PBS) or sodium carbonate buffer, pH 9.6 to make working solutions. The peptides were applied as lines on a nylon membrane (Pall, Portsmouth, United Kingdom), after which the membrane was treated with casein to block unoccupied binding sites. The membrane was subsequently cut into strips perpendicular to the direction of the peptide lines. Each strip was then incubated with a serum sample diluted 1 to 100, obtained from an HCV-infected individual. Antibody binding was detected by incubating the strips with goat anti-human immunoglobulin antibodies conjugated to the enzyme alkaline phosphatase. After removing unbound conjugate by washing, a substrate solution containing 5-bromo-4-chloro-3-indolylphosphate and nitro blue tetrazolium was added.

Positive reactions are visible as colored lines corresponding to the positions of the peptides which are specifically recognized. The reaction patterns of thirty-six different sera are tabulated in Table 1. The results shown in Table 1 are further summarized in Table 2.

B. Use of peptides in an enzyme-linked immunosorbent assay (ELISA).

Peptide stock solutions were diluted in sodium carbonate buffer, pH 9.6 and used to coat microtiter plates at a peptide concentration of 2 micrograms per milliliter. A mixture consisting of peptides II, III, V, IX, and XVIII was also used to coat plates. Following coating, the plates were blocked with casein. Fifteen HCV-antibody-positive sera and control sera from seven uninfected blood donors were diluted 1 to 20 and incubated in wells of the peptide-coated plates. Antibody binding was detected by incubating the plates with goat anti-human immunoglobulin antibodies conjugated to the enzyme horseradish peroxidase. Following removal of unbound conjugate by washing, a solution containing H₂O₂ and 3,3',5,5'-tetramethylbenzidine was added. Reactions were stopped after a suitable interval by addition of sulfuric acid. Positive reactions gave rise to a yellow color which was quantified using a conventional microtiter plate reader. The results of these determinations are tabulated in Table 3. To correct for any aspecific binding which could be attributable to the physical or chemical properties of the peptides themselves, a cut-off value was determined for each peptide individually. This cut-off absorbance value was calculated as the average optical density of the negative samples plus 0.200. Samples giving absorbance values higher than the cut-off values are considered positive. The results for the fifteen positive serum samples are further summarized in Table 4.

While it is evident that some of the peptides are recognized by a large percentage of sera from HCV-infected individuals, it is also clear that no single peptide is recognized by all sera. In contrast, the peptide mixture was recognized by all fifteen sera and, for six of the fifteen sera, the optical densities obtained were equal to or higher than those obtained for any of the peptides individually. These

results serve to illustrate the advantages of using mixtures of peptides for the detection of anti-HCV antibodies.

C. Binding of antibodies in sera from HCV-infected patients to various individual peptides and peptide mixtures in an ELISA.

Five peptides were used individually and in seven different combinations to coat microtiter plates. The plates were subsequently incubated with dilutions of fifteen HCV antibody-positive sera in order to evaluate the relative merits of using mixtures as compared to individual peptides for antibody detection. The mixtures used and the results obtained are shown in Figure 2.

In general, the mixtures functioned better than individual peptides. This was particularly evident for mixture 12 (peptides I, III, V, IX, and XVIII) which was recognized by all twelve of the sera tested. These results underscore the advantages of using mixtures of peptides in diagnostic tests for the detection of antibodies to HCV.

D. Use of a mixture of peptides in an ELISA assay for the detection of anti-HCV antibodies.

A mixture of peptides II, III, V, IX, and XVIII was prepared and used to coat microtiter plates according to the same procedure used to test the individual peptides. A total of forty-nine sera were tested from patients with clinically diagnosed but undifferentiated chronic non A non B hepatitis as well as forty-nine sera from healthy blood donors. Detection of antibody binding was accomplished using goat anti-human immunoglobulin antibodies conjugated to horseradish peroxidase. The resulting optical density values are given in Table 5. These results indicate that the mixture of peptides is not recognized by antibodies in sera from healthy donors (0/49 reactives) but is recognized by a large proportion (41/49, or 84%) of the sera from patients with chronic NANBH. These results demonstrate that the peptides described can be used effectively as mixtures for the diagnosis of HCV infection.

E. Detection of anti-HCV antibodies in sera from patients with acute NANB infection using individual peptides bound to nylon membranes and a mixture of peptides in an ELISA assay, and comparison with a commercially available kit.

Peptides were applied to nylon membranes or mixed and used to coat microtiter plates as previously described. The peptide mixture consisted of peptides II, III, V, IX, and XVIII. Sera obtained from twenty-nine patients with acute non-A, non-B hepatitis were then tested for the presence of antibodies to hepatitis C virus. These same sera were also evaluated using a commercially available kit (Ortho, Emeryville, CA, USA).

The results of this comparative study are given in Table 6. In order to be able to compare the peptide-based ELISA with the commercially available kit, the results for both tests are also expressed as signal to noise ratios (S/N) which were calculated by dividing the measured optical density obtained for each sample by the cut-off value. A signal-to-noise ratio greater or equal to 1.0 is taken to represent a positive reaction. For the commercially available kit, the cut-off value was calculated according to the manufacturer's instructions. The cut-off value for the peptide-based ELISA was calculated as the average optical density of five negative samples plus 0.200.

The scale used to evaluate antibody recognition of nylon-bound peptides was the same as that given in Table 1. Of the twenty-nine samples tested, twenty-five (86%) were positive in the peptide-based ELISA and recognized one or more nylon-bound peptides. In contrast, only fourteen of the twenty-nine sera scored positive in the commercially available ELISA. These results serve to illustrate the advantages of using peptide mixtures for the detection of anti-HIV antibodies as well as the need to include in the mixtures peptides which contain amino acid sequences derived from different regions of the HCV polyprotein.

This Sheet Has Been Replaced



Table 1. Recognition of peptides bound to nylon membranes by sera from persons infected by HCV.

		PEPTIDE								
Serum nr.		I	II	III	IV	V	VI	VII	VIII	IX
1				3	1					1
2										
3	1	0.5	2	1			0.5		2	0.5
4										
6			2	1	0.5					
7	0.5	1	2	1	0.5				3	2
8	0.5	1	3	1	1			1	1	
10		1	0.5						3	1
13	0.5	0.5	2		0.5					
15				0.5					2	1
16	2	1	0.5	0.5	1	0.5			2	0.5
18	1	1	3	0.5			2	0.5		
23	0.5		1	1				0.5		
24	1	0.5	2	1	0.5	0.5	0.5	0.5	2	
25			1	0.5					2	0.5
26									1	
27	0.5	0.5	1						3	2
29		0.5	3	2	1	1	0.5			
30		0.5	0.5	1	1	0.5				
31		1	0.5							
32		1	2							
33									0.5	
34		1	1	1					3	1
35	1	1	2	1	1	1	0.5			
36	1		2	1		1				
37	1	1								
44		1	2	1	0.5					
46		0.5	2	0.5	0.5			0.5	2	
47		0.5		0.5			0.5		1	
48	1	2		2				0.5	2	
49		1	1	0.5	0.5	0.5				
50		1	2	1					2	0.5
51			2	0.5	0.5			0.5		
52			2	0.5				0.5		
54			2		0.5	0.5		1	0.5	
56	ND	ND	ND	ND	ND	ND	ND	ND		2

Blank: no reaction; 0.5: weakly positive; 1: clearly positive;
2: strong reaction; 3: intense reaction; ND: not determined

This Sheet Has Been Replaced



TABLE 1 continued

	X	XI	XII	XIII	XIV	XV	XVI	XVII	XVIII	XIX
5		0.5					2	2	1	1
		0.5		1	2		2	1		
	1	0.5								
				1						
							2			
10	2	1		1		2		0.5	1	1
	2	1		1			1	1	0.5	
	1				0.5	2	2		2	2
	1			1	0.5		0.5			
15	0.5				1		0.5			
	1					2	2	1	2	
							1	0.5		
	1	0.5		0.5		0.5				
							1			
20	0.5	2		1	1		2			
							0.5			
							1	2	1	0.5
	2	1		1	1	2	2	1	1	1
25										
	0.5	0.5				0.5				
	1	0.5				1	1			
	1	0.5							0.5	
							1	1	0.5	
30										
						2		0.5		
35										
									1	1
				0.5		1	1		1	0.5
	0.5	0.5		0.5				1		
	1	1		1			1	1	0.5	0.5
40							1	1	0.5	
	0.5									
	1	1		1			1	1	1	
		0.5	1	2	1					

45

50

This Sheet Has Been Replaced



55

Table 2

Summary of antibody binding to nylon-bound HCV peptides by sera from infected patients.		
Peptide	No. reactive sera	% reactive sera
I	13/35	37
II	22/35	63
III	27/35	77
IV	24/35	69
V	14/35	40
VI	11/35	31
VII	11/35	31
VIII	19/36	53
IX	9/36	25
X	17/36	47
XI	15/36	42
XII	1/36	3
XIII	13/36	36
XIV	7/36	19
XV	9/36	25
XVI	20/36	56
XVII	14/36	39
XVIII	14/36	39
XIX	8/36	22

This Sheet Has Been Replaced



Table 3. Comparison of Individual Peptides in an ELISA Assay
for the Detection of Antibodies to HCV.

sample ident	peptide								
	I	II	III	IV	V	VI	VII	VIII	IX
1	0.786	1.119	1.284	0.265	0.042	0.04	0.05	0.571	0.659
2	0.044	0.039	0.11	0.041	0.037	0.038	0.039	0.479	0.78
3	0.815	0.944	0.825	0.399	0.654	0.487	0.32	0.705	0.965
7	1.122	1.23	0.588	0.682	0.659	0.182	0.107	0.907	1.42
8	1.155	1.159	1.2	0.508	1.272	0.433	0.623	0.61	0.863
10	1.089	1.236	1.083	0.044	0.508	0.042	0.073	1.49	1.529
11	0.048	0.051	0.476	0.052	0.119	0.039	0.1	0.634	0.711
15	0.224	0.602	0.813	0.093	0.068	0.077	0.147	0.807	1.225
23	0.62	0.8	0.924	0.568	0.759	0.442	0.683	0.089	0.121
24	1.042	1.132	1.026	0.518	0.916	0.302	0.253	1.013	1.364
49	0.624	0.73	0.884	0.171	0.372	0.055	0.04	0.084	0.064
13	0.76	0.857	0.815	0.087	0.422	0.098	0.045	0.473	0.489
31	0.84	1.114	0.445	0.672	0.046	0.041	0.042	0.184	0.15
47	1.303	1.53	1.236	0.751	0.83	0.629	0.073	0.545	0.739
56	1.169	1.301	1.364	1.269	1.374	0.85	1.066	1.45	1.523
bd A28	0.054	0.043	0.139	0.045	0.135	0.042	0.041	0.086	0.115
bd A169	0.041	0.042	0.134	0.044	0.038	0.04	0.041	0.061	0.07
bd A170	0.04	0.044	0.117	0.04	0.036	0.04	0.04	0.081	0.05
bd A171	0.041	0.046	0.148	0.043	0.037	0.045	0.045	0.077	0.065
bd A166	0.047	0.046	0.124	0.044	0.038	0.042	0.041	0.056	0.066
bd A165	0.041	0.046	0.123	0.043	0.035	0.051	0.042	0.051	0.091
AVG	0.044	0.045	0.131	0.043	0.053	0.043	0.042	0.069	0.076
STD	0.005	0.002	0.011	0.002	0.037	0.004	0.002	0.013	0.021
cut off	0.109	0.101	0.214	0.099	0.214	0.105	0.098	0.158	0.189

EP 0 644 202 A1

This Sheet Has Been Replaced



TABLE 3 continued

X	XI	XII	XIII	XIV	XV	XVI	XVII	XVIII	XIX
0.048	0.04	0.043	0.068	0.044	0.041	1.063	0.956	1.383	1.346
0.169	0.563	0.039	0.042	0.515	0.039	0.64	0.319	0.154	0.49
0.468	0.668	0.041	0.093	0.341	0.043	0.292	0.038	0.046	0.038
0.663	0.646	0.041	0.235	0.068	0.575	0.042	0.041	0.872	1.271
0.752	1.175	0.046	0.42	0.102	0.068	0.552	0.671	0.417	0.058
0.689	0.834	0.041	0.044	0.314	0.793	0.886	0.037	1.335	1.356
0.199	0.967	0.125	0.454	0.088	0.111	0.274	0.093	0.838	0.065
0.315	0.688	0.046	0.154	0.202	0.065	0.372	0.097	0.155	0.077
0.422	0.896	0.041	0.049	0.101	0.068	0.311	0.038	0.052	0.05
0.236	0.397	0.054	0.123	0.076	0.051	0.418	0.053	0.1	0.085
0.209	0.731	0.044	0.113	0.039	0.044	0.299	0.038	0.192	0.041
0.529	0.735	0.043	0.044	0.186	0.043	0.086	0.037	0.066	0.04
0.255	0.69	0.041	0.04	0.061	0.136	0.292	0.038	0.224	0.501
0.044	0.041	0.041	0.041	0.498	0.04	0.268	0.042	1.288	1.206
0.079	1.069	0.058	0.568	0.038	0.039	0.218	0.036	0.087	0.039
0.044	0.042	0.044	0.052	0.043	0.043	0.307	0.042	0.045	0.061
0.043	0.042	0.041	0.04	0.041	0.041	0.255	0.038	0.056	0.042
0.04	0.039	0.04	0.038	0.038	0.144	0.292	0.036	0.058	0.039
0.043	0.041	0.043	0.039	0.04	0.045	0.286	0.037	0.05	0.04
0.041	0.041	0.042	0.04	0.041	0.041	0.207	0.039	0.046	0.041
0.041	0.04	0.042	0.039	0.043	0.039	0.253	0.034	0.06	0.098
0.042	0.041	0.042	0.041	0.041	0.059	0.267	0.038	0.053	0.054
0.001	0.001	0.001	0.005	0.002	0.038	0.033	0.002	0.006	0.021
0.095	0.094	0.095	0.106	0.097	0.223	0.416	0.084	0.121	0.167

EP 0 644 202 A1

This Sheet Has Been Replaced



Table 4

Summary of antibody-binding to individual peptides in an ELISA assay.		
Peptide	No. reactive sera	% reactive sera
I	13	87
II	13	87
III	14	93
IV	10	67
V	10	67
VI	7	47
VII	8	53
VIII	13	87
IX	12	80
X	13	87
XI	13	87
XII	1	7
XIII	7	47
XIV	8	53
XV	2	13
XVI	5	33
XVII	4	27
XVIII	10	67
XIX	6	40

This Sheet Has Been Replaced



Table 5 Use of a peptide mixture for the detection of antibodies to HCV in sera from chronic NANBH patients and comparison to sera from healthy blood donors.

Chronic NANB Sera		Control Sera	
Serum nr.	Optical Density	Serum nr.	Optical Density
101	0.041	1	0.049
102	1.387	2	0.047
103	1.578	3	0.049
104	1.804	4	0.046
105	1.393	5	0.049
107	1.604	6	0.045
108	1.148	7	0.043
109	1.714	8	0.053
110	1.692	9	0.049
112	0.919	10	0.047
113	1.454	11	0.060
114	0.936	12	0.044
115	0.041	13	0.049
116	1.636	14	0.051
118	1.242	15	0.056
119	1.568	16	0.050
120	1.290	17	0.049
121	1.541	18	0.055
122	1.422	19	0.054
123	1.493	20	0.058
124	1.666	21	0.050
125	1.644	22	0.044
126	1.409	23	0.043
127	1.625	24	0.045
128	1.061	25	0.046
129	1.553	26	0.049

This Sheet Has Been Replaced



Table 5
continued

5	130	1.709	27	0.050
	131	0.041	28	0.047
10	132	0.044	29	0.050
	133	1.648	30	0.053
15	134	0.043	31	0.051
	135	1.268	32	0.053
	136	1.480	33	0.055
20	138	0.628	34	0.064
	139	0.042	35	0.063
	140	0.040	36	0.057
25	141	0.039	38	0.048
	142	1.659	39	0.045
	143	1.457	40	0.046
30	144	0.722	41	0.046
	145	1.256	42	0.051
	146	0.373	43	0.057
35	147	1.732	44	0.050
	148	1.089	45	0.050
	149	1.606	46	0.045
40	150	1.725	47	0.041
	151	1.449	48	0.064
	154	1.639	49	0.040
45	155	1.775	50	0.036

This Sheet Has Been Replaced



Table 6. Comparison of anti-HCV antibody detection by nylon-bound peptides, a peptide-based ELISA, and a commercially available kit.

Serum nr.	Nylon-bound peptides										
	I	III	IV	V	VI	VIII	XI	XIV	XV	XVI	XVIII
191	0	0	0	0	0	0	0	0	0	0	0
192	0	0	0	0	0	0	0	0	0	0	0
193	0	0	0	0	0	0	0	0	0	0	0
194	0	0	0	0	0	0	0	0	0	0	1
195	1	2	2	3	0	0	0.5	0.5	1	3	0
196	1	2	1	2	0.5	0.5	0.5	0.5	0.5	2	0
197	1	2	1	2	0	0.5	0.5	0.5	1	2	0
198	1	2	2	2	0	0	0	0	1	2	1
211	0.5	1	0.5	0.5	0	2	2	0	2	0	0
213	0	0	0	1	0	0	0	0	0	0	0
214	0	0	0	1	0	0	0	0	0	0	0
215	0	0	0	1	0	0	0	0	0	0	0
216	0	0	0	0.5	0	0	0	0	0	0	0
217	0	0	0	1	0	0	0	0	0	0	1
219	0.5	1	1	2	1	0.5	1	0	0.5	0.5	1
220	0.5	1	1	2	1	0.5	0	0	0	0	0
221	0	0	0	0.5	0	0	2	0.5	0.5	0	0
222	1	1	1	1	0	0	3	0.5	0.5	0	0
223	1	1	1	1	0	0.5	3	0.5	0.5	0	0
224	1	1	2	1	0	0.5	0.5	0.5	0	0	2
225	0	0	0	0	0	2	3	2	0.5	0.5	1
226	0.5	0	0	0	0	2	2	0.5	0.5	0.5	2
227	0	0	0	0	0	2	2	2	0	0	2
229	0.5	0	0.5	0.5	0	2	2	2	0	0	3
234	0.5	0.5	0	0.5	0	0	3	1	3	1	0
235	0	0	0	0.5	0	0	0	0	0	0	0
236	0	0	0	0.5	0	0	0	0	0	0	0
237	0	0	0	1	0	0	0	0	0	0	0
238	0	0	0	1	0	1	1	0	0	0	0

0: no reaction; 0.5: weakly positive; 1: clearly positive;
2: strong reaction; 3: intense reaction;

This Sheet Has Been Replaced



Table 6 continued

Serum nr.	Optical density Peptide EUSA	S/N	Optical density Commercial EUSA	S/N
191	0.045	0.18	0.295	0.47
192	0.042	0.17	0.289	0.46
193	0.039	0.16	0.197	0.32
194	0.044	0.18	0.183	0.29
195	1.692	6.77	3.000*	4.82*
196	1.569	6.28	0.386	0.62
197	1.523	6.09	0.447	0.72
198	1.578	6.31	0.354	0.57
211	1.606	6.42	3.000*	4.82*
213	0.369	1.48	0.127	0.20
214	0.444	1.78	0.101	0.16
215	0.637	2.55	0.101	0.16
216	0.812	3.25	0.092	0.15
217	1.320	5.28	0.875	1.40
219	1.547	6.19	3.000*	4.82*
220	1.536	6.14	3.000*	4.82*
221	1.428	5.71	0.327	0.52
222	1.362	5.45	3.000*	4.82*
223	1.316	5.26	3.000*	4.82*
224	1.304	5.22	3.000*	4.82*
225	1.178	4.71	2.398	3.85
226	1.256	5.14	3.000*	4.82*
227	1.335	5.34	3.000*	4.82*
228	1.400	5.60	3.000*	4.82*
234	1.481	5.92	3.000*	4.82*
235	0.351	1.40	0.257	0.41
236	0.475	1.50	0.245	0.39
237	1.134	4.54	0.351	0.56
238	1.056	4.38	1.074	1.72

Cut-off: 0.250

Cut-off: 0.623

* O.D. exceeded 3.000 and was out of range. The values given are therefore minimum values.

This Sheet Has Been Replaced



SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

- (A) NAME: Innogenetics sa.
- (B) STREET: Industriepark Zwijnaarde 7, box 4
- (C) CITY: Ghent
- (E) COUNTRY: Belgium
- (F) POSTAL CODE (ZIP): B-9052
- (G) TELEPHONE: 00 32 9 241 07 11
- (H) TELEFAX: 00 32 9 241 07 99

(ii) TITLE OF INVENTION: Synthetic antigens for the detection of antibodies to Hepatitis C virus

(iii) NUMBER OF SEQUENCES: 20

(iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.25 (EPO)

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 1
- (D) OTHER INFORMATION: /label= bound-moiety
/note= "H, one or more N-terminal amino acid(s, or (a) chemical group(s) used for amino-terminal linking of the amino acid sequence to a solid phase or carrier"

This Sheet Has Been Replaced



(ix) FEATURE:

(A) NAME/KEY: Modified-site

(B) LOCATION: 20

(D) OTHER INFORMATION: /label= bound-moiety

/note= "Z-X : Z is a bond, (an) amino acid(s), or
 (a) chemical group(s) used for carboxy-terminal linking
 the amino acid sequence to a solid phase or carrier"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

Met Ser Thr Ile Pro Lys Pro Gln Arg Lys Thr Lys Arg Asn Thr Asn

1 5 10 15

Arg Arg Pro Gln

20

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

(A) NAME/KEY: Modified-site

(B) LOCATION: 1

(D) OTHER INFORMATION: /label= bound-moiety

/note= "H, one or more N-terminal amino acid(s), or
 (a) chemical group(s) used for amino-terminal linking
 of the amino acid sequence to a solid phase or
 carrier"

(ix) FEATURE:

(A) NAME/KEY: Modified-site

(B) LOCATION: 20

(D) OTHER INFORMATION: /label= bound-moiety

/note= "Z-X : Z is a bond, (an) amino acid(s), or (a)
 chemical group(s) used for carboxy-terminal
 linking the amino acid sequence to a solid phase or
 carrier"

This Sheet Has Been Replaced



(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Pro Gln Arg Lys Thr Lys Arg Asn Thr Asn Arg Arg Pro Gln Asp Val
1 5 10 15

Lys Phe Pro Gly
20

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 11 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 1
- (D) OTHER INFORMATION: /label= bound-moiety
/note= "H, one or more N-terminal amino acid(s),
or (a) chemical group(s) used for amino-terminal
linking of the amino acid sequence to a solid phase or
carrier"

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 11
- (D) OTHER INFORMATION: /label= bound-moiety
/note= "Z-X : Z is a bond, (an) amino acid(s), or (a)
chemical group(s) used for carboxy-terminal
linking the amino acid sequence to a solid phase or
carrier"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

Gln Arg Lys Thr Lys Arg Asn Thr Asn Arg Arg
1 5 10

This Sheet Has Been Replaced



(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
 (B) LOCATION: 1
 (D) OTHER INFORMATION: /label= bound-moiety
 /note= "H, one or more N-terminal amino acid(s),
 or (a) chemical group(s) used for amino-terminal
 linking of the amino acid sequence to a solid phase or
 carrier"

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
 (B) LOCATION: 20
 (D) OTHER INFORMATION: /label= bound-moiety
 /note= "Z-X : Z is a bond, (an) amino acid(s), or (a)
 chemical group(s) used for carboxy-terminal
 linking the amino acid sequence to a solid phase or
 carrier"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

Arg	Asn	Thr	Asn	Arg	Arg	Pro	Gln	Asp	Val	Lys	Phe	Pro	Gly	Gly	Gly
1				5					10					15	
Gln	Ile	Val	Gly												
				20											

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

This Sheet Has Been Replaced



(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

(A) NAME/KEY: Modified-site
(B) LOCATION: 1
(D) OTHER INFORMATION: /label= bound-moiety
/note= "H, one or more N-terminal amino acid(s),
or (a) chemical group(s) used for amino-terminal
linking of the amino acid sequence to a solid phase or
carrier"

(ix) FEATURE:

(A) NAME/KEY: Modified-site
(B) LOCATION: 20
(D) OTHER INFORMATION: /label= bound-moiety
/note= "Z-X : Z is a bond, (an) amino acid(s), or (a)
chemical group(s) used for carboxy-terminal
linking the amino acid sequence to a solid phase or
carrier"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

Leu	Pro	Arg	Arg	Gly	Pro	Arg	Leu	Gly	Val	Arg	Ala	Thr	Arg	Lys	Thr
1				5				10					15		
Ser	Glu	Arg	Ser												
				20											

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

(A) NAME/KEY: Modified-site
(B) LOCATION: 1
(D) OTHER INFORMATION: /label= bound-moiety
/note= "H, one or more N-terminal amino acid(s),
or (a) chemical group(s) used for amino-terminal
linking of the amino acid sequence to a solid phase or
carrier"

This Sheet Has Been Replaced



(ix) FEATURE:

(A) NAME/KEY: Modified-site

(B) LOCATION: 20

(D) OTHER INFORMATION: /label= bound-moiety

/note= "Z-X : Z is a bond, (an) amino acid(s), or (a)
chemical group(s) used for carboxy-terminal
linking the amino acid sequence to a solid phase or
carrier"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

Thr Arg Lys Thr Ser Glu Arg Ser Gln Pro Arg Gly Arg Arg Gln Pro
1 5 10 15

Ile Pro Lys Val
20

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

(A) NAME/KEY: Modified-site

(B) LOCATION: 1

(D) OTHER INFORMATION: /label= bound-moiety

/note= "H, one or more N-terminal amino acid(s),
or (a) chemical group(s) used for amino-terminal
linking of the amino acid sequence to a solid phase or
carrier"

(ix) FEATURE:

(A) NAME/KEY: Modified-site

(B) LOCATION: 20

(D) OTHER INFORMATION: /label= bound-moiety

/note= "Z-X : Z is a bond, (an) amino acid(s), or (a)
chemical group(s) used for carboxy-terminal
linking the amino acid sequence to a solid phase or
carrier"

This Sheet Has Been Replaced



(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

Arg Arg Gln Pro Ile Pro Lys Val Arg Arg Pro Glu Gly Arg Thr Trp
1 5 10 15

Ala Gln Pro Gly
20

(2) INFORMATION FOR SEQ ID NO: 8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 1
- (D) OTHER INFORMATION: /label= bound-moiety
/note= "H, one or more N-terminal amino acid(s),
or (a) chemical group(s) used for amino-terminal
linking of the amino acid sequence to a solid phase or
carrier"

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 20
- (D) OTHER INFORMATION: /label= bound-moiety
/note= "Z-X : Z is a bond, (an) amino acid(s), or (a)
chemical group(s) used for carboxy-terminal
linking the amino acid sequence to a solid phase or
carrier"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

Gly Arg Thr Trp Ala Gln Pro Gly Tyr Pro Trp Pro Leu Tyr Gly Asn
1 5 10 15

Glu Gly Cys Gly
20

This Sheet Has Been Replaced



(2) INFORMATION FOR SEQ ID NO: 9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
 (B) LOCATION: 1
 (D) OTHER INFORMATION: /label= bound-moiety
 /note= "H, one or more N-terminal amino acid(s),
 or (a) chemical group(s) used for amino-terminal
 linking of the amino acid sequence to a solid phase or
 carrier"

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
 (B) LOCATION: 20
 (D) OTHER INFORMATION: /label= bound-moiety
 /note= "Z-X : Z is a bond, (an) amino acid(s), or (a)
 chemical group(s) used for carboxy-terminal
 linking the amino acid sequence to a solid phase or
 carrier"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

Leu	Ser	Gly	Lys	Pro	Ala	Ile	Ile	Pro	Asp	Arg	Glu	Val	Leu	Tyr	Arg
1				5				10					15		
Glu	Phe	Asp	Glu												
				20											

(2) INFORMATION FOR SEQ ID NO: 10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

This Sheet Has Been Replaced



(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

(A) NAME/KEY: Modified-site

(B) LOCATION: 1

(D) OTHER INFORMATION: /label= bound-moiety

/note= "H, one or more N-terminal amino acid(s),
or (a) chemical group(s) used for amino-terminal
linking of the amino acid sequence to a solid phase or
carrier"

(ix) FEATURE:

(A) NAME/KEY: Modified-site

(B) LOCATION: 20

(D) OTHER INFORMATION: /label= bound-moiety

/note= "Z-X : Z is a bond, (an) amino acid(s), or (a)
chemical group(s) used for carboxy-terminal
linking the amino acid sequence to a solid phase or
carrier"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

Ile	Ile	Pro	Asp	Arg	Glu	Val	Leu	Tyr	Arg	Glu	Phe	Asp	Glu	Met	Glu
1				5					10					15	
Glu Cys Ser Gln															
20															

(2) INFORMATION FOR SEQ ID NO: 11:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

(A) NAME/KEY: Modified-site

(B) LOCATION: 1

(D) OTHER INFORMATION: /label= bound-moiety

This Sheet Has Been Replaced



/note= "H, one or more N-terminal amino acid(s),
or (a) chemical group(s) used for amino-terminal
linking of the amino acid sequence to a solid phase or
carrier"

(ix) FEATURE:

(A) NAME/KEY: Modified-site

(B) LOCATION: 20

(D) OTHER INFORMATION: /label= bound-moiety

/note= "Z-X : Z is a bond, (an) amino acid(s), or (a)
chemical group(s) used for carboxy-terminal
linking the amino acid sequence to a solid phase or
carrier"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

Asp	Glu	Met	Glu	Glu	Cys	Ser	Gln	His	Leu	Pro	Tyr	Ile	Glu	Gln	Gly
1				5					10				15		
Met Met Leu Ala															
20															

(2) INFORMATION FOR SEQ ID NO: 12:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

(A) NAME/KEY: Modified-site

(B) LOCATION: 1

(D) OTHER INFORMATION: /label= bound-moiety

/note= "H, one or more N-terminal amino acid(s),
or (a) chemical group(s) used for amino-terminal
linking of the amino acid sequence to a solid phase or
carrier"

This Sheet Has Been Replaced



(ix) FEATURE:

- (A) NAME/KEY: Modified-site
 (B) LOCATION: 20
 (D) OTHER INFORMATION: /label= bound-moiety
 /note= "Z-X : Z is a bond, (an) amino acid(s), or (a) chemical group(s) used for carboxy-terminal linking the amino acid sequence to a solid phase or carrier"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

Ser Gln His Leu Pro Tyr Ile Glu Gln Gly Met Met Leu Ala Glu Gln
 1 5 10 15

Phe Lys Gln Gln
 20

(2) INFORMATION FOR SEQ ID NO: 13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
 (B) LOCATION: 1
 (D) OTHER INFORMATION: /label= bound-moiety
 /note= "H, one or more N-terminal amino acid(s), or (a) chemical group(s) used for amino-terminal linking of the amino acid sequence to a solid phase or carrier"

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
 (B) LOCATION: 20
 (D) OTHER INFORMATION: /label= bound-moiety
 /note= "Z-X : Z is a bond, (an) amino acid(s), or (a) chemical group(s) used for carboxy-terminal linking the amino acid sequence to a solid phase or carrier"

This Sheet Has Been Replaced



(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

5 Ile Glu Gln Gly Met Met Leu Ala Glu Gln Phe Lys Gln Lys Ala Leu
1 5 10 15
Gly Leu Leu Gln
20

10

(2) INFORMATION FOR SEQ ID NO: 14:

(i) SEQUENCE CHARACTERISTICS:

15 (A) LENGTH: 20 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

20

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

25 (A) NAME/KEY: Modified-site
(B) LOCATION: 1
(D) OTHER INFORMATION: /label= bound-moiety
/note= "H, one or more N-terminal amino acid(s),
30 or (a) chemical group(s) used for amino-terminal
linking of the amino acid sequence to a solid phase or
carrier"

35 (ix) FEATURE:

(A) NAME/KEY: Modified-site
(B) LOCATION: 20
(D) OTHER INFORMATION: /label= bound-moiety
40 /note= "Z-X : Z is a bond, (an) amino acid(s), or (a)
chemical group(s) used for carboxy-terminal
linking the amino acid sequence to a solid phase or
carrier"

45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

Leu Ala Glu Gln Phe Lys Gln Lys Ala Leu Gly Leu Leu Gln Thr Ala
1 5 10 15
50 Ser Arg Gln Ala
20

55

This Sheet Has Been Replaced



(2) INFORMATION FOR SEQ ID NO: 15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 1
- (D) OTHER INFORMATION: /label= bound-moiety
/note= "H, one or more N-terminal amino acid(s),
or (a) chemical group(s) used for amino-terminal
linking of the amino acid sequence to a solid phase or
carrier"

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 20
- (D) OTHER INFORMATION: /label= bound-moiety
/note= "Z-X : Z is a bond, (an) amino acid(s), or (a)
chemical group(s) used for carboxy-terminal
linking the amino acid sequence to a solid phase or
carrier"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

Gln Lys Ala Leu Gly Leu Leu Gln Thr Ala Ser Arg Gln Ala Glu Val
 1 5 10 15
 Ile Ala Pro Ala
 20

(2) INFORMATION FOR SEQ ID NO: 16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

This Sheet Has Been Replaced



(ii) MOLECULE TYPE: peptide

5

(ix) FEATURE:

(A) NAME/KEY: Modified-site

(B) LOCATION: 1

(D) OTHER INFORMATION: /label= bound-moiety

10

/note= "H, one or more N-terminal amino acid(s),
or (a) chemical group(s) used for amino-terminal
linking of the amino acid sequence to a solid phase or
carrier"

15

(ix) FEATURE:

(A) NAME/KEY: Modified-site

(B) LOCATION: 20

(D) OTHER INFORMATION: /label= bound-moiety

20

/note= "Z-X : Z is a bond, (an) amino acid(s), or (a)
chemical group(s) used for carboxy-terminal
linking the amino acid sequence to a solid phase or
carrier"

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:

Glu Asp Glu Arg Glu Ile Ser Val Pro Ala Glu Ile Leu Arg Lys Ser
1 5 10 15

30

Arg Arg Phe Ala
20

35

(2) INFORMATION FOR SEQ ID NO: 17:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 amino acids

(B) TYPE: amino acid

40

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

45

(ix) FEATURE:

(A) NAME/KEY: Modified-site

(B) LOCATION: 1

50

(D) OTHER INFORMATION: /label= bound-moiety

/note= "H, one or more N-terminal amino acid(s),

55

This Sheet Has Been Replaced



or (a) chemical group(s) used for amino-terminal
linking of the amino acid sequence to a solid phase or
carrier"

(ix) FEATURE:

(A) NAME/KEY: Modified-site

(B) LOCATION: 20

(D) OTHER INFORMATION: /label= bound-moiety

/note= "Z-X : Z is a bond, (an) amino acid(s), or (a)
chemical group(s) used for carboxy-terminal
linking the amino acid sequence to a solid phase or
carrier"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17:

Leu Arg Lys Ser Arg Arg Phe Ala Gln Ala Leu Pro Val Trp Ala Arg

1 5 10 15

Pro Asp Tyr Asn

20

(2) INFORMATION FOR SEQ ID NO: 18:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

(A) NAME/KEY: Modified-site

(B) LOCATION: 1

(D) OTHER INFORMATION: /label= bound-moiety

/note= "H, one or more N-terminal amino acid(s),
or (a) chemical group(s) used for amino-terminal
linking of the amino acid sequence to a solid phase or
carrier"

(ix) FEATURE:

(A) NAME/KEY: Modified-site

(B) LOCATION: 20

(D) OTHER INFORMATION: /label= bound-moiety

This Sheet Has Been Replaced



/note= "Z-X : Z is a bond, (an) amino acid(s), or (a) chemical group(s) used for carboxy-terminal linking the amino acid sequence to a solid phase or carrier"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18:

Val Trp Ala Arg Pro Asp Tyr Asn Pro Pro Leu Val Glu Thr Trp Lys
 1 5 10 15
 Lys Pro Asp Tyr
 20

(2) INFORMATION FOR SEQ ID NO: 19:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

(A) NAME/KEY: Modified-site

(B) LOCATION: 1

(D) OTHER INFORMATION: /label= bound-moiety

/note= "H, one or more N-terminal amino acid(s), or (a) chemical group(s) used for amino-terminal linking of the amino acid sequence to a solid phase or carrier"

(ix) FEATURE:

(A) NAME/KEY: Modified-site

(B) LOCATION: 20

(D) OTHER INFORMATION: /label= bound-moiety

/note= "Z-X : Z is a bond, (an) amino acid(s), or (a) chemical group(s) used for carboxy-terminal linking the amino acid sequence to a solid phase or carrier"

This Sheet Has Been Replaced





(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19:

Glu Thr Trp Lys Lys Pro Asp Tyr Glu Pro Pro Val Val His Gly Cys

1 5 10 15

Pro Leu Pro Pro

20

(2) INFORMATION FOR SEQ ID NO: 20:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

(A) NAME/KEY: Modified-site

(B) LOCATION: 1

(D) OTHER INFORMATION: /label= bound-moiety

/note= "H, one or more N-terminal amino acid(s), --
or (a) chemical group(s) used for amino-terminal
linking of the amino acid sequence to a solid phase or
carrier"

(ix) FEATURE:

(A) NAME/KEY: Modified-site

(B) LOCATION: 20

(D) OTHER INFORMATION: /label= bound-moiety

/note= "Z-X : Z is a bond, (an) amino acid(s), or (a)
chemical group(s) used for carboxy-terminal
linking the amino acid sequence to a solid phase or carrier"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 20:

Val His Gly Cys Pro Leu Pro Pro Pro Lys Ser Pro Pro Val Pro Pro

1 5 10 15

Pro Arg Lys Lys

20

Claims

1. A peptide composition comprising at least one peptide containing fewer than 50 amino acids, with said peptide being selected from the following group of amino acid sequences:



(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19:

Glu Thr Trp Lys Lys Pro Asp Tyr Glu Pro Pro Val Val His Gly Cys
1 5 10 15

Pro Leu Pro Pro
20

(2) INFORMATION FOR SEQ ID NO: 20:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

(A) NAME/KEY: Modified-site

(B) LOCATION: 1

(D) OTHER INFORMATION: /label= bound-moiety

/note= "H, one or more N-terminal amino acid(s),
or (a) chemical group(s) used for amino-terminal
linking of the amino acid sequence to a solid phase or
carrier"

(ix) FEATURE:

(A) NAME/KEY: Modified-site

(B) LOCATION: 20

(D) OTHER INFORMATION: /label= bound-moiety

/note= "Z-X : Z is a bond, (an) amino acid(s), or (a)
chemical group(s) used for carboxy-terminal
linking the amino acid sequence to a solid phase or carrier"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 20:

Val His Gly Cys Pro Leu Pro Pro Pro Lys Ser Pro Pro Val Pro Pro
1 5 10 15

Pro Arg Lys Lys
20

Claims

1. A peptide composition comprising at least one peptide containing fewer than 50 amino acids, with said peptide being selected from the following group of amino acid sequences:



- (XV) Y-Glu-Asp-Glu-Arg-Glu-Ile-Ser-Val-Pro-Ala-Glu-Ile-Leu-Arg-Lys-Ser-Arg-Arg-Phe-Ala-Z-X (SEQ ID NO 16),
- (XVI) Y-Leu-Arg-Lys-Ser-Arg-Arg-Phe-Ala-Gln-Ala-Leu-Pro-Val-Trp-Ala-Arg-Pro-Asp-Tyr-Asn-Z-X (SEQ ID NO 17),
- (XVII) Y-Val-Trp-Ala-Arg-Pro-Asp-Tyr-Asn-Pro-Pro-Leu-Val-Glu-Thr-Trp-Lys-Lys-Pro-Asp-Tyr-Z-X (SEQ ID NO 18),
- (XVIII) Y-Glu-Thr-Trp-Lys-Lys-Pro-Asp-Tyr-Glu-Pro-Pro-Val-Val-His-Gly-Cys-Pro-Leu-Pro-Pro-Z-X (SEQ ID NO 19),
- (XIX) Y-Val-His-Gly-Cys-Pro-Leu-Pro-Pro-Pro-Lys-Ser-Pro-Pro-Val-Pro-Pro-Pro-Arg-Lys-Lys-Z-X (SEQ ID NO 20),

wherein Y is H, one or more N-terminal amino acids, or (a) chemical group(s) used for amino-terminal linking of the amino acid sequence to a solid phase or carrier, Z is a bond, (an) amino acid(s), or (a) chemical group(s) used for carboxy-terminal linking the amino acid sequence to a solid phase or carrier; and X is NH₂, OH, or a linkage involving either of these groups, or,

- (a) variants of each of the above peptides having an amino acid sequence which contains insertions, deletions, conservative or non-conservative mutations of one or more amino acids with respect to each of amino acid sequences as represented by SEQ ID NO 16 to 20, provided that said variant peptides are capable of providing for immunological competition with at least one strain of HCV;
- (b) fragments of each of the above peptides or variants thereof comprising at least five contiguous amino acids of any of said amino acid sequences as represented by SEQ ID NO 16 to 20, provided that said variant peptides are capable of providing for immunological competition with at least one strain of HCV;
- (c) peptide structures comprising two or more of the above peptides or variants of the peptides joined together;
- (d) cyclic peptides comprising at least one of the above peptides or variants thereof; provided said peptides are different from any of the following peptides:

This Sheet Has Been Replaced



Ser-Phe-Asp-Pro-Leu-Val-Ala-Glu-Glu-Asp-Glu-Arg-Glu-Ile-Ser-Val
 Glu-Arg-Glu-Ile-Ser-Val-Pro-Ala-Glu-Ile-Leu-Arg-Lys-Ser-Arg-Arg
 5 Arg-Phe-Ala-Gln-Ala-Leu-Pro-Val-Trp-Ala-Arg
 Arg-Pro-Asp-Tyr-Asn-Pro-Pro-Leu-Val-Glu-Thr-Trp-Lys-Lys-Pro-Asp-Tyr-Glu-
 Pro-Pro-Val
 10 Val-Val-His-Gly-Cys-Pro-Leu-Pro-Pro-Pro-Lys-Ser-Pro-Pro-Val-Pro-Pro-Pro-
 Arg-Lys-Lys

- 15 2. A peptide composition according to claim 1, further characterized in that said peptide(s) contain(s) fewer than 35 amino acids.
3. A peptide according to claim 1, further characterized in that said peptide(s) contain(s) fewer than 25
 20 amino acids.
4. A peptide composition according to any of claims 1 to 4, further characterized in that it also comprises at least one peptide containing fewer than 50 amino acids, with said peptide being selected from the following group of amino acid sequences:

(I) Y-Met-Ser-Thr-Ile-Pro-Lys-Pro-Gln-Arg-Lys-Thr-Lys-Arg-Asn-Thr-Asn-
 30 Arg-Arg-Pro-Gln-Z-X (SEQ ID NO 1),

(II) Y-Pro-Gln-Arg-Lys-Thr-Lys-Arg-Asn-Thr-Asn-Arg-Arg-Pro-Gln-Asp-
 35 Val-Lys-Phe-Pro-Gly-Z-X (SEQ ID NO 2),

45 This Sheet Has Been Replaced





- 5 (III) Y-Arg-Asn-Thr-Asn-Arg-Arg-Pro-Gln-Asp-Val-Lys-Phe-Pro-Gly-Gly-Gly-Gln-Ile-Val-Gly-Z-X (SEQ ID NO 4),
- 10 (IV) Y-Leu-Pro-Arg-Arg-Gly-Pro-Arg-Leu-Gly-Val-Arg-Ala-Thr-Arg-Lys-Thr-Ser-Glu-Arg-Ser-Z-X (SEQ ID NO 5),
- 15 (V) Y-Thr-Arg-Lys-Thr-Ser-Glu-Arg-Ser-Gln-Pro-Arg-Gly-Arg-Arg-Gln-Pro-Ile-Pro-Lys-Val-Z-X (SEQ ID NO 6),
- 20 (VI) Y-Arg-Arg-Gln-Pro-Ile-Pro-Lys-Val-Arg-Arg-Pro-Glu-Gly-Arg-Thr-Trp-Ala-Gln-Pro-Gly-Z-X (SEQ ID NO 7),
- 25 (VII) Y-Gly-Arg-Thr-Trp-Ala-Gln-Pro-Gly-Tyr-Pro-Trp-Pro-Leu-Tyr-Gly-Asn-Glu-Gly-Cys-Gly-Z-X (SEQ ID NO 8),
- 30 (VIII) Y-Leu-Ser-Gly-Lys-Pro-Ala-Ile-Ile-Pro-Asp-Arg-Glu-Val-Leu-Tyr-Arg-Glu-Phe-Asp-Gly-Z-X (SEQ ID NO 9),
- 35 (IX) Y-Ile-Ile-Pro-Asp-Arg-Glu-Val-Leu-Tyr-Arg-Glu-Phe-Asp-Glu-Met-Glu-Glu-Cys-Ser-Gln-Z-X (SEQ ID NO 10),
- 40 (X) Y-Asp-Glu-Met-Glu-Glu-Cys-Ser-Gln-His-Leu-Pro-Tyr-Ile-Glu-Gln-Gly-Met-Met-Leu-Ala-Z-X (SEQ ID NO 11),
- 45 (XI) Y-Ser-Gln-His-Leu-Pro-Tyr-Ile-Glu-Gln-Gly-Met-Met-Leu-Ala-Glu-Gln-Phe-Lys-Gln-Lys-Z-X (SEQ ID NO 12),
- 50 (XII) Y-Ile-Glu-Gln-Gly-Met-Met-Leu-Ala-Glu-Gln-Phe-Lys-Gln-Lys-Ala-Leu-Gly-Leu-Leu-Gln-Z-X (SEQ ID NO 13),
- 55 (XIII) Y-Leu-Ala-Glu-Gln-Phe-Lys-Gln-Lys-Ala-Leu-Gly-Leu-Leu-Gln-Thr-Ala-Ser-Arg-Gln-Ala-Z-X (SEQ ID NO 14),
- (XIV) Y-Gln-Lys-Ala-Leu-Gly-Leu-Leu-Gln-Thr-Ala-Ser-Arg-Gln-Ala-Glu-Val-Ile-Ala-Pro-Ala-Z-X (SEQ ID NO 15),





wherein Y is H, one or more N-terminal amino acids, or (a) chemical group(s) used for amino-terminal linking of the amino acid sequence to a solid phase or carrier, Z is a bond, (an) amino acid(s), or (a) chemical group(s) used for carboxy-terminal linking the amino acid sequence to a solid phase or carrier; and X is NH₂, OH, or a linkage involving either of these groups,

or,

(a) variants of each of the above peptides having an amino acid sequence which contains insertions, deletions, conservative or non-conservative mutations of one or more amino acids with respect to each of amino acid sequences as represented in SEQ ID NO 1 to 15, provided that said variant peptides are capable of providing for immunological competition with at least one strain of HCV;

(b) fragments of each of the above peptides or variants thereof comprising at least five contiguous amino acids of any of said amino acid sequences as represented in SEQ ID NO 1 to 15, provided that said variant peptides are capable of providing for immunological competition with at least one strain of HCV;

(c) peptide structures comprising two or more of the above peptides or variants of the peptides joined together;

(d) cyclic peptides comprising at least one of the above peptides or variants thereof.

5. A peptide composition according to any of claims 1 to 4, further characterized in that it contains at least one of the following mixtures of peptides having the amino acid sequence as defined in any of claims 1 or 4:

A. II (SEQ ID NO 2), III (SEQ ID NO 4), V (SEQ ID NO 6), IX (SEQ ID NO 10), and XVIII (SEQ ID NO 19),

B. I (SEQ ID NO 1), II (SEQ ID NO 2), V (SEQ ID NO 6), IX (SEQ ID NO 10), XI (SEQ ID NO 12), XVI (SEQ ID NO 17), and XVIII (SEQ ID NO 19),

C. II (SEQ ID NO 2), III (SEQ ID NO 4), IV (SEQ ID NO 5), V (SEQ ID NO 6), VIII (SEQ ID NO 9), XI (SEQ ID NO 12), XVI (SEQ ID NO 17), and XVIII (SEQ ID NO 19),

D. II (SEQ ID NO 2), IX (SEQ ID NO 10), and XVIII (SEQ ID NO 19),

E. II (SEQ ID NO 2), III (SEQ ID NO 4), IV (SEQ ID NO 5), and V (SEQ ID NO 6),

F. VIII (SEQ ID NO 9), IX (SEQ ID NO 10), XI (SEQ ID NO 12), XIII (SEQ ID NO 14), and XIX (SEQ ID NO 15),

G. XV (SEQ ID NO 16), XVI (SEQ ID NO 17), XVII (SEQ ID NO 18), XVIII (SEQ ID NO 19), and XIX (SEQ ID NO 20).

6. A peptide composition or peptide according to claim any of claims 1 to 5, further characterized in that said peptides are coupled N-terminally, C-terminally or internally to a carrier molecule for the purpose of raising antibodies or facilitating the adsorption of said peptides to a solid phase.

7. A peptide composition or peptide according to any of claims 1 to 6, further characterized in that said peptides contain a detectable label.

8. Use of a peptide composition according to any of claims 1 to 7, for the incorporation into an immunoassay for detecting the presence of antibodies to Hepatitis C virus, present in a body fluid.

9. A method for the *in vitro* detection of antibodies to hepatitis C virus present in a body fluid such as serum or plasma, comprising at least the steps of: (a) contacting said body fluid of a person to be diagnosed with a peptide composition or peptide according to any of claims 1 to 7, and, (b) detecting the immunological complex formed between said antibodies and the peptide(s) used.

10. A kit for the detection of anti-hepatitis C virus antibodies in a body fluid, comprising at least the following components:

- a peptide composition or peptide according to any of claims 1 to 7,
- a means for detecting an immunological complex formed between said peptides and said antibodies.

11. A method according to claim 9, further characterized in that said peptides are applied as lines on a nylon membrane, and with said nylon membrane being preferably cut into strips perpendicular to the direction of the peptide lines, thus allowing each strip to be incubated with an appropriately diluted serum sample from an individual.



12. A kit according to claim 10, further characterized in that said peptides, separately or in combination, are used to coat the wells of microtiter plates.

5 13. Use of a peptide composition or a peptide according to any of claims 1 to 7 to raise antibodies against HCV.

10

15

20

25

30

35

This Sheet Has Been Replaced



40

45



* C L H S *

50

55