



- (51) **International Patent Classification:**
C07K 1/30 (2006.01) *C07K 16/00* (2006.01)
- (21) **International Application Number:**
PCT/EP2013/003552
- (22) **International Filing Date:**
25 November 2013 (25.11.2013)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
12008475.1 20 December 2012 (20.12.2012) EP
- (71) **Applicant:** MERCK PATENT GMBH [DE/DE]; Frankfurter Strasse 250, 64293 Darmstadt (DE).
- (72) **Inventors:** BAUER, Johann; Huegelstrasse 71, 64283 Darmstadt (DE). RAPP, Almut; Taunusstrasse 36, 64289 Darmstadt (DE). STANISLAWSKI, Bernd; Dehnhardtstrasse 20, 60433 Frankfurt am Main (DE). CAPITO, Florian; Riedeselstrasse 68, 64283 Darmstadt (DE).
- (81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.


- (84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) **Title:** COPOLYMERS FOR PROTEIN PRECIPITATION

Fig. 1



(57) **Abstract:** The present invention relates to the isolation of recombinant and/or biotherapeutic proteins for capture or clarification from cell culture fluid using copolymers. The copolymers used according to the process of the present invention comprise hydrophobic and anionic residues.