

[Name of Document] Specification

Title of Invention: Pharmaceutical Composition for Edaravone Oral Administration And

Method For Administering Same

Technical Field

[0001]


The present invention relates to a pharmaceutical composition containing edaravone as an active ingredient for oral administration or for intragastric administration.

Background Art

[0002]

Edaravone is 3-methyl-1-phenyl-2-pyrazolin-5-one represented by a formula below and is known for its pharmaceutical applications as a therapeutic agent or a brain function normalizing agent for oxidative stress diseases, such as amyotrophic lateral sclerosis (ALS) (Patent Documents 1 and 2).

[Chem. 1]



[0003]

ALS is an intractable disease with an initial symptom including weakness of the hands,

dyskinesia of the fingers, and fasciculation of the upper limbs, and followed by a symptom such as muscle atrophy, muscle weakness, bulbar palsy, and muscular fasciculation, ALS leads to respiratory failure. Based on the site of onset, ALS is classified into an upper limb type, a bulbar type, a lower limb type, and a mixed type, all of which affect muscle groups throughout the body as symptoms progress. Although the etiology of ALS has not yet been fully elucidated, the following hypotheses have been presented as the major etiology: (1) autoimmunity (appearance of autoantibodies against a Ca channel), (2) excitatory amino acid excess and poisoning (increase in extracellular glutamic acid and impaired glutamic acid transport), (3) oxidative stress disorder (neuronopathy due to Cu/Zn superoxide dismutase (SOD) gene abnormality and free radicals), (4) cytoskeletal damage (accumulation of neurofilaments and appearance of inclusion bodies in motor nerve cells), and (5) neurotrophic factor deficiency.

[0004]

Currently, edaravone is typically used as a therapeutic agent for ALS, and in this application, it is only provided in the form of an injectable agent. Thus, an oral formulation is desired to be provided, as it is less burdensome for patients and caregivers and is preferred in terms of QOL. To meet such a demand, an oral formulation of edaravone is being developed through clinical and other studies (Patent Document 3).

Citation List

Patent Literature

[0005]

Patent Document 1: JP 05-031523 B

Patent Document 2: JP 3758164 B

Patent Document 3: WO 2020/091036

Non-Patent Literature

[0006]

Non-Patent Document 1: Pharmaceutical Research, Vol. 22, No. 1, 11-23, 2005

Summary of Invention

Technical Problem

[0007]

Generally, with respect to therapeutic agents for an oxidative stress disease, the same agent may have different conditions under which the agent can effectively exert its effect depending on the administration method of the agent. In addition, an oral formulation of edaravone is still under development, and conditions under which the same effect as that of the injection can be exerted has not been elucidated.

[0008]

Thus, an object of the present invention is to provide a pharmaceutical composition containing edaravone as an active ingredient for oral administration and equivalent administration.

Solution to Problem

[0009]

A pharmaceutical composition according to an embodiment of the present invention is characterized by containing edaravone as an active ingredient, the pharmaceutical composition being administered to a subject orally or intragastrically with any one first time interval selected from the group consisting of 1) to 3) below:

- 1) in a case where the subject has consumed a high-fat meal, 8 hours or longer after the consumption;
- 2) in a case where the subject has consumed a standard meal, 4 hours or longer after the consumption; and
- 3) in a case where the subject has consumed a light meal, 2 hours or longer after the consumption.

[0010]

A method of administering an edaravone-containing pharmaceutical composition according to an embodiment of the present invention is characterized by administering a pharmaceutical composition containing edaravone as an active ingredient to a subject orally or intragastrically with any one first time interval selected from the group consisting of 1) to 3) below:

- 1) in a case where the subject has consumed a high-fat meal, 8 hours or longer after the consumption;
- 2) in a case where the subject has consumed a standard meal, 4 hours or longer after the

consumption; and

3) in a case where the subject has consumed a light meal, 2 hours or longer after the consumption.

[0011]

The present inventors have found the present invention through the circumstances described below. These descriptions, however, do not particularly limit the present invention.

[0012]

Food-drug interactions can have a significant effect on safety and efficacy of drugs. For example, administration of a drug together with a meal may increase systemic absorption of the drug and change pharmacokinetics, and may result in improvement in efficacy and increase in the incidence of side effects. Alternatively, administration of a drug together with a meal may reduce systemic absorption of the drug and reduce efficacy.

[0013]

On the other hand, Wu et al. reported, for drugs exhibiting high solubility and high permeability and classified as BCS Class 1, meal consumption has little effect on pharmacokinetics, such as bioavailability, when the drugs are administered orally (Non-Patent Document 1).

[0014]

The present inventors found that a volume of water required for dissolving 1 g of edaravone is 500 mL (Drug Interview Form of Radicut (Revised in June 2017 (Revision 20)), which

means a volume of water required for dissolving 105 mg of edaravone is calculated as 52.5 mL, and thus edaravone is classified as a high solubility compound. In addition, the present inventors also determined that edaravone is classified as a high permeability compound from the results of a membrane permeability test using Caco-2 cells shown in Example 3 described below. Thus, edaravone is regarded as a BCS Class 1 drug with high solubility and high permeability in the report of Wu et al. These findings led to a presumption that edaravone is a compound whose pharmacokinetics are not affected by meal consumption.

[0015]

Nevertheless, consumption of a meal prior to administration was found to affect the pharmacokinetics of edaravone as shown in Examples described below. Thus, as a result of further diligent research, the present inventors newly found that not only the presence or absence of a meal but also the type of meal can cause a difference in effects on the pharmacokinetics of edaravone, and further found that a suitable administration timing depends on each type of meal, and completed the present invention.

Advantageous Effects of Invention

[0016]

An embodiment of the present invention can optimize, for each type of meal, a first time interval after a meal to administration and thus, for example, can avoid effects of a meal on pharmacokinetics such as bioavailability in both oral administration and intragastric administration of edaravone. Thus, the pharmaceutical composition according to an

embodiment of the present invention can exhibit an excellent therapeutic effect on oxidative stress diseases as a formulation for oral administration (hereinafter also referred to as an “oral formulation”) or a formulation for intragastric administration (hereinafter also referred to as an “intragastric formulation”) similarly to an injection administered intravenously which is the administration method known in the art. In addition, the pharmaceutical composition according to an embodiment of the present invention can be administered orally or intragastrically and thus can further reduce the burden on patients and caregivers.

Brief Description of Drawings

[0017]

FIG. 1 shows spaghetti plots comparing the C_{max} after administration of edaravone oral formulation between a fasted group, a group administered 8 hours after high-fat meal consumption, a group administered 4 hours after standard meal consumption, and a group administered 2 hours after light meal consumption. The circles with solid lines represent individual values, and the crosses with dotted lines represent mean values.

FIG. 2 shows spaghetti plots comparing the AUC_{0-1h} and $AUC_{1h-\infty}$ after oral administration of an edaravone oral formulation between a high-fat meal group consuming a high-fat meal 1 hour after the administration and a fasted group administered under fasted conditions. The circles with solid lines represent individual values, and the crosses with dotted lines represent mean values.

FIG. 3 shows a box plot (study #9) and individual plots (studies #5, #6, and #9) for the

$AUC_{0-\infty}$ and C_{max} after administration of an edaravone oral formulation. The circles represent individual values, and the crosses represent mean values. In the box plot, the top and bottom of the whiskers represent the maximum and minimum values, and the upper limit, middle, and lower limit of the box represent the 25th, 50th, and 75th percentile values, respectively.

FIG. 4 shows a graph indicating an individual value plot of the $AUC_{0-\infty}$ vs. edaravone doses and a curve regressed by a 4-parameter logistic model. The circles represent individual values in a study #7, and the squares represent individual values in a study #8. An $AUC_{0-\infty}$ of edaravone upon intravenous administration is indicated by the dotted line using a value upon administration at 60 mg/60 min.

Description of Embodiments

[0018]

Embodiments of the present invention are as follows.

[0019]

For a pharmaceutical composition according to an embodiment of the present invention, for example, a second time interval from edaravone administration to next meal consumption is 30 minutes or longer.

[0020]

For the pharmaceutical composition according to an embodiment of the present invention, for example, a second time interval from edaravone administration to next meal consumption is 1

hour or longer.

[0021]

The pharmaceutical composition according to an embodiment of the present invention is, for example, for oral administration.

[0022]

The pharmaceutical composition according to an embodiment of the present invention is, for example, for intragastric administration.

[0023]

For the pharmaceutical composition according to an embodiment of the present invention, for example, the intragastric administration is administration via a tube.

[0024]

For the pharmaceutical composition according to an embodiment of the present invention, for example, the tube is a percutaneous endoscopic gastrostomy tube (PEG tube) or a nasopharyngeal gastric tube (NG tube).

[0025]

The pharmaceutical composition according to an embodiment of the present invention is, for example, for treating an oxidative stress disease.

[0026]

For the pharmaceutical composition according to an embodiment of the present invention, for example, the oxidative stress disease is any one of diseases selected from the group

consisting of amyotrophic lateral sclerosis, Parkinson's disease, spinocerebellar degeneration, muscular dystrophy, Alzheimer's disease, cerebral infarction, multiple sclerosis, systemic scleroderma, and stomatitis.

[0027]

For the pharmaceutical composition according to an embodiment of the present invention, for example, the oxidative stress disease is amyotrophic lateral sclerosis.

[0028]

For the pharmaceutical composition according to an embodiment of the present invention, for example, the first time interval is a time interval during an administration period of intermittent administration or daily administration.

[0029]

For the pharmaceutical composition according to an embodiment of the present invention, for example, a dose per administration is from 90 to 120 mg of edaravone.

[0030]

For the pharmaceutical composition according to an embodiment of the present invention, for example, a dose per administration is from 90 to 105 mg of edaravone.

[0031]

For the pharmaceutical composition according to an embodiment of the present invention, for example, a daily administration frequency is once, and a dose per administration is 105 mg of edaravone.

[0032]

In a method of administering a pharmaceutical composition according to an embodiment of the present invention, for example, a second time interval from edaravone administration to next meal consumption is set to 30 minutes or longer.

[0033]

In the method of administering a pharmaceutical composition according to an embodiment of the present invention, for example, a second time interval from edaravone administration to next meal consumption is set to 1 hour or longer.

[0034]

In the method of administering a pharmaceutical composition according to an embodiment of the present invention, for example, the pharmaceutical composition is administered orally.

[0035]

In the method of administering a pharmaceutical composition according to an embodiment of the present invention, for example, the pharmaceutical composition is administered intragastrically.

[0036]

In the method of administering a pharmaceutical composition according to an embodiment of the present invention, for example, the intragastric administration is administration via a tube.

[0037]

In the method of administering a pharmaceutical composition according to an embodiment of the present invention, for example, the tube is a percutaneous endoscopic gastrostomy tube (PEG tube) or a nasopharyngeal gastric tube (NG tube).

[0038]

In the pharmaceutical composition according to an embodiment of the present invention, for example, the pharmaceutical composition is administered to treat an oxidative stress disease.

[0039]

In the method of administering a pharmaceutical composition according to an embodiment of the present invention, for example, the oxidative stress disease is any one of diseases selected from the group consisting of amyotrophic lateral sclerosis, Parkinson's disease, spinocerebellar degeneration, muscular dystrophy, Alzheimer's disease, cerebral infarction, multiple sclerosis, systemic scleroderma, and stomatitis.

[0040]

In the method of administering a pharmaceutical composition according to an embodiment of the present invention, for example, the oxidative stress disease is amyotrophic lateral sclerosis.

[0041]

In the method of administering a pharmaceutical composition according to an embodiment of the present invention, for example, administration of the pharmaceutical composition includes intermittent administration or daily administration, and the pharmaceutical

composition is administered with the first time interval during an administration period of the intermittent administration or daily administration.

[0042]

In the method of administering a pharmaceutical composition according to an embodiment of the present invention, for example, a dose per administration of edaravone is from 90 to 120 mg.

[0043]

In the method of administering a pharmaceutical composition according to an embodiment of the present invention, for example, a dose per administration of edaravone is from 90 to 105 mg.

[0044]

In the method of administering a pharmaceutical composition according to an embodiment of the present invention, for example, a daily dose frequency is once, and a dose per administration is 105 mg of edaravone.

[0045]

Hereinafter, embodiments of the present invention will be described with reference to specific examples, but the present invention is not limited to these examples. In addition, the pharmaceutical composition according to an embodiment of the present invention can be used in the method of administration according to an embodiment of the present invention, and a description of the pharmaceutical composition according to an embodiment of the

present invention and a description of the method of administration according to an embodiment of the present invention can be incorporated into each other.

[0046]

In the present invention, the “first time interval” means a time duration from consumption of a meal to administration, for example, it may be used interchangeably with “time interval before administration” or “time interval after meal or before administration”, and the “second time interval” means a time duration from administration to consumption of a meal, for example, it may be used interchangeably with “time interval after administration” or “time interval after administration or before meal”. In the “first time interval” and the “second time interval”, the first and second notations are notations for formally distinguishing between the two types of time intervals before and after administration, and do not limit the order, for example.

[0047]

The pharmaceutical composition according to an embodiment of the present invention and the method of administration according to an embodiment of the present invention are characterized by the conditions for the administration timing in which the pharmaceutical composition is administered with any first time interval of 1) to 3) according to the type of meal as described above, and other conditions and processes are not particularly limited. Specifically, the effect of the present invention, that is, the excellent effect of avoiding influence from a meal on the pharmacokinetics of edaravone is achieved by the

administration timing. Thus, for example, the edaravone content and composition in the pharmaceutical composition according to an embodiment of the present invention is not particularly limited. In an embodiment of the present invention, the type of meal can be classified, for example, based on calories.

[0048]

In the present specification, “oral administration” means administering a drug from the mouth, and “intra gastric administration” means administering a drug directly to the stomach. In addition, the pharmaceutical composition according to an embodiment of the present invention is a pharmaceutical composition for use in such oral administration or intra gastric administration. In the use of the pharmaceutical composition according to an embodiment of the present invention and in the administration method according to an embodiment of the present invention, for example, the type of instrument, device, or the like when used for oral administration or intra gastric administration is not particularly limited. For intra gastric administration, for example, a tube, such as a feeding tube (hereinafter, it may be referred to as a tube for tube feeding, a tube for feeding, or a feed tube), can be used, and the tube is also referred to as, for example, a catheter. The intra gastric administration via the tube is also referred to as, for example, transluminal administration or intra gastric transluminal administration. Examples of the intra gastric administration include administration via a gastrostomy tube and nasogastric administration. The administration via a gastrostomy tube is, for example, a method of perforating the stomach by making a perforation in the surface

of the abdomen to deliver nutrients directly to the stomach (performing gastrostomy), passing the tube through the perforation (gastric fistula), and administering a drug directly into the stomach. Examples of the tube used for the administration via a gastrostomy tube include a percutaneous endoscopic gastrostomy tube (PEG tube) (hereinafter, it may be referred to as a gastric fistula tube, a feeding tube, a connecting tube for feeding, a connecting tube, a Percutaneous Endoscopic Gastrostomy (PEG) catheter, or a Percutaneous Endoscopic Gastrostomy (PEG) tube). In addition, the nasogastric administration is, for example, a method of passing a tube from the nose into the stomach and administering a drug directly into the stomach. Examples of the tube used for the nasogastric administration include a nasopharyngeal gastric tube (NG tube) (hereinafter, referred to as a nasogastric tube, a nasoesophageal feeding tube, a nasal probe, a nasal tube, a nasotracheal tube, a feeding catheter, a nasoesophageal feeding catheter, a nasogastric catheter, a gastric catheter, or a Nasopharyngeal Gastric (NG) tube). The pharmaceutical composition according to an embodiment of the present invention is also referred to as, for example, a “transluminal formulation” when used for intragastric administration via the tube.

[0049]

As described above, the pharmaceutical composition according to an embodiment of the present invention is characterized by containing edaravone as an active ingredient, the pharmaceutical composition being administered orally or intragastrically with any one of the optimal first time intervals selected from the group consisting of 1) to 3) described above for

each type of meal.

[0050]

The composition of the pharmaceutical composition according to an embodiment of the present invention is any composition containing edaravone as an active ingredient, and other constitution is not particularly limited. The pharmaceutical composition according to an embodiment of the present invention, for example, may contain only an active ingredient or may contain an active ingredient and an additional excipient. The active ingredient may be, for example, only edaravone or may be edaravone in combination with another drug. In the pharmaceutical composition according to an embodiment of the present invention, the form of edaravone is not particularly limited, and examples include a particulate form (hereinafter also referred to as an “edaravone particle”). The additional excipient is not particularly limited, and examples include dispersants, such as polyvinyl alcohols and sucrose fatty acid esters; thickeners, such as xanthan gums and powdered tragacanth; sweeteners, such as sorbitol; stabilizers, such as sodium hydrogen sulfite and L-cysteine hydrochloride; pH adjusters, such as phosphoric acid, acetic acid, and sodium hydroxide; defoamers, such as simethicone emulsions; fillers, such as glucose and starch; solvents; and perfumes. Examples of the solvent include aqueous solvents, such as water, buffers, and saline; and oil solvents, such as olive oil.

[0051]

Specific examples of the pharmaceutical composition according to an embodiment of the

present invention include suspensions containing an edaravone particle, a dispersant, and water. The suspension is also referred to as, for example, an edaravone suspension for oral administration (which may be hereinafter referred to as the “edaravone oral formulation”) or an edaravone suspension for intragastric administration (which may be hereinafter referred to as the “edaravone intragastric formulation”). Hereinafter, for the pharmaceutical composition according to an embodiment of the present invention, the oral formulation can be interchangeably read as the intragastric formulation, and the intragastric formulation can be interchangeably read as the oral formulation. The pharmaceutical composition according to an embodiment of the present invention is characterized by the administration timing as described above, and the composition is not limited. Thus, for the details of the composition of the pharmaceutical composition according to an embodiment of the present invention, the contents described in WO 2020/091036 can be incorporated herein, and the pharmaceutical composition according to an embodiment of the present invention can be manufactured by the method described in WO 2020/091036. Subjects of the administration of the pharmaceutical composition according to an embodiment of the present invention is, for example, humans or non-human animals and preferably humans. Non-human animals are, for example, mice, rats, rabbits, or horses.

[0052]

The form of the pharmaceutical composition according to an embodiment of the present invention is not particularly limited and can be, for example, a form suitable for oral

administration, a form suitable for intragastric administration, or a form applicable to both oral administration and intragastric administration. The form is, for example, a liquid, a gel, or a solid. The liquid is exemplified by solutions and suspensions, the gel is exemplified by jellies, and the solid is exemplified by tablets, capsules, powders, fine granules, and granules. In an embodiment of the present invention, the preferred form is a liquid, and in particular, a suspension is preferred.

[0053]

Examples of the embodiment of the pharmaceutical composition of the present invention include compositions (1) to (5) exemplified below. The embodiment is preferably the composition (1) but is not limited to this.

(1) Edaravone particle, polyvinyl alcohol (dispersant), xanthan gum (thickener), sorbitol (sweetener), sodium hydrogen sulfite (stabilizer), L-cysteine hydrochloride (stabilizer), phosphoric acid (pH adjuster), sodium hydroxide (pH adjuster), simethicone emulsion (defoamer), and water

(2) Edaravone particle, polyvinyl alcohol (dispersant), powdered tragacanth (thickener), sorbitol (sweetener), sodium hydrogen sulfite (stabilizer), L-cysteine hydrochloride (stabilizer), phosphoric acid (pH adjuster), sodium hydroxide (pH adjuster), simethicone emulsion (defoamer), and water

(3) Edaravone particle, sucrose fatty acid ester (dispersant), xanthan gum (thickener), sucrose (sweetener), sodium hydrogen sulfite (stabilizer), L-cysteine hydrochloride (stabilizer),

acetic acid (pH adjuster), sodium hydroxide (pH adjuster), simethicone emulsion (defoamer), and water

(4) Edaravone particle, polyvinyl alcohol (dispersant), xanthan gum (thickener), sorbitol (sweetener), sodium hydrogen sulfite (stabilizer), phosphoric acid (pH adjuster), sodium hydroxide (pH adjuster), simethicone emulsion (defoamer), and water

(5) Edaravone particle, polyvinyl alcohol (dispersant), xanthan gum (thickener), sorbitol (sweetener), sodium hydrogen sulfite (stabilizer), L-cysteine hydrochloride (stabilizer), phosphoric acid (pH adjuster), sodium hydroxide (pH adjuster), simethicone emulsion (defoamer), perfume, and water

[0054]

More specifically, examples of the composition (1) include a suspension of the following composition. The suspension of the following composition has, for example, a total volume of 5 mL, an edaravone content of 105 mg, and a form of, for example, a liquid. However, the present invention is not limited to this.

Edaravone particle: 2.1% (w/v)

Polyvinyl alcohol (dispersant): 0.1% (w/v)

Xanthan gum (thickener): 0.3% (w/v)

Sorbitol (sweetener): 40% (w/v)

Sodium hydrogen sulfite (stabilizer): 0.1% (w/v)

L-Cysteine hydrochloride (stabilizer): 0.05% (w/v)

Sodium Hydroxide (pH adjuster): q.s.

Phosphoric acid (pH adjuster): q.s.

Simethicone emulsion (defoamer): 0.05% (w/v)

Dispersion medium: water

pH: from 2.5 to 6.0

[0055]

The pharmaceutical composition according to an embodiment of the present invention can avoid effects of meal consumption on the pharmacokinetics of edaravone in oral administration or intragastric administration of edaravone by varying the timing of administration according to the type of meal as described above, that is, setting the first time interval from consumption of a meal to administration to the condition 1), 2), or 3) described above. In an embodiment of the present invention, “avoid effects of meal consumption on the pharmacokinetics of edaravone” means, for example, maintaining the same pharmacokinetics as the pharmacokinetics when a pharmaceutical composition containing edaravone is administered under fasting conditions (e.g., fasting 10 hours or longer). Specifically, this means, for example, that in administration of the same pharmaceutical composition containing edaravone by the same method of oral administration or intragastric administration, the pharmacokinetics administered under the condition 1), 2), or 3) described above according to a meal maintains the same pharmacokinetics as the pharmacokinetics administered under the fasting conditions described above. In addition, “the same

pharmacokinetics” means, for example, that the target pharmacokinetics is completely identical or in a range not significantly different. Specific examples include measuring the concentration of edaravone in plasma of a subject, for example, at a predetermined period before and after administration of the pharmaceutical composition containing edaravone. Then, “the same pharmacokinetics” means that comparison of the maximum plasma concentration (C_{max}) and area under the curve (AUC) of edaravone with the C_{max} and AUC in fasting for 10 hours or longer indicates no significant change and also the same level of distribution of individual values, or indicates statistically the same level to common criteria for bioequivalence (e.g., a least squares mean ratio and its 90% confidence interval are in the range of 0.8 to 1.25). Thus, according to an embodiment of the present invention, oral administration or intragastric administration of the pharmaceutical composition containing edaravone with an optimal administration interval of 1) to 3) described above according to the type of meal can maintain the same pharmacokinetics as the pharmacokinetics when the pharmaceutical composition is administered in fasting conditions, for example, regardless of the formula of the pharmaceutical composition.

[0056]

The pharmaceutical composition according to an embodiment of the present invention can be administered to a subject orally or intragastrically with any one of the first time intervals selected from the group consisting of 1) to 3) below, for example, according to the type of meal upon consuming a meal:

1) in a case where the subject has consumed a high-fat meal, 8 hours or longer after the consumption;

2) in a case where the subject has consumed a standard meal, 4 hours or longer after the consumption; and

3) in a case where the subject has consumed a light meal, 2 hours or longer after the consumption.

[0057]

In a certain embodiment, the pharmaceutical composition according to an embodiment of the present invention can be administered orally or intragastrically with a first time interval selected from 2) or 3) described above, for example, according to the type of meal upon consuming a meal.

[0058]

Furthermore, the pharmaceutical composition according to an embodiment of the present invention can further effectively maintain the pharmacokinetics of edaravone, for example, by adjusting the second time interval until the next meal consumption after administration.

The pharmaceutical composition according to an embodiment of the present invention may be administered with a second time interval from administration of the composition to the next meal consumption of, for example, 30 minutes or longer, preferably 1 hour or longer or 2 hours or longer. Specifically, the pharmaceutical composition according to an embodiment of the present invention may be administered with a time interval of, for example, 30 minutes

or longer, 1 hour or longer, or 2 hours or longer until the next consumption of a meal. In an embodiment of the present invention, the length of the time second interval after administration is not limited, for example, by the type of the next meal.

[0059]

In a certain embodiment, the pharmaceutical composition according to an embodiment of the present invention can be administered orally or intragastrically with any one of the first time intervals selected from the group consisting of 1) to 3) described above and with a second time interval of 30 minutes or longer until the next meal consumption, for example, according to the type of meal upon consuming a meal.

[0060]

In a certain embodiment, the pharmaceutical composition according to an embodiment of the present invention can be administered orally or intragastrically with any one of the first time intervals selected from the group consisting of 1) to 3) described above and with a second time interval of 1 hour or longer until the next meal consumption, for example, according to the type of meal upon consuming a meal.

[0061]

In a certain embodiment, the pharmaceutical composition according to an embodiment of the present invention can be administered orally or intragastrically with any one of the first time intervals selected from the group consisting of 1) to 3) described above and with a second time interval of 2 hours or longer until the next meal consumption, for example, according to

the type of meal upon consuming a meal.

[0062]

In a certain embodiment, the pharmaceutical composition according to an embodiment of the present invention can be administered orally or intragastrically with a first time interval selected from the group consisting of 2) or 3) described above and with a second time interval of 30 minutes or longer until the next meal consumption, for example, according to the type of meal upon consuming a meal.

[0063]

In a certain embodiment, the pharmaceutical composition according to an embodiment of the present invention can be administered orally or intragastrically with a first time interval selected from 2) or 3) described above and with a second time interval of 1 hour or longer until the next meal consumption, for example, according to the type of meal upon consuming a meal.

[0064]

In a certain embodiment, the pharmaceutical composition according to an embodiment of the present invention can be administered orally or intragastrically with a first time interval selected from 2) or 3) described above and with a second time interval of 2 hours or longer until the next meal consumption, for example, according to the type of meal upon consuming a meal.

[0065]

In an embodiment of the present invention, the high-fat meal, the standard meal, and the light meal can be classified, for example, based on common classification criteria. Specifically, those can be classified, for example, according to total calories of a single meal, and more specifically, those can be classified, for example, according to the total calories and percentage of calories from lipid in the total calories.

[0066]

The high-fat meal is primarily a meal classified as a high-fat meal and may contain, for example, a meal with calories approximating the upper limit side and lower limit side of the common reference value classified as a high fat meal. Specifically, the high-fat meal in an embodiment of the present invention may contain, for example, in addition to a high-fat meal, a meal with calories less than reference calories for a high-fat meal and exceeding the reference calories for a standard meal in common classification criteria or may contain a meal with calories exceeding the reference calories for a high-fat meal. Applying thus the first time interval condition 1) described above not only to the high-fat meal but also to a meal with calories exceeding the reference calories for a standard meal can effectively maintain the pharmacokinetics of edaravone administered orally or intragastrically also when a meal corresponding to a border between a high-fat meal and a standard meal is consumed. In addition, the same applies to a meal with calories exceeding the reference calories for a high-fat meal.

[0067]

A common criterion for a high-fat meal is, for example, total calories per meal of 800 to 1000 kilocalories, and examples include a meal with total calories of 800 to 1000 kilocalories and calorie intake from lipid of 50%, or a meal with total calories of 800 to 1000 kilocalories and calorie intake from lipid of 500 to 600 kilocalories and calorie intake from carbohydrate of 250 kilocalories. In an embodiment of the present invention, in addition to the high-fat meal described above, for example, a meal with total calories greater than 500 kilocalories and less than 800 kilocalories can also be classified as the high-fat meal. In addition, in an embodiment of the present invention, for example, a meal with total calories greater than 1000 kilocalories can also be classified as the high-fat meal. Furthermore, in an embodiment of the present invention, the high-fat meal described above may be, for example, a meal described as the high-fat meal in the FDA guideline (Assessing the Effects of Food on Drugs in INDs and NDAs - Clinical Pharmacology Considerations) or a meal described as the high-fat meal in the EMA guideline (Guideline on the investigation of drug interactions).

[0068]

In an embodiment of the present invention, for the unit of total calories, “kilocalories”, one kilocalorie is defined as energy required to raise the temperature of one liter of water by 1°C. In addition, kilocalories in the present specification are synonymous with “calories” used for meals as described in the website of the U.S. Department of Agriculture (USDA) (<https://www.nutrition.gov/expert-q-a>) (see the section “What is the difference between calories and kilocalories?”). In the present specification, the calorific value is described in

the unit “kilocalories” based on the former definition as described above. For example, the calorific value of 800 to 1000 kilocalories described in the present specification means a calorific value of 800 to 1000 calories according to the description of the U.S. Department of Agriculture (USDA).

[0069]

The standard meal is primarily a meal classified as a standard meal and may contain, for example, a meal with calories approximating the lower limit side of the common reference value classified as a standard meal. In addition, in the present specification, the standard meal may be referred to as a low-fat meal or a normal meal. Specifically, the standard meal in an embodiment of the present invention may contain, for example, in addition to a standard meal, a meal with calories less than reference calories for a standard meal and exceeding the reference calories for a light meal in common classification criteria. Applying thus the first time interval condition 2) described above not only to the standard meal but also to a meal with calories exceeding the reference calories for a light meal can effectively maintain the pharmacokinetics of edaravone administered orally or intragastrically also when a meal corresponding to a border between a standard meal and a light meal is consumed.

[0070]

A common criterion for a standard meal is, for example, total calories per meal of 400 to 500 kilocalories, and examples include a meal with total calories of 400 to 500 kilocalories and calorie intake from lipid of 25%, or a meal with total calories of 400 to 500 kilocalories and

calorie intake from lipid of 150 kilocalories. In an embodiment of the present invention, in addition to the standard meal described above, for example, a meal with total calories greater than 250 kilocalories and less than 400 kilocalories can also be classified as the standard meal. Furthermore, in an embodiment of the present invention, the standard meal described above may be, for example, a meal described as the low-fat meal in the FDA guideline (Assessing the Effects of Food on Drugs in INDs and NDAs - Clinical Pharmacology Considerations) or a meal described as the moderate meal in the EMA guideline (Guideline on the investigation of drug interactions).

[0071]

The light meal is primarily a meal classified as a light meal and may contain, for example, a meal with calories approximating the lower limit side of the common reference value classified as a light meal. In addition, in the present specification, the light meal may be referred to as a caloric supplement, a calorie supplement, a calorie supplement agent, a dietary supplement, a nutritional supplement, a nutritional meal, an enteral nutrition agent, a protein amino acid formulation, an elemental diet, a digestive enteral nutrition agent, a digestive nutrition agent, a semi-digestion enteral nutrition agent, a semi-digestion nutrition agent, a nutritional supplement, a nutritional supplement drink, a nutritional food, or a nutritional functional food. Specifically, the light meal in an embodiment of the present invention may contain, for example, in addition to a light meal, a meal with calories less than reference calories for a light meal in common classification criteria. Applying thus the first

time interval condition 3) described above not only to the light meal but also to a meal with calories below the reference calories for a light meal can effectively maintain the pharmacokinetics of edaravone administered orally or intragastrically also when a meal with calories below the reference calories for a light meal is consumed.

[0072]

A common criterion for a light meal is, for example, a meal with a total calories per meal of 250 kilocalories. Specific examples of the light meal include liquid type nutritional supplements with approximately 250 kilocalories (enteral nutritional supplement: Ensure Liquid (trade name)). This is used, for example, for nutritional retention for patients after surgery and tubal alimentation for patients with difficulty in meal consumption over a long period. In addition, in an embodiment of the present invention, in addition to the light meal with approximately 250 kilocalories described above, for example, a meal with total calories less than 250 kilocalories can also be classified as the light meal.

[0073]

The method of administering the pharmaceutical composition according to an embodiment of the present invention is oral administration or intragastric administration, and may be either. In administering the pharmaceutical composition according to an embodiment of the present invention to a patient over multiple times, the administration may be intragastric administration only, oral administration only, a combination of intragastric administration and oral administration, a change from oral administration to intragastric administration, or a

change from intragastric administration to oral administration. In general, intragastric administration is used as a measure for a case where nutrition cannot be consumed through the mouth because of dysphagia or the like. Thus, for example, for more effective treatment according to symptoms, the administration may be changed from oral administration to intragastric administration. In addition, in the intragastric administration, for example, the administration is intragastric administration via a tube (also referred to as transluminal administration or transluminal intragastric administration) as described above, and as a specific example, the administration may be administration via a percutaneous endoscopic gastrostomy tube (PEG tube), administration via a nasopharyngeal gastric tube (NG tube), a combination of both, or a change from one to the other.

[0074]

A material of the catheter or tube is not particularly limited, and examples include a silicone, a polyvinyl chloride, or a polyurethane. In the catheter or tube, the material of the button is, for example, a silicone, a polyurethane, a silicone rubber, or the like, and examples of the material of the tubing include a polyvinyl chloride or a silicone rubber/polyacetal. In the catheter or tube, examples of a combination of the materials of the button and tubing include a combination of a silicone button and a polyvinyl chloride tubing, a combination of a polyurethane button and a polyvinyl chloride tubing, or a combination of a silicone rubber button and a silicone rubber/polyacetal tubing.

[0075]

In a preferred embodiment of the present invention, the active ingredient according to an embodiment of the present invention, edaravone, is administered orally or via a feeding tube in the form of the pharmaceutical composition according to an embodiment of the present invention. For the feeding tube, for example, a nasopharyngeal gastric tube (which may be referred to as an “NG tube”) or a percutaneous endoscopic gastrostomy tube (which may be referred to as a “PEG tube”) can be used, and the pharmaceutical composition is administered via this feeding tube. Examples of the feeding tube include those made of a silicone, a polyvinyl chloride (PVC), and/or a polyurethane.

[0076]

In a preferred embodiment of the present invention, when the active ingredient according to an embodiment of the present invention, edaravone, is administered in the form of the pharmaceutical composition according to an embodiment of the present invention, the pharmaceutical composition is taken in a fasting state. In this embodiment, a meal is preferably not consumed 1 hour after administration, and consumption of a high-fat meal (from 800 to 1000 calories (synonymous with from 800 to 1000 kilocalories as defined in the present specification), fat 50%) is desirably avoided 8 hours before administration, consumption of a low-fat meal (from 400 to 500 calories (synonymous with from 400 to 500 kilocalories as defined in the present specification), fat 25%) is desirably avoided 4 hours before administration, or consumption of a caloric supplement (250 calories (synonymous with 250 kilocalories as defined in the present specification), e.g., ENSURE LIQUID (trade

name)) is desirably avoided 2 hours before administration.

[0077]

In another preferred embodiment of the present invention, when the active ingredient according to an embodiment of the present invention, edaravone, is administered in the form of the pharmaceutical composition according to an embodiment of the present invention, preferably, a patient is informed that edaravone needs to be taken in a fasting state in the morning, a meal at bedtime the night before each administration is not taken, and a meal is not consumed 1 hour after administration. In addition, for postprandial administration, a patient having consumed a high-fat meal (1000 calories (synonymous with 1000 kilocalories as defined in the present specification), 50% fat) is preferably fasted for 8 hours before administration, and a patient having consumed a low-fat meal (standard meal) (400 calories (synonymous with 400 kilocalories as defined in the present specification), 25% fat) is preferably fasted for 4 hours before administration. As an alternative option, a patient may consume a caloric supplement (250 calories (synonymous with 250 kilocalories as defined in the present specification) (e.g., a protein drink)) 2 hours before administration.

[0078]

In administering the pharmaceutical composition according to an embodiment of the present invention, consumption of water is not particularly limited. In oral administration, the pharmaceutical composition according to an embodiment of the present invention when it is solid is, for example, preferably administered together with water, and the pharmaceutical

composition according to an embodiment of the present invention when it is liquid or gel may be administered, for example, without water or together with water. In the intragastric administration via a tube, for example, water is preferably injected from the tube (e.g., the PEG tube or NG tube) to wash in the pharmaceutical composition according to an embodiment of the present invention after administration.

[0079]

The pharmaceutical composition according to an embodiment of the present invention avoids effects of a meal on the pharmacokinetics of edaravone upon oral administration or intragastric administration of edaravone as described above. Thus, the application of the pharmaceutical composition according to an embodiment of the present invention is oral administration or intragastric administration of edaravone, and the target disease and symptom are any of those on which edaravone can act directly or indirectly and not particularly limited. Specifically, the target disease and symptom are diseases or symptoms that can be treated, for example, by scavenging free radicals by edaravone.

[0080]

The disease is, for example, an oxidative stress disease, specifically examples include neurodegenerative diseases with motor dysfunction, such as ALS, Parkinson's disease, and spinocerebellar degeneration; muscle diseases, such as muscular dystrophy; intracerebral neurodegenerative diseases exhibiting cognitive impairment, such as Alzheimer's disease; vascular disorders, such as cerebral infarction; systemic inflammatory diseases, such as

multiple sclerosis and systemic scleroderma; or local inflammatory diseases, such as stomatitis. Of these, the disease is preferably ALS. In an embodiment of the present invention, treatment of the disease includes the meaning of, for example, alleviation of disease progression, cure of the disease, prevention of morbidity, and prevention of recurrence, and may mean treatment for symptoms due to the disease described above. In an embodiment of the present invention, the treatment for symptoms includes the meaning of, for example, suppression of progression of symptoms, alleviation of symptoms, cure of symptoms, prevention of the occurrence of symptoms, and prevention of recurrence of symptoms. The symptom is, for example, a dysfunction in the oxidative stress disease, and specific examples include motor dysfunction and cognitive impairment. In an embodiment of the present invention, the treatment of the disease can be interchangeably read, for example, as treatment of a disease or treatment of a symptom associated with the disease, and the therapeutic agent can be read, for example, as the therapeutic agent for a disease or the therapeutic agent for a symptom associated with the disease (specific examples include progression inhibitors for symptoms). In an embodiment of the present invention, the pharmaceutical preparation according to an embodiment of the present invention may be used, for example, for any one target or two or more targets with regard to the treatment of the disease or the symptom.

[0081]

The method of administering a pharmaceutical composition according to an embodiment of

the present invention can be described as, for example, a method of treating diseases on which edaravone acts directly or indirectly, and a specific example is the method of treating the oxidative stress disease.

[0082]

The administration of the pharmaceutical composition according to an embodiment of the present invention is not particularly limited except for administering orally or administering intragastrically under the condition 1), 2), or 3) as described above, and for example, reference can be made to administration conditions for the use of an injection containing edaravone. Specifically, when the pharmaceutical composition according to an embodiment of the present invention is used for the treatment of an oxidative stress disease (preferably ALS), reference can be made, for example, to the administration method described in WO 2020/091036 or the administration method using an edaravone injection currently used in clinical practice. The administration method may be daily administration or intermittent administration, and specific examples of the administration method include intermittent administration methods used for the treatment of ALS. That is, in the present invention, the first time interval and the second time interval may be, for example, time intervals during an administration period of intermittent administration or daily administration. The intermittent administration method is an intermittent administration, that is, for example, a method of administration in which an administration period and an interruption period are combined as one unit and this unit is repeated two or more times. When an administration period and an

interruption period are combined as one unit and the unit is repeated two or more times, the last period is the interruption period, but the last interruption period may or may not be provided. That is, for example, when the administration period and the interruption period are combined as one unit and the unit is repeated twice, the periods may be combined, for example, as “the administration period, the interruption period, the administration period, and the interruption period” or “the administration period, the interruption period, and the administration period” without providing the last interruption period.

[0083]

In an embodiment of the present invention, the interruption period is, for example, a period in which the drug is not administered for consecutive several days or more and preferably 7 days or 14 days. The administration period is, for example, for 14 days, and in this case, the drug may be administered for consecutive 14 days or in 10 days in 14 days during the administration period. Here, 10 days in 14 days mean, for example, any 10 days in consecutive 14 days, and 10 days when the drug is administered may be consecutive 10 days or nonconsecutive 10 days divided by one or more periods (e.g., 1 day to 4 days) when the drug is not administered. For the administration period, a preferred period can be selected, for example, by observing the patient's condition. More specifically, examples include a method of providing an initial administration period of 14 days, followed by an interruption period of 14 days, and then further repeating an administration period of 10 days in 14 days and an interruption period of 14 days. The number of times of repeating the administration

period of 10 days in 14 days and the interruption period of 14 days is not particularly limited and is, for example, one or more times.

[0084]

The pharmaceutical composition according to an embodiment of the present invention when used for the treatment of an oxidative stress disease (preferably ALS) may be administered to the patient repeatedly, for example, every day or almost every day during the administration period (also referred to as the daily administration). The dose per administration of the pharmaceutical composition according to an embodiment of the present invention can be appropriately selected based on, for example, age and status (e.g., severity of the disease) of the patient. The dose of the pharmaceutical composition according to an embodiment of the present invention can be determined, for example, as a dose of edaravone contained in the pharmaceutical composition. As a typical specific example, for the dose per administration of the pharmaceutical composition according to an embodiment of the present invention in terms of a dose of edaravone for adults, the lower limit is, for example, 60 mg or 90 mg, the upper limit is, for example, 400 mg, 140 mg, 120 mg, or 105 mg, and the range is, for example, from 60 mg to 400 mg, preferably from 90 to 140 mg, more preferably from 90 to 120 mg, and even more preferably from 90 mg to 105 mg, and the dose is particularly preferably 90 mg, 100 mg, 105 mg, or 120 mg, and in particular, preferably 100 mg or 105 mg, and most preferably 105 mg.

[0085]

In addition, when the pharmaceutical composition according to an embodiment of the present invention is used for the treatment of an oxidative stress disease (preferably ALS), for example, the dose per administration during the administration period of an intermittent administration method is the same as the dose described above.

[0086]

For edaravone injection, sold under the brand name “Radicut” (trade name) in Japan, which is currently used as a therapeutic agent for ALS in a clinical setting, the dose per administration thereof is set to 60 mg as edaravone. For the pharmaceutical composition according to an embodiment of the present invention, which is for oral administration or intragastric administration, to attain the same effect as in the intravenous injection of the edaravone injection at 60 mg per administration, the dose of the subject pharmaceutical composition is preferably, for example, set to 90 to 105 mg as edaravone, and more specifically, it is preferably set to 105 mg as edaravone per administration.

[0087]

An administration frequency of the pharmaceutical composition according to an embodiment of the present invention is not particularly limited whether, for example, it is for oral administration or for intragastric administration, and the administration frequency is not particularly limited whether it is for everyday administration or for intermittent administration. The administration frequency can be selected to a preferred frequency, for example, by observing the patient's condition. As a specific example, the number of times of

administration is preferably once or twice per day and more preferably once per day in consideration of, for example, the burden of patients.

Examples

[0088]

Next, the present invention will be specifically described with reference to examples and study examples, but the present invention is not limited to these. Hereinafter, an edaravone oral formulation was used as the pharmaceutical composition according to an embodiment of the present invention, and evaluation using oral administration or intragastric administration via a tube was carried out. The edaravone oral formulation may also be referred to as the edaravone intragastric formulation when used for intragastric administration.

[0089]

Example 1

Effects of meal consumption before administration of edaravone oral formulation on pharmacokinetics of edaravone

[0090]

Effects of meal consumption before administration of the edaravone oral formulation on the pharmacokinetics of edaravone were evaluated. Specifically, clinical studies #1, #2, and #3 described below were conducted on separate days in Japanese healthy adult male subjects. In each of clinical studies #1, #2, and #3, one dose each of edaravone oral formulations (edaravone suspensions) of formulas #1, #2, and #3 shown in Table 1 below were used

respectively and administered orally once during the study period. In addition, the number of subjects in each clinical study is shown in Table 2 described below. For each clinical study, a reference study was conducted (reference example), in which the edaravone oral formulation was administered in fasted conditions as described below, and in each clinical study and the corresponding reference study, the edaravone oral formulation was each administered to the same subjects.

[0091]

[Table 1]

Formulas used in clinical studies #1, #2, and #3

	Study #1 amount	Study #2 amount	Study #3 amount
Edaravone	200 mg	100 mg	105 mg
Polyvinyl alcohol	10 mg	5 mg	5 mg
Xanthan gum		15 mg	15 mg
Sodium hydrogen sulfite		5 mg	5 mg
L-Cysteine hydrochloride hydrate		2.5 mg	2.5 mg
Sodium hydroxide		q.s.	q.s.
Phosphoric acid		q.s.	q.s.
Simethicone emulsion		2.5 mg	2.5 mg
D-Sorbitol		2000 mg	2000 mg
Purified water	q.s.	q.s.	q.s.
Total	10 mL	5 mL	5 mL

[0092]

For the high-fat meal, a meal with total calories per meal of 800 to 1000 kilocalories and calorie intake from lipid of 50% was used. For the standard meal, a meal with total calories per meal of 400 to 500 kilocalories and calorie intake from lipid of 25% was used. For the light meal, a liquid-type nutritional supplement with total calories per meal of 250 kilocalories (enteral nutritional supplement: Ensure Liquid (trade name)) was used.

[0093]

Clinical study #1: comparative example for first time interval condition 1)

The edaravone oral formulation (formula #1) was administered orally to the subjects 30 minutes after high-fat meal consumption. Blood samples were each collected before the administration, and 15 minutes, 30 minutes, 1 hour, 1.5 hours, 2 hours, 4 hours, 6 hours, 8 hours, 12 hours, 24 hours, 36 hours, and 48 hours after the administration, and the concentrations of unchanged edaravone in plasma were measured. On the other hand, as a reference example, the edaravone oral formulation was administered orally to the same subjects in fasted conditions, blood samples were collected, and the concentrations of unchanged edaravone in plasma were measured in the same manner. For the administration in fasted conditions, the edaravone oral formulation was administered orally after a lapse of 10 hours or more from the previous meal consumption (the same applies hereinafter). None of the subjects consumed a next meal until the blood sample collection 4 hours after the administration was completed. The resulting PK profiles are shown in Table 2 below.

[0094]

Clinical study #2: comparative example for first time interval condition 1)

The edaravone oral formulation (formula #2) was administered orally to the subjects 4 hours after high-fat meal consumption. Blood samples were each collected before the administration, and 5 minutes, 15 minutes, 30 minutes, 1 hour, 1.5 hours, 2 hours, 4 hours, 6 hours, 8 hours, 12 hours, 24 hours, 36 hours, and 48 hours after the administration, and the concentrations of unchanged edaravone in plasma were measured. On the other hand, as a reference example, the edaravone oral formulation was administered orally to the same subjects in fasted conditions, blood samples were collected, and the concentrations of unchanged edaravone in plasma were measured in the same manner. For the administration in fasted conditions, the edaravone oral formulation was administered orally after a lapse of 10 hours or longer from the previous meal consumption. None of the subjects consumed a next meal until the blood sample collection 4 hours after the administration was completed. The resulting PK profiles are shown in Table 2 below.

[0095]

Clinical study #3: examples for first time interval conditions 1), 2), and 3), and comparative example for first time interval condition 2)

As examples, the edaravone oral formulation (formula #3) was administered orally to each subject of 8 hours after high-fat meal consumption, 4 hours after standard meal consumption, and 2 hours after light meal consumption. Blood samples were each collected before the administration, and 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 1.5 hours, 2

hours, 4 hours, 6 hours, 8 hours, 10 hours, 12 hours, 24 hours, 36 hours, and 48 hours after the administration, and the concentrations of unchanged edaravone in plasma were measured. In addition, as a comparative example, the edaravone oral formulation was administered orally to the same subjects 2 hours after standard meal consumption, blood samples were collected, and the concentrations of unchanged edaravone in plasma were measured in the same manner. On the other hand, as a reference example, the edaravone oral formulation was administered orally to the same subjects in fasted conditions, blood samples were collected, and the concentrations of unchanged edaravone in plasma were measured in the same manner. For the administration in fasted conditions, the edaravone oral formulation was administered orally after a lapse of 10 hours or longer from the previous meal consumption. None of the subjects consumed a next meal until the blood sample collection 4 hours after the administration was completed. The resulting PK parameters are shown in Table 2 below.

[0096]

[Table 2]

PK parameters of edaravone due to meal consumption before administration of edaravone oral formulation

Study	Meal condition	Edaravone dose (mg)	Statistics	$t_{\max}$ (h)	$C_{\max}$ (ng/mL)	$AUC_{0-\infty}$ (ng•h/mL)	$t_{1/2}$ (h)
Study #1 (n = 6)	Fasted	200	Mean	0.42	4933	6312.88	9.05
			SD	0.13	1268	1246.42	2.37
Study #1 (comparative example) (n = 5)	30 min after high-fat meal	200	Mean	1.35	899 (82%↓)	2466.38 (61%↓)	5.23
			SD	1.56	463.9	824.73	1.71
Study #2 (n = 9)	Fasted	100	Mean	0.36	1810	1646.73	9.33
			SD	0.25	849.8	432.96	4.87
Study #2 (comparative example) (n = 9)	4 h after high-fat meal	100	Mean	0.47	1012 (44% ↓)	1247.4 (24%↓)	7.66
			SD	0.32	603.3	425.22	4.12
Study #3 (n = 16)	Fasted	105	Mean	0.38	2318	2165	8.17
			SD	0.13	1229	673	2.29

Study #3 (example) (n = 16)	8 h after high-fat meal	105	Mean	0.36	2525 (→)	2209 (→)	7.38
			SD	0.19	1337	658	1.97
Study #3 (example) (n = 16)	4 h after standard meal	105	Mean	0.44	2020 (→)	2073 (→)	9.05
			SD	0.19	1114	641	5.07
Study #3 (example) (n = 16)	2 h after light meal	105	Mean	0.41	1898 (→)	1955 (→)	7.31
			SD	0.2	865.9	523	4.3
Study #3 (comparative example) (n = 16)	2 h after standard meal	105	Mean	0.63	1276 (45%↓)	1717 (21%↓)	11.25
			SD	0.32	805.6	463	8.35

[0097]

For the effects of high-fat meal consumption, the following results were obtained. First, a comparison was made between the comparative example of #2 (#2: 4 h after high-fat meal) in which the edaravone oral formulation was administered 4 hours after high-fat meal consumption and the example of #3 (#3: 8 h after high-fat meal) in which the edaravone oral formulation was administered 8 hours after high-fat meal consumption. As shown in Table 2 above, in the comparative example of #2 (#2: 4 h after high-fat meal), as a result of the administration 4 hours after high-fat meal consumption, a decrease in the C_{max} by 44% (percentage of decrease, the same applies hereinafter) and a decrease in the AUC by 24% were observed in comparison with the administration in fasted conditions (#2: Fasted). In contrast to this, as shown in Table 2, in the example of the study #3 (#3: 8 h after high-fat meal), with the administration 8 hours after high-fat meal consumption satisfying the first time interval condition 1) described above, no decrease in the C_{max} and AUC was observed in comparison with the administration in fasted conditions (#3: Fasted). The formulas of the edaravone oral formulations used in the comparative example of #2 and the example of #3 are almost identical, and thus, an embodiment of the present invention was found to be able to maintain the pharmacokinetics of edaravone in a manner similar to the administration in fasted conditions by the timing of administration after meal consumption. In addition, in the comparative example of #1 (#1: 30 min after high-fat meal), the time from meal consumption to the administration of the edaravone oral formulation was further shortened to 30 minutes

after high-fat meal consumption. As a result, a decrease in the C_{max} by 82% and a decrease in the AUC by 61% were observed in comparison with the administration in fasted conditions (#1: Fasted), and the percentages of the decreases were greater than those in the comparative example of #2. The content of edaravone in the edaravone oral formulation of the formula #1 used in the comparative example of #1 was about twice those in the edaravone oral formulations used in the comparative example of #2 and the example of #3. This revealed that the effect of the present invention is not compensated, for example, by the increased edaravone content in the edaravone oral formulation but is obtained by the first time interval from meal consumption to the administration of the edaravone oral formulation according to the type of meal.

[0098]

For the effects of the standard meal consumption and light meal consumption, the following results were obtained. That is, as shown in Table 2 above, both in the example of the study #3 (#3: 4-h after standard meal) with the administration 4 hours after the standard meal consumption satisfying the first time interval condition 2) and in the example of the study #3 (#3: 2-h after standard meal) with the administration 2 hours after the light meal consumption satisfying the first time interval condition 3), no decrease in the C_{max} and AUC was observed in comparison with the administration in fasted conditions (#3: Fasted). No decrease in the C_{max} and AUC was thus observed in the example of the study #3 above, and this was also supported by results of an analysis of variance shown below.

[0099]

That is, from the results of the edaravone PK parameters, an analysis of variance was conducted on the AUC and C_{max} in consideration of the meal condition. This result is shown in Table 3 below. In addition, results of summary statistics for the C_{max} are shown in Table 4 below, and spaghetti plots are shown in FIG. 1. The spaghetti plots in FIG. 1 show results from individual subjects, in which results from the same subjects are connected with lines between results in the administration in fasted conditions and results in the postprandial administration.

[0100]

[Table 3]

Ratios of PK parameters determined by analysis of variance and 90% confidence intervals

Parameter (Unit)	Meal condition	Least-square mean	Ratio [90% Confidence interval] (dose with meal/fasted)
C _{max} (ng/mL)	Fasted	2024	-
	8 hours after high-fat meal	2192	1.083 [0.821-1.429]
	4 hours after standard meal	1764	0.872 [0.661-1.150]
	2 hours after light meal	1659	0.820 [0.621-1.082]
AUC _{0-∞} (ng•h/mL)	Fasted	2072	-
	8 hours after high-fat meal	2123	1.025 [0.931-1.128]
	4 hours after standard meal	1987	0.959 [0.871-1.056]
	2 hours after light meal	1885	0.910 [0.827-1.002]

[Table 4]

Summary statistics for C_{max}

Meal condition	Mean	SD	CV%	Min	Median	Max
Fasted	2318	1229	53	797	1992	4816
8 hours after high-fat meal	2525	1337	52.9	595	2237	5270
4 hours after standard meal	2020	1114	55.2	642	1736	4286
2 hours after light meal	1898	866	45.6	388	1957	3237

[0101]

As shown in Table 3 above, the analysis of variance indicated that for the least squares mean ratio of the AUC of the example administered under each first time interval condition after the meal consumption to the AUC of the reference example administered in fasted conditions (Fasted), the 90% confidence interval was in the range of 0.8 to 1.25, which is a common criterion for equivalence. This result indicates that the edaravone oral formulation administered 8 hours after high-fat meal consumption, 4 hours after standard meal consumption, and 2 hours after light meal consumption can exert an effect of maintaining the pharmacokinetics similar to those in the administration in fasted conditions without effects of meals on the AUC. In addition, as shown in Table 3 above, for the C_{max}, although the 90% confidence intervals of the ratio (numerical values in parentheses in the column of the ratio) did not fall within the range of 0.8 to 1.25, which is the above criterion, because of large coefficients of variation (ratios of SD to the mean value), the point estimates of the ratio (the

value outside the parentheses in the Confidence column) indicated no significant change and were within the range of the above criterion. Furthermore, as shown in the summary statistics of the C_{max} in Table 4 above and the spaghetti plots in FIG. 1, the administration in fasted conditions and the postprandial administration resulted in values of each subject distributed in the range of about 1000 to about 5000 with a median of about 2000 in a similar manner regardless of the conditions of each postprandial administration, showing no change in comparison with the administration in fasted conditions.

[0102]

The example of the study #3 thus indicated that administering the edaravone oral formulation 8 hours after high-fat meal consumption satisfying the condition 1), 4 hours after standard meal consumption satisfying the condition 2), and 2 hours after light meal consumption satisfying the condition 3) can achieve the C_{max} and AUC similar to those in the oral administration in fasted conditions and can avoid effects of the meal on the pharmacokinetics of edaravone.

[0103]

As described above, the examples confirmed that administering after meal consumption in the timing of the first time interval conditions 1), 2), and 3) can maintain pharmacokinetics similar to those in administering the same edaravone oral formulation in fasted conditions. In addition, these results supported that the pharmaceutical composition according to an embodiment of the present invention is any composition containing edaravone as an active

ingredient, the composition itself is not particularly limited, and even in using the pharmaceutical composition of the same composition, the effect of the present invention can be obtained by setting conditions for administration as described above according to the type of meal.

[0104]

Example 2

Effects of meal consumption after administration of edaravone oral formulation on pharmacokinetics of edaravone 1

[0105]

Effects of meal consumption 1 hour after administration of the edaravone oral formulation on the pharmacokinetics of edaravone were evaluated. Specifically, a clinical study #4 described below was conducted on the same subjects in Japanese healthy adult male subjects. In the clinical study #4, the edaravone oral formulation of the formula #2 in Table 1 above was used and administered orally once during the study period. In addition, the same high-fat meal as that used in Example 1 above was used.

[0106]

Clinical study #4

As a post-administration consumption group, the same subjects in fasted conditions after a lapse of 10 hours or longer from meal consumption were orally administered with the edaravone oral formulation, and the high-fat meal was consumed 1 hour after the

administration. Blood samples were each collected before the administration, and 5 minutes, 15 minutes, 30 minutes, 1 hour, 1.5 hours, 2 hours, 4 hours, 6 hours, 8 hours, 12 hours, 24 hours, 36 hours, and 48 hours after the administration, and the concentrations of unchanged edaravone in plasma were measured. The consumption of the high-fat meal 1 hour after the administration was carried out after the blood sample collection 1 hour after the administration, and a subsequent meal was not consumed until the blood sample collection 4 hours after the administration was completed. On the other hand, as a fasted group for reference, the edaravone oral formulation was administered orally to the same subjects in fasted conditions, blood samples were collected, and the concentrations of unchanged edaravone in plasma were measured in the same manner. The subjects in the fasted group did not consume a meal either until the blood sample collection 4 hours after the administration was completed. The resulting PK parameters are shown in Table 5 below.

[0107]

[Table 5]

PK parameters of edaravone due to meal consumption after administration of edaravone oral formulation

Study	Meal condition	Edaravone dose (mg)	Statistics	$t_{\max}$ (h)	$C_{\max}$ (ng/mL)	$AUC_{0-\infty}$ (ng•h/mL)	$t_{1/2}$ (h)
Study #4 (n = 9)	Fasted	100	Mean	0.36	1810	1646.73	9.33
			SD	0.25	849.8	432.96	4.87
Study #4 (n = 9)	1 h before high-fat meal	100	Mean	0.6	1502 (→)	1475.31 (→)	9.65
			SD	0.44	1272	658.1	5.41

[0108]

A comparison was made between the post-administration consumption group, in which the high-fat meal was consumed 1 hour after the administration of the edaravone oral formulation, and the fasted group, in which no meal was consumed for another 4 hours or longer after the administration in fasted conditions. As a result, as shown in Table 5 above, the post-administration consumption group showed only a slight decrease in the C_{max} by about 17.0% and the AUC by about 10.4% in comparison with the fasted group, and no significant difference was observed. Thus, the meal consumption 1 hour after the oral administration was found to have no effect on the pharmacokinetics of edaravone.

[0109]

In addition, the $AUC_{0-\infty}$ after the administration of the edaravone oral formulation was evaluated separately on the results from 0 hour to 1 hour after the administration (0 to 1 h, before the consumption of a meal resumed after the administration) (AUC_{0-1h}) and the results 1 hour after the administration (1 h to ∞ h, after the consumption of a meal resumed after the administration) ($AUC_{1h-\infty}$) for the post-administration consumption group (1 h before high-fat meal) and the fasted group (Fasted). These results are shown in Table 6 below and spaghetti plots in FIG. 2. As shown in Table 6 below and FIG. 2, in the post-administration consumption group (1 h before high-fat meal), although a slight decrease in the AUC_{0-1h} occurred from the edaravone administration to 1 hour before the consumption of a meal resumed after the administration in comparison with the administration in the fasted group

(Fasted), no change in the $AUC_{1h-\infty}$ was noted 1 hour and longer after the edaravone administration. From this, the result of the slight decrease in the AUC_{0-1h} in the post-administration consumption group is not considered due to decreases in the C_{max} and AUC due to effects of the meal but is considered to be a result due to fluctuations in the same subjects (i.e., day-to-day fluctuations due to acquiring the data of fasted conditions and the meal consumption from the same subjects on separate days).

[0110]

[Table 6]

AUC of edaravone when high-fat meal is consumed 1 hour after administration of edaravone oral formulation

	Meal condition	AUC_{0-1h} (ng•h/mL)	$AUC_{1h-\infty}$ (ng•h/mL)
Study #4 (n = 9)	Fasted	890 ± 349	756 ± 232
Study #4 (n = 9)	1 h before high-fat meal	772 ± 519	703 ± 202

[0111]

The above results indicated that when the high-fat meal was consumed 1 hour after the administration of the edaravone oral formulation, no change in the pharmacokinetics of edaravone was noted in comparison with the administration in fasted conditions, and thus a meal after the administration has no effect on the pharmacokinetics. The results of the

evaluation using the high-fat meal thus indicated no effect on the pharmacokinetics of edaravone. Thus, regardless of the high-fat meal, consumption of the standard meal and the light meal 1 hour after the administration of the edaravone oral formulation is considered to cause no change in the pharmacokinetics of edaravone in comparison with the administration in fasted conditions, and the meal is considered to have no effect on the pharmacokinetics.

[0112]

Example 3

Evaluation of edaravone in Caco-2 cell membrane permeability test

[0113]

Caco-2 cells were seeded on a 24-well plate, cultured at 37°C for 21 to 23 days in a CO₂ incubator, and then a Caco-2 cell monolayer membrane permeability test was conducted. After checking the membrane resistance value of the cell layer after the culture, the culture medium in the plate was replaced with HBSS (pH 7.4). After preincubation at 37°C for 1 hour, edaravone and control compounds at predetermined concentrations were each added to the apical side and incubated at 37°C for 2 hours. For the control compounds, [³H]propranolol as a high membrane permeability control and [¹⁴C]mannitol as a low membrane permeability control were used. Then, the concentrations of edaravone and the control compounds on the basal side were each measured, and the permeability coefficients (P_{app}) were calculated. These results are shown in Table 7 below. The data of P_{app} are indicated by mean ± SD. As shown in Table 7 below, edaravone exhibited membrane

permeability equal to or higher than that of propranolol used as the high membrane

permeability control compound.

[0114]

[Table 7]

Permeability coefficient of edaravone

Compound	Concentration ($\mu\text{mol/L}$)	P_{app} ($\times 10^{-6}$ cm/sec) Apical to basal (n = 3)
Edaravone	100	32.0 ± 1.7
[^3H]Propranolol	10	28.9 ± 1.1
[^{14}C]Mannitol	10	0.530 ± 0.007

[0115]

Cell membrane permeability is known to be constant regardless of the concentration of the substance when no transporter is involved. The control compounds, propranolol and mannitol, are not substrates for transporters P-gp and BCRP expressed in Caco-2 cells, and thus the values of the permeability coefficient are not considered to change depending on the concentrations used in the test. Thus, further, the following study was conducted to confirm that edaravone is not a substrate for transporters P-gp and BCRP as well.

[0116]

That is, Caco-2 cells were seeded on a 96-well plate, cultured at 37°C for 10 to 11 days in a CO₂ incubator, and then a transcellular transport study was conducted. Then, a ratio of P_{app} from the apical side to the basal side to P_{app} from the basal side to the apical side was

calculated. The efflux ratio was calculated as the ratio of the mean values of P_{app} calculated from three measurements. This result is shown in Table 8 below. As shown in Table 8 below, the efflux ratio of edaravone did not exceed 2 and was not affected by addition of typical inhibitors of P-gp and/or BCRP. Thus, this indicated that edaravone is not a substrate for P-gp and BCRP.

[0117]

[Table 8]

Ratios of permeability coefficient of edaravone in Caco-2 cells in both directions in presence or absence of typical inhibitors of P-gp and BCRP

Compound (Added concentration)	Inhibitor (Added concentration)	Efflux ratio
Edaravone (10 $\mu\text{mol/L}$)	-	0.9
	Quinidine (30 $\mu\text{mol/L}$)	1
	Novobiocin (30 $\mu\text{mol/L}$)	0.8
	Quinidine (30 $\mu\text{mol/L}$) + Novobiocin (30 $\mu\text{mol/L}$)	0.6

Quinidine: P-gp inhibitor

Novobiocin: BCRP inhibitor

[0118]

Literatures by Sun et al. (Pharmaceutical Research, Vol. 19, No. 10, 2002) and Yamashita et al. (European Journal of Pharmaceutical Sciences 10 (2000) 195-204) indicate that the

permeability coefficient of a drug by the Caco-2 cell membrane permeability test correlates with the absorption rate of the drug in humans. In addition, literatures by Paterson et al. (Pharmacologia Clinica 2, 127-133 (1970)) and Borgstrom et al. (Journal of Pharmacokinetics and Biopharmaceutics, Vol 9 No. 4, 1981) indicate that when administered orally to humans, almost all of propranolol is absorbed in the intestine.

[0119]

As described above, edaravone exhibited a permeability coefficient equal to or higher than that of propranolol. Thus, edaravone when administered orally to humans was considered to exhibit high absorption rate and was found to be classified as a high permeability compound. The report by Wu et al. (Non-Patent Document 1) describes that high permeability compounds are less susceptible to effects of a meal on pharmacokinetics as described above. On the contrary, however, despite the fact that edaravone has a high permeability, its pharmacokinetics was affected by a meal as also described in Examples 1 and 2 above. The pharmacokinetics of edaravone is thus affected by a meal; further, the effect varies with the type of meal; and the effect on the pharmacokinetics can be avoided by setting the timing of administration to the conditions 1), 2), and 3) described above according to the type of meal. These have been found by the present inventors for the first time.

[0120]

Example 4

Pharmacokinetics of edaravone in ALS patients

[0121]

To evaluate the pharmacokinetics of edaravone in ALS patients, a clinical study #5 shown below was conducted in ALS patients who can live daily life independently (the patient group).

[0122]

Clinical study #5

The edaravone oral formulation was administered orally to the patient group in fasted conditions after a lapse of 10 hours or longer from consumption of a meal. In the patient group, ALS patients who had been already administered with an edaravone intravenous infusion formulation were orally administered with the edaravone oral formulation after a lapse of 48 hours or longer from the time of completion of the previous administration of the intravenous infusion formulation. The same edaravone oral formulation as the formula #3 in Table 1 above was used. Then, blood samples were each collected before the administration, and 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 1.5 hours, 2 hours, 4 hours, 6 hours, 8 hours, 10 hours, 12 hours, and 24 hours after the administration, and the concentrations of unchanged edaravone in plasma were measured. The patient group did not consume a meal after the administration until the blood sample collection 4 hours after the administration was completed.

[0123]

Clinical study #9

On the other hand, a clinical study #9 shown below was conducted in healthy adult male and female subjects (42 in total) (the reference group). That is, the edaravone oral formulation of the formula #3 was administered orally to the reference group in fasted conditions after a lapse of 10 hours or longer. Then, blood samples were each collected before the administration, and 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 1.5 hours, 2 hours, 4 hours, 6 hours, 8 hours, 10 hours, 12 hours, 24 hours, 36 hours, and 48 hours after the administration, and the concentrations of unchanged edaravone in plasma were measured. The reference group did not consume a meal after the administration either until the blood sample collection 4 hours after the administration was completed.

[0124]

The resulting PK parameters were corrected for body weight of each subject adjusted to 70 kg by a proportional calculation for each of the patient group and the reference group. The results of the PK parameters before correction of the patient group and the reference group are shown in Table 9 below. In addition, the results of the PK parameters after correction of the patient group are shown as box-whisker plots of #9 (the reference group), and individual plots of #5 (the patient group) and #9 (the reference group) in FIG. 3. In FIG. 3, X represents a mean value. As shown in Table 9 below, the difference in the C_{max} before weight correction was 15%, and the difference in the AUC before weight correction was 1.5% between the patient group and the reference group, but as shown in FIG. 3, the difference in the C_{max} after weight correction was 9.8%, and the difference in the AUC after weight

correction was 5.8%, both being within 10%. On the other hand, the mean value $\pm$ standard error of the C_{max} after weight correction of the patient group was 1607 $\pm$ 231, and the mean value $\pm$ standard error of the AUC after weight correction of the patient group was 1461 $\pm$ 166, both with a large standard error. Thus, the difference in the C_{max} and the difference in the AUC before weight correction between the patient group and the reference group seen in Table 9 below are considered due to the difference in body weight between the patient group and the reference group, and the difference in the C_{max} and the difference in the AUC after weight correction between the patient group and the reference group observed in FIG. 3 are each considered to be within the range of error.

[0125]

[Table 9]

PK Parameters of edaravone in ALS patients

subjects	Edaravone dose (mg)	Statistics	t _{max} (h)	AUC _{inf} (ng•h/mL)	C _{max} (ng/mL)
healthy (n = 42)	105	Mean	0.44	1762	1656
		SD	0.17	540	734
ALS patients (n = 9)	105	Mean	0.31	1736	1903
		SD	0.15	811	978

[0126]

As described above, the result of comparing the C_{max} and AUC of edaravone after the

administration of the edaravone oral formulation between ALS patients and healthy adults revealed that the pharmacokinetics of edaravone is similar between ALS patients and healthy adults and that sufficient plasma concentration of edaravone is maintained in ALS patients as well as in healthy adults. From this, it can be said that oral administration of edaravone itself does not affect the plasma concentration regardless of whether the subject is a patient.

[0127]

Example 5

Pharmacokinetics of edaravone in administration via percutaneous endoscopic gastrostomy tube (PEG tube)

[0128]

Pharmacokinetics of edaravone in administration of the edaravone oral formulation via a PEG tube was evaluated. Specifically, a clinical study #6 described below was conducted in ALS patients with a percutaneous endoscopic gastrostomy tube (PEG tube) (the patient group). In the clinical study #6, the edaravone oral formulation of the formula #3 shown in Table 1 was used.

[0129]

Clinical study #6

The edaravone oral formulation was administered via a PEG tube to the patient group in fasted conditions after a lapse of 10 hours or longer from consumption of a meal. That is, first, 100 mL of lukewarm water was injected through a PEG tube 1 hour before the

administration of the edaravone oral formulation, and 10 mL of lukewarm water was injected through a PEG tube immediately before the administration. Then, the edaravone oral formulation was administered to the PEG tube, and further 10 mL of water was injected through the PEG tube three times to wash in the edaravone oral formulation. In the patient group, ALS patients who had been already administered with an edaravone intravenous infusion formulation were administered with the edaravone oral formulation via a PEG tube after a lapse of 48 hours or longer from the time of completion of the previous administration of the intravenous infusion formulation. Blood samples were each collected before the administration, and 15 minutes, 30 minutes, 1 hour, 2 hours, 4 hours, 8 hours, and 24 hours after the administration, and the concentrations of unchanged edaravone in plasma were measured. The patient group did not consume a meal until the blood sample collection 4 hours after the administration via a PEG tube was completed. For the reference example, the results of the reference group in Example 4, in which the edaravone oral formulation of the same formula #3 was administered orally and blood samples were collected, were utilized.

[0130]

The resulting PK parameters were corrected for body weight of each subject in the patient group adjusted to 70 kg by a proportional calculation. The results of the PK parameters before correction of the patient group are shown in Table 10 below. In addition, the results of the PK parameters after correction of the patient group are shown in individual plots as #6 in FIG. 3. A comparison was made between the patient group administered via a PEG tube in

Table 10 below and the reference group administered orally in Table 9 above. As a result, a difference in the C_{max} before weight correction of 30% and a difference in the AUC before weight correction of 33% were observed between the patient group administered via a PEG tube and the reference group administered orally, but as shown in FIG. 3, the difference in the C_{max} after weight correction was 8.3%, and the difference in the AUC after weight correction was 9.5%, both being within 10%. On the other hand, as shown in FIG. 3, the mean value $\pm$ standard error of the C_{max} after weight correction of the patient group administered via a PEG tube was 1585 ± 251 , and the mean value $\pm$ standard error of the AUC after weight correction of the patient group was 1698 ± 228 , both with a large standard error. Thus, the difference in the C_{max} and the difference in the AUC before weight correction between the patient group administered via a PEG tube and the reference group administered orally seen in Table 10 below are considered due to the difference in body weight between the patient group administered via a PEG tube and the reference group administered orally, and the difference in the C_{max} and the difference in the AUC after weight correction between the patient group administered via a PEG tube and the reference group administered orally seen in FIG. 3 are each considered to be within the range of error.

[0131]

[Table 10]

PK Parameters of edaravone in ALS patients in administration via a PEG-tube

Edaravone dose	Statistics	t _{max} (h)	AUC _{inf} (ng•h/mL)	C _{max} (ng/mL)
-------------------	------------	-------------------------	---------------------------------	-----------------------------

(mg)				
105	Mean	0.33	2344	2163
	SD	0.13	971	902

[0132]

The results of the oral administration in Example 4 above and the results of the administration via a PEG tube in this Example 5 indicated that the C_{max} and AUC of edaravone after administration of the edaravone oral formulation via a PEG tube were similar to the results after the oral administration of the same edaravone oral formulation. This suggests no difference in the pharmacokinetics of edaravone between administration via a PEG tube and oral administration and revealed that sufficient plasma concentration of edaravone is maintained in administration via a PEG tube as well as in oral administration.

[0133]

Example 6

Dose of edaravone in oral formulation exhibiting bioavailability equivalent to that of edaravone intravenous administration

[0134]

The dose of edaravone in the oral formulation exhibiting bioavailability equivalent to that of the 60-mg edaravone intravenous (IV) administration was investigated. Specifically, clinical studies #7 and #8 described below were conducted in healthy adult male subjects for each subject group. In clinical studies #7 and #8, one dose each of edaravone oral formulations

(edaravone suspensions) of formulas #7 and #8 shown in Table 11 below were used

respectively and administered orally once during the study period.

[0135]

[Table 11]

Formulas used in studies #7 and #8

	Study #7					Study #8
	amount					amount
Edaravone	30 mg	60 mg	120 mg	200 mg	300 mg	100 mg
Polyvinyl alcohol	50 mg	50 mg	10 mg	10 mg	10 mg	5 mg
Xanthan gum						15 mg
Sodium hydrogen sulfite						5 mg
L-Cysteine hydrochloride hydrate						2.5 mg
Sodium hydroxide						q.s.
Phosphoric acid						q.s.
Simethicone emulsion						2.5 mg
D-Sorbitol						2000 mg
Purified water	q.s.	q.s.	q.s.	q.s.	q.s.	q.s.
Total	50 mL	50 mL	10 mL	10 mL	10 mL	5 mL

[0136]

Clinical study #7

Any of the edaravone oral formulations (formula #7) with an edaravone content of 30 mg, 60 mg, 120 mg, 200 mg, or 300 mg was administered orally to subjects in fasted conditions after

a lapse of 10 hours or longer from consumption of a meal. The edaravone oral formulations with each content were each administered orally to 6 subjects. Blood samples were each collected before the administration, and 15 minutes, 30 minutes, 1 hour, 1.5 hours, 2 hours, 4 hours, 6 hours, 8 hours, 12 hours, 24 hours, 36 hours, and 48 hours after the administration, the concentrations of unchanged edaravone were measured, and the AUCs were calculated. The subjects did not consume a meal after the administration until the blood sample collection 4 hours after the administration was completed.

[0137]

Clinical study #8

The edaravone oral formulations (formula #8) with an edaravone content of 100 mg was administered orally to 9 subjects in fasted conditions after a lapse of 10 hours or longer from consumption of a meal. Blood samples were each collected before the administration, and 5 minutes, 15 minutes, 30 minutes, 1 hour, 1.5 hours, 2 hours, 4 hours, 6 hours, 8 hours, 12 hours, 24 hours, 36 hours, and 48 hours after the administration, the concentrations of unchanged edaravone in plasma were measured, and the AUCs were calculated. The subjects did not consume a meal after the administration either until the blood sample collection 4 hours after the administration was completed.

[0138]

These results are shown in FIG. 4 as an individual value plot of the doses of edaravone and the AUC and a 4-parameter logistic model indicating the relationship between both. In the

figure, the horizontal axis is the dose of edaravone and the vertical axis is the AUC. In addition, the value of the AUC (1738 ng•h/mL) of the approved 60-mg edaravone intravenous administration formulation is shown in FIG. 4 with a dotted line. As shown in FIG. 4, the amount of edaravone indicating an AUC similar to the AUC of the 60-mg edaravone intravenous administration formulation was calculated to be within the range of 100 to 105 mg. From this result, the edaravone oral formulation (including the meaning of the edaravone intragastric formulation) maintains an effect similar to that of the approved 60-mg edaravone intravenous administration formulation, for example, by setting the amount of edaravone contained in one administration to 100 to 105 mg and administering orally or via a PEG tube with the timing of administration in an embodiment of the present invention described above and can provide medical care with a reduced burden on patients and caregivers.

[0139]

Example 7

Effects of meal consumption after administration of edaravone oral formulation on pharmacokinetics of edaravone 2

[0140]

Effects of meal consumption 30 minutes after administration of the edaravone oral formulation on the pharmacokinetics of edaravone were evaluated. Specifically, a clinical study #10 described below was conducted in Japanese healthy adult male subjects. In the

clinical study #10, the edaravone oral formulation of the formula #7 (edaravone content of 120 mg) in Table 11 above was used and administered orally once during the study period. In addition, the same standard meal as that used in Example 1 above was used.

[0141]

Clinical study #10

As a post-administration consumption group, subjects in fasted conditions after a lapse of 10 hours or longer from meal consumption were orally administered with the edaravone oral formulation, and the standard meal was consumed 30 minutes after the administration. Blood samples were each collected before the administration, and 15 minutes, 30 minutes, 1 hour, 1.5 hours, 2 hours, 4 hours, 6 hours, 8 hours, 12 hours, and 24 hours after the administration, and the concentrations of unchanged edaravone in plasma were measured. The consumption of the standard meal 30 minutes after the administration was carried out after the blood sample collection 30 minutes after the administration, and a subsequent meal was not consumed until the blood sample collection 4 hours after the administration was completed.

[0142]

On the other hand, as a fasted group for reference, the edaravone oral formulation was administered orally to different subjects in fasted conditions, blood samples were collected, and the concentrations of unchanged edaravone in plasma were measured in the same manner. The subjects in the fasted group did not consume a meal either until the blood sample collection 4 hours after the administration was completed. The resulting PK

parameters are shown in Table 12 below.

[0143]

[Table 12]

PK parameters of edaravone due to meal consumption after administration of edaravone oral formulation

Dose (mg)	Meal condition	Statistics	$t_{\max}$ (h)	$C_{\max}$ (ng/mL)	AUC_{0-24h} (ng•h/mL)	$t_{1/2}$ (h)
120	Fasted (n = 6)	Mean	0.38	1735	2241.77	5.13
		SD	0.14	738	910.63	1.46
	30 min before low-fat meal (n = 6)	Mean	0.42	1953	1916.91	4.6
		SD	0.13	838.3	383.34	0.41

[0144]

A comparison was made between the post-administration consumption group, in which the standard meal was consumed 30 minutes after the administration of the edaravone oral formulation, and the fasted group, in which no meal was consumed for another 4 hours or longer after the administration in fasted conditions. As a result, as shown in Table 12 above, the post-administration consumption group indicated only a slight decrease in the $C_{\max}$ by about 12.6% and the AUC by about 14.5% in comparison with the fasted group, and no significant difference was observed. Thus, the meal consumption 30 minutes after the oral administration was found to have no effect on the pharmacokinetics of edaravone.

[0145]

From the above results, when the standard meal was consumed 30 minutes after the administration of the edaravone oral formulation, no significant change was considered to occur in comparison with the fasted conditions. In addition, edaravone was administered as a suspension and thus is presumed to have been rapidly excreted from the stomach, and most of edaravone is presumed to have reached the upper part of the small intestine, which is considered to be the absorption site, by 30 minutes after the administration. Thus, a difference in the type of meal consumed 30 minutes after the administration is considered to have no effect on edaravone. This suggested that in oral administration of edaravone, edaravone is not affected by a meal after a lapse of 30 minutes from the administration regardless of the type of meal after the administration.

[0146]

Example 8

Pharmacokinetics of edaravone in administration via percutaneous endoscopic gastrostomy tube (PEG tube) 2

Pharmacokinetics of edaravone in administration of the edaravone oral formulation via a PEG tube was evaluated in 36 healthy adults. Specifically, performing gastrostomy in healthy adults is difficult, thus a nasopharyngeal gastric tube (NG tube), which allows the administration into the stomach via a tube in the same manner, was used as a model of administration via a percutaneous endoscopic gastrostomy tube (PEG tube), and a clinical study #11 described below was conducted. In the clinical study #11, the edaravone oral

formulation of the formula #3 shown in Table 1 was used.

[0147]

Clinical study #11

The edaravone oral formulation (#3) was administered via a nasopharyngeal gastric tube (NG tube) to the subject group in fasted conditions after a lapse of 10 hours or longer from consumption of a meal. In the administration via an NG tube, first, 100 mL of lukewarm water was injected through an NG tube 1 hour before the administration of the edaravone oral formulation, and 10 mL of lukewarm water was injected through an NG tube immediately before the administration. Then, the edaravone oral formulation was administered to the NG tube, and further 30 mL of water was injected through the NG tube to wash in the edaravone oral formulation. Blood samples were each collected before the administration, and 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 1.5 hours, 2 hours, 4 hours, 6 hours, 8 hours, 10 hours, 12 hours, 24 hours, 36 hours, and 48 hours after the administration, and the concentrations of unchanged edaravone in plasma were measured. As a reference example, the same edaravone oral formulation was administered orally to the same subjects, blood samples were collected, and the concentrations of unchanged edaravone in plasma were measured in the same manner. For the oral administration, the edaravone oral formulation was administered orally after a lapse of 10 hours or longer from the previous meal consumption. None of the subjects consumed a next meal until the blood sample collection 4 hours after the administration was completed. The resulting PK parameters are shown in

Table 13 below. In addition, results of an analysis of variance conducted on the AUC and C_{max} in consideration of the meal condition are shown in Table 14 below.

[0148]

[Table 13]

PK parameters of edaravone after oral administration and administration via NG tube of edaravone oral formulation

Administration route	PK parameter	t _{max} ^[a] (h)	C _{max} (ng/mL)	AUC _{0-t} (ng•h/mL)	AUC _{0-∞} (ng•h/mL)	t _{1/2} (h)
Oral (N = 36)	Mean	0.5	2470	2612	2657	21.52
	CV%	0.250, 0.500	34	30.1	30.1	157.2
Nasopharyngeal gastric tube (N = 36)	Mean	0.25	2775	2592	2617	13.53
	CV%	0.083, 1.000	52.1	33.4	33.2	102.1

[a]: median, minimum, and maximum

[0149]

[Table 14]

Ratios of PK parameters determined by analysis of variance and 90% confidence intervals

Parameter	Units	Least-square mean	Ratio	90% Confidence interval
-----------	-------	-------------------	-------	-------------------------

		Nasopharyngeal gastric tube	Oral		Lower	Upper
C _{max}	ng/mL	2431	2310	1.052	0.9028	1.2267
AUC _{0-∞}	ng•h/mL	2501	2551	0.981	0.9307	1.0333

[0150]

As shown in Table 13, each PK parameter was very similar regardless of the route of administration, indicating that pharmacokinetics is maintained regardless of the route of administration. As shown in Table 14 above, the analysis of variance indicated that for the least squares mean ratio of the AUC and C_{max} of the example administered via an NG tube to the AUC and C_{max} of the reference example administered orally, the 90% confidence interval was in the range of 0.8 to 1.25, which is a common criterion for equivalence. This result shows that administration of edaravone into the stomach via an NG tube does not affect the pharmacokinetic parameters AUC and C_{max} and can exert an effect of maintaining the pharmacokinetics similar to those in oral administration.

[0151]

The results of the oral administration in this Example 8 and the results of the administration via a PEG tube in the above Example 5 indicated that the C_{max} and AUC of edaravone after intragastric administration of the edaravone oral formulation via an NG tube were equivalent to the results after the oral administration of the same edaravone oral formulation. This suggests no difference in the pharmacokinetics of edaravone between intragastric

administration via a catheter and oral administration and revealed that sufficient plasma concentration of edaravone is maintained in intragastric administration via a catheter as well as in oral administration. This result with more subjects supports the finding in Example 5 that plasma concentration of edaravone administered via a PEG tube to ALS patients is maintained in a manner similar to the oral administration.

[0152]

Example 9

Investigation of adsorption of edaravone to catheter in vitro

In administration using a catheter, catheters of various materials are used in medical facilities. Thus, whether administration is feasible with a catheter of any material was investigated by evaluating adsorption properties of catheters. That is, the edaravone oral formulation (#3) was passed through a catheter, and then the percentage of edaravone collected by washing in with water was evaluated. The types of percutaneous endoscopic gastrostomy tubes (PEG tubes) Nos. 1 to 3 used for the evaluation are shown in Table 15.

[0153]

[Table 15]

Types of PEG tubes used for adsorption evaluation

	No. 1	No. 2	No. 3
Material	Button: silicone Tubing: polyvinyl chloride	Button: polyurethane Tubing: polyvinyl chloride	Balloon: silicone rubber Tubing: silicone rubber/polyacetal

Type	Bumps/button	Bumps/button	Balloon/button
Size	<u>Bump</u>	<u>Bump</u>	<u>Balloon</u>
	Length: 5.0 cm	Length: 5.5 cm	Length: 4.4 cm
	Diameter: 24 Fr	Diameter: 24 Fr	Diameter: 24 Fr
	<u>Tubing</u>	<u>Tubing</u>	<u>Tubing</u>
	Length: 60 cm	Length: 30 cm	Length: 60 cm
	Diameter: outer 6.0 mm	Diameter: outer 5.5 mm	Diameter: outer 5.5 mm
	inner 4.0 mm	inner 3.7 mm	inner 3.5 mm

[0154]

Specifically, the edaravone oral formulation was loaded into a syringe for catheter administration to give an edaravone amount of 105 mg, and the edaravone oral formulation was passed from the syringe to each of the catheters. Water at a predetermined temperature was then injected into the catheter to wash in the edaravone oral formulation. The edaravone oral formulation that passed through the catheter and the wash-in liquid were collected and mixed, and an amount of edaravone contained in the mixed liquid was measured. The edaravone oral formulation was washed in under any of the following conditions: with 10 mL of water three times, with 30 mL of water at 25°C or 40°C once, or with 50 mL of water at 25°C or 40°C once. Three measurements were made for each catheter, and the mean and SD of recovery% were calculated. These results are shown in Table 16.

[0155]

[Table 16]

Adsorption evaluation

Wash-in conditions	Type of PEG tube		
	No. 1	No. 2	No. 3
10 mL 3 times	98.0% (0.6)	100.0% (1.5)	99.1% (0.6)
30 mL once (Water temperature at 25°C)	97.1% (1.5)	98.6% (1.7)	97.9% (1.3)
30 mL once (Water temperature at 40°C)	101.9% (2.1)	104.1% (0.7)	103.0% (0.7)
50 mL once (Water temperature at 25°C)	96.5% (1.8)	97.0% (0.9)	97.5% (0.4)
50 mL once (Water temperature at 40°C)	97.6% (0.8)	97.9% (0.5)	95.5% (0.6)

Shown as mean (SD) of N = 3

[0156]

As shown in Table 16 above, almost all amount of edaravone was recovered by each specified washing in with water regardless of the material of the catheter used. No adsorption of edaravone was thus observed on the catheters regardless of the material, thus showing that transcatheter administration of the edaravone oral formulation is feasible regardless of the material of the catheter.

[0157]

The present invention has been described above with reference to embodiments and examples, but the present invention is not limited to the above embodiments. Various

changes that can be understood by those skilled in the art can be made to the constitution and details of the present invention within the scope of the present invention.

[0158]

The present application claims priority based on JP 2020-188514, filed on November 12, 2020, the entire disclosure of which is incorporated herein.

Industrial Applicability

[0159]

The pharmaceutical composition containing edaravone of the present invention can avoid effects of meal consumption on the pharmacokinetics by administration with an optimal administration interval for each type of meal and can exert excellent effects as the oral formulation or intragastric formulation of edaravone, and thus is useful, for example, for the treatment of oxidative stress diseases.

Claims

[Claim 1]

A pharmaceutical composition containing edaravone as an active ingredient, the pharmaceutical composition being administered to a subject orally or intragastrically with any one first time interval selected from the group consisting of 1) to 3):

1) in a case where the subject has consumed a high-fat meal, 8 hours or longer after the consumption;

2) in a case where the subject has consumed a standard meal, 4 hours or longer after the consumption; and

3) in a case where the subject has consumed a light meal, 2 hours or longer after the consumption.

[Claim 2]

The pharmaceutical composition according to claim 1, wherein a second time interval from edaravone administration to next meal consumption is 30 minutes or longer.

[Claim 3]

The pharmaceutical composition according to claim 1 or 2, wherein the second time interval from edaravone administration to next meal consumption is 1 hour or longer.

[Claim 4]

The pharmaceutical composition according to any one of claims 1 to 3, which is for oral administration.

[Claim 5]

The pharmaceutical composition according to any one of claims 1 to 3, which is for intragastric administration.

[Claim 6]

The pharmaceutical composition according to any one of claims 1 to 5, wherein the intragastric administration is administration via a tube.

[Claim 7]

The pharmaceutical composition according to claim 6, wherein the tube is a percutaneous endoscopic gastrostomy tube (PEG tube).

[Claim 8]

The pharmaceutical composition according to claim 6, wherein the tube is a nasopharyngeal gastric tube (NG tube).

[Claim 9]

The pharmaceutical composition according to any one of claims 1 to 8, which is for treating an oxidative stress disease.

[Claim 10]

The pharmaceutical composition according to claim 9, wherein the oxidative stress disease is any one of diseases selected from the group consisting of amyotrophic lateral sclerosis, Parkinson's disease, spinocerebellar degeneration, muscular dystrophy, Alzheimer's disease, cerebral infarction, multiple sclerosis, systemic scleroderma, and stomatitis.

[Claim 11]

The pharmaceutical composition according to claim 9 or 10, wherein the oxidative stress disease is amyotrophic lateral sclerosis.

[Claim 12]

The pharmaceutical composition according to any one of claims 1 to 11, wherein the first time interval is a time interval during an administration period of intermittent administration or daily administration.

[Claim 13]

The pharmaceutical composition according to any one of claims 1 to 12, wherein a dose per administration is from 90 to 120 mg of edaravone.

[Claim 14]

The pharmaceutical composition according to any one of claims 1 to 12, wherein a dose per administration is from 90 to 105 mg of edaravone.

[Claim 15]

The pharmaceutical composition according to any one of claims 1 to 14, wherein a daily administration frequency is once, and a dose per administration is 105 mg of edaravone.

[Claim 16]

A method of administering an edaravone-containing pharmaceutical composition which contains edaravone as an active ingredient, the method comprising administering the pharmaceutical composition to a subject orally or intragastrically with any one first time

interval selected from the group consisting of 1) to 3):

1) in a case where the subject has consumed a high-fat meal, 8 hours or longer after the consumption;

2) in a case where the subject has consumed a standard meal, 4 hours or longer after the consumption; and

3) in a case where the subject has consumed a light meal, 2 hours or longer after the consumption.

[Claim 17]

The method according to claim 16, wherein a second time interval from administration of the pharmaceutical composition to next meal consumption is 30 minutes or longer.

[Claim 18]

The method according to claim 16 or 17, wherein the second time interval from administration of the pharmaceutical composition to next meal consumption is 1 hour or longer.

[Claim 19]

The method according to any one of claims 16 to 18, wherein the pharmaceutical composition is administered orally.

[Claim 20]

The method according to any one of claims 16 to 18, wherein the pharmaceutical composition is administered intragastrically.

[Claim 21]

The method according to claim 20, wherein the intragastric administration is administration via a tube.

[Claim 22]

The method according to claim 21, wherein the tube is a percutaneous endoscopic gastrostomy tube (PEG tube).

[Claim 23]

The method according to claim 21, wherein the tube is a nasopharyngeal gastric tube (NG tube).

[Claim 24]

The method according to any one of claims 16 to 23, wherein the pharmaceutical composition is administered to treat an oxidative stress disease.

[Claim 25]

The method according to claim 24, wherein the oxidative stress disease is any one of diseases selected from the group consisting of amyotrophic lateral sclerosis, Parkinson's disease, spinocerebellar degeneration, muscular dystrophy, Alzheimer's disease, cerebral infarction, multiple sclerosis, systemic scleroderma, and stomatitis.

[Claim 26]

The method according to claim 24 or 25, wherein the oxidative stress disease is amyotrophic lateral sclerosis.

[Claim 27]

The method according to any one of claims 16 to 26, wherein administration of the pharmaceutical composition includes intermittent administration or daily administration; and the pharmaceutical composition is administered with the first time interval during an administration period of the intermittent administration or daily administration.

[Claim 28]

The method according to any one of claims 16 to 27, wherein a dose per administration of the pharmaceutical composition is from 90 to 120 mg of edaravone.

[Claim 29]

The method according to any one of claims 16 to 27, wherein a dose per administration of the pharmaceutical composition is from 90 to 105 mg of edaravone.

[Claim 30]

The method according to any one of claims 16 to 29, wherein a daily administration frequency of the pharmaceutical composition is once, and a dose per administration of the pharmaceutical composition is 105 mg of edaravone.