


Description


This invention relates to an optical isomer of an imidazole derivative.
Medetomidine which has the formula :




I

is known as a selective and potent α_2 -receptor agonist. It has been described, e.g. in European Patent Publication No. 72615, as an antihypertensive agent and in the European Patent Publication No. 187471 as a veterinary sedative-analgesic agent.

The optically active d- and l-enantiomers of medetomidine, and their non-toxic pharmaceutically acceptable acid addition salts are new compounds. These compounds may be represented by the formulae :



II



III

Racemic medetomidine is separated into the enantiomers II and III by conversion of the racemate into a mixture of diastereoisomers and separating the latter by fractional crystallization. Since medetomidine is a base, it may be converted into a diastereoisomer salt mixture by reaction with an optically active acid such as (+)-tartaric acid. Other useful optically active acids are, e.g., (-)-malic acid, (-)-mandelic acid and (+)-camphor-10-sulfonic acid. (+)-Tartaric acid is especially useful for the resolution. The separation of the diastereoisomers is performed by repeated crystallizing from an alcohol such as methanol or ethanol or a mixture of them.

Once the diastereoisomers have been separated the acid addition salts can be converted back to the free bases by making their aqueous solutions alkaline with sodium hydroxide and by extracting the liberated base in an appropriate organic solvent such as methylene chloride.

The d- and l-enantiomers of medetomidine react with organic and inorganic acids to form the corresponding acid addition salts, which have the same therapeutic activities as the bases. They can thus form many pharmaceutically usable acid addition salts, as, for instance, chlorides, bromides, sulfates, nitrates, phosphates, sulfonates, formates, tartrates, maleates, citrates, benzoates, salicylates or ascorbates.

The d- and l-enantiomers of medetomidine are selective and potent α_2 -receptor agonists.

Adrenergic receptors are physiologically important binding sites which are specific to noradrenaline and adrenaline and located on the surface of the cell membrane. The adrenoceptors of the sympathetic nervous system have been classified into two different subtypes, alpha- (α) and beta- (β) receptors, which can be further divided into two subgroups, viz α_1 and α_2 and β_1 and β_2 . Of these receptor types, β_1 , β_2 and α_1 are mainly located post-synaptically on the surface of, e.g., smooth muscle and thus mediate, e.g., smooth muscle contraction or relaxation ; whereas α_2 receptors are mainly located presynaptically on the terminals of noradrenergic nerves. If α_2 -receptors are stimulated by noradrenaline under physiological conditions noradrenaline release is blocked, i.e. there is a negative feed-back phenomenon. This negative feed-back phenomenon may also be induced by certain synthetic α_2 -agonists like medetomidine and some of its near derivatives.

In animal experiments, the d- and l- enantiomers of the present invention and especially the d-enantiomer, have proved to possess highly enhanced α_2 -selectivity and potency compared to the racemic mixture (i.e. medetomidine). The d-enantiomer can be expected to be of value, e.g., as a sedative-analgesic, anxiolytic or antihypertensive agent. Furthermore, it can be used as a pharmacological tool in the study of the physiology and pharmacology of α_2 -adrenoceptors.

The pharmacological activity of the compounds of the invention was determined as follows :

1. Alpha-2 agonism in vitro

α_2 -agonism was determined by means of isolated, electrically stimulated mouse vas deferens preparation (Marshall et al., Br. J. Pharmac. 62, 147-151, 1978). In this model, an α_2 -agonist is able to block electrically induced muscular contractions by activating the presynaptic α_2 -adrenoceptors and thus diminishing the secretion on the motor transmitter. Known α_2 -agonists like detomidine, medetomidine and clonidine were used as reference substances. Results are shown in Table 1, where the α_2 -agonist effect is presented as the pD_2 -value (negative logarithm of the molar concentration of the compound producing 50 percent of maximal inhibition.)

Table 1.

Compound	α_2 -agonism <u>in vitro</u> (mouse vas deferens). pD_2
d-enantiomer	9.3
l-enantiomer	6.0 (partial agonist)
medetomidine	9.0
detomidine	8.5
clonidine	8.5

These results show that the α_2 -agonist activity of medetomidine is limited to the d-enantiomer. The d-enantiomer shows an enhanced α_2 -agonist activity compared to the other agents studied.

2. α_2/α_1 -selectivity in vitro

The selectivity of the d-enantiomer as an α_2 -agonist was studied in receptor binding experiments using rat brain membranes. The ability of the d-isomer and the reference compounds to compete with 3H -clonidine (for α_2 -receptors) and 3H -prazosin (for α_1 -receptors) was studied essentially as described by Virtanen and Nyman in Eur. J. Pharmac. 108, 163-9, 1985. Results of the test are presented in Table 2, where the ability of the studied agents to compete with 3H -clonidine and 3H -prazosin binding is expressed as the IC_{50} -value (molar concentration of the competing ligand needed to displace 50 percent of the radioactive ligand).

Table 2.

Compound	³ H-clonidine displacement IC ₅₀ , nM	³ H-prazosin displacement IC ₅₀ , nM	α_2/α_1 - selectivity
d-enantiomer	1.2	55019	45849
l-enantiomer	46	189975	4129
medetomidine	3.3	16700	5060
detomidine	3.7	242	65
clonidine	6.4	6200	969

The results show that the d-enantiomer is an extremely selective α_2 -agonist compared to medetomidine and the other reference compounds.

3. Sedative analgesic effects

The sedative-analgesic properties of the compounds were studied in the spontaneous motility and writhing-test in the mouse. Spontaneous motility of mice and rats was measured using the Animex-activity meter. The test compounds were administered i.p. 30 minutes before the measuring periods of two minutes.

In the writhing test the compounds studied and saline were administered s.c. to rats, and 45 min. later 1 ml of 1% acetic acid was administered i.p. The number of writhes was recorded in the following 25 min. period (Koster et al., Fed. Proc. 18 : 412, 1959).

Results are shown in Tables 3 and 4.

Table 3: ED₅₀-values of the studied compounds in reducing spontaneous motility in mice

Compound	ED ₅₀ (mg/kg s.c.)
d-enantiomer	0.02
l-enantiomer	>10
medetomidine	0.05
detomidine	0.3
clonidine	0.3

Table 4: ED₅₀-values of the studied compounds in acetic acid-induced writhing test in mice

Compound	ED ₅₀ (mg/kg s.c.)
d-enantiomer	0.01
l-enantiomer	>10
medetomidine	0.02
detomidine	0.02
clonidine	0.03

These results show that the d-enantiomer has an enhanced sedative/analgesic property compared to the racemic mixture (medetomidine) and other reference compounds. The sedative/analgesic effects of medetomidine are confined to the d-enantiomer.

4. Anxiolytic effects

The anxiolytic effects of the compounds was studied using a method described by Handley & Mithoni : Naunyn-Schmiedeberg, Arch. Pharmacol. 327, 1-5, 1984. In this test the manner of exploration of open and enclosed arms in an elevated t-maze by a rat is examined. It has been shown that anxiolytic drugs increase the relative exploration of open arms. A rat is placed in the center of the t-maze and the number of open and enclosed entries is recorded during 5 minutes. Results obtained are shown in Table 5.

Table 5.

Drug/dose, mg/kg	Mean number of entries (n=6)			
	open	closed	total	open/total
NaCl	3.4	8.6	12.0	0.28
d-enantiomer				
0.0003	4.8	10.6	14.0	0.20
0.001	3.2	10.6	13.8	0.23
0.003	4.0	9.5	13.5	0.29
0.01	5.8	8.8	14.6	0.39
0.03	2.5	3.0	5.5	0.45
diazepam				
1	5.2	10.5	15.7	0.33

The results show that the d-enantiomer has an anxiolytic profile in the elevated t-maze test.

It is well known that anxiety states connected to withdrawal symptoms are due to noradrenergic hyperactivity. Therefore such symptoms can be successfully treated with drugs reducing the level of noradrenaline, e.g. clonidine. Experiments in the rat indicate that the d-enantiomer is able to reduce noradrenaline release and thus sympathetic tone both in the central and peripheral nervous systems. This has clearly been demonstrated by measuring CSF-concentrations of MHPG-SO₄ (the principal metabolite of central noradrenaline) in the rat after d-enantiomer administration. The results are shown in Table 6.

Table 6.

d-enantiomer dose μg/kg	CSF MHPG-SO ₄ (% of control) (4 h after d-enantiomer adm.)
0	100
3	-10
10	-20
30	-30
100	-65

5. Antihypertensive effects

The antihypertensive properties of the compounds of the invention have been studied as follows : Sprague-Dawley rats of normal weight were first anesthetized with urethane. After this, the femoral artery was connected by a polyethylene tube to a blood pressure transducer. The test substance was then injected into the femoral vein and the blood pressure and the pulse frequency were registered with a recorder. Results are shown in Table 7.

Table 7: Antihypertensive effects of the d-enantiomer in anesthetized rats

Dose, mg/kg	Decrease in BP, %	Decrease in heart rate, %
0.001	-8	-21
0.003	-23	-40
0.01	-43	-47
0.03	-45	-48
0.1	-45	-50

The results show that the d-enantiomer possesses clear anti-hypertensive and bradycardia effects.

The d-enantiomer, and its non-toxic, pharmaceutically acceptable acid addition salts or mixtures thereof may be administered parenterally, intravenously or orally. Typically, an effective amount of the compound is combined with a suitable pharmaceutical carrier. As used herein, the term "effective amount" encompasses those amounts which yield the desired activity without causing adverse side-effects. The precise amount employed in a particular situation is dependent upon numerous factors such as method of administration, type of mammal, condition for which the derivative is administered, and of course the structure of the derivative.

The pharmaceutical carriers which are typically employed with the compounds of the present invention may be solid or liquid and are generally selected with the planned manner of administration in mind. Thus, for example, solid carriers include lactose, sucrose, gelatin and agar, while liquid carriers include water, syrup, peanut oil and olive oil. Other suitable carriers are well known to those skilled in the art of pharmaceutical formulations. The combination of the derivative and the carrier may be fashioned into numerous acceptable forms, such as tablets, capsules, suppositories, solutions, emulsions, and powders.

The following Example illustrates the separation of the new enantiomers.

Example

14 g of medetomidine (base) were dissolved in 50 ml of methanol. 10.5 g of (+)-tartaric acid were dissolved in 50 ml of methanol. The solutions were mixed and the solvent was evaporated to a volume of 50 ml. The mixture was put into an ice bath and 9 g of white precipitate was obtained. The precipitate was suspended in 25 ml of ethanol, the mixture was kept in ultrasonic sound for 14 min and filtered. The precipitate was dissolved in a mixture of 20 ml abs. ethanol and 60 ml methanol by heating on a steam bath. After cooling, 5 g of precipitate (degree of rotation + 55°) was obtained. After recrystallization from 60 ml of methanol, 4.1 g of product was obtained, degree of rotation + 60°. Recrystallization was repeated until the degree of rotation did not increase any longer. The d-enantiomer tartrate was dissolved in water, the solution was made alkaline and the d-enantiomer was dissolved in an organic solvent e.g. dichloromethane or diethyl ether. The degree of rotation of the d-enantiomer base was + 75°.

The l-enantiomers may be isolated from the mother liquors.

Claims

Claims for the following Contracting States : AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE

1. The d-enantiomer of medetomidine and its non-toxic pharmaceutically acceptable acid addition salts.
2. The d-enantiomer of medetomidine for use in therapy as a selective α_2 -receptor agonist.
3. A pharmaceutical composition comprising the d-enantiomer of medetomidine or a non-toxic pharmaceutically acceptable acid addition salt thereof in association with a pharmaceutical carrier.

Table 7: Antihypertensive effects of the d-enantiomer in anesthetized rats

Dose, mg/kg	Decrease in BP, %	Decrease in heart rate, %
0.001	-8	-21
0.003	-23	-40
0.01	-43	-47
0.03	-45	-48
0.1	-45	-50

The results show that the d-enantiomer possesses clear anti-hypertensive and bradycardia effects.

The d-enantiomer, and its non-toxic, pharmaceutically acceptable acid addition salts or mixtures thereof may be administered parenterally, intravenously or orally. Typically, an effective amount of the compound is combined with a suitable pharmaceutical carrier. As used herein, the term "effective amount" encompasses those amounts which yield the desired activity without causing adverse side-effects. The precise amount employed in a particular situation is dependent upon numerous factors such as method of administration, type of mammal, condition for which the derivative is administered, and of course the structure of the derivative.

The pharmaceutical carriers which are typically employed with the compounds of the present invention may be solid or liquid and are generally selected with the planned manner of administration in mind. Thus, for example, solid carriers include lactose, sucrose, gelatin and agar, while liquid carriers include water, syrup, peanut oil and olive oil. Other suitable carriers are well known to those skilled in the art of pharmaceutical formulations. The combination of the derivative and the carrier may be fashioned into numerous acceptable forms, such as tablets, capsules, suppositories, solutions, emulsions, and powders.

The following Example illustrates the separation of the new enantiomers.

Example

14 g of medetomidine (base) were dissolved in 50 ml of methanol. 10.5 g of (+)-tartaric acid were dissolved in 50 ml of methanol. The solutions were mixed and the solvent was evaporated to a volume of 50 ml. The mixture was put into an ice bath and 9 g of white precipitate was obtained. The precipitate was suspended in 25 ml of ethanol, the mixture was kept in ultrasonic sound for 14 min and filtered. The precipitate was dissolved in a mixture of 20 ml abs. ethanol and 60 ml methanol by heating on a steam bath. After cooling, 5 g of precipitate (degree of rotation + 55°) was obtained. After recrystallization from 60 ml of methanol, 4.1 g of product was obtained, degree of rotation + 60°. Recrystallization was repeated until the degree of rotation did not increase any longer. The d-enantiomer tartrate was dissolved in water, the solution was made alkaline and the d-enantiomer was dissolved in an organic solvent e.g. dichloromethane or diethyl ether. The degree of rotation of the d-enantiomer base was + 75°.

The l-enantiomers may be isolated from the mother liquors.

Claims

Claims for the following Contracting States : AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE

1. The d-enantiomer of medetomidine and its non-toxic pharmaceutically acceptable acid addition salts.
2. The d-enantiomer of medetomidine for use in therapy as a selective α_2 -receptor agonist.
3. A pharmaceutical composition comprising the d-enantiomer of medetomidine or a non-toxic pharmaceutically acceptable acid addition salt thereof in association with a pharmaceutical carrier.



Claims for the following Contracting States : ES, GR

1. A method for separating the d-enantiomer of medetomidine which comprises converting racemic medetomidine into a diastereoisomer salt mixture by reaction with an optically active acid, and then separating the mixture of diastereoisomeric salts by fractional crystallization.
2. Method according to claim 1 in which the optically active acid used is (+)-tartaric acid.
3. A method of producing a pharmaceutical composition comprising admixing the d-enantiomer of medetomidine or a non-toxic pharmaceutically acceptable acid addition salt thereof with a pharmaceutical carrier.

Patentansprüche

Patentansprüche für folgende Vertragsstaaten : AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE

1. Das d-Enantiomer von Medetomidin und dessen nichttoxische pharmazeutisch annehmbare Säureadditionssalze.
2. Das d-Enantiomer von Medetomidin für die Verwendung bei der Therapie als ein selektiver α_2 -Rezeptor-Agonist.
3. Pharmazeutische Zusammensetzung umfassend das d-Enantiomer von Medetomidin oder ein nichttoxisches pharmazeutisches annehmbares Säureadditionssalz davon zusammen mit einem pharmazeutischen Träger.

Patentansprüche für folgende Vertragsstaaten : ES, GR

1. Verfahren zum Abtrennen des d-Enantiomers von Medetomidin, umfassend das Überführen von racemischem Medetomidin in ein diastereoisomere Salzmischung durch Umsetzung mit einer optischen aktiven Säure und dann das Trennen der Mischung der diastereoisomeren Salze durch fraktionierte Kristallisation.
2. Verfahren gemäß Anspruch 1, bei dem die verwendete optisch aktive Säure (+)-Weinsäure ist.
3. Verfahren zur Herstellung einer pharmazeutischen Zusammensetzung, umfassend das Abmischen des d-Enantiomers von Medetomidin oder eines nichttoxischen pharmazeutisch annehmbaren Säureadditionssalzes davon mit einem pharmazeutischen Träger.

Revendications

Revendications pour les Etats contractants suivants : AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE

1. Enantiomère d de la médétomidine et ses sels d'addition d'acide pharmaceutiquement acceptables et non toxiques.
2. Enantiomère d de la médétomidine, destiné à être employé dans des soins en tant qu'agoniste sélectif des récepteurs α_2 .
3. Composition pharmaceutique comprenant l'énantiomère d de la médétomidine ou un de ses sels d'addition d'acide pharmaceutiquement acceptables et non toxiques, associé avec un véhicule pharmaceutique.

Revendications pour les Etats contractants suivants : ES, GR

1. Procédé de séparation de l'énantiomère d de la médétomidine, qui comporte la transformation de la médétomidine sous forme de mélange racémique en un mélange de sels diastéréoisomères, par réaction avec un acide optiquement actif, et ensuite la séparation du mélange de sels diastéréoisomères par cristallisation fractionnée.
2. Procédé conforme à la revendication 1, dans lequel l'acide optiquement actif utilisé est l'acide (+)-tartrique.
3. Procédé de production d'une composition pharmaceutique, qui comprend le fait de mélanger de l'énantiomère d de la médétomidine, ou de l'un de ses sels d'addition d'acide pharmaceutiquement acceptables et non toxiques, avec un véhicule pharmaceutique.

